


Proportion of HIV/AIDS Cases and Population among Adolescents 13 to 19 Years of Age, by Race/Ethnicity Diagnosed in 2006—33 States




Note. Data include persons with a diagnosis of HIV infection regardless of their AIDS status at diagnosis. Data from 33 states with confidential name-based HIV infection reporting since at least 2003. Data have been adjusted for reporting delays.
*Includes 22 persons of unknown race or multiple races.




Proportion of HIV/AIDS Cases and Population among Young Adults 20 to 24 Years of Age, by Race/Ethnicity Diagnosed in 2006—33 States

HIV/AIDS cases
N=3,886*



Population, 33 states
N = 13,424,369



White, not Hispanic
Black, not Hispanic
Hispanic


Asian/Pacific Islander
American Indian/
Alaska Native



Note. Data include persons with a diagnosis of HIV infection regardless of their AIDS status at diagnosis. Data from 33 states with confidential name-based HIV infection reporting since at least 2003. Data have been adjusted for reporting delays.
*Includes 22 persons of unknown race or multiple races.



Proportion of HIV/AIDS Cases among Adults and Adolescents, by Sex and Age Group Diagnosed in 2006—33 States



Note. Data include persons with a diagnosis of HIV infection regardless of their AIDS status at diagnosis.
Data from 33 states with confidential name-based HIV infection reporting since at least 2003.
Data based on person's age at diagnosis.
Data have been adjusted for reporting delays.



HIV/AIDS Cases among Male Adolescents and Young Adults, by Transmission Category 2003–2006—33 States

Transmission	13–19 years		20–24 years	
	N	%	N	%
Male-to-male sexual contact	2,286	81	7,756	78
Injection drug use (IDU)	146	5	645	7
Male-to-male sexual contact and IDU	100	4	454	5
High-risk heterosexual contact*	277	10	1,013	10
Other/not identified [†]	9	<1	20	<1
Total	2,817	100	9,888	100



Note. Data include persons with a diagnosis of HIV infection regardless of AIDS status at diagnosis. Data from 33 states with confidential name-based HIV infection reporting since at least 2003. Data have been adjusted for reporting delays.

*Heterosexual contact with a person known to have, or to be at high risk for, HIV infection.

†Includes hemophilia, blood transfusion, perinatal exposure, and risk factor not reported or not identified.



HIV/AIDS Cases among Female Adolescents and Young Adults, by Transmission Category 2003–2006—33 States

Transmission Category	13–19 years		20–24 years	
	N	%	N	%
Injection drug use	233	13	597	14
High-risk heterosexual contact*	1,546	86	3,492	85
Other/not Identified†	15	<1	32	<1
Total	1,793	100	4,121	100

Note. Data include persons with a diagnosis of HIV infection regardless of AIDS status at diagnosis.
Data from 33 states with confidential name-based HIV infection reporting since at least 2003.


Data have been adjusted for reporting delays and cases without risk factor information were proportionally redistributed.

*Heterosexual contact with a person known to have, or to be at high risk for, HIV infection.

†Includes blood transfusion, perinatal exposure, and risk factor not reported or not identified.




HIV/AIDS Cases among Adolescents 13 to 19 Years of Age, 2006—33 States N=1,373



Note. Data include persons with a diagnosis of HIV infection regardless of their AIDS status at diagnosis. Data from 33 states with confidential name-based HIV infection reporting since at least 2003. Data have been adjusted for reporting delays. Data based on person's age at diagnosis.



Adolescents 13 to 19 Years of Age Living with HIV/AIDS, 2006—33 States N=5,678




Note. Data include persons with a diagnosis of HIV infection regardless of their AIDS status at diagnosis. Data from 33 states with confidential name-based HIV infection reporting since at least 2003. Data have been adjusted for reporting delays. Data based on age as of December 31, 2006.



Reported AIDS Cases among Adolescents 13 to 19 Years of Age, by Sex, 1985–2006—United States and Dependent Areas

N=6,642




Note. Data based on person's age at diagnosis.



Reported AIDS Cases among Young Adults 20 to 24 Years of Age, by Sex, 1985–2006—United States and Dependent Areas



N=36,293



Note. Data based on person's age at diagnosis.



Proportion of AIDS Cases and Population among Adolescents 13 to 19 Years of Age, by Race/Ethnicity Reported in 2006—50 States and DC



White, not Hispanic
 Black, not Hispanic
 Hispanic



Asian/Pacific Islander
 American Indian/
Alaska Native



Note. Data based on person's age at AIDS diagnosis.
*Includes 6 persons of unknown race or multiple races.



Proportion of AIDS Cases and Population among Young Adults 20 to 24 Years of Age, by Race/Ethnicity Reported in 2006—50 States and DC




- White, not Hispanic
- Black, not Hispanic
- Hispanic
- Asian/Pacific Islander
- American Indian/
Alaska Native

Note. Data based on person's age at AIDS diagnosis.
* Includes 18 persons of unknown race or multiple races.



Proportion of AIDS Cases among Adults and Adolescents, by Sex and Age Group, Reported in 2006—United States and Dependent Areas



Note. Data based on person's age at diagnosis.



AIDS Cases among Male Adolescents and Young Adults, by Transmission Category 2002–2006—United States and Dependent Areas

Transmission category	13–19 years		20–24 years	
	N	%	N	%
Male-to-male sexual contact	736	60	3,983	72
Injection drug use (IDU)	86	7	480	9
Male-to-male sexual contact and IDU	38	3	325	6
High-risk heterosexual*	147	12	706	13
Other/not identified [†]	225	18	58	1
Total	1,231	100	5,552	100

Note. Data have been adjusted for reporting delays and cases without risk factor information were proportionally redistributed.

*Heterosexual contact with a person known to have, or to be at high risk for, HIV infection.

[†] Includes hemophilia, blood transfusion, perinatal exposure, and risk factor not reported or not identified.



AIDS Cases among Female Adolescents and Young Adults, by Transmission Category 2002–2006—United States and Dependent Areas

Transmission category	13–19 years		20–24 years	
	N	%	N	%
Injection drug	108	11	416	16
High-risk heterosexual contact*	549	59	2,072	81
Other/not identified †	279	30	62	2
Total	936	100	2,550	100



Note. Data have been adjusted for reporting delays and cases without risk factor information were proportionally redistributed.


*Heterosexual contact with a person known to have, or to be at high risk for, HIV infection.

†Includes blood transfusion, perinatal exposure, and risk factor not reported or not identified.



Reported AIDS Cases among Adolescents 13 to 19 Years of Age, 2006—United States and Dependent Areas


N=581



Note. Data based on person's age at diagnosis.



Adolescents 13 to 19 Years of Age Reported to be Living with AIDS, 2006—United States and Dependent Areas N=3,116*



Note. Data based on person's age as of December 31, 2006.
*Includes 13 persons whose state of residence is unknown or missing.

