



Characteristic Coastal Habitats Choosing Spill Response Alternatives



National Oceanic and Atmospheric Administration

National Ocean Service

Office of Response and Restoration

Hazardous Materials Response Division



National Oceanic and Atmospheric Administration • **National Ocean Service**
Office of Response and Restoration • **Hazardous Materials Response Division**
November 2000

Characteristic Coastal Habitats is based upon information contained in *Environmental Considerations for Marine Oil Spill Response*, published by the American Petroleum Institute, the National Oceanic and Atmospheric Administration, the U.S. Coast Guard, and the U.S. Environmental Protection Agency.

CONTENTS

INTRODUCTION	4
HOW TO USE THIS DOCUMENT	5
Absence of Oil for Intertidal and Ice Environments	6
Absence of Oil for On-Water and Subtidal Environments	10

HABITAT

INTERTIDAL

Exposed Rocky Shores	14
Exposed, Solid Man-made Structures	17
Exposed, Wave-cut Platforms	20
Sand Beaches	23
Tundra Cliffs	26
Mixed Sand and Gravel Beaches	29
Gravel Beaches	32
Riprap	35
Exposed Tidal Flats	38
Sheltered Rocky Shores and Scarps	41
Sheltered, Solid Man-Made Structures	45
Peat Shores	48
Sheltered Tidal Flats	51
Salt to Brackish Marshes	54
Mangroves	57
Inundated Lowland Tundra	60

SUBTIDAL

Coral Reefs	63
Seagrasses	66
Kelp	69
Soft Bottom	72
Mixed and Hard Bottom	75

ICE

Accessible and Inaccessible	78
-----------------------------	----

ON-WATER

Offshore	82
Bays and Estuaries	85

Introduction

Oil on coastal waters, shorelines, or subtidal habitats can harm the environment, intrude on recreational activities, cause economic hardship, disrupt commercial activities, and be expensive to clean up. Decisions about if, where, when, and how to remove oil from coastal habitats affect each of these potential spill consequences.

Sound cleanup decisions depend on accurate information about the types of habitats that the oil affects, the degree of oiling, and the location of oiling. *Characteristic Coastal Habitats* illustrates typical physical and biological attributes of North American coastal habitats at risk from oil spills. The text describes each habitat and discusses both how oil is likely to behave there and considerations for treating oil.

The *Characteristic Coastal Habitats* collection is designed as a companion to *Environmental Considerations for Marine Oil Spill Response*, published by the American Petroleum Institute, the National Oceanic and Atmospheric Administration, the U.S. Coast Guard, and the U.S. Environmental Protection Agency. The Response Method table for each habitat is based on information contained in *Environmental Considerations for Marine Oil Spill Response*. Refer to that publication for complete information on proper use and caveats regarding the guidelines presented in the Response Method tables.

The *Characteristic Coastal Habitats* collection is a useful aid for training people who will be participating in cleanup assessment as part of an Environmental Unit within the Incident Command System. It also complements NOAA's *Shoreline Assessment Manual* and *Shoreline Assessment Job Aid*. Visit the Office of Response and Restoration's home page at <http://response.restoration.noaa.gov> for copies of the manual and job aid. *Characteristic Coastal Habitats* is available in a digital version at this website.

How to use this document

This document summarizes the technical rationale for selecting response methods for five categories of oil in specific habitats. As a companion to *Environmental Considerations for Marine Oil Spill Response*, *Characteristic Coastal Habitats* can help you select appropriate response options to minimize the adverse environmental impacts of a marine oil spill. The guide discusses on-water, shallow subtidal, shoreline intertidal, and ice habitats. Specific response options include natural recovery; mechanical, chemical, and biological treatment; and in-situ burning.

When choosing effective response options, including natural recovery, you must consider trade-offs affecting the option's potential environmental impact, its appropriateness for the habitat, and timing of its application. *Environmental Considerations for Marine Oil Spill Response* discusses these considerations in detail; consult it before using this guideline. Remember that the benefits and impacts of response options depend upon incident-specific conditions and affect the options' suitability for use in a habitat during any spill. For example, dovetailing multiple methods simultaneously throughout an incident might produce a more effective response and fewer adverse environmental impacts.

Each section of this guide includes information about methods currently in use during oil spill responses in marine environments. The two tables following this section present information on the relative environmental impact of methods in the *absence* of oil for each habitat. These tables will help you understand the impact of a response option independent of oil effects. Following this section is the Habitat section with descriptions of intertidal, shallow subtidal, ice, and on-water habitats. For each category there is a description of habitat type and a table describing the relative environmental impact of the response method and oil in that habitat, for five different oil categories. These tables will help you understand the impact of the response option *and* oil together for each habitat.

Shoreline Intertidal and Ice Environments (1 of 2)

The following categories compare the relative environmental impact of each response method in a specific environment and habitat. The codes in each table mean:

- A = The least adverse habitat impact.
- B = Some adverse habitat impact.
- C = Significant adverse habitat impact.
- D = The most adverse habitat impact.
- I = Insufficient information - impact or effectiveness of the method could not be evaluated.
- = Not applicable.

Response Method

	Exposed Rocky Shores	Exposed Solid Man-made Structures	Exposed Wave-cut Platforms	Sand Beaches*	Tundra Cliffs	Mixed Sand and Gravel Beaches*	Gravel Beaches	Riprap	Exposed Tidal Flats
Natural Recovery	A	A	A	A	A	A	A	A	A
Booming	-	-	-	-	-	-	-	-	-
Skimming	-	-	-	-	-	-	-	-	-
Barriers/Berms	-	-	-	B	D	B	C	-	C
Physical Herding	-	-	-	-	-	-	-	-	B
Manual Oil Removal/Cleaning	B	B	B	B	D	B	B	A	C
Mechanical Oil Removal	-	-	-	C	D	C	C	C	D
Sorbents	B	B	B	B	C	B	B	A	B
Vacuum	B	-	B	B	D	B	B	A	B
Debris Removal	B	-	B	B	A	B	B	A	B
Sediment Reworking/Tilling	-	-	-	C	D	C	C	-	D
Vegetation Cutting/Removal	-	-	-	C	D	C	-	-	D
Flooding (deluge)	-	-	B	B	C	B	B	B	B

This table provides information on the relative environmental impacts of response methods in the absence of oil in shoreline intertidal and ice environments.

Consult the *Environmental Considerations for Marine Oil Spill Response* document referenced on page 5 before using this table.

* - Special biological need consideration - if birds and turtles are nesting, the ranking would be "D."

Sheltered Rocky Shores & Scarps	Sheltered Solid Man-made Structures	Peat Shores	Sheltered Tidal Flats	Marshes	Mangroves	Inundated Lowland Tundra	Accessible Ice	Inaccessible Ice
A	A	A	A	A	A	A	A	A
-	-	-	-	-	-	-	B	-
-	-	-	-	-	-	-	B	B
-	-	-	C	C	C	-	B	-
-	-	-	-	-	-	-	B	B
B	B	B	D	D	C	D	B	-
-	-	D	-	D	-	D	B	B
B	A	B	B	C	C	C	B	B
B	-	B	B	C	C	D	B	B
B	A	B	C	C	C	D	B	B
-	-	B	-	D	-	-	-	-
D	-	C	D	D	-	D	-	-
B	-	B	B	B	B	C	-	-

Shoreline Intertidal and Ice Environments (2 of 2)

The following categories are used to compare the relative environmental impact of each response method in the specific environment and habitat. The codes in each table mean:

- A = The least adverse habitat impact.
- B = Some adverse habitat impact.
- C = Significant adverse habitat impact.
- D = The most adverse habitat impact.
- I = Insufficient information - impact or effectiveness of the method could not be evaluated.
- = Not applicable.

Response Method (cont.)

	Exposed Rocky Shores	Exposed Solid Man-made Structures	Exposed Wave-cut Platforms	Sand Beaches *	Tundra Cliffs	Mixed Sand and Gravel Beaches *	Gravel Beaches	Riprap	Exposed Tidal Flats
Low-pressure, Ambient Water Flushing	B	B	B	B	D	B	B	B	C
High-pressure, Ambient Water Flushing	C	C	C	—	—	C	C	C	—
Low-pressure, Hot Water Flushing	D	D	D	D	—	D	D	D	—
High-pressure, Hot Water Flushing	D	D	D	—	—	D	D	D	—
Steam Cleaning	D	D	D	—	—	D	D	D	—
Sand Blasting	D	D	D	—	—	—	—	D	—
Dispersants	—	—	—	—	—	—	—	—	—
Emulsion-treating Agents	—	—	—	—	—	—	—	—	—
Elasticity Modifiers	—	—	—	—	—	—	—	—	—
Herdng Agents	—	—	—	—	—	—	—	—	—
Solidifiers	—	—	B	B	C	B	B	B	B
Shoreline Cleaning Agents	B	B	B	—	—	C	B	B	—
Nutrient Enrichment	—	—	—	B	C	B	B	B	I
Natural Microbe Seeding	—	—	—	I	I	I	I	I	I
In-situ Burning	—	—	D	C	—	C	C	D	—

This table provides information on the relative environmental impacts of response methods in the absence of oil in shoreline intertidal and ice environments.

Consult the *Environmental Considerations for Marine Oil Spill Response* document referenced on page 5 before using this table.

* - Special biological need consideration - if birds and turtles are nesting, the ranking would be "D."

	Sheltered Rocky Shores & Scarps	Sheltered Solid Man-made Structures	Peat Shores	Sheltered Tidal Flats	Marshes	Mangroves	Inundated Lowland Tundra	Accessible Ice	Inaccessible Ice
	B	B	B	C	B	B	D	B	B
	C	C	-	-	-	-	-	-	-
	D	D	-	-	-	-	-	B	B
	D	D	-	-	-	-	-	-	-
	D	D	-	-	-	-	-	B	B
	D	D	-	-	-	-	-	-	-
	-	-	-	-	-	-	-	B	B
	-	-	-	-	-	-	-	B	B
	-	-	-	-	-	-	-	B	B
	-	-	-	-	-	-	-	B	-
	B	-	-	B	C	C	C	B	B
	B	B	-	-	B	I	-	-	-
	B	I	B	I	B	I	I	I	I
	I	I	I	I	I	I	I	I	I
	D	-	-	-	C	-	D	B	B

On-water and Shallow Subtidal Environments

The following categories are used to compare the relative environmental impact of each response method in the specific environment and habitat. The codes in this table mean:

- A = The least adverse habitat impact.
- B = Some adverse habitat impact.
- C = Significant adverse habitat impact.
- D = The most adverse habitat impact.
- I = Insufficient information - impact or effectiveness of the method could not be evaluated.
- = Not applicable.

Response Method	Offshore	Bays and Estuaries	Coral Reefs	Seagrasses	Kelp	Soft Bottom	Mixed and Hard Bottom
Natural Recovery	A	A	A	A	A	A	A
Booming	B	B	C	B	B	A	A
Skimming	B	B	B	B	B	A	A
Barriers/Berms	-	-	-	-	-	-	-
Physical Herding	B	B	B	B	B	B	B
Manual Oil Removal/Cleaning	-	B	D	C	B	B	C
Mechanical Oil Removal	-	-	D	D	D	C	-
Sorbents	B	B	D	B	B	B	C
Vacuum	-	-	C	C	B	B	C
Debris Removal	B	B	B	B	B	B	B
Sediment Reworking/Tilling	-	-	-	-	-	-	-
Vegetation Cutting/Removal	-	-	-	B	C	-	-
Flooding (deluge)	-	-	-	-	-	-	-

This table provides information on the relative environmental impacts of response methods in the absence of oil in on-water and shallow subtidal environments.

Consult the *Environmental Considerations for Marine Oil Spill Response* document referenced on page 5 before using this table.


* - Special biological need consideration - if birds and turtles are nesting, the ranking would be "D."

Response Method (cont.)	Offshore	Bays and Estuaries	Coral Reefs	Seagrasses	Kelp	Soft Bottom	Mixed and Hard Bottom
Low-pressure, Ambient Water Flushing	-	-	B	-	-	-	-
High-pressure, Ambient Water Flushing	-	-	-	-	-	-	-
Low-pressure, Hot Water Flushing	-	-	-	-	-	-	-
High-pressure, Hot Water Flushing	-	-	-	-	-	-	-
Steam Cleaning	-	-	-	-	-	-	-
Sand Blasting	-	-	-	-	-	-	-
Dispersants	B	B	D	D	D	D	C
Emulsion-treating Agents	B	B	I	I	I	I	I
Elasticity Modifiers	B	B	I	I	I	I	I
Herding Agents	B	B	I	I	I	I	I
Solidifiers	B	B	I	I	I	I	I
Shoreline Cleaning Agents	-	-	-	-	-	-	-
Nutrient Enrichment	-	-	-	-	-	-	-
Natural Microbe Seeding	-	-	-	-	-	-	-
In-situ Burning	B	B	B	B	B	B	B

HABITAT

Exposed Rocky Shores

INTERTIDAL



INTERTIDAL

Exposed Rocky Shores

Description

- The intertidal zone is steep (>30° slope) and narrow with very little width.
- Sediment accumulations are uncommon because waves remove debris that has slumped from the eroding cliffs.
- There is strong vertical zonation of intertidal biological communities.
- Species density and diversity vary greatly, but barnacles, snails, mussels, polychaetes, and macroalgae can be abundant.

Predicted Oil Behavior

- Oil is held offshore by waves reflecting off the steep, hard surfaces.
- Any oil that is deposited is rapidly removed from exposed faces.
- The most resistant oil would remain as a patchy band at or above the high-tide line.
- Impacts to intertidal communities are expected to be short-term. An exception would be where heavy concentrations of a light refined product came ashore very quickly.

Response Considerations

- Cleanup is usually not required.
- Access can be difficult and dangerous.

Exposed Rocky Shores

INTERTIDAL

Oil Category Descriptions


- I - Gasoline products
- II - Diesel-like products and light crudes
- III - Medium grade crudes and intermediate products
- IV - Heavy crudes and residual products
- V - Non-floating oil products

The following categories are used to compare the relative environmental impact of each response method in the specific environment and habitat for each oil type. The codes in each table mean:

- A = The least adverse habitat impact.
- B = Some adverse habitat impact.
- C = Significant adverse habitat impact.
- D = The most adverse habitat impact.
- I = Insufficient information - impact or effectiveness of the method could not be evaluated.
- = Not applicable.

Response Method	Oil Category				
	I	II	III	IV	V
Natural Recovery	A	A	A	A	A
Barriers/Berms	-	-	-	-	-
Manual Oil Removal/Cleaning	-	-	B	B	B
Mechanical Oil Removal	-	-	-	-	-
Sorbents	-	B	A	A	A
Vacuum	-	A	A	A	A
Debris Removal	-	A	A	A	A
Sediment Reworking/Tilling	-	-	-	-	-
Vegetation Cutting/Removal	-	-	-	-	-
Flooding (deluge)	-	-	-	-	-
Low-pressure, Ambient Water Flushing	-	A	A	B	B
High-pressure, Ambient Water Flushing	-	B	B	B	B
Low-pressure, Hot Water Flushing	-	-	C	C	C
High-pressure, Hot Water Flushing	-	-	C	C	C
Steam Cleaning	-	-	D	D	D
Sand Blasting	-	-	D	D	D
Solidifiers	-	-	-	-	-
Shoreline Cleaning Agents	-	-	C	C	C
Nutrient Enrichment	-	-	-	-	-
Natural Microbe Seeding	-	-	-	-	-
In-situ Burning	-	-	-	-	-

Consult the *Environmental Considerations for Marine Oil Spill Response* document referenced on page 5 before using this table.



Description

- These are solid, man-made structures such as seawalls, groins, revetments, piers, and port facilities.
- Many structures are constructed of concrete, wood, or metal.
- They are built to protect the shore from erosion by waves, boat wakes, and currents, and thus are exposed to rapid natural removal processes.
- Often there is no exposed substrate at low tide, but multiple habitats may be present.
- Attached animals and plants are sparse to moderate.

Predicted Oil Behavior

- Oil is held offshore by waves reflecting off the steep, hard surfaces in exposed settings.
- Oil readily adheres to the dry, rough surfaces, but it does not adhere to wet substrates.
- The most resistant oil would remain as a patchy band at or above the high-tide line.

Response Considerations

- Cleanup is usually not required.
- High-pressure water spraying may be conducted to remove risks of contamination of people or vessels or to improve aesthetics.

INTERTIDAL


Exposed, Solid Man-made Structures

Response Method	Oil Category				
	I	II	III	IV	V
Oil Category Descriptions					
I - Gasoline products					
II - Diesel-like products and light crudes					
III - Medium grade crudes and intermediate products					
IV - Heavy crudes and residual products					
V - Non-floating oil products					
The following categories are used to compare the relative environmental impact of each response method in the specific environment and habitat for each oil type. The codes in this table mean:					
A = The least adverse habitat impact.					
B = Some adverse habitat impact.					
C = Significant adverse habitat impact.					
D = The most adverse habitat impact.					
I = Insufficient information - impact or effectiveness of the method could not be evaluated.					
-- = Not applicable.					
Natural Recovery	A	A	A	A	A
Barriers/Berms	-	-	-	-	-
Manual Oil Removal/Cleaning	-	-	B	B	B
Mechanical Oil Removal	-	-	-	-	-
Sorbents	-	B	A	A	A
Vacuum	-	-	-	-	-
Debris Removal	-	-	-	-	-
Sediment Reworking/Tilling	-	-	-	-	-
Vegetation Cutting/Removal	-	-	-	-	-
Flooding (deluge)	-	-	-	-	-
Low-pressure, Ambient Water Flushing	-	A	A	B	B
High-pressure, Ambient Water Flushing	-	B	B	B	B
Low-pressure, Hot Water Flushing	-	-	C	C	C
High-pressure, Hot Water Flushing	-	-	C	C	C
Steam Cleaning	-	-	D	D	D
Sand Blasting	-	-	D	D	D
Solidifiers	-	-	-	-	-
Shoreline Cleaning Agents	-	-	B	B	B
Nutrient Enrichment	-	-	-	-	-
Natural Microbe Seeding	-	-	-	-	-
In-situ Burning	-	-	-	-	-

Consult the *Environmental Considerations for Marine Oil Spill Response* document referenced on page 5 before using this table.

Exposed, Wave-cut Platforms

INTERTIDAL



Description

- These shores consist of a bedrock shelf or platform of variable width and very gentle slope.
- The surface of the platform is irregular; tide pools are common.
- Along headlands, they have only small accumulation of sediments, mostly at the high-tide line.
- They often co-occur with gravel beaches; the gravel beach can be either at the upper or the lower half of the intertidal zone, depending on the nature of the bedrock outcrop.
- Species density and diversity vary greatly, but barnacles, snails, mussels, and macroalgae are often abundant.

Predicted Oil Behavior

- Oil will not adhere to the wet rock surface, but could penetrate crevices or sediment veneers.
- Oil persistence is usually short-term, except in wave shadows or where the oil was deposited high above normal wave activity.

Response Considerations

- Cleanup is usually not required.
- Where the high-tide area is accessible, it may be feasible to manually remove heavy oil accumulations and oiled debris.

Exposed, Wave-cut Platforms


INTERTIDAL

Response Method	Oil Category				
	I	II	III	IV	V
Oil Category Descriptions					
I - Gasoline products					
II - Diesel-like products and light crudes					
III - Medium grade crudes and intermediate products					
IV - Heavy crudes and residual products					
V - Non-floating oil products					
Natural Recovery	A	A	A	A	A
Barriers/Berms	-	-	-	-	-
Manual Oil Removal/Cleaning	-	B	B	B	B
Mechanical Oil Removal	-	-	-	-	-
Sorbents	-	B	A	A	A
Vacuum	-	A	A	A	A
Debris Removal	-	A	A	A	A
Sediment Reworking/Tilling	-	-	-	-	-
Vegetation Cutting/Removal	-	-	-	-	-
Flooding (deluge)	-	A	A	B	B
Low-pressure, Ambient Water Flushing	-	A	A	B	B
High-pressure, Ambient Water Flushing	-	B	B	B	B
Low-pressure, Hot Water Flushing	-	D	C	C	C
High-pressure, Hot Water Flushing	-	D	C	C	C
Steam Cleaning	-	-	D	D	D
Sand Blasting	-	-	D	D	D
Solidifiers	-	C	C	-	-
Shoreline Cleaning Agents	-	-	C	C	C
Nutrient Enrichment	-	-	-	-	-
Natural Microbe Seeding	-	I	I	I	I
In-situ Burning	-	D	D	D	-

The following categories are used to compare the relative environmental impact of each response method in the specific environment and habitat for each oil type. The codes in each table mean:

- A = The least adverse habitat impact.
- B = Some adverse habitat impact.
- C = Significant adverse habitat impact.
- D = The most adverse habitat impact.
- I = Insufficient information - impact or effectiveness of the method could not be evaluated.
- = Not applicable.

Consult the *Environmental Considerations for Marine Oil Spill Response* document referenced on page 5 before using this table.



Description

- These beaches are flat to moderately sloping and relatively hard-packed.
- There can be heavy accumulations of wrack.
- They are used by birds and turtles.
- Upper beach fauna include ghost crabs and amphipods; lower beach fauna can be moderate, but highly variable.

Predicted Oil Behavior

- Light oil accumulations will be deposited as oily swashes or bands along the upper intertidal zone.
- Heavy oil accumulations will cover the entire beach surface; oil will be lifted off the lower beach with the rising tide.
- Maximum penetration of oil into fine- to medium-grained sand is about 10-15 cm, up to 25 cm in coarse-grained sand. Maximum penetration of oil into fine to medium-grained sand beaches is about 10-15 cm, and about 25 cm into coarse-grained sand beaches.
- Burial of oiled layers by clean sand can be rapid (within one day), and burial to depths as much as one meter is possible if the oil comes ashore at the beginning of a depositional period.
- Organisms living in the beach sediment may be killed by smothering or lethal oil concentrations in the interstitial water.
- Biological impacts include temporary declines in infauna, which can affect important shorebird foraging areas.

Response Considerations

- These beaches are among the easiest shoreline types to clean.
- Cleanup should concentrate on removing oil and oily debris from the upper swash zone once most of the oil has come ashore.
- Manual cleanup, rather than road graders and front-end loaders, is advised to minimize volume of sand removed from the shore and requiring disposal.
- All efforts should focus on preventing vehicular and foot traffic from mixing oil deeper into the sediments.
- Mechanical reworking of lightly oiled sediments from the high-tide line to the upper intertidal zone can be effective along exposed beaches.

INTERTIDAL


Sand Beaches

Response Method	Oil Category				
	I	II	III	IV	V
Oil Category Descriptions					
I – Gasoline products					
II – Diesel-like products and light crudes					
III – Medium grade crudes and intermediate products					
IV – Heavy crudes and residual products					
V – Non-floating oil products					
The following categories are used to compare the relative environmental impact of each response method in the specific environment and habitat for each oil type. The codes in each table mean:					
A = The least adverse habitat impact.					
B = Some adverse habitat impact.					
C = Significant adverse habitat impact.					
D = The most adverse habitat impact.					
I = Insufficient information - impact or effectiveness of the method could not be evaluated.					
— = Not applicable.					
Natural Recovery	A	B	B	C	D
Barriers/Berms	B	B	B	B	B
Manual Oil Removal/Cleaning	D	B	A	A	A
Mechanical Oil Removal	D	B	B	B	B
Sorbents	–	B	A	A	B
Vacuum	–	–	B	A	A
Debris Removal	–	A	A	A	A
Sediment Reworking/Tilling	D	B	B	B	B
Vegetation Cutting/Removal	–	C	C	C	C
Flooding (deluge)	A	A	A	B	C
Low-pressure, Ambient Water Flushing	B	B	B	B	C
High-pressure, Ambient Water Flushing	–	–	–	–	–
Low-pressure, Hot Water Flushing	–	–	C	C	C
High-pressure, Hot Water Flushing	–	–	–	–	–
Steam Cleaning	–	–	–	–	–
Sand Blasting	–	–	–	–	–
Solidifiers	–	–	B	–	–
Shoreline Cleaning Agents	–	–	C	C	C
Nutrient Enrichment	–	A	A	B	C
Natural Microbe Seeding	–	I	I	I	I
In-situ Burning	–	–	C	C	C

Consult the *Environmental Considerations for Marine Oil Spill Response* document referenced on page 5 before using this table.

Tundra Cliffs

INTERTIDAL



INTERTIDAL

Description

- These are erosional features with tundra vegetation overlying peat and exposed ground ice or permafrost.
- Cliff heights range from less than 1 meter to as much as 5-10 meters.
- There may be a narrow beach present or just a vertical scarp.
- As the cliffs erode at rates of 0.5-4 meters/year, the vegetation and peat accumulate as fragmented and irregular blocks at the base of the cliff until they are reworked by waves.
- The vegetation on the tundra is a living plant community that is sensitive to disturbances.
- Large numbers of migratory birds can use these shorelines during the summer months.

Predicted Oil Behavior

- Oil could be stranded onshore only during the ice-free summer season.
- Oil is not likely to adhere to exposed ground ice, unless air temperatures are below freezing.
- Oil persistence on the vegetation and peat substrates would be short in most cases, due to natural cliff erosion, provided that the oil is not stranded at the onset of freeze-up.
- If the oil mixes with the peaty substrate or accumulated peat, it could create sheens until the oiled area erodes.
- Biological risks would be greatest to birds feeding along oiled cliffs in summer.

Response Considerations

- Natural peat can be used as a sorbent as long as it is taken from beach peat deposits and not the living tundra.
- Manual or mechanical removal of oil or oiled tundra/peat may be the most practical method if oil removal is required, though the peat substrate is soft and readily trampled.
- Hot-water washing or even low-pressure flushing is not appropriate because they may accelerate thermal and mechanical erosion of the ice in the cliff, triggering unexpected block falls, slumping, or mud flows.
- The cliffs are commonly undercut and naturally unstable, so worker safety is a primary concern.
- * Cleanup occurs only in the short arctic summer, a very limited window of intense ecological activity.

Response Method	Oil Category				
	I	II	III	IV	V
Natural Recovery	A	B	B	B	B
Barriers/Berms	B	B	B	B	B
Manual Oil Removal/Cleaning	D	B	B	B	B
Mechanical Oil Removal	C	C	C	C	C
Sorbents	-	B	A	A	B
Vacuum	-	-	B	A	A
Debris Removal	-	B	B	B	B
Sediment Reworking/Tilling	D	B	B	B	C
Vegetation Cutting/Removal	D	D	D	D	D
Flooding (deluge)	A	A	A	B	C
Low-pressure, Ambient Water Flushing	C	B	B	B	C
High-pressure, Ambient Water Flushing	-	-	-	-	-
Low-pressure, Hot Water Flushing	-	-	-	-	-
High-pressure, Hot Water Flushing	-	-	-	-	-
Steam Cleaning	-	-	-	-	-
Sand Blasting	-	-	-	-	-
Solidifiers	-	-	B	-	-
Shoreline Cleaning Agents	-	-	-	-	-
Nutrient Enrichment	-	B	B	C	C
Natural Microbe Seeding	-	I	I	I	I
In-situ Burning	-	-	-	-	-

Consult the *Environmental Considerations for Marine Oil Spill Response* document referenced on page 5 before using this table.

Oil Category Descriptions

- I - Gasoline products
- II - Diesel-like products and light crudes
- III - Medium grade crudes and intermediate products
- IV - Heavy crudes and residual products
- V - Non-floating oil products

The following categories are used to compare the relative environmental impact of each response method in the specific environment and habitat for each oil type. The codes in each table mean:

- A = The least adverse habitat impact.
- B = Some adverse habitat impact.
- C = Significant adverse habitat impact.
- D = The most adverse habitat impact.
- I = Insufficient information - impact or effectiveness of the method could not be evaluated.
- = Not applicable.

Description

- Because of the mixed sediment sizes on these moderately sloping beaches, there may be zones of pure sand, pebbles, or cobbles.
- There can be large-scale changes in the sediment distribution patterns depending upon season, because of the transport of the sand fraction offshore during storms.
- Desiccation and sediment mobility on exposed beaches cause low densities of attached animals and plants.
- The presence of attached algae, mussels, and barnacles indicates beaches that are relatively sheltered, with the more stable substrate supporting a richer biota.

Predicted Oil Behavior

- During small spills, oil will be deposited along and above the high-tide swash.
- Large spills will spread across the entire intertidal area.
- Oil penetration into the beach sediments may be up to 50 cm; however, the sand fraction can be quite mobile, and oil behavior is much like on a sand beach if the sand fraction exceeds about 40 percent.
- Burial of oil may be deep at and above the high-tide line, where oil tends to persist, particularly where beaches are only intermittently exposed to waves.
- In sheltered pockets on the beach, pavements of asphalted sediments can form if oil accumulations are not removed, because most of the oil remains on the surface.

Response Considerations

- Remove heavy accumulations of pooled oil from the upper beachface.
- All oiled debris should be removed; sediment removal should be limited as much as possible.
- Low-pressure flushing can be used to float oil away from the sediments for recovery by skimmers or sorbents. High-pressure spraying should be avoided because of potential for transporting contaminated finer sediments (sand) to the lower intertidal or subtidal zones.
- Mechanical reworking of oiled sediments from the high-tide zone to the upper intertidal zone can be effective in areas regularly exposed to wave activity. However, oiled sediments should not be relocated below the mid-tide zone.
- In-place tilling may be used to reach deeply buried oil layers in the mid-tide zone on exposed beaches.

INTERTIDAL

Mixed Sand and Gravel Beaches

Response Method	Oil Category				
	I	II	III	IV	V
Natural Recovery	A	B	B	C	C
Barriers/Berms	C	C	C	B	B
Manual Oil Removal/Cleaning	D	C	B	A	A
Mechanical Oil Removal	D	C	B	B	B
Sorbents	–	A	A	B	B
Vacuum	–	–	B	B	B
Debris Removal	–	A	A	A	A
Sediment Reworking/Tilling	D	B	B	B	B
Vegetation Cutting/Removal	–	C	C	C	C
Flooding (deluge)	A	A	B	C	C
Low-pressure, Ambient Water Flushing	B	A	A	B	C
High-pressure, Ambient Water Flushing	–	–	C	D	D
Low-pressure, Hot Water Flushing	–	–	C	C	C
High-pressure, Hot Water Flushing	–	–	D	D	D
Steam Cleaning	–	–	D	D	D
Sand Blasting	–	–	–	–	–
Solidifiers	–	–	B	–	–
Shoreline Cleaning Agents	–	–	C	C	C
Nutrient Enrichment	–	A	A	B	C
Natural Microbe Seeding	–	I	I	I	I
In-situ Burning	–	–	C	C	C

Oil Category Descriptions

- I – Gasoline products
- II – Diesel-like products and light crudes
- III – Medium grade crudes and intermediate products
- IV – Heavy crudes and residual products
- V – Non-floating oil products

The following categories are used to compare the relative environmental impact of each response method in the specific environment and habitat for each oil type. The codes in each table mean:

- A = The least adverse habitat impact.
- B = Some adverse habitat impact.
- C = Significant adverse habitat impact.
- D = The most adverse habitat impact.
- I = Insufficient information - impact or effectiveness of the method could not be evaluated.
- = Not applicable.

Consult the *Environmental Considerations for Marine Oil Spill Response* document referenced on page 5 before using this table.

Description

- Gravel beaches can be very steep, with multiple wave-built berms forming the upper beach.
- The degree of exposure to wave energy can be highly variable among gravel beaches.
- Density of animals and plants in the upper intertidal zone is low on exposed beaches, but can be high on sheltered gravel beaches and on the lower intertidal zone of all beaches.

Predicted Oil Behavior

- Stranded oil is likely to penetrate deeply into gravel beaches because of their high permeability.
- Long-term persistence will be controlled by the depth of routine reworking by the waves.
- Along sheltered portions of the shorelines, chronic sheening and the formation of asphalt pavements is likely where accumulations are heavy.

Response Considerations

- Heavy accumulations of pooled oil should be removed quickly from the upper beach.
- All oiled debris should be removed.
- Sediment removal should be limited as much as possible.
- Low- to high-pressure flushing can be effective if all released oil is recovered with skimmers or sorbents.
- Mechanical reworking of oiled sediments from the high-tide line to the lower beachface can be effective in areas regularly exposed to wave activity; the presence of multiple storm berms is evidence of wave activity.
- In-place tilling may be used to reach deeply buried oil layers along the mid-tide zone on exposed beaches.

Gravel Beaches

INTERTIDAL

Response Method	Oil Category				
	I	II	III	IV	V
Natural Recovery	A	A	B	B	B
Barriers/Berms	–	B	B	B	B
Manual Oil Removal/Cleaning	D	C	B	B	A
Mechanical Oil Removal	D	D	C	C	C
Sorbents	–	A	A	B	B
Vacuum	–	–	B	B	B
Debris Removal	–	A	A	A	A
Sediment Reworking/Tilling	D	B	B	B	B
Vegetation Cutting/Removal	–	–	–	–	–
Flooding (deluge)	A	A	B	C	C
Low-pressure, Ambient Water Flushing	A	A	A	B	C
High-pressure, Ambient Water Flushing	–	–	B	B	B
Low-pressure, Hot Water Flushing	–	–	C	B	B
High-pressure, Hot Water Flushing	–	–	C	C	C
Steam Cleaning	–	–	D	D	D
Sand Blasting	–	–	–	–	–
Solidifiers	–	–	B	–	–
Shoreline Cleaning Agents	–	–	B	B	B
Nutrient Enrichment	–	A	A	B	B
Natural Microbe Seeding	–	I	I	I	I
In-situ Burning	–	–	C	C	C


Oil Category Descriptions

- I – Gasoline products
- II – Diesel-like products and light crudes
- III – Medium grade crudes and intermediate products
- IV – Heavy crudes and residual products
- V – Non-floating oil products

The following categories are used to compare the relative environmental impact of each response method in the specific environment and habitat for each oil type. The codes in each table mean:

- A = The least adverse habitat impact.
- B = Some adverse habitat impact.
- C = Significant adverse habitat impact.
- D = The most adverse habitat impact.
- I = Insufficient information - impact or effectiveness of the method could not be evaluated.
- = Not applicable.

Consult the *Environmental Considerations for Marine Oil Spill Response* document referenced on page 5 before using this table.



Riprap

INTERTIDAL

Description

- Riprap structures are composed of cobble- to boulder-sized blocks of granite, limestone, concrete, or other materials.
- Riprap structures are used as revetment and groins for shoreline protection, and as breakwaters and jetties around inlets and marinas.
- Attached biota are generally sparse on exposed riprap.
- They are common in highly developed waterfront areas.

Predicted Oil Behavior

- Deep penetration of oil between the blocks is likely.
- Oil adheres readily to the rough surfaces of the blocks.
- Uncleaned oil can cause chronic leaching until the oil hardens.

Response Considerations

- When the oil is fresh and liquid, high pressure spraying and/or water flooding may be effective if all liberated oil is recovered.
- Heavy and weathered oils are more difficult to remove, requiring scraping and high-pressure, hot-water flushing.


INTERTIDAL

Response Method	Oil Category				
	I	II	III	IV	V
Oil Category Descriptions					
I - Gasoline products					
II - Diesel-like products and light crudes					
III - Medium grade crudes and intermediate products					
IV - Heavy crudes and residual products					
V - Non-floating oil products					
The following categories are used to compare the relative environmental impact of each response method in the specific environment and habitat for each oil type. The codes in each table mean:					
A = The least adverse habitat impact.					
B = Some adverse habitat impact.					
C = Significant adverse habitat impact.					
D = The most adverse habitat impact.					
I = Insufficient information - impact or effectiveness of the method could not be evaluated.					
— = Not applicable.					
Natural Recovery	A	A	B	B	B
Barriers/Berms	—	—	—	—	—
Manual Oil Removal/Cleaning	—	A	A	A	A
Mechanical Oil Removal	—	—	B	C	C
Sorbents	—	A	A	B	B
Vacuum	—	—	A	A	A
Debris Removal	—	A	A	A	A
Sediment Reworking/Tilling	—	—	—	—	—
Vegetation Cutting/Removal	—	—	—	—	—
Flooding (deluge)	A	A	B	C	C
Low-pressure, Ambient Water Flushing	A	A	B	C	C
High-pressure, Ambient Water Flushing	A	A	B	B	C
Low-pressure, Hot Water Flushing	—	C	C	C	C
High-pressure, Hot Water Flushing	—	C	C	C	C
Steam Cleaning	—	—	D	D	D
Sand Blasting	—	—	D	D	D
Solidifiers	—	B	B	—	—
Shoreline Cleaning Agents	—	—	B	B	B
Nutrient Enrichment	—	A	A	B	B
Natural Microbe Seeding	—	I	I	I	I
In-situ Burning	—	—	D	D	—

Consult the *Environmental Considerations for Marine Oil Spill Response* document referenced on page 5 before using this table.

Exposed Tidal Flats

INTERTIDAL



INTERTIDAL Description

- Exposed tidal flats are broad intertidal areas composed primarily of sand and minor amounts of gravel.
- The presence of sand indicates that tidal currents and waves are strong enough to mobilize the sediments.
- They are usually associated with another shoreline type on the landward side of the flat, though they can occur as separate shoals; they are commonly associated with tidal inlets.
- Biological use can be very high, with large numbers of infauna, heavy use by birds for roosting and foraging, and use by foraging fish.

Predicted Oil Behavior

- Oil does not usually adhere to the surface of exposed tidal flats, but rather moves across the flat and accumulates at the high-tide line.
- Deposition of oil on the flat may occur on a falling tide if concentrations are heavy.
- Oil does not penetrate water-saturated sediments, but may penetrate coarse-grained sand and coat gravel.
- Biological damage may be severe, primarily to infauna, thereby reducing food sources for birds and other predators.

Response Considerations

- Currents and waves can be very effective in natural removal of the oil.
- The use of heavy machinery should be restricted to prevent oil mixing into the sediments.

Exposed Tidal Flats

INTERTIDAL

Response Method	Oil Category				
	I	II	III	IV	V
Natural Recovery	A	A	A	A	A
Barriers/Berms	B	B	B	B	B
Manual Oil Removal/Cleaning	-	C	B	B	B
Mechanical Oil Removal	-	D	D	D	D
Sorbents	-	A	A	B	B
Vacuum	-	C	B	B	B
Debris Removal	-	B	B	B	B
Sediment Reworking/Tilling	-	-	C	C	C
Vegetation Cutting/Removal	-	D	D	D	D
Flooding (deluge)	-	A	A	A	B
Low-pressure, Ambient Water Flushing	-	B	B	C	C
High-pressure, Ambient Water Flushing	-	-	-	-	-
Low-pressure, Hot Water Flushing	-	-	-	-	-
High-pressure, Hot Water Flushing	-	-	-	-	-
Steam Cleaning	-	-	-	-	-
Sand Blasting	-	-	-	-	-
Solidifiers	-	C	C	-	-
Shoreline Cleaning Agents	-	-	-	-	-
Nutrient Enrichment	-	I	I	I	I
Natural Microbe Seeding	-	I	I	I	I
In-situ Burning	-	-	-	-	-

Oil Category Descriptions

- I - Gasoline products
- II - Diesel-like products and light crudes
- III - Medium grade crudes and intermediate products
- IV - Heavy crudes and residual products
- V - Non-floating oil products


The following categories are used to compare the relative environmental impact of each response method in the specific environment and habitat for each oil type. The codes in each table mean:


- A = The least adverse habitat impact.
- B = Some adverse habitat impact.
- C = Significant adverse habitat impact.
- D = The most adverse habitat impact.
- I = Insufficient information - impact or effectiveness of the method could not be evaluated.
- = Not applicable.

Consult the *Environmental Considerations for Marine Oil Spill Response* document referenced on page 5 before using this table.

INTERTIDAL

Sheltered Rocky Shores





INTERTIDAL

Description

- Sheltered rocky shores are characterized by a rocky substrate that can vary widely in permeability. Of particular concern are rocky shores that have a semi-permeable veneer of angular rubble overlying the bedrock.
- Sheltered clay scarps are characterized by a steep, usually vertical scarp in hard-packed and stiff clay. Vegetation usually occurs landward of the scarp.

Predicted oil behavior

- Oil will adhere readily to dry, rough, rocky surfaces, particularly at the high-tide line, forming a distinct oil band.
- The lower intertidal zone of rocky shores is usually algae-covered and stays wet, preventing oil from adhering.
- Oil will not adhere to the wet clay sediment surface, but could penetrate dry sediment.
- Stranded oil will persist because of the low-energy setting.

Response Considerations

- Low-pressure flushing of rocky shores at ambient temperatures is most effective when the oil is fresh and still liquid.
- Extreme care must be taken during flushing operations in the upper intertidal zone to prevent oily effluents from impacting biologically rich lower tidal levels.
- Do not cut oiled, attached algae; use sorbents to recover oil as it is remobilized by tidal action.
- Where the high-water area of scarps is accessible, it might be feasible to manually remove heavy oil accumulations and oiled debris.
- The muddy substrate of scarps cannot support heavy equipment, and even foot traffic could disrupt the sediments and mix oil deeper.

Response Method	Oil Category				
	I	II	III	IV	V
Natural Recovery	A	A	B	B	B
Barriers/Berms	-	-	-	-	-
Manual Oil Removal/Cleaning	-	C	B	C	C
Mechanical Oil Removal	-	-	-	-	-
Sorbents	A	A	B	C	C
Vacuum	-	B	B	B	C
Debris Removal	-	A	A	A	A
Sediment Reworking/Tilling	-	-	-	-	-
Vegetation Cutting/Removal	-	-	D	D	D
Flooding (deluge)	-	A	A	B	C
Low-pressure, Ambient Water Flushing	-	A	A	B	C
High-pressure, Ambient Water Flushing	-	C	B	B	C
Low-pressure, Hot Water Flushing	-	-	D	D	D
High-pressure, Hot Water Flushing	-	-	D	D	D
Steam Cleaning	-	-	D	D	D
Sand Blasting	-	-	D	D	D
Solidifiers	-	C	C	-	-
Shoreline Cleaning Agents	-	-	B	B	B
Nutrient Enrichment	-	A	B	C	C
Natural Microbe Seeding	-	I	I	I	I
In-situ Burning	-	D	C	C	C

Oil Category Descriptions

- I - Gasoline products
- II - Diesel-like products and light crudes
- III - Medium grade crudes and intermediate products
- IV - Heavy crudes and residual products
- V - Non-floating oil products

The following categories are used to compare the relative environmental impact of each response method in the specific environment and habitat for each oil type. The codes in each table mean:

- A = The least adverse habitat impact.
- B = Some adverse habitat impact.
- C = Significant adverse habitat impact.
- D = The most adverse habitat impact.
- I = Insufficient information - impact or effectiveness of the method could not be evaluated.
- = Not applicable.

Consult the *Environmental Considerations for Marine Oil Spill Response* document referenced on page 5 before using this table.

Description

- These are structures such as seawalls, groins, revetments, piers, and port facilities, constructed of concrete, wood, or metal.
- Most structures are designed to protect a single lot, thus their composition, design, and condition are highly variable.
- Often there is no exposed beach at low tide, but multiple habitats may be present.
- There can be dense attachments of animal and plant life.
- They are common in developed waterfront areas.

Predicted Oil Behavior

- Oil will adhere readily to the rough surface, particularly along the high-tide line, forming a distinct oil band.
- The lower intertidal zone usually stays wet (particularly if algae-covered), preventing oil from adhering to the surface.

Response Considerations

- Seawalls are usually cleaned for aesthetic reasons or to prevent leaching of oil.
- Low- to high-pressure spraying at ambient water temperatures is most effective when the oil is fresh.

INTERTIDAL

Sheltered, Solid Man-made Structures

Response Method	Oil Category				
	I	II	III	IV	V
Natural Recovery	A	A	B	B	B
Barriers/Berms	-	-	-	-	-
Manual Oil Removal/Cleaning	-	B	B	B	B
Mechanical Oil Removal	-	-	-	-	-
Sorbents	-	A	A	B	B
Vacuum	-	-	-	-	-
Debris Removal	-	A	A	A	A
Sediment Reworking/Tilling	-	-	-	-	-
Vegetation Cutting/Removal	-	-	-	-	-
Flooding (deluge)	-	-	-	-	-
Low-pressure, Ambient Water Flushing	-	A	B	C	C
High-pressure, Ambient Water Flushing	-	B	B	C	C
Low-pressure, Hot Water Flushing	-	-	C	C	C
High-pressure, Hot Water Flushing	-	-	C	C	C
Steam Cleaning	-	-	D	D	D
Sand Blasting	-	-	D	D	D
Solidifiers	-	-	-	-	-
Shoreline Cleaning Agents	-	-	B	B	B
Nutrient Enrichment	-	I	I	I	I
Natural Microbe Seeding	-	I	I	I	I
In-situ Burning	-	-	-	-	-

Oil Category Descriptions

- I - Gasoline products
- II - Diesel-like products and light crudes
- III - Medium grade crudes and intermediate products
- IV - Heavy crudes and residual products
- V - Non-floating oil products


The following categories are used to compare the relative environmental impact of each response method in the specific environment and habitat for each oil type. The codes in each table mean:

- A = The least adverse habitat impact.
- B = Some adverse habitat impact.
- C = Significant adverse habitat impact.
- D = The most adverse habitat impact.
- I = Insufficient information - impact or effectiveness of the method could not be evaluated.
- = Not applicable.

Consult the *Environmental Considerations for Marine Oil Spill Response* document referenced on page 5 before using this table.

Peat Shores

INTERTIDAL



Description

- This shoreline type includes exposed peat scarps, eroded peat, and peat slurries.
- Exposed peat scarps occur where the peat is frozen.
- They are highly erosional (>1 meter/year), resulting from wave action, ice scour, and melting of the frozen peat.
- The intertidal zone is often very complex, with slumped peat blocks and a thin (and temporary) sand layer on the peat.
- Eroded peat occurs as a peat mat or veneer in a dewatered state, deposited on a sand or gravel beach; it is usually less than 20 cm thick and considered to be relatively transient.
- Peat slurries (which have the appearance of coffee grounds) are up to 50 cm thick and 10 meters wide.
- Peat slurries are found at the foot of eroding peat scarps and in depositional areas; they are relatively permanent features that move along the shore with the currents.
- Peat shorelines comprise about 70 percent of the Beaufort Sea coast of Alaska.
- The intertidal zone of this shoreline type is not particularly important as biological habitat.

Predicted Oil Behavior

- Oil could be stranded onshore only during the ice-free summer season.
- Oil penetration and persistence are expected to be very low in frozen peat scarps.
- Light oil can penetrate peat slurries, especially when the peat is dry.
- Peat resists penetration by heavy oils, even when dry.
- Peat slurry reacts with oil like loose granular sorbent and will partially contain and prevent the oil from spreading.

Response Considerations

- The peat substrate is soft, thus cleanup will be difficult; trampling is less of concern where peat is frozen or work is conducted from boats.
- Substrate disruption is of limited concern because of high erosion rates so long as adjacent tundra is not disturbed.
- Peat slurry may be used as a natural sorbent; sorption will be more effective with liquid and fresh oils.
- With high erosion rates, stranded oil will have a short residence time.
- Tundra cliffs are commonly undercut and naturally unstable, so safety is a primary concern during response operations.
- Hot-water washing or even low-pressure flushing activities are not appropriate because large quantities of peat could be eroded from the treatment area.

Response Method	Oil Category				
	I	II	III	IV	V
Natural Recovery	A	A	A	A	A
Barriers/Berms	-	-	-	-	-
Manual Oil Removal/Cleaning	C	B	B	B	B
Mechanical Oil Removal	D	D	D	D	D
Sorbents	-	A	A	B	B
Vacuum	-	B	B	B	B
Debris Removal	C	B	B	B	B
Sediment Reworking/Tilling	C	C	B	B	B
Vegetation Cutting/Removal	D	D	C	C	C
Flooding (deluge)	C	B	B	C	D
Low-pressure, Ambient Water Flushing	C	B	B	C	D
High-pressure, Ambient Water Flushing	-	-	-	-	-
Low-pressure, Hot Water Flushing	-	-	-	-	-
High-pressure, Hot Water Flushing	-	-	-	-	-
Steam Cleaning	-	-	-	-	-
Sand Blasting	-	-	-	-	-
Solidifiers	-	-	-	-	-
Shoreline Cleaning Agents	-	-	-	-	-
Nutrient Enrichment	-	B	B	C	C
Natural Microbe Seeding	-	I	I	I	I
In-situ Burning	-	-	-	-	-

Oil Category Descriptions

- I - Gasoline products
- II - Diesel-like products and light crudes
- III - Medium grade crudes and intermediate products
- IV - Heavy crudes and residual products
- V - Non-floating oil products

The following categories are used to compare the relative environmental impact of each response method in the specific environment and habitat for each oil type. The codes in each table mean:

- A = The least adverse habitat impact.
- B = Some adverse habitat impact.
- C = Significant adverse habitat impact.
- D = The most adverse habitat impact.
- I = Insufficient information - impact or effectiveness of the method could not be evaluated.
- = Not applicable.

Consult the *Environmental Considerations for Marine Oil Spill Response* document referenced on page 5 before using this table.

Description

- Sheltered tidal flats are composed primarily of mud with minor amounts of sand and shell.
- They are usually present in calm-water habitats, sheltered from major wave activity, and backed by marshes.
- The sediments are very soft and cannot support even light foot traffic in many areas.
- There can be large concentrations of bivalves, worms, and other invertebrates in the sediments.
- They are heavily used by birds for feeding.

Predicted Oil Behavior

- Oil does not usually adhere to the surface of sheltered tidal flats, but rather moves across the flat and accumulates at the high-tide line.
- Deposition of oil on the flat may occur on a falling tide if concentrations are heavy.
- Oil will not penetrate the water-saturated sediments, but could penetrate burrows and desiccation cracks or other crevices in muddy sediments.
- In areas of high suspended sediment concentrations, the oil and sediments could mix, resulting in the deposition of contaminated sediments on the flats.
- Biological impacts may be severe.

Response Considerations

- These are high-priority areas for protection since cleanup options are limited.
- Cleanup of the flat surface is very difficult because of the soft substrate; many methods may be restricted.
- Low-pressure flushing and deployment of sorbents from shallow-draft boats may be attempted.

INTERTIDAL

Sheltered Tidal Flats

Response Method	Oil Category				
	I	II	III	IV	V
Natural Recovery	A	A	B	B	B
Barriers/Berms	B	B	B	B	B
Manual Oil Removal/Cleaning	-	D	C	C	C
Mechanical Oil Removal	-	-	-	-	-
Sorbents	-	A	A	B	B
Vacuum	-	C	B	B	B
Debris Removal	-	B	B	B	B
Sediment Reworking/Tilling	-	-	-	-	-
Vegetation Cutting/Removal	-	-	D	D	D
Flooding (deluge)	-	B	B	B	C
Low-pressure, Ambient Water Flushing	-	C	C	D	D
High-pressure, Ambient Water Flushing	-	-	-	-	-
Low-pressure, Hot Water Flushing	-	-	-	-	-
High-pressure, Hot Water Flushing	-	-	-	-	-
Steam Cleaning	-	-	-	-	-
Sand Blasting	-	-	-	-	-
Solidifiers	-	C	C	-	-
Shoreline Cleaning Agents	-	-	-	-	-
Nutrient Enrichment	-	I	I	I	I
Natural Microbe Seeding	-	I	I	I	I
In-situ Burning	-	-	-	-	-


Oil Category Descriptions

- I - Gasoline products
- II - Diesel-like products and light crudes
- III - Medium grade crudes and intermediate products
- IV - Heavy crudes and residual products
- V - Non-floating oil products

The following categories are used to compare the relative environmental impact of each response method in the specific environment and habitat for each oil type. The codes in each table mean:

- A = The least adverse habitat impact.
- B = Some adverse habitat impact.
- C = Significant adverse habitat impact.
- D = The most adverse habitat impact.
- I = Insufficient information - impact or effectiveness of the method could not be evaluated.
- = Not applicable.

Consult the *Environmental Considerations for Marine Oil Spill Response* document referenced on page 5 before using this table.



INTERTIDAL

Description

- Intertidal wetlands contain emergent, herbaceous vegetation, including both tidal and muted tidal marshes. Depending on location and interannual variations in rainfall and runoff, associated vegetation may include species tolerant or adapted to salt, brackish, or even tidal freshwater conditions.
- The marsh width may vary from a narrow fringe to extensive areas.
- Sediments are composed of organic muds except where sand is abundant on the margins of exposed areas.
- Exposed areas are located along bays with wide fetches and along heavily trafficked waterways.
- Sheltered areas are not exposed to significant wave or boat wake activity.
- Abundant resident flora and fauna with numerous species and high use by birds, fish, and shellfish.

Predicted Oil Behavior

- Oil adheres readily to intertidal vegetation.
- The band of coating will vary widely, depending upon the water level at the time of oiling.
- Large slicks will persist through multiple tidal cycles and will coat the entire stem from the high-tide line to the base.
- Heavy oil coating will be restricted to the outer fringe of thick vegetation, although lighter oils can penetrate deeper, to the limit of tidal influence.
- Medium to heavy oils do not readily adhere to or penetrate the fine sediments, but can pool on the surface or in animal burrows and root cavities.
- Light oils can penetrate the top few centimeters of sediment; under some circumstances oil can penetrate burrows and cracks up to one meter.

Response Considerations

- Under light oiling, the best practice is to let the area recover naturally.
- Natural removal processes and rates should be evaluated before conducting cleanup.
- Heavily pooled oil can be removed by vacuum, sorbents, or low-pressure flushing. During flushing, care must be taken to prevent transporting oil to sensitive areas down slope or along shore.
- Cleanup activities should be carefully supervised to avoid damaging vegetation.
- Any cleanup activity must not mix the oil deeper into the sediments. Trampling of the plants and disturbance of soft sediments must be minimized.
- Aggressive cleanup methods should only be considered when other resources (migratory birds, endangered species) are at greater risk from oiled vegetation left in place.

Response Method	Oil Category				
	I	II	III	IV	V
Natural Recovery	A	A	B	B	B
Barriers/Berms	B	B	B	B	B
Manual Oil Removal/Cleaning	D	D	C	C	C
Mechanical Oil Removal	D	D	D	D	D
Sorbents	-	A	A	A	B
Vacuum	-	B	B	B	B
Debris Removal	-	B	B	B	B
Sediment Reworking/Tilling	D	D	D	D	D
Vegetation Cutting/Removal	D	D	C	C	C
Flooding (deluge)	B	B	B	B	B
Low-pressure, Ambient Water Flushing	B	B	B	B	B
High-pressure, Ambient Water Flushing	-	-	-	-	-
Low-pressure, Hot Water Flushing	-	-	-	-	-
High-pressure, Hot Water Flushing	-	-	-	-	-
Steam Cleaning	-	-	-	-	-
Sand Blasting	-	-	-	-	-
Solidifiers	-	C	C	-	-
Shoreline Cleaning Agents	-	-	B	B	I
Nutrient Enrichment	-	A	B	B	B
Natural Microbe Seeding	-	I	I	I	I
In-situ Burning	-	B	B	B	C


Oil Category Descriptions

- I - Gasoline products
- II - Diesel-like products and light crudes
- III - Medium grade crudes and intermediate products
- IV - Heavy crudes and residual products
- V - Non-floating oil products

The following categories are used to compare the relative environmental impact of each response method in the specific environment and habitat for each oil type. The codes in each table mean:

- A = The least adverse habitat impact.
- B = Some adverse habitat impact.
- C = Significant adverse habitat impact.
- D = The most adverse habitat impact.
- I = Insufficient information - impact or effectiveness of the method could not be evaluated.
- = Not applicable.

Consult the *Environmental Considerations for Marine Oil Spill Response* document referenced on page 5 before using this table.



Description

- The roots and trunks are intertidal, with only the lowest leaves inundated by high tide.
- The width of the forest can range from one tree, to many kilometers.
- The substrate can be sand, mud, leaf litter, or peat, often as a veneer over bedrock.
- Wrack accumulations can be very heavy.
- They are highly productive, serve as nursery habitat, and support a great diversity and abundance of animal and plant species.

Predicted Oil Behavior

- Oil can wash through mangroves if oil comes ashore at high tide.
- If there is a berm or shoreline present, oil tends to concentrate and penetrate into the berm sediments or accumulated wrack/litter.
- Heavy and emulsified oil can be trapped in thickets of red mangrove prop roots.
- Oil readily adheres to prop roots, tree trunks, and pneumatophores.
- Reoiling from resuspended or released oil residues may cause additional injury over time.
- Oiled trees start to show evidence of effects (leaf yellowing) weeks after oiling; tree mortality may take months, especially for heavy oils.

Response Considerations

- Oiled wrack can be removed once the threat of oiling has passed. Wrack can actually protect the trees from direct oil contact.
- Sorbent boom can be placed in front of oiled forests to recover oil released naturally.
- In most cases, no other cleanup activities are recommended.
- Where thick oil accumulations are not being naturally removed, low-pressure flushing or vacuum may be attempted at the outer fringe.
- No attempt should be made to clean interior mangroves, except where access to the oil is possible from terrestrial areas.
- It is extremely important to prevent disturbance of the substrate by foot traffic; thus most activities should be conducted from boats.

INTERTIDAL

Mangroves

Response Method	Oil Category				
	I	II	III	IV	V
Natural Recovery	A	A	A	A	A
Barriers/Berms	B	B	B	B	B
Manual Oil Removal/Cleaning	-	D	C	C	C
Mechanical Oil Removal	-	-	-	-	-
Sorbents	-	A	A	A	B
Vacuum	-	B	B	B	B
Debris Removal	-	A	A	A	A
Sediment Reworking/Tilling	-	-	-	-	-
Vegetation Cutting/Removal	-	-	-	-	-
Flooding (deluge)	-	B	B	B	B
Low-pressure, Ambient Water Flushing	-	B	C	C	C
High-pressure, Ambient Water Flushing	-	-	-	-	-
Low-pressure, Hot Water Flushing	-	-	-	-	-
High-pressure, Hot Water Flushing	-	-	-	-	-
Steam Cleaning	-	-	-	-	-
Sand Blasting	-	-	-	-	-
Solidifiers	-	C	C	-	-
Shoreline Cleaning Agents	-	-	I	I	I
Nutrient Enrichment	-	I	I	I	I
Natural Microbe Seeding	-	I	I	I	I
In-situ Burning	-	-	-	-	-

Oil Category Descriptions

- I - Gasoline products
- II - Diesel-like products and light crudes
- III - Medium grade crudes and intermediate products
- IV - Heavy crudes and residual products
- V - Non-floating oil products


The following categories are used to compare the relative environmental impact of each response method in the specific environment and habitat for each oil type. The codes in each table mean:

- A = The least adverse habitat impact.
- B = Some adverse habitat impact.
- C = Significant adverse habitat impact.
- D = The most adverse habitat impact.
- I = Insufficient information - impact or effectiveness of the method could not be evaluated.
- = Not applicable.

Consult the *Environmental Considerations for Marine Oil Spill Response* document referenced on page 5 before using this table.

Inundated Lowland Tundra

INTERTIDAL



Description

- This shoreline type occurs where very low-lying sections of the Arctic shoreline have been recently flooded by the sea, due to subsidence.
- Also includes areas that are not normally in the intertidal zone but can be frequently inundated by salt water during spring tides or wind-induced surges.
- They have complex and convoluted shorelines comprised of tundra, vegetated flats, river banks, peat mats, brackish lagoons, and small streams.
- These shorelines have high ice content; the surface material is mostly peat with little mineral sediments.
- Where present, the vegetation is salt-tolerant and may be more adapted to drier conditions than the salt marshes.
- The tundra is a living plant community and provides important feeding areas for migrating birds in the summer.

Predicted Oil Behavior

- Oil could be stranded onshore only during the ice-free summer season.
- During storm surges, spilled oil could strand hundreds of meters inland.
- During the summer months, the surface sediments/peat deposits are usually water-saturated, so stranded oil is likely to remain on the surface.
- Physical removal rates of medium to heavy oils will be slow.

Response Considerations

- In summer, the substrate will be too soft to support foot or vehicular traffic; any work will require construction of walkways or roads.
- In winter, such work will be less damaging when the load-bearing capacity of these low-lying areas is increased.
- Excessive physical disruption can completely alter the substrate, hydrology, and vegetation patterns for many years.
- Avoid raking and trampling oil into living plants.
- Peat may be used as a natural sorbent; sorption will be more effective with liquid and fresh oils.
- Low-pressure, ambient-water flood and/or flushing could raise the local water table to float and direct oil towards a boomed area for collection.
- If salt-tolerant species are present, seawater may be used; use fresh water only if freshwater species are present.
- Consider burning only where there is an insulating water layer to protect roots and prevent deeper penetration into the substrate. Peat with a high water content may make burning ineffective, leaving a persistent surface residue that is more difficult to remove than the spilled oil.

Response Method	Oil Category				
	I	II	III	IV	V
Natural Recovery	A	A	A	B	B
Barriers/Berms	-	-	-	-	-
Manual Oil Removal/Cleaning	D	C	C	C	C
Mechanical Oil Removal	D	C	C	C	C
Sorbents	-	C	C	C	-
Vacuum	-	B	B	B	C
Debris Removal	-	C	C	C	C
Sediment Reworking/Tilling	-	-	-	-	-
Vegetation Cutting/Removal	D	D	D	D	D
Flooding (deluge)	C	C	C	D	-
Low-pressure, Ambient Water Flushing	-	D	D	-	-
High-pressure, Ambient Water Flushing	-	-	-	-	-
Low-pressure, Hot Water Flushing	-	-	-	-	-
High-pressure, Hot Water Flushing	-	-	-	-	-
Steam Cleaning	-	-	-	-	-
Sand Blasting	-	-	-	-	-
Solidifiers	-	C	C	-	-
Shoreline Cleaning Agents	-	-	-	-	-
Nutrient Enrichment	-	I	I	I	I
Natural Microbe Seeding	-	I	I	I	I
In-situ Burning	-	C	C	C	-

Consult the *Environmental Considerations for Marine Oil Spill Response* document referenced on page 5 before using this table.

Oil Category Descriptions


- I - Gasoline products
- II - Diesel-like products and light crudes
- III - Medium grade crudes and intermediate products
- IV - Heavy crudes and residual products
- V - Non-floating oil products

The following categories are used to compare the relative environmental impact of each response method in the specific environment and habitat for each oil type. The codes in each table mean:

- A = The least adverse habitat impact.
- B = Some adverse habitat impact.
- C = Significant adverse habitat impact.
- D = The most adverse habitat impact.
- I = Insufficient information - impact or effectiveness of the method could not be evaluated.
- = Not applicable.

SUBTIDAL

Coral Reefs



Description

- Coral reefs are structures created and maintained by the establishment and growth of populations of stony coral and coralline algae.
- Coral reefs are mostly subtidal in nature, although the most shallow portions of some reefs can be exposed during very low tides.
- Broad, pavement-like platforms formed by reefs when they reach sea level are a special concern.
- Many coral species spawn simultaneously over a very short time period (days), a behavior that makes the entire recruitment class very vulnerable.

Predicted Oil Behavior

- Coral reefs vary widely in sensitivity to spilled oil, depending on the water depth, oil type, and duration of exposure.
- There are three primary exposure pathways: direct contact with floating oil; exposure to dissolved and dispersed oil in the water column; and contamination of the substrate by oil deposited on the seafloor.
- Reef-associated community of fish, crustaceans, sea urchins, etc. can experience significant mortality.

Response Considerations

- Caution is needed when deploying and anchoring booms near reefs to prevent physical damage to the reef.
- Foot and vehicular traffic should not be allowed across a reef flat; access must be from the seaward side via boats.
- The use of dispersants directly over shallow reefs is likely to have significant impacts to the reef community. Their use in offshore areas can reduce impacts to highly sensitive intertidal environments.
- In situ burning outside of the immediate vicinity of reefs can protect sensitive intertidal environments. Burn residues can sink; the potential effects of these residues will depend on the composition and amount of oil.

SUBTIDAL

Coral Reefs

Response Method	Oil Category				
	I	II	III	IV	V
Natural Recovery	A	A	A	A	B
Booming	-	B	B	B	-
Skimming	-	B	B	B	-
Physical Herding	-	-	-	-	-
Manual Oil Removal/Cleaning	-	-	B	B	B
Mechanical Oil Removal	-	-	-	D	D
Sorbents	-	A	A	A	B
Vacuum	-	-	B	B	B
Debris Removal	-	-	-	-	-
Vegetation Cutting/Removal	-	-	-	-	-
Low-pressure, Ambient Water	B	B	B	C	C
Dispersants	-	C	C	C	-
In-situ Burning	-	B	B	B	-

Oil Category Descriptions

- I - Gasoline products
- II - Diesel-like products and light crudes
- III - Medium grade crudes and intermediate products
- IV - Heavy crudes and residual products
- V - Non-floating oil products

The following categories are used to compare the relative environmental impact of each response method in the specific environment and habitat for each oil type. The codes in each table mean:

- A = The least adverse habitat impact.
- B = Some adverse habitat impact.
- C = Significant adverse habitat impact.
- D = The most adverse habitat impact.
- I = Insufficient information - impact or effectiveness of the method could not be evaluated.
- = Not applicable.

Consult the *Environmental Considerations for Marine Oil Spill Response* document referenced on page 5 before using this table.



Description

- Seagrasses are highly productive habitats that occur on intertidal flats and in shallow coastal waters worldwide from arctic to tropical climates.
- Water temperature, light penetration, sediment type, salinity, and wave or current energy control seagrass distribution.
- Seagrasses provide a food source for green turtles, manatees, and waterfowl, who graze on seagrasses.
- Seagrasses are used by fish and shellfish as nursery areas.

Predicted Oil Behavior

- Oil will usually pass over subtidal seagrass beds, with no direct contamination.
- Oil that is heavier than seawater can become trapped in the beds, coating the leaves and sediments.
- Oil readily adheres to the vegetation, and the oiled blades are quickly defoliated when intertidal beds are oiled.
- Floating oil stranded on adjacent beaches can pick up sediment and then get eroded and deposited in adjacent beds.

Response Considerations

- Be careful when deploying and anchoring booms to prevent physical damage to seagrass beds.
- Be careful to prevent sediment suspension and mixing with the oil, and disturbance of roots and vegetation by foot traffic and boat activity.
- Do not cut seagrass unless species like sea turtles, manatees, or waterfowl are at significant risk of contacting or ingesting oil.
- Dispersant use directly over subtidal seagrass beds may impact the highly sensitive communities. However, use in offshore areas can reduce impacts to highly sensitive intertidal environments.
- In situ burning can be considered outside the immediate vicinity of seagrass beds to protect sensitive intertidal environments. Burn residues can sink; the potential effects of residues will depend on the composition and amount of the oil to be burned.

Seagrasses

SUBTIDAL

Response Method	Oil Category				
	I	II	III	IV	V
Natural Recovery	A	A	A	B	B
Booming	B	B	B	B	-
Skimming	-	B	B	B	-
Physical Herding	-	B	B	B	-
Manual Oil Removal/Cleaning	-	-	B	B	B
Mechanical Oil Removal	-	-	D	D	D
Sorbents	-	A	A	A	B
Vacuum	-	-	B	B	B
Debris Removal	-	-	B	B	B
Vegetation Cutting/Removal	-	-	C	C	C
Low-pressure, Ambient Water Flushing	-	-	-	-	-
Dispersants	-	C	C	C	-
In-situ Burning	-	B	B	B	-

Oil Category Descriptions

- I - Gasoline products
- II - Diesel-like products and light crudes
- III - Medium grade crudes and intermediate products
- IV - Heavy crudes and residual products
- V - Non-floating oil products


The following categories are used to compare the relative environmental impact of each response method in the specific environment and habitat for each oil type. The codes in each table mean:

- A = The least adverse habitat impact.
- B = Some adverse habitat impact.
- C = Significant adverse habitat impact.
- D = The most adverse habitat impact.
- I = Insufficient information - impact or effectiveness of the method could not be evaluated.
- = Not applicable.

Consult the *Environmental Considerations for Marine Oil Spill Response* document referenced on page 5 before using this table.

SUBTIDAL

Kelp



Description

- Kelps are very large brown algae that grow on hard subtidal substrates in cold temperate regions.
- Kelps have a holdfast that attaches to the substrate, a stem-like or trunk-like stipe, and large, flattened, leaf-like blades called fronds.
- Because kelps require constant water motion to provide nutrients, they are located in relatively high-energy settings.
- Kelp forests support a diverse animal community of fish, invertebrates, and marine mammals as well as important algal communities.

Predicted Oil Behavior

- Kelp has a mucous coating that prevents oil from adhering directly to the vegetation on the water surface.
- Oil can be trapped in the dense surface canopy, increasing the persistence of oil within the kelp environment.
- Oil persistence in kelp increases the risks of exposure to organisms concentrated in kelp forest habitats.

Response Considerations

- Cleanup efforts are often hampered by the difficulty of recovering oil from the dense canopy.
- Heavy oils could accumulate in sheltered pockets on the bottom, refloat during storms and re-expose resources to the oil.
- Use caution when anchoring vessels and boom to minimize mechanical damage to the kelp.
- Cutting kelp abruptly changes the light regime to the seafloor below.
- Cutting can be more appropriate for some kelp (*Macrocystis* and *Cystoseria*) than for others (*Nereocystis*).
- The impact of dispersed oil is likely to be greater on the community of organisms associated with the kelp habitat than on the kelp itself.
- In situ burning would be conditional on the absence or removal of mammals and birds in the immediate area. The kelp canopy might act as a natural boom against and within which oil can concentrate to burnable thicknesses.

Response Method	Oil Category				
	I	II	III	IV	V
Natural Recovery	A	A	A	B	B
Booming	-	B	B	B	-
Skimming	-	B	B	B	-
Physical Herding	-	B	B	B	-
Manual Oil Removal/Cleaning	-	-	-	-	-
Mechanical Oil Removal	-	-	-	-	-
Sorbents	-	A	A	A	-
Vacuum	-	-	-	-	-
Debris Removal	-	-	-	-	-
Vegetation Cutting/Removal	-	-	B	B	-
Low-pressure, Ambient Water Flushing	-	-	-	-	-
Dispersants	-	C	C	C	-
In-situ Burning	-	B	B	B	-

Oil Category Descriptions


- I - Gasoline products
- II - Diesel-like products and light crudes
- III - Medium grade crudes and intermediate products
- IV - Heavy crudes and residual products
- V - Non-floating oil products

The following categories are used to compare the relative environmental impact of each response method in the specific environment and habitat for each oil type. The codes in each table mean:

- A = The least adverse habitat impact.
- B = Some adverse habitat impact.
- C = Significant adverse habitat impact.
- D = The most adverse habitat impact.
- I = Insufficient information - impact or effectiveness of the method could not be evaluated.
- = Not applicable.

Consult the *Environmental Considerations for Marine Oil Spill Response* document referenced on page 5 before using this table.

SUBTIDAL



Description

- Soft-bottom, subtidal habitats consist of various percentages of sand, silt, and clay, occurring in sheltered bays and estuaries, and deeper offshore areas.
- The presence of fine-grained sediments indicates that the substrate is not exposed to significant wave or tidal energy.
- Biological resources associated with this habitat include shrimp, crabs, clams, fish, and the pelagic and benthic communities that support them (e.g., plankton, worms, amphipods, isopods).

Predicted Oil Behavior

- This habitat is not often exposed to spilled oil. The greatest risk of exposure is from the sinking oil or the sorption of dispersed oil onto suspended sediments that are then deposited on the bottom.
- Significant natural dispersion of oil and sediments into the water column occurs only during large storms and nearshore oil spills.
- Shoreline cleanup can suspend oil and fine-grained sediments, causing deposition of oily sediments in nearshore habitats.
- Concerns about seafood contamination from dispersed oil or oiled sediments can become a significant issue. Real, potential, or fear of contamination can close seafood harvesting activities.

Response Considerations

- Removal might be needed where significant amounts of oil have sunk and formed mats or concentrations of tarballs on the sediment surface.
- Special efforts will be needed to control suspended sediments and resuspended oil during recovery operations.
- Dispersants can be used over soft subtidal habitats in order to protect more sensitive intertidal environments. Effects on biota are less for applications in deep water or high dilution rates.
- In situ burning can be used to protect sensitive intertidal environments. When burned, some oils can produce a sinkable residue; the potential effects of these residues will depend on the composition and amount of oil to be burned.

Response Method	Oil Category				
	I	II	III	IV	V
Natural Recovery	A	A	A	B	B
Booming	A	A	A	A	-
Skimming	-	A	A	A	-
Physical Herding	-	B	B	B	-
Manual Oil Removal/Cleaning	-	-	B	B	B
Mechanical Oil Removal	-	-	-	C	C
Sorbents	-	A	A	A	B
Vacuum	-	-	B	B	B
Debris Removal	-	-	-	-	-
Vegetation Cutting/Removal	-	-	-	-	-
Low-pressure, Ambient Water Flushing	-	-	-	-	-
Dispersants	-	C	C	C	-
In-situ Burning	-	B	B	B	-

Oil Category Descriptions

- I - Gasoline products
- II - Diesel-like products and light crudes
- III - Medium grade crudes and intermediate products
- IV - Heavy crudes and residual products
- V - Non-floating oil products


The following categories are used to compare the relative environmental impact of each response method in the specific environment and habitat for each oil type. The codes in each table mean:

- A = The least adverse habitat impact.
- B = Some adverse habitat impact.
- C = Significant adverse habitat impact.
- D = The most adverse habitat impact.
- I = Insufficient information - impact or effectiveness of the method could not be evaluated.
- = Not applicable.

Consult the *Environmental Considerations for Marine Oil Spill Response* document referenced on page 5 before using this table.

SUBTIDAL

Mixed and Hard Bottom



Description

- This habitat consists of subtidal substrates composed of rock, boulders, or cobbles, though there can be patches of sand veneer covering a hard bottom.
- There may be rich, diverse communities of attached and associated algae and animals; often there is little open space.
- Some of these habitats form a relief (reef or bank) several meters high that attracts a diversity of fish.

Predicted Oil Behavior

- Mixed and hard-bottom habitats are usually considered to have low sensitivity to oil spills.
- Oil in the water column seldom reaches toxic levels and benthic organisms have little exposure.
- There is little risk of deposition of oil or oiled sediments in these habitats.
- There could be a short-term exposure as oiled sediments are transported through the habitat into deeper areas.
- Concerns about seafood contamination from dispersed oil or oiled sediments can become a significant issue. Real, potential, or fear of contamination can close seafood harvesting activities.

Response Considerations

- Natural cleansing is expected to occur quickly, especially in the higher-energy environments.
- Avoid anchoring booms in known sensitive areas, such as unique live-bottom areas.
- Dispersants can be used directly over these habitats to protect sensitive intertidal areas. The deeper the water, the greater the dilution, and hence the lesser effect it will have on the mixed and hard-bottom habitats.
- In situ burning can be used directly over these habitats to protect sensitive intertidal environments. When burned, some oils can produce a sinkable residue; the potential effects of these residues will depend on the composition and amount of oil to be burned.

SUBTIDAL

Mixed and Hard Bottom

Response Method	Oil Category				
	I	II	III	IV	V
Natural Recovery	A	A	A	B	B
Booming	-	B	B	B	-
Skimming	-	A	A	A	-
Physical Herding	-	A	A	A	-
Manual Oil Removal/Cleaning	-	-	B	B	B
Mechanical Oil Removal	-	-	-	-	-
Sorbents	-	A	A	A	B
Vacuum	-	-	B	B	B
Debris Removal	-	-	-	B	B
Vegetation Cutting/Removal	-	-	-	-	-
Low-pressure, Ambient Water Flushing	-	-	-	-	-
Dispersants	-	B	B	B	-
In-situ Burning	-	B	B	B	-

Oil Category Descriptions

- I - Gasoline products
- II - Diesel-like products and light crudes
- III - Medium grade crudes and intermediate products
- IV - Heavy crudes and residual products
- V - Non-floating oil products


The following categories are used to compare the relative environmental impact of each response method in the specific environment and habitat for each oil type. The codes in each table mean:

- A = The least adverse habitat impact.
- B = Some adverse habitat impact.
- C = Significant adverse habitat impact.
- D = The most adverse habitat impact.
- I = Insufficient information - impact or effectiveness of the method could not be evaluated.
- = Not applicable.

Consult the *Environmental Considerations for Marine Oil Spill Response* document referenced on page 5 before using this table.

Accessible and Inaccessible Ice

ICE



Description

- Ice forms on the sea surface during winter in cold climates and can persist for several months.
- Most sea surface ice is floating but can be frozen to the bottom or stranded in intertidal areas during low tide.
- *Accessible ice* can safely support the personnel and equipment suitable for response to a particular oil spill on, in, under, or adjacent to solid ice.
- *Inaccessible ice* cannot safely support response personnel and response equipment.

Predicted Oil Behavior

- Ice along the shoreline or in the adjacent nearshore water can act as a natural barrier, reducing the amount of oil that might otherwise make contact with the shoreline substrate.
- During the ice growth phase, oil in or under the ice can become encapsulated within the ice.
- During a thaw, or if the surface of the ice is melting and wet, oil is unlikely to adhere to the ice surface and will tend to remain on the water surface or in leads.
- In the spring, before the ice becomes inaccessible, oil in or below sea ice will often migrate through brine channels to the surface.

Response Considerations

- The ice habitat presents unique safety issues in terms of cold, ice stability, and wildlife interactions.
- Oil spills on, in, under, or adjacent to brash ice, small or fast moving floes, or other ice types which are “inaccessible” must be treated from the air or from vessels working in, or alongside, the ice.
- Some methods, including flooding, debris removal, sediment reworking, vegetation cutting and removal, high-pressure flushing, sand blasting, solidifiers, and shoreline cleaning agents, are not considered suitable for use in these environments.

Accessible and Inaccessible Ice

ICE

Relative environmental impacts

Accessible Ice

Response Method	Oil Category				
	I	II	III	IV	V
Natural Recovery	A	B	B	C	C
Booming	-	B	B	B	-
Skimming	-	A	A	A	-
Barriers/Berms	B	B	B	-	-
Physical Herding	B	B	B	B	-
Manual Oil Removal/Cleaning	-	A	A	A	A
Mechanical Oil Removal	-	B	B	B	B
Sorbents	-	B	B	B	-
Vacuum	-	A	A	A	A
Low-pressure, Ambient Water Flushing	B	B	B	B	C
Low-pressure, Hot Water Flushing	-	B	B	B	C
Steam Cleaning	-	B	B	B	C
Dispersants	-	B	B	-	-
Emulsion-treating Agents	-	I	I	I	I
Elasticity Modifiers	-	A	-	-	-
Herding Agents	I	I	I	-	-
Nutrient Enrichment	-	I	I	I	I
Natural Microbe Seeding	-	I	I	I	I
In-situ Burning	B	B	B	B	-

Oil Category Descriptions

- I - Gasoline products
- II - Diesel-like products and light crudes
- III - Medium grade crudes and intermediate products
- IV - Heavy crudes and residual products
- V - Non-floating oil products

The following categories are used to compare the relative environmental impact of each response method in the specific environment and habitat for each oil type. The codes in each table mean:

- A = The least adverse habitat impact.
- B = Some adverse habitat impact.
- C = Significant adverse habitat impact.
- D = The most adverse habitat impact.
- I = Insufficient information - impact or effectiveness of the method could not be evaluated.
- = Not applicable.

Consult the *Environmental Considerations for Marine Oil Spill Response* document referenced on page 5 before using this table.

ICE

Relative environmental impacts

Inaccessible Ice

Oil Category Descriptions

- I - Gasoline products
- II - Diesel-like products and light crudes
- III - Medium grade crudes and intermediate products
- IV - Heavy crudes and residual products
- V - Non-floating oil products


The following categories are used to compare the relative environmental impact of each response method in the specific environment and habitat for each oil type. This method may cause:

- A = The least adverse habitat impact.
- B = Some adverse habitat impact.
- C = Significant adverse habitat impact.
- D = The most adverse habitat impact.
- I = Insufficient information - impact or effectiveness of the method could not be evaluated.
- = Not applicable.

Accessible and Inaccessible Ice

Response Method	Oil Category				
	I	II	III	IV	V
Natural Recovery	A	A	B	B	B
Booming	-	B	B	B	-
Skimming	-	A	A	A	-
Barriers/Berms	-	-	-	-	-
Physical Herding	B	B	B	B	-
Manual Oil Removal/Cleaning	-	-	-	-	-
Mechanical Oil Removal	-	B	B	B	B
Sorbents	-	B	B	B	-
Vacuum	-	A	A	A	-
Low-pressure, Ambient Water Flushing	-	B	B	B	-
Low-pressure, Hot Water Flushing	-	B	B	B	-
Steam Cleaning	-	-	-	-	-
Dispersants	-	B	B	-	-
Emulsion-treating Agents	-	I	I	I	I
Elasticity Modifiers	-	B	-	-	-
Herding Agents	I	I	I	-	-
Nutrient Enrichment	-	I	I	I	I
Natural Microbe Seeding	-	I	I	I	I
In-situ Burning	B	B	B	-	-

Use this table only after consulting the *Environmental Considerations for Marine Oil Spill Response* document referenced on page 5.



ON-WATER

Description

- Offshore waters are those where the water depth is > 30 feet (10 meters) with no surrounding land.
- Evaluation of environmental impacts to open water habitats is focused on water column organisms and those which inhabit or use the sea surface.
- Animals include marine mammals, sea turtles, pelagic birds, and many commercially and recreationally important fish and pelagic invertebrates.
- Organism densities in this habitat are low on average.
- Localized high densities can occur in areas such as convergence zones and upwelling areas.
- Pelagic birds are at greatest risk when large numbers are concentrated for feeding, migration, overwintering, or breeding.
- Biological resources in the water column are less vulnerable to spills than those at the water surface.
- The sea surface microlayer is important for biochemical processes; the organisms most vulnerable to exposure are poor or passive swimmers (planktonic forms).

Predicted Oil Behavior

- Spilled oil transport is controlled more by wind and ocean currents than by tides and mixing with freshwater outflows.
- Most of the soluble and toxic components of the spilled oil are lost through weathering within hours and days.
- Dissolved or dispersed oil concentrations are likely to be greatest in the top few meters.

Response Considerations

- Response activities are focused on removing oil from the water surface.
- Spill response is not conducted from a shoreline, but from water-based vessels or aircraft.
- Weather and sea conditions can significantly hamper response operations.
- Category V oils are likely to submerge and most of the response methods can only be used on the surface of the water.
- Special equipment might be needed for some products (e.g., containment booms which extend at least 9 ft.).
- Use of certain response options is seasonally limited to protect sensitive life histories.

Response Method	Oil Category				
	I	II	III	IV	V
Natural Recovery	A	A	B	B	B
Booming-Containment	-	A	A	A	-
Booming-Deflection/Exclusion	A	A	A	A	-
Skimming	-	A	A	A	-
Physical Herding	B	B	B	B	-
Manual Oil Removal/Cleaning	-	-	-	-	-
Sorbents	-	B	B	B	-
Debris Removal	-	A	A	A	-
Dispersants	B	A	A	A	-
Emulsion-treating Agents	-	B	B	B	-
Elasticity Modifiers	-	B	B	-	-
Herding Agents	-	B	B	-	-
Solidifiers	-	B	B	-	-
In-situ Burning	-	A	A	A	-


Oil Category Descriptions

- I - Gasoline products
- II - Diesel-like products and light crudes
- III - Medium grade crudes and intermediate products
- IV - Heavy crudes and residual products
- V - Non-floating oil products

The following categories are used to compare the relative environmental impact of each response method in the specific environment and habitat for each oil type. The codes in each table mean:

- A = The least adverse habitat impact.
- B = Some adverse habitat impact.
- C = Significant adverse habitat impact.
- D = The most adverse habitat impact.
- I = Insufficient information - impact or effectiveness of the method could not be evaluated.
- = Not applicable.

Consult the *Environmental Considerations for Marine Oil Spill Response* document referenced on page 5 before using this table.



Description

- Near coastal waters partially surrounded by land and more sheltered than offshore habitats.
- Limited circulation and flushing, with depths frequently <30 feet.
- Suspended sediment concentrations can be high.
- Highly sensitive to oil spills, particularly where flushing rates are low and the probability of contact increases.
- Many species spawn in these habitats during spring, and their sensitive early life stages can persist in shallow waters.
- Large numbers of migratory or wintering waterfowl, wading, and diving birds are often found here. Bays and estuaries are also home to marine mammals and sea turtles.
- Estuaries and bays are used by commercially or recreationally important finfish, shellfish, and other organisms that migrate seasonally.

Predicted Oil Behavior

- Oil can impact bottom habitats (benthic organisms) when water is shallow.
- Stranded oil on nearby shorelines can become a prolonged source for oil re-released to the water column.
- Tides and fresh water can substantially influence spilled oil movement.

Response Considerations

- Reducing impacts to organisms that live on or in the sea surface is often a high priority.
- Reducing the extent of impacts to sensitive nearshore subtidal or intertidal habitats should be considered.
- Spill response is not conducted from a shoreline, but from water-based vessels or aircraft.
- Use of certain response options is seasonally limited to protect sensitive life histories.
- Adverse effects to birds would be greatest during migration and overwintering when the birds form large flocks.

Response Method	Oil Category				
	I	II	III	IV	V
Oil Category Descriptions					
I - Gasoline products					
II - Diesel-like products and light crudes					
III - Medium grade crudes and intermediate products					
IV - Heavy crudes and residual products					
V - Non-floating oil products					
Natural Recovery	A	B	B	C	C
Booming-Containment	-	A	A	B	-
Booming-Deflection/Exclusion	A	A	A	B	-
Skimming	-	A	A	A	-
Physical Herding	B	B	B	B	-
Manual Oil Removal/Cleaning	-	-	C	B	B
Sorbents	-	B	B	B	-
Debris Removal	-	A	A	A	B
Dispersants	B	B	B	B	-
Emulsion-treating Agents	-	B	B	B	-
Elasticity Modifiers	-	B	B	-	-
Herding Agents	-	B	B	-	-
Solidifiers	-	B	B	-	-
In-situ Burning	-	A	A	B	-

The following categories are used to compare the relative environmental impact of each response method in the specific environment and habitat for each oil type. The codes in each table mean:

- A = The least adverse habitat impact.
- B = Some adverse habitat impact.
- C = Significant adverse habitat impact.
- D = The most adverse habitat impact.
- I = Insufficient information - impact or effectiveness of the method could not be evaluated.
- = Not applicable.

Consult the *Environmental Considerations for Marine Oil Spill Response* document referenced on page 5 before using this table.



U.S. Department of Commerce
National Oceanic and Atmospheric Administration

Norman Y. Mineta
Secretary, U.S. Department of Commerce

D. James Baker
Under Secretary of Commerce for
Oceans and Atmosphere and
Administrator, National Oceanic and
Atmospheric Administration

Scott B. Gudes
Deputy Under Secretary of Commerce for
Oceans and Atmosphere (Acting)

Margaret A. Davidson
Acting Assistant Administrator for
Ocean Services and Coastal Zone Management,
NOAA National Ocean Service