

**NORA manufacturing sector and unspecified and other pneumoconioses:
Proportionate mortality ratio (PMR) adjusted for age, sex, and race by industry and
occupation, U.S. residents age 15 and over, selected states and years, 1990–1999**

Bureau of Census Industry/Occupation		Number of Deaths*	PMR	95% Confidence Interval	
				LCL	UCL
CIC	Industry				
252	Structural clay products	7	4.5	1.8	9.2
262	Miscellaneous nonmetallic mineral and stone products	7	3.0	1.2	6.3
271	Iron and steel foundries	10	2.8	1.3	5.1
	All other industries within this sector	212			
	TOTAL	236			
COC	Occupation				
783	Welders and cutters	14	1.9	1.0	3.2
	All other occupations within this sector	218			
	Occupation not reported within this sector	4			
	TOTAL	236			

NORA - National Occupational Research Agenda CIC - Census Industry Code COC - Census Occupation Code

n.e.c. - not elsewhere classified LCL - lower confidence limit UCL - upper confidence limit

* The total number of unspecified and other pneumoconioses deaths was 1,440 in these same selected states and years, and the comparable number of unspecified and other pneumoconioses deaths in the entire United States for this same time period was 3,384.

NOTE: See appendices for source description, methods, ICD codes, industry and occupation codes, industrial sector groupings, and list of selected states and years.

SOURCE: National Center for Health Statistics multiple cause-of-death data.