

2005Table 6-1. All pneumoconioses: Number of deaths by sex, race, and age, and median age at death, U.S. residents age 15 and over, 1993–2002

Year	No. of Deaths	Underlying Cause (%)	Sex		Race			Age Group (yrs)								Median Age (yrs)
			Male	Female	White	Black	Other	15–24	25–34	35–44	45–54	55–64	65–74	75–84	85+	
1993	3,238	35.3	3,170	68	3,049	173	16	-	4	14	56	277	953	1,410	524	77.0
1994	3,127	34.9	3,053	74	2,964	154	9	-	-	10	45	233	894	1,434	511	77.0
1995	3,151	35.4	3,092	59	2,983	160	8	-	4	10	68	245	913	1,337	574	77.0
1996	3,115	35.8	3,031	84	2,937	164	14	-	1	11	51	210	883	1,364	595	78.0
1997	2,928	37.1	2,851	77	2,780	139	9	-	2	12	48	191	751	1,313	611	78.0
1998	2,791	39.4	2,720	71	2,651	127	13	-	3	11	41	189	733	1,226	588	78.0
1999	2,745	39.5	2,683	62	2,592	136	17	-	1	9	38	177	683	1,210	627	78.0
2000	2,864	39.9	2,783	81	2,704	144	16	2	4	14	42	170	651	1,236	745	79.0
2001	2,747	40.4	2,674	73	2,585	148	14	2	-	11	47	194	632	1,195	666	78.0
2002	2,718	40.3	2,631	87	2,570	137	11	1	3	10	41	186	599	1,213	665	79.0
TOTAL	29,424	37.7	28,688	736	27,815	1,482	127	5	22	112	477	2,072	7,692	12,938	6,106	78.0

- indicates no deaths listed.

NOTE: Slight variations in numbers may exist when compared to previous WoRLD Surveillance Reports. See appendices for source description, methods, and ICD codes.

SOURCE: National Center for Health Statistics multiple cause-of-death data.