

Polyphosphate kinase as a nucleoside diphosphate kinase in *Escherichia coli* and *Pseudomonas aeruginosa*

AKIO KURODA* AND ARTHUR KORNBERG†

Department of Biochemistry, Stanford University School of Medicine, Stanford, CA 94305-5307

Contributed by Arthur Kornberg, November 15, 1996

ABSTRACT Generation of a wide variety of nucleoside (and deoxynucleoside) triphosphates (NTPs) from their cognate nucleoside diphosphates (NDPs) is of critical importance in virtually every aspect of cellular life. Their function is fulfilled largely by the ubiquitous and potent nucleoside diphosphate kinase (NDK), most commonly using ATP as the donor. Considerable interest is attached to the consequence to a cell in which the NDK activity becomes deficient or overabundant. We have discovered an additional and possibly auxiliary NDK-like activity in the capacity of polyphosphate kinase (PPK) to use inorganic polyphosphate as the donor in place of ATP, thereby converting GDP and other NDPs to NTPs. This reaction was observed with the PPK activity present in crude membrane fractions from *Escherichia coli* and *Pseudomonas aeruginosa* as well as with the purified PPK from *E. coli*; the activity was absent from the membrane fractions obtained from *E. coli* mutants lacking the *ppk* gene. The order of substrate specificity for PPK was: ADP > GDP > UDP, CDP; activity with ADP was 2–60 times greater than with GDP, depending on the reaction condition. Although the transfer of a phosphate from polyphosphate to GDP by PPK to produce GTP was the predominant reaction, the enzyme also transferred a pyrophosphate group to GDP to form the linear guanosine 5' tetraphosphate.

Supplies of the ribo- and deoxyribonucleoside triphosphates (NTPs, dNTPs) are crucial in virtually all biosynthetic and regulatory activities. Some notable examples in eukaryotic systems are apparent in developmental control (1–5), signal transduction (6, 7), and tumor metastasis (8, 9). The source of the various NTPs is the phosphorylation of the nucleoside diphosphates (NDPs) by the ubiquitous nucleoside diphosphate kinase (NDK), generally with ATP as the donor molecule (10). The cellular consequences of varying the level of NDK have been examined recently in *Escherichia coli* (11). Mutants lacking NDK grew at a normal rate but showed an increased mutation frequency, presumably due to the imbalance of dNTP pools (11). The residual NDK activity to support growth in the mutants was supplied by adenylate kinase (12), an enzyme previously thought to be specific for adenosine nucleotides.

Among the numerous functions observed or considered for inorganic polyphosphate (polyP) are roles as an ATP substitute and as an energy reservoir (13). A linear polymer of hundreds of phosphate linked by high-energy, phosphoanhydride bonds, polyP is made from the terminal phosphate of ATP in a readily reversible reaction, the phosphorylation of ADP by polyP. polyP has been demonstrated to substitute for ATP in the kinase-catalyzed phosphorylation of sugars (14,

15), protein (16), AMP (17), and ADP [by polyP kinase (PPK); ref. 18].

Although PPK of *E. coli* is specific for ATP as the nucleoside triphosphate in the formation of polyP, this report will show that in the reverse reaction, polyP can serve as a donor not only for ADP but also for GDP and other NDPs, thus fulfilling the function of a NDK. The PPK activity observed in crude extracts of *Pseudomonas aeruginosa* has properties similar to those of *E. coli* PPK.

MATERIALS AND METHODS

Reagents and Proteins. Sources were as follows: [γ - 32 P]ATP, Amersham; all nonradiolabeled NDPs and NTPs, guanosine 5'-tetraphosphate (ppppG), and bovine serum albumin, Sigma; and polyethylenimine cellulose (PEI)-TLC plates, Merck.

Bacterial Strains. The parental *E. coli* MG1655 (wild type) and the null mutant, $\Delta ppk ppk::kan$ were generously provided by M. Cashel (National Institutes of Health). *P. aeruginosa* PAO1 was furnished by A. Chakrabarty (University of Illinois).

Preparation of Crude Membrane Fractions. Cells were grown at 37°C on Luria-Bertani (LB) medium and harvested. The cell pellet was resuspended in 100 μ l of buffer A (50 mM Tris-Cl, pH 7.5/10% sucrose), frozen in liquid nitrogen, and stored at -80°C . Lysozyme (250 $\mu\text{g}/\text{ml}$) was added to the thawed cell suspension and incubated at 0°C for 30 min. The cells were lysed by exposure to 37°C for 4 min, followed by immediate chilling in ice water and subjected to centrifugation [65,000 rpm in a model TA100-1 centrifuge (Beckman) for 20 min]. The pellet was washed with buffer A and dispersed by sonication in buffer A containing 5 mM MgCl_2 , 10 $\mu\text{g}/\text{ml}$ RNase, and 10 $\mu\text{g}/\text{ml}$ DNase.

polyP-Dependent NDK Assay. The reaction mixture contained a crude membrane fraction or purified PPK (7×10^7 units per mg of protein) in PPK buffer {50 mM Hepes, pH 7.2/4 mM MgCl_2 /40 mM $(\text{NH}_4)_2\text{SO}_4$ unless otherwise indicated/0.1 mM [^{32}P]polyP/1 mM NDP}. [^{32}P]polyP was made as described previously (18). The mixture was incubated at 37°C for the time indicated, and 1- μ l samples were spotted onto a PEI-TLC plate, which was developed with 0.75 M or 1.5 M KH_2PO_4 (pH 3.5). The plate was dried after developing, exposed to film, and then visualized in a PhosphorImager scanner (Molecular Dynamics). The ratio of the image intensity of the NTP spot to the total (polyP plus NTP) was calculated. One unit of activity generated 1 pmol of [^{32}P]NTP per min.

Two-Dimensional TLC. Identification of ppppG was performed as described by Bochner and Ames (19). The solvent

The publication costs of this article were defrayed in part by page charge payment. This article must therefore be hereby marked "advertisement" in accordance with 18 U.S.C. §1734 solely to indicate this fact.

Copyright © 1997 by THE NATIONAL ACADEMY OF SCIENCES OF THE USA
0027-8424/97/94439-4\$2.00/0

PNAS is available online at <http://www.pnas.org>.

Abbreviations: NDK, nucleoside diphosphate kinase; polyP, inorganic polyphosphate; PPK, polyP kinase; ppppG, guanosine 5'-tetraphosphate; PPX, exopolyphosphatase.

*Present address: Department of Fermentation Technology, Hiroshima University, 1-4-1 Kagamiyama, Higashi-Hiroshima, Hiroshima 739, Japan.

†To whom reprint requests should be addressed.




FIG. 1. NDK activity of purified *E. coli* PPK. The reaction mixture (10 μ l) containing 29 ng of purified PPK, PPK buffer, 1 mM (d)NDP, and 0.1 mM [32 P]polyP was incubated at 37°C for 30 min. The TLC plate was developed with 1.5 M KH_2PO_4 (pH 3.5) and autoradiographed.

in the first dimension was 0.75 M Tris/0.45 M HCl/0.5 M LiCl, and the solvent in the second dimension contained 43.5 g of NH_4SO_4 , 0.4 g of NH_4HSO_4 , and 4 g of Na_2EDTA per 100 ml.

Digestion of ppppG by Yeast Exopolyphosphatase (PPX). After development of ppppG on PEI-TLC, the spot containing ppppG was scratched from the plate and then incubated with 0.5 ml of 0.25 M NH_4HCO_3 for 30 min at room temperature. The PEI cellulose was pelleted by centrifugation in a microfuge, the supernatant was removed, and the elution process was repeated. The two supernatant fractions were pooled, concentrated in a lyophilizer overnight, and suspended in water. Yeast PPX (6.7×10^7 units per mg of protein) was used for digestion of ppppG in the PPX buffer (20).

RESULTS

PPK Catalyzes the Synthesis of NTPs by Transfer of a Phosphate Group from polyP to NDPs. Purified PPK from *E. coli* can transfer a phosphate group from polyP to all NDPs, including ADP, GDP, CDP, UDP, dADP, dGDP, dCDP, and TDP (Fig. 1). The efficiency of NDP substrates was as follows: ADP, dADP > dGDP, GDP, TDP > UDP, CDP, dCDP. The kinetic properties of PPK for various NDPs is summarized in Table 1. The specificity of PPK for nucleoside diphosphates was reduced in the absence of $(\text{NH}_4)_2\text{SO}_4$ and in the presence of 1 mM CaCl_2 in the reaction mixture (Table 2). Zn^{2+} and Cu^{2+} reduced the total activity, but Co^{2+} and Mn^{2+} had no significant effect (data not shown).

Table 1. K_m and k_{cat} values of *E. coli* PPK

NDP	K_m , mM	k_{cat} , min^{-1}	k_{cat}/K_m , $\text{mM}^{-1}\cdot\text{min}^{-1}$
ADP	0.25	630	2500
GDP	0.63	5.3	8.4
CDP	>3	10	<3
UDP	>3	7	<2

Table 2. NDK specific activity of PPK

NDP*	<i>E. coli</i> PPK (purified)				Membrane	
	-(NH_4) $_2\text{SO}_4$		-(NH_4) $_2\text{SO}_4$, + Ca^{2+}		<i>E. coli</i>	<i>P. aeruginosa</i>
ADP	370	26	25	4.3	1.3	1.7
GDP	5.7	3.4	3.9	2.0	0.032	0.23
CDP	1.7	1.2	1.8	1.3	0.007	0.021
UDP	1.7	1.4	2.1	1.3	0.006	0.027

Reaction buffer is based on 50 mM Hepes (pH 7.2), 4 mM MgCl_2 , and 40 mM $(\text{NH}_4)_2\text{SO}_4$ (18). $(\text{NH}_4)_2\text{SO}_4$ was removed from the buffer, or 1 mM CaCl_2 was added if indicated. All values are in pmol of NTP per min per mg of protein $\times 10^{-4}$.

*[32 P]polyP (0.1 mM) was added to each NDP substrate. NDPs were at a concentration of 1 mM.

polyP-Dependent NDK Activity in the Crude Membrane Fraction. Crude membrane fractions from *E. coli* contain virtually all the PPK activity catalyzing the synthesis of ATP from ADP and polyP (Fig. 2A). GTP synthesis activity from GDP and polyP was also detected in this fraction (Fig. 2A), as were the less efficient phosphate transfers to CDP and UDP (Fig. 2 and Table 2). Inasmuch as no NTP spot was detected in reactions without NDP (Fig. 2A), it seems likely that the phosphate group transfer from polyP to the NDPs was direct. Most significantly, membrane fractions of the *ppk ppx* deletion mutant showed no polyP-dependent NDK activity (Fig. 2A).

PPK Transfers Pyrophosphate Group from polyP to GDP. Among the products of the PPK reaction of polyP with GDP was a compound more highly phosphorylated than GTP (Fig. 1). The ratio of GTP to the new spot was $\approx 10:1$. By two-dimensional TLC (Fig. 3), the radioactive spots corresponded to the positions of GTP and ppppG. Inasmuch as PPK failed to make ppppG from GTP and polyP (data not shown), it




FIG. 2. polyP-dependent NDK activity in the *E. coli* membrane fraction. The reaction mixture (10 μ l) containing the membrane fraction, PPK buffer, 1 mM NDP, and 0.1 mM [32 P]polyP was incubated at 37°C for 40 min. *E. coli* membrane fractions from MG1655 (wild type; 1.6 μ g) and $\Delta ppk ppx$ mutant (2.1 μ g) were used. The TLC plates were developed with (A) 1.5 M KH_2PO_4 (pH 3.5) and (B) 0.75 M KH_2PO_4 (pH 3.5).




FIG. 3. Autoradiogram of the products of PPK action on [³²P]polyP and GDP. Unlabeled GTP, guanosine 5'-diphosphate 3'-diphosphate (ppGpp), and ppppG were visualized by UV (outlined). The two-dimensional TLC analysis is described in *Materials and Methods*.

seems likely that a pyrophosphate group was transferred to GDP from polyP. Yeast PPX (20) cleaved ppppG to P_i and GTP (Fig. 4), as it was known to hydrolyze ppppA to P_i and ATP (S.-J. Liu, personal communication). *E. coli* PPX (21) also degraded ppppG, yet both the yeast and *E. coli* enzymes share no amino acid homologies (data not shown).

polyP-Dependent NDK Activity in *P. aeruginosa*. Since PPK activity is abundant and membrane attached in *P. aeruginosa* as in *E. coli*, it was of immediate interest to determine whether here too the PPK could use NDPs other than ADP as substrates. Membrane fractions of *P. aeruginosa* PAO1 possessed a polyP-dependent NDP kinase activity (Fig. 5 and Table 2), with a relative specificity for GDP almost 10 times higher than that of *E. coli* PPK (Table 2). GDP was phosphorylated in the presence of ADP as well as in its absence (Fig. 5). Because of the need for GTP in the major build up of alginate (22) and the concomitant accumulation of polyP at this stage (H.-Y. Kim and A. Kornberg,



FIG. 4. Degradation of ppppG by PPX. α -[³²P]ppppG (20 pmol) was digested by yeast PPX (500 units) at 37°C for 60 min. The TLC plate was developed with 1.5 M KH₂PO₄ (pH 3.5) and autoradiographed.





FIG. 5. NDK activity of *P. aeruginosa* membranes. The reaction mixture (10 μ l) containing *P. aeruginosa* membrane fraction (0.73 μ g), PPK buffer, 1 mM NDP, and 0.1 mM [³²P]polyP was incubated at 37°C for 60 min. The TLC plate was developed with 0.75 M KH₂PO₄ (pH 3.5) and autoradiographed.

unpublished data), the capacity of PPK to regenerate GTP from polyP may supply an important function.

DISCUSSION

In this report, we demonstrate that PPK can transfer a phosphate group from polyP to NDP. Thus PPK may provide an NDK-like activity:



Unlike NDK, the reaction catalyzed by PPK is not reversible, since PPK can synthesize polyP only from ATP. NDP kinase activity of PPK in *E. coli* lysates is only 0.06–0.3 nmol/min per mg of protein, a value only 0.01% that of NDK (10). Although the contribution of PPK as a NDP kinase may seem minor, the membrane localization of PPK may be significant, since levels of polyP can reach 20 mM in the stationary phase. A mutant of *E. coli* that lacks NDK grows at a normal rate but shows an elevated mutation rate (11). Tracking the residual NDP kinase activity in the mutant revealed adenylate kinase to have this activity (12). Surprisingly, adenylate kinase can transfer phosphate to an NDP but with 1–5% the specific activity of NDK (12).

Although ammonium sulfate increased the activity of *E. coli* PPK in polyP synthesis from ATP (18), it alters the preferential order of NDPs as substrates in the NDK reaction. Without ammonium sulfate present, the preference for ADP is markedly reduced (Table 2). Ammonium sulfate increased the *k*_{cat} value for ADP 20-fold (data not shown), while Ca²⁺ ion appeared to reduce the preference for ADP. The strong chelation of Ca²⁺ by polyP and the role of Ca²⁺-polyP combined with poly- β -hydroxybutyrate in membrane complexes

(23) directs more attention to Ca^{2+} and its influence on PPK activities.

With polyP and GDP as substrates, the predominant reaction catalyzed by PPK was the transfer of a phosphate group to synthesize GTP. About 10% of the time, PPK produced ppppG. Inasmuch as the ppppG was not produced from GTP, we presume that a pyrophosphate was transferred to GDP. A highly phosphorylated compound (likely inosine 5'-tetraphosphate) was also detected in the reaction of IDP and polyP, but nucleoside 5'-tetraphosphate was hardly detected in reactions with ADP, CDP, or UDP. The absence of ppppG among the products of PPK activity in the crude membrane fraction may be due to the presence of PPX, which degrades ppppG. Other known sources of nucleoside 5-tetraphosphates are yeast phosphoglycerate kinase, which forms adenosine 5'-tetraphosphate (24) and ppppG (25) from 1,3-bis-phosphoglycerate, and ATP or GTP. The physiological role of tetraphosphates is unknown. Adenosine, guanosine, and uridine tetraphosphates are strong competitive inhibitors, with nanomolar K_i values, of dinucleoside tetraphosphatase (26), which cleaves dinucleoside tetraphosphates to nucleoside tri- and monophosphates. Should diadenosine tetraphosphate prove to have important roles in metabolic regulation, changes in the level of ppppN could modulate its concentration and physiological effects. Another enzyme known to be inhibited by nucleotide 5'-tetraphosphates, with a micromolar K_i value, is the soluble guanylate cyclase (27).

The finding of a polyP-dependent NDP kinase activity in *P. aeruginosa* as well as in *E. coli* may suggest that this property is a general feature of PPK. The specificity for GDP of the *P. aeruginosa* PPK was almost 10 times higher than that of *E. coli*. The need for *P. aeruginosa* to produce a large amount of alginate dependent on GTP (by way of the alginate precursor, GDP-mannose; ref. 22) and the large accumulations of polyP at this stage of the life cycle suggest that an NDK role for PPK may prove significant.

Drs. M. Cashel and A. Chakrabarty generously provided strains. We thank Dr. N. N. Rao for technical help and LeRoy Bertsch for critical reading of this manuscript. A. Kuroda is the recipient of a fellowship from the Human Frontier Science Program. This work was supported by a grant from the National Institutes of Health.

- Biggs, J., Hersperger, E., Steeg, P. S., Liotta, L. A. & Shearn, A. (1990) *Cell* **63**, 933-940.
- Munoz-Dorado, J., Inouye, M. & Inouye, S. (1990) *J. Biol. Chem.* **265**, 2702-2706.
- Wallet, V., Mutzel, R., Troll, H., Barzu, O., Wurster, B., Veron, M. & Lacombe, M. L. (1990) *J. Natl. Cancer Inst.* **82**, 1199-1202.
- Okabe-Kado, J., Kasukabe, T., Hozumi, M., Honma, Y., Kimura, N., Baba, H., Urano, T. & Shiku, H. (1995) *FEBS Lett.* **363**, 311-315.
- Venturelli, D., Martinez, R., Melotti, P., Casella, I., Peschle, C., Cucco, C., Spampinato, G., Darzynkiewicz, Z. & Calabretta, B. (1995) *Proc. Natl. Acad. Sci. USA* **92**, 7435-7439.
- Bominaar, A. A., Molijin, A. C., Pestel, M., Veron, M. & Van, H. P. J. (1993) *EMBO J.* **12**, 2275-2279.
- Izumiya, H. & Yamamoto, M. (1995) *J. Biol. Chem.* **270**, 27859-27864.
- Steeg, P. S., Bevilacqua, G., Kopper, L., Thorgeirsson, U. P., Talmadge, J. E., Liotta, L. A. & Sobel, M. E. (1988) *J. Natl. Cancer Inst.* **80**, 200-204.
- Rosengard, A. M., Krutzsch, H. C., Shearn, A., Biggs, J. R., Barker, E., Margulies, I. M., King, C. R., Liotta, L. A. & Steeg, P. S. (1989) *Nature (London)* **342**, 177-180.
- Parks, R. E. & Agarwal, R. P. (1973) in *The Enzymes*, ed. Boyer, P. D. (Academic, New York), pp. 307-333.
- Lu, Q., Zhang, X., Almula, N., Mathews, C. K. & Inouye, M. (1995) *J. Mol. Biol.* **254**, 337-341.
- Lu, Q. & Inouye, M. (1996) *Proc. Natl. Acad. Sci. USA* **93**, 5720-5725.
- Kornberg, A. (1995) *J. Bacteriol.* **177**, 491-496.
- Kulaev, I. S. (1979) *The Biochemistry of Inorganic Polyphosphates* (Wiley, New York).
- Clark, J. E., Beegen, H. & Wood, H. G. (1986) *J. Bacteriol.* **168**, 1212-1219.
- Skorko, R. (1989) *Biochimie* **71**, 1089-1093.
- Bonting, C. F., Kortstee, G. J. & Zehnder, A. J. (1991) *J. Bacteriol.* **173**, 6484-6488.
- Ahn, K. & Kornberg, A. (1990) *J. Biol. Chem.* **265**, 11734-11739.
- Bochner, B. R. & Ames, B. N. (1982) *J. Biol. Chem.* **257**, 9759-9769.
- Wurst, H. & Kornberg, A. (1994) *J. Biol. Chem.* **269**, 10996-11001.
- Akiyama, M., Crooke, E. & Kornberg, A. (1993) *J. Biol. Chem.* **268**, 633-639.
- May, T. B. & Chakrabarty, A. M. (1994) *Trends Microbiol.* **2**, 151-157.
- Castuma, C. E., Huang, R., Kornberg, A. & Reusch, R. N. (1995) *J. Biol. Chem.* **270**, 12980-12983.
- Small, G. D. & Cooper, C. (1966) *Biochemistry* **5**, 26-33.
- Garcia-Diaz, M., Canales, J., Sillero, M. A., Sillero, A. & Cameselle, J. C. (1989) *Biochem. Int.* **19**, 1253-1264.
- Moreno, A., Lobaton, C. D., Sillero, M. A. G. & Sillero, A. (1982) *Int. J. Biochem.* **14**, 629-634.
- Ignarro, L. J., Gross, R. A. & Gross, D. M. (1976) *J. Cyclic Nucleotide Res.* **2**, 337-346.