

Making Health Care Safer: A Critical Analysis of Patient Safety Practices

Prepared for:

Agency for Healthcare Research and Quality
U.S. Department of Health and Human Services
2101 East Jefferson Street
Rockville, MD 20852
www.ahrq.gov

Contract No. 290-97-0013

Prepared by:

University of California at San Francisco (UCSF)–Stanford University
Evidence-based Practice Center

Editorial Board

Kaveh G. Shojania, M.D. (UCSF)
Bradford W. Duncan, M.D. (Stanford)
Kathryn M. McDonald, M.M. (Stanford)
Robert M. Wachter, M.D. (UCSF)

Managing Editor

Amy J. Markowitz, JD

Robert M. Wachter, MD

Project Director

Kathryn M. McDonald, MM
UCSF–Stanford EPC Coordinator

AHRQ Publication 01-E058

July 20, 2001 (Revised printing, indexed)

Preface

The Agency for Healthcare Research and Quality (AHRQ), formerly the Agency for Health Care Policy and Research (AHCPR), through its Evidence-based Practice Centers (EPCs), sponsors the development of evidence reports and technology assessments to assist public- and private-sector organizations in their efforts to improve the quality of health care in the United States. The reports and assessments provide organizations with comprehensive, science-based information on common, costly medical conditions and new health care technologies. The EPCs systematically review the relevant scientific literature on topics assigned to them by AHRQ and conduct additional analyses when appropriate prior to developing their reports and assessments.

To bring the broadest range of experts into the development of evidence reports and health technology assessments, AHRQ encourages the EPCs to form partnerships and enter into collaborations with other medical and research organizations. The EPCs work with these partner organizations to ensure that the evidence reports and technology assessments they produce will become building blocks for health care quality improvement projects throughout the Nation. The reports undergo peer review prior to their release.

AHRQ expects that the EPC evidence reports and technology assessments will inform individual health plans, providers, and purchasers as well as the health care system as a whole by providing important information to help improve health care quality.

We welcome written comments on this evidence report. They may be sent to: Director, Center for Practice and Technology Assessment, Agency for Healthcare Research and Quality, 6010 Executive Blvd., Suite 300, Rockville, MD 20852.

John M. Eisenberg, M.D.
Director
Agency for Healthcare
Research and Quality

Douglas B. Kamerow, M.D.
Director, Center for Practice and
Technology Assessment
Agency for Healthcare
Research and Quality

The authors of this report are responsible for its content. Statements in the report should not be construed as endorsement by the Agency for Healthcare Research and Quality or the U.S. Department of Health and Human Services of a particular drug, device, test, treatment, or other clinical service.

Structured Abstract

Objectives: Patient safety has received increased attention in recent years, but mostly with a focus on the epidemiology of errors and adverse events, rather than on practices that reduce such events. This project aimed to collect and critically review the existing evidence on practices relevant to improving patient safety.

Search Strategy and Selection Criteria: Patient safety practices were defined as those that reduce the risk of adverse events related to exposure to medical care across a range of diagnoses or conditions. Potential patient safety practices were identified based on preliminary surveys of the literature and expert consultation. This process resulted in the identification of 79 practices for review. The practices focused primarily on hospitalized patients, but some involved nursing home or ambulatory patients. Protocols specified the inclusion criteria for studies and the structure for evaluation of the evidence regarding each practice. Pertinent studies were identified using various bibliographic databases (e.g., MEDLINE, PsycINFO, ABI/INFORM, INSPEC), targeted searches of the Internet, and communication with relevant experts.

Data Collection and Analysis: Included literature consisted of controlled observational studies, clinical trials and systematic reviews found in the peer-reviewed medical literature, relevant non-health care literature and “gray literature.” For most practices, the project team required that the primary outcome consist of a clinical endpoint (i.e., some measure of morbidity or mortality) or a surrogate outcome with a clear connection to patient morbidity or mortality. This criterion was relaxed for some practices drawn from the non-health care literature. The evidence supporting each practice was summarized using a prospectively determined format. The project team then used a predefined consensus technique to rank the practices according to the strength of evidence presented in practice summaries. A separate ranking was developed for research priorities.

Main Results: Practices with the strongest supporting evidence are generally clinical interventions that decrease the risks associated with hospitalization, critical care, or surgery. Many patient safety practices drawn primarily from nonmedical fields (e.g., use of simulators, bar coding, computerized physician order entry, crew resource management) deserve additional research to elucidate their value in the health care environment. The following 11 practices were rated most highly in terms of strength of the evidence supporting more widespread implementation.

- Appropriate use of prophylaxis to prevent venous thromboembolism in patients at risk;
- Use of perioperative beta-blockers in appropriate patients to prevent perioperative morbidity and mortality;
- Use of maximum sterile barriers while placing central intravenous catheters to prevent infections;

- Appropriate use of antibiotic prophylaxis in surgical patients to prevent postoperative infections;
- Asking that patients recall and restate what they have been told during the informed consent process;
- Continuous aspiration of subglottic secretions (CASS) to prevent ventilator-associated pneumonia;
- Use of pressure relieving bedding materials to prevent pressure ulcers;
- Use of real-time ultrasound guidance during central line insertion to prevent complications;
- Patient self-management for warfarin (Coumadin™) to achieve appropriate outpatient anticoagulation and prevent complications;
- Appropriate provision of nutrition, with a particular emphasis on early enteral nutrition in critically ill and surgical patients; and
- Use of antibiotic-impregnated central venous catheters to prevent catheter-related infections.

Conclusions: An evidence-based approach can help identify practices that are likely to improve patient safety. Such practices target a diverse array of safety problems. Further research is needed to fill the substantial gaps in the evidentiary base, particularly with regard to the generalizability of patient safety practices heretofore tested only in limited settings and to promising practices drawn from industries outside of health care.

This document is in the public domain and may be used and reprinted without permission except those copyrighted materials noted for which further reproduction is prohibited without the specific permission of copyright holders.

Suggested Citation:

Shojania KG, Duncan BW, McDonald KM, et al., eds. Making Health Care Safer: A Critical Analysis of Patient Safety Practices. Evidence Report/Technology Assessment No. 43 (Prepared by the University of California at San Francisco–Stanford Evidence-based Practice Center under Contract No. 290-97-0013), AHRQ Publication No. 01-E058, Rockville, MD: Agency for Healthcare Research and Quality. July 2001.

CONTENTS

Summary	1
Evidence Report	9
PART I. OVERVIEW	11
Chapter 1. An Introduction to the Report	13
1.1. General Overview	13
1.2. How to Use this Report	18
1.3. Acknowledgments	20
Chapter 2. Drawing on Safety Practices from Outside Health Care	25
Chapter 3. Evidence-based Review Methodology	29
PART II. REPORTING AND RESPONDING TO PATIENT SAFETY PROBLEMS	39
Chapter 4. Incident Reporting	41
Chapter 5. Root Cause Analysis	51
PART III. PATIENT SAFETY PRACTICES & TARGETS	57
Section A. Adverse Drug Events (ADEs)	58
Chapter 6. Computerized Physician Order Entry (CPOE) with Clinical Decision Support Systems (CDSSs)	59
Chapter 7. The Clinical Pharmacist’s Role in Preventing Adverse Drug Events	71
Chapter 8. Computer Adverse Drug Event (ADE) Detection and Alerts	79
Chapter 9. Protocols for High-Risk Drugs: Reducing Adverse Drug Events Related to Anticoagulants	87
Chapter 10. Unit-Dose Drug Distribution Systems	101
Chapter 11. Automated Medication Dispensing Devices	111
Section B. Infection Control	118
Chapter 12. Practices to Improve Handwashing Compliance	119
Chapter 13. Impact of Barrier Precautions in Reducing the Transmission of Serious Nosocomial Infections	127
Chapter 14. Impact of Changes in Antibiotic Use Practices on Nosocomial Infections and Antimicrobial Resistance – <i>Clostridium Difficile</i> and Vancomycin-resistant Enterococcus (VRE)	141
Chapter 15. Prevention of Nosocomial Urinary Tract Infections	149
Subchapter 15.1. Use of Silver Alloy Urinary Catheters	149
Subchapter 15.2. Use of Suprapubic Catheters	155
Chapter 16. Prevention of Intravascular Catheter-Associated Infections	163
Subchapter 16.1. Use of Maximum Barrier Precautions During Central Venous Catheter Insertion	165
Subchapter 16.2. Use of Central Venous Catheters Coated with Antibacterial or Antiseptic Agents	169
Subchapter 16.3. Use of Chlorhexidine Gluconate at the Central Venous Catheter Insertion Site	175
Subchapter 16.4. Other Practices	178

Chapter 17. Prevention of Ventilator-Associated Pneumonia.....	185
Subchapter 17.1. Patient Positioning: Semi-recumbent Positioning and Continuous Oscillation	185
Subchapter 17.2. Continuous Aspiration of Subglottic Secretions	190
Subchapter 17.3. Selective Digestive Tract Decontamination	193
Subchapter 17.4. Sucralfate and Prevention of VAP	198
Section C. Surgery, Anesthesia, and Perioperative Medicine	204
Chapter 18. Localizing Care to High-Volume Centers	205
Chapter 19. Learning Curves for New Procedures – the Case of Laparoscopic Cholecystectomy	213
Chapter 20. Prevention of Surgical Site Infections	221
Subchapter 20.1. Prophylactic Antibiotics.....	221
Subchapter 20.2. Perioperative Normothermia.....	231
Subchapter 20.3. Supplemental Perioperative Oxygen.....	234
Subchapter 20.4. Perioperative Glucose Control	237
Chapter 21. Ultrasound Guidance of Central Vein Catheterization	245
Chapter 22. The Retained Surgical Sponge.....	255
Chapter 23. Pre-Anesthesia Checklists To Improve Patient Safety	259
Chapter 24. The Impact Of Intraoperative Monitoring On Patient Safety	265
Chapter 25. Beta-blockers and Reduction of Perioperative Cardiac Events	271
Section D. Safety Practices for Hospitalized or Institutionalized Elders.....	280
Chapter 26. Prevention of Falls in Hospitalized and Institutionalized Older People.....	281
Subchapter 26.1. Identification Bracelets for High-Risk Patients	284
Subchapter 26.2. Interventions that Decrease the Use of Physical Restraints	287
Subchapter 26.3. Bed Alarms.....	291
Subchapter 26.4. Special Hospital Flooring Materials to Reduce Injuries from Patient Falls	293
Subchapter 26.5. Hip Protectors to Prevent Hip Fracture	295
Chapter 27. Prevention of Pressure Ulcers in Older Patients	301
Chapter 28. Prevention of Delirium in Older Hospitalized Patients	307
Chapter 29. Multidisciplinary Geriatric Consultation Services.....	313
Chapter 30. Geriatric Evaluation and Management Units for Hospitalized Patients	323
Section E. General Clinical Topics.....	331
Chapter 31. Prevention of Venous Thromboembolism	333
Chapter 32. Prevention of Contrast-Induced Nephropathy	349
Chapter 33. Nutritional Support	359
Chapter 34. Prevention of Clinically Significant Gastrointestinal Bleeding in Intensive Care Unit Patients	368
Chapter 35. Reducing Errors in the Interpretation of Plain Radiographs and Computed Tomography Scans	376
Chapter 36. Pneumococcal Vaccination Prior to Hospital Discharge	385
36.1. Vaccine Effectiveness	386
36.2. Vaccine Delivery Methods.....	388

Chapter 37. Pain Management.....	396
Subchapter 37.1. Use of Analgesics in the Acute Abdomen	396
Subchapter 37.2. Acute Pain Services.....	400
Subchapter 37.3. Prophylactic Antiemetics During Patient-controlled Analgesia Therapy	404
Subchapter 37.4. Non-pharmacologic Interventions for Postoperative Plan	406
Section F. Organization, Structure, and Culture	411
Chapter 38. “Closed” Intensive Care Units and Other Models of Care for Critically Ill Patients.....	413
Chapter 39. Nurse Staffing, Models of Care Delivery, and Interventions	423
Chapter 40. Promoting a Culture of Safety	447
Section G. Systems Issues and Human Factors	458
Chapter 41. Human Factors and Medical Devices	459
Subchapter 41.1. The Use of Human Factors in Reducing Device-related Medical Errors	459
Subchapter 41.2. Refining the Performance of Medical Device Alarms	462
Subchapter 41.3. Equipment Checklists in Anesthesia	466
Chapter 42. Information Transfer.....	471
Subchapter 42.1. Information Transfer Between Inpatient and Outpatient Pharmacies ..	471
Subchapter 42.2. Sign-Out Systems for Cross-Coverage	476
Subchapter 42.3. Discharge Summaries and Follow-up	479
Subchapter 42.4. Notifying Patients of Abnormal Results	482
Chapter 43. Prevention of Misidentifications.....	487
Subchapter 43.1. Bar Coding	487
Subchapter 43.2. Strategies to Avoid Wrong-Site Surgery.....	494
Chapter 44. Crew Resource Management and its Applications in Medicine	501
Chapter 45. Simulator-Based Training and Patient Safety.....	511
Chapter 46. Fatigue, Sleepiness, and Medical Errors.....	519
Chapter 47. Safety During Transport of Critically Ill Patients.....	534
Subchapter 47.1. Interhospital Transport	536
Subchapter 47.2. Intrahospital Transport	538
Section H. Role of the Patient.....	544
Chapter 48. Procedures For Obtaining Informed Consent	546
Chapter 49. Advance Planning For End-of-Life Care	556
Chapter 50. Other Practices Related to Patient Participation	569
PART IV. PROMOTING AND IMPLEMENTING SAFETY PRACTICES	573
Chapter 51. Practice Guidelines	575
Chapter 52. Critical Pathways	581
Chapter 53. Clinical Decision Support Systems.....	589
Chapter 54. Educational Techniques Used in Changing Provider Behavior.....	595
Chapter 55. Legislation, Accreditation, and Market-Driven and Other Approaches to Improving Patient Safety	601

PART V. ANALYZING THE PRACTICES611
Chapter 56. Methodology for Summarizing the Evidence for the Practices613
Chapter 57. Practices Rated by Strength of Evidence619
Chapter 58. Practices Rated by Research Priority627
Chapter 59. Listing of All Practices, Categorical Ratings, and Comments.....633
Appendix: List of Contributors649
Index655

Summary

Overview

Patient safety has become a major concern of the general public and of policymakers at the State and Federal levels. This interest has been fueled, in part, by news coverage of individuals who were the victims of serious medical errors and by the publication in 1999 of the Institute of Medicine's (IOM's) report *To Err is Human: Building a Safer Health System*. In its report, IOM highlighted the risks of medical care in the United States and shocked the sensibilities of many Americans, in large part through its estimates of the magnitude of medical-errors-related deaths (44,000 to 98,000 deaths per year) and other serious adverse events. The report prompted a number of legislative and regulatory initiatives designed to document errors and begin the search for solutions. But Americans, who now wondered whether their next doctor's or hospital visit might harm rather than help them, began to demand concerted action.

Three months after publication of the IOM report, an interagency Federal government group, the Quality Interagency Coordination Task Force (QuIC), released its response, *Doing What Counts for Patient Safety: Federal Actions to Reduce Medical Errors and Their Impact*. That report, prepared at the President's request, both inventoried on-going Federal actions to reduce medical errors and listed more than 100 action items to be undertaken by Federal agencies.

An action promised by the Agency for Healthcare Research and Quality (AHRQ), the Federal agency leading efforts to research and promote patient safety, was "the development and dissemination of evidence-based, best safety practices to provider organizations." To initiate the work to be done in fulfilling this promise, AHRQ commissioned the University of California at San Francisco (UCSF) – Stanford University Evidence-based Practice Center (EPC) in January 2001 to review the scientific literature regarding safety improvement. To accomplish this, the EPC established an Editorial Board that oversaw development of this report by teams of content experts who served as authors.

Defining Patient Safety Practices

Working closely with AHRQ and the National Forum for Quality Measurement and Reporting (the National Quality Forum, or NQF)—a public-private partnership formed in 1999 to promote a national health care quality agenda the EPC began its work by defining a patient safety practice as *a type of process or structure whose application reduces the probability of adverse events resulting from exposure to the health care system across a range of diseases and procedures*.

This definition is consistent with the dominant conceptual framework in patient safety, which holds that systemic change will be far more productive in reducing medical errors than will targeting and punishing individual providers. The definition's focus on actions that cut across diseases and procedures also allowed the research team to distinguish patient safety activities from the more targeted quality improvement practices (e.g., practices designed to increase the use of beta-blockers in patients who are admitted to the hospital after having a myocardial infarction). The editors recognize, however, that this distinction is imprecise.

This evidence-based review also focuses on hospital care as a starting point because the risks associated with hospitalization are significant, the strategies for improvement are better documented there than in other health care settings, and the importance of patient trust is paramount. The report, however, also considers evidence regarding other sites of care, such as nursing homes, ambulatory care, and patient self-management.

The results of this EPC study will be used by the NQF to identify a set of proven patient safety practices that should be used by hospitals. Identification of these practices by NQF will allow patients throughout the nation to evaluate the actions their hospitals and/or health care facilities have taken to improve safety.

Reporting The Evidence

As is typical for evidence-based reviews, the goal was to provide a critical appraisal of the evidence on the topic. This information would then be available to others to ensure that no practice unsupported by evidence would be endorsed and that no practice substantiated by a high level of proof would lack endorsement. Readers familiar with the state of the evidence regarding quality improvement in areas of health care where this has been a research priority (e.g., cardiovascular care) may be surprised and even disappointed, by the paucity of high quality evidence in other areas of health care for many patient safety practices. One reason for this is the relative youth of the field. Just as there had been little public recognition of the risks of health care prior to the first IOM report, there has been relatively little attention paid to such risks – and strategies to mitigate them – among health professionals and researchers.

Moreover, there are a number of methodologic reasons why research in patient safety is particularly challenging. Many practices (e.g., the presence of computerized physician order entry systems, modifying nurse staffing levels) cannot be the subject of double-blind studies because their use is evident to the participants. Second, capturing all relevant outcomes, including “near misses” (such as a nurse catching an excessive dosage of a drug just before it is administered to a patient) and actual harm, is often very difficult. Third, many effective practices are multidimensional, and sorting out precisely which part of the intervention works is often quite challenging. Fourth, many of the patient safety problems that generate the most concern (wrong-site surgery, for example) are uncommon enough that demonstrating the success of a “safety practice” in a statistically meaningful manner with respect to outcomes is all but impossible.

Finally, establishing firm epidemiologic links between presumed (and accepted) causes and adverse events is critical, and frequently difficult. For instance, in studying an intuitively plausible “risk factor” for errors, such as “fatigue,” analyses of errors commonly reveal the presence of fatigued providers (because many health care providers work long hours and/or late at night). The question is whether or not fatigue is over-represented among situations that lead to errors. The point is not that the problem of long work-hours should be ignored, but rather that strong epidemiologic methods need to be applied before concluding that an intuitive cause of errors is, in fact, causal.

Researchers now believe that most medical errors cannot be prevented by perfecting the technical work of individual doctors, nurses, or pharmacists. Improving patient safety often involves the coordinated efforts of multiple members of the health care team, who may adopt strategies from outside health care. The report reviews several practices whose evidence came

from the domains of commercial aviation, nuclear safety, and aerospace, and the disciplines of human factors engineering and organizational theory. Such practices include root cause analysis, computerized physician order entry and decision support, automated medication dispensing systems, bar coding technology, aviation-style preoperative checklists, promoting a “culture of safety,” crew resource management, the use of simulators in training, and integrating human factors theory into the design of medical devices and alarms. In reviewing these practices, the research team sought to be flexible regarding standards of evidence, and included research evidence that would not have been considered for medical interventions. For example, the randomized trial that is appropriately hailed as the “gold standard” in clinical medicine is not used in aviation, as this design would not capture all relevant information. Instead, detailed case studies and industrial engineering research approaches are utilized.

Methodology

To facilitate identification and evaluation of potential patient safety practices, the Editorial Board divided the content for the project into different *domains*. Some cover “content areas,” including traditional clinical areas such as adverse drug events, nosocomial infections, and complications of surgery, but also less traditional areas such as fatigue and information transfer. Other domains consist of practices drawn from broad (primarily nonmedical) disciplines likely to contain promising approaches to improving patient safety (e.g., information technology, human factors research, organizational theory). Once this list was created—with significant input from patient safety experts, clinician–researchers, AHRQ, and the NQF Safe Practices Committee—the editors selected teams of authors with expertise in the relevant subject matter and/or familiarity with the techniques of evidence-based review and technology appraisal.

The authors were given explicit instructions regarding search strategies for identifying safety practices for evaluation (including explicit inclusion and exclusion criteria) and criteria for assessing each practice’s level of evidence for efficacy or effectiveness in terms of study design and study outcomes. Some safety practices did not meet the inclusion criteria because of the paucity of evidence regarding efficacy or effectiveness but were included in the report because an informed reader might reasonably expect them to be evaluated or because of the depth of public and professional interest in them. For such high profile topics (such as bar coding to prevent misidentifications), the researchers tried to fairly present the practice’s background, the experience with the practice thus far, and the evidence (and gaps in the evidence) regarding the practice’s value.

For each practice, authors were instructed to research the literature for information on:

- prevalence of the problem targeted by the practice;
- severity of the problem targeted by the practice;
- the current utilization of the practice;
- evidence on efficacy and/or effectiveness of the practice;
- the practice’s potential for harm;
- data on cost, if available; and
- implementation issues.

The report presents the salient elements of each included study (e.g., study design, population/setting, intervention details, results), and highlights any important weaknesses and

biases of these studies. Authors were *not* asked to formally synthesize or combine the evidence across studies (e.g., perform a meta-analysis) as part of their task.

The Editorial Board and the Advisory Panel reviewed the list of domains and practices to identify gaps in coverage. Submitted chapters were reviewed by the Editorial Board and revised by the authors, aided by feedback from the Advisory Panel. Once the content was finalized, the editors analyzed and ranked the practices using a methodology summarized below.

Summarizing the Evidence and Rating the Practices

Because the report is essentially an anthology of a diverse and extensive group of patient safety practices with highly variable relevant evidence, synthesizing the findings was challenging, but necessary to help readers use the information. Two of the most obvious uses for this report are: 1) to inform efforts of providers and health care organizations to improve the safety of the care they provide, and 2) to inform AHRQ, other research agencies, and foundations about potential fruitful investments for their research support. Other uses of the information are likely. In fact, the National Quality Forum plans to use this report to help identify a list of patient safety practices that consumers and others should know about as they choose among the health care provider organizations to which they have access.

In an effort to assist both health care organizations interested in taking substantive actions to improve patient safety and research funders seeking to spend scarce resources wisely, AHRQ asked the EPC to rate the evidence and rank the practices by opportunity for safety improvement and by research priority. This report, therefore, contains two lists.

To create these lists, the editors aimed to separate the practices that are most promising or effective from those that are least so on a range of dimensions, without implying any ability to calibrate a finely gradated scale for those practices in between. The editors also sought to present the ratings in an organized, accessible way while highlighting the limitations inherent in their rating schema. Proper metrics for more precise comparisons (e.g., cost-effectiveness analysis) require more data than are currently available in the literature.

Three major categories of information were gathered to inform the rating exercise:

- *Potential Impact of the Practice*: based on prevalence and severity of the patient safety target, and current utilization of the practice;
- *Strength of the Evidence Supporting the Practice*: including an assessment of the relative weight of the evidence, effect size, and need for vigilance to reduce any potential negative collateral effects of the practice; and
 - *Implementation*: considering costs, logistical barriers, and policy issues.

For all of these data inputs into the practice ratings, the primary goal was to find the best available evidence from publications and other sources. Because the literature has not been previously organized with an eye toward addressing each of these areas, most of the estimates could be improved with further research, and some are informed by only general and somewhat speculative knowledge. In the summaries, the editors have attempted to highlight those assessments made with limited data.

The four-person editorial team independently rated each of the 79 practices using general scores (e.g., High, Medium, Low) for a number of dimensions, including those italicized in the

section above. The editorial team convened for 3 days in June, 2001 to compare scores, discuss disparities, and come to consensus about ratings for each category.

In addition, each member of the team considered the totality of information on potential impact and support for a practice to score each of these factors on a 0 to 10 scale (creating a “Strength of the Evidence” list). For these ratings, the editors took the perspective of a leader of a large health care enterprise (e.g., a hospital or integrated delivery system) and asked the question, “If I wanted to improve patient safety at my institution over the next 3 years and resources were not a significant consideration, how would I grade this practice?” For this rating, the Editorial Board explicitly chose *not* to formally consider the difficulty or cost of implementation in the rating. Rather, the rating simply reflected the strength of the evidence regarding the effectiveness of the practice and the probable impact of its implementation on reducing adverse events related to health care exposure. If the patient safety target was rated as “High” impact and there was compelling evidence (i.e., “High” relative study strength) that a particular practice could significantly reduce (e.g., “Robust” effect size) the negative consequences of exposure to the health care system (e.g., hospital-acquired infections), raters were likely to score the practice close to 10. If the studies were less convincing, the effect size was less robust, or there was a need for a “Medium” or “High” degree of vigilance because of potential harms, then the rating would be lower.

At the same time, the editors also rated the usefulness of conducting more research on each practice, emphasizing whether there appeared to be questions that a research program might have a reasonable chance of addressing successfully (creating a “Research Priority” list). Here, they asked themselves, “If I were the leader of a large agency or foundation committed to improving patient safety, and were considering allocating funds to promote additional research, how would I grade this practice?” If there was a simple gap in the evidence that could be addressed by a research study or if the practice was multifaceted and implementation could be eased by determining the specific elements that were effective, then the research priority was high. (For this reason, some practices are highly rated on both the “Strength of the Evidence” and “Research Priority” lists.) If the area was one of high potential impact (i.e., large number of patients at risk for morbid or mortal adverse events) and a practice had been inadequately researched, then it would also receive a relatively high rating for research need. Practices might receive low research scores if they held little promise (e.g., relatively few patients are affected by the safety problem addressed by the practice *or* a significant body of knowledge already demonstrates the practice’s lack of utility). Conversely, a practice that was clearly effective, low cost, and easy to implement would not require further research and would also receive low research scores.

In rating both the strength of the evidence and the research priority, the purpose was not to report precise 0 to 10 scores, but to develop general “zones” or practice groupings. This is important because better methods are available for making comparative ratings *when the data inputs are available*. The relative paucity of the evidence dissuaded the editors from using a more precise, sophisticated, but ultimately unfeasible, approach.

Clear Opportunities for Safety Improvement

The following 11 patient safety practices were the most highly rated (of the 79 practices reviewed in detail in the full report and ranked in the Executive Summary Addendum, AHRQ Publication No. 01-E057b) in terms of strength of the evidence supporting more widespread implementation. Practices appear in descending order, with the most highly rated practices listed first. Because of the imprecision of the ratings, the editors did not further divide the practices, nor indicate where there were ties.

- Appropriate use of prophylaxis to prevent venous thromboembolism in patients at risk;
- Use of perioperative beta-blockers in appropriate patients to prevent perioperative morbidity and mortality;
- Use of maximum sterile barriers while placing central intravenous catheters to prevent infections;
- Appropriate use of antibiotic prophylaxis in surgical patients to prevent perioperative infections;
- Asking that patients recall and restate what they have been told during the informed consent process;
- Continuous aspiration of subglottic secretions (CASS) to prevent ventilator-associated pneumonia;
- Use of pressure relieving bedding materials to prevent pressure ulcers;
- Use of real-time ultrasound guidance during central line insertion to prevent complications;
- Patient self-management for warfarin (Coumadin™) to achieve appropriate outpatient anticoagulation and prevent complications;
- Appropriate provision of nutrition, with a particular emphasis on early enteral nutrition in critically ill and surgical patients; and
- Use of antibiotic-impregnated central venous catheters to prevent catheter-related infections.

This list is generally weighted toward clinical rather than organizational matters, and toward care of the very, rather than the mildly or chronically ill. Although more than a dozen practices considered were general safety practices that have been the focus of patient safety experts for decades (i.e., computerized physician order entry, simulators, creating a “culture of safety,” crew resource management), most research on patient safety has focused on more clinical areas. The potential application of practices drawn from outside health care has excited the patient safety community, and many such practices have apparent validity. However, clinical research has been promoted by the significant resources applied to it through Federal, foundation, and industry support. Since this study went where the evidence took it, more clinical practices rose to the top as potentially ready for implementation.

Clear Opportunities for Research

Until recently, patient safety research has had few champions, and even fewer champions with resources to bring to bear. The recent initiatives from AHRQ and other funders are a promising shift in this historical situation, and should yield important benefits.

In terms of the research agenda for patient safety, the following 12 practices rated most highly, as follows:

- Improved perioperative glucose control to decrease perioperative infections;
- Localizing specific surgeries and procedures to high volume centers;
- Use of supplemental perioperative oxygen to decrease perioperative infections;
- Changes in nursing staffing to decrease overall hospital morbidity and mortality;
- Use of silver alloy-coated urinary catheters to prevent urinary tract infections;
- Computerized physician order entry with computerized decision support systems to decrease medication errors and adverse events primarily due to the drug ordering process;
- Limitations placed on antibiotic use to prevent hospital-acquired infections due to antibiotic-resistant organisms;
- Appropriate use of antibiotic prophylaxis in surgical patients to prevent perioperative infections;
- Appropriate use of prophylaxis to prevent venous thromboembolism in patients at risk;
- Appropriate provision of nutrition, with a particular emphasis on early enteral nutrition in critically ill and post-surgical patients;
- Use of analgesics in the patient with an acutely painful abdomen without compromising diagnostic accuracy; and
- Improved handwashing compliance (via education/behavior change; sink technology and placement; or the use of antimicrobial washing substances).

Of course, the vast majority of the 79 practices covered in this report would benefit from additional research. In particular, some practices with longstanding success outside of medicine (e.g., promoting a culture of safety) deserve further analysis, but were not explicitly ranked due to their unique nature and the present weakness of the evidentiary base in the health care literature.

Conclusions

This report represents a first effort to approach the field of patient safety through the lens of evidence-based medicine. Just as *To Err is Human* sounded a national alarm regarding patient safety and catalyzed other important commentaries regarding this vital problem, this review seeks to plant a seed for future implementation and research by organizing and evaluating the relevant literature. Although all those involved tried hard to include all relevant practices and to review all pertinent evidence, inevitably some of both were missed. Moreover, the effort to grade and rank practices, many of which have only the beginnings of an evidentiary base, was admittedly ambitious and challenging. It is hoped that this report provides a template for future clinicians, researchers, and policy makers as they extend, and inevitably improve upon, this work.

In the detailed reviews of the practices, the editors have tried to define (to the extent possible from the literature) the associated costs—financial, operational, and political. However, these considerations were not factored into the summary ratings, nor were judgments made regarding the appropriate expenditures to improve safety. Such judgments, which involve complex tradeoffs between public dollars and private ones, and between saving lives by improving patient safety versus doing so by investing in other health care or non-health care practices, will obviously be critical. However, the public reaction to the IOM report, and the media and legislative responses that followed it, seem to indicate that Americans are highly concerned about the risks of medical errors and would welcome public and private investment to decrease them. It seems logical to infer that Americans value safety during a hospitalization just as highly as safety during a transcontinental flight.

Evidence Report

