

S U D A A N

Software for the Statistical Analysis of Correlated Data
Copyright Research Triangle Institute February

2005

Release 9.0.1

Number of observations read	:	19759	Weighted count	:278652243
Observations in subpopulation	:	9471	Weighted count:	198419259
Denominator degrees of freedom	:	29		

For Subpopulation: RIDAGEYR >= 20

Standardized estimates

Age-adjusted means & standard errors of body mass index: NHANES 1999-2002

by: Variable, Gender, Race Ethnicity.

for: Variable = Body Mass Index (kg/m**2).

Gender			
Race Ethnicity	Sample Size	Adjusted Mean	Standard Error
<hr/>			
Total			
Total	9064	28.02	0.134
NH-White	4384	27.77	0.163
NH-Black	1732	29.57	0.193
Mex-Am	2181	28.52	0.198
Other	767	27.67	0.334
Male			
Total	4262	27.84	0.130
NH-White	2103	27.92	0.168
NH-Black	809	27.53	0.185
Mex-Am	1018	28.01	0.189
Other	332	27.12	0.296
Female			
Total	4802	28.19	0.167
NH-White	2281	27.62	0.207
NH-Black	923	31.18	0.276
Mex-Am	1163	29.06	0.317
Other	435	28.09	0.425

Age adjusted by the direct method to the year 2000 Census population projections using the age groups 20-39, 40-59, and 60+