

S U D A A N

Software for the Statistical Analysis of Correlated Data
Copyright Research Triangle Institute February

2005

Release 9.0.1

Number of observations read	:	19759	Weighted count	:278652243
Observations in subpopulation	:	9471	Weighted count:	198419259
Denominator degrees of freedom	:	29		

Variance Estimation Method: Taylor Series (WR)

For Subpopulation: RIDAGEYR >= 20

Chi-square test for blood pressure cuff size: NHANES 1999-2002

by: Gender - Adjudicated., Coded cuff size.

Gender - Adjudicated.		Coded cuff size		
		Total	Infant/- Child	Adult
Total	Sample Size	9094	285	3452
	Row Percent	100.00	3.15	36.53
	Col Percent	100.00	100.00	100.00
Male	Sample Size	4312	63	1387
	Row Percent	100.00	1.50	28.77
	Col Percent	48.10	22.87	37.87
Female	Sample Size	4782	222	2065
	Row Percent	100.00	4.67	43.73
	Col Percent	51.90	77.13	62.13

Variance Estimation Method: Taylor Series (WR)

For Subpopulation: RIDAGEYR >= 20

Chi-square test for blood pressure cuff size: NHANES 1999-2002

by: Gender - Adjudicated., Coded cuff size.

Gender - Adjudicated.		Coded cuff size	
		Large	Thigh
Total	Sample Size	4411	946
	Row Percent	49.12	11.21
	Col Percent	100.00	100.00
Male	Sample Size	2409	453
	Row Percent	57.74	12.00
	Col Percent	56.54	51.50
Female	Sample Size	2002	493
	Row Percent	41.12	10.47
	Col Percent	43.46	48.50

Variance Estimation Method: Taylor Series (WR)

For Subpopulation: RIDAGEYR >= 20

Chi-square test for blood pressure cuff size: NHANES 1999-2002

Chi Square Test of Independence for Gender - Adjudicated, and Coded cuff size

	ChiSq	274.74
	P-value ChiSq	0.0000
	Degrees of	
	Freedom ChiSq	3
	LLChiSq	257.33
	P-value LLChiSq	0.0000
	Degrees of	
	Freedom LLChiSq	3
