

Table 94. Select Family Planning and Medical Service Use by Women, 15 to 44 Years of Age: 2002

[61,561 represents 61,561,000. Based on the National Survey of Family Growth, see Appendix III]

Characteristic	Number (1,000)	Percent using—							
		Family planning services			Medical services				
		At least one family planning service	Birth control method	Birth control checkup or test	At least one medical service	Pregnancy test	Pap smear	Pelvic exam	Counseling/test/treatment for STD ¹
All women²	61,561	41.7	33.9	23.6	69.1	19.7	64.4	59.7	12.6
15 to 19 years old	9,834	39.9	31.1	22.0	40.6	18.3	34.6	27.0	15.2
15 to 17 years old	5,819	31.8	22.2	15.8	28.0	11.4	23.2	17.9	11.1
18 to 19 years old	4,016	51.6	43.9	31.0	58.9	28.2	51.2	40.2	21.1
20 to 24 years old	9,840	63.3	54.0	35.7	75.7	31.5	69.7	60.6	22.3
25 to 29 years old	9,249	55.4	46.3	30.2	75.9	30.2	70.7	66.0	16.6
30 to 34 years old	10,272	47.0	39.1	27.2	78.1	22.2	72.7	69.7	12.2
35 to 39 years old	10,853	30.5	23.9	18.6	71.5	13.6	68.3	65.9	6.9
40 to 44 years old	11,512	19.5	14.0	10.8	71.8	5.9	69.4	67.0	4.4
Currently married	28,327	39.5	31.5	21.3	77.2	21.1	73.1	69.8	8.1
Currently cohabiting	5,570	50.4	43.2	30.2	77.2	31.0	72.2	64.7	20.3
Never married, not cohabiting	21,568	44.4	36.4	25.4	55.8	16.5	50.7	43.9	15.9
Formerly married, not cohabiting	6,096	34.5	28.0	22.0	71.0	14.1	66.0	64.0	14.4
White only, Non-Hispanic	39,498	43.1	36.4	25.4	70.1	17.3	66.1	63.2	11.8
Black only, Non-Hispanic	8,250	39.4	30.6	21.8	74.4	23.7	69.1	58.1	15.7
Hispanic ³	9,107	39.7	28.9	20.6	63.4	24.3	57.1	48.5	12.5

¹ STD stands for sexually transmitted disease. ² Includes other races not shown separately. ³ Persons of Hispanic origin may be any race.

Source: U.S. National Center for Health Statistics, *Advance Data*, Number 350, "Use of Contraception and Use of Family Planning Services in the United States: 1982–2002," December 10, 2004.