

Table 177. Reported AIDS Cases for Adults and Adolescents by Transmission Category and Sex: 2006

[Provisional. For cases reported in the year shown. Includes Puerto Rico, Virgin Islands, Guam, and U.S. Pacific Islands. Acquired immunodeficiency syndrome (AIDS) is a specific group of diseases or conditions which are indicative of severe immunosuppression related to infection with the human immunodeficiency virus (HIV). Data are subject to retrospective changes and may differ from those data in Table 175. For information on HIV death rates, go to Table 121]

| Transmission category | 2006 | | | Cumulative through 2006 ¹ | | |
|---|---------------|---------------|---------------|--------------------------------------|----------------|----------------|
| | Total | Male | Female | Total | Male | Female |
| Persons 13 years old and over, total | 38,916 | 28,378 | 10,537 | 983,343 | 791,626 | 191,714 |
| Male-to-male sexual contact | 13,775 | 13,775 | (X) | 429,897 | 429,897 | (X) |
| Injection drug use | 5,239 | 3,413 | 1,826 | 230,763 | 163,070 | 67,693 |
| Male-to-male sexual contact and injection drug use. . . . | 1,603 | 1,603 | (X) | 65,524 | 65,524 | (X) |
| Hemophilia/coagulation disorder | 54 | 43 | 11 | 5,524 | 5,176 | 348 |
| High-risk heterosexual contact. | 7,836 | 2,997 | 4,839 | 134,796 | 49,610 | 85,186 |
| Sex with injection drug user. | 1,036 | 330 | 706 | 37,756 | 11,683 | 26,073 |
| Sex with bisexual male. | 234 | (X) | 234 | 5,162 | (X) | 5,162 |
| Sex with person with hemophilia | 12 | 1 | 11 | 586 | 84 | 502 |
| Sex with HIV-infected person, transfusion recipient . . | 43 | 22 | 21 | 1,346 | 553 | 793 |
| Sex with HIV-infected person, risk factor not specified | 6,511 | 2,644 | 3,867 | 89,946 | 37,290 | 52,656 |
| Receipt of blood transfusion, blood components, or tissue | 131 | 63 | 68 | 9,389 | 5,213 | 4,176 |
| Other/risk not reported or identified | 10,278 | 6,484 | 3,793 | 107,450 | 73,136 | 34,311 |

X Not applicable. ¹ Includes persons with a diagnosis of AIDS, reported from the beginning of the epidemic through 2006. Cumulative total includes persons with characteristics unknown.

Source: U.S. Centers for Disease Control and Prevention, Atlanta, GA, *HIV/AIDS Surveillance Report, 2006*, Volume 18; <<http://www.cdc.gov/hiv/topics/surveillance/resources/reports/2006report/default.htm>> (revised 1 April 2008).