

Detection of Neuron-Specific Enolase in Formalin-Fixed, Paraffin-Embedded Rat Tissue

Reagent and Antibody Information

[1X Wash Buffer](#)

[3% Hydrogen Peroxide](#)

[1% BSA Diluent](#)

[1X Citrate Buffer](#)

[DAB Chromagen](#)

[Hematoxylin](#)

Primary Antibody: Mouse Anti-Neuron Specific Enolase Antibody

Lab Vision / Thermo Fisher Scientific

Fremont, CA 94539

www.labvision.com

1-800-828-1628

Catalog # MS-1717-P1

Negative Control Serum: Normal Mouse Serum

Jackson ImmunoResearch Laboratories, Inc.

West Grove, PA 19390

www.jacksonimmuno.com

1-800-367-5296

Catalog # 015-000-001

Staining Kit: LSAB 2 System-HRP

Dakocytomation Corporation

Carpinteria CA 93013

www.dakousa.com

1-800-235-5763

Code No. K0609

Note: This kit includes reagents needed for the secondary antibody (link) and label complex.

Staining Procedure

Positive Control Tissue: Islets of the pancreas
Stain Localization: Cytoplasmic / nucleat

1. Deparaffinize and hydrate slides through the following solutions:

Solution	Repetitions	Time
Xylene	2 times	5 minutes
100% Ethanol	2 times	3 minutes
95% Ethanol	2 times	3 minutes
1X Wash Buffer	2 times	5 minutes

2. Quench endogenous peroxidase by placing the slides in 3% hydrogen peroxide for 15 minutes.

3. Rinse the slides in 2 changes of 1X Wash Buffer for 5 minutes each.

4. Heat-Induced Epitope Retrieval Using The Decloaker

Add 500 ml of distilled water to the pan inside the decloaker.

Place a full rack of slides into a Tissue Tek® container with 200 ml of 1X citrate buffer

(Insert blank slides into any empty slots in the rack to ensure even heating of slides)

Place the container stably inside the pan and decloak for 5 minutes. *Maximum Pressure* _____

Depressurize for 10 minutes.

Remove pan top and cool for 10 minutes. *Temperature Before Cooling Slides* _____

Rinse the slides in 2 changes of distilled water for 3 minutes each time.

5. Rinse the slides in 2 changes of 1X Wash Buffer for 5 minutes each time.

6. Apply the primary antibody at a 1:800 dilution, and incubate for 10 minutes at room temperature.

Lot # _____ Exp Date _____

For negative control slides, dilute the protein concentration of the normal mouse serum to match that of the primary antibody. Make a 1:800 dilution from this normalized serum, and apply to the slides.

Incubate for 10 minutes at room temperature.

Lot # _____ Date Reconstituted _____

7. Rinse the slides in 2 changes of 1X Wash Buffer for 5 minutes each.

LSAB 2 Kit

Lot # _____ Exp Date _____

8. Apply the Link (yellow bottle) from the LSAB 2 Kit, and incubate for 10 minutes at room temperature.

9. Rinse the slides in 2 changes of 1X Wash Buffer for 5 minutes each.

10. Apply the Label (red bottle) from the LSAB 2 Kit, and incubate for 10 minutes at room temperature.

11. Rinse the slides in 2 changes of 1X Wash Buffer for 5 minutes each
12. Apply the DAB chromagen, and incubate in the dark for 6 minutes at room temperature.
(Add 1 drop of DAB per ml of substrate)
Lot #_____ Exp Date_____ New Kit: yes / no
13. Rinse the slides in tap water 3 minutes.
14. Counterstain with Harris Hematoxylin for 20 seconds.
15. Rinse the slides in tap water until water is clear.
16. Gently agitate slides in 1X Wash Buffer until they turn blue.
17. Dehydrate through the following solutions:

Solution	Repetitions	Time
95% Ethanol	1 change	3 minutes
100% Ethanol	3 changes	3 minutes
Xylene	2 changes	5 minutes

18. Coverslip

Updated 07/29/04