Detection of Progesterone Receptor in Formalin Fixed, Paraffin-Embedded Rat Tissue

Reagent and Antibody Information

1X Wash Buffer
3% Hydrogen Peroxide
1% BSA Diluent
1X Citrate Buffer
DAB Chromagen
Hematoxylin

Staining Kit: Vectastain Elite ABC Kit (Mouse IgG)
Vector Laboratories, Inc.
Burlingame, CA 94010
www.vectorlabs.com
1-800-227-6666
Catalog # PK-6102

Note: This kit contains all reagents necessary to make the blocking reagent, secondary antibody and label complex.

Avidin / Biotin Blocking Kit Vector Laboratories, Inc. Burlingame, CA 94010 www.vectorlabs.com 1-800-227-6666 Catalog # SP-2001

<u>Primary Antibody</u>: <u>Mouse Anti-Progestrone Receptor Antibody</u> Beckman Coulter, Inc. Fullerton, CA 92834

1-800-458-5060 Catalog # IM1546

Negative Control Serum: Normal Mouse Serum

Jackson Immunoresearch Laboratories, Inc.

West Grove, PA 19390 www.jacksonimmuno.com

1-800-367-5296 Catalog # 015-000-001

Staining Procedure

Positive Control Tissue: Female reproductive tract

Stain localization: Nuclear

1. Deparaffinize and hydrate slides through the following solutions:

Solution	Repetitions	Time
Xylene	2 times	5 minutes
100% Ethanol	2 times	3 minutes
95% Ethanol	2 times	3 minutes
1X Wash Buffer	2 times	5 minutes

- 2. Quench endogenous peroxidase by placing the slides in 3% hydrogen peroxide for 15 minutes.
- 3. Rinse the slides in 2 changes of 1X Wash Buffer for 5 minutes each.

4.	Heat-Induced Epitope Retrieval Using The Decloaker
	Add 500 ml of distilled water to the pan inside the decloaker.
	Place a full rack of slides into a Tissue Tek® container with 200 ml of 1X citrate buffer
	(Insert blank slides into any empty slots in the rack to ensure even heating of slides)
	Place the container stably inside the pan and decloak for 5 minutes. Maximum Pressure
	Depressurize for 10 minutes.
	Remove pan top and cool for 10 minutes. <i>Temperature Before Cooling Slides</i>
	Rinse the slides in 2 changes of distilled water for 3 minutes each time.
5.	Rinse the slides in 2 changes of 1X Wash Buffer for 5 minutes each time.
	Vectastain Mouse Elite Staining Kit

v cetastam mouse Ente st	anning rait			
Exp Date	New Kit:	yes	/	no

5.	6. Apply the block from the Mouse Elite Kit, and incubate for 20 minutes at room temperature					
	DO NOT RINSE THE SLIDES. CONTINUE TO AVIDIN-BIOTIN BLOCK.					
7.	Avidin / Biotin Blocking Kit					
	Lot # Exp. Date New Kit: yes / no					
	Apply avidin block for 15 minutes at room temperature.					
	Quick rinse in 1X Wash Buffer.					
	Apply biotin block for 15 minutes at room temperature.					
	DO NOT RINSE SLIDES WITH BUFFER BEFORE ADDING PRIMARY ANTIBODY.					
	ONLY WIPE EXCESS BUFFER.					
3.	Apply primary antibody at a 1:600 dilution, and incubate for 1 hour at room temperature.					
	Lot # Exp Date					

the primary antibody. Make a 1 Incubate for 1 hour at room tem	te the protein concentration of the normal mouse serum to match that of :600 dilution from this normalized serum, and apply to the slides. perature. e Reconstituted
9. Rinse the slides in 2 changes of	1X Wash Buffer for 5 minutes each.
10. Apply the secondary antibody temperature.	from Mouse Elite Kit, and incubate for 30 minutes at room
11. Rinse the slides in 2 changes o	f 1X Wash Buffer for 5 minutes each.
12. Apply the label complex from (Prepare at least 30 minutes pr	the Mouse Elite Kit, and incubate for 30 minutes at room temperature. ior to use.)
13. Rinse the slides in 2 changes o	f 1X Wash Buffer for 5 minutes each.
(Add 1 drop of DAB per ml of	nd incubate in the dark for 6 minutes at room temperature. substrate) Date New Kit: yes / no
15. Rinse the slides in tap water 3	minutes.
16. Counterstain with Harris Hema	atoxylin for 20 seconds.
17. Rinse the slides in tap water ur	ntil water is clear.

19. Dehydrate through the following solutions:

Solution	Repetitions	Time
95% Ethanol	1 time	3 minutes
100% Ethanol	3 times	3 minutes
Xylene	2 times	5 minutes

18. Gently agitate slides in 1X Wash Buffer until they turn blue.

20. Coverslip

Updated 10/20/04