DWARF SPERM WHALE (Kogia sima): Hawaiian Stock

STOCK DEFINITION AND GEOGRAPHIC RANGE

Dwarf sperm whales are found throughout the world in tropical to warm-temperate waters (Nagorsen 1985). One sighting in an unspecified locality, one stranding on Oahu (Tomich 1986), and one stranding on Lanai (Nitta 1991) constitute the only evidence that this species inhabits Hawaiian waters (Tomich 1986). Two sightings of pygmy or dwarf sperm whales were made between Hawaii and Maui during 1993-98 aerial surveys within about 25 nmi of the main Hawaiian Islands (Figure 1; Mobley et al. 1999). The difficulty of detecting and identifying it at sea, as well as its confusion with the pygmy sperm whale, may partially explain the paucity of records. For the Marine Mammal Protection Act (MMPA) stock assessment reports, there is a single Pacific management stock of dwarf sperm whales including only animals found within the U.S. Exclusive Economic Zone of the Hawaiian Islands. Rice (1998) recently argued that the species name simus, is incorrect and should be replaced by sima. This change is not taxonomic, but merely reflects rules of Latin usage.




Figure 1. Pygmy or dwarf sperm whale sighting locations during 1993-98 aerial surveys within about 25 nmi of the main Hawaiian Islands (see Appendix 2 for details on timing and location of survey effort). Outer line indicates approximate boundary of survey area.

POPULATION SIZE

Wade and Gerrodette (1993) provided an estimate for the eastern tropical Pacific, but no data are available to estimate population size for this species in the central Pacific. As part of the Marine Mammal Research Program of the Acoustic Thermometry of Ocean Climate (ATOC) study, a total of twelve aerial surveys were conducted within about 25 nmi of the main Hawaiian Islands in 1993, 1995 and 1998. Two sightings of five pygmy or dwarf sperm whales were made; however these sightings were excluded during recent abundance analyses (Mobley et al. 2000), because they were made during poor observation conditions. Therefore, no abundance estimate is available for dwarf sperm whales within Hawaiian waters.

Minimum Population Estimate

No data are available for a minimum population estimate.

Current Population Trend

No data are available on current population trend.

CURRENT AND MAXIMUM NET PRODUCTIVITY RATES

No data are available on current or maximum net productivity rate.

POTENTIAL BIOLOGICAL REMOVAL

No PBR can be calculated for this species at this time.

HUMAN-CAUSED MORTALITY AND SERIOUS INJURY

Fishery Information

No estimate of annual human-caused mortality and serious injury is available as there are no reports of direct or incidental takes of dwarf sperm whales in Hawaiian waters (Nitta and Henderson 1993). However, mortality of other cetacean species has been observed in Hawaiian fisheries, and the gear types used in these fisheries are responsible for marine mammal mortality and serious injury in other fisheries throughout U.S. waters. Gillnets are used in Hawaiian waters and appear to capture marine mammals wherever they are used, and float lines from lobster traps and longlines can be expected to occasionally entangle whales (Perrin et al. 1994).

Interactions with cetaceans have been reported for all Hawaiian pelagic fisheries (Nitta and Henderson 1993), but no interactions with dwarf sperm whales have been documented. None were observed hooked in the Hawaiian longline fishery between 1994 and 1998, with approximately 4.4% of all effort (measured as the number of hooks fished) observed (Kleiber 1999).

STATUS OF STOCK

The status of dwarf sperm whales in Hawaiian waters relative to OSP is unknown, and there are insufficient data to evaluate trends in abundance. They are not listed as "threatened" or "endangered" under the Endangered Species Act (1973), nor as "depleted" under the MMPA. Although information on dwarf sperm whales in Hawaiian waters is limited, this stock would not be considered strategic under the 1994 amendments to the MMPA because there has been no reported fisheries related mortality. The total fishery mortality and serious injury for dwarf sperm whales is zero and therefore can be considered to be insignificant and approaching zero mortality and serious injury rate. The increasing levels of anthropogenic noise in the world's oceans has been suggested to be a habitat concern for whales, particularly for deep-diving whales like dwarf sperm whales that feed in the oceans" "sound channel".

REFERENCES

- Kleiber, P. 1999. Estimates of marine mammal takes in the Hawaiian longline fishery. (Unpublished). Southwest Fisheries Science Center, NMFS, 2570 Dole St, Honolulu, HI, 96822-2396.
- Mobley, J. R., Jr., R. A. Grotefendt, P. H. Forestell, and A. Frankel. 1999. Results of aerial surveys of marine mammals in the major Hawaiian Islands (1993-98): Report to the Acoustic Thermometry of Ocean Climate Marine Mammal Research Program (ATOC MMRP) (unpublished). 39 pp.
- Mobley, J. R., Jr, S. S. Spitz, K. A. Forney, R. A. Grotefendt, and P. H. Forestall. 2000. Distribution and abundance of odontocete species in Hawaiian waters: preliminary results of 1993-98 aerial surveys Admin. Rep. LJ-00-14C. Southwest Fisheries Science Center, National Marine Fisheries Service, P.O. Box 271, La Jolla, CA 92038. 26 pp.

Nagorsen, D. 1985. Kogia simus. Mamm. Spec. 239:1-6.

- Nitta, E. 1991. The marine mammal stranding network for Hawaii: an overview. *In*: J.E. Reynolds III, D.K. Odell (eds.), Marine Mammal Strandings in the United States, pp.56-62. NOAA Tech. Rep. NMFS 98, 157 pp.
- Nitta, E. and J. R. Henderson. 1993. A review of interactions between Hawaii's fisheries and protected species. Mar. Fish. Rev. 55(2):83-92.
- Perrin, W.F., G. P. Donovan and J. Barlow. 1994. Gillnets and Cetaceans. Rep. Int. Whal. Commn., Special Issue 15, 629 pp.
- Rice, D.W. 1998. Marine Mammals of the World Systematics and Distribution. Special Publication Number 4 The
- Society of Marine Mammalogy. Douglas Wartzok (ed.). 231 pp.
- Tomich, P. Q. 1986. Mammals in Hawaii: A Synopsis and Notational Bibliography. Bishop Museum Press, Hawaii, 375 pp.
- Wade, P. R. and T. Gerrodette. 1993. Estimates of cetacean abundance and distribution in the eastern tropical Pacific. Rep. Int. Whal. Commn. 43:477-493.