

**NORA services sector: Number and percent of deaths with selected work-related respiratory condition, U.S. residents age 15 and over, selected states and years, 1990–1999**

| <b>Condition</b>                        | <b>Number of Deaths</b> | <b>Percent</b> |
|---|-------------------------|----------------|
| All Pneumoconioses                      | 731                     | 0.1            |
| Asbestosis                              | 434                     | 0.0            |
| Coal Workers' Pneumoconiosis            | 152                     | 0.0            |
| Malignant Mesothelioma (all sites)*     | 135                     | 0.0            |
| Unspecified and Other Pneumoconioses    | 84                      | 0.0            |
| Silicosis                               | 60                      | 0.0            |
| Hypersensitivity Pneumonitis            | 12                      | 0.0            |
| Byssinosis                              | 9                       | 0.0            |
| All other conditions within this sector | 1,419,930               | 99.9           |
| <b>TOTAL</b>                            | <b>1,421,547</b>        | <b>100.0</b>   |

NORA - National Occupational Research Agenda

\* data for malignant mesothelioma are for 1999 only.

NOTE: The sum of the individual and all other condition death totals may be greater than the total number of deaths in the selected states and years because some decedents have more than one condition listed on their death certificates. Percentages may not total to 100% due to rounding. See appendices for source description, methods, ICD codes, industry and occupation codes, industrial sector groupings, and list of selected states and years.

SOURCE: National Center for Health Statistics multiple cause-of-death data.