

NATIONAL TOXICOLOGY PROGRAM
Technical Report Series
No. 273



TOXICOLOGY AND CARCINOGENESIS
STUDIES OF
TRICHLOROETHYLENE
(CAS NO. 79-01-6)
IN FOUR STRAINS OF RATS
(ACI, AUGUST, MARSHALL, OSBORNE-MENDEL)
(GAVAGE STUDIES)

U.S. DEPARTMENT OF HEALTH AND HUMAN SERVICES
Public Health Service
National Institutes of Health

NATIONAL TOXICOLOGY PROGRAM

The National Toxicology Program (NTP), established in 1978, develops and evaluates scientific information about potentially toxic and hazardous chemicals. This knowledge can be used for protecting the health of the American people and for the primary prevention of disease. By bringing together the relevant programs, staff, and resources from the U.S. Public Health Service, DHHS, the National Toxicology Program has centralized and strengthened activities relating to toxicology research, testing and test development/validation efforts, and the dissemination of toxicological information to the public and scientific communities and to the research and regulatory agencies.

The NTP is made up of four charter DHHS agencies: the National Cancer Institute (NCI), National Institutes of Health; the National Institute of Environmental Health Sciences (NIEHS), National Institutes of Health; the National Center for Toxicological Research (NCTR), Food and Drug Administration; and the National Institute for Occupational Safety and Health (NIOSH), Centers for Disease Control. In July 1981, the Carcinogenesis Bioassay Testing Program, NCI, was transferred to the NIEHS.

**NTP TECHNICAL REPORT
ON THE
TOXICOLOGY AND CARCINOGENESIS
STUDIES OF TRICHLOROETHYLENE**

(CAS NO. 79-01-6)

**IN FOUR STRAINS OF RATS
(ACI, AUGUST, MARSHALL, OSBORNE-MENDEL)
(GAVAGE STUDIES)**

John H. Mennear, Ph.D., Chemical Manager



**NATIONAL TOXICOLOGY PROGRAM
P.O. Box 12233
Research Triangle Park, NC 27709**

April 1988

NTP TR 273

NIH Publication No. 88-2529

**U.S. DEPARTMENT OF HEALTH AND HUMAN SERVICES
Public Health Service
National Institutes of Health**

NOTE TO THE READER

This study was performed under the direction of the National Institute of Environmental Health Sciences as a function of the National Toxicology Program. The studies described in this Technical Report have been conducted in compliance with NTP chemical health and safety requirements and must meet or exceed all applicable Federal, state, and local health and safety regulations. Animal care and use were in accordance with the U.S. Public Health Service Policy on Humane Care and Use of Animals. All NTP toxicology and carcinogenesis studies are subjected to a data audit before being presented for public peer review.

Although every effort is made to prepare the Technical Reports as accurately as possible, mistakes may occur. Readers are requested to identify any mistakes so that corrective action may be taken. Further, anyone who is aware of related ongoing or published studies not mentioned in this report is encouraged to make this information known to the NTP. Comments and questions about the National Toxicology Program Technical Reports on Toxicology and Carcinogenesis Studies should be directed to Dr. J.E. Huff, National Toxicology Program, P.O. Box 12233, Research Triangle Park, NC 27709 (919-541-3780).


These NTP Technical Reports are available for sale from the National Technical Information Service, U.S. Department of Commerce, 5285 Port Royal Road, Springfield, VA 22161 (703-487-4650). Single copies of this Technical Report are available without charge (and while supplies last) from the NTP Public Information Office, National Toxicology Program, P.O. Box 12233, Research Triangle Park, NC 27709.

CONTENTS

| | PAGE |
|---|-------------|
| NOTE TO THE READER | 2 |
| ABSTRACT | 5 |
| EXPLANATION OF LEVELS OF EVIDENCE OF CARCINOGENIC ACTIVITY | 9 |
| CONTRIBUTORS | 10 |
| PEER REVIEW PANEL (DECEMBER 1985) | 12 |
| SUMMARY OF PEER REVIEW COMMENTS (DECEMBER 1985) | 13 |
| PEER REVIEW PANEL (AUGUST 1986) | 15 |
| SUMMARY OF PEER REVIEW COMMENTS (AUGUST 1986) | 16 |
| I. INTRODUCTION | 19 |
| II. MATERIALS AND METHODS | 27 |
| PROCUREMENT AND CHARACTERIZATION OF TRICHLOROETHYLENE | 28 |
| PREPARATION AND CHARACTERIZATION OF DOSE MIXTURES | 28 |
| THIRTEEN-WEEK STUDIES | 28 |
| TWO-YEAR STUDIES | 29 |
| STUDY DESIGN | 29 |
| SOURCE AND SPECIFICATIONS OF ANIMALS | 29 |
| ANIMAL MAINTENANCE | 31 |
| CLINICAL EXAMINATIONS AND PATHOLOGY | 31 |
| STATISTICAL METHODS | 32 |
| III. RESULTS | 35 |
| THIRTEEN-WEEK STUDIES | 36 |
| TWO-YEAR STUDIES | 37 |
| BODY WEIGHTS AND CLINICAL SIGNS | 37 |
| SURVIVAL | 37 |
| PATHOLOGY AND STATISTICAL ANALYSES OF RESULTS | 56 |
| NONNEOPLASTIC LESIONS OF THE KIDNEY | 56 |
| NEOPLASTIC LESIONS | 57 |
| NEGATIVE TRENDS AND LOWER INCIDENCES | 62 |
| IV. DISCUSSION AND CONCLUSIONS | 67 |
| V. REFERENCES | 73 |

APPENDIXES

| | PAGE |
|------------|---|
| APPENDIX A | SUMMARY OF LESIONS IN MALE ACI RATS IN THE TWO-YEAR GAVAGE STUDY OF TRICHLOROETHYLENE 79 |
| APPENDIX B | SUMMARY OF LESIONS IN FEMALE ACI RATS IN THE TWO-YEAR GAVAGE STUDY OF TRICHLOROETHYLENE 99 |
| APPENDIX C | SUMMARY OF LESIONS IN MALE AUGUST RATS IN THE TWO-YEAR GAVAGE STUDY OF TRICHLOROETHYLENE 119 |
| APPENDIX D | SUMMARY OF LESIONS IN FEMALE AUGUST RATS IN THE TWO-YEAR GAVAGE STUDY OF TRICHLOROETHYLENE 141 |
| APPENDIX E | SUMMARY OF LESIONS IN MALE MARSHALL RATS IN THE TWO-YEAR GAVAGE STUDY OF TRICHLOROETHYLENE 163 |
| APPENDIX F | SUMMARY OF LESIONS IN FEMALE MARSHALL RATS IN THE TWO-YEAR GAVAGE STUDY OF TRICHLOROETHYLENE 185 |
| APPENDIX G | SUMMARY OF LESIONS IN MALE OSBORNE-MENDEL RATS IN THE TWO-YEAR GAVAGE STUDY OF TRICHLOROETHYLENE 207 |
| APPENDIX H | SUMMARY OF LESIONS IN FEMALE OSBORNE-MENDEL RATS IN THE TWO-YEAR GAVAGE STUDY OF TRICHLOROETHYLENE 229 |
| APPENDIX I | GENETIC TOXICOLOGY OF TRICHLOROETHYLENE 251 |
| APPENDIX J | CHEMICAL CHARACTERIZATION OF TRICHLOROETHYLENE 257 |
| APPENDIX K | IDENTIFICATION OF FOREIGN MATERIAL FOUND IN TRICHLOROETHYLENE, LOT NO. TB05-206AA 271 |
| APPENDIX L | PREPARATION AND CHARACTERIZATION OF DOSE MIXTURES 277 |
| APPENDIX M | METHODS OF ANALYSIS OF DOSE MIXTURES 281 |
| APPENDIX N | RESULTS OF ANALYSIS OF DOSE MIXTURES 285 |
| APPENDIX O | SENTINEL ANIMAL PROGRAM 289 |
| APPENDIX P | SURVIVAL AND MEAN BODY WEIGHTS OF ACI, AUGUST, AND MARSHALL RATS IN THE THIRTEEN-WEEK GAVAGE STUDIES OF TRICHLOROETHYLENE 293 |
| APPENDIX Q | AUDIT SUMMARY 297 |



TRICHLOROETHYLENE

CAS No. 79-01-6

C_2HCl_3 Molecular weight 131.4

Synonyms: Acetylene trichloride; 1-chloro-2,2-dichloroethylene; 1,1-dichloro-2-chloroethylene; ethinyl trichloride; ethylene trichloride; 1,1,2-trichloroethylene; trichloroethene

Trade names of formulations: Algylen; Anamenth; Benzinol; Blacosolv; Blancosolv; Cecolene; Chlorilen; Chlorylea; Chlorylen; Chorylen; Circosolv; Crawhaspol; Densinfluat; Dow-Tri; Dukeron; Fleck-Flip; Flock Flip; Fluate; Gemalgene; Germalgene; Lanadin; Lethurin; Narcogen; Narkogen; Narkosoid; Nialk; Perma-A-Chlor; Perm-A-Clor; Petzinol; Philex; Threthylen; Threthylene; Trethylene; Tri; Triad; Trial; Triasol; Trichloran; Trichloren; Triclene; Tri-Clene; Trielene; Trielin; Triklone; Trilen; Trilene; Triline; Trimar; Triol; TRI-plus; TRI-plus M; Vestrol; Vitran; Westrosol

ABSTRACT

Trichloroethylene is an industrial solvent used primarily for vapor degreasing and cold cleaning. It was selected for study because of its industrial use and potential for human exposure. (An estimated 3.5 million workers are exposed to trichloroethylene.) In an earlier study (NCI TR 2), trichloroethylene (stabilized with epichlorohydrin and 1,2-epoxybutane) administered by gavage caused hepatocellular carcinomas in male and female B6C3F₁ mice. Trichloroethylene administration did not increase the incidence of tumors in male or female Osborne-Mendel rats. However, the survival of dosed rats was reduced, thereby compromising the sensitivity of the study to detect a carcinogenic effect.

The studies described in this report were conducted to compare the sensitivities of four strains of rats (ACI, August, Marshall, and Osborne-Mendel) to diisopropylamine-stabilized trichloroethylene. The results of the present studies demonstrate that long-term administration of trichloroethylene produces nephrotoxicity in four strains of rats and that the susceptibilities of these strains to the nephrotoxic effects of the chemical are similar. Because of chemically induced toxicity, reduced survival, and incomplete documentation of experimental data, the studies are considered inadequate for either comparing or assessing trichloroethylene-induced carcinogenesis in these strains of rats.

Toxicology and carcinogenesis studies of trichloroethylene (more than 99% pure, stabilized with 8 ppm diisopropylamine) were conducted by administering the chemical in corn oil by gavage at doses of 0, 500, or 1,000 mg/kg per day, 5 days per week, for 103 weeks to groups of 50 male and 50 female ACI, August, Marshall, and Osborne-Mendel rats. The doses were selected on the basis of results from 13-week gavage studies in which groups of 10 male and 10 female ACI, August, and Marshall rats received daily doses of trichloroethylene (male: 125-2,000 mg/kg; female: 63-1,000 mg/kg). Doses for Osborne-Mendel rats were selected to conform with doses used in an earlier carcinogenicity study in that strain (NCI TR 2).

In the 13-week studies, male ACI and August rats receiving 2,000 mg/kg trichloroethylene and male and female Marshall rats receiving 1,835 mg/kg had final mean body weights 12%-17% lower than those of the vehicle controls. All other dose groups had body weights comparable to those of the vehicle controls. Three male August rats dosed with 2,000 mg/kg died. Histopathologic evaluation of tissues revealed no lesions attributable to trichloroethylene administration in the 13-week studies. This absence of histopathologic findings did not accurately predict the nephrotoxic effects of long-term administration of trichloroethylene to rats.

Body Weight and Survival in the Two-Year Studies: In the 2-year studies, all dosed groups exhibited some reduction in mean body weights relative to the vehicle controls. Survival relative to vehicle controls was significantly reduced in 7/16 dosed groups (see following table). Also, the survival of high dose male Marshall rats was reduced by a large number of accidental deaths. Nephrotoxicity, reduced survival, and central nervous system toxicity (characterized by sedation, loss of consciousness, tremors, and convulsions) showed that the doses of trichloroethylene selected for the 2-year studies were too high.

SUMMARY OF SURVIVAL AND FINAL MEAN BODY WEIGHTS OF ACI, AUGUST, MARSHALL, AND OSBORNE-MENDEL RATS IN THE TWO-YEAR GAVAGE STUDIES OF TRICHLOROETHYLENE

| Group | Survival (a) | Final Mean Body Weight (percent of vehicle control) | | |
|-----------------------|--------------|---|-----------|-------|
| | | Male | Female | |
| ACI | | | | |
| Untreated control | 39/49 | -- | 37/49 | -- |
| Vehicle control | 38/50 | -- | 35/50 | -- |
| 500 mg/kg | (b) 19/50 | 89.0 | 20/50 | 91.7 |
| 1,000 mg/kg | (b) 11/50 | 87.5 | (b) 19/50 | 93.0 |
| August | | | | |
| Untreated control | 24/50 | -- | 26/50 | -- |
| Vehicle control | 21/50 | -- | 23/50 | -- |
| 500 mg/kg | 13/50 | 93.5 | 26/50 | 94.5 |
| 1,000 mg/kg | 16/49 | 87.7 | 25/50 | 92.4 |
| Marshall | | | | |
| Untreated control | 32/50 | -- | 31/50 | -- |
| Vehicle control | 26/50 | -- | 30/50 | -- |
| 500 mg/kg | (b) 12/50 | 93.3 | (b) 12/50 | 96.0 |
| 1,000 mg/kg | (c) 6/50 | 96.8 | (b) 10/50 | 89.9 |
| Osborne-Mendel | | | | |
| Untreated control | 18/50 | -- | 19/50 | -- |
| Vehicle control | 22/50 | -- | 20/50 | -- |
| 500 mg/kg | 17/50 | 97.5 | 11/50 | 92.7 |
| 1,000 mg/kg | 15/50 | 88.4 | (b) 7/50 | 100.7 |

(a) Proportion of survivors after 2 years

(b) Survival was significantly ($P < 0.05$) reduced relative to vehicle controls.

(c) Twenty-five animals were accidentally killed.

Renal Effects in the Two-Year Studies: Trichloroethylene caused tubular cell cytomegaly in 82%-100% of all dosed animals. In addition, trichloroethylene produced toxic nephropathy (which was distinguishable from age-related nephropathy) in 17%-80% of the dosed animals. Cytomegaly, karyomegaly, or toxic nephropathy was not found in untreated or vehicle control animals. Trichloroethylene administration was also associated with increased incidences of renal tubular cell adenomas and adenocarcinomas. The incidences of renal lesions are shown in the following table.

Other Pathologic Effects in the Two-Year Studies: An increased incidence of interstitial cell tumors of the testis was observed in high dose male Marshall rats (untreated control, 16/46; vehicle control, 17/46; low dose, 21/48; high dose, 32/48; P = 0.002). The incidences of pheochromocytomas of the adrenal gland were significantly reduced in male ACI, female August, female Marshall, and male and female Osborne-Mendel rats.

Genetic Toxicology: Trichloroethylene did not cause mutations in *Salmonella typhimurium* strains TA98, TA100, TA1535, or TA1537 with or without metabolic activation. In Chinese hamster ovary cells, trichloroethylene did not induce chromosomal aberrations; the results for sister chromatid exchanges were considered positive. Trichloroethylene was mutagenic to mouse L5178Y lymphoma cells in the presence of rat liver S9.

SUMMARY OF INCIDENCES OF RENAL LESIONS IN THE TWO-YEAR GAVAGE STUDIES OF TRICHLOROETHYLENE

| Lesion | Male | | | | Female | | | |
|-----------------------------|-------------------|-----------------|-----------|-------------|-------------------|-----------------|-----------|-------------|
| | Untreated Control | Vehicle Control | 500 mg/kg | 1,000 mg/kg | Untreated Control | Vehicle Control | 500 mg/kg | 1,000 mg/kg |
| ACI | | | | | | | | |
| No. kidneys examined | 49 | 50 | 49 | 49 | 49 | 48 | 47 | 43 |
| Cytomegaly | 0 | 0 | 40 | 48 | 0 | 0 | 43 | 42 |
| Toxic nephropathy | 0 | 0 | 18 | 18 | 0 | 0 | 21 | 19 |
| Tubular cell adenoma | 0 | 0 | 0 | 0 | 0 | 0 | 2 | 0 |
| Tubular cell adenocarcinoma | 0 | 0 | 1 | 0 | 0 | 0 | (a) 1 | 1 |
| August | | | | | | | | |
| No. kidneys examined | 50 | 50 | 50 | 49 | 50 | 49 | 48 | 50 |
| Cytomegaly | 0 | 0 | 46 | 46 | 0 | 0 | 46 | 50 |
| Toxic nephropathy | 0 | 0 | 10 | 31 | 0 | 0 | 8 | 29 |
| Tubular cell adenoma | 0 | 0 | 1 | 1 | 0 | 1 | 2 | 0 |
| Tubular cell adenocarcinoma | 0 | 0 | 1 | 0 | 0 | 0 | 2 | 0 |
| Marshall | | | | | | | | |
| No. kidneys examined | 49 | 49 | 50 | 47 | 49 | 50 | 48 | 44 |
| Cytomegaly | 0 | 0 | 48 | 47 | 0 | 0 | 46 | 43 |
| Toxic nephropathy | 0 | 0 | 18 | 23 | 1 | 0 | 30 | 30 |
| Tubular cell adenoma | 2 | 0 | 1 | 0 | 1 | 1 | 1 | 0 |
| Tubular cell adenocarcinoma | 0 | 0 | 0 | 1 | 0 | 0 | 1 | 1 |
| Osborne-Mendel | | | | | | | | |
| No. kidneys examined | 50 | 50 | 50 | 50 | 50 | 50 | 50 | 49 |
| Cytomegaly | 0 | 0 | 48 | 49 | 0 | 0 | 48 | 49 |
| Toxic nephropathy | 0 | 0 | 39 | 35 | 0 | 0 | 30 | 39 |
| Tubular cell adenoma | 0 | 0 | 6 | 1 | 1 | 0 | 0 | 1 |
| Tubular cell adenocarcinoma | 0 | 0 | 0 | 1 | 0 | 0 | 0 | 0 |

(a) Adenocarcinoma, NOS

Data Audit: Audits of the experimental data for these 2-year studies of trichloroethylene were conducted by the National Toxicology Program (Appendix Q). The results of the audits revealed evidence that the doses of trichloroethylene were too high. In addition, there was insufficient documentation of animal breeding, clinical observations, environmental conditions, and analytical chemistry data. Also, individual animal identification was not always verifiable.

Conclusions: Under the conditions of these 2-year gavage studies of trichloroethylene in male and female ACI, August, Marshall, and Osborne-Mendel rats, trichloroethylene administration caused renal tubular cell cytomegaly and toxic nephropathy in both sexes of the four strains. However, these are considered to be *inadequate studies of carcinogenic activity** because of chemically induced toxicity, reduced survival, and deficiencies in the conduct of the studies. Despite these limitations, tubular cell neoplasms of the kidney were observed in rats exposed to trichloroethylene and interstitial cell neoplasms of the testis were observed in Marshall rats exposed to trichloroethylene.

**SUMMARY OF THE TWO-YEAR GAVAGE STUDIES OF TRICHLOROETHYLENE IN MALE AND FEMALE
ACI, AUGUST, MARSHALL, AND OSBORNE-MENDEL RATS**

Doses

0, 500, and 1,000 mg/kg trichloroethylene in corn oil, 5 d/wk

Level of evidence of carcinogenic activity

Inadequate study

As indicated by

Chemically induced toxicity, reduced survival, and deficiencies in study conduct

Other toxic effects

Renal tubular cytomegaly, toxic nephropathy

Neoplastic effects

Uncommon renal tubular cell neoplasms in exposed rats and testicular interstitial cell tumors in male Marshall rats

*Explanation of Levels of Evidence of Carcinogenic Activity is on page 9.

Summaries of the Peer Review comments and the public discussions on this Technical Report appear on pages 13-14 and 16-17.

EXPLANATION OF LEVELS OF EVIDENCE OF CARCINOGENIC ACTIVITY

These studies are designed and conducted to characterize and evaluate the toxicologic potential, including carcinogenic activity, of selected chemicals in laboratory animals (usually two species, rats and mice). Chemicals selected for NTP toxicology and carcinogenesis studies are chosen primarily on the bases of human exposure, level of production, and chemical structure. Selection per se is not an indicator of a chemical's carcinogenic potential.

Negative results, in which the study animals do not have a greater incidence of neoplasia than control animals, do not necessarily mean that a chemical is not a carcinogen, inasmuch as the experiments are conducted under a limited set of conditions. Positive results demonstrate that a chemical is carcinogenic for laboratory animals under the conditions of the study and indicate that exposure to the chemical has the potential for hazard to humans.

The National Toxicology Program describes the results of individual experiments on a chemical agent and notes the strength of the evidence for conclusions regarding each study. Other organizations, such as the International Agency for Research on Cancer, assign a strength of evidence for conclusions based on an examination of all available evidence including: animal studies such as those conducted by the NTP, epidemiologic studies, and estimates of exposure. Thus, the actual determination of risk to humans from chemicals found to be carcinogenic in laboratory animals requires a wider analysis that extends beyond the purview of these studies.

Five categories of evidence of carcinogenic activity are used in the Technical Report series to summarize the strength of the evidence observed in each experiment: two categories for positive results ("Clear Evidence" and "Some Evidence"); one category for uncertain findings ("Equivocal Evidence"); one category for no observable effects ("No Evidence"); and one category for experiments that because of major flaws cannot be evaluated ("Inadequate Study"). These categories of interpretative conclusions were first adopted in June 1983 and then revised in March 1986 for use in the Technical Reports series to incorporate more specifically the concept of actual weight of evidence of carcinogenic activity. For each separate experiment (male rats, female rats, male mice, female mice), one of the following quintet is selected to describe the findings. These categories refer to the strength of the experimental evidence and not to either potency or mechanism.

- **Clear Evidence of Carcinogenic Activity** is demonstrated by studies that are interpreted as showing a dose-related (i) increase of malignant neoplasms, (ii) increase of a combination of malignant and benign neoplasms, or (iii) marked increase of benign neoplasms if there is an indication from this or other studies of the ability of such tumors to progress to malignancy.
- **Some Evidence of Carcinogenic Activity** is demonstrated by studies that are interpreted as showing a chemically related increased incidence of neoplasms (malignant, benign, or combined) in which the strength of the response is less than that required for clear evidence.
- **Equivocal Evidence of Carcinogenic Activity** is demonstrated by studies that are interpreted as showing a marginal increase of neoplasms that may be chemically related.
- **No Evidence of Carcinogenic Activity** is demonstrated by studies that are interpreted as showing no chemically related increases in malignant or benign neoplasms.
- **Inadequate Study of Carcinogenic Activity** is demonstrated by studies that because of major qualitative or quantitative limitations cannot be interpreted as valid for showing either the presence or absence of carcinogenic activity.

When a conclusion statement for a particular experiment is selected, consideration must be given to key factors that would extend the actual boundary of an individual category of evidence. This should allow for incorporation of scientific experience and current understanding of long-term carcinogenesis studies in laboratory animals, especially for those evaluations that may be on the borderline between two adjacent levels. These considerations should include:

- The adequacy of the experimental design and conduct;
- Occurrence of common versus uncommon neoplasia;
- Progression (or lack thereof) from benign to malignant neoplasia as well as from preneoplastic lesions;
- Some benign neoplasms have the capacity to regress but others (of the same morphologic type) progress. At present, it is impossible to identify the difference. Therefore, where progression is known to be a possibility, the most prudent course is to assume that benign neoplasms of those types have the potential to become malignant;
- Combining benign and malignant tumor incidences known or thought to represent stages of progression in the same organ or tissue;
- Latency in tumor induction;
- Multiplicity in site-specific neoplasia;
- Metastases;
- Supporting information from proliferative lesions (hyperplasia) in the same site of neoplasia or in other experiments (same lesion in another sex or species);
- The presence or absence of dose relationships;
- The statistical significance of the observed tumor increase;
- The concurrent control tumor incidence as well as the historical control rate and variability for a specific neoplasm;
- Survival-adjusted analyses and false positive or false negative concerns;
- Structure-activity correlations; and
- In some cases, genetic toxicology.

These considerations together with the definitions as written should be used as composite guidelines for selecting one of the five categories. Additionally, the following concepts (as patterned from the International Agency for Research on Cancer Monographs) have been adopted by the NTP to give further clarification of these issues:

The term *chemical carcinogenesis* generally means the induction by chemicals of neoplasms not usually observed, the induction by chemicals of more neoplasms than are generally found, or the earlier induction by chemicals of neoplasms that are commonly observed. Different mechanisms may be involved in these situations. Etymologically, the term *carcinogenesis* means induction of cancer, that is, of malignant neoplasms; however, the commonly accepted meaning is the induction of various types of neoplasms or of a combination of malignant and benign neoplasms. In the Technical Reports, the words *tumor* and *neoplasm* are used interchangeably.

CONTRIBUTORS

The NTP Technical Report on the Toxicology and Carcinogenesis Studies of Trichloroethylene in Four Strains of Rats is based on the 13-week studies that began in March or April 1977 and ended in June or July 1977 at Papanicolaou Research Institute and on the 2-year studies that began between December 1978 and November 1979 and ended between November 1980 and November 1981 at Papanicolaou Cancer Research Institute (Miami, Florida).

National Toxicology Program (Evaluated Experiment, Interpreted Results, and Reported Findings)

John H. Mennear, Ph.D., Chemical Manager

Jack Bishop, Ph.D.

Gary A. Boorman, D.V.M., Ph.D.

Joseph K. Haseman, Ph.D.

James Huff, Ph.D.

C.W. Jameson, Ph.D.

E.E. McConnell, D.V.M.

G.N. Rao, D.V.M., Ph.D.

B.A. Schwetz, D.V.M., Ph.D.

NTP Pathology Working Group (Evaluated ACI Rat Slides and Prepared Pathology Report on 2/25/82)

G. Boorman, D.V.M., Ph.D. (NTP) (Chair)

P. Hildebrandt, D.V.M. (Tracor Jitco, Inc.)

R. Kovatch, D.V.M. (Tracor Jitco, Inc.)

E. McConnell, D.V.M. (NTP)

(Evaluated August Rat Slides and Prepared Pathology Report on 7/27/82)

M. Anver, D.V.M. (Clement Associates) (Chair)

G. Boorman, D.V.M., Ph.D. (NTP)

S. Eustis, D.V.M., Ph.D. (NTP)

R. Kovatch, D.V.M. (Tracor Jitco, Inc.)

(Evaluated Marshall Rat Slides and Prepared Pathology Report on 9/2/81)

G. Boorman, D.V.M., Ph.D. (NTP) (Chair)

B. Gupta, B.V.Sc., Ph.D. (NTP)

L. Lomax, D.V.M. (NTP)

E. McConnell, D.V.M. (NTP)

F. Voelker, D.V.M.

Burroughs Wellcome Laboratories

(Evaluated Osborne-Mendel Rat Slides and Prepared Pathology Report on 8/13/82)

G. Boorman, D.V.M., Ph.D. (NTP) (Chair)

S. Eustis, D.V.M., Ph.D. (NTP)

P. Hildebrandt, D.V.M. (Tracor Jitco, Inc.)

R. Sauer, V.M.D. (Gillette Research)

(Special Review of Kidney Lesions on 12/13/83)

H. Solleveld, D.V.M., Ph.D. (NTP) (Chair)

G. Boorman, D.V.M., Ph.D. (NTP)

R. Goyer, M.D. (NIEHS)

A. Macklin, D.V.M., Ph.D.

Burroughs Wellcome Laboratories

R. Maronpot, D.V.M. (NTP)

C. Montgomery, D.V.M. (NTP)

K. Morgan, Ph.D.

Chemical Industry Institute of Technology

Principal Contributors at Papanicolaou Cancer Research Institute (Conducted Studies and Evaluated Tissues)

Norman Altman, D.V.M.

Ernesto Bernal, D.V.M., Pathologist (ACI, August,
and Osborne-Mendel rats)

Fazel Ahmed, Ph.D.

Herman Seibold, D.V.M., Pathologist
(Marshall rats)

CONTRIBUTORS (Continued)

**Principal Contributors at Experimental Pathology Laboratories, Inc.
(Conducted Pathology Quality Assurance)**

| | |
|---------------------|-----------------------|
| J. Hardisty, D.V.M. | ACI Rats |
| C.E. Gilmore, Ph.D. | August Rats |
| M. Hamlin, D.V.M. | Marshall Rats |
| L. Ackerman, D.V.M. | Marshall Rats |
| M. Robl, D.V.M. | Osborne-Mendel Rats |
| J. Gauchat | Pathology Coordinator |

**Principal Contributors at Carltech Associates, Inc.
(Contractor for Technical Report Preparation)**

| | |
|-----------------------------|-------------------|
| William D. Theriault, Ph.D. | John Warner, M.S. |
| Abigail C. Jacobs, Ph.D. | |

PEER REVIEW PANEL (DECEMBER 1985)

The members of the Peer Review Panel who evaluated the draft Technical Report on trichloroethylene on December 9, 1985, are listed below. Panel members serve as independent scientists, not as representatives of any institution, company, or governmental agency. In this capacity, Panel members have five major responsibilities: (a) to ascertain that all relevant literature data have been adequately cited and interpreted, (b) to determine if the design and conditions of the NTP studies were appropriate, (c) to ensure that the Technical Report presents the experimental results and conclusions fully and clearly, (d) to judge the significance of the experimental results by scientific criteria, and (e) to assess the evaluation of the evidence of carcinogenicity and other observed toxic responses.

National Toxicology Program Board of Scientific Counselors Technical Reports Review Subcommittee

Jerry B. Hook, Ph.D. (Chair)

Vice President, Preclinical Research and Development
Smith Kline & French Laboratories, Philadelphia, Pennsylvania

Frederica Perera, Dr. P.H.
Division of Environmental Sciences
School of Public Health, Columbia University
New York, New York

James Swenberg, D.V.M., Ph.D.
(Principal Reviewer)
Head, Department of Biochemical
Toxicology and Pathobiology
Chemical Industry Institute of Toxicology
Research Triangle Park, North Carolina

Ad Hoc Subcommittee Panel of Experts

John J. Crowley, Ph.D. (Principal Reviewer)
Division of Public Health Science
The Fred Hutchinson Cancer Research Center
Seattle, Washington

Franklin E. Mirer, Ph.D.
Director, Health and Safety Department
International Union, United Auto
Workers, Detroit, Michigan

Kim Hooper, Ph.D. (Principal Reviewer)
Hazard Evaluation System and
Information Services
Department of Health Services
State of California
Berkeley, California

I.F.H. Purchase, B.V.Sc., Ph.D., F.R.C. Path.
Director, Central Toxicology Laboratory
Imperial Chemical Industries, PLC
Alderley Park, England

Thomas C. Jones, D.V.M.
Professor, Comparative Pathology
New England Regional Primate Research Center
Harvard Medical School
Southborough, Massachusetts

Robert A. Scala, Ph.D.
Senior Scientific Advisor, Medicine and
Environmental Health Department
Exxon Corporation
East Millstone, New Jersey

Richard J. Kociba, D.V.M., Ph.D.
Dow Chemical USA
Midland, Michigan

Steven R. Tannenbaum, Ph.D.
Professor, Department of Nutrition and
Food Science
Massachusetts Institute of Technology
Cambridge, Massachusetts

David Kotelchuck, Ph.D.*
Environmental Health Science Program
Hunter School of Health Sciences
New York, New York

Bruce W. Turnbull, Ph.D.
Professor and Associate Director
College of Engineering
Cornell University
Ithaca, New York

*Unable to attend

**SUMMARY OF PEER REVIEW COMMENTS
ON THE TOXICOLOGY AND CARCINOGENESIS STUDIES OF
TRICHLOROETHYLENE (DECEMBER 1985)**

On December 9, 1985, the draft Technical Report on the toxicology and carcinogenesis studies of trichloroethylene received peer review by the National Toxicology Program Board of Scientific Counselors' Technical Reports Review Subcommittee and associated Panel of Experts. The review meeting was held at the National Institute of Environmental Health Sciences, Research Triangle Park, North Carolina.

Dr. J. Mennear, NTP Chemical Manager, introduced the toxicology and carcinogenesis studies of trichloroethylene by reviewing the experimental designs, results, and proposed conclusions:

These 2-year gavage studies of trichloroethylene in male and female ACI, August, Marshall, and Osborne-Mendel rats are considered to be inadequate studies of carcinogenicity because of insufficient survival in dosed animals and incomplete documentation of the conduct of the studies. However, under the conditions of these studies trichloroethylene administration was strongly associated with renal tubular cell cytomegaly and karyomegaly and toxic nephropathy in both sexes of the four strains. In addition, an increased incidence of renal tubular cell neoplasms in male Osborne-Mendel rats, and possibly in female ACI and female August rats, and an increased incidence of testicular interstitial cell tumors in male Marshall rats may have been associated with the administration of trichloroethylene.

Dr. Hooper, a principal reviewer, agreed with the proposed conclusions. He discussed deficiencies in study design, conduct, and recordkeeping. With regard to the large numbers of "accidental" or "gavage-related" deaths, he argued that gavage trauma is lethal when accompanied by the toxicity of trichloroethylene. Dr. Hooper proposed that inhalation studies in both rats and mice be designed to confirm or deny the assertion that trichloroethylene is a tissue- and species-specific carcinogen (mouse liver). He asked that reference be included to a 1983 Japanese inhalation study in which trichloroethylene was reported to produce pulmonary adenocarcinomas in female ICR mice.

As a second principal reviewer, Dr. Swenberg agreed with the conclusions but stated that these studies were inadequately designed, failing to utilize a previous study with Osborne-Mendel rats in dose selection. Beginning with the Abstract, it should be stated that the doses used exceeded the maximum tolerated dose in all four strains of rats. He detailed a number of suggested revisions. However, he said that the last sentence regarding kidney tumors should be deleted as the findings described have not been reproduced in several negative studies and are difficult to assess due to the severe renal toxicity.

As a third principal reviewer, Dr. Crowley also suggested that the last sentence should be deleted from the conclusions. He also asked for clarification concerning the relationship of gavage trauma with chemical toxicity in causation of "accidental" deaths. Dr. Mennear responded that the mortality originally termed "gavage-related deaths" was likely to be associated with toxicity of trichloroethylene.

Dr. E. McConnell, NIEHS, said that despite the deficiencies in these studies, the NTP considered the toxic renal lesions to be real and consistent with effects seen with other halogenated solvents.

In other discussion, Dr. Tannenbaum pointed out that the major metabolite of trichloroethylene in humans and rats is trichloroacetic acid, and this compound could be the cause of the nephrotoxicity, thus raising the question of whether there might be a threshold effect. Dr. Mirer cautioned against minimizing the importance of the renal toxicity in view of the fact that the doses used were in the same range as the occupationally permitted exposure levels in air. Dr. Perera spoke against deleting the last sentence of the conclusions, arguing that if the nontumor toxic effects should not be dismissed despite the deficiencies of the studies, then neither should the neoplastic effects. Dr. McConnell said that the toxic renal lesions were observed at very high incidence whereas the neoplastic changes were found at very low incidence. Dr. Swenberg stated that conclusions about carcinogenicity cannot be drawn from an inadequate study of carcinogenicity.

Dr. Scala said that the audit report findings suggest these studies are flawed and should not be published as an NTP Technical Report. The data on the kidney lesions could be reported separately. Dr. Jones said that the information should be made readily available. Dr. Purchase thought it difficult to recommend publication or not based on the present report. He suggested that the report be redrafted and brought back to the Panel.

Dr. Hooper moved that the report be deferred for extensive revision and then brought back to the Panel. Dr. Swenberg seconded the motion. Dr. Kociba requested that more information from the Audit Report be included in the redraft. Dr. Swenberg asked that information also be provided on the findings from an independent audit by the Halogenated Solvent Industry Alliance (HSIA). Dr. J. Huff, NIEHS, said that a revised Technical Report could be ready for consideration at the summer 1986 meeting. Dr. Hook summarized the motion to say that the Technical Report would be rewritten, adding a more extensive summary of the findings from audits conducted by both the NTP and HSIA and returned to the Panel for review. The Panel approved the motion by nine affirmative votes with two abstentions (Dr. Kociba and Dr. Purchase).

PEER REVIEW PANEL (AUGUST 1986)

The members of the Peer Review Panel who evaluated the draft Technical Report on trichloroethylene on August 19, 1986, are listed below. Panel members serve as independent scientists, not as representatives of any institution, company, or governmental agency. In this capacity, Panel members have five major responsibilities: (a) to ascertain that all relevant literature data have been adequately cited and interpreted, (b) to determine if the design and conditions of the NTP studies were appropriate, (c) to ensure that the Technical Report presents the experimental results and conclusions fully and clearly, (d) to judge the significance of the experimental results by scientific criteria, and (e) to assess the evaluation of the evidence of carcinogenicity and other observed toxic responses.

National Toxicology Program Board of Scientific Counselors Technical Reports Review Subcommittee

Robert A. Scala, Ph.D. (Chair)

Senior Scientific Advisor, Medicine and Environmental Health Department
Research and Environmental Health Division, Exxon Corporation
East Millstone, New Jersey

Michael A. Gallo, Ph.D.

Associate Professor, Director of Toxicology
Department of Environmental and Community
Medicine, UMDNJ - Rutgers Medical School
Piscataway, New Jersey

Frederica Perera, Dr. P.H.

Division of Environmental Sciences
School of Public Health, Columbia
University
New York, New York

Ad Hoc Subcommittee Panel of Experts

Charles C. Capen, D.V.M., Ph.D.

Department of Veterinary Pathobiology
Ohio State University
Columbus, Ohio

Franklin E. Mirer, Ph.D.

Director, Health and Safety Department
International Union, United Auto
Workers, Detroit, Michigan

Vernon M. Chinchilli, Ph.D.

Department of Biostatistics
Medical College of Virginia
Virginia Commonwealth University
Richmond, Virginia

James A. Popp, D.V.M., Ph.D. (Principal
Reviewer) Head, Department of

Experimental Pathology and Toxicology
Chemical Industry Institute of Toxicology
Research Triangle Park, North Carolina

John J. Crowley, Ph.D. (Principal Reviewer)

Division of Public Health Science
The Fred Hutchinson Cancer Research Center
Seattle, Washington

I.F.H. Purchase, B.V.Sc., Ph.D., F.R.C. Path.

Director, Central Toxicology Laboratory
Imperial Chemical Industries, PLC
Alderley Park, England

Kim Hooper, Ph.D. (Principal Reviewer)

Hazard Evaluation System and
Information Services
Department of Health Services
State of California
Berkeley, California

Andrew Sivak, Ph.D.

Vice President, Biomedical Science
Arthur D. Little, Inc.
Cambridge, Massachusetts

Donald H. Hughes, Ph.D.

Scientific Coordinator, Regulatory Services
Division, The Procter and Gamble Company
Cincinnati, Ohio

**SUMMARY OF PEER REVIEW COMMENTS
ON THE TOXICOLOGY AND CARCINOGENESIS STUDIES OF
TRICHLOROETHYLENE (AUGUST 1986)**

On August 19, 1986, the draft Technical Report on the toxicology and carcinogenesis studies of trichloroethylene received peer review by the National Toxicology Program Board of Scientific Counselors' Technical Reports Review Subcommittee and associated Panel of Experts. The review meeting was held at the National Institute of Environmental Health Sciences, Research Triangle Park, North Carolina.

Dr. J. Mennear, NTP, introduced the studies by reviewing the experimental design, results, and proposed conclusions (inadequate studies of carcinogenic activity in the four strains of rats).

Dr. Hooper, a principal reviewer, stated that this version was improved considerably over the previous draft, especially in justifying dose selection, comparing renal toxicity in five rat strains (including F344), eliminating mention of gavage trauma as the major cause of decreased survival, and summarizing audit findings and deficiencies in the execution of the studies. Among many comments, Dr. Hooper suggested that significant carcinogenic effects should be included in the summary, even though the overall studies were judged inadequate for assessing carcinogenicity, and he cited specific increases in renal and testicular tumors. He asked for a clarifying discussion on what is meant by "accidental" deaths, since some of the excessive mortality could be due to anesthetic or toxic properties of the chemical. Dr. Hooper suggested that a more balanced discussion be given to the carcinogenic effects of chlorinated aliphatics, including findings from inhalation studies in which the carcinogenic responses appeared to be broader in terms of site.

As a second principal reviewer, Dr. Popp basically agreed with the conclusion that these were inadequate studies but felt that the report should more clearly and specifically state the basis for the studies' inadequacies. Dr. Popp indicated that statements in the text about significant increases in renal tumors could lead to a misunderstanding that these were positive studies. He thought that these statements could be better qualified.

As a third principal reviewer, Dr. Crowley questioned whether the report yet comes to grips with how serious the data problems are. For example, the results of the data audit indicate that analyses by dose group and evaluations of dose response may be potentially misleading. He said that if these were inadequate studies, the significant tumor findings should be downplayed or not presented at all. Dr. Crowley noted a possible exception, testicular tumors in Marshall rats, which did not involve a compromised organ and were statistically significant regardless of dose identification.

Most of the discussion was concerned with the weight that should be given to the renal and testicular tumors observed and whether statistical significances should be given within the context of inadequate studies. One viewpoint as supported by Dr. Purchase, Dr. Crowley, and Dr. Popp was that conclusions about carcinogenic activity from inadequate studies are unwarranted, as are statements about the statistical significance of low incidence rates in animals with uncertain identification. A second viewpoint, supported by Dr. Hooper, Dr. Mirer, and Dr. Perera, was that, although conclusions cannot be drawn from inadequate studies, more emphasis could be given to the tumor data if it is believed that there is a probable association between chemical administration and increased tumor incidence. Dr. J. Huff, NIEHS, stated that NTP staff agreed that these renal tumors were related to trichloroethylene administration, and he reported that combining data from male and female animals for both vehicle control and dosed groups shows only 2 renal tumors in vehicle controls versus 26 in dosed animals. Dr. S. Eustis, NIEHS, noted that there was an overemphasis regarding the purported misidentification of animals. Dr. J. Selkirk, NIEHS, said that the audit summary reinforced the idea

that animals were indeed identifiable and could be separated easily into dosed and vehicle control animals.

Dr. Hooper moved that the Technical Report on trichloroethylene be accepted with the conclusions as written, inadequate studies of carcinogenic activity in the four strains of rats, and with the addition of a statement to the summary that there were increased incidences of renal tubular cell tumors observed in dosed rats and an increased incidence of interstitial cell tumors of the testes in dosed Marshall rats. Dr. Mirer seconded the motion, and it was approved by six reviewers with three dissenting (Dr. Capen, Dr. Crowley, and Dr. Popp) and one abstaining (Dr. Purchase).

I. INTRODUCTION

Use

Chemical and Physical Properties of

Trichloroethylene

Exposure and Production


Teratogenicity and Embryotoxicity

Genetic Toxicology

Carcinogenicity

Study Rationale

I. INTRODUCTION



TRICHLOROETHYLENE

CAS No. 79-01-6

C_2HCl_3 Molecular weight 131.4

Synonyms: Acetylene trichloride; 1-chloro-2,2-dichloroethylene; 1,1-dichloro-2-chloroethylene; ethinyl trichloride; ethylene trichloride; 1,1,2-trichloroethylene; trichloroethene

Trade names of formulations: Algylen; Anamenth; Benzinol; Blacosolv; Blancosolv; Cecolene; Chlorilen; Chlorylea; Chlorylen; Chorylen; Circosolv; Crawhaspol; Densinfluat; Dow-Tri; Dukeron; Fleck-Flip; Flock Flip; Fluate; Gemalgene; Germalgene; Lanadin; Lethurin; Narcogen; Narkogen; Narkosoid; Nialk; Perma-A-Chlor; Perm-A-Clor; Petzinol; Philex; Threthylen; Threthylene; Trethylene; Tri; Triad; Trial; Triasol; Trichloran; Trichloren; Triclene; Tri-Clene; Trielene; Trielin; Triklone; Trilen; Trilene; Triline; Trimar; Triol; TRI-plus; TRI-plus M; Vestrol; Vitran; Westrosol

Use

Trichloroethylene is an industrial solvent used primarily for vapor degreasing and cold cleaning of fabricated metal parts. Trichloroethylene has been used as a carrier solvent for the active ingredients of insecticides and fungicides; as a solvent for waxes, fats, resins, and oils; as a spot remover; as an anesthetic for medical, dental, and veterinary use; and as an extractant for spice oleoresins and for caffeine from coffee. Trichloroethylene may be found in printing inks, varnishes, adhesives, paints, lacquers, spot removers, rug cleaners, disinfectants, and cosmetic cleansing fluids. Trichloroethylene may be used as a chain terminator in polyvinyl chloride production and as an intermediate in the production of pentachloroethane (Kirk-Othmer, 1963, 1979; IARC, 1979; Defalque, 1961; Wetterhahn, 1972; USCFR, 1976; Valle-Riestra, 1974). Trichloroethylene is no longer used with food, drugs, or cosmetics (IARC, 1979; Food Chemical News, 1978). Before 1976, tolerances for trichloroethylene in decaffeinated ground coffee were set at 25 ppm (USCFR, 1976). Trichloroethylene is no longer used in the decaffeination of coffee.

Chemical and Physical Properties of Trichloroethylene

Selected physical and chemical properties of trichloroethylene are presented in Table 1. Seven reviews on trichloroethylene are available (Huff, 1971; Waters et al., 1976, 1977; Mercier, 1977; Lyman, 1978; IARC, 1979; v. Apeldoorn, 1984; WHO, 1985).

Exposure and Production

The general population is exposed to trichloroethylene at low concentrations (in the parts per billion range) in air, water, and food. Reduction of the use of the chemical in anesthesia, solvent extraction and fumigation of foodstuffs, and dry-cleaning of textiles has reduced exposure from these sources. Exposure during the production of trichloroethylene is relatively low because of the nature of the process. Industrial uses, such as metal degreasing, can involve high exposures. The respiratory route is the principal route of exposure with dermal exposure being an additional route. Oral intake is insignificant in industrial settings (WHO, 1985). In 1981, production of trichloroethylene was 117 million kg

TABLE 1. SELECTED CHEMICAL AND PHYSICAL PROPERTIES OF TRICHLOROETHYLENE

Description: Colorless liquid
Boiling point: 87° C
Density: 1.4642 g/ml at 20° C
Refractive index: $n_D = 1.4773$ at 20° C
Spectroscopy data: $\lambda_{\text{vap}} < 200$ nm
Solubility: Water, 0.1% w/v at 20° C; miscible with acetone, ethanol, diethyl ether, chloroform, and oils
Volatility: Vapor pressure, 77 mm at 25° C
Vapor density: 4.54 (air = 1)
Stability: Nonflammable; when pure and containing a stabilizer, it is stable in presence of air, moisture, and light and in contact with metals up to 130° C. When heated with ozone, it decomposes rapidly into products such as hydrogen chloride, phosgene, carbon monoxide, and chlorine peroxide. At 700° C and above, the vapor decomposes to give a mixture of dichloroethylene, tetrachloroethylene, carbon tetrachloride, chloroform, and methyl chloride. Upon contact with certain metals, high temperatures, open flame, or ultraviolet light, it decomposes almost instantly to phosgene and/or hydrogen chloride, chlorine, and dichloroacetyl chloride. In the presence of alkali, trichloroethylene decomposes to dichloroacetylene.
Reactivity: The most important reaction of trichloroethylene is its oxidative breakdown by atmospheric oxygen, greatly accelerated by elevation of temperature and exposure to light, especially ultraviolet light; not hydrolyzed by water under normal conditions; reacts with alkali under pressure at 150° C to produce glycolic acid and with sulfuric acid to give monochloroacetic acid
Conversion factor: 1 ppm in air is equivalent to 5.37 mg/m³

(USITC, 1982). An estimated 3.5 million workers are exposed to trichloroethylene (Page, 1979).

The recommended threshold limit value for industrial exposure to trichloroethylene is 50 ppm (ACGIH, 1985-86), and the Federal OSHA standard for trichloroethylene is 100 ppm. The California standard is set at 25 ppm.

Teratogenicity and Embryotoxicity

Trichloroethylene has been studied for teratogenic and embryo/fetotoxicologic potential in mice, rats, and rabbits. The results of these studies have not revealed consistent adverse effects in these species.

Mice: Swiss-Webster mice were exposed by inhalation to trichloroethylene at 300 ppm in air 7 hours per day on days 6-15 of gestation (Schwetz et al., 1975). The mice were observed daily throughout pregnancy, and maternal weights were recorded on days 6, 10, and 18 of gestation. No effects were observed on the average number of implantation sites per litter, litter size, incidence of fetal resorptions, fetal sex ratios, or fetal body measurements on day 18 of gestation. The incidences of gross and microscopic anomalies were not significantly greater among exposed than among control litters (Leong et al., 1975; Schwetz et al., 1975).

Rats: Exposure of Sprague Dawley and Charles River rats to trichloroethylene at 300 ppm produced no evidence of teratogenicity or maternal toxicity (Leong et al., 1975; Schwetz et al., 1975). Inhalation exposure of Sprague Dawley rats to trichloroethylene (500 ppm trichloroethylene, 7 hours per day, 5 days per week) during a 3-week pregestational period and on days 0-18 or days 6-18 of gestation caused no maternal or embryonal/fetal toxicity (Beliles et al., 1980). These authors concluded that neither the frequency nor character of the macro- or microscopic findings in the dosed groups indicated an adverse effect.

Female Long-Evans rats exposed to trichloroethylene at 1,800 ppm via inhalation 6 hours per day, 5 days per week, for 2 weeks before mating and/or on days 0-20 of gestation did not exhibit any signs of maternal toxicity (Dorfmueller et al., 1979). There was no indication of embryonic toxicity, and no significant compound-related effects were found in the number of corpora lutea, implantation sites per litter, fetal body weights, resorbed fetuses per litter, or sex ratios. No significant compound-related effects were observed in the analysis of total soft-tissue anomalies. However, prenatal exposure at 1,800 ppm trichloroethylene caused an elevation of incomplete ossification of the sternum, possibly indicative of delays in maturation but not considered to be a major malformation. The study

I. INTRODUCTION

indicated no compound-related effects on general postnatal behavior, but there was a small depression of postnatal weight gains in offspring of the premating exposure group.

Rabbits: Female New Zealand white rabbits were exposed to trichloroethylene at 500 ppm in air 7 hours per day, 5 days per week, for 3 weeks pregestation and on days 0-21 or on days 7-21 of gestation (Beliles et al., 1980). The results revealed no evidence of maternal toxicity or embryotoxicity. The authors did report the occurrence of external hydrocephalus in a few fetuses in one of the study groups (four fetuses per two litters), but no definitive conclusion was made from these findings.

Genetic Toxicology

Trichloroethylene has been reported to be weakly mutagenic or nonmutagenic for *Salmonella typhimurium* strain TA100 (Baden et al., 1979; Bartsch et al., 1979; Simmon et al., 1977; Waskell, 1978). Cerna and Kypenova (1977) reported that trichloroethylene was mutagenic (without metabolic activation) in *S. typhimurium* strains TA1535 and TA1538 and in host-mediated assays with TA1950, TA1951, and TA1952. Greim et al. (1975) reported that microsomally activated reagent-grade trichloroethylene was slightly mutagenic for *Escherichia coli* K12, but Loprieno et al. (1979) found no mutagenic activity in a series of short-term tests using a distilled pure sample of trichloroethylene. Some of this variability may be due to differences in testing protocol, and some may be due to the presence of various stabilizers used in trichloroethylene. NTP studies demonstrated that epichlorohydrin, one chemical frequently used to stabilize trichloroethylene, is clearly mutagenic in Salmonella strains TA100 and TA1535, inducing base-substitution mutations (Canter et al., 1986). In contrast, diisopropylamine, which was the stabilizer in the trichloroethylene used in the NTP 13-week and 2-year rodent studies, is not mutagenic in Salmonella strains TA100, TA1535, TA1537, or TA98 when tested in a preincubation protocol with or without metabolic activation from Aroclor 1254-induced male Sprague Dawley rat or Syrian hamster liver S9 at doses up to 10,000 µg/plate (Mortelmans et al., 1986). Slacik-Erben et al.

(1980) studied trichloroethylene (99.5% pure) in a dominant lethal test with male Han/BGA NMRI mice and found no mutagenic activity. The trichloroethylene used in that study was stabilized with 100 mg/liter triethanolamine, a nonmutagenic chemical (Mortelmans et al., 1986; Gulati et al., 1985; Yoon et al., 1985),

In studies conducted by the NTP (Appendix I), trichloroethylene, stabilized with diisopropylamine, did not cause mutations in *S. typhimurium* strains TA98, TA100, TA1535, or TA1537, with or without metabolic activation by Aroclor 1254-induced male Sprague Dawley rat or male Syrian hamster liver S9 by a liquid-incubation procedure (Mortelmans et al., 1986). Trichloroethylene did not induce chromosomal aberrations in Chinese hamster ovary cells with or without metabolic activation; the results for sister chromatid exchanges were considered positive (Gulati et al., 1985). Trichloroethylene was mutagenic to mouse L5178Y lymphoma cells with but not without activation by Aroclor 1254-induced male F344 rat liver S9.

Carcinogenicity

Evidence for a carcinogenic effect of industrial-grade (greater than 99% pure) trichloroethylene was reported by the National Cancer Institute after the completion of studies in B6C3F₁ mice and Osborne-Mendel rats (NCI, 1976). Trichloroethylene was administered by gavage for 78 weeks; rats were then observed for an additional 32 weeks and mice for 12 weeks. In mice, time-weighted-average gavage doses of 1,170 and 2,340 mg/kg in males and 870 and 1,740 mg/kg in females were associated with significant increases in the incidences of hepatocellular carcinomas. Trichloroethylene administration was associated with increased incidences of alveolar/bronchiolar adenomas and carcinomas in male and female mice (male: vehicle control, 0/20; low dose, 5/50; high dose, 1/48; female: 1/20; 5/50; 7/47). These incidences were not statistically significant and were not considered to be evidence of carcinogenicity in mice. In Osborne-Mendel rats, time-weighted-average gavage doses of 550 and 1,100 mg/kg (both sexes) did not increase the incidences of primary tumors. However, as in several other carcinogenicity studies of chlorinated ethanes and

I. INTRODUCTION

ethylenes (hexachloroethane, NCI, 1978a; 1,1,2,2-tetrachloroethane, NCI, 1978b; 1,1,2-trichloroethane, NCI, 1978c; tetrachloroethylene, NCI, 1977; pentachloroethane, NTP, 1983), the survival of the rats was compromised by the dose regimen.

The results of most of these carcinogenicity studies have been summarized (Weisburger, 1977) and reviewed (IARC, 1979). The International Agency for Research on Cancer (IARC) considered the trichloroethylene studies in Osborne-Mendel rats to be inadequate for evaluation and the studies in B6C3F₁ mice to provide limited evidence of carcinogenicity; that is, trichloroethylene was found to be carcinogenic in one species (IARC, 1979). IARC evaluated trichloroethylene as being "carcinogenic to mice after its oral administration, producing liver and lung neoplasms" (IARC, 1982).

The interpretation of the earlier trichloroethylene study (NCI, 1976) was complicated by the presence of certain additives, particularly epichlorohydrin (0.09%) in the study material. Epichlorohydrin had been previously shown to induce local sarcomas in mice following subcutaneous injection (Van Duuren et al., 1974) and was subsequently shown to cause nasal carcinomas in rats after inhalation exposure (Laskin et al., 1980). Further, epichlorohydrin is a mutagen for *S. typhimurium* strain TA100 (Simmon, 1977). Therefore, although the carcinogenicity of industrial-grade trichloroethylene (containing epichlorohydrin) in B6C3F₁ mice was firmly established, unequivocal statements regarding the carcinogenicity of pure trichloroethylene in mice could not be made.

Results of long-term inhalation studies with purified trichloroethylene (less than 0.25 ppm of each of five chlorinated hydrocarbon impurities as analyzed by gas chromatography/mass spectrophotometry, stabilized with 15 ppm triethanolamine) have been reported (Henschler et al., 1980). In these studies, male and female Wistar rats, NMRI mice, and Syrian hamsters were exposed to air containing up to 500 ppm trichloroethylene for 18 months (6 hours per day, 5 days per week). This regimen did not produce compound-related increases in primary tumors in these species. The investigators did report an

increase in the incidence of malignant lymphomas in female mice (control, 9/29; low dose, 17/30; high dose, 18/28), but the relationship of this lesion to trichloroethylene exposure was considered questionable because of the high incidence of lymphomas in control mice.

In an inhalation experiment, Fukuda et al. (1983) exposed female ICR mice and SD rats to 50, 150, or 450 ppm trichloroethylene (containing 0.019% epichlorohydrin) 7 hours per day, 5 days per week for 104 weeks. These workers reported increased incidences of pulmonary adenocarcinomas in dosed mice (control, 1/49; 50 ppm, 3/50; 150 ppm, 8/50; 450 ppm, 7/46). No increase in liver tumors was reported. There were no increases in the incidences of tumors in female rats.

Henschler and coworkers compared the carcinogenicity of trichloroethylene stabilized with triethanolamine (0.0015%), epichlorohydrin (0.8%), 1,2-epoxybutane (0.8%), or epichlorohydrin plus 1,2-epoxybutane (0.25% and 0.25%) to that of industrial-grade trichloroethylene in groups of 50 male and 50 female ICR/HA-Swiss mice (Henschler et al., 1984). The study compounds were administered by gavage in corn oil 5 days per week. The original doses were 2.4 and 1.8 g/kg of trichloroethylene in males and females, respectively. Because of dose-related toxicity, administration was stopped after 34 weeks and then resumed during week 41 with doses that were one-half the original doses. The animals were not dosed during week 65, and dosing was terminated after week 68. Animals were observed until week 106. Amine-stabilized trichloroethylene did not cause an increased incidence of tumors in either sex. Trichloroethylene stabilized with epichlorohydrin, 1,2-epoxybutane, or both and industrial-grade trichloroethylene caused increases in the incidences of forestomach papillomas and squamous cell carcinomas. Although these results indicate that both epichlorohydrin- and 1,2-epoxybutane-stabilized trichloroethylene are carcinogenic in mice, no conclusion regarding amine-stabilized trichloroethylene can be reached because the shortened duration of administration (61 weeks) reduced the sensitivity of the studies for detecting carcinogenesis.

I. INTRODUCTION

In a more recent study (NTP TR 243, in preparation), epichlorohydrin-free trichloroethylene, administered by gavage, was found to produce hepatocellular neoplasms in male and female B6C3F₁ mice and renal tubular cell adenomas and adenocarcinomas in male F344/N rats. The interpretation of the rat portion of the studies was confounded by the fact that both doses used (500 and 1,000 mg/kg per day) produced significant nontumor renal pathologic effects and significantly reduced the survival of male rats; therefore, the male rat study was judged to be inadequate to evaluate the presence or absence of a carcinogenic response to trichloroethylene. The most striking effect that trichloroethylene produced in rats was toxic nephrosis; it was found in 98% of the dosed males, in all of the dosed females, and in none of the controls. The lesion was first noticed in dosed rats that died early in the studies and was shown microscopically as frank enlargement of the nucleus and cytoplasm of scattered individual tubular cells that had brush borders and were located near the corticomedullary junction. The renal lesion was progressive, as shown by the fact that, as exposure time increased, the affected tubular cells were enlarged and the number of affected tubules and tubular cells was increased. Occasionally, some tubules were enlarged or dilated to the extent that they were difficult to identify as tubules. In animals that survived longer, there was a decrease in the number of enlarged cells, the corresponding tubules were dilated, and portions of the basement membrane had a stripped appearance. Periodic acid Schiff's stain was not useful in attempts to determine if the apparently stripped basement membrane was in fact naked or covered by a thin cytoplasmic membrane extending from the one or more remaining cytomegalic tubular cells. In the most advanced stage, the lesion extended to the subcapsular cortex, where enlarged tubular cells were readily found. Development of cytomegaly did not completely overshadow development of the normal aging rat nephropathy, which was also present.

Although Fukuda et al. (1983) reported increased incidences of pulmonary adenocarcinomas in mice that inhaled trichloroethylene for up to 104 weeks, most of the evidence for the carcinogenicity of trichloroethylene, like that of the

chlorinated ethanes and ethylenes studied earlier (hexachloroethane, NCI, 1978a; 1,1,2,2-tetrachloroethane, NCI, 1978b; 1,1,2-trichloroethane, NCI, 1978c; tetrachloroethylene, NCI, 1977; pentachloroethane, NTP, 1983) comes from data obtained in gavage studies in mice. The interpretation of chemically induced carcinogenicity for these materials is based on increases in the incidences of hepatocellular neoplasms in male and female B6C3F₁ mice. Because this is a relatively common tumor in male mice of this strain (seen in approximately 32% of corn oil vehicle control males and 7% of vehicle control females), the significance of the lesion is frequently debated. Also, the reason for the apparent relative insensitivity of Osborne-Mendel and F344/N rats to the carcinogenic effects of members of this chemical class remains unknown. Possibly increased tumor incidences were not seen in rats because the animals did not survive long enough to develop the lesions.

Inter- or intraspecies differences in susceptibility to the effects of chemicals can be mediated through inherited pharmacokinetic mechanisms. Stott et al. (1982) reported that B6C3F₁ mice metabolize more trichloroethylene (per body weight) than do Osborne-Mendel rats. A similar difference in trichloroethylene metabolism between B6C3F₁ mice and Sprague Dawley rats has been reported (Parchman and Magee, 1982). Prout et al. (1985) compared the metabolism of trichloroethylene in B6C3F₁ and Swiss Webster mice and Osborne-Mendel and Wistar-derived rats. These investigators reported that in mice the metabolism of a single oral dose of trichloroethylene was linear over a dose range of 10-2,000 mg/kg. In rats, however, metabolism was independent of dose at 1,000 mg/kg and above. Consequently, at doses in excess of 1,000 mg/kg, mice are exposed to relatively higher concentrations of trichloroethylene metabolites. If a carcinogenic effect of trichloroethylene requires biotransformation of the parent molecule to a reactive metabolite, B6C3F₁ mice might be expected to be more susceptible than Osborne-Mendel or Sprague Dawley rats because the mice are exposed to higher concentrations of metabolites. Trichloroethylene epoxide has been suggested as an electrophilic metabolite of trichloroethylene (Van Duuren, 1975), but Miller and Guengerich (1982) have reported the results

of in vitro experiments which suggest that the epoxide may not be an intermediate in trichloroethylene metabolism.

Elcomb et al. (1985) administered 10 daily doses of 500, 1,000, or 1,500 mg/kg trichloroethylene by gavage to Osborne-Mendel and Wistar-derived rats and B6C3F₁ and Swiss Webster mice. The dose regimen increased liver weight and decreased hepatic DNA content in both species and increased hepatic DNA synthesis in mice but not in rats. Also, trichloroethylene increased hepatic peroxisomes in mice but not in rats. A relationship between peroxisome proliferation and hepatocarcinogenesis has been suggested (Reddy and Lalwani, 1983). If such a relationship

exists, the insensitivity of rats to trichloroethylene-induced peroxisome proliferation might explain the species differences in hepatocellular responses to the chemical.

Study Rationale

To determine the effects of trichloroethylene in various strains of rats, the National Cancer Institute initiated a series of 2-year toxicology and carcinogenesis studies of trichloroethylene in several strains. The results obtained with F344/N rats and B6C3F₁ mice have been reported (NTP TR 243, in preparation), and the present report describes the results of studies in four additional strains of rats (ACI 9935, August 28807, Marshall 520, and Osborne-Mendel).

II. MATERIALS AND METHODS

**PROCUREMENT AND CHARACTERIZATION OF
TRICHLOROETHYLENE**

**PREPARATION AND CHARACTERIZATION OF
DOSE MIXTURES**

THIRTEEN-WEEK STUDIES

TWO-YEAR STUDIES

Study Design

Source and Specifications of Animals

Animal Maintenance

Clinical Examinations and Pathology

Statistical Methods

II. MATERIALS AND METHODS

PROCUREMENT AND CHARACTERIZATION OF TRICHLOROETHYLENE

High purity "Hi-Tri" trichloroethylene was obtained in two lots from Missouri Solvents (Kansas City, Missouri).

Purity and identity analyses were conducted at Midwest Research Institute (Kansas City, Missouri) (Appendix J). The results of the elemental analyses for both lots were consistent with the theoretical values. Twelve impurities having an area totaling less than 0.04% of the area of the major peak were detected in lot no. TB05-206AA by one gas chromatography system. Eight impurities having an area less than 0.02% that of the major peak were detected in a second system. One impurity with an area of 0.02% that of the major peak was detected in lot no. TB08-039AA.

The infrared and nuclear magnetic resonance spectra of both lots were consistent with the literature spectra. "Hi-Tri" trichloroethylene contains 8 ppm of an amine stabilizer but no epichlorohydrin or 1,2-epoxybutane, as determined by gas chromatography/mass spectrometry.

Throughout the course of these studies, the trichloroethylene was stored at 4° C. Papanicolaou Cancer Research Institute periodically analyzed the study chemical versus a standard maintained at -20° C by gas chromatography with a 10% OV-101 glass column at 70° C. The chemical showed no decrease in purity over the course of the studies, even though a white flocculent material was noticed in the July 1979 reanalysis of lot no. TB05-206AA. A 5-gallon can of this material was returned to Midwest Research Institute for attempted purification. An analysis of the precipitate, which was present at a concentration of 25-30 ppm, is presented in Appendix K. Midwest Research Institute shipped the filtered trichloroethylene back to Papanicolaou Cancer Research Institute in October 1979.

A new lot (TB08-039AA) was received at Papanicolaou in December 1979 and was used immediately.

PREPARATION AND CHARACTERIZATION OF DOSE MIXTURES

Weighed amounts of trichloroethylene were mixed with corn oil to give the desired concentrations of stock solutions (Table 2; Appendix L). Doses were administered at a constant volume. Three of the rat strains received 1.0 ml per dose. Osborne-Mendel rats received 5 ml/kg body weight for the first 7 weeks.

Trichloroethylene in corn oil (1% w/v) was found to be stable for 7 days at room temperature (Appendix L). Later, stock solutions of trichloroethylene in corn oil were found to be stable at 4° C for 8 weeks. Stock solutions were prepared once per week for the first 13 weeks of the 2-year studies and once per month for the remainder of the studies. Stock solutions were analyzed for trichloroethylene content by gas chromatography (Appendix M). Because 201/234 of the dose solutions analyzed were within $\pm 10\%$ of the target concentrations, it is estimated that solutions were prepared within specifications 86% of the time (Table 3; Appendix N). Of the 33 dose mixtures varying from target concentrations by more than 10%, 24 were within $\pm 15\%$ of the target concentrations, 7 were within 20% of the target concentrations, and the remaining 2 were within 25% of the target concentrations.

THIRTEEN-WEEK STUDIES

Thirteen-week studies were conducted to evaluate the cumulative toxicity of trichloroethylene and to determine the doses to be used in the 2-year studies.

Male and female ACI, August, and Marshall rats were obtained from Papanicolaou Cancer Research Institute. August and ACI rats were assigned to cages according to a table of random numbers. Cages were then assigned to dosed and vehicle control groups according to another table of random numbers.

Rats were housed five per cage in polycarbonate cages. Purina Lab Chow® and water (via an

TABLE 2. PREPARATION AND STORAGE OF DOSE MIXTURES IN THE GAVAGE STUDIES OF TRICHLOROETHYLENE

| Thirteen-Week Studies | Two-Year Studies |
|---|--|
| Preparation Trichloroethylene mixed with 100 ml corn oil to prepare stock solution for 1 wk | Trichloroethylene mixed with corn oil in mixing cylinder to prepare stock solution |
| Maximum Storage Time Generally 10 d | 1 wk for the first 13 wk, then 1 mo |
| Storage Conditions 2°-5° C | 2°-5° C |

TABLE 3. SUMMARY OF RESULTS OF ANALYSIS OF DOSE MIXTURES IN THE TWO-YEAR GAVAGE STUDIES OF TRICHLOROETHYLENE

| | |
|---|------------|
| Mean (percent of target concentration) | 102.2 |
| Standard deviation | 6.61 |
| Coefficient of variation (percent) | 6.5 |
| Range (percent of target concentration) | 77.0-121.1 |
| Number of samples | 234 |

automatic watering system) were available ad libitum.

Groups of 10 male ACI and August rats were administered 0, 125, 250, 500, 1,000, or 2,000 mg/kg trichloroethylene in corn oil by gavage 5 days per week for 13 weeks. Groups of 10 female August and ACI rats were administered 0, 62.5, 125, 250, 500, or 1,000 mg/kg on the same schedule. Groups of 10 male Marshall rats were administered 0, 268, 308, 495, 932, or 1,834 mg/kg and groups of 10 female Marshall rats were administered 0, 134, 153, 248, 466, or 918 mg/kg on the same schedule. Further experimental details are summarized in Table 4.

Animals were checked twice daily; moribund animals were killed. Individual animal weights were recorded weekly. Clinical examinations were performed weekly.

At the end of the 13-week studies, survivors were killed. A necropsy was performed on all animals except those excessively autolyzed or cannibalized. Groups examined are listed in Table 4.

TWO-YEAR STUDIES

Study Design

Groups of 50 rats of each sex of four strains (ACI, August, Marshall, Osborne-Mendel) were administered 0, 500, or 1,000 mg/kg trichloroethylene in corn oil by gavage 5 days per week for 103 weeks. Groups of 50 rats of each sex and strain served as untreated controls.

Source and Specifications of Animals

The ACI 9935, August 28807, and Marshall 520 rats used in these studies were produced under strict barrier conditions at Papanicolaou Research Institute under a contract to the Carcinogenesis Program. The Osborne-Mendel rats were produced under similar conditions at CAMM Research Laboratory. Breeding stock for the foundation colonies at the production facilities originated at the National Institutes of Health Repository. The animals were progeny of defined microflora-associated parents that were transferred from isolators to barrier-maintained rooms. Animals were transferred to the study

TABLE 4. EXPERIMENTAL DESIGN AND MATERIALS AND METHODS IN THE GAVAGE STUDIES OF TRICHLOROETHYLENE

| Thirteen-Week Studies | Two-Year Studies |
|--|---|
| EXPERIMENTAL DESIGN | |
| Study Laboratory Papanicolaou Cancer Research Institute (Miami, FL) | Papanicolaou Cancer Research Institute (Miami, FL) |
| Size of Study Groups 10 males and 10 females of each strain | 50 males and 50 females of each strain |
| Doses ACI and August rats: male--0, 125, 250, 500, 1,000, or 2,000 mg/kg trichloroethylene in corn oil by gavage; ACI and August rats: female--0, 62.5, 125, 250, 500, or 1,000 mg/kg; Marshall rats: male--0, 268, 308, 495, 932, or 1,834 mg/kg; female--0, 134, 153, 248, 466, or 918 mg/kg; 1 ml dose volume | 0, 500, or 1,000 mg/kg trichloroethylene in corn oil by gavage; groups of untreated controls also included; 1 ml dose vol (except Osborne-Mendel for the first 7 wk, dose vol--5 ml/kg) |
| Date of First Dose ACI: 3/28/77 | ACI: 2/1/79; August: 10/15/79; Marshall: 12/11/78; Osborne-Mendel: 11/30/79 |
| Date of Last Dose | ACI: 1/22/81; August: 10/2/81; Marshall: 11/28/80; Osborne-Mendel: 11/30/81 |
| Duration of Dosing 5 d/wk for 13 wk | 5 d/wk for 103 wk |
| Type and Frequency of Observation Marshall rats observed 2 × d; weighed 1 × wk | Observed 2 × d; palpated weekly; weighed 1 × wk for 12-15 wk and then 1 × mo |
| Necropsy and Histologic Examination Necropsy performed on all animals; all vehicle control and high dose animals examined microscopically | Necropsy performed on all animals; histologic exam performed on all animals; tissues examined include: gross lesions and tissue masses, skin, mesenteric lymph nodes, mammary gland, salivary glands, thigh muscle, lungs and mainstem bronchi, heart, thyroid gland, parathyroids, esophagus, stomach, duodenum, ileum, colon, liver, vertebrae with bone marrow, thymus, larynx, trachea, pancreas, spleen, kidneys, adrenal glands, urinary bladder, brain, pituitary gland, spinal cord, eyes, seminal vesicles/prostate/testes or ovaries/uterus |
| ANIMALS AND ANIMAL MAINTENANCE | |
| Strain and Species ACI rats; August rats; Marshall rats | ACI 9935 rats; August 28807 rats; Marshall 520 rats; Osborne-Mendel rats |
| Animal Source Papanicolaou Cancer Research Institute | Papanicolaou Cancer Research Institute; except Osborne-Mendel: CAMM Research Lab Animals (Wayne, NJ) |
| Time Held Before Study | ACI: 3.5 wk; August: 2 wk; Marshall: 4 wk; Osborne-Mendel: 3 wk |
| Age When Placed on Study | ACI: 6.5 wk; August: 8 wk; Marshall: 7 wk; Osborne-Mendel: 8 wk |
| Age When Killed | ACI and August: 17-18 wk; Marshall: 110-111 wk |
| Trichloroethylene, NTP TR 273 | 30 |

TABLE 4. EXPERIMENTAL DESIGN AND MATERIALS AND METHODS IN THE GAVAGE STUDIES OF TRICHLOROETHYLENE (Continued)

| Thirteen-Week Studies | Two-Year Studies |
|---|---|
| ANIMALS AND ANIMAL MAINTENANCE (Continued) | |
| Necropsy Dates ACI: 6/27/77; August: 7/19/77; Marshall: 6/13/77 | ACI: 1/29/81-2/17/81; August: 10/12/81-10/15/81; Marshall: 12/8/80-12/16/80; Osborne-Mendel: 11/30/81-12/2/81 |
| Method of Animal Distribution August and ACI: assigned to cages and then to dose groups according to tables of random numbers; Marshall: nonsystematized randomization | Two-step randomization |
| Feed Purina Lab Chow®; available ad libitum | Purina Lab Chow 5001 Pellets® (O.K. Feed Store, Miami, FL); available ad libitum |
| Bedding Wood chips | Sani-chip hardwood (Pinewood Products Co., O.K. Feed Store, Miami, FL) |
| Water Edstrom automatic watering system (Waterford, WI); available ad libitum | Edstrom automatic watering system (Waterford, WI); available ad libitum |
| Cages Polycarbonate; changed 2 × wk | Polycarbonate (Lab Products, Rochelle Park, NJ); changed 2 × wk |
| Cage Filters Filter covers (Monsanto Co., St. Louis, MO) | Cerex-spun nylon (Florida Filters, Miami, FL); changed 2 × mo |
| Animals per Cage 5 | 5 |
| Animal Room Environment Temp--not recorded; hum--not recorded; fluorescent light 12 h/d; 18-20 room air changes/h | Av temp--23° C; humidity--52%-78%; fluorescent light 12 h/d; 10-15 room air changes/h |

laboratory at 3-6 weeks of age and were quarantined for 2-4 weeks. Thereafter, a complete necropsy was performed on five animals of each sex and species to assess their health status. The rodents were placed on study at 6-8 weeks of age. The health of the animals was monitored during the course of the studies according to the protocols of the NTP Sentinel Animal Program (Appendix O).

Animal Maintenance

Rats were housed five per cage in polycarbonate cages. Food and water were available ad libitum. Details of animal maintenance are summarized in Table 4.

Clinical Examinations and Pathology

All animals were observed twice daily, and clinical signs were recorded once per week. Body weights by cage were recorded once per week for the first 12-15 weeks of the studies and once per month thereafter. Mean body weights were calculated for each group. Animals found moribund and those surviving to the end of the studies were humanely killed. A necropsy was performed on all animals including those found dead, unless they were excessively autolyzed or cannibalized, missexed, or found missing. Thus, the number of animals from which particular organs or tissues were examined microscopically

II. MATERIALS AND METHODS

varies and is not necessarily equal to the number of animals that were placed on study.

During necropsy, all organs and tissues were examined for grossly visible lesions. Tissues were preserved in 10% neutral buffered formalin, embedded in paraffin, sectioned, and stained with hematoxylin and eosin. Tissues examined microscopically are listed in Table 4.

When the pathology evaluation was completed, the slides, paraffin blocks, and residual wet tissues were sent to the NCI/NTP Archives for inventory, slide/block match, and wet tissue audit. The slides, individual animal data records, and pathology tables were sent to an independent quality assessment laboratory. The individual animal records and tables were compared for accuracy, slides and tissue counts were verified, and histotechnique was evaluated. All tumor diagnoses, all target tissues, and all tissues from a randomly selected 10% of the animals were evaluated by a quality assessment pathologist. The quality assessment report and slides were submitted to the Pathology Working Group (PWG) Chairperson, who reviewed all target tissues and those about which there was a disagreement between the laboratory and quality assessment pathologists.

Representative slides selected by the Chairperson were reviewed by the PWG, which includes the laboratory pathologist, without knowledge of previously rendered diagnoses. When the consensus diagnosis of the PWG differed from that of the laboratory pathologist, the laboratory pathologist was asked to reconsider the original diagnosis. This procedure has been described, in part, by Maronpot and Boorman (1982) and Boorman et al. (1985). The final diagnoses represent a consensus of contractor pathologists and the NTP Pathology Working Group. For subsequent analysis of pathology data, the diagnosed lesions for each tissue type are combined according to the guidelines of McConnell et al. (1986).

Slides/tissues are generally not evaluated in a blind fashion (i.e., without knowledge of dose group) unless the lesions in question are subtle or unless there is an inconsistent diagnosis of lesions by the laboratory pathologist. Nonneoplastic lesions are not examined routinely by the

quality assessment pathologist or PWG unless they are considered part of the toxic effect of the chemical.

Statistical Methods

Data Recording: Data on this experiment were recorded in the Carcinogenesis Bioassay Data System (Linhart et al., 1974). The data elements include descriptive information on the chemicals, animals, experimental design, survival, body weight, and individual pathologic results, as recommended by the International Union Against Cancer (Berenblum, 1969).

Survival Analyses: The probability of survival was estimated by the product-limit procedure of Kaplan and Meier (1958) and is presented in the form of graphs. Animals were censored from the survival analyses at the time they were found to be missing or dead of other than natural causes; animals dying from natural causes were not censored. Statistical analyses for a possible dose-related effect on survival used the method of Cox (1972) for testing two groups for equality and Tarone's (1975) life table test for a dose-related trend. All reported P values for the survival analysis are two-sided.

Calculation of Incidence: The incidence of neoplastic or nonneoplastic lesions is given as the ratio of the number of animals bearing such lesions at a specific anatomic site to the number of animals in which that site was examined. In most instances, the denominators include only those animals for which the site was examined histologically. However, when macroscopic examination was required to detect lesions (e.g., skin or mammary tumors) prior to histologic sampling or when lesions could have appeared at multiple sites (e.g., lymphomas), the denominators consist of the number of animals on which a necropsy was performed.

Analysis of Tumor Incidence: Three statistical methods are used to analyze tumor incidence data. The two that adjust for intercurrent mortality employ the classical method for combining contingency tables developed by Mantel and Haenszel (1959). Tests of significance included pairwise comparisons of high dose and low dose groups with vehicle controls and tests for overall dose-response trends.

II. MATERIALS AND METHODS

For studies in which compound administration has little effect on survival, the results of the three alternative analyses will generally be similar. When differing results are obtained by the three methods, the final interpretation of the data will depend on the extent to which the tumor under consideration is regarded as being the cause of death. Continuity-corrected tests are used in the analysis of tumor incidence, and reported P values are one-sided.

Life Table Analysis--The first method of analysis assumed that all tumors of a given type observed in animals dying before the end of the study were "fatal"; i.e., they either directly or indirectly caused the death of the animal. According to this approach, the proportions of tumor-bearing animals in the dosed and vehicle control groups were compared at each point in time at which an animal died with a tumor of interest. The denominators of these proportions were the total number of animals at risk in each group. These results, including the data from animals killed at the end of the study, were then combined by the Mantel-Haenszel method to obtain an overall P value. This method of adjusting for intercurrent mortality is the life table method of Cox (1972) and of Tarone (1975). The underlying variable considered by this analysis is time to death due to tumor. If the tumor is rapidly lethal, then time to death due to tumor closely approximates time to tumor onset. In this case, the life table test also provides a comparison of the time-specific tumor incidence.

Incidental Tumor Analysis--The second method of analysis assumed that all tumors of a given type observed in animals that died before the

end of the study were "incidental"; i.e., they were merely observed at necropsy in animals dying of an unrelated cause. According to this approach, the proportions of tumor-bearing animals in dosed and vehicle control groups were compared in each of five time intervals: weeks 0-52, weeks 53-78, weeks 79-92, week 93 to the week before the terminal-kill period, and the terminal-kill period. The denominators of these proportions were the number of animals actually examined for tumors during the time interval. The individual time interval comparisons were then combined by the previously described method to obtain a single overall result. (See Haseman, 1984, for the computational details of both methods.)

Unadjusted Analyses--Primarily, survival-adjusted methods are used to evaluate tumor incidence. In addition, the results of the Fisher exact test for pairwise comparisons and the Cochran-Armitage linear trend test (Armitage, 1971; Gart et al., 1979) are given in the appendixes containing the analyses of primary tumor incidence. These two tests are based on the overall proportion of tumor-bearing animals and do not adjust for survival differences.

Historical Control Data: Although the concurrent control group is always the first and most appropriate control group used for evaluation, there are certain instances in which historical control data can be helpful in the overall assessment of tumor incidence. Consequently, control tumor incidences from the NTP historical control data base (Haseman et al., 1984, 1985) are included for those tumors appearing to show compound-related effects.

III. RESULTS

THIRTEEN-WEEK STUDIES

TWO-YEAR STUDIES

Body Weights and Clinical Signs

Survival

Pathology and Statistical Analyses of Results

Nonneoplastic Lesions of the Kidney

Neoplastic Lesions

Negative Trends and Lower Incidences

III. RESULTS

THIRTEEN-WEEK STUDIES

Thirteen-week studies were conducted on male and female ACI, August, and Marshall rats. Data from previous studies of Osborne-Mendel rats (NCI, 1976) were used to set doses for the 2-year studies of the Osborne-Mendel strain. The results of the current 13-week studies are condensed, for comparative purposes, and presented in Tables 5 and 6. More detailed information is presented in Appendix P.

Doses used for ACI and August rats ranged from 125 to 2,000 mg/kg for males and 62.5 to 1,000 mg/kg for females. Doses for Marshall rats ranged from 268 to 1,834 mg/kg for males and 134 to 918 mg/kg for females.

With the exception of three male August rats that received 2,000 mg/kg, all animals survived to the end of the 13-week experimental period. In male rats, the highest dose administered reduced final mean body weights (relative to vehicle controls) in all three strains by more than 10%. These reductions ranged from 12% in Marshall rats to 17% in ACI rats. In females, final mean body weights relative to vehicle controls were somewhat depressed; however, none was depressed in excess of 8% (August). Body weight changes (percent) in female rats administered the highest dose (1,000 mg/kg in ACI and August; 918 mg/kg in Marshall) were comparable

to those in males administered similar doses (1,000 mg/kg in ACI and August; 932 mg/kg in Marshall). No behavioral effects attributable to trichloroethylene administration were recorded. Similarly, the administration of the chemical for 13 weeks was not associated with histopathologic changes.

Dose Selection Rationale: Doses selected for the 2-year studies in ACI, August, Marshall, and Osborne-Mendel rats (both sexes) were 0, 500, and 1,000 mg/kg in corn oil by gavage, administered 5 days per week. The selection of these doses was based on survival (all rats dosed with 1,000 mg/kg trichloroethylene for 13 weeks survived for the entire study), body weight gains (in male rats, mean body weights of dosed animals at the end of the 13-week studies were from 91% to 96% those of vehicle controls; in females, final mean body weights of dosed animals were from 93% to 96% those of vehicle controls), and absence of histopathologic changes produced in ACI, August, and Marshall rats during the 13-week studies. Doses for the Osborne-Mendel rats (also 0, 500, and 1,000 mg/kg per day) were similar to the doses used in the earlier 2-year studies (NCI, 1976). The doses in those studies, expressed as time-weighted averages, were 550 and 1,100 mg/kg and significantly reduced survival in each sex. Both untreated control and corn oil vehicle control groups were used in the present studies.

TABLE 5. COMPARISON OF EFFECTS OF TRICHLOROETHYLENE ON SURVIVAL AND BODY WEIGHTS IN MALE ACI, AUGUST, AND MARSHALL RATS IN THE THIRTEEN-WEEK GAVAGE STUDIES (a)

| ACI | | | August | | | Marshall | | |
|--------------|----------|--|--------------|----------|--|--------------|----------|--|
| Dose (mg/kg) | Survival | Final Body Weight Relative to Veh. Cont. (percent) | Dose (mg/kg) | Survival | Final Body Weight Relative to Veh. Cont. (percent) | Dose (mg/kg) | Survival | Final Body Weight Relative to Veh. Cont. (percent) |
| 0 | 10/10 | -- | 0 | 10/10 | -- | 0 | 10/10 | -- |
| 125 | 10/10 | 99.6 | 125 | 10/10 | 101.2 | 268 | 10/10 | 103.2 |
| 250 | 10/10 | 94.2 | 250 | 10/10 | 96.7 | 308 | 10/10 | 104.9 |
| 500 | 10/10 | 97.7 | 500 | 10/10 | 95.9 | 495 | 10/10 | 100.8 |
| 1,000 | 10/10 | 90.7 | 1,000 | 10/10 | 95.9 | 932 | 10/10 | 93.9 |
| 2,000 | 10/10 | 82.6 | 2,000 | 7/10 | 84.6 | 1,834 | 10/10 | 87.9 |

(a) Trichloroethylene was mixed with corn oil and administered by gavage 5 d/wk for 13 weeks.

TABLE 6. COMPARISON OF EFFECTS OF TRICHLOROETHYLENE ON SURVIVAL AND BODY WEIGHTS IN FEMALE ACI, AUGUST, AND MARSHALL RATS IN THE THIRTEEN-WEEK GAVAGE STUDIES (a)

| ACI | | | August | | | Marshall | | |
|--------------|----------|--|--------------|----------|--|--------------|----------|--|
| Dose (mg/kg) | Survival | Final Body Weight Relative to Veh. Cont. (percent) | Dose (mg/kg) | Survival | Final Body Weight Relative to Veh. Cont. (percent) | Dose (mg/kg) | Survival | Final Body Weight Relative to Veh. Cont. (percent) |
| 0 | 10/10 | -- | 0 | 10/10 | -- | 0 | 10/10 | -- |
| 62.5 | 10/10 | 99.4 | 62.5 | 10/10 | 101.4 | 134 | 10/10 | 101.2 |
| 125 | 10/10 | 94.8 | 125 | 10/10 | 100.5 | 153 | 10/10 | 102.9 |
| 250 | 10/10 | 98.9 | 250 | 10/10 | 100.5 | 248 | 10/10 | 102.3 |
| 500 | 10/10 | 92.0 | 500 | 10/10 | 98.6 | 466 | 10/10 | 97.1 |
| 1,000 | 10/10 | 93.1 | 1,000 | 10/10 | 92.8 | 918 | 10/10 | 95.9 |

(a) Trichloroethylene was mixed with corn oil and administered by gavage 5 d/wk for 13 weeks.

TWO-YEAR STUDIES

Body Weights and Clinical Signs

Detailed data for each strain are presented in Tables 7 through 10 and in Figures 1 through 4. The final body weights (relative to vehicle controls) and percent of animals surviving to terminal kill are summarized in Table 11.

The final mean body weights of all dosed groups except for high dose female Osborne-Mendel rats were somewhat lower than those of the vehicle controls. Trichloroethylene at the 1,000 mg/kg dose reduced mean final body weight by 10% or more in ACI, August, and Osborne-Mendel males and Marshall females. Final mean body weights in other high dose groups ranged from 3% lower in Marshall males to 8% lower in August females. Final mean body weight depression at the 500 mg/kg dose exceeded 10% only in the ACI males. Final body weight decrements in other low dose groups ranged from 3% in Osborne-Mendel males to 8% in ACI females.

Clinical signs of central nervous system toxicity were observed in dosed animals. Male and

female rats of all strains exhibited sporadic ataxia, lethargy, convulsions, and hindlimb paralysis after dosing. On several occasions, animals in all dosed groups lost consciousness. These effects were generally transient; however, later in the studies, some animals convulsed before dosing and during weighing periods.

Survival

Estimates of the probabilities of survival for each strain of male and female rats at the doses used in these studies and for untreated and vehicle controls are shown in Table 12 and in the Kaplan and Meier curves in Figures 5, 7, 9, and 11. Unadjusted survival curves are presented in Figures 6, 8, 10, and 12; in these curves, animals killed accidentally have not been censored.

Survival was significantly reduced in ACI males and high dose females, in high dose female Osborne-Mendel rats, and in female Marshall rats. Survival of male Marshall rats at both doses was reduced because of mortality that occurred during the administration of the chemical.

TABLE 7. MEAN BODY WEIGHTS AND SURVIVAL OF ACI RATS IN THE TWO-YEAR GAVAGE STUDIES OF TRICHLOROETHYLENE

| Weeks on Study | Vehicle Control | | 500 mg/kg | | | 1,000 mg/kg | | |
|----------------|-----------------|------------------|-----------------|--------------------------------|------------------|-----------------|--------------------------------|------------------|
| | Av. Wt. (grams) | No. of Survivors | Av. Wt. (grams) | Wt. (percent of veh. controls) | No. of Survivors | Av. Wt. (grams) | Wt. (percent of veh. controls) | No. of Survivors |
| MALE | | | | | | | | |
| 0 | 169 | 50 | 158 | 93 | 50 | 162 | 96 | 50 |
| 1 | 188 | 50 | 182 | 97 | 50 | 183 | 97 | 50 |
| 2 | 205 | 50 | 198 | 97 | 50 | 199 | 97 | 50 |
| 3 | 220 | 50 | 213 | 97 | 50 | 211 | 96 | 49 |
| 4 | 233 | 50 | 227 | 97 | 50 | 221 | 95 | 49 |
| 5 | 247 | 50 | 237 | 96 | 49 | 231 | 94 | 48 |
| 6 | 252 | 50 | 242 | 96 | 49 | 239 | 95 | 47 |
| 7 | 260 | 50 | 255 | 98 | 48 | 249 | 96 | 47 |
| 8 | 270 | 50 | 258 | 96 | 47 | 252 | 93 | 47 |
| 9 | 275 | 50 | 264 | 96 | 47 | 260 | 95 | 47 |
| 10 | 280 | 50 | 253 | 90 | 47 | 260 | 93 | 46 |
| 11 | 288 | 50 | 259 | 90 | 47 | 257 | 89 | 46 |
| 12 | 283 | 49 | 268 | 95 | 45 | 262 | 93 | 45 |
| 13 | 283 | 49 | 267 | 94 | 45 | 264 | 93 | 42 |
| 14 | 283 | 49 | 260 | 92 | 45 | 256 | 90 | 42 |
| 15 | 291 | 49 | 260 | 89 | 45 | 256 | 88 | 42 |
| 17 | 296 | 49 | 266 | 90 | 45 | 268 | 91 | 42 |
| 21 | 296 | 49 | 276 | 93 | 45 | 274 | 93 | 38 |
| 26 | 327 | 49 | 297 | 91 | 41 | 299 | 91 | 30 |
| 30 | 342 | 49 | 311 | 91 | 40 | 306 | 89 | 35 |
| 34 | 339 | 49 | 316 | 93 | 36 | 310 | 91 | 32 |
| 38 | .. | .. | 308 | .. | 36 | 316 | .. | 30 |
| 39 | 340 | 48 | .. | .. | .. | .. | .. | .. |
| 42 | .. | .. | 318 | .. | 36 | 307 | .. | 30 |
| 45 | 344 | 47 | 325 | 94 | 33 | 314 | 91 | 28 |
| 50 | 350 | 46 | 327 | 93 | 32 | 314 | 90 | 26 |
| 54 | .. | .. | 319 | .. | 29 | 307 | .. | 23 |
| 58 | 343 | 46 | 320 | 93 | 29 | 310 | 90 | 23 |
| 61 | 345 | 45 | 325 | 94 | 29 | 311 | 90 | 22 |
| 65 | 350 | 45 | 320 | 91 | 27 | 313 | 89 | 19 |
| 69 | 357 | 45 | 328 | 92 | 26 | 322 | 90 | 19 |
| 73 | 357 | 44 | 327 | 92 | 26 | 316 | 89 | 18 |
| 77 | 361 | 43 | 330 | 91 | 26 | 323 | 89 | 14 |
| 81 | 364 | 40 | 344 | 95 | 26 | 322 | 88 | 14 |
| 85 | 358 | 39 | 323 | 90 | 25 | 319 | 89 | 14 |
| 89 | 355 | 39 | 319 | 90 | 24 | 316 | 89 | 13 |
| 94 | 361 | 39 | 316 | 88 | 21 | 314 | 87 | 13 |
| 97 | 359 | 39 | 313 | 87 | 20 | 309 | 86 | 12 |
| 98 | 357 | 39 | 312 | 87 | 20 | 315 | 88 | 12 |
| 101 | 353 | 38 | 313 | 89 | 20 | 311 | 88 | 12 |
| 103 | 353 | 38 | 314 | 89 | 19 | 309 | 88 | 11 |
| FEMALE | | | | | | | | |
| 0 | 129 | 50 | 127 | 98 | 50 | 126 | 98 | 50 |
| 1 | 139 | 50 | 139 | 100 | 50 | 137 | 99 | 50 |
| 2 | 148 | 50 | 141 | 95 | 50 | 144 | 97 | 50 |
| 3 | 153 | 50 | 150 | 98 | 50 | 151 | 99 | 50 |
| 4 | 157 | 50 | 157 | 100 | 50 | 156 | 99 | 50 |
| 5 | 169 | 50 | 164 | 97 | 50 | 162 | 96 | 49 |
| 6 | 173 | 50 | 166 | 96 | 50 | 165 | 95 | 49 |
| 7 | 171 | 50 | 173 | 101 | 50 | 171 | 100 | 49 |
| 8 | 182 | 50 | 172 | 95 | 50 | 172 | 95 | 49 |
| 9 | 172 | 50 | 174 | 101 | 50 | 175 | 102 | 48 |
| 10 | 181 | 50 | 161 | 89 | 50 | 172 | 95 | 47 |
| 11 | 180 | 50 | 175 | 97 | 49 | 177 | 98 | 47 |
| 12 | 187 | 50 | 178 | 95 | 49 | 181 | 97 | 46 |
| 13 | 189 | 50 | 180 | 95 | 48 | 181 | 96 | 46 |
| 14 | 184 | 49 | 182 | 99 | 48 | 178 | 97 | 46 |
| 15 | 184 | 49 | 182 | 99 | 48 | 178 | 97 | 46 |
| 17 | 185 | 49 | 181 | 98 | 47 | 186 | 101 | 43 |
| 21 | 188 | 49 | 186 | 99 | 45 | 186 | 99 | 43 |
| 26 | 206 | 48 | 198 | 96 | 45 | 201 | 98 | 41 |
| 30 | 218 | 48 | 200 | 92 | 45 | 205 | 94 | 39 |
| 34 | 216 | 48 | 201 | 93 | 44 | 207 | 96 | 39 |
| 38 | .. | .. | 202 | .. | 44 | 204 | .. | 39 |
| 39 | 212 | 48 | .. | .. | .. | .. | .. | .. |
| 42 | .. | .. | 205 | .. | 44 | 205 | .. | 38 |
| 45 | 213 | 47 | 211 | 99 | 44 | 208 | 98 | 36 |
| 50 | 217 | 47 | 212 | 98 | 41 | 213 | 98 | 36 |
| 54 | .. | .. | 203 | .. | 40 | 205 | .. | 33 |
| 58 | 224 | 45 | 209 | 93 | 40 | 214 | 96 | 31 |
| 61 | 226 | 45 | 213 | 94 | 40 | 217 | 96 | 31 |
| 65 | 225 | 45 | 210 | 93 | 35 | 214 | 95 | 27 |
| 69 | 230 | 44 | 213 | 93 | 33 | 220 | 96 | 27 |
| 73 | 232 | 44 | 214 | 92 | 31 | 211 | 91 | 25 |
| 77 | 238 | 44 | 222 | 93 | 29 | 223 | 94 | 24 |
| 81 | 240 | 44 | 211 | 88 | 29 | 221 | 92 | 24 |
| 85 | 237 | 43 | 217 | 92 | 28 | 221 | 93 | 22 |
| 89 | 239 | 43 | 219 | 92 | 23 | 224 | 94 | 21 |
| 94 | 246 | 42 | 219 | 89 | 21 | 227 | 92 | 21 |
| 97 | 244 | 39 | 221 | 91 | 21 | 226 | 93 | 21 |
| 98 | 243 | 39 | 218 | 90 | 21 | 229 | 94 | 20 |
| 101 | 242 | 36 | 222 | 92 | 20 | 227 | 94 | 20 |
| 103 | 242 | 35 | 222 | 92 | 20 | 225 | 93 | 19 |





FIGURE 1. GROWTH CURVES FOR ACI RATS ADMINISTERED TRICHLOROETHYLENE IN CORN OIL BY GAVAGE FOR TWO YEARS

TABLE 8. MEAN BODY WEIGHTS AND SURVIVAL OF AUGUST RATS IN THE TWO-YEAR GAVAGE STUDIES OF TRICHLOROETHYLENE

| Weeks on Study | Vehicle Control | | 500 mg/kg | | | 1,000 mg/kg | | |
|----------------|-----------------|------------------|-----------------|--------------------------------|------------------|-----------------|--------------------------------|------------------|
| | Av. Wt. (grams) | No. of Survivors | Av. Wt. (grams) | Wt. (percent of veh. controls) | No. of Survivors | Av. Wt. (grams) | Wt. (percent of veh. controls) | No. of Survivors |
| MALE | | | | | | | | |
| 0 | 162 | 50 | 167 | 103 | 50 | 163 | 101 | 50 |
| 1 | 193 | 50 | 186 | 96 | 50 | 183 | 95 | 50 |
| 2 | 227 | 50 | 223 | 98 | 50 | 216 | 95 | 50 |
| 3 | 241 | 50 | 209 | 87 | 50 | 232 | 96 | 50 |
| 4 | 256 | 50 | 238 | 93 | 50 | 242 | 95 | 50 |
| 5 | 234 | 50 | 248 | 106 | 50 | 250 | 107 | 50 |
| 6 | 245 | 49 | 257 | 105 | 50 | 262 | 107 | 49 |
| 7 | 266 | 49 | 270 | 102 | 49 | 273 | 103 | 48 |
| 8 | 271 | 49 | 279 | 103 | 49 | 277 | 102 | 46 |
| 10 | 292 | 49 | 287 | 98 | 48 | 288 | 99 | 46 |
| 12 | 297 | 49 | 295 | 99 | 48 | 289 | 97 | 46 |
| 13 | 303 | 49 | 300 | 99 | 48 | 296 | 98 | 46 |
| 16 | 319 | 49 | 320 | 100 | 45 | 308 | 97 | 45 |
| 20 | 322 | 49 | 318 | 99 | 44 | 320 | 99 | 45 |
| 25 | 345 | 49 | 336 | 97 | 42 | 330 | 96 | 43 |
| 30 | 354 | 49 | 351 | 99 | 40 | 329 | 93 | 40 |
| 33 | 366 | 49 | 359 | 98 | 38 | 345 | 94 | 37 |
| 38 | 381 | 49 | 370 | 97 | 36 | 351 | 92 | 37 |
| 41 | 391 | 49 | 372 | 95 | 36 | 356 | 91 | 37 |
| 45 | 396 | 49 | 370 | 93 | 36 | 357 | 90 | 36 |
| 49 | 399 | 49 | -- | -- | -- | -- | -- | -- |
| 50 | -- | -- | 375 | -- | 36 | 362 | -- | 34 |
| 54 | 407 | 47 | 382 | 94 | 35 | 364 | 89 | 34 |
| 57 | 407 | 47 | -- | -- | -- | -- | -- | -- |
| 58 | -- | -- | 382 | -- | 35 | 371 | -- | 33 |
| 62 | 413 | 46 | 380 | 92 | 35 | 366 | 89 | 30 |
| 65 | 418 | 48 | 380 | 91 | 33 | 363 | 87 | 30 |
| 69 | -- | -- | 389 | -- | 29 | 371 | -- | 26 |
| 70 | 416 | 42 | -- | -- | -- | -- | -- | -- |
| 73 | 421 | 40 | 385 | 91 | 28 | 374 | 89 | 23 |
| 76 | 417 | 38 | 382 | 92 | 27 | 359 | 86 | 23 |
| 79 | 412 | 36 | 399 | 97 | 25 | 368 | 89 | 23 |
| 83 | 409 | 36 | 377 | 92 | 24 | 365 | 89 | 22 |
| 87 | 408 | 33 | 364 | 89 | 23 | 362 | 89 | 20 |
| 91 | 405 | 31 | 372 | 92 | 19 | 362 | 89 | 19 |
| 96 | 399 | 29 | 363 | 91 | 17 | 356 | 89 | 18 |
| 100 | 391 | 25 | 364 | 93 | 16 | 336 | 86 | 16 |
| 104 | 382 | 21 | 357 | 93 | 13 | 335 | 88 | 16 |
| FEMALE | | | | | | | | |
| 0 | 127 | 50 | 124 | 98 | 50 | 127 | 100 | 50 |
| 1 | 137 | 50 | 131 | 96 | 50 | 134 | 98 | 50 |
| 2 | 148 | 50 | 147 | 99 | 50 | 148 | 100 | 50 |
| 3 | 157 | 50 | 140 | 89 | 50 | 156 | 99 | 50 |
| 4 | 169 | 50 | 161 | 95 | 50 | 152 | 90 | 50 |
| 5 | 153 | 50 | 166 | 108 | 50 | 167 | 109 | 50 |
| 6 | 163 | 50 | 166 | 102 | 50 | 165 | 101 | 50 |
| 7 | 170 | 50 | 173 | 102 | 50 | 175 | 103 | 50 |
| 8 | 170 | 50 | 177 | 104 | 50 | 169 | 99 | 50 |
| 10 | 176 | 50 | 175 | 99 | 50 | 179 | 102 | 50 |
| 12 | 181 | 50 | 179 | 99 | 50 | 179 | 99 | 50 |
| 13 | 179 | 50 | 180 | 101 | 48 | 182 | 102 | 50 |
| 16 | 188 | 50 | 184 | 98 | 48 | 188 | 100 | 49 |
| 20 | 189 | 50 | 193 | 102 | 45 | 194 | 103 | 46 |
| 25 | 196 | 50 | 194 | 99 | 45 | 197 | 101 | 44 |
| 30 | 189 | 50 | 197 | 104 | 45 | 191 | 101 | 43 |
| 33 | 202 | 50 | 198 | 98 | 45 | 201 | 100 | 42 |
| 38 | 203 | 50 | 203 | 100 | 45 | 203 | 100 | 41 |
| 41 | 204 | 50 | 205 | 100 | 45 | 206 | 101 | 41 |
| 45 | 207 | 50 | 205 | 99 | 44 | 206 | 100 | 40 |
| 49 | 209 | 49 | -- | -- | -- | -- | -- | -- |
| 50 | -- | -- | 206 | -- | 43 | 208 | -- | 38 |
| 54 | 212 | 48 | 207 | 98 | 43 | 209 | 99 | 38 |
| 57 | 214 | 48 | -- | -- | -- | -- | -- | -- |
| 58 | -- | -- | 209 | -- | 42 | 209 | -- | 35 |
| 62 | 216 | 46 | 208 | 96 | 42 | 208 | 96 | 32 |
| 65 | 220 | 45 | 212 | 96 | 41 | 207 | 94 | 31 |
| 69 | -- | -- | 216 | -- | 40 | 219 | -- | 30 |
| 70 | 222 | 45 | -- | -- | -- | -- | -- | -- |
| 73 | 227 | 43 | 217 | 98 | 38 | 217 | 96 | 29 |
| 76 | 226 | 43 | 218 | 98 | 38 | 214 | 95 | 29 |
| 79 | 231 | 42 | 223 | 97 | 36 | 217 | 94 | 29 |
| 83 | 237 | 40 | 224 | 95 | 35 | 217 | 92 | 28 |
| 87 | 235 | 38 | 226 | 98 | 34 | 219 | 93 | 27 |
| 91 | 232 | 35 | 228 | 97 | 34 | 219 | 94 | 26 |
| 96 | 241 | 33 | 225 | 93 | 32 | 221 | 92 | 26 |
| 100 | 242 | 29 | 225 | 93 | 30 | 219 | 90 | 26 |
| 104 | 238 | 23 | 225 | 95 | 26 | 220 | 92 | 25 |





FIGURE 2. GROWTH CURVES FOR AUGUST RATS ADMINISTERED TRICHLOROETHYLENE IN CORN OIL BY GAVAGE FOR TWO YEARS

TABLE 9. MEAN BODY WEIGHTS AND SURVIVAL OF MARSHALL RATS IN THE TWO-YEAR GAVAGE STUDIES OF TRICHLOROETHYLENE

| Weeks on Study | Vehicle Control | | 500 mg/kg | | | 1,000 mg/kg | | |
|----------------|-----------------|------------------|-----------------|--------------------------------|------------------|-----------------|--------------------------------|------------------|
| | Av. Wt. (grams) | No. of Survivors | Av. Wt. (grams) | Wt. (percent of veh. controls) | No. of Survivors | Av. Wt. (grams) | Wt. (percent of veh. controls) | No. of Survivors |
| MALE | | | | | | | | |
| 1 | 196 | 50 | 199 | 102 | 50 | 189 | 96 | 50 |
| 2 | 217 | 50 | 217 | 100 | 50 | 200 | 92 | 50 |
| 3 | 232 | 50 | 231 | 100 | 50 | 218 | 94 | 50 |
| 4 | 241 | 50 | 238 | 99 | 50 | 225 | 93 | 50 |
| 5 | 251 | 50 | 246 | 98 | 50 | 232 | 92 | 50 |
| 6 | 260 | 50 | 238 | 92 | 50 | 240 | 92 | 49 |
| 7 | 260 | 50 | 254 | 98 | 50 | 243 | 93 | 49 |
| 8 | 248 | 50 | 250 | 101 | 49 | 242 | 98 | 49 |
| 9 | 262 | 50 | 258 | 98 | 49 | 247 | 94 | 49 |
| 10 | 269 | 50 | 265 | 99 | 49 | 255 | 95 | 49 |
| 11 | 273 | 50 | 269 | 99 | 46 | 256 | 94 | 49 |
| 12 | 274 | 50 | 271 | 99 | 45 | 257 | 94 | 48 |
| 16 | 292 | 50 | 281 | 96 | 43 | 272 | 93 | 48 |
| 20 | 295 | 50 | 296 | 100 | 41 | 281 | 95 | 46 |
| 24 | 302 | 49 | 302 | 100 | 36 | 287 | 95 | 46 |
| 28 | 311 | 49 | 307 | 99 | 35 | 281 | 90 | 43 |
| 31 | -- | -- | 313 | -- | 34 | 296 | -- | 41 |
| 32 | 324 | 49 | -- | -- | -- | -- | -- | -- |
| 35 | 323 | 49 | 322 | 100 | 33 | 303 | 94 | 39 |
| 39 | 323 | 49 | 326 | 101 | 32 | 308 | 95 | 39 |
| 43 | 319 | 48 | 319 | 100 | 31 | 308 | 97 | 37 |
| 47 | 311 | 48 | 312 | 100 | 29 | 304 | 98 | 35 |
| 51 | 299 | 47 | 310 | 104 | 27 | 298 | 100 | 32 |
| 54 | 309 | 45 | 304 | 98 | 25 | 300 | 97 | 31 |
| 58 | 313 | 42 | 311 | 99 | 24 | 299 | 96 | 24 |
| 62 | 313 | 41 | 306 | 98 | 23 | 296 | 95 | 24 |
| 66 | -- | -- | 291 | -- | 19 | 282 | -- | 23 |
| 67 | 310 | 40 | -- | -- | -- | -- | -- | -- |
| 70 | 315 | 38 | 296 | 94 | 14 | 296 | 94 | 21 |
| 74 | 314 | 38 | 294 | 94 | 12 | 292 | 93 | 18 |
| 78 | 315 | 38 | 300 | 95 | 12 | 300 | 95 | 16 |
| 82 | 308 | 36 | 291 | 94 | 12 | 298 | 97 | 14 |
| 86 | 317 | 33 | 300 | 95 | 12 | 298 | 94 | 12 |
| 90 | 315 | 31 | 307 | 97 | 12 | 302 | 96 | 9 |
| 94 | 315 | 28 | 299 | 95 | 12 | 299 | 95 | 9 |
| 96 | 312 | 28 | 302 | 97 | 12 | 307 | 98 | 9 |
| 98 | 313 | 26 | 299 | 96 | 12 | 304 | 97 | 9 |
| 100 | 313 | 26 | 292 | 93 | 12 | 303 | 97 | 7 |
| 102 | 315 | 26 | -- | -- | -- | -- | -- | -- |
| 103 | -- | -- | 300 | -- | 12 | 307 | -- | 7 |
| FEMALE | | | | | | | | |
| 1 | 144 | 50 | 140 | 97 | 50 | 143 | 99 | 50 |
| 2 | 155 | 50 | 149 | 96 | 50 | 152 | 98 | 50 |
| 3 | 161 | 50 | 155 | 96 | 50 | 157 | 98 | 50 |
| 4 | 166 | 50 | 161 | 97 | 50 | 161 | 97 | 50 |
| 5 | 171 | 50 | 161 | 94 | 50 | 161 | 94 | 50 |
| 6 | 169 | 50 | 167 | 99 | 50 | 165 | 98 | 50 |
| 7 | 174 | 50 | 169 | 97 | 50 | 170 | 98 | 50 |
| 8 | 168 | 50 | 172 | 102 | 50 | 172 | 102 | 50 |
| 9 | 178 | 50 | 174 | 98 | 50 | 171 | 98 | 48 |
| 10 | 182 | 50 | 179 | 98 | 50 | 178 | 98 | 48 |
| 11 | 179 | 50 | 181 | 101 | 50 | 177 | 99 | 48 |
| 12 | 187 | 50 | 180 | 96 | 50 | 178 | 95 | 48 |
| 16 | 196 | 50 | 190 | 97 | 49 | 189 | 96 | 48 |
| 20 | 205 | 50 | 200 | 98 | 48 | 199 | 97 | 48 |
| 24 | 217 | 50 | 202 | 93 | 48 | 200 | 92 | 48 |
| 28 | 216 | 50 | 208 | 96 | 48 | 199 | 92 | 48 |
| 31 | -- | -- | 213 | -- | 48 | 206 | -- | 42 |
| 32 | 229 | 50 | -- | -- | -- | -- | -- | -- |
| 35 | 233 | 50 | 225 | 97 | 47 | 219 | 94 | 40 |
| 39 | 237 | 50 | 230 | 97 | 47 | 222 | 94 | 40 |
| 43 | 241 | 50 | 229 | 95 | 47 | 223 | 93 | 39 |
| 47 | 234 | 50 | 227 | 97 | 47 | 218 | 93 | 39 |
| 51 | 235 | 49 | 232 | 99 | 46 | 222 | 94 | 39 |
| 54 | 241 | 49 | 228 | 95 | 45 | 222 | 92 | 38 |
| 58 | 245 | 49 | 236 | 96 | 44 | 224 | 91 | 33 |
| 62 | 244 | 49 | 232 | 95 | 43 | 222 | 91 | 31 |
| 66 | -- | -- | 240 | -- | 40 | 224 | -- | 30 |
| 67 | 243 | 49 | -- | -- | -- | -- | -- | -- |
| 70 | 255 | 47 | 238 | 93 | 37 | 228 | 89 | 27 |
| 74 | 253 | 47 | 238 | 94 | 35 | 228 | 90 | 23 |
| 78 | 257 | 44 | 244 | 95 | 27 | 231 | 90 | 21 |
| 82 | 283 | 44 | 239 | 91 | 24 | 224 | 85 | 19 |
| 86 | 259 | 43 | 240 | 93 | 23 | 228 | 88 | 18 |
| 90 | 258 | 40 | 240 | 93 | 21 | 236 | 91 | 13 |
| 94 | 257 | 37 | 237 | 92 | 20 | 231 | 90 | 13 |
| 96 | 252 | 36 | 236 | 94 | 18 | 231 | 92 | 13 |
| 98 | 248 | 35 | 238 | 95 | 17 | 227 | 92 | 11 |
| 100 | 247 | 31 | 237 | 96 | 14 | 222 | 90 | 10 |
| 102 | 247 | 30 | -- | -- | -- | -- | -- | -- |
| 103 | -- | -- | 235 | -- | 13 | 227 | -- | 10 |





FIGURE 3. GROWTH CURVES FOR MARSHALL RATS ADMINISTERED TRICHLOROETHYLENE IN CORN OIL BY GAVAGE FOR TWO YEARS

TABLE 10. MEAN BODY WEIGHTS AND SURVIVAL OF OSBORNE-MENDEL RATS IN THE TWO-YEAR GAVAGE STUDIES OF TRICHLOROETHYLENE

| Weeks on Study | Vehicle Control | | 500 mg/kg | | | 1,000 mg/kg | | |
|----------------|-----------------|------------------|-----------------|--------------------------------|------------------|-----------------|--------------------------------|------------------|
| | Av. Wt. (grams) | No. of Survivors | Av. Wt. (grams) | Wt. (percent of veh. controls) | No. of Survivors | Av. Wt. (grams) | Wt. (percent of veh. controls) | No. of Survivors |
| MALE | | | | | | | | |
| 0 | 197 | 50 | 197 | 100 | 50 | 199 | 101 | 50 |
| 1 | 252 | 50 | 246 | 98 | 50 | 247 | 98 | 50 |
| 2 | 285 | 50 | 272 | 95 | 50 | 269 | 94 | 50 |
| 3 | 307 | 50 | 280 | 91 | 50 | 289 | 94 | 50 |
| 4 | 329 | 50 | 315 | 96 | 50 | 312 | 95 | 49 |
| 5 | 338 | 50 | 338 | 100 | 50 | 337 | 100 | 49 |
| 6 | 352 | 50 | 347 | 99 | 50 | 345 | 98 | 49 |
| 7 | 364 | 50 | 360 | 99 | 50 | 353 | 97 | 48 |
| 8 | 372 | 50 | 370 | 99 | 50 | 348 | 94 | 49 |
| 9 | 375 | 50 | 376 | 100 | 50 | 359 | 96 | 49 |
| 10 | 390 | 50 | 387 | 99 | 49 | 365 | 94 | 49 |
| 11 | 394 | 50 | 393 | 100 | 49 | 373 | 95 | 49 |
| 12 | 403 | 50 | 385 | 96 | 48 | 376 | 93 | 48 |
| 13 | 402 | 50 | 397 | 99 | 48 | 387 | 96 | 48 |
| 17 | 415 | 50 | 412 | 99 | 48 | 391 | 94 | 48 |
| 20 | 437 | 50 | 432 | 99 | 48 | 400 | 92 | 48 |
| 23 | 446 | 50 | 441 | 99 | 48 | 416 | 93 | 47 |
| 26 | 450 | 50 | 440 | 98 | 48 | 420 | 93 | 46 |
| 29 | 470 | 50 | 456 | 97 | 47 | 433 | 92 | 45 |
| 33 | 476 | 49 | 467 | 98 | 45 | 435 | 91 | 45 |
| 37 | 478 | 49 | 472 | 99 | 45 | 442 | 92 | 43 |
| 41 | 489 | 49 | 472 | 97 | 45 | 445 | 91 | 42 |
| 46 | 493 | 49 | 482 | 98 | 44 | 444 | 90 | 39 |
| 50 | 502 | 49 | 481 | 96 | 44 | 446 | 89 | 37 |
| 53 | 511 | 49 | | | | | | |
| 54 | | | 494 | | 44 | 468 | | 35 |
| 56 | 498 | 47 | | | | | | |
| 57 | | | 499 | | 44 | 466 | | 33 |
| 61 | 520 | 47 | 502 | 97 | 44 | 467 | 90 | 33 |
| 64 | 532 | 47 | 507 | 95 | 43 | 470 | 88 | 32 |
| 67 | 533 | 46 | 503 | 94 | 42 | 461 | 86 | 31 |
| 69 | 530 | 44 | | | | | | |
| 70 | | | 509 | | 40 | 469 | | 31 |
| 73 | 534 | 43 | 509 | 95 | 40 | 467 | 87 | 30 |
| 77 | 539 | 41 | 495 | 92 | 40 | 458 | 85 | 29 |
| 81 | 533 | 40 | 503 | 94 | 36 | 461 | 86 | 27 |
| 84 | 530 | 38 | | | | | | |
| 85 | | | 504 | | 32 | 458 | | 24 |
| 90 | 532 | 35 | 503 | 95 | 28 | 452 | 85 | 23 |
| 98 | 519 | 31 | 491 | 95 | 26 | 440 | 85 | 21 |
| 97 | 517 | 28 | 490 | 95 | 25 | 438 | 85 | 19 |
| 100 | 499 | 24 | 478 | 96 | 18 | 410 | 82 | 17 |
| 104 | 476 | 22 | 464 | 97 | 17 | 421 | 88 | 14 |
| FEMALE | | | | | | | | |
| 0 | 166 | 50 | 166 | 100 | 50 | 167 | 101 | 50 |
| 1 | 190 | 50 | 186 | 98 | 50 | 187 | 98 | 50 |
| 2 | 207 | 50 | 203 | 98 | 50 | 206 | 100 | 50 |
| 3 | 224 | 50 | 211 | 94 | 50 | 210 | 94 | 50 |
| 4 | 227 | 50 | 230 | 101 | 50 | 230 | 101 | 50 |
| 5 | 234 | 50 | 240 | 103 | 50 | 241 | 103 | 50 |
| 6 | 238 | 50 | 240 | 101 | 50 | 240 | 101 | 50 |
| 7 | 244 | 49 | 246 | 101 | 50 | 251 | 103 | 50 |
| 8 | 247 | 49 | 250 | 101 | 50 | 248 | 100 | 50 |
| 9 | 249 | 49 | 253 | 102 | 50 | 252 | 101 | 50 |
| 10 | 256 | 49 | 256 | 100 | 48 | 255 | 100 | 50 |
| 11 | 259 | 49 | 262 | 101 | 48 | 258 | 100 | 50 |
| 12 | 263 | 49 | 266 | 101 | 48 | 266 | 101 | 50 |
| 13 | 265 | 49 | 262 | 99 | 48 | 268 | 101 | 50 |
| 17 | 275 | 48 | 277 | 101 | 48 | 276 | 100 | 49 |
| 20 | 278 | 48 | 283 | 103 | 48 | 279 | 101 | 47 |
| 23 | 283 | 48 | 286 | 101 | 48 | 279 | 99 | 47 |
| 26 | 290 | 48 | 285 | 98 | 48 | 284 | 98 | 47 |
| 29 | 290 | 48 | 290 | 100 | 48 | 284 | 98 | 46 |
| 33 | 298 | 48 | 298 | 100 | 48 | 296 | 99 | 46 |
| 37 | 293 | 46 | 300 | 102 | 46 | 293 | 100 | 46 |
| 38 | 293 | 46 | | | | | | |
| 41 | 297 | 45 | 296 | 100 | 45 | 286 | 96 | 46 |
| 46 | 296 | 45 | 295 | 100 | 44 | 288 | 97 | 46 |
| 50 | 301 | 44 | 298 | 99 | 44 | 292 | 97 | 45 |
| 53 | 303 | 43 | | | | | | |
| 54 | | | 297 | | 44 | 289 | | 45 |
| 56 | 302 | 43 | | | | | | |
| 57 | | | 296 | | 44 | 297 | | 45 |
| 61 | 310 | 43 | 304 | 98 | 42 | 300 | 97 | 43 |
| 64 | 307 | 41 | 301 | 98 | 41 | 299 | 97 | 43 |
| 67 | 307 | 41 | 305 | 99 | 41 | 300 | 98 | 42 |
| 69 | 308 | 40 | | | | | | |
| 70 | | | 312 | | 41 | 306 | | 41 |
| 73 | 313 | 39 | 310 | 99 | 39 | 308 | 98 | 39 |
| 77 | 315 | 37 | 300 | 95 | 33 | 298 | 95 | 37 |
| 81 | 316 | 35 | 309 | 98 | 32 | 312 | 99 | 29 |
| 84 | 317 | 32 | | | | | | |
| 85 | | | 305 | | 31 | 313 | | 28 |
| 90 | 322 | 31 | 302 | 94 | 23 | 306 | 95 | 22 |
| 98 | 319 | 30 | 296 | 93 | 21 | 297 | 93 | 21 |
| 97 | 317 | 28 | 301 | 95 | 17 | 306 | 97 | 18 |
| 100 | 319 | 22 | 302 | 95 | 15 | 307 | 96 | 15 |
| 104 | 303 | 18 | 281 | 93 | 10 | 305 | 101 | 7 |





FIGURE 4. GROWTH CURVES FOR OSBORNE-MENDEL RATS ADMINISTERED TRICHLOROETHYLENE IN CORN OIL BY GAVAGE FOR TWO YEARS

TABLE 11. COMPARISON OF EFFECTS ON SURVIVAL AND FINAL BODY WEIGHTS IN ACI, AUGUST, MARSHALL, AND OSBORNE-MENDEL RATS IN THE TWO-YEAR GAVAGE STUDIES OF TRICHLOROETHYLENE (a)

| Group | Final Weight Relative to Vehicle Control (percent) | Survival (percent) | Final Weight Relative to Vehicle Control (percent) | Survival (percent) |
|-------------------|--|---------------------|--|--------------------|
| | | ACI (b) | August (c) | |
| MALE | | | | |
| Untreated control | 102.8 | 78 | 105.2 | 48 |
| Vehicle control | -- | 76 | -- | 42 |
| 500 mg/kg | 89.0 | (d) 38 | 93.5 | 26 |
| 1,000 mg/kg | 87.5 | (d) 22 | 87.7 | 32 |
| FEMALE | | | | |
| Untreated control | 111.6 | 74 | 96.6 | 52 |
| Vehicle control | -- | 70 | -- | 46 |
| 500 mg/kg | 91.7 | 40 | 94.5 | 52 |
| 1,000 mg/kg | 93.0 | (d) 38 | 92.4 | 50 |
| | | Marshall (e) | Osborne-Mendel (c) | |
| MALE | | | | |
| Untreated control | 99.4 | 64 | 109.2 | 42 |
| Vehicle control | -- | 52 | -- | 44 |
| 500 mg/kg | 93.3 | (d) 24 | 97.5 | 34 |
| 1,000 mg/kg | 96.8 | 12 | 88.4 | 30 |
| FEMALE | | | | |
| Untreated control | 103.2 | 62 | 107.9 | 38 |
| Vehicle control | -- | 60 | -- | 40 |
| 500 mg/kg | 96.0 | (d) 24 | 92.7 | 22 |
| 1,000 mg/kg | 89.9 | (d) 20 | 100.7 | (d) 14 |

(a) Trichloroethylene was mixed in corn oil and administered by gavage 5 days per week.

(b) Final body weight recorded week 103

(c) Final body weight recorded week 104

(d) Survival was significantly ($P < 0.05$) reduced relative to vehicle controls.

(e) Final body weight recorded week 100

TABLE 12. SURVIVAL OF RATS IN THE TWO-YEAR GAVAGE STUDIES OF TRICHLOROETHYLENE (a)

| | Untreated Control | Vehicle Control | 500 mg/kg | 1,000 mg/kg |
|---|-------------------|-----------------|-----------|-------------|
| MALE ACI RATS (b) | | | | |
| Nonaccidental deaths before termination (c) | 10 | 12 | 20 | 21 |
| Accidentally killed | 0 | 0 | 11 | 18 |
| Animals missing | 1 | 0 | 0 | 0 |
| Killed at termination | 36 | 37 | 19 | 11 |
| Died during termination period | 3 | 1 | 0 | 0 |
| Survival P values (d) | -- | 0.001 | 0.019 | 0.002 |
| FEMALE ACI RATS (b) | | | | |
| Nonaccidental deaths before termination (c) | 12 | 13 | 16 | 20 |
| Accidentally killed | 0 | 2 | 14 | 12 |
| Animals missing | 1 | 0 | 0 | 0 |
| Killed at termination | 36 | 33 | 20 | 17 |
| Died during termination period | 1 | 2 | 0 | (e) 2 |
| Survival P values (d) | -- | 0.021 | 0.202 | 0.034 |
| MALE AUGUST RATS (f) | | | | |
| Nonaccidental deaths before termination (c) | 25 | 23 | 25 | 22 |
| Accidentally killed | 1 | 6 | 12 | 11 |
| Animals missexed | 0 | 0 | 0 | 1 |
| Killed at termination | 24 | 21 | 13 | 15 |
| Died during termination period | 0 | 0 | 0 | 1 |
| Survival P values (d) | -- | 0.252 | 0.163 | 0.302 |
| FEMALE AUGUST RATS (f) | | | | |
| Nonaccidental deaths before termination (c) | 24 | 26 | 18 | 12 |
| Accidentally killed | 0 | 1 | 6 | 13 |
| Killed at termination | 26 | 23 | 26 | 24 |
| Died during termination period | 0 | 0 | 0 | 1 |
| Survival P values (d) | -- | 0.136 | 0.371 | 0.186 |
| MALE MARSHALL RATS (g) | | | | |
| Nonaccidental deaths before termination (c) | 16 | 22 | 26 | 19 |
| Accidentally killed | 1 | 2 | 12 | 25 |
| Animals missing | 1 | 0 | 0 | 0 |
| Killed at termination | 32 | 26 | 12 | 6 |
| Survival P values (d) | -- | 0.137 | 0.007 | 0.123 |
| FEMALE MARSHALL RATS (g) | | | | |
| Nonaccidental deaths before termination (c) | 18 | 17 | 24 | 22 |
| Accidentally killed | 1 | 3 | 14 | 18 |
| Killed at termination | 31 | 30 | 12 | 10 |
| Survival P values (d) | -- | 0.002 | 0.011 | 0.005 |
| MALE OSBORNE-MENDEL RATS (h) | | | | |
| Nonaccidental deaths before termination (c) | 29 | 27 | 27 | 28 |
| Accidentally killed | 0 | 1 | 6 | 7 |
| Killed at termination | 18 | 22 | 17 | 14 |
| Died during termination period | 3 | 0 | 0 | 1 |
| Survival P values (d) | -- | 0.173 | 0.657 | 0.195 |
| FEMALE OSBORNE-MENDEL RATS (h) | | | | |
| Nonaccidental deaths before termination (c) | 31 | 22 | 33 | 37 |
| Accidentally killed | 0 | 8 | 6 | 6 |
| Killed at termination | 19 | 18 | 10 | 7 |
| Died during termination period | 0 | 2 | 1 | 0 |
| Survival P values (d) | -- | 0.008 | 0.052 | 0.009 |

(a) Fifty animals initially in each study

(b) Terminal-kill period: weeks 104-106

(c) Includes animals killed in a moribund condition

(d) The result of the life table trend test is in the vehicle control column, and the results of the life table pairwise comparisons with the vehicle controls are in the dosed columns.

(e) Includes one accidentally killed animal; this animal is also included among those accidentally killed.

(f) Terminal-kill period: week 104

(g) Terminal-kill period: male--weeks 104-105; female--week 104

(h) Terminal-kill period: weeks 104-105





FIGURE 5. KAPLAN-MEIER SURVIVAL CURVES FOR ACI RATS ADMINISTERED TRICHLOROETHYLENE IN CORN OIL BY GAVAGE FOR TWO YEARS





FIGURE 6. UNADJUSTED SURVIVAL CURVES FOR ACI RATS ADMINISTERED TRICHLOROETHYLENE IN CORN OIL BY GAVAGE FOR TWO YEARS





FIGURE 7. KAPLAN-MEIER SURVIVAL CURVES FOR AUGUST RATS ADMINISTERED TRICHLOROETHYLENE IN CORN OIL BY GAVAGE FOR TWO YEARS





FIGURE 8. UNADJUSTED SURVIVAL CURVES FOR AUGUST RATS ADMINISTERED TRICHLOROETHYLENE IN CORN OIL BY GAVAGE FOR TWO YEARS





FIGURE 9. KAPLAN-MEIER SURVIVAL CURVES FOR MARSHALL RATS ADMINISTERED TRICHLOROETHYLENE IN CORN OIL BY GAVAGE FOR TWO YEARS





FIGURE 10. UNADJUSTED SURVIVAL CURVES FOR MARSHALL RATS ADMINISTERED TRICHLOROETHYLENE IN CORN OIL BY GAVAGE FOR TWO YEARS





FIGURE 11. KAPLAN-MEIER SURVIVAL CURVES FOR OSBORNE-MENDEL RATS ADMINISTERED TRICHLOROETHYLENE IN CORN OIL BY GAVAGE FOR TWO YEARS





FIGURE 12. UNADJUSTED SURVIVAL CURVES FOR OSBORNE-MENDEL RATS ADMINISTERED TRICHLOROETHYLENE IN CORN OIL BY GAVAGE FOR TWO YEARS

III. RESULTS

Pathology and Statistical Analyses of Results

This section describes the significant or noteworthy changes in the incidences of rats with neoplastic or nonneoplastic lesions of the kidney, testis, hematopoietic system, subcutaneous tissue, adrenal gland, pituitary gland, mammary gland, thyroid gland, and uterus.

Lesions in male rats are summarized in Appendixes A, C, E, and G. Lesions in female rats are summarized in Appendixes B, D, F, and H. Histopathologic findings on neoplasms are summarized in Tables A1, C1, E1, and G1 (male rats) and B1, D1, F1, and H1 (female rats). Tables A2, C2, E2, and G2 (male rats) and B2, D2, F2, and H2 (female rats) give the survival and tumor status for individual rats. Tables A3, C3, E3, and G3 (male rats) and B3, D3, F3, and H3 (female rats) contain the statistical analyses of those primary tumors that occurred with an incidence of at least 5% in the vehicle control or in one of the dosed groups. The statistical analyses used are discussed in Chapter II (Statistical Methods) and Tables A3, C3, E3, and G3 (male rats) and B3, D3, F3, and H3 (female rats) (footnotes). Findings on nonneoplastic lesions are summarized in Table A4, C4, E4, and G4 (male rats) and B4, D4, F4, and H4 (female rats). Statistical comparisons were made between vehicle control and dosed groups.

Nonneoplastic Lesions of the Kidney

Administration of trichloroethylene caused karyomegaly of renal tubular epithelial cells and toxic nephropathy in all four strains of rats (Tables 13 and 14). The inner cortex and outer stripe of the outer medulla were the primary areas affected. The epithelial cells of the pars recta (pars descendens) of the proximal tubule were enlarged and contained nuclei several times their normal

size (karyomegaly). The enlarged nuclei were often hyperchromatic and irregular to oblong in shape. In kidneys with more extensive changes, tubular epithelial cells located in more superficial areas of the cortex were also affected. Other lesions observed in severely affected kidneys were diagnosed as toxic nephropathy to distinguish them from the common spontaneous nephropathy of aging rats. Toxic nephropathy occurred only in dosed rats and consisted of dilated tubules lined by elongated and flattened epithelial cells. The degree of flattening was often severe and was roughly proportional to the degree of dilation. The lumens of the affected tubules were empty or contained wisps of eosinophilic material.

Tubular cell hyperplasia occurred at low incidences in dosed male and female Osborne-Mendel rats and less frequently in untreated or vehicle control rats. One animal in each of the high dose male and female August and female Marshall and low dose male Marshall groups also had renal tubular cell hyperplasia. Tubular cell hyperplasia generally consisted of one or two tubules with stratified epithelium that partially or completely filled the tubular lumens. The affected tubular epithelial cells were large with abundant eosinophilic or basophilic cytoplasm and vesicular nuclei containing prominent nucleoli. Cells in mitosis were variable in number or absent. This lesion was distinctly different from the background tubular regeneration that is a component of spontaneous chronic progressive rat nephropathy. The latter lesion consists of cortical tubules lined by a single layer of cuboidal cells with small amounts of basophilic cytoplasm, vesicular nuclei, and variable mitoses.

No additional nonneoplastic lesions were observed which could be attributed to trichloroethylene administration.

TABLE 13. INCIDENCES OF RENAL CYTOMEGLAY IN ACI, AUGUST, MARSHALL, AND OSBORNE-MENDEL RATS IN THE TWO-YEAR GAVAGE STUDIES OF TRICHLOROETHYLENE (a)

| Group | ACI | August | Marshall | Osborne-Mendel |
|-----------------|-------------|--------------|--------------|----------------|
| MALE | | | | |
| Vehicle control | 0/50 | 0/50 | 0/49 | 0/50 |
| 500 mg/kg | 40/49 (82%) | 46/50 (92%) | 48/50 (96%) | 48/50 (96%) |
| 1,000 mg/kg | 48/49 (98%) | 46/49 (94%) | 47/47 (100%) | 49/50 (98%) |
| FEMALE | | | | |
| Vehicle control | 0/48 | 0/49 | 0/50 | 0/50 |
| 500 mg/kg | 43/47 (91%) | 46/48 (96%) | 46/48 (96%) | 48/50 (96%) |
| 1,000 mg/kg | 42/43 (98%) | 50/50 (100%) | 43/44 (98%) | 49/49 (100%) |

(a) All incidences in dosed groups significantly greater ($P < 0.01$) than those in the vehicle controls

TABLE 14. INCIDENCES OF TOXIC NEPHROPATHY IN ACI, AUGUST, MARSHALL, AND OSBORNE-MENDEL RATS IN THE TWO-YEAR GAVAGE STUDIES OF TRICHLOROETHYLENE (a)

| Group | ACI | August | Marshall | Osborne-Mendel |
|-----------------|-------------|-------------|-------------|----------------|
| MALE | | | | |
| Vehicle control | 0/50 | 0/50 | 0/49 | 0/50 |
| 500 mg/kg | 18/49 (37%) | 10/50 (20%) | 18/50 (36%) | 39/50 (78%) |
| 1,000 mg/kg | 18/49 (37%) | 31/49 (63%) | 23/47 (49%) | 35/50 (70%) |
| FEMALE | | | | |
| Vehicle control | 0/48 | 0/49 | 0/50 | 0/50 |
| 500 mg/kg | 21/47 (45%) | 8/48 (17%) | 30/48 (63%) | 30/50 (60%) |
| 1,000 mg/kg | 19/43 (44%) | 29/50 (58%) | 30/44 (68%) | 39/49 (80%) |

(a) All incidences in dosed groups significantly greater ($P < 0.01$) than those in the vehicle controls

Neoplastic Lesions

Kidney: Neoplastic lesions of the kidney were characterized according to the following criteria. Tubular cell adenomas were discrete masses of epithelial cells with cytologic features similar to foci of tubular hyperplasia. The adenomas exhibited loss of tubular structure, although the tumor cells were arranged in irregular clusters incompletely separated by basement membranes. Tubular cell adenocarcinomas were larger than the adenomas, less discrete, and more heterogeneous in growth pattern. These

tumor cells generally showed more pleomorphism than did the tumor cells of the tubular cell adenomas.

Proliferative renal cortical lesions were diagnosed in all strains of rats. The incidences of these lesions are summarized in Table 15, and the statistical analyses for the incidences in male Osborne-Mendel rats are shown in Table 16. Dosing with trichloroethylene significantly increased the incidence of renal tubular cell adenomas in male Osborne-Mendel rats at the low, but not the high, dose level.

TABLE 15. INCIDENCES OF RENAL CORTICAL PROLIFERATIVE LESIONS IN ACI, AUGUST, MARSHALL, AND OSBORNE-MENDEL RATS IN THE TWO-YEAR GAVAGE STUDIES OF TRICHLOROETHYLENE

| Group | | Tubular Cell Hyperplasia | Tubular Cell Adenoma | Tubular Cell Adenocarcinoma |
|-----------------------|-------------------|-----------------------------|-------------------------|--------------------------------|
| MALE | | | | |
| ACI | Untreated control | 0/49 | 0/49 | 0/49 |
| | Vehicle control | 0/50 | 0/50 | 0/50 |
| | 500 mg/kg | 0/49 | 0/49 | 1/49 |
| | 1,000 mg/kg | 0/49 | 0/49 | 0/49 |
| August | Untreated control | 0/50 | 0/50 | 0/50 |
| | Vehicle control | 0/50 | 0/50 | 0/50 |
| | 500 mg/kg | 0/50 | 1/50 | 1/50 |
| | 1,000 mg/kg | 1/49 | 1/49 | 0/49 |
| Marshall | Untreated control | 0/49 | 2/49 | 0/49 |
| | Vehicle control | 0/49 | 0/49 | 0/49 |
| | 500 mg/kg | 1/50 | 1/50 | 0/50 |
| | 1,000 mg/kg | 0/47 | 0/47 | 1/47 |
| Osborne-Mendel | Untreated control | 0/50 | 0/50 | 0/50 |
| | Vehicle control | 0/50 | 0/50 | 0/50 |
| | 500 mg/kg | 5/50 | 6/50 | 0/50 |
| | 1,000 mg/kg | 3/50 | 1/50 | 1/50 |
| FEMALE | | | | |
| ACI | Untreated control | 0/49 | 0/49 | 0/49 |
| | Vehicle control | 0/48 | 0/48 | 0/48 |
| | 500 mg/kg | 0/47 | 2/47 | (a) 1/47 |
| | 1,000 mg/kg | 0/43 | 0/43 | 1/43 |
| August | Untreated control | 0/50 | 0/50 | 0/50 |
| | Vehicle control | 0/49 | 1/49 | 0/49 |
| | 500 mg/kg | 0/48 | 2/48 | 2/48 |
| | 1,000 mg/kg | 1/50 | 0/50 | 0/50 |
| Marshall | Untreated control | 1/49 | 1/49 | 0/49 |
| | Vehicle control | 1/50 | 1/50 | 0/50 |
| | 500 mg/kg | 0/48 | 1/48 | 1/48 |
| | 1,000 mg/kg | 1/44 | 0/44 | 1/44 |
| Osborne-Mendel | Untreated control | 0/50 | 1/50 | 0/50 |
| | Vehicle control | 1/50 | 0/50 | 0/50 |
| | 500 mg/kg | 1/50 | 0/50 | 0/50 |
| | 1,000 mg/kg | 3/49 | 1/49 | 0/49 |

(a) Adenocarcinoma, NOS

TABLE 16. ANALYSIS OF TUBULAR CELL RENAL LESIONS IN MALE OSBORNE-MENDEL RATS IN THE TWO-YEAR GAVAGE STUDY OF TRICHLOROETHYLENE (a)

| | Untreated Control | Vehicle Control | 500 mg/kg | 1,000 mg/kg |
|--------------------------------------|-------------------|-----------------|------------|-------------|
| Hyperplasia | | | | |
| Overall Rates | 0/50 (0%) | 0/50 (0%) | 5/50 (10%) | 3/50 (6%) |
| Adenoma | | | | |
| Overall Rates | 0/50 (0%) | 0/50 (0%) | 6/50 (12%) | 1/50 (2%) |
| Adjusted Rates | 0.0% | 0.0% | 32.2% | 6.7% |
| Terminal Rates | 0/21 (0%) | 0/22 (0%) | 5/17 (29%) | 1/15 (7%) |
| Life Table Test | | P=0.246 | P=0.007 | P=0.424 |
| Incidental Tumor Tests | | P=0.243 | P=0.007 | P=0.424 |
| Adenoma or Adenocarcinoma (b) | | | | |
| Overall Rates | 0/50 (0%) | 0/50 (0%) | 6/50 (12%) | 2/50 (4%) |
| Adjusted Rates | 0.0% | 0.0% | 32.2% | 10.9% |
| Terminal Rates | 0/21 (0%) | 0/22 (0%) | 5/17 (29%) | 1/15 (7%) |
| Life Table Tests | | P=0.125 | P=0.007 | P=0.158 |
| Incidental Tumor Tests | | P=0.122 | P=0.007 | P=0.158 |

(a) The statistical analyses used are discussed in Chapter II (Statistical Methods) and Appendix G, Table G3.

(b) No historical data are available.

Testis: Interstitial cell tumors occurred with a positive trend in male ACI rats; the incidence in the high dose group was not significantly greater than that in the vehicle controls by the incidental tumor test (Table 17). The overall incidences in the dosed groups were lower than that in the vehicle controls, but if incidences are based on those animals surviving until the appearance of the first tumor (week 75), an

increasing trend is evident (vehicle control, 36/43; low dose, 23/26; high dose, 17/17). In male Marshall rats, interstitial cell tumors and interstitial cell tumors or malignant interstitial cell tumors (combined) occurred with positive trends, and the incidences in the high dose group were significantly greater than those in the vehicle controls (Table 18).

TABLE 17. ANALYSIS OF TESTICULAR LESIONS IN MALE ACI RATS IN THE TWO-YEAR GAVAGE STUDY OF TRICHLOROETHYLENE

| | Untreated Control | Vehicle Control | 500 mg/kg | 1,000 mg/kg |
|--------------------------------------|-------------------|-----------------|-------------|--------------|
| Interstitial Cell Hyperplasia | | | | |
| Overall Rates | 4/47 (9%) | 5/49 (10%) | 5/49 (10%) | 3/49 (6%) |
| Interstitial Cell Tumor | | | | |
| Overall Rates | 38/47 (81%) | 36/49 (73%) | 23/49 (47%) | 17/49 (35%) |
| Adjusted Rates | 90.4% | 94.7% | 95.8% | 100.0% |
| Terminal Rates | 34/38 (89%) | 36/38 (95%) | 18/19 (95%) | 11/11 (100%) |
| Incidental Tumor Tests | | P=0.019 | P=0.223 | P=0.074 |

TABLE 18. ANALYSIS OF TESTICULAR LESIONS IN MALE MARSHALL RATS IN THE TWO-YEAR GAVAGE STUDY OF TRICHLOROETHYLENE

| | Untreated Control | Vehicle Control | 500 mg/kg | 1,000 mg/kg |
|--|-------------------|-----------------|-------------|-------------|
| Interstitial Cell Hyperplasia | | | | |
| Overall Rates | 2/46 (4%) | 1/46 (2%) | 6/48 (13%) | 5/48 (10%) |
| Interstitial Cell Tumor | | | | |
| Overall Rates | 16/46 (35%) | 17/46 (37%) | 21/48 (44%) | 31/48 (65%) |
| Adjusted Rates | 46.9% | 55.7% | 95.1% | 100.0% |
| Terminal Rates | 14/32 (44%) | 13/26 (50%) | 11/12 (92%) | 6/6 (100%) |
| Incidental Tumor Tests | | P<0.001 | P<0.001 | P<0.001 |
| Interstitial Cell Tumor, Malignant | | | | |
| Overall Rates | 0/46 (0%) | 0/46 (0%) | 0/48 (0%) | 1/48 (2%) |
| Interstitial Cell Tumor or Interstitial Cell Tumor, Malignant | | | | |
| Overall Rates | 16/46 (35%) | 17/46 (37%) | 21/48 (44%) | 32/48 (67%) |
| Adjusted Rates | 46.9% | 55.7% | 95.1% | 100.0% |
| Terminal Rates | 14/32 (44%) | 13/26 (50%) | 11/12 (92%) | 6/6 (100%) |
| Incidental Tumor Tests | | P<0.001 | P<0.001 | P<0.001 |

III. RESULTS

Hematopoietic System: Leukemia occurred in female August rats with a statistically significant positive trend, but the incidences in the dosed groups were not significantly greater than that in the vehicle controls in the pairwise comparisons (Table 19).

Subcutaneous Tissue: Sarcomas occurred in male August rats with a significant positive trend, but the incidences in the dosed groups were not significantly greater than that in the

vehicle controls in the pairwise comparisons (Table 20).

Adrenal Gland: Adrenal cortical adenomas occurred in female Osborne-Mendel rats with a significant positive trend by the life table test, and the incidence in the high dose group was significantly greater than that in the vehicle controls by the life table test (Table 21). The incidental tumor test, a more appropriate statistical test for this generally nonlethal neoplasm, revealed no statistically significant differences.

TABLE 19. ANALYSIS OF LEUKEMIA IN FEMALE AUGUST RATS IN THE TWO-YEAR GAVAGE STUDY OF TRICHLOROETHYLENE

| | Untreated Control | Vehicle Control | 500 mg/kg | 1,000 mg/kg |
|------------------------|-------------------|-----------------|-----------|-------------|
| Overall Rates | 0/50 (0%) | 1/50 (2%) | 0/50 (0%) | 5/50 (10%) |
| Adjusted Rates | 0.0% | 2.2% | 0.0% | 18.2% |
| Terminal Rates | 0/26 (0%) | 0/23 (0%) | 0/26 (0%) | 3/25 (12%) |
| Life Table Tests | | P=0.027 | P=0.523N | P=0.078 |
| Incidental Tumor Tests | | P=0.020 | P=0.469N | P=0.059 |

TABLE 20. ANALYSIS OF SUBCUTANEOUS TISSUE SARCOMAS IN MALE AUGUST RATS IN THE TWO-YEAR GAVAGE STUDY OF TRICHLOROETHYLENE

| | Untreated Control | Vehicle Control | 500 mg/kg | 1,000 mg/kg |
|------------------------|-------------------|-----------------|-----------|---------------|
| Overall Rates | 1/50 (2%) | 0/50 (0%) | 1/50 (2%) | (a) 3/49 (6%) |
| Adjusted Rates | 3.8% | 0.0% | 2.9% | 15.0% |
| Terminal Rates | 0/24 (0%) | 0/21 (0%) | 0/13 (0%) | 1/16 (6%) |
| Life Table Tests | | P=0.033 | P=0.440 | P=0.064 |
| Incidental Tumor Tests | | P=0.032 | P=0.519 | P=0.050 |

(a) Includes one sarcoma, unclear primary or metastatic

TABLE 21. ANALYSIS OF ADRENAL GLAND LESIONS IN FEMALE OSBORNE-MENDEL RATS IN THE TWO-YEAR GAVAGE STUDY OF TRICHLOROETHYLENE

| | Untreated Control | Vehicle Control | 500 mg/kg | 1,000 mg/kg |
|-------------------------|-------------------|-----------------|-------------|-------------|
| Hyperplasia | | | | |
| Overall Rates | 1/49 (2%) | 1/50 (2%) | 4/50 (8%) | 0/49 (0%) |
| Cortical Adenoma | | | | |
| Overall Rates | 13/49 (27%) | 16/50 (32%) | 13/50 (26%) | 19/49 (39%) |
| Adjusted Rates | 54.8% | 55.6% | 66.3% | 92.7% |
| Terminal Rates | 9/19 (47%) | 9/20 (45%) | 6/11 (55%) | 6/7 (86%) |
| Life Table Tests | | P=0.008 | P=0.365 | P=0.011 |
| Incidental Tumor Tests | | P=0.100 | P=0.484N | P=0.127 |

III. RESULTS

Negative Trends and Lower Incidences

Adrenal Gland: Pheochromocytomas and pheochromocytomas or malignant pheochromocytomas (combined) occurred with significant negative trends in all strains and sexes except female ACI, male August, and male Marshall rats (Tables 22 to 25).

Several statistically significant negative trends and lower incidences of neoplasms were detected in various strains (Appendixes A-H).

Pituitary Gland: Adenomas occurred with significant negative trends in male and female August rats and in female Marshall rats.

Mammary Gland: Fibromas or fibroadenomas (combined) occurred in female August rats with a significant negative trend by life table analysis. The incidence of fibroadenomas in low dose female Marshall rats was significantly lower than that in the vehicle controls by the incidental tumor test.

Thyroid Gland: C-Cell adenomas occurred with significant negative trends in male August and male and female Osborne-Mendel rats.

Uterus: Endometrial stromal polyps occurred with a significant negative trend in female August rats, and the incidence of endometrial stromal polyps in low dose Osborne-Mendel females was significantly lower than that in the vehicle controls by the incidental tumor test.

TABLE 22. ANALYSIS OF ADRENAL GLAND LESIONS IN MALE ACI RATS IN THE TWO-YEAR GAVAGE STUDY OF TRICHLOROETHYLENE

| | Untreated Control | Vehicle Control | 500 mg/kg | 1,000 mg/kg |
|-------------------------|-------------------|-----------------|-----------|-------------|
| Hyperplasia | | | | |
| Overall Rates | 0/48 (0%) | 1/48 (2%) | 0/44 (0%) | 1/45 (2%) |
| Pheochromocytoma | | | | |
| Overall Rates | 4/48 (8%) | 8/48 (17%) | 0/44 (0%) | 0/45 (0%) |
| Adjusted Rates | 10.3% | 21.1% | 0.0% | 0.0% |
| Terminal Rates | 4/39 (10%) | 8/38 (21%) | 0/18 (0%) | 0/11 (0%) |
| Life Table Tests | | P=0.017N | P=0.047N | P=0.117N |
| Incidental Tumor Tests | | P=0.017N | P=0.047N | P=0.117N |

TABLE 23. ANALYSIS OF ADRENAL GLAND LESIONS IN AUGUST RATS IN THE TWO-YEAR GAVAGE STUDIES OF TRICHLOROETHYLENE

| | Untreated Control | Vehicle Control | 500 mg/kg | 1,000 mg/kg |
|--|-------------------|-----------------|-------------|-------------|
| MALE | | | | |
| Hyperplasia | | | | |
| Overall Rates | 4/49 (8%) | 2/50 (4%) | 3/49 (6%) | 1/47 (2%) |
| Pheochromocytoma | | | | |
| Overall Rates | 16/49 (33%) | 10/50 (20%) | 5/49 (10%) | 2/47 (4%) |
| Adjusted Rates | 63.7% | 41.0% | 33.9% | 12.5% |
| Terminal Rates | 15/24 (63%) | 7/21 (33%) | 4/13 (31%) | 2/16 (13%) |
| Life Table Tests | | P=0.039N | P=0.455N | P=0.049N |
| Incidental Tumor Tests | | P=0.053N | P=0.460N | P=0.073N |
| Pheochromocytoma, Malignant | | | | |
| Overall Rates | 0/49 (0%) | 0/50 (0%) | 1/49 (2%) | 1/47 (2%) |
| Pheochromocytoma or Pheochromocytoma, Malignant | | | | |
| Overall Rates | 16/49 (33%) | 10/50 (20%) | 6/49 (12%) | 3/47 (6%) |
| Adjusted Rates | 63.7% | 41.0% | 41.3% | 18.8% |
| Terminal Rates | 15/24 (63%) | 7/21 (33%) | 5/13 (38%) | 3/16 (19%) |
| Life Table Tests | | P=0.090N | P=0.602N | P=0.104N |
| Incidental Tumor Tests | | P=0.119N | P=0.610N | P=0.150N |
| FEMALE | | | | |
| Hyperplasia | | | | |
| Overall Rates | 10/50 (20%) | 5/48 (10%) | 12/48 (25%) | 0/50 (0%) |
| Pheochromocytoma | | | | |
| Overall Rates | 6/50 (12%) | 9/48 (19%) | 2/48 (4%) | 0/50 (0%) |
| Adjusted Rates | 20.5% | 33.2% | 7.7% | 0.0% |
| Terminal Rates | 4/26 (15%) | 6/23 (26%) | 2/26 (8%) | 0/25 (0%) |
| Life Table Tests | | P<0.001N | P=0.019N | P=0.003N |
| Incidental Tumor Tests | | P=0.001N | P=0.034N | P=0.009N |

TABLE 24. ANALYSIS OF ADRENAL GLAND LESIONS IN MARSHALL RATS IN THE TWO-YEAR GAVAGE STUDIES OF TRICHLOROETHYLENE

| | Untreated Control | Vehicle Control | 500 mg/kg | 1,000 mg/kg |
|--|-------------------|-----------------|-------------|-------------|
| MALE | | | | |
| Hyperplasia | | | | |
| Overall Rates | 8/48 (17%) | 6/48 (13%) | 5/43 (12%) | 3/43 (7%) |
| Pheochromocytoma or Pheochromocytoma, Malignant | | | | |
| Overall Rates | 23/48 (48%) | 25/48 (52%) | 12/43 (28%) | 12/43 (28%) |
| Adjusted Rates | 65.4% | 75.1% | 84.8% | 90.9% |
| Terminal Rates | 19/31 (61%) | 18/26 (69%) | 9/11 (82%) | 5/6 (83%) |
| Life Table Tests | | P=0.035 | P=0.353 | P=0.037 |
| Incidental Tumor Tests | | P=0.239 | P=0.158 | P=0.288 |
| FEMALE | | | | |
| Hyperplasia | | | | |
| Overall Rates | 16/47 (34%) | 4/49 (8%) | 5/47 (11%) | 2/43 (5%) |
| Pheochromocytoma | | | | |
| Overall Rates | 30/47 (64%) | 39/49 (80%) | 15/47 (32%) | 9/43 (21%) |
| Adjusted Rates | 74.8% | 92.7% | 76.5% | 67.0% |
| Terminal Rates | 21/31 (68%) | 26/29 (90%) | 8/12 (67%) | 6/10 (60%) |
| Life Table Tests | | P=0.050N | P=0.263N | P=0.067N |
| Incidental Tumor Tests | | P<0.001N | P=0.002N | P<0.001N |
| Pheochromocytoma, Malignant | | | | |
| Overall Rates | 4/47 (9%) | 4/49 (8%) | 0/47 (0%) | 1/43 (2%) |
| Pheochromocytoma or Pheochromocytoma, Malignant | | | | |
| Overall Rates | 32/47 (68%) | 40/49 (82%) | 15/47 (32%) | 9/43 (21%) |
| Adjusted Rates | 79.8% | 95.1% | 76.5% | 67.0% |
| Terminal Rates | 23/31 (74%) | 27/29 (93%) | 8/12 (67%) | 6/10 (60%) |
| Life Table Tests | | P=0.036N | P=0.222N | P=0.050N |
| Incidental Tumor Tests | | P<0.001N | P<0.001N | P<0.001N |

TABLE 25. ANALYSIS OF ADRENAL GLAND LESIONS IN OSBORNE-MENDEL RATS IN THE TWO-YEAR GAVAGE STUDIES OF TRICHLOROETHYLENE

| | Untreated Control | Vehicle Control | 500 mg/kg | 1,000 mg/kg |
|--|-------------------|-----------------|------------|-------------|
| MALE | | | | |
| Hyperplasia | | | | |
| Overall Rates | 6/50 (12%) | 4/50 (8%) | 3/49 (6%) | 5/50 (10%) |
| Pheochromocytoma | | | | |
| Overall Rates | (a) 9/50 (18%) | (b) 13/50 (26%) | 6/49 (12%) | 3/50 (6%) |
| Adjusted Rates | 34.8% | 42.9% | 31.6% | 18.7% |
| Terminal Rates | 6/21 (29%) | 7/22 (32%) | 5/17 (29%) | 2/15 (13%) |
| Life Table Tests | | P=0.028N | P=0.154N | P=0.052N |
| Incidental Tumor Tests | | P=0.023N | P=0.128N | P=0.043N |
| FEMALE | | | | |
| Hyperplasia | | | | |
| Overall Rates | 1/49 (2%) | 1/50 (2%) | 2/50 (4%) | 2/49 (4%) |
| Pheochromocytoma | | | | |
| Overall Rates | 4/49 (8%) | 8/50 (16%) | 4/50 (8%) | 1/49 (2%) |
| Adjusted Rates | 16.1% | 30.4% | 18.8% | 6.7% |
| Terminal Rates | 2/19 (11%) | 4/20 (20%) | 1/11 (9%) | 0/7 (0%) |
| Life Table Tests | | P=0.080N | P=0.394N | P=0.118N |
| Incidental Tumor Tests | | P=0.009N | P=0.178N | P=0.020N |
| Pheochromocytoma, Malignant | | | | |
| Overall Rates | 0/49 (0%) | 1/50 (2%) | 2/50 (4%) | 0/49 (0%) |
| Pheochromocytoma or Pheochromocytoma, Malignant | | | | |
| Overall Rates | 4/49 (8%) | 9/50 (18%) | 6/50 (12%) | 1/49 (2%) |
| Adjusted Rates | 16.1% | 33.1% | 26.0% | 6.7% |
| Terminal Rates | 2/19 (11%) | 4/20 (20%) | 1/11 (9%) | 0/7 (0%) |
| Life Table Tests | | P=0.073N | P=0.564N | P=0.084N |
| Incidental Tumor Tests | | P=0.004N | P=0.252N | P=0.009N |

(a) One pheochromocytoma, malignant, was also observed.

(b) One pheochromocytoma, malignant, was also observed in an animal bearing a benign pheochromocytoma.

IV. DISCUSSION AND CONCLUSIONS

Thirteen-Week Studies

Two-Year Studies

Conclusions

IV. DISCUSSION AND CONCLUSIONS

The possibility of a strain difference in rat susceptibility to trichloroethylene prompted the National Cancer Institute to initiate a series of 2-year toxicology and carcinogenesis studies of trichloroethylene in five strains of rats. The results obtained with F344/N rats have been reported separately (NTP TR 243, in preparation), and the present report describes the results of studies in four additional strains (ACI, August, Marshall, and Osborne-Mendel). For comparative purposes, some data from the study in F344/N rats have been summarized in this discussion.

Thirteen-Week Studies

The results of the 13-week gavage studies of trichloroethylene in ACI, August, and Marshall rats were similar to those of previous 8- or 13-week studies in Osborne-Mendel or F344/N rats, respectively (NCI, 1976; NTP TR 243, in preparation). In males, trichloroethylene at the highest dose given, 2,000 mg/kg (1,834 mg/kg in Marshall males), consistently reduced final body weights by 12%-17%. Three of 10 male August rats died in the 2,000 mg/kg group. All male F344/N rats dosed with 2,000 mg/kg for 13 weeks survived (NTP TR 243, in preparation), and 5/5 Osborne-Mendel males dosed with 3,160 mg/kg per day, 5 days per week for 8 weeks, survived (NCI, 1976).

The highest dose studied in females for 13 weeks in the present studies (1,000 mg/kg in ACI and August; 918 mg/kg in Marshall) was not lethal but caused from 4% to 7% reductions in final body weights. In female F344/N rats, the 1,000 mg/kg dose produced a 3% decrease in final body weight (NTP TR 243, in preparation), and 3,160 mg/kg per day, 5 days per week for 8 weeks reduced final body weights of female Osborne-Mendel rats by 13% (NCI, 1976).

As in the earlier studies in rats, the chronic nephrotoxic effect of trichloroethylene observed in the 2-year studies was not predicted by the results of the short-term studies. In the studies in F344/N rats (NTP TR 243, in preparation), 13 weeks of dosing with trichloroethylene produced minimal to mild cytomegaly and karyomegaly of the tubular epithelial cells in the inner renal

cortex in both sexes. These changes were so subtle that they were diagnosed only during a reevaluation of tissues which was prompted by the detection of more pronounced renal effects during the 2-year studies.

Two-Year Studies

The audits of the experimental data from the present studies revealed insufficient documentation of animal breeding, clinical observations, environmental conditions, and analytical chemistry data (Appendix Q). In addition, the production of central nervous system toxicity (characterized by sedation, loss of consciousness, tremors, and convulsive seizures) and reduced survival indicate that the doses selected for these studies were too high. For these reasons, these studies were considered inadequate to evaluate the presence or absence of carcinogenic potential of trichloroethylene in ACI, August, Marshall, and Osborne-Mendel rats. The studies do, however, demonstrate a clear nephrotoxic effect of trichloroethylene in rats.

Survival and body weights: All dosed groups except high dose female Osborne-Mendel rats exhibited some degree of final body weight depression. This depression was 10%-12% in five of the dose groups (all high dose males except Marshall, low dose ACI males, and high dose Marshall females). Final mean body weight decrements in other groups ranged from 3% in low dose Osborne-Mendel males to 8% in low dose ACI females. In the earlier studies in F344/N rats, trichloroethylene at 1,000 mg/kg reduced mean body weight gain of males by 13% and at 500 and 1,000 mg/kg reduced body weight gains of females by 12% and 18%, respectively.

Trichloroethylene administered at both doses significantly reduced the survival of ACI males and Marshall females. Trichloroethylene at 1,000 mg/kg also reduced the survival of female Osborne-Mendel and ACI rats. The low dose, but not the high dose, reduced the survival of male Marshall rats. In the earlier gavage studies of trichloroethylene in F344/N rats, both the 500 and 1,000 mg/kg doses significantly reduced the survival of males, whereas neither dose reduced the survival of females. It is not clear

IV. DISCUSSION AND CONCLUSIONS

whether the excessive mortality observed in many dosed groups was caused by gavage-related trauma, the anesthetic properties of the chemical, nephrotoxicity, or a combination of these factors.

The absence of a dose-response relationship in survival of male Marshall rats may be related to the high accidental death rate in the high dose group. Fifty percent of the animals in this group were accidentally killed, and animals that are considered to have died from accidental causes are not included in the Kaplan-Meier probability of survival calculation after the time of death. The incidences of accidental deaths also were high in other dosed groups. A total of 24% of all dosed animals died of accidental causes. This rate contrasts with an overall accidental death rate of 5% in vehicle control animals. All groups of dosed animals were affected, and the overall combined incidences were dose related (male: vehicle control, 9/200; low dose, 41/200; high dose, 61/200; female: 14/200; 40/200; 49/200). High accidental death rates were observed in other NTP studies in which trichloroethylene or tetrachloroethylene was administered by gavage. As suggested earlier (NTP TR 243, in preparation), perhaps some effect of trichloroethylene and related chemicals predisposes repeatedly dosed animals to gavage accidents.

The survival of untreated and vehicle control Osborne-Mendel rats of each sex was poor relative to the other three strains and to historical survival of F344/N rats. However, survival of Osborne-Mendel control animals in the present study was similar to that of controls in the NCI studies (NCI, 1976) (male, 40/100, and female, 37/100, in the present study vs male, 26/100, and female, 51/100, in the earlier study).

Renal toxicity (nonproliferative changes): Earlier studies of trichloroethylene administered by gavage (NCI, 1976; NTP TR 243, in preparation) identified the kidney of Osborne-Mendel and F344/N rats and B6C3F₁ mice as a target organ for the production of nonneoplastic pathologic changes characterized as cytomegaly, karyomegaly, and toxic nephrosis of the tubular

epithelial cells in the inner renal cortex. The identical lesion was reported in male and female F344/N rats exposed to tetrachloroethylene by inhalation (NTP, 1986). Neither Henschler et al. (1980) nor Fukuda et al. (1983) reported the presence of nonproliferative renal lesions in mice, rats, or hamsters exposed to trichloroethylene by inhalation. The results of the present studies show that the kidneys of ACI, August, and Marshall rats are affected similarly. The incidences of these lesions in five strains of rats are summarized in Table 26.

The nonneoplastic renal changes diagnosed in these studies were clearly attributable to the administration of trichloroethylene, since none of the untreated or vehicle control animals had the lesions. In the case of renal cytomegaly, the incidence of the change ranged from 82% to 100% in the dosed groups. Cytomegalic changes were diagnosed in dosed rats that died after as few as 26 weeks of exposure to trichloroethylene. The severity of the change appeared to be directly proportional to the duration of dosing.

Toxic nephropathy, which was clearly distinguishable from the spontaneous nephropathy of aging rats, was observed at increased incidences in dosed rats of each sex and strain but was noted only infrequently in animals that died before week 52. This lesion also appeared to increase in severity with longer exposure.

Renal toxicity (proliferative changes): Trichloroethylene produced a statistically significant increase in the incidence of renal tubular cell adenomas or adenocarcinomas (combined) in low dose male Osborne-Mendel rats. In addition, tubular cell hyperplasia was present in 5/50 low dose and 3/50 high dose Osborne-Mendel male rats. No proliferative tubular cell lesions were diagnosed in either control group. Although not statistically significant, proliferative lesions also were diagnosed in dosed male and female ACI and August rats, whereas no such lesions were detected in any of the control groups. Proliferative lesions of the renal tubular cells were diagnosed in dosed Marshall male and female rats, but these changes were also present in untreated and vehicle control rats.

TABLE 26. INCIDENCES OF RENAL CORTICAL LESIONS IN ACI, AUGUST, MARSHALL, OSBORNE-MENDEL, AND F344/N RATS IN THE TWO-YEAR GAVAGE STUDIES OF TRICHLOROETHYLENE

| | Group | Toxic Nephropathy | Tubular Cell Hyperplasia | Tubular Cell Adenoma | Tubular Cell Adenocarcinoma |
|-----------------------|-------------------|-------------------|--------------------------|----------------------|-----------------------------|
| MALE | | | | | |
| ACI | Untreated control | 0/49 | 0/49 | 0/49 | 0/49 |
| | Vehicle control | 0/50 | 0/50 | 0/50 | 0/50 |
| | 500 mg/kg | 18/49 | 0/49 | 0/49 | 1/49 |
| | 1,000 mg/kg | 18/49 | 0/49 | 0/49 | 0/49 |
| August | Untreated control | 0/50 | 0/50 | 0/50 | 0/50 |
| | Vehicle control | 0/50 | 0/50 | 0/50 | 0/50 |
| | 500 mg/kg | 10/50 | 0/50 | 1/50 | 1/50 |
| | 1,000 mg/kg | 31/49 | 1/49 | 1/49 | 0/49 |
| Marshall | Untreated control | 0/49 | 0/49 | 2/49 | 0/49 |
| | Vehicle control | 0/49 | 0/49 | 0/49 | 0/49 |
| | 500 mg/kg | 18/50 | 1/50 | 1/50 | 0/50 |
| | 1,000 mg/kg | 23/47 | 0/47 | 0/47 | 1/47 |
| Osborne-Mendel | Untreated control | 0/50 | 0/50 | 0/50 | 0/50 |
| | Vehicle control | 0/50 | 0/50 | 0/50 | 0/50 |
| | 500 mg/kg | 39/50 | 5/50 | 6/50 | 0/50 |
| | 1,000 mg/kg | 35/50 | 3/50 | 1/50 | 1/50 |
| F344/N (a) | Untreated control | 0/49 | 0/49 | 0/49 | 0/49 |
| | Vehicle control | 0/48 | 0/48 | 0/48 | 0/48 |
| | 500 mg/kg | 48/49 | 0/49 | 2/49 | 0/49 |
| | 1,000 mg/kg | 48/49 | 1/49 | 0/49 | 3/49 |
| FEMALE | | | | | |
| ACI | Untreated control | 0/49 | 0/49 | 0/49 | 0/49 |
| | Vehicle control | 0/48 | 0/48 | 0/48 | 0/48 |
| | 500 mg/kg | 21/47 | 0/47 | 2/47 | (b) 1/47 |
| | 1,000 mg/kg | 19/43 | 0/43 | 0/43 | 1/43 |
| August | Untreated control | 0/50 | 0/50 | 0/50 | 0/50 |
| | Vehicle control | 0/49 | 0/49 | 1/49 | 0/49 |
| | 500 mg/kg | 8/48 | 0/48 | 2/48 | 2/48 |
| | 1,000 mg/kg | 29/50 | 1/50 | 0/50 | 0/50 |
| Marshall | Untreated control | 0/49 | 1/49 | 1/49 | 0/49 |
| | Vehicle control | 0/50 | 1/50 | 1/50 | 0/50 |
| | 500 mg/kg | 30/48 | 0/48 | 1/48 | 1/48 |
| | 1,000 mg/kg | 30/44 | 1/44 | 0/44 | 1/44 |
| Osborne-Mendel | Untreated control | 0/50 | 0/50 | 1/50 | 0/50 |
| | Vehicle control | 0/50 | 1/50 | 0/50 | 0/50 |
| | 500 mg/kg | 30/50 | 1/50 | 0/50 | 0/50 |
| | 1,000 mg/kg | 39/49 | 3/49 | 1/49 | 0/49 |
| F344/N (a) | Untreated control | 0/49 | 0/49 | 0/49 | 0/49 |
| | Vehicle control | 0/50 | 0/50 | 0/50 | 0/50 |
| | 500 mg/kg | 48/49 | 0/49 | 0/49 | 0/49 |
| | 1,000 mg/kg | 48/48 | 0/48 | 0/48 | 1/48 |

(a) Data for F344/N rats obtained from NTP Technical Report No. 243; toxic nephropathy was designated as cytomegaly in that report.

(b) Adenocarcinoma, NOS

IV. DISCUSSION AND CONCLUSIONS

Increased incidences of renal tubular cell adenomas or adenocarcinomas (combined) were found in five male F344/N rats dosed with trichloroethylene (NTP TR 243, in preparation), and a tubular cell adenocarcinoma was found in a low dose (time-weighted-average dose of 549 mg/kg of trichloroethylene) male Osborne-Mendel rat (NCI, 1976). The incidences of these lesions in the five strains of rats are shown in Table 26. Henschler et al. (1980) reported a total of four renal tumors (adenomas, cystadenomas, adenocarcinomas) in 60 male WIST rats exposed to trichloroethylene by inhalation at 100 or 500 ppm. No renal tumors were reported in male control rats.

Because the rats could not always be unequivocally assigned to a high or low dose group, tumor incidences in dosed groups of the same sex and strain were pooled and compared with incidences in the corresponding vehicle control groups. By this analysis, the incidence of renal tubular cell tumors in dosed male Osborne-Mendel rats was still elevated (tubular cell adenomas: vehicle control, 0/50; pooled dosed, 7/100; adenomas or adenocarcinomas (combined): vehicle control, 0/50; pooled dosed, 8/100). For all five rat strains studied, a total of 32 renal tubular cell neoplasms were observed in dosed animals, compared with 3 in all the vehicle control groups and 3 in all the untreated control groups.

Nonneoplastic kidney effects are common responses of F344 rats to the long-term administration of chlorinated ethanes and ethylenes. The NTP has also noted these effects in gavage studies of pentachloroethane (NTP, 1983) and trichloroethylene (NTP TR 243, in preparation) and in inhalation studies of tetrachloroethylene (NTP, 1986). Since these changes appear consistently in dosed rats but not in controls, they are considered to be due to trichloroethylene administration. This spectrum of kidney lesions appears to be similar to that described in male rats exposed to petroleum products (Mehlman et al., 1984). Unlike the petroleum-induced lesions, however, those produced by chlorinated ethanes and ethylenes appear in both male and female rats as well as in both sexes of mice (NTP TR 243, in preparation).

Testicular interstitial cell tumors: The incidences of testicular interstitial cell tumors were increased in dosed male ACI and Marshall rats. In ACI rats, these lesions showed a dose-related increase when based on animals surviving until the appearance of the first tumor. In Marshall rats, the incidence of interstitial cell tumors or malignant interstitial cell tumors (combined) in the high dose group was significantly increased and may be due to trichloroethylene. For ACI rats, the increased incidence at the high dose was significant only by life table analysis. This marginal effect was not considered to be chemically related.

Other tumors: Leukemia in female August rats and subcutaneous tissue sarcomas in male August rats occurred with positive trends, but the incidences in dosed groups were not significantly elevated relative to vehicle controls. In earlier studies in F344/N and Osborne-Mendel rats, the incidences of leukemia and subcutaneous neoplasms were not influenced by the administration of trichloroethylene.

Negative trends: Several negative trends in tumor incidences were observed in these studies. The majority of these findings were discounted either because there were nonsignificant differences in pairwise comparisons or because only the incidence in the low dose group was significantly reduced relative to the vehicle controls.

There were, however, more consistent negative trends in the incidences of adrenal gland pheochromocytomas in male ACI, female Marshall, and male and female August and Osborne-Mendel rats. We are unable to explain this effect. In the earlier study in F344/N rats, the incidence of pheochromocytomas appeared to be somewhat decreased in males (vehicle control, 4/45; low dose, 3/42; high dose, 1/44), but neither trend nor pairwise comparisons revealed statistically significant differences. No pheochromocytomas were diagnosed in any group of Osborne-Mendel rats dosed with trichloroethylene in an earlier study (NCI, 1976).

The results of these comparative studies of the toxicity of orally administered trichloroethylene

IV. DISCUSSION AND CONCLUSIONS

in four strains of rats show that there are few notable differences in the responsiveness of the strains to toxic doses of the chemical. Further, the responses of these four strains of rats were found to be similar to the response of F344/N rats.

The experimental and tabulated data for the NTP Technical Report on trichloroethylene were examined for accuracy, consistency, and compliance with Good Laboratory Practice requirements. As summarized in Appendix Q, the audits revealed problems with the conduct of the studies and with collection and documentation of the experimental data. Discrepancies were found that influenced the final interpretation of the results of these studies. The NTP also has reviewed a partial draft of an independent audit sponsored by the Halogenated Solvent Industry Alliance (HSIA) for which the findings were

similar to those of the NTP audit. The findings of all audits were taken into consideration during the interpretation of the results of these studies and the preparation of this report.

Conclusions: Under the conditions of these 2-year gavage studies of trichloroethylene in male and female ACI, August, Marshall, and Osborne-Mendel rats, trichloroethylene administration caused renal tubular cell cytomegaly and toxic nephropathy in both sexes of the four strains. However, these are considered to be *inadequate studies of carcinogenic activity** because of chemically induced toxicity, reduced survival, and deficiencies in the conduct of the studies. Despite these limitations, tubular cell neoplasms of the kidney were observed in rats exposed to trichloroethylene and interstitial cell neoplasms of the testis were observed in Marshall rats exposed to trichloroethylene.

*Explanation of Levels of Evidence of Carcinogenic Activity is on page 9. Summaries of the Peer Review comments and the public discussions on this Technical Report appear on pages 13-14 and 16-17.

V. REFERENCES

V. REFERENCES

1. American Conference of Governmental Industrial Hygienists (ACGIH) (1985-86) Threshold Limit Values and Biological Exposure Indices for 1985-86.
2. Armitage, P. (1971) *Statistical Methods in Medical Research*. New York: John Wiley & Sons, Inc., pp. 362-365.
3. Bachman, K.; Zimmerli, A.; Simons, E. (1950) Vapor-liquid equilibria at subatmospheric pressures. Trichloroethylene-tetrachloroethylene and ethyl phenyl acetate-ethyl α -phenyl butyrate. *Ind. Eng. Chem.* 42:2569-2575.
4. Baden, J.; Kelly, M.; Mazze, R.; Simmon, V. (1979) Mutagenicity of inhalation anaesthetics--trichloroethylene, divinyl ether, nitrous oxide and cyclopropane. *Br. J. Anaesth.* 51:417-421.
5. Bartsch, H.; Malavielle, C.; Barbin, A.; Planche, G. (1979) Mutagenic and alkylating metabolites of haloethylenes, chlorobutadienes and dichlorobutenes produced by rodent or human liver tissues: Evidence for oxirane formation by P450-linked microsomal mono-oxygenases. *Arch. Toxicol.* 41:249-277.
6. Beliles, R.; Brusick, D.; Mecler, F. (1980) Teratogenic-Mutagenic Risk of Workplace Contaminants: Trichloroethylene, Perchloroethylene and Carbon Disulfide. Contract No. 210-77-047. Litton Bionetics, Inc., Kensington, MD.
7. Berenblum, I., Ed. (1969) *Carcinogenicity Testing: A Report of the Panel on Carcinogenicity of the Cancer Research Commission of UICC, Vol. 2*. Geneva: International Union Against Cancer.
8. Boorman, G.; Montgomery, C., Jr.; Eustis, S.; Wolfe, M.; McConnell, E.; Hardisty, J. (1985) Quality assurance in pathology for rodent carcinogenicity studies. Milman, H.; Weisburger, E., Eds.: *Handbook of Carcinogen Testing*. Park Ridge, NJ: Noyes Publications, pp. 345-357.
9. Canter, D.; Zeiger, E.; Haworth, S.; Lawlor, T.; Mortelmans, K.; Speck, W. (1986) Comparative mutagenicity of aliphatic epoxides in *Salmonella*. *Mutat. Res.* 172:105-138.
10. Cerna, M.; Kypenova, H. (1977) Mutagenic activity of chloroethylenes analyzed by screening system tests. *Mutat. Res.* 46:214-215.
11. Clive, D.; Johnson, K.; Spector, J.; Batson, A.; Brown, M. (1979) Validation and characterization of the L5178Y/TK^{+/-} mouse lymphoma mutagen assay system. *Mutat. Res.* 59:61-108.
12. Cox, D. (1972) Regression models and life tables. *J.R. Stat. Soc.* B34:187-220.
13. Defalque, R. (1961) Pharmacology and toxicity of trichloroethylene. *Clin. Pharmacol. Therap.* 2:665-688.
14. Dorfmüller, M.; Henne, S.; York, R.; Bornschein, R.; Manson, J. (1979) Evaluation of teratogenicity and behavioral toxicity with inhalation exposure of maternal rats to trichloroethylene. *Toxicology* 14:153-166.
15. Elcombe, C.; Rose, M.; Pratt, I. (1985) Biochemical, histological, and ultrastructural changes in rat and mouse liver following the administration of trichloroethylene: Possible relevance to species differences in hepatocarcinogenicity. *Toxicol. Appl. Pharmacol.* 79:365-376.
16. Food Chemical News (1978) January 3, pp. 41-42.
17. Fukuda, K.; Takemoto, K.; Tsuruta, H. (1983) Inhalation carcinogenicity of trichloroethylene in mice and rats. *Ind. Health* 21:243-254.
18. Gallant, R. (1966) Physical properties of hydrocarbons. VI. Chlorinated ethylenes. *Hydrocarbon Process. Petrol. Refiner* 45:153-160.
19. Gart, J.; Chu, K.; Tarone, R. (1979) Statistical issues in interpretation of chronic bioassay tests for carcinogenicity. *J. Natl. Cancer Inst.* 62:957-974.
20. Goto, K.; Maeda, S.; Kano, Y.; Sugimura, T. (1978) Factors involved in differential Giemsa-staining of sister chromatids. *Chromosoma* 66:351-359.

V. REFERENCES

21. Greim, H.; Bonse, G.; Radwan, Z.; Reichert, D.; Henschler, D. (1975) Mutagenicity in vitro and potential carcinogenicity of chlorinated ethylenes as a function of metabolic oxirane formation. *Biochem. Pharmacol.* 24:2013-2017.
22. Gulati, D.; Sabharwal, P.; Shelby, M. (1985) Tests for the induction of chromosomal aberrations and sister chromatid exchanges in cultured Chinese hamster ovary (CHO) cells. Ashby, J.; de Serres, F.; Draper, M.; Ishidate, M., Jr.; Margolin, B.; Matter, B.; Shelby, M., Eds.: *Evaluation of Short-Term Tests for Carcinogens*. Amsterdam: Elsevier Science Publishers, Inc., pp. 413-426.
23. Haseman, J. (1984) Statistical issues in the design, analysis and interpretation of animal carcinogenicity studies. *Environ. Health Perspect.* 58:385-392.
24. Haseman J.; Huff, J.; Boorman, G. (1984) Use of historical control data in carcinogenicity studies in rodents. *Toxicol. Pathol.* 12:126-135.
25. Haseman, J.; Huff, J.; Rao, G.; Arnold, J.; Boorman, G.; McConnell, E. (1985) Neoplasms observed in untreated and corn oil gavage control groups of F344/N rats and (C57BL/6N × C3H/HeN)F₁ (B6C3F₁) mice. *J. Natl. Cancer Inst.* 75:975-984.
26. Haworth, S.; Lawlor, T.; Mortelmans, K.; Speck, W.; Zeiger, E. (1983) Salmonella mutagenicity test results for 250 chemicals. *Environ. Mutagen. (Suppl. 1)* 5:3-142.
27. Henschler, D.; Romen, W.; Elsasser, H.; Reichert, D.; Eder, E.; Radway, Z. (1980) Carcinogenicity study of trichloroethylene by long-term inhalation in three animal species. *Arch. Toxicol.* 43:237-248.
28. Henschler, D.; Elsasser, H.; Romen, W.; Eder, E. (1984) Carcinogenicity study of trichloroethylene, with and without epoxide stabilizers, in mice. *J. Cancer Res. Clin. Oncol.* 107:149-156.
29. Huff, J. (1971) New evidence on the old problems of trichloroethylene. *Ind. Med.* 40:25-33.
30. International Agency for Research on Cancer (IARC) (1979) *IARC Monographs on the Carcinogenic Risk of Chemicals to Humans. Trichloroethylene*. Lyon, France: World Health Organization 20:429-572.
31. International Agency for Research on Cancer (IARC) (1982) *Trichloroethylene. IARC Monographs on the Evaluation of the Carcinogenic Risk of Chemicals to Humans. Some Chemicals, Industrial Processes and Industries Associated with Cancer in Humans, IARC Monographs, Vols. 1 to 29. IARC Monographs Supplement 4*. Lyon, France: IARC, pp. 247-249.
32. Kaplan, E.; Meier, P. (1958) Nonparametric estimation of incomplete observations. *J. Am. Stat. Assoc.* 53:457-481.
33. Kirk-Othmer *Encyclopedia of Chemical Technology* (1963) Vol. 2, 2nd ed. New York: John Wiley & Sons Inc., p. 400.
34. Kirk-Othmer *Encyclopedia of Chemical Technology* (1979) Vol. 3, 3rd ed. New York: John Wiley & Sons Inc., pp. 745-753.
35. Lacher, J.; Hummel, L.; Bohmfalk, E.; Park, J. (1950) The near ultraviolet absorption spectra of some fluorinated derivatives of methane and ethylene. *J. Am. Chem. Soc.* 72:5486-5489.
36. Laskin, S.; Sellakumar, A.; Kuschner, M.; Nelson, N.; La Mendola, S.; Rusch, G.; Katz, G.; Dulak, N.; Albert, R. (1980) Inhalation carcinogenicity of epichlorohydrin in non-inbred Sprague-Dawley rats. *J. Natl. Cancer Inst.* 65:751-757.
37. Leong, B.; Schwetz, B.; Gehring, P. (1975) Embryo and fetotoxicity of inhaled trichloroethylene, perchloroethylene, methyl chloroform and methylene chloride in mice and rats. *Toxicol. Appl. Pharmacol.* 33:136 (abstract no. 34).
38. Linhart, M.; Cooper, J.; Martin, R.; Page, N.; Peters, J. (1974) *Carcinogenesis Bioassay Data System. Comput. Biomed. Res.* 7:230-248.

V. REFERENCES

39. Loprieno, N.; Barale, R.; Presciuttini, S.; Rossi, A.; Sbrana, I.; Stretti, G.; Zaccaro, L.; Abbondandolo, A.; Bonatti, S.; Fiorio, R. (1979) Comparative data with different test systems using micro-organisms and mammalian cells on references and environmental mutagens. *Mutat. Res.* 64:119.
40. Lyman, W. (1978) Report on trichloroethylene. Berkowitz, J., Ed.: *Literature Review--Problem Definition Studies on Selected Chemicals*. Cambridge, MA: Arthur D. Little, Inc., pp. 19-69.
41. Mantel, N.; Haenszel, W. (1959) Statistical aspects of the analysis of data from retrospective studies of disease. *J. Natl. Cancer Inst.* 22:719-748.
42. Maronpot, R.; Boorman, G. (1982) Interpretation of rodent hepatocellular proliferative alterations and hepatocellular tumors in chemical safety assessment. *Toxicol. Pathol.* 10:71-80.
43. McConnell, E.; Solleveld, H.; Swenberg, J.; Boorman, G. (1986) Guidelines for combining neoplasms for evaluation of rodent carcinogenesis. *J. Natl. Cancer Inst.* 76:283-289.
44. Mehlman, M.; Hemstreet, G.; Thorpe, J.; Weaver, N., Eds. (1984) Renal effects of petroleum hydrocarbons. *Advances in Modern Environmental Toxicology*, Vol. 7. Princeton, NJ: Princeton Scientific Publishers, Inc.
45. Mercier (1977) *Criteria (Exposure/Effect Relationships) for Organo-Solvents, V/F/177-4*. Luxembourg, Commission of the European Communities, pp. 123-190.
46. Miller, R.; Guengerich, F. (1982) Oxidation of TCE by liver microsomal cytochrome p-450: Evidence for chlorine migration in a transition state not involving trichloroethylene oxide. *Biochemistry* 21:1090-1097.
47. Mortelmans, K.; Haworth, S.; Lawlor, T.; Speck, W.; Tainer, B.; Zeiger, E. (1986) Salmonella mutagenicity tests: II. Results from the testing of 270 chemicals. *Environ. Mutagen.* 8(Suppl. 7):1-119.
48. National Cancer Institute (NCI) (1976) Bioassay of Trichloroethylene for Possible Carcinogenicity. NCI TR 2. U.S. Department of Health, Education, and Welfare, Bethesda, MD.
49. National Cancer Institute (NCI) (1977) Bioassay of Tetrachloroethylene for Possible Carcinogenicity. NCI TR 13. U.S. Department of Health, Education, and Welfare, Bethesda, MD.
50. National Cancer Institute (NCI) (1978a) Bioassay of Hexachloroethane for Possible Carcinogenicity. NCI TR 68. U.S. Department of Health, Education, and Welfare, Bethesda, MD.
51. National Cancer Institute (NCI) (1978b) Bioassay of 1,1,2,2-Tetrachloroethane for Possible Carcinogenicity. NCI TR 27. U.S. Department of Health, Education, and Welfare, Bethesda, MD.
52. National Cancer Institute (NCI) (1978c) Bioassay of 1,1,2-Trichloroethane for Possible Carcinogenicity. NCI TR 74. U.S. Department of Health, Education, and Welfare, Bethesda, MD.
53. National Toxicology Program (NTP) (1983) Carcinogenesis Bioassay of Pentachloroethane in F344/N Rats and B6C3F₁ Mice. NTP TR 232. U.S. Department of Health and Human Services, Bethesda, MD. 149 p.
54. National Toxicology Program (NTP) (1986) Toxicology and Carcinogenesis Studies of Tetrachloroethylene in F344/N Rats and B6C3F₁ Mice (Inhalation Studies). NTP TR 311. U.S. Department of Health and Human Services, Public Health Service, National Institutes of Health. 197 p.
55. National Toxicology Program (NTP) (in preparation) Toxicology and Carcinogenesis Studies of Trichloroethylene in F344/N Rats and B6C3F₁ Mice. NTP TR 243. U.S. Department of Health and Human Services, Public Health Service, National Institutes of Health.
56. Page, N. (1979) Assessment of trichloroethylene as an occupational carcinogen. Davis, W.; Rosenfeld, C., Eds.: *Carcinogenic Risks: Strategies for Intervention*. IARC Scientific Publications No. 25, INSERM Symposium Series, Vol. 74. Lyon, France: International Agency for Research on Cancer.

V. REFERENCES

57. Parchman, L.; Magee, P. (1982) Metabolism of 14 carbon trichloroethylene to $^{14}\text{CO}_2$ and interaction of a metabolite with liver DNA in rats and mice. *J. Toxicol. Environ. Health* 9:797-813.
58. Perry, P.; Wolff, S. (1974) New Giemsa method for the differential staining of sister chromatids. *Nature (London)* 251:156-158.
59. Prout, M.; Provan, W.; Green, T. (1985) Species differences in response to trichloroethylene. I. Pharmacokinetics in rats and mice. *Toxicol. Appl. Pharmacol.* 79:389-400.
60. Reddy, J.; Lalwani, N. (1983) Carcinogenesis by hepatic peroxisome proliferators: Evaluation of the risks of hypolipidemic drugs and industrial plasticizers to humans. *CRC Crit. Rev. Toxicol.* 13:1-58.
61. Sadtler Research Laboratories, Sadtler Standard Spectra. Philadelphia: Sadtler Research Laboratories; IR No. 185; NMR No. 9266.
62. Schwetz, B.; Leong, B.; Gehring, P. (1975) The effect of maternally inhaled trichloroethylene, perchloroethylene, methyl chloroform, and methylene chloride on embryonal and fetal development in mice and rats. *Toxicol. Appl. Pharmacol.* 32:84-96.
63. Simmon, V. (1977) Structural correlations of carcinogenic and mutagenic alkyl halides. Asher, I.; Zervos, C., Eds.: *Symposium on Structural Correlates of Carcinogenesis and Mutagenesis*. U.S. Food and Drug Administration, Office of Science, Rockville, MD, pp. 163-171.
64. Simmon, V.; Kauhanen, K.; Tardiff, R. (1977) Mutagenic activity of chemicals identified in drinking water. *Dev. Toxicol. Environ. Sci.* 2:249-258.
65. Slacik-Erben, R.; Roll, R.; Franke, G.; Vehleke, H. (1980) Trichloroethylene vapours do not produce dominant lethal mutations in male mice. *Arch. Toxicol.* 45:37-44
66. Stott, W.; Quast, J.; Watanabe, P. (1982) Pharmacokinetics and macromolecular interactions of trichloroethylene in mice and rats. *Toxicol. Appl. Pharmacol.* 62:137-151.
67. Tarone, R. (1975) Tests for trend in life table analysis. *Biometrika* 62:679-682.
68. U.S. Code of Federal Regulations (USCFR) (1976) 21CFR 8.305.
69. U.S. International Trade Commission (USITC) (1982) *Synthetic Organic Chemicals, United States Production and Sales 1981*. USITC Publication No. 1292. Washington, DC: Government Printing Office.
70. v. Apeldoorn, M. (1984) *Toxicological Evaluation of Trichloroethene*. Report 668310-001. National Institute of Public Health and Environmental Hygiene, The Netherlands.
71. Valle-Riestra, J. (1974) Food processing with chlorinated solvents. *Food Technol.* 28:25-32.
72. Van Duuren, B. (1975) On the possible mechanism of carcinogenic action of vinyl chloride. *Ann. N. Y. Acad. Sci.* 246:258-268.
73. Van Duuren, B.; Goldschmidt, B.; Katz, C.; Seidman, I.; Paul, J. (1974) Carcinogenic activity of alkylating agents. *J. Natl. Cancer Inst.* 53:695-700.
74. Waskell, L. (1978) A study of the mutagenicity of anesthetics and their metabolites. *Mutat. Res.* 57:141-153.
75. Waters, E.; Gerstner, H.; Huff, J. (1976) Trichloroethylene. I. An Impact Overview. II. An Abstracted Literature Collection. Oak Ridge National Laboratory ORNL/TIRC-76/2.
76. Waters, E.; Gerstner, H.; Huff, J. (1977) Trichloroethylene. I. An overview. *J. Toxic. Environ. Health* 2:671-707.
77. Weisburger, E. (1977) Carcinogenicity studies on halogenated hydrocarbons. *Environ. Health Perspect.* 21:7-16.
78. Wetterhahn, J. (1972) Eye make-up. Balsam, M.; Sagarin, E., Eds.: *Cosmetics Science and Technology*. New York: John Wiley & Sons Inc., p. 419.
79. World Health Organization (WHO) (1985) *Environmental Health Criteria Document on Trichloroethylene*.
80. Yoon, J.; Mason, J.; Valencia, R.; Woodruff, R.; Zimmering, S. (1985) Chemical mutagenesis testing in *Drosophila*. IV. Results of 45 coded compounds tested for the National Toxicology Program. *Environ. Mutagen.* 7:349-367.

APPENDIX A

SUMMARY OF LESIONS IN MALE ACI RATS IN THE TWO-YEAR GAVAGE STUDY OF TRICHLOROETHYLENE

| | PAGE | |
|----------|--|----|
| TABLE A1 | SUMMARY OF THE INCIDENCE OF NEOPLASMS IN MALE ACI RATS IN THE TWO-YEAR GAVAGE STUDY OF TRICHLOROETHYLENE | 81 |
| TABLE A2 | INDIVIDUAL ANIMAL TUMOR PATHOLOGY OF MALE ACI RATS IN THE TWO-YEAR GAVAGE STUDY OF TRICHLOROETHYLENE | 84 |
| TABLE A3 | ANALYSIS OF PRIMARY TUMORS IN MALE ACI RATS IN THE TWO-YEAR GAVAGE STUDY OF TRICHLOROETHYLENE | 92 |
| TABLE A4 | SUMMARY OF THE INCIDENCE OF NONNEOPLASTIC LESIONS IN MALE ACI RATS IN THE TWO-YEAR GAVAGE STUDY OF TRICHLOROETHYLENE | 94 |

TABLE A1. SUMMARY OF THE INCIDENCE OF NEOPLASMS IN MALE ACI RATS IN THE TWO-YEAR GAVAGE STUDY OF TRICHLOROETHYLENE

| | CONTROL (UNTR) | CONTROL (VEH) | LOW DOSE | HIGH DOSE |
|---|----------------|---------------|----------|-----------|
| ANIMALS INITIALLY IN STUDY | 50 | 50 | 50 | 50 |
| ANIMALS MISSING | 1 | | | |
| ANIMALS NECROPSIED | 49 | 50 | 49 | 50 |
| ANIMALS EXAMINED HISTOPATHOLOGICALLY | 49 | 50 | 49 | 50 |
| INTEGUMENTARY SYSTEM | | | | |
| *Skin | (49) | (50) | (49) | (50) |
| Squamous cell carcinoma | | 1 (2%) | | |
| Basal cell carcinoma | | 1 (2%) | | |
| RESPIRATORY SYSTEM | | | | |
| *Tracheal lumen | (49) | (50) | (49) | (50) |
| Fibrosarcoma, metastatic | | 1 (2%) | | |
| #Lung | (48) | (49) | (47) | (46) |
| Transitional cell carcinoma, metastatic | | | 1 (2%) | |
| Alveolar/bronchiolar adenoma | | | 2 (4%) | |
| Tubular cell adenoca, metastatic | | | 1 (2%) | |
| Fibrosarcoma, metastatic | | 1 (2%) | | |
| HEMATOPOIETIC SYSTEM | | | | |
| *Multiple organs | (49) | (50) | (49) | (50) |
| Leukemia, NOS | | 1 (2%) | | |
| Monocytic leukemia | 2 (4%) | 2 (4%) | | |
| #Thymus | (3) | (7) | (12) | (24) |
| Malignant lymphoma, NOS | | | 1 (8%) | |
| CIRCULATORY SYSTEM | | | | |
| *Chin | (49) | (50) | (49) | (50) |
| Hemangiosarcoma | 1 (2%) | | | |
| DIGESTIVE SYSTEM | | | | |
| #Liver | (49) | (50) | (49) | (49) |
| Hepatocellular carcinoma | | 1 (2%) | 1 (2%) | 1 (2%) |
| #Pancreas | (46) | (50) | (47) | (44) |
| Acinar cell adenoma | | | 1 (2%) | |
| URINARY SYSTEM | | | | |
| #Kidney | (49) | (50) | (49) | (49) |
| Tubular cell adenocarcinoma | | | 1 (2%) | |
| #Kidney/pelvis | (49) | (50) | (49) | (49) |
| Transitional cell papilloma | 1 (2%) | | | |
| Transitional cell carcinoma | 3 (6%) | | 1 (2%) | 1 (2%) |
| #Urinary bladder | (44) | (44) | (47) | (42) |
| Squamous cell papilloma | | | | 1 (2%) |
| Transitional cell papilloma | 1 (2%) | | 1 (2%) | |
| Transitional cell carcinoma | | 1 (2%) | 1 (2%) | |
| ENDOCRINE SYSTEM | | | | |
| #Pituitary | (48) | (39) | (33) | (29) |
| Adenoma, NOS | 8 (17%) | 6 (15%) | 4 (12%) | 3 (10%) |
| #Adrenal | (48) | (48) | (44) | (45) |
| Cortical adenoma | 8 (17%) | 6 (13%) | 1 (2%) | |
| Pheochromocytoma | 4 (8%) | 8 (17%) | | |

TABLE A1. SUMMARY OF THE INCIDENCE OF NEOPLASMS IN MALE ACI RATS IN THE TWO-YEAR GAVAGE STUDY OF TRICHLOROETHYLENE (Continued)

| | CONTROL (UNTR) | CONTROL (VEH) | LOW DOSE | HIGH DOSE |
|-------------------------------------|----------------|---------------|----------|-----------|
| ENDOCRINE SYSTEM (Continued) | | | | |
| #Thyroid | (48) | (46) | (40) | (39) |
| Follicular cell adenoma | | 1 (2%) | | |
| C-cell adenoma | 1 (2%) | | | |
| #Pancreatic islets | (46) | (50) | (47) | (44) |
| Islet cell adenoma | | 1 (2%) | | |
| REPRODUCTIVE SYSTEM | | | | |
| #Prostate | (48) | (49) | (45) | (43) |
| Carcinoma, NOS | | | 1 (2%) | |
| Adenoma, NOS | | 3 (6%) | | |
| Adenocarcinoma, NOS | | 1 (2%) | | |
| #Testis | (47) | (49) | (49) | (49) |
| Interstitial cell tumor | 38 (81%) | 36 (73%) | 23 (47%) | 17 (35%) |
| NERVOUS SYSTEM | | | | |
| #Brain/meninges | (49) | (50) | (47) | (47) |
| Granular cell tumor, NOS | | 1 (2%) | | |
| SPECIAL SENSE ORGANS | | | | |
| None | | | | |
| MUSCULOSKELETAL SYSTEM | | | | |
| None | | | | |
| BODY CAVITIES | | | | |
| *Thorax | (49) | (50) | (49) | (50) |
| Sarcoma, NOS | 1 (2%) | | | |
| *Peritoneum | (49) | (50) | (49) | (50) |
| Mesothelioma, NOS | 1 (2%) | | | |
| *Tunica vaginalis | (49) | (50) | (49) | (50) |
| Mesothelioma, NOS | 1 (2%) | | | 1 (2%) |
| ALL OTHER SYSTEMS | | | | |
| Neck | | | | |
| Neurilemoma, malignant | | 1 | | |
| Tail | | | | |
| Fibrosarcoma | | 1 | | |
| ANIMAL DISPOSITION SUMMARY | | | | |
| Animals initially in study | 50 | 50 | 50 | 50 |
| Natural death | 9 | 8 | 18 | 18 |
| Moribund sacrifice | 4 | 5 | 2 | 3 |
| Terminal sacrifice | 36 | 37 | 19 | 11 |
| Dosing accident | | | 1 | 1 |
| Accidentally killed, NOS | | | 10 | 17 |
| Animal missing | 1 | | | |

TABLE A1. SUMMARY OF THE INCIDENCE OF NEOPLASMS IN MALE ACI RATS IN THE TWO-YEAR GAVAGE STUDY OF TRICHLOROETHYLENE (Continued)

| | CONTROL (UNTR) | CONTROL (VEH) | LOW DOSE | HIGH DOSE |
|--|----------------|---------------|----------|-----------|
| TUMOR SUMMARY | | | | |
| Total animals with primary tumors** | 45 | 44 | 27 | 17 |
| Total primary tumors | 70 | 72 | 38 | 24 |
| Total animals with benign tumors | 42 | 39 | 24 | 17 |
| Total benign tumors | 61 | 61 | 32 | 21 |
| Total animals with malignant tumors | 7 | 9 | 6 | 2 |
| Total malignant tumors | 7 | 10 | 6 | 2 |
| Total animals with secondary tumors## | | 1 | 2 | |
| Total secondary tumors | | 2 | 2 | |
| Total animals with tumors uncertain-- benign or malignant | 2 | 1 | | 1 |
| Total uncertain tumors | 2 | 1 | | 1 |

* Number of animals receiving complete necropsy examination; all gross lesions including masses examined microscopically.

** Primary tumors: all tumors except secondary tumors

Number of animals examined microscopically at this site

Secondary tumors: metastatic tumors or tumors invasive into an adjacent organ

TABLE A2. INDIVIDUAL ANIMAL TUMOR PATHOLOGY OF MALE ACI RATS IN THE TWO-YEAR GAVAGE STUDY OF TRICHLOROETHYLENE: UNTREATED CONTROL

| ANIMAL NUMBER | 646 | 851 | 652 | 655 | 653 | 672 | 677 | 677 | 677 | 677 | 688 | 689 | 689 | 689 | 689 | 689 | 689 | 689 | 689 | 689 | 689 | 689 | 689 | 689 | 689 | 689 | 689 | 689 | 689 | 689 | 689 | 689 | 689 | 689 | 689 | 689 | 689 | 689 | 689 | 689 | |
|--------------------------------|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| WEEKS ON STUDY | 107 | 107 | 114 | 110 | 117 | 117 | 117 | 117 | 117 | 117 | 117 | 117 | 117 | 117 | 117 | 117 | 117 | 117 | 117 | 117 | 117 | 117 | 117 | 117 | 117 | 117 | 117 | 117 | 117 | 117 | 117 | 117 | 117 | 117 | 117 | 117 | 117 | 117 | 117 | 117 | 117 |
| RESPIRATORY SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Lungs and bronchi | + | + | + | + | + | - | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | | |
| Trachea | + | + | + | + | + | - | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| HEMATOPOIETIC SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Bone marrow | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| Spleen | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| Lymph nodes | + | + | - | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| Thymus | - | - | - | - | - | - | - | - | - | - | + | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | | |
| CIRCULATORY SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Heart | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| DIGESTIVE SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Salivary gland | + | + | + | - | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| Liver | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| Bile duct | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| Gallbladder & common bile duct | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | | |
| Pancreas | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| Esophagus | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| Stomach | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| Small intestine | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| Large intestine | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| URINARY SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Kidney | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| Kidney/pelvis | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| Transitional cell papilloma | | | | X | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Transitional cell carcinoma | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Urinary bladder | + | + | + | + | + | - | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| Transitional cell papilloma | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| ENDOCRINE SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Pituitary | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| Adenoma, NOS | X | | | | X | | | | | | | | X | | | | | | | | | | | | | | | | | | | | | | | | | | | X | |
| Adrenal | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| Cortical adenoma | X | | | X | | X | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Pheochromocytoma | | | | | | | | | | | | X | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Thyroid | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| C-cell adenoma | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Parathyroid | + | + | + | - | + | - | + | - | - | - | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| REPRODUCTIVE SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Mammary gland | + | N | + | + | + | + | + | + | + | + | N | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | | |
| Testis | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| Interstitial cell tumor | | X | | X | | X | X | X | X | X | X | X | X | X | X | X | X | X | X | X | X | X | X | X | X | X | X | X | X | X | X | X | X | X | X | X | X | X | X | | |
| Prostate | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| NERVOUS SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Brain | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | | |
| BODY CAVITIES | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Pleura | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | | |
| Sarcoma, NOS | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Peritoneum | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | | |
| Mesothelioma, NOS | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Tunica vaginalis | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| Mesothelioma, NOS | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| ALL OTHER SYSTEMS | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Multiple organs, NOS | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | | |
| Monocytic leukemia | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Chin | | | | | | | | | | | X | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Hemangiosarcoma | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | X | |

+: Tissue examined microscopically
 -: Required tissue not examined microscopically
 X: Tumor incidence
 N: Necropsy, no autolysis, no microscopic examination
 S: Animal missexed

: No tissue information submitted
 C: Necropsy, no histology due to protocol
 A: Autolysis
 M: Animal missing
 B: No necropsy performed

TABLE A2. INDIVIDUAL ANIMAL TUMOR PATHOLOGY OF MALE ACI RATS: UNTREATED CONTROL (Continued)

| ANIMAL NUMBER | 7 6 0 | 7 6 1 | 7 6 4 | 7 7 2 | 7 7 5 | 7 7 7 | 7 7 8 | 7 7 8 | 7 7 8 | 7 7 9 | 7 7 9 | 8 8 0 | 8 8 0 | 8 8 0 | 8 8 0 | 8 8 1 | 8 8 1 | 8 8 1 | 8 8 1 | 8 8 2 | 8 8 2 | 8 8 2 | 8 8 2 | 8 8 3 | 8 8 3 | 8 8 4 | 8 8 4 | 8 8 5 | 8 8 5 | 8 8 6 | 8 8 6 | 8 8 7 | TOTAL TISSUES TUMORS | | | |
|--------------------------------|---------|---------|---------|---------|---------|---------|---------|---------|---------|---------|---------|---------|---------|---------|---------|---------|---------|---------|---------|---------|---------|---------|---------|---------|---------|---------|---------|---------|---------|---------|---------|---------|----------------------|-----|-----|---|
| WEEKS ON STUDY | 1 0 7 | 1 0 7 | 1 1 1 | 1 1 1 | 1 1 1 | 1 1 1 | 1 1 1 | 1 1 1 | 1 1 1 | 1 1 1 | 1 1 1 | 1 1 1 | 1 1 1 | 1 1 1 | 1 1 1 | 1 1 1 | 1 1 1 | 1 1 1 | 1 1 1 | 1 1 1 | 1 1 1 | 1 1 1 | 1 1 1 | 1 1 1 | 1 1 1 | 1 1 1 | 1 1 1 | 1 1 1 | 1 1 1 | 1 1 1 | 1 1 1 | 1 1 1 | 1 1 1 | | | |
| RESPIRATORY SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Lungs and bronchi | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | 48 | |
| Trachea | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | 46 | |
| HEMATOPOIETIC SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Bone marrow | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | 47 | |
| Spleen | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | 49 | |
| Lymph nodes | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | 40 | |
| Thymus | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | 3 | |
| CIRCULATORY SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Heart | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | 49 | |
| DIGESTIVE SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Salivary gland | + | + | + | + | + | - | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | 46 | |
| Liver | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | 49 | |
| Bile duct | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | 49 | |
| Gallbladder & common bile duct | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | *49 | |
| Pancreas | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | 46 | |
| Esophagus | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | 49 | |
| Stomach | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | 49 | |
| Small intestine | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | 48 | |
| Large intestine | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | 48 | |
| URINARY SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Kidney | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | 49 | |
| Kidney/pelvis | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | 49 | |
| Transitional cell papilloma | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | 1 | |
| Transitional cell carcinoma | | | | | | | | | X | | | | | | | | | | | | | | | | | | | | | | | | | X | 3 | |
| Urinary bladder | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | 44 | |
| Transitional cell papilloma | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | 1 | |
| ENDOCRINE SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Pituitary | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | 48 | |
| Adenoma, NOS | | | X | | | | | | | | | X | X | | X | | | | | | | | | | | | | | | | | | | | 8 | |
| Adrenal | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | 48 | |
| Cortical adenoma | | | X | X | | | | | | X | | | | | | | | | | | | | | | | | | | | | | | | | 8 | |
| Pheochromocytoma | X | | | | | X | | | | X | | | | | | | | | | | | | | | | | | | | | | | | | 4 | |
| Thyroid | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | 48 | |
| C-cell adenoma | | | | | | | | | | | | X | | | | | | | | | | | | | | | | | | | | | | | 1 | |
| Parathyroid | + | + | - | - | + | + | + | + | + | + | + | - | + | + | - | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | 33 | |
| REPRODUCTIVE SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Mammary gland | + | + | N | + | + | + | + | + | + | N | + | + | N | N | + | + | N | + | + | + | N | + | + | N | + | + | N | N | N | | | | | *49 | | |
| Testis | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | 47 | |
| Interstitial cell tumor | X | X | X | X | X | X | X | X | X | X | X | X | X | X | X | X | X | X | X | X | X | X | X | X | X | X | X | X | X | X | X | X | X | X | 38 | |
| Prostate | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | 48 | |
| NERVOUS SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Brain | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | 49 | |
| BODY CAVITIES | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Pleura | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | *49 | |
| Sarcoma, NOS | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | 1 |
| Peritoneum | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | *49 | |
| Mesothelioma, NOS | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | 1 |
| Tunica vaginalis | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | *49 | |
| Mesothelioma, NOS | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | 1 |
| ALL OTHER SYSTEMS | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Multiple organs, NOS | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | *49 | |
| Monocytic leukemia | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | 2 |
| Chin | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Hemangiosarcoma | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | 1 |

* Animals necropsied

TABLE A2. INDIVIDUAL ANIMAL TUMOR PATHOLOGY OF MALE ACI RATS IN THE TWO-YEAR GAVAGE STUDY OF TRICHLOROETHYLENE: VEHICLE CONTROL

| ANIMAL NUMBER | 644 | 649 | 645 | 680 | 688 | 682 | 688 | 688 | 687 | 677 | 673 | 680 | 684 | 687 | 689 | 692 | 697 | 711 | 712 | 715 | 711 | 713 | 716 | 712 | 718 | 713 | 718 | 710 | | | |
|--------------------------------|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|---|---|
| WEEKS ON STUDY | 105 | 105 | 105 | 000 | 008 | 008 | 005 | 005 | 005 | 007 | 005 | 005 | 005 | 001 | 008 | 005 | 006 | 004 | 005 | 001 | 005 | 005 | 005 | 005 | 005 | 005 | 005 | 005 | 005 | | |
| INTEGUMENTARY SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Skin | N | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | | |
| Squamous cell carcinoma | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Basal cell carcinoma | | | | | | | X | | | | | | | | | | | | | | | | | | | | | | | | |
| RESPIRATORY SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Lungs and bronchi | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| Fibrosarcoma, metastatic | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Trachea | + | + | + | - | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| HEMATOPOIETIC SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Bone marrow | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| Spleen | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| Lymph nodes | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| Thymus | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | |
| CIRCULATORY SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Heart | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| DIGESTIVE SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Salivary gland | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| Liver | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| Hepatocellular carcinoma | | | | | | | X | | | | | | | | | | | | | | | | | | | | | | | | |
| Bile duct | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| Gallbladder & common bile duct | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | |
| Pancreas | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| Esophagus | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| Stomach | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| Small intestine | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| Large intestine | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| URINARY SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Kidney | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| Urinary bladder | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| Transitional cell carcinoma | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | X |
| ENDOCRINE SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Pituitary | + | + | + | - | - | + | + | + | - | + | + | + | + | + | + | + | + | - | - | - | - | - | + | + | + | + | + | + | + | + | |
| Adenoma, NOS | | | | X | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Adrenal | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| Cortical adenoma | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Pheochromocytoma | | | | | | | | X | | | X | | | | | | | | | | | | | | | | | | | | |
| Thyroid | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| Follicular cell adenoma | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Parathyroid | + | - | - | - | + | - | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| Pancreatic islets | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| Islet cell adenoma | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| REPRODUCTIVE SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Mammary gland | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | |
| Testis | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| Interstitial cell tumor | X | X | X | X | X | X | X | X | X | X | X | X | X | X | X | X | X | X | X | X | X | X | X | X | X | X | X | X | X | X | |
| Prostate | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| Adenoma, NOS | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Adenocarcinoma, NOS | X | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| NERVOUS SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Brain | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| Granular cell tumor, NOS | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| ALL OTHER SYSTEMS | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Multiple organs, NOS | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | |
| Leukemia, NOS | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Monocytic leukemia | | | | | X | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Neck, NOS | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Neurilemoma, malignant | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Tail | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Fibrosarcoma | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Tracheal lumen | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Fibrosarcoma, metastatic | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |

TABLE A2. INDIVIDUAL ANIMAL TUMOR PATHOLOGY OF MALE ACI RATS: VEHICLE CONTROL
(Continued)

| ANIMAL NUMBER | 7 3 7 | 7 4 3 | 7 4 6 | 7 4 8 | 7 5 9 | 7 6 5 | 7 6 8 | 7 6 9 | 7 7 1 | 7 7 9 | 7 8 4 | 7 8 8 | 7 9 4 | 7 9 5 | 7 9 6 | 8 0 0 | 8 0 3 | 8 0 5 | 8 0 6 | 8 0 9 | 8 2 6 | 8 3 0 | 8 3 1 | 8 3 3 | 8 3 4 | 8 4 2 | TOTAL TISSUES TUMORS |
|--------------------------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|----------------------|
| WEEKS ON STUDY | 1 0 5 | 1 0 5 | 1 0 5 | 1 0 6 | 1 0 5 | 1 0 5 | 1 0 4 | 1 0 5 | 1 0 6 | 1 0 8 | 1 0 9 | 1 0 6 | 1 0 6 | 1 0 5 | |
| INTEGUMENTARY SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Skin | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Squamous cell carcinoma | | | | | | | | | | | | | | | | | | | | | | | | | | *50 | |
| Basal cell carcinoma | | | | | | | | | | | | | | | | | | | | | | | | | | 1 | |
| RESPIRATORY SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Lungs and bronchi | | | | | | | | | | | | | | | | | | | | | | | | | | 49 | |
| Fibrosarcoma, metastatic | | | | | | | | | | | | | | | | | | | | | | | | | | 1 | |
| Trachea | | | | | | | | | | | | | | | | | | | | | | | | | | 48 | |
| HEMATOPOIETIC SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Bone marrow | | | | | | | | | | | | | | | | | | | | | | | | | | 50 | |
| Spleen | | | | | | | | | | | | | | | | | | | | | | | | | | 50 | |
| Lymph nodes | | | | | | | | | | | | | | | | | | | | | | | | | | 45 | |
| Thymus | | | | | | | | | | | | | | | | | | | | | | | | | | 7 | |
| CIRCULATORY SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Heart | | | | | | | | | | | | | | | | | | | | | | | | | | 49 | |
| DIGESTIVE SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Salivary gland | | | | | | | | | | | | | | | | | | | | | | | | | | 48 | |
| Liver | | | | | | | | | | | | | | | | | | | | | | | | | | 50 | |
| Hepatocellular carcinoma | | | | | | | | | | | | | | | | | | | | | | | | | | 1 | |
| Bile duct | | | | | | | | | | | | | | | | | | | | | | | | | | 50 | |
| Gallbladder & common bile duct | | | | | | | | | | | | | | | | | | | | | | | | | | *50 | |
| Pancreas | | | | | | | | | | | | | | | | | | | | | | | | | | 50 | |
| Esophagus | | | | | | | | | | | | | | | | | | | | | | | | | | 49 | |
| Stomach | | | | | | | | | | | | | | | | | | | | | | | | | | 49 | |
| Small intestine | | | | | | | | | | | | | | | | | | | | | | | | | | 49 | |
| Large intestine | | | | | | | | | | | | | | | | | | | | | | | | | | 48 | |
| URINARY SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Kidney | | | | | | | | | | | | | | | | | | | | | | | | | | 50 | |
| Urinary bladder | | | | | | | | | | | | | | | | | | | | | | | | | | 44 | |
| Transitional cell carcinoma | | | | | | | | | | | | | | | | | | | | | | | | | | 1 | |
| ENDOCRINE SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Pituitary | | | | | | | | | | | | | | | | | | | | | | | | | | 39 | |
| Adenoma, NOS | | | | | | | | | | | | | | | | | | | | | | | | | | 8 | |
| Adrenal | | | | | | | | | | | | | | | | | | | | | | | | | | 48 | |
| Cortical adenoma | | | | | | | | | | | | | | | | | | | | | | | | | | 6 | |
| Pheochromocytoma | | | | | | | | | | | | | | | | | | | | | | | | | | 8 | |
| Thyroid | | | | | | | | | | | | | | | | | | | | | | | | | | 48 | |
| Follicular cell adenoma | | | | | | | | | | | | | | | | | | | | | | | | | | 1 | |
| Parathyroid | | | | | | | | | | | | | | | | | | | | | | | | | | 30 | |
| Pancreatic islets | | | | | | | | | | | | | | | | | | | | | | | | | | 50 | |
| Islet cell adenoma | | | | | | | | | | | | | | | | | | | | | | | | | | 1 | |
| REPRODUCTIVE SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Mammary gland | | | | | | | | | | | | | | | | | | | | | | | | | | *50 | |
| Testis | | | | | | | | | | | | | | | | | | | | | | | | | | 49 | |
| Interstitial cell tumor | | | | | | | | | | | | | | | | | | | | | | | | | | 36 | |
| Prostate | | | | | | | | | | | | | | | | | | | | | | | | | | 49 | |
| Adenoma, NOS | | | | | | | | | | | | | | | | | | | | | | | | | | 3 | |
| Adenocarcinoma, NOS | | | | | | | | | | | | | | | | | | | | | | | | | | 1 | |
| NERVOUS SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Brain | | | | | | | | | | | | | | | | | | | | | | | | | | 50 | |
| Granular cell tumor, NOS | | | | | | | | | | | | | | | | | | | | | | | | | | 1 | |
| ALL OTHER SYSTEMS | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Multiple organs, NOS | | | | | | | | | | | | | | | | | | | | | | | | | | *50 | |
| Leukemia, NOS | | | | | | | | | | | | | | | | | | | | | | | | | | 1 | |
| Monocytic leukemia | | | | | | | | | | | | | | | | | | | | | | | | | | 2 | |
| Neck, NOS | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Neurilemoma, malignant | | | | | | | | | | | | | | | | | | | | | | | | | | 1 | |
| Tail | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Fibrosarcoma | | | | | | | | | | | | | | | | | | | | | | | | | | 1 | |
| Tracheal lumen | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Fibrosarcoma, metastatic | | | | | | | | | | | | | | | | | | | | | | | | | | 1 | |

* Animals necropsied

TABLE A2. INDIVIDUAL ANIMAL TUMOR PATHOLOGY OF MALE ACI RATS IN THE TWO-YEAR GAVAGE STUDY OF TRICHLOROETHYLENE: LOW DOSE

| ANIMAL NUMBER | 648 | 657 | 658 | 659 | 666 | 668 | 669 | 670 | 675 | 677 | 678 | 679 | 680 | 681 | 682 | 683 | 684 | 685 | 686 | 687 | 688 | 689 | 690 | 691 | 692 | 693 | 694 | 695 | 696 | 697 | 698 | 699 | | | | |
|---|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|---|---|---|---|
| WEEKS ON STUDY | 104 | 050 | 001 | 004 | 007 | 004 | 004 | 009 | 012 | 011 | 014 | 014 | 014 | 011 | 012 | 013 | 014 | 014 | 017 | 017 | 017 | 017 | 017 | 017 | 017 | 017 | 017 | 017 | 017 | 017 | 017 | 017 | | | | |
| RESPIRATORY SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Lungs and bronchi | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | | | |
| Transitional cell carcinoma, metastatic | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Alveolar/bronchiolar adenoma | X | | X | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Tubular cell adenocarcinoma, metastatic | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Trachea | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | | |
| HEMATOPOIETIC SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Bone marrow | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | | |
| Spleen | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | | |
| Lymph nodes | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| Thymus | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | | |
| Malignant lymphoma, NOS | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| CIRCULATORY SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Heart | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | | |
| DIGESTIVE SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Salivary gland | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| Liver | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| Hepatocellular carcinoma | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Bile duct | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + |
| Gallbladder & common bile duct | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | |
| Pancreas | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + |
| Acinar cell adenoma | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Esophagus | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + |
| Stomach | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + |
| Small intestine | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + |
| Large intestine | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + |
| URINARY SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Kidney | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| Tubular cell adenocarcinoma | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Kidney/pelvis | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + |
| Transitional cell carcinoma | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Urinary bladder | + | - | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + |
| Transitional cell papilloma | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Transitional cell carcinoma | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| ENDOCRINE SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Pituitary | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| Adenoma, NOS | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Adrenal | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + |
| Cortical adenoma | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Thyroid | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + |
| Parathyroid | + | - | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + |
| REPRODUCTIVE SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Mammary gland | + | N | N | N | N | N | + | N | N | + | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | |
| Testis | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + |
| Interstitial cell tumor | X | | | | | | X | X | | X | X | X | X | X | X | X | X | X | X | X | X | X | X | X | X | X | X | X | X | X | X | X | X | X | X | |
| Prostate | + | + | - | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + |
| Carcinoma, NOS | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| NERVOUS SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Brain | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |

**TABLE A2. INDIVIDUAL ANIMAL TUMOR PATHOLOGY OF MALE ACI RATS: LOW DOSE
(Continued)**

| ANIMAL NUMBER | 7 | 7 | 7 | 7 | 7 | 7 | 7 | 7 | 7 | 7 | 7 | 7 | 7 | 7 | 7 | 7 | 8 | 8 | 8 | 8 | 8 | 8 | 8 | 8 | 8 | 8 | 8 | |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|----------------------|
| | 0 | 1 | 0 | 0 | 1 | 0 | 1 | 0 | 0 | 1 | 0 | 0 | 0 | 0 | 0 | 1 | 0 | 0 | 0 | 0 | 0 | 1 | 0 | 0 | 0 | 0 | 0 | |
| WEEKS ON STUDY | 4 | 2 | 5 | 9 | 2 | 5 | 5 | 7 | 8 | 9 | 7 | 4 | 6 | 8 | 8 | 3 | 9 | 9 | 8 | 0 | 5 | 7 | 1 | 2 | 6 | 9 | 1 | |
| RESPIRATORY SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | | TOTAL: ISSUES TUMORS |
| Lungs and bronchi | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| Transitional cell carcinoma, metastatic | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Alveolar/bronchiolar adenoma | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Tubular cell adenocarcinoma, metastatic | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Trachea | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + |
| HEMATOPOIETIC SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | | TOTAL: ISSUES TUMORS |
| Bone marrow | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| Spleen | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + |
| Lymph nodes | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + |
| Thymus | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + |
| Malignant lymphoma, NOS | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| CIRCULATORY SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | | TOTAL: ISSUES TUMORS |
| Heart | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| DIGESTIVE SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | | TOTAL: ISSUES TUMORS |
| Salivary gland | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| Liver | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + |
| Hepatocellular carcinoma | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Bile duct | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + |
| Gallbladder & common bile duct | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N |
| Pancreas | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + |
| Acinar cell adenoma | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Esophagus | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + |
| Stomach | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + |
| Small intestine | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + |
| Large intestine | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + |
| URINARY SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | | TOTAL: ISSUES TUMORS |
| Kidney | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| Tubular cell adenocarcinoma | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Kidney/pelvis | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + |
| Transitional cell carcinoma | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Urinary bladder | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + |
| Transitional cell papilloma | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Transitional cell carcinoma | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| ENDOCRINE SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | | TOTAL: ISSUES TUMORS |
| Pituitary | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| Adenoma, NOS | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Adrenal | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + |
| Cortical adenoma | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Thyroid | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + |
| Parathyroid | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + |
| REPRODUCTIVE SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | | TOTAL: ISSUES TUMORS |
| Mammary gland | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| Testis | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + |
| Interstitial cell tumor | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Prostate | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + |
| Carcinoma, NOS | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| NERVOUS SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | | TOTAL: ISSUES TUMORS |
| Brain | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |

* Animals necropsied

TABLE A2. INDIVIDUAL ANIMAL TUMOR PATHOLOGY OF MALE ACI RATS IN THE TWO-YEAR GAVAGE STUDY OF TRICHLOROETHYLENE: HIGH DOSE

| ANIMAL NUMBER | 6 | 8 | 8 | 6 | 6 | 8 | 6 | 6 | 6 | 6 | 6 | 6 | 6 | 6 | 7 | 7 | 7 | 7 | 7 | 7 | 7 | 7 | 7 | 7 | 7 |
|--------------------------------|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| | 4 | 4 | 4 | 5 | 5 | 5 | 6 | 7 | 7 | 8 | 8 | 9 | 9 | 9 | 9 | 0 | 0 | 0 | 1 | 1 | 1 | 2 | 2 | 3 | 3 |
| WEEKS ON STUDY | 3 | 5 | 7 | 0 | 3 | 6 | 1 | 4 | 8 | 1 | 9 | 1 | 4 | 5 | 8 | 4 | 6 | 9 | 4 | 7 | 8 | 4 | 5 | 3 | 6 |
| RESPIRATORY SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | |
| Lungs and bronchi | + | - | - | + | + | + | + | + | + | + | + | + | + | + | + | + | + | - | A | + | + | + | + | + | + |
| Trachea | + | - | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | A | + | - | + | + | + | + |
| HEMATOPOIETIC SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | |
| Bone marrow | + | + | + | + | + | + | + | + | + | + | - | - | + | + | + | + | + | A | - | + | + | + | + | + | + |
| Spleen | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | A | + | + | + | + | + | + | + |
| Lymph nodes | + | - | - | - | + | - | + | - | - | + | + | + | + | - | + | + | + | A | + | - | + | + | + | + | - |
| Thymus | - | + | + | + | - | + | - | + | - | + | + | + | - | - | - | + | A | - | + | - | + | + | - | - | + |
| CIRCULATORY SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | |
| Heart | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | A | + | + | + | + | + | + | + |
| DIGESTIVE SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | |
| Salivary gland | + | + | + | + | - | + | + | + | - | + | + | + | + | - | + | + | + | A | + | - | + | + | + | + | - |
| Liver | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | A | + | + | + | + | + | + | + |
| Hepatocellular carcinoma | | | | | | | | | | | | | | | | | | | | | | | | | |
| Bile duct | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | A | + | + | + | + | + | + | + |
| Gallbladder & common bile duct | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N |
| Pancreas | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | A | + | + | + | + | + | + | + |
| Esophagus | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | A | + | + | + | + | + | + | + |
| Stomach | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | A | + | + | + | + | + | + | + |
| Small intestine | + | + | + | + | + | + | + | + | - | + | + | + | + | + | + | + | + | A | + | + | + | + | + | + | + |
| Large intestine | + | + | + | + | + | + | - | + | + | - | + | + | - | + | + | + | + | A | + | + | + | + | + | + | + |
| URINARY SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | |
| Kidney | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | A | + | + | + | + | + | + | + |
| Kidney/pelvis | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | A | + | + | + | + | + | + | + |
| Transitional cell carcinoma | | | | | | | | | | | | | | | | | | | | | | | | | |
| Urinary bladder | + | + | + | - | + | + | + | + | + | + | + | + | + | + | + | + | + | A | + | + | + | + | + | + | + |
| Squamous cell papilloma | | | | | | | | | | | | | | | | | | | | | | | | | |
| ENDOCRINE SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | |
| Pituitary | + | + | - | - | + | + | + | + | + | - | + | - | + | + | + | + | + | A | + | + | - | + | - | + | - |
| Adenoma, NOS | X | | | | | | | | | | | | | | | | | | | | | X | | | |
| Adrenal | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | A | - | - | + | + | + | + | |
| Thyroid | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | A | - | - | + | + | + | + | |
| Parathyroid | - | + | - | + | + | - | + | + | - | - | - | - | + | - | - | + | A | - | - | - | - | + | - | - | |
| REPRODUCTIVE SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | |
| Mammary gland | + | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N |
| Testis | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + |
| Interstitial cell tumor | X | | | | | X | X | X | | | | | | X | X | | | | | | | X | X | X | X |
| Prostate | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | A | + | + | + | + | + | + | + |
| NERVOUS SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | |
| Brain | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | A | + | + | + | + | + | + | + |
| BODY CAVITIES | | | | | | | | | | | | | | | | | | | | | | | | | |
| Tunica vaginalis | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + |
| Mesothelioma, NOS | | | | | | | | | | | | | | | | | | | | | | | | | |

TABLE A3. ANALYSIS OF PRIMARY TUMORS IN MALE ACI RATS IN THE TWO-YEAR GAVAGE STUDY OF TRICHLOROETHYLENE

| | Untreated Control | Vehicle Control | 500 mg/kg | 1,000 mg/kg |
|--|-------------------|-----------------|-------------|--------------|
| Hematopoietic System: Leukemia | | | | |
| Overall Rates (a) | 2/49 (4%) | 3/50 (6%) | 0/49 (0%) | 0/50 (0%) |
| Adjusted Rates (b) | 4.6% | 7.2% | 0.0% | 0.0% |
| Terminal Rates (c) | 0/39 (0%) | 1/38 (3%) | 0/19 (0%) | 0/11 (0%) |
| Life Table Tests (d) | | P=0.139N | P=0.239N | P=0.384N |
| Incidental Tumor Tests (d) | | P=0.093N | P=0.121N | P=0.423N |
| Cochran-Armitage Trend Test (d) | | P=0.038N | | |
| Fisher Exact Test | | | P=0.125N | P=0.121N |
| Pituitary: Adenoma | | | | |
| Overall Rates (a) | 8/48 (17%) | 6/39 (15%) | 4/33 (12%) | 3/29 (10%) |
| Adjusted Rates (b) | 19.9% | 16.9% | 19.5% | 28.7% |
| Terminal Rates (c) | 6/38 (16%) | 5/34 (15%) | 3/19 (16%) | 2/9 (22%) |
| Life Table Tests (d) | | P=0.257 | P=0.522 | P=0.315 |
| Incidental Tumor Tests (d) | | P=0.478 | P=0.547N | P=0.607 |
| Cochran-Armitage Trend Test (d) | | P=0.329N | | |
| Fisher Exact Test | | | P=0.480N | P=0.409N |
| Adrenal: Cortical Adenoma | | | | |
| Overall Rates (a) | 8/48 (17%) | 6/48 (13%) | 1/44 (2%) | 0/45 (0%) |
| Adjusted Rates (b) | 19.9% | 15.8% | 5.6% | 0.0% |
| Terminal Rates (c) | 7/39 (18%) | 6/38 (16%) | 1/18 (6%) | 0/11 (0%) |
| Life Table Tests (d) | | P=0.080N | P=0.260N | P=0.191N |
| Incidental Tumor Tests (d) | | P=0.080N | P=0.260N | P=0.191N |
| Cochran-Armitage Trend Test (d) | | P=0.006N | | |
| Fisher Exact Test | | | P=0.070N | P=0.016N |
| Adrenal: Pheochromocytoma | | | | |
| Overall Rates (a) | 4/48 (8%) | 8/48 (17%) | 0/44 (0%) | 0/45 (0%) |
| Adjusted Rates (b) | 10.3% | 21.1% | 0.0% | 0.0% |
| Terminal Rates (c) | 4/39 (10%) | 8/38 (21%) | 0/18 (0%) | 0/11 (0%) |
| Life Table Tests (d) | | P=0.017N | P=0.047N | P=0.117N |
| Incidental Tumor Tests (d) | | P=0.017N | P=0.047N | P=0.117N |
| Cochran-Armitage Trend Test (d) | | P=0.001N | | |
| Fisher Exact Test | | | P=0.004N | P=0.004N |
| Prostate: Adenoma | | | | |
| Overall Rates (a) | 0/48 (0%) | 3/49 (6%) | 0/45 (0%) | 0/43 (0%) |
| Adjusted Rates (b) | 0.0% | 7.9% | 0.0% | 0.0% |
| Terminal Rates (c) | 0/39 (0%) | 3/38 (8%) | 0/19 (0%) | 0/11 (0%) |
| Life Table Tests (d) | | P=0.153N | P=0.266N | P=0.403N |
| Incidental Tumor Tests (d) | | P=0.153N | P=0.266N | P=0.403N |
| Cochran-Armitage Trend Test (d) | | P=0.046N | | |
| Fisher Exact Test | | | P=0.137N | P=0.147N |
| Prostate: Adenoma, Adenocarcinoma, or Carcinoma | | | | |
| Overall Rates (a) | 0/48 (0%) | 4/49 (8%) | 1/45 (2%) | 0/43 (0%) |
| Adjusted Rates (b) | 0.0% | 10.5% | 5.3% | 0.0% |
| Terminal Rates (c) | 0/39 (0%) | 4/38 (11%) | 1/19 (5%) | 0/11 (0%) |
| Life Table Tests (d) | | P=0.176N | P=0.435N | P=0.311N |
| Incidental Tumor Tests (d) | | P=0.176N | P=0.435N | P=0.311N |
| Cochran-Armitage Trend Test (d) | | P=0.034N | | |
| Fisher Exact Test | | | P=0.208N | P=0.076N |
| Testis: Interstitial Cell Tumor | | | | |
| Overall Rates (a) | 38/47 (81%) | 36/49 (73%) | 23/49 (47%) | 17/49 (35%) |
| Adjusted Rates (b) | 90.4% | 94.7% | 95.8% | 100.0% |
| Terminal Rates (c) | 34/38 (89%) | 36/38 (95%) | 18/19 (95%) | 11/11 (100%) |
| Life Table Tests (d) | | P<0.001 | P=0.024 | P<0.001 |
| Incidental Tumor Tests (d) | | P=0.019 | P=0.223 | P=0.074 |
| Cochran-Armitage Trend Test (d) | | P<0.001N | | |
| Fisher Exact Test | | | P=0.006N | P<0.001N |

TABLE A3. ANALYSIS OF PRIMARY TUMORS IN MALE ACI RATS IN THE TWO-YEAR GAVAGE STUDY OF TRICHLOROETHYLENE (Continued)

- (a) Number of tumor-bearing animals/number of animals examined at the site
- (b) Kaplan-Meier estimated tumor incidence at the end of the study after adjusting for intercurrent mortality
- (c) Observed tumor incidence at terminal kill
- (d) Beneath the control incidence are the P values associated with the trend test. Beneath the dosed group incidence are the P values corresponding to pairwise comparisons between that dosed group and the vehicle controls. The life table analysis regards tumors in animals dying prior to terminal kill as being (directly or indirectly) the cause of death. The incidental tumor test regards these lesions as nonfatal. The Cochran-Armitage and Fisher exact tests compare directly the overall incidence rates. A negative trend or lower incidence in a dosed group is indicated by (N).

TABLE A4. SUMMARY OF THE INCIDENCE OF NONNEOPLASTIC LESIONS IN MALE ACI RATS IN THE TWO-YEAR GAVAGE STUDY OF TRICHLOROETHYLENE

| | CONTROL (UNTR) | CONTROL (VEH) | LOW DOSE | HIGH DOSE |
|--------------------------------------|----------------|---------------|----------|-----------|
| ANIMALS INITIALLY IN STUDY | 50 | 50 | 50 | 50 |
| ANIMALS MISSING | 1 | | | |
| ANIMALS NECROPSIED | 49 | 50 | 49 | 50 |
| ANIMALS EXAMINED HISTOPATHOLOGICALLY | 49 | 50 | 49 | 50 |
| INTEGUMENTARY SYSTEM | | | | |
| *Subcutaneous tissue | (49) | (50) | (49) | (50) |
| Inflammation, acute | | | 1 (2%) | 1 (2%) |
| Necrosis, NOS | | | 1 (2%) | |
| RESPIRATORY SYSTEM | | | | |
| *Nasal cavity | (49) | (50) | (49) | (50) |
| Inflammation, chronic | | | | 1 (2%) |
| Inflammation, chronic focal | | | | 1 (2%) |
| *Tracheal lumen | (49) | (50) | (49) | (50) |
| Hemorrhage | 3 (6%) | 1 (2%) | 1 (2%) | 5 (10%) |
| *Larynx | (49) | (50) | (49) | (50) |
| Inflammation, NOS | | | 1 (2%) | |
| Inflammation, necrotizing | | | | 1 (2%) |
| #Trachea | (46) | (46) | (43) | (45) |
| Hemorrhage | | | | 2 (4%) |
| Inflammation, necrotizing | | | 1 (2%) | |
| #Lung | (48) | (49) | (47) | (46) |
| Emphysema, alveolar | | | | 1 (2%) |
| Congestion, NOS | | 1 (2%) | | 3 (7%) |
| Edema, NOS | 1 (2%) | | | 2 (4%) |
| Hemorrhage | 1 (2%) | 4 (8%) | 4 (9%) | 14 (30%) |
| Inflammation, NOS | | 23 (47%) | 6 (13%) | 2 (4%) |
| Inflammation, focal | | 4 (8%) | 1 (2%) | 2 (4%) |
| Inflammation, acute focal | | | 1 (2%) | |
| Inflammation, chronic focal | | 1 (2%) | 3 (6%) | |
| Foreign material, NOS | | 13 (27%) | 5 (11%) | 3 (7%) |
| Pigmentation, NOS | | 1 (2%) | | |
| #Lung/alveoli | (48) | (49) | (47) | (46) |
| Hemorrhage | | | 1 (2%) | 1 (2%) |
| HEMATOPOIETIC SYSTEM | | | | |
| #Bone marrow | (47) | (50) | (45) | (45) |
| Hyperplasia, NOS | 1 (2%) | | | |
| Hyperplasia, granulocytic | | 1 (2%) | 1 (2%) | |
| #Spleen | (49) | (50) | (48) | (44) |
| Congestion, NOS | | | | 1 (2%) |
| Hemorrhage | 1 (2%) | | 1 (2%) | |
| Hemosiderosis | | | 1 (2%) | |
| Hyperplasia, lymphoid | 1 (2%) | | | |
| Hematopoiesis | | | 1 (2%) | |
| #Mediastinal lymph node | (40) | (45) | (35) | (30) |
| Hemorrhage | | | 1 (3%) | |
| #Mesenteric lymph node | (40) | (45) | (35) | (30) |
| Hemorrhage | | 1 (2%) | | 2 (7%) |
| #Thymus | (3) | (7) | (12) | (24) |
| Multiple cysts | | | | 2 (8%) |
| Hemorrhage | | | | 3 (13%) |
| CIRCULATORY SYSTEM | | | | |
| #Trachea | (46) | (46) | (43) | (45) |
| Embolus, septic | | 1 (2%) | | |

TABLE A4. SUMMARY OF THE INCIDENCE OF NONNEOPLASTIC LESIONS IN MALE ACI RATS IN THE TWO-YEAR GAVAGE STUDY OF TRICHLOROETHYLENE (Continued)

| | CONTROL (UNTR) | CONTROL (VEH) | LOW DOSE | HIGH DOSE |
|---------------------------------------|----------------|---------------|----------|-----------|
| CIRCULATORY SYSTEM (Continued) | | | | |
| #Lung | (48) | (49) | (47) | (46) |
| Thrombosis, NOS | | | | 1 (2%) |
| #Heart | (49) | (49) | (48) | (48) |
| Inflammation, focal | 1 (2%) | 1 (2%) | 1 (2%) | |
| Inflammation, acute/chronic | | 1 (2%) | | |
| Inflammation, chronic | | 6 (12%) | | |
| Inflammation, chronic focal | 2 (4%) | 2 (4%) | | |
| Fibrosis, focal | | | 1 (2%) | |
| Necrosis, focal | 2 (4%) | | | |
| Calcification, focal | 1 (2%) | | | |
| *Aorta | (49) | (50) | (49) | (50) |
| Mineralization | | 1 (2%) | | |
| Calcification, NOS | | | | 1 (2%) |
| Calcification, metastatic | 1 (2%) | | | |
| *Pulmonary artery | (49) | (50) | (49) | (50) |
| Calcification, metastatic | 1 (2%) | | | |
| DIGESTIVE SYSTEM | | | | |
| #Liver | (49) | (50) | (49) | (49) |
| Hemorrhage | | 1 (2%) | 1 (2%) | 1 (2%) |
| Inflammation, acute focal | 1 (2%) | | | |
| Inflammation, chronic focal | 1 (2%) | | | |
| Necrosis, focal | | 1 (2%) | 2 (4%) | |
| Metamorphosis, fatty | 1 (2%) | 2 (4%) | | |
| Cytoplasmic vacuolization | | 1 (2%) | | |
| Basophilic cyto change | 1 (2%) | 2 (4%) | | 1 (2%) |
| Clear cell change | 2 (4%) | 1 (2%) | 1 (2%) | 1 (2%) |
| Cytologic alteration, NOS | | 1 (2%) | | |
| Hepatocytomegaly | 1 (2%) | | | |
| Angiectasis | | | 1 (2%) | |
| #Liver/centrilobular | (49) | (50) | (49) | (49) |
| Necrosis, NOS | 1 (2%) | | | |
| #Bile duct | (49) | (50) | (49) | (49) |
| Dilatation, NOS | | 1 (2%) | | |
| Inflammation, chronic | | 1 (2%) | | |
| Hyperplasia, NOS | 1 (2%) | | | |
| #Pancreas | (46) | (50) | (47) | (44) |
| Hemorrhage | | | 1 (2%) | |
| Inflammation, chronic | | | | 1 (2%) |
| Inflammation, granulomatous | 1 (2%) | | | |
| Atrophy, focal | | | | 1 (2%) |
| Atrophy, granular | 1 (2%) | | | |
| #Pancreatic duct | (46) | (50) | (47) | (44) |
| Degeneration, cystic | 1 (2%) | | | |
| #Pancreatic acinus | (46) | (50) | (47) | (44) |
| Atrophy, NOS | | 1 (2%) | | |
| Atrophy, focal | 2 (4%) | 1 (2%) | | |
| Atrophy, granular | | 1 (2%) | | |
| *Esophageal lumen | (49) | (50) | (49) | (50) |
| Hemorrhage | 7 (14%) | 1 (2%) | 6 (12%) | 3 (6%) |
| #Esophagus | (49) | (49) | (47) | (47) |
| Dilatation, NOS | 1 (2%) | | | |
| Hemorrhage | 1 (2%) | | | |
| Inflammation, necrotizing | | | | 1 (2%) |
| Abscess, chronic | | | | 1 (2%) |
| #Stomach | (49) | (49) | (48) | (46) |
| Calcification, metastatic | 1 (2%) | | | |
| Hyperplasia, epithelial | | | | 2 (4%) |
| #Gastric mucosa | (49) | (49) | (48) | (46) |
| Hyperplasia, NOS | | 1 (2%) | | |

TABLE A4. SUMMARY OF THE INCIDENCE OF NONNEOPLASTIC LESIONS IN MALE ACI RATS IN THE TWO-YEAR GAVAGE STUDY OF TRICHLOROETHYLENE (Continued)

| | CONTROL (UNTR) | CONTROL (VEH) | LOW DOSE | HIGH DOSE |
|-------------------------------------|----------------|---------------|----------|-----------|
| DIGESTIVE SYSTEM (Continued) | | | | |
| #Forestomach | (49) | (49) | (48) | (46) |
| Hyperplasia, epithelial | 1 (2%) | | | |
| Hyperkeratosis | 1 (2%) | | | |
| #Duodenum | (48) | (49) | (48) | (46) |
| Hemorrhage | | | | 1 (2%) |
| #Colon | (48) | (48) | (46) | (43) |
| Parasitism | | | 1 (2%) | 1 (2%) |
| URINARY SYSTEM | | | | |
| #Kidney | (49) | (50) | (49) | (49) |
| Mineralization | 39 (80%) | 28 (56%) | 14 (29%) | 9 (18%) |
| Cast, NOS | | 1 (2%) | | |
| Hydronephrosis | 5 (10%) | 4 (8%) | 5 (10%) | 5 (10%) |
| Hemorrhage | | | 3 (6%) | |
| Inflammation, NOS | | | | 1 (2%) |
| Pyelonephritis, acute | | | 1 (2%) | 2 (4%) |
| Inflammation, acute | 1 (2%) | | | |
| Pyelonephritis, acute/chronic | | 1 (2%) | 1 (2%) | |
| Inflammation, chronic | | 1 (2%) | | |
| Pyelonephritis, chronic | 1 (2%) | | | |
| Nephropathy | 48 (98%) | 49 (98%) | 40 (82%) | 43 (88%) |
| Nephropathy, toxic | | | 18 (37%) | 18 (37%) |
| Necrosis, NOS | | 1 (2%) | | |
| Necrosis, focal | | 1 (2%) | | |
| Infarct, NOS | | | | 1 (2%) |
| Calcification, focal | 2 (4%) | 5 (10%) | 7 (14%) | 11 (22%) |
| Cytomegaly | | | 40 (82%) | 48 (98%) |
| Hyperplasia, epithelial | 2 (4%) | | 1 (2%) | |
| Angiectasis | 6 (12%) | 1 (2%) | | 1 (2%) |
| #Kidney/cortex | (49) | (50) | (49) | (49) |
| Cyst, NOS | | | | 1 (2%) |
| #Kidney/tubule | (49) | (50) | (49) | (49) |
| Necrosis, NOS | | | 1 (2%) | |
| #Kidney/pelvis | (49) | (50) | (49) | (49) |
| Dilatation, NOS | 1 (2%) | | | |
| Hemorrhage | 1 (2%) | | | |
| Inflammation, NOS | | | | 1 (2%) |
| Inflammation, focal | | | | 1 (2%) |
| Inflammation, suppurative | | | | 1 (2%) |
| Inflammation, acute | 1 (2%) | | 1 (2%) | 2 (4%) |
| Inflammation, chronic | 1 (2%) | | | |
| Hyperplasia, epithelial | 19 (39%) | 15 (30%) | 12 (24%) | 6 (12%) |
| #Urinary bladder | (44) | (44) | (47) | (42) |
| Cast, NOS | | 1 (2%) | | |
| Hemorrhage | 2 (5%) | 2 (5%) | 3 (6%) | 1 (2%) |
| Inflammation, suppurative | 1 (2%) | | | |
| Inflammation, acute | | | 2 (4%) | 1 (2%) |
| Inflammation, acute diffuse | 1 (2%) | | | |
| Inflammation, acute suppurative | | | 1 (2%) | 1 (2%) |
| Inflammation, acute/chronic | | | 1 (2%) | |
| Inflammation, chronic | 1 (2%) | | 1 (2%) | 2 (5%) |
| Hyperplasia, epithelial | 2 (5%) | 1 (2%) | 3 (6%) | 1 (2%) |
| Hyperplasia, papillary | 1 (2%) | | | |

TABLE A4. SUMMARY OF THE INCIDENCE OF NONNEOPLASTIC LESIONS IN MALE ACI RATS IN THE TWO-YEAR GAVAGE STUDY OF TRICHLOROETHYLENE (Continued)

| | CONTROL (UNTR) | CONTROL (VEH) | LOW DOSE | HIGH DOSE |
|-----------------------------------|----------------|---------------|----------|-----------|
| ENDOCRINE SYSTEM | | | | |
| #Pituitary | (48) | (39) | (33) | (29) |
| Hamartoma | | 1 (3%) | | |
| Pigmentation, NOS | 2 (4%) | 1 (3%) | | |
| Hyperplasia, focal | 1 (2%) | 1 (3%) | 1 (3%) | |
| #Adrenal | (48) | (48) | (44) | (45) |
| Accessory structure | 1 (2%) | | | |
| Hemorrhage | 1 (2%) | | | |
| Inflammation, acute focal | 1 (2%) | | | |
| #Adrenal cortex | (48) | (48) | (44) | (45) |
| Hemorrhage | | | 1 (2%) | |
| Degeneration, NOS | | 1 (2%) | | |
| Metamorphosis, fatty | 1 (2%) | | | |
| Lipoidosis | 2 (4%) | | 2 (5%) | |
| Hyperplasia, focal | | 1 (2%) | | 1 (2%) |
| #Adrenal medulla | (48) | (48) | (44) | (45) |
| Hyperplasia, focal | | 1 (2%) | | 1 (2%) |
| #Thyroid | (48) | (46) | (40) | (39) |
| Hemorrhage | | | 1 (3%) | |
| Inflammation, necrotizing | | | 1 (3%) | |
| Hyperplasia, C-cell | 1 (2%) | | | 1 (3%) |
| REPRODUCTIVE SYSTEM | | | | |
| *Mammary gland | (49) | (50) | (49) | (50) |
| Hyperplasia, epithelial | | | | 1 (2%) |
| Lactation | 3 (6%) | | 1 (2%) | 1 (2%) |
| #Prostate | (48) | (49) | (45) | (43) |
| Dilatation, NOS | | | 1 (2%) | |
| Hemorrhage | | 2 (4%) | 6 (13%) | 3 (7%) |
| Inflammation, NOS | | | 1 (2%) | 2 (5%) |
| Inflammation, focal | | 2 (4%) | | |
| Inflammation, suppurative | 2 (4%) | | 3 (7%) | 2 (5%) |
| Inflammation, acute | | | 3 (7%) | 1 (2%) |
| Inflammation, acute diffuse | 1 (2%) | | | |
| Inflammation, acute suppurative | 1 (2%) | | 1 (2%) | 1 (2%) |
| Abscess, NOS | | | | 1 (2%) |
| Inflammation, acute/chronic | | | 1 (2%) | 1 (2%) |
| Inflammation, chronic | 2 (4%) | | 1 (2%) | |
| Inflammation, chronic focal | | 2 (4%) | | |
| Inflammation, chronic suppurative | | 1 (2%) | | 2 (5%) |
| Abscess, chronic | | | | 1 (2%) |
| Corpora amylacea | | 1 (2%) | | |
| Foreign material, NOS | | 1 (2%) | | |
| Pigmentation, NOS | | 1 (2%) | | |
| Atrophy, NOS | 1 (2%) | | | |
| Hyperplasia, epithelial | | 2 (4%) | | |
| Hyperplasia, focal | | 1 (2%) | | 1 (2%) |
| *Seminal vesicle | (49) | (50) | (49) | (50) |
| Congenital hypoplasia | | | 1 (2%) | |
| Dilatation, NOS | | | | 1 (2%) |
| Hemorrhage | | 2 (4%) | 4 (8%) | 1 (2%) |
| Inflammation, NOS | | | | 1 (2%) |
| Inflammation, acute | | | 3 (6%) | 1 (2%) |
| Inflammation, acute focal | 1 (2%) | | | |
| Inflammation, acute suppurative | 1 (2%) | | 1 (2%) | |
| Inflammation, acute/chronic | | | | 1 (2%) |
| Inflammation, chronic suppurative | | 1 (2%) | | |
| Abscess, chronic | | | | 1 (2%) |
| Fibrosis | 1 (2%) | | | |
| Hypoplasia, NOS | 1 (2%) | | | 1 (2%) |
| Atrophy, NOS | 1 (2%) | | | |

TABLE A4. SUMMARY OF THE INCIDENCE OF NONNEOPLASTIC LESIONS IN MALE ACI RATS IN THE TWO-YEAR GAVAGE STUDY OF TRICHLOROETHYLENE (Continued)

| | CONTROL (UNTR) | CONTROL (VEH) | LOW DOSE | HIGH DOSE |
|--|----------------|---------------|----------|-----------|
| REPRODUCTIVE SYSTEM (Continued) | | | | |
| #Testis | (47) | (49) | (49) | (49) |
| Necrosis, focal | | | 1 (2%) | |
| Infarct, acute | | | 1 (2%) | |
| Calcification, focal | | | 1 (2%) | |
| Atrophy, NOS | 1 (2%) | 8 (16%) | 4 (8%) | 4 (8%) |
| Atrophy, focal | | 1 (2%) | 3 (6%) | |
| Atrophy, diffuse | 1 (2%) | 1 (2%) | 3 (6%) | 2 (4%) |
| Aspermatogenesis | | 1 (2%) | 1 (2%) | 2 (4%) |
| Hypospermatogenesis | 1 (2%) | | | |
| Hyperplasia, interstitial cell | 4 (9%) | 5 (10%) | 5 (10%) | 3 (6%) |
| NERVOUS SYSTEM | | | | |
| #Brain/meninges | (49) | (50) | (47) | (47) |
| Pigmentation, NOS | | | | 1 (2%) |
| #Brain | (49) | (50) | (47) | (47) |
| Corpora amylacea | | | | 1 (2%) |
| SPECIAL SENSE ORGANS | | | | |
| None | | | | |
| MUSCULOSKELETAL SYSTEM | | | | |
| *Skeletal muscle | (49) | (50) | (49) | (50) |
| Abscess, NOS | 1 (2%) | | | |
| *Muscle of neck | (49) | (50) | (49) | (50) |
| Inflammation, NOS | | | | 1 (2%) |
| Necrosis, focal | | | | 1 (2%) |
| BODY CAVITIES | | | | |
| *Mediastinum | (49) | (50) | (49) | (50) |
| Inflammation, focal | | 1 (2%) | | |
| *Pleura | (49) | (50) | (49) | (50) |
| Inflammation, focal | 1 (2%) | | | |
| *Mesentery | (49) | (50) | (49) | (50) |
| Inflammation, granulomatous | | | 1 (2%) | |
| ALL OTHER SYSTEMS | | | | |
| *Multiple organs | (49) | (50) | (49) | (50) |
| Hemorrhage | | 1 (2%) | | 1 (2%) |
| Axilla | | | | |
| Abscess, chronic | | 1 | | |
| SPECIAL MORPHOLOGY SUMMARY | | | | |
| Animal missing/no necropsy | 1 | | | |
| Auto/necropsy/histo perf | | | 1 | 1 |
| Autolysis/no necropsy | | | 1 | |

* Number of animals receiving complete necropsy examination; all gross lesions including masses examined microscopically.
 # Number of animals examined microscopically at this site

APPENDIX B

SUMMARY OF LESIONS IN FEMALE ACI RATS IN THE TWO-YEAR GAVAGE STUDY OF TRICHLOROETHYLENE

| | PAGE | |
|----------|--|-----|
| TABLE B1 | SUMMARY OF THE INCIDENCE OF NEOPLASMS IN FEMALE ACI RATS IN THE TWO-YEAR GAVAGE STUDY OF TRICHLOROETHYLENE | 101 |
| TABLE B2 | INDIVIDUAL ANIMAL TUMOR PATHOLOGY OF FEMALE ACI RATS IN THE TWO-YEAR GAVAGE STUDY OF TRICHLOROETHYLENE | 104 |
| TABLE B3 | ANALYSIS OF PRIMARY TUMORS IN FEMALE ACI RATS IN THE TWO-YEAR GAVAGE STUDY OF TRICHLOROETHYLENE | 112 |
| TABLE B4 | SUMMARY OF THE INCIDENCE OF NONNEOPLASTIC LESIONS IN FEMALE ACI RATS IN THE TWO-YEAR GAVAGE STUDY OF TRICHLOROETHYLENE | 114 |

TABLE B1. SUMMARY OF THE INCIDENCE OF NEOPLASMS IN FEMALE ACI RATS IN THE TWO-YEAR GAVAGE STUDY OF TRICHLOROETHYLENE

| | CONTROL (UNTR) | CONTROL (VEH) | LOW DOSE | HIGH DOSE |
|---|----------------|---------------|----------|-----------|
| ANIMALS INITIALLY IN STUDY | 50 | 50 | 50 | 50 |
| ANIMALS MISSING | 1 | | | |
| ANIMALS NECROPSIED | 49 | 50 | 50 | 46 |
| ANIMALS EXAMINED HISTOPATHOLOGICALLY | 49 | 49 | 50 | 45 |
| INTEGUMENTARY SYSTEM | | | | |
| *Subcutaneous tissue | (49) | (50) | (50) | (46) |
| Adenoma, NOS | 1 (2%) | | | |
| Sarcoma, NOS | | 1 (2%) | | 1 (2%) |
| RESPIRATORY SYSTEM | | | | |
| #Lung | (49) | (49) | (47) | (42) |
| Carcinoma, NOS, metastatic | 1 (2%) | | | |
| Transitional cell carcinoma, metastatic | | | 1 (2%) | |
| Alveolar/bronchiolar adenoma | | | 1 (2%) | |
| Tubular cell adenocarcinoma, metastatic | | | | 1 (2%) |
| Endometrial stromal sarcoma, metastatic | | | | 1 (2%) |
| HEMATOPOIETIC SYSTEM | | | | |
| *Multiple organs | (49) | (50) | (50) | (46) |
| Monocytic leukemia | 2 (4%) | | | |
| #Lymph node | (43) | (47) | (36) | (28) |
| Carcinoma, NOS, metastatic | 1 (2%) | | | |
| Sarcoma, NOS, metastatic | | | | 1 (4%) |
| #Liver | (49) | (49) | (46) | (39) |
| Leukemia, NOS | | | | 1 (3%) |
| #Thymus | (10) | (7) | (19) | (14) |
| Thymoma, malignant | | 1 (14%) | | |
| Malignant lymphoma, lymphocytic type | 1 (10%) | 1 (14%) | 2 (11%) | |
| CIRCULATORY SYSTEM | | | | |
| #Uterus | (48) | (49) | (48) | (40) |
| Angiosarcoma | 1 (2%) | | | |
| DIGESTIVE SYSTEM | | | | |
| #Liver | (49) | (49) | (46) | (39) |
| Neoplastic nodule | 1 (2%) | 2 (4%) | | |
| URINARY SYSTEM | | | | |
| #Kidney | (49) | (48) | (47) | (43) |
| Adenocarcinoma, NOS | | | 1 (2%) | |
| Tubular cell adenoma | | | 2 (4%) | |
| Tubular cell adenocarcinoma | | | | 1 (2%) |
| Sarcoma, NOS | 1 (2%) | | | |
| Nephroblastoma | | 1 (2%) | | |
| #Kidney/pelvis | (49) | (48) | (47) | (43) |
| Transitional cell papilloma | | | 2 (4%) | |
| Transitional cell carcinoma | 1 (2%) | 4 (8%) | 3 (6%) | 1 (2%) |

TABLE B1. SUMMARY OF THE INCIDENCE OF NEOPLASMS IN FEMALE ACI RATS IN THE TWO-YEAR GAVAGE STUDY OF TRICHLOROETHYLENE (Continued)

| | CONTROL (UNTR) | CONTROL (VEH) | LOW DOSE | HIGH DOSE |
|-------------------------------|----------------|---------------|----------|-----------|
| ENDOCRINE SYSTEM | | | | |
| #Pituitary | (47) | (42) | (36) | (32) |
| Carcinoma, NOS | | | | 1 (3%) |
| Adenoma, NOS | 24 (51%) | 18 (43%) | 7 (19%) | 7 (22%) |
| Adenocarcinoma, NOS | | 1 (2%) | | |
| Chromophobe adenoma | | 1 (2%) | | |
| Chromophobe carcinoma | | 1 (2%) | | |
| #Adrenal | (49) | (48) | (44) | (40) |
| Cortical adenoma | 1 (2%) | 2 (4%) | 1 (2%) | 3 (8%) |
| Cortical carcinoma | | 1 (2%) | | |
| Pheochromocytoma | 1 (2%) | 3 (6%) | | 2 (5%) |
| Pheochromocytoma, malignant | | | 1 (2%) | |
| #Thyroid | (46) | (47) | (44) | (38) |
| Follicular cell adenoma | | | | 1 (3%) |
| REPRODUCTIVE SYSTEM | | | | |
| *Mammary gland | (49) | (50) | (50) | (46) |
| Adenoma, NOS | | 1 (2%) | | 1 (2%) |
| Adenocarcinoma, NOS | 1 (2%) | 1 (2%) | 1 (2%) | |
| Papillary adenoma | 1 (2%) | | | |
| Fibroadenoma | 1 (2%) | 1 (2%) | | |
| #Uterus | (48) | (49) | (48) | (40) |
| Carcinoma, NOS | 1 (2%) | | | |
| Adenocarcinoma, NOS | 1 (2%) | | | |
| Endometrial stromal polyp | 1 (2%) | 4 (8%) | 3 (6%) | 2 (5%) |
| Endometrial stromal sarcoma | | | | 1 (3%) |
| #Cervix uteri | (48) | (49) | (48) | (40) |
| Endometrial stromal sarcoma | | | 1 (2%) | |
| #Uterus/endometrium | (48) | (49) | (48) | (40) |
| Carcinoma, NOS | | | | 1 (3%) |
| #Ovary | (48) | (46) | (46) | (38) |
| Luteoma | 1 (2%) | | | |
| NERVOUS SYSTEM | | | | |
| #Brain | (49) | (48) | (49) | (42) |
| Granular cell tumor, NOS | | 1 (2%) | | 1 (2%) |
| SPECIAL SENSE ORGANS | | | | |
| *Zymbal gland | (49) | (50) | (50) | (46) |
| Adenosquamous carcinoma | | 1 (2%) | | |
| MUSCULOSKELETAL SYSTEM | | | | |
| *Skeletal muscle | (49) | (50) | (50) | (46) |
| Sarcoma, NOS | | | | 1 (2%) |
| BODY CAVITIES | | | | |
| None | | | | |

TABLE B1. SUMMARY OF THE INCIDENCE OF NEOPLASMS IN FEMALE ACI RATS IN THE TWO-YEAR GAVAGE STUDY OF TRICHLOROETHYLENE (Continued)

| | CONTROL (UNTR) | CONTROL (VEH) | LOW DOSE | HIGH DOSE |
|--|----------------|---------------|----------|-----------|
| ALL OTHER SYSTEMS | | | | |
| *Multiple organs | (49) | (50) | (50) | (46) |
| Sarcoma, NOS | | | 1 (2%) | |
| Periorbital region | | | | |
| Sarcoma, NOS | 1 | | | |
| ANIMAL DISPOSITION SUMMARY | | | | |
| Animals initially in study | 50 | 50 | 50 | 50 |
| Natural death | 11 | 5 | 12 | 17 |
| Moribund sacrifice | 2 | 10 | 4 | 4 |
| Terminal sacrifice | 36 | 33 | 20 | 17 |
| Dosing accident | | | 2 | 3 |
| Accidentally killed, NOS | | 2 | 12 | 9 |
| Animal missing | 1 | | | |
| TUMOR SUMMARY | | | | |
| Total animals with primary tumors** | 34 | 33 | 21 | 18 |
| Total primary tumors | 42 | 46 | 26 | 25 |
| Total animals with benign tumors | 28 | 27 | 15 | 14 |
| Total benign tumors | 31 | 30 | 16 | 16 |
| Total animals with malignant tumors | 9 | 12 | 10 | 8 |
| Total malignant tumors | 10 | 13 | 10 | 8 |
| Total animals with secondary tumors## | 1 | | 1 | 3 |
| Total secondary tumors | 2 | | 1 | 3 |
| Total animals with tumors uncertain--benign or malignant | 1 | 3 | | 1 |
| Total uncertain tumors | 1 | 3 | | 1 |

* Number of animals receiving complete necropsy examination; all gross lesions including masses examined microscopically.

** Primary tumors: all tumors except secondary tumors

Number of animals examined microscopically at this site

Secondary tumors: metastatic tumors or tumors invasive into an adjacent organ

TABLE B2. INDIVIDUAL ANIMAL TUMOR PATHOLOGY OF FEMALE ACI RATS IN THE TWO-YEAR GAVAGE STUDY OF TRICHLOROETHYLENE: UNTREATED CONTROL

| ANIMAL NUMBER | 05 | 06 | 07 | 08 | 09 | 10 | 11 | 12 | 13 | 14 | 15 | 16 | 17 | 18 | 19 | 20 | 21 | 22 | 23 | 24 | 25 | 26 | 27 | 28 | 29 | 30 | | |
|--------------------------------------|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|---|--|
| WEEKS ON STUDY | 106 | 104 | 106 | 106 | 106 | 106 | 105 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | | |
| INTEGUMENTARY SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Subcutaneous tissue | + | + | + | + | + | + | M | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| Adenoma, NOS | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| RESPIRATORY SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Lungs and bronchi | + | + | + | + | + | + | M | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| Carcinoma, NOS, metastatic | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Trachea | + | + | + | + | + | + | M | + | + | + | + | - | + | + | + | + | + | + | + | + | + | + | + | + | + | + | - | |
| HEMATOPOIETIC SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Bone marrow | + | + | + | + | + | + | M | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| Spleen | + | + | + | + | + | + | M | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| Lymph nodes | + | + | + | + | + | + | M | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| Carcinoma, NOS, metastatic | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Thymus | - | - | - | - | - | - | M | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | |
| Malignant lymphoma, lymphocytic type | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| CIRCULATORY SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Heart | + | + | + | + | + | + | M | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| DIGESTIVE SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Salivary gland | + | + | + | + | + | + | M | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| Liver | + | + | + | + | + | + | M | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| Neoplastic nodule | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Bile duct | + | + | + | + | + | + | M | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| Gallbladder & common bile duct | N | N | N | N | N | N | M | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | |
| Pancreas | + | + | + | + | + | + | M | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| Esophagus | + | + | + | + | + | + | M | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| Stomach | + | + | + | + | + | + | M | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| Small intestine | + | + | + | + | + | + | M | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| Large intestine | + | + | + | + | + | + | M | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| URINARY SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Kidney | + | + | + | + | + | + | M | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| Sarcoma, NOS | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Kidney/pelvis | + | + | + | + | + | + | M | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| Transitional cell carcinoma | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Urinary bladder | + | + | + | + | + | + | M | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| ENDOCRINE SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Pituitary | + | + | + | + | + | + | M | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| Adenoma, NOS | | X | X | | X | X | | | | | X | X | | X | X | | X | | X | | X | | X | | X | | | |
| Adrenal | + | + | + | + | + | + | M | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| Cortical adenoma | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Pheochromocytoma | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Thyroid | + | - | + | + | + | + | M | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | - | |
| Parathyroid | | | | | | | M | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | - | |
| REPRODUCTIVE SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Mammary gland | + | + | + | + | + | + | M | + | + | + | + | N | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| Adenocarcinoma, NOS | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Papillary adenoma | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Fibroadenoma | X | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Uterus | + | + | + | + | + | + | M | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| Carcinoma, NOS | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Adenocarcinoma, NOS | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Endometrial stromal polyp | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Angiosarcoma | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Ovary | + | + | + | + | + | + | M | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| Luteoma | | | | | | | | | | | | | | | | | | | | | | | | | | | X | |
| NERVOUS SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Brain | + | + | + | + | + | + | M | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| ALL OTHER SYSTEMS | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Multiple organs, NOS | N | N | N | N | N | N | M | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | |
| Monocytic leukemia | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Periorbital region | | | | | | | M | | | | | | | | | | | | | | | | | | | | | |
| Sarcoma, NOS | | | | | | | | | | | | | | | | | | | | | | | | | | | | |

+: Tissue examined microscopically
 -: Required tissue not examined microscopically
 X: Tumor incidence
 N: Necropsy, no autolysis, no microscopic examination
 S: Animal missexed

: No tissue information submitted
 C: Necropsy, no histology due to protocol
 A: Autolysis
 M: Animal missing
 B: No necropsy performed

**TABLE B2. INDIVIDUAL ANIMAL TUMOR PATHOLOGY OF FEMALE ACI RATS:
UNTREATED CONTROL (Continued)**

| ANIMAL NUMBER | 9 0 2 | 9 0 6 | 9 1 4 | 9 1 5 | 9 1 9 | 9 2 6 | 9 3 6 | 9 4 0 | 9 4 8 | 9 5 0 | 9 5 8 | 9 6 9 | 9 7 0 | 9 7 5 | 9 7 6 | 9 7 7 | 9 7 8 | 9 8 1 | 9 8 4 | 9 9 0 | 9 9 1 | 9 9 2 | 9 9 4 | 9 9 6 | TOTAL TISSUES TUMORS |
|--------------------------------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|---------------|----------------------|
| WEEKS ON STUDY | 0 5 4 | 1 0 6 | 0 9 9 | 1 0 6 | 1 0 6 | 1 0 6 | 0 7 4 | 1 0 6 | 1 0 6 | 0 7 6 | 1 0 6 | |
| INTEGUMENTARY SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | |
| Subcutaneous tissue | + | | | | | | | | | | | | | | | | | | | | | | | *49 1 | |
| Adenoma, NOS | + | | | | | | | | | | | | | | | | | | | | | | | | |
| RESPIRATORY SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | |
| Lungs and bronchi | + | | | | | | | | | | | | | | | | | | | | | | | 49 1 48 | |
| Carcinoma, NOS, metastatic | + | | | | | | | | | | | | | | | | | | | | | | | | |
| Trachea | - | | | | | | | | | | | | | | | | | | | | | | | | |
| HEMATOPOIETIC SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | |
| Bone marrow | + | | | | | | | | | | | | | | | | | | | | | | | 49 48 | |
| Spleen | + | | | | | | | | | | | | | | | | | | | | | | | | |
| Lymph nodes | + | | | | | | | | | | | | | | | | | | | | | | | 43 1 | |
| Carcinoma, NOS, metastatic | - | | | | | | | | | | | | | | | | | | | | | | | | |
| Thymus | - | | | | | | | | | | | | | | | | | | | | | | | 10 1 | |
| Malignant lymphoma, lymphocytic type | - | | | | | | | | | | | | | | | | | | | | | | | | |
| CIRCULATORY SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | |
| Heart | + | | | | | | | | | | | | | | | | | | | | | | | 49 | |
| DIGESTIVE SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | |
| Salivary gland | + | | | | | | | | | | | | | | | | | | | | | | | 45 49 | |
| Liver | + | | | | | | | | | | | | | | | | | | | | | | | | |
| Neoplastic nodule | + | | | | | | | | | | | | | | | | | | | | | | | 1 49 | |
| Bile duct | - | | | | | | | | | | | | | | | | | | | | | | | | |
| Gallbladder & common bile duct | N | | | | | | | | | | | | | | | | | | | | | | | *49 48 | |
| Pancreas | + | | | | | | | | | | | | | | | | | | | | | | | | |
| Esophagus | + | | | | | | | | | | | | | | | | | | | | | | | 47 49 | |
| Stomach | + | | | | | | | | | | | | | | | | | | | | | | | | |
| Small intestine | + | | | | | | | | | | | | | | | | | | | | | | | 49 48 | |
| Large intestine | + | | | | | | | | | | | | | | | | | | | | | | | | |
| URINARY SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | |
| Kidney | + | | | | | | | | | | | | | | | | | | | | | | | 49 1 | |
| Sarcoma, NOS | + | | | | | | | | | | | | | | | | | | | | | | | | |
| Kidney/pelvis | + | | | | | | | | | | | | | | | | | | | | | | | 49 1 | |
| Transitional cell carcinoma | - | | | | | | | | | | | | | | | | | | | | | | | | |
| Urinary bladder | + | | | | | | | | | | | | | | | | | | | | | | | 47 | |
| ENDOCRINE SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | |
| Pituitary | + | | | | | | | | | | | | | | | | | | | | | | | 47 24 | |
| Adenoma, NOS | + | | | | | | | | | | | | | | | | | | | | | | | | |
| Adrenal | + | | | | | | | | | | | | | | | | | | | | | | | 49 1 | |
| Cortical adenoma | + | | | | | | | | | | | | | | | | | | | | | | | | |
| Pheochromocytoma | + | | | | | | | | | | | | | | | | | | | | | | | 1 | |
| Thyroid | - | | | | | | | | | | | | | | | | | | | | | | | 46 31 | |
| Parathyroid | - | | | | | | | | | | | | | | | | | | | | | | | | |
| REPRODUCTIVE SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | |
| Mammary gland | + | | | | | | | | | | | | | | | | | | | | | | | *49 1 | |
| Adenocarcinoma, NOS | + | | | | | | | | | | | | | | | | | | | | | | | | |
| Papillary adenoma | + | | | | | | | | | | | | | | | | | | | | | | | 1 1 | |
| Fibroadenoma | + | | | | | | | | | | | | | | | | | | | | | | | | |
| Uterus | + | | | | | | | | | | | | | | | | | | | | | | | 48 1 | |
| Carcinoma, NOS | - | | | | | | | | | | | | | | | | | | | | | | | | |
| Adenocarcinoma, NOS | + | | | | | | | | | | | | | | | | | | | | | | | 1 1 | |
| Endometrial stromal polyp | + | | | | | | | | | | | | | | | | | | | | | | | | |
| Angiosarcoma | + | | | | | | | | | | | | | | | | | | | | | | | 1 | |
| Ovary | + | | | | | | | | | | | | | | | | | | | | | | | 48 1 | |
| Luteoma | + | | | | | | | | | | | | | | | | | | | | | | | | |
| NERVOUS SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | |
| Brain | + | | | | | | | | | | | | | | | | | | | | | | | 49 | |
| ALL OTHER SYSTEMS | | | | | | | | | | | | | | | | | | | | | | | | | |
| Multiple organs, NOS | N | | | | | | | | | | | | | | | | | | | | | | | *49 2 | |
| Monocytic leukemia | N | | | | | | | | | | | | | | | | | | | | | | | | |
| Periorbital region | N | | | | | | | | | | | | | | | | | | | | | | | 1 | |
| Sarcoma, NOS | X | | | | | | | | | | | | | | | | | | | | | | | | |

* Animals necropsied

TABLE B2. INDIVIDUAL ANIMAL TUMOR PATHOLOGY OF FEMALE ACI RATS IN THE TWO-YEAR GAVAGE STUDY OF TRICHLOROETHYLENE: VEHICLE CONTROL

| ANIMAL NUMBER | 0 | 0 | 0 | 0 | 0 | 8 | 8 | 8 | 8 | 8 | 8 | 8 | 8 | 8 | 8 | 8 | 8 | 8 | 8 | 8 | 8 | 8 | 8 | 9 |
|--------------------------------------|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| WEEKS ON STUDY | 0 | 2 | 2 | 3 | 4 | 4 | 4 | 4 | 5 | 5 | 5 | 5 | 6 | 6 | 6 | 6 | 7 | 7 | 8 | 8 | 8 | 8 | 8 | 9 |
| | 0 | 0 | 1 | 2 | 5 | 4 | 5 | 9 | 0 | 6 | 8 | 9 | 0 | 6 | 9 | 0 | 8 | 2 | 3 | 5 | 9 | 0 | 2 | 7 |
| INTEGUMENTARY SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | |
| Subcutaneous tissue | | | | | | | | | | | | | | | | | | | | | | | | |
| Sarcoma, NOS | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + |
| RESPIRATORY SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | |
| Lungs and bronchi | | | | | | | | | | | | | | | | | | | | | | | | |
| Trachea | + | + | + | + | + | - | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + |
| HEMATOPOIETIC SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | |
| Bone marrow | | | | | | | | | | | | | | | | | | | | | | | | |
| Spleen | | | | | | | | | | | | | | | | | | | | | | | | |
| Lymph nodes | | | | | | | | | | | | | | | | | | | | | | | | |
| Thymus | - | - | - | - | + | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - |
| Thymoma, malignant | | | | | | | | | | | | | | | | | | | | | | | | |
| Malignant lymphoma, lymphocytic type | | | | | | | | | | | | | | | | | | | | | | | | |
| CIRCULATORY SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | |
| Heart | | | | | | | | | | | | | | | | | | | | | | | | |
| DIGESTIVE SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | |
| Salivary gland | | | | | | | | | | | | | | | | | | | | | | | | |
| Liver | | | | | | | | | | | | | | | | | | | | | | | | |
| Neoplastic nodule | | | | | | | | | | | | | | | | | | | | | | | | |
| Bile duct | | | | | | | | | | | | | | | | | | | | | | | | |
| Gallbladder & common bile duct | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N |
| Pancreas | | | | | | | | | | | | | | | | | | | | | | | | |
| Esophagus | | | | | | | | | | | | | | | | | | | | | | | | |
| Stomach | | | | | | | | | | | | | | | | | | | | | | | | |
| Small intestine | | | | | | | | | | | | | | | | | | | | | | | | |
| Large intestine | | | | | | | | | | | | | | | | | | | | | | | | |
| URINARY SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | |
| Kidney | | | | | | | | | | | | | | | | | | | | | | | | |
| Nephroblastoma | | | | | | | | | | | | | | | | | | | | | | | | |
| Kidney/pelvis | | | | | | | | | | | | | | | | | | | | | | | | |
| Transitional cell carcinoma | | | | | | | | | | | | | | | | | | | | | | | | |
| Urinary bladder | | | | | | | | | | | | | | | | | | | | | | | | |
| ENDOCRINE SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | |
| Pituitary | | | | | | | | | | | | | | | | | | | | | | | | |
| Adenoma, NOS | | | | | | | | | | | | | | | | | | | | | | | | |
| Adenocarcinoma, NOS | | | | | | | | | | | | | | | | | | | | | | | | |
| Chromophobe adenoma | | | | | | | | | | | | | | | | | | | | | | | | |
| Chromophobe carcinoma | | | | | | | | | | | | | | | | | | | | | | | | |
| Adrenal | | | | | | | | | | | | | | | | | | | | | | | | |
| Cortical adenoma | | | | | | | | | | | | | | | | | | | | | | | | |
| Cortical carcinoma | | | | | | | | | | | | | | | | | | | | | | | | |
| Pheochromocytoma | | | | | | | | | | | | | | | | | | | | | | | | |
| Thyroid | | | | | | | | | | | | | | | | | | | | | | | | |
| Parathyroid | | | | | | | | | | | | | | | | | | | | | | | | |
| REPRODUCTIVE SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | |
| Mammary gland | | | | | | | | | | | | | | | | | | | | | | | | |
| Adenoma, NOS | | | | | | | | | | | | | | | | | | | | | | | | |
| Adenocarcinoma, NOS | | | | | | | | | | | | | | | | | | | | | | | | |
| Fibroadenoma | | | | | | | | | | | | | | | | | | | | | | | | |
| Uterus | | | | | | | | | | | | | | | | | | | | | | | | |
| Endometrial stromal polyp | | | | | | | | | | | | | | | | | | | | | | | | |
| Ovary | | | | | | | | | | | | | | | | | | | | | | | | |
| NERVOUS SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | |
| Brain | | | | | | | | | | | | | | | | | | | | | | | | |
| Granular cell tumor, NOS | | | | | | | | | | | | | | | | | | | | | | | | |
| SPECIAL SENSE ORGANS | | | | | | | | | | | | | | | | | | | | | | | | |
| Zymbal gland | | | | | | | | | | | | | | | | | | | | | | | | |
| Adenosquamous carcinoma | N | N | N | N | N | + | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N |

TABLE B2. INDIVIDUAL ANIMAL TUMOR PATHOLOGY OF FEMALE ACI RATS: VEHICLE CONTROL (Continued)

| ANIMAL NUMBER | 9 0 5 | 9 0 7 | 9 0 8 | 9 1 8 | 9 2 8 | 9 3 9 | 9 4 1 | 9 4 6 | 9 4 6 | 9 5 4 | 9 5 5 | 9 5 5 | 9 6 0 | 9 6 4 | 9 6 6 | 9 6 6 | 9 7 3 | 9 8 0 | 9 8 3 | 9 8 5 | 9 8 6 | 9 9 3 | 9 9 5 | TOTAL: TISSUES TUMORS | |
|--------------------------------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-----------------------------|-------------|
| WEEKS ON STUDY | 1 0 6 | 1 0 6 | 1 0 6 | 1 0 6 | 2 6 8 | 2 4 1 | 2 0 6 | 2 0 6 | 2 0 6 | 3 0 6 | 3 0 6 | 3 0 6 | 3 0 6 | 4 0 6 | 4 0 6 | 4 0 6 | 5 0 6 | 5 0 6 | 6 0 6 | 6 0 6 | 6 0 6 | 6 0 6 | 6 0 6 | 6 0 6 | 6 0 6 |
| INTEGUMENTARY SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | |
| Subcutaneous tissue | + | + | N | + | + | + | + | + | + | + | + | + | + | + | + | + | + | N | + | + | + | + | + | + | *50 |
| Sarcoma, NOS | | | | | | | | | | | | | X | | | | | | | | | | | | 1 |
| RESPIRATORY SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | |
| Lungs and bronchi | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | A | + | + | + | + | + | + | 49 |
| Trachea | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | A | + | + | + | + | + | + | 47 |
| HEMATOPOIETIC SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | |
| Bone marrow | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | A | + | + | - | + | + | + | 48 |
| Spleen | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | A | + | + | + | + | + | + | 49 |
| Lymph nodes | + | + | + | - | + | + | + | + | + | + | + | + | + | + | + | + | + | A | - | + | + | + | + | + | 47 |
| Thymus | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | A | - | - | + | - | - | - | 7 |
| Thymoma, malignant | | | | | | | | | | | | | | | | | | | | | | | | 1 | |
| Malignant lymphoma, lymphocytic type | | | | | | | | | | | | | | | | | | | | | | | | 1 | |
| CIRCULATORY SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | |
| Heart | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | A | + | + | - | + | + | + | 48 |
| DIGESTIVE SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | |
| Salivary gland | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | A | + | + | - | + | + | + | 45 |
| Liver | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | A | + | + | + | + | + | + | 49 |
| Neoplastic nodule | | | | | | | | | | | | | | | | | | | | | | | | 2 | |
| Bile duct | N | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | A | + | + | + | + | + | + | 49 |
| Gallbladder & common bile duct | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | *50 |
| Pancreas | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | A | + | + | + | + | + | + | 48 |
| Esophagus | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | A | + | + | + | + | + | + | 47 |
| Stomach | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | A | + | + | - | + | + | + | 47 |
| Small intestine | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | A | + | + | + | + | + | + | 49 |
| Large intestine | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | A | - | + | + | + | + | + | 47 |
| URINARY SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | |
| Kidney | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | A | + | + | + | + | + | + | 48 |
| Nephroblastoma | | | | | | | | | | | | | | | | | | | X | | | | | 1 | |
| Kidney/pelvis | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | A | + | + | + | + | + | + | 48 |
| Transitional cell carcinoma | | | | | | | | | | | | | | | | | | | X | | | | | 4 | |
| Urinary bladder | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | A | + | + | - | + | + | + | 43 |
| ENDOCRINE SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | |
| Pituitary | + | + | + | - | + | + | + | + | + | + | + | + | + | + | + | + | + | A | + | + | - | - | + | + | 42 |
| Adenoma, NOS | X | X | | | | | | | | X | | | X | X | X | | | | X | | | | | 18 | |
| Adenocarcinoma, NOS | | | | | | | | | | X | | | | | | | | | | | | | | 1 | |
| Chromophobe adenoma | | | | | | | | | | | | | | | | | | | | | | X | | 1 | |
| Chromophobe carcinoma | | | | | | | | | | | | | | | | | | | | | | | | 1 | |
| Adrenal | + | + | - | + | + | + | + | + | + | + | + | + | + | + | + | + | + | A | + | + | - | + | + | + | 48 |
| Cortical adenoma | | | | | | | | | | | | | | | | | | | | | | | | 2 | |
| Cortical carcinoma | | | | | | | | | | | | | | | | | | | | | | | | 1 | |
| Pheochromocytoma | | | | X | | | | | | | | | | | | | | | X | | | | | 3 | |
| Thyroid | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | A | + | + | + | - | + | + | 47 |
| Parathyroid | - | - | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | A | + | + | + | + | + | + | 35 |
| REPRODUCTIVE SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | |
| Mammary gland | + | + | N | + | + | + | + | + | + | + | + | + | + | + | + | + | + | N | N | + | + | N | + | + | *50 |
| Adenoma, NOS | | | | | | | | | | | | | | | | | | | | | | | | 1 | |
| Adenocarcinoma, NOS | | | | | | | | | | | | | | | | | | | | | | | | 1 | |
| Fibroadenoma | | | | | | | | | | | | | | | | | | | | | X | | | 1 | |
| Uterus | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | A | + | + | + | + | + | + | 49 |
| Endometrial stromal polyp | | | | | | | | | | | | | | | | | | | | | | | | 4 | |
| Ovary | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | A | - | + | + | + | + | + | 46 |
| NERVOUS SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | |
| Brain | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | A | + | + | - | + | + | + | 48 |
| Granular cell tumor, NOS | | | | | | | | | | | | | | | | | | | | | | | | 1 | |
| SPECIAL SENSE ORGANS | | | | | | | | | | | | | | | | | | | | | | | | | |
| Zymbal gland | N | N | N | N | + | + | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | *50 |
| Adenosquamous carcinoma | | | | | X | | | | | | | | | | | | | | | | | | | 1 | |

* Animals necropsied

**TABLE B2. INDIVIDUAL ANIMAL TUMOR PATHOLOGY OF FEMALE ACI RATS: LOW DOSE
(Continued)**

| ANIMAL NUMBER | 8 9 | 8 9 | 9 0 | 9 0 | 9 1 | 9 1 | 9 1 | 9 1 | 9 2 | 9 2 | 9 2 | 9 3 | 9 3 | 9 3 | 9 3 | 9 4 | 9 4 | 9 4 | 9 5 | 9 6 | 9 7 | 9 7 | 9 8 | 9 8 | 9 9 | 9 9 | TOTAL TISSUES TUMORS | |
|---|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|----------------------|--------|
| WEEKS ON STUDY | 6 8 | 1 2 | 0 5 | 0 4 | 0 4 | 0 4 | 0 5 | 0 1 | 0 4 | 0 8 | 0 8 | 0 9 | 0 0 | 0 2 | 0 5 | 0 0 | 0 4 | 0 4 | 0 5 | 0 0 | 0 7 | 0 4 | 0 5 | 0 8 | 0 8 | 0 4 | 0 1 | 0 4 |
| RESPIRATORY SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Lungs and bronchi | + | + | + | + | + | + | + | + | + | - | + | + | + | + | + | + | + | + | + | + | + | + | + | - | + | + | + | |
| Transitional cell carcinoma, metastatic | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Alveolar/bronchiolar adenoma | | | X | | | | | | | | | | | | | | | | | | | | | | | | | |
| Trachea | + | + | + | + | + | + | + | + | + | + | + | + | + | - | + | A | - | + | + | + | - | + | - | + | + | + | + | |
| HEMATOPOIETIC SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Bone marrow | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | A | + | + | + | + | + | + | + | - | + | + | + | |
| Spleen | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | A | + | + | + | + | + | + | + | - | + | + | + | |
| Lymph nodes | - | + | + | + | - | + | + | + | + | - | + | - | + | - | + | A | + | + | + | + | + | + | + | - | + | + | + | |
| Thymus | - | + | - | - | - | - | + | + | + | + | + | + | + | + | + | A | + | + | + | - | + | + | - | - | - | + | - | |
| Malignant lymphoma, lymphocytic type | | | | | | | | | | X | | | | | | | | | | | X | | | | | | | |
| CIRCULATORY SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Heart | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | - | + | + | |
| DIGESTIVE SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Salivary gland | + | + | + | + | + | + | + | + | + | - | + | + | + | + | + | A | + | + | + | + | - | + | + | + | + | - | + | |
| Liver | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | A | + | + | + | + | + | + | + | - | + | + | + | |
| Bile duct | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | A | + | + | + | + | + | + | + | + | + | + | + | |
| Gallbladder & common bile duct | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | |
| Pancreas | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | A | + | + | + | + | + | + | + | - | + | + | + | |
| Esophagus | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | A | + | + | + | + | + | + | + | + | + | + | + | |
| Stomach | - | + | + | + | + | + | + | + | + | + | + | + | + | + | + | A | + | + | + | + | + | + | + | - | + | + | + | |
| Small intestine | + | - | + | + | + | + | + | + | + | + | + | + | + | + | + | A | + | + | + | + | + | + | + | + | + | + | + | |
| Large intestine | + | - | + | + | + | + | + | + | + | + | + | + | + | + | + | A | + | + | + | + | + | + | + | + | + | + | + | |
| URINARY SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Kidney | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | - | + | + | |
| Adenocarcinoma, NOS | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Tubular cell adenoma | | | | | | | | | | | | | | | | | | | | | | | X | | | | | |
| Kidney/pelvis | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | - | + | + | |
| Transitional cell papilloma | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Transitional cell carcinoma | | | | | | | | | | | | | | | | | | | | | | | | | | | X | |
| Urinary bladder | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | A | + | + | + | - | + | + | + | + | + | + | - | |
| ENDOCRINE SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Pituitary | + | - | + | + | + | + | + | + | + | - | + | X | - | + | A | + | + | + | + | + | + | + | + | - | + | + | + | |
| Adenoma, NOS | | | | | | | | | | | | | | | | | | | | | | | X | | | | X | |
| Adrenal | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | A | + | + | + | + | + | + | + | + | - | + | + | |
| Cortical adenoma | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Pheochromocytoma, malignant | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Thyroid | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | A | + | + | + | + | + | + | + | + | + | + | + | |
| Parathyroid | - | + | + | - | - | + | + | + | - | - | + | - | + | - | + | A | + | - | + | + | - | - | - | - | - | + | + | |
| REPRODUCTIVE SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Mammary gland | + | + | + | + | + | + | + | + | + | N | + | + | + | + | N | N | + | + | + | + | + | + | N | + | + | + | N | |
| Adenocarcinoma, NOS | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Uterus | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | A | + | + | + | + | + | + | + | + | + | + | + | |
| Endometrial stromal polyp | | | | | | | | | | | X | | | | | | | | | | | X | | | | | | |
| Endometrial stromal sarcoma | | | | | | | | | | | | | X | | | | | | | | | | | | | | | |
| Ovary | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | A | + | + | + | + | + | + | + | + | + | + | + | |
| NERVOUS SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Brain | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | A | + | + | + | + | + | + | + | + | + | + | + | |
| ALL OTHER SYSTEMS | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Multiple organs, NOS | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | |
| Sarcoma, NOS | | | | | | | | | | | | | | | | | | | | | | | | | | | | |

* Animals necropsied

TABLE B2. INDIVIDUAL ANIMAL TUMOR PATHOLOGY OF FEMALE ACI RATS IN THE TWO-YEAR GAVAGE STUDY OF TRICHLOROETHYLENE: HIGH DOSE

| ANIMAL NUMBER | 01 | 02 | 03 | 04 | 05 | 06 | 07 | 08 | 09 | 10 | 11 | 12 | 13 | 14 | 15 | 16 | 17 | 18 | 19 | 20 | 21 | 22 | 23 | 24 | 25 | 26 | 27 | 28 | 29 | 30 | |
|---|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|-----|-----|-----|-----|
| WEEKS ON STUDY | 73 | 74 | 75 | 76 | 77 | 78 | 79 | 80 | 81 | 82 | 83 | 84 | 85 | 86 | 87 | 88 | 89 | 90 | 91 | 92 | 93 | 94 | 95 | 96 | 97 | 98 | 99 | 100 | 101 | 102 | 103 |
| INTEGUMENTARY SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Subcutaneous tissue | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| Sarcoma, NOS | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| RESPIRATORY SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Lungs and bronchi | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| Tubular cell adenocarcinoma, metastatic | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Endometrial stromal sarcoma, metastatic | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Trachea | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| HEMATOPOIETIC SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Bone marrow | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| Spleen | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| Lymph nodes | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| Sarcoma, NOS, metastatic | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Thymus | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| CIRCULATORY SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Heart | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| DIGESTIVE SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Salivary gland | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| Liver | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| Leukemia, NOS | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Bile duct | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| Gallbladder & common bile duct | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | |
| Pancreas | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| Esophagus | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| Stomach | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| Small intestine | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| Large intestine | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| URINARY SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Kidney | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| Tubular cell adenocarcinoma | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Kidney/pelvis | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| Transitional cell carcinoma | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Urinary bladder | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| ENDOCRINE SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Pituitary | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| Carcinoma, NOS | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Adenoma, NOS | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Adrenal | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| Cortical adenoma | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Pheochromocytoma | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Thyroid | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| Follicular cell adenoma | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Parathyroid | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| REPRODUCTIVE SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Mammary gland | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| Adenoma, NOS | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Uterus | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| Carcinoma, NOS | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Endometrial stromal polyp | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Endometrial stromal sarcoma | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Ovary | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| NERVOUS SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Brain | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| Granular cell tumor, NOS | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| MUSCULOSKELETAL SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Muscle | N | N | + | N | N | + | N | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| Sarcoma, NOS | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |

TABLE B3. ANALYSIS OF PRIMARY TUMORS IN FEMALE ACI RATS IN THE TWO-YEAR GAVAGE STUDY OF TRICHLOROETHYLENE

| | Untreated Control | Vehicle Control | 500 mg/kg | 1,000 mg/kg |
|---|-------------------|-----------------|------------|-------------|
| Kidney: Transitional Cell Carcinoma | | | | |
| Overall Rates (a) | 1/49 (2%) | 4/48 (8%) | 3/47 (6%) | 1/43 (2%) |
| Adjusted Rates (b) | 2.7% | 11.8% | 12.6% | 5.3% |
| Terminal Rates (c) | 1/37 (3%) | 4/34 (12%) | 2/20 (10%) | 1/19 (5%) |
| Life Table Tests (d) | | P=0.346N | P=0.549 | P=0.388N |
| Incidental Tumor Tests (d) | | P=0.264N | P=0.642N | P=0.388N |
| Cochran-Armitage Trend Test (d) | | P=0.160N | | |
| Fisher Exact Test | | | P=0.512N | P=0.217N |
| Kidney: Transitional Cell Papilloma or Carcinoma | | | | |
| Overall Rates (a) | 1/49 (2%) | 4/48 (8%) | 5/47 (11%) | 1/43 (2%) |
| Adjusted Rates (b) | 2.7% | 11.8% | 20.6% | 5.3% |
| Terminal Rates (c) | 1/37 (3%) | 4/34 (12%) | 3/20 (15%) | 1/19 (5%) |
| Life Table Tests (d) | | P=0.427N | P=0.216 | P=0.388N |
| Incidental Tumor Tests (d) | | P=0.299N | P=0.407 | P=0.388N |
| Cochran-Armitage Trend Test (d) | | P=0.193N | | |
| Fisher Exact Test | | | P=0.486 | P=0.217N |
| Kidney: Adenoma or Adenocarcinoma | | | | |
| Overall Rates (a) | 0/49 (0%) | 0/48 (0%) | 3/47 (6%) | 1/43 (2%) |
| Adjusted Rates (b) | 0.0% | 0.0% | 13.9% | 4.3% |
| Terminal Rates (c) | 0/37 (0%) | 0/34 (0%) | 2/20 (10%) | 0/19 (0%) |
| Life Table Tests (d) | | P=0.200 | P=0.044 | P=0.375 |
| Incidental Tumor Tests (d) | | P=0.310 | P=0.104 | P=0.581 |
| Cochran-Armitage Trend Test (d) | | P=0.343 | | |
| Fisher Exact Test | | | P=0.117 | P=0.473 |
| Pituitary: Adenoma | | | | |
| Overall Rates (a) | 24/47 (51%) | 19/42 (45%) | 7/36 (19%) | 7/32 (22%) |
| Adjusted Rates (b) | 59.8% | 52.1% | 33.7% | 36.8% |
| Terminal Rates (c) | 21/37 (57%) | 16/33 (48%) | 6/19 (32%) | 7/19 (37%) |
| Life Table Tests (d) | | P=0.095N | P=0.144N | P=0.149N |
| Incidental Tumor Tests (d) | | P=0.068N | P=0.073N | P=0.109N |
| Cochran-Armitage Trend Test (d) | | P=0.015N | | |
| Fisher Exact Test | | | P=0.014N | P=0.032N |
| Pituitary: Adenoma, Adenocarcinoma, or Carcinoma | | | | |
| Overall Rates (a) | 24/47 (51%) | 21/42 (50%) | 7/36 (19%) | 8/32 (25%) |
| Adjusted Rates (b) | 59.8% | 55.9% | 33.7% | 40.0% |
| Terminal Rates (c) | 21/37 (57%) | 17/33 (52%) | 6/19 (32%) | 7/19 (37%) |
| Life Table Tests (d) | | P=0.085N | P=0.076N | P=0.146N |
| Incidental Tumor Tests (d) | | P=0.052N | P=0.029N | P=0.085N |
| Cochran-Armitage Trend Test (d) | | P=0.011N | | |
| Fisher Exact Test | | | P=0.005N | P=0.025N |
| Adrenal: Cortical Adenoma | | | | |
| Overall Rates (a) | 1/49 (2%) | 2/48 (4%) | 1/44 (2%) | 3/40 (7%) |
| Adjusted Rates (b) | 2.7% | 5.7% | 5.0% | 15.8% |
| Terminal Rates (c) | 1/37 (3%) | 2/35 (6%) | 1/20 (5%) | 3/19 (16%) |
| Life Table Tests (d) | | P=0.179 | P=0.692N | P=0.235 |
| Incidental Tumor Tests (d) | | P=0.179 | P=0.692N | P=0.235 |
| Cochran-Armitage Trend Test (d) | | P=0.329 | | |
| Fisher Exact Test | | | P=0.533N | P=0.413 |
| Adrenal: Cortical Adenoma or Carcinoma | | | | |
| Overall Rates (a) | 1/49 (2%) | 3/48 (6%) | 1/44 (2%) | 3/40 (7%) |
| Adjusted Rates (b) | 2.7% | 8.6% | 5.0% | 15.8% |
| Terminal Rates (c) | 1/37 (3%) | 3/35 (9%) | 1/20 (5%) | 3/19 (16%) |
| Life Table Tests (d) | | P=0.314 | P=0.519N | P=0.363 |
| Incidental Tumor Tests (d) | | P=0.314 | P=0.519N | P=0.363 |
| Cochran-Armitage Trend Test (d) | | P=0.514 | | |
| Fisher Exact Test | | | P=0.342N | P=0.571 |

TABLE B3. ANALYSIS OF PRIMARY TUMORS IN FEMALE ACI RATS IN THE TWO-YEAR GAVAGE STUDY OF TRICHLOROETHYLENE (Continued)

| | Untreated Control | Vehicle Control | 500 mg/kg | 1,000 mg/kg |
|---|-------------------|-----------------|------------|-------------|
| Adrenal: Pheochromocytoma | | | | |
| Overall Rates (a) | 1/49 (2%) | 3/48 (6%) | 0/44 (0%) | 2/40 (5%) |
| Adjusted Rates (b) | 2.7% | 7.9% | 0.0% | 10.5% |
| Terminal Rates (c) | 1/37 (3%) | 2/35 (6%) | 0/20 (0%) | 2/19 (11%) |
| Life Table Tests (d) | | P=0.594 | P=0.227N | P=0.599 |
| Incidental Tumor Tests (d) | | P=0.582N | P=0.139N | P=0.650 |
| Cochran-Armitage Trend Test (d) | | P=0.456N | | |
| Fisher Exact Test | | | P=0.138N | P=0.587N |
| Adrenal: Pheochromocytoma or Pheochromocytoma, Malignant | | | | |
| Overall Rates (a) | 1/49 (2%) | 3/48 (6%) | 1/44 (2%) | 2/40 (5%) |
| Adjusted Rates (b) | 2.7% | 7.9% | 3.6% | 10.5% |
| Terminal Rates (c) | 1/37 (3%) | 2/35 (6%) | 0/20 (0%) | 2/19 (11%) |
| Life Table Tests (d) | | P=0.541 | P=0.492N | P=0.599 |
| Incidental Tumor Tests (d) | | P=0.578N | P=0.267N | P=0.650 |
| Cochran-Armitage Trend Test (d) | | P=0.472N | | |
| Fisher Exact Test | | | P=0.342N | P=0.587N |
| Mammary Gland: Adenoma, Papillary Adenoma, Fibroadenoma, or Adenocarcinoma | | | | |
| Overall Rates (a) | 3/49 (6%) | 3/50 (6%) | 1/50 (2%) | 1/46 (2%) |
| Adjusted Rates (b) | 8.1% | 7.5% | 4.2% | 5.3% |
| Terminal Rates (c) | 3/37 (8%) | 1/35 (3%) | 0/20 (0%) | 1/19 (5%) |
| Life Table Tests (d) | | P=0.428N | P=0.547N | P=0.555N |
| Incidental Tumor Tests (d) | | P=0.466N | P=0.545N | P=0.658N |
| Cochran-Armitage Trend Test (d) | | P=0.222N | | |
| Fisher Exact Test | | | P=0.309N | P=0.341N |
| Uterus: Endometrial Stromal Polyp | | | | |
| Overall Rates (a) | 1/48 (2%) | 4/49 (8%) | 3/48 (6%) | 2/40 (5%) |
| Adjusted Rates (b) | 2.7% | 10.7% | 15.0% | 10.5% |
| Terminal Rates (c) | 1/37 (3%) | 3/35 (9%) | 3/20 (15%) | 2/19 (11%) |
| Life Table Tests (d) | | P=0.580N | P=0.505 | P=0.641N |
| Incidental Tumor Tests (d) | | P=0.546 | P=0.434 | P=0.668 |
| Cochran-Armitage Trend Test (d) | | P=0.348N | | |
| Fisher Exact Test | | | P=0.512N | P=0.440N |
| Uterus: Endometrial Stromal Polyp or Sarcoma | | | | |
| Overall Rates (a) | 1/48 (2%) | 4/49 (8%) | 4/48 (8%) | 3/40 (7%) |
| Adjusted Rates (b) | 2.7% | 10.7% | 18.3% | 13.8% |
| Terminal Rates (c) | 1/37 (3%) | 3/35 (9%) | 3/20 (15%) | 2/19 (11%) |
| Life Table Tests (d) | | P=0.370 | P=0.321 | P=0.491 |
| Incidental Tumor Tests (d) | | P=0.428 | P=0.369 | P=0.594 |
| Cochran-Armitage Trend Test (d) | | P=0.535N | | |
| Fisher Exact Test | | | P=0.631 | P=0.613N |

(a) Number of tumor-bearing animals/number of animals examined at the site

(b) Kaplan-Meier estimated tumor incidence at the end of the study after adjusting for intercurrent mortality

(c) Observed tumor incidence at terminal kill

(d) Beneath the control incidence are the P values associated with the trend test. Beneath the dosed group incidence are the P values corresponding to pairwise comparisons between that dosed group and the vehicle controls. The life table analysis regards tumors in animals dying prior to terminal kill as being (directly or indirectly) the cause of death. The incidental tumor test regards these lesions as nonfatal. The Cochran-Armitage and Fisher exact tests compare directly the overall incidence rates. A negative trend or lower incidence in a dosed group is indicated by (N).

TABLE B4. SUMMARY OF THE INCIDENCE OF NONNEOPLASTIC LESIONS IN FEMALE ACI RATS IN THE TWO-YEAR GAVAGE STUDY OF TRICHLOROETHYLENE

| | CONTROL (UNTR) | CONTROL (VEH) | LOW DOSE | HIGH DOSE |
|--------------------------------------|----------------|---------------|----------|-----------|
| ANIMALS INITIALLY IN STUDY | 50 | 50 | 50 | 50 |
| ANIMALS MISSING | 1 | | | |
| ANIMALS NECROPSIED | 49 | 50 | 50 | 46 |
| ANIMALS EXAMINED HISTOPATHOLOGICALLY | 49 | 49 | 50 | 45 |
| INTEGUMENTARY SYSTEM | | | | |
| *Subcutaneous tissue | (49) | (50) | (50) | (46) |
| Inflammation, acute suppurative | | 1 (2%) | | |
| Inflammation, acute/chronic | | | | 1 (2%) |
| Inflammation, chronic | 1 (2%) | | | |
| RESPIRATORY SYSTEM | | | | |
| *Nasal cavity | (49) | (50) | (50) | (46) |
| Inflammation, suppurative | 1 (2%) | | | 1 (2%) |
| *Tracheal lumen | (49) | (50) | (50) | (46) |
| Hemorrhage | | 1 (2%) | 2 (4%) | 1 (2%) |
| #Trachea | (46) | (47) | (41) | (40) |
| Hemorrhage | | | 1 (2%) | |
| #Peritracheal tissue | (46) | (47) | (41) | (40) |
| Inflammation, acute | | | 1 (2%) | |
| #Lung/bronchus | (49) | (49) | (47) | (42) |
| Inflammation, acute focal | 1 (2%) | | | |
| #Lung | (49) | (49) | (47) | (42) |
| Congestion, NOS | 1 (2%) | | | 2 (5%) |
| Edema, NOS | 1 (2%) | | | |
| Hemorrhage | 2 (4%) | 6 (12%) | 2 (4%) | 1 (2%) |
| Inflammation, NOS | | 12 (24%) | 7 (15%) | 3 (7%) |
| Inflammation, focal | | 3 (6%) | 8 (17%) | 1 (2%) |
| Inflammation, multifocal | | 2 (4%) | | |
| Pneumonia, lipid | | | 1 (2%) | |
| Inflammation, acute | | | | 1 (2%) |
| Inflammation, acute focal | | | 1 (2%) | |
| Abscess, NOS | | 1 (2%) | | |
| Inflammation, chronic | | | 1 (2%) | |
| Inflammation, chronic focal | | 1 (2%) | 5 (11%) | 1 (2%) |
| Inflammation, granulomatous | | | 1 (2%) | 2 (5%) |
| Inflammation granulomatous focal | 1 (2%) | 2 (4%) | 2 (4%) | |
| Granuloma, foreign body | | | 1 (2%) | |
| Parasitism | 1 (2%) | | | |
| Foreign material, NOS | | 7 (14%) | 3 (6%) | |
| Pigmentation, NOS | | 1 (2%) | | 2 (5%) |
| Hemosiderosis | | | 1 (2%) | |
| Alveolar macrophages | | | 1 (2%) | |
| #Lung/alveoli | (49) | (49) | (47) | (42) |
| Hemorrhage | | 1 (2%) | | |
| HEMATOPOIETIC SYSTEM | | | | |
| #Spleen | (48) | (49) | (46) | (42) |
| Hemosiderosis | 3 (6%) | 3 (6%) | | 1 (2%) |
| Depletion, lymphoid | | 1 (2%) | | |
| Hematopoiesis | 1 (2%) | | 2 (4%) | 2 (5%) |
| #Splenic capsule | (48) | (49) | (46) | (42) |
| Hemorrhage | 1 (2%) | | | |
| Fibrosis | 1 (2%) | | | |
| #Lymph node | (43) | (47) | (36) | (28) |
| Necrosis, focal | | 1 (2%) | | |
| Pigmentation, NOS | | | | 1 (4%) |

TABLE B4. SUMMARY OF THE INCIDENCE OF NONNEOPLASTIC LESIONS IN FEMALE ACI RATS IN THE TWO-YEAR GAVAGE STUDY OF TRICHLOROETHYLENE (Continued)

| | CONTROL (UNTR) | CONTROL (VEH) | LOW DOSE | HIGH DOSE |
|---|----------------|---------------|----------|-----------|
| HEMATOPOIETIC SYSTEM (Continued) | | | | |
| #Mesenteric lymph node | (43) | (47) | (36) | (28) |
| Hemorrhage | 1 (2%) | | | |
| Inflammation, NOS | | 1 (2%) | | |
| #Liver | (49) | (49) | (46) | (39) |
| Hematopoiesis | | 1 (2%) | 1 (2%) | 1 (3%) |
| #Jejunum | (49) | (49) | (45) | (41) |
| Hyperplasia, lymphoid | | 1 (2%) | | |
| #Thymus | (10) | (7) | (19) | (14) |
| Cyst, NOS | 2 (20%) | | 5 (26%) | 2 (14%) |
| Multiple cysts | | | 4 (21%) | |
| Hemorrhage | 1 (10%) | | | |
| CIRCULATORY SYSTEM | | | | |
| #Heart | (49) | (48) | (49) | (42) |
| Mineralization | 1 (2%) | | | |
| Inflammation, acute focal | 1 (2%) | | | |
| Inflammation, chronic | 1 (2%) | | 1 (2%) | |
| Inflammation, chronic focal | | | 1 (2%) | |
| Fibrosis, focal | 2 (4%) | 1 (2%) | | |
| Calcification, metastatic | 1 (2%) | | | |
| DIGESTIVE SYSTEM | | | | |
| #Liver | (49) | (49) | (46) | (39) |
| Traumatic abnormality | | | | 1 (3%) |
| Congestion, NOS | | 1 (2%) | | |
| Hemorrhage | 1 (2%) | 1 (2%) | | |
| Inflammation, focal | | | 1 (2%) | |
| Necrosis, NOS | | | 1 (2%) | |
| Necrosis, focal | 3 (6%) | | | |
| Metamorphosis, fatty | 1 (2%) | 3 (6%) | 4 (9%) | 5 (13%) |
| Pigmentation, NOS | | | 1 (2%) | |
| Hemosiderosis | | 1 (2%) | | |
| Basophilic cyto change | | | | 1 (3%) |
| Clear cell change | | 1 (2%) | 5 (11%) | 3 (8%) |
| Cytologic alteration, NOS | | 1 (2%) | | |
| Hepatocytomegaly | | 3 (6%) | | 6 (15%) |
| #Liver/centrilobular | (49) | (49) | (46) | (39) |
| Necrosis, NOS | | 1 (2%) | | |
| #Bile duct | (49) | (49) | (46) | (39) |
| Inflammation, acute | | | | 1 (3%) |
| Inflammation, acute/chronic | | 1 (2%) | | |
| Hyperplasia, NOS | 2 (4%) | | | 1 (3%) |
| Hyperplasia, focal | 1 (2%) | | | |
| Hyperplasia, diffuse | | 1 (2%) | | |
| #Pancreas | (48) | (48) | (46) | (42) |
| Cytoplasmic vacuolization | | 1 (2%) | | |
| *Esophageal lumen | (49) | (50) | (50) | (46) |
| Hemorrhage | 3 (6%) | 2 (4%) | 8 (16%) | 8 (17%) |
| #Esophagus | (47) | (47) | (49) | (43) |
| Inflammation, acute | | 1 (2%) | | |
| Inflammation, acute focal | | 1 (2%) | | |
| Inflammation, granulomatous | | 1 (2%) | | |
| Granuloma, foreign body | | | 1 (2%) | |
| #Esophageal mucosa | (47) | (47) | (49) | (43) |
| Necrosis, NOS | | 1 (2%) | | |
| #Stomach | (49) | (47) | (45) | (42) |
| Mineralization | 1 (2%) | | | |
| Necrosis, focal | 1 (2%) | | | |
| Calcification, metastatic | | 1 (2%) | | |
| Hyperplasia, epithelial | 1 (2%) | | | |

TABLE B4. SUMMARY OF THE INCIDENCE OF NONNEOPLASTIC LESIONS IN FEMALE ACI RATS IN THE TWO-YEAR GAVAGE STUDY OF TRICHLOROETHYLENE (Continued)

| | CONTROL (UNTR) | CONTROL (VEH) | LOW DOSE | HIGH DOSE |
|-------------------------------------|----------------|---------------|----------|-----------|
| DIGESTIVE SYSTEM (Continued) | | | | |
| #Forestomach | (49) | (47) | (45) | (42) |
| Hyperplasia, epithelial | | 3 (6%) | 1 (2%) | |
| #Duodenum | (49) | (49) | (45) | (41) |
| Hemorrhage | | | 1 (2%) | |
| #Ileum | (49) | (49) | (45) | (41) |
| Parasitism | | | | 1 (2%) |
| #Colon | (48) | (47) | (45) | (38) |
| Hemorrhage | 1 (2%) | | | |
| Parasitism | | 1 (2%) | 1 (2%) | |
| URINARY SYSTEM | | | | |
| #Urinary bladder/cavity | (47) | (43) | (45) | (35) |
| Hemorrhage | 1 (2%) | 1 (2%) | | |
| #Kidney | (49) | (48) | (47) | (43) |
| Mineralization | 38 (78%) | 38 (79%) | 21 (45%) | 16 (37%) |
| Hydronephrosis | 3 (6%) | 3 (6%) | 4 (9%) | 4 (9%) |
| Cyst, NOS | | | 1 (2%) | |
| Hemorrhage | 1 (2%) | | 1 (2%) | |
| Inflammation, acute/chronic | | | 1 (2%) | |
| Pyelonephritis, acute/chronic | 1 (2%) | | | |
| Inflammation, chronic | | | 1 (2%) | |
| Nephropathy | 46 (94%) | 47 (98%) | 42 (89%) | 40 (93%) |
| Nephropathy, toxic | | | 21 (45%) | 19 (44%) |
| Nephrosis, hemoglobinuric | | 1 (2%) | | |
| Infarct, acute | | | 1 (2%) | |
| Calcification, NOS | | | 1 (2%) | |
| Calcification, focal | 5 (10%) | 5 (10%) | 14 (30%) | 15 (35%) |
| Calcification, metastatic | 1 (2%) | | | |
| Hemosiderosis | | 1 (2%) | | |
| Cytomegaly | | | 43 (91%) | 42 (98%) |
| Hyperplasia, epithelial | 2 (4%) | | 1 (2%) | 1 (2%) |
| Angiectasis | 6 (12%) | 6 (13%) | 1 (2%) | 1 (2%) |
| #Renal papilla | (49) | (48) | (47) | (43) |
| Mineralization | | 1 (2%) | | |
| #Kidney/tubule | (49) | (48) | (47) | (43) |
| Degeneration, NOS | | | 1 (2%) | |
| #Kidney/pelvis | (49) | (48) | (47) | (43) |
| Hemorrhage | | 1 (2%) | | |
| Hematoma, NOS | | | 1 (2%) | |
| Inflammation, suppurative | | | | 1 (2%) |
| Inflammation, acute | | | 1 (2%) | |
| Hyperplasia, epithelial | 22 (45%) | 22 (46%) | 19 (40%) | 11 (26%) |
| Angiectasis | | 1 (2%) | | 2 (5%) |
| #Urinary bladder | (47) | (43) | (45) | (35) |
| Hemorrhage | | 1 (2%) | 1 (2%) | |
| Inflammation, NOS | | | 1 (2%) | |
| Hyperplasia, epithelial | | | 2 (4%) | |
| ENDOCRINE SYSTEM | | | | |
| #Pituitary | (47) | (42) | (36) | (32) |
| Cyst, NOS | 1 (2%) | 1 (2%) | 1 (3%) | 1 (3%) |
| Pigmentation, NOS | | | | 1 (3%) |
| Hyperplasia, focal | | 2 (5%) | | |
| Angiectasis | | 2 (5%) | 1 (3%) | |

TABLE B4. SUMMARY OF THE INCIDENCE OF NONNEOPLASTIC LESIONS IN FEMALE ACI RATS IN THE TWO-YEAR GAVAGE STUDY OF TRICHLOROETHYLENE (Continued)

| | CONTROL (UNTR) | CONTROL (VEH) | LOW DOSE | HIGH DOSE |
|-------------------------------------|----------------|---------------|----------|-----------|
| ENDOCRINE SYSTEM (Continued) | | | | |
| #Adrenal | (49) | (48) | (44) | (40) |
| Necrosis, focal | 1 (2%) | 1 (2%) | | |
| Lipoidosis | | | | 1 (3%) |
| Pigmentation, NOS | 1 (2%) | | | |
| Angiectasis | | | 1 (2%) | |
| #Adrenal cortex | (49) | (48) | (44) | (40) |
| Hemorrhage | | 1 (2%) | | |
| Lipoidosis | | 1 (2%) | | |
| Cytomegaly | 1 (2%) | 1 (2%) | | 1 (3%) |
| #Adrenal medulla | (49) | (48) | (44) | (40) |
| Hyperplasia, focal | | 3 (6%) | 1 (2%) | |
| #Thyroid | (46) | (47) | (44) | (38) |
| Cyst, NOS | | | | 1 (3%) |
| Follicular cyst, NOS | 1 (2%) | | | |
| Inflammation, chronic | 1 (2%) | | | |
| Hyperplasia, C-cell | | 1 (2%) | | |
| #Parathyroid | (31) | (35) | (25) | (27) |
| Hyperplasia, NOS | | 1 (3%) | | |
| REPRODUCTIVE SYSTEM | | | | |
| *Mammary gland | (49) | (50) | (50) | (46) |
| Lactation | 1 (2%) | 2 (4%) | | |
| #Uterus | (48) | (49) | (48) | (40) |
| Hydrometra | 1 (2%) | | 1 (2%) | 2 (5%) |
| Hemorrhage | | | 1 (2%) | |
| Hematometra | 1 (2%) | | | |
| #Uterus/endometrium | (48) | (49) | (48) | (40) |
| Accessory structure | | | | 1 (3%) |
| Cyst, NOS | | 1 (2%) | | |
| Hemorrhagic cyst | | 1 (2%) | | |
| Hyperplasia, cystic | 1 (2%) | 6 (12%) | | |
| Decidual alteration, NOS | | | 1 (2%) | |
| #Ovary | (48) | (46) | (46) | (38) |
| Cyst, NOS | | 1 (2%) | | |
| Follicular cyst, NOS | 3 (6%) | | 2 (4%) | 1 (3%) |
| NERVOUS SYSTEM | | | | |
| #Brain/meninges | (49) | (48) | (49) | (42) |
| Pigmentation, NOS | | 1 (2%) | | |
| *Central canal spinal cord | (49) | (50) | (50) | (46) |
| Retention fluid | | 1 (2%) | | |
| #Brain | (49) | (48) | (49) | (42) |
| Hydrocephalus, internal | 1 (2%) | | | |
| Hemorrhage | 1 (2%) | 1 (2%) | | |
| Inflammation, acute focal | 1 (2%) | | | |
| Inflammation, chronic | | 1 (2%) | | |
| Malacia | | | 1 (2%) | |
| *Spinal cord | (49) | (50) | (50) | (46) |
| Hemorrhage | 1 (2%) | | | |
| SPECIAL SENSE ORGANS | | | | |
| *Eye/cornea | (49) | (50) | (50) | (46) |
| Inflammation, acute suppurative | 1 (2%) | | | |

TABLE B4. SUMMARY OF THE INCIDENCE OF NONNEOPLASTIC LESIONS IN FEMALE ACI RATS IN THE TWO-YEAR GAVAGE STUDY OF TRICHLOROETHYLENE (Continued)

| | CONTROL (UNTR) | CONTROL (VEH) | LOW DOSE | HIGH DOSE |
|-----------------------------------|----------------|---------------|----------|-----------|
| MUSCULOSKELETAL SYSTEM | | | | |
| *Skeletal muscle | (49) | (50) | (50) | (46) |
| Abscess, chronic | | | | 1 (2%) |
| *Muscle of neck | (49) | (50) | (50) | (46) |
| Inflammation, chronic | | | 1 (2%) | |
| BODY CAVITIES | | | | |
| *Mediastinum | (49) | (50) | (50) | (46) |
| Inflammation, NOS | | 1 (2%) | | |
| Abscess, chronic | | | | 1 (2%) |
| *Peritoneum | (49) | (50) | (50) | (46) |
| Inflammation, NOS | | | | 1 (2%) |
| *Pleura | (49) | (50) | (50) | (46) |
| Inflammation, acute focal | | | | 1 (2%) |
| ALL OTHER SYSTEMS | | | | |
| Site unknown | | | | |
| Inflammation, granulomatous | | 1 | | |
| SPECIAL MORPHOLOGY SUMMARY | | | | |
| No lesion reported | | | 2 | 1 |
| Animal missing/no necropsy | 1 | | | |
| Auto/necropsy/histo perf | | | 2 | |
| Auto/necropsy/no histo | | 1 | | 1 |
| Autolysis/no necropsy | | | | 4 |

* Number of animals receiving complete necropsy examination; all gross lesions including masses examined microscopically.

Number of animals examined microscopically at this site

APPENDIX C

SUMMARY OF LESIONS IN MALE AUGUST RATS IN THE TWO-YEAR GAVAGE STUDY OF TRICHLOROETHYLENE

| | PAGE | |
|----------|---|-----|
| TABLE C1 | SUMMARY OF THE INCIDENCE OF NEOPLASMS IN MALE AUGUST RATS IN THE TWO-YEAR GAVAGE STUDY OF TRICHLOROETHYLENE | 121 |
| TABLE C2 | INDIVIDUAL ANIMAL TUMOR PATHOLOGY OF MALE AUGUST RATS IN THE TWO-YEAR GAVAGE STUDY OF TRICHLOROETHYLENE | 124 |
| TABLE C3 | ANALYSIS OF PRIMARY TUMORS IN MALE AUGUST RATS IN THE TWO-YEAR GAVAGE STUDY OF TRICHLOROETHYLENE | 132 |
| TABLE C4 | SUMMARY OF THE INCIDENCE OF NONNEOPLASTIC LESIONS IN MALE AUGUST RATS IN THE TWO-YEAR GAVAGE STUDY OF TRICHLOROETHYLENE | 134 |

TABLE C1. SUMMARY OF THE INCIDENCE OF NEOPLASMS IN MALE AUGUST RATS IN THE TWO-YEAR GAVAGE STUDY OF TRICHLOROETHYLENE

| | CONTROL (UNTR) | CONTROL (VEH) | LOW DOSE | HIGH DOSE |
|---|----------------|---------------|----------|-----------|
| ANIMALS INITIALLY IN STUDY | 50 | 50 | 50 | 50 |
| ANIMALS NECROPSIED | 50 | 50 | 50 | 49 |
| ANIMALS EXAMINED HISTOPATHOLOGICALLY | 50 | 50 | 50 | 49 |
| INTEGUMENTARY SYSTEM | | | | |
| *Skin | (50) | (50) | (50) | (49) |
| Squamous cell carcinoma | 1 (2%) | | 1 (2%) | |
| Basal cell carcinoma | 1 (2%) | | | |
| *Subcutaneous tissue | (50) | (50) | (50) | (49) |
| Sarcoma, NOS | 1 (2%) | | 1 (2%) | 2 (4%) |
| Sarcoma, NOS, unclear primary or metastatic | | | | 1 (2%) |
| Fibroma | 1 (2%) | | | 1 (2%) |
| RESPIRATORY SYSTEM | | | | |
| #Lung | (50) | (50) | (50) | (49) |
| Squamous cell carcinoma | | | 1 (2%) | |
| Alveolar/bronchiolar adenoma | 1 (2%) | | | |
| Sarcoma, NOS, metastatic | | 1 (2%) | | |
| HEMATOPOIETIC SYSTEM | | | | |
| *Multiple organs | (50) | (50) | (50) | (49) |
| Malignant lymphoma, NOS | | | 1 (2%) | 2 (4%) |
| Malignant lymphoma, histiocytic type | 1 (2%) | | | |
| Leukemia, NOS | 2 (4%) | | | |
| Monocytic leukemia | | 1 (2%) | | |
| CIRCULATORY SYSTEM | | | | |
| *Skeletal muscle | (50) | (50) | (50) | (49) |
| Hemangioma | | | 1 (2%) | |
| #Heart | (50) | (50) | (50) | (49) |
| Carcinoma, NOS | 1 (2%) | | | |
| Carcinoma, NOS, metastatic | 1 (2%) | | | |
| DIGESTIVE SYSTEM | | | | |
| #Salivary gland | (47) | (50) | (49) | (46) |
| Sarcoma, NOS, unclear primary or metastatic | | | | 1 (2%) |
| #Liver | (50) | (50) | (50) | (48) |
| Bile duct adenoma | 1 (2%) | | | |
| Neoplastic nodule | | | 1 (2%) | 1 (2%) |
| #Stomach | (50) | (50) | (50) | (49) |
| Papilloma, NOS | | | | 1 (2%) |
| #Forestomach | (50) | (50) | (50) | (49) |
| Squamous cell papilloma | | 1 (2%) | | |
| URINARY SYSTEM | | | | |
| #Kidney | (50) | (50) | (50) | (49) |
| Squamous cell carcinoma | | | 1 (2%) | |
| Tubular cell adenoma | | | 1 (2%) | 1 (2%) |
| Tubular cell adenocarcinoma | | | 1 (2%) | |
| #Urinary bladder | (50) | (50) | (47) | (44) |
| Transitional cell papilloma | 1 (2%) | 1 (2%) | | |
| Transitional cell carcinoma | | 1 (2%) | | |

TABLE C1. SUMMARY OF THE INCIDENCE OF NEOPLASMS IN MALE AUGUST RATS IN THE TWO-YEAR GAVAGE STUDY OF TRICHLOROETHYLENE (Continued)

| | CONTROL (UNTR) | CONTROL (VEH) | LOW DOSE | HIGH DOSE |
|--------------------------------|----------------|---------------|----------|-----------|
| ENDOCRINE SYSTEM | | | | |
| #Pituitary | (46) | (44) | (41) | (41) |
| Carcinoma, NOS | | 3 (7%) | | |
| Adenoma, NOS | 15 (33%) | 23 (52%) | 6 (15%) | 4 (10%) |
| Chromophobe adenoma | 1 (2%) | | 1 (2%) | |
| #Adrenal | (49) | (50) | (49) | (47) |
| Cortical adenoma | 6 (12%) | 5 (10%) | 3 (6%) | 5 (11%) |
| Pheochromocytoma | 15 (31%) | 10 (20%) | 5 (10%) | 2 (4%) |
| Pheochromocytoma, malignant | | | 1 (2%) | 1 (2%) |
| #Adrenal medulla | (49) | (50) | (49) | (47) |
| Pheochromocytoma | 1 (2%) | | | |
| Neurilemoma | | 1 (2%) | | |
| #Thyroid | (49) | (46) | (44) | (45) |
| C-cell adenoma | 3 (6%) | 7 (15%) | | |
| C-cell carcinoma | | 1 (2%) | | |
| #Parathyroid | (40) | (26) | (32) | (24) |
| Adenoma, NOS | | | 1 (3%) | |
| #Pancreatic islets | (49) | (50) | (49) | (48) |
| Islet cell adenoma | 5 (10%) | 5 (10%) | | 1 (2%) |
| REPRODUCTIVE SYSTEM | | | | |
| *Mammary gland | (50) | (50) | (50) | (49) |
| Adenoma, NOS | | | | 1 (2%) |
| Fibroma | | | 1 (2%) | |
| Fibroadenoma | 1 (2%) | 1 (2%) | | |
| *Preputial gland | (50) | (50) | (50) | (49) |
| Carcinoma, NOS | | 1 (2%) | | 1 (2%) |
| Adenoma, NOS | 1 (2%) | | | |
| #Testis | (50) | (50) | (50) | (49) |
| Interstitial cell tumor | 36 (72%) | 34 (68%) | 30 (60%) | 26 (53%) |
| NERVOUS SYSTEM | | | | |
| #Brain | (50) | (50) | (50) | (49) |
| Granular cell tumor, NOS | 1 (2%) | 1 (2%) | | 1 (2%) |
| Granular cell tumor, malignant | | | 1 (2%) | |
| Astrocytoma | | 1 (2%) | | |
| Oligodendroglioma | 1 (2%) | | | |
| *Spinal cord | (50) | (50) | (50) | (49) |
| Astrocytoma | 1 (2%) | | | |
| SPECIAL SENSE ORGANS | | | | |
| None | | | | |
| MUSCULOSKELETAL SYSTEM | | | | |
| None | | | | |

TABLE C1. SUMMARY OF THE INCIDENCE OF NEOPLASMS IN MALE AUGUST RATS IN THE TWO-YEAR GAVAGE STUDY OF TRICHLOROETHYLENE (Continued)

| | CONTROL (UNTR) | CONTROL (VEH) | LOW DOSE | HIGH DOSE |
|--|----------------|---------------|----------|-----------|
| BODY CAVITIES | | | | |
| *Mediastinum | (50) | (50) | (50) | (49) |
| Sarcoma, NOS, unclear primary or metast | | 1(2%) | | |
| ALL OTHER SYSTEMS | | | | |
| None | | | | |
| ANIMAL DISPOSITION SUMMARY | | | | |
| Animals initially in study | 50 | 50 | 50 | 50 |
| Natural death | 16 | 10 | 17 | 17 |
| Moribund sacrifice | 9 | 13 | 8 | 6 |
| Terminal sacrifice | 24 | 21 | 13 | 15 |
| Dosing accident | 1 | 2 | 3 | 7 |
| Accidentally killed, NOS | | 4 | 9 | 4 |
| Animal missexed | | | | 1 |
| TUMOR SUMMARY | | | | |
| Total animals with primary tumors** | 43 | 45 | 32 | 27 |
| Total primary tumors | 98 | 98 | 58 | 52 |
| Total animals with benign tumors | 40 | 43 | 32 | 27 |
| Total benign tumors | 88 | 88 | 49 | 42 |
| Total animals with malignant tumors | 8 | 8 | 7 | 5 |
| Total malignant tumors | 9 | 8 | 8 | 6 |
| Total animals with secondary tumors## | 1 | 1 | | |
| Total secondary tumors | 1 | 1 | | |
| Total animals with tumors uncertain--benign or malignant | 1 | 1 | 1 | 2 |
| Total uncertain tumors | 1 | 1 | 1 | 2 |
| Total animals with tumors uncertain--primary or metastatic | | 1 | | 1 |
| Total uncertain tumors | | 1 | | 2 |

* Number of animals receiving complete necropsy examination; all gross lesions including masses examined microscopically.

** Primary tumors: all tumors except secondary tumors

Number of animals examined microscopically at this site

Secondary tumors: metastatic tumors or tumors invasive into an adjacent organ

TABLE C2. INDIVIDUAL ANIMAL TUMOR PATHOLOGY OF MALE AUGUST RATS: UNTREATED CONTROL (Continued)

| ANIMAL NUMBER | 2 6 1 | 2 6 8 | 2 7 0 | 2 7 3 | 2 7 4 | 2 7 6 | 2 7 7 | 2 8 1 | 2 8 3 | 2 8 5 | 2 8 7 | 2 9 5 | 2 9 9 | 3 0 0 | 3 0 0 | 3 0 0 | 3 0 0 | 3 0 0 | 3 0 1 | 3 0 1 | 3 0 1 | 3 0 2 | 3 0 5 | 3 0 6 | 3 0 7 | 3 0 9 | 3 1 0 | 3 1 1 | 3 1 2 | 3 1 3 | 3 1 3 | 3 1 4 | 3 1 8 | 3 1 8 | 3 1 0 | TOTAL TISSUES TUMORS |
|--------------------------------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|----------------------|
| WEEKS ON STUDY | 0 6 | 1 4 | 1 4 | 0 9 | 0 3 | 0 6 | 1 4 | 1 4 | 1 4 | 1 4 | 1 4 | 0 7 | 1 4 | 1 3 | 1 4 | 0 9 | 1 0 | 0 6 | 0 7 | 0 9 | 1 0 | 0 4 | 0 4 | 0 5 | 0 7 | 0 9 | 0 7 | 0 6 | 1 0 | 0 4 | 0 6 | 1 0 | 0 8 | 0 3 | | |
| INTEGUMENTARY SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Skin | | | | | | | | | | | | | | | | | | | | | | | | | | | | | *50 | | | | | | | |
| Squamous cell carcinoma | | | | | | | | | | | | | | | | | | | | | | | | | | | | | 1 | | | | | | | |
| Basal cell carcinoma | | | | | | | | | | | | | | | | | | | | | | | | | | | | | 1 | | | | | | | |
| Subcutaneous tissue | | | | | | | | | | | | | | | | | | | | | | | | | | | | | *50 | | | | | | | |
| Sarcoma, NOS | | | | | | | | | | | | | | | | | | | | | | | | | | | | | 1 | | | | | | | |
| Fibroma | | | | | | | | | | | | | | | | | | | | | | | | | | | | | 1 | | | | | | | |
| RESPIRATORY SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Lungs and bronchi | | | | | | | | | | | | | | | | | | | | | | | | | | | | | 50 | | | | | | | |
| Alveolar/bronchiolar adenoma | | | | | | | | | | | | | | | | | | | | | | | | | | | | | 1 | | | | | | | |
| Trachea | | | | | | | | | | | | | | | | | | | | | | | | | | | | | 49 | | | | | | | |
| HEMATOPOIETIC SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Bone marrow | | | | | | | | | | | | | | | | | | | | | | | | | | | | | 50 | | | | | | | |
| Spleen | | | | | | | | | | | | | | | | | | | | | | | | | | | | | 49 | | | | | | | |
| Lymph nodes | | | | | | | | | | | | | | | | | | | | | | | | | | | | | 48 | | | | | | | |
| Thymus | | | | | | | | | | | | | | | | | | | | | | | | | | | | | 6 | | | | | | | |
| CIRCULATORY SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Heart | | | | | | | | | | | | | | | | | | | | | | | | | | | | | 50 | | | | | | | |
| Carcinoma, NOS | | | | | | | | | | | | | | | | | | | | | | | | | | | | | 1 | | | | | | | |
| Carcinoma, NOS, metastatic | | | | | | | | | | | | | | | | | | | | | | | | | | | | | 1 | | | | | | | |
| DIGESTIVE SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Salivary gland | | | | | | | | | | | | | | | | | | | | | | | | | | | | | 47 | | | | | | | |
| Liver | | | | | | | | | | | | | | | | | | | | | | | | | | | | | 50 | | | | | | | |
| Bile duct adenoma | | | | | | | | | | | | | | | | | | | | | | | | | | | | | 1 | | | | | | | |
| Bile duct | | | | | | | | | | | | | | | | | | | | | | | | | | | | | 50 | | | | | | | |
| Gallbladder & common bile duct | | | | | | | | | | | | | | | | | | | | | | | | | | | | | *50 | | | | | | | |
| Pancreas | | | | | | | | | | | | | | | | | | | | | | | | | | | | | 49 | | | | | | | |
| Esophagus | | | | | | | | | | | | | | | | | | | | | | | | | | | | | 50 | | | | | | | |
| Stomach | | | | | | | | | | | | | | | | | | | | | | | | | | | | | 50 | | | | | | | |
| Small intestine | | | | | | | | | | | | | | | | | | | | | | | | | | | | | 49 | | | | | | | |
| Large intestine | | | | | | | | | | | | | | | | | | | | | | | | | | | | | 49 | | | | | | | |
| URINARY SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Kidney | | | | | | | | | | | | | | | | | | | | | | | | | | | | | 50 | | | | | | | |
| Urinary bladder | | | | | | | | | | | | | | | | | | | | | | | | | | | | | 50 | | | | | | | |
| Transitional cell papilloma | | | | | | | | | | | | | | | | | | | | | | | | | | | | | 1 | | | | | | | |
| ENDOCRINE SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Pituitary | | | | | | | | | | | | | | | | | | | | | | | | | | | | | 46 | | | | | | | |
| Adenoma, NOS | | | | | | | | | | | | | | | | | | | | | | | | | | | | | 15 | | | | | | | |
| Chromophobe adenoma | | | | | | | | | | | | | | | | | | | | | | | | | | | | | 1 | | | | | | | |
| Adrenal | | | | | | | | | | | | | | | | | | | | | | | | | | | | | 49 | | | | | | | |
| Cortical adenoma | | | | | | | | | | | | | | | | | | | | | | | | | | | | | 6 | | | | | | | |
| Pheochromocytoma | | | | | | | | | | | | | | | | | | | | | | | | | | | | | 16 | | | | | | | |
| Thyroid | | | | | | | | | | | | | | | | | | | | | | | | | | | | | 49 | | | | | | | |
| C-cell adenoma | | | | | | | | | | | | | | | | | | | | | | | | | | | | | 3 | | | | | | | |
| Parathyroid | | | | | | | | | | | | | | | | | | | | | | | | | | | | | 40 | | | | | | | |
| Pancreatic islets | | | | | | | | | | | | | | | | | | | | | | | | | | | | | 49 | | | | | | | |
| Islet cell adenoma | | | | | | | | | | | | | | | | | | | | | | | | | | | | | 5 | | | | | | | |
| REPRODUCTIVE SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Mammary gland | | | | | | | | | | | | | | | | | | | | | | | | | | | | | *50 | | | | | | | |
| Fibroadenoma | | | | | | | | | | | | | | | | | | | | | | | | | | | | | 1 | | | | | | | |
| Testis | | | | | | | | | | | | | | | | | | | | | | | | | | | | | 50 | | | | | | | |
| Interstitial cell tumor | | | | | | | | | | | | | | | | | | | | | | | | | | | | | 36 | | | | | | | |
| Prostate | | | | | | | | | | | | | | | | | | | | | | | | | | | | | 50 | | | | | | | |
| Preputial/clitoral gland | | | | | | | | | | | | | | | | | | | | | | | | | | | | | *50 | | | | | | | |
| Adenoma, NOS | | | | | | | | | | | | | | | | | | | | | | | | | | | | | 1 | | | | | | | |
| NERVOUS SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Brain | | | | | | | | | | | | | | | | | | | | | | | | | | | | | 50 | | | | | | | |
| Granular cell tumor, NOS | | | | | | | | | | | | | | | | | | | | | | | | | | | | | 1 | | | | | | | |
| Oligodendroglioma | | | | | | | | | | | | | | | | | | | | | | | | | | | | | 1 | | | | | | | |
| Spinal cord | | | | | | | | | | | | | | | | | | | | | | | | | | | | | *50 | | | | | | | |
| Astrocytoma | | | | | | | | | | | | | | | | | | | | | | | | | | | | | 1 | | | | | | | |
| ALL OTHER SYSTEMS | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Multiple organs, NOS | | | | | | | | | | | | | | | | | | | | | | | | | | | | | *50 | | | | | | | |
| Malignant lymphoma, histiocytic type | | | | | | | | | | | | | | | | | | | | | | | | | | | | | 1 | | | | | | | |
| Leukemia, NOS | | | | | | | | | | | | | | | | | | | | | | | | | | | | | 2 | | | | | | | |

* Animals necropsied

TABLE C2. INDIVIDUAL ANIMAL TUMOR PATHOLOGY OF MALE AUGUST RATS IN THE TWO-YEAR GAVAGE STUDY OF TRICHLOROETHYLENE: VEHICLE CONTROL

| ANIMAL NUMBER | 1 2 | 1 4 | 1 4 | 1 8 | 1 0 | 1 8 | 1 6 | 1 7 | 1 7 | 1 7 | 1 7 | 1 7 | 1 8 | 1 8 | 1 9 | 2 0 | 2 0 | 2 0 | 2 1 | 2 1 | 2 1 | 2 1 | 2 2 | | |
|---|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--|--|
| WEEKS ON STUDY | 0 7 | 0 3 | 1 4 | 0 9 | 1 3 | 1 4 | 1 4 | 1 4 | 0 1 | 0 3 | 0 2 | 1 4 | 1 4 | 1 4 | 1 4 | 1 4 | 1 4 | 0 6 | 0 0 | 1 0 | 0 8 | 0 2 | 0 8 | 0 9 | 0 7 | 0 6 | 0 9 | 0 6 | 0 4 | | | |
| RESPIRATORY SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Lungs and bronchi | + | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Sarcoma, NOS, metastatic | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Trachea | + | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| HEMATOPOIETIC SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Bone marrow | + | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Spleen | + | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Lymph nodes | + | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Thymus | - | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| CIRCULATORY SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Heart | + | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| DIGESTIVE SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Salivary gland | + | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Liver | + | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Bile duct | + | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Gallbladder & common bile duct | N | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Pancreas | + | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Esophagus | + | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Stomach | + | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Squamous cell papilloma | X | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Small intestine | + | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Large intestine | + | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| URINARY SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Kidney | + | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Urinary bladder | + | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Transitional cell papilloma | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Transitional cell carcinoma | X | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| ENDOCRINE SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Pituitary | + | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Carcinoma, NOS | + | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Adenoma, NOS | X | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Adrenal | + | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Cortical adenoma | + | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Pheochromocytoma | X | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Neurilemoma | + | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Thyroid | + | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| C-cell adenoma | + | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| C-cell carcinoma | X | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Parathyroid | + | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Pancreatic islets | + | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Islet cell adenoma | X | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| REPRODUCTIVE SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Mammary gland | N | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Fibroadenoma | + | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Testis | + | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Interstitial cell tumor | X | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Prostate | + | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Preputial/clitoral gland | N | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Carcinoma, NOS | N | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| NERVOUS SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Brain | + | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Granular cell tumor, NOS | + | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Astrocytoma | X | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| BODY CAVITIES | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Mediastinum | N | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Sarcoma, NOS, unclear primary or metastatic | X | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| ALL OTHER SYSTEMS | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Multiple organs, NOS | N | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Monocytic leukemia | X | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |

TABLE C2. INDIVIDUAL ANIMAL TUMOR PATHOLOGY OF MALE AUGUST RATS: VEHICLE CONTROL (Continued)

| ANIMAL NUMBER | 222 | 223 | 224 | 225 | 226 | 227 | 228 | 229 | 230 | 231 | 232 | 233 | 234 | 235 | 236 | 237 | 238 | 239 | 240 | 241 | 242 | 243 | 244 | 245 | 246 | 247 | 248 | 249 | 250 | TOTAL ISSUES | |
|--|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|--------------|---|
| WEEKS ON STUDY | 50 | 99 | 68 | 65 | 98 | 83 | 44 | 95 | 07 | 57 | 78 | 66 | 66 | 11 | 11 | 00 | 00 | 11 | 11 | 11 | 11 | 11 | 11 | 11 | 11 | 11 | 11 | 11 | 50 | | |
| RESPIRATORY SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Lungs and bronchi | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | 50 | |
| Sarcoma, NOS, metastatic | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | 1 | |
| Trachea | - | + | + | + | + | + | + | + | + | + | + | + | + | - | + | + | + | + | + | + | + | + | + | + | + | + | + | + | 48 | | |
| HEMATOPOIETIC SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Bone marrow | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | 50 | |
| Spleen | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | 50 | |
| Lymph nodes | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | 45 | |
| Thymus | - | - | - | - | - | - | - | - | - | - | - | - | - | + | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | 1 | |
| CIRCULATORY SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Heart | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | 50 | |
| DIGESTIVE SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Salivary gland | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | 50 | |
| Liver | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | 50 | |
| Bile duct | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | 50 | |
| Gallbladder & common bile duct | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | *50 | |
| Pancreas | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | 50 | |
| Esophagus | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | 48 | |
| Stomach | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | 50 | |
| Squamous cell papilloma | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | 1 | |
| Small intestine | + | + | + | - | + | + | + | + | + | + | + | + | + | - | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | 48 | |
| Large intestine | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | 50 | |
| URINARY SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Kidney | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | 50 | |
| Urinary bladder | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | 50 | |
| Transitional cell papilloma | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | 1 | |
| Transitional cell carcinoma | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | 1 | |
| ENDOCRINE SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Pituitary | + | + | - | - | + | + | + | + | + | + | + | + | + | - | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | 44 | |
| Carcinoma, NOS | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | 3 | |
| Adenoma, NOS | X | + | + | + | X | | | X | | | | | X | X | X | | | | X | | X | | X | | X | | X | | 23 | | |
| Adrenal | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | 50 | | |
| Cortical adenoma | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | 5 | |
| Pheochromocytoma | | | | | | X | | | | | | | | | | | | | | | | | | | | | | | | 10 | |
| Neurilemoma | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | 1 | |
| Thyroid | + | + | + | + | + | + | + | + | + | + | + | + | + | - | + | + | + | + | + | + | + | + | + | + | + | + | + | + | 46 | | |
| C-cell adenoma | | | | | | X | | | | | | | | | | | | | | | | | | | | | | | | 7 | |
| C-cell carcinoma | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | 1 | |
| Parathyroid | - | + | - | - | + | + | + | + | + | - | - | + | - | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | 26 | |
| Pancreatic islets | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | 50 | |
| Islet cell adenoma | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | 5 | |
| REPRODUCTIVE SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Mammary gland | N | N | N | N | N | + | + | N | + | N | + | N | N | + | + | N | + | + | + | + | + | + | + | + | + | + | + | + | + | *50 | |
| Fibroadenoma | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | 1 |
| Testis | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | 50 | |
| Interstitial cell tumor | | X | | | X | | | X | X | | | | X | | X | X | X | X | X | X | X | X | X | X | X | X | X | X | X | 34 | |
| Prostate | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | 50 | |
| Preputial/clitoral gland | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | *50 | |
| Carcinoma, NOS | | | | | | X | | | | | | | | | | | | | | | | | | | | | | | | 1 | |
| NERVOUS SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Brain | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | 50 | |
| Granular cell tumor, NOS | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | 1 |
| Astrocytoma | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | 1 |
| BODY CAVITIES | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Mediastinum | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | *50 | |
| Sarcoma, NOS, unclear prim or metastat | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | 1 |
| ALL OTHER SYSTEMS | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Multiple organs, NOS | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | *50 | |
| Monocytic leukemia | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | 1 |

* Animals necropsied

TABLE C2. INDIVIDUAL ANIMAL TUMOR PATHOLOGY OF MALE AUGUST RATS IN THE TWO-YEAR GAVAGE STUDY OF TRICHLOROETHYLENE: LOW DOSE

| ANIMAL NUMBER | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 2 | 2 | 2 | 2 | 2 | 2 | 2 | 2 | 2 | 2 |
|--------------------------------|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| WEEKS ON STUDY | 4 | 4 | 4 | 4 | 4 | 5 | 6 | 6 | 7 | 7 | 7 | 7 | 8 | 8 | 8 | 8 | 8 | 9 | 0 | 0 | 0 | 0 | 0 | 1 | 2 | 3 | 3 | 3 |
| | 1 | 0 | 0 | 0 | 0 | 1 | 0 | 0 | 1 | 0 | 0 | 0 | 0 | 1 | 0 | 0 | 1 | 1 | 0 | 0 | 1 | 0 | 0 | 1 | 0 | 0 | 1 | 0 |
| | 4 | 9 | 6 | 8 | 3 | 4 | 5 | 0 | 0 | 1 | 4 | 2 | 2 | 4 | 7 | 0 | 4 | 4 | 4 | 4 | 3 | 4 | 4 | 2 | 8 | 4 | 0 | 6 |
| INTEGUMENTARY SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Skin | + | + | + | + | + | + | + | N | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + |
| Squamous cell carcinoma | | | | | | | | | | | | | | | | | | | | | | | | | | | X | + |
| Subcutaneous tissue | + | + | + | + | + | + | + | N | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + |
| Sarcoma, NOS | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| RESPIRATORY SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Lungs and bronchi | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + |
| Squamous cell carcinoma | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Trachea | + | + | - | + | + | + | - | - | + | + | + | + | - | + | + | - | + | + | + | + | + | + | + | + | + | + | + | + |
| HEMATOPOIETIC SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Bone marrow | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + |
| Spleen | + | + | + | + | + | + | + | + | + | + | + | + | - | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + |
| Lymph nodes | + | + | + | + | - | + | + | + | + | + | + | - | + | + | + | + | + | + | + | - | + | + | + | + | + | + | + | - |
| Thymus | - | - | + | + | - | - | - | + | - | - | + | - | + | - | - | + | - | - | + | - | - | - | - | - | - | - | - | - |
| CIRCULATORY SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Heart | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + |
| DIGESTIVE SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Salivary gland | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + |
| Liver | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + |
| Neoplastic nodule | X | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Bile duct | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + |
| Gallbladder & common bile duct | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N |
| Pancreas | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + |
| Esophagus | + | + | - | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + |
| Stomach | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + |
| Small intestine | + | + | + | + | - | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + |
| Large intestine | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + |
| URINARY SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Kidney | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + |
| Squamous cell carcinoma | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Tubular cell adenoma | | | | | | | | | | | | | X | | | | | | | | | | | | | | | |
| Tubular cell adenocarcinoma | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Urinary bladder | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | X | + | + | + | + | + | + | + | + | + |
| ENDOCRINE SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Pituitary | - | + | + | A | + | + | - | - | + | + | + | + | + | - | - | - | + | + | + | + | + | + | + | + | + | + | + | - |
| Adenoma, NOS | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Chromophobe adenoma | | | | | | | | | | | | X | | | | | | | | | | | | | | | | X |
| Adrenal | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + |
| Cortical adenoma | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Pheochromocytoma | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Pheochromocytoma, malignant | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Thyroid | + | + | - | + | + | + | - | - | + | + | + | - | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + |
| Parathyroid | + | + | - | + | + | + | - | - | + | + | + | - | - | + | + | - | + | + | - | + | + | - | + | + | - | - | + | + |
| Adenoma, NOS | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| REPRODUCTIVE SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Mammary gland | + | + | + | N | N | + | N | N | + | + | N | + | N | + | N | N | + | + | N | + | + | N | + | + | + | N | + | + |
| Fibroma | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Testis | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + |
| Interstitial cell tumor | X | X | X | | X | | | | X | | | X | X | | X | X | | X | X | | X | X | | X | X | | X | X |
| Prostate | + | + | + | A | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + |
| NERVOUS SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Brain | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + |
| Granular cell tumor, malignant | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| MUSCULOSKELETAL SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Muscle | N | + | N | + | N | N | + | + | N | N | N | N | + | N | N | + | N | N | N | N | N | N | N | N | N | N | N | N |
| Hemangioma | | X | | | | | | | | | | | | | | | | | | | | | | | | | | |
| ALL OTHER SYSTEMS | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Multiple organs, NOS | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N |
| Malignant lymphoma, NOS | | | | | | | | | | | | | | | | | | | | | | | | | | | | |

TABLE C2. INDIVIDUAL ANIMAL TUMOR PATHOLOGY OF MALE AUGUST RATS: LOW DOSE
(Continued)

| ANIMAL NUMBER | 21 41 | 22 52 | 23 57 | 24 58 | 25 62 | 26 66 | 27 67 | 28 71 | 29 72 | 30 75 | 31 82 | 32 84 | 33 90 | 34 91 | 35 97 | 36 98 | 37 99 | 38 00 | 39 01 | 40 06 | 41 08 | 42 09 | 43 12 | 44 17 | 45 20 | 46 22 | 47 24 | 48 25 | 49 26 | 50 30 | TOTAL TISSUES TUMORS | |
|--------------------------------|----------|----------|----------|----------|----------|----------|----------|----------|----------|----------|----------|----------|----------|----------|----------|----------|----------|----------|----------|----------|----------|----------|----------|----------|----------|----------|----------|----------|----------|----------|----------------------|-----|
| WEEKS ON STUDY | 07 | 08 | 08 | 09 | 10 | 10 | 11 | 11 | 11 | 12 | 12 | 13 | 13 | 14 | 14 | 15 | 15 | 16 | 16 | 16 | 18 | 18 | 19 | 19 | 20 | 20 | 21 | 21 | 22 | 22 | 23 | 23 |
| INTEGUMENTARY SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Skin | + | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | *50 |
| Squamous cell carcinoma | + | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | 1 |
| Subcutaneous tissue | + | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | *50 |
| Sarcoma, NOS | X | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | 1 |
| RESPIRATORY SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Lungs and bronchi | + | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | 50 |
| Squamous cell carcinoma | X | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | 1 |
| Trachea | + | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | 42 |
| HEMATOPOIETIC SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Bone marrow | + | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | 49 |
| Spleen | + | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | 48 |
| Lymph nodes | + | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | 43 |
| Thymus | - | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | 9 |
| CIRCULATORY SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Heart | + | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | 50 |
| DIGESTIVE SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Salivary gland | + | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | 49 |
| Liver | + | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | 50 |
| Neoplastic nodule | + | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | 1 |
| Bile duct | + | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | 50 |
| Gallbladder & common bile duct | N | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | *50 |
| Pancreas | + | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | 49 |
| Esophagus | + | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | 48 |
| Stomach | + | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | 50 |
| Small intestine | + | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | 48 |
| Large intestine | + | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | 50 |
| URINARY SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Kidney | + | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | 50 |
| Squamous cell carcinoma | X | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | 1 |
| Tubular cell adenoma | + | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | 1 |
| Tubular cell adenocarcinoma | + | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | 1 |
| Urinary bladder | + | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | 47 |
| ENDOCRINE SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Pituitary | + | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | 41 |
| Adenoma, NOS | - | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | 6 |
| Chromophobe adenoma | + | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | 1 |
| Adrenal | + | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | 49 |
| Cortical adenoma | + | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | 3 |
| Pheochromocytoma | X | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | 5 |
| Pheochromocytoma, malignant | X | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | 1 |
| Thyroid | + | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | 44 |
| Parathyroid | + | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | 32 |
| Adenoma, NOS | X | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | 1 |
| REPRODUCTIVE SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Mammary gland | N | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | *50 |
| Fibroma | + | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | 1 |
| Testis | + | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | 50 |
| Interstitial cell tumor | X | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | 30 |
| Prostate | + | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | 48 |
| NERVOUS SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Brain | + | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | 50 |
| Granular cell tumor, malignant | X | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | 1 |
| MUSCULOSKELETAL SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Muscle | + | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | *50 |
| Hemangioma | N | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | 1 |
| ALL OTHER SYSTEMS | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Multiple organs, NOS | N | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | *50 |
| Malignant lymphoma, NOS | N | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | 1 |

* Animals necropsied

TABLE C2. INDIVIDUAL ANIMAL TUMOR PATHOLOGY OF MALE AUGUST RATS: HIGH DOSE (Continued)

| ANIMAL NUMBER | 2030 | 2031 | 2032 | 2033 | 2034 | 2035 | 2036 | 2037 | 2038 | 2039 | 2040 | 2041 | 2042 | 2043 | 2044 | 2045 | 2046 | 2047 | 2048 | 2049 | TOTAL TISSUES TUMORS | | | | | | |
|---------------------------------------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|----------------------|---|---|-----|-----|-----|-----|
| WEEKS ON STUDY | 4 | 4 | 3 | 7 | 9 | 4 | 4 | 9 | 9 | 4 | 7 | 9 | 8 | 5 | 4 | 8 | 4 | 5 | 0 | 1 | 4 | 2 | 4 | 4 | | | |
| INTEGUMENTARY SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Subcutaneous tissue | | | | | | | | | | | | | | | | | | | | | | | | *49 | | | |
| Sarcoma, NOS | | | | | | | | | | | | | | | | | | | | | | | | | 2 | | |
| Sarcoma, NOS, unclear prim or metasta | | | | | | | | | | | | | | | | | | | | | | | | | 1 | | |
| Fibroma | | | | | | | | | | | | | | | | | | | | | | | | | 1 | | |
| RESPIRATORY SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Lungs and bronchi | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | | 49 | | |
| Trachea | + | + | - | - | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | 42 | | |
| HEMATOPOIETIC SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Bone marrow | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | | 49 | | |
| Spleen | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | | 49 | | |
| Lymph nodes | + | + | - | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | | 33 | | |
| Thymus | - | - | + | + | - | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | | 8 | | |
| CIRCULATORY SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Heart | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | | 49 | | |
| DIGESTIVE SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Salivary gland | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | | 46 | | |
| Sarcoma, NOS, unclear prim or metasta | | | | | | | | | | | | | | | | | | | | | | | | | 1 | | |
| Liver | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | | 48 | | |
| Neoplastic nodule | | | | | | | | | | | | | | | | | | | | | | | | | 1 | | |
| Bile duct | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | | 48 | | |
| Gallbladder & common bile duct | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | | *49 | | |
| Pancreas | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | | 48 | | |
| Esophagus | + | + | + | - | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | | 47 | | |
| Stomach | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | | 49 | | |
| Papilloma, NOS | | | | | | | | | | | | | | | | | | | | | | | | | 1 | | |
| Small intestine | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | | 44 | | |
| Large intestine | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | | 44 | | |
| URINARY SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Kidney | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | | 49 | | |
| Tubular cell adenoma | | | | | | | | | | | | | | | | | | | | | | | | | X | | |
| Urinary bladder | + | - | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | | 44 | | |
| ENDOCRINE SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Pituitary | + | + | - | - | - | + | - | + | + | + | - | + | + | + | - | - | + | + | + | + | + | + | + | | 41 | | |
| Adenoma, NOS | X | | | | | X | | | | | | | | | | | | | | | | | | | X | | |
| Adrenal | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | | 47 | | |
| Cortical adenoma | | | | | | | | | | | | | | | | | | | | | | | | | 5 | | |
| Phochromocytoma | | | | | | | | | | | | | | | | | | | | | | | | | 2 | | |
| Phochromocytoma, malignant | | | | | | | | | | | | | | | | | | | | | | | | | 1 | | |
| Thyroid | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | | 45 | | |
| Parathyroid | + | + | + | - | + | + | - | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | | 24 | | |
| Pancreatic islets | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | | 48 | | |
| Islet cell adenoma | | | | | | | X | | | | | | | | | | | | | | | | | | 1 | | |
| REPRODUCTIVE SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Mammary gland | N | N | N | N | + | + | N | + | N | N | N | + | N | + | N | N | N | N | N | + | N | N | + | + | N | | *49 |
| Adenoma, NOS | | | | | | | | | | | | | | | | | | | | | | | | | | | 1 |
| Testis | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | | 49 | |
| Interstitial cell tumor | X | X | | | | X | X | X | X | X | X | | | | X | X | | | X | | X | X | X | X | | 28 | |
| Prostate | + | - | + | - | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | | 48 | |
| Preputial/clitoral gland | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | | *49 | |
| Carcinoma, NOS | X | | | | | | | | | | | | | | | | | | | | | | | | | 1 | |
| NERVOUS SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Brain | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | | 49 | |
| Granular cell tumor, NOS | | | | | | | | | | | X | | | | | | | | | | | | | | | 1 | |
| ALL OTHER SYSTEMS | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Multiple organs, NOS | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | | *49 | |
| Malignant lymphoma, NOS | | | | | | | | | | | | | | | | | | | | | | | | | | X | |
| | | | | | | | | | | | | | | | | | | | | | | | | | | 2 | |

* Animals necropsied

TABLE C3. ANALYSIS OF PRIMARY TUMORS IN MALE AUGUST RATS IN THE TWO-YEAR GAVAGE STUDY OF TRICHLOROETHYLENE

| | Untreated Control | Vehicle Control | 500 mg/kg | 1,000 mg/kg |
|---|-------------------|-----------------|------------|-------------|
| Subcutaneous Tissue: Sarcoma or Sarcoma, Unclear Primary or Metastatic | | | | |
| Overall Rates (a) | 1/50 (2%) | 0/50 (0%) | 1/50 (2%) | 3/49 (6%) |
| Adjusted Rates (b) | 3.8% | 0.0% | 2.9% | 15.0% |
| Terminal Rates (c) | 0/24 (0%) | 0/21 (0%) | 0/13 (0%) | 1/16 (6%) |
| Life Table Tests (d) | | P=0.033 | P=0.440 | P=0.064 |
| Incidental Tumor Tests (d) | | P=0.032 | P=0.519 | P=0.050 |
| Cochran-Armitage Trend Test (d) | | P=0.058 | | |
| Fisher Exact Test | | | P=0.500 | P=0.117 |
| Pituitary: Adenoma | | | | |
| Overall Rates (a) | 16/46 (35%) | 23/44 (52%) | 7/41 (17%) | 4/41 (10%) |
| Adjusted Rates (b) | 55.5% | 70.3% | 34.9% | 26.7% |
| Terminal Rates (c) | 11/23 (48%) | 12/21 (57%) | 2/11 (18%) | 4/15 (27%) |
| Life Table Tests (d) | | P=0.001N | P=0.069N | P=0.002N |
| Incidental Tumor Tests (d) | | P<0.001N | P=0.010N | P<0.001N |
| Cochran-Armitage Trend Test (d) | | P<0.001N | | |
| Fisher Exact Test | | | P=0.001N | P<0.001N |
| Pituitary: Carcinoma | | | | |
| Overall Rates (a) | 0/46 (0%) | 3/44 (7%) | 0/41 (0%) | 0/41 (0%) |
| Adjusted Rates (b) | 0.0% | 14.3% | 0.0% | 0.0% |
| Terminal Rates (c) | 0/23 (0%) | 3/21 (14%) | 0/11 (0%) | 0/15 (0%) |
| Life Table Tests (d) | | P=0.074N | P=0.252N | P=0.183N |
| Incidental Tumor Tests (d) | | P=0.074N | P=0.252N | P=0.183N |
| Cochran-Armitage Trend Test (d) | | P=0.042N | | |
| Fisher Exact Test | | | P=0.134N | P=0.134N |
| Pituitary: Adenoma or Carcinoma | | | | |
| Overall Rates (a) | 16/46 (35%) | 26/44 (59%) | 7/41 (17%) | 4/41 (10%) |
| Adjusted Rates (b) | 55.5% | 80.2% | 34.9% | 26.7% |
| Terminal Rates (c) | 11/23 (48%) | 15/21 (71%) | 2/11 (18%) | 4/15 (27%) |
| Life Table Tests (d) | | P<0.001N | P=0.027N | P<0.001N |
| Incidental Tumor Tests (d) | | P<0.001N | P=0.002N | P<0.001N |
| Cochran-Armitage Trend Test (d) | | P<0.001N | | |
| Fisher Exact Test | | | P<0.001N | P<0.001N |
| Adrenal: Cortical Adenoma | | | | |
| Overall Rates (a) | 6/49 (14%) | 5/50 (10%) | 3/49 (6%) | 5/47 (11%) |
| Adjusted Rates (b) | 23.5% | 20.0% | 20.4% | 29.4% |
| Terminal Rates (c) | 5/24 (21%) | 3/21 (14%) | 2/13 (15%) | 4/16 (25%) |
| Life Table Tests (d) | | P=0.356 | P=0.627N | P=0.414 |
| Incidental Tumor Tests (d) | | P=0.271 | P=0.633N | P=0.333 |
| Cochran-Armitage Trend Test (d) | | P=0.533 | | |
| Fisher Exact Test | | | P=0.369N | P=0.590 |
| Adrenal: Pheochromocytoma | | | | |
| Overall Rates (a) | 16/49 (33%) | 10/50 (20%) | 5/49 (10%) | 2/47 (4%) |
| Adjusted Rates (b) | 63.7% | 41.0% | 33.9% | 12.5% |
| Terminal Rates (c) | 15/24 (63%) | 7/21 (33%) | 4/13 (31%) | 2/16 (13%) |
| Life Table Tests (d) | | P=0.039N | P=0.455N | P=0.049N |
| Incidental Tumor Tests (d) | | P=0.053N | P=0.460N | P=0.073N |
| Cochran-Armitage Trend Test (d) | | P=0.012N | | |
| Fisher Exact Test | | | P=0.140N | P=0.018N |
| Adrenal: Pheochromocytoma or Pheochromocytoma, Malignant | | | | |
| Overall Rates (a) | 16/49 (33%) | 10/50 (20%) | 6/49 (12%) | 3/47 (6%) |
| Adjusted Rates (b) | 63.7% | 41.0% | 41.3% | 18.8% |
| Terminal Rates (c) | 15/24 (63%) | 7/21 (33%) | 5/13 (38%) | 3/16 (19%) |
| Life Table Tests (d) | | P=0.090N | P=0.602N | P=0.104N |
| Incidental Tumor Tests (d) | | P=0.119N | P=0.610N | P=0.150N |
| Cochran-Armitage Trend Test (d) | | P=0.033N | | |
| Fisher Exact Test | | | P=0.220N | P=0.046N |

TABLE C3. ANALYSIS OF PRIMARY TUMORS IN MALE AUGUST RATS IN THE TWO-YEAR GAVAGE STUDY OF TRICHLOROETHYLENE (Continued)

| | Untreated Control | Vehicle Control | 500 mg/kg | 1,000 mg/kg |
|--|-------------------|-----------------|-------------|--------------|
| Thyroid: C-Cell Adenoma | | | | |
| Overall Rates (a) | 3/49 (6%) | 7/46 (15%) | 0/44 (0%) | 0/45 (0%) |
| Adjusted Rates (b) | 13.0% | 27.1% | 0.0% | 0.0% |
| Terminal Rates (c) | 3/23 (13%) | 4/21 (19%) | 0/13 (0%) | 0/16 (0%) |
| Life Table Tests (d) | | P=0.005N | P=0.041N | P=0.030N |
| Incidental Tumor Tests (d) | | P=0.005N | P=0.035N | P=0.030N |
| Cochran-Armitage Trend Test (d) | | P=0.001N | | |
| Fisher Exact Test | | | P=0.007N | P=0.007N |
| Thyroid: C-Cell Adenoma or Carcinoma | | | | |
| Overall Rates (a) | 3/49 (6%) | 8/46 (17%) | 0/44 (0%) | 0/45 (0%) |
| Adjusted Rates (b) | 13.0% | 31.4% | 0.0% | 0.0% |
| Terminal Rates (c) | 3/23 (13%) | 5/21 (24%) | 0/13 (0%) | 0/16 (0%) |
| Life Table Tests (d) | | P=0.003N | P=0.027N | P=0.018N |
| Incidental Tumor Tests (d) | | P=0.003N | P=0.023N | P=0.018N |
| Cochran-Armitage Trend Test (d) | | P<0.001N | | |
| Fisher Exact Test | | | P=0.003N | P=0.003N |
| Pancreatic Islets: Islet Cell Adenoma | | | | |
| Overall Rates (a) | 5/49 (10%) | 5/50 (10%) | 0/49 (0%) | 1/48 (2%) |
| Adjusted Rates (b) | 19.8% | 22.0% | 0.0% | 6.3% |
| Terminal Rates (c) | 4/24 (17%) | 4/21 (19%) | 0/13 (0%) | 1/16 (6%) |
| Life Table Tests (d) | | P=0.079N | P=0.088N | P=0.181N |
| Incidental Tumor Tests (d) | | P=0.094N | P=0.098N | P=0.210N |
| Cochran-Armitage Trend Test (d) | | P=0.040N | | |
| Fisher Exact Test | | | P=0.030N | P=0.112N |
| Testis: Interstitial Cell Tumor | | | | |
| Overall Rates (a) | 36/50 (72%) | 34/50 (68%) | 30/50 (60%) | 26/49 (53%) |
| Adjusted Rates (b) | 97.2% | 97.1% | 96.7% | 100.0% |
| Terminal Rates (c) | 23/24 (96%) | 20/21 (95%) | 12/13 (92%) | 16/16 (100%) |
| Life Table Tests (d) | | P=0.279 | P=0.049 | P=0.385 |
| Incidental Tumor Tests (d) | | P=0.129 | P=0.055 | P=0.179 |
| Cochran-Armitage Trend Test (d) | | P=0.078N | | |
| Fisher Exact Test | | | P=0.267N | P=0.094N |

(a) Number of tumor-bearing animals/number of animals examined at the site

(b) Kaplan-Meier estimated tumor incidence at the end of the study after adjusting for intercurrent mortality

(c) Observed tumor incidence at terminal kill

(d) Beneath the control incidence are the P values associated with the trend test. Beneath the dosed group incidence are the P values corresponding to pairwise comparisons between that dosed group and the vehicle controls. The life table analysis regards tumors in animals dying prior to terminal kill as being (directly or indirectly) the cause of death. The incidental tumor test regards these lesions as nonfatal. The Cochran-Armitage and Fisher exact tests compare directly the overall incidence rates. A negative trend or lower incidence in a dosed group is indicated by (N).

**TABLE C4. SUMMARY OF THE INCIDENCE NONNEOPLASTIC LESIONS OF MALE AUGUST RATS
IN THE TWO-YEAR GAVAGE STUDY OF TRICHLOROETHYLENE**

| | CONTROL (UNTR) | CONTROL (VEH) | LOW DOSE | HIGH DOSE |
|---|----------------|---------------|----------|-----------|
| ANIMALS INITIALLY IN STUDY | 50 | 50 | 50 | 50 |
| ANIMALS NECROPSIED | 50 | 50 | 50 | 49 |
| ANIMALS EXAMINED HISTOPATHOLOGICALLY | 50 | 50 | 50 | 49 |
| INTEGUMENTARY SYSTEM | | | | |
| *Subcutaneous tissue | (50) | (50) | (50) | (49) |
| Hemorrhage | 1 (2%) | | | |
| Abscess, chronic | | | 1 (2%) | 1 (2%) |
| RESPIRATORY SYSTEM | | | | |
| *Tracheal lumen | (50) | (50) | (50) | (49) |
| Hemorrhage | | 1 (2%) | 1 (2%) | |
| #Trachea | (49) | (48) | (42) | (42) |
| Inflammation, acute | 1 (2%) | | | |
| Foreign material, NOS | 2 (4%) | 1 (2%) | 2 (5%) | 3 (7%) |
| #Tracheal submucosa | (49) | (48) | (42) | (42) |
| Hemorrhage | | | | 1 (2%) |
| #Lung | (50) | (50) | (50) | (49) |
| Aspiration, foreign body | | | 1 (2%) | |
| Emphysema, alveolar | | | | 2 (4%) |
| Congestion, NOS | | | 3 (6%) | 8 (16%) |
| Edema, NOS | | | 1 (2%) | 4 (8%) |
| Hemorrhage | 2 (4%) | 4 (8%) | 13 (26%) | 12 (24%) |
| Inflammation, NOS | 2 (4%) | 1 (2%) | 2 (4%) | |
| Inflammation, focal | 4 (8%) | 7 (14%) | 1 (2%) | 2 (4%) |
| Inflammation, multifocal | 1 (2%) | 2 (4%) | 1 (2%) | 1 (2%) |
| Inflammation, interstitial | 1 (2%) | | | |
| Pneumonia, aspiration | | 1 (2%) | | |
| Bronchopneumonia, acute | 3 (6%) | | 1 (2%) | 1 (2%) |
| Inflammation, acute | 4 (8%) | 1 (2%) | 2 (4%) | 5 (10%) |
| Inflammation, acute/chronic | | | 1 (2%) | |
| Inflammation, chronic | 1 (2%) | 3 (6%) | | 1 (2%) |
| Pneumonia, interstitial chronic | | | | 1 (2%) |
| Inflammation, chronic focal | 1 (2%) | 4 (8%) | 4 (8%) | 2 (4%) |
| Inflammation, granulomatous | 5 (10%) | 4 (8%) | 2 (4%) | 3 (6%) |
| Inflammation, granulomatous focal | 6 (12%) | 19 (38%) | 16 (32%) | 12 (24%) |
| Foreign material, NOS | 2 (4%) | 3 (6%) | 10 (20%) | 8 (16%) |
| Alveolar macrophages | | | | 1 (2%) |
| #Lung/alveoli | (50) | (50) | (50) | (49) |
| Hemorrhage | 1 (2%) | | 1 (2%) | |
| HEMATOPOIETIC SYSTEM | | | | |
| #Bone marrow | (50) | (50) | (49) | (49) |
| Hyperplasia, granulocytic | | | | 1 (2%) |
| #Spleen | (49) | (50) | (48) | (49) |
| Congestion, NOS | | | 2 (4%) | 2 (4%) |
| Hemorrhage | | | 1 (2%) | |
| Hyperplasia, lymphoid | | 1 (2%) | | |
| Hematopoiesis | 6 (12%) | 5 (10%) | 2 (4%) | 8 (16%) |
| #Lymph node | (48) | (45) | (43) | (33) |
| Congestion, NOS | | | 2 (5%) | |
| #Pancreatic lymph node | (48) | (45) | (43) | (33) |
| Hemorrhage | | | 1 (2%) | |
| #Mesenteric lymph node | (48) | (45) | (43) | (33) |
| Congestion, NOS | | | | 2 (6%) |
| Hemorrhage | 1 (2%) | | 2 (5%) | 2 (6%) |
| #Liver | (50) | (50) | (50) | (48) |
| Hematopoiesis | 1 (2%) | | | |

TABLE C4. SUMMARY OF THE INCIDENCE OF NONNEOPLASTIC LESIONS IN MALE AUGUST RATS
IN THE TWO-YEAR GAVAGE STUDY OF TRICHLOROETHYLENE (Continued)

| | CONTROL (UNTR) | CONTROL (VEH) | LOW DOSE | HIGH DOSE |
|---|----------------|---------------|----------|-----------|
| HEMATOPOIETIC SYSTEM (Continued) | | | | |
| #Thymus | (6) | (1) | (9) | (8) |
| Hemorrhage | | | 2 (22%) | 1 (13%) |
| Karyorrhexis | | | 1 (11%) | |
| CIRCULATORY SYSTEM | | | | |
| #Heart | (50) | (50) | (50) | (49) |
| Mineralization | | | | 1 (2%) |
| Hemorrhage | | | | 1 (2%) |
| Inflammation, focal | 1 (2%) | | | |
| Inflammation, chronic | 1 (2%) | 1 (2%) | | |
| Inflammation, chronic focal | 1 (2%) | 1 (2%) | | |
| Fibrosis, focal | 1 (2%) | | | |
| Degeneration, NOS | | 1 (2%) | | |
| #Myocardium | (50) | (50) | (50) | (49) |
| Degeneration, NOS | 1 (2%) | | | |
| #Endocardium | (50) | (50) | (50) | (49) |
| Hyperplasia, focal | 1 (2%) | | | |
| *Pulmonary artery | (50) | (50) | (50) | (49) |
| Calcification, focal | | 1 (2%) | 1 (2%) | 1 (2%) |
| DIGESTIVE SYSTEM | | | | |
| #Salivary gland | (47) | (50) | (49) | (46) |
| Inflammation, chronic focal | | | | 1 (2%) |
| Atrophy, focal | | 2 (4%) | 1 (2%) | |
| #Liver | (50) | (50) | (50) | (48) |
| Congestion, NOS | 1 (2%) | | | 1 (2%) |
| Congestion, passive | | | | 1 (2%) |
| Congestion, acute passive | | | | 1 (2%) |
| Hemorrhage | 1 (2%) | | | |
| Inflammation, NOS | | 1 (2%) | | |
| Inflammation, focal | | | | 2 (4%) |
| Inflammation, acute | 1 (2%) | | | |
| Inflammation, chronic | | | | 1 (2%) |
| Inflammation, granulomatous focal | | | | 1 (2%) |
| Fibrosis, focal | | | | 1 (2%) |
| Necrosis, focal | 3 (6%) | 4 (8%) | 2 (4%) | 5 (10%) |
| Necrosis, diffuse | | 1 (2%) | | 1 (2%) |
| Necrosis, hemorrhagic | 1 (2%) | | | |
| Necrosis, central | 3 (6%) | 2 (4%) | 2 (4%) | 1 (2%) |
| Necrosis, peripheral | | 1 (2%) | | |
| Metamorphosis, fatty | 9 (18%) | 4 (8%) | 1 (2%) | 2 (4%) |
| Lipoidosis | | 2 (4%) | | |
| Cytoplasmic change, NOS | | 4 (8%) | | |
| Basophilic cyto change | | | | 1 (2%) |
| Focal cellular change | 2 (4%) | 2 (4%) | 1 (2%) | |
| Eosinophilic cyto change | | | | 1 (2%) |
| Clear cell change | 6 (12%) | | | |
| Cytologic alteration, NOS | 1 (2%) | | 1 (2%) | |
| Hepatocytomegaly | 2 (4%) | | | |
| #Liver/centrilobular | (50) | (50) | (50) | (48) |
| Necrosis, NOS | | 1 (2%) | | |
| Necrosis, hemorrhagic | 1 (2%) | | | |
| Metamorphosis, fatty | | | 1 (2%) | |
| #Liver/hepatocytes | (50) | (50) | (50) | (48) |
| Necrosis, NOS | 1 (2%) | | | |

TABLE C4. SUMMARY OF THE INCIDENCE OF NONNEOPLASTIC LESIONS IN MALE AUGUST RATS IN THE TWO-YEAR GAVAGE STUDY OF TRICHLOROETHYLENE (Continued)

| | CONTROL (UNTR) | CONTROL (VEH) | LOW DOSE | HIGH DOSE |
|-------------------------------------|----------------|---------------|----------|-----------|
| DIGESTIVE SYSTEM (Continued) | | | | |
| #Bile duct | (50) | (50) | (50) | (48) |
| Inflammation, chronic | 1 (2%) | 4 (8%) | 1 (2%) | |
| Inflammation, chronic focal | 1 (2%) | | | |
| Fibrosis | | 1 (2%) | | |
| Sclerosis | | | 1 (2%) | |
| Hyperplasia, NOS | 5 (10%) | 6 (12%) | 1 (2%) | |
| Hyperplasia, focal | | 1 (2%) | | |
| #Pancreas | (49) | (50) | (49) | (48) |
| Hemorrhage | 1 (2%) | 1 (2%) | 1 (2%) | |
| Inflammation, focal | | | | 1 (2%) |
| Inflammation, acute focal | | 1 (2%) | | |
| Atrophy, NOS | 1 (2%) | | | |
| Atrophy, focal | 2 (4%) | 1 (2%) | 2 (4%) | 4 (8%) |
| #Pancreatic acinus | (49) | (50) | (49) | (48) |
| Atrophy, NOS | 20 (41%) | 2 (4%) | 2 (4%) | |
| Atrophy, focal | 12 (24%) | 29 (58%) | 17 (35%) | 13 (27%) |
| Atrophy, diffuse | 1 (2%) | 3 (6%) | 2 (4%) | |
| *Esophageal lumen | (50) | (50) | (50) | (49) |
| Hemorrhage | 1 (2%) | 2 (4%) | 2 (4%) | 2 (4%) |
| #Esophagus | (50) | (48) | (48) | (47) |
| Dilatation, NOS | | 2 (4%) | | |
| Inflammation, interstitial | 1 (2%) | | | |
| Inflammation, acute | | 1 (2%) | | |
| Inflammation, granulomatous | | 1 (2%) | | |
| #Stomach | (50) | (50) | (50) | (49) |
| Edema, NOS | | | | 1 (2%) |
| Inflammation, acute/chronic | | | 1 (2%) | |
| #Gastric mucosa | (50) | (50) | (50) | (49) |
| Erosion | | 1 (2%) | | |
| Hyperkeratosis | | | | 1 (2%) |
| #Gastric submucosa | (50) | (50) | (50) | (49) |
| Inflammation, focal | | 1 (2%) | | |
| Inflammation, chronic | 1 (2%) | 1 (2%) | | |
| Inflammation, granulomatous | 2 (4%) | | | |
| #Forestomach | (50) | (50) | (50) | (49) |
| Ulcer, NOS | | 1 (2%) | | |
| Inflammation, acute/chronic | | 1 (2%) | | |
| Degeneration, NOS | | 1 (2%) | | |
| Hyperplasia, epithelial | 5 (10%) | 1 (2%) | | |
| Hyperkeratosis | 2 (4%) | | | 1 (2%) |
| #Colon | (49) | (50) | (50) | (44) |
| Hemorrhage | | | 1 (2%) | |
| Parasitism | 2 (4%) | 1 (2%) | | 1 (2%) |
| *Rectum | (50) | (50) | (50) | (49) |
| Ulcer, acute | | | 1 (2%) | |
| URINARY SYSTEM | | | | |
| #Kidney | (50) | (50) | (50) | (49) |
| Mineralization | 1 (2%) | | | |
| Cast, NOS | | 1 (2%) | 2 (4%) | |
| Hydronephrosis | 1 (2%) | 1 (2%) | | |
| Cyst, NOS | 2 (4%) | 3 (6%) | | 1 (2%) |
| Inflammation, chronic focal | 2 (4%) | 1 (2%) | | |
| Nephropathy | 43 (86%) | 47 (94%) | 32 (64%) | 36 (73%) |
| Nephropathy, toxic | | | 10 (20%) | 31 (63%) |
| Infarct, healed | | | 1 (2%) | |
| Pigmentation, NOS | 1 (2%) | | | |
| Cytomegaly | | | 46 (92%) | 46 (94%) |
| Hyperplasia, tubular cell | | | | 1 (2%) |

TABLE C4. SUMMARY OF THE INCIDENCE OF NONNEOPLASTIC LESIONS IN MALE AUGUST RATS IN THE TWO-YEAR GAVAGE STUDY OF TRICHLOROETHYLENE (Continued)

| | CONTROL (UNTR) | CONTROL (VEH) | LOW DOSE | HIGH DOSE |
|-----------------------------------|----------------|---------------|----------|-----------|
| URINARY SYSTEM (Continued) | | | | |
| #Kidney/tubule | (50) | (50) | (50) | (49) |
| Dilatation, NOS | | 1 (2%) | | 1 (2%) |
| Degeneration, NOS | 1 (2%) | 1 (2%) | | 1 (2%) |
| Atypia, NOS | | | 1 (2%) | 1 (2%) |
| #Kidney/pelvis | (50) | (50) | (50) | (49) |
| Hyperplasia, epithelial | 3 (6%) | 1 (2%) | | |
| #Urinary bladder | (50) | (50) | (47) | (44) |
| Hemorrhage | | | 1 (2%) | 1 (2%) |
| #Urinary bladder/mucosa | (50) | (50) | (47) | (44) |
| Hyperplasia, papillary | 1 (2%) | | | |
| ENDOCRINE SYSTEM | | | | |
| #Pituitary | (46) | (44) | (41) | (41) |
| Cyst, NOS | 3 (7%) | 1 (2%) | | |
| Hemorrhage | | | | 1 (2%) |
| Hyperplasia, NOS | 1 (2%) | | | |
| Hyperplasia, focal | 2 (4%) | 2 (5%) | 4 (10%) | 1 (2%) |
| Angiectasis | 2 (4%) | | 1 (2%) | |
| #Adrenal | (49) | (50) | (49) | (47) |
| Congestion, NOS | 1 (2%) | | | |
| Inflammation, focal | | | 1 (2%) | |
| Inflammation, chronic focal | | | 1 (2%) | |
| Necrosis, cortical | | | | 1 (2%) |
| Lipoidosis | | 1 (2%) | | |
| #Adrenal cortex | (49) | (50) | (49) | (47) |
| Lipoidosis | 2 (4%) | 1 (2%) | | |
| Focal cellular change | | 1 (2%) | | |
| Hyperplasia, NOS | | 1 (2%) | | |
| Hyperplasia, focal | | 4 (8%) | | |
| #Adrenal medulla | (49) | (50) | (49) | (47) |
| Cytomegaly | 1 (2%) | | | |
| Hyperplasia, NOS | 1 (2%) | 1 (2%) | 1 (2%) | |
| Hyperplasia, focal | 3 (6%) | 1 (2%) | 2 (4%) | 1 (2%) |
| #Thyroid | (49) | (46) | (44) | (45) |
| Follicular cyst, NOS | | | 1 (2%) | |
| Inflammation, granulomatous focal | | | | 1 (2%) |
| Hyperplasia, C-cell | 1 (2%) | 3 (7%) | | |
| #Pancreatic islets | (49) | (50) | (49) | (48) |
| Hyperplasia, NOS | | 1 (2%) | | |
| Hyperplasia, focal | | | 1 (2%) | |
| REPRODUCTIVE SYSTEM | | | | |
| *Mammary gland | (50) | (50) | (50) | (49) |
| Pigmentation, NOS | | | | 1 (2%) |
| Hyperplasia, NOS | 1 (2%) | | 1 (2%) | |
| Lactation | 2 (4%) | 3 (6%) | 1 (2%) | 1 (2%) |
| #Prostate | (50) | (50) | (48) | (46) |
| Edema, NOS | | | | 1 (2%) |
| Hemorrhage | | | 1 (2%) | 1 (2%) |
| Inflammation, NOS | | | 1 (2%) | |
| Inflammation, focal | | 2 (4%) | | |
| Inflammation, acute | 1 (2%) | | | 1 (2%) |
| Inflammation, acute focal | | | 1 (2%) | |
| Inflammation, acute/chronic | | 1 (2%) | | |
| Inflammation, chronic | | | 1 (2%) | 1 (2%) |
| Inflammation, chronic focal | 1 (2%) | 2 (4%) | 1 (2%) | |
| Fibrosis, focal | | | | 1 (2%) |
| Hyperplasia, epithelial | | 1 (2%) | | |

TABLE C4. SUMMARY OF THE INCIDENCE OF NONNEOPLASTIC LESIONS IN MALE AUGUST RATS IN THE TWO-YEAR GAVAGE STUDY OF TRICHLOROETHYLENE (Continued)

| | CONTROL (UNTR) | CONTROL (VEH) | LOW DOSE | HIGH DOSE |
|--|----------------|---------------|----------|-----------|
| REPRODUCTIVE SYSTEM (Continued) | | | | |
| #Prostatic duct | (50) | (50) | (48) | (46) |
| Polyp, NOS | 1 (2%) | | | |
| *Seminal vesicle | (50) | (50) | (50) | (49) |
| Hemorrhage | | | 1 (2%) | |
| Inflammation, NOS | | | 1 (2%) | |
| #Testis | (50) | (50) | (50) | (49) |
| Edema, interstitial | 1 (2%) | | | |
| Hemorrhage | | | 1 (2%) | 1 (2%) |
| Fibrosis, diffuse | | 1 (2%) | | |
| Degeneration, NOS | | | 1 (2%) | 2 (4%) |
| Syncytial alteration | | | | 1 (2%) |
| Atrophy, NOS | 5 (10%) | 11 (22%) | 5 (10%) | 5 (10%) |
| Atrophy, diffuse | 1 (2%) | 1 (2%) | | |
| Hyperplasia, interstitial cell | 7 (14%) | 9 (18%) | 6 (12%) | 10 (20%) |
| #Testis/tubule | (50) | (50) | (50) | (49) |
| Degeneration, NOS | 1 (2%) | | | |
| Atrophy, focal | 1 (2%) | | | 1 (2%) |
| NERVOUS SYSTEM | | | | |
| #Brain/meninges | (50) | (50) | (50) | (49) |
| Fibrosis | 1 (2%) | | | |
| #Brain | (50) | (50) | (50) | (49) |
| Hydrocephalus, internal | 1 (2%) | 2 (4%) | | |
| Cyst, NOS | | 1 (2%) | | |
| Hemorrhage | 1 (2%) | | | 2 (4%) |
| SPECIAL SENSE ORGANS | | | | |
| *Harderian gland | (50) | (50) | (50) | (49) |
| Inflammation, focal | | | | 1 (2%) |
| Inflammation, chronic | | | 1 (2%) | |
| Inflammation, chronic focal | | | 2 (4%) | |
| *Middle ear | (50) | (50) | (50) | (49) |
| Inflammation, acute suppurative | | | 1 (2%) | |
| MUSCULOSKELETAL SYSTEM | | | | |
| *Muscle of neck | (50) | (50) | (50) | (49) |
| Inflammation, granulomatous | | | 1 (2%) | |
| BODY CAVITIES | | | | |
| *Mediastinum | (50) | (50) | (50) | (49) |
| Inflammation, acute suppurative | | | 1 (2%) | |
| ALL OTHER SYSTEMS | | | | |
| *Multiple organs | (50) | (50) | (50) | (49) |
| Hemorrhage | | 1 (2%) | | |
| Inflammation, hemorrhagic | | | | 1 (2%) |
| Adipose tissue | | | | |
| Inflammation, NOS | | | 1 | |

**TABLE C4. SUMMARY OF THE INCIDENCE OF NONNEOPLASTIC LESIONS IN MALE AUGUST RATS
IN THE TWO-YEAR GAVAGE STUDY OF TRICHLOROETHYLENE (Continued)**

| | CONTROL (UNTR) | CONTROL (VEH) | LOW DOSE | HIGH DOSE |
|-----------------------------------|----------------|---------------|----------|-----------|
| SPECIAL MORPHOLOGY SUMMARY | | | | |
| Animal missexed/no necropsy | | | | 1 |
| Auto/necropsy/histo perf | | | 1 | |

* Number of animals receiving complete necropsy examination; all gross lesions including masses examined microscopically.
Number of animals examined microscopically at this site

APPENDIX D

SUMMARY OF LESIONS IN FEMALE AUGUST RATS IN THE TWO-YEAR GAVAGE STUDY OF TRICHLOROETHYLENE

| | PAGE | |
|----------|---|-----|
| TABLE D1 | SUMMARY OF THE INCIDENCE OF NEOPLASMS IN FEMALE AUGUST RATS IN THE TWO-YEAR GAVAGE STUDY OF TRICHLOROETHYLENE | 143 |
| TABLE D2 | INDIVIDUAL ANIMAL TUMOR PATHOLOGY OF FEMALE AUGUST RATS IN THE TWO-YEAR GAVAGE STUDY OF TRICHLOROETHYLENE | 146 |
| TABLE D3 | ANALYSIS OF PRIMARY TUMORS IN FEMALE AUGUST RATS IN THE TWO-YEAR GAVAGE STUDY OF TRICHLOROETHYLENE | 154 |
| TABLE D4 | SUMMARY OF THE INCIDENCE OF NONNEOPLASTIC LESIONS IN FEMALE AUGUST RATS IN THE TWO-YEAR GAVAGE STUDY OF TRICHLOROETHYLENE | 157 |

TABLE D1. SUMMARY OF THE INCIDENCE OF NEOPLASMS IN FEMALE AUGUST RATS IN THE TWO-YEAR GAVAGE STUDY OF TRICHLOROETHYLENE

| | CONTROL (UNTR) | CONTROL (VEH) | LOW DOSE | HIGH DOSE |
|--|----------------|---------------|----------|-----------|
| ANIMALS INITIALLY IN STUDY | 50 | 50 | 50 | 50 |
| ANIMALS NECROPSIED | 50 | 50 | 50 | 50 |
| ANIMALS EXAMINED HISTOPATHOLOGICALLY | 50 | 50 | 50 | 50 |
| INTEGUMENTARY SYSTEM | | | | |
| *Skin | (50) | (50) | (50) | (50) |
| Squamous cell carcinoma | 1 (2%) | | | |
| Malignant melanoma | | | 1 (2%) | |
| *Subcutaneous tissue | (50) | (50) | (50) | (50) |
| Sarcoma, NOS | 1 (2%) | 1 (2%) | | |
| RESPIRATORY SYSTEM | | | | |
| #Lung | (50) | (50) | (50) | (50) |
| Undifferentiated carcinoma, metastatic | | 1 (2%) | | |
| Squamous cell carcinoma, metastatic | 1 (2%) | | | |
| Alveolar/bronchiolar adenoma | | | 1 (2%) | |
| Sarcoma, NOS, metastatic | 1 (2%) | | | |
| HEMATOPOIETIC SYSTEM | | | | |
| *Multiple organs | (50) | (50) | (50) | (50) |
| Malignant lymphoma, NOS | | | 2 (4%) | |
| Leukemia, NOS | | 1 (2%) | | 1 (2%) |
| Monocytic leukemia | | | | 2 (4%) |
| #Liver | (50) | (48) | (48) | (50) |
| Leukemia, NOS | | | | 2 (4%) |
| CIRCULATORY SYSTEM | | | | |
| None | | | | |
| DIGESTIVE SYSTEM | | | | |
| #Liver | (50) | (48) | (48) | (50) |
| Neoplastic nodule | | 2 (4%) | | |
| URINARY SYSTEM | | | | |
| #Kidney | (50) | (49) | (48) | (50) |
| Tubular cell adenoma | | 1 (2%) | 2 (4%) | |
| Tubular cell adenocarcinoma | | | 2 (4%) | |
| Sarcoma, NOS, metastatic | 1 (2%) | | | |
| ENDOCRINE SYSTEM | | | | |
| #Pituitary | (49) | (49) | (47) | (43) |
| Adenoma, NOS | 42 (86%) | 32 (65%) | 26 (55%) | 18 (42%) |
| Chromophobe adenoma | | 3 (6%) | | |
| Meningioma | | | | 1 (2%) |
| #Adrenal | (50) | (48) | (48) | (50) |
| Cortical adenoma | 4 (8%) | 6 (13%) | 6 (13%) | 5 (10%) |
| Cortical carcinoma | | 1 (2%) | | |
| Pheochromocytoma | 6 (12%) | 9 (19%) | 2 (4%) | |
| #Thyroid | (50) | (49) | (49) | (50) |
| Follicular cell adenoma | 1 (2%) | | 2 (4%) | |
| C-cell adenoma | | | 3 (6%) | 1 (2%) |
| C-cell carcinoma | | | 1 (2%) | |

TABLE D1. SUMMARY OF THE INCIDENCE OF NEOPLASMS IN FEMALE AUGUST RATS IN THE TWO-YEAR GAVAGE STUDY OF TRICHLOROETHYLENE (Continued)

| | CONTROL (UNTR) | CONTROL (VEH) | LOW DOSE | HIGH DOSE |
|-------------------------------------|----------------|---------------|----------|-----------|
| ENDOCRINE SYSTEM (Continued) | | | | |
| #Pancreatic islets | (50) | (49) | (48) | (48) |
| Islet cell adenoma | 1 (2%) | | 1 (2%) | 1 (2%) |
| Islet cell carcinoma | | | 1 (2%) | |
| REPRODUCTIVE SYSTEM | | | | |
| *Mammary gland | (50) | (50) | (50) | (50) |
| Undifferentiated carcinoma | | 1 (2%) | | |
| Adenoma, NOS | 3 (6%) | 4 (8%) | 4 (8%) | |
| Adenocarcinoma, NOS | | 1 (2%) | | 1 (2%) |
| Fibroma | 1 (2%) | 2 (4%) | | |
| Fibroadenoma | 13 (26%) | 10 (20%) | 8 (16%) | 4 (8%) |
| *Clitoral gland | (50) | (50) | (50) | (50) |
| Carcinoma, NOS | | | | 1 (2%) |
| #Uterus | (50) | (49) | (48) | (49) |
| Sarcoma, NOS | 2 (4%) | 1 (2%) | | |
| Leiomyosarcoma | | | 1 (2%) | |
| Endometrial stromal polyp | 1 (2%) | 3 (6%) | | |
| Carcinosarcoma | 1 (2%) | | | |
| #Ovary | (49) | (47) | (47) | (46) |
| Luteoma | | | 1 (2%) | |
| NERVOUS SYSTEM | | | | |
| *Peripheral nerve | (50) | (50) | (50) | (50) |
| Neurofibrosarcoma | | | 1 (2%) | |
| #Brain | (50) | (49) | (50) | (50) |
| Granular cell tumor, NOS | 1 (2%) | 1 (2%) | 1 (2%) | 1 (2%) |
| Granular cell tumor, malignant | 1 (2%) | | 2 (4%) | |
| Ependymoma | | | | 1 (2%) |
| Astrocytoma | | | 1 (2%) | |
| SPECIAL SENSE ORGANS | | | | |
| None | | | | |
| MUSCULOSKELETAL SYSTEM | | | | |
| None | | | | |
| BODY CAVITIES | | | | |
| None | | | | |
| ALL OTHER SYSTEMS | | | | |
| *Multiple organs | (50) | (50) | (50) | (50) |
| Adenoma, NOS | 1 (2%) | | | |
| Thigh | | | | |
| Sarcoma, NOS | 1 | | | |

TABLE D1. SUMMARY OF THE INCIDENCE OF NEOPLASMS IN FEMALE AUGUST RATS IN THE TWO-YEAR GAVAGE STUDY OF TRICHLOROETHYLENE (Continued)

| | CONTROL (UNTR) | CONTROL (VEH) | LOW DOSE | HIGH DOSE |
|--|----------------|---------------|----------|-----------|
| ANIMAL DISPOSITION SUMMARY | | | | |
| Animals initially in study | 50 | 50 | 50 | 50 |
| Natural death | 4 | 4 | 5 | 6 |
| Moribund sacrifice | 20 | 22 | 13 | 7 |
| Terminal sacrifice | 26 | 23 | 26 | 24 |
| Dosing accident | | 1 | 3 | 6 |
| Accidentally killed, nda | | | 1 | |
| Accidentally killed, NOS | | | 2 | 7 |
| TUMOR SUMMARY | | | | |
| Total animals with primary tumors** | 45 | 43 | 35 | 29 |
| Total primary tumors | 81 | 79 | 69 | 39 |
| Total animals with benign tumors | 44 | 40 | 32 | 24 |
| Total benign tumors | 73 | 70 | 56 | 29 |
| Total animals with malignant tumors | 5 | 6 | 11 | 9 |
| Total malignant tumors | 7 | 6 | 12 | 9 |
| Total animals with secondary tumors## | 2 | 1 | | |
| Total secondary tumors | 3 | 1 | | |
| Total animals with tumors uncertain-- benign or malignant | 1 | 3 | 1 | 1 |
| Total uncertain tumors | 1 | 3 | 1 | 1 |

* Number of animals receiving complete necropsy examination; all gross lesions including masses examined microscopically.

** Primary tumors: all tumors except secondary tumors

Number of animals examined microscopically at this site

Secondary tumors: metastatic tumors or tumors invasive into an adjacent organ

TABLE D2. INDIVIDUAL ANIMAL TUMOR PATHOLOGY OF FEMALE AUGUST RATS IN THE TWO-YEAR GAVAGE STUDY OF TRICHLOROETHYLENE: VEHICLE CONTROL

| ANIMAL NUMBER | 345 | 347 | 358 | 359 | 362 | 364 | 366 | 367 | 368 | 371 | 372 | 373 | 374 | 375 | 377 | 379 | 380 | 381 | 382 | 383 | 384 | 385 | 386 | 387 | 388 | 389 | 390 | 391 | 392 | 393 | 394 | 395 | 396 | 397 | 398 | 399 | 400 | | | |
|--|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|----|---|---|
| WEEKS ON STUDY | 65 | 04 | 10 | 14 | 14 | 02 | 06 | 04 | 04 | 08 | 04 | 02 | 04 | 00 | 02 | 03 | 04 | 04 | 01 | 01 | 00 | 00 | 01 | 01 | 00 | 00 | 01 | 01 | 00 | 00 | 01 | 01 | 00 | 00 | 01 | 01 | 00 | 09 | | |
| INTEGUMENTARY SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Subcutaneous tissue | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Sarcoma, NOS | + | + | + | + | + | + | + | N | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | | |
| RESPIRATORY SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Lungs and bronchi | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Undifferentiated carcinoma, metastatic | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| Trachea | + | + | + | + | + | + | + | - | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + |
| HEMATOPOIETIC SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Bone marrow | + | + | + | + | + | + | + | - | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| Spleen | - | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + |
| Lymph nodes | - | + | - | + | + | - | - | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + |
| Thymus | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | |
| CIRCULATORY SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Heart | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| DIGESTIVE SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Salivary gland | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Liver | - | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| Neoplastic nodule | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Bile duct | - | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + |
| Gallbladder & common bile duct | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | |
| Pancreas | - | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + |
| Esophagus | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + |
| Stomach | - | + | + | + | + | - | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + |
| Small intestine | - | + | + | + | + | - | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + |
| Large intestine | - | + | + | + | + | - | - | - | + | + | + | - | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + |
| URINARY SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Kidney | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Tubular cell adenoma | - | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| Urinary bladder | + | + | - | + | + | - | - | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + |
| ENDOCRINE SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Pituitary | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Adenoma, NOS | - | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| Chromophobe adenoma | X | X | | | | X | X | X | X | X | X | X | X | X | X | X | X | X | X | X | X | X | X | X | X | X | X | X | X | X | X | X | X | X | X | X | X | X | | |
| Adrenal | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Cortical adenoma | - | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| Cortical carcinoma | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Pheochromocytoma | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Thyroid | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| Parathyroid | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| REPRODUCTIVE SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Mammary gland | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Undifferentiated carcinoma | N | + | + | + | + | N | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | | |
| Adenoma, NOS | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Adenocarcinoma, NOS | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Fibroma | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Fibroadenoma | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Uterus | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| Sarcoma, NOS | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Endometrial stromal polyp | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Ovary | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| NERVOUS SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Brain | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Granular cell tumor, NOS | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| ALL OTHER SYSTEMS | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Multiple organs, NOS | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | | |
| Leukemia, NOS | X | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |

TABLE D2. INDIVIDUAL ANIMAL TUMOR PATHOLOGY OF FEMALE AUGUST RATS: VEHICLE CONTROL (Continued)

| ANIMAL NUMBER | 4 5 6 | 4 5 6 | 4 6 2 | 4 6 4 | 4 6 8 | 4 7 2 | 4 8 4 | 4 8 5 | 4 8 6 | 4 8 7 | 4 8 8 | 4 8 8 | 4 9 9 | 5 9 9 | 5 0 1 | 5 1 1 | 5 1 1 | 5 1 1 | 5 1 1 | 5 1 1 | 5 2 1 | 5 2 4 | 5 2 6 | 5 2 9 | 5 2 9 | 5 3 1 | 5 3 3 | 5 3 3 | TOTAL TISSUES TUMORS |
|--------------------------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|----------------------------|
| WEEKS ON STUDY | 0 7 6 | 0 9 4 | 1 0 4 | 1 0 4 | 1 5 2 | 1 0 4 | 1 0 4 | 1 0 3 | 1 0 1 | 1 0 4 | 1 0 4 | 1 0 4 | 1 0 0 | 1 0 1 | 1 0 3 | 1 0 6 | 1 0 9 | 1 0 4 | 1 0 4 | 1 0 1 | 1 0 4 | 1 0 9 | 1 0 5 | 1 0 9 | 1 0 9 | 1 0 9 | 1 0 0 | 1 0 4 | |
| INTEGUMENTARY SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Subcutaneous tissue | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Sarcoma, NOS | X | N | + | + | + | + | + | + | + | N | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | *50 1 |
| RESPIRATORY SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Lungs and bronchi | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Undiff. carcinoma, metastatic | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Trachea | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| HEMATOPOIETIC SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Bone marrow | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Spleen | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Lymph nodes | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Thymus | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| CIRCULATORY SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Heart | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| DIGESTIVE SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Salivary gland | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Liver | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Neoplastic nodule | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Bile duct | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Gallbladder & common bile duct | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Pancreas | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Esophagus | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Stomach | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Small intestine | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Large intestine | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| URINARY SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Kidney | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Tubular cell adenoma | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Urinary bladder | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| ENDOCRINE SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Pituitary | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Adenoma, NOS | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Chromophobe adenoma | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Adrenal | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Cortical adenoma | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Cortical carcinoma | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Pheochromocytoma | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Thyroid | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Parathyroid | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| REPRODUCTIVE SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Mammary gland | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Undifferentiated carcinoma | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Adenoma, NOS | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Adenocarcinoma, NOS | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Fibroma | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Fibroadenoma | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Uterus | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Sarcoma, NOS | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Endometrial stromal polyp | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Ovary | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| NERVOUS SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Brain | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Granular cell tumor, NOS | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| ALL OTHER SYSTEMS | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Multiple organs, NOS | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Leukemia, NOS | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |

* Animals necropsied

TABLE D2. INDIVIDUAL ANIMAL TUMOR PATHOLOGY OF FEMALE AUGUST RATS IN THE TWO-YEAR GAVAGE STUDY OF TRICHLOROETHYLENE: LOW DOSE

| ANIMAL NUMBER | 3/4/8 | 3/6/6 | 3/7/3 | 3/7/4 | 3/7/6 | 3/8/2 | 3/8/4 | 3/8/5 | 3/8/6 | 3/8/7 | 3/8/9 | 3/9/0 | 4/0/4 | 4/0/5 | 4/0/8 | 4/0/9 | 4/1/3 | 4/1/6 | 4/1/1 | 4/1/7 | 4/2/2 | 4/2/2 | 4/2/3 | 4/2/3 | 4/2/4 | 4/2/4 | 4/2/5 | | |
|--------------------------------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|---|---|
| WEEKS ON STUDY | 0/2 | 0/4 | 1/4 | 0/0 | 1/4 | 1/4 | 1/4 | 1/4 | 1/4 | 1/4 | 0/4 | 0/7 | 1/4 | 1/4 | 1/4 | 0/6 | 1/2 | 1/4 | 1/4 | 0/2 | 1/4 | 1/4 | 1/4 | 1/4 | 1/4 | 1/4 | 1/4 | | |
| INTEGUMENTARY SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Skin | N | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| Malignant melanoma | | | | | | | | | | | | | | | | | X | | | | | | | | | | | | |
| RESPIRATORY SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Lungs and bronchi | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + |
| Alveolar/bronchiolar adenoma | | | | | | | | | | | | | | | | | | X | | | | | | | | | | | |
| Trachea | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + |
| HEMATOPOIETIC SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Bone marrow | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + |
| Spleen | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + |
| Lymph nodes | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + |
| Thymus | - | + | - | - | - | - | - | - | - | - | + | - | - | - | - | - | - | - | - | - | + | - | - | - | - | - | - | - | - |
| CIRCULATORY SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Heart | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + |
| DIGESTIVE SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Salivary gland | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + |
| Liver | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + |
| Bile duct | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + |
| Gallbladder & common bile duct | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N |
| Pancreas | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + |
| Esophagus | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + |
| Stomach | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + |
| Small intestine | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + |
| Large intestine | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + |
| URINARY SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Kidney | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + |
| Tubular cell adenoma | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Tubular cell adenocarcinoma | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Urinary bladder | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + |
| ENDOCRINE SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Pituitary | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + |
| Adenoma, NOS | | X | | | X | X | X | | X | | | | X | X | | X | X | X | X | | | | | | | X | X | X | X |
| Adrenal | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + |
| Cortical adenoma | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Pheochromocytoma | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Thyroid | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + |
| Follicular cell adenoma | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| C-cell adenoma | | X | | | | | | | | | | | | | | | | | | | | | | | | | | | X |
| C-cell carcinoma | | | | | | | | X | | | | | | | | | | | | | | | | | | | | | |
| Parathyroid | - | - | - | + | - | + | - | X | - | + | - | - | + | - | + | + | - | + | + | - | + | + | - | + | - | + | - | + | + |
| Pancreatic islets | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + |
| Islet cell adenoma | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Islet cell carcinoma | | | | | | | | | | | | | | | | | | | | | | | | | | | | | X |
| REPRODUCTIVE SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Mammary gland | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + |
| Adenoma, NOS | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Fibrosarcoma | | | | | | X | | | X | | | | | | | | X | X | | | | | | X | | X | X | | |
| Uterus | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + |
| Leiomyosarcoma | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Ovary | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + |
| Luteoma | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| NERVOUS SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Nerves | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N |
| Neurofibrosarcoma | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Brain | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + |
| Granular cell tumor, NOS | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Granular cell tumor, malignant | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Astrocytoma | | | | | | | | | X | | | | | | | | X | | | | | | | | | X | | | |
| ALL OTHER SYSTEMS | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Multiple organs, NOS | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N |
| Malignant lymphoma, NOS | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |

TABLE D2. INDIVIDUAL ANIMAL TUMOR PATHOLOGY OF FEMALE AUGUST RATS: LOW DOSE
(Continued)

| ANIMAL NUMBER | 4 8 | 4 9 | 4 1 | 4 5 | 4 7 | 4 0 | 4 1 | 4 0 | 4 4 | 4 6 | 4 2 | 4 3 | 4 9 | 4 0 | 4 1 | 4 3 | 4 3 | 5 4 | 5 0 | 5 2 | 5 2 | 5 3 | 5 3 | 5 3 | 5 3 | 5 4 | 5 8 | 5 1 | TOTAL: TISSUES TUMORS | |
|--------------------------------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|-----------------------------|----------|
| WEEKS ON STUDY | 0 2 | 1 4 | 1 4 | 0 8 | 1 4 | 0 7 | 1 4 | 0 4 | 0 2 | 0 4 | 0 4 | 0 0 | 0 0 | 0 4 | 0 8 | 0 6 | 0 4 | 0 9 | 0 4 | 0 2 | 0 0 | 0 1 | 0 0 | 0 9 | 0 4 | 0 4 | 0 3 | 0 0 | | |
| INTEGUMENTARY SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Skin | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Malignant melanoma | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | *50 1 |
| RESPIRATORY SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Lungs and bronchi | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Alveolar/bronchiolar adenoma | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | 50 1 |
| Trachea | + | + | + | + | + | - | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | - | + | + | + | + | 47 |
| HEMATOPOIETIC SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Bone marrow | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | 49 |
| Spleen | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | 48 |
| Lymph nodes | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | 49 |
| Thymus | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | 4 |
| CIRCULATORY SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Heart | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | 50 |
| DIGESTIVE SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Salivary gland | + | + | + | + | + | - | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | 47 |
| Liver | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | 48 |
| Bile duct | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | 48 |
| Gallbladder & common bile duct | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | *50 |
| Pancreas | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | 48 |
| Esophagus | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | 49 |
| Stomach | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | 48 |
| Small intestine | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | 47 |
| Large intestine | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | 48 |
| URINARY SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Kidney | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | 48 |
| Tubular cell adenoma | X | | | | X | | | | | | | | | | | | | | | | | | | | | | | | 2 | |
| Tubular cell adenocarcinoma | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | 2 |
| Urinary bladder | + | + | + | + | + | + | - | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | 47 |
| ENDOCRINE SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Pituitary | + | + | + | + | + | + | + | + | - | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | 47 |
| Adenoma, NOS | X | X | X | X | X | X | | | | X | X | X | X | X | | | | | X | | | | | | | X | X | | 28 | |
| Adrenal | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | 48 |
| Cortical adenoma | | | X | | | | | | | | | | | | | | | | | | | | | | | | | | 6 | |
| Pheochromocytoma | | | | | | X | | | | | | | | | | | | | | | | | | | | | | | 2 | |
| Thyroid | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | 49 |
| Follicular cell adenoma | | | | | | | | | | | | | | | | | | | | | | | | | | | | | 2 | |
| C-cell adenoma | | | | | | | | | | | | | | | | | | | | | | | | | | X | | | 3 | |
| C-cell carcinoma | | | | | | | | | | | | | | | | | | | | | | | | | | | | | 1 | |
| Parathyroid | + | - | - | - | - | + | - | + | - | + | + | + | - | - | - | + | + | - | - | - | + | + | - | - | + | + | + | + | 23 | |
| Pancreatic islets | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | 48 |
| Islet cell adenoma | | | | | | | | | | | | | | | | | | | | | | | | | | | | | 1 | |
| Islet cell carcinoma | | | | | | | | | | | | | | | | | | | | | | | | | | | | | 1 | |
| REPRODUCTIVE SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Mammary gland | + | + | + | + | + | + | N | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | *50 |
| Adenoma, NOS | | | | | | | | | | | | X | | | | | | | | | | | | | | | | | 4 | |
| Fibroadenoma | X | X | | | | | | | | | | | | | | | | | | | | | | | | | | X | 8 | |
| Uterus | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | 48 |
| Leiomyosarcoma | | | | | | | | | | | | | | | | | | | | | | | | | | | | | 1 | |
| Ovary | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | 47 |
| Luteoma | | | | | | | | | | | | | | | | | | | | | | | | | | | | X | 1 | |
| NERVOUS SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Nerves | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | *50 |
| Neurofibrosarcoma | | | | | | | | | | | | | | | | | | | | | | | | | | | | X | 1 | |
| Brain | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | 50 |
| Granular cell tumor, NOS | | | | | | | | | | | | | | | | | | | | | | | | | | | | | 1 | |
| Granular cell tumor, malignant | | | | | | | | | | | | | | | | | | | | | | | | | | | | | 2 | |
| Astrocytoma | | | | | | | | | | | | X | | | | | | | | | | | | | | | | | 1 | |
| ALL OTHER SYSTEMS | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Multiple organs, NOS | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | *50 |
| Malignant lymphoma, NOS | | | | | | | | | | | | | | | | | | | | | | | | | | | | | 2 | |

* Animals necropsied

TABLE D2. INDIVIDUAL ANIMAL TUMOR PATHOLOGY OF FEMALE AUGUST RATS IN THE TWO-YEAR GAVAGE STUDY OF TRICHLOROETHYLENE: HIGH DOSE

| ANIMAL NUMBER | 3 | 3 | 3 | 3 | 3 | 3 | 3 | 3 | 3 | 3 | 3 | 3 | 3 | 3 | 3 | 3 | 3 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | |
|--------------------------------|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|--|
| WEEKS ON STUDY | 4 | 4 | 4 | 4 | 5 | 5 | 5 | 6 | 6 | 6 | 7 | 7 | 8 | 8 | 8 | 9 | 9 | 9 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | |
| | 3 | 4 | 9 | 0 | 2 | 6 | 1 | 3 | 9 | 7 | 8 | 0 | 1 | 3 | 1 | 8 | 9 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | |
| RESPIRATORY SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Lungs and bronchi | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| Trachea | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| HEMATOPOIETIC SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Bone marrow | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| Spleen | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| Lymph nodes | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| Thymus | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | |
| CIRCULATORY SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Heart | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| DIGESTIVE SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Salivary gland | + | + | + | + | + | + | - | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| Liver | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| Leukemia, NOS | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Bile duct | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| Gallbladder & common bile duct | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | |
| Pancreas | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| Esophagus | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| Stomach | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| Small intestine | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| Large intestine | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| URINARY SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Kidney | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| Urinary bladder | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| ENDOCRINE SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Pituitary | - | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| Adenoma, NOS | | X | X | | | X | | | | | | | | | | | | | | | | | | | | | | |
| Meningioma | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Adrenal | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| Cortical adenoma | | | | | | | X | | X | | | | | | | | | | | | | | | | | | | |
| Thyroid | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| C-cell adenoma | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Parathyroid | - | + | + | - | - | + | - | + | - | - | - | - | - | - | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| Pancreatic islets | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| Islet cell adenoma | | | | | | | X | | | | | | | | | | | | | | | | | | | | | |
| REPRODUCTIVE SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Mammary gland | + | + | + | N | + | + | + | + | + | + | + | N | + | N | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| Adenocarcinoma, NOS | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Fibroadenoma | | | | X | | | | | | | | | | | | | | | | | | | | | | | | |
| Preputial/clitoral gland | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | |
| Carcinoma, NOS | | | | | | | X | | | | | | | | | | | | | | | | | | | | | |
| Uterus | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| Ovary | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| NERVOUS SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Brain | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| Granular cell tumor, NOS | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Ependymoma | | | | | | | X | | | | | | | | | | | | | | | | | | | | | |
| ALL OTHER SYSTEMS | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Multiple organs, NOS | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | |
| Leukemia, NOS | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Monocytic leukemia | | | | | | | | | | | | | | | | | | | | | | | | | | | X | |

**TABLE D2. INDIVIDUAL ANIMAL TUMOR PATHOLOGY OF FEMALE AUGUST RATS: HIGH DOSE
(Continued)**

| ANIMAL NUMBER | 4 2 0 | 4 2 5 | 4 2 9 | 4 3 6 | 4 4 3 | 4 5 0 | 4 6 6 | 4 6 7 | 4 6 9 | 4 7 7 | 4 8 0 | 4 8 1 | 4 8 8 | 4 9 2 | 4 9 5 | 5 0 0 | 5 0 2 | 5 0 4 | 5 1 6 | 5 2 8 | 5 3 5 | 5 3 7 | 5 3 9 | 5 4 9 | 5 4 0 | TOTAL: TISSUES TUMORS |
|--------------------------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-----------------------------|
| WEEKS ON STUDY | 0 4 9 | 1 0 4 | 0 7 1 | 0 1 9 | 1 1 4 | 1 0 4 | 1 0 4 | 1 0 4 | 1 0 4 | 1 0 4 | 0 3 3 | 1 0 4 | 1 0 4 | 0 8 0 | 0 2 9 | 0 5 6 | 1 0 4 | 1 0 2 | 0 8 4 | 1 0 4 | 1 0 4 | 0 5 6 | 1 0 4 | 1 0 4 | 1 0 4 | |
| RESPIRATORY SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Lungs and bronchi | + | | | | | | | | | | | | | | | | | | | | | | | | 50 | |
| Trachea | + | | | | | | | | | | | | | | | | | | | | | | | | 47 | |
| HEMATOPOIETIC SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Bone marrow | + | | | | | | | | | | | | | | | | | | | | | | | | 50 | |
| Spleen | + | | | | | | | | | | | | | | | | | | | | | | | | 49 | |
| Lymph nodes | + | | | | | | | | | | | | | | | | | | | | | | | | 44 | |
| Thymus | + | | | | | | | | | | | | | | | | | | | | | | | | 10 | |
| CIRCULATORY SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Heart | + | | | | | | | | | | | | | | | | | | | | | | | | 50 | |
| DIGESTIVE SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Salivary gland | + | | | | | | | | | | | | | | | | | | | | | | | | 47 | |
| Liver | + | | | | | | | | | | | | | | | | | | | | | | | | 50 | |
| Leukemia, NOS | + | | | | | | | | | | | | | | | | | | | | | | | | 2 | |
| Bile duct | + | | | | | | | | | | | | | | | | | | | | | | | | 50 | |
| Gallbladder & common bile duct | N | | | | | | | | | | | | | | | | | | | | | | | | *50 | |
| Pancreas | + | | | | | | | | | | | | | | | | | | | | | | | | 48 | |
| Esophagus | + | | | | | | | | | | | | | | | | | | | | | | | | 49 | |
| Stomach | + | | | | | | | | | | | | | | | | | | | | | | | | 48 | |
| Small intestine | + | | | | | | | | | | | | | | | | | | | | | | | | 46 | |
| Large intestine | + | | | | | | | | | | | | | | | | | | | | | | | | 47 | |
| URINARY SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Kidney | + | | | | | | | | | | | | | | | | | | | | | | | | 50 | |
| Urinary bladder | + | | | | | | | | | | | | | | | | | | | | | | | | 44 | |
| ENDOCRINE SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Pituitary | + | | | | | | | | | | | | | | | | | | | | | | | | 43 | |
| Adenoma, NOS | X | | | | | | | | | | | | | | | | | | | | | | | | 18 | |
| Meningioma | + | | | | | | | | | | | | | | | | | | | | | | | | 1 | |
| Adrenal | + | | | | | | | | | | | | | | | | | | | | | | | | 50 | |
| Cortical adenoma | X | | | | | | | | | | | | | | | | | | | | | | | | 5 | |
| Thyroid | + | | | | | | | | | | | | | | | | | | | | | | | | 50 | |
| C-cell adenoma | X | | | | | | | | | | | | | | | | | | | | | | | | 1 | |
| Parathyroid | + | | | | | | | | | | | | | | | | | | | | | | | | 28 | |
| Pancreatic islets | + | | | | | | | | | | | | | | | | | | | | | | | | 48 | |
| Islet cell adenoma | + | | | | | | | | | | | | | | | | | | | | | | | | 1 | |
| REPRODUCTIVE SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Mammary gland | N | | | | | | | | | | | | | | | | | | | | | | | | *50 | |
| Adenocarcinoma, NOS | + | | | | | | | | | | | | | | | | | | | | | | | | 1 | |
| Fibroadenoma | X | | | | | | | | | | | | | | | | | | | | | | | | 4 | |
| Preputial/clitoral gland | N | | | | | | | | | | | | | | | | | | | | | | | | *50 | |
| Carcinoma, NOS | N | | | | | | | | | | | | | | | | | | | | | | | | 1 | |
| Uterus | + | | | | | | | | | | | | | | | | | | | | | | | | 49 | |
| Ovary | + | | | | | | | | | | | | | | | | | | | | | | | | 46 | |
| NERVOUS SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Brain | + | | | | | | | | | | | | | | | | | | | | | | | | 50 | |
| Granular cell tumor, NOS | X | | | | | | | | | | | | | | | | | | | | | | | | 1 | |
| Ependymoma | + | | | | | | | | | | | | | | | | | | | | | | | | 1 | |
| ALL OTHER SYSTEMS | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Multiple organs, NOS | N | | | | | | | | | | | | | | | | | | | | | | | | *50 | |
| Leukemia, NOS | X | | | | | | | | | | | | | | | | | | | | | | | | 1 | |
| Monocytic leukemia | X | | | | | | | | | | | | | | | | | | | | | | | | 2 | |

* Animals necropsied

TABLE D3. ANALYSIS OF PRIMARY TUMORS IN FEMALE AUGUST RATS IN THE TWO-YEAR GAVAGE STUDY OF TRICHLOROETHYLENE

| | Untreated Control | Vehicle Control | 500 mg/kg | 1,000 mg/kg |
|---|-------------------|-----------------|-------------|-------------|
| Hematopoietic System: Leukemia | | | | |
| Overall Rates (a) | 0/50 (0%) | 1/50 (2%) | 0/50 (0%) | 5/50 (10%) |
| Adjusted Rates (b) | 0.0% | 2.2% | 0.0% | 18.2% |
| Terminal Rates (c) | 0/26 (0%) | 0/23 (0%) | 0/26 (0%) | 3/25 (12%) |
| Life Table Tests (d) | | P=0.027 | P=0.523N | P=0.078 |
| Incidental Tumor Tests (d) | | P=0.020 | P=0.469N | P=0.059 |
| Cochran-Armitage Trend Test (d) | | P=0.037 | | |
| Fisher Exact Test | | | P=0.500N | P=0.102 |
| Kidney: Tubular Cell Adenoma or Adenocarcinoma | | | | |
| Overall Rates (a) | 0/50 (0%) | 1/49 (2%) | 4/48 (8%) | 0/50 (0%) |
| Adjusted Rates (b) | 0.0% | 4.3% | 13.6% | 0.0% |
| Terminal Rates (c) | 0/26 (0%) | 1/23 (4%) | 2/26 (8%) | 0/25 (0%) |
| Life Table Tests (d) | | P=0.390N | P=0.210 | P=0.483N |
| Incidental Tumor Tests (d) | | P=0.570N | P=0.123 | P=0.483N |
| Cochran-Armitage Trend Test (d) | | P=0.384N | | |
| Fisher Exact Test | | | P=0.174 | P=0.495N |
| Pituitary Gland: Chromophobe Adenoma | | | | |
| Overall Rates (a) | 0/49 (0%) | 3/49 (6%) | 0/47 (0%) | 0/43 (0%) |
| Adjusted Rates (b) | 0.0% | 6.4% | 0.0% | 0.0% |
| Terminal Rates (c) | 0/26 (0%) | 0/23 (0%) | 0/26 (0%) | 0/25 (0%) |
| Life Table Tests (d) | | P=0.062N | P=0.150N | P=0.197N |
| Incidental Tumor Tests (d) | | P=0.015N | P=0.058N | P=0.053N |
| Cochran-Armitage Trend Test (d) | | P=0.044N | | |
| Fisher Exact Test | | | P=0.129N | P=0.147N |
| Pituitary Gland: Adenoma | | | | |
| Overall Rates (a) | 42/49 (86%) | 35/49 (71%) | 26/47 (55%) | 18/43 (42%) |
| Adjusted Rates (b) | 97.7% | 84.9% | 76.0% | 66.6% |
| Terminal Rates (c) | 25/26 (96%) | 17/23 (74%) | 18/26 (69%) | 16/25 (64%) |
| Life Table Tests (d) | | P=0.001N | P=0.053N | P=0.003N |
| Incidental Tumor Tests (d) | | P=0.033N | P=0.173N | P=0.055N |
| Cochran-Armitage Trend Test (d) | | P=0.003N | | |
| Fisher Exact Test | | | P=0.077N | P=0.004N |
| Adrenal Gland: Cortical Adenoma | | | | |
| Overall Rates (a) | 4/50 (8%) | 6/48 (13%) | 6/48 (13%) | 5/50 (10%) |
| Adjusted Rates (b) | 14.0% | 22.4% | 23.1% | 20.0% |
| Terminal Rates (c) | 3/26 (12%) | 4/23 (17%) | 6/26 (23%) | 5/25 (20%) |
| Life Table Tests (d) | | P=0.401N | P=0.550N | P=0.480N |
| Incidental Tumor Tests (d) | | P=0.520N | P=0.604N | P=0.599 |
| Cochran-Armitage Trend Test (d) | | P=0.409N | | |
| Fisher Exact Test | | | P=0.621 | P=0.471N |
| Adrenal Gland: Cortical Adenoma or Carcinoma | | | | |
| Overall Rates (a) | 4/50 (8%) | 7/48 (15%) | 6/48 (13%) | 5/50 (10%) |
| Adjusted Rates (b) | 14.0% | 26.5% | 23.1% | 20.0% |
| Terminal Rates (c) | 3/26 (12%) | 5/23 (22%) | 6/26 (23%) | 5/25 (20%) |
| Life Table Tests (d) | | P=0.281N | P=0.420N | P=0.353N |
| Incidental Tumor Tests (d) | | P=0.384N | P=0.472N | P=0.534N |
| Cochran-Armitage Trend Test (d) | | P=0.296N | | |
| Fisher Exact Test | | | P=0.500N | P=0.351N |
| Adrenal Gland: Pheochromocytoma | | | | |
| Overall Rates (a) | 6/50 (12%) | 9/48 (19%) | 2/48 (4%) | 0/50 (0%) |
| Adjusted Rates (b) | 20.5% | 33.2% | 7.7% | 0.0% |
| Terminal Rates (c) | 4/26 (15%) | 6/23 (26%) | 2/26 (8%) | 0/25 (0%) |
| Life Table Tests (d) | | P<0.001N | P=0.019N | P=0.003N |
| Incidental Tumor Tests (d) | | P=0.001N | P=0.034N | P=0.009N |
| Cochran-Armitage Trend Test (d) | | P<0.001N | | |
| Fisher Exact Test | | | P=0.025N | P=0.001N |

TABLE D3. ANALYSIS OF PRIMARY TUMORS IN FEMALE AUGUST RATS IN THE TWO-YEAR GAVAGE STUDY OF TRICHLOROETHYLENE (Continued)

| | Untreated Control | Vehicle Control | 500 mg/kg | 1,000 mg/kg |
|---|-------------------|-----------------|-------------|-------------|
| Thyroid: C-Cell Adenoma | | | | |
| Overall Rates (a) | 0/50 (0%) | 0/49 (0%) | 3/49 (6%) | 1/50 (2%) |
| Adjusted Rates (b) | 0.0% | 0.0% | 12.0% | 4.0% |
| Terminal Rates (c) | 0/26 (0%) | 0/23 (0%) | 3/25 (12%) | 1/25 (4%) |
| Life Table Tests (d) | | P=0.403 | P=0.134 | P=0.517 |
| Incidental Tumor Tests (d) | | P=0.403 | P=0.134 | P=0.517 |
| Cochran-Armitage Trend Test (d) | | P=0.385 | | |
| Fisher Exact Test | | | P=0.121 | P=0.505 |
| Thyroid: C-Cell Adenoma or Carcinoma | | | | |
| Overall Rates (a) | 0/50 (0%) | 0/49 (0%) | 4/49 (8%) | 1/50 (2%) |
| Adjusted Rates (b) | 0.0% | 0.0% | 16.0% | 4.0% |
| Terminal Rates (c) | 0/26 (0%) | 0/23 (0%) | 4/25 (16%) | 1/25 (4%) |
| Life Table Tests (d) | | P=0.418 | P=0.071 | P=0.517 |
| Incidental Tumor Tests (d) | | P=0.418 | P=0.071 | P=0.517 |
| Cochran-Armitage Trend Test (d) | | P=0.398 | | |
| Fisher Exact Test | | | P=0.059 | P=0.505 |
| Mammary Gland: Adenoma | | | | |
| Overall Rates (a) | 3/50 (6%) | 4/50 (8%) | 4/50 (8%) | 0/50 (0%) |
| Adjusted Rates (b) | 8.2% | 12.5% | 14.4% | 0.0% |
| Terminal Rates (c) | 0/26 (0%) | 1/23 (4%) | 3/26 (12%) | 0/25 (0%) |
| Life Table Tests (d) | | P=0.069N | P=0.612N | P=0.090N |
| Incidental Tumor Tests (d) | | P=0.135N | P=0.562 | P=0.132N |
| Cochran-Armitage Trend Test (d) | | P=0.060N | | |
| Fisher Exact Test | | | P=0.643 | P=0.059N |
| Mammary Gland: Adenoma, Adenocarcinoma, or Carcinoma | | | | |
| Overall Rates (a) | 3/50 (6%) | 6/50 (12%) | 4/50 (8%) | 1/50 (2%) |
| Adjusted Rates (b) | 8.2% | 16.3% | 14.4% | 4.0% |
| Terminal Rates (c) | 0/26 (0%) | 1/23 (4%) | 3/26 (12%) | 1/25 (4%) |
| Life Table Tests (d) | | P=0.058N | P=0.367N | P=0.097N |
| Incidental Tumor Test (d) | | P=0.079N | P=0.429N | P=0.075N |
| Cochran-Armitage Trend Test (d) | | P=0.042N | | |
| Fisher Exact Test | | | P=0.370N | P=0.056N |
| Mammary Gland: Fibroadenoma | | | | |
| Overall Rates (a) | 13/50 (26%) | 10/50 (20%) | 8/50 (16%) | 4/50 (8%) |
| Adjusted Rates (b) | 38.2% | 33.1% | 25.9% | 14.3% |
| Terminal Rates (c) | 6/26 (23%) | 4/23 (17%) | 4/26 (15%) | 3/25 (12%) |
| Life Table Tests (d) | | P=0.072N | P=0.353N | P=0.090N |
| Incidental Tumor Test (d) | | P=0.356N | P=0.563 | P=0.385N |
| Cochran-Armitage Trend Test (d) | | P=0.060 | | |
| Fisher Exact Test | | | P=0.398N | P=0.074N |
| Mammary Gland: Fibroma or Fibroadenoma | | | | |
| Overall Rates (a) | 14/50 (28%) | 12/50 (24%) | 8/50 (16%) | 4/50 (8%) |
| Adjusted Rates (b) | 39.8% | 36.9% | 25.9% | 14.3% |
| Terminal Rates (c) | 6/26 (23%) | 4/23 (17%) | 4/26 (15%) | 3/25 (12%) |
| Life Table Tests (d) | | P=0.031N | P=0.210N | P=0.045N |
| Incidental Tumor Tests (d) | | P=0.240N | P=0.512N | P=0.304N |
| Cochran-Armitage Trend Test (d) | | P=0.020N | | |
| Fisher Exact Test | | | P=0.227N | P=0.027N |
| Mammary Gland: All Tumors (e) | | | | |
| Overall Rates (a) | 17/50 (34%) | 17/50 (34%) | 12/50 (24%) | 5/50 (10%) |
| Adjusted Rates (b) | 44.9% | 45.7% | 38.1% | 18.2% |
| Terminal Rates (c) | 6/26 (23%) | 5/23 (22%) | 7/26 (27%) | 4/25 (16%) |
| Life Table Tests (d) | | P=0.008N | P=0.188N | P=0.013N |
| Incidental Tumor Tests (d) | | P=0.061N | P=0.440N | P=0.058N |
| Cochran-Armitage Trend Test (d) | | P=0.003N | | |
| Fisher Exact Test | | | P=0.189N | P=0.004N |

TABLE D3. ANALYSIS OF PRIMARY TUMORS IN FEMALE AUGUST RATS IN THE TWO-YEAR GAVAGE STUDY OF TRICHLOROETHYLENE (Continued)

| | Untreated Control | Vehicle Control | 500 mg/kg | 1,000 mg/kg |
|--|-------------------|-----------------|------------|-------------|
| Uterus: Endometrial Stromal Polyp | | | | |
| Overall Rates (a) | 1/50 (2%) | 3/49 (6%) | 0/48 (0%) | 0/49 (0%) |
| Adjusted Rates (b) | 3.8% | 13.0% | 0.0% | 0.0% |
| Terminal Rates (c) | 1/26 (4%) | 3/23 (13%) | 0/26 (0%) | 0/25 (0%) |
| Life Table Tests (d) | | P=0.030N | P=0.098N | P=0.105N |
| Incidental Tumor Tests (d) | | P=0.030N | P=0.098N | P=0.105N |
| Cochran-Armitage Trend Test (d) | | P=0.038N | | |
| Fisher Exact Test | | | P=0.125N | P=0.121N |
| Brain: Granular Cell Tumor | | | | |
| Overall Rates (a) | 2/50 (4%) | 1/49 (2%) | 3/50 (6%) | 1/50 (2%) |
| Adjusted Rates (b) | 7.7% | 4.5% | 11.5% | 4.0% |
| Terminal Rates (c) | 2/26 (8%) | 1/22 (5%) | 3/26 (12%) | 1/25 (4%) |
| Life Table Tests (d) | | P=0.567N | P=0.365 | P=0.734N |
| Incidental Tumor Tests (d) | | P=0.567N | P=0.365 | P=0.734N |
| Cochran-Armitage Trend Test (d) | | P=0.603N | | |
| Fisher Exact Test | | | P=0.316 | P=0.747N |

(a) Number of tumor-bearing animals/number of animals examined at the site

(b) Kaplan-Meier estimated tumor incidence at the end of the study after adjusting for intercurrent mortality

(c) Observed tumor incidence at terminal kill

(d) Beneath the control incidence are the P values associated with the trend test. Beneath the dosed group incidence are the P values corresponding to pairwise comparisons between that dosed group and the vehicle controls. The life table analysis regards tumors in animals dying prior to terminal kill as being (directly or indirectly) the cause of death. The incidental tumor test regards these lesions as nonfatal. The Cochran-Armitage and Fisher exact tests compare directly the overall incidence rates. A negative trend or lower incidence in a dosed group is indicated by (N).

(e) Includes undifferentiated carcinoma, adenoma, adenocarcinoma, fibroma, and fibroadenoma

TABLE D4. SUMMARY OF THE INCIDENCE OF NONNEOPLASTIC LESIONS IN FEMALE AUGUST RATS IN THE TWO-YEAR GAVAGE STUDY OF TRICHLOROETHYLENE

| | CONTROL (UNTR) | CONTROL (VEH) | LOW DOSE | HIGH DOSE |
|--------------------------------------|----------------|---------------|----------|-----------|
| ANIMALS INITIALLY IN STUDY | 50 | 50 | 50 | 50 |
| ANIMALS NECROPSIED | 50 | 50 | 50 | 50 |
| ANIMALS EXAMINED HISTOPATHOLOGICALLY | 50 | 50 | 50 | 50 |
| INTEGUMENTARY SYSTEM | | | | |
| *Skin | (50) | (50) | (50) | (50) |
| Ulcer, NOS | 1 (2%) | 1 (2%) | | |
| Inflammation, chronic | 1 (2%) | 2 (4%) | | |
| Ulcer, chronic | | 1 (2%) | | |
| Inflammation, chronic diffuse | | 1 (2%) | | |
| Inflammation, granulomatous focal | 1 (2%) | | | |
| *Subcutaneous tissue | (50) | (50) | (50) | (50) |
| Hemorrhage | | | | 1 (2%) |
| Inflammation, necrotizing | | | 1 (2%) | |
| Abscess, NOS | | | 1 (2%) | |
| Inflammation, granulomatous | | | | 1 (2%) |
| Foreign material, NOS | | | | 2 (4%) |
| RESPIRATORY SYSTEM | | | | |
| *Nasal cavity | (50) | (50) | (50) | (50) |
| Inflammation, acute suppurative | | | 1 (2%) | |
| *Tracheal lumen | (50) | (50) | (50) | (50) |
| Hemorrhage | | | 1 (2%) | 4 (8%) |
| *Larynx | (50) | (50) | (50) | (50) |
| Inflammation, chronic | | 1 (2%) | | |
| Foreign material, NOS | | 1 (2%) | | |
| #Trachea | (50) | (48) | (47) | (47) |
| Inflammation, chronic | | 1 (2%) | | |
| Foreign material, NOS | | | 1 (2%) | 4 (9%) |
| #Lung | (50) | (50) | (50) | (50) |
| Bronchiectasis | | 1 (2%) | | 1 (2%) |
| Atelectasis | | | | 1 (2%) |
| Congestion, NOS | | | 2 (4%) | 2 (4%) |
| Edema, NOS | | | | 1 (2%) |
| Hemorrhage | 1 (2%) | 2 (4%) | 4 (8%) | 3 (6%) |
| Inflammation, NOS | | | 2 (4%) | 1 (2%) |
| Inflammation, focal | 1 (2%) | 2 (4%) | 3 (6%) | 2 (4%) |
| Inflammation, multifocal | | 1 (2%) | 4 (8%) | |
| Pneumonia, aspiration | | 1 (2%) | | |
| Bronchopneumonia, acute | | 1 (2%) | 2 (4%) | |
| Inflammation, acute | 1 (2%) | | | 1 (2%) |
| Inflammation, acute focal | | 1 (2%) | | |
| Inflammation, chronic | | 1 (2%) | | 1 (2%) |
| Inflammation, chronic focal | | 3 (6%) | 3 (6%) | 3 (6%) |
| Inflammation, granulomatous | 2 (4%) | 1 (2%) | 2 (4%) | 3 (6%) |
| Inflammation, granulomatous focal | 11 (22%) | 12 (24%) | 18 (36%) | 17 (34%) |
| Foreign material, NOS | 4 (8%) | 4 (8%) | 4 (8%) | 10 (20%) |
| Hemosiderosis | | | 1 (2%) | |

TABLE D4. SUMMARY OF THE INCIDENCE OF NONNEOPLASTIC LESIONS IN FEMALE AUGUST RATS IN THE TWO-YEAR GAVAGE STUDY OF TRICHLOROETHYLENE (Continued)

| | CONTROL (UNTR) | CONTROL (VEH) | LOW DOSE | HIGH DOSE |
|-----------------------------------|----------------|---------------|----------|-----------|
| HEMATOPOIETIC SYSTEM | | | | |
| #Bone marrow | (50) | (47) | (49) | (50) |
| Hyperplasia, granulocytic | | | | 1 (2%) |
| #Spleen | (50) | (49) | (48) | (49) |
| Congestion, NOS | | 1 (2%) | 1 (2%) | 2 (4%) |
| Hemosiderosis | 3 (6%) | 1 (2%) | 2 (4%) | 2 (4%) |
| Hyperplasia, lymphoid | | 1 (2%) | | |
| Hematopoiesis | 14 (28%) | 16 (33%) | 15 (31%) | 8 (16%) |
| #Lymph node | (47) | (43) | (49) | (44) |
| Congestion, NOS | 1 (2%) | | | |
| Abscess, NOS | | | 1 (2%) | |
| #Mesenteric lymph node | (47) | (43) | (49) | (44) |
| Congestion, NOS | 1 (2%) | | 1 (2%) | |
| Hemorrhage | | 1 (2%) | | 2 (5%) |
| #Liver | (50) | (48) | (48) | (50) |
| Hematopoiesis | | | 1 (2%) | |
| #Thymus | | | (4) | (10) |
| Hemorrhage | | | 1 (25%) | 1 (10%) |
| CIRCULATORY SYSTEM | | | | |
| *Multiple organs | (50) | (50) | (50) | (50) |
| Periarteritis | | | 1 (2%) | |
| #Heart | (50) | (50) | (50) | (50) |
| Thrombosis, NOS | | | | 1 (2%) |
| Inflammation, chronic focal | | | | 1 (2%) |
| Inflammation, chronic diffuse | | | | 1 (2%) |
| *Pulmonary artery | (50) | (50) | (50) | (50) |
| Calcification, focal | | 1 (2%) | | |
| *Sup. panc-duod. artery | (50) | (50) | (50) | (50) |
| Thrombosis, NOS | | | 1 (2%) | |
| Degeneration, hyaline | 1 (2%) | | | |
| #Pancreas | (50) | (49) | (48) | (48) |
| Periarteritis | | 1 (2%) | | |
| DIGESTIVE SYSTEM | | | | |
| #Salivary gland | (50) | (48) | (47) | (47) |
| Inflammation, chronic | 1 (2%) | | | |
| Inflammation, granulomatous focal | | | | 1 (2%) |
| #Liver | (50) | (48) | (48) | (50) |
| Congestion, NOS | | | | 1 (2%) |
| Congestion, chronic passive | | 1 (2%) | | |
| Hemorrhage | 2 (4%) | | | |
| Inflammation, focal | | | 1 (2%) | 1 (2%) |
| Inflammation, acute focal | 1 (2%) | | | 1 (2%) |
| Inflammation, acute/chronic | | | 1 (2%) | |
| Inflammation, chronic | | | 1 (2%) | |
| Inflammation, chronic focal | | 2 (4%) | | |
| Necrosis, NOS | | | | 1 (2%) |
| Necrosis, focal | | 3 (6%) | 2 (4%) | 2 (4%) |
| Necrosis, central | 1 (2%) | | 1 (2%) | 1 (2%) |
| Metamorphosis, fatty | 1 (2%) | 3 (6%) | | |
| Lipoidosis | 2 (4%) | | 2 (4%) | |
| Pigmentation, NOS | | | 1 (2%) | |
| Mitotic alteration | | | | 1 (2%) |
| Focal cellular change | 1 (2%) | | | 1 (2%) |
| Clear cell change | | | | 1 (2%) |
| Hepatocytomegaly | 2 (4%) | 5 (10%) | 3 (6%) | 1 (2%) |
| Angiectasis | | | 1 (2%) | |
| Regeneration, NOS | | | 1 (2%) | 1 (2%) |
| #Liver/midlobular | (50) | (48) | (48) | (50) |
| Lipoidosis | | | 1 (2%) | |

TABLE D4. SUMMARY OF THE INCIDENCE OF NONNEOPLASTIC LESIONS IN FEMALE AUGUST RATS IN THE TWO-YEAR GAVAGE STUDY OF TRICHLOROETHYLENE (Continued)

| | CONTROL (UNTR) | CONTROL (VEH) | LOW DOSE | HIGH DOSE |
|-------------------------------------|----------------|---------------|----------|-----------|
| DIGESTIVE SYSTEM (Continued) | | | | |
| #Bile duct | (50) | (48) | (48) | (50) |
| Dilatation, NOS | 1 (2%) | | | |
| Inflammation, chronic | | 1 (2%) | | 2 (4%) |
| Fibrosis | 1 (2%) | | | |
| Sclerosis | 1 (2%) | | | |
| Hyperplasia, NOS | 1 (2%) | | 1 (2%) | 5 (10%) |
| Hyperplasia, focal | | | 1 (2%) | 1 (2%) |
| #Pancreas | (50) | (49) | (48) | (48) |
| Inflammation, acute/chronic | | | 2 (4%) | |
| Inflammation, chronic | 1 (2%) | | | 1 (2%) |
| Fibrosis, multifocal | | | 1 (2%) | |
| Atrophy, focal | 4 (8%) | 2 (4%) | 4 (8%) | 4 (8%) |
| #Pancreatic duct | (50) | (49) | (48) | (48) |
| Inflammation, acute | 1 (2%) | | | |
| #Pancreatic acinus | (50) | (49) | (48) | (48) |
| Atrophy, NOS | | 4 (8%) | 2 (4%) | |
| Atrophy, focal | 21 (42%) | 27 (55%) | 24 (50%) | 25 (52%) |
| Atrophy, diffuse | 7 (14%) | 2 (4%) | 2 (4%) | |
| *Esophageal lumen | (50) | (50) | (50) | (50) |
| Hemorrhage | | 2 (4%) | | 4 (8%) |
| #Esophagus | (50) | (49) | (49) | (49) |
| Penetrating wound | | 1 (2%) | | |
| Inflammation, NOS | | | | 1 (2%) |
| Foreign material, NOS | | | | 1 (2%) |
| Hyperkeratosis | | | 1 (2%) | |
| #Periesophageal tissue | (50) | (49) | (49) | (49) |
| Inflammation, NOS | | | | 1 (2%) |
| Inflammation, acute | | | | 1 (2%) |
| #Stomach | (50) | (48) | (48) | (48) |
| Edema, NOS | 1 (2%) | | | |
| Inflammation, NOS | | 1 (2%) | | |
| #Gastric submucosa | (50) | (48) | (48) | (48) |
| Inflammation, granulomatous | 1 (2%) | | | |
| #Forestomach | (50) | (48) | (48) | (48) |
| Hyperplasia, epithelial | | 1 (2%) | 1 (2%) | |
| Hyperkeratosis | | 1 (2%) | | |
| #Colon | (50) | (43) | (48) | (47) |
| Hemorrhage | | 1 (2%) | | |
| Parasitism | 3 (6%) | 2 (5%) | 1 (2%) | 1 (2%) |
| URINARY SYSTEM | | | | |
| #Kidney | (50) | (49) | (48) | (50) |
| Calculus, gross observation only | 1 (2%) | | | |
| Mineralization | 5 (10%) | 3 (6%) | 3 (6%) | |
| Cast, NOS | 1 (2%) | | | |
| Hydronephrosis | 4 (8%) | 1 (2%) | 3 (6%) | |
| Cyst, NOS | 2 (4%) | 2 (4%) | 1 (2%) | 1 (2%) |
| Inflammation, acute focal | | | | 1 (2%) |
| Inflammation, chronic | 1 (2%) | | | |
| Inflammation, granulomatous | | | 1 (2%) | |
| Fibrosis, diffuse | 1 (2%) | | | |
| Nephropathy | 47 (94%) | 37 (76%) | 42 (88%) | 42 (84%) |
| Nephropathy, toxic | | | 8 (17%) | 29 (58%) |
| Calcification, focal | | 1 (2%) | | 2 (4%) |
| Cytomegaly | | | 46 (96%) | 50 (100%) |
| Hyperplasia, tubular cell | | | | 1 (2%) |
| #Kidney/tubule | (50) | (49) | (48) | (50) |
| Dilatation, NOS | | 1 (2%) | | |
| Degeneration, NOS | | | 1 (2%) | |
| Atypia, NOS | | | 1 (2%) | |

TABLE D4. SUMMARY OF THE INCIDENCE OF NONNEOPLASTIC LESIONS IN FEMALE AUGUST RATS IN THE TWO-YEAR GAVAGE STUDY OF TRICHLOROETHYLENE (Continued)

| | CONTROL (UNTR) | CONTROL (VEH) | LOW DOSE | HIGH DOSE |
|-----------------------------------|----------------|---------------|----------|-----------|
| URINARY SYSTEM (Continued) | | | | |
| #Kidney/pelvis | (50) | (49) | (48) | (50) |
| Dilatation, NOS | | | | 1 (2%) |
| Hemorrhage | 3 (6%) | | 2 (4%) | |
| Hyperplasia, epithelial | 10 (20%) | 1 (2%) | 4 (8%) | 3 (6%) |
| #Urinary bladder | (48) | (46) | (47) | (44) |
| Inflammation, NOS | 1 (2%) | | | |
| Inflammation, acute diffuse | | | | 1 (2%) |
| Hyperplasia, epithelial | 1 (2%) | 1 (2%) | | 1 (2%) |
| Angiectasis | | | 1 (2%) | |
| ENDOCRINE SYSTEM | | | | |
| #Pituitary | (49) | (49) | (47) | (43) |
| Cyst, NOS | 1 (2%) | | 1 (2%) | 1 (2%) |
| Congestion, NOS | 1 (2%) | | | |
| Hyperplasia, NOS | 1 (2%) | | | |
| Hyperplasia, focal | | 1 (2%) | 4 (9%) | 3 (7%) |
| Angiectasis | | | 2 (4%) | |
| #Adrenal | (50) | (48) | (48) | (50) |
| Cyst, NOS | | 1 (2%) | | |
| Congestion, NOS | | | 2 (4%) | |
| Hemorrhage | | | 1 (2%) | |
| Necrosis, cortical | 1 (2%) | | | |
| Lipoidosis | | 2 (4%) | | |
| Angiectasis | 1 (2%) | 1 (2%) | 2 (4%) | 1 (2%) |
| #Adrenal/capsule | (50) | (48) | (48) | (50) |
| Inflammation, chronic | | | 1 (2%) | |
| #Adrenal cortex | (50) | (48) | (48) | (50) |
| Congestion, NOS | | | 1 (2%) | |
| Hemorrhage | | | 1 (2%) | |
| Inflammation, focal | | | 1 (2%) | |
| Lipoidosis | 3 (6%) | 1 (2%) | | 1 (2%) |
| Focal cellular change | | 1 (2%) | 1 (2%) | |
| Cytomegaly | | 1 (2%) | 1 (2%) | |
| Hyperplasia, NOS | 1 (2%) | 1 (2%) | | |
| Hyperplasia, focal | | 4 (8%) | 1 (2%) | 1 (2%) |
| #Adrenal medulla | (50) | (48) | (48) | (50) |
| Focal cellular change | 1 (2%) | 1 (2%) | | |
| Hyperplasia, NOS | 2 (4%) | | | |
| Hyperplasia, focal | 8 (16%) | 5 (10%) | 12 (25%) | |
| #Thyroid | (50) | (49) | (49) | (50) |
| Inflammation, chronic focal | 1 (2%) | | | |
| Hyperplasia, C-cell | 1 (2%) | 2 (4%) | 2 (4%) | 1 (2%) |
| REPRODUCTIVE SYSTEM | | | | |
| *Mammary gland | (50) | (50) | (50) | (50) |
| Cyst, NOS | | | | 1 (2%) |
| Degeneration, cystic | | | 1 (2%) | |
| Hyperplasia, NOS | 1 (2%) | | | |
| Hyperplasia, focal | | | | 1 (2%) |
| Hyperplasia, cystic | | | | 1 (2%) |
| Lactation | 30 (60%) | 21 (42%) | 20 (40%) | 7 (14%) |
| *Vaginal mucosa | (50) | (50) | (50) | (50) |
| Degeneration, NOS | 3 (6%) | 3 (6%) | | |

TABLE D4. SUMMARY OF THE INCIDENCE OF NONNEOPLASTIC LESIONS IN FEMALE AUGUST RATS IN THE TWO-YEAR GAVAGE STUDY OF TRICHLOROETHYLENE (Continued)

| | CONTROL (UNTR) | CONTROL (VEH) | LOW DOSE | HIGH DOSE |
|--|----------------|---------------|----------|-----------|
| REPRODUCTIVE SYSTEM (Continued) | | | | |
| #Uterus | (50) | (49) | (48) | (49) |
| Dilatation, NOS | | | | 4 (8%) |
| Hydrometra | 1 (2%) | 1 (2%) | 4 (8%) | 2 (4%) |
| Hemorrhage | | | 1 (2%) | |
| Inflammation, NOS | | | 1 (2%) | |
| Inflammation, acute | 1 (2%) | | | |
| Inflammation, acute/chronic | | | | 1 (2%) |
| Metaplasia, squamous | | 1 (2%) | | |
| #Uterus/endometrium | (50) | (49) | (48) | (49) |
| Cyst, NOS | | 1 (2%) | | |
| Inflammation, acute | | | | 1 (2%) |
| Degeneration, NOS | | | | 1 (2%) |
| Hyperplasia, NOS | | 1 (2%) | | |
| Hyperplasia, cystic | 1 (2%) | | | |
| #Ovary | (49) | (47) | (47) | (46) |
| Cyst, NOS | 2 (4%) | | 3 (6%) | |
| Follicular cyst, NOS | 1 (2%) | | 2 (4%) | 3 (7%) |
| NERVOUS SYSTEM | | | | |
| #Brain | (50) | (49) | (50) | (50) |
| Hydrocephalus, internal | | | | 1 (2%) |
| Hemorrhage | | 1 (2%) | 3 (6%) | 1 (2%) |
| SPECIAL SENSE ORGANS | | | | |
| *Middle ear | (50) | (50) | (50) | (50) |
| Inflammation, acute suppurative | | | 1 (2%) | |
| MUSCULOSKELETAL SYSTEM | | | | |
| None | | | | |
| BODY CAVITIES | | | | |
| *Mediastinum | (50) | (50) | (50) | (50) |
| Abscess, NOS | | | 1 (2%) | |
| *Pleura | (50) | (50) | (50) | (50) |
| Inflammation, acute/chronic | | 1 (2%) | | |
| *Pericardium | (50) | (50) | (50) | (50) |
| Inflammation, acute/chronic | | 1 (2%) | | |
| ALL OTHER SYSTEMS | | | | |
| None | | | | |
| SPECIAL MORPHOLOGY SUMMARY | | | | |
| None | | | | |

* Number of animals receiving complete necropsy examination; all gross lesions including masses examined microscopically.
 # Number of animals examined microscopically at this site

APPENDIX E

SUMMARY OF LESIONS IN MALE MARSHALL RATS IN THE TWO-YEAR GAVAGE STUDY OF TRICHLOROETHYLENE

| | PAGE | |
|----------|---|-----|
| TABLE E1 | SUMMARY OF THE INCIDENCE OF NEOPLASMS IN MALE MARSHALL RATS IN THE TWO-YEAR GAVAGE STUDY OF TRICHLOROETHYLENE | 165 |
| TABLE E2 | INDIVIDUAL ANIMAL TUMOR PATHOLOGY OF MALE MARSHALL RATS IN THE TWO-YEAR GAVAGE STUDY OF TRICHLOROETHYLENE | 168 |
| TABLE E3 | ANALYSIS OF PRIMARY TUMORS IN MALE MARSHALL RATS IN THE TWO-YEAR GAVAGE STUDY OF TRICHLOROETHYLENE | 176 |
| TABLE E4 | SUMMARY OF THE INCIDENCE OF NONNEOPLASTIC LESIONS IN MALE MARSHALL RATS IN THE TWO-YEAR GAVAGE STUDY OF TRICHLOROETHYLENE | 178 |

TABLE E1. SUMMARY OF THE INCIDENCE OF NEOPLASMS IN MALE MARSHALL RATS IN THE TWO-YEAR GAVAGE STUDY OF TRICHLOROETHYLENE

| | CONTROL (UNTR) | CONTROL (VEH) | LOW DOSE | HIGH DOSE |
|---|----------------|---------------|----------|-----------|
| ANIMALS INITIALLY IN STUDY | 50 | 50 | 50 | 50 |
| ANIMALS MISSING | 1 | | | |
| ANIMALS NECROPSIED | 49 | 50 | 50 | 50 |
| ANIMALS EXAMINED HISTOPATHOLOGICALLY | 49 | 50 | 50 | 50 |
| INTEGUMENTARY SYSTEM | | | | |
| *Skin | (49) | (50) | (50) | (50) |
| Trichoepithelioma | 2 (4%) | | 1 (2%) | 1 (2%) |
| *Subcutaneous tissue | (49) | (50) | (50) | (50) |
| Sarcoma, NOS | 1 (2%) | | | |
| Teratoma, benign | | | 1 (2%) | |
| RESPIRATORY SYSTEM | | | | |
| #Lung | (47) | (49) | (50) | (47) |
| Neoplasm, NOS, unclear primary or metas | | | 1 (2%) | |
| Alveolar/bronchiolar adenoma | 1 (2%) | | | |
| Tubular cell adenocarcinoma, metastatic | | | | 1 (2%) |
| Pheochromocytoma, metastatic | 1 (2%) | 3 (6%) | 1 (2%) | 1 (2%) |
| HEMATOPOIETIC SYSTEM | | | | |
| *Multiple organs | (49) | (50) | (50) | (50) |
| Malignant lymphoma, histiocytic type | | | | 1 (2%) |
| CIRCULATORY SYSTEM | | | | |
| #Spleen | (46) | (48) | (43) | (46) |
| Hemangiosarcoma | | | 1 (2%) | |
| #Heart | (47) | (49) | (50) | (48) |
| Tubular cell adenocarcinoma, metastatic | | | | 1 (2%) |
| Pheochromocytoma, metastatic | | 1 (2%) | | |
| DIGESTIVE SYSTEM | | | | |
| #Liver | (47) | (49) | (50) | (47) |
| Carcinoma, NOS, metastatic | | 1 (2%) | | |
| #Cecum | (41) | (45) | (43) | (43) |
| Adenocarcinoma in adenomatous polyp | 1 (2%) | | | |
| URINARY SYSTEM | | | | |
| #Kidney | (49) | (49) | (50) | (47) |
| Tubular cell adenoma | 2 (4%) | | 1 (2%) | |
| Tubular cell adenocarcinoma | | | | 1 (2%) |
| Interstitial cell tumor, invasive | | | | 1 (2%) |
| #Urinary bladder | (41) | (45) | (41) | (44) |
| Transitional cell carcinoma | 1 (2%) | | | |
| ENDOCRINE SYSTEM | | | | |
| #Pituitary | (36) | (21) | (25) | (28) |
| Chromophobe adenoma | 1 (3%) | 2 (10%) | | |
| #Adrenal | (48) | (48) | (43) | (43) |
| Cortical adenoma | | 2 (4%) | | |
| Pheochromocytoma | 19 (40%) | 21 (44%) | 12 (28%) | 11 (26%) |
| Pheochromocytoma, malignant | 4 (8%) | 5 (10%) | 1 (2%) | 2 (5%) |
| #Thyroid | (41) | (41) | (41) | (40) |
| C-cell adenoma | 8 (20%) | 4 (10%) | 5 (12%) | 2 (5%) |
| C-cell carcinoma | 4 (10%) | 1 (2%) | | 1 (3%) |

TABLE E1. SUMMARY OF THE INCIDENCE OF NEOPLASMS IN MALE MARSHALL RATS IN THE TWO-YEAR GAVAGE STUDY OF TRICHLOROETHYLENE (Continued)

| | CONTROL (UNTR) | CONTROL (VEH) | LOW DOSE | HIGH DOSE |
|-------------------------------------|----------------|---------------|----------|-----------|
| ENDOCRINE SYSTEM (Continued) | | | | |
| #Parathyroid | (15) | (16) | (21) | (18) |
| Adenoma, NOS | | 2 (13%) | | 1 (6%) |
| #Pancreatic islets | (42) | (49) | (46) | (46) |
| Islet cell adenoma | 1 (2%) | 1 (2%) | | |
| REPRODUCTIVE SYSTEM | | | | |
| #Testis | (46) | (46) | (48) | (48) |
| Interstitial cell tumor | 16 (35%) | 17 (37%) | 21 (44%) | 31 (65%) |
| Interstitial cell tumor, malignant | | | | 1 (2%) |
| NERVOUS SYSTEM | | | | |
| #Brain | (45) | (48) | (50) | (47) |
| Glioma, NOS | | | | 1 (2%) |
| Astrocytoma | | | | 1 (2%) |
| *Spinal cord | (49) | (50) | (50) | (50) |
| Meningioma | | 1 (2%) | | |
| SPECIAL SENSE ORGANS | | | | |
| None | | | | |
| MUSCULOSKELETAL SYSTEM | | | | |
| *Skull | (49) | (50) | (50) | (50) |
| Mixed mesenchymal tumor, benign | | 1 (2%) | | |
| *Vertebra | (49) | (50) | (50) | (50) |
| Meningioma, invasive | | 1 (2%) | | |
| BODY CAVITIES | | | | |
| *Mediastinum | (49) | (50) | (50) | (50) |
| Carcinoma, NOS | | 1 (2%) | | |
| *Peritoneum | (49) | (50) | (50) | (50) |
| Sarcoma, NOS | | 1 (2%) | | |
| Mesothelioma, NOS | | 2 (4%) | 1 (2%) | 1 (2%) |
| Mesothelioma, malignant | | | | 2 (4%) |
| *Peritoneal mesothelium | (49) | (50) | (50) | (50) |
| Mesothelioma, NOS | 1 (2%) | | | |
| *Tunica vaginalis | (49) | (50) | (50) | (50) |
| Mesothelioma, NOS | 1 (2%) | | 1 (2%) | |
| ALL OTHER SYSTEMS | | | | |
| None | | | | |
| ANIMAL DISPOSITION SUMMARY | | | | |
| Animals initially in study | 50 | 50 | 50 | 50 |
| Natural death | 13 | 14 | 20 | 16 |
| Moribund sacrifice | 3 | 8 | 6 | 3 |
| Terminal sacrifice | 32 | 26 | 12 | 6 |
| Dosing accident | | 1 | | |
| Accidentally killed, NOS | 1 | 1 | 12 | 25 |
| Animal missing | 1 | | | |

TABLE E1. SUMMARY OF THE INCIDENCE OF NEOPLASMS IN MALE MARSHALL RATS IN THE TWO-YEAR GAVAGE STUDY OF TRICHLOROETHYLENE (Continued)

| | CONTROL (UNTR) | CONTROL (VEH) | LOW DOSE | HIGH DOSE |
|--|----------------|---------------|----------|-----------|
| TUMOR SUMMARY | | | | |
| Total animals with primary tumors** | 38 | 37 | 23 | 32 |
| Total primary tumors | 63 | 61 | 46 | 57 |
| Total animals with benign tumors | 35 | 35 | 22 | 32 |
| Total benign tumors | 50 | 50 | 41 | 46 |
| Total animals with malignant tumors | 11 | 9 | 2 | 7 |
| Total malignant tumors | 11 | 9 | 2 | 10 |
| Total animals with secondary tumors## | 1 | 5 | 1 | 3 |
| Total secondary tumors | 1 | 6 | 1 | 4 |
| Total animals with tumors uncertain-- benign or malignant | 2 | 2 | 2 | 1 |
| Total uncertain tumors | 2 | 2 | 2 | 1 |
| Total animals with tumors uncertain-- primary or metastatic | | | 1 | |
| Total uncertain tumors | | | 1 | |

* Number of animals receiving complete necropsy examination; all gross lesions including masses examined microscopically.

** Primary tumors: all tumors except secondary tumors

Number of animals examined microscopically at this site

Secondary tumors: metastatic tumors or tumors invasive into an adjacent organ

TABLE E2. INDIVIDUAL ANIMAL TUMOR PATHOLOGY OF MALE MARSHALL RATS IN THE TWO-YEAR GAVAGE STUDY OF TRICHLOROETHYLENE: UNTREATED CONTROL

| ANIMAL NUMBER | 696 | 898 | 699 | 701 | 705 | 711 | 718 | 720 | 733 | 736 | 741 | 744 | 746 | 750 | 752 | 756 | 764 | 772 | 777 | 786 | 788 | 792 | 799 | 802 | | |
|-------------------------------------|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|----|--|
| WEEKS ON STUDY | 05 | 15 | 04 | 10 | 10 | 10 | 10 | 05 | 10 | 10 | 10 | 08 | 10 | 10 | 10 | 06 | 10 | 04 | 10 | 09 | 09 | 10 | 10 | 05 | 10 | |
| INTEGUMENTARY SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Skin | + | + | + | + | + | N | + | + | + | + | + | + | + | + | + | + | + | + | + | N | + | + | + | + | + | |
| Trichoepithelioma | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Subcutaneous tissue | + | + | + | + | + | N | + | + | + | + | + | + | + | + | + | + | + | + | + | N | + | + | + | + | + | |
| Sarcoma, NOS | | | | | | | | | | | | | X | | | | | | | | | | | | | |
| RESPIRATORY SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Lungs and bronchi | + | + | + | + | + | + | - | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| Alveolar/bronchiolar adenoma | | | | | | X | | | | | | | | | | | | | | | | | | | | |
| Pheochromocytoma, metastatic | | | | | | | | | | | | | | | | | | | | | | X | | | | |
| Trachea | + | + | - | + | + | + | + | - | + | + | - | - | + | + | + | + | + | + | + | + | + | + | - | + | + | |
| HEMATOPOIETIC SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Bone marrow | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| Spleen | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| Lymph nodes | + | + | - | - | + | + | - | + | + | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | |
| Thymus | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | |
| CIRCULATORY SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Heart | + | + | + | + | + | + | - | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| DIGESTIVE SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Salivary gland | + | + | + | + | + | + | - | + | + | + | - | + | + | + | + | + | + | + | + | + | + | + | + | - | + | |
| Liver | + | + | + | + | + | + | - | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| Bile duct | + | + | + | + | + | + | - | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| Gallbladder & common bile duct | + | + | + | + | + | + | - | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| Pancreas | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | |
| Esophagus | + | + | - | + | + | + | - | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| Stomach | + | + | + | + | + | + | - | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| Small intestine | + | + | + | + | + | + | - | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| Large intestine | + | + | - | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| Adenocarcinoma in adenomatous polyp | | | | | | | | | | | | | | | | | | | | | | | | | | |
| URINARY SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Kidney | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| Tubular cell adenoma | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Urinary bladder | + | + | - | + | + | + | + | + | + | + | + | - | + | + | + | + | + | + | + | + | + | + | + | + | - | |
| Transitional cell carcinoma | | | | | | | | | | | | | | | | | | | | | | | | | | |
| ENDOCRINE SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Pituitary | - | + | + | + | + | + | + | - | + | + | + | - | + | + | + | + | - | - | + | - | + | + | + | + | - | |
| Chromophobe adenoma | | | | | | X | | | | | | | | | | | | | | | | | | | | |
| Adrenal | + | + | + | + | + | + | + | - | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| Pheochromocytoma | X | X | | X | X | | | X | | X | X | X | X | X | X | X | X | X | X | X | X | X | X | X | X | |
| Pheochromocytoma, malignant | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Thyroid | + | + | - | + | + | + | - | + | + | + | - | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| C-cell adenoma | X | X | | X | X | | | - | + | + | + | - | X | + | + | + | + | + | + | X | + | + | + | - | | |
| C-cell carcinoma | | | | | | X | | | | | | | | | | | | | | | | | | X | | |
| Parathyroid | + | - | - | + | + | - | + | - | + | + | - | - | + | - | - | - | - | - | - | - | - | - | + | + | | |
| Pancreatic islets | + | + | - | + | + | + | + | - | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| Islet cell adenoma | | | | | | | | | | X | | | | | | | | | | | | | | | | |
| REPRODUCTIVE SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Mammary gland | + | + | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | |
| Testis | + | + | - | + | + | + | + | + | + | + | + | - | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| Interstitial cell tumor | | | | X | | | | | | | | | X | | | | | | | | X | | X | | | |
| Prostate | + | + | - | + | + | + | + | + | + | + | + | - | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| NERVOUS SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Brain | + | + | + | + | + | + | + | - | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| BODY CAVITIES | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Peritoneum | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | |
| Mesothelioma, NOS | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Tunica vaginalis | + | + | N | + | + | + | + | + | + | + | + | + | N | + | + | + | + | + | + | + | + | + | + | + | + | |
| Mesothelioma, NOS | | | | | | | | | | | | | | | | | | | | | | | | | | |

+: Tissue examined microscopically
 -: Required tissue not examined microscopically
 X: Tumor incidence
 N: Necropsy, no autolysis, no microscopic examination
 S: Animal missexed

: No tissue information submitted
 C: Necropsy, no histology due to protocol
 A: Autolysis
 M: Animal missing
 B: No necropsy performed

TABLE E2. INDIVIDUAL ANIMAL TUMOR PATHOLOGY OF MALE MARSHALL RATS IN THE TWO-YEAR GAVAGE STUDY OF TRICHLOROETHYLENE: VEHICLE CONTROL

| ANIMAL NUMBER | 693 | 694 | 695 | 697 | 700 | 701 | 702 | 703 | 704 | 705 | 706 | 707 | 708 | 709 | 710 | 711 | 712 | 713 | 714 | 715 | 716 | 717 | 718 | 719 | 720 | | |
|---------------------------------|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|--|
| WEEKS ON STUDY | 105 | 088 | 105 | 105 | 105 | 007 | 105 | 105 | 105 | 002 | 105 | 105 | 105 | 009 | 105 | 006 | 009 | 105 | 105 | 105 | 105 | 105 | 105 | 105 | 009 | 007 | |
| RESPIRATORY SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Lungs and bronchi | + | + | + | + | + | + | + | + | + | - | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| Pheochromocytoma, metastatic | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| Trachea | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | - | - | |
| HEMATOPOIETIC SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Bone marrow | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| Spleen | + | + | + | + | + | + | + | + | + | - | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| Lymph nodes | + | - | - | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| Thymus | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | |
| CIRCULATORY SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Heart | + | + | + | + | + | + | + | + | + | - | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| Pheochromocytoma, metastatic | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| DIGESTIVE SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Salivary gland | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| Liver | + | + | + | + | + | + | + | + | + | - | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| Carcinoma, NOS, metastatic | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| Bile duct | + | + | + | + | + | + | + | + | + | - | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| Gallbladder & common bile duct | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | |
| Pancreas | + | + | + | + | + | + | + | + | + | - | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| Esophagus | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| Stomach | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| Small intestine | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| Large intestine | + | + | + | + | + | + | + | + | + | - | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| URINARY SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Kidney | + | + | + | + | + | + | + | + | + | - | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| Urinary bladder | + | + | + | + | + | + | - | + | + | - | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| ENDOCRINE SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Pituitary | + | + | - | - | - | + | + | - | - | + | - | - | - | + | + | + | - | - | - | - | - | - | - | + | - | | |
| Chromophobe adenoma | | X | | | | | | | | | | | | | X | | | | | | | | | | | | |
| Adrenal | + | + | + | + | + | + | + | + | + | - | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | | |
| Cortical adenoma | | | | | | | | | | | | | | | | | | | | | | | X | | | | |
| Pheochromocytoma | X | X | X | X | | | X | | | X | X | | | | | | X | X | X | X | | X | X | X | X | | |
| Pheochromocytoma, malignant | | | | | | | | X | | | | | | | | | | | | | | | | | | | |
| Thyroid | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | | |
| C-cell adenoma | | | | | | | | | | | | | | | | | | | | | | X | X | X | | | |
| C-cell carcinoma | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Parathyroid | - | - | - | + | + | + | + | - | - | - | - | - | - | - | + | + | - | + | - | + | + | - | - | - | - | | |
| Adenoma, NOS | | | | X | | X | | | | | | | | | | | | | | | | | | | | | |
| Pancreatic islets | + | + | + | + | + | + | + | + | + | - | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | | |
| Islet cell adenoma | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| REPRODUCTIVE SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Mammary gland | + | N | N | N | + | N | N | N | N | N | + | N | + | N | N | N | + | + | N | N | N | N | N | N | N | | |
| Testis | + | + | + | + | + | + | + | + | + | - | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | | |
| Interstitial cell tumor | | X | X | | | | X | | | X | X | X | | | | | X | X | X | X | | X | X | X | X | | |
| Prostate | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | | |
| NERVOUS SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Brain | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | | |
| Spinal cord | + | + | + | + | + | + | + | + | + | N | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | | |
| Meningioma | | | | | | | | | | | | | | | | | | | | | | X | | | | | |
| MUSCULOSKELETAL SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Bone | N | N | N | N | N | + | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | | |
| Mixed mesenchymal tumor, benign | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Meningioma, invasive | | | | | | | | | | | | | | | | | | | | | | | X | | | | |
| BODY CAVITIES | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Mediastinum | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | | |
| Carcinoma, NOS | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Peritoneum | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | | |
| Sarcoma, NOS | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Mesothelioma, NOS | | | | | | | | | | | | | | | | | | | | | | | | | X | | |

TABLE E2. INDIVIDUAL ANIMAL TUMOR PATHOLOGY OF MALE MARSHALL RATS IN THE TWO-YEAR GAVAGE STUDY OF TRICHLOROETHYLENE: LOW DOSE

| ANIMAL NUMBER | 7 0 | 7 0 | 7 0 | 7 0 | 7 1 | 7 1 | 7 1 | 7 2 | 7 2 | 7 2 | 7 3 | 7 3 | 7 3 | 7 4 | 7 4 | 7 4 | 7 4 | 7 5 | 7 5 | 7 5 | 7 6 | 7 6 | 7 8 | 7 8 | 7 8 | | |
|--|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|--|
| WEEKS ON STUDY | 0 8 8 | 1 0 4 | 0 1 1 | 0 0 4 | 1 6 9 | 1 0 4 | 1 6 5 | 0 0 4 | 1 0 0 | 1 1 0 | 2 0 0 | 3 0 3 | 3 0 4 | 3 0 9 | 4 0 0 | 4 0 4 | 4 0 7 | 4 0 2 | 5 0 0 | 5 0 5 | 5 0 7 | 6 1 1 | 6 0 4 | 6 0 4 | 8 0 6 | 8 0 2 | |
| INTEGUMENTARY SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Skin | + | + | + | + | + | + | + | N | + | + | + | N | + | + | N | N | + | N | + | + | N | + | + | + | + | + | |
| Trichoepithelioma | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Subcutaneous tissue | + | + | + | + | + | + | + | N | + | + | + | N | + | + | N | N | + | N | + | + | N | + | + | + | + | + | |
| Teratoma, benign | | | | | | | | | | | | | X | | | | | | | | | | | | X | | |
| RESPIRATORY SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Lungs and bronchi | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| Neoplasm, NOS, unclear primary or metastatic | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Pheochromocytoma, metastatic | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Trachea | + | + | + | + | - | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| HEMATOPOIETIC SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Bone marrow | + | + | - | + | + | + | + | + | - | - | + | + | + | - | - | - | + | + | + | + | - | + | - | + | + | | |
| Spleen | + | - | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | | |
| Hemangiosarcoma | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Lymph nodes | - | - | + | + | - | - | + | - | - | - | + | - | - | - | - | - | + | + | + | + | - | + | - | + | + | | |
| Thymus | - | - | + | - | + | - | + | - | + | - | + | - | - | - | - | + | + | + | + | - | + | - | - | + | + | | |
| CIRCULATORY SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Heart | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| DIGESTIVE SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Salivary gland | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| Liver | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| Bile duct | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| Gallbladder & common bile duct | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | |
| Pancreas | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| Esophagus | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| Stomach | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| Small intestine | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| Large intestine | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| URINARY SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Kidney | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| Tubular cell adenoma | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Urinary bladder | + | + | + | + | + | - | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| ENDOCRINE SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Pituitary | + | + | - | + | + | + | + | - | - | - | + | + | - | - | - | + | - | - | - | - | - | + | + | + | - | | |
| Adrenal | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | | |
| Pheochromocytoma | | | | | X | | X | X | X | | | | X | | | | | | | | | | | | | | |
| Pheochromocytoma, malignant | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Thyroid | - | + | + | + | - | + | + | X | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| C-cell adenoma | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Parathyroid | - | - | + | - | - | + | + | X | + | - | + | + | X | - | - | + | - | - | - | - | - | - | + | + | + | | |
| REPRODUCTIVE SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Mammary gland | + | + | N | + | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | + | N | N | + | + | N | |
| Testis | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| Interstitial cell tumor | X | X | | X | | | X | X | | | | X | X | X | | | | | | | X | | X | X | X | | |
| Prostate | + | + | + | + | + | + | + | + | - | - | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| NERVOUS SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Brain | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| BODY CAVITIES | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Peritoneum | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | |
| Mesothelioma, NOS | | | | | X | | | | | | | | | | | | | | | | | | | | | | |
| Tunica vaginalis | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| Mesothelioma, NOS | | | | | | | | | | | | | | | | | | | | | | | | | | X | |

TABLE E2. INDIVIDUAL ANIMAL TUMOR PATHOLOGY OF MALE MARSHALL RATS: LOW DOSE
(Continued)

| ANIMAL NUMBER | 7 8 | 7 9 | 7 9 | 8 0 | 8 0 | 8 0 | 8 0 | 8 1 | 8 1 | 8 1 | 8 2 | 8 2 | 8 2 | 8 3 | 8 3 | 8 3 | 8 4 | 8 4 | 8 5 | 8 5 | 8 6 | 8 6 | 8 7 | 8 7 | 8 7 | 8 8 | 8 8 | 8 9 | 8 9 | TOTAL TISSUES TUMORS |
|-------------------------------------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|----------------------|
| WEEKS ON STUDY | 0 1 | 0 1 | 1 4 | 0 1 | 0 8 | 0 4 | 1 5 | 0 4 | 1 2 | 1 4 | 0 8 | 0 6 | 0 8 | 0 3 | 0 7 | 0 5 | 0 4 | 0 7 | 0 2 | 0 2 | 0 3 | 0 6 | 0 3 | 0 1 | 0 7 | 0 1 | 0 8 | 0 2 | 0 0 | |
| INTEGUMENTARY SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Skin | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Trichoepithelioma | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Subcutaneous tissue | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Teratoma, benign | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| RESPIRATORY SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Lungs and bronchi | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Neoplasm, NOS, unclear prim or meta | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Pheochromocytoma, metastatic | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Trachea | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| HEMATOPOIETIC SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Bone marrow | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Spleen | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Hemangiosarcoma | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Lymph nodes | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Thymus | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| CIRCULATORY SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Heart | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| DIGESTIVE SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Salivary gland | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Liver | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Bile duct | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Gallbladder & common bile duct | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Pancreas | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Esophagus | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Stomach | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Small intestine | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Large intestine | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| URINARY SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Kidney | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Tubular cell adenoma | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Urinary bladder | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| ENDOCRINE SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Pituitary | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Adrenal | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Pheochromocytoma | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Pheochromocytoma, malignant | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Thyroid | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| C-cell adenoma | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Parathyroid | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| REPRODUCTIVE SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Mammary gland | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Testis | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Interstitial cell tumor | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Prostate | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| NERVOUS SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Brain | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| BODY CAVITIES | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Peritoneum | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Mesothelioma, NOS | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Tunica vaginalis | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Mesothelioma, NOS | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |

* Animals necropsied

TABLE E2. INDIVIDUAL ANIMAL TUMOR PATHOLOGY OF MALE MARSHALL RATS IN THE TWO-YEAR GAVAGE STUDY OF TRICHLOROETHYLENE: HIGH DOSE

| ANIMAL NUMBER | 0001 | 0002 | 0003 | 0004 | 0005 | 0006 | 0007 | 0008 | 0009 | 0010 | 0011 | 0012 | 0013 | 0014 | 0015 | 0016 | 0017 | 0018 | 0019 | 0020 | 0021 | 0022 | 0023 | 0024 | 0025 | 0026 | 0027 | 0028 | 0029 | 0030 | |
|---|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|----|
| WEEKS ON STUDY | 58 | 57 | 68 | 25 | 76 | 80 | 04 | 25 | 05 | 05 | 09 | 03 | 04 | 07 | 01 | 09 | 02 | 06 | 04 | 07 | 08 | 00 | 06 | 03 | 08 | 01 | 03 | 04 | 05 | 07 | 02 |
| INTEGUMENTARY SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Skin | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Trichoepithelioma | + | + | + | N | + | + | + | + | + | + | + | + | + | + | N | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| RESPIRATORY SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Lungs and bronchi | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Tubular cell adenocarcinoma, metastatic | + | + | + | + | + | A | + | + | + | + | + | + | + | + | + | A | + | + | + | + | + | + | X | + | + | - | + | + | + | + | |
| Pheochromocytoma, metastatic | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Trachea | - | + | + | + | + | + | + | + | + | - | + | + | + | + | - | A | + | + | + | + | + | + | + | + | + | - | + | + | + | - | |
| HEMATOPOIETIC SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Bone marrow | + | + | + | + | + | + | + | + | + | + | + | + | - | - | A | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| Spleen | + | + | + | + | + | A | + | + | + | + | + | + | + | + | + | A | + | + | + | + | + | + | + | + | - | + | + | + | + | - | |
| Lymph nodes | + | + | - | - | - | A | + | + | - | + | + | - | + | + | + | A | - | + | + | + | + | + | + | + | - | + | + | + | + | - | |
| Thymus | - | + | - | - | - | A | - | - | + | + | + | - | + | + | A | - | + | + | + | - | - | - | - | - | - | + | - | + | + | + | |
| CIRCULATORY SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Heart | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Tubular cell adenocarcinoma, metastatic | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | A | + | + | + | + | + | + | + | + | + | - | + | + | + | + | |
| DIGESTIVE SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Salivary gland | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Liver | + | + | + | + | + | A | + | + | + | + | + | + | + | + | + | A | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| Bile duct | + | + | + | + | + | A | + | + | + | + | + | + | + | + | + | A | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| Gallbladder & common bile duct | + | + | + | + | + | A | + | + | + | + | + | + | + | + | + | A | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| Pancreas | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | |
| Esophagus | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | A | + | + | + | + | + | + | + | + | + | - | + | + | + | - | |
| Stomach | + | + | + | + | + | A | + | + | + | + | + | + | + | + | + | A | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| Small intestine | + | + | + | + | + | A | + | + | + | + | + | + | + | + | + | A | + | + | + | + | + | + | + | + | + | - | + | + | + | + | |
| Large intestine | + | + | + | + | + | A | + | + | + | + | + | + | + | + | + | A | + | + | + | + | + | + | + | + | + | - | + | + | + | + | |
| URINARY SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Kidney | + | + | + | + | + | A | + | + | + | + | + | + | + | + | + | A | + | + | + | + | + | + | + | + | + | - | + | + | + | + | |
| Tubular cell adenocarcinoma | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Interstitial cell tumor, invasive | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Urinary bladder | + | + | + | + | + | A | + | + | + | + | + | + | + | + | + | A | + | + | + | + | + | + | + | + | + | - | + | + | + | + | |
| ENDOCRINE SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Pituitary | + | + | + | + | + | A | - | - | - | + | - | - | - | - | - | A | + | + | + | + | + | + | + | + | + | + | + | + | + | - | |
| Adrenal | + | + | + | + | + | A | + | + | + | + | + | + | + | + | + | A | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| Pheochromocytoma | X | + | + | + | + | X | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | X | + | - | + | + | + | |
| Pheochromocytoma, malignant | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Thyroid | + | + | + | + | + | + | + | + | + | - | - | + | + | + | + | A | + | + | + | + | + | + | + | + | + | + | + | + | + | - | |
| C-cell adenoma | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| C-cell carcinoma | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Parathyroid | + | - | - | - | + | A | + | - | - | - | - | - | - | - | - | A | - | - | - | - | - | - | - | - | - | - | - | - | - | | |
| Adenoma, NOS | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| REPRODUCTIVE SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Mammary gland | N | N | N | N | + | + | + | N | N | + | N | N | N | N | + | N | + | N | + | N | N | N | N | + | N | N | N | N | + | N | |
| Testis | + | + | + | + | + | A | + | + | + | + | + | + | + | + | + | A | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| Interstitial cell tumor | X | X | X | X | X | X | X | X | X | X | X | X | X | X | X | X | X | X | X | X | X | X | X | X | X | X | X | X | X | X | |
| Interstitial cell tumor, malignant | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Prostate | + | + | + | + | - | A | + | + | + | + | + | + | + | + | + | A | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| NERVOUS SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Brain | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | A | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| Glioma, NOS | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Astrocytoma | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| BODY CAVITIES | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Peritoneum | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | |
| Mesothelioma, NOS | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Mesothelioma, malignant | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| ALL OTHER SYSTEMS | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Multiple organs, NOS | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | |
| Malignant lymphoma, histiocytic type | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |

TABLE E3. ANALYSIS OF PRIMARY TUMORS IN MALE MARSHALL RATS IN THE TWO-YEAR GAVAGE STUDY OF TRICHLOROETHYLENE

| | Untreated Control | Vehicle Control | 500 mg/kg | 1,000 mg/kg |
|--|-------------------|-----------------|-------------|-------------|
| Adrenal: Pheochromocytoma | | | | |
| Overall Rates (a) | 19/48 (40%) | 21/48 (44%) | 12/43 (28%) | 11/43 (26%) |
| Adjusted Rates (b) | 55.6% | 69.1% | 84.8% | 89.4% |
| Terminal Rates (c) | 16/31 (52%) | 17/26 (65%) | 9/11 (82%) | 5/6 (83%) |
| Life Table Tests (d) | | P=0.014 | P=0.147 | P=0.021 |
| Incidental Tumor Tests (d) | | P=0.095 | P=0.119 | P=0.127 |
| Cochran-Armitage Trend Test (d) | | P=0.040N | | |
| Fisher Exact Test | | | P=0.088N | P=0.055N |
| Adrenal: Pheochromocytoma, Malignant | | | | |
| Overall Rates (a) | 4/48 (8%) | 5/48 (10%) | 1/43 (2%) | 2/43 (5%) |
| Adjusted Rates (b) | 12.1% | 16.2% | 9.1% | 21.4% |
| Terminal Rates (c) | 3/31 (10%) | 2/26 (8%) | 1/11 (9%) | 0/6 (0%) |
| Life Table Tests (d) | | P=0.517 | P=0.420N | P=0.524 |
| Incidental Tumor Tests (d) | | P=0.440N | P=0.695 | P=0.500N |
| Cochran-Armitage Trend Test (d) | | P=0.165N | | |
| Fisher Exact Test | | | P=0.129N | P=0.266N |
| Adrenal: Pheochromocytoma or Pheochromocytoma, Malignant | | | | |
| Overall Rates (a) | 23/48 (48%) | 25/48 (52%) | 12/43 (28%) | 12/43 (28%) |
| Adjusted Rates (b) | 65.4% | 75.1% | 84.8% | 90.9% |
| Terminal Rates (c) | 19/31 (61%) | 18/26 (69%) | 9/11 (82%) | 5/6 (83%) |
| Life Table Tests (d) | | P=0.035 | P=0.353 | P=0.037 |
| Incidental Tumor Tests (d) | | P=0.239 | P=0.158 | P=0.288 |
| Cochran-Armitage Trend Test (d) | | P=0.010N | | |
| Fisher Exact Test | | | P=0.016N | P=0.016N |
| Thyroid: C-Cell Adenoma | | | | |
| Overall Rates (a) | 8/41 (20%) | 4/41 (10%) | 5/41 (12%) | 2/40 (5%) |
| Adjusted Rates (b) | 25.0% | 14.0% | 41.7% | 40.0% |
| Terminal Rates (c) | 8/32 (25%) | 3/25 (12%) | 5/12 (42%) | 2/5 (40%) |
| Life Table Tests (d) | | P=0.129 | P=0.113 | P=0.385 |
| Incidental Tumor Tests (d) | | P=0.172 | P=0.153 | P=0.513 |
| Cochran-Armitage Trend Test (d) | | P=0.292N | | |
| Fisher Exact Test | | | P=0.500 | P=0.350N |
| Thyroid: C-Cell Adenoma or Carcinoma | | | | |
| Overall Rates (a) | 12/41 (29%) | 5/41 (12%) | 5/41 (12%) | 3/40 (7%) |
| Adjusted Rates (b) | 37.5% | 17.9% | 41.7% | 47.5% |
| Terminal Rates (c) | 12/32 (38%) | 4/25 (16%) | 5/12 (42%) | 2/5 (40%) |
| Life Table Tests (d) | | P=0.078 | P=0.178 | P=0.206 |
| Incidental Tumor Tests (d) | | P=0.170 | P=0.228 | P=0.424 |
| Cochran-Armitage Trend Test (d) | | P=0.308N | | |
| Fisher Exact Test | | | P=0.631N | P=0.370N |
| Testis: Interstitial Cell Tumor | | | | |
| Overall Rates (a) | 16/46 (35%) | 17/46 (37%) | 21/48 (44%) | 31/48 (65%) |
| Adjusted Rates (b) | 46.9% | 55.7% | 95.1% | 100.0% |
| Terminal Rates (c) | 14/32 (44%) | 13/26 (50%) | 11/12 (92%) | 6/6 (100%) |
| Life Table Tests (d) | | P<0.001 | P<0.001 | P<0.001 |
| Incidental Tumor Tests (d) | | P<0.001 | P<0.001 | P<0.001 |
| Cochran-Armitage Trend Test (d) | | P=0.005 | | |
| Fisher Exact Test | | | P=0.323 | P=0.007 |
| Testis: Interstitial Cell Tumor or Interstitial Cell Tumor, Malignant | | | | |
| Overall Rates (a) | 16/46 (35%) | 17/46 (37%) | 21/48 (44%) | 32/48 (67%) |
| Adjusted Rates (b) | 46.9% | 55.7% | 95.1% | 100.0% |
| Terminal Rates (c) | 14/32 (44%) | 13/26 (50%) | 11/12 (92%) | 6/6 (100%) |
| Life Table Tests (d) | | P<0.001 | P<0.001 | P<0.001 |
| Incidental Tumor Tests (d) | | P<0.001 | P<0.001 | P<0.001 |
| Cochran-Armitage Trend Test (d) | | P=0.003 | | |
| Fisher Exact Test | | | P=0.323 | P=0.004 |

TABLE E3. ANALYSIS OF PRIMARY TUMORS IN MALE MARSHALL RATS IN THE TWO-YEAR GAVAGE STUDY OF TRICHLOROETHYLENE (Continued)

| | Untreated Control | Vehicle Control | 500 mg/kg | 1,000 mg/kg |
|---------------------------------|-------------------|-----------------|------------|-------------|
| All Sites: Mesothelioma | | | | |
| Overall Rates (a) | 2/49 (4%) | 2/50 (4%) | 2/50 (4%) | 3/50 (6%) |
| Adjusted Rates (b) | 5.5% | 6.0% | 16.7% | 24.5% |
| Terminal Rates (c) | 1/32 (3%) | 0/26 (0%) | 2/12 (17%) | 0/6 (0%) |
| Life Table Tests (d) | | P=0.063 | P=0.356 | P=0.127 |
| Incidental Tumor Tests (d) | | P=0.244 | P=0.090 | P=0.496 |
| Cochran-Armitage Trend Test (d) | | P=0.406 | | |
| Fisher Exact Test | | | P=0.691N | P=0.500 |

(a) Number of tumor-bearing animals/number of animals examined at the site

(b) Kaplan-Meier estimated tumor incidence at the end of the study after adjusting for intercurrent mortality

(c) Observed tumor incidence at terminal kill

(d) Beneath the control incidence are the P values associated with the trend test. Beneath the dosed group incidence are the P values corresponding to pairwise comparisons between that dosed group and the vehiclecontrols. The life table analysis regards tumors in animals dying prior to terminal kill as being (directly or indirectly) the cause of death. The incidental tumor test regards these lesions as nonfatal. The Cochran-Armitage and Fisher exact tests compare directly the overall incidence rates. A negative trend or lower incidence in a dosed group is indicated by (N).

TABLE E4. SUMMARY OF THE INCIDENCE OF NONNEOPLASTIC LESIONS IN MALE MARSHALL RATS IN THE TWO-YEAR GAVAGE STUDY OF TRICHLOROETHYLENE

| | CONTROL (UNTR) | CONTROL (VEH) | LOW DOSE | HIGH DOSE |
|--------------------------------------|----------------|---------------|----------|-----------|
| ANIMALS INITIALLY IN STUDY | 50 | 50 | 50 | 50 |
| ANIMALS MISSING | 1 | | | |
| ANIMALS NECROPSIED | 49 | 50 | 50 | 50 |
| ANIMALS EXAMINED HISTOPATHOLOGICALLY | 49 | 50 | 50 | 50 |
| INTEGUMENTARY SYSTEM | | | | |
| *Skin | (49) | (50) | (50) | (50) |
| Epidermal inclusion cyst | | 1 (2%) | | |
| Hemorrhage | | 1 (2%) | | |
| Abscess, chronic | 1 (2%) | | | |
| *Subcutaneous tissue | (49) | (50) | (50) | (50) |
| Granuloma, NOS | | 1 (2%) | | |
| RESPIRATORY SYSTEM | | | | |
| *Nasal cavity | (49) | (50) | (50) | (50) |
| Vegetable foreign body | | | 1 (2%) | |
| Impaction, NOS | | | | 1 (2%) |
| Hemorrhage | | | 1 (2%) | |
| Inflammation, NOS | | 2 (4%) | | |
| Inflammation, suppurative | | 1 (2%) | 4 (8%) | 2 (4%) |
| Inflammation, acute | | | 1 (2%) | |
| Inflammation, acute suppurative | | | | 1 (2%) |
| #Trachea | (41) | (38) | (41) | (41) |
| Wound, NOS | | | | 1 (2%) |
| Lacerated wound | | 1 (3%) | | 2 (5%) |
| Penetrating wound | | | | 1 (2%) |
| Foreign material, NOS | | 1 (3%) | 1 (2%) | 2 (5%) |
| #Peritracheal tissue | (41) | (38) | (41) | (41) |
| Abscess, chronic | | 1 (3%) | | |
| #Lung | (47) | (49) | (50) | (47) |
| Emphysema, alveolar | | 1 (2%) | 1 (2%) | |
| Collapse | | 1 (2%) | 1 (2%) | |
| Congestion, NOS | 11 (23%) | 8 (16%) | 21 (42%) | 19 (40%) |
| Edema, NOS | | 1 (2%) | 3 (6%) | 5 (11%) |
| Hemorrhage | 4 (9%) | 6 (12%) | 12 (24%) | 15 (32%) |
| Bronchopneumonia, NOS | 1 (2%) | | | |
| Bronchopneumonia, focal | | 1 (2%) | | |
| Inflammation, focal | 19 (40%) | 29 (59%) | 14 (28%) | 6 (13%) |
| Pneumonia, lipid | | 1 (2%) | | |
| Pneumonia, aspiration | | | 1 (2%) | 1 (2%) |
| Bronchopneumonia, acute | | 1 (2%) | | |
| Inflammation, chronic focal | 1 (2%) | 6 (12%) | 14 (28%) | 10 (21%) |
| Inflammation, granulomatous focal | 3 (6%) | 1 (2%) | | 4 (9%) |
| Granuloma, pyogenic | | 1 (2%) | | 1 (2%) |
| Fibrosis, focal | | | | 1 (2%) |
| Foreign material, NOS | | | 4 (8%) | 11 (23%) |
| Hemosiderosis | | 1 (2%) | | |
| Alveolar macrophages | | | 1 (2%) | |
| Hyperplasia, adenomatous | | 1 (2%) | | |
| HEMATOPOIETIC SYSTEM | | | | |
| *Harderian gland | (49) | (50) | (50) | (50) |
| Mastocytosis | | | | 1 (2%) |
| *Blood | (49) | (50) | (50) | (50) |
| Leukocytosis, NOS | 1 (2%) | | | |
| Hypochromasia | 1 (2%) | | | |
| #Bone marrow | (46) | (49) | (36) | (45) |
| Hyperplasia, granulocytic | 1 (2%) | 1 (2%) | | 1 (2%) |

TABLE E4. SUMMARY OF THE INCIDENCE OF NONNEOPLASTIC LESIONS IN MALE MARSHALL RATS IN THE TWO-YEAR GAVAGE STUDY OF TRICHLOROETHYLENE (Continued)

| | CONTROL (UNTR) | CONTROL (VEH) | LOW DOSE | HIGH DOSE |
|---|----------------|---------------|----------|-----------|
| HEMATOPOIETIC SYSTEM (Continued) | | | | |
| #Spleen | (46) | (48) | (43) | (46) |
| Hamartoma | | 1 (2%) | | |
| Congestion, NOS | | | 2 (5%) | |
| Hemosiderosis | 2 (4%) | 1 (2%) | 1 (2%) | 1 (2%) |
| Atrophy, NOS | | 1 (2%) | | |
| Hematopoiesis | 4 (9%) | 1 (2%) | | 2 (4%) |
| Erythropoiesis | | | | 1 (2%) |
| Myelopoiesis | | 1 (2%) | | |
| #Lymph node | (36) | (33) | (25) | (27) |
| Plasmacytosis | 1 (3%) | | | |
| #Mandibular lymph node | (36) | (33) | (25) | (27) |
| Hemorrhage | 1 (3%) | 1 (3%) | | |
| #Mesenteric lymph node | (36) | (33) | (25) | (27) |
| Hemorrhage | 7 (19%) | 4 (12%) | | |
| Inflammation, chronic focal | | | | 1 (4%) |
| Fibrosis, focal | 1 (3%) | | | |
| #Liver | (47) | (49) | (50) | (47) |
| Hematopoiesis | | 1 (2%) | | |
| #Peyers patch | (43) | (44) | (48) | (44) |
| Hyperplasia, lymphoid | 1 (2%) | | | |
| #Thymus | | (7) | (18) | (21) |
| Hemorrhage | | | 2 (11%) | 6 (29%) |
| Involution, NOS | | | 1 (6%) | |
| CIRCULATORY SYSTEM | | | | |
| #Lung | (47) | (49) | (50) | (47) |
| Perivasculitis | | | 1 (2%) | 2 (4%) |
| #Heart | (47) | (49) | (50) | (48) |
| Thrombosis, NOS | | | | 1 (2%) |
| Thrombus, organized | | | 1 (2%) | |
| Inflammation, focal | | 1 (2%) | 1 (2%) | |
| Inflammation, interstitial | 1 (2%) | 1 (2%) | | |
| Inflammation, chronic | 1 (2%) | | | |
| Fibrosis, focal | 3 (6%) | 2 (4%) | 6 (12%) | |
| Endocardiosis | | 1 (2%) | | |
| #Endocardium | (47) | (49) | (50) | (48) |
| Endocarditis, verrucous | 1 (2%) | | | |
| #Cardiac valve | (47) | (49) | (50) | (48) |
| Thrombus, organized | | | | 1 (2%) |
| Fibrosis | 1 (2%) | | | |
| *Artery | (49) | (50) | (50) | (50) |
| Mineralization | | 1 (2%) | | |
| Periarteritis | | 1 (2%) | | |
| *Pulmonary artery | (49) | (50) | (50) | (50) |
| Calcification, focal | 1 (2%) | 4 (8%) | | |
| DIGESTIVE SYSTEM | | | | |
| #Salivary gland | (40) | (43) | (42) | (41) |
| Inflammation, chronic | | | 1 (2%) | |
| #Liver | (47) | (49) | (50) | (47) |
| Cyst, NOS | 1 (2%) | | | |
| Multilocular cyst | 1 (2%) | | | |
| Congestion, NOS | 1 (2%) | 2 (4%) | 1 (2%) | 3 (6%) |
| Congestion, chronic passive | 1 (2%) | | | |
| Hemorrhage | | | 3 (6%) | 1 (2%) |
| Inflammation, NOS | 22 (47%) | 31 (63%) | 6 (12%) | |
| Inflammation, focal | | | 2 (4%) | 1 (2%) |
| Inflammation, acute/chronic | | 1 (2%) | | |

TABLE E4. SUMMARY OF THE INCIDENCE OF NONNEOPLASTIC LESIONS IN MALE MARSHALL RATS IN THE TWO-YEAR GAVAGE STUDY OF TRICHLOROETHYLENE (Continued)

| | CONTROL (UNTR) | CONTROL (VEH) | LOW DOSE | HIGH DOSE |
|-----------------------------------|----------------|---------------|----------|-----------|
| DIGESTIVE SYSTEM | | | | |
| #Liver (Continued) | (47) | (49) | (50) | (47) |
| Degeneration, NOS | | | 1 (2%) | |
| Necrosis, focal | 1 (2%) | | 5 (10%) | |
| Necrosis, hemorrhagic | 1 (2%) | | | |
| Necrosis, central | 2 (4%) | | | |
| Metamorphosis, fatty | 7 (15%) | 3 (6%) | 7 (14%) | |
| Nuclear enlargement | | | 1 (2%) | |
| Basophilic cyto change | 1 (2%) | | | |
| Clear cell change | 5 (11%) | 4 (8%) | 3 (6%) | 2 (4%) |
| Cell size alteration | 1 (2%) | | | |
| Depletion, glycogen | | 1 (2%) | | |
| #Liver/periportal | (47) | (49) | (50) | (47) |
| Atrophy, NOS | 1 (2%) | | | |
| #Pancreas | (42) | (49) | (46) | (46) |
| Inflammation, chronic focal | 1 (2%) | | | 1 (2%) |
| Necrosis, hemorrhagic | 1 (2%) | | | |
| Atrophy, NOS | 1 (2%) | | | |
| Atrophy, focal | 1 (2%) | | | |
| *Pharynx | (49) | (50) | (50) | (50) |
| Granuloma, foreign body | | 1 (2%) | | |
| #Esophagus | (45) | (47) | (44) | (47) |
| Lacerated wound | | 1 (2%) | 1 (2%) | 2 (4%) |
| Penetrating wound | | | | 1 (2%) |
| Inflammation, granulomatous focal | | | 1 (2%) | |
| Perforation, inflammatory | | | | 1 (2%) |
| #Gastric mucosa | (47) | (47) | (48) | (47) |
| Calcification, metastatic | | 1 (2%) | | |
| #Small intestine | (43) | (44) | (48) | (44) |
| Impaction, NOS | | | 1 (2%) | |
| #Colon | (41) | (45) | (43) | (43) |
| Parasitism | 3 (7%) | 3 (7%) | 2 (5%) | 6 (14%) |
| *Rectum | (49) | (50) | (50) | (50) |
| Parasitism | | 1 (2%) | | |
| URINARY SYSTEM | | | | |
| #Kidney | (49) | (49) | (50) | (47) |
| Congenital hydronephrosis | 10 (20%) | 7 (14%) | 5 (10%) | 1 (2%) |
| Hydronephrosis | 1 (2%) | 1 (2%) | 3 (6%) | 1 (2%) |
| Cyst, NOS | 1 (2%) | | | 1 (2%) |
| Congestion, NOS | 1 (2%) | | 1 (2%) | |
| Hemorrhage | 2 (4%) | | | |
| Pyelonephritis, NOS | | 1 (2%) | 1 (2%) | |
| Glomerulonephritis, focal | | 1 (2%) | | |
| Pyelonephritis, focal | 1 (2%) | | | |
| Glomerulonephritis, acute | | 1 (2%) | | |
| Glomerulonephritis, subacute | 1 (2%) | | | |
| Glomerulonephritis, chronic | | 1 (2%) | | |
| Inflammation, granulomatous focal | | | 1 (2%) | |
| Nephropathy | 45 (92%) | 49 (100%) | 47 (94%) | 46 (98%) |
| Nephropathy, toxic | | | 18 (36%) | 23 (49%) |
| Nephrosis, NOS | 1 (2%) | | | |
| Necrosis, medullary | | | 1 (2%) | |
| Calcification, focal | 21 (43%) | 14 (29%) | 8 (16%) | 5 (11%) |
| Pigmentation, NOS | | | | 1 (2%) |
| Cytomegaly | | | 48 (96%) | 47 (100%) |
| Hyperplasia, tubular cell | | | 1 (2%) | |
| #Renal papilla | (49) | (49) | (50) | (47) |
| Inflammation, acute | | | 1 (2%) | |

TABLE E4. SUMMARY OF THE INCIDENCE OF NONNEOPLASTIC LESIONS IN MALE MARSHALL RATS
IN THE TWO-YEAR GAVAGE STUDY OF TRICHLOROETHYLENE (Continued)

| | CONTROL (UNTR) | CONTROL (VEH) | LOW DOSE | HIGH DOSE |
|-----------------------------------|----------------|---------------|----------|-----------|
| BODY CAVITIES | | | | |
| *Thorax | (49) | (50) | (50) | (50) |
| Inflammation, necrotizing | | 1 (2%) | | |
| *Peritoneal cavity | (49) | (50) | (50) | (50) |
| Hemoperitoneum | | | 1 (2%) | |
| *Pleural cavity | (49) | (50) | (50) | (50) |
| Hemorrhage | | | | 1 (2%) |
| ALL OTHER SYSTEMS | | | | |
| *Multiple organs | (49) | (50) | (50) | (50) |
| Hemorrhage | | | 1 (2%) | |
| SPECIAL MORPHOLOGY SUMMARY | | | | |
| No lesion reported | | | | 1 |
| Animal missing/no necropsy | 1 | | | |
| Auto/necropsy/histo perf | 1 | | | 2 |

* Number of animals receiving complete necropsy examination; all gross lesions including masses examined microscopically.

Number of animals examined microscopically at this site

TABLE E4. SUMMARY OF THE INCIDENCE OF NONNEOPLASTIC LESIONS IN MALE MARSHALL RATS IN THE TWO-YEAR GAVAGE STUDY OF TRICHLOROETHYLENE (Continued)

| | CONTROL (UNTR) | CONTROL (VEH) | LOW DOSE | HIGH DOSE |
|-----------------------------------|----------------|---------------|----------|-----------|
| URINARY SYSTEM (Continued) | | | | |
| #Kidney/pelvis | (49) | (49) | (50) | (47) |
| Inflammation, NOS | 1 (2%) | | | |
| Inflammation, suppurative | 1 (2%) | | | |
| Inflammation, acute | 1 (2%) | 2 (4%) | 1 (2%) | |
| Hyperplasia, epithelial | 2 (4%) | | | |
| #Urinary bladder | (41) | (45) | (41) | (44) |
| Calculus, unknown gross or micro | | 1 (2%) | | |
| Hemorrhage | 1 (2%) | | 2 (5%) | |
| Inflammation, NOS | 1 (2%) | | | |
| Inflammation, hemorrhagic | | 1 (2%) | | |
| Inflammation, acute | | | 1 (2%) | |
| Inflammation, acute focal | | 1 (2%) | | |
| Inflammation, acute hemorrhagic | | | | 1 (2%) |
| #Urinary bladder/mucosa | (41) | (45) | (41) | (44) |
| Hyperplasia, papillary | | 1 (2%) | | |
| *Urethra | (49) | (50) | (50) | (50) |
| Inflammation, acute | | | 1 (2%) | |
| ENDOCRINE SYSTEM | | | | |
| #Pituitary | (36) | (21) | (25) | (28) |
| Cyst, NOS | | | 1 (4%) | 1 (4%) |
| Hyperplasia, chromophobe cell | 3 (8%) | | | |
| #Adrenal | (48) | (48) | (43) | (43) |
| Hemorrhage | | 1 (2%) | | |
| Hyperplasia, focal | | 1 (2%) | | |
| Angiectasis | | 2 (4%) | | |
| #Adrenal cortex | (48) | (48) | (43) | (43) |
| Hemorrhage | 1 (2%) | 2 (4%) | | 1 (2%) |
| Degeneration, lipoid | 2 (4%) | 1 (2%) | | 1 (2%) |
| Hyperplasia, focal | 2 (4%) | 1 (2%) | 1 (2%) | |
| Angiectasis | | | 1 (2%) | |
| #Adrenal medulla | (48) | (48) | (43) | (43) |
| Hyperplasia, NOS | | 1 (2%) | | |
| Hyperplasia, focal | 8 (17%) | 5 (10%) | 5 (12%) | 3 (7%) |
| #Thyroid | (41) | (41) | (41) | (40) |
| Cyst, NOS | | 1 (2%) | | |
| Follicular cyst, NOS | | | 2 (5%) | |
| Hyperplasia, C-cell | 6 (15%) | 10 (24%) | 4 (10%) | 1 (3%) |
| REPRODUCTIVE SYSTEM | | | | |
| *Mammary gland | (49) | (50) | (50) | (50) |
| Hyperplasia, cystic | 1 (2%) | | | |
| #Prostate | (44) | (47) | (44) | (47) |
| Retention of content | | | | 1 (2%) |
| Hemorrhage | | | 1 (2%) | |
| Inflammation, NOS | 4 (9%) | 5 (11%) | 2 (5%) | 3 (6%) |
| Inflammation, focal | 30 (68%) | 29 (62%) | 14 (32%) | 16 (34%) |
| Inflammation, multifocal | | | | 1 (2%) |
| Inflammation, suppurative | 1 (2%) | | 3 (7%) | |
| Inflammation, hemorrhagic | | 1 (2%) | | |
| Inflammation, acute | 1 (2%) | | 2 (5%) | |
| Inflammation, acute suppurative | | 2 (4%) | | 1 (2%) |
| Inflammation, acute hemorrhagic | | | | 1 (2%) |
| Inflammation, acute | 1 (2%) | | 2 (5%) | |
| Inflammation, acute suppurative | | 2 (4%) | | 1 (2%) |
| Inflammation, acute hemorrhagic | | | | 1 (2%) |
| Inflammation, acute/chronic | 2 (5%) | 2 (4%) | | 2 (4%) |
| Inflammation, chronic | | | | 1 (2%) |
| Inflammation, chronic focal | | | 1 (2%) | |

TABLE E4. SUMMARY OF THE INCIDENCE OF NONNEOPLASTIC LESIONS IN MALE MARSHALL RATS IN THE TWO-YEAR GAVAGE STUDY OF TRICHLOROETHYLENE (Continued)

| | CONTROL (UNTR) | CONTROL (VEH) | LOW DOSE | HIGH DOSE |
|-----------------------------------|----------------|---------------|----------|-----------|
| REPRODUCTIVE SYSTEM | | | | |
| #Prostate (Continued) | (44) | (47) | (44) | (47) |
| Abscess, chronic | | | | 1 (2%) |
| Inflammation, granulomatous focal | | 1 (2%) | | |
| Granuloma, pyogenic | | | | 1 (2%) |
| Corpora amylacea | 37 (84%) | 38 (81%) | 19 (43%) | 29 (62%) |
| Hyperplasia, NOS | | 1 (2%) | 1 (2%) | |
| Hyperplasia, papillary | | | 1 (2%) | |
| *Seminal vesicle | (49) | (50) | (50) | (50) |
| Congenital hypoplasia | 1 (2%) | | 2 (4%) | |
| Retention of content | | | | 1 (2%) |
| Hemorrhage | | | 2 (4%) | |
| Inflammation, NOS | | 1 (2%) | | |
| Inflammation, focal | 1 (2%) | 1 (2%) | | |
| Inflammation, suppurative | 1 (2%) | 1 (2%) | 2 (4%) | |
| Inflammation, hemorrhagic | | 1 (2%) | | |
| Inflammation, acute | | | 1 (2%) | 1 (2%) |
| Inflammation, acute suppurative | | 2 (4%) | | |
| Inflammation, acute hemorrhagic | | | | 1 (2%) |
| Abscess, NOS | | | 1 (2%) | |
| Inflammation, acute/chronic | | 2 (4%) | | |
| Inflammation chronic suppurative | 1 (2%) | | | |
| Atrophy, NOS | | | | 2 (4%) |
| #Testis | (46) | (46) | (48) | (48) |
| Agenesis | | 2 (4%) | | |
| Hemorrhage | | 1 (2%) | | |
| Lipogranuloma | | | 1 (2%) | |
| Calcification, focal | | 1 (2%) | | |
| Syncytial alteration | | 1 (2%) | | |
| Atrophy, NOS | | | 1 (2%) | |
| Atrophy, focal | 3 (7%) | 1 (2%) | 1 (2%) | |
| Aspermatogenesis | 1 (2%) | 2 (4%) | | 1 (2%) |
| Hypospermatogenesis | | | 1 (2%) | |
| Hyperplasia, interstitial cell | 2 (4%) | 1 (2%) | 6 (13%) | 5 (10%) |
| Dysplasia, NOS | 1 (2%) | | | |
| #Rete testis | (46) | (46) | (48) | (48) |
| Hyperplasia, NOS | | | | 1 (2%) |
| *Epididymis | (49) | (50) | (50) | (50) |
| Inflammation, focal | 1 (2%) | | | |
| Inflammation, interstitial | | | 1 (2%) | |
| NERVOUS SYSTEM | | | | |
| *Spinal cord | (49) | (50) | (50) | (50) |
| Hemorrhage | | 1 (2%) | | |
| SPECIAL SENSE ORGANS | | | | |
| *Harderian gland | (49) | (50) | (50) | (50) |
| Inflammation, focal | | | | 1 (2%) |
| Inflammation, acute focal | | | 1 (2%) | |
| Inflammation, chronic | | | 3 (6%) | 1 (2%) |
| Inflammation, chronic focal | | 2 (4%) | 7 (14%) | 8 (16%) |
| *Middle ear | (49) | (50) | (50) | (50) |
| Inflammation, suppurative | | | 2 (4%) | |
| MUSCULOSKELETAL SYSTEM | | | | |
| *Muscle hip/thigh | (49) | (50) | (50) | (50) |
| Degeneration, NOS | | | 1 (2%) | |
| Atrophy, serous | | 1 (2%) | | |

APPENDIX F

SUMMARY OF LESIONS IN FEMALE MARSHALL RATS IN THE TWO-YEAR GAVAGE STUDY OF TRICHLOROETHYLENE

| | PAGE | |
|----------|--|-----|
| TABLE F1 | SUMMARY OF THE INCIDENCE OF NEOPLASMS IN FEMALE MARSHALL RATS IN THE TWO-YEAR GAVAGE STUDY OF TRICHLOROETHYLENE | 187 |
| TABLE F2 | INDIVIDUAL ANIMAL TUMOR PATHOLOGY OF FEMALE MARSHALL RATS IN THE TWO-YEAR GAVAGE STUDY OF TRICHLOROETHYLENE | 190 |
| TABLE F3 | ANALYSIS OF PRIMARY TUMORS IN FEMALE MARSHALL RATS IN THE TWO-YEAR GAVAGE STUDY OF TRICHLOROETHYLENE | 198 |
| TABLE F4 | SUMMARY OF THE INCIDENCE OF NONNEOPLASTIC LESIONS IN FEMALE MARSHALL RATS IN THE TWO-YEAR GAVAGE STUDY OF TRICHLOROETHYLENE | 201 |

TABLE F1. SUMMARY OF THE INCIDENCE OF NEOPLASMS IN FEMALE MARSHALL RATS IN THE TWO-YEAR GAVAGE STUDY OF TRICHLOROETHYLENE

| | CONTROL (UNTR) | CONTROL (VEH) | LOW DOSE | HIGH DOSE |
|--------------------------------------|----------------|---------------|----------|-----------|
| ANIMALS INITIALLY IN STUDY | 50 | 50 | 50 | 50 |
| ANIMALS NECROPSIED | 50 | 50 | 50 | 48 |
| ANIMALS EXAMINED HISTOPATHOLOGICALLY | 50 | 50 | 50 | 48 |
| INTEGUMENTARY SYSTEM | | | | |
| *Skin | (50) | (50) | (50) | (48) |
| Squamous cell carcinoma | | | 1 (2%) | |
| Basal cell tumor | | 1 (2%) | | |
| Basal cell carcinoma | | 1 (2%) | | |
| Trichoepithelioma | | 1 (2%) | | |
| RESPIRATORY SYSTEM | | | | |
| #Lung | (50) | (49) | (49) | (46) |
| Neoplasm, NOS, metastatic | | | 1 (2%) | |
| Squamous cell carcinoma, metastatic | | | 1 (2%) | |
| Pheochromocytoma, metastatic | 3 (6%) | 3 (6%) | | 1 (2%) |
| Sarcoma, NOS, metastatic | | | 1 (2%) | |
| HEMATOPOIETIC SYSTEM | | | | |
| *Multiple organs | (50) | (50) | (50) | (48) |
| Malignant lymphoma, NOS | 1 (2%) | | | |
| Malignant lymphoma, undiffer type | | | 1 (2%) | |
| Malignant lymphoma, lymphocytic type | | 1 (2%) | | |
| Malignant lymphoma, histiocytic type | 1 (2%) | | | |
| Lymphocytic leukemia | | 1 (2%) | | |
| CIRCULATORY SYSTEM | | | | |
| *Axilla | (50) | (50) | (50) | (48) |
| Hemangiosarcoma | | 1 (2%) | | |
| #Lung | (50) | (49) | (49) | (46) |
| Hemangiosarcoma, metastatic | | 1 (2%) | | |
| DIGESTIVE SYSTEM | | | | |
| *Buccal mucosa | (50) | (50) | (50) | (48) |
| Undifferentiated carcinoma | | | 1 (2%) | |
| #Stomach | (49) | (49) | (46) | (43) |
| Squamous cell papilloma | | | | 1 (2%) |
| URINARY SYSTEM | | | | |
| #Kidney | (49) | (50) | (48) | (44) |
| Tubular cell adenoma | 1 (2%) | 1 (2%) | 1 (2%) | |
| Tubular cell adenocarcinoma | | | 1 (2%) | 1 (2%) |
| #Kidney/pelvis | (49) | (50) | (48) | (44) |
| Transitional cell papilloma | | | 2 (4%) | 1 (2%) |
| Transitional cell carcinoma | 1 (2%) | 1 (2%) | | 2 (5%) |
| #Urinary bladder | (45) | (46) | (40) | (40) |
| Transitional cell papilloma | | | 1 (3%) | |

TABLE F1. SUMMARY OF THE INCIDENCE OF NEOPLASMS IN FEMALE MARSHALL RATS IN THE TWO-YEAR GAVAGE STUDY OF TRICHLOROETHYLENE (Continued)

| | CONTROL (UNTR) | CONTROL (VEH) | LOW DOSE | HIGH DOSE |
|---------------------------------|----------------|---------------|----------|-----------|
| ENDOCRINE SYSTEM | | | | |
| #Pituitary | (42) | (45) | (38) | (42) |
| Carcinoma, NOS | 1 (2%) | | | |
| Chromophobe adenoma | 22 (52%) | 18 (40%) | 8 (21%) | 6 (14%) |
| Chromophobe carcinoma | 1 (2%) | 1 (2%) | | |
| Acidophil adenoma | | 1 (2%) | | |
| Acidophil carcinoma | 1 (2%) | | | |
| #Adrenal | (47) | (49) | (47) | (43) |
| Cortical adenoma | 4 (9%) | 5 (10%) | 2 (4%) | 3 (7%) |
| Pheochromocytoma | 30 (64%) | 39 (80%) | 15 (32%) | 9 (21%) |
| Pheochromocytoma, malignant | 4 (9%) | 4 (8%) | | 1 (2%) |
| #Thyroid | (48) | (43) | (46) | (43) |
| Adenoma, NOS | 1 (2%) | | | |
| C-cell adenoma | 12 (25%) | 10 (23%) | 5 (11%) | 4 (9%) |
| C-cell carcinoma | 1 (2%) | | 2 (4%) | 1 (2%) |
| #Pancreatic islets | (48) | (49) | (46) | (42) |
| Islet cell adenoma | 2 (4%) | | | |
| REPRODUCTIVE SYSTEM | | | | |
| *Mammary gland | (50) | (50) | (50) | (48) |
| Adenocarcinoma, NOS | 1 (2%) | 3 (6%) | | |
| Fibroadenoma | 7 (14%) | 11 (22%) | 2 (4%) | 4 (8%) |
| *Vagina | (50) | (50) | (50) | (48) |
| Papilloma, NOS | | | | 1 (2%) |
| #Uterus | (47) | (50) | (45) | (44) |
| Endometrial stromal polyp | 6 (13%) | 3 (6%) | 5 (11%) | 2 (5%) |
| #Uterus/endometrium | (47) | (50) | (45) | (44) |
| Carcinoma, NOS | | | | 2 (5%) |
| #Ovary | (47) | (48) | (45) | (44) |
| Granulosa cell carcinoma | | 1 (2%) | | |
| NERVOUS SYSTEM | | | | |
| #Brain | (50) | (50) | (50) | (48) |
| Chromophobe carcinoma, invasive | | 1 (2%) | | |
| Acidophil carcinoma, invasive | 1 (2%) | | | |
| SPECIAL SENSE ORGANS | | | | |
| None | | | | |
| MUSCULOSKELETAL SYSTEM | | | | |
| *Mandible | (50) | (50) | (50) | (48) |
| Sarcoma, NOS | | | 1 (2%) | |
| BODY CAVITIES | | | | |
| *Thorax | (50) | (50) | (50) | (48) |
| Fibrosarcoma | | | 1 (2%) | |
| *Mediastinum | (50) | (50) | (50) | (48) |
| Nonchromaffin paraganglioma | | | 1 (2%) | |
| ALL OTHER SYSTEMS | | | | |
| None | | | | |

TABLE F1. SUMMARY OF THE INCIDENCE OF NEOPLASMS IN FEMALE MARSHALL RATS IN THE TWO-YEAR GAVAGE STUDY OF TRICHLOROETHYLENE (Continued)

| | CONTROL (UNTR) | CONTROL (VEH) | LOW DOSE | HIGH DOSE |
|---------------------------------------|----------------|---------------|----------|-----------|
| ANIMAL DISPOSITION SUMMARY | | | | |
| Animals initially in study | 50 | 50 | 50 | 50 |
| Natural death | 10 | 7 | 14 | 19 |
| Moribund sacrifice | 8 | 10 | 10 | 3 |
| Terminal sacrifice | 31 | 30 | 12 | 10 |
| Accidentally killed, NOS | 1 | 3 | 14 | 18 |
| TUMOR SUMMARY | | | | |
| Total animals with primary tumors** | 46 | 47 | 31 | 22 |
| Total primary tumors | 97 | 104 | 50 | 38 |
| Total animals with benign tumors | 45 | 46 | 28 | 19 |
| Total benign tumors | 85 | 90 | 42 | 31 |
| Total animals with malignant tumors | 10 | 11 | 8 | 7 |
| Total malignant tumors | 12 | 14 | 8 | 7 |
| Total animals with secondary tumors## | 4 | 5 | 3 | 1 |
| Total secondary tumors | 4 | 5 | 3 | 1 |

* Number of animals receiving complete necropsy examination; all gross lesions including masses examined microscopically.

** Primary tumors: all tumors except secondary tumors

Number of animals examined microscopically at this site

Secondary tumors: metastatic tumors or tumors invasive into an adjacent organ

TABLE F2. INDIVIDUAL ANIMAL TUMOR PATHOLOGY OF FEMALE MARSHALL RATS: VEHICLE CONTROL (Continued)

| ANIMAL NUMBER | 902 | 906 | 920 | 928 | 930 | 933 | 935 | 936 | 939 | 944 | 946 | 949 | 955 | 960 | 964 | 966 | 971 | 972 | 974 | 975 | 976 | 977 | 983 | 989 | 996 | 999 | TOTAL TISSUES TUMORS | |
|--------------------------------------|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|----------------------|-----|
| WEEKS ON STUDY | 104 | 011 | 104 | 008 | 104 | 104 | 104 | 104 | 100 | 008 | 000 | 109 | 104 | 104 | 004 | 109 | 104 | 104 | 104 | 104 | 104 | 100 | 109 | 004 | 006 | 104 | | |
| INTEGUMENTARY SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Skin | N | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | *50 | |
| Basal cell tumor | | | | | | | | | | | | | | | | | | | | | | | | | | | 1 | |
| Basal cell carcinoma | | | | | | | | | | | | | | | | | | | | | | | | | | | 1 | |
| Trichoepithelioma | | | | | | | | | | | | | | | X | X | | | | | | | | | | | 1 | |
| RESPIRATORY SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Lungs and bronchi | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | 49 |
| Pheochromocytoma, metastatic | | | | | | | | | | | | | | | | | | | | | | | | | | | 3 | |
| Hemangiosarcoma, metastatic | | | | | | | | | | | | | | | | | | | | | | | | | | | 1 | |
| Trachea | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | 39 | |
| HEMATOPOIETIC SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Bone marrow | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | 50 |
| Spleen | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | 49 | |
| Lymph nodes | - | + | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | 31 | |
| Thymus | - | + | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | 7 | |
| CIRCULATORY SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Heart | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | 49 |
| DIGESTIVE SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Salivary gland | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | 46 |
| Liver | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | 49 |
| Bile duct | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | 49 |
| Gallbladder & common bile duct | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | *50 |
| Pancreas | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | 49 |
| Esophagus | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | 50 |
| Stomach | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | 49 |
| Small intestine | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | 46 |
| Large intestine | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | 45 |
| URINARY SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Kidney | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | 50 |
| Tubular cell adenoma | | | | | | | | | | | | | | | | | | | | | | | | | | | 1 | |
| Kidney/pelvis | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | 50 |
| Transitional cell carcinoma | | | | | | | | | | | | | | | | | | | | | | | | | | | 1 | |
| Urinary bladder | + | + | + | - | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | 46 |
| ENDOCRINE SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Pituitary | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | 45 |
| Chromophobe adenoma | X | X | | | | | | | | | | | | | | | | | | | | | | | | | 18 | |
| Chromophobe carcinoma | | | | | | | | | | | | | | | | | | | | | | | | | | | 1 | |
| Acidophil adenoma | | | | | | | | | | | | | | | | | | | | | | | | | | | 1 | |
| Adrenal | - | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | 49 |
| Cortical adenoma | | X | X | | | | | | | | | | | | | | | | | | | | | | | | 5 | |
| Pheochromocytoma | | | X | X | X | X | X | X | X | X | X | X | X | X | X | X | X | X | X | X | X | X | X | X | X | X | X | 39 |
| Pheochromocytoma, malignant | | | | | | | | | | | | | | | | | | | | | | | | | | | 4 | |
| Thyroid | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | 43 |
| C-cell adenoma | | | | | X | | | | | | | | | | | | | | | | | | | | | | 10 | |
| Parathyroid | - | + | + | - | + | + | - | + | - | - | - | - | - | - | + | + | - | - | - | - | - | - | - | - | - | - | 22 | |
| REPRODUCTIVE SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Mammary gland | + | + | N | + | N | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | *50 |
| Adenocarcinoma, NOS | X | | | | | | | | | | | | | | | | | | | | | | | | | | 3 | |
| Fibroadenoma | | | | X | | | | | | | | | | | | | | | | | | | | | | | 11 | |
| Uterus | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | 50 |
| Endometrial stromal polyp | | | | X | | | | | | | | | | | | | | | | | | | | | | | 3 | |
| Ovary | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | 48 |
| Granulosa cell carcinoma | | | | | | | | | | | | | | | | | | | | | | | | | | | 1 | |
| NERVOUS SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Brain | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | 50 |
| Chromophobe carcinoma, invasive | | | | | | | | | | | | | | | | | | | | | | | | | | | 1 | |
| ALL OTHER SYSTEMS | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Multiple organs, NOS | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | *50 |
| Malignant lymphoma, lymphocytic type | | | | | | | | | | | | | | | | | | | | | | | | | | | 1 | |
| Lymphocytic leukemia | | | | | | | | | | | | | | | | | | | | | | | | | | | 1 | |
| Axilla, NOS | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Hemangiosarcoma | | | | | | | | | | | | | | | | | | | | | | | | | | | 1 | |

* Animals necropsied

TABLE F2. INDIVIDUAL ANIMAL TUMOR PATHOLOGY OF FEMALE MARSHALL RATS: LOW DOSE (Continued)

| ANIMAL NUMBER | 9 1 3 | 9 1 5 | 9 1 8 | 9 2 1 | 9 2 2 | 9 2 3 | 9 3 1 | 9 3 2 | 9 3 3 | 9 4 3 | 9 4 4 | 9 5 5 | 9 7 7 | 9 0 0 | 9 1 1 | 9 2 2 | 9 4 4 | 9 5 5 | 9 6 6 | 9 7 7 | 9 8 8 | 9 8 8 | 9 8 8 | 9 8 8 | TOTAL: TISSUES TUMORS | | |
|-------------------------------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-----------------------------|-------------|----|
| WEEKS ON STUDY | 0 8 1 | 0 6 8 | 0 6 8 | 0 4 8 | 1 0 4 | 1 0 4 | 1 0 4 | 0 9 7 | 0 6 2 | 0 7 6 | 1 0 4 | 0 7 7 | 0 7 1 | 0 1 2 | 1 0 4 | 1 0 4 | 1 0 4 | 0 0 4 | 1 0 3 | 1 0 3 | 0 1 7 | 1 0 4 | 0 7 8 | 1 0 4 | 1 0 3 | 0 0 0 | |
| INTEGUMENTARY SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Skin | + | + | + | + | + | N | + | + | + | + | + | + | + | + | N | + | + | + | + | + | + | + | + | + | + | *50 1 | |
| Squamous cell carcinoma | | | | | | | | | | | | | | | | | | | | | | | | | X | | |
| RESPIRATORY SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Lungs and bronchi | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | 49 1 | |
| Neoplasm, NOS, metastatic | | | | | | | | | | | | | | | | | | | | | | | X | | | | |
| Squamous cell carcinoma, metastatic | | | | | | | | | | | | | | | | | | | | | | | | | | X | |
| Sarcoma, NOS, metastatic | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Trachea | - | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | - | + | + | - | - | 45 |
| HEMATOPOIETIC SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Bone marrow | + | + | + | + | + | + | + | + | + | + | + | + | + | - | + | + | + | + | + | + | - | + | + | + | + | 48 | |
| Spleen | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | 47 | |
| Lymph nodes | + | - | + | - | - | + | + | + | + | - | + | + | + | - | - | - | - | - | - | - | - | - | - | - | - | 29 | |
| Thymus | + | + | - | - | - | - | - | - | - | + | - | - | + | - | - | - | - | - | - | - | - | - | - | - | - | 16 | |
| CIRCULATORY SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Heart | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | 50 | |
| DIGESTIVE SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Oral cavity | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | *50 1 | |
| Undifferentiated carcinoma | X | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Salivary gland | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | 47 | |
| Liver | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | - | + | + | + | + | 48 | |
| Bile duct | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | - | + | + | + | + | 48 | |
| Gallbladder & common bile duct | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | *50 | |
| Pancreas | + | + | + | + | + | + | + | + | + | + | + | + | + | - | + | + | + | + | + | + | + | + | + | + | + | 46 | |
| Esophagus | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | 50 | |
| Stomach | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | 46 | |
| Small intestine | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | 45 | |
| Large intestine | + | + | - | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | - | + | + | + | + | 43 | |
| URINARY SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Kidney | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | 48 | |
| Tubular cell adenoma | | | | | | | | | | | | | | | | | | | | | | X | | | | 1 | |
| Tubular cell adenocarcinoma | | | | | | | | | | | | | | | | | | | | | | | | | | 1 | |
| Kidney/pelvis | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | - | + | + | + | + | 48 | |
| Transitional cell papilloma | | | | | | | | | X | | | | | | | | | | | | | | | | | 2 | |
| Urinary bladder | - | + | + | + | + | + | + | + | + | + | + | + | + | - | + | + | + | + | + | + | - | + | + | - | + | 40 | |
| Transitional cell papilloma | | | | | | | | | | | | | | | | | | | | | | | | | | 1 | |
| ENDOCRINE SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Pituitary | - | + | + | - | + | + | + | + | + | + | + | + | - | - | + | - | + | + | + | + | - | - | + | + | + | 38 | |
| Chromophobe adenoma | | | | | | | X | | | | | | | | X | | X | | | | | | | X | | 8 | |
| Adrenal | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | - | + | + | + | + | 47 | |
| Cortical adenoma | | | | | | | X | | | | X | | | | | | | | | | | | | | | 2 | |
| Pheochromocytoma | X | X | | | | X | | | | X | | | | X | | X | | | | | | | X | X | X | 15 | |
| Thyroid | + | + | + | + | + | + | + | + | + | + | + | + | - | + | + | + | + | + | + | + | + | + | + | + | + | 46 | |
| C-cell adenoma | | | | | X | | | | | X | | | | | | | | | | | | X | | | | 5 | |
| C-cell carcinoma | | | | | | | | | | | | | | | | | | | | | | | | | | 2 | |
| Parathyroid | + | + | + | - | + | - | - | + | + | - | + | - | + | - | + | - | + | + | + | + | - | - | - | - | - | 27 | |
| REPRODUCTIVE SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Mammary gland | + | + | + | N | + | N | + | + | N | + | + | + | + | N | N | + | + | N | + | + | + | + | + | + | + | *50 2 | |
| Fibroadenoma | | | | | | | | | | | | | | | | | | | | | | X | | X | | | |
| Uterus | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | - | + | + | + | + | 45 | |
| Endometrial stromal polyp | | | X | | X | | | | | | | X | | | | | | | | | | | X | | | 5 | |
| Ovary | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | - | + | + | + | + | 45 | |
| NERVOUS SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Brain | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | 50 | |
| MUSCULOSKELETAL SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Bone | N | + | N | N | N | N | N | N | + | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | *50 1 | |
| Sarcoma, NOS | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| BODY CAVITIES | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Pleura | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | *50 1 | |
| Fibrosarcoma | | | | | | | | | | | | | | | | | | | | | | | X | | | | |
| Mediastinum | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | *50 1 | |
| Nonchromaffin paraganglioma | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| ALL OTHER SYSTEMS | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Multiple organs, NOS | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | *50 1 | |
| Malignant lymphoma, undiffer type | | | | | | | | | | | | | | | | | | | | | | | | | | | |

* Animals necropsied

TABLE F2. INDIVIDUAL ANIMAL TUMOR PATHOLOGY OF FEMALE MARSHALL RATS IN THE TWO-YEAR GAVAGE STUDY OF TRICHLOROETHYLENE: HIGH DOSE

| ANIMAL NUMBER | 06 | 07 | 09 | 03 | 01 | 01 | 01 | 02 | 02 | 02 | 03 | 03 | 03 | 04 | 04 | 04 | 04 | 05 | 05 | 05 | 06 | 06 | 06 | 06 | 07 | 07 | 08 | 08 | 08 | 08 | 08 | 08 | 08 |
|--------------------------------|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|
| WEEKS ON STUDY | 10 | 06 | 06 | 07 | 07 | 03 | 08 | 08 | 07 | 07 | 05 | 00 | 04 | 05 | 06 | 00 | 09 | 08 | 00 | 07 | 08 | 00 | 01 | 04 | 07 | 08 | 00 | 01 | 01 | 08 | 03 | 00 | |
| RESPIRATORY SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Lungs and bronchi | + | + | + | + | + | A | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| Pheochromocytoma, metastatic | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Trachea | + | + | - | + | + | A | + | + | + | + | + | + | + | + | + | - | + | - | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| HEMATOPOIETIC SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Bone marrow | + | + | + | + | + | A | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | - | + | + | |
| Spleen | + | + | + | + | + | A | + | + | + | + | + | + | + | + | + | - | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| Lymph nodes | - | - | + | + | - | A | - | + | - | - | + | - | - | - | - | - | + | - | - | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| Thymus | - | + | - | - | - | A | - | + | + | - | + | - | + | + | + | - | + | - | - | + | + | + | + | + | + | + | + | + | + | + | + | - | |
| CIRCULATORY SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Heart | + | + | + | + | + | A | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| DIGESTIVE SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Salivary gland | + | + | + | + | + | A | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| Liver | + | + | + | + | + | A | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| Bile duct | + | + | + | + | + | A | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| Gallbladder & common bile duct | N | N | N | N | N | A | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | |
| Pancreas | + | + | + | + | + | A | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| Esophagus | + | + | + | + | + | A | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| Stomach | - | + | + | + | + | A | + | + | + | + | + | + | + | + | + | - | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| Squamous cell papilloma | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Small intestine | + | + | + | + | + | A | + | + | + | + | + | + | + | + | + | - | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| Large intestine | + | + | + | + | + | A | + | + | + | - | + | + | + | + | + | - | + | + | + | + | + | + | + | + | + | + | + | + | + | - | + | + | |
| URINARY SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Kidney | + | + | + | + | + | A | + | + | + | + | + | + | + | + | + | - | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| Tubular cell adenocarcinoma | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Kidney/pelvis | + | + | + | + | + | A | + | + | + | + | + | + | + | + | + | - | + | + | + | + | + | + | + | + | + | + | + | + | + | X | + | + | |
| Transitional cell papilloma | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Transitional cell carcinoma | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Urinary bladder | + | + | + | + | + | A | + | + | + | + | + | + | + | + | + | - | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| ENDOCRINE SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Pituitary | - | + | + | + | - | A | + | - | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| Chromophobe adenoma | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Adrenal | + | + | + | + | + | A | + | + | + | + | + | + | + | + | + | - | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| Cortical adenoma | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Pheochromocytoma | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Pheochromocytoma, malignant | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Thyroid | + | + | + | + | + | A | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| C-cell adenoma | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| C-cell carcinoma | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Parathyroid | - | - | + | - | - | A | - | - | + | - | - | + | - | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | | |
| REPRODUCTIVE SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Mammary gland | + | + | + | N | + | A | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | | |
| Fibroadenoma | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Vagina | N | N | N | N | N | A | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | |
| Papilloma, NOS | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Uterus | + | + | + | + | + | A | + | + | + | + | + | + | + | + | + | - | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | | |
| Carcinoma, NOS | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Endometrial stromal polyp | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Ovary | + | + | + | + | + | A | + | + | + | + | + | + | + | + | + | - | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | | |
| NERVOUS SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Brain | + | + | + | + | + | A | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | | |

TABLE F3. ANALYSIS OF PRIMARY TUMORS IN FEMALE MARSHALL RATS IN THE TWO-YEAR GAVAGE STUDY OF TRICHLOROETHYLENE

| | Untreated Control | Vehicle Control | 500 mg/kg | 1,000 mg/kg |
|---|-------------------|-----------------|-------------|-------------|
| Kidney: Transitional Cell Papilloma or Carcinoma | | | | |
| Overall Rates (a) | 1/49 (2%) | 1/50 (2%) | 2/48 (4%) | 3/44 (7%) |
| Adjusted Rates (b) | 3.2% | 3.3% | 6.7% | 15.5% |
| Terminal Rates (c) | 1/31 (3%) | 1/30 (3%) | 0/12 (0%) | 1/10 (10%) |
| Life Table Tests (d) | | P=0.060 | P=0.302 | P=0.095 |
| Incidental Tumor Tests (d) | | P=0.205 | P=0.515 | P=0.278 |
| Cochran-Armitage Trend Test (d) | | P=0.184 | | |
| Fisher Exact Test | | | P=0.485 | P=0.262 |
| Pituitary: Chromophobe Adenoma | | | | |
| Overall Rates (a) | 22/42 (52%) | 18/45 (40%) | 8/38 (21%) | 6/42 (14%) |
| Adjusted Rates (b) | 64.4% | 51.7% | 36.3% | 39.2% |
| Terminal Rates (c) | 15/26 (58%) | 11/26 (42%) | 2/11 (18%) | 2/9 (22%) |
| Life Table Tests (d) | | P=0.412N | P=0.482N | P=0.506N |
| Incidental Tumor Tests (d) | | P=0.036N | P=0.097N | P=0.079N |
| Cochran-Armitage Trend Test (d) | | P=0.004N | | |
| Fisher Exact Test | | | P=0.052N | P=0.007N |
| Pituitary: Chromophobe Adenoma or Carcinoma | | | | |
| Overall Rates (a) | 23/42 (55%) | 19/45 (42%) | 8/38 (21%) | 6/42 (14%) |
| Adjusted Rates (b) | 65.5% | 53.1% | 36.3% | 39.2% |
| Terminal Rates (c) | 15/26 (58%) | 11/26 (42%) | 2/11 (18%) | 2/9 (22%) |
| Life Table Tests (d) | | P=0.357N | P=0.427N | P=0.458N |
| Incidental Tumor Tests (d) | | P=0.023N | P=0.062N | P=0.057N |
| Cochran-Armitage Trend Test (d) | | P=0.002N | | |
| Fisher Exact Test | | | P=0.034N | P=0.004N |
| Pituitary: Adenoma | | | | |
| Overall Rates (a) | 22/42 (52%) | 19/45 (42%) | 8/38 (21%) | 6/42 (14%) |
| Adjusted Rates (b) | 64.4% | 54.9% | 36.3% | 39.2% |
| Terminal Rates (c) | 15/26 (58%) | 12/26 (46%) | 2/11 (18%) | 2/9 (22%) |
| Life Table Tests (d) | | P=0.354N | P=0.429N | P=0.453N |
| Incidental Tumor Tests (d) | | P=0.026N | P=0.075N | P=0.062N |
| Cochran-Armitage Trend Test (d) | | P=0.002N | | |
| Fisher Exact Test | | | P=0.034N | P=0.004N |
| Pituitary: Adenoma or Carcinoma | | | | |
| Overall Rates (a) | 25/42 (60%) | 20/45 (44%) | 8/38 (21%) | 6/42 (14%) |
| Adjusted Rates (b) | 67.3% | 56.2% | 36.3% | 39.2% |
| Terminal Rates (c) | 15/26 (58%) | 12/26 (46%) | 2/11 (18%) | 2/9 (22%) |
| Life Table Tests (d) | | P=0.304N | P=0.376N | P=0.408N |
| Incidental Tumor Tests (d) | | P=0.016N | P=0.047N | P=0.044N |
| Cochran-Armitage Trend Test (d) | | P=0.001N | | |
| Fisher Exact Test | | | P=0.021N | P=0.002N |
| Adrenal: Cortical Adenoma | | | | |
| Overall Rates (a) | 4/47 (9%) | 5/49 (10%) | 2/47 (4%) | 3/43 (7%) |
| Adjusted Rates (b) | 12.1% | 15.9% | 13.4% | 18.8% |
| Terminal Rates (c) | 3/31 (10%) | 4/29 (14%) | 1/12 (8%) | 1/10 (10%) |
| Life Table Tests (d) | | P=0.346 | P=0.620N | P=0.394 |
| Incidental Tumor Tests (d) | | P=0.551 | P=0.514N | P=0.649N |
| Cochran-Armitage Trend Test (d) | | P=0.333N | | |
| Fisher Exact Test | | | P=0.235N | P=0.433N |
| Adrenal: Pheochromocytoma | | | | |
| Overall Rates (a) | 30/47 (64%) | 39/49 (80%) | 15/47 (32%) | 9/43 (21%) |
| Adjusted Rates (b) | 74.8% | 92.7% | 76.5% | 67.0% |
| Terminal Rates (c) | 21/31 (68%) | 26/29 (90%) | 8/12 (67%) | 6/10 (60%) |
| Life Table Tests (d) | | P=0.050N | P=0.263N | P=0.067N |
| Incidental Tumor Tests (d) | | P<0.001N | P=0.002N | P<0.001N |
| Cochran-Armitage Trend Test (d) | | P<0.001N | | |
| Fisher Exact Test | | | P<0.001N | P<0.001N |

TABLE F3. ANALYSIS OF PRIMARY TUMORS IN FEMALE MARSHALL RATS IN THE TWO-YEAR GAVAGE STUDY OF TRICHLOROETHYLENE (Continued)

| | Untreated Control | Vehicle Control | 500 mg/kg | 1,000 mg/kg |
|---|-------------------|-----------------|-------------|-------------|
| Adrenal: Pheochromocytoma, Malignant | | | | |
| Overall Rates (a) | 4/47 (9%) | 4/49 (8%) | 0/47 (0%) | 1/43 (2%) |
| Adjusted Rates (b) | 12.9% | 13.8% | 0.0% | 10.0% |
| Terminal Rates (c) | 4/31 (13%) | 4/29 (14%) | 0/12 (0%) | 1/10 (10%) |
| Life Table Tests (d) | | P=0.354N | P=0.222N | P=0.593N |
| Incidental Tumor Tests (d) | | P=0.354N | P=0.222N | P=0.593N |
| Cochran-Armitage Trend Test (d) | | P=0.100N | | |
| Fisher Exact Test | | | P=0.064N | P=0.224N |
| Adrenal: Pheochromocytoma or Pheochromocytoma, Malignant | | | | |
| Overall Rates (a) | 32/47 (68%) | 40/49 (82%) | 15/47 (32%) | 9/43 (21%) |
| Adjusted Rates (b) | 79.8% | 95.1% | 76.5% | 67.0% |
| Terminal Rates (c) | 23/31 (74%) | 27/29 (93%) | 8/12 (67%) | 6/10 (60%) |
| Life Table Tests (d) | | P=0.036N | P=0.222N | P=0.050N |
| Incidental Tumor Tests (d) | | P<0.001N | P<0.001N | P<0.001N |
| Cochran-Armitage Trend Test (d) | | P<0.001N | | |
| Fisher Exact Test | | | P<0.001N | P<0.001N |
| Thyroid: C-Cell Adenoma | | | | |
| Overall Rates (a) | 12/48 (25%) | 10/43 (23%) | 5/46 (11%) | 4/43 (9%) |
| Adjusted Rates (b) | 37.3% | 31.8% | 45.5% | 26.2% |
| Terminal Rates (c) | 11/31 (35%) | 9/30 (30%) | 5/11 (45%) | 2/10 (20%) |
| Life Table Tests (d) | | P=0.432 | P=0.394 | P=0.561 |
| Incidental Tumor Tests (d) | | P=0.552N | P=0.423 | P=0.473N |
| Cochran-Armitage Trend Test (d) | | P=0.046N | | |
| Fisher Exact Test | | | P=0.101N | P=0.071N |
| Thyroid: C-Cell Adenoma or Carcinoma | | | | |
| Overall Rates (a) | 13/48 (27%) | 10/43 (23%) | 7/46 (15%) | 5/43 (12%) |
| Adjusted Rates (b) | 40.4% | 31.8% | 50.1% | 35.5% |
| Terminal Rates (c) | 12/31 (39%) | 9/30 (30%) | 5/11 (45%) | 3/10 (30%) |
| Life Table Tests (d) | | P=0.219 | P=0.150 | P=0.364 |
| Incidental Tumor Tests (d) | | P=0.403 | P=0.249 | P=0.593 |
| Cochran-Armitage Trend Test (d) | | P=0.096N | | |
| Fisher Exact Test | | | P=0.244N | P=0.128N |
| Mammary Gland: Adenocarcinoma | | | | |
| Overall Rates (a) | 1/50 (2%) | 3/50 (6%) | 0/50 (0%) | 0/50 (0%) |
| Adjusted Rates (b) | 3.0% | 10.0% | 0.0% | 0.0% |
| Terminal Rates (c) | 0/31 (0%) | 3/30 (10%) | 0/12 (0%) | 0/10 (0%) |
| Life Table Tests (d) | | P=0.157N | P=0.320N | P=0.366N |
| Incidental Tumor Tests (d) | | P=0.157N | P=0.320N | P=0.366N |
| Cochran-Armitage Trend Test (d) | | P=0.037N | | |
| Fisher Exact Test | | | P=0.121N | P=0.121N |
| Mammary Gland: Fibroadenoma | | | | |
| Overall Rates (a) | 7/50 (14%) | 11/50 (22%) | 2/50 (4%) | 4/48 (8%) |
| Adjusted Rates (b) | 19.8% | 28.2% | 16.7% | 24.2% |
| Terminal Rates (c) | 5/31 (16%) | 5/30 (17%) | 2/12 (17%) | 0/10 (0%) |
| Life Table Tests (d) | | P=0.418N | P=0.159N | P=0.601N |
| Incidental Tumor Tests (d) | | P=0.087N | P=0.024N | P=0.120N |
| Cochran-Armitage Trend Test (d) | | P=0.023N | | |
| Fisher Exact Test | | | P=0.007N | P=0.054N |
| Mammary Gland: Fibroadenoma or Adenocarcinoma | | | | |
| Overall Rates (a) | 8/50 (16%) | 14/50 (28%) | 2/50 (4%) | 4/48 (8%) |
| Adjusted Rates (b) | 22.2% | 36.8% | 16.7% | 24.2% |
| Terminal Rates (c) | 5/31 (16%) | 8/30 (27%) | 2/12 (17%) | 0/10 (0%) |
| Life Table Tests (d) | | P=0.228N | P=0.076N | P=0.429N |
| Incidental Tumor Tests (d) | | P=0.032N | P=0.009N | P=0.062N |
| Cochran-Armitage Trend Test (d) | | P=0.003N | | |
| Fisher Exact Test | | | P=0.001N | P=0.011N |

TABLE F3. ANALYSIS OF PRIMARY TUMORS IN FEMALE MARSHALL RATS IN THE TWO-YEAR GAVAGE STUDY OF TRICHLOROETHYLENE (Continued)

| | Untreated Control | Vehicle Control | 500 mg/kg | 1,000 mg/kg |
|--|-------------------|-----------------|------------|-------------|
| Uterus: Endometrial Stromal Polyp | | | | |
| Overall Rates (a) | 6/47 (13%) | 3/50 (6%) | 5/45 (11%) | 2/44 (5%) |
| Adjusted Rates (b) | 17.0% | 8.2% | 23.2% | 20.0% |
| Terminal Rates (c) | 3/30 (10%) | 1/30 (3%) | 2/12 (17%) | 2/10 (20%) |
| Life Table Tests (d) | | P=0.244 | P=0.107 | P=0.409 |
| Incidental Tumor Tests (d) | | P=0.533 | P=0.386 | P=0.505 |
| Cochran-Armitage Trend Test (d) | | P=0.489N | | |
| Fisher Exact Test | | | P=0.300 | P=0.561 |

(a) Number of tumor-bearing animals/number of animals examined at the site

(b) Kaplan-Meier estimated tumor incidence at the end of the study after adjusting for intercurrent mortality

(c) Observed tumor incidence at terminal kill

(d) Beneath the control incidence are the P values associated with the trend test. Beneath the dosed group incidence are the P values corresponding to pairwise comparisons between that dosed group and the vehicle controls. The life table analysis regards tumors in animals dying prior to terminal kill as being (directly or indirectly) the cause of death. The incidental tumor test regards these lesions as nonfatal. The Cochran-Armitage and Fisher exact tests compare directly the overall incidence rates. A regular trend or lower incidence in a dosed group is indicated by (N).

TABLE F4. SUMMARY OF THE INCIDENCE OF NONNEOPLASTIC LESIONS IN FEMALE MARSHALL RATS IN THE TWO-YEAR GAVAGE STUDY OF TRICHLOROETHYLENE

| | CONTROL (UNTR) | CONTROL (VEH) | LOW DOSE | HIGH DOSE |
|--------------------------------------|----------------|---------------|----------|-----------|
| ANIMALS INITIALLY IN STUDY | 50 | 50 | 50 | 50 |
| ANIMALS NECROPSIED | 50 | 50 | 50 | 48 |
| ANIMALS EXAMINED HISTOPATHOLOGICALLY | 50 | 50 | 50 | 48 |
| INTEGUMENTARY SYSTEM | | | | |
| *Subcutaneous tissue | (50) | (50) | (50) | (48) |
| Inflammation, necrotizing | | | | 1 (2%) |
| Foreign material, NOS | | | 1 (2%) | 1 (2%) |
| RESPIRATORY SYSTEM | | | | |
| *Nasal cavity | (50) | (50) | (50) | (48) |
| Inflammation, NOS | | | 1 (2%) | |
| Inflammation, suppurative | | 2 (4%) | 6 (12%) | 1 (2%) |
| Inflammation, acute/chronic | | | | 1 (2%) |
| Inflammation chronic suppurative | | | 1 (2%) | |
| *Larynx | (50) | (50) | (50) | (48) |
| Granuloma, foreign body | | | 1 (2%) | |
| #Trachea | (47) | (39) | (45) | (38) |
| Wound, NOS | | | 1 (2%) | |
| Lacerated wound | | | | 1 (3%) |
| Penetrating wound | | | | 1 (3%) |
| Inflammation, acute focal | 1 (2%) | | | |
| Foreign material, NOS | 1 (2%) | | 2 (4%) | 3 (8%) |
| #Peritracheal tissue | (47) | (39) | (45) | (38) |
| Abscess, chronic | | | 1 (2%) | |
| #Lung | (50) | (49) | (49) | (46) |
| Emphysema, alveolar | | | | 2 (4%) |
| Collapse | 2 (4%) | | 3 (6%) | 1 (2%) |
| Congestion, NOS | 5 (10%) | 6 (12%) | 10 (20%) | 12 (26%) |
| Congestion, chronic passive | 1 (2%) | | | |
| Edema, NOS | 1 (2%) | 1 (2%) | 3 (6%) | 2 (4%) |
| Hemorrhage | 5 (10%) | 6 (12%) | 9 (18%) | 10 (22%) |
| Bronchopneumonia, NOS | | 2 (4%) | 1 (2%) | |
| Bronchopneumonia, focal | | | 2 (4%) | |
| Inflammation, focal | 1 (2%) | 19 (39%) | 10 (20%) | 2 (4%) |
| Inflammation, interstitial | | 1 (2%) | | |
| Inflammation, necrotizing | | | 1 (2%) | |
| Bronchopneumonia, acute | 2 (4%) | 2 (4%) | 2 (4%) | 1 (2%) |
| Inflammation, acute focal | | 1 (2%) | | |
| Abscess, NOS | | | 1 (2%) | |
| Inflammation, chronic focal | | 6 (12%) | 11 (22%) | 7 (15%) |
| Inflammation, granulomatous focal | 3 (6%) | 6 (12%) | 1 (2%) | 11 (24%) |
| Foreign material, NOS | 1 (2%) | 1 (2%) | 5 (10%) | 9 (20%) |
| Hemosiderosis | | 1 (2%) | | |
| Russell body | 1 (2%) | | | |
| Alveolar macrophages | | 1 (2%) | 1 (2%) | 1 (2%) |
| Hyperplasia, alveolar epithelium | 1 (2%) | | | 1 (2%) |
| HEMATOPOIETIC SYSTEM | | | | |
| #Bone marrow | (50) | (50) | (48) | (44) |
| Hyperplasia, granulocytic | | | 2 (4%) | |
| #Spleen | (49) | (49) | (47) | (43) |
| Congestion, NOS | | 1 (2%) | 1 (2%) | |
| Hemosiderosis | 4 (8%) | 2 (4%) | 2 (4%) | |
| Hematopoiesis | 6 (12%) | 12 (24%) | 5 (11%) | 3 (7%) |
| Erythropoiesis | 1 (2%) | | | |
| #Mandibular lymph node | (31) | (31) | (29) | (23) |
| Hemorrhage | | | 1 (3%) | 1 (4%) |

TABLE F4. SUMMARY OF THE INCIDENCE OF NONNEOPLASTIC LESIONS IN FEMALE MARSHALL RATS IN THE TWO-YEAR GAVAGE STUDY OF TRICHLOROETHYLENE (Continued)

| | CONTROL (UNTR) | CONTROL (VEH) | LOW DOSE | HIGH DOSE |
|---|----------------|---------------|----------|-----------|
| HEMATOPOIETIC SYSTEM (Continued) | | | | |
| #Mesenteric lymph node | (31) | (31) | (29) | (23) |
| Hemorrhage | 2 (6%) | | 2 (7%) | |
| Inflammation, chronic focal | 1 (3%) | | | |
| #Liver | (50) | (49) | (48) | (46) |
| Hematopoiesis | | | 1 (2%) | |
| #Thymus | (6) | (7) | (16) | (21) |
| Multiple cysts | | 1 (14%) | | |
| Hemorrhage | | | 1 (6%) | |
| Involution, NOS | | 1 (14%) | | |
| CIRCULATORY SYSTEM | | | | |
| *Multiple organs | (50) | (50) | (50) | (48) |
| Arteriosclerosis | | | 1 (2%) | |
| #Lung | (50) | (49) | (49) | (46) |
| Thrombosis, NOS | | 1 (2%) | | |
| Perivasculitis | 1 (2%) | | 1 (2%) | |
| #Heart | (50) | (49) | (50) | (46) |
| Thrombus, mural | | 1 (2%) | | |
| Thrombus, fibrin | | | 1 (2%) | |
| Inflammation, focal | 1 (2%) | | | |
| Fibrosis | 2 (4%) | | | |
| Fibrosis, focal | 2 (4%) | | 1 (2%) | 1 (2%) |
| Periarteritis | | 1 (2%) | | |
| Calcification, focal | 1 (2%) | 1 (2%) | | |
| *Aorta | (50) | (50) | (50) | (48) |
| Polyangiitis | | | | 1 (2%) |
| Calcification, metastatic | | | | 1 (2%) |
| *Pulmonary artery | (50) | (50) | (50) | (48) |
| Calcification, focal | 5 (10%) | 1 (2%) | | |
| #Pancreas | (48) | (49) | (46) | (42) |
| Periarteritis | | | 1 (2%) | |
| *Mesentery | (50) | (50) | (50) | (48) |
| Periarteritis | | | 1 (2%) | |
| DIGESTIVE SYSTEM | | | | |
| #Salivary gland | (47) | (46) | (47) | (44) |
| Inflammation, NOS | | | | 1 (2%) |
| Inflammation, acute | | | 1 (2%) | |
| Inflammation, chronic | | | | 1 (2%) |
| Atrophy, focal | | 1 (2%) | | |
| #Liver | (50) | (49) | (48) | (46) |
| Congestion, NOS | 2 (4%) | 2 (4%) | 1 (2%) | 1 (2%) |
| Congestion, passive | 1 (2%) | 1 (2%) | | 1 (2%) |
| Congestion, acute passive | | | 1 (2%) | |
| Inflammation, focal | 1 (2%) | 1 (2%) | 1 (2%) | |
| Necrosis, NOS | 1 (2%) | 1 (2%) | | |
| Necrosis, focal | 2 (4%) | | 2 (4%) | 1 (2%) |
| Necrosis, central | 1 (2%) | 1 (2%) | 1 (2%) | 2 (4%) |
| Metamorphosis, fatty | 1 (2%) | 1 (2%) | 3 (6%) | 1 (2%) |
| Mitotic alteration | | 1 (2%) | | |
| Basophilic cyto change | 1 (2%) | | | |
| Eosinophilic cyto change | 1 (2%) | | | |
| Clear cell change | 2 (4%) | 7 (14%) | | 2 (4%) |
| Atrophy, NOS | 1 (2%) | | | |
| Depletion, glycogen | 1 (2%) | | | |

TABLE F4. SUMMARY OF THE INCIDENCE OF NONNEOPLASTIC LESIONS IN FEMALE MARSHALL RATS IN THE TWO-YEAR GAVAGE STUDY OF TRICHLOROETHYLENE (Continued)

| | CONTROL (UNTR) | CONTROL (VEH) | LOW DOSE | HIGH DOSE |
|-------------------------------------|----------------|---------------|----------|-----------|
| DIGESTIVE SYSTEM (Continued) | | | | |
| #Esophagus | (49) | (50) | (50) | (46) |
| Lacerated wound | | 3 (6%) | 3 (6%) | 6 (13%) |
| Diverticulum | | | | 1 (2%) |
| Impaction, NOS | 2 (4%) | | | |
| Inflammation, granulomatous | | 1 (2%) | | |
| #Stomach | (49) | (49) | (46) | (43) |
| Calcification, metastatic | | | | 1 (2%) |
| #Gastric mucosa | (49) | (49) | (46) | (43) |
| Inflammation, focal | | | 1 (2%) | |
| Hyperkeratosis | | | 1 (2%) | |
| #Colon | (44) | (45) | (43) | (41) |
| Hemorrhage | | | | 1 (2%) |
| Parasitism | 3 (7%) | 3 (7%) | 2 (5%) | 4 (10%) |
| *Rectum | (50) | (50) | (50) | (48) |
| Parasitism | | | | 1 (2%) |
| URINARY SYSTEM | | | | |
| #Kidney | (49) | (50) | (48) | (44) |
| Congenital hydronephrosis | 23 (47%) | 21 (42%) | 17 (35%) | 12 (27%) |
| Calculus, unknown gross or micro | | | 1 (2%) | 1 (2%) |
| Hydronephrosis | | | | 5 (11%) |
| Hemorrhage | | 1 (2%) | | |
| Pyelonephritis, NOS | | | | 1 (2%) |
| Pyelonephritis, acute | | | 1 (2%) | |
| Pyelonephritis, healed | 1 (2%) | | | |
| Nephropathy | 49 (100%) | 50 (100%) | 47 (98%) | 39 (89%) |
| Nephropathy, toxic | | | 30 (63%) | 30 (68%) |
| Degeneration, hyaline | 1 (2%) | | | |
| Calcification, focal | 19 (39%) | 11 (22%) | 12 (25%) | 13 (30%) |
| Cytomegaly | | | 46 (96%) | 43 (98%) |
| Hyperplasia, tubular cell | 1 (2%) | 1 (2%) | | 1 (2%) |
| Angiectasis | | | | 1 (2%) |
| #Renal papilla | (49) | (50) | (48) | (44) |
| Necrosis, focal | | | 1 (2%) | |
| #Kidney/pelvis | (49) | (50) | (48) | (44) |
| Calculus, unknown gross or micro | 6 (12%) | | | |
| Dilatation, NOS | | | 1 (2%) | |
| Hemorrhage | | 1 (2%) | | |
| Hematoma, NOS | | 1 (2%) | | |
| Calcium deposit | 1 (2%) | | | |
| Hyperplasia, epithelial | 2 (4%) | | 2 (4%) | 1 (2%) |
| Metaplasia, squamous | 1 (2%) | | | |
| #Urinary bladder | (45) | (46) | (40) | (40) |
| Calculus, unknown gross or micro | | | | 1 (3%) |
| Hemorrhage | | | | 1 (3%) |
| Hyperplasia, epithelial | | | | 1 (3%) |
| Metaplasia, squamous | 1 (2%) | | | 1 (3%) |
| ENDOCRINE SYSTEM | | | | |
| #Pituitary | (42) | (45) | (38) | (42) |
| Cyst, NOS | 1 (2%) | | 1 (3%) | |
| Multiple cysts | | 1 (2%) | | |
| Hemorrhage | | 2 (4%) | | 1 (2%) |
| Hyperplasia, focal | 1 (2%) | | | |
| Hyperplasia, chromophobe cell | 2 (5%) | 2 (4%) | 4 (11%) | 2 (5%) |

TABLE F4. SUMMARY OF THE INCIDENCE OF NONNEOPLASTIC LESIONS IN FEMALE MARSHALL RATS IN THE TWO-YEAR GAVAGE STUDY OF TRICHLOROETHYLENE (Continued)

| | CONTROL (UNTR) | CONTROL (VEH) | LOW DOSE | HIGH DOSE |
|-------------------------------------|----------------|---------------|----------|-----------|
| ENDOCRINE SYSTEM (Continued) | | | | |
| #Adrenal | (47) | (49) | (47) | (43) |
| Cyst, NOS | | 1 (2%) | | |
| Hemorrhage | | 1 (2%) | 1 (2%) | 1 (2%) |
| Hematoma, NOS | 1 (2%) | 1 (2%) | 1 (2%) | 1 (2%) |
| Degeneration, lipid | | 1 (2%) | 1 (2%) | 1 (2%) |
| Hyperplasia, focal | 1 (2%) | | | |
| Angiectasis | 2 (4%) | 5 (10%) | 9 (19%) | 6 (14%) |
| #Adrenal cortex | (47) | (49) | (47) | (43) |
| Hemorrhage | 6 (13%) | 3 (6%) | 7 (15%) | 5 (12%) |
| Hematoma, NOS | | | | 1 (2%) |
| Degeneration, lipid | 4 (9%) | 3 (6%) | 4 (9%) | 3 (7%) |
| Necrosis, focal | | | 1 (2%) | |
| Metamorphosis, fatty | | | | 1 (2%) |
| Lipoidosis | | | 1 (2%) | |
| Atrophy, NOS | | | | 1 (2%) |
| Atrophy, diffuse | | 1 (2%) | | |
| Hyperplasia, focal | 9 (19%) | 5 (10%) | 4 (9%) | 3 (7%) |
| Hyperplasia, diffuse | | 1 (2%) | | |
| Angiectasis | 4 (9%) | 1 (2%) | 5 (11%) | 2 (5%) |
| #Adrenal medulla | (47) | (49) | (47) | (43) |
| Hyperplasia, focal | 16 (34%) | 4 (8%) | 5 (11%) | 2 (5%) |
| #Thyroid | (48) | (43) | (46) | (43) |
| Follicular cyst, NOS | | | 1 (2%) | |
| Hyperplasia, C-cell | 11 (23%) | 6 (14%) | 5 (11%) | 6 (14%) |
| Angiectasis | 1 (2%) | | | |
| #Parathyroid | (27) | (22) | (27) | (22) |
| Hyperplasia, NOS | | | | 1 (5%) |
| REPRODUCTIVE SYSTEM | | | | |
| *Mammary gland | (50) | (50) | (50) | (48) |
| Galactocele | | 1 (2%) | 1 (2%) | |
| Hyperplasia, cystic | | 1 (2%) | | |
| Lactation | 4 (8%) | | 1 (2%) | |
| #Uterus | (47) | (50) | (45) | (44) |
| Hydrometra | 1 (2%) | | | 3 (7%) |
| Hematometra | | 1 (2%) | 1 (2%) | |
| Pyometra | | 1 (2%) | | |
| Hemosiderosis | 2 (4%) | 1 (2%) | 1 (2%) | |
| #Cervix uteri | (47) | (50) | (45) | (44) |
| Cyst, NOS | 1 (2%) | 2 (4%) | | |
| Multiple cysts | 1 (2%) | | | |
| Inflammation, focal | | | 1 (2%) | |
| #Uterus/endometrium | (47) | (50) | (45) | (44) |
| Multilocular cyst | | | | 1 (2%) |
| Inflammation, NOS | | | | 1 (2%) |
| Hemosiderosis | | | | 3 (7%) |
| #Ovary | (47) | (48) | (45) | (44) |
| Cyst, NOS | 4 (9%) | 4 (8%) | 2 (4%) | 2 (5%) |
| Luteinized follic cyst | | | 1 (2%) | |
| Multiple cysts | | 1 (2%) | | |
| Parovarian cyst | | | 1 (2%) | |
| Hemosiderosis | | 1 (2%) | | |

TABLE F4. SUMMARY OF THE INCIDENCE OF NONNEOPLASTIC LESIONS IN FEMALE MARSHALL RATS IN THE TWO-YEAR GAVAGE STUDY OF TRICHLOROETHYLENE (Continued)

| | CONTROL (UNTR) | CONTROL (VEH) | LOW DOSE | HIGH DOSE |
|-----------------------------------|----------------|---------------|----------|-----------|
| NERVOUS SYSTEM | | | | |
| #Brain | (50) | (50) | (50) | (48) |
| Hemorrhage | 2 (4%) | 1 (2%) | 1 (2%) | |
| Calcification, focal | | | | 1 (2%) |
| SPECIAL SENSE ORGANS | | | | |
| *Eye | (50) | (50) | (50) | (48) |
| Retinopathy | | | | 1 (2%) |
| Cataract | | | | 1 (2%) |
| *Harderian gland | (50) | (50) | (50) | (48) |
| Inflammation, acute/chronic | | | | 2 (4%) |
| Inflammation, chronic | | | 2 (4%) | |
| Inflammation, chronic focal | 1 (2%) | | 2 (4%) | 4 (8%) |
| Atrophy, focal | 1 (2%) | | | |
| MUSCULOSKELETAL SYSTEM | | | | |
| None | | | | |
| BODY CAVITIES | | | | |
| *Pleural cavity | (50) | (50) | (50) | (48) |
| Hydrothorax | | 1 (2%) | | |
| *Pericardium | (50) | (50) | (50) | (48) |
| Inflammation, chronic | | | | 1 (2%) |
| ALL OTHER SYSTEMS | | | | |
| Lumbar region | | | | |
| Fracture, NOS | 1 | | | |
| Lower leg | | | | |
| Abscess, chronic | | | 1 | |
| SPECIAL MORPHOLOGY SUMMARY | | | | |
| Auto/necropsy/histo perf | | | | 1 |
| Autolysis/no necropsy | | | | 2 |

* Number of animals receiving complete necropsy examination; all gross lesions including masses examined microscopically.

Number of animals examined microscopically at this site

APPENDIX G

SUMMARY OF LESIONS IN MALE OSBORNE-MENDEL RATS IN THE TWO-YEAR GAVAGE STUDY OF TRICHLOROETHYLENE

| | PAGE | |
|----------|---|-----|
| TABLE G1 | SUMMARY OF THE INCIDENCE OF NEOPLASMS IN MALE OSBORNE-MENDEL RATS IN THE TWO-YEAR GAVAGE STUDY OF TRICHLOROETHYLENE | 209 |
| TABLE G2 | INDIVIDUAL ANIMAL TUMOR PATHOLOGY OF MALE OSBORNE-MENDEL RATS IN THE TWO-YEAR GAVAGE STUDY OF TRICHLOROETHYLENE | 212 |
| TABLE G3 | ANALYSIS OF PRIMARY TUMORS IN MALE OSBORNE-MENDEL RATS IN THE TWO-YEAR GAVAGE STUDY OF TRICHLOROETHYLENE | 220 |
| TABLE G4 | SUMMARY OF THE INCIDENCE OF NONNEOPLASTIC LESIONS IN MALE OSBORNE-MENDEL RATS IN THE TWO-YEAR GAVAGE STUDY OF TRICHLOROETHYLENE | 223 |

TABLE G1. SUMMARY OF THE INCIDENCE OF NEOPLASMS IN MALE OSBORNE-MENDEL RATS IN THE TWO-YEAR GAVAGE STUDY OF TRICHLOROETHYLENE

| | CONTROL (UNTR) | CONTROL (VEH) | LOW DOSE | HIGH DOSE |
|--------------------------------------|----------------|---------------|----------|-----------|
| ANIMALS INITIALLY IN STUDY | 50 | 50 | 50 | 50 |
| ANIMALS NECROPSIED | 50 | 50 | 50 | 50 |
| ANIMALS EXAMINED HISTOPATHOLOGICALLY | 50 | 50 | 50 | 50 |
| INTEGUMENTARY SYSTEM | | | | |
| *Skin | (50) | (50) | (50) | (50) |
| Squamous cell carcinoma | 1 (2%) | | 1 (2%) | |
| Keratoacanthoma | | | 1 (2%) | 2 (4%) |
| Fibroma | | | | |
| *Subcutaneous tissue | (50) | (50) | (50) | (50) |
| Sarcoma, NOS | 2 (4%) | | 2 (4%) | |
| Fibroma | 3 (6%) | 4 (8%) | 3 (6%) | |
| RESPIRATORY SYSTEM | | | | |
| *Nasal cavity | (50) | (50) | (50) | (50) |
| Carcinoma, NOS | 1 (2%) | | | |
| #Lung | (50) | (50) | (50) | (50) |
| Alveolar/bronchiolar adenoma | | 1 (2%) | | 1 (2%) |
| Alveolar/bronchiolar carcinoma | | 1 (2%) | | |
| Sarcoma, NOS, metastatic | | | 1 (2%) | |
| HEMATOPOIETIC SYSTEM | | | | |
| *Multiple organs | (50) | (50) | (50) | (50) |
| Leukemia, NOS | | | | 1 (2%) |
| Monocytic leukemia | 1 (2%) | 1 (2%) | | 1 (2%) |
| *Thorax | (50) | (50) | (50) | (50) |
| Plasma cell myeloma | | | | 1 (2%) |
| #Pancreas | (50) | (49) | (50) | (50) |
| Malignant lymphoma, NOS | 1 (2%) | | | |
| #Ileum | (48) | (49) | (50) | (48) |
| Malignant lymphoma, NOS | | 1 (2%) | | |
| CIRCULATORY SYSTEM | | | | |
| None | | | | |
| DIGESTIVE SYSTEM | | | | |
| #Salivary gland | (47) | (48) | (49) | (47) |
| Adenoma, NOS | | 1 (2%) | | |
| #Liver | (50) | (50) | (50) | (49) |
| Bile duct adenoma | | 1 (2%) | 1 (2%) | 1 (2%) |
| Neoplastic nodule | 1 (2%) | 1 (2%) | | 1 (2%) |
| Hepatocellular carcinoma | | | | 1 (2%) |
| #Pancreas | (50) | (49) | (50) | (50) |
| Acinar cell adenoma | | | | 1 (2%) |
| #Stomach | (50) | (50) | (50) | (50) |
| Squamous cell carcinoma | 2 (4%) | | 1 (2%) | 1 (2%) |
| #Forestomach | (50) | (50) | (50) | (50) |
| Papillomatosis | 1 (2%) | | | |
| Squamous cell carcinoma | 3 (6%) | 3 (6%) | 2 (4%) | |
| #Duodenum | (48) | (49) | (50) | (48) |
| Adenocarcinoma, NOS | 1 (2%) | | | |
| #Jejunum | (48) | (49) | (50) | (48) |
| Carcinoma, NOS | | | 1 (2%) | |

**TABLE G1. SUMMARY OF THE INCIDENCE OF NEOPLASMS IN MALE OSBORNE-MENDEL RATS
IN THE TWO-YEAR GAVAGE STUDY OF TRICHLOROETHYLENE (Continued)**

| | CONTROL (UNTR) | CONTROL (VEH) | LOW DOSE | HIGH DOSE |
|-------------------------------|----------------|---------------|----------|-----------|
| URINARY SYSTEM | | | | |
| #Kidney | (50) | (50) | (50) | (50) |
| Tubular cell adenoma | | | 6 (12%) | 1 (2%) |
| Tubular cell adenocarcinoma | | | | 1 (2%) |
| Liposarcoma | | 1 (2%) | | |
| Nephroblastoma | | | | 1 (2%) |
| ENDOCRINE SYSTEM | | | | |
| #Pituitary | (46) | (42) | (45) | (38) |
| Carcinoma, NOS | 1 (2%) | | | |
| Adenoma, NOS | 16 (35%) | 11 (26%) | 7 (16%) | 10 (26%) |
| #Adrenal | (50) | (50) | (49) | (50) |
| Cortical adenoma | 11 (22%) | 9 (18%) | 11 (22%) | 11 (22%) |
| Cortical carcinoma | | 1 (2%) | | |
| Pheochromocytoma | 9 (18%) | 13 (26%) | 6 (12%) | 3 (6%) |
| Pheochromocytoma, malignant | 1 (2%) | 1 (2%) | | |
| #Thyroid | (48) | (49) | (48) | (48) |
| Follicular cell adenoma | | | | 1 (2%) |
| C-cell adenoma | 4 (8%) | 8 (16%) | 4 (8%) | 1 (2%) |
| C-cell carcinoma | | 1 (2%) | | |
| #Thyroid follicle | (48) | (49) | (48) | (48) |
| Cystadenoma, NOS | | | 1 (2%) | |
| #Parathyroid | (33) | (38) | (30) | (33) |
| Adenoma, NOS | 2 (6%) | | | |
| #Pancreatic islets | (50) | (49) | (50) | (50) |
| Islet cell adenoma | 1 (2%) | 3 (6%) | 2 (4%) | |
| Islet cell carcinoma | | 1 (2%) | | |
| REPRODUCTIVE SYSTEM | | | | |
| *Mammary gland | (50) | (50) | (50) | (50) |
| Cystadenoma, NOS | | | | 1 (2%) |
| Fibroma | 1 (2%) | | | |
| Fibroadenoma | 1 (2%) | | | |
| #Prostate | (49) | (49) | (50) | (50) |
| Adenoma, NOS | | 1 (2%) | | 1 (2%) |
| #Testis | (50) | (49) | (50) | (50) |
| Interstitial cell tumor | 1 (2%) | | | 1 (2%) |
| NERVOUS SYSTEM | | | | |
| *Nerve tract | (50) | (50) | (50) | (50) |
| Neurilemoma | | | 1 (2%) | |
| #Brain | (49) | (49) | (50) | (50) |
| Granular cell tumor, NOS | | 1 (2%) | | |
| #Cerebellum | (49) | (49) | (50) | (50) |
| Medulloblastoma | | | | 1 (2%) |
| SPECIAL SENSE ORGANS | | | | |
| None | | | | |
| MUSCULOSKELETAL SYSTEM | | | | |
| *Skeletal muscle | (50) | (50) | (50) | (50) |
| Sarcoma, NOS | | | 1 (2%) | |

**TABLE G1. SUMMARY OF THE INCIDENCE OF NEOPLASMS IN MALE OSBORNE-MENDEL RATS
IN THE TWO-YEAR GAVAGE STUDY OF TRICHOLORETHYLENE (Continued)**

| | CONTROL (UNTR) | CONTROL (VEH) | LOW DOSE | HIGH DOSE |
|--|----------------|---------------|----------|-----------|
| BODY CAVITIES | | | | |
| *Tunica vaginalis | (50) | (50) | (50) | (50) |
| Mesothelioma, NOS | 1 (2%) | | | |
| ALL OTHER SYSTEMS | | | | |
| *Multiple organs | (50) | (50) | (50) | (50) |
| Squamous cell carcinoma | | 1 (2%) | | |
| Sarcoma, NOS, metastatic | 1 (2%) | | | |
| Forearm | | | | |
| Mesenchymoma, malignant | | | 1 | |
| ANIMAL DISPOSITION SUMMARY | | | | |
| Animals initially in study | 50 | 50 | 50 | 50 |
| Natural death | 22 | 19 | 18 | 22 |
| Moribund sacrifice | 10 | 8 | 9 | 7 |
| Terminal sacrifice | 18 | 22 | 17 | 14 |
| Dosing accident | | | 6 | 2 |
| Accidentally killed, NOS | | 1 | | 5 |
| TUMOR SUMMARY | | | | |
| Total animals with primary tumors** | 39 | 37 | 35 | 29 |
| Total primary tumors | 66 | 66 | 52 | 44 |
| Total animals with benign tumors | 34 | 35 | 29 | 22 |
| Total benign tumors | 50 | 52 | 44 | 35 |
| Total animals with malignant tumors | 13 | 11 | 8 | 8 |
| Total malignant tumors | 14 | 12 | 8 | 8 |
| Total animals with secondary tumors## | 1 | | 1 | |
| Total secondary tumors | 1 | | 1 | |
| Total animals with tumors uncertain-- benign or malignant | 2 | 2 | | 1 |
| Total uncertain tumors | 2 | 2 | | 1 |

* Number of animals receiving complete necropsy examination; all gross lesions including masses examined microscopically.

** Primary tumors: all tumors except secondary tumors

Number of animals examined microscopically at this site

Secondary tumors: metastatic tumors or tumors invasive into an adjacent organ

TABLE G2. INDIVIDUAL ANIMAL TUMOR PATHOLOGY OF MALE OSBORNE-MENDEL RATS IN THE TWO-YEAR GAVAGE STUDY OF TRICHLOROETHYLENE: UNTREATED CONTROL

| ANIMAL NUMBER | WEEKS ON STUDY | | | | | | | | | | | | | | | | | | | | | | | | | | |
|--------------------------------|----------------|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|--|
| | 1 | 1 | 0 | 1 | 0 | 1 | 0 | 1 | 0 | 1 | 0 | 1 | 0 | 1 | 0 | 1 | 1 | 1 | 1 | 0 | 0 | 0 | 0 | 0 | 1 | 1 | |
| | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | |
| | 3 | 8 | 9 | 0 | 1 | 7 | 8 | 0 | 1 | 2 | 4 | 4 | 4 | 5 | 5 | 5 | 6 | 7 | 7 | 7 | 8 | 8 | 8 | 9 | 9 | 0 | |
| INTEGUMENTARY SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Skin | + | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Squamous cell carcinoma | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Subcutaneous tissue | + | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Sarcoma, NOS | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Fibroma | | | | | | | | | | | | | | X | | | | | | | | | | | | | |
| RESPIRATORY SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Lungs and bronchi | + | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Trachea | + | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Nasal cavity | + | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Carcinoma, NOS | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | |
| HEMATOPOIETIC SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Bone marrow | + | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Spleen | + | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Lymph nodes | + | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Thymus | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | |
| CIRCULATORY SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Heart | + | | | | | | | | | | | | | | | | | | | | | | | | | | |
| DIGESTIVE SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Salivary gland | + | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Liver | + | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Neoplastic nodule | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Bile duct | + | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Gallbladder & common bile duct | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | |
| Pancreas | + | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Malignant lymphoma, NOS | + | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Esophagus | + | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Stomach | + | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Papillomatosis | + | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Squamous cell carcinoma | | | | | | | | | | | | | | X | | | | | | | | | | | | | |
| Small intestine | | | | | | | | | | | | | | X | | | | | | | | | | | | | |
| Adenocarcinoma, NOS | | | | | | | | | | | | | | X | | | | | | | | | | | | | |
| Large intestine | + | | | | | | | | | | | | | | | | | | | | | | | | | | |
| URINARY SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Kidney | + | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Urinary bladder | + | | | | | | | | | | | | | | | | | | | | | | | | | | |
| ENDOCRINE SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Pituitary | + | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Carcinoma, NOS | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Adenoma, NOS | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Adrenal | + | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Cortical adenoma | X | X | | | | | | | | | | | | | | | | | | | | | | | | | |
| Pheochromocytoma | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Pheochromocytoma, malignant | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Thyroid | + | | | | | | | | | | | | | | | | | | | | | | | | | | |
| C-cell adenoma | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Parathyroid | + | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Adenoma, NOS | X | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Pancreatic islets | + | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Islet cell adenoma | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| REPRODUCTIVE SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Mammary gland | N | + | N | N | + | + | + | + | + | + | N | + | + | N | + | N | + | + | + | + | N | N | + | + | N | N | |
| Fibroma | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Fibroadenoma | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Testis | + | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Interstitial cell tumor | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Prostate | + | | | | | | | | | | | | | | | | | | | | | | | | | | |
| NERVOUS SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Brain | + | | | | | | | | | | | | | | | | | | | | | | | | | | |
| BODY CAVITIES | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Tunica vaginalis | + | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Mesothelioma, NOS | X | | | | | | | | | | | | | | | | | | | | | | | | | | |
| ALL OTHER SYSTEMS | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Multiple organs, NOS | N | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Sarcoma, NOS, metastatic | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Monocytic leukemia | | | | | | | | | | | | | | | | | | | | | | | | | | | |

+: Tissue examined microscopically
 -: Required tissue not examined microscopically
 X: Tumor incidence
 N: Necropsy, no autolysis, no microscopic examination
 S: Animal missexed

: No tissue information submitted
 C: Necropsy, no histology due to protocol
 A: Autolysis
 M: Animal missing
 B: No necropsy performed

**TABLE G2. INDIVIDUAL ANIMAL TUMOR PATHOLOGY OF MALE OSBORNE-MENDEL RATS:
UNTREATED CONTROL (Continued)**

| ANIMAL NUMBER | 1 0 7 | 1 1 1 | 1 1 6 | 1 1 8 | 1 2 1 | 1 2 4 | 1 3 3 | 1 3 5 | 1 4 9 | 1 4 2 | 1 5 2 | 1 5 6 | 1 5 7 | 1 6 3 | 1 6 5 | 1 6 9 | 1 7 6 | 1 7 7 | 1 8 8 | 1 8 8 | 1 9 5 | 1 9 7 | 1 9 9 | 1 9 9 | TOTAL TISSUES TUMORS |
|--------------------------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|----------------------------|
| WEEKS ON STUDY | 4 | 5 | 2 | 7 | 6 | 7 | 4 | 5 | 5 | 5 | 8 | 4 | 1 | 3 | 1 | 8 | 7 | 5 | 6 | 5 | 2 | 5 | 5 | 2 | 5 |
| INTEGUMENTARY SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | |
| Skin | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | *50 |
| Squamous cell carcinoma | | | | | | | | | | | | | | | | | | | X | | | | | | 1 |
| Subcutaneous tissue | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | *50 |
| Sarcoma, NOS | | | | | | | | | | | | | | | | | | | X | | | | | | 2 |
| Fibroma | | | | | | | | | | | | | | X | X | | | | | | | | | | 3 |
| RESPIRATORY SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | |
| Lungs and bronchi | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | 50 |
| Trachea | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | 50 |
| Nasal cavity | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | *50 |
| Carcinoma, NOS | | | | | | | | | | | | | X | | | | | | | | | | | | 1 |
| HEMATOPOIETIC SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | |
| Bone marrow | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | 50 |
| Spleen | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | 50 |
| Lymph nodes | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | 49 |
| Thymus | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | 3 |
| CIRCULATORY SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | |
| Heart | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | 50 |
| DIGESTIVE SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | |
| Salivary gland | + | + | - | - | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | 47 |
| Liver | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | 50 |
| Neoplastic nodule | | | | | | | | | | | | | | | | | | | X | | | | | | 1 |
| Bile duct | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | 50 |
| Gallbladder & common bile duct | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | *50 |
| Pancreas | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | 50 |
| Malignant lymphoma, NOS | | | | | | | | | | | | | | | | | | | X | | | | | | 1 |
| Esophagus | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | 50 |
| Stomach | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | 50 |
| Papillomatosis | | | | X | | | | | | | | | | | | | | | | | | | | | 1 |
| Squamous cell carcinoma | | | | | | X | | | | | | | | | | | | | | | | | | | 5 |
| Small intestine | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | 48 |
| Adenocarcinoma, NOS | | X | | | | | | | | | | | | | | | | | | | | | | | 1 |
| Large intestine | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | 48 |
| URINARY SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | |
| Kidney | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | 50 |
| Urinary bladder | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | 50 |
| ENDOCRINE SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | |
| Pituitary | + | + | - | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | 46 |
| Carcinoma, NOS | | | | | | | | | | | | | | | | | | | X | | | | | | 1 |
| Adenoma, NOS | | | | | | | X | X | X | | | | | X | X | | | X | X | X | | | | X | 16 |
| Adrenal | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | 50 |
| Cortical adenoma | X | | | X | | | X | | X | X | | | X | | | X | | | | | | | | 11 | |
| Pheochromocytoma | | X | | X | | | X | | X | | | X | | | | X | | | | | | X | | 9 | |
| Pheochromocytoma, malignant | | | | | | | X | | | | | | | | | | | | | | | | | 1 | |
| Thyroid | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | 48 |
| C-cell adenoma | | X | | | | | X | | | X | | | X | | | X | | | | | | | | 4 | |
| Parathyroid | + | + | - | - | + | + | - | + | + | - | + | + | - | + | + | - | + | + | - | + | + | + | + | + | 33 |
| Adenoma, NOS | | | | | | | | | | | | | | | | | | | | | | | | | 2 |
| Pancreatic islets | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | 50 |
| Islet cell adenoma | | | | | | | | | | | | | | | | | | | | | | | | | 1 |
| REPRODUCTIVE SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | |
| Mammary gland | + | + | N | + | N | + | N | + | + | N | + | + | N | + | + | N | + | + | + | + | + | + | + | N | *50 |
| Fibroma | | | | | | | | | | | | | | | | | | | | | | X | | | 1 |
| Fibroadenoma | | | | | | | | | | | | | | | | | | | | | | | | | 1 |
| Testis | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | 50 |
| Interstitial cell tumor | | | | | | | | | | | | | | | | | | | | | | | | | 1 |
| Prostate | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | 49 |
| NERVOUS SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | |
| Brain | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | 49 |
| BODY CAVITIES | | | | | | | | | | | | | | | | | | | | | | | | | |
| Tunica vaginalis | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | *50 |
| Mesothelioma, NOS | | | | | | | | | | | | | | | | | | | | | | | | | 1 |
| ALL OTHER SYSTEMS | | | | | | | | | | | | | | | | | | | | | | | | | |
| Multiple organs, NOS | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | *50 |
| Sarcoma, NOS, metastatic | | | | | | | | | | | | | | | | | | | | | | | | | 1 |
| Monocytic leukemia | | | | | | | | | | | | | | X | | | | | | | | | | | 1 |

* Animals necropsied

TABLE G2. INDIVIDUAL ANIMAL PATHOLOGY OF MALE OSBORNE-MENDEL RATS IN THE TWO-YEAR GAVAGE STUDY OF TRICHLOROETHYLENE: VEHICLE CONTROL

| ANIMAL NUMBER | 07 | 04 | 05 | 09 | 04 | 07 | 00 | 03 | 04 | 04 | 05 | 05 | 05 | 05 | 06 | 07 | 09 | 01 | 02 | 04 | 00 | 07 | 07 | 07 | 08 | 08 | 08 | | | | |
|--------------------------------|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|
| WEEKS ON STUDY | 10 | 10 | 08 | 00 | 11 | 11 | 00 | 03 | 06 | 00 | 11 | 11 | 11 | 11 | 00 | 11 | 00 | 11 | 00 | 11 | 00 | 11 | 00 | 07 | 06 | 08 | 00 | 11 | 11 | 11 | 00 |
| INTEGUMENTARY SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Subcutaneous tissue | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| Fibroma | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| RESPIRATORY SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Lungs and bronchi | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| Alveolar/bronchiolar adenoma | | | | | | X | | | | | | | | | | | | | | | | | | | | | | | | | |
| Alveolar/bronchiolar carcinoma | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Trachea | + | + | + | + | + | + | + | + | A | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | X | |
| HEMATOPOIETIC SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Bone marrow | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| Spleen | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| Lymph nodes | + | + | + | + | + | + | + | + | A | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| Thymus | - | - | - | - | - | - | - | - | A | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | |
| CIRCULATORY SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Heart | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| DIGESTIVE SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Salivary gland | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| Adenoma, NOS | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Liver | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| Bile duct adenoma | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Neoplastic nodule | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Bile duct | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| Gallbladder & common bile duct | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | |
| Pancreas | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| Esophagus | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| Stomach | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| Squamous cell carcinoma | X | | X | | X | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Small intestine | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| Malignant lymphoma, NOS | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Large intestine | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| URINARY SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Kidney | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| Liposarcoma | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Urinary bladder | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| ENDOCRINE SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Pituitary | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| Adenoma, NOS | X | | | X | | | | | A | | X | | | | | | | | | | | | | | | | | | | | |
| Adrenal | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| Cortical adenoma | X | X | X | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Cortical carcinoma | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Pheochromocytoma | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Pheochromocytoma, malignant | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Thyroid | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| C-cell adenoma | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| C-cell carcinoma | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Parathyroid | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| Pancreatic islets | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| Islet cell adenoma | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Islet cell carcinoma | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| REPRODUCTIVE SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Mammary gland | + | N | + | + | N | + | N | + | + | + | + | + | + | + | N | + | N | N | N | + | + | N | + | N | N | + | + | + | + | | |
| Testis | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| Prostate | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| Adenoma, NOS | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| NERVOUS SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Brain | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| Granular cell tumor, NOS | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| ALL OTHER SYSTEMS | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Multiple organs, NOS | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | |
| Squamous cell carcinoma | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Monocytic leukemia | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |

TABLE G2. INDIVIDUAL ANIMAL TUMOR PATHOLOGY OF MALE OSBORNE-MENDEL RATS: VEHICLE CONTROL (Continued)

| ANIMAL NUMBER | 0 | 0 | 0 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 2 | TOTAL TISSUES TUMORS |
|--------------------------------|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|----------------------|
| WEEKS ON STUDY | 9 | 9 | 9 | 0 | 0 | 1 | 1 | 1 | 2 | 2 | 2 | 2 | 3 | 3 | 4 | 4 | 5 | 5 | 6 | 6 | 6 | 7 | 8 | 9 | |
| INTEGUMENTARY SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | |
| Subcutaneous tissue | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| Fibroma | X | | | | | | | | X | | | | | | | | | | | | | | | | *50 4 |
| RESPIRATORY SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | |
| Lungs and bronchi | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | 50 |
| Alveolar/bronchiolar adenoma | | | | | | | | | | | | | | | | | | | | | | | | | 1 |
| Alveolar/bronchiolar carcinoma | | | | | | | | | | | | | | | | | | | | | | | | | 1 |
| Trachea | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | 49 |
| HEMATOPOIETIC SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | |
| Bone marrow | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | - | 49 |
| Spleen | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | 49 |
| Lymph nodes | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | 48 |
| Thymus | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | + | 3 |
| CIRCULATORY SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | |
| Heart | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | 50 |
| DIGESTIVE SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | |
| Salivary gland | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | - | 48 |
| Adenoma, NOS | | | | | | | | | | | | | | | | | | | | | | | | X | 1 |
| Liver | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | 50 |
| Bile duct adenoma | | | | | | | | | | | | | | | | | | | | | | | | | 1 |
| Neoplastic nodule | | | | X | | | | | | | | | | | | | | | | | | | | | 1 |
| Bile duct | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | 50 |
| Gallbladder & common bile duct | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | *50 |
| Pancreas | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | 49 |
| Esophagus | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | 50 |
| Stomach | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | 50 |
| Squamous cell carcinoma | | | | | | | | | | | | | | | | | | | | | | | | | 3 |
| Small intestine | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | - | 49 |
| Malignant lymphoma, NOS | | | | | | | | | | | | | | | | | | | | | | | X | | 1 |
| Large intestine | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | - | 47 |
| URINARY SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | |
| Kidney | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | 50 |
| Liposarcoma | | | | | | | | | | | | | | | | | | | | | | | | | 1 |
| Urinary bladder | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | - | 46 |
| ENDOCRINE SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | |
| Pituitary | + | + | + | + | + | - | - | + | + | - | + | + | + | + | + | + | + | + | + | + | + | + | + | - | 42 |
| Adenoma, NOS | | | X | | X | | | | | | X | | | X | | | X | | | | | X | X | | 11 |
| Adrenal | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | 50 |
| Cortical adenoma | | | | X | X | | | | | | X | X | | | | | | | | | | | | | 9 |
| Cortical carcinoma | | | | | | | | | | | | | | | | | | X | | | | | | | 1 |
| Pheochromocytoma | | | | | | | | | | | | | | | | | | | X | | | | | | 13 |
| Pheochromocytoma, malignant | X | X | X | | | | | | X | X | | | | | X | | | | | | | | | | 1 |
| Thyroid | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | 49 |
| C-cell adenoma | | | | X | | | | | | | | | X | | | | | X | X | | | | | | 8 |
| C-cell carcinoma | | | | | | | | | | | | | | | | | | | | | | | | | 1 |
| Parathyroid | + | + | - | + | + | + | - | + | + | - | + | + | + | + | + | + | + | + | + | + | + | + | + | - | 38 |
| Pancreatic islets | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | 49 |
| Islet cell adenoma | | | | | | | | | | | | | | | | | | | | | | | | | 3 |
| Islet cell carcinoma | | | | | | | | | | | | | | | | | | | | | | | | X | 1 |
| REPRODUCTIVE SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | |
| Mammary gland | + | + | + | N | + | + | N | + | + | N | N | + | N | + | N | + | + | + | + | N | N | N | N | + | *50 |
| Testis | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | - | 49 |
| Prostate | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | 49 |
| Adenoma, NOS | | | | | | | | | | | | | | | | | | | | | | | | | 1 |
| NERVOUS SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | |
| Brain | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | - | 49 |
| Granular cell tumor, NOS | | | | | | | | | | | | | | | | | | | | | | | | X | 1 |
| ALL OTHER SYSTEMS | | | | | | | | | | | | | | | | | | | | | | | | | |
| Multiple organs, NOS | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | *50 |
| Squamous cell carcinoma | | | | | | | | | | | | | | | | | | | | | | | | | 1 |
| Monocytic leukemia | | | | | | | | | | | | | | | | | | | | | | | | | 1 |

* Animals necropsied

TABLE G3. ANALYSIS OF PRIMARY TUMORS IN MALE OSBORNE-MENDEL RATS IN THE TWO-YEAR GAVAGE STUDY OF TRICHLOROETHYLENE

| | Untreated Control | Vehicle Control | 500 mg/kg | 1,000 mg/kg |
|---|-------------------|-----------------|------------|-------------|
| Subcutaneous Tissue: Fibroma | | | | |
| Overall Rates (a) | 3/50 (6%) | 4/50 (8%) | 3/50 (6%) | 0/50 (0%) |
| Adjusted Rates (b) | 9.8% | 13.6% | 14.0% | 0.0% |
| Terminal Rates (c) | 0/21 (0%) | 1/22 (5%) | 2/17 (12%) | 0/15 (0%) |
| Life Table Tests (d) | | P=0.112N | P=0.612N | P=0.129N |
| Incidental Tumor Tests (d) | | P=0.095N | P=0.547N | P=0.130N |
| Cochran-Armitage Trend Test (d) | | P=0.049N | | |
| Fisher Exact Test | | | P=0.500N | P=0.059N |
| Integumentary System: Fibroma | | | | |
| Overall Rates (a) | 3/50 (6%) | 4/50 (8%) | 4/50 (8%) | 2/50 (4%) |
| Adjusted Rates (b) | 9.8% | 13.6% | 19.8% | 11.6% |
| Terminal Rates (c) | 0/21 (0%) | 1/22 (5%) | 3/17 (18%) | 1/15 (7%) |
| Life Table Tests (d) | | P=0.468N | P=0.519 | P=0.528N |
| Incidental Tumor Tests (d) | | P=0.451N | P=0.582 | P=0.548N |
| Cochran-Armitage Trend Test (d) | | P=0.274N | | |
| Fisher Exact Test | | | P=0.643 | P=0.339N |
| Integumentary System: Fibroma or Sarcoma | | | | |
| Overall Rates (a) | 4/50 (8%) | 4/50 (8%) | 6/50 (12%) | 2/50 (4%) |
| Adjusted Rates (b) | 15.0% | 13.6% | 24.7% | 11.6% |
| Terminal Rates (c) | 2/21 (10%) | 1/22 (5%) | 3/17 (18%) | 1/15 (7%) |
| Life Table Tests (d) | | P=0.509N | P=0.255 | P=0.528N |
| Incidental Tumor Tests (d) | | P=0.489N | P=0.340 | P=0.548N |
| Cochran-Armitage Trend Test (d) | | P=0.290N | | |
| Fisher Exact Test | | | P=0.370 | P=0.339N |
| Stomach: Squamous Cell Carcinoma | | | | |
| Overall Rates (a) | (e) 5/50 (10%) | 3/50 (6%) | 3/50 (6%) | 1/50 (2%) |
| Adjusted Rates (b) | 18.4% | 11.5% | 7.3% | 2.1% |
| Terminal Rates (c) | 2/22 (9%) | 2/22 (9%) | 0/17 (0%) | 0/15 (0%) |
| Life Table Tests (d) | | P=0.342N | P=0.583 | P=0.421N |
| Incidental Tumor Tests (d) | | P=0.053N | P=0.431N | P=0.230N |
| Cochran-Armitage Trend Test (d) | | P=0.238N | | |
| Fisher Exact Test | | | P=0.661 | P=0.309N |
| Kidney: Tubular Cell Adenoma | | | | |
| Overall Rates (a) | 0/50 (0%) | 0/50 (0%) | 6/50 (12%) | 1/50 (2%) |
| Adjusted Rates (b) | 0.0% | 0.0% | 32.2% | 6.7% |
| Terminal Rates (c) | 0/21 (0%) | 0/22 (0%) | 5/17 (29%) | 1/15 (7%) |
| Life Table Tests (d) | | P=0.246 | P=0.007 | P=0.424 |
| Incidental Tumor Tests (d) | | P=0.243 | P=0.007 | P=0.424 |
| Cochran-Armitage Trend Test (d) | | P=0.406 | | |
| Fisher Exact Test | | | P=0.013 | P=0.500 |
| Kidney: Tubular Cell Adenoma or Adenocarcinoma | | | | |
| Overall Rates (a) | 0/50 (0%) | 0/50 (0%) | 6/50 (12%) | 2/50 (4%) |
| Adjusted Rates (b) | 0.0% | 0.0% | 32.2% | 10.9% |
| Terminal Rates (c) | 0/21 (0%) | 0/22 (0%) | 5/17 (29%) | 1/15 (7%) |
| Life Table Tests (d) | | P=0.125 | P=0.007 | P=0.158 |
| Incidental Tumor Tests (d) | | P=0.122 | P=0.007 | P=0.158 |
| Cochran-Armitage Trend Test (d) | | P=0.252 | | |
| Fisher Exact Test | | | P=0.013 | P=0.247 |
| Pituitary: Adenoma | | | | |
| Overall Rates (a) | (f) 16/46 (35%) | 11/42 (26%) | 7/45 (16%) | 10/38 (26%) |
| Adjusted Rates (b) | 55.4% | 42.0% | 35.0% | 50.6% |
| Terminal Rates (c) | 9/21 (43%) | 8/22 (36%) | 5/17 (29%) | 6/15 (40%) |
| Life Table Tests (d) | | P=0.277 | P=0.419N | P=0.294 |
| Incidental Tumor Tests (d) | | P=0.287 | P=0.334N | P=0.300 |
| Cochran-Armitage Trend Test (d) | | P=0.542N | | |
| Fisher Exact Test | | | P=0.169N | P=0.595 |

TABLE G3. ANALYSIS OF PRIMARY TUMORS IN MALE OSBORNE-MENDEL RATS IN THE TWO-YEAR GAVAGE STUDY OF TRICHLOROETHYLENE (Continued)

| | Untreated Control | Vehicle Control | 500 mg/kg | 1,000 mg/kg |
|---|-------------------|-----------------|-------------|-------------|
| Adrenal: Cortical Adenoma | | | | |
| Overall Rates (a) | 11/50 (22%) | 9/50 (18%) | 11/49 (22%) | 11/50 (22%) |
| Adjusted Rates (b) | 40.3% | 32.4% | 41.1% | 54.4% |
| Terminal Rates (c) | 7/21 (33%) | 5/22 (23%) | 3/17 (18%) | 7/15 (47%) |
| Life Table Tests (d) | | P=0.094 | P=0.215 | P=0.115 |
| Incidental Tumor Tests (d) | | P=0.109 | P=0.327 | P=0.160 |
| Cochran-Armitage Trend Test (d) | | P=0.356 | | |
| Fisher Exact Test | | | P=0.382 | P=0.402 |
| Adrenal: Cortical Adenoma or Carcinoma | | | | |
| Overall Rates (a) | 11/50 (22%) | 10/50 (20%) | 11/49 (22%) | 11/50 (22%) |
| Adjusted Rates (b) | 40.3% | 34.4% | 41.1% | 54.4% |
| Terminal Rates (c) | 7/21 (33%) | 5/22 (23%) | 3/17 (18%) | 7/15 (47%) |
| Life Table Tests (d) | | P=0.137 | P=0.286 | P=0.165 |
| Incidental Tumor Tests (d) | | P=0.161 | P=0.435 | P=0.221 |
| Cochran-Armitage Trend Test (d) | | P=0.452 | | |
| Fisher Exact Test | | | P=0.479 | P=0.500 |
| Adrenal: Pheochromocytoma | | | | |
| Overall Rates (a) | (g) 9/50 (18%) | (h) 13/50 (26%) | 6/49 (12%) | 3/50 (6%) |
| Adjusted Rates (b) | 34.8% | 42.9% | 31.6% | 18.7% |
| Terminal Rates (c) | 6/21 (29%) | 7/22 (32%) | 5/17 (29%) | 2/15 (13%) |
| Life Table Tests (d) | | P=0.028N | P=0.154N | P=0.052N |
| Incidental Tumor Tests (d) | | P=0.023N | P=0.128N | P=0.043N |
| Cochran-Armitage Trend Test (d) | | P=0.004N | | |
| Fisher Exact Test | | | P=0.068N | P=0.006N |
| Thyroid: C-Cell Adenoma | | | | |
| Overall Rates (a) | 4/48 (8%) | 8/49 (16%) | 4/48 (8%) | 1/48 (2%) |
| Adjusted Rates (b) | 19.0% | 33.9% | 21.4% | 6.7% |
| Terminal Rates (c) | 4/21 (19%) | 7/22 (32%) | 3/17 (18%) | 1/15 (7%) |
| Life Table Tests (d) | | P=0.038N | P=0.312N | P=0.055N |
| Incidental Tumor Tests (d) | | P=0.038N | P=0.299N | P=0.057N |
| Cochran-Armitage Trend Test (d) | | P=0.011N | | |
| Fisher Exact Test | | | P=0.188N | P=0.017N |
| Thyroid: C-Cell Adenoma or Carcinoma | | | | |
| Overall Rates (a) | 4/48 (8%) | 9/49 (18%) | 4/48 (8%) | 1/48 (2%) |
| Adjusted Rates (b) | 19.0% | 36.3% | 21.4% | 6.7% |
| Terminal Rates (c) | 4/21 (19%) | 7/22 (32%) | 3/17 (18%) | 1/15 (7%) |
| Life Table Tests (d) | | P=0.022N | P=0.224N | P=0.038N |
| Incidental Tumor Tests (d) | | P=0.022N | P=0.211N | P=0.039N |
| Cochran-Armitage Trend Test (d) | | P=0.005N | | |
| Fisher Exact Test | | | P=0.124N | P=0.008N |
| Pancreatic Islets: Islet Cell Adenoma | | | | |
| Overall Rates (a) | 1/50 (2%) | 3/49 (6%) | 2/50 (4%) | 0/50 (0%) |
| Adjusted Rates (b) | 4.8% | 12.0% | 11.8% | 0.0% |
| Terminal Rates (c) | 1/21 (5%) | 1/22 (5%) | 2/17 (12%) | 0/15 (0%) |
| Life Table Tests (d) | | P=0.152N | P=0.602N | P=0.203N |
| Incidental Tumor Tests (d) | | P=0.155N | P=0.590N | P=0.213N |
| Cochran-Armitage Trend Test (d) | | P=0.079N | | |
| Fisher Exact Test | | | P=0.490N | P=0.117N |
| Pancreatic Islets: Islet Cell Adenoma or Carcinoma | | | | |
| Overall Rates (a) | 1/50 (2%) | 4/49 (8%) | 2/50 (4%) | 0/50 (0%) |
| Adjusted Rates (b) | 4.8% | 16.1% | 11.8% | 0.0% |
| Terminal Rates (c) | 1/21 (5%) | 2/22 (9%) | 2/17 (12%) | 0/15 (0%) |
| Life Table Tests (d) | | P=0.082N | P=0.448N | P=0.127N |
| Incidental Tumor Tests (d) | | P=0.083N | P=0.435N | P=0.133N |
| Cochran-Armitage Trend Test (d) | | P=0.035N | | |
| Fisher Exact Test | | | P=0.329N | P=0.056N |

TABLE G3. ANALYSIS OF PRIMARY TUMORS IN MALE OSBORNE-MENDEL RATS IN THE TWO-YEAR GAVAGE STUDY OF TRICHLOROETHYLENE (Continued)

- (a) Number of tumor-bearing animals/number of animals examined at the site
- (b) Kaplan-Meier estimated tumor incidence at the end of the study after adjusting for intercurrent mortality
- (c) Observed tumor incidence at terminal kill
- (d) Beneath the control incidence are the P values associated with the trend test. Beneath the dosed group incidence are the P values corresponding to pairwise comparisons between that dosed group and the vehicle controls. The life table analysis regards tumors in animals dying prior to terminal kill as being (directly or indirectly) the cause of death. The incidental tumor test regards these lesions as nonfatal. The Cochran-Armitage and Fisher exact tests compare directly the overall incidence rates. A negative trend or lower incidence in a dosed group is indicated by (N).
- (e) One instance of papillomatosis was also observed.
- (f) One carcinoma, NOS, was also observed.
- (g) One malignant pheochromocytoma was also observed.
- (h) One animal also had a malignant pheochromocytoma.

TABLE G4. SUMMARY OF THE INCIDENCE OF NONNEOPLASTIC LESIONS IN MALE OSBORNE-MENDEL RATS IN THE TWO-YEAR GAVAGE STUDY OF TRICHLOROETHYLENE

| | CONTROL (UNTR) | CONTROL (VEH) | LOW DOSE | HIGH DOSE |
|--------------------------------------|----------------|---------------|----------|-----------|
| ANIMALS INITIALLY IN STUDY | 50 | 50 | 50 | 50 |
| ANIMALS NECROPSIED | 50 | 50 | 50 | 50 |
| ANIMALS EXAMINED HISTOPATHOLOGICALLY | 50 | 50 | 50 | 50 |
| INTEGUMENTARY SYSTEM | | | | |
| *Skin | (50) | (50) | (50) | (50) |
| Cyst, NOS | | 1 (2%) | | |
| *Subcutaneous tissue | (50) | (50) | (50) | (50) |
| Hemorrhage | | 1 (2%) | 1 (2%) | |
| Inflammation, NOS | | | | 1 (2%) |
| Inflammation, acute | | 1 (2%) | | |
| Abscess, NOS | | | 1 (2%) | |
| Inflammation, granulomatous | | | 1 (2%) | |
| RESPIRATORY SYSTEM | | | | |
| *Tracheal lumen | (50) | (50) | (50) | (50) |
| Hemorrhage | | | 1 (2%) | 1 (2%) |
| *Bronchial lumen | (50) | (50) | (50) | (50) |
| Hemorrhage | | | | 1 (2%) |
| *Larynx | (50) | (50) | (50) | (50) |
| Inflammation, acute necrotizing | | | 1 (2%) | |
| #Lung | (50) | (50) | (50) | (50) |
| Vegetable foreign body | | | | 2 (4%) |
| Emphysema, alveolar | | | | 1 (2%) |
| Atelectasis | | | 1 (2%) | |
| Congestion, NOS | 4 (8%) | 1 (2%) | 3 (6%) | 11 (22%) |
| Edema, NOS | 2 (4%) | 1 (2%) | 1 (2%) | 3 (6%) |
| Hemorrhage | 8 (16%) | 5 (10%) | 6 (12%) | 2 (4%) |
| Inflammation, NOS | | 1 (2%) | | |
| Inflammation, focal | | 2 (4%) | | 2 (4%) |
| Bronchopneumonia, acute | | 1 (2%) | | |
| Inflammation, acute | | | 1 (2%) | 3 (6%) |
| Inflammation, acute necrotizing | | | 1 (2%) | |
| Inflammation, chronic | 1 (2%) | 1 (2%) | 1 (2%) | |
| Inflammation, chronic focal | 2 (4%) | | 1 (2%) | |
| Inflammation, granulomatous | | | | 1 (2%) |
| Inflammation, granulomatous focal | | 1 (2%) | 5 (10%) | 6 (12%) |
| Foreign material, NOS | | | 1 (2%) | 2 (4%) |
| Hyperplasia, epithelial | | 1 (2%) | | |
| #Lung/alveoli | (50) | (50) | (50) | (50) |
| Hemorrhage | | | | 1 (2%) |
| HEMATOPOIETIC SYSTEM | | | | |
| #Bone marrow | (50) | (49) | (50) | (50) |
| Hyperplasia, NOS | 2 (4%) | | | |
| #Spleen | (50) | (49) | (50) | (50) |
| Hemorrhage | 1 (2%) | | 1 (2%) | |
| Hyperplasia, reticulum cell | | | | 1 (2%) |
| Hyperplasia, lymphoid | | 1 (2%) | | 1 (2%) |
| Hematopoiesis | 4 (8%) | 3 (6%) | 4 (8%) | 4 (8%) |
| #Lymph node | (49) | (48) | (49) | (44) |
| Hemorrhage | | | 3 (6%) | |
| Inflammation, NOS | | 1 (2%) | | |
| Inflammation, chronic | | | 1 (2%) | |
| Necrosis, focal | | | 1 (2%) | |
| Angiectasis | | | 1 (2%) | |

TABLE G4. SUMMARY OF THE INCIDENCE OF NONNEOPLASTIC LESIONS IN MALE OSBORNE-MENDEL RATS IN THE TWO-YEAR GAVAGE STUDY OF TRICHLOROETHYLENE (Continued)

| | CONTROL (UNTR) | CONTROL (VEH) | LOW DOSE | HIGH DOSE |
|---|----------------|---------------|----------|-----------|
| HEMATOPOIETIC SYSTEM (Continued) | | | | |
| #Pancreatic lymph node | (49) | (48) | (49) | (44) |
| Edema, NOS | 1 (2%) | | | |
| Hemorrhage | | | | 1 (2%) |
| Inflammation, NOS | | 1 (2%) | | |
| Inflammation, chronic | | 1 (2%) | | |
| Hyperplasia, NOS | | | 1 (2%) | |
| Angiectasis | 1 (2%) | 1 (2%) | | |
| Hyperplasia, lymphoid | | 1 (2%) | | |
| #Mesenteric lymph node | (49) | (48) | (49) | (44) |
| Hemorrhage | 1 (2%) | 2 (4%) | | 2 (5%) |
| #Liver | (50) | (50) | (50) | (49) |
| Hematopoiesis | 1 (2%) | | | |
| #Colon | (48) | (47) | (49) | (48) |
| Hyperplasia, lymphoid | 2 (4%) | | | |
| #Thymus | (3) | (3) | (8) | (13) |
| Hemorrhage | 1 (33%) | 2 (67%) | 2 (25%) | 2 (15%) |
| Inflammation, chronic | | | | 1 (8%) |
| Hyperplasia, lymphoid | | | | 1 (8%) |
| CIRCULATORY SYSTEM | | | | |
| #Lung | (50) | (50) | (50) | (50) |
| Perivasculitis | | | 1 (2%) | |
| #Heart | (50) | (50) | (50) | (50) |
| Thrombosis, NOS | | 2 (4%) | | |
| Inflammation, focal | | 3 (6%) | | |
| Inflammation, interstitial | | | 1 (2%) | 1 (2%) |
| Inflammation, active chronic | | 1 (2%) | | |
| Inflammation, chronic focal | 1 (2%) | 2 (4%) | | |
| Fibrosis | | 1 (2%) | | |
| Fibrosis, focal | | 1 (2%) | | 1 (2%) |
| Endocardiosis | 1 (2%) | | | |
| Calcification, focal | 1 (2%) | | | |
| #Auricular appendage | (50) | (50) | (50) | (50) |
| Thrombosis, NOS | 1 (2%) | | | 1 (2%) |
| #Endocardium | (50) | (50) | (50) | (50) |
| Inflammation with fibrosis | | 1 (2%) | | |
| *Coronary artery | (50) | (50) | (50) | (50) |
| Calcification, NOS | | 1 (2%) | | |
| *Pulmonary artery | (50) | (50) | (50) | (50) |
| Calcification, focal | | 1 (2%) | 2 (4%) | |
| #Pancreas | (50) | (49) | (50) | (50) |
| Periarteritis | 1 (2%) | | | |
| *Esophageal lumen | (50) | (50) | (50) | (50) |
| Thrombosis, NOS | | | 1 (2%) | |
| #Testis | (50) | (49) | (50) | (50) |
| Periarteritis | 1 (2%) | 1 (2%) | | |
| DIGESTIVE SYSTEM | | | | |
| #Salivary gland | (47) | (48) | (49) | (47) |
| Inflammation, chronic | 1 (2%) | | 1 (2%) | 1 (2%) |
| Inflammation, chronic focal | 1 (2%) | 1 (2%) | 1 (2%) | |
| Degeneration, NOS | | | | 2 (4%) |
| Metamorphosis, fatty | 1 (2%) | | | |
| Cytoplasmic vacuolization | | 1 (2%) | 1 (2%) | |
| Atrophy, focal | | 1 (2%) | | |
| Atrophy, diffuse | 1 (2%) | | | |

TABLE G4. SUMMARY OF THE INCIDENCE OF NONNEOPLASTIC LESIONS IN MALE OSBORNE-MENDEL RATS IN THE TWO-YEAR GAVAGE STUDY OF TRICHLOROETHYLENE (Continued)

| | CONTROL (UNTR) | CONTROL (VEH) | LOW DOSE | HIGH DOSE |
|-------------------------------------|----------------|---------------|----------|-----------|
| DIGESTIVE SYSTEM (Continued) | | | | |
| #Liver | (50) | (50) | (50) | (49) |
| Congestion, NOS | 2 (4%) | 1 (2%) | | 1 (2%) |
| Hemorrhage | | 1 (2%) | | 1 (2%) |
| Inflammation, focal | | | 2 (4%) | |
| Inflammation, chronic | | | 1 (2%) | |
| Fibrosis, multifocal | 1 (2%) | | | |
| Necrosis, focal | 1 (2%) | 2 (4%) | 5 (10%) | 2 (4%) |
| Necrosis, central | 2 (4%) | | | 1 (2%) |
| Metamorphosis, fatty | | 1 (2%) | 1 (2%) | |
| Lipoidosis | 1 (2%) | 1 (2%) | | |
| Cytoplasmic change, NOS | 12 (24%) | 13 (26%) | 16 (32%) | 13 (27%) |
| Basophilic cyto change | | | 1 (2%) | 1 (2%) |
| Focal cellular change | 1 (2%) | 1 (2%) | | 1 (2%) |
| Clear cell change | | | 1 (2%) | |
| Cytologic alteration, NOS | | | 1 (2%) | 1 (2%) |
| Angiectasis | | 1 (2%) | | |
| #Liver/centrilobular | (50) | (50) | (50) | (49) |
| Necrosis, NOS | | 1 (2%) | | |
| Metamorphosis, fatty | 1 (2%) | | | |
| #Bile duct | (50) | (50) | (50) | (49) |
| Dilatation, NOS | 1 (2%) | | 2 (4%) | |
| Inflammation, chronic | 1 (2%) | 2 (4%) | 1 (2%) | |
| Fibrosis, focal | | | 1 (2%) | |
| Hyperplasia, NOS | 1 (2%) | 6 (12%) | 3 (6%) | |
| Hyperplasia, cystic | | | 1 (2%) | |
| #Pancreas | (50) | (49) | (50) | (50) |
| Dilatation/ducts | | | | 1 (2%) |
| Inflammation, fibrinous | | 1 (2%) | | |
| Inflammation, chronic | | 1 (2%) | 1 (2%) | |
| Inflammation, granulomatous | 1 (2%) | 1 (2%) | | |
| #Pancreatic acinus | (50) | (49) | (50) | (50) |
| Atrophy, NOS | 1 (2%) | | 1 (2%) | |
| Atrophy, focal | 6 (12%) | 6 (12%) | 3 (6%) | |
| *Esophageal lumen | (50) | (50) | (50) | (50) |
| Hemorrhage | 1 (2%) | | 1 (2%) | |
| #Stomach | (50) | (50) | (50) | (50) |
| Mineralization | | | 1 (2%) | |
| Cyst, NOS | | | | 1 (2%) |
| Inflammation, NOS | | | 1 (2%) | |
| Inflammation, chronic | 1 (2%) | | | |
| Inflammation, chronic focal | | 1 (2%) | 1 (2%) | |
| #Gastric mucosa | (50) | (50) | (50) | (50) |
| Inflammation, NOS | | | 1 (2%) | |
| Erosion | 1 (2%) | | | |
| Hyperkeratosis | | | | 1 (2%) |
| #Forestomach | (50) | (50) | (50) | (50) |
| Edema, NOS | | 1 (2%) | | |
| Ulcer, NOS | 1 (2%) | | | |
| Inflammation, chronic | 1 (2%) | | | |
| Hyperplasia, epithelial | | 2 (4%) | | |
| Hyperkeratosis | 1 (2%) | 1 (2%) | | |
| #Colon | (48) | (47) | (49) | (48) |
| Parasitism | 8 (17%) | 11 (23%) | 11 (22%) | 10 (21%) |
| #Cecum | (48) | (47) | (49) | (48) |
| Parasitism | 1 (2%) | | | |
| *Rectum | (50) | (50) | (50) | (50) |
| Parasitism | | | | 1 (2%) |

TABLE G4. SUMMARY OF THE INCIDENCE OF NONNEOPLASTIC LESIONS IN MALE OSBORNE-MENDEL RATS IN THE TWO-YEAR GAVAGE STUDY OF TRICHLOROETHYLENE (Continued)

| | CONTROL (UNTR) | CONTROL (VEH) | LOW DOSE | HIGH DOSE |
|----------------------------------|----------------|---------------|----------|-----------|
| URINARY SYSTEM | | | | |
| #Kidney | (50) | (50) | (50) | (50) |
| Mineralization | 1 (2%) | 1 (2%) | 1 (2%) | |
| Cast, NOS | 2 (4%) | | 1 (2%) | |
| Hydronephrosis | | 1 (2%) | | 1 (2%) |
| Cyst, NOS | | | 1 (2%) | |
| Congestion, NOS | 1 (2%) | | | |
| Hemorrhage | 1 (2%) | | | |
| Inflammation, acute | 1 (2%) | | | 1 (2%) |
| Nephropathy | 50 (100%) | 50 (100%) | 44 (88%) | 47 (94%) |
| Nephropathy, toxic | | | 39 (78%) | 35 (70%) |
| Calcification, focal | | | | 1 (2%) |
| Cytomegaly | | | 48 (96%) | 49 (98%) |
| Hyperplasia, tubular cell | | | 5 (10%) | 3 (6%) |
| Hyperplasia, cystic | 1 (2%) | | | |
| #Kidney/medulla | (50) | (50) | (50) | (50) |
| Hyperplasia, epithelial | 1 (2%) | 2 (4%) | | |
| #Kidney/tubule | (50) | (50) | (50) | (50) |
| Dilatation, NOS | | | 1 (2%) | |
| #Kidney/pelvis | (50) | (50) | (50) | (50) |
| Hemorrhage | | | 1 (2%) | |
| Hyperplasia, epithelial | 1 (2%) | | 2 (4%) | |
| #Urinary bladder | (50) | (46) | (50) | (49) |
| Calculus, gross observation only | | | | 1 (2%) |
| Cast, NOS | | | | 1 (2%) |
| Hemorrhage | | 2 (4%) | | |
| Inflammation, NOS | | | | 1 (2%) |
| Inflammation, acute/chronic | | 1 (2%) | | |
| Hyperplasia, epithelial | | | 1 (2%) | |
| ENDOCRINE SYSTEM | | | | |
| #Pituitary | (46) | (42) | (45) | (38) |
| Cyst, NOS | 2 (4%) | 2 (5%) | 1 (2%) | |
| Abscess, NOS | 1 (2%) | | | |
| Hyperplasia, focal | 3 (7%) | 2 (5%) | 1 (2%) | |
| #Adrenal | (50) | (50) | (49) | (50) |
| Cyst, NOS | 2 (4%) | | | |
| Hemorrhage | | 1 (2%) | | 1 (2%) |
| Inflammation, chronic focal | | | 1 (2%) | |
| Necrosis, NOS | | 1 (2%) | | |
| Necrosis, focal | | | 1 (2%) | |
| Metamorphosis, fatty | 1 (2%) | | | |
| Lipoidosis | 6 (12%) | 12 (24%) | 8 (16%) | 5 (10%) |
| Cytoplasmic vacuolization | 1 (2%) | | | |
| Focal cellular change | | 1 (2%) | | |
| Hyperplasia, NOS | | 1 (2%) | | |
| Angiectasis | 1 (2%) | 2 (4%) | 4 (8%) | 2 (4%) |
| #Adrenal cortex | (50) | (50) | (49) | (50) |
| Hemorrhage | | | | 1 (2%) |
| Metamorphosis, fatty | 1 (2%) | | | |
| Lipoidosis | 1 (2%) | | 3 (6%) | 1 (2%) |
| Focal cellular change | 1 (2%) | | | 1 (2%) |
| Cytologic alteration, NOS | | | 1 (2%) | |
| Hyperplasia, NOS | | 1 (2%) | | |
| Hyperplasia, focal | | | 1 (2%) | |
| #Adrenal medulla | (50) | (50) | (49) | (50) |
| Hyperplasia, NOS | 2 (4%) | 1 (2%) | | 1 (2%) |
| Hyperplasia, focal | 4 (8%) | 3 (6%) | 3 (6%) | 4 (8%) |

TABLE G4. SUMMARY OF THE INCIDENCE OF NONNEOPLASTIC LESIONS IN MALE OSBORNE-MENDEL RATS IN THE TWO-YEAR STUDY OF TRICHOLOETHYLENE (Continued)

| | CONTROL (UNTR) | CONTROL (VEH) | LOW DOSE | HIGH DOSE |
|-------------------------------------|----------------|---------------|----------|-----------|
| ENDOCRINE SYSTEM (Continued) | | | | |
| #Thyroid | (48) | (49) | (48) | (48) |
| Cyst, NOS | | | 2 (4%) | 4 (8%) |
| Follicular cyst, NOS | | | | 1 (2%) |
| Hyperplasia, C-cell | 1 (2%) | 5 (10%) | 5 (10%) | 3 (6%) |
| Hyperplasia, follicular cell | 2 (4%) | | | |
| #Parathyroid | (33) | (38) | (30) | (33) |
| Hyperplasia, NOS | | 1 (3%) | 1 (3%) | |
| #Pancreatic islets | (50) | (49) | (50) | (50) |
| Hyperplasia, focal | 1 (2%) | | | |
| REPRODUCTIVE SYSTEM | | | | |
| *Preputial gland | (50) | (50) | (50) | (50) |
| Abscess, NOS | 1 (2%) | | | |
| #Prostate | (49) | (49) | (50) | (50) |
| Cyst, NOS | | 1 (2%) | | |
| Hemorrhage | | | | 2 (4%) |
| Inflammation, focal | | | | 1 (2%) |
| Inflammation, acute | 1 (2%) | | | |
| Inflammation, acute focal | 2 (4%) | 1 (2%) | | |
| Inflammation, acute suppurative | | | | 1 (2%) |
| Inflammation, acute necrotizing | | 1 (2%) | | |
| Inflammation, acute/chronic | 1 (2%) | 1 (2%) | | |
| Inflammation, chronic | | 1 (2%) | | 1 (2%) |
| Inflammation, chronic focal | | 1 (2%) | 1 (2%) | |
| Inflammation, granulomatous | | 1 (2%) | | |
| Hyperplasia, NOS | | 1 (2%) | | |
| *Seminal vesicle | (50) | (50) | (50) | (50) |
| Inflammation, acute focal | | | | 1 (2%) |
| *Coagulating gland | (50) | (50) | (50) | (50) |
| Hyperplasia, focal | | | | 1 (2%) |
| #Testis | (50) | (49) | (50) | (50) |
| Atrophy, NOS | 1 (2%) | 7 (14%) | 1 (2%) | 2 (4%) |
| Atrophy, focal | | | 1 (2%) | |
| Atrophy, diffuse | | 1 (2%) | | |
| Hyperplasia, interstitial cell | 1 (2%) | | | |
| NERVOUS SYSTEM | | | | |
| #Brain | (49) | (49) | (50) | (50) |
| Cyst, NOS | | | | 1 (2%) |
| Congestion, NOS | | 1 (2%) | | |
| Hemorrhage | 1 (2%) | 2 (4%) | | 2 (4%) |
| *Spinal cord | (50) | (50) | (50) | (50) |
| Hemorrhage | | | | 1 (2%) |
| Calcification, focal | | | | 1 (2%) |
| *Pineal body | (50) | (50) | (50) | (50) |
| Degeneration, NOS | | | 1 (2%) | |
| SPECIAL SENSE ORGANS | | | | |
| *Eye/lacrimal gland | (50) | (50) | (50) | (50) |
| Inflammation, chronic | | 1 (2%) | 1 (2%) | |
| Inflammation, chronic focal | | | 1 (2%) | |
| Atrophy, NOS | | | 1 (2%) | |
| MUSCULOSKELETAL SYSTEM | | | | |
| *Pharyngeal muscle | (50) | (50) | (50) | (50) |
| Inflammation, acute necrotizing | | | 1 (2%) | |

TABLE G4. SUMMARY OF THE INCIDENCE OF NONNEOPLASTIC LESIONS IN MALE OSBORNE-MENDEL RATS IN THE TWO-YEAR GAVAGE STUDY OF TRICHLOROETHYLENE (Continued)

| | CONTROL (UNTR) | CONTROL (VEH) | LOW DOSE | HIGH DOSE |
|-----------------------------------|----------------|---------------|----------|-----------|
| BODY CAVITIES | | | | |
| *Thorax | (50) | (50) | (50) | (50) |
| Abscess, chronic | | | | 1 (2%) |
| *Mediastinum | (50) | (50) | (50) | (50) |
| Hemorrhage | | | | 1 (2%) |
| Abscess, NOS | | 1 (2%) | | |
| Inflammation, active chronic | | 1 (2%) | | |
| *Abdominal cavity | (50) | (50) | (50) | (50) |
| Necrosis, fat | 1 (2%) | | | |
| *Epicardium | (50) | (50) | (50) | (50) |
| Inflammation, chronic | | | | 1 (2%) |
| ALL OTHER SYSTEMS | | | | |
| *Multiple organs | (50) | (50) | (50) | (50) |
| Hemorrhage | | | | 1 (2%) |
| SPECIAL MORPHOLOGY SUMMARY | | | | |
| None | | | | |

* Number of animals receiving complete necropsy examination; all gross lesions including masses examined microscopically.
 # Number of animals examined microscopically at this site

APPENDIX H

SUMMARY OF LESIONS IN FEMALE OSBORNE-MENDEL RATS IN THE TWO-YEAR GAVAGE STUDY OF TRICHLOROETHYLENE

| | PAGE | |
|----------|---|-----|
| TABLE H1 | SUMMARY OF THE INCIDENCE OF NEOPLASMS IN FEMALE OSBORNE-MENDEL RATS IN THE TWO-YEAR GAVAGE STUDY OF TRICHLOROETHYLENE | 231 |
| TABLE H2 | INDIVIDUAL ANIMAL TUMOR PATHOLOGY OF FEMALE OSBORNE-MENDEL RATS IN THE TWO-YEAR GAVAGE STUDY OF TRICHLOROETHYLENE | 234 |
| TABLE H3 | ANALYSIS OF PRIMARY TUMORS IN FEMALE OSBORNE-MENDEL RATS IN THE TWO-YEAR GAVAGE STUDY OF TRICHLOROETHYLENE | 242 |
| TABLE H4 | SUMMARY OF THE INCIDENCE OF NONNEOPLASTIC LESIONS IN FEMALE OSBORNE-MENDEL RATS IN THE TWO-YEAR GAVAGE STUDY OF TRICHLOROETHYLENE | 245 |

TABLE H1. SUMMARY OF THE INCIDENCE OF NEOPLASMS IN FEMALE OSBORNE-MENDEL RATS
IN THE TWO-YEAR GAVAGE STUDY OF TRICHLOROETHYLENE

| | CONTROL (UNTR) | CONTROL (VEH) | LOW DOSE | HIGH DOSE |
|---|----------------|---------------|----------------|----------------|
| ANIMALS INITIALLY IN STUDY | 50 | 50 | 50 | 50 |
| ANIMALS NECROPSIED | 50 | 50 | 50 | 50 |
| ANIMALS EXAMINED HISTOPATHOLOGICALLY | 50 | 50 | 50 | 50 |
| INTEGUMENTARY SYSTEM | | | | |
| *Subcutaneous tissue Fibroma | (50) | (50) | (50) 2 (4%) | (50) 1 (2%) |
| RESPIRATORY SYSTEM | | | | |
| #Lung | (50) | (50) | (50) | (50) |
| Carcinoma, NOS, metastatic | | | 1 (2%) | |
| Squamous cell carcinoma, metastatic | | | 1 (2%) | |
| Alveolar/bronchiolar adenoma | | | | 1 (2%) |
| Alveolar/bronchiolar carcinoma | | | 1 (2%) | |
| Osteosarcoma, metastatic | | | | 1 (2%) |
| HEMATOPOIETIC SYSTEM | | | | |
| *Multiple organs | (50) | (50) | (50) | (50) |
| Malignant lymphoma, NOS | 1 (2%) | 1 (2%) | | 2 (4%) |
| Malignant lymphoma, lymphocytic type | | | 1 (2%) | |
| Monocytic leukemia | 2 (4%) | | 2 (4%) | |
| *Mediastinum | (50) | (50) | (50) | (50) |
| Malignant lymphoma, undiffer type | | 1 (2%) | | |
| #Liver | (50) | (50) | (50) | (49) |
| Leukemia, NOS | | | 1 (2%) | |
| CIRCULATORY SYSTEM | | | | |
| #Liver | (50) | (50) | (50) | (49) |
| Angioma | | | 1 (2%) | |
| DIGESTIVE SYSTEM | | | | |
| #Liver | (50) | (50) | (50) | (49) |
| Carcinoma, NOS, metastatic | | | 1 (2%) | |
| Bile duct carcinoma | | | 1 (2%) | |
| Neoplastic nodule | 1 (2%) | | | 2 (4%) |
| #Pancreas | (50) | (50) | (48) | (49) |
| Squamous cell carcinoma, metastatic | | | 1 (2%) | |
| Adenocarcinoma, NOS, metastatic | | | 1 (2%) | |
| #Stomach | (50) | (48) | (50) | (49) |
| Squamous cell carcinoma | | | | 1 (2%) |
| #Forestomach | (50) | (48) | (50) | (49) |
| Papillomatosis | 1 (2%) | | | |
| #Duodenum | (50) | (48) | (49) | (49) |
| Adenocarcinoma, NOS | | 1 (2%) | | |
| URINARY SYSTEM | | | | |
| #Kidney | (50) | (50) | (50) | (49) |
| Tubular cell adenoma | 1 (2%) | | | 1 (2%) |
| Nephroblastoma | | | 1 (2%) | |
| #Urinary bladder | (48) | (48) | (49) | (50) |
| Adenocarcinoma, NOS, metastatic | | | | 1 (2%) |

TABLE H1. SUMMARY OF THE INCIDENCE OF NEOPLASMS IN FEMALE OSBORNE-MENDEL RATS
IN THE TWO-YEAR GAVAGE STUDY OF TRICHLOROETHYLENE (Continued)

| | CONTROL (UNTR) | CONTROL (VEH) | LOW DOSE | HIGH DOSE |
|-----------------------------|----------------|---------------|----------|-----------|
| ENDOCRINE SYSTEM | | | | |
| #Pituitary | (47) | (46) | (47) | (46) |
| Carcinoma, NOS | | 1 (2%) | | |
| Adenoma, NOS | 13 (28%) | 13 (28%) | 9 (19%) | 8 (17%) |
| Papillary adenoma | | 1 (2%) | | |
| Chromophobe adenoma | 1 (2%) | | | |
| #Adrenal | (49) | (50) | (50) | (49) |
| Cortical adenoma | 13 (27%) | 16 (32%) | 13 (26%) | 19 (39%) |
| Pheochromocytoma | 3 (6%) | 8 (16%) | 4 (8%) | 1 (2%) |
| Pheochromocytoma, malignant | | 1 (2%) | 1 (2%) | |
| #Adrenal medulla | (49) | (50) | (50) | (49) |
| Pheochromocytoma | 1 (2%) | | | |
| Pheochromocytoma, malignant | | | 1 (2%) | |
| #Thyroid | (49) | (49) | (49) | (49) |
| Follicular cell adenoma | 1 (2%) | 1 (2%) | | |
| C-cell adenoma | 5 (10%) | 15 (31%) | 5 (10%) | 1 (2%) |
| #Parathyroid | (37) | (32) | (32) | (32) |
| Adenoma, NOS | | | | 1 (3%) |
| #Pancreatic islets | (50) | (50) | (48) | (49) |
| Islet cell adenoma | | 3 (6%) | 1 (2%) | |
| REPRODUCTIVE SYSTEM | | | | |
| *Mammary gland | (50) | (50) | (50) | (50) |
| Adenoma, NOS | | 1 (2%) | 1 (2%) | 1 (2%) |
| Adenocarcinoma, NOS | | 1 (2%) | 1 (2%) | 1 (2%) |
| Fibroma | | 1 (2%) | 1 (2%) | 1 (2%) |
| Mixed tumor, benign | | | 1 (2%) | |
| Fibroadenoma | 17 (34%) | 14 (28%) | 7 (14%) | 11 (22%) |
| *Vagina | (50) | (50) | (50) | (50) |
| Squamous cell carcinoma | | | | 1 (2%) |
| #Uterus | (50) | (49) | (49) | (50) |
| Carcinoma, NOS | 1 (2%) | 1 (2%) | 2 (4%) | 1 (2%) |
| Carcinoma, NOS, metastatic | | | 1 (2%) | |
| Papillomatosis | | | 1 (2%) | |
| Squamous cell papilloma | | 1 (2%) | | |
| Squamous cell carcinoma | | | 1 (2%) | |
| Adenoma, NOS | 1 (2%) | | | |
| Adenocarcinoma, NOS | | 1 (2%) | 5 (10%) | 2 (4%) |
| Multiple polyposis | | | 1 (2%) | |
| Sarcoma, NOS | 1 (2%) | | | |
| Endometrial stromal polyp | 4 (8%) | 6 (12%) | 1 (2%) | 6 (12%) |
| Carcinosarcoma | | | | 1 (2%) |
| #Uterus/endometrium | (50) | (49) | (49) | (50) |
| Carcinoma, NOS | | | | 1 (2%) |
| Adenocarcinoma, NOS | | 1 (2%) | | |
| #Ovary | (49) | (48) | (49) | (49) |
| Carcinoma, NOS | | | 1 (2%) | |
| Papillary adenoma | | | 1 (2%) | |
| Granulosa cell tumor | | | 1 (2%) | |
| Granulosa cell carcinoma | | | | 1 (2%) |
| NERVOUS SYSTEM | | | | |
| *Nerve tract | (50) | (50) | (50) | (50) |
| Neurilemoma | 1 (2%) | | | |
| SPECIAL SENSE ORGANS | | | | |
| None | | | | |

**TABLE H1. SUMMARY OF THE INCIDENCE OF NEOPLASMS IN FEMALE OSBORNE-MENDEL RATS
IN THE TWO-YEAR GAVAGE STUDY OF TRICHLOROETHYLENE (Continued)**

| | CONTROL (UNTR) | CONTROL (VEH) | LOW DOSE | HIGH DOSE |
|--|----------------|---------------|----------|-----------|
| MUSCULOSKELETAL SYSTEM | | | | |
| *Maxilla | (50) | (50) | (50) | (50) |
| Osteosarcoma | | | | 1 (2%) |
| *Skeletal muscle | (50) | (50) | (50) | (50) |
| Rhabdomyosarcoma | | | | 1 (2%) |
| BODY CAVITIES | | | | |
| *Abdominal cavity | (50) | (50) | (50) | (50) |
| Bile duct carcinoma, metastatic | | | 1 (2%) | |
| ALL OTHER SYSTEMS | | | | |
| *Multiple organs | (50) | (50) | (50) | (50) |
| Carcinoma, NOS | | | 2 (4%) | |
| Undifferentiated carcinoma | | | 1 (2%) | |
| Carcinosarcoma, metastatic | | | | 1 (2%) |
| Mesothelioma, malignant | 1 (2%) | | | |
| Lower leg | | | | |
| Sarcoma, NOS | | | | 1 |
| Site unknown | | | | |
| Carcinoma, NOS | | 1 | | |
| ANIMAL DISPOSITION SUMMARY | | | | |
| Animals initially in study | 50 | 50 | 50 | 50 |
| Natural death | 14 | 7 | 16 | 16 |
| Moribund sacrifice | 17 | 17 | 18 | 21 |
| Terminal sacrifice | 19 | 18 | 10 | 7 |
| Dosing accident | | 6 | 4 | 5 |
| Accidentally killed, NOS | | 2 | 2 | 1 |
| TUMOR SUMMARY | | | | |
| Total animals with primary tumors** | 44 | 40 | 36 | 37 |
| Total primary tumors | 69 | 90 | 72 | 68 |
| Total animals with benign tumors | 40 | 39 | 29 | 32 |
| Total benign tumors | 62 | 80 | 49 | 52 |
| Total animals with malignant tumors | 6 | 8 | 19 | 12 |
| Total malignant tumors | 6 | 10 | 22 | 14 |
| Total animals with secondary tumors## | | | 5 | 3 |
| Total secondary tumors | | | 7 | 3 |
| Total animals with tumors uncertain-- benign or malignant | 1 | | 1 | 2 |
| Total uncertain tumors | 1 | | 1 | 2 |

* Number of animals receiving complete necropsy examination; all gross lesions including masses examined microscopically.

** Primary tumors: all tumors except secondary tumors

Number of animals examined microscopically at this site

Secondary tumors: metastatic tumors or tumors invasive into an adjacent organ

TABLE H2. INDIVIDUAL ANIMAL TUMOR PATHOLOGY OF FEMALE OSBORNE-MENDEL RATS: VEHICLE CONTROL (Continued)

| ANIMAL NUMBER | 3 | 3 | 3 | 3 | 3 | 3 | 3 | 3 | 3 | 3 | 3 | 3 | 3 | 3 | 3 | 3 | 3 | 3 | 3 | 3 | 3 | 3 | 3 | 4 | |
|-----------------------------------|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---------------------|
| | 1 | 2 | 2 | 2 | 3 | 3 | 3 | 3 | 3 | 5 | 5 | 6 | 6 | 7 | 7 | 7 | 7 | 8 | 8 | 9 | 9 | 9 | 9 | 9 | 0 |
| WEEKS ON STUDY | 0 | 3 | 8 | 0 | 0 | 0 | 0 | 8 | 7 | 1 | 9 | 0 | 0 | 0 | 0 | 5 | 6 | 0 | 8 | 0 | 0 | 0 | 9 | 0 | 8 |
| | 0 | 3 | 0 | 6 | 0 | 5 | 7 | 8 | 9 | 0 | 5 | 4 | 1 | 4 | 2 | 1 | 5 | 3 | 5 | 5 | 0 | 4 | 5 | 4 | |
| RESPIRATORY SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | TOTAL ISSUES |
| Lungs and bronchi | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | 50 |
| Trachea | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | 50 |
| HEMATOPOIETIC SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | |
| Bone marrow | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | 50 |
| Spleen | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | 49 |
| Lymph nodes | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | 46 |
| Thymus | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | 4 |
| CIRCULATORY SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | |
| Heart | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | 49 |
| DIGESTIVE SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | |
| Salivary gland | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | 50 |
| Liver | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | 50 |
| Bile duct | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | 50 |
| Gallbladder & common bile duct | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | *50 |
| Pancreas | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | 50 |
| Esophagus | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | 48 |
| Stomach | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | 48 |
| Small intestine | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | 48 |
| Adenocarcinoma, NOS | + | - | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | 1 |
| Large intestine | + | - | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | 48 |
| URINARY SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | |
| Kidney | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | 50 |
| Urinary bladder | + | - | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | 48 |
| ENDOCRINE SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | |
| Pituitary | + | + | + | - | + | + | + | + | + | + | + | + | + | + | + | - | - | + | + | + | + | + | + | + | 46 |
| Carcinoma, NOS | | | | | | | | | | | | | | | | | | | | | | | | | 1 |
| Adenoma, NOS | | | | | | X | | | | | | | X | | | | X | | X | | | | | X | 13 |
| Papillary adenoma | | | | | | | | | | | | | | | | | | | | | | | | | 1 |
| Adrenal | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | 50 |
| Cortical adenoma | | | | | | X | | X | | | | | | | | | | | | | | | | | 16 |
| Pheochromocytoma | X | | | | | | X | | | | | | | | | X | | | | | | X | X | | 8 |
| Pheochromocytoma, malignant | | | | | | | | | | | | | | | | | | | | | | | | | 1 |
| Thyroid | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | 49 |
| Follicular cell adenoma | | | | | | | | | | | | | | | | | | | | | | | | | 1 |
| C-cell adenoma | X | | | | | | | | | | | | | | | X | | | X | X | | | | | 15 |
| Parathyroid | + | - | + | - | + | + | + | + | - | + | + | + | + | + | + | - | - | + | - | - | + | - | + | + | 32 |
| Pancreatic islets | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | 50 |
| Islet cell adenoma | | | | | | | | | | | | | | X | | | | | | | | | | | 3 |
| REPRODUCTIVE SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | |
| Mammary gland | + | N | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | *50 |
| Adenoma, NOS | | | | | | | | | | | | | | | | | | | | | | | X | | 1 |
| Adenocarcinoma, NOS | | | | | | | | | | | | | | | | | | | | | | | | | 1 |
| Fibroma | | | | | | | | | | | | | | | | | | | | | | | | | 1 |
| Fibroadenoma | | | | | | | X | X | | | | | | X | | X | X | X | | | X | | X | | 14 |
| Uterus | + | - | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | 49 |
| Carcinoma, NOS | | | | | | | X | | | | | | | | | | | | | | | | | | 1 |
| Squamous cell papilloma | | | | | | | | | | | | | | | | | | | | | | | | | 1 |
| Adenocarcinoma, NOS | | | | | | | | | | | | | | | | | | | | | | | | X | 2 |
| Endometrial stromal polyp | | | | | | | | | | | | | | | | X | X | | | | | | | | 8 |
| Ovary | + | - | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | 48 |
| NERVOUS SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | |
| Brain | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | 50 |
| BODY CAVITIES | | | | | | | | | | | | | | | | | | | | | | | | | |
| Mediastinum | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | *50 |
| Malignant lymphoma, undiffer type | | | | | | | | | | | | | X | | | | | | | | | | | | 1 |
| ALL OTHER SYSTEMS | | | | | | | | | | | | | | | | | | | | | | | | | |
| Multiple organs, NOS | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | *50 |
| Malignant lymphoma, NOS | | | | | | | | | | | | | | | | | | | | | | | | | 1 |
| Site unknown | | | | | | | | | | | | | | | | | | | | | | | | | |
| Carcinoma, NOS | | | | | | | | | | | | | | | | | | | | | | | | | 1 |

* Animals necropsied

TABLE H2. INDIVIDUAL ANIMAL TUMOR PATHOLOGY OF FEMALE OSBORNE-MENDEL RATS: HIGH DOSE (Continued)

| ANIMAL NUMBER | 3 0 3 | 3 0 5 | 3 0 7 | 3 2 5 | 3 2 6 | 3 2 7 | 3 3 0 | 3 4 0 | 3 4 1 | 3 4 2 | 3 4 3 | 3 4 6 | 3 6 1 | 3 6 3 | 3 6 6 | 3 7 0 | 3 7 1 | 3 7 3 | 3 7 7 | 3 8 0 | 3 8 6 | 3 8 9 | 3 9 4 | 3 9 5 | 3 9 8 | 3 9 9 | TOTAL TISSUES TUMORS |
|---------------------------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|----------------------------|
| WEEKS ON STUDY | 0 0 | 1 3 | 0 8 | 0 3 | 1 4 | 0 9 | 0 9 | 0 0 | 0 2 | 0 0 | 0 7 | 0 3 | 0 4 | 0 0 | 0 5 | 0 9 | 1 4 | 1 4 | 0 6 | 1 3 | 0 5 | 1 1 | 0 5 | 1 5 | 0 8 | 0 5 | |
| INTEGUMENTARY SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Subcutaneous tissue | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Fibroma | + | + | + | + | + | + | | X | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | *50 1 |
| RESPIRATORY SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Lungs and bronchi | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Alveolar/bronchiolar adenoma | + | + | + | + | + | + | | X | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | 50 1 |
| Osteosarcoma, metastatic | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Trachea | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | 47 |
| HEMATOPOIETIC SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Bone marrow | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Spleen | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Lymph nodes | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Thymus | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | 49 49 46 3 |
| CIRCULATORY SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Heart | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | 49 |
| DIGESTIVE SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Salivary gland | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Liver | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Neoplastic nodule | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | 47 49 2 |
| Bile duct | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Gallbladder & common bile duct | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | 49 *50 |
| Pancreas | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | 49 |
| Esophagus | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | 50 |
| Stomach | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | 49 |
| Squamous cell carcinoma | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Small intestine | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | 1 49 |
| Large intestine | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | 50 |
| URINARY SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Kidney | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Tubular cell adenoma | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | 49 1 |
| Urinary bladder | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Adenocarcinoma, NOS, metastatic | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | 50 1 |
| ENDOCRINE SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Pituitary | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Adenoma, NOS | + | + | + | + | + | - | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | 46 8 |
| Adrenal | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Cortical adenoma | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | 49 19 |
| Pheochromocytoma | X | X | | X | X | | X | X | | X | X | | | | | | | | X | | X | X | | | | 1 | |
| Thyroid | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| C-cell adenoma | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | 49 1 |
| Parathyroid | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Adenoma, NOS | + | + | + | + | + | - | + | + | + | + | + | + | + | - | - | - | + | + | + | - | + | - | + | + | + | + | 32 1 |
| REPRODUCTIVE SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Mammary gland | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Adenoma, NOS | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | *50 1 |
| Adenocarcinoma, NOS | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Fibroma | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Fibroadenoma | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Vagina | X | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | 11 *50 |
| Squamous cell carcinoma | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Uterus | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | 1 |
| Carcinoma, NOS | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Adenocarcinoma, NOS | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Endometrial stromal polyp | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Carcinosarcoma | X | | | | | | X | | | | | | X | | | | | X | | | | | X | | | 2 2 6 | |
| Ovary | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Granulosa cell carcinoma | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | 1 49 |
| NERVOUS SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Brain | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | 49 |
| MUSCULOSKELETAL SYSTEM | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Bone | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Osteosarcoma | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | *50 1 |
| Muscle | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Rhabdomyosarcoma | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | *50 1 |
| ALL OTHER SYSTEMS | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Multiple organs, NOS | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Carcinosarcoma, metastatic | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Malignant lymphoma, NOS | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Lower leg, NOS | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Sarcoma, NOS | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | *50 1 2 1 |

* Animals necropsied

TABLE H3. ANALYSIS OF PRIMARY TUMORS IN FEMALE OSBORNE-MENDEL RATS IN THE TWO-YEAR GAVAGE STUDY OF TRICHLOROETHYLENE

| | Untreated Control | Vehicle Control | 500 mg/kg | 1,000 mg/kg |
|---|-------------------|-----------------|-------------|-------------|
| Hematopoietic System: Leukemia | | | | |
| Overall Rates (a) | 2/50 (4%) | 0/50 (0%) | 3/50 (6%) | 0/50 (0%) |
| Adjusted Rates (b) | 6.5% | 0.0% | 13.9% | 0.0% |
| Terminal Rates (c) | 0/19 (0%) | 0/20 (0%) | 1/11 (9%) | 0/7 (0%) |
| Life Table Tests (d) | | P=0.528 | P=0.089 | (e) |
| Incidental Tumor Tests (d) | | P=0.596 | P=0.109 | (e) |
| Cochran-Armitage Trend Test (d) | | P=0.640 | | |
| Fisher Exact Test | | | P=0.121 | (e) |
| Pituitary: Adenoma | | | | |
| Overall Rates (a) | 14/47 (30%) | (f) 14/46 (30%) | 9/47 (19%) | 8/46 (17%) |
| Adjusted Rates (b) | 62.4% | 56.5% | 57.9% | 50.6% |
| Terminal Rates (c) | 11/19 (58%) | 10/20 (50%) | 5/11 (45%) | 2/7 (29%) |
| Life Table Tests (d) | | P=0.387 | P=0.519 | P=0.458 |
| Incidental Tumor Tests (d) | | P=0.353N | P=0.466N | P=0.364N |
| Cochran-Armitage Trend Test (d) | | P=0.084N | | |
| Fisher Exact Test | | | P=0.154N | P=0.111N |
| Pituitary: Adenoma or Carcinoma | | | | |
| Overall Rates (a) | 14/47 (30%) | 15/46 (33%) | 9/47 (19%) | 8/46 (17%) |
| Adjusted Rates (b) | 62.4% | 58.5% | 57.9% | 50.6% |
| Terminal Rates (c) | 11/19 (58%) | 10/20 (50%) | 5/11 (45%) | 2/7 (29%) |
| Life Table Tests (d) | | P=0.475 | P=0.599 | P=0.541 |
| Incidental Tumor Tests (d) | | P=0.250N | P=0.355N | P=0.258N |
| Cochran-Armitage Trend Test (d) | | P=0.054N | | |
| Fisher Exact Test | | | P=0.106N | P=0.074N |
| Adrenal: Cortical Adenoma | | | | |
| Overall Rates (a) | 13/49 (27%) | 16/50 (32%) | 13/50 (26%) | 19/49 (39%) |
| Adjusted Rates (b) | 54.8% | 55.6% | 66.3% | 92.7% |
| Terminal Rates (c) | 9/19 (47%) | 9/20 (45%) | 6/11 (55%) | 6/7 (86%) |
| Life Table Tests (d) | | P=0.008 | P=0.365 | P=0.011 |
| Incidental Tumor Tests (d) | | P=0.100 | P=0.484N | P=0.127 |
| Cochran-Armitage Trend Test (d) | | P=0.272 | | |
| Fisher Exact Test | | | P=0.330N | P=0.310 |
| Adrenal: Pheochromocytoma | | | | |
| Overall Rates (a) | 4/49 (8%) | 8/50 (16%) | 4/50 (8%) | 1/49 (2%) |
| Adjusted Rates (b) | 16.1% | 30.4% | 18.8% | 6.7% |
| Terminal Rates (c) | 2/19 (11%) | 4/20 (20%) | 1/11 (9%) | 0/7 (0%) |
| Life Table Tests (d) | | P=0.080N | P=0.394N | P=0.118N |
| Incidental Tumor Tests (d) | | P=0.009N | P=0.178N | P=0.020N |
| Cochran-Armitage Trend Test (d) | | P=0.011N | | |
| Fisher Exact Test | | | P=0.178N | P=0.017N |
| Adrenal: Pheochromocytoma or Pheochromocytoma, Malignant | | | | |
| Overall Rates (a) | 4/49 (8%) | 9/50 (18%) | 6/50 (12%) | 1/49 (2%) |
| Adjusted Rates (b) | 16.1% | 33.1% | 26.0% | 6.7% |
| Terminal Rates (c) | 2/19 (11%) | 4/20 (20%) | 1/11 (9%) | 0/7 (0%) |
| Life Table Tests (d) | | P=0.073N | P=0.564N | P=0.084N |
| Incidental Tumor Tests (d) | | P=0.004N | P=0.252N | P=0.009N |
| Cochran-Armitage Trend Test (d) | | P=0.008N | | |
| Fisher Exact Test | | | P=0.288N | P=0.009N |
| Thyroid: C-Cell Adenoma | | | | |
| Overall Rates (a) | 5/49 (10%) | 15/49 (31%) | 5/49 (10%) | 1/49 (2%) |
| Adjusted Rates (b) | 24.2% | 60.7% | 32.7% | 6.7% |
| Terminal Rates (c) | 4/19 (21%) | 11/20 (55%) | 2/11 (18%) | 0/7 (0%) |
| Life Table Tests (d) | | P=0.006N | P=0.127N | P=0.013N |
| Incidental Tumor Tests (d) | | P<0.001N | P=0.053N | P=0.003N |
| Cochran-Armitage Trend Test (d) | | P<0.001N | | |
| Fisher Exact Test | | | P=0.011N | P<0.001N |

TABLE H3. ANALYSIS OF PRIMARY TUMORS IN FEMALE OSBORNE-MENDEL RATS IN THE TWO-YEAR GAVAGE STUDY OF TRICHLOROETHYLENE (Continued)

| | Untreated Control | Vehicle Control | 500 mg/kg | 1,000 mg/kg |
|---|-------------------|-----------------|------------|-------------|
| Pancreatic Islets: Islet Cell Adenoma | | | | |
| Overall Rates (a) | 0/50 (0%) | 3/50 (6%) | 1/48 (2%) | 0/49 (0%) |
| Adjusted Rates (b) | 0.0% | 14.1% | 3.2% | 0.0% |
| Terminal Rates (c) | 0/19 (0%) | 2/20 (10%) | 0/11 (0%) | 0/7 (0%) |
| Life Table Tests (d) | | P=0.153N | P=0.456N | P=0.294N |
| Incidental Tumor Tests (d) | | P=0.078N | P=0.365N | P=0.221N |
| Cochran-Armitage Trend Test (d) | | P=0.063N | | |
| Fisher Exact Test | | | P=0.324N | P=0.125N |
| Mammary Gland: Fibroadenoma | | | | |
| Overall Rates (a) | 17/50 (34%) | 14/50 (28%) | 7/50 (14%) | 11/50 (22%) |
| Adjusted Rates (b) | 50.0% | 45.1% | 39.4% | 56.3% |
| Terminal Rates (c) | 5/19 (26%) | 6/20 (30%) | 2/11 (18%) | 2/7 (29%) |
| Life Table Tests (d) | | P=0.335 | P=0.291N | P=0.350 |
| Incidental Tumor Tests (d) | | P=0.289N | P=0.104N | P=0.330N |
| Cochran-Armitage Trend Test (d) | | P=0.271N | | |
| Fisher Exact Test | | | P=0.070N | P=0.322N |
| Mammary Gland: Fibroma or Fibroadenoma | | | | |
| Overall Rates (a) | 17/50 (34%) | 15/50 (30%) | 8/50 (16%) | 12/50 (24%) |
| Adjusted Rates (b) | 50.0% | 46.9% | 42.2% | 57.7% |
| Terminal Rates (c) | 5/19 (26%) | 6/20 (30%) | 2/11 (18%) | 2/7 (29%) |
| Life Table Tests (d) | | P=0.320 | P=0.324N | P=0.338 |
| Incidental Tumor Tests (d) | | P=0.240N | P=0.096N | P=0.262N |
| Cochran-Armitage Trend Test (d) | | P=0.277N | | |
| Fisher Exact Test | | | P=0.077N | P=0.326N |
| Mammary Gland: All Tumors (g) | | | | |
| Overall Rates (a) | 17/50 (34%) | 16/50 (32%) | 9/50 (18%) | 14/50 (28%) |
| Adjusted Rates (b) | 50.0% | 48.2% | 48.6% | 69.5% |
| Terminal Rates (c) | 5/19 (26%) | 6/20 (30%) | 3/11 (27%) | 3/7 (43%) |
| Life Table Tests (d) | | P=0.202 | P=0.352N | P=0.229 |
| Incidental Tumor Tests (d) | | P=0.372N | P=0.106N | P=0.367N |
| Cochran-Armitage Trend Test (d) | | P=0.366N | | |
| Fisher Exact Test | | | P=0.083N | P=0.414N |
| Uterus: Adenocarcinoma | | | | |
| Overall Rates (a) | (h) 0/50 (0%) | 2/49 (4%) | 5/49 (10%) | 2/50 (4%) |
| Adjusted Rates (b) | 0.0% | 9.3% | 32.7% | 19.3% |
| Terminal Rates (c) | 0/19 (0%) | 1/20 (5%) | 3/11 (27%) | 1/7 (14%) |
| Life Table Tests (d) | | P=0.234 | P=0.080 | P=0.422 |
| Incidental Tumor Tests (d) | | P=0.426 | P=0.151 | P=0.580 |
| Cochran-Armitage Trend Test (d) | | P=0.573N | | |
| Fisher Exact Test | | | P=0.218 | P=0.684N |
| Uterus: Endometrial Stromal Polyp | | | | |
| Overall Rates (a) | 4/50 (8%) | 6/49 (12%) | 1/49 (2%) | 6/50 (12%) |
| Adjusted Rates (b) | 14.5% | 19.8% | 4.3% | 24.9% |
| Terminal Rates (c) | 2/19 (11%) | 2/20 (10%) | 0/11 (0%) | 0/7 (0%) |
| Life Table Tests (d) | | P=0.365 | P=0.122N | P=0.386 |
| Incidental Tumor Tests (d) | | P=0.379N | P=0.039N | P=0.414N |
| Cochran-Armitage Trend Test (d) | | P=0.558N | | |
| Fisher Exact Test | | | P=0.056N | P=0.606N |
| Multiple Organs: Carcinoma | | | | |
| Overall Rates (a) | 0/50 (0%) | (i) 1/50 (2%) | 3/50 (6%) | 0/50 (0%) |
| Adjusted Rates (b) | 0.0% | 3.8% | 10.9% | 0.0% |
| Terminal Rates (c) | 0/19 (0%) | 0/20 (0%) | 0/11 (0%) | 0/7 (0%) |
| Life Table Tests (d) | | P=0.482N | P=0.227 | P=0.602N |
| Incidental Tumor Tests (d) | | P=0.203N | P=0.434 | P=0.433N |
| Cochran-Armitage Trend Test (d) | | P=0.378N | | |
| Fisher Exact Test | | | P=0.309 | P=0.500N |

TABLE H3. ANALYSIS OF PRIMARY TUMORS IN FEMALE OSBORNE-MENDEL RATS IN THE TWO-YEAR GAVAGE STUDY OF TRICHLOROETHYLENE (Continued)

- (a) Number of tumor-bearing animals/number of animals examined at the site
- (b) Kaplan-Meier estimated tumor incidence at the end of the study after adjusting for intercurrent mortality
- (c) Observed tumor incidence at terminal kill
- (d) Beneath the control incidence are the P values associated with the trend test. Beneath the dosed group incidence are the P values corresponding to pairwise comparisons between that dosed group and the vehicle controls. The life table analysis regards tumors in animals dying prior to terminal kill as being (directly or indirectly) the cause of death. The incidental tumor test regards these lesions as nonfatal. The Cochran-Armitage and Fisher exact tests compare directly the overall incidence rates. A negative trend or lower incidence in a dosed group is indicated by (N).
- (e) No P value is reported because no tumors were observed in the 1,000 mg/kg and vehicle control groups.
- (f) Incidence includes one papillary adenoma.
- (g) Includes adenoma, adenocarcinoma, fibroma, and fibroadenoma. A mixed tumor, benign, was observed in a low dose animal that also had a fibroadenoma.
- (h) One adenoma, NOS, was present.
- (i) Carcinoma, NOS; site unknown.

TABLE H4. SUMMARY OF THE INCIDENCE OF NONNEOPLASTIC LESIONS IN FEMALE OSBORNE-MENDEL RATS IN THE GAVAGE STUDY OF TRICHLOROETHYLENE

| | CONTROL (UNTR) | CONTROL (VEH) | LOW DOSE | HIGH DOSE |
|--------------------------------------|----------------|---------------|----------|-----------|
| ANIMALS INITIALLY IN STUDY | 50 | 50 | 50 | 50 |
| ANIMALS NECROPSIED | 50 | 50 | 50 | 50 |
| ANIMALS EXAMINED HISTOPATHOLOGICALLY | 50 | 50 | 50 | 50 |
| INTEGUMENTARY SYSTEM | | | | |
| *Subcutaneous tissue | (50) | (50) | (50) | (50) |
| Abscess, chronic | | 2 (4%) | | 1 (2%) |
| RESPIRATORY SYSTEM | | | | |
| *Nasal cavity | (50) | (50) | (50) | (50) |
| Ulcer, NOS | | 1 (2%) | | |
| *Tracheal lumen | (50) | (50) | (50) | (50) |
| Vegetable foreign body | 1 (2%) | | | |
| #Trachea | (49) | (50) | (47) | (47) |
| Inflammation, NOS | | | 1 (2%) | |
| Inflammation, chronic | | | 1 (2%) | 1 (2%) |
| Inflammation, chronic focal | | | | 1 (2%) |
| Hyperplasia, epithelial | | | 1 (2%) | |
| #Lung | (50) | (50) | (50) | (50) |
| Vegetable foreign body | | 3 (6%) | 3 (6%) | |
| Emphysema, alveolar | | 1 (2%) | 1 (2%) | 2 (4%) |
| Collapse | | 1 (2%) | | |
| Congestion, NOS | 1 (2%) | 2 (4%) | 4 (8%) | 5 (10%) |
| Edema, NOS | 1 (2%) | 1 (2%) | 3 (6%) | 3 (6%) |
| Hemorrhage | 2 (4%) | 3 (6%) | 6 (12%) | 3 (6%) |
| Inflammation, NOS | 1 (2%) | | | 1 (2%) |
| Bronchopneumonia, focal | | 1 (2%) | | |
| Inflammation, focal | 3 (6%) | 5 (10%) | 1 (2%) | 1 (2%) |
| Inflammation, necrotizing | | 1 (2%) | | |
| Bronchopneumonia, acute | | | 1 (2%) | |
| Inflammation, acute/chronic | | | 1 (2%) | |
| Inflammation, chronic | | 2 (4%) | | 2 (4%) |
| Inflammation, chronic focal | 1 (2%) | 1 (2%) | 1 (2%) | |
| Inflammation, granulomatous | 1 (2%) | 4 (8%) | 2 (4%) | 2 (4%) |
| Inflammation, granulomatous focal | 3 (6%) | 1 (2%) | 4 (8%) | 8 (16%) |
| Inflammation necro granulomatous | | 1 (2%) | | |
| Infection, bacterial | 1 (2%) | | | |
| Proteinosis, alveolar | | 1 (2%) | | |
| Calcification, metastatic | | 1 (2%) | | |
| Foreign material, NOS | | | 1 (2%) | 1 (2%) |
| #Lung/alveoli | (50) | (50) | (50) | (50) |
| Histiocytosis | | 1 (2%) | | |
| HEMATOPOIETIC SYSTEM | | | | |
| #Bone marrow | (50) | (50) | (50) | (49) |
| Hyperplasia, granulocytic | | | | 1 (2%) |
| #Spleen | (49) | (49) | (48) | (49) |
| Congestion, NOS | | | 1 (2%) | |
| Hemorrhage | 1 (2%) | | | |
| Abscess, NOS | | 1 (2%) | | |
| Inflammation, granulomatous focal | | 1 (2%) | | |
| Necrosis, focal | | | 2 (4%) | |
| Hemosiderosis | 2 (4%) | | 1 (2%) | 1 (2%) |
| Depletion, lymphoid | | | 1 (2%) | |
| Hyperplasia, lymphoid | | | 1 (2%) | |
| Hematopoiesis | 11 (22%) | 8 (16%) | 14 (29%) | 11 (22%) |

TABLE H4. SUMMARY OF THE INCIDENCE OF NONNEOPLASTIC LESIONS IN FEMALE OSBORNE-MENDEL RATS IN THE TWO-YEAR GAVAGE STUDY OF TRICHLOROETHYLENE (Continued)

| | CONTROL (UNTR) | CONTROL (VEH) | LOW DOSE | HIGH DOSE |
|---|----------------|---------------|----------|-----------|
| HEMATOPOIETIC SYSTEM (Continued) | | | | |
| #Lymph node | (47) | (46) | (45) | (46) |
| Congestion, NOS | 1 (2%) | | | |
| Hemorrhage | 1 (2%) | | | |
| Degeneration, cystic | 1 (2%) | | | |
| #Mandibular lymph node | (47) | (46) | (45) | (46) |
| Hemorrhage | | | 1 (2%) | |
| Plasmacytosis | | 1 (2%) | | |
| #Mesenteric lymph node | (47) | (46) | (45) | (46) |
| Inflammation, chronic | | | 1 (2%) | |
| Hyperplasia, reticulum cell | | | 1 (2%) | |
| #Liver | (50) | (50) | (50) | (49) |
| Hematopoiesis | 1 (2%) | 2 (4%) | 1 (2%) | |
| #Thymus | (3) | (4) | (8) | (3) |
| Congestion, NOS | | | 1 (13%) | |
| Hemorrhage | | | 2 (25%) | 1 (33%) |
| CIRCULATORY SYSTEM | | | | |
| #Lung | (50) | (50) | (50) | (50) |
| Thrombosis, NOS | 1 (2%) | | | |
| #Heart | (50) | (49) | (50) | (49) |
| Hemorrhage | | | 1 (2%) | |
| Inflammation, chronic | | 1 (2%) | | |
| Inflammation, chronic focal | | 4 (8%) | 1 (2%) | |
| Endocardiosis | | | 1 (2%) | |
| Necrosis, focal | | 1 (2%) | | |
| *Coronary artery | (50) | (50) | (50) | (50) |
| Calcification, metastatic | 1 (2%) | | | |
| *Pulmonary artery | (50) | (50) | (50) | (50) |
| Mineralization | | 1 (2%) | | |
| Calcification, focal | 1 (2%) | 1 (2%) | | |
| Calcification, metastatic | 1 (2%) | | | |
| #Liver | (50) | (50) | (50) | (49) |
| Perivasculitis | | | 1 (2%) | |
| #Pancreas | (50) | (50) | (48) | (49) |
| Periarteritis | 1 (2%) | | | |
| DIGESTIVE SYSTEM | | | | |
| #Salivary gland | (46) | (50) | (50) | (47) |
| Atrophy, NOS | | 1 (2%) | | |
| Atrophy, focal | | | | 1 (2%) |
| #Liver | (50) | (50) | (50) | (49) |
| Congestion, NOS | | | 1 (2%) | |
| Hemorrhage | | | | 1 (2%) |
| Inflammation, focal | | | | 2 (4%) |
| Inflammation, active chronic | | | | 1 (2%) |
| Inflammation, chronic focal | 1 (2%) | | | |
| Peliosis hepatitis | 1 (2%) | | | |
| Necrosis, focal | 1 (2%) | | 5 (10%) | |
| Necrosis, diffuse | | 1 (2%) | | |
| Necrosis, central | | | 1 (2%) | |
| Metamorphosis, fatty | 1 (2%) | 1 (2%) | 1 (2%) | 1 (2%) |
| Lipoidosis | | | | 2 (4%) |
| Pigmentation, NOS | 1 (2%) | | | |
| Cytoplasmic change, NOS | 2 (4%) | 1 (2%) | 1 (2%) | 2 (4%) |
| Basophilic cyto change | 1 (2%) | | | |
| Focal cellular change | 1 (2%) | | | 1 (2%) |
| Clear cell change | | 1 (2%) | | 2 (4%) |
| Cytologic alteration, NOS | | 1 (2%) | | |
| Hepatocytomegaly | | | | 1 (2%) |

TABLE H4. SUMMARY OF THE INCIDENCE OF NONNEOPLASTIC LESIONS IN FEMALE OSBORNE-MENDEL RATS IN THE TWO-YEAR GAVAGE STUDY OF TRICHLOROETHYLENE (Continued)

| | CONTROL (UNTR) | CONTROL (VEH) | LOW DOSE | HIGH DOSE |
|-------------------------------------|----------------|---------------|----------|-----------|
| DIGESTIVE SYSTEM (Continued) | | | | |
| #Liver/centrilobular | (50) | (50) | (50) | (49) |
| Necrosis, NOS | | | | 1 (2%) |
| #Bile duct | (50) | (50) | (50) | (49) |
| Dilatation, NOS | | | | 1 (2%) |
| Inflammation, chronic | 1 (2%) | | 2 (4%) | 1 (2%) |
| Hyperplasia, NOS | 5 (10%) | 3 (6%) | 2 (4%) | 1 (2%) |
| Hyperplasia, focal | | 1 (2%) | 1 (2%) | |
| Hyperplasia, diffuse | | | 1 (2%) | |
| #Pancreas | (50) | (50) | (48) | (49) |
| Hemorrhage | | 1 (2%) | | |
| Inflammation, acute/chronic | | | 1 (2%) | |
| Inflammation, chronic | | | 2 (4%) | 1 (2%) |
| Inflammation, chronic focal | | | | 1 (2%) |
| Atrophy, focal | | 1 (2%) | | |
| #Pancreatic acinus | (50) | (50) | (48) | (49) |
| Atrophy, NOS | 1 (2%) | | | |
| Atrophy, focal | 3 (6%) | 2 (4%) | 3 (6%) | 3 (6%) |
| *Esophageal lumen | (50) | (50) | (50) | (50) |
| Hemorrhage | 2 (4%) | 4 (8%) | | 1 (2%) |
| #Esophagus | (50) | (48) | (50) | (50) |
| Hyperkeratosis | 1 (2%) | | | |
| #Thoracic esophagus | (50) | (48) | (50) | (50) |
| Granulation tissue | | | | 1 (2%) |
| #Stomach | (50) | (48) | (50) | (49) |
| Inflammation, suppurative | | | | 1 (2%) |
| Calcification, metastatic | | 1 (2%) | | |
| #Gastric mucosa | (50) | (48) | (50) | (49) |
| Inflammation, NOS | 1 (2%) | | | |
| Hyperplasia, epithelial | 1 (2%) | | | |
| Hyperkeratosis | 1 (2%) | | | |
| #Forestomach | (50) | (48) | (50) | (49) |
| Inflammation, NOS | 1 (2%) | | | |
| Hyperplasia, epithelial | 1 (2%) | 1 (2%) | | |
| Hyperkeratosis | 1 (2%) | | | |
| #Jejunum | (50) | (48) | (49) | (49) |
| Hemorrhage | | 1 (2%) | | |
| #Colon | (50) | (48) | (47) | (50) |
| Parasitism | 7 (14%) | 5 (10%) | 7 (15%) | 7 (14%) |
| URINARY SYSTEM | | | | |
| #Kidney | (50) | (50) | (50) | (49) |
| Mineralization | 8 (16%) | 3 (6%) | 3 (6%) | |
| Cast, NOS | 4 (8%) | 7 (14%) | 2 (4%) | 1 (2%) |
| Hydronephrosis | 2 (4%) | | 1 (2%) | 4 (8%) |
| Cyst, NOS | 2 (4%) | 1 (2%) | | |
| Hemorrhage | | 1 (2%) | | |
| Inflammation, acute focal | 1 (2%) | | | |
| Inflammation, acute/chronic | | 1 (2%) | | |
| Inflammation, chronic | | | | 1 (2%) |
| Inflammation, chronic focal | 1 (2%) | | | |
| Inflammation, granulomatous focal | | 1 (2%) | | |
| Fibrosis | | | | 1 (2%) |
| Nephropathy | 43 (86%) | 43 (86%) | 41 (82%) | 47 (96%) |
| Nephropathy, toxic | | | 30 (60%) | 39 (80%) |
| Necrosis, focal | 1 (2%) | | | |
| Calcification, focal | 1 (2%) | 4 (8%) | 2 (4%) | 2 (4%) |
| Calcification, metastatic | | 1 (2%) | | |
| Cytomegaly | | | 48 (96%) | 49 (100%) |
| Hyperplasia, tubular cell | | 1 (2%) | 1 (2%) | 3 (6%) |
| #Kidney/cortex | (50) | (50) | (50) | (49) |
| Degeneration, NOS | 1 (2%) | | | |

TABLE H4. SUMMARY OF THE INCIDENCE OF NONNEOPLASTIC LESIONS IN FEMALE OSBORNE-MENDEL RATS IN THE TWO-YEAR GAVAGE STUDY OF TRICHLOROETHYLENE (Continued)

| | CONTROL (UNTR) | CONTROL (VEH) | LOW DOSE | HIGH DOSE |
|-----------------------------------|----------------|---------------|----------|-----------|
| URINARY SYSTEM (Continued) | | | | |
| #Kidney/medulla | (50) | (50) | (50) | (49) |
| Hyperplasia, NOS | 1 (2%) | | | |
| Hyperplasia, epithelial | 6 (12%) | 5 (10%) | 2 (4%) | |
| #Kidney/tubule | (50) | (50) | (50) | (49) |
| Dilatation, NOS | | | 1 (2%) | |
| #Kidney/pelvis | (50) | (50) | (50) | (49) |
| Hemorrhage | 2 (4%) | | | |
| Inflammation, acute | 1 (2%) | 1 (2%) | | |
| Hyperplasia, epithelial | 2 (4%) | 4 (8%) | 5 (10%) | 6 (12%) |
| #Urinary bladder | (48) | (48) | (49) | (50) |
| Hemorrhage | | 1 (2%) | | |
| Inflammation, acute | | 1 (2%) | | |
| Inflammation, acute/chronic | 1 (2%) | 1 (2%) | | |
| Inflammation, chronic | 1 (2%) | | | |
| Hyperplasia, epithelial | 1 (2%) | 1 (2%) | | |
| ENDOCRINE SYSTEM | | | | |
| #Pituitary | (47) | (46) | (47) | (46) |
| Cyst, NOS | | 1 (2%) | 2 (4%) | 1 (2%) |
| Congestion, NOS | | 1 (2%) | | |
| Hemorrhage | | | 1 (2%) | |
| Hyperplasia, focal | 1 (2%) | 2 (4%) | 1 (2%) | 2 (4%) |
| #Adrenal | (49) | (50) | (50) | (49) |
| Congestion, NOS | | | | 1 (2%) |
| Hemorrhage | 1 (2%) | 1 (2%) | 2 (4%) | |
| Degeneration, NOS | | | | 2 (4%) |
| Degeneration, cystic | | 1 (2%) | | |
| Necrosis, focal | | | 1 (2%) | 1 (2%) |
| Metamorphosis, fatty | | | 1 (2%) | |
| Lipoidosis | 2 (4%) | 2 (4%) | | |
| Angiectasis | 3 (6%) | 6 (12%) | 9 (18%) | 12 (24%) |
| #Adrenal cortex | (49) | (50) | (50) | (49) |
| Degeneration, NOS | | | | 1 (2%) |
| Lipoidosis | | | 3 (6%) | 1 (2%) |
| Focal cellular change | 1 (2%) | | 1 (2%) | |
| Atrophy, NOS | 1 (2%) | | | |
| Hyperplasia, NOS | | | 1 (2%) | |
| Hyperplasia, focal | 1 (2%) | 1 (2%) | 3 (6%) | |
| Angiectasis | 1 (2%) | | | |
| #Adrenal medulla | (49) | (50) | (50) | (49) |
| Hyperplasia, NOS | | | | 1 (2%) |
| Hyperplasia, focal | 1 (2%) | 1 (2%) | 2 (4%) | 1 (2%) |
| #Thyroid | (49) | (49) | (49) | (49) |
| Cyst, NOS | | 2 (4%) | 3 (6%) | 6 (12%) |
| Inflammation, chronic | | 1 (2%) | | |
| Hyperplasia, C-cell | 4 (8%) | 1 (2%) | 2 (4%) | 1 (2%) |
| REPRODUCTIVE SYSTEM | | | | |
| *Mammary gland | (50) | (50) | (50) | (50) |
| Cyst, NOS | 1 (2%) | | | 1 (2%) |
| Hemorrhage | 1 (2%) | | | |
| Degeneration, NOS | | 1 (2%) | | |
| Hyperplasia, NOS | 1 (2%) | | | |
| Lactation | 11 (22%) | 4 (8%) | 4 (8%) | 7 (14%) |
| *Vagina | (50) | (50) | (50) | (50) |
| Hyperplasia, epithelial | | | 1 (2%) | 1 (2%) |
| *Vaginal mucosa | (50) | (50) | (50) | (50) |
| Hyperplasia, cystic | 1 (2%) | | | |

TABLE H4. SUMMARY OF THE INCIDENCE OF NONNEOPLASTIC LESIONS IN FEMALE OSBORNE-MENDEL RATS IN THE TWO-YEAR GAVAGE STUDY OF TRICHLOROETHYLENE (Continued)

| | CONTROL (UNTR) | CONTROL (VEH) | LOW DOSE | HIGH DOSE |
|--|----------------|---------------|----------|-----------|
| REPRODUCTIVE SYSTEM (Continued) | | | | |
| #Uterus | (50) | (49) | (49) | (50) |
| Dilatation, NOS | 4 (8%) | 7 (14%) | 2 (4%) | 6 (12%) |
| Hydrometra | 6 (12%) | 2 (4%) | 8 (16%) | 6 (12%) |
| Epidermal inclusion cyst | | 1 (2%) | | |
| Hemorrhage | 3 (6%) | | 1 (2%) | |
| Hematometra | 1 (2%) | 1 (2%) | 4 (8%) | |
| Inflammation, NOS | 1 (2%) | 2 (4%) | 1 (2%) | |
| Ulcer, NOS | | | 4 (8%) | 1 (2%) |
| Pyometra | | 1 (2%) | 3 (6%) | 1 (2%) |
| Inflammation, acute | 3 (6%) | | 1 (2%) | 2 (4%) |
| Inflammation, acute focal | 1 (2%) | | | |
| Inflammation, acute necrotizing | | | 1 (2%) | |
| Inflammation, acute/chronic | 1 (2%) | | 2 (4%) | 3 (6%) |
| Inflammation, chronic | | | 1 (2%) | 1 (2%) |
| Necrosis, hemorrhagic | | | | 1 (2%) |
| Pigmentation, NOS | | | 1 (2%) | |
| Hyperplasia, epithelial | 1 (2%) | 1 (2%) | | 1 (2%) |
| Hyperkeratosis | | 2 (4%) | 1 (2%) | 1 (2%) |
| Metaplasia, NOS | | 1 (2%) | 1 (2%) | |
| Metaplasia, squamous | 1 (2%) | 3 (6%) | 7 (14%) | 4 (8%) |
| #Uterus/endometrium | (50) | (49) | (49) | (50) |
| Cyst, NOS | | | 1 (2%) | |
| Ulcer, NOS | | 1 (2%) | | |
| Inflammation, acute | | | 1 (2%) | 1 (2%) |
| Hyperplasia, NOS | | 1 (2%) | 2 (4%) | 3 (6%) |
| Hyperplasia, cystic | 12 (24%) | 8 (16%) | 9 (18%) | 4 (8%) |
| Metaplasia, NOS | 3 (6%) | 1 (2%) | 1 (2%) | |
| Metaplasia, squamous | 1 (2%) | 2 (4%) | 1 (2%) | 2 (4%) |
| #Ovary | (49) | (48) | (49) | (49) |
| Cyst, NOS | 3 (6%) | 4 (8%) | 1 (2%) | 5 (10%) |
| Follicular cyst, NOS | 1 (2%) | 2 (4%) | 1 (2%) | 2 (4%) |
| Inflammation, suppurative | | 1 (2%) | | |
| Inflammation, acute | 1 (2%) | 1 (2%) | | |
| NERVOUS SYSTEM | | | | |
| #Brain | (50) | (50) | (50) | (49) |
| Cyst, NOS | | 1 (2%) | | |
| Hemorrhage | | 1 (2%) | 2 (4%) | |
| Abscess, NOS | | | 1 (2%) | |
| Inflammation, chronic focal | | | 1 (2%) | |
| *Spinal cord | (50) | (50) | (50) | (50) |
| Hemorrhage | | | 1 (2%) | |
| SPECIAL SENSE ORGANS | | | | |
| *Harderian gland | (50) | (50) | (50) | (50) |
| Inflammation, chronic | | 1 (2%) | | 1 (2%) |
| Inflammation, chronic focal | | | | 1 (2%) |
| MUSCULOSKELETAL SYSTEM | | | | |
| None | | | | |

TABLE H4. SUMMARY OF THE INCIDENCE OF NONNEOPLASTIC LESIONS IN FEMALE OSBORNE-MENDEL RATS IN THE TWO-YEAR GAVAGE STUDY OF TRICHLOROETHYLENE (Continued)

| | CONTROL (UNTR) | CONTROL (VEH) | LOW DOSE | HIGH DOSE |
|-----------------------------------|----------------|---------------|----------|-----------|
| BODY CAVITIES | | | | |
| *Peritoneum | (50) | (50) | (50) | (50) |
| Inflammation, acute focal | | 1 (2%) | | |
| *Pleura | (50) | (50) | (50) | (50) |
| Inflammation, acute | | | 1 (2%) | |
| *Pericardium | (50) | (50) | (50) | (50) |
| Inflammation, chronic | | 1 (2%) | | |
| Inflammation, chronic focal | | 1 (2%) | | |
| ALL OTHER SYSTEMS | | | | |
| *Multiple organs | (50) | (50) | (50) | (50) |
| Hemorrhage | | 1 (2%) | | |
| SPECIAL MORPHOLOGY SUMMARY | | | | |
| Auto/necropsy/histo perf | | 1 | | |

* Number of animals receiving complete necropsy examination; all gross lesions including masses examined microscopically.

Number of animals examined microscopically at this site

APPENDIX I

GENETIC TOXICOLOGY OF

TRICHLOROETHYLENE

| | PAGE |
|--|------|
| TABLE I1 MUTAGENICITY OF TRICHLOROETHYLENE IN <i>SALMONELLA TYPHIMURIUM</i> | 252 |
| TABLE I2 MUTAGENICITY OF TRICHLOROETHYLENE IN L5178Y MOUSE LYMPHOMA CELLS IN THE ABSENCE OF S9 | 253 |
| TABLE I3 MUTAGENICITY OF TRICHLOROETHYLENE IN L5178Y MOUSE LYMPHOMA CELLS IN THE PRESENCE OF S9 | 254 |
| TABLE I4 INDUCTION OF SISTER CHROMATID EXCHANGES IN CHINESE HAMSTER OVARY CELLS BY TRICHLOROETHYLENE | 255 |
| TABLE I5 INDUCTION OF CHROMOSOMAL ABERRATIONS IN CHINESE HAMSTER OVARY CELLS BY TRICHLOROETHYLENE | 255 |

TABLE II. MUTAGENICITY OF TRICHLOROETHYLENE IN *SALMONELLA TYPHIMURIUM*

| Strain | Dose ($\mu\text{g}/\text{plate}$) | Revertants/plate (a,b) | | |
|--------|--|------------------------|--------------------|-------------------|
| | | -S9 | +S9 (rat) | +S9 (hamster) |
| TA100 | 0 | 148 \pm 4.6 | 136 \pm 9.3 | 139 \pm 5.0 |
| | 10 | 151 \pm 2.0 | 131 \pm 5.5 | 136 \pm 10.8 |
| | 33 | 137 \pm 5.2 | 114 \pm 11.9 | 152 \pm 6.1 |
| | 100 | 143 \pm 6.1 | 115 \pm 10.9 | 143 \pm 11.3 |
| | 333 | 128 \pm 4.2 | 135 \pm 6.6 | 130 \pm 0.9 |
| | 1,000 | (c) 123 \pm 2.9 | (c) 134 \pm 13.7 | (c) 141 \pm 5.2 |
| TA1535 | 0 | 29 \pm 2.0 | 11 \pm 0.6 | 14 \pm 1.3 |
| | 10 | 28 \pm 0.6 | 16 \pm 1.5 | 14 \pm 1.2 |
| | 33 | 27 \pm 1.0 | 11 \pm 2.2 | 14 \pm 1.0 |
| | 100 | 22 \pm 2.9 | 11 \pm 0.3 | 12 \pm 1.7 |
| | 333 | 24 \pm 0.9 | 17 \pm 3.5 | 15 \pm 3.2 |
| | 1,000 | (c) 11 \pm 4.3 | (c) 8 \pm 3.5 | (c) 10 \pm 0.9 |
| TA98 | 0 | 19 \pm 2.1 | 24 \pm 3.8 | 28 \pm 1.2 |
| | 10 | 21 \pm 2.0 | 21 \pm 1.2 | 28 \pm 1.3 |
| | 33 | 16 \pm 2.3 | 29 \pm 0.3 | 38 \pm 2.3 |
| | 100 | 18 \pm 1.5 | 26 \pm 4.5 | 28 \pm 3.5 |
| | 333 | 19 \pm 4.0 | 25 \pm 2.6 | 24 \pm 0.0 |
| | 1,000 | (c) 17 \pm 3.0 | (c) 25 \pm 3.8 | (c) 25 \pm 2.2 |
| TA1537 | 0 | 7 \pm 1.5 | 6 \pm 1.0 | 4 \pm 0.9 |
| | 10 | 7 \pm 2.0 | 8 \pm 1.2 | 8 \pm 0.3 |
| | 33 | 7 \pm 0.7 | 7 \pm 2.7 | 9 \pm 0.9 |
| | 100 | 6 \pm 0.6 | 6 \pm 1.5 | 7 \pm 1.5 |
| | 333 | 9 \pm 0.7 | 6 \pm 1.9 | 7 \pm 2.7 |
| | 1,000 | (c) 5 \pm 0.6 | (c) 6 \pm 1.8 | (c) 6 \pm 0.3 |

(a) The S9 fractions were prepared from the liver of Aroclor 1254-induced male Sprague-Dawley rats and male Syrian hamsters. Cells and study compound or dimethyl sulfoxide were incubated for 20 minutes at 37° C in the presence of either S9 or buffer. After the addition of soft agar, the contents of each tube were poured onto minimal medium, and the plates were incubated at 37° C for 48 hours (Haworth et al., 1983). The experiment was performed twice, each in triplicate; because the results were similar, data from only one experiment are shown.

(b) Mean \pm standard error

(c) Slight toxicity

TABLE 12. MUTAGENICITY OF TRICHLOROETHYLENE IN L5178Y MOUSE LYMPHOMA CELLS IN THE ABSENCE OF S9 (a)

| Compound | Concentration | Total Mutant Clones | Cloning Efficiency (percent) | Relative Total Growth (percent) | Mutation Frequency (mutants/10 ⁶ clonable cells) |
|---------------------------|---------------|---------------------|------------------------------|---------------------------------|---|
| DMSO | 1% | 68 | 114.8 | 110 | 20 |
| | | 74 | 94.0 | 98 | 26 |
| | | 94 | 105.5 | 99 | 30 |
| | | 68 | 111.7 | 91 | 20 |
| Ethylmethane sulfonate | 250 µg/kg | 1,188 | 67.8 | 33.8 | 584 |
| Trichloroethylene (nl/ml) | 25.0 | 69 | 103.2 | 95.0 | 22 |
| | | 84 | 98.7 | 85.6 | 28 |
| | | 56 | 94.8 | 89.2 | 20 |
| | 50.0 | 49 | 95.0 | 92.5 | 17 |
| | | 74 | 91.8 | 66.1 | 27 |
| | | 83 | 110.5 | 73.3 | 25 |
| | 100.0 | 46 | 99.8 | 62.5 | 15 |
| | | 53 | 97.3 | 80.9 | 18 |
| | | 107 | 124.3 | 66.3 | 29 |
| | 200.0 | 63 | 90.3 | 55.7 | 23 |
| | | 72 | 104.7 | 45.2 | 23 |
| | | 58 | 79.8 | 17.4 | 24 |

(a) Experiments were performed twice, and all doses were tested in duplicate or triplicate. Because the results were similar, data from only one experiment are shown. The protocol was basically that of Clive et al. (1979). Cells (6×10^5 /ml) were treated for 4 hours at 37° C in medium, washed, resuspended in medium, and incubated for 48 hours at 37° C. After expression, 3×10^6 cells were plated in medium supplemented with trifluorothymidine for selection of cells that were mutant at the thymidine kinase (TK) locus, and 600 cells were plated in nonselective medium to determine the percentage of viable cells.

TABLE 13. MUTAGENICITY OF TRICHLOROETHYLENE IN L5178Y MOUSE LYMPHOMA CELLS IN THE PRESENCE OF S9 (a)

| Compound | Concentration | Total Mutant Clones | Cloning Efficiency (percent) | Relative Total Growth (percent) | Mutation Frequency (mutants/10 ⁶ clonable cells) |
|---------------------------|---------------|---------------------|------------------------------|---------------------------------|---|
| DMSO | 1% | 227 | 107.2 | 102 | 71 |
| | | 178 | 93.0 | 114 | 64 |
| | | 233 | 113.0 | 102 | 69 |
| | | 261 | 105.8 | 98 | 82 |
| 3-Methylcholanthrene | 2.5 µg/ml | 802 | 83.2 | 35.2 | 321 |
| | | 813 | 90.5 | 44.6 | 299 |
| | | 775 | 79.5 | 21.5 | 325 |
| Trichloroethylene (nl/ml) | 25.0 | 331 | 98.7 | 65.6 | 112 |
| | | 265 | 96.0 | 73.7 | 92 |
| | | 308 | 104.7 | 73.9 | 98 |
| | 50.0 | 342 | 105.8 | 55.8 | 108 |
| | | 311 | 108.3 | 60.6 | 96 |
| | | 284 | 108.0 | 59.9 | 88 |
| | 100.0 | 403 | 89.5 | 48.0 | 150 |
| | | 382 | 99.8 | 39.2 | 128 |
| | | 257 | 81.7 | 38.7 | 105 |
| | 200.0 | 498 | 95.0 | 17.2 | 175 |
| | | 448 | 94.3 | 19.9 | 158 |
| | | 441 | 99.8 | 15.9 | 147 |

(a) Experiments were performed twice, and all doses were tested in duplicate. Because the results were similar, data from only one experiment are shown. The protocol was basically that of Clive et al. (1979). Cells (6×10^5 /ml) were treated for 4 hours at 37° C in medium, washed, resuspended in medium, and incubated for 48 hours at 37° C. After expression, 3×10^6 cells were plated in medium supplemented with trifluorothymidine for selection of cells that were mutant at the thymidine kinase (TK) locus, and 600 cells were plated in nonselective medium to determine the percentage of viable cells. S9 was prepared from the liver of Aroclor 1254-induced male F344 rats.

TABLE 14. INDUCTION OF SISTER CHROMATID EXCHANGES IN CHINESE HAMSTER OVARY CELLS BY TRICHLOROETHYLENE (a)

| | - S9 (b) | | + S9 (c) | |
|-------------------|--------------|---------------|------------------|---------------|
| | Dose (µg/ml) | SCEs/Cell (d) | Dose (µg/ml) | SCEs/Cell (d) |
| DMSO (10 µl) | | 7.6 | DMSO (10 µl) | 8.4 |
| Trichloroethylene | | | | |
| | 499 | 7.7 | 401 | 10.1 |
| | 596 | 8.2 | 499 | 10.1 |
| | 700 | 9.1 | 596 | 9.5 |
| Mitomycin C | 0.005 | 20.4 | Cyclophosphamide | 20.0 |
| | | | 1.5 | |

(a) SCE = sister chromatid exchange

(b) In the absence of S9, Chinese hamster ovary cells were incubated with study compound or solvent for 2 hours at 37° C. Then 10 µM bromodeoxyuridine (BrdU) was added, and incubation was continued for 24 hours. Cells were washed, fresh medium containing BrdU (10 µM) and colcemid (0.1 µg/ml) was added, and incubation was continued for 2-3 hours.

(c) In the presence of S9, Chinese hamster ovary cells were incubated with study compound or solvent for 2 hours at 37° C. Then cells were washed, and medium containing 10 µM BrdU was added. Cells were incubated for a further 26 hours, with colcemid (0.1 µg/ml) present for the final 2-3 hours. S9 was from the liver of Aroclor 1254-induced male Sprague-Dawley rats.

(d) Cells were then collected by mitotic shake-off, treated for 3 minutes with KCl (75 mM), washed twice with fixative, and dropped onto slides and air-dried. Staining was by a modified technique (after Perry and Wolff, 1974; Goto et al., 1978).

TABLE 15. INDUCTION OF CHROMOSOMAL ABERRATIONS IN CHINESE HAMSTER OVARY CELLS BY TRICHLOROETHYLENE (a)

| | - S9 (b) | | + S9 (c) | |
|-------------------|--------------|--|------------------|--|
| | Dose (µg/ml) | Abs/100 Cells (percent cells with abs) | Dose (µg/ml) | Abs/100 Cells (percent cells with abs) |
| DMSO (10 µl) | | 7 (7) | DMSO (10 µl) | 2 (2) |
| Trichloroethylene | | | | |
| | 745 | 2 (2) | 499 | 6 (5) |
| | 801 | 4 (4) | 700 | 7 (5) |
| | | | 745 | 8 (6) |
| | 14,900 | 4 (3) | 14,900 | 2 (2) |
| Mitomycin C | 0.500 | 24 (18) | Cyclophosphamide | 94 (35) |
| | | | 50 | |

(a) Abs = aberrations

(b) In the absence of S9, Chinese hamster ovary cells were incubated with study compound or solvent for 8-10 hours at 37° C. Cells were then washed, and fresh medium containing colcemid (0.1 µg/ml) was added. After a further 2-3 hours of incubation, cells were harvested by mitotic shake-off, fixed, and stained in 6% Giemsa.

(c) In the presence of S9, cells were incubated with study compound or solvent for 2 hours at 37° C. Cells were then washed, medium was added, and incubation was continued for 8-10 hours. Colcemid (0.1 µg/ml) was added for the last 2-3 hours of incubation; then cells were harvested and fixed as above. S9 was from the liver of Aroclor 1254-induced male Sprague-Dawley rats.

APPENDIX J

CHEMICAL CHARACTERIZATION

OF TRICHLOROETHYLENE

APPENDIX J. CHEMICAL CHARACTERIZATION

I. Identity and Purity Determinations of Lot No. TB05-206AA Performed by the Analytical Chemistry Laboratory

A. Physical properties

| 1. Boiling point: | <u>Determined</u> | <u>Literature Values</u> |
|-------------------------|---|---|
| | 86.0° ± 0.8(δ)° C at 737 mm (visual, micro boiling point); 84.5°-87° C (Dupont 900 DTA) | 86.7° C at 760 mm (Gallant, 1966) |
| 2. Index of refraction: | $n_D^{20}: 1.4766 \pm 0.0002(\delta)$ | $n_D^{20}: 1.4776$ (Bachman et al., 1950) |
| 3. Density: | $d_{23}^{22}: 1.46315 \pm 0.00002(\delta) \text{ g/ml}$ | $d^{23}: 1.458$ (read from graph) (Gallant, 1966) |

B. Spectral data

1. Infrared

| | | |
|-------------|---|---|
| Instrument: | Beckman IR-12 | |
| Cell: | 0.015-mm liquid cell, sodium chloride windows | |
| Results: | See Figure J-1 | Consistent with literature spectrum (Sadler Standard Spectra) |

2. Ultraviolet/visible

| | | |
|----------------|--|---|
| Instrument: | Cary 118 | |
| Concentration: | 1 mg/ml | 0.0002% |
| Solvent: | Methanol | Methanol |
| Results: | No absorbance between 800 and 350 nm. No maximum between 208 and 350 nm, but a gradual increase in absorbance toward the solvent cutoff at 208 nm. | No maximum observed in the near ultraviolet range (Lacher et al., 1950) |




FIGURE J-1. INFRARED ABSORPTION SPECTRUM OF TRICHLOROETHYLENE (LOT NO. TB05-206)

APPENDIX J. CHEMICAL CHARACTERIZATION

| | | | |
|--|------------------------------------|---|-----------|
| 3. Nuclear magnetic resonance | <u>Determined</u> | <u>Literature Values</u> | |
| Instrument: | Varian HA-100 | | |
| Solvent: | Neat, with added tetramethylsilane | | |
| Assignments: | See Figure J-2 | Consistent with literature spectrum (Sadler Standard Spectra) | |
| Chemical shift (δ): | s, 6.34 ppm | | |
| Integration ratios: | 1.00 | | |
| C. Water analysis (Karl Fischer): | 0.0097% \pm 0.0020(δ)% | | |
| D. Elemental analysis | | | |
| Element | C | H | Cl |
| | <hr/> | | |
| Theory | 18.28 | 0.77 | 80.95 |
| Determined | 18.31 | 0.78 | 80.69 |
| | 18.45 | 0.80 | 80.85 |
| | <hr/> | | |




FIGURE J-2. NUCLEAR MAGNETIC RESONANCE SPECTRUM OF TRICHLOROETHYLENE (LOT NO. TB05-206AA)

APPENDIX J. CHEMICAL CHARACTERIZATION

E. Gas chromatography

Instrument: Tracor MT 220

Detector: Flame ionization

Inlet temperature: 200° C

Detector temperature: 255° C

System 1

Column: 10% Carbowax 20M-TPA on 80/100 Chromosorb W(AW), 1.8 m × 4 mm ID, glass

Oven temperature program: 50° C, 5 min; 50°-200° C at 10° C/min

Results: Major peak and 12 impurities. The total area of the impurities is less than 0.04% of the major peak.

| <u>Peak No.</u> | <u>Retention Time (min)</u> | <u>Retention Time Relative to Major Peak</u> | <u>Area (percent of major peak)</u> |
|-----------------|-----------------------------|--|-------------------------------------|
| 1 | 0.3 | 0.10 | trace, < 0.001 |
| 2 | 1.2 | 0.40 | trace, < 0.001 |
| 3 | 1.4 | 0.47 | 0.002 |
| 4 | 2.0 | 0.67 | 0.001 |
| 5 | 2.1 | 0.70 | trace, < 0.001 |
| 6 | 2.4 | 0.80 | 0.001 |
| 7 | 3.0 | 1.00 | 100 |
| 8 | 7.4 | 2.47 | 0.001 |
| 9 | 7.9 | 2.63 | 0.02 |
| 10 | 9.8 | 3.27 | trace, < 0.001 |
| 11 | 10.3 | 3.43 | 0.003 |
| 12 | 11.8 | 3.93 | trace, < 0.001 |
| 13 | 12.4 | 4.13 | 0.003 |

APPENDIX J. CHEMICAL CHARACTERIZATION

System 2

Column: 20% SP2100/0.1% Carbowax 1500 on 80/100 Supelcoport, 1.8 m × 4 mm ID, glass

Oven temperature program: 50° C, 5 min; 50°-170° C at 10° C/min

Results: Major peak and eight impurities. All impurities have areas less than 0.01% of the major peak. The total area of the impurities is less than 0.02% of the major peak.

| <u>Peak No.</u> | <u>Retention Time (min)</u> | <u>Retention Time Relative to Major Peak</u> | <u>Area (percent of major peak)</u> |
|-----------------|-----------------------------|--|-------------------------------------|
| 1 | 2.6 | 0.50 | 0.0005 |
| 2 | 3.8 | 0.73 | 0.004 |
| 3 | 5.2 | 1.00 | 100 |
| 4 | 7.4 | 1.42 | 0.002 |
| 5 | 7.7 | 1.48 | 0.002 |
| 6 | 8.8 | 1.69 | 0.004 |
| 7 | 9.4 | 1.81 | 0.0005 |
| 8 | 10.2 | 1.96 | 0.0005 |
| 9 | 13.9 | 2.67 | 0.002 |

APPENDIX J. CHEMICAL CHARACTERIZATION

II. Identity and Purity of Lot No. TB08-039AA

| A. Spectral data | <u>Determined</u> | <u>Literature Values</u> |
|--|---|--|
| 1. Infrared | | |
| Instrument: | Beckman IR-12 | |
| Cell: | Thin film between silver chloride plates | |
| Results: | See Figure J-3 | Spectrum consistent with literature spectrum (Sadler Standard Spectra) |
| 2. Ultraviolet/visible | | |
| Instrument: | Cary 118 | |
| Solvent: | Methanol | Methanol |
| Results: | No absorbance between 800 and 350 nm at a concentration of 1% (v/v). No absorbance maximum between 350 and 215 nm but a gradual increase in absorbance toward 215 nm at a concentration of 0.0004% (v/v). | No maximum observed in the near ultraviolet range. (Lacher et al., 1950) |
| 3. Nuclear magnetic resonance | | |
| Instrument: | Varian HA-100 | |
| Solvent: | Neat, with added tetramethylsilane | |
| Assignments: | See Figure J-4 | Spectrum consistent with literature spectrum (Sadler Standard Spectra) |
| Chemical shift (δ): | s, 6.43 ppm | |




FIGURE J-3. INFRARED ABSORPTION SPECTRUM OF TRICHLOROETHYLENE (LOT NO. TB08-039AA)




FIGURE J-4. NUCLEAR MAGNETIC RESONANCE SPECTRUM OF TRICHLOROETHYLENE
(LOT NO. TB08-039AA)

APPENDIX J. CHEMICAL CHARACTERIZATION

B. Elemental analysis

| Element | C | H | Cl |
|------------|----------------|--------------|----------------|
| Theory | 18.28 | 0.77 | 80.95 |
| Determined | 18.29 18.15 | 0.80 0.81 | 80.78 80.95 |

C. Water analysis (Karl Fischer): < 0.003%

D. Gas chromatography

Instrument: Varian 3700
Detector: Flame ionization
Inlet temperature: 200° C
Carrier gas: Nitrogen, 70 ml/min

System 1

Column: 10% Carbowax 20M-TPA on 80/100 Chromosorb W (AW), 1.8 m × 4 mm ID, glass

Oven temperature program: 50° C, for 5 min; 50°-200° C at 10° C/min

Sample injected: Neat liquid (3.5 µl) and solutions of 1.0% and 0.5% (v/v) trichloroethylene in *o*-dichlorobenzene to quantitate the major peak and check for detector overload

Results: Major peak and one impurity after the major peak with an area 0.02% of the major peak.

| <u>Peak No.</u> | <u>Retention Time (min)</u> | <u>Retention Time Relative to Major Peak</u> | <u>Area (percent of major peak)</u> |
|-----------------|-----------------------------|--|-------------------------------------|
| 1 | 3.9 | 1.00 | 100 |
| 2 | 11.2 | 2.87 | 0.02 |

APPENDIX J. CHEMICAL CHARACTERIZATION

System 2

Detector temperature: 240° C

Column: 20% SP2100/0.1% Carbowax 1500 on 100/120 Supelcoport, 1.8 m × 4 mm ID, glass

Oven temperature program: 50° C, for 5 min; 50°-170° C at 10° C/min

Samples injected: Neat liquid (3 µl) and solutions of 1.0% and 0.5% trichloroethylene in *o*-dichlorobenzene to quantitate the major peak and check for detector overload.

Results: Major peak and one impurity after the major peak with an area of 0.02% of the major peak.

| <u>Peak No.</u> | <u>Retention Time (min)</u> | <u>Retention Time Relative to Major Peak</u> | <u>Area (percent of major peak)</u> |
|-----------------|-----------------------------|--|-------------------------------------|
| 1 | 6.3 | 1.00 | 100 |
| 2 | 8.3 | 1.32 | 0.02 |

III. Quantitation of Impurity Present in Lot No. TB05-206AA

Analysis by gas chromatography

A. System

Instrument: Varian 3700

Detector: Flame ionization

Inlet temperature: 200° C

Detector temperature: 250° C

Carrier gas: Nitrogen, 70 ml/min

Column: 80/100 Carbopack C/0.1% SP1000; 1.8 m × 4 mm ID, glass

Oven temperature program: 50° C, isothermal

B. Results: Trichloroethylene, when injected as a neat liquid on the above system, had a retention time of 16.0 minutes and contained a peak with a retention time of 7.2 minutes with a shoulder at 7.6 minutes. The shoulder was enhanced by addition of an epichlorohydrin standard to the sample (0.001%, v/v, relative to trichloroethylene).

C. Conclusions: Calculation of the amount of epichlorohydrin present in the unspiked sample by the standard addition method showed that, if epichlorohydrin was present in this lot of trichloroethylene, it was at a level less than or equal to 0.001% (v/v).

APPENDIX J. CHEMICAL CHARACTERIZATION

IV. Identification of the Impurity in Lot No. TB05-206AA

Analysis by gas chromatography/mass spectrometry

A. System

Instrument: Varian 311-A Mass Spectrometer interfaced via a single-stage glass jet separator to a Varian 2700 Gas Chromatograph. Data handled by an Incos 2300 Data System

Column: 80/100 Carbowax C/0.1% SP1000; 1.8 m × 2 mm ID, glass

Column oven temperature: Ambient

Heated zone temperatures

Inlet: 170°C

Transfer line: 300°C

Helium separator: 275°C

Carrier: Helium, 30 ml/min

Scan range: 5-350 amu

Scan times (sec): up: 2.50 top: 0.00
down: 0.00 bottom: 0.50

Accelerator voltage: 3000

Electron multiplier voltage: -2000

Resolution: 801

Sample injected: 5 µl of neat trichloroethylene

- B. Results:** The impurity corresponding to the peak reported in III.B. (with a retention time of 7.2 minutes) eluted in 9.65 minutes on this system. The spectrum obtained from this peak and a literature spectrum of *n*-pentane are given below.

| <u>Spectrum of Impurity</u> | | <u>Literature Spectrum of <i>n</i>-Pentane</u> | |
|-----------------------------|---|--|---|
| <u>m/e</u> | <u>Relative Abundance (percent of m/e 43)</u> | <u>m/e</u> | <u>Relative Abundance (percent of m/e 43)</u> |
| 43 | 100 | 43 | 100 |
| 41 | 62 | 42 | 59 |
| 42 | 61 | 41 | 41 |
| 39 | 26 | 27 | 34 |
| 27 | 25 | 29 | 25 |
| 29 | 17 | 39 | 14 |
| 57 | 11 | 57 | 13 |
| 72 | 7 | 72 | 9 |

- C. Conclusions:** The impurity with the retention time of 7.2 minutes, reported in III.B., was identified on the basis of its mass spectrum as *n*-pentane.

APPENDIX J. CHEMICAL CHARACTERIZATION

V. Chemical Stability at the Study Laboratory

A. **Storage conditions:** The chemical was stored at -20°C .

B. Bulk analysis

Instrument: Varian 3700 with CDS 111 Data System

Detector: Flame ionization

Temperature

Detector: 200°C

Injector: 130°C

Oven: 70°C , isothermal

Carrier gas: Nitrogen, 60 ml/min

Column: 10% OV-101 on 100/120 Supelcoport

Sample injected: Neat liquid (1 μl)

C. Results

| <u>Date of Analysis</u> | <u>Lot No.</u> | <u>Purity (percent)</u> | |
|-----------------------------|----------------|-------------------------|------------------|
| | | <u>Bulk</u> | <u>Reference</u> |
| 04/12/79 | TB05206AA | -- | 99.96 |
| 07/31/79 | | 99.15 | |
| 12/17/79 | | 99.96 | 99.99 |
| 04/23/80 | | 99.96 | 99.96 |
| 08/19/80 | | 99.15 | 100.00 |
| 11/20/80 | TB08039AA | -- | 99.95 |
| 12/03/80 | | 99.95 | 99.94 |
| 03/13/81 | | 99.95 | 99.95 |
| 03/17/81 | | 99.96 | 99.95 |
| 06/22/81 | | 99.95 | 99.94 |
| 12/21/81 | | 99.97 | 99.98 |

D. **Conclusion:** No notable degradation was observed.

APPENDIX K

IDENTIFICATION OF FOREIGN MATERIAL FOUND IN TRICHLOROETHYLENE, LOT NO. TB05-206AA

APPENDIX K. IDENTIFICATION OF FOREIGN MATERIAL

I. Analysis

- A. Undissolved solids:** The original sample of this lot of trichloroethylene was received in a 55-gallon drum and then transferred to 5-gallon drums. In July 1979, two drums were returned to Midwest Research Institute for analysis of flocculent material. The contents of two of these drums was filtered through ashless filter paper. The material was air dried and then weighed. Ten gallons (85 kg) of trichloroethylene was found to contain 260 mg of undissolved solid material. This represents undissolved solids at a level of 3 ppm. The empty drums were then cut open and visually inspected. None of the solids had remained in the drums. The drums were found to be uncoated on the inside and patches of light corrosion were found, probably due to the action of hydrochloric acid.
- B. Dissolved solids:** The amount of dissolved solids was determined by evaporating 100-ml aliquots of filtered trichloroethylene to dryness and weighing the residue. The trichloroethylene was found to contain dissolved solids at a level of $25.6 \pm 1.7(8)$ ppm.
- C. Melting point determination:** A Büchi Model 510[®] melting point apparatus was used to determine the melting characteristics of the foreign material. No melting was observed. At 110° C, the sample began to darken and continued to darken until complete decomposition was evident at 290° C.

D. Elemental analysis (a)

| <u>Element</u> | <u>Percent Found in Foreign Material</u> |
|----------------|--|
| C | 41.29 |
| H | 4.21 |
| N | <0.05 |
| Cl | 0.95 |

(a) Analysis performed by Galbraith Laboratories, Inc.,
2323 Sycamore Drive, Knoxville, Tennessee 37921

APPENDIX K. IDENTIFICATION OF FOREIGN MATERIAL

E. Spark-source mass spectrometry (a)

| | | | | | |
|-----------|----------|-----------|----------|------------|--------|
| Uranium | <1.2 | Iodine | 2.9 | Calcium | >1.0% |
| Thorium | <2.4 | Tellurium | | Potassium | >0.5% |
| Bismuth | 2.5 | Tin | 3.1 | Chlorine | ≈1,800 |
| Lead | 31 | Indium | Internal | Sulfur | ≈1,600 |
| Thallium | <3.0 | | standard | Phosphorus | ≈4,000 |
| Mercury | NR | Yttrium | 6.2 | Silicon | >0.5% |
| Rhenium | Internal | Strontium | 39 | Aluminum | 600 |
| | standard | Bromine | 17 | Magnesium | >1.0% |
| Tungsten | <1.2 | Arsenic | <1.2 | Sodium | >1.0% |
| Hafnium | <2.8 | Zinc | 380 | Fluorine | 300 |
| Erbium | <1.8 | Copper | 990 | Oxygen | NR |
| Samarium | <1.2 | Nickel | 1.4 | Nitrogen | NR |
| Neodymium | <1.8 | Cobalt | 1.3 | Carbon | NR |
| Cerium | 2.0 | Iron | ≈3,100 | Boron | 10 |
| Lanthanum | 1.6 | Manganese | 360 | Lithium | 8.6 |
| Barium | 3.4 | Chromium | 38 | | |
| Cesium | | Titanium | 58 | | |

(a) Analysis performed by Camp Dresser and McKee, Inc., 11455 W. 48th Avenue, Wheat Ridge, Colorado 80033. Data are expressed in parts per million (by weight) of foreign material unless otherwise noted. All elements for which values are not entered are < 1.0 ppm.
NR = not reported

F. Free acid titration: Titration of aliquots of trichloroethylene with 0.01 N sodium hydroxide indicated the presence of $6.14 \pm 0.25(6)$ ppm free acid (as hydrochloric acid).

G. Infrared spectroscopy

Instrument: Beckman IR-12

Cell: Thin film on silver chloride plates

Sample preparation: Toluene suspension of precipitated material

Results: The absorbances at 2,530, 1,795, 1,440, and 700 cm^{-1} are characteristic of calcium carbonate. The other absorbances at 2,930, 2,860, 967, and 912 cm^{-1} are compatible with an unsaturated hydrocarbon (Figure K-1).

H. Direct inlet mass spectrometry: The 70 eV mass spectrum (Table K1) was obtained from the foreign material. The spectrum indicates that the material was a mixture. Alkane and alkene fragmentation series were clearly evident. However, it was felt that the spectrum obtained may not be representative of the compound because of the nonvolatile and thermally unstable nature of the material (see Section I.C.). Spectra obtained from low density polyethylene, silicon rubber, and Teflon[®] did not correspond to the spectrum of the foreign material.




FIGURE K-1. INFRARED ABSORPTION SPECTRUM OF PRECIPITATE FOUND IN TRICHLOROETHYLENE (LOT NO. TB05-206AA)

TABLE K1. ANALYSIS BY DIRECT INLET MASS SPECTROMETRY OF FOREIGN MATERIAL FOUND IN TRICHLOROETHYLENE

| m/e | Relative Abundance (percent of m/e 69) | m/e | Relative Abundance (percent of m/e 69) |
|------------|---|------------|---|
| 69 | 100 | 137 | 18 |
| 55 | 94 | 121 | 18 |
| 98 | 92 | 99 | 18 |
| 83 | 75 | 59 | 17 |
| 57 | 60 | 54 | 17 |
| 97 | 60 | 87 | 17 |
| 81 | 56 | 125 | 17 |
| 129 | 55 | 108 | 16 |
| 73 | 52 | 151 | 15 |
| 95 | 51 | 80 | 15 |
| 43 | 50 | 93 | 15 |
| 41 | 49 | 42 | 14 |
| 67 | 48 | 79 | 14 |
| 44 | 41 | 124 | 14 |
| 71 | 38 | 39 | 14 |
| 60 | 36 | 239 | 13 |
| 85 | 36 | 94 | 13 |
| 84 | 35 | 107 | 13 |
| 311 | 35 | 138 | 13 |
| 96 | 35 | 152 | 12 |
| 135 | 34 | 171 | 12 |
| 70 | 32 | 53 | 12 |
| 111 | 32 | 126 | 11 |
| 109 | 30 | 101 | 11 |
| 56 | 30 | 29 | 11 |
| 112 | 26 | 143 | 11 |
| 313 | 26 | 72 | 11 |
| 236 | 24 | 61 | 10 |
| 123 | 22 | 89 | 10 |
| 339 | 21 | 91 | 10 |
| 110 | 21 | 113 | 10 |
| 116 | 21 | 149 | 10 |
| 237 | 20 | 157 | 10 |
| 185 | 19 | 165 | 10 |
| | | 115 | 10 |

APPENDIX K. IDENTIFICATION OF FOREIGN MATERIAL

II. Summary of Analytical Data

The precipitated material was present in trichloroethylene at a level of 3 ppm. Nonvolatile and nonfilterable residue was present at a level of $25.6 \pm 1.7(\delta)$ ppm. The results of elemental analysis indicated that the precipitate contained 41.29% carbon, 4.21% hydrogen, 0.95% chlorine, and no detectable nitrogen. Spark-source mass spectrometry indicated the presence of calcium, magnesium, and sodium at levels greater than 1% and potassium at levels greater than 0.5%. The dried precipitate was a fibrous-type material that decomposed on heating. Decomposition began at 110° C and was completed at 290° C. The infrared spectroscopy (see Figure K-1) indicated that the material was a mixture of calcium carbonate and probably an unsaturated hydrocarbon. Mass spectroscopy also indicated that the material was a mixture probably containing alkane and alkene materials.

APPENDIX L

PREPARATION AND CHARACTERIZATION

OF DOSE MIXTURES

APPENDIX L. PREPARATION AND CHARACTERIZATION

I. Stability of Trichloroethylene in Corn Oil at Room Temperature

- A. Sample preparation:** A 1% (w/v) solution of trichloroethylene in corn oil was prepared for each day of the study as follows: 10 ml of corn oil was transferred into a 50-ml Hypo-vial, the vial was sealed, and then approximately 95 mg of trichloroethylene (measured exactly for each sample) was added via a 100- μ l syringe. The samples were shaken and stored at room temperature from 1 to 7 days.
- B. Extraction and analysis:** Each sample was extracted with 20 ml of methanol, which was injected into the sample vial via a 10-ml syringe. Samples for analysis were withdrawn directly from the top methanol layer in the vial and analyzed by gas chromatography with the following system.

Instrument: Tracor MT 220

Column: Chromosorb 102, 60/80 mesh, 1.8 m \times 2 mm ID, glass

Detection: Flame ionization

Oven temperature: 160 $^{\circ}$ C, isothermal

Detector temperature: 260 $^{\circ}$ C

Inlet temperature: 200 $^{\circ}$ C

Retention time of compound: 4.15 min

C. Results

| <u>End of Day</u> | <u>Average Percent Trichloroethylene in Chemical/Vehicle Mixture (a)</u> |
|-------------------|--|
| 1 | 1.00 \pm 0.05 |
| 2 | 0.96 \pm 0.05 |
| 3 | 0.99 \pm 0.05 |
| 4 | 0.98 \pm 0.05 |
| 5 | 1.00 \pm 0.05 |
| 6 | 0.98 \pm 0.05 |
| 7 | 0.99 \pm 0.05 |

(a) Corrected for an average spiked recovery yield of 61.7% \pm 0.9%.
Target concentration of chemical/vehicle mixture, 1.00%.

- D. Conclusion:** Trichloroethylene mixed with corn oil is stable for 7 days at room temperature.

APPENDIX L. PREPARATION AND CHARACTERIZATION

II. Stability of Trichloroethylene in Corn Oil at 4° C

A. **Sample preparation and analysis:** Sample preparation and analysis procedures were the same as for the room temperature study.

B. Results

| Number of Weeks After Mixing | Target Concentration (a) | |
|---|---------------------------------|--------------|
| | 59.5 | 235.5 |
| 1 | 60.4 | 240 |
| 2 | 59.2 | 236.9 |
| 3 | 56.9 | 236.9 |
| 4 | 58.4 | 236.9 |
| 5 | 56.9 | 235.3 |
| 6 | 57.3 | 235.3 |
| 7 | 55.4 | 227.6 |
| 8 | 55.8 | 239.6 |

(a) Milligrams trichloroethylene/milliliter sample

C. **Conclusions:** Trichloroethylene mixed with corn oil is stable for 4 weeks at 4° C.

APPENDIX M

METHODS OF ANALYSIS OF DOSE MIXTURES

APPENDIX M. METHODS OF ANALYSIS

I. Study Laboratory

A. System

Instrument: Varian 3700 with CDS III Integrator

Detector: Flame ionization

Detector temperature: 200° C

Injector temperature: 130° C

Oven temperature: 70° C (isothermal)

Carrier: Nitrogen, 30 ml/min

Column: 10% OV-101 on 100/120 Supelcoport; 10 foot × 1/8 inch stainless steel until 4/28/80, then 15 foot × 1/4 inch glass column

- B. Procedure:** Initially, the concentration of trichloroethylene in dose mixtures was calculated from the percent recovery after the trichloroethylene/corn oil mixture was extracted into methanol. The recovery for this method was usually about 65% and was replaced on March 16, 1979, with a more reliable procedure, the internal-standard method.

A 2-ml aliquot of a trichloroethylene/corn oil mixture was diluted to 25 ml with an internal-standard solution containing 1.5 mg octane/ml chloroform. A 3- μ l aliquot of this mixture was injected into the gas chromatograph. If the sample was too concentrated (i.e., if the peak went offscale), the solution was rediluted as stated above with 2- to 25-ml portions of internal-standard solution.

APPENDIX M. METHODS OF ANALYSIS

II. Analysis Performed at Midwest Research Institute

A. Extraction and analysis: The samples were allowed to equilibrate to room temperature; aliquots (approximately 5 g) of the chemical/vehicle mixture were transferred into 50-ml septum vials and accurately weighed. Methanol (5 ml) was added to each aliquot. Blanks and standards were prepared by weighing corn oil (5 g) into septum vials and spiking each sample with methanol or trichloroethylene in methanol (5 ml). Two standard solutions of trichloroethylene were prepared, and the standards were diluted further with methanol and added to the corn oil samples to give standards bracketing the concentration range of the sample. The samples were extracted immediately after spiking as follows.

Methanol (10-20 ml) containing *n*-butyl alcohol (6 or 11.34 mg/ml) or *n*-hexyl alcohol (1.8 or 2.5 mg/ml) as an internal standard was added to each sample. The septum vials were sealed and the compound extracted into the methanol by mechanical mixing on a vortex mixer for 30 seconds followed by 30 seconds in an ultrasonic vibratory bath. The vials were centrifuged for 3 minutes to separate the corn oil/methanol layers and sampled directly from the top methanol layer for analysis by gas chromatography.

Instrumental parameters

Instrument: Tracor MT-220 with Hewlett-Packard 3380A integrator or Varian 3700 with autosampler and Varian CDS III-C integrator

Column: GP 20% SP2100/0.1% Carbowax 1500, on 100/120 mesh Supelcoport, 1.8 m × 4 mm ID, glass, silanized

Detection: Flame ionization

| | <u>Tracor MT-220</u> | <u>Varian 3700</u> |
|---|----------------------|--|
| Temperatures | | |
| Inlet: | 230° C | 200° C |
| Oven: | 65° C, isothermal | 80° C, isothermal |
| Detector: | 230° C | 250° C |
| Carrier gas: | Nitrogen, 70 ml/min | Nitrogen, 30 ml/min |
| Volume of solution injected: | 5 µl | 3 µl |
| Retention time of trichloroethylene: | 3.7-4.1 min | 2.3-5.5 min |
| Retention time of internal standard: | 2.8-3.1 min | 4.2 min (<i>n</i> -butyl alcohol) 6.5-10.6 min (hexyl alcohol) |

B. Quality control procedures: Analyses were performed in triplicate for the study chemical/vehicle sample and in duplicate for the blank corn oil sample. Spiked samples were prepared from two separately weighed standard solutions. The chemical/vehicle samples, the blank corn oil samples, and the spiked samples were all extracted and prepared for analysis in the same manner. Blank samples showed no interference from the corn oil at the retention time of the major component. A calibration curve was established with the spiked sample extracts.

APPENDIX M. METHODS OF ANALYSIS

III. Analysis Performed by Raltech Scientific Services, Inc.

A. Standard preparation: Trichloroethylene (6.25 g) was weighed into a 25-ml volumetric flask and diluted to volume with methanol. This stock solution was 250 mg/ml. Dilutions of the stock standard were made to obtain working standards of 5, 25, 50, and 125 mg/ml. Dilutions were made in methanol. *n*-Butanol (5 g) was weighed into a 500-ml volumetric flask and diluted to volume with methanol to obtain a 10 mg/ml internal standard solution.

B. Sample preparation and analysis: Sample aliquots (5 g) were transferred into 50-ml septum vials along with 5 ml of methanol and 20 ml of the methanol internal standard. The septum vials were sealed with Teflon[®]-faced septa, mixed on a Vortex mixer for 1 minute, and placed in an ultrasonic bath for 30 seconds. The vials were centrifuged for 5 minutes, and the upper methanol layer was sampled directly by syringe for gas chromatographic analysis.

C. Instrumental operating parameters

Instrument: Hewlett-Packard 5730 A Gas Chromatograph

Detector: Flame ionization

Column: 1.8 m × 4 mm ID glass on-column injection

Column packing: 20% SP2100/0.1% Carbowax 1500 by weight on 100/120 Supelcoport

Carrier gas: Nitrogen, 50 ml/min

Temperatures

Column: 70° C

Detector: 250° C

Injector: 200° C

Injection volume: 5 µl

Data were generated and analyzed on a Hewlett-Packard 3350 laboratory automation system.

D. Quality assurance: The sample was analyzed in triplicate. Duplicate blanks were prepared by adding 5 ml of methanol and 20 ml of methanol internal standard to 5 g of blank corn oil and extracting. Aliquots (5 ml) of the working trichloroethylene standards, ranging from 5 to 50 mg/ml, were added to 5 g of blank corn oil along with 20 ml of methanol internal standard. These extracted standards were equivalent to 5-250 mg/g of trichloroethylene in corn oil.

APPENDIX N

RESULTS OF ANALYSIS OF DOSE MIXTURES

| | | PAGE |
|----------|--|------|
| TABLE N1 | RESULTS OF ANALYSIS OF DOSE MIXTURES IN THE TWO-YEAR GAVAGE STUDIES OF TRICHLOROETHYLENE | 286 |
| TABLE N2 | RESULTS OF REFEREE ANALYSIS OF DOSE MIXTURES IN THE TWO-YEAR GAVAGE STUDIES OF TRICHLOROETHYLENE | 288 |

TABLE N1. RESULTS OF ANALYSIS OF DOSE MIXTURES IN THE TWO-YEAR GAVAGE STUDIES OF TRICHLOROETHYLENE

| Date Mixed | Concentration (mg/ml) | | Percent D/T | Date Mixed | Concentration (mg/ml) | | Percent D/T |
|------------|-----------------------|--------------------|-------------|------------|-----------------------|--------------------|-------------|
| | Target (T) | Determined (D) (a) | | | Target (T) | Determined (D) (a) | |
| 11/16/78 | 333.5 | 322.1 | 96.6 | 09/26/79 | 619.8 | 597.4 | 96.4 |
| 12/08/78 | 164.3 | 164.4 | 100.1 | 10/12/79 | 615.9 | 581.7 | 94.4 |
| 12/14/78 | 72.5 | 66.6 | 91.9 | 10/15/79 | 325.3 | 319.1 | 98.1 |
| | 170.0 | 175.4 | 103.2 | 10/17/79 | 164.8 | 157.1 | 95.3 |
| | 334.5 | 273.2 | 81.7 | | 730.0 | 664.0 | 91.0 |
| 12/21/78 | 189.5 | 145.9 | 77.0 | 10/19/79 | 365.4 | 351.0 | 96.1 |
| 12/28/78 | 200.0 | 210.6 | 105.3 | 11/21/79 | 499.5 | 479.6 | 96.0 |
| 01/04/79 | 218.4 | 214.7 | 98.3 | 12/10/79 | 546.0 | 506.8 | 92.8 |
| 01/11/79 | 78.5 | 77.2 | 98.3 | 12/14/79 | 553.3 | 512.6 | 92.6 |
| | 356.6 | 354.8 | 99.5 | 12/17/79 | 628.8 | 582.6 | 92.7 |
| | 225.4 | 214.7 | 95.3 | 12/20/79 | 566.1 | 522.8 | 92.4 |
| 01/18/79 | 232.0 | 238.2 | 102.7 | 01/02/80 | 600.8 | 602.4 | 100.3 |
| 01/25/79 | 240.2 | 214.7 | 89.4 | 01/09/80 | 663.6 | 625.7 | 94.3 |
| 01/31/79 | 243.0 | 236.6 | 97.4 | | 577.2 | 566.7 | 98.2 |
| | 161.8 | 150.8 | 93.2 | 01/16/80 | 628.4 | 592.6 | 94.3 |
| 02/07/79 | 357.5 | 364.1 | 101.8 | 01/24/80 | 696.2 | 666.7 | 95.8 |
| | 74.9 | 69.0 | 92.1 | 02/06/80 | 70.6 | 79.4 | 112.5 |
| | 242.2 | 251.2 | 103.7 | | 70.6 | 70.3 | (b) 99.6 |
| | 182.8 | 175.4 | 96.0 | | 74.4 | 84.8 | 114.0 |
| 02/14/79 | 246.9 | 255.1 | 103.3 | | 74.4 | 76.8 | (b) 103.2 |
| | 198.5 | 191.8 | 96.6 | 02/07/80 | 339.4 | 364.8 | 107.5 |
| 02/21/79 | 510.0 | 537.1 | 105.3 | | 182.5 | 184.6 | 101.2 |
| | 423.0 | 434.5 | 102.7 | | 218.7 | 240.1 | 109.8 |
| 02/28/79 | 512.3 | 519.2 | 101.3 | | 218.7 | 203.1 | (b) 92.9 |
| | 441.2 | 441.6 | 100.1 | | 122.3 | 140.3 | 114.7 |
| | 699.8 | 727.7 | 104.0 | | 122.3 | 119.5 | (b) 97.7 |
| | 514.9 | 525.1 | 102.0 | 02/14/80 | 306.9 | 335.3 | 109.3 |
| 03/07/79 | 462.9 | 460.9 | 99.6 | | 160.5 | 172.9 | 107.7 |
| | 156.0 | 168.4 | 107.9 | | 204.5 | 225.4 | 110.2 |
| 03/14/79 | 477.0 | 513.7 | 107.7 | | 204.5 | 187.4 | (b) 91.6 |
| 03/21/79 | 497.6 | 499.2 | 100.3 | | 101.6 | 113.7 | 111.9 |
| 03/28/79 | 492.9 | 517.8 | 105.1 | | 101.6 | 98.5 | (b) 96.9 |
| 04/04/79 | 728.4 | 765.1 | 105.0 | 02/22/80 | 308.3 | 337.7 | 109.5 |
| | 156.9 | 165.8 | 105.7 | | 183.7 | 206.4 | 112.4 |
| | 68.0 | 69.8 | 102.6 | | 183.7 | 173.2 | (b) 94.3 |
| | 544.2 | 570.6 | 104.9 | | 159.9 | 178.6 | 111.7 |
| | 519.8 | 545.9 | 105.0 | | 159.9 | 149.8 | (b) 93.7 |
| 04/11/79 | 519.0 | 511.7 | 98.6 | | 91.7 | 105.3 | 114.8 |
| 04/18/79 | 512.9 | 542.7 | 105.8 | | 91.7 | 94.5 | (b) 103.1 |
| 04/25/79 | 524.5 | 543.0 | 103.5 | 02/27/80 | 116.1 | 135.5 | 116.7 |
| 05/01/79 | 729.8 | 736.9 | 101.0 | | 116.1 | 119.4 | (b) 102.8 |
| | 161.9 | 175.8 | 108.6 | | 592.6 | 619.2 | 104.5 |
| | 561.6 | 601.4 | 107.1 | | 153.2 | 175.7 | 114.7 |
| | 524.5 | 562.0 | 107.1 | | 153.2 | 145.8 | (b) 95.2 |
| 05/30/79 | 751.6 | 756.1 | 100.6 | 02/28/80 | 130.6 | 138.5 | 106.0 |
| | 573.0 | 602.1 | 105.1 | | 130.5 | 118.7 | (b) 91.0 |
| | 535.2 | 547.1 | 102.2 | | 198.3 | 205.8 | 103.8 |
| | 156.5 | 164.5 | 105.1 | 03/05/80 | 330.4 | 358.9 | 108.6 |
| 06/27/79 | 161.2 | 166.6 | 103.3 | | 75.8 | 88.0 | 116.1 |
| | 730.2 | 731.0 | 100.1 | | 75.8 | 74.3 | (b) 98.0 |
| | 561.2 | 584.5 | 104.2 | | 69.4 | 78.6 | 113.3 |
| | 548.2 | 567.8 | 103.6 | | 69.4 | 69.5 | (b) 100.1 |
| 07/19/79 | 591.0 | 567.0 | 95.9 | | 218.7 | 258.2 | 118.1 |
| 07/25/79 | 755.0 | 778.1 | 103.1 | | 218.7 | 204.7 | (b) 93.6 |
| | 164.4 | 164.6 | 100.1 | 03/12/80 | 214.3 | 222.7 | 103.9 |
| 07/31/79 | 597.4 | 573.6 | 96.0 | 03/19/80 | 319.5 | 344.6 | 107.9 |
| 08/15/79 | 606.5 | 578.6 | 95.4 | | 319.5 | 342.2 | 107.1 |
| 08/21/79 | 85.4 | 84.7 | 99.2 | | 158.8 | 175.1 | 110.3 |
| | 716.2 | 680.5 | 95.0 | | 158.8 | 148.4 | (b) 93.5 |
| 08/27/79 | 612.5 | 557.5 | 91.0 | 03/26/80 | 224.0 | 248.1 | 110.8 |
| 09/13/79 | 615.8 | 572.1 | 92.9 | | 224.0 | 211.1 | (b) 94.2 |
| 09/19/79 | 732.8 | 662.0 | 90.3 | | 282.1 | 314.2 | 111.4 |
| | 164.8 | 161.5 | 98.0 | | 282.1 | 261.4 | (b) 107.9 |

TABLE N1. RESULTS OF ANALYSIS OF DOSE MIXTURES IN THE TWO-YEAR GAVAGE STUDIES OF TRICHLOROETHYLENE (Continued)

| Date Mixed | Concentration (mg/ml) | | Percent D/T | Date Mixed | Concentration (mg/ml) | | Percent D/T |
|------------|-----------------------|--------------------|-------------|------------|-----------------------|--------------------|-------------|
| | Target (T) | Determined (D) (a) | | | Target (T) | Determined (D) (a) | |
| 03/27/80 | 391.6 | 421.0 | 107.5 | 10/08/80 | 149.3 | 176.9 | 118.5 |
| | 275.5 | 305.4 | 110.9 | | 149.3 | 154.5 | (b) 103.5 |
| | 275.5 | 254.4 | (b) 92.3 | | 118.6 | 138.7 | 116.9 |
| 04/02/80 | 174.2 | 168.4 | 96.7 | | 118.6 | 126.5 | (b) 106.7 |
| | 69.7 | 72.2 | 103.6 | 10/17/80 | 240.8 | 266.0 | 110.5 |
| | 66.0 | 68.9 | 104.4 | | 240.8 | 238.1 | (b) 98.9 |
| 04/08/80 | 121.9 | 137.0 | 112.4 | | 147.7 | 162.9 | 110.3 |
| 04/09/80 | 162.6 | 170.9 | 105.1 | | 147.7 | 146.7 | (b) 99.3 |
| | 106.4 | 111.0 | 104.3 | 10/23/80 | 159.3 | 165.2 | 103.7 |
| 04/16/80 | 198.4 | 199.7 | 100.7 | | 109.6 | 110.0 | 100.4 |
| | 96.8 | 101.1 | 104.4 | 10/29/80 | 191.0 | 193.5 | 101.3 |
| 04/22/80 | 215.7 | 220.1 | 102.0 | | 103.5 | 107.0 | 103.4 |
| | 141.8 | 146.0 | 103.0 | 11/05/80 | 304.0 | 303.5 | 99.8 |
| | 148.1 | 140.8 | 95.1 | | 226.6 | 227.4 | 100.4 |
| 04/23/80 | 118.9 | 120.3 | 101.2 | 11/13/80 | 240.4 | 235.5 | 98.0 |
| 04/30/80 | 225.4 | 230.3 | 102.2 | | 149.1 | 156.0 | 104.6 |
| | 65.1 | 68.0 | 104.5 | 11/18/80 | 313.9 | 316.3 | 100.8 |
| | 73.8 | 79.6 | 107.9 | | 226.7 | 232.1 | 102.4 |
| | 327.3 | 330.2 | 100.9 | 11/25/80 | 370.6 | 363.6 | 98.1 |
| 05/07/80 | 313.4 | 324.0 | 103.4 | | 209.4 | 215.8 | 103.1 |
| | 213.7 | 221.8 | 103.8 | 12/10/80 | 247.2 | 250.9 | 101.5 |
| | 416.4 | 429.5 | 103.1 | | 148.4 | 155.0 | 104.4 |
| | 278.7 | 279.9 | 100.4 | 12/17/80 | 157.0 | 164.3 | 104.6 |
| 05/13/80 | 216.5 | 220.7 | 101.9 | | 110.6 | 115.6 | 104.5 |
| | 329.1 | 323.2 | 98.2 | 12/23/80 | 189.9 | 218.4 | 115.0 |
| 05/22/80 | 292.2 | 286.1 | 97.9 | | 189.9 | 215.3 | (b) 113.4 |
| | 228.4 | 233.8 | 102.4 | | 104.1 | 113.8 | 109.3 |
| 06/04/80 | 106.2 | 113.1 | 106.5 | 01/07/81 | 465.9 | 454.5 | 97.6 |
| | 163.8 | 164.8 | 100.6 | | 297.5 | 297.6 | 100.0 |
| 06/11/80 | 99.1 | 96.4 | 97.3 | 01/20/81 | 363.2 | 354.8 | 97.7 |
| | 179.5 | 173.9 | 96.9 | | 207.0 | 200.9 | 97.1 |
| 06/18/80 | 149.7 | 153.6 | 102.6 | 02/17/81 | 194.3 | 205.3 | 105.7 |
| | 121.7 | 128.0 | 105.2 | | 107.8 | 119.9 | 111.2 |
| 06/25/80 | 228.0 | 238.4 | 104.6 | | 107.8 | 120.7 | (b) 112.0 |
| | 145.1 | 151.4 | 104.3 | | 107.8 | 114.8 | (b) 106.5 |
| 07/02/80 | 315.8 | 323.0 | 102.3 | | 370.7 | 370.7 | 100.0 |
| | 210.7 | 220.0 | 104.4 | | 218.7 | 232.7 | 106.4 |
| 07/09/80 | 350.9 | 355.7 | 101.4 | 02/18/81 | 253.6 | 266.5 | 105.1 |
| | 203.3 | 213.4 | 105.0 | | 150.6 | 161.1 | 107.0 |
| 07/16/80 | 298.0 | 279.5 | 93.8 | 02/24/81 | 108.0 | 122.7 | 113.6 |
| | 223.8 | 211.6 | 94.5 | | 108.0 | 113.6 | (b) 105.2 |
| 07/23/80 | 435.4 | 403.1 | 92.6 | 02/25/81 | 108.0 | 116.8 | 108.1 |
| | 295.8 | 275.4 | 93.1 | 03/17/81 | 373.6 | 360.5 | 96.5 |
| 07/30/80 | 185.8 | 198.8 | 107.0 | | 217.8 | 214.3 | 98.4 |
| | 102.5 | 105.6 | 103.0 | | 461.3 | 449.1 | 97.4 |
| 08/13/80 | 149.8 | 168.2 | 112.3 | | 299.5 | 295.9 | 98.8 |
| | 149.8 | 162.1 | (b) 108.2 | 04/28/81 | 192.4 | 189.2 | 98.3 |
| | 120.5 | 137.8 | 114.4 | | 192.4 | 177.9 | (b,c) 92.5 |
| | 120.3 | 132.3 | (b) 110.0 | | 111.1 | 117.0 | 105.4 |
| 08/20/80 | 236.2 | 245.9 | 104.1 | | 111.1 | 110.0 | (b,c) 99.0 |
| | 149.8 | 159.3 | 106.3 | | 254.8 | 246.7 | 96.8 |
| 08/27/80 | 171.9 | 163.2 | 94.9 | | 254.8 | 244.8 | (b,c) 96.1 |
| | 356.5 | 371.5 | 104.2 | | 155.1 | 160.6 | 103.5 |
| 09/03/80 | 206.4 | 216.9 | 105.1 | | 155.1 | 154.6 | (b,c) 99.7 |
| 09/10/80 | 301.9 | 293.0 | 97.1 | 05/26/81 | 458.3 | 475.6 | 103.8 |
| | 235.8 | 253.2 | 107.4 | | 297.9 | 312.2 | 104.8 |
| 09/17/80 | 444.7 | 453.3 | 101.9 | | 364.9 | 383.1 | 105.0 |
| | 286.0 | 300.5 | 105.1 | | 217.5 | 224.4 | 103.2 |
| 09/24/80 | 319.0 | 335.6 | 105.2 | 06/23/81 | 182.3 | 172.3 | 94.5 |
| | 220.9 | 232.0 | 105.0 | | 113.6 | 99.4 | (d) 87.5 |
| 10/01/80 | 187.7 | 227.3 | 121.1 | | 113.6 | 103.3 | (b,d) 90.9 |
| | 187.7 | 193.6 | (b) 103.1 | | 252.1 | 229.9 | 91.2 |
| | 102.9 | 122.1 | 118.7 | | 154.1 | 147.8 | 95.9 |
| | 102.9 | 102.7 | (b) 99.8 | | 361.7 | 339.2 | 93.8 |

TABLE N1. RESULTS OF ANALYSIS OF DOSE MIXTURES IN THE TWO-YEAR GAVAGE STUDIES OF TRICHLOROETHYLENE (Continued)

| Date Mixed | Concentration (mg/ml) | | Percent D/T | Date Mixed | Concentration (mg/ml) | | Percent D/T |
|------------------------------------|-----------------------|--------------------|-------------|------------|-----------------------|--------------------|-------------|
| | Target (T) | Determined (D) (a) | | | Target (T) | Determined (D) (a) | |
| 06/23/81 | 219.4 | 217.0 | 98.9 | | 251.3 | 254.1 | 101.1 |
| | 154.1 | 142.8 | 92.7 | | 251.3 | 246.2 | (b,c)98.0 |
| 07/01/81 | 110.0 | 111.7 | 101.5 | 09/15/81 | 151.3 | 148.9 | 98.4 |
| 07/21/81 | 361.9 | 345.9 | 95.6 | | 151.3 | 144.6 | (b,c)95.6 |
| | 218.8 | 203.8 | 93.1 | 336.4 | 318.2 | 94.6 | |
| 08/18/81 | 458.0 | 446.2 | 97.4 | 219.2 | 205.0 | 93.5 | |
| | 312.4 | 295.8 | 94.7 | 440.4 | 415.6 | 94.4 | |
| | 181.4 | 172.9 | 95.3 | 297.1 | 284.9 | 95.9 | |
| | 181.4 | 173.2 | (b,c)95.5 | 245.2 | 237.6 | 96.9 | |
| | 112.6 | 113.6 | 100.9 | 150.6 | 148.5 | 98.6 | |
| | 112.6 | 112.9 | (b,c)100.3 | | | | |
| Mean (percent of target) | | | | 102.2 | | | |
| Standard deviation | | | | 6.61 | | | |
| Coefficient of variation (percent) | | | | 6.5 | | | |
| Range (percent of target) | | | | 77.0-121.1 | | | |
| Number of samples | | | | 234 | | | |

- (a) Results of duplicate analysis
- (b) Repeat analysis; not included in mean
- (c) Different sample preparation method used
- (d) Not used in the studies

TABLE N2. RESULTS OF REFEREE ANALYSIS OF DOSE MIXTURES IN THE TWO-YEAR GAVAGE STUDIES OF TRICHLOROETHYLENE

| Date Mixed | Target Concentration (mg/ml) | Determined Concentration | |
|------------|------------------------------|--------------------------|------------------------|
| | | Study Laboratory (a) | Referee Laboratory (b) |
| 03/19/80 | 158.8 | 175.1 | 160.4 |
| 03/17/81 | 217.8 | 214.3 | 221.1 |
| 06/23/81 | 154.1 | 147.8 | 150.0 |
| 09/15/81 | 336.4 | 318.2 | 329.0 |

- (a) Results of duplicate analysis
- (b) Results of triplicate analysis

APPENDIX O

SENTINEL ANIMAL PROGRAM

| | PAGE |
|--|------|
| TABLE O1 MURINE VIRUS ANTIBODY DETERMINATIONS FOR RATS IN THE TWO-YEAR GAVAGE STUDIES OF TRICHLOROETHYLENE | 291 |

APPENDIX O. SENTINEL ANIMAL PROGRAM

I. Methods

Rodents used in the Carcinogenesis Program of the National Toxicology Program are produced in optimally clean facilities to eliminate potential pathogens that may affect study results. The Sentinel Animal Program is part of the periodic monitoring of animal health that occurs during the toxicologic evaluation of chemical compounds. Under this program, the disease state of the rodents is monitored via viral serology on sera from extra (sentinel) animals in the study rooms. These animals are untreated, and these animals and the study animals are both subject to identical environmental conditions. The sentinel animals come from the same production source and weanling groups as the animals used for the studies of chemical compounds.

Fifteen rats of each strain and sex are selected at the time of randomization and allocation of the animals to the various study groups. Five animals of each designated sentinel group are killed at 6, 12, and 18 months on study. Data from animals surviving 24 months are collected from 5/50 randomly selected vehicle control animals of each sex and species. The blood from each animal is collected and clotted, and the serum is separated. The serum is cooled on ice and shipped to Microbiological Associates' Comprehensive Animal Diagnostic Service for determination of the viral antibody titers. The following tests are performed:

Hemagglutination Inhibition

PVM (pneumonia virus of mice)
Sendai (August, 18 mo;
Marshall, 24 mo)
KRV (Kilham rat virus)
H-1 (Toolan's H-1 virus)
Reo 3 (reovirus type 3)
(Marshall, 24 mo)
Poly (polyoma virus)
(Marshall, 24 mo)
Ectro (infectious ectromelia)
(Marshall, 24 mo)

Complement Fixation

RCV (rat coronavirus)
Sendai
M. Ad. (mouse adenovirus)
(Marshall, 24 mo)
MHV (mouse
hepatitis virus)
(Marshall, 24 mo)

II. Results

Results are presented in Table O1.

TABLE O1. MURINE VIRUS ANTIBODY DETERMINATIONS FOR RATS IN THE TWO-YEAR GAVAGE STUDIES OF TRICHLOROETHYLENE (a)

| Interval (months) | No. of Animals | Positive Serologic Reaction for |
|-----------------------|--------------------------------|---------------------------------|
| ACI | | |
| 18 | 10/10 1/10 10/10 8/10 | PVM KRV Sendai RCV |
| August | | |
| 6 | 7/10 9/10 | PVM RCV |
| 18 | 7/10 | PVM |
| Marshall | | |
| 18 | 9/9 9/9 9/9 | PVM Sendai RCV |
| 24 | 9/9 8/9 3/9 | PVM Sendai RCV |
| Osborne-Mendel | | |
| 12 | -- | None positive |

(a) Blood samples were taken from sentinel animals at 6, 12, and 18 months after the start of dosing and from the vehicle control animals just before they were killed; samples were sent to Microbiological Associates (Bethesda, MD) for determination of antibody titers.

APPENDIX P

SURVIVAL AND MEAN BODY WEIGHTS OF ACI, AUGUST, AND MARSHALL RATS IN THE THIRTEEN-WEEK GAVAGE STUDIES OF TRICHLOROETHYLENE

| | PAGE |
|---|------|
| TABLE P1 SURVIVAL AND MEAN BODY WEIGHTS OF ACI RATS IN THE THIRTEEN-WEEK GAVAGE STUDIES OF TRICHLOROETHYLENE | 294 |
| TABLE P2 SURVIVAL AND MEAN BODY WEIGHTS OF AUGUST RATS IN THE THIRTEEN-WEEK GAVAGE STUDIES OF TRICHLOROETHYLENE | 294 |
| TABLE P3 SURVIVAL AND MEAN BODY WEIGHTS OF MARSHALL RATS IN THE THIRTEEN-WEEK GAVAGE STUDIES OF TRICHLOROETHYLENE | 295 |

TABLE P1. SURVIVAL AND MEAN BODY WEIGHTS OF ACI RATS IN THE THIRTEEN-WEEK GAVAGE STUDIES OF TRICHLOROETHYLENE

| Dose (mg/kg) | Survival (a) | Mean Body Weights (grams) | | | Final Weight Relative to Vehicle Controls (percent) |
|-----------------|--------------|---------------------------|-------|--------|---|
| | | Initial | Final | Change | |
| MALE | | | | | |
| 0 | 10/10 | 46 | 258 | +212 | -- |
| 125 | 10/10 | 67 | 257 | +190 | 99.6 |
| 250 | 10/10 | 57 | 243 | +186 | 94.2 |
| 500 | 10/10 | 55 | 252 | +197 | 97.7 |
| 1,000 | 10/10 | 52 | 234 | +182 | 90.7 |
| 2,000 | 10/10 | 52 | 213 | +161 | 82.6 |
| FEMALE | | | | | |
| 0 | 10/10 | 45 | 174 | +129 | -- |
| 62.5 | 10/10 | 63 | 173 | +110 | 99.4 |
| 125 | 10/10 | 51 | 165 | +114 | 94.8 |
| 250 | 10/10 | 51 | 172 | +121 | 98.9 |
| 500 | 10/10 | 45 | 160 | +115 | 92.0 |
| 1,000 | 10/10 | 37 | 162 | +125 | 93.1 |

(a) Number surviving/number in group

TABLE P2. SURVIVAL AND MEAN BODY WEIGHTS OF AUGUST RATS IN THE THIRTEEN-WEEK GAVAGE STUDIES OF TRICHLOROETHYLENE

| Dose (mg/kg) | Survival (a) | Mean Body Weights (grams) | | | Final Weight Relative to Vehicle Controls (percent) |
|-----------------|--------------|---------------------------|-------|--------|---|
| | | Initial | Final | Change | |
| MALE | | | | | |
| 0 | 10/10 | 79 | 338 | +259 | -- |
| 125 | 10/10 | 88 | 342 | +254 | 101.2 |
| 250 | 10/10 | 94 | 327 | +233 | 96.7 |
| 500 | 10/10 | 91 | 324 | +233 | 95.9 |
| 1,000 | 10/10 | 94 | 324 | +230 | 95.9 |
| 2,000 | 7/10 | 81 | 286 | +205 | 84.6 |
| FEMALE | | | | | |
| 0 | 10/10 | 78 | 207 | +129 | -- |
| 62.5 | 10/10 | 80 | 210 | +130 | 101.4 |
| 125 | 10/10 | 82 | 208 | +126 | 100.5 |
| 250 | 10/10 | 78 | 208 | +130 | 100.5 |
| 500 | 10/10 | 76 | 204 | +128 | 98.6 |
| 1,000 | 10/10 | 73 | 192 | +119 | 92.8 |

(a) Number surviving/number in group

TABLE P3. SURVIVAL AND MEAN BODY WEIGHTS OF MARSHALL RATS IN THE THIRTEEN-WEEK GAVAGE STUDIES OF TRICHLOROETHYLENE

| Dose (mg/kg) | Survival (a) | Mean Body Weights (grams) | | | Final Weight Relative to Vehicle Controls (percent) |
|-----------------|--------------|---------------------------|-------|--------|---|
| | | Initial | Final | Change | |
| MALE | | | | | |
| 0 | 10/10 | 112 | 247 | +135 | -- |
| 268 | 10/10 | 140 | 255 | +115 | 103.2 |
| 308 | 10/10 | 134 | 259 | +125 | 104.9 |
| 495 | 10/10 | 118 | 249 | +131 | 100.8 |
| 932 | 10/10 | 111 | 232 | +121 | 93.9 |
| 1,834 | 10/10 | 128 | 217 | +89 | 87.9 |
| FEMALE | | | | | |
| 0 | 10/10 | 95 | 171 | +76 | -- |
| 134 | 10/10 | 114 | 173 | +59 | 101.2 |
| 153 | 10/10 | 109 | 176 | +67 | 102.9 |
| 248 | 10/10 | 108 | 175 | +67 | 102.3 |
| 466 | 10/10 | 96 | 166 | +70 | 97.1 |
| 918 | 10/10 | 94 | 164 | +70 | 95.9 |

(a) Number surviving/number in group

APPENDIX Q

AUDIT SUMMARY

APPENDIX Q. AUDIT SUMMARY

The experimental data, documents, and pathology materials for the NTP Technical Report on the 2-year toxicology and carcinogenesis studies of trichloroethylene administered by gavage in corn oil to ACI, August, Marshall, and Osborne-Mendel rats were examined for completeness, consistency, and accuracy. These studies, initiated by the National Cancer Institute under a prime contract to Tracor Jitco, Inc., were conducted at the Papanicolaou Cancer Research Institute, Miami, Florida, between December 1978 and November 1981. The experiments were started before the NTP required compliance with Good Laboratory Practice (GLP) regulations of the Food and Drug Administration (implemented by the NTP October 1, 1981). The following three contractors conducted the audits:

(1) Immuquest Laboratories, Inc., Rockville, Maryland: Toxicology and pathology data for studies in *ACI and August rats* (October through November 1983) and chemistry data for all studies (November 1983). Personnel conducting the audits were Pamela Errico, M.A.; Caroline Reese; Karen Witkin, Ph.D.; and Ronald Schueler, D.V.M.

(2) Argus Research Laboratories, Inc., Horsham, Pennsylvania: Toxicology and pathology data for studies in *Marshall rats* (March 1984). Personnel conducting the audit were Jane Goeke, Ph.D., and F. Garner, D.V.M.

(3) Clement Associates, Inc., Washington, DC: Pathology data for studies conducted in *Osborne-Mendel rats* (March 1984). The audit was conducted by Miriam Anver, D.V.M., Ph.D. The toxicology portion of the studies in Osborne-Mendel rats was not audited.

The audit reports are on file at the National Institute of Environmental Health Sciences, Research Triangle Park, North Carolina. The audits included a review of:

- (1) Inlife toxicology data for 5%-10% of the study animals.
- (2) All records concerning dosing, clinical signs, and mortality of study animals.
- (3) All Individual Animal Data Records to examine for correspondence between necropsy observations and histologic findings.
- (4) Wet tissues from a random 10% sample of the study animals to verify animal identification and to examine for untrimmed masses.
- (5) All slides and blocks of tissue from both sexes of all strains in the high dose and vehicle control groups to examine for proper match and inventory.
- (6) All available analytical chemistry and dose mixture data.

The audit of the inlife phase of these studies, which were conducted before implementation of GLP requirements, identified a number of omissions in study documentation. These included: (1) absence of complete study protocol, (2) absence of animal breeding and husbandry records, and (3) absence of environmental condition records. The clinical observation records showed inconsistent recording of palpable masses and other clinical signs, missing or incorrectly recorded body weights, and occasional discrepancies between mortality and clinical records which, taken collectively, leave open the possibility that some animals may have been misidentified during the studies. The records also documented clinical observation of toxicity associated with administration of doses. In addition, there were incidents recorded where malfunctions in automated watering systems resulted in flooding of animal cages.

The contract governing the conduct of these studies did not require the retention of animal carcasses containing individual animal identifying markers; however, examination of the residual, formalin-fixed tissues (wet tissues) revealed that ears but not feet for most animals had been preserved. Ears contained the first two digits of the three- and four-digit identification numbers used. Review of the wet tissues for 240 rats showed that 237 of them had an ear punch that matched the bag label. The slide/block audit showed good match up, but sections taken from a number of blocks appeared to

APPENDIX Q. AUDIT SUMMARY

be incomplete (i.e., the blocks did not always have corresponding full-face sections). Instances of gross observations without corresponding microscopic diagnoses and untrimmed potential lesions in wet tissues which were detected by the audit were reviewed by NTP staff and were either resolved satisfactorily or believed to have no impact on interpretation of the study results.

Original records documenting the chemical analyses performed by the study laboratory were not available to fully validate the chemistry portion of the studies. Those records that were available indicated that samples taken from seven chemical/vehicle preparations were shipped to Midwest Research Institute (MRI) for analysis. The analyses performed by MRI were well documented and showed good agreement with those reported by the study laboratory. The bulk chemical purity was documented by the MRI records. Audit of the study laboratory's chemical use log indicated that dose preparation calculations were valid and that sufficient chemical was available for the conduct of the studies. Comparison of the chemical use log with the dose preparation log showed that the amount of chemical used matched the amount checked out.

These studies were not required to be conducted under GLP standards. Nevertheless, retrospective audit of the available documents and materials shows that recordkeeping was less than adequate to fully document all of the procedures followed and the original data generated. The Tracor Jitco Statement of Work, applicable under the terms of the contract governing this study, is presumed to have been followed in lieu of a formal study protocol and system for documenting procedures actually followed. The most significant audit findings from review of the existing study records and data involved inconsistent recording of inlife records of clinical observations and palpable masses and the absence of complete carcass identification information in wet tissues. Thus, the possibility that some animals may have been mixed up during the conduct of the studies cannot be excluded. Because of the characteristic kidney lesions produced by trichloroethylene, it is unlikely that dosed and vehicle control animals were confused. It is recognized, however, that total confidence cannot be placed in the differentiation of all high dose from low dose animals. Incidents of misdosing (animals dosed twice in 1 day or given incorrect doses) were reported. Tissue accountability for the kidney was generally adequate. Despite incomplete chemistry records, the original data that are available indicate that animals were dosed with trichloroethylene and that dose mixtures were prepared properly.

The NIEHS/NTP concludes that the incomplete documentation for these studies reveals certain gaps and inconsistencies that have been considered in the interpretations given in the Technical Report.

NATIONAL TOXICOLOGY PROGRAM TECHNICAL REPORTS
PUBLISHED AS OF JANUARY 1988

| TR No. | CHEMICAL | TR No. | CHEMICAL |
|--------|---|--------|---|
| 201 | 2,3,7,8-Tetrachlorodibenzo- <i>p</i> -dioxin (Dermal) | 263 | 1,2-Dichloropropane |
| 206 | Dibromochloropropane | 267 | Propylene Oxide |
| 207 | Cytembena | 269 | Telone II® |
| 208 | FD & C Yellow No. 6 | 271 | HC Blue No. 1 |
| 209 | 2,3,7,8-Tetrachlorodibenzo- <i>p</i> -dioxin (Gavage) | 272 | Propylene |
| 210 | 1,2-Dibromoethane (Inhalation) | 274 | Tris(2-ethylhexyl)phosphate |
| 211 | C.I. Acid Orange 10 | 275 | 2-Chloroethanol |
| 212 | Di(2-ethylhexyl)adipate | 276 | 8-Hydroxyquinoline |
| 213 | Butylbenzyl Phthalate | 281 | H.C. Red No. 3 |
| 214 | Caprolactam | 282 | Chlorodibromomethane |
| 215 | Bisphenol A | 284 | Diallylphthalate (Rats) |
| 216 | 11-Aminoundecanoic Acid | 285 | C.I. Basic Red 9 Monohydrochloride |
| 217 | Di(2-ethylhexyl)phthalate | 287 | Dimethyl Hydrogen Phosphite |
| 219 | 2,6-Dichloro- <i>p</i> -phenylenediamine | 288 | 1,3-Butadiene |
| 220 | C.I. Acid Red 14 | 289 | Benzene |
| 221 | Locust Bean Gum | 291 | Isophorone |
| 222 | C.I. Disperse Yellow 3 | 293 | HC Blue No. 2 |
| 223 | Eugenol | 294 | Chlorinated Trisodium Phosphate |
| 224 | Tara Gum | 295 | Chrysotile Asbestos (Rats) |
| 225 | D & C Red No. 9 | 296 | Tetrakis(hydroxymethyl)phosphonium Sulfate and Tetrakis(hydroxymethyl)phosphonium Chloride |
| 226 | C.I. Solvent Yellow 14 | 298 | Dimethyl Morpholinophosphoramidate |
| 227 | Gum Arabic | 299 | C.I. Disperse Blue 1 |
| 228 | Vinylidene Chloride | 300 | 3-Chloro-2-methylpropene |
| 229 | Guar Gum | 301 | <i>o</i> -Phenylphenol |
| 230 | Agar | 303 | 4-Vinylcyclohexene |
| 231 | Stannous Chloride | 304 | Chlorendic Acid |
| 232 | Pentachloroethane | 305 | Chlorinated Paraffins (C ₂₃ , 43% chlorine) |
| 233 | 2-Biphenylamine Hydrochloride | 306 | Dichloromethane |
| 234 | Allyl Isothiocyanate | 307 | Ephedrine Sulfate |
| 235 | Zearalenone | 308 | Chlorinated Paraffins (C ₁₂ , 60% chlorine) |
| 236 | D-Mannitol | 309 | Decabromodiphenyl Oxide |
| 237 | 1,1,1,2-Tetrachloroethane | 310 | Marine Diesel Fuel and JP-5 Navy Fuel |
| 238 | Ziram | 311 | Tetrachloroethylene (Inhalation) |
| 239 | Bis(2-chloro-1-methylethyl)ether | 312 | <i>n</i> -Butyl Chloride |
| 240 | Propyl Gallate | 314 | Methyl Methacrylate |
| 242 | Diallyl Phthalate (Mice) | 315 | Oxytetracycline Hydrochloride |
| 244 | Polybrominated Biphenyl Mixture | 316 | 1-Chloro-2-methylpropene |
| 245 | Melamine | 317 | Chlorpheniramine Maleate |
| 247 | L-Ascorbic Acid | 318 | Ampicillin Trihydrate |
| 248 | 4,4'-Methylenedianiline Dihydrochloride | 319 | 1,4-Dichlorobenzene |
| 249 | Amosite Asbestos | 321 | Bromodichloromethane |
| 250 | Benzyl Acetate | 322 | Phenylephrine Hydrochloride |
| 251 | Toluene Diisocyanate | 323 | Dimethyl Methylphosphonate |
| 252 | Geranyl Acetate | 324 | Boric Acid |
| 253 | Allyl Isovalerate | 325 | Pentachloronitrobenzene |
| 255 | 1,2-Dichlorobenzene | 326 | Ethylene Oxide |
| 257 | Diglycidyl Resorcinol Ether | 327 | Xylenes (Mixed) |
| 259 | Ethyl Acrylate | 328 | Methyl Carbamate |
| 261 | Chlorobenzene | | |

These NTP Technical Reports are available for sale from the National Technical Information Service, U.S. Department of Commerce, 5285 Port Royal Road, Springfield, VA 22161 (703-487-4650). Single copies of this Technical Report are available without charge (and while supplies last) from the NTP Public Information Office, National Toxicology Program, P.O. Box 12233, Research Triangle Park, NC 27709.