

**Appendix Table I. Number and percent of persons uninsured, with public or private coverage, based on the 1990 and 2000 census population estimates, by age group: United States, 2000-2002**

Health insurance coverage and age group	2000		2001		2002	
	1990-based <sup>1</sup>	2000-based <sup>2</sup>	1990-based <sup>1</sup>	2000-based <sup>2</sup>	1990-based <sup>1</sup>	2000-based <sup>2</sup>
<b>Number of uninsured in millions</b>						
All ages	40.4	41.3	39.1	40.2	40.4	41.5
Under 65 years	39.9	40.8	38.7	39.8	40.0	41.1
18-64 years	31.2	32.0	30.9	31.9	32.4	33.5
Under 18 years	8.8	8.9	7.8	7.9	7.6	7.6
<b>Percent (95% confidence interval)</b>						
<b>Uninsured</b>						
All ages	14.7 (14.3-15.1)	14.9 (14.5-15.3)	14.1 (13.7-14.6)	14.3 (13.8-14.8)	14.5 (14.1-14.9)	14.7 (14.3-15.1)
Under 65 years						
Crude	16.6 (16.1-17.1)	16.8 (16.3-17.2)	15.9 (15.4-16.4)	16.2 (15.7-16.7)	16.3 (15.8-16.7)	16.5 (16.0-16.9)
Age-adjusted	16.6 (16.1-17.1)	16.8 (16.3-17.3)	16.0 (15.5-16.5)	16.2 (15.7-16.7)	16.4 (15.9-16.9)	16.6 (16.1-17.1)
18-64 years	18.4 (17.9-18.9)	18.7 (18.1-19.2)	18.1 (17.5-18.6)	18.3 (17.8-18.8)	18.8 (18.3-19.3)	19.1 (18.6-19.6)
Under 18 years	12.2 (11.6-12.8)	12.3 (11.7-12.9)	10.8 (10.1-11.5)	11.0 (10.3-11.7)	10.4 (9.8-11.0)	10.5 (9.9-11.1)
<b>Public</b>						
Under 65 years	12.9 (12.4-13.4)	12.9 (12.4-13.4)	13.6 (13.1-14.1)	13.6 (13.1-14.1)	15.2 (14.6-15.8)	15.2 (14.6-15.8)
18-64 years	9.0 (8.7-9.4)	9.1 (8.7-9.4)	9.4 (9.0-9.8)	9.4 (9.0-9.8)	10.2 (9.8-10.7)	10.3 (9.9-10.7)
Under 18 years	21.8 (20.8-22.8)	22.0 (21.0-23.0)	23.4 (22.4-24.3)	23.6 (22.6-24.5)	26.8 (25.7-27.9)	27.1 (26.0-28.2)
<b>Private</b>						
Under 65 years	72.0 (71.4-72.7)	71.8 (71.1-72.5)	71.9 (71.2-72.6)	71.6 (70.9-72.3)	70.1 (69.3-70.9)	69.8 (69.0-70.6)
18-64 years	74.0 (73.4-74.6)	73.8 (73.2-74.4)	73.9 (73.3-74.6)	73.7 (73.1-74.4)	72.6 (71.9-73.2)	72.3 (71.6-72.9)
Under 18 years	67.5 (66.4-68.5)	67.1 (66.1-68.2)	67.1 (66.0-68.2)	66.7 (66.4-68.6)	64.3 (63.1-65.5)	63.9 (62.7-65.1)

<sup>1</sup>The estimates are calculated based on data with sample weights derived from the 1990 census.

<sup>2</sup>The estimates are calculated based on data with sample weights derived from the 2000 census.

**Appendix Table II. Percent of persons under age 65 years without health insurance coverage, with public or private coverage, based on the 1990 and 2000 census population estimates, by age and poverty status: United States, 2000-2002**

Health insurance coverage, age group, and poverty status	2000		2001		2002	
	1990-based <sup>1</sup>	2000-based <sup>2</sup>	1990-based <sup>1</sup>	2000-based <sup>2</sup>	1990-based <sup>1</sup>	2000-based <sup>2</sup>
<b>Uninsured</b>						
<b>Percent (standard error) uninsured</b>						
<b>Under age 18 years</b>						
Total	12.2 (0.32)	12.3 (0.32)	10.8 (0.34)	11.0 (0.34)	10.4 (0.32)	10.5 (0.32)
Poor	20.5 (1.04)	20.6 (1.04)	18.7 (1.26)	18.8 (1.24)	15.8 (0.98)	15.9 (0.97)
Near poor	21.2 (0.93)	21.4 (0.93)	16.8 (0.85)	17.0 (0.85)	15.6 (0.84)	15.7 (0.84)
Not poor	5.3 (0.30)	5.3 (0.30)	4.4 (0.26)	4.4 (0.26)	5.3 (0.36)	5.3 (0.36)
Unknown	14.8 (0.71)	15.0 (0.72)	15.3 (0.84)	15.5 (0.84)	14.0 (0.76)	14.1 (0.76)
<b>18-64 years</b>						
Total	18.4 (0.26)	18.7 (0.27)	18.1 (0.26)	18.3 (0.27)	18.8 (0.26)	19.1 (0.26)
Poor	40.4 (1.04)	41.1 (1.05)	38.8 (1.19)	39.5 (1.19)	36.4 (1.09)	37.0 (1.09)
Near poor	37.0 (0.77)	37.4 (0.77)	35.1 (0.78)	35.6 (0.78)	35.6 (0.76)	36.2 (0.77)
Not poor	9.9 (0.24)	10.0 (0.24)	9.8 (0.22)	9.9 (0.22)	10.9 (0.25)	11.0 (0.25)
Unknown	21.3 (0.52)	21.5 (0.53)	21.7 (0.51)	22.1 (0.52)	22.9 (0.56)	23.2 (0.56)
<b>Public</b>						
<b>Percent (standard error) with public coverage</b>						
<b>Under age 18 years</b>						
Total	21.8 (0.50)	22.0 (0.50)	23.3 (0.50)	23.6 (0.50)	26.8 (0.54)	27.1 (0.54)
Poor	61.8 (1.49)	61.8 (1.48)	65.2 (1.48)	65.2 (1.47)	69.0 (1.35)	69.0 (1.33)
Near poor	32.4 (1.15)	32.4 (1.13)	36.9 (1.24)	37.0 (1.23)	42.1 (1.19)	42.2 (1.18)
Not poor	7.3 (0.40)	7.4 (0.39)	8.1 (0.39)	8.1 (0.39)	8.8 (0.44)	8.9 (0.45)
Unknown	21.9 (0.85)	22.1 (0.85)	22.8 (0.94)	23.1 (0.94)	30.3 (0.99)	30.7 (0.99)
<b>18-64 years</b>						
Total	9.0 (0.19)	9.1 (0.19)	9.4 (0.21)	9.4 (0.21)	10.2 (0.23)	10.3 (0.23)
Poor	31.3 (1.01)	31.1 (1.00)	31.0 (1.11)	30.8 (1.10)	32.6 (1.11)	32.5 (1.10)
Near poor	15.2 (0.54)	15.2 (0.54)	17.9 (0.62)	17.8 (0.62)	18.4 (0.67)	18.3 (0.66)
Not poor	4.5 (0.19)	4.5 (0.19)	4.7 (0.20)	4.8 (0.20)	5.1 (0.22)	5.1 (0.22)
Unknown	9.2 (0.33)	9.1 (0.33)	9.4 (0.33)	9.4 (0.33)	11.2 (0.35)	11.2 (0.35)
<b>Private</b>						
<b>Percent (standard error) with private coverage</b>						
<b>Under age 18 years</b>						
Total	67.5 (0.53)	67.1 (0.53)	67.1 (0.57)	66.7 (0.57)	64.3 (0.61)	63.9 (0.61)
Poor	19.7 (1.23)	19.5 (1.21)	18.3 (1.14)	18.1 (1.12)	17.3 (1.09)	17.2 (1.08)
Near poor	49.0 (1.24)	48.8 (1.23)	48.6 (1.24)	48.4 (1.23)	45.2 (1.31)	44.9 (1.29)
Not poor	88.4 (0.47)	88.4 (0.47)	88.5 (0.40)	88.4 (0.40)	87.0 (0.54)	86.9 (0.54)
Unknown	64.6 (0.99)	64.2 (0.99)	62.7 (1.16)	62.2 (1.16)	56.7 (1.20)	56.3 (1.19)
<b>18-64 years</b>						
Total	74.0 (0.31)	73.8 (0.32)	73.9 (0.33)	73.7 (0.33)	72.6 (0.35)	72.3 (0.35)
Poor	29.7 (1.19)	29.2 (1.16)	31.3 (1.45)	31.7 (1.41)	32.3 (1.54)	31.8 (1.50)
Near poor	49.5 (0.83)	49.3 (0.83)	48.7 (0.83)	48.4 (0.82)	48.0 (0.85)	47.5 (0.85)
Not poor	87.1 (0.27)	87.1 (0.27)	86.9 (0.28)	86.8 (0.28)	85.8 (0.30)	85.7 (0.30)
Unknown	70.9 (0.61)	70.6 (0.61)	70.2 (0.61)	69.9 (0.61)	67.2 (0.62)	66.9 (0.62)

<sup>1</sup>The estimates are calculated based on data with sample weights derived from the 1990 census.

<sup>2</sup>The estimates are calculated based on data with sample weights derived from the 2000 census.

**Appendix Table III. Estimates of selected measures based on the 1990 and 2000 census population estimates: United States, 2000-2002**

Indicator	2000		2001		2002	
	1990 based <sup>1</sup>	2000 based <sup>2</sup>	1990 based <sup>1</sup>	2000 based <sup>2</sup>	1990 based <sup>1</sup>	2000 based <sup>2</sup>
<b>Percent (95% confidence interval)</b>						
With a usual place to go for medical care	87.1 (86.7-87.5)	87.0 (86.5-87.4)	88.2 (87.7-88.6)	88.0 (87.6-88.5)	87.8 (87.3-88.2)	87.6 (87.1-88.0)
Failed to obtain needed care due to cost	4.5 (4.3-4.7)	4.5 (4.3-4.7)	4.7 (4.5-4.9)	4.7 (4.5-4.9)	4.7 (4.5-4.9)	4.7 (4.5-4.9)
Had influenza vaccination in the past year	64.3 (62.9-65.7)	64.4 (63.0-65.9)	63.0 (61.6-64.4)	63.1 (61.7-64.5)	65.6 (64.1-67.0)	65.7 (64.3-67.2)
Had pneumococcal vaccination in the past year	52.9 (51.3-54.5)	53.1 (51.5-54.7)	53.8 (52.2-55.4)	54.0 (52.4-55.6)	55.7 (54.2-57.2)	56.0 (54.5-57.5)
With obesity	21.8 (21.2-22.4)	21.8 (21.2-22.4)	23.0 (22.4-23.5)	23.0 (22.4-23.6)	23.9 (23.3-24.6)	23.9 (23.3-24.6)
Engaged in regular leisure-time physical activity	32.0 (31.3-32.7)	31.9 (31.2-32.6)	32.1 (31.3-32.8)	32.1 (31.3-32.8)	32.0 (31.2-32.8)	31.9 (31.1-32.7)
Current smoking	23.3 (22.7-23.9)	23.2 (22.5-23.8)	22.8 (22.2-23.4)	22.7 (22.1-23.3)	22.5 (21.9-23.1)	22.4 (21.7-23.0)
Had 5 or more drinks in one day at least once in the past year	19.3 (18.7-20.0)	19.2 (18.6-19.9)	20.1 (19.5-20.7)	20.0 (19.4-20.6)	19.9 (19.3-20.6)	19.9 (19.2-20.5)
Ever had HIV test	32.2 (31.5-32.9)	32.1 (31.4-32.8)	33.3 (32.6-34.0)	33.2 (32.5-34.0)	34.5 (33.8-35.3)	34.5 (33.8-35.2)
With excellent or very good health status	68.4 (67.9-69.0)	68.3 (67.7-68.8)	68.6 (68.1-69.2)	68.4 (67.9-69.0)	67.6 (67.0-68.2)	67.4 (66.8-68.0)
With need for personal care	6.3 (5.9-6.8)	6.4 (5.9-6.9)	6.5 (6.0-7.0)	6.5 (6.0-7.0)	6.2 (5.6-6.7)	6.2 (5.7-6.7)
Serious psychological distress	2.7 (2.5-2.9)	2.7 (2.5-2.9)	3.2 (2.9-3.4)	3.2 (2.9-3.4)	3.0 (2.8-3.3)	3.0 (2.8-3.2)
Diagnosed Diabetes	5.9 (5.6-6.2)	5.9 (5.6-6.2)	6.4 (6.1-6.7)	6.4 (6.1-6.7)	6.5 (6.2-6.8)	6.5 (6.2-6.8)
Asthma episodes in the past year	40.1 (37.9-42.4)	40.0 (37.7-42.2)	43.4 (41.0-45.8)	43.2 (40.8-45.5)	42.8 (40.4-45.2)	42.6 (40.2-44.9)

<sup>1</sup>The estimates are calculated based on data with sample weights derived from the 1990 census.

<sup>2</sup>The estimates are calculated based on data with sample weights derived from the 2000 census.