


**Figure 14.1. Prevalence of diagnosed diabetes among adults aged 18 years and over: United States, 1997–2005**




NOTES: Prevalence of diagnosed diabetes is based on self-report of ever having been diagnosed with diabetes by a doctor or other health professional. Persons reporting “borderline” diabetes status and women reporting diabetes only during pregnancy were not coded as having diabetes in the analyses. The analyses excluded persons with unknown diabetes status (about 0.1% of respondents each year). Beginning with the 2003 data, the National Health Interview Survey (NHIS) transitioned to weights derived from the 2000 census. In this Early Release, estimates for 2000–2002 were recalculated using weights derived from the 2000 census. See “About This Release” and Table III in the Appendix for more details.

DATA SOURCE: Sample Adult Core component of the 1997–2005 NHIS. Data are based on household interviews of a sample of the civilian noninstitutionalized population.

■ In 2005, 7.4% (95% confidence interval = 7.10–7.78%) of U.S. adults aged 18 years and over had ever been diagnosed as having diabetes, which was higher than, but not significantly different from, the 2004 estimate of 7.0%.

■ From 1997 to 2005, there was an increasing trend in the prevalence of diagnosed diabetes among U.S. adults aged 18 years and over, from 5.1% in 1997 to 7.4% in 2005.

**Figure 14.2. Prevalence of diagnosed diabetes among adults aged 18 years and over, by age group and sex: United States, 2005**




NOTES: Prevalence of diagnosed diabetes is based on self-report of ever having been diagnosed with diabetes by a doctor or other health professional. Persons reporting "borderline" diabetes status and women reporting diabetes only during pregnancy were not coded as having diabetes in the analyses. The analyses excluded 24 persons (0.1%) with unknown diabetes status.

DATA SOURCE: Sample Adult Core component of the 2005 National Health Interview Survey. Data are based on household interviews of a sample of the civilian noninstitutionalized population.

■ For both sexes combined, the prevalence of diagnosed diabetes increased with age, with the highest rate among adults aged 65 years and over (17.0%) and the lowest rate among adults aged 18-44 years (2.4%).

■ For the age groups 45-54 years and 65 years and over, the prevalence of diagnosed diabetes was higher among men than women.

**Figure 14.3. Age-sex-adjusted prevalence of diagnosed diabetes among adults aged 18 years and over, by race/ethnicity: United States, 2005**



NOTES: Prevalence of diagnosed diabetes is based on self-report of ever having been diagnosed with diabetes by a doctor or other health professional. Persons reporting "borderline" diabetes status and women reporting diabetes only during pregnancy were not coded as having diabetes in the analyses. The analyses excluded 24 persons (0.1%) with unknown diabetes status. Estimates are age-sex adjusted to the 2000 projected U.S. standard population using four age groups: 18–44 years, 45–54 years, 55–64 years, and 65 years and over.

DATA SOURCE: Sample Adult Core component of the 2005 National Health Interview Survey. Data are based on household interviews of a sample of the civilian noninstitutionalized population.

■ The age-sex-adjusted prevalence of diagnosed diabetes was 9.8% for Hispanic persons, 6.5% for non-Hispanic white persons, and 11.3% for non-Hispanic black persons.

■ The prevalence of diagnosed diabetes was higher among non-Hispanic black persons and Hispanic persons than among non-Hispanic white persons.

## Data tables for figures 14.1–14.3:

### Data table for figure 14.1. Prevalence of diagnosed diabetes among adults aged 18 years and over: United States, 1997–2005

Year	Percent (95% confidence interval)	
	Crude <sup>1</sup>	Age-adjusted <sup>2</sup>
1997	5.1 (4.9-5.4)	5.3 (5.1-5.6)
1998	5.3 (5.0-5.6)	5.4 (5.1-5.7)
1999	5.4 (5.1-5.7)	5.5 (5.2-5.8)
2000	5.9 (5.6-6.2)	6.0 (5.7-6.3)
2001	6.4 (6.1-6.7)	6.4 (6.1-6.7)
2002	6.5 (6.2-6.8)	6.5 (6.2-6.8)
2003	6.6 (6.3-6.9)	6.5 (6.2-6.9)
2004	7.0 (6.7-7.4)	6.9 (6.6-7.3)
2005	7.4 (7.10-7.78)	7.3 (6.95-7.57)

<sup>1</sup>Crude estimates are presented in the figure.

<sup>2</sup>Estimates are age adjusted to the 2000 projected U.S. standard population using four age groups: 18–44 years, 45–54 years, 55–64 years, and 65 years and over.

NOTES: Beginning with the 2003 data, the National Health Interview Survey (NHIS) transitioned to weights derived from the 2000 census. In this Early Release, estimates for 2000–2002 were recalculated using weights derived from the 2000 census. See “About This Release” and Table III in the Appendix for more details.

DATA SOURCE: NHIS, 1997–2005. Data are based on household interviews of a sample of the civilian noninstitutionalized population.

**Data table for figure 14.2. Prevalence of diagnosed diabetes among adults aged 18 years and over, by age group and sex: United States, 2005**

Age and sex	Percent	95% confidence interval
<b>18–44 years</b>		
Total	2.4	2.08-2.70
Male	2.3	1.84-2.72
Female	2.5	2.05-2.95
<b>45–54 years</b>		
Total	7.7	6.89-8.42
Male	8.5	7.26-9.66
Female	6.9	5.95-7.82
<b>55–64 years</b>		
Total	14.5	13.31-15.68
Male	14.9	13.10-16.74
Female	14.1	12.49-15.72
<b>65 years and over</b>		
Total	17.0	15.96-18.07
Male	18.4	16.85-19.93
Female	16.0	14.51-17.47
<b>18 years and over: crude<sup>1</sup></b>		
Total	7.4	7.10-7.78
Male	7.5	7.06-8.00
Female	7.4	6.88-7.82
<b>18 years and over: age-adjusted<sup>2</sup></b>		
Total	7.3	6.95-7.57
Male	7.6	7.18-8.08
Female	7.0	6.52-7.39

<sup>1</sup>Crude estimates are presented in the figure.

<sup>2</sup>Estimates are age adjusted to the 2000 projected U.S. standard population using four age groups: 18–44 years, 45–54 years, 55–64 years, and 65 years and over.

DATA SOURCE: National Health Interview Survey, 2005. Data are based on household interviews of a sample of the civilian noninstitutionalized population.

**Data table for figure 14.3. Age-sex-adjusted prevalence of diagnosed diabetes among adults aged 18 years and over, by race/ethnicity: United States, 2005**

Race/ethnicity	Percent <sup>1</sup>	95% confidence interval
<b>Hispanic or Latino</b>	9.8	8.71-10.89
<b>Not Hispanic or Latino:</b>		
<b>White, single race</b>	6.5	6.17-6.88
<b>Black, single race</b>	11.3	10.31-12.28

<sup>1</sup>Estimates are age-sex adjusted to the 2000 projected U.S. standard population using four age groups: 18–44 years, 45–54 years, 55–64 years, and 65 years and over.

DATA SOURCE: National Health Interview Survey, 2005. Data are based on household interviews of a sample of the civilian noninstitutionalized population.