

Prepared in cooperation with
the Bureau of Reclamation

Glen Canyon Dam Adaptive Management Program Budget and Annual Work Plan— Fiscal Year 2008

By the Grand Canyon
Monitoring and
Research Center

Planning Document



Cover: Don Becker, U.S. Geological Survey; Paul Alley, U.S. Geological Survey; Arizona Game and Fish Department; Amy Draut, U.S. Geological Survey.



Prepared in cooperation with the Bureau of Reclamation

Glen Canyon Dam Adaptive Management Program Budget and Annual Work Plan— Fiscal Year 2008

By Grand Canyon Monitoring and Research Center

Final Planning Document—April 13, 2008

**U.S. Department of the Interior
U.S. Geological Survey**

Contents

Contents	iii
Chapter 1. Bureau of Reclamation, Upper Colorado Regional Office, Glen Canyon Dam Adaptive Management Program Annual Budget and Work Plan—Fiscal Year 2008	1
Introduction	1
PROJECT TITLE AND ID: A.1. Personnel Costs	2
PROJECT TITLE AND ID: A.2. AMWG Member Travel Reimbursement	3
PROJECT TITLE AND ID: A.3. Reclamation Travel.....	4
PROJECT TITLE AND ID: A.4. Facilitation Contract	5
PROJECT TITLE AND ID: A.5. Public Outreach.....	6
PROJECT TITLE AND ID: A.6. Other.....	7
PROJECT TITLE AND ID: B.1. Personnel Costs	8
PROJECT TITLE AND ID: B.2. TWG Member Travel Reimbursement.....	9
PROJECT TITLE AND ID: B.3. Reclamation Travel.....	10
PROJECT TITLE AND ID: B.4. TWG Chair Reimbursement.....	11
PROJECT TITLE AND ID: B.5. Other.....	12
PROJECT TITLE AND ID: C.1. Compliance Documents	13
PROJECT TITLE AND ID: C.2. Administrative Support for NPS Permitting.....	14
PROJECT TITLE AND ID: C.3. Contract Administration	15
PROJECT TITLE AND ID: C.4. Experimental Carryover Funds.....	16
PROJECT TITLE AND ID: C.5. Integrated Tribal Resources Monitoring	17
PROJECT TITLE AND ID: D.1. Programmatic Agreement, Reclamation Administrative Costs.....	18
PROJECT TITLE AND ID: D.2. NPS Support for Archaeological Site Assessment	19
PROJECT TITLE AND ID: D.4. Glen and Grand Canyon Treatment Plan Implementation.....	20
PROJECT TITLE AND ID: E. Tribal Consultation—Sole-Source Contracts with Tribes	22
Chapter 2. U.S. Geological Survey, Southwest Biological Science Center, Grand Canyon Monitoring and Research Center Annual Budget and Work Plan—Fiscal Year 2008.....	23
Introduction	23
Project Descriptions.....	37
GCDAMP Goal 1: Protect or improve the aquatic food base so that it will support viable populations of desired species at higher trophic levels.	38
BIO 1.R1.08: Aquatic Food Base	38
BIO 1.R3.08: Diet, Drift, and Predation Data Analysis.....	43
BIO 1.R4.08: Impacts of Various Flow Regimes on the Aquatic Food Base	47
GCDAMP Goal 2: Maintain or attain viable populations of existing native fish, remove jeopardy from humpback chub and razorback sucker, and prevent adverse modification to their critical habitat.	52
BIO 2.R1.08: Little Colorado River Humpback Chub Monitoring Lower 15 km	52
BIO 2.R2.08: Little Colorado River Humpback Chub Monitoring Lower 1,200 m	52
BIO 2.R3.08: Humpback Chub Monitoring Above Chute Falls.....	52
BIO 2.R4.08: Monitoring Mainstem Fishes	59
BIO 2.R5.08: Nonnative Control Planning	62

BIO 2.R6.08: Nonnative Control Pilot Testing	62
BIO 2.R7.08: Stock Assessment of Native Fish in Grand Canyon.....	66
BIO 2.R8.08: Abundance Estimation Procedures	69
BIO 2.R9.08: Investigate Factors Affecting the Survival Rate of Juvenile Native Fishes in the Mainstem Colorado River	71
BIO 2.R11.08: Native Fishes Habitat Data Analysis.....	74
BIO 2.R12.08: Trammel Net Effects	77
BIO 2.R13.08: Remote PIT Tag Reading.....	77
BIO 2.R14.08: Test Sonic Tags	77
GCDAMP Goal 4: Maintain a naturally reproducing population of rainbow trout above the Paria River, to the extent practicable and consistent with the maintenance of viable populations of native fish.	81
BIO.4.M1.08: Status and Trends of Lees Ferry Trout.....	81
BIO 4.E1.08: Monitoring Rainbow Trout Redds and Larvae	84
GCDAMP Goal 5: Maintain or attain viable populations of Kanab ambersnail.	88
BIO 5.R1.08: Monitor Kanab ambersnail	88
GCDAMP Goal 6: Protect or improve the biotic riparian and spring communities, including threatened and endangered species and their critical habitat.	91
BIO 6.R1.08: Vegetation Mapping	91
BIO 6.R2.08: Vegetation Transects	91
BIO 6.R3.08: Vegetation Synthesis.....	96
GCDAMP Goal 7: Establish water temperature, quality, and flow dynamics to achieve the Adaptive Management Program ecosystem goals.	100
BIO 7.R1.08: Water Quality Monitoring of Lake Powell and the Glen Canyon Dam Tailwater.....	100
PHY 7.M1.08: Core Monitoring of Downstream Integrated Quality of Water	105
PHY 7.R1.08: Modeling Support Linked with Downstream Integrated Quality-of-Water Monitoring Project	111
GCDAMP Goal 8: Maintain or attain levels of sediment storage within the main channel and along shorelines to achieve the Adaptive Management Program ecosystem goals.	115
PHY 8.M1.08: Core Monitoring for the Sediment Budget and Sandbar Status throughout the CRE Utilizing Direct Topographic/Bathymetric Measurements and Remote Sensing	115
GCDAMP Goal 9: Maintain or improve the quality of recreational experiences for users of the Colorado River ecosystem, within the framework of GCDAMP ecosystem goals.....	123
REC 9.R1.08/PHYS 8.M2.08: Sandbar and Campable Area Monitoring	123
REC 9.R3.08: Compile Campsite Inventory and GIS Atlas	128
GCDAMP Goal 10: Maintain power production capacity and energy generation, and increase where feasible and advisable, within the framework of the Adaptive Management ecosystem goals.....	131
HYD 10.M1.08: Monitor Power Generation and Market Values under Current and Future Dam Operations	131
GCDAMP Goal 11: Preserve, protect, manage, and treat cultural resources for the inspiration and benefit of past, present, and future generations.	135
CUL 11.R1.08: Research and Development towards Core Monitoring, Phase II	135
CUL 11.R2.08: Implement Tribal Monitoring Projects	141
GCDAMP Goal 12: Maintain a high-quality monitoring, research, and adaptive management program...	144
DASA 12.D1.08: Preparation for Monitoring Data Acquisition	144
DASA 12.D2.08: Grand Canyon Integrated Oracle Database Management System.....	148

DASA 12.D3.08: Library Operations	151
DASA 12.D4.08: Legacy Analog Data Conversion.....	153
DASA 12.D5.08: GIS General Support for Integrated Analyses and Projects, GIS Lead	155
DASA 12.D6.08: Integrated Analysis and Modeling—Mapping Shoreline Habitat Changes	158
Logistics, Support, and Control.....	162
SUP 12.S1.08: Logistics Base Costs.....	162
SUP 12.S2.08: Survey Operations.....	164
SUP 12.S3.08: Control Network	167
PLAN 12.P1.08: Enhancing the Conceptual Ecosystem Model to Identify Critical Ecosystem Interactions and Data Gap.....	170
PLAN 12.P2.08: GCDAMP Effectiveness Workshop Followup.....	173
PLAN 12.P3.08: Maintain a high-quality monitoring, research, and adaptive management program	176
Administrative Operations	183
ADM 12.A1.08: Administrative Operations	183
ADM 12.A2.08: Program Planning and Management.....	186
ADM 12.A3.08: AMWG/TWG Meeting Travel Funds	189
ADM 12.A4.08: Independent Reviews	191
ADM 12.A6.08: Science Symposium.....	191
ADM 12.A5.08: GCMRC Component of SBSC Systems Administration Support.....	195
References Cited	198
Glossary.....	205
Appendix A. Key Science Questions Addressed in the FY2007–11 Science Program.....	207
Appendix B. GCDAMP Fiscal Year 2008 Budget Explanatory Material.....	211
Appendix C. GCDAMP Fiscal Year 2008 Budget.....	213

Chapter 1. Bureau of Reclamation, Upper Colorado Regional Office, Glen Canyon Dam Adaptive Management Program Annual Budget and Work Plan—Fiscal Year 2008

Introduction

The Glen Canyon Dam Adaptive Management Program (GCDAMP) was established in 1997 as an outcome of the Record of Decision on the Operation of Glen Canyon Dam Final Environmental Impact Statement. The purpose of the program is to conduct research and monitoring that is used to develop recommendations to the Secretary of the Interior on the operation of the dam and other authorities under the Secretary.

The Glen Canyon Dam Adaptive Management Work Group (AMWG), a Federal Advisory Committee, is chaired by a designee, appointed by the Secretary of the Interior. Membership is appointed by the Secretary of the Interior with representation from each of the cooperating agencies, Colorado River Basin States, environmental groups, recreation interests, and contractors for Federal power from Glen Canyon Dam. The formation of an advisory committee has provided a forum of discussion for bringing key issues to resolution. The Secretary of the Interior has been mandated to operate the dam and regulate the river in such a manner as to meet the many and varied statutory goals mandated by Congress. The AMWG makes it possible for the Secretary to bring all these varied interests to a consensus on how to protect downstream resources and strike a wise balance on river operations. The Technical Work Group (TWG) provides recommendations to the AMWG based on scientific findings of the GCDAMP.

The Bureau of Reclamation (Reclamation) Upper Colorado Regional Office has responsibility for the administrative activities associated with the AMWG and the U.S. Geological Survey's (USGS) Grand Canyon Monitoring and Research Center (GCMRC) has responsibility for the scientific monitoring and research of the GCDAMP (see chapter 2, this report). Chapter 1 presents the fiscal year 2008 (FY2008) budget and annual work plan for administrative activities overseen by Reclamation.

PROJECT TITLE AND ID: A.1. Personnel Costs

General project description: This project represents Reclamation’s staff costs to perform the daily work activities required to operate the Adaptive Management Work Group (AMWG). The work includes completing assignments resulting from AMWG meetings, consulting with stakeholders on a variety of GCDAMP issues relating to the operation of Glen Canyon Dam (GCD), disseminating pertinent information to the AMWG, preparing and tracking budget expenses, and updating Reclamation’s Web page.

Project goals and objectives: The primary goal is to perform all work associated with the AMWG in a timely and efficient manner, while using the funds available as prudently as possible. Secondary goals include increasing each stakeholder’s awareness of significant budget and legislative issues related to the GCDAMP, improving working relationships with the AMWG members/alternates, finding constructive ways to resolve differences, and addressing individual concerns in an open and accepting forum of discussion.

Expected results: Personnel costs will not exceed what has been proposed in the budget for FY2008 and Reclamation staff will provide budget information to the AMWG on a regular basis. Completed work products will be of high quality and promptly distributed to AMWG members/alternates and interested parties. Budget reports will be presented in a format conducive to AMWG needs.

Budget: FY2008 = \$159,267

Reclamation Project A.1. Personnel Costs—Funding History					
Activity	2004	2005	2006	2007	2008
Outside USBR science/labor	—	—	—	—	—
Logistics field support	—	—	—	—	—
Project-related travel/training	—	—	—	—	—
Operations/supplies	—	—	—	—	—
USBR salaries	110,230	113,537	116,375	119,866	123,462
Subtotal	110,230	113,537	116,375	119,866	123,462
DOI customer burden (29%)	40,770	41,993	43,043	34,762	35,805
Project Total	151,000	155,530	159,418	154,628	159,267
% Total Outsourced	—	—	—	—	

PROJECT TITLE AND ID: A.2. AMWG Member Travel Reimbursement

General project description: This project covers the costs to reimburse AMWG members or alternates to attend regularly scheduled AMWG meetings.

Project goals and objectives: The primary goal for reimbursing travel expenses to AMWG members or alternates is to encourage their attendance at all meetings. Because the meetings are often scheduled in Phoenix, Arizona, many members must incur air or privately owned vehicle (POV) travel, and by having Reclamation reimburse those and other related travel costs, e.g., hotel, per diem, rental car, etc., opportunities are increased for more members to participate in a variety of AMWG assignments. Also, because Reclamation can purchase airline tickets at the Federal Government rate, there are additional cost savings to the program.

Expected results: The GCDAMP will benefit by having all the AMWG members participating in regularly scheduled meetings. As a collective body, they address and resolve concerns associated with the operation of GCD and make recommendations to the Secretary of the Interior for continued science efforts performed below the GCD.

Budget: FY2008 = \$16,683

Reclamation Project A.2. AMWG Member Travel Reimbursement—Funding History					
Activity	2004	2005	2006	2007	2008
Outside USBR science/labor	—	—	—	—	—
Logistics field support	—	—	—	—	—
Project-related travel/training	10,000	13,000	15,725	16,197	16,683
Operations/supplies	—	—	—	—	—
USBR salaries	—	—	—	—	—
Subtotal	10,000	13,000	15,725	16,197	16,683
DOI customer burden (29%)	—	—	—	—	—
Project Total	10,000	13,000	15,725	16,197	16,683
% Total Outsourced	—	—	—	—	—

PROJECT TITLE AND ID: A.3. Reclamation Travel

General project description: This project covers travel expenses Reclamation staff incur to attend AMWG and ad hoc group meetings. In order to work on AMWG/ad hoc assignments, the meetings are often held in Phoenix, Arizona. As such, Reclamation staff must make additional trips throughout the year in completion of those assignments.

Project goals and objectives: The primary goal is for Reclamation staff to be able to travel to meetings and participate in completing AMWG/TWG assignments. By doing so, the program benefits from greater interaction among its members as well as continued improvement and commitment to operating GCD in the best manner possible and obtaining the results from science work being done in the canyon.

Expected results: Reclamation staff will be involved with the AMWG/TWG members in completing work assignments and resolving issues that affect the GCDAMP. They will develop better working relationships with all involved and work toward consensus on a variety of sensitive issues.

Budget: FY2008 = \$13,792

Bureau of Reclamation Project A.3. Reclamation Travel—Funding History					
Activity	2004	2005	2006	2007	2008
Outside USBR science/labor	—	—	—	—	—
Logistics field support	—	—	—	—	—
Project-related travel/training	18,000	15,540	13,000	13,390	13,792
Operations/supplies	—	—	—	—	—
USBR salaries	—	—	—	—	—
Subtotal	18,000	15,540	13,000	13,390	13,792
DOI customer burden (29%)	—	—	—	—	—
Project Total	18,000	15,540	13,000	13,390	13,792
% Total Outsourced	—	—	—	—	—

PROJECT TITLE AND ID: A.4. Facilitation Contract

General project description: This project represents the work assigned to one individual under contract to Reclamation to facilitate at Adaptive Management Work Group meetings. This person may also assist AMWG ad hoc groups in completing AMWG assignments.

Project goals and objectives: The facilitator’s primary responsibility is to keep the AMWG meetings organized and help the members reach consensus on important issues. The facilitator creates a setting in which all members and the public are able to express their views.

Results: The facilitator will create an atmosphere in which the members and other participants at AMWG meetings feel comfortable expressing their individual viewpoints. The facilitator will bring the AMWG members to consensus on pertinent issues affecting the GCDAMP.

Budget: FY2008 = \$25,750

Reclamation Project A.4. Facilitation Contract—Funding History					
Activity	2004	2005	2006	2007	2008
USBR reimbursements	21,000	21,000	25,000	25,000	25,750
Logistics field support	—	—	—	—	—
Project-related travel/training	—	—	—	—	—
Operations/supplies	—	—	—	—	—
USBR salaries	—	—	—	—	—
Subtotal	21,000	21,000	25,000	25,000	25,750
DOI customer burden (29%)	—	—	—	—	—
Project Total	21,000	21,000	25,000	25,000	25,750
% Total Outsourced	—	—	—	—	—

PROJECT TITLE AND ID: A.5. Public Outreach

General project description: This project covers the expenses for Reclamation staff and the Public Outreach Ad Hoc Group (POAHG) to develop materials for the GCDAMP public outreach efforts.

Project goals and objectives: Reclamation Public Affairs staff and the POAHG will work jointly in developing materials to inform and educate the public on the goals and administration of the GCDAMP. They will keep other GCDAMP members advised of progress and expenditures.

Expected results: Products will include fact sheets, Web site information, tribal outreach materials, video B-roll, special events, conference participation, and other pertinent means of advising the public and program members on the achievements of the GCDAMP. The POAHG will maintain accurate records of payments made against the contracts and will keep Reclamation staff informed of discrepancies or concerns.

Budget: FY2008 = \$53,045 (The AMWG approved carryover of \$25,000 but not to exceed a total budget of \$75,000 each fiscal year.)

Reclamation Project A.5. Public Outreach—Funding History					
Activity	2004	2005	2006	2007	2008
Outside USBR science/labor	—	—	—	—	—
Logistics field Support	—	—	—	—	—
Project-related travel/training	—	—	—	—	—
Operations/supplies	—	—	—	—	—
USBR salaries	0	0	50,000	51,500	37,662
Subtotal	0	0	50,000	51,500	37,662
DOI customer burden (29%)	—	—	—	—	15,383
Project Total	0	0	50,000	51,500	53,045
% Total Outsourced	—	—	—	—	—

PROJECT TITLE AND ID: A.6. Other

General project description: This project represents some of the other “miscellaneous” expenses incurred in operation of the AMWG. Some examples follow:

- Overnight mailings of AMWG meeting packets.
- Copying of reports.
- Purchasing meeting materials (cassette tapes, markers, paper, software upgrades for GCDAMP Web site posting, etc.).
- Equipment (audio recording/transcribing machines).

In addition to the above, training courses are often required for staff to keep current on environmental issues, Federal Advisory Committee Act changes, computer technology improvements, etc. Also included in this category are monetary awards given to Reclamation staff who have contributed significantly to the success of the GCDAMP.

Project goals and objectives: The primary goal is to limit spending on “other” items as much as possible. By doing so, more money can be applied to science and research.

Expected results: Other expenses will be kept to a minimum in an effort to reduce the administrative portion of the GCDAMP budget.

Budget: FY2008 = \$7,612

Reclamation Project A.6. Other—Funding History					
Activity	2004	2005	2006	2007	2008
Outside USBR science/labor	—	—	—	—	—
Logistics field support	—	—	—	—	—
Project-related travel/training/awards	6,000	5,000	5,000	5,390	5,612
Operations/supplies	3,000	2,000	2,175	2,000	2,000
USBR salaries	—	—	—	—	—
Subtotal	9,000	7,000	7,175	7,390	7,612
DOI customer burden (29%)	—	—	—	—	—
Project Total	9,000	7,000	7,175	7,390	7,612
% Total Outsourced	—	—	—	—	—

Note: Because many of the AMWG and TWG meetings are held at the Bureau of Indian Affairs (BIA) office in downtown Phoenix, Arizona, there is a cost savings of approximately \$12,800 for not having to use hotel conference rooms; room costs range between \$600 and \$800 per day. Also, because BIA has been able to host many of the AMWG and TWG meetings, they provide use of their copiers and other equipment needed for the meetings at a savings of at least \$1,000 a year to the program.

PROJECT TITLE AND ID: B.1. Personnel Costs

This project represents Reclamation staff costs to perform the daily work activities required to operate the Technical Work Group (TWG), a subgroup of the AMWG. The work includes completing assignments resulting from TWG meetings, consulting with stakeholders on a variety of GCDAMP issues relating to the operation of GCD, disseminating pertinent information to the TWG, preparing and tracking budget expenses, and updating Reclamation’s Web page.

Project goals and objectives: This project represents Reclamation staff costs to perform the daily work activities required to operate the TWG. The work includes completing assignments resulting from AMWG or TWG meetings, consulting with stakeholders on a variety of GCDAMP issues relating to the operation of GCD, disseminating pertinent information to the TWG, preparing and tracking budget expenses, and updating Reclamation’s Web page.

Expected results: Personnel costs will not exceed what has been proposed in the budget and Reclamation staff will provide budget information to the TWG on a regular basis. Completed work products will be promptly distributed to TWG members/alternates and interested parties.

Budget: FY2008 = \$72,777

Reclamation Project B.1. Personnel Costs—Funding History					
Activity	2004	2005	2006	2007	2008
Outside USBR science/labor	—	—	—	—	—
Logistics field support	—	—	—	—	—
Project-related travel/training	—	—	—	—	—
Operations/supplies	—	—	—	—	—
USBR salaries	50,370	51,881	53,178	54,773	56,416
Subtotal	50,370	51,881	53,178	54,773	56,416
DOI customer burden (29%)	18,630	19,189	19,669	15,884	16,361
Project Total	69,000	71,070	72,847	70,657	72,777
% Total Outsourced	—	—	—	—	

PROJECT TITLE AND ID: B.2. TWG Member Travel Reimbursement

General project description: This project covers the costs to reimburse TWG members or alternates to attend regularly scheduled TWG meetings.

Project goals and objectives: The primary goal for reimbursing travel expenses to TWG members or alternates is to encourage their attendance at all meetings. Because the meetings are often scheduled in Phoenix, Arizona, many members must incur air or personal vehicle travel. By reimbursing those and other related travel costs, e.g., hotel, per diem, rental car, etc., opportunities are increased for more members to participate in a variety of AMWG/TWG assignments.

Expected results: The GCD Adaptive Management Program will benefit from having all the TWG members participate in regularly scheduled meetings. As a collective body, they address and resolve concerns associated with the operation of GCD and make recommendations to the AMWG for continued research in the canyon.

Budget: FY2008 = \$22,877

Reclamation Project B.2. TWG Member Travel Reimbursement—Funding History					
Activity	2004	2005	2006	2007	2008
Outside USBR science/labor	—	—	—	—	—
Logistics field support	—	—	—	—	—
Project-related travel/training	15,000	15,540	20,836	22,211	22,877
Operations/supplies	—	—	—	—	—
USBR salaries	—	—	—	—	—
Subtotal	15,000	15,540	20,836	22,211	22,877
DOI customer burden (29%)	—	—	—	—	—
Project Total	15,000	15,540	20,836	22,211	22,877
% Total Outsourced	—	—	—	—	—

PROJECT TITLE AND ID: B.3. Reclamation Travel

General project description: This project covers travel expenses Reclamation staff will incur to prepare and attend TWG meetings as well as ad hoc group meetings which result from AMWG/TWG assignments. In order to work on those assignments, the meetings are often held in Phoenix, Arizona, because it is centrally located to those entities/States represented on the AMWG/TWG. This often requires Reclamation staff to make additional trips throughout the year in completion of AMWG/TWG assignments.

Project goals and objectives: The primary goal is for Reclamation staff to be able to travel to meetings and participate in completing AMWG/TWG assignments. By doing so, the program benefits from greater interaction among its members as well as continued improvement and commitment to operating GCD in the best manner possible and for obtaining the necessary results from science work done in the canyon.

Expected results: Reclamation staff will continue to be involved in meeting with the AMWG/TWG members in completing work assignments and resolving issues that affect the operation of GCD. They will develop better working relationships with all involved and work toward consensus on a variety of GCDAMP issues.

Budget: FY2008 = \$16,866

Reclamation Project B.3. Reclamation Travel—Funding History					
Activity	2004	2005	2006	2007	2008
Outside USBR science/labor	—	—	—	—	—
Logistics field support	—	—	—	—	—
Project-related travel/training	17,000	15,510	15,898	16,375	16,866
Operations/supplies	—	—	—	—	—
USBR salaries	—	—	—	—	—
Subtotal	17,000	15,510	15,898	16,375	16,866
DOI customer burden (29%)	—	—	—	—	—
Project Total	17,000	15,510	15,898	16,375	16,866
% Total Outsourced	—	—	—	—	—

PROJECT TITLE AND ID: B.4. TWG Chair Reimbursement

General project description: This project represents the work assigned to one individual under contract to Reclamation to act as chairperson at TWG meetings. This person may also work on AMWG/TWG ad hoc group assignments.

Project goals and objectives: The chairperson’s primary responsibility is to conduct regularly scheduled TWG meetings. The chairperson also participates in ad hoc group assignments and works closely with Reclamation and GCMRC in setting meeting agendas. The chairperson follows up on TWG and ad hoc group assignments and ensures that information is shared with the members and alternates in a timely manner.

Expected results: The chairperson creates an atmosphere in which the members and other participants at TWG meetings feel comfortable expressing their individual viewpoints. The chairperson will bring the TWG members to consensus on sensitive issues with the ultimate goal of making recommendations to AMWG that incorporate the best scientific information available to the GCDAMP. The chairperson will follow up on action items and make assignments as necessary to accomplish TWG objectives.

Budget: FY2008 = \$23,520

Reclamation Project B.4 TWG Chair Reimbursement—Funding History					
Activity	2004	2005	2006	2007	2008
Outside USBR science/labor	21,000	21,630	22,171	22,836	23,520
Logistics field support	—	—	—	—	—
Project-related travel/training	—	—	—	—	—
Operations/supplies	—	—	—	—	—
USBR salaries	—	—	—	—	—
Subtotal	21,000	21,630	22,171	22,836	23,520
DOI customer burden (29%)	—	—	—	—	—
Project Total	21,000	21,630	22,171	22,836	23,520
% Total Outsourced	—	—	—	—	—

PROJECT TITLE AND ID: B.5. Other

General project description: This project represents some of the other “miscellaneous” expenses incurred in operation of the TWG, as follows, for example:

- Overnight mailings of TWG meeting packets.
- Copying of reports.
- Purchasing meeting materials (cassette tapes, markers, paper, etc.).
- Equipment (audio recording/transcribing machines).

Project goals and objectives: The primary goal is to limit spending on “other” items as much as possible. By doing so, more money can be spent on science and research.

Expected results: Other expenses will be kept to a minimum in an effort to keep within the GCDAMP budget.

Budget: FY2008 = \$2,175

Reclamation Project B.5. Other—Funding History					
Activity	2004	2005	2006	2007	2008
Outside USBR science/labor	—	—	—	—	—
Logistics field support	—	—	—	—	—
Project-related travel/training	—	—	—	—	—
Operations/supplies	2,000	2,000	2,050	2,112	2,175
USBR salaries	—	—	—	—	—
Subtotal	2,000	2,000	2,050	2,112	2,175
DOI customer burden (29%)	—	—	—	—	—
Project Total	2,000	2,000	2,050	2,112	2,175
% Total Outsourced	—	—	—	—	—

Note: Because many of the AMWG and TWG meetings are held at the Bureau of Indian Affairs office in downtown Phoenix, Arizona, there is a cost savings of approximately \$12,800 for not having to use hotel conference rooms: room costs range between \$600 and \$800 per day. Also, because BIA has been able to host many of the AMWG and TWG meetings, they provide use of their copiers and other equipment needed for the meetings at a savings of at least \$1,000 a year to the program.

PROJECT TITLE AND ID: C.1. Compliance Documents

General project description: This project covers the costs for preparing compliance documents for GCDAMP-proposed actions in order to comply with the Endangered Species Act (ESA), National Environmental Policy Act (NEPA), and National Historic Preservation Act (NHPA). In FY2007 much of this funding will be used for compliance documents for the Long-Term Experimental Plan. This will include changes in dam releases and nonflow actions, perhaps including testing of a temperature control device on GCD.

Project goals and objectives: Reclamation staff will keep informed on changes to ESA, NEPA, and NHPA, and will consult with AMWG stakeholders to ensure appropriate compliance is undertaken for actions taken in support of the GCDAMP.

Expected results: Reclamation staff will be involved in all compliance issues related to the GCD Adaptive Management Program. They will utilize travel expenses to meet with the GCDAMP stakeholders to resolve any differences.

Budget: FY2008 = \$271,531 (2/3 Reclamation, 1/3 outside)

Reclamation Project C.1. Compliance Documents—Funding History					
Activity	2004	2005	2006	2007	2008
Outside USBR science/labor	—	—	—	—	90,510
Logistics field support	—	—	—	—	—
Project-related travel/training	—	—	—	—	—
Operations/supplies	—	—	—	—	—
USBR salaries	26,000	26,780	22,450	263,622	128,525
Subtotal	26,000	26,780	22,450	263,622	128,525
DOI customer burden (29%)	—	—	—	—	52,496
Project Total	26,000	26,780	22,450	263,622	271,531
% Total Outsourced	—	—	—	—	—

PROJECT TITLE AND ID: C.2. Administrative Support for NPS Permitting

General project description: This project provides funding to support the Grand Canyon National Park permitting of research and monitoring projects conducted under the GCDAMP. Grand Canyon National Park employs a permitting specialist and staff who review all proposals for projects to be completed in the Park under the auspices of the GCDAMP. The program provides these funds to offset the administrative burden of the Park in providing these services.

Project goals and objectives: The primary goal is to ensure that projects conducted under the GCDAMP are reviewed and permitted by the National Park Service (NPS).

Expected results: Projects conducted under the GCDAMP will receive permits from the NPS in a timely manner.

Budget: FY2008 = \$113,300

Reclamation Project C.2. Administrative Support for NPS Permitting—Funding History					
Activity	2004	2005	2006	2007	2008
Outside USBR science/labor				—	—
Logistics field support				—	—
Project-related travel/training				—	—
Operations/supplies					
USBR salaries				—	—
Subtotal			100,000	110,000	113,300
DOI customer burden (29%)				—	—
Project Total			100,000	110,000	113,300
% Total Outsourced				—	—

PROJECT TITLE AND ID: C.3. Contract Administration

General project description: This project covers the expenses for Reclamation staff to prepare and monitor contracts associated with the GCDAMP. Specifically, these contracts are for AMWG facilitation, TWG chairperson reimbursement, tribal participation, and programmatic agreement work.

Project goals and objectives: Reclamation contract specialists will accurately apply funds spent on individual contracts to ensure costs do not exceed contract limits. They will keep other Reclamation staff informed as to those charges so accurate reporting can be made to both AMWG and TWG members.

Expected results: Contract specialists will ensure that individual contractors are fulfilling the requirements of their contracts. They will maintain accurate records of payments made against the contracts and will keep Reclamation staff informed of discrepancies or concerns. Work will be completed on time and within the limits of the contract.

Budget: FY2008 = \$33,385

Reclamation Project C.3. Contract Administration—Funding History					
Activity	2004	2005	2006	2007	2008
Outside USBR science/labor	—	—	—	—	—
Logistics field support	—	—	—	—	—
Project-related travel/training	—	—	—	—	—
Operations/supplies	—	—	—	—	—
USBR salaries	25,000	25,750	24,394	32,413	23,703
Subtotal	25,000	25,750	24,394	32,413	23,703
DOI customer burden (29%)	—	—	—	—	9,682
Project Total	25,000	25,750	24,394	32,413	33,385
% Total Outsourced	—	—	—	—	—

PROJECT TITLE AND ID: C.4. Experimental Carryover Funds

General project description: This budget item reserves funds for conducting experiments under the GCDAMP. The estimated need for a large-scale beach/habitat-building flows (BHBF) experiment based on past experience is approximately \$1.5 million. This amount will be reserved over the course of several years so that the effects on annual budget and work plan are minimized.

Project goals and objectives: As above.

Expected results: The funds will be available to conduct a large-scale experiment when conditions are appropriate.

Budget: FY2008 = \$500,000

Reclamation Project C.4 Experimental Carryover Funds—Funding History					
Activity	2004	2005	2006	2007	2008
Outside USBR science/labor	—	—	—	—	—
Logistics field Support	—	—	—	—	—
Project-related travel/training	—	—	—	—	—
Operations/supplies	—	—	—	—	—
USBR salaries					
Subtotal					
DOI customer burden (29%)					
Project Total			424,675	500,000	500,000
% Total Outsourced	—	—	—	—	—

PROJECT TITLE AND ID: C.5. Integrated Tribal Resources Monitoring

General project description: Funding is provided for identification of TCPs and implementation of pilot projects for protocols developed in the FY2007 resources monitoring agreed to by the TWG as part of core-monitoring development.

Project goals and objectives: Primary goal is to evaluate effects of dam operations and other actions under the authority of the Secretary of the Interior on resources of value to Native American tribes.

Expected results: Reports detailing their activities, findings, and monitoring results from implementing protocols as part of core monitoring.

Budget: FY2008 = 136,475

Reclamation Project C.5 Integrated Tribal Resources Monitoring—Funding History					
Activity	2004	2005	2006	2007	2008
Outside USBR science/labor	—	—	—	—	—
Logistics field support	—	—	—	—	—
Project-related travel/training	—	—	—	—	—
Operations/supplies	—	—	—	—	—
USBR salaries	—	—			
Subtotal	—	—	125,000	132,500	136,475
DOI customer burden (29%)	—	—			
Project Total	—	—	125,000	132,500	136,475
% Total Outsourced	—	—	—	—	—

PROJECT TITLE AND ID: D.1. Programmatic Agreement, Reclamation Administrative Costs

General project description: Reclamation’s regional archaeologist administers the programmatic agreement (PA) and tribal contracts. This project funds salary, travel, and indirect costs of program administration. The costs integrate the PA and tribal consultation into the larger GCDAMP.

Project goals and objectives:

- Management of five \$95k (FY2006 funds) tribal sole-source contracts for participation in the GCDAMP. Initiation of second option year for five \$95,000 (FY2007 funds) tribal sole-source contracts. Management of five \$25,000 (FY2006 funds) tribal sole-source contracts.
- Modification of the extant Utah State University (USU)/Zuni Cultural Resource Enterprise and Cooperative Ecosystem Study Unit to add the remaining \$95,000 (FY2006 funds) of the original AMWG treatment plan funding (\$250,000) for purposes of emergency treatment of at-risk archaeological sites.
- Manage one PA meeting and attend TWG and AMWG meetings.

Expected results: The major product is administration of the Glen and Grand Canyon treatment plans, accountability for the tribal contracts, and use of both appropriated dollars and power revenues.

Budget: FY2008 = \$57,354

Reclamation Project D.1 Programmatic Agreement, Reclamation Administrative Costs—Funding History					
Activity	2004	2005	2006	2007	2008
Outside USBR science/labor	—	—	—	—	—
Logistics field support	—	—	—	—	—
Project-related travel/training	—	—	—	—	—
Operations/supplies	—	—	—	—	—
USBR salaries	43,000	51,500	54,107	71,892	40,721
Subtotal	43,000	51,500	54,107	71,892	40,721
DOI customer burden (29%)	—	—	—	—	16,633
Project Total	43,000	51,500	54,107	71,892	57,354
% Total Outsourced	—	—	—	—	—

PROJECT TITLE AND ID: D.2. NPS Support for Archaeological Site Assessment

General project description: This funding is to provide support for NPS involvement in the assessment task for GCMRC project 11.R1.08 (See GCMRC project 11.R1.08 description, this report).

Project goals and objectives: The project goals, objectives, and expected results are provided in the project description (see GCMRC project 11.R1.08 description, this report).

Budget: FY2008 = \$69,525 moved to GCRMC (below the line, added to line 108)

Reclamation Project D.2 NPS Support for Archaeological Site Assessment—Funding History					
Activity	2004	2005	2006	2007	2008
Outside USBR science/labor	—	—	—	—	—
Logistics field support	—	—	—	—	—
Project-related travel/training	—	—	—	—	—
Operations/supplies	—	—	—	—	—
USBR salaries				67,500	0
Subtotal				67,500	0
DOI customer burden (29%)				0	0
Project Total				67,500	0
% Total Outsourced	—	—	—	—	—

PROJECT TITLE AND ID: D.4. Glen and Grand Canyon Treatment Plan Implementation

General project description: In consultation with Grand Canyon NPS, the Arizona State Historic Preservation Office (SHPO) and the remainder of the PA signatories, Reclamation completed a scope-of-work for the development of a treatment plan for the cultural resources of Grand Canyon. This work was completed by March 2007 under a Cooperative Ecosystem Studies Unit agreement with Utah State University. An analogous set of treatment plan recommendations was completed in FY2006 (based on FY2004 and FY2005 funding) by the Navajo Nation Archaeological Department. Treatment of individual properties may include in situ preservation measures, nature and extent testing, full data recovery, or additional documentation/recordation. The determination of appropriate treatment will be based on consultation with NPS, the Arizona SHPO, and Southwestern tribal entities (tribal consultation will not be restricted to PA signatories). This consultation is taking place during FY2007 with treatment plan implementation scheduled to begin in FY2008.

Project goals and objectives:

- Development of a treatment plan Memorandum of Agreement (MOA) through consultation with SHPO, NPS, Tribes, and other stakeholders.
- Government-to-government consultation with tribal councils based upon the treatment plan recommendations.
- Onsite assessment by PA members and tribal elders of a selected sample of historic properties.
- Consequent formulation of an MOA for mitigation with Reclamation, NPS, and the SHPO as principal signatories.
- Field work to be initiated in winter 2007 and completed in spring 2008. Four sites have been selected for treatment in FY2008.

Expected results: Prioritization, based on significance, of all affected Glen and Grand Canyon properties and completion of an MOA for treatment of adverse effects. Detailed and comprehensive reports on consultant activities, results, and recommendations. Evaluation and implementation of mitigative measures or total data recovery, following the Secretary of the Interior Standards and Guidelines for Historic Preservation and guidance of the Advisory Council on Historic Preservation.

Budget: FY2008 = \$300,000 (\$75,000 = sole-source contract for logistical assistance during the consultation; \$70,000 = Park service for logistical support and consulting with them)

**Reclamation Project D.4 Glen and Grand Canyon Treatment Plan Implementation—
Funding History**

Activity	2004	2005	2006	2007	2008
Outside USBR science/labor	371,000	676,340	270,000	145,000	300,000
Logistics field support	—	—	—	—	—
Project-related travel/training	—	—	—	—	—
Operations/supplies	—	—	—	—	—
USBR salaries	—	—	—	—	—
Subtotal	371,000	676,340	270,000	145,000	300,000
DOI customer burden (29%)	—	—	—	—	—
Project Total	371,000	676,340	270,000	145,000	300,000
% Total Outsourced	—	—	—	—	—

PROJECT TITLE AND ID: E. Tribal Consultation—Sole-Source Contracts with Tribes

General project description: Government-to-government consultation will be maintained among the five GCDAMP tribes (Hopi Tribe, Hualapai Tribe, Southern Paiute Consortium, Pueblo of Zuni, Navajo Nation) and five DOI agencies (U.S. Geological Survey [USGS], NPS, Reclamation, U.S. Fish and Wildlife Service [USFWS], and BIA).

Project goals and objectives: The purpose of the continued funding of tribal contracts is to ensure tribal viewpoints are integrated into continuing GCDAMP dialogs, votes, and in the final recommendations made to the Secretary of the Interior.

Expected results: The most important product is the incorporation of tribal perspectives into the recommendations forwarded to the Secretary. In addition, the tribes prepare annual reports on activities funded under the contracts. Continued funding of government-to-government consultation through the agreements ensures enhanced communication and understanding of the GCDAMP issues and concerns.

Budget: FY2008 = \$475,000

Reclamation Project E. Tribal Consultation—Sole-Source Contracts with Tribes—Funding History					
Activity	2004	2005	2006	2007	2008
Outside USBR science/labor	320,000	477,375	477,375	475,000	475,000
Logistics field support	—	—	—	—	—
Project-related travel/training	—	—	—	—	—
Operations/supplies	—	—	—	—	—
USBR salaries	—	—	—	—	—
Subtotal	320,000	477,375	477,375	475,000	475,000
DOI customer burden (29%)	—	—	—	—	—
Project Total	320,000	477,375	477,375	475,000	475,000
% Total Outsourced	100%	100%	100%	100%	100%

Chapter 2. U.S. Geological Survey, Southwest Biological Science Center, Grand Canyon Monitoring and Research Center Annual Budget and Work Plan—Fiscal Year 2008


Introduction

The Glen Canyon Dam Adaptive Management Program (GCDAMP) is a science-based process for continually improving management practices related to the operation of Glen Canyon Dam (GCD) by emphasizing learning through monitoring, research, and experimentation. The U.S. Geological Survey's (USGS) Grand Canyon Monitoring and Research Center (GCMRC) has responsibility for the scientific monitoring and research of the GCDAMP. GCMRC staff worked cooperatively with GCDAMP participants to identify the scope, objectives, and budget for the monitoring and research projects for fiscal year 2008 (FY2008) presented in the Grand Canyon Monitoring and Research Center Fiscal Year 2008 Budget and Annual Work Plan (AWP). As was the case in FY2007, the AWP for FY2008 is a transitional plan designed to fund the GCDAMP Science Program for 1 year while consideration is given to the development of the Long-Term Experimental Plan (LTEP), a science and funding plan for a temperature control device (TCD), and the development of a recovery program for the humpback chub (*Gila cypha*) (HBC) in Grand Canyon. Beginning in FY2009, the expectations are that biennial work plans (BWP) will be developed as noted below. Other major components of the science planning process include the following:

- Final Draft Glen Canyon Dam Adaptive Management Program Strategic Plan (AMPSP): A long-term plan drafted by GCDAMP participants in cooperation with the GCMRC that identifies the Adaptive Management Work Group (AMWG) vision and mission statement, principles, goals, management objectives, information needs, and management actions.
- GCMRC Strategic Science Plan (SSP): Developed by the GCMRC in cooperation with GCDAMP participants to identify strategies for providing science information during a 5-year period to respond, consistent with the AMPSP, to goals, management objectives, and priority questions of the GCDAMP participants. The SSP was approved by the Secretary of the Interior in May 2006.
- GCMRC Monitoring and Research Plan (MRP): Developed by the GCMRC in cooperation with the GCDAMP to specify research and monitoring activities for the next 5 years consistent with the strategies and priorities in the SSP. The MRP identifies the objectives associated with each strategic science question (SSQ) and related monitoring, experimental research, and research and development projects. The November 14, 2006, draft of the MRP was approved as a working document by the AMWG in December 2006 to help guide the development of the FY2008–09 work plan for the GCDAMP.

Figure 1 depicts the flow of information in the science planning and implementation process. Annually, the GCMRC will report on accomplishments related to projects included in the biennial work plan and evaluate how science has advanced knowledge relative to GCDAMP goals and management objectives. At 5-year intervals, GCMRC shall formally synthesize new scientific information in the form of an updated The State of the Colorado River Ecosystem in Grand Canyon (SCORE) report (Gloss and others, 2005). In addition, the Knowledge Assessment Report (KAR) (Melis and others, 2006) will be revised to identify knowledge gaps related to the effects of various treatments/management actions on resources of interest to the GCDAMP (e.g., the effect of dam operations on humpback chub recruitment). Information from the Knowledge Assessment (KA) will be used to identify key strategic questions associated with priority GCDAMP information needs or questions. Priority information needs and science questions will be evaluated by scientists and managers to determine what revisions to the science program are needed. This includes development of revised SSP and MRP documents and a new

Figure 1. Collaborative science planning and implementation process. The Glen Canyon Dam Adaptive Management Program and Department of the Interior have lead responsibility for the shaded boxes. The Grand Canyon Monitoring and Research Center has lead responsibility for the boxes not shaded.



experimental research plan. The BWP will be updated annually to address new information needs and to develop new work plans for the second year of the 2-year planning cycle. All these activities will be carried out collaboratively by scientists and GCDAMP participants. Involvement will be provided through the AMWG, TWG, appropriate ad hoc groups, and the Science Advisors Board (SAB).

Purpose

The purpose of the AWP is to describe the core-monitoring, long-term experimental, research and development, and other related activities that will be implemented in FY2008 to address priority goals, questions, and information needs specified by the GCDAMP.

Overview of the GCMRC Strategic Science Plan and Monitoring and Research Plan

The AWP is designed to implement and be consistent with the draft GCMRC SSP and MRP dated May 5, 2006, and June 21, 2006, respectively. The principal elements of the MRP and SSP that are addressed by the FY2008 AWP include:

1. **Adaptive environmental assessment and management (AEAM) approach**—The GCMRC science program will be based on the AEAM approach to natural resources management that was developed by Holling (1978) and Walters (1986) and articulated in the AMPSP.
2. **Collaborative science planning process**—The GCMRC will use the planning process described above and illustrated in figure 1 to develop and update science plans and related work plans.

3. **Priority strategic science questions**—The GCDAMP priority questions and the associated SSQs provide the primary (but not exclusive) basis for designing the science program (appendix A).
4. **Interdisciplinary integrated river science**—Increased emphasis will be provided on employing an interdisciplinary, integrated science approach over the next 5 years. Principal elements of this approach involve
 - aligning GCMRC staffing/organization to facilitate integrated, interdisciplinary science;
 - enhancing the Colorado River conceptual ecosystem model (CEM) to identify critical ecosystem interactions and data gaps; and
 - initiating an effort to gather and evaluate baseline data and develop modeling capabilities to assist in planning and evaluating a proposed GCD TCD.
5. **Bridging science and management**—The GCMRC will develop and implement a collaborative plan/assessment among scientists and GCDAMP participants to improve the effectiveness of the GCDAMP and better integrate the use of scientific information into the GCDAMP process. The plan/assessment will address (1) the feasibility of developing/using decision-support tools to facilitate integration of scientific information in the science planning and GCDAMP decisionmaking processes including resource tradeoff assessments, and (2) strategies/approaches for improving the effectiveness of the GCDAMP process. In FY2008, the GCMRC will convene a workshop for scientists and GCDAMP participants to develop an action plan for addressing priority issues, needs, or opportunities related to the effectiveness of the GCDAMP, and the use of scientific information in the GCDAMP process.

Overview of Annual Work Plan and Budget

The FY2008 AWP was developed based on the Monitoring and Research Plan to Support the Glen Canyon Dam Adaptive Management Program (hereafter MRP). In December 2006, the AMWG approved the MRP as a working document to help guide the preparation of the FY2008–09 work plan and budget.

The proposed budget provides for the continued implementation of projects included in the approved GCDAMP Fiscal Year 2007 Annual Budget and Work Plan. The only new projects identified in FY2008 are the following:

- The Grand Canyon archaeological site treatment plan (\$300,000 in the Bureau of Reclamation [Reclamation] Department of the Interior [DOI] portion of the budget).
- The long-term sediment storage monitoring project, which will be reviewed by the TWG for core-monitoring status (\$95,000).

To achieve a balanced budget, a number of projects had to be scaled back to accommodate the increased funding being requested for the new projects noted above and for other nondiscretionary increases in costs for continuing projects.

Several new projects that were identified in the MRP to start in FY2008 will need to be deferred because no funding was available after the continuing projects were funded. These include the following:

- Expanding/updating the CEM
- Hiring a visiting ecosystem scientist to pursue specific integrated ecosystem science strategies
- Compiling and analyzing existing recreation safety data (deferred from FY2007)
- Evaluating the relative importance and effects of different flows on the recreation experience

Unfunded projects identified in the MRP should be the highest priority for any discretionary funds that may become available in FY2008.

In addition, the MRP identified several new initiatives to be undertaken in FY2008 to address priority research and monitoring information needs in areas outside of the Colorado River ecosystem (CRE). These projects, which were proposed for funding by USGS or other non-GCDAMP sources, include the following:

- Lake Powell modeling and data synthesis;
- LCR gaging, water quality/quantity synthesis, and contaminant risk assessment; and
- Assessment of climate change and drought effects on GCD operations.

USGS funding to address these needs was not included in the President's proposed FY2008 budget request, and therefore, these activities are not included in the FY2008 AWP.

The FY2008 AWP does not account for the GCMRC's support for the development of the Long-Term Experimental Plan (LTEP) environmental impact statement (EIS). The FY2008 work plan may need to be adjusted after Reclamation provides more detailed direction on the scope of GCMRC involvement in the EIS process.

Table 1 summarizes core-monitoring, research and development, and experimental activities in the FY2008 annual work plan for the GCMRC. Activities address GCDAMP goals 1–11, including related science questions and information needs. Priority and related SSQs are paraphrased from the Draft GCMRC Strategic Science Plan (appendix A) and the core-monitoring information needs developed by the Science Planning Group (SPG). Three categories of activities are identified:

1. **Core-monitoring activities**—Core monitoring is consistent, long-term repeated measurements using scientifically accepted protocols to measure status and trends of key resources. Core-monitoring activities are those that have been pilot tested for one to several years, undergone a protocols evaluation panel (PEP) evaluation and peer review, and have been formally approved by the GCDAMP for core-monitoring status. In FY2008, the monitoring activities associated with the status of HBC in the LCR and mainstem Colorado River are scheduled for PEP evaluation by the GCMRC and the TWG for core-monitoring status.
2. **Research and development activities**—Activities aimed at (1) addressing specific hypotheses or information needs related to a priority GCDAMP resource(s) and (2) developing/testing new technologies or monitoring procedures. Examples of research and development (R&D) activities in the FY2008 work plan include:
 - linking whole-system carbon cycling to food webs in the Colorado River—the project that will provide the basis for the food base monitoring program;
 - investigating new, more effective technologies for sampling fish populations, such as remote passive integrated transponder (PIT) tag reading technology and sonic tag technology;
 - advancing the development of downstream flow, temperature, and suspended-sediment models; and
 - evaluating the quality of historical remote-sensing imagery for change detection.
3. **Experimental activities**—A suite of flow and nonflow treatments and/or management actions designed to improve conditions of target resources (HBC, sediment, etc.) while allowing for an understanding of the relationship between treatments/management actions and the target resources. The LTEP has yet to be finalized by the GCDAMP. Several long-term experimental options are currently being evaluated by Reclamation pursuant to the National Environmental Policy Act in coordination with the GCDAMP. The LTEP will be implemented following issuance of a Record of Decision (ROD) in FY2009.

Table 1. Summary of core-monitoring, research and development, and experimental activities in the fiscal year 2008 (FY2008) annual work plan for the Grand Canyon Monitoring and Research Center (GCMRC). Several long-term experimental options currently under discussion are not reflected in the table; additional experimental options will be developed upon approval by the U.S. Department of the Interior. Activities address Glen Canyon Dam Adaptive Management Program (GCDAMP) goals 1–12 in relation to science questions and information needs. Priority and related strategic science questions are paraphrased from the Draft GCMRC Strategic Science Plan (appendix A). Information needs are paraphrased from the GCDAMP Strategic Plan. Abbreviations are as follows: SSQ = strategic science question, CMIN = core-monitoring information need, RIN = research information need, and SA = GCDAMP Science Advisors summary questions.

GCDAMP goal	Priority science questions and information needs (questions from Strategic Science Plan and Monitoring and Research Plan in italics)	Core-monitoring activities	Experimental activities	Research and development activities
<p><i>1. Food base</i></p>	<p><i>AMWG Priority: 1, 3, and 5</i></p> <p>SSQ 1-5. What are the important pathways that link lower trophic levels with fish and how will they link to dam operations?</p> <p>SSQ 1-6. Are fish populations, trends, or indicators from fish, such as growth, condition, and body composition, correlated with patterns in invertebrate flux?</p> <p>SSQ 5-2. Is invertebrate flux affected by water quality (e.g., temperature, nutrient concentrations, turbidity) and dam operations?</p>			<p><i>FY2006–09: Determine carbon budget to understand how energy is exchanged among organisms in the Colorado River; develop monitoring techniques and metrics for key organisms</i></p> <p><i>FY2008: Diet, drift, and predation data analysis</i></p>
<p><i>2. Humpback chub (HBC) and other native fishes (A.)</i></p>	<p><i>AMWG Priority: 1, 3, and 5</i></p> <p>SSQ 1-1. To what extent are adult populations of native fish controlled by production of young fish from tributaries, spawning and incubation in the mainstem, survival of young-of-year (YoY) and juvenile stages in the mainstem, or by changes in growth and maturation in the adult population as influenced by mainstem conditions?</p> <p>SSQ 1-4. Can long-term decreases in abundance of rainbow trout be sustained with a reduced level of effort of mechanical removal or will recolonization from tributaries and from downstream and upstream of the removal reach require that mechanical removal be an ongoing management action? This question also applies to future removal programs targeting other nonnative species.</p> <p><i>CMIN 2.1.2 Determine and track abundance and distribution of all size classes of HBC in the Little Colorado River (LCR) and the mainstem.</i></p>	<p><i>FY2007–08: Monitor status and trends of HBC in LCR and mainstem using existing protocols</i></p>		<p><i>FY2006 and ongoing: Stock assessment</i></p> <p><i>FY2007–08: Gear efficiency/sampling evaluation</i></p> <p><i>FY2007–11: Statistical review of existing HBC monitoring protocols and habitat data</i></p> <p><i>FY2007–11: Evaluate protocols for warmwater and coldwater nonnative fish monitoring, removal, and control; effects on native fish</i></p>

Table 1. Summary of core-monitoring, research and development, and experimental activities in the fiscal year 2008 (FY2008) annual work plan for the Grand Canyon Monitoring and Research Center (GCMRC). Several long-term experimental options currently under discussion are not reflected in the table; additional experimental options will be developed upon approval by the U.S. Department of the Interior. Activities address Glen Canyon Dam Adaptive Management Program (GCDAMP) goals 1–12 in relation to science questions and information needs. Priority and related strategic science questions are paraphrased from the Draft GCMRC Strategic Science Plan (appendix A). Information needs are paraphrased from the GCDAMP Strategic Plan. Abbreviations are as follows: SSQ = strategic science question, CMIN = core-monitoring information need, RIN = research information need, and SA = GCDAMP Science Advisors summary questions.—Continued

GCDAMP goal	Priority science questions and information needs (questions from Strategic Science Plan and Monitoring and Research Plan in italics)	Core-monitoring activities	Experimental activities	Research and development activities
<p>2. HBC and other native fishes (B.)</p>	<p>AMWG Priority: 1, 3, and 5</p> <p><i>SSQ 1-2. Does a decrease in the abundance of rainbow trout and other cold- and warmwater nonnatives in Marble and eastern Grand Canyons result in an improvement in the recruitment rate of juvenile HBC to the adult population?</i></p> <p><i>SSQ 1-4. Can long-term decreases in abundance of rainbow trout in Marble and eastern Grand Canyons be sustained with a reduced level of effort of mechanical removal or will recolonization from tributaries and from downstream and upstream of the removal reach require that mechanical removal be an ongoing management action?</i></p> <p><i>SSQ 5-6. Do the potential benefits of improved rearing habitat (warmer, more stable, more backwater and vegetated shorelines, more food) outweigh negative impacts due to increases in nonnative fish abundance?</i></p> <p>CMIN 2.4.1 Determine and track the abundance and distribution of nonnative predatory fish species in the CRE and their impacts on native fish.</p> <p>RIN 2.4.1: What are the most effective strategies and control methods to limit nonnative fish predation and competition on native fish?</p> <p>RIN 2.4.3: To what degree, which species, and where in the system are exotic fish a detriment to the existence of native fish through predation or competition?</p>	<p>FY2007–08: Continue mainstem monitoring of fish community</p>		<p>FY2007–10: Develop and test nonnative fish management plan</p> <p>FY2007–11: Develop abundance estimation framework that allows scientists to better estimate nonnative fish numbers in mechanical removal reaches</p> <p>FY2007–10: Develop bioenergetic model to predict changes in fish communities in response to environmental changes</p>
<p>2. HBC and other native fishes (C.)</p>	<p>AMWG Priority: 1, 3, and 5</p> <p>SSQ 1-1. To what extent are adult populations of native fish controlled by production of young fish from tributaries, spawning and incubation in the mainstem, survival of YoY and juvenile stages in the mainstem, or by changes in growth and maturation in the adult population as influenced by mainstem conditions?</p> <p>SSQ 1-7. Which tributary and mainstem habitats are most important to native fishes and how can these habitats best be made useable and maintained?</p> <p>SA 1. What are the most limiting factors to successful HBC adult recruitment in the mainstem: spawning success, predation on YoY and juveniles, habitat (water, temperature), pathogens, adult maturation, food availability, competition?</p>			<p><i>FY2007–10: Review data and literature on HBC in upper basin to see if HBC habitat can be identified, protected, and recreated below GCD</i></p>

Table 1. Summary of core-monitoring, research and development, and experimental activities in the fiscal year 2008 (FY2008) annual work plan for the Grand Canyon Monitoring and Research Center (GCMRC). Several long-term experimental options currently under discussion are not reflected in the table; additional experimental options will be developed upon approval by the U.S. Department of the Interior. Activities address Glen Canyon Dam Adaptive Management Program (GCDAMP) goals 1–12 in relation to science questions and information needs. Priority and related strategic science questions are paraphrased from the Draft GCMRC Strategic Science Plan (appendix A). Information needs are paraphrased from the GCDAMP Strategic Plan. Abbreviations are as follows: SSQ = strategic science question, CMIN = core-monitoring information need, RIN = research information need, and SA = GCDAMP Science Advisors summary questions.—Continued

GCDAMP goal	Priority science questions and information needs (questions from Strategic Science Plan and Monitoring and Research Plan in italics)	Core-monitoring activities	Experimental activities	Research and development activities
<p>2. HBC and other native fishes (D.)</p>	<p>AMWG Priority: 1, 3, and 5</p> <p><i>SSQ 1-8. How can native and nonnative fishes best be monitored while minimizing impacts from capture and handling or sampling?</i></p>			<p>FY2007–09: Develop alternative, noninvasive HBC monitoring gear to reduce stress on fish (e.g., DIDSON camera, remote passive integrated transponder (PIT) tag reading, and sonic tags)</p> <p>FY2007–09. Evaluate the effects of trammel net sampling</p>
<p>3. Extirpated species</p>		<p>No projects</p>	<p>FY2007–11: Evaluation and planning of temperature control device</p>	<p>No projects</p>
<p>4. Rainbow trout (RBT)</p>	<p>AMWG Priority: 3</p> <p><i>SSQ 3-6: What Glen Canyon Dam operations (ramping rates, daily flow range, etc.) maximize trout fishing opportunities and catchability?</i></p> <p>CMIN 4.1.2 Determine annual proportional stock density of rainbow trout in the Lees Ferry reach.</p> <p>CMIN 4.1.4 Determine annual standard condition (Kn) and relative weight of rainbow trout in the Lees Ferry reach.</p>	<p>FY2007–11: Monitor status and trends of Lees Ferry RBT population</p> <p>FY2008: Review/evaluate RBT monitoring for core-monitoring status</p>		

Table 1. Summary of core-monitoring, research and development, and experimental activities in the fiscal year 2008 (FY2008) annual work plan for the Grand Canyon Monitoring and Research Center (GCMRC). Several long-term experimental options currently under discussion are not reflected in the table; additional experimental options will be developed upon approval by the U.S. Department of the Interior. Activities address Glen Canyon Dam Adaptive Management Program (GCDAMP) goals 1–12 in relation to science questions and information needs. Priority and related strategic science questions are paraphrased from the Draft GCMRC Strategic Science Plan (appendix A). Information needs are paraphrased from the GCDAMP Strategic Plan. Abbreviations are as follows: SSQ = strategic science question, CMIN = core-monitoring information need, RIN = research information need, and SA = GCDAMP Science Advisors summary questions.—Continued

GCDAMP goal	Priority science questions and information needs (questions from Strategic Science Plan and Monitoring and Research Plan in italics)	Core-monitoring activities	Experimental activities	Research and development activities
5. Kanab ambersnail (KAS)	AMWG Priority: 3 CMIN 5.1.1 Determine and track the abundance and distribution of KAS at Vaseys Paradise. CMIN 5.2.1 Determine and track the size and composition of the habitat used by KAS at Vaseys Paradise.	FY2008: KAS habitat monitoring.		
6. Springs /riparian	AMWG Priority: 4 <i>SSQ 2-1. Do dam-controlled flows affect (increase or decrease) rates of erosion and vegetation growth at archaeological sites and TCP sites, and if so, how?</i> <i>SSQ 3-2. How important are backwaters and vegetated shoreline habitats to the overall growth and survival of YoY and juvenile native fish? Does the long-term benefit of increasing these habitats outweigh short-term potential costs?</i> CMIN 6.1.1., 6.6.1., 6.2.1., 6.5.1. Determine and track the abundance, composition, distribution, and area of terrestrial native and nonnative vegetation species in the CRE.			FY2008: Terrestrial monitoring FY2008 and ongoing: Terrestrial mapping FY2007–11: Vegetation synthesis project
7. Quality-of-water	AMWG Priority: 1, 3, and 5 <i>SSQ 3-5. How is invertebrate flux affected by water quality (e.g., temperature, nutrient concentrations, turbidity) and dam operations?</i> <i>SSQ 5-1. How do dam release temperatures, flows (average and fluctuating component), meteorology, canyon orientation and geometry, and reach morphology interact to determine mainstem and nearshore water temperatures throughout the CRE)?</i> <i>SSQ 5-3. To what extent do temperature and fluctuations in flow limit spawning and incubation success for native fish?</i> CMIN 7.3.1. What are the status and trends of water quality releases from Glen Canyon Dam?	FY2007–09: Lake Powell monitoring using existing protocols FY2007–11: Downstream integrated quality-of-water (IQW) monitoring (including suspended-sediment flux)		FY2007–11: Advanced development of downstream flow, temperature, and suspended-sediment models

Table 1. Summary of core-monitoring, research and development, and experimental activities in the fiscal year 2008 (FY2008) annual work plan for the Grand Canyon Monitoring and Research Center (GCMRC). Several long-term experimental options currently under discussion are not reflected in the table; additional experimental options will be developed upon approval by the U.S. Department of the Interior. Activities address Glen Canyon Dam Adaptive Management Program (GCDAMP) goals 1–12 in relation to science questions and information needs. Priority and related strategic science questions are paraphrased from the Draft GCMRC Strategic Science Plan (appendix A). Information needs are paraphrased from the GCDAMP Strategic Plan. Abbreviations are as follows: SSQ = strategic science question, CMIN = core-monitoring information need, RIN = research information need, and SA = GCDAMP Science Advisors summary questions.—Continued

GCDAMP goal	Priority science questions and information needs (questions from Strategic Science Plan and Monitoring and Research Plan in italics)	Core-monitoring activities	Experimental activities	Research and development activities
<p>8. <i>Sediment (fine and coarse sediment)</i></p>	<p><i>AMWG Priority: 1,2,3, and 4</i> <i>SSQ 4-1. Is there a “Flow-Only” operation (i.e., a strategy for dam releases, including managing tributary inputs with (beach/habitat-building flows (BHBFs), without sediment augmentation) that will restore and maintain sandbar habitats over decadal timescales?</i></p>	<p>FY2007–11: Implementation of recommendations from the final sediment transport modeling review-protocols evaluation panel (SEDS-PEP) (summer 2006) FY2008: Fine sediment “SED TREND” monitoring —detection of trends in lower elevation channel sand deposits through annual reach-scale topographic measurements of sand storage between suspended-sediment flux monitoring stations. In FY2008, the reach between river mile 0 and 30 will be mapped using multibeam acoustic bathymetry methods for comparison with 2000-01 measurements. FY2008: Coarse sediment—no core-monitoring activities are scheduled at present until the next remote-sensing overflight occurs in FY2009.</p>		<p>FY2007–11: Map change in nearshore habitat resulting from 2004 BHBF; convert existing overflight analog images to digital to facilitate research</p>

Table 1. Summary of core-monitoring, research and development, and experimental activities in the fiscal year 2008 (FY2008) annual work plan for the Grand Canyon Monitoring and Research Center (GCMRC). Several long-term experimental options currently under discussion are not reflected in the table; additional experimental options will be developed upon approval by the U.S. Department of the Interior. Activities address Glen Canyon Dam Adaptive Management Program (GCDAMP) goals 1–12 in relation to science questions and information needs. Priority and related strategic science questions are paraphrased from the Draft GCMRC Strategic Science Plan (appendix A). Information needs are paraphrased from the GCDAMP Strategic Plan. Abbreviations are as follows: SSQ = strategic science question, CMIN = core-monitoring information need, RIN = research information need, and SA = GCDAMP Science Advisors summary questions.—Continued

GCDAMP goal	Priority science questions and information needs (Questions from Strategic Science Plan and Monitoring and Research Plan in italics)	Core-monitoring activities	Experimental activities	Research and development activities
9. Recreation (A)	<p>AMWG Priority: 3 and 4</p> <p><i>SSQ 3-9. How do varying flows positively or negatively affect campsite attributes that are important to visitor experience?</i></p> <p>CMIN 9.3.1. Determine and track the size, quality, and distribution of camping beaches by reach and stage level in Glen and Grand Canyons.</p>	<p>FY2007–11: Monitor change in sandbar campable area, topography, and volume (see above, project linked to sandbar monitoring)</p>		<p>FY2007–08: Complete campsite inventory and Geographic Information Systems (GIS) atlas</p> <p>FY2007–08: Evaluate use of field data vs. remotely sensed data for campable area monitoring</p>
9. Recreation (B)	<p>AMWG Priority: 3</p> <p><i>SSQ 3-7. How do dam-controlled flows affect visitors’ recreational experiences, and what is/are the optimal flows for maintaining a high-quality recreational experience in the CRE?</i></p> <p><i>SSQ 3-8. What are the drivers for recreational experiences in the CRE, and how important are flows relative to other drivers in shaping recreational experience outcomes?</i></p> <p><i>SSQ 3-10. How can safety and navigability be reliably measured relative to flows?</i></p> <p><i>SSQ 3-11. How do varying flows positively or negatively affect visitor safety, health and navigability of the rapids?</i></p> <p><i>SSQ 3-12. How do varying flows positively or negatively affect group encounter rates, campsite competition, and other social parameters that are known to be important variables of visitor experience?</i></p>			<p>FY2008: Compile and analyze existing safety data</p>
10. Hydropower	<p>AMWG Priority: 3</p> <p><i>SSQ 3-3. What are annual hydropower replacement costs of the modified low fluctuating flow (MLFF) since 1996?</i></p> <p><i>SSQ 3-4. What are the projected hydropower costs associated with the various alternative flow regimes being discussed for future experimental science (as defined in the next phase of experimental design)?</i></p> <p>CMIN 10.1.1. Determine and track the marketable capacity and energy produced through dam operations in relation to the various release scenarios (daily fluctuation limit, upramp and downramp limits, maximum flow limit of 25,000 cfs minimum flow limit of 5,000 cfs).</p>	<p>FY2007–11: Monitor power generation and market values under current and future dam operations</p>		

Table 1. Summary of core-monitoring, research and development, and experimental activities in the fiscal year 2008 (FY2008) annual work plan for the Grand Canyon Monitoring and Research Center (GCMRC). Several long-term experimental options currently under discussion are not reflected in the table; additional experimental options will be developed upon approval by the U.S. Department of the Interior. Activities address Glen Canyon Dam Adaptive Management Program (GCDAMP) goals 1–12 in relation to science questions and information needs. Priority and related strategic science questions are paraphrased from the Draft GCMRC Strategic Science Plan (appendix A). Information needs are paraphrased from the GCDAMP Strategic Plan. Abbreviations are as follows: SSQ = strategic science question, CMIN = core-monitoring information need, RIN = research information need, and SA = GCDAMP Science Advisors summary questions.—Continued

GCDAMP goal	Priority science questions and information needs (Questions from Strategic Science Plan and Monitoring and Research Plan in italics)	Core-monitoring activities	Experimental activities	Research and development activities
<p>11. Cultural</p>	<p>AMWG Priority:2, 3, and 4</p> <p><i>SSQ 2-1. Do dam-controlled flows affect (increase or decrease) rates of erosion and vegetation growth at archaeological sites and TCP sites in the CRE, and if so, how?</i></p> <p><i>SSQ 2-4. How effective are various treatments (e.g., check dams, vegetation management, etc.) in slowing rates of erosion at archaeological sites over the long term?</i></p> <p><i>SSQ 2-7. Are dam-controlled flows affecting TCPs and other tribally-valued resources, and if so, in what respects?</i></p> <p>CMIN 11.1.1 Determine the condition and integrity of archaeological sites and TCPs in the CRE through tracking rates of erosion, visitor impacts, and other relevant variables. (SPG revised CMIN)</p> <p>CMIN 11:2.1 Determine the condition of traditionally important resources and locations using tribal perspectives and values. (SPG revised CMIN)</p>			<p>FY2008: Research and development towards core monitoring (development of protocols for archaeological sites and TCPs)</p> <p>FY2008: Implement Technical Work Group (TWG) approved tribal monitoring projects</p>
<p>12. High-quality monitoring, research, and Adaptive Management Program</p> <p>(A.) Data acquisition, storage, and analysis</p>	<p>AMWG Priority: 1,2, 3, 4, and 5</p>	<p>FY2007–11: Remote-sensing activities related to the preparation, acquisition, and storage of 2009 terrestrial resource monitoring data</p>	<p>No projects</p>	<p>FY2007–11: Convert existing analog images (especially overflight imagery) and reports to digital (see also goal 8)</p> <p>FY2007–11: Shoreline habitat and change detection mapping (see goals 2 and 8)</p>

There are a number of projects and activities associated with GCDAMP goal 12—the maintenance of a high-quality monitoring, research, and adaptive management program—presented in this annual work plan. In general, these activities are aimed at effective management and administration of the GCMRC science program, logistical support for field activities, data management and analysis, independent peer review, and developing an action plan to improve the effectiveness of the GCDAMP. These support activities fall into eight categories:

1. Data acquisition, storage, and analysis
 - Acquire remote-sensing data (bank funding for FY2009 acquisition)
 - Maintain, update, and enhance Oracle database
 - Convert analogue data (report and imagery) to digital format
 - Provision of Geographic Information Systems (GIS) support
 - Support library functions
 - Map shoreline habitat changes over a 5-year period
 - Survey operations support
2. Logistical support for field activities/river trips
3. Development of work plan for enhancing the CEM
4. Workshop to develop an action plan for improving GCDAMP effectiveness
5. Administrative support for the GCMRC
6. GCMRC program planning and management
7. Independent peer review and science advisor support
 - Provide followup activities for the GCDAMP effectiveness workshop
 - Supply risk assessment of LTEP alternatives
 - Review/assess integrated, interdisciplinary science approaches
 - Organize 2008 science symposium
8. Secure Southwest Biological Science Center (SBSC) information technology (IT) support

A summary of the anticipated FY2008 funding by funding source is provided in table 2 and figure 2 summarizes GCMRC's FY2008 budget by GCDAMP goal. A breakout of the projects included as part of goal 12 is summarized in figure 3. The budget for each project in the work plan is included in the project descriptions and summarized for the entire budget in the separate budget attachment.

Table 2. Total anticipated funding to support the GCMRC in fiscal year 2008 (FY2008).

FUNDING SOURCES	FISCAL YEAR 2008
Power Revenues Under Cap—Estimated USGS Portion(1)	\$7,532,825
USGS Appropriations—Assistance with Burden Costs (Cost Share)	\$1,000,000
Reclamation Operations and Maintenance (Water Quality Monitoring of Lake Powell and Tailwaters Agreement)	\$226,148
TOTAL ESTIMATED AVAILABLE FUNDS FOR FY2008:	\$8,758,973

Figure 2. Budget comparison of Grand Canyon Monitoring and Research Center fiscal year (FY) 2007 approved budget and FY2008 preliminary budget by Glen Canyon Dam Adaptive Management Program goal.





Figure 3. A breakout of the projects included as part of goal 12.



Project Descriptions

Detailed descriptions of each activity included in the AWP are described in the following section. Activities are presented based on the GCDAMP goal they are designed to address. Activities included in the AWP will be carried out in an integrated, interdisciplinary fashion. Integration efforts are described as an element of each description below.

Since its inception, the GCDAMP has attempted to ensure appropriate science program continuity and balance across all goals adopted by the program. The current focus of the GCDAMP is on SSQs associated with high priority AMWG information needs. Other GCDAMP goals will still be pursued, but with less intensity until priority issues of concern are resolved and monies can be reprogrammed or obtained through alternative sources. The AWP, with the exception of GCDAMP goal 3 (restore extirpated species), includes at least one activity to address each GCDAMP goal.

GCDAMP Goal 1: Protect or improve the aquatic food base so that it will support viable populations of desired species at higher trophic levels.

BIO 1.R1.08: Aquatic Food Base

Start Date

September 2005

End Date

September 2009

Principal Investigator(s)

Robert Hall, Ph.D., Aquatic Biologist, University of Wyoming; Emma Rosi-Marshall, Ph.D., Aquatic Biologist, Loyola University, Chicago; Colden Baxter, Ph.D., Fisheries Biologist, Idaho State University; and Theodore Kennedy, Ph.D., Aquatic Biologist, Grand Canyon Monitoring and Research Center

Geographic Scope

Systemwide with monthly sampling at accessible sites (Glen Canyon, about river mile (RM) -15-0, and Diamond Creek, about RM 225) and quarterly sampling at less accessible sites (Marble Canyon, about RM 30; below Little Colorado River (LCR) confluence, about RM 61; Randy's Rock, about RM 126; and below Havasu Creek, about RM 163). Three of these sites are known HBC aggregations.

Project Goals/Tasks

The overall goal of this project is to determine the role that food is playing in the distribution, condition, and abundance of fishes throughout the entire system. Quantifying the density and production of basal resources (i.e., algae, terrestrial leaf litter, etc.) and invertebrates will determine the amount of energy that is available to support production of fishes. The trophic basis of production calculations, where the types and amounts of different food items eaten by invertebrates and fishes are quantified, will determine the relative contribution of basal resources, invertebrates, and other food items to fish production. The results of this work will establish the degree to which native fishes are limited by food resources, by either low production at the base of the food web or via shunting of energy to nonnative animals such as New Zealand mudsnails or rainbow trout (RBT). This information, in turn, provides guidance to managers considering various management options.

The objectives that are addressed by this project include

- determining the important energy sources and pathways that support fishes, especially native species and trout,
- quantifying the abundance of basal resources using a carbon budget framework to determine potential available energy for higher trophic levels,
- identifying composition and quantity of drifting organic matter and invertebrates,
- incorporating knowledge into bioenergetics model and trophic basis of production calculations, and

- developing core-monitoring strategies for the aquatic food base in the Colorado River from GCD to Diamond Creek.

Need for Project

The aquatic PEP (Anders and others, 2001) and Science Advisor (Palmer, 2004) review of food base monitoring and research both recommended major changes in the GCMRC food base program. Specifically, Anders and others (2001) made the following remarks and recommendations:

The food base program needs to be critically reviewed because the current level of understanding about the linkages between lower trophic levels and food availability of native fishes is not adequate to interpret food base data in relation to the management goal.

Since there are scientific as well as statistical uncertainties associated with any approach for study[ing] the relation of food base to trends in abundance of fish populations the best approach is likely a fully integrated one, utilizing data on the abundance of prey available to fish in the GCE, the apparent food habits as indicated by stomach content analysis, and indicators from the fish themselves, including isotopes, growth and condition, and body composition.

Because the food habits of specific life stages of most native species are not well known, a broad look at the potentially available food is required for a monitoring program. The best indicator of potential energy available is a measure of production—both primary and secondary—which is a measure of organic matter creation over time (mass/area/time).

These recommendations formed the basis for the food base request for proposals (RFP) released by the GCMRC in May 2005. The research proposal submitted by Dr. Hall and others that was awarded a cooperative agreement by the GCMRC closely followed the recommendations laid out in the PEP and SA reviews and the food base RFP. The GCMRC continues to lead and monitor the project progress.

Strategic Science Questions

Primary SSQ addressed:

SSQ 1-5. What are the important pathways, and the rate of flux among them, that link lower trophic levels with fish and how will they link to dam operations?

Additional SSQs addressed:

SSQ 1-6. Are trends in the abundance of fish populations, or indicators from fish such as growth, condition, and body composition (e.g., lipids), correlated with patterns in invertebrate flux?

SSQ 5-2. How is invertebrate flux affected by water quality (e.g., temperature, nutrient concentrations, turbidity) and dam operations?

Links/Relationships to Other Projects

Physical Sciences

Four of our six study reaches are fine-grained integrated sediment transport (FIST) and integrated water-quality (IWQ) monitoring sites. We will use bathymetry, bed classification, sediment transport, and water quality data to determine how the physical environment affects the standing mass, distribution, and production of basal resources and invertebrates. We will work closely with the Physical Science and Modeling Program, relying on their infrastructure and capabilities, to estimate inputs of organic matter from the Paria River during base flow and flooding events. Finally, the temperature model that is being developed by the Physical Science and Modeling

Program will be a valuable tool for estimating systemwide growth rates of algae and invertebrates because temperature is an important determinant of algae and invertebrate growth rates.

Fisheries

Ongoing fisheries monitoring data on the distribution and relative density of common native and nonnative fishes will be used to determine rates of energy flow to fishes in the system. Where possible, we will also rely on existing fisheries monitoring efforts to obtain the fish stomachs and tissue samples required for gut content and stable isotope analysis, respectively.

Terrestrial Resources

Ongoing vegetation mapping efforts will be used to estimate rates of allochthonous inputs to the mainstem Colorado River, a potentially significant basal resource supporting invertebrate and fish growth.

Information Needs Addressed

This project focuses on quantifying food availability, and determining which food resources are most important to invertebrates and fishes, in the CRE in Glen and Grand Canyons. The distribution of multiple sampling sites over multiple years will allow a number of research information and core-monitoring information needs to be directly addressed, as enumerated below in the research information needs (RIN) and core-monitoring information needs (CMIN).

Primary information needs addressed:

RIN: 1.1. What are the fundamental trophic interactions in the aquatic ecosystem?

RIN 1.4. What is the current carbon budget for the Colorado River ecosystem?

Other information needs addressed:

CMIN 1.2.1. Determine and track the composition and biomass of benthic organisms between Glen Canyon Dam and the Paria River in conjunction with measurements of flow, nutrients, water temperature, and light regime.

CMIN 1.3.1. Determine and track the composition and biomass of primary producers below the Paria River.

General Methods

Quantify Basal Resources Using a Carbon Budget Framework

That is, quantify inputs, standing stock, and transport of organic matter throughout the river. (RIN 1.4)

Primary production and ecosystem respiration will be quantified using whole-tream metabolism calculations: Use diel changes in dissolved oxygen concentration, a byproduct of algal photosynthesis, to determine rates of algae production for mile-long reaches of the river. Use nighttime sags in dissolved oxygen concentration to determine ecosystem respiration, a measure of basal resource (both leaf litter and algae) consumption. If quantity of carbon consumed during respiration exceeds quantity of carbon produced by algal photosynthesis, this indicates allochthonous inputs may be an important basal resource fueling the aquatic food web. Data collected monthly at Glen Canyon and Diamond Creek and four times per year along the river corridor.

Allochthonous Inputs

Allochthonous inputs originate from riparian vegetation, tributaries, and Lake Powell. Allochthonous inputs from riparian vegetation have been quantified by Ralston and Kennedy (unpub. data). Use ISCO automated water samplers (only at Paria River and LCR) to collect samples of fine organic matter during flooding events. We will

also sample coarse organic matter on the Paria River during flooding events using large plankton nets. Collections also occur monthly on the Paria River and four times per year at major downstream tributaries. Water samples and plankton nets will be used to quantify the concentration of dissolved nutrients, dissolved organic matter, and plankton coming from Lake Powell. Samples will be collected monthly.

Standing Stocks

The standing stock of algae and organic matter will be quantified using a Hess sampler and by scraping algae off rocks. These data will provide a measure of basal resource availability within each reach. Collections will occur monthly at Glen Canyon and Diamond Creek and four times per year at downstream locations.

Transported Organic Matter and Invertebrates

The amount of organic matter and invertebrates transported into and out of each reach will determine the extent to which downstream reaches are linked to upstream processes. Depth-integrated water samples will be used to quantify transported organic matter and invertebrates.

Determine Important Trophic Pathways Linking Basal Resources with Fishes

Stable isotope and diet analysis of invertebrates and fish. Collect diet information from gut content studies of invertebrates and fishes. Collect standards of food items (e.g., algae, benthic invertebrates, terrestrial invertebrates) for signatures for use in stable isotope analysis. Samples collected four times per year along the river corridor.

Determine Flux along Trophic Pathways

Invertebrate density, production, and growth measurements. Sample all benthic habitats (i.e., cobble bars, cliff faces, boulders, talus slopes, sandy bottom, etc.) to quantify density of invertebrates. Habitat-specific density estimates will be made using shoreline and bed-classification data from the Physical Science and Modeling Program. Growth measurements for the most common invertebrates (e.g., New Zealand mudsnails, Gammarus, chironomids, simuliids) in controlled chambers. Production of invertebrates will be calculated using density estimates coupled with growth measurements. Invertebrate density will be estimated monthly at Glen Canyon and Diamond Creek and four times per year at downstream locations. Growth measurements will be taken four times per year at Glen Canyon and Diamond Creek.

Fish density and production estimates. Density estimates for small-bodied and juvenile fishes will be determined quarterly using the multipass depletion method. Density estimates for larger bodied fishes will be derived using existing fisheries monitoring data. Production estimates will be attempted using existing fisheries data and literature values.

Bioenergetics modeling and trophic basis of production calculations. Invertebrate and fish production data will be coupled with diet information (derived from both gut content and stable isotope analysis) to determine the relative contribution of basal resources to invertebrate and fish production.

Products/Reports

Publications

We anticipate at least six publications in peer-reviewed journals will be produced during this project. Tentative subjects for these publications include:

- measuring air-water gas exchange and whole-system metabolism in a large, regulated river (proof-of-concept paper);

- assessing the seasonal and spatial variation in organic matter inputs to the Colorado River, Grand Canyon (synthesis paper of metabolism, allochthonous inputs, lake inputs, tributary inputs, etc.);
- determining spatial variation of secondary production of invertebrates in the Colorado River;
- analyzing the spatial variation in the relative importance of basal resources to invertebrate and fish production in the Colorado River;
- linking whole-river carbon flows with food webs in the Colorado River; and
- determining impacts of New Zealand mudsnails on invertebrate and fish production in the Colorado River.

Reports

A final report summarizing major results and recommendations will be submitted by December 2009.

Monitoring Protocols

A report describing potential monitoring protocols will be submitted at the close of the project. Some potential monitoring tools that will be evaluated during the course of the project include measurement of primary production and ecosystem respiration using whole stream metabolism methods, production measurements of significant invertebrate taxa (e.g., Gammarus, simuliids, and New Zealand mudsnails), fish diet analysis, and organic and invertebrate drift measurements.

Budget

BIO 1.R1.08	
Aquatic Food Base (FY2007–09)	
	Fiscal year 2008
GCMRC personnel costs (19% burden)	114,845
GCMRC project-related travel/training (19% burden)	3,000
GCMRC operations/supplies (19% burden)	3,010
GCMRC equipment purchase/replacement (19% burden)	6,200
AMP logistical support (19% burden)	112,000
Outside GCMRC and contract science labor (19% and/or other burden rate)	—
Cooperative/interagency agreements (6.09% GCMRC burden plus cooperator's burden)	216,000
Project Subtotal	455,055
DOI customer burden (combined 6.09%, 19% and/or other rates)	58,575
Project Total (Gross)	513,630
Percent outsourced (outside of GCMRC; includes 50% of logistics)	60%

BIO 1.R3.08: Diet, Drift, and Predation Data Analysis

Start Date

February 2007

End Date

March 2008

Principal Investigator(s)

Colden Baxter, Ph.D., Fisheries Biologist, Idaho State University; and Michael Yard, Ph.D., Fisheries Biologist, Idaho State University

Geographic Scope

The mechanical removal reaches of the mainstem Colorado River, RM 50–70

Project Goals/Tasks

This project uses diet and drifting organic matter data collected in 2002 and 2003 to (1) determine the important energy sources and pathways that support fishes, especially trout, (2) identify variable food availability in the drift (flux) along trophic pathways, (3) incorporate knowledge into bioenergetics model and trophic basis for production calculations, and (4) document primary production and drift of fish food items in response to varying flow regimens.

Rainbow and Brown Trout Diet Analysis

The purpose of this study was to describe quantitatively the diet proportions (density and weight) of RBT and brown trout, and determine if biotic and environmental factors influenced food resource use patterns. Objectives were to determine if there were differential use of prey items (fish and invertebrates), item sizes, and abundance (biomass and density). Additional objectives were to determine if differential use of food resources were due to interactions from differences between biotic and physical factors. These factors included differences among prey and predator densities, predator and prey size differences, food resource availability, flow discharge, and suspended sediment loads. Diet analysis was also to include specific indexes representing electivity, diet overlap, and diet breadth between RBT and brown trout.

Food Resource Availability

Drift samples collected concurrent with fish removal efforts provided a means for determining density and biomass estimates of prey items available to foraging fish. Results from this analysis were to be used as part of food electivity which characterized food resource use in relation to availability. Other study objectives were to determine if food resource availability differed spatially and/or temporally due to variability in seasonal production, flow discharge, and sediment discharge.

Incidence of Piscivory

The primary goal of this study was to better understand fish interactions occurring among different environmental factors that potentially contribute to predatory behavior within and among different fish species, sampling periods, and spatial strata. The biotic factors include differences in prey and predator densities, predator and prey size-classes, and food resource availability. Physical factors include differences in flow, water clarity, and temperature.

To assess and account for the separate effects associated with mechanical treatments, as well as the natural variability occurring in the ecosystem, large sample sizes were required to determine if diet composition and mean incidence of predation varied significantly among sampling trips, seasons, and years.

Specific tasks completed to date are field work, sample enumeration and biomass determination, and data entry. However, these data have not been assessed for data omission, data entry errors, or data completely compiled into a database design. Only preliminary analysis has been conducted to date and results have not been documented in the form of reports or manuscripts. Therefore, this proposal identifies separate tasks required to complete each of the three study projects. A sequential order is suggested for completing each of the necessary tasks, specific to each project. Tasks will include database development, data entry, literature search, data analysis, manuscript development, and documentation of metadata (see methods, below).

Need for Project

Over the past two decades, research has been directed toward understanding causal mechanisms limiting the phytoplankton community (aquatic food base), and, more recently, monitoring these resource trends in the CRE (Blinn and others, 1995; Shaver and others, 1997; Benenati and others, 1998). Although this bottom-up perspective has provided greater understanding of resource availability, very little dietary use information is known (although often presumed) (Maddux, 1987; McKinney and others, 1999) regarding the utilization of different food resources by the higher trophic levels (Shannon and others, 2001). In this ecosystem the importance of aquatic food resources has been implicitly recognized; however, it remains uncertain whether or not the availability of aquatic as well as terrestrial invertebrates are spatially and/or temporally limited in their availability to higher trophic levels.

Interactions with nonnative fish are implicated in the decline and extinction of native fishes throughout the Colorado River Basin (Tyus and Saunders III, 2000). The cumulative effect from piscivory is known to structure fish communities, especially species that have been compromised by changes in habitat and demographic characteristics that result in low abundance and recruitment levels (e.g., HBC). While it is difficult to determine what is the primary factor most responsible for the decline in HBC recruitment (Coggins and others, 2006), negative interactions (predation and competition) with nonnative coldwater salmonids such as RBT and brown trout are one possible factor that is scientifically testable.

An experimental manipulation was used to test the nonnative fish predation/competition hypothesis. Trout were mechanically removed from selected reaches near the LCR inflow area. This mechanical removal study had multiple study projects. Over a 2-year period (2003–04), approximately 16,000 fish were caught and assessed for the incidence of predation. Diet and drifting organic matter were both sampled. Sampling design, field collection, processing, and preservation methods used are explained in greater detail by Coggins and others (2002).

This proposal has been specifically developed to provide a 1-year approach that completes study projects that were designed to assess nonnative fish diet utilization and food resource availability to provide a better understanding of predatory and possible competitive interactions with native fishes.

Strategic Science Questions

Primary SSQs addressed:

SSQ 1-5. What are the important pathways, and the rate of flux among them, that link lower trophic levels with fish and how will they link to dam operations?

SSQ 1-6. Are trends in the abundance of fish populations, or indicators from fish such as growth, condition, and body composition (e.g., lipids), correlated with patterns in invertebrate flux?

Links/Relationships to Other Projects

How the available aquatic food base is utilized by fishes is important for managers to understand as they consider various flow regimens. A more complete understanding of fish diets will help managers decide what primary and secondary production should be targeted as management scenarios are considered. Results of this project will support management of the RBT population as a sport species below GCD and as a predator/competitor in Grand Canyon.

Information Needs Addressed

Primary information need addressed:

RIN 1.1. What are the fundamental trophic interactions in the aquatic ecosystem?

Diet analysis of rainbow and brown trout will provide a comprehensive look at what fish are eating in the system.

Other information need addressed:

RIN 1.5.3. How has the value and availability of drift as a food source for humpback chub changed with the implementation of Record of Decision operations?

Drift samples will be compared with data from Valdez and Ryel (1995) also collected at the LCR confluence, to determine whether value and availability has changed with implementation of the Record of Decision (ROD).

General Methods

Database Development

Presently these data exist as a series of separate files (Microsoft® Excel) found in spreadsheet form. These files are currently archived at the GCMRC. Data contained in spreadsheets need to be imported into a common database (Microsoft® Access), and relationally linked to other field collection data (locality, sampling period, and sample bottle number). These data need to be checked for data entry errors, duplications, relational links, and omissions. Data omissions will be determined by conducting a series of cross comparisons with sample bottle numbers against common fields in the GCMRC fish database containing data from the mechanical trout removal study. This linkage is critical in relating specific data (species, size, sex, location, and date) to stomach contents. Identified errors will be resolved by reentry of data from original data sheets.

Data Entry

Preliminary assessment of data entry efforts for the incidence of piscivory indicates that data entry is only partially complete (60 percent) for this project. This will require determining which sample data are missing for specific sampling periods, locating the appropriate data sheets, and entering the data. Estimated time required for this task is identified in the summary budget.

Literature Search

This project will be initiated with a comprehensive review of the most current literature. The time and costs required for conducting the search, review process, and photocopying appropriate publications are identified in the summary budget under other direct costs.

Data Analysis

The selected contractor will have limited use of available statistical software (SAS, Inc.) currently licensed to the GCMRC to conduct appropriate statistical analysis. This approach will result in a net savings to the GCMRC because the purchase of additional software will not be necessary.

Products/Reports

Draft Manuscript Development

A draft manuscript will be developed and subjected to peer review in April 2008. The budget includes costs to prepare the submittal draft and to modify the report in accordance with reviewers' comments.

Metadata Completion

Final data will be transferred to the GCMRC in Microsoft® Access database structure. Documentation of field collection methodologies and analysis will be developed as well as information concerning data fields. These are to be provided to the GCMRC as specified in their standard metadata format structure.

Budget

BIO 1.R3.08	
Diet, Drift, and Predation Data Analysis (FY2007)	
	Fiscal year 2008
GCMRC personnel costs (19% burden)	—
GCMRC project-related travel/training (19% burden)	—
GCMRC operations/supplies (19% burden)	—
GCMRC equipment purchase/replacement (19% burden)	—
AMP logistical support (19% burden)	—
Outside GCMRC and contract science labor (19% and/or other burden rate)	—
Cooperative/interagency agreements (6.09% GCMRC burden plus cooperator's burden)	—
Project Subtotal	—
DOI customer burden (combined 6.09%, 19% and/or other rates)	—
Project Total (Gross)	—
Percent outsourced (outside of GCMRC; includes 50% of logistics)	0%

*No funding needed in FY2008

BIO 1.R4.08: Impacts of Various Flow Regimes on the Aquatic Food Base

Start Date

2008

End Date

2010

Principal Investigator(s)

Theodore Kennedy, Ph.D., Aquatic Biologist, USGS Grand Canyon Monitoring and Research Center

Geographic Scope

Three sites (Glen Canyon about RM -15-0, Diamond Creek about RM 225, and LCR confluence about RM 61)

Project Goals/Tasks

This project will be done in close association with research project BIO 1.R1.08, which will quantify, on a monthly basis, the density and production of basal resources (i.e., algae, terrestrial leaf litter, etc.) and invertebrates, and will determine the amount of energy that is available to support production of fishes (e.g., monitoring). In addition, short-term experiments should be conducted to obtain a greater understanding of the responses of the aquatic food base to specific aspects of different flow regimes. The primary goal of such experiments would be to verify the range of flow magnitudes and flow fluctuations that could occur without significantly impacting the long-term sustainability of the aquatic food base. This would, in turn, help identify ways to accommodate power generation and water storage purposes at GCD while protecting downstream aquatic resources.

Short-term experiments would be conducted over the course of several years in order to capitalize on the availability of specific hydrological conditions. During experiments, measurements would be made of algal and invertebrate standing crop, together with levels of invertebrate drift. To the extent possible, data collection methods should be consistent with the methods described in BIO 1.R1.08. Aspects of hydrographs that should be examined include fluctuation levels, ramp rates, and minimum flow levels. In addition, it may be desirable to evaluate effects of the proposed flow regime on fish (especially trout) activity levels, foraging success, and survival as related to fluctuation patterns and food conditions (e.g., concentrations and composition of the drift).

Benthic production is the source for food base items in the drift, so it is important to assess both how the source of drifting material and the quantity of drifting material is affected by dam operations. These measurements should be made in three locations, Glen Canyon, Diamond Creek, and the confluence with the LCR. The effects of dam operations on benthic production and drift are likely very different in the tailwater relative to downstream reaches. The food base program (BIO 1.R1.08) will continue to make monthly measurements of benthic production and drift at Glen Canyon and Diamond Creek, which will provide an excellent set of baseline data that can be used to compare results from experimental dam operations. The LCR confluence should be included in addition to the sampling at Glen Canyon and Diamond Creek because of its relevance to the availability of food for HBC. Adding sampling at the LCR confluence will significantly increase the cost of this research because it will require three full sets of sampling equipment and a downstream river trip.

The objectives that are addressed by this project include (1) quantifying the abundance of the aquatic food base on substrates in response to changes in the flow regime; (2) identifying the composition and quantity of drifting organic matter and invertebrates in response to changes in the flow regime; (3) determining the effects of ramp rates on invertebrate productivity, standing crop, and drift; (4) determining the effects on drift of short-term 5,000-cfs flow fluctuations during a fall steady flow experiment; and (5) determining the effects of short-term flow reductions and subsequent (e.g., 2 hours later) increase in flow on drift rates.

Need for Project

The food base in any aquatic system is an important factor that directly affects fish community dynamics including abundance, reproduction and recruitment, condition, and even distribution. Much of the diet of trout and HBC consists of food items that have been suspended and are drifting in the water column (Valdez and Ryel 1995). The drifting food base in the CRE is generally composed of freely floating aquatic invertebrates and *Cladophora glomerata* (a long filamentous green algae) that are available to fish for consumption. Primary production at Lees Ferry is dominated by *Cladophora*, which acts as a substrate for various types of epiphytic diatoms which provide a food source for chironomids and simuliids (aquatic insect larvae) and for the shrimp-like amphipod, *Gammarus lacustris* (Pinney 1991). The nutritional value of *Cladophora* to fish is enhanced by the presence of lipid-rich epiphytic diatoms, and diatoms have been shown to provide an important source of energy for RBT (Leibfried, 1988). Studies indicate that the abundance of invertebrates used as food by trout and other fishes in the CRE is generally proportional to the abundance of *Cladophora*.

In order to understand the current condition of the aquatic food base, measurements of epiphytic diatoms, aquatic invertebrates, and algal abundance in the Colorado River downstream of GCD should be conducted. The GCMRC's food base monitoring program (BIO 1.R1.08) would largely fill this need on a monthly basis. However, the response of these benthic and drifting resources to various flow management regimes remains uncertain. Thus, this research project should be conducted to identify the responses of the benthic and drifting food base to various aspects of the proposed flow regime. This adds an important component to the food base research program under BIO 1.R1.08 which may help to identify flow regimes likely to contribute to the recovery of HBC populations in Grand Canyon.

Strategic Science Questions

Primary SSQ addressed:

SSQ 1-5. What are the important pathways, and the rate of flux among them, that link lower trophic levels with fish and how will they link to dam operations?

Additional SSQs addressed:

SSQ 3-5. How is invertebrate flux affected by water quality (e.g., temperature, nutrient concentrations, turbidity) and dam operations?

SSQ 1-6. Are trends in the abundance of fish populations, or indicators from fish such as growth, condition, and body composition (e.g., lipids), correlated with patterns in invertebrate flux?

Links/Relationships to Other Projects

Under Research Project BIO 1.R1.08—Aquatic Food Base four broad tasks would be performed: (1) quantify basal resources using a carbon budget framework, (2) determine important trophic pathways linking basal resources to fish, (3) estimate fish density and production, and (4) model bioenergetics and the trophic basis of production calculations. We will work closely with this project, relying on much of their infrastructure and capabilities, to estimate primary and secondary biomass, productivity, and drift. This project builds upon the aquatic food base program by carrying more intensive observations during various experimental flow regimes with the intent of distinguishing the effects of various flow changes compared to “base” conditions.

Information Needs Addressed

This project focuses on quantifying food availability (drift) in the CRE in Glen and Grand Canyons due to experimental changes in flow from GCD. The distribution of multiple sampling sites over multiple years will allow a number of research information and CMINs to be directly addressed, as enumerated below:

Primary information needs addressed:

RIN: 1.1. What are the fundamental trophic interactions in the aquatic ecosystem?

RIN: 2.1.2. Quantify sources of mortality for humpback chub <51 mm in rearing habitats in the LCR and mainstem and how these sources of mortality are related to dam operations.

RIN 12.9.2. What is the best combination of dam operations and other management actions to achieve the vision, mission, goals, and objectives of the GCDAMP management objectives?

Other information needs addressed:

RIN 1.4. What is the current carbon budget for the Colorado River ecosystem?

CMIN 1.2.1. Determine and track the composition and biomass of benthic organisms between Glen Canyon Dam and the Paria River in conjunction with measurements of flow, nutrients, water temperature, and light regime.

CMIN 1.3.1. Determine and track the composition and biomass of primary producers below the Paria River.

General Methods

Quantify Algae Production

The food base project has been successful in verifying that algae/macrophyte production in the Colorado River can be measured using open-system measurements. This technique involves constructing an oxygen budget for a reach of river, accounting for all inputs (algae production, air-water gas exchange) and outputs (respiration) of oxygen.

Primary production and ecosystem respiration will be quantified using whole stream metabolism calculations: Use diel changes in dissolved oxygen concentration, a byproduct of algal photosynthesis, to determine rates of algae production for mile-long reaches of the river. Use nighttime sags in dissolved oxygen concentration to determine ecosystem respiration, a measure of basal resource (both leaf litter and algae) consumption. If quantity of carbon consumed during respiration exceeds quantity of carbon produced by algal photosynthesis, this indicates allochthonous inputs may be an important basal resource fueling the aquatic food web. Data collected monthly at Glen Canyon and Diamond Creek and four times per year along the river corridor.

Quantify Benthic Biomass/Drift

Standing Stocks

The standing stock of algae and organic matter will be quantified using a Hess sampler and by scraping algae off rocks. These data will provide a measure of basal resource availability within each reach. Collections will occur monthly at Glen Canyon and Diamond Creek and four times per year at the confluence with the LCR.

Invertebrate Density, Production, and Growth Measurements

Sample all benthic habitats (i.e., cobble bars, cliff faces, boulders, talus slopes, sandy bottom, etc.) to quantify density of invertebrates. Habitat-specific density estimates will be made using shoreline and bed classification data from the Physical Science and Modeling Program. Growth measurements for the most common invertebrates (e.g., New Zealand mudsnails, Gammarus, chironomids, simuliids) in controlled chambers. Production of invertebrates will be calculated using density estimates coupled with growth measurements. Invertebrate density will be estimated monthly at Glen Canyon and Diamond Creek and four times per year at downstream locations. Growth measurements will be taken four times per year at Glen Canyon and Diamond Creek.

Transported Organic Matter and Invertebrates

The amount of organic matter and invertebrates transported into and out of each reach will determine the extent to which downstream reaches are linked to upstream processes. Depth-integrated water samples will be used to quantify transported organic matter and invertebrates.

Products/Reports

Tentative subjects for publications include (1) the response of primary production and secondary production of invertebrates in the Colorado River to various flow regimes from GCD, and (2) the effect of various flow regimes from GCD on the availability of drifting food base for HBC and trout.

Reports

A final report summarizing major results and recommendations will be submitted at the close of the project.

Monitoring Protocols

A report describing potential monitoring protocols will be submitted at the close of the project. Some potential monitoring tools that will be evaluated during the course of the project include (1) measurement of primary production and ecosystem respiration using whole-stream metabolism methods and (2) organic and invertebrate drift measurements.

Budget

We already have four sondes equipped with dissolved oxygen sensors, but would need to purchase two additional sondes in order for these measurements to be made simultaneously at Glen Canyon and Diamond Creek. The Clark-type dissolved oxygen sensors that we are currently using for making these measurements can be calibrated with a good degree of precision and accuracy, but the calibration of those sensors is only reliable for 5–7 days because the sensors tend to “drift.” Optical dissolved oxygen sensors are a relatively new technology that can be calibrated more accurately and precisely than Clark-type sensors, and they hold their calibration for over a month. With the purchase of optical dissolved oxygen sensors, we will be able to make continuous, and more accurate and precise, measurements of algae production for the duration of any flow experiments.

We already have all of the equipment needed to make measurements of benthic biomass (i.e., algae, invertebrates, terrestrial leaf litter), but we only have one set of this equipment. We will need to purchase an additional set of equipment so that these measurements can be made simultaneously at both Glen Canyon and Diamond Creek. We already have enough wet suits and winches (needed for ponar dredge) to accommodate simultaneous sampling at two or three locations.

Budget

BIO 1.R4.08	
Impacts of Various Flow Regimes on the Aquatic Food Base (FY2008–09)	
	Fiscal year 2008
GCMRC personnel costs (19% burden)	—
GCMRC project-related travel/training (19% burden)	—
GCMRC operations/supplies (19% burden)	14,096
GCMRC equipment purchase/replacement (19% burden)	19,600
AMP logistical support (19% burden)	—
Outside GCMRC and contract science labor (19% and/or other burden rate)	—
Cooperative/interagency agreements (6.09% GCMRC burden plus cooperator's burden)	46,566
Project Subtotal	80,262
DOI customer burden (combined 6.09%, 19% and/or other rates)	9,238
Project Total (Gross)	89,500
Percent outsourced (outside of GCMRC; includes 50% of logistics)	58%

NOTE: \$17,500 of this funding is from FY2007 carry forward recommended by the AMWG August 30, 2007, Motion 5. \$72,700 is funded from the FY2008 GCDAMP power revenue budget.

GCDAMP Goal 2: Maintain or attain viable populations of existing native fish, remove jeopardy from humpback chub and razorback sucker, and prevent adverse modification to their critical habitat.

BIO 2.R1.08: Little Colorado River Humpback Chub Monitoring Lower 15 km (HBC Population Estimates)

BIO 2.R2.08: Little Colorado River Humpback Chub Monitoring Lower 1,200 m

BIO 2.R3.08: Humpback Chub Monitoring Above Chute Falls

Start Date

Ongoing

End Date

Ongoing

Principal Investigator(s)

U.S. Fish and Wildlife Service (BIO 2.R1.08 and BIO 2.R3.08) and the Arizona Game and Fish Department (BIO 2.R2.08), with support from the Grand Canyon Monitoring and Research Center (M.E. Andersen, L.G. Coggins, Ph.D., staff)

Geographic Scope

Little Colorado River

Project Goals/Tasks

- Elucidate critical physical and biotic factors that may be limiting to, or supportive of, the HBC and other native fish populations in Grand Canyon. Seek methods that reduce, eliminate, or control limiting factors.
- Identify habitat characteristics that are most important to all life stages of HBC and seek methods that maintain, and possibly replicate, suitable habitats.
- Determine and refine the most appropriate method(s) for estimating the population size of HBC and other Grand Canyon fishes, including sampling design, gear selection, and development of remote monitoring methods. The method(s) developed and selected should be consistent with the second edition of the Colorado River Endangered Fishes Recovery Goals. (The U.S. Fish and Wildlife Service (USFWS) has scheduled revision of the goals to be initiated in 2007.)
- Improve understanding of dam operations on (young-of-year (YoY) and juvenile HBC survival and habitat use.
- Establish core-monitoring protocols for HBC in Grand Canyon.

The specific goal of the suite of tasks identified in this project description is to provide current evaluations of the HBC population in the LCR. The specific projects that will be conducted in FY2008 are (1) estimating population of HBC in the LCR, (2) monitoring HBC above Chute Falls, and (3) monitoring HBC in lowest 1,200 meters of LCR.

Specific objectives of the projects include:

- obtaining population estimates of HBC ≥ 150 mm and ≥ 200 mm in the lower 15 km of the LCR and in the LCR above Chute Falls;
- providing other information related to physical parameters of the LCR (i.e., temperature and turbidity), length frequency data, community composition, sexual condition and characteristics of native fish (sex, ripe, tuberculate, etc.), frequency of external parasites (i.e., primarily *Lernaea cyprinacea*), and predation; and
- collecting data in support of stock assessment models (e.g., mark-recapture tagging data, length frequency data).

Need for Project

Because the LCR is the primary tributary where young HBC are produced, a rigorous stock assessment of this endangered species is needed to allow managers to assess the condition of the population and its response to management actions. These projects will conduct this assessment in FY2008. Reviews by peer scientists, statistical data analysis, and historical review of existing data will provide the basis for directing how monitoring of HBC will be conducted in future years. A PEP will be convened to address this issue and core-monitoring needs in FY2008.

Strategic Science Questions

Primary SSQ addressed:

SSQ 1-1. To what extent are adult populations of native fish controlled by production of young fish from tributaries, spawning and incubation in the mainstem, survival of YoY and juvenile stages in the mainstem, or by changes in growth and maturation in the adult population as influenced by mainstem conditions?

Additional science question addressed by these projects:

SSQ 1-2. Does a decrease in the abundance of rainbow trout and other cold- and warmwater nonnatives in Marble and eastern Grand Canyons result in an improvement in the recruitment rate of juvenile HBC to the adult population?

GCDAMP SAs have summarized the SSQs with the following question (the projects outlined here specifically address this question, especially their evaluation of annual spawning success):

SA 1. What are the most limiting factors to successful HBC adult recruitment in the mainstem: spawning success, predation on YoY and juveniles, habitat (water, temperature), pathogens, adult maturation, food availability, competition?

Links/Relationships to Other Projects

The HBC is the only remaining member of the genus *Gila* inhabiting the Colorado River between GCD and Grand Wash Cliffs. This species was listed as endangered by the FWS in 1967 and is protected under the ESA of 1973. HBC distribution in Grand Canyon has been characterized as occurring in discrete locations or aggregations (Valdez and Ryel, 1995). Of these nine aggregations (30 Mile, RM 29.8–31.3; LCR inflow, RM 57–65.4; Lava Canyon to Hance, RM 65.7–76.3; Bright Angel Creek Inflow, RM 83.8–92.2; Shinumo Creek Inflow, RM 108.1–108.6; Stephen Aisle, RM 114.9–120.1; Middle Granite Gorge, RM 126.1–129.0; Havasu Creek Inflow, RM 155.8–156.7; and Pumpkin Spring, RM 212.5–213.2), only the LCR inflow is recognized as a population in that it consistently demonstrates some level of successful recruitment (Kaeding and Zimmerman,

1983; Valdez and Ryel, 1995; Gorman and Stone, 1999). The current paradigm is that the remaining eight aggregations exist as a result of either downstream transport of juvenile HBC from the LCR inflow aggregation, or relict fish (30 Mile population) produced in years immediately following construction of GCD (Valdez and Ryel, 1995). However, limited movement between the LCR inflow and both the 30 Mile and Havasu Creek Inflow aggregations has been observed.

Improvement of the status of the HBC will be necessary for the species to be considered for down listing or delisting. The GCDAMP can contribute to an improved status for HBC, thereby decreasing the amount of effort required of the GCDAMP on behalf of this species. The most recent iteration of the recovery goals for this species, now scheduled for review and revision beginning in 2007, required a minimum of 2,100 adults in the Grand Canyon, a steady or increasing trend in the population, and control of environmental threats, among other requirements. One potential element of conservation of HBC in Grand Canyon may be a GCD flow release regimen that supports this species. These flows can be expected to impact many of the elements of the canyon resources, including sediment, cultural resources, and recreation. Therefore, releases that benefit one resource, HBC in this example, must be consistent with conservation of other resources. Conservation of LCR resources, especially water and protection from catastrophic events, whether accomplished through the GCDAMP process or by other means, would be important not only to protecting the spawning HBC population in the LCR but other organisms found there.

Information Needs Addressed

Primary information needs addressed:

CMIN 2.1.2 Determine and track recruitment (identify life stage), abundance, and distribution of HBC in the LCR.

CMIN 2.1.1. Determine and track year-class strength of HBC between 51-150 mm in the LCR and the main channel.

General Methods

Annual Spring (March and April) HBC Abundance Assessments in the Lower 15 km of the LCR

This monitoring effort provides relative abundance assessments of the spawning and resident populations of HBC in the LCR below Chute Falls. It will be conducted concurrent with mainstem sampling to provide a more ideal sampling design in support of model refinement and use and stock assessment. Hoop nets are deployed to capture fishes for this effort. Evaluation of relative trends of other fishes, especially native bluehead suckers and flannelmouth suckers, is a desirable side benefit of this sampling. Relative abundances of nonnative fishes in the LCR are also developed from this sampling.

Annual Fall (September and October) HBC Abundance Assessments in the Lower 15 km of the LCR

This program has been ongoing since 2000 and annually produces assessments of the abundance of HBC >150 mm total length (Coggins and Van Haverbeke, 2001; Van Haverbeke and Coggins, 2003; Van Haverbeke, 2003; Van Haverbeke, 2004). The fall sampling is aimed primarily at providing an estimate of the abundance of subadult fishes rearing in the LCR. These efforts rely on multiple event mark-recapture analysis of PIT tag data to produce abundance estimates using closed population models. In FY2008 these data will be combined with concurrent mainstem sampling (see above) results to support use of the age-structured mark-recapture (ASMR) model to assess HBC population numbers. Two 12-day trips into the LCR are conducted to collect the data used to construct these estimates in the fall (September and October). Sampling is predominantly conducted using hoop nets evenly distributed throughout the lower 15 km of the LCR. Other types of sampling gear are not used in the LCR because they have been shown to be less efficient at capturing HBC >150 mm total length in the LCR.

Annual Spring Relative Abundance Assessment in the Lower 1,200 m of the LCR

This program was established by the Arizona Game and Fish Department (AZGFD) in 1987 and has operated continuously through 2004 with the exception of the years 2000–01 (Ward and Persons, unpub. data). This program annually produces assessments of the relative abundance (i.e., catch per unit effort [CPUE]) of all size classes of HBC, flannelmouth suckers, bluehead suckers, speckled dace, and a host of nonnative fishes in the lower 1,200 m of the LCR. Data is collected during a 30–40-day period in spring (April and May) using hoop nets set in standardized locations distributed throughout the reach. In general, this effort represents the longest and most consistent relative abundance dataset available to infer trends in the LCR HBC population. Importantly, it provides an independent comparison to the mark-recapture-based assessments. The statistical power of this portion of the monitoring program has not yet been assessed, but statistically significant differences in relative abundance are apparent in current data.

Above Chute Falls

Two trips are conducted above Chute Falls in the LCR to initiate a stock assessment program of translocated individuals, and potential offspring. These trips will occur during late May when the LCR discharge is at base flow to provide an annual abundance estimate of HBC within this region. In addition to the annual population estimates, this data can be incorporated into open population models for HBC being developed by the GCMRC. Moreover, because we have and will continue to implant these fish with PIT tags (Biomark, Inc.), it is likely that some individuals will eventually be recaptured in the lower LCR corridor and/or Colorado River, which would increase our knowledge of migration patterns.

During the LCR trip, personnel will reside at the established translocation camp located at 16.2 river kilometers (rkm) on Navajo lands. This camp has an established helicopter landing pad and offers high ground protection from most floods. Baited hoop nets (0.5–0.6-m diameter, 1.0-m length, 6-mm mesh, single 10-cm throat) will be set from shorelines to capture and PIT tag HBC as part of a mark-recapture program to estimate the abundance of individuals ≥ 150 mm in the upper 13.6 km of the LCR.

Personnel will be responsible for fishing baited hoop nets in the LCR corridor above Chute Falls (13.6 rkm), which is the upstream extent of the current downstream LCR monitoring. Approximately 50 hoop nets will be fished throughout this upper reach from 13.6 rkm to 18.0 rkm, with the average spacing between nets approximately 100–150 m. Each hoop net will be positioned in favorable habitat suspected of yielding good catches of HBC. Nets will be repositioned as needed. On average, each hoop net will be checked once every 24 hours. Each net will be baited near its cod end by attaching a nylon mesh bag (30 by 30-cm, 6-mm mesh) containing AquaMax™ Grower 600 for Carnivorous Species (Purina Mills Inc., Brentwood, MO). All captured HBC will be examined for colored elastomer tags and PIT tags. Those individuals not previously PIT tagged, but of sufficient size to be tagged without injury, will be held overnight either offshore in an aerated tank or in the LCR in a secured holding pen to allow time for digestion of any consumed bait, whereby they will be tagged and released.

The overall reach will be broken down into two subreaches and each subreach fished for 3 days. The upper reach designation will be from 18.0 to 15.0 rkm (undesignated point below Blue Spring to first travertine dam above Chute Falls). Currently 18 rkm is the highest point in which HBC have been located above Chute Falls. The lower subreach will extend from 15.0 to 13.6 rkm (first dam above Chute Falls to Lower Atomizer Falls where lower LCR monitoring begins). The lower subreach is relatively small because of the time constraints needed to maneuver around major travertine dams so that we can sufficiently sample the myriad of adult HBC habitats (deep pools, large boulders, etc.) existing within this subreach. In addition to fishing baited hoop nets and PIT tagging HBC as detailed above, personnel will be responsible for:

- measuring and recording the fork and total lengths, sex, sexual condition, and sexual characteristics for all captured native fishes (except speckled dace);
- measuring and recording the total length, sex, and sexual condition of all other captured fish;
- recording the stomach contents of all captured large-bodied nonnative fish except common carp;

- recording the location, shoreline habitat, hydraulic unit, and set and pull time, and map locations for each hoop net set; and
- making daily turbidity with the Hach 2100 turbidimeter, water temperature measurements, and CO₂ using titration.

Management Plan

Once the initial stock assessment has been completed, FWS will draft a management plan that will direct any future management action above Chute Falls. This document will evaluate the benefits or disadvantages of additional translocations and, if possible, provide a trigger for when additional movements of fish should be performed.

Quality Control

Quality control relative to data delivery will be assured through the use of standardized data collecting, recording, and electronic entry procedures. These include use of standardized fish handling protocols, field data collection forms, and computerized data entry routines. Additionally, various automated summary reports of submitted data are being developed to aid in identifying errors in electronic versions of submitted data. Copies of original field data sheets are held by the GCMRC library so that future problems encountered with fish databases may be checked against field data sheets. Electronic copies of data are submitted to the GCMRC on a CD/DVD format. Data must meet the GCMRC's data standards.

Analysis of the Little Colorado River Monitoring Program

The value of four LCR sampling occasions, monitoring above Chute Falls, and monitoring of the lower 1,200 m of the LCR will be included in the 2008 PEP regarding monitoring of the Grand Canyon HBC population.

Products/Reports

The FWS will deliver two trip reports annually, including data collected, to the GCMRC. The trip reports will be summarized and analyzed in a final report delivered to the GCMRC in January of the following year. These reports address the lower 15-km monitoring and the monitoring above Chute Falls. The AZGFD will deliver one annual report on the results of their monitoring of the lower 1,200 m to the GCMRC.

A report addressing the current statistical rigor of the sampling methods in the LCR will be produced by October 2008. Program and external review of the sampling for HBC in the LCR will be conducted in FY2008. Any recommended and accepted monitoring changes will be implemented in FY2009.

Budget

BIO 2.R1.08	
LCR HBC Monitoring Lower 15 km (HBC Population Est; Ongoing)	
	Fiscal year 2008
GCMRC personnel costs (19% burden)	13,328
GCMRC project-related travel/training (19% burden)	—
GCMRC operations/supplies (19% burden)	—
GCMRC equipment purchase/replacement (19% burden)	—
AMP logistical support (19% burden)	19,500
Outside GCMRC and contract science labor (19% and/or other burden rate)	—
Cooperative/interagency agreements (6.09% GCMRC burden plus cooperator's burden)	347,455
Project Subtotal	380,283
DOI customer burden (combined 6.09%, 19% and/or other rates)	27,397
Project Total (Gross)	407,680
Percent outsourced (outside of GCMRC; includes 50% of logistics)	94%

BIO 2.R2.08	
LCR HBC Monitoring Lower 1,200 m; Ongoing)	
	Fiscal year 2008
GCMRC personnel costs (19% burden)	—
GCMRC project-related travel/training (19% burden)	—
GCMRC operations/supplies (19% burden)	5,000
GCMRC equipment purchase/replacement (19% burden)	—
AMP logistical support (19% burden)	8,500
Outside GCMRC and contract science labor (19% and/or other burden rate)	—
Cooperative/interagency agreements (6.09% GCMRC burden plus cooperator's burden)	53,750
Project Subtotal	67,250
DOI customer burden (combined 6.09%, 19% and/or other rates)	5,838
Project Total (Gross)	73,088
Percent outsourced (outside of GCMRC; includes 50% of logistics)	86%

BIO 2.R3.08	
HBC Monitoring Above Chute Falls; Ongoing)	
	Fiscal year 2008
GCMRC personnel costs (19% burden)	2,857
GCMRC project-related travel/training (19% burden)	—
GCMRC operations/supplies (19% burden)	—
GCMRC equipment purchase/replacement (19% burden)	—
AMP logistical support (19% burden)	15,000
Outside GCMRC and contract science labor (19% and/or other burden rate)	—
Cooperative/interagency agreements (6.09% GCMRC burden plus cooperator's burden)	55,050
Project Subtotal	72,907
DOI customer burden (combined 6.09%, 19% and/or other rates)	6,745
Project Total (Gross)	79,652
Percent outsourced (outside of GCMRC; includes 50% of logistics)	86%

BIO 2.R4.08: Monitoring Mainstem Fishes

Start Date

Ongoing

End Date

Ongoing

Principal Investigator(s)

Arizona Game and Fish Department, with support from Grand Canyon Monitoring and Research Center (M.E. Andersen, L.G. Coggins, Ph.D., staff)

Geographic Scope

The mainstem Colorado River in Grand Canyon between Lees Ferry and upper Lake Mead

Project Goals/Tasks

The objectives that are addressed by this project are as follows:

- Determine and refine the most appropriate method(s) for estimating the population size of HBC and other Grand Canyon fishes, including sampling design, gear selection, and development of remote monitoring methods. The method(s) developed and selected should be consistent with the second edition of the Colorado River Endangered Fishes Recovery Goals. (The FWS has scheduled revision of the goals to be initiated in 2007).
- Improve understanding of dam operations on YoY and juvenile HBC survival and habitat use.
- Establish core-monitoring protocols for HBC in Grand Canyon.
- Monitor small-bodied fishes (native and nonnative species) in backwater habitats, in conjunction with monitoring of Kanab ambersnail (KAS) (BIO 5.R1.08).

The goals of this project are to provide status and trend information on the abundance and recruitment of the fish community in Grand Canyon. It is one of the projects that will be the subject of a PEP in FY2009.

Need for Project

Native fish populations in Grand Canyon are key resources of concern influencing decisions on both the operation of GCD and other nonflow-related actions. To inform these decisions, it is imperative that accurate and timely information on the status of fish populations, particularly the endangered HBC, is available to managers. A suite of adaptive experimental management actions are being contemplated to better understand the mechanisms controlling the population dynamics of native fishes, and to identify policies that are consistent with the attainment of management goals. The assessments generated from this project provide a baseline from which to assess the effects of implemented experimental actions. This information is therefore crucial to (1) inform the program as to attainment of identified goals, (2) provide baseline status and trend information to be used as a backdrop to further understand mechanisms controlling native fish population dynamics, and (3) evaluate the efficacy of particular management policies in attaining program goals. The results of this project are potentially useful in assessing changes to Federal ESA listing status of native fishes in the Colorado River.

Strategic Science Questions

Primary SSQ addressed:

SSQ 1-1. To what extent are adult populations of native fish controlled by production of young fish from tributaries, spawning and incubation in the mainstem, survival of YoY and juvenile stages in the mainstem, or by changes in growth and maturation in the adult population as influenced by mainstem conditions?

Additional SSQs addressed:

SSQ 1-4. Can long-term decreases in abundance of rainbow trout in Marble and eastern Grand Canyons be sustained with a reduced level of effort of mechanical removal or will recolonization from tributaries and from downstream and upstream of the removal reach require that mechanical removal be an ongoing management action? This question also applies to future removal programs targeting other nonnative species.

SSQ 1-8. How can native and nonnative fishes best be monitored while minimizing impacts from capture and handling or sampling?

The AWP SAs have articulated the following summary science questions that are addressed by this project:

SA 1. What are the most limiting factors to successful HBC adult recruitment in the mainstem: spawning success, predation on YoY and juveniles, habitat (water, temperature), pathogens, adult maturation, food availability, competition?

SA 2. What are the most probably positive and negative impacts of warming the Colorado River on HBC adults and juveniles?

Links/Relationships to Other Projects

Understanding the factors influencing the dynamics of the Grand Canyon native fish populations, especially the endangered HBC, is important to evaluating the effects of management and conservation activities, especially GCD operations. To discover these factors, a combination of large-scale manipulations (e.g., experimental removal of nonnative fish or long-term implementation of contrasting flow regimes) and smaller scale process-oriented research (e.g., assessment of juvenile fish growth rates under various temperature regimes or availability of particular food items) will likely prove most efficient in determining the key mechanisms regulating native fish populations. In each of these endeavors, it is critical that baseline trends in population abundance and recruitment be known. It is only with this knowledge that it is possible to assess population level impacts of large-scale manipulations. Though it is informative to assess the effects of experimental management on processes thought to be important like growth or survival at particular life stages, this is not ultimately sufficient to determine efficacy of particular management actions. Linkages between these processes and ultimate recruitment to populations must be established. Again, these linkages can only be made if baseline trends in population abundance and recruitment are available.

Information Needs Addressed

Primary information needs addressed:

CMIN 2.1.2. Determine and track year abundance and distribution of all size classes of HBC between in the LCR and the mainstem.

RIN 2.4.2. Determine if suppression of nonnative predators and competitors increases native fish populations.

The mainstem sampling described in this project description will provide an evaluation of the trend of HBC abundance, especially those greater than 150 mm, through calculation of CPUE. HBC who are identified by a PIT tag provide catch data to the ASMR model, further supporting evaluation of abundance for this species. Mainstem hoop net sampling, shown to be of value for assessing catch rates of HBC, especially those less than 150 mm, during the mechanical removal project (2003–06) will be employed to help address science questions regarding success or failure of HBC to recruit in the mainstem. It will be valuable to compare the results of mainstem sampling for smaller size classes to the same results from the LCR for evaluation of year-class survivorship in the mainstem.

General Methods

Mainstem fish monitoring, including the monitoring below Diamond Creek, has used electrofishing to provide an overall view of the status and trends of native and nonnative fishes in the Colorado River between Lees Ferry and Lake Mead. The electrofishing gear is not without its limitations, particularly its lack of effectiveness at sampling deep water habitats. However, it remains the most important tool for providing an overall assessment of the mainstem fish community and its use will be retained in FY2008. Two mainstem electrofishing trips will be conducted, one during March and again in September. The September trip will also include sampling below Diamond Creek. Concurrent with LCR sampling in the spring (March and April), two mainstem monitoring trips utilizing trammel nets and hoop nets will be conducted to capture and tag HBC in the LCR inflow reach. These data, as well as data collected during the spring LCR sampling efforts, will be used to conduct a concurrent closed population mark-recapture estimate of HBC abundance. This monitoring sampling design will be assessed as part of the PEP scheduled for 2008.

Products/Reports

Annual reports detailing the findings of each of the above activities will be prepared and submitted to the GCMRC for internal and/or external review as center policy dictates. As warranted, project findings will be prepared and submitted for publication in the primary peer-reviewed literature. These data will be utilized in the 2008 PEP.

Budget

BIO 2.R4.08	
Monitoring Mainstem Fishes (includes Diamond Down; Ongoing)	
	Fiscal year 2008
GCMRC personnel costs (19% burden)	63,618
GCMRC project-related travel/training (19% burden)	4,967
GCMRC operations/supplies (19% burden)	14,000
GCMRC equipment purchase/replacement (19% burden)	12,000
AMP logistical support (19% burden)	92,700
Outside GCMRC and contract science labor (19% and/or other burden rate)	—
Cooperative/interagency agreements (6.09% GCMRC burden plus cooperator's burden)	278,600
Project Subtotal	465,885
DOI customer burden (combined 6.09%, 19% and/or other rates)	52,551
Project Total (Gross)	518,436
Percent outsourced (outside of GCMRC; includes 50% of logistics)	70%

BIO 2.R5.08: Nonnative Control Planning

BIO 2.R6.08: Nonnative Control Pilot Testing

Start Date

September 2007

End Date

September 2011

Investigator(s)

Grand Canyon Monitoring and Research Center (K.D. Hilwig, M.E. Andersen, L.G. Coggins, Ph.D., staff), in cooperation with the U.S. Fish and Wildlife Service and Arizona Game and Fish Department

Geographic Scope

The Colorado River ecosystem in Grand Canyon

Project Goals/Tasks

These projects seek to address the following objectives:

- To elucidate critical physical and biotic factors that may be limiting to, or supportive of, the HBC and other native fish populations in Grand Canyon and to seek methods that reduce, eliminate, or control limiting factors
- To determine and refine the most appropriate method(s) for estimating the population size of HBC and other Grand Canyon fishes, including sampling design, gear selection, and development of remote monitoring methods (The method(s) developed and selected should be consistent with the second edition of the Colorado River Endangered Fishes Recovery Goals. The FWS has scheduled revision of the goals to be initiated in 2007.)

The specific goal of the tasks identified in this project description is to evaluate threats to native fishes resulting from nonnative fishes, to develop a plan to control those species that pose the greatest threats to natives, and to test implementation of this plan. This project is expected to be complete in September 2011.

Need for Project

Nonnative fishes are among the greatest threats to native fishes in western North America rivers. Nonnative fishes may threaten native fishes by direct predation, by competing for available food and other resources, and by habitat modification. Nonnative fishes were introduced into Grand Canyon not later than early in the twentieth century. While native fishes survived these initial introductions at least long enough to be described by early researchers, other system stressors, especially the modification of natural flows as a result of dam installation, appear to have increased the threats to native fishes from nonnative fishes.

The GCDAMP has recognized nonnative fishes as a threat that needs to be addressed, and proceeded with implementation of a RBT and other nonnative fish control experiment around the LCR inflow reach over the last 4 years. The work described in this work plan builds on that effort. As the Colorado River mainstem becomes warmer due to climate effects, the potential for an increased threat from warmwater-adapted nonnative fishes increases. There is an immediate need to begin investigating what species pose the greatest threats to natives, how those species might be controlled, and to test control approaches for efficacy.

Strategic Science Questions

Primary SSQs addressed:

SSQ 1-2. Does a decrease in the abundance of rainbow trout and other cold- and warmwater nonnatives in Marble and eastern Grand Canyons result in an improvement in the recruitment rate of juvenile humpback chub to the adult population?

SSQ 1-4. Can long-term decreases in abundance of rainbow trout in Marble and eastern Grand Canyons be sustained with a reduced level of effort of mechanical removal or will recolonization from tributaries and from downstream and upstream of the removal reach require that mechanical removal be an ongoing management action? This question also applies to future removal programs targeting other nonnative species.

SSQ 5-6. Do the potential benefits of improved rearing habitat (warmer, more stable, more backwater and vegetated shorelines, more food) outweigh negative impacts due to increases in nonnative fish abundance?

The AMP SAs have articulated the following summary science questions that are addressed by this project:

SA 1. What are the most limiting factors to successful humpback chub adult recruitment in the mainstem: spawning success, predation on YoY and juveniles, habitat (water, temperature), pathogens, adult maturation, food availability, competition?

SA 2. What are the most probably positive and negative impacts of warming the Colorado River on humpback chub adults and juveniles?

Links/Relationships to Other Programs

Understanding the status and trends of the Grand Canyon fish populations, especially the endangered HBC, is important to evaluating the effects of management and conservation activities, especially GCD operations. If HBC populations are stable or increasing, then dam operations are unlikely to be having a negative effect on the population, and may be supporting population stability and growth. If the populations are decreasing, the operations may be having a negative impact and may need to be critically evaluated, along with other physical and biotic factors, especially nonnative fish populations.

One of the management approaches that have been proposed to support HBC and other native fishes in Grand Canyon is the installation of a selective withdrawal structure on the GCD so that water of various temperatures, especially warmer water from the reservoir's epilimnion, may be released. A potential concern with this approach is that warmer mainstem temperatures may also favor warmer water nonnatives, increasing the risk from these species to natives. This project will help address the potential threat from nonnatives and how it may be addressed, thereby helping evaluate the impact of a selective withdrawal structure.

Information Needs Addressed

Primary information needs addressed:

CMIN 2.4.1 Determine and track the abundance and distribution of nonnative predatory fish species in the Colorado River.

RIN 2.4.1. What are the most effective strategies and control methods to limit nonnative fish predation and competition on native fish?

RIN 2.4.3. To what degree, which species, and where in the system are exotic fish a detriment to the existence of native fish through predation or competition?

RIN 2.4.4. What are the target population levels, body size, and age structure for nonnative fish in the Colorado River ecosystem that limit their levels to those commensurate with the viability of native fish populations?

General Methods

A project manager was hired in October 2007 to begin working on this project full time. She is reviewing relevant literature, especially the history of fish introductions in Grand Canyon, life histories, and habitat used by those species, and case histories of nonnative control in other big river systems. Currently, the manager is developing a comprehensive nonnative control plan, due for completion by September 2011 and a short-term response plan due for completion in 2007. The field study planned for 2008 will be directed by the results of the pilot project scheduled for 2007. Beginning in 2007, a brief annual progress report will be delivered which will include the results of annual control method and gear testing projects. The 2007 project will test three gear types for effectiveness at capturing channel catfish and the use of sonic tags to track channel catfish.

Products/Reports

Brief annual reports will be produced each year of the project. One experimental trip is anticipated each year. Each experimental trip will be preceded by a complete trip plan and followed by a complete trip report. These field studies will supplement literature studies to be incorporated into a comprehensive nonnative control document scheduled for completion in September 2011.

Budget

BIO 2.R5.08	
Nonnative Control Planning (FY2007–10)	
	Fiscal year 2008
GCMRC personnel costs (19% burden)	89,550
GCMRC project-related travel/training (19% burden)	1,500
GCMRC operations/supplies (19% burden)	560
GCMRC equipment purchase/replacement (19% burden)	—
AMP logistical support (19% burden)	—
Outside GCMRC and contract science labor (19% and/or other burden rate)	—
Cooperative/interagency agreements (6.09% GCMRC burden plus cooperator's burden)	—
Project Subtotal	91,610
DOI customer burden (combined 6.09%, 19% and/or other rates)	17,406
Project Total (Gross)	109,016
Percent outsourced (outside of GCMRC; includes 50% of logistics)	0%

BIO 2.R6.08	
Nonnative Control Pilot Testing (FY2007–10)	
	Fiscal year 2008
GCMRC personnel costs (19% burden)	15,556
GCMRC project-related travel/training (19% burden)	500
GCMRC operations/supplies (19% burden)	—
GCMRC equipment purchase/replacement (19% burden)	10,000
AMP logistical support (19% burden)	32,000
Outside GCMRC and contract science labor (19% and/or other burden rate)	—
Cooperative/interagency agreements (6.09% GCMRC burden plus cooperator's burden)	49,479
Project Subtotal	107,535
DOI customer burden (combined 6.09%, 19% and/or other rates)	14,044
Project Total (Gross)	121,579
Percent outsourced (outside of GCMRC; includes 50% of logistics)	61%

BIO 2.R7.08: Stock Assessment of Native Fish in Grand Canyon

Start Date

October 2006

End Date

Ongoing

Principal Investigator(s)

Lewis G. Coggins, Ph.D., Grand Canyon Monitoring and Research Center

Geographic Scope

The mainstem Colorado River in Grand Canyon

Project Goals/Tasks

The objective addressed by this project is to determine and refine the most appropriate method(s) for estimating the population size of HBC and other Grand Canyon fishes, including sampling design, gear selection, and development of remote monitoring methods. The method(s) developed and selected should be consistent with the second edition of the Colorado River Endangered Fishes Recovery Goals. (The USFWS has scheduled revision of the goals to be initiated in 2007.)

The specific goals of the tasks identified in this project description are to annually update and refine stock assessment models for HBC, and to attempt to develop stock assessment models for flannelmouth sucker and bluehead sucker.

Need for Project

Native fish populations in Grand Canyon are key resources of concern influencing decisions on both the operation of GCD and other nonflow related actions. To inform these decisions, it is imperative that accurate and timely information on the status of native fish populations, particularly the endangered HBC, be available to managers. Additionally, a suite of adaptive experimental management actions are being contemplated to better understand the mechanisms controlling the population dynamics of native fishes, and to identify policies that are consistent with the attainment of management goals. The assessments generated from this project will be used, in part, to assess the effects of implemented experimental actions. This information is therefore crucial to (1) inform the program as to attainment of identified goals, (2) provide baseline status and trend information to be used as a backdrop to further understand mechanisms controlling native fish population dynamics, and (3) evaluate the efficacy of particular management policies in attaining program goals. Finally, results from this project are potentially useful in assessing changes to Federal Endangered Species Act listing status of native fishes in the Colorado River.

Strategic Science Questions

Primary SSQ addressed:

SSQ 1-1. To what extent are adult populations of native fish controlled by production of young fish from tributaries, spawning and incubation in the mainstem, survival of YoY and juvenile stages in the mainstem, or by changes in growth and maturation in the adult population as influenced by mainstem conditions?

Additional SSQ addressed:

SSQ 1-8. How can native and nonnative fishes best be monitored while minimizing impacts from capture and handling or sampling?

The AMP SAs have articulated the following science question, which is partially addressed by this project:

SA 1. What are the most limiting factors to successful humpback chub adult recruitment in the mainstem: spawning success, predation on YoY and juveniles, habitat (water, temperature), pathogens, adult maturation, food availability, competition?

Links/Relationships to Other Projects

Understanding the factors influencing the dynamics of the Grand Canyon native fish populations, especially the endangered HBC, is important to evaluating the effects of management and conservation activities, especially GCD operations. To discover these factors, a combination of large-scale manipulations (e.g., experimental removal of nonnative fish or long-term implementation of contrasting flow regimes) and smaller scale process-oriented research (e.g., assessment of juvenile fish growth rates under various temperature regimes or availability of particular food items) will likely prove most efficient in determining the key mechanisms regulating native fish populations. In each of these endeavors, it is critical that baseline trends in population abundance and recruitment be known. It is only with this knowledge that it is possible to assess population-level impacts of large-scale manipulations. Additionally, though it is informative to assess the effects of experimental management on processes thought to be important like growth or survival at particular life stages, this is not ultimately sufficient to determine efficacy of particular management actions. Linkages between these processes and ultimate recruitment to populations must be established. Again, these linkages can only be made if baseline trends in population abundance and recruitment are available.

Information Needs Addressed

RIN most directly addressed:

RIN 2.2.2. Determine if a population dynamics model can effectively predict response of native fish under different flow regimes and environmental conditions.

The activities in this project will refine and apply modeling to investigation of native and nonnative fish populations, allowing for comparison with various environmental factors, including flow regimes. Other RINs that ask questions about fish responses to environmental conditions that can be partially addressed with accurate modeling of the populations include the following:

RIN 2.2.8. What combination of dam release patterns and nonnative fish control facilitates successful spawning and recruitment of humpback chub in the Colorado River ecosystem?

RIN: 2.2.12. What are the impacts of research activities on mortality, recruitment, and the population size of humpback chub?

RIN 2.4.2. Determine if suppression of nonnative predators and competitors increases native fish populations.

General Methods

To provide HBC status and trend information, the GCMRC mark-recapture database will be annually updated with most recent data collected during routine monitoring efforts. Following this update, the HBC mark-recapture database will be reanalyzed using (where appropriate) both open and closed mark-recapture-based abundance estimators to provide the most current information on HBC status and trend. In particular, we will rely heavily on the ASMR models to determine trends in HBC abundance and recruitment. Ultimately we will consider the

performance of a suite of assessment models to infer the current status of the HBC in Grand Canyon. Finally, we will evaluate the applicability of similar techniques as described above to assessing stocks of flannelmouth sucker and bluehead sucker.

Products/Reports

- Annual assessment results will be presented to the TWG/AMWG as requested via oral reports.
- Biennially, native fish assessments will be compiled in peer-reviewed reports.

Budget

BIO 2.R7.08	
Stock Assessment of Native Fish in Grand Canyon (FY2007–Ongoing)	
	Fiscal year 2008
GCMRC personnel costs (19% burden)	34,033
GCMRC project-related travel/training (19% burden)	750
GCMRC operations/supplies (19% burden)	—
GCMRC equipment purchase/replacement (19% burden)	—
AMP logistical support (19% burden)	—
Outside GCMRC and contract science labor (19% and/or other burden rate)	—
Cooperative/interagency agreements (6.09% GCMRC burden plus cooperator's burden)	—
Project Subtotal	34,783
DOI customer burden (combined 6.09%, 19% and/or other rates)	6,609
Project Total (Gross)	41,392
Percent outsourced (outside of GCMRC; includes 50% of logistics)	0%

BIO 2.R8.08: Abundance Estimation Procedures

Start Date

October 2006

End Date

Ongoing

Principal Investigator(s)

Lewis G. Coggins, Ph.D., Grand Canyon Monitoring and Research Center

Geographic Scope

The mechanical removal reaches in the Colorado River in Grand Canyon

Project Goals/Tasks

The objectives addressed by this project are the following:

- To elucidate critical physical and biotic factors that may be limiting to, or supportive of, the HBC and other native fish populations in Grand Canyon and to seek methods that reduce, eliminate, or control limiting factors
- To explore the use of alternative estimation methods to evaluate the size of nonnative fish populations in the removal reach because the refinement of these estimates allows a more precise description of the magnitude of experimental mechanical removal treatment effects and may also help to evaluate factors affecting capture probability of electrofishing methods

The goal of this project is to evaluate the utility of Bayesian hierarchical models to estimate the abundance of nonnative fish (primarily RBT).

Need for Project

Precise and unbiased estimates of the abundance of RBT in the removal reaches of the Colorado River are necessary to evaluate both the magnitude and efficiency of removal efforts. These estimates allow computation of (1) the magnitude of the treatment effect (i.e., what percentage of nonnative fishes have been removed from the removal reach?), (2) the efficacy of the removal program (e.g., what percentage of fish is removed with each depletion pass?), and (3) the rate that fish immigrate back into the removal reach. In general, these estimates are the fundamental metrics of interest in the mechanical removal project.

Science Questions

Primary SSQ addressed:

SSQ 1-4. Can long-term decreases in abundance of rainbow trout in Marble and eastern Grand Canyons be sustained with a reduced level of effort of mechanical removal or will recolonization from tributaries and from downstream and upstream of the removal reach require that mechanical removal be an ongoing management action? This question also applies to future removal programs targeting other nonnative species.

Links/Relationships to Other Projects

The work outlined in this project has a direct linkage to the mechanical removal project through the estimation of abundance and the other metrics described above. Additionally, deriving a general relationship between turbidity and vulnerability of fish to capture is potentially extremely useful to the electrofishing-based elements of the fish

monitoring program. Because our monitoring program currently relies on electrofishing catch rate to index the abundance of rainbow and brown trout, patterns in catch rate are possibly a result of both changes in abundance and turbidity-induced changes in vulnerability. If it becomes possible to estimate the relationship between turbidity and vulnerability, we could essentially “correct” both the historic and future catch rate estimates to obtain a less biased index of abundance.

Information Needs Addressed

RIN 2.2.8. What combination of dam release patterns and nonnative fish control facilitates successful spawning and recruitment of HBC in the Colorado River ecosystem?

This project contributes to resolution of this RIN by helping to quantify the number of nonnative fishes that must be removed from the system to allow rebound of HBC population numbers.

General Methods

Currently, the traditional Zippin abundance estimator is used to estimate the abundance of nonnative fish (primarily RBT) in the mechanical removal reaches of the Colorado River. Though accepted and widely applied, this estimator makes the strict assumption that the vulnerability of fish among depletion passes is constant. Because large changes in turbidity are commonly observed within and among removal trips, this assumption is questionable. A more contemporary Bayesian estimation framework allows relaxation of this assumption if the relationship between a covariate (e.g., turbidity or sediment concentration) and vulnerability can be estimated. Additionally, this framework may allow more efficient use of the available data by allowing model-based aggregation of site-specific estimates. The program Bayesian Inference using the Gibbs sampler (BUGS) will be used to fit models to our removal data.

Products/Reports

This work will appear as part of the Coggins dissertation and/or publications in the primary literature.

Budget

BIO 2.R8.08	
Abundance Estimation Procedures (FY2007–Ongoing)	
	Fiscal year 2008
GCMRC personnel costs (19% burden)	34,033
GCMRC project-related travel/training (19% burden)	750
GCMRC operations/supplies (19% burden)	—
GCMRC equipment purchase/replacement (19% burden)	—
AMP logistical support (19% burden)	—
Outside GCMRC and contract science labor (19% and/or other burden rate)	—
Cooperative/interagency agreements (6.09% GCMRC burden plus cooperator's burden)	—
Project Subtotal	34,783
DOI customer burden (combined 6.09%, 19% and/or other rates)	6,609
Project Total (Gross)	41,392
Percent outsourced (outside of GCMRC; includes 50% of logistics)	0%

BIO 2.R9.08: Investigate Factors Affecting the Survival Rate of Juvenile Native Fishes in the Mainstem Colorado River

Start Date

October 2006

End Date

September 2010

Principal Investigator(s)

Lewis G. Coggins, Ph.D., Grand Canyon Monitoring and Research Center

Geographic Scope

The mainstem Colorado River in Grand Canyon

Project Goals/Tasks

The objectives addressed by this project are the following:

- To improve understanding of factors influencing survival of YoY and juvenile native fishes
- To identify habitat characteristics (biotic and abiotic) that are important to juvenile life stages of native fishes, particularly HBC

This project was titled Bioenergetic Modeling for FY2007. However, we have retitled this project to more closely describe the purpose, rather than the method, of the project. Additionally, although bioenergetic models are one potential tool to evaluate the effect of dam operations, water temperature, and biotic interactions on survival rate of young native fishes, we are also investigating the use of other models to achieve this goal.

Need for Project

Informed predictions of ecosystem responses from well-constructed models to particular biotic and abiotic perturbations are useful for a number of reasons. First, they are useful as a policy screening mechanism to select potential experimental management actions or treatments that have a high probability of achieving desired resource responses, or eliminating from consideration those that have low success probability. Second, they can be used to predict consequences of unintended actions such as introduction of nonnative fishes not presently in the system. Lastly, they can be used to evaluate hypotheses about the relative importance of factors influencing the survival rate of juvenile native fish and the fish community as a whole.

Science Questions

SSQ 1-4. Can long-term decreases in abundance of rainbow trout in Marble and eastern Grand Canyons be sustained with a reduced level of effort of mechanical removal or will recolonization from tributaries and from downstream and upstream of the removal reach require that mechanical removal be an ongoing management action? This question also applies to future removal programs targeting other nonnative species.

SA 1. What are the most limiting factors to successful humpback chub adult recruitment in the mainstem: spawning success, predation on YOY and juveniles, habitat (water, temperature), pathogens, adult maturation, food availability, competition?

Links/Relationships to Other Projects

Adaptive management, as described in the DOI handbook, requires predictive models to evaluate potential management actions or experimental policies relative to resource response and learning. These predictive models can take many forms such as bioenergetic models or more mechanistic observational models. We will attempt to use monitoring data on juvenile native fish near the mouth of the LCR to model survival rate of those fish as a function of dam operations, water temperature, and nonnative fish abundance. Additionally, we will continue to evaluate the utility of a specific kind of bioenergetic model (ecopath) to investigate linkages to all elements of the aquatic ecosystem. These linkages will foster better collaboration between terrestrial, aquatic food base, and fisheries investigations by making these linkages explicit in a common modeling framework. Using the ecosim functionality, which allows policy simulations, this model could be used in a planning context at all levels of the program with regard to questions about the aquatic ecosystem.

Information Needs Addressed

RIN 2.4.2. Determine if suppression of nonnative predators and competitors increases native fish populations.

This project is aimed at providing information on the relative magnitude of effects of dam operations, water temperature, and nonnative fish abundance on the survival of juvenile native fish in the mainstem Colorado River.

General Methods

We will construct a mechanistic model to describe the abundance of juvenile native fish in the mainstem Colorado River below the confluence of the LCR. We will populate the model with the relative abundance measurements collected during mechanical removal and select monitoring trips during 2003–04. We will attempt to relate apparent survival of these fish to changes in dam operations, water temperature, and nonnative fish abundance. Additionally, we may populate an ecopath model (<http://www.ecopath.org/>) using data available from previous studies conducted in Grand Canyon as well as the relevant scientific literature to provide auxiliary information on the magnitude of mortality effects from nonnative fishes. Of particular importance will be the diet data collected associated with the mechanical removal project.

Products/Reports

This work will appear as part of the Coggins dissertation and subsequent publications in the primary literature.

Budget

BIO 2.R9.08	
Bioenergetics Modeling (FY2007–10)	
	Fiscal year 2008
GCMRC personnel costs (19% burden)	34,033
GCMRC project-related travel/training (19% burden)	—
GCMRC operations/supplies (19% burden)	750
GCMRC equipment purchase/replacement (19% burden)	—
AMP logistical support (19% burden)	—
Outside GCMRC and contract science labor (19% and/or other burden rate)	—
Cooperative/interagency agreements (6.09% GCMRC burden plus cooperator's burden)	—
Project Subtotal	34,783
DOI customer burden (combined 6.09%, 19% and/or other rates)	6,609
Project Total (Gross)	41,392
Percent outsourced (outside of GCMRC; includes 50% of logistics)	0%

BIO 2.R11.08: Native Fishes Habitat Data Analysis

Start Date

October 2006

End Date

September 2010

Principal Investigator(s)

Matthew E. Andersen, Grand Canyon Monitoring and Research Center

Geographic Scope

The mainstem Colorado River in Grand Canyon

Project Goals/Tasks

The objectives addressed by this project are the following:

- To elucidate critical physical and biotic factors that may be limiting to, or supportive of, the HBC and other native fish populations in Grand Canyon and to seek methods that reduce, eliminate, or control limiting factors
- To identify habitat characteristics that are most important to all life stages of HBC and to seek methods that maintain, and possibly replicate, suitable habitats
- To improve understanding of dam operations on YoY and juvenile HBC survival and habitat use

This project will use available literature to help determine the specific habitat preferences for different life history stages of native fishes, especially the endangered HBC. The available literature, including databases, will be analyzed with multivariate statistics in order to develop indicators of what habitat characteristics are most important for HBC and other natives.

Need for Project

A great deal of peer-reviewed literature, gray literature, and database information addresses specific aspects of habitat preferences/usage by different life stages of Grand Canyon native fishes, especially HBC. Scientists and managers trying to provide GCD flow recommendations have repeatedly tried to informally assimilate and synthesize the available data, but the data remain so scattered that these attempts are difficult. It is not uncommon for different individuals, reading different literature sources, to come to different conclusions regarding what native fish in Grand Canyon need. The lack of synthetic, statistically robust information makes recommendations to dam operators less than compelling. This project initiates a multiyear effort to synthesize data and subject it to rigorous statistical methods to help guide habitat maintenance/creation recommendations to dam operators and natural resource managers.

Strategic Science Questions

Primary SSQ addressed:

SSQ 1-1. To what extent are adult populations of native fish controlled by production of young fish from tributaries, spawning and incubation in the mainstem, survival of YoY and juvenile stages in the mainstem, or by changes in growth and maturation in the adult population as influenced by mainstem conditions?

Additional SSQs addressed:

SSQ 1-7. Which tributary and mainstem habitats are most important to native fishes and how can these habitats best be made useable and maintained?

SSQ 5-3. To what extent do temperature and fluctuations in flow limit spawning and incubation success for native fish?

SSQ 5-4. What is the relative importance of increased water temperature, shoreline stability, and food availability on the survival and growth of YoY and juvenile native fish?

The GCDAMP SAs have articulated the following science questions, which are addressed by this project:

SA 1. What are the most limiting factors to successful HBC adult recruitment in the mainstem: spawning success, predation on YoY and juveniles, habitat (water, temperature), pathogens, adult maturation, food availability, competition?

SA 2. What are the most probably positive and negative impacts of warming the Colorado River on HBC adults and juveniles?

Links/Relationships to Other Projects

Understanding the status and trends of the Grand Canyon fish populations, especially the endangered HBC, is important to evaluating the effects of management and conservation activities, especially GCD operations. If HBC populations are stable or increasing, then dam operations are unlikely to be having a negative effect on the population, and may be supporting population stability and growth. If the populations are decreasing, the operations may be having a negative impact and may need to be critically evaluated, along with other physical and biotic factors, especially nonnative fish populations.

Because of the diversity of individuals and available literature regarding HBC and other native fishes habitat preferences, recommendations for dam operations can be diverse and are not always well supported. Well-intentioned dam operators and natural resource managers often need to make decisions but currently do not have comprehensive, synthetic information available on which to base their decisions. Consequently, counterproductive dam and resource management decisions may be made. This project seeks to address these information needs and reduce the potential for negative or counterproductive management actions.

Information Needs Addressed

Primary information needs addressed:

RIN 2.1.4. What habitats enhance recruitment of native fish in the LCR and mainstem? What are the physical and biological characteristics of those habitats?

RIN 2.2.5. What are the appropriate habitat conditions for HBC spawning? Where are these found? Can they be created in the mainstem?

This project addresses these RINs by investigating what available data indicate are habitat characteristics that support native fish spawning and recruitment and where the data indicate these habitats are found. Improved definition of habitat characteristics that support native fish spawning and recruitment allows for investigation into what would be required to create such habitats.

General Methods

M.E. Andersen, GCMRC Supervisory Biologist, will pursue this project with input from other internal and external scientists. He will attempt to bring in all available data regarding HBC habitat preferences, and those of

other native fishes, as available. Considerable effort is anticipated to bring together disparate data sources into a single format that can be subjected to statistical analysis. The multivariate statistical package CANOCO, Version 4.5, and some supporting literature, has been purchased by the GCMRC for this purpose. Andersen acquired additional training in multivariate methods in FY2007 in support of this project.

Products/Reports

- A brief annual report describing project progress will be produced by September 2007.
- A more comprehensive 2-year report describing project progress, needs, and recommendations will be produced by September 2008.
- At this time it is anticipated that this project will lead to preparation of at least one manuscript that will be submitted for consideration for publication.

Budget

In FY2008, this project will be pursued by M.E. Andersen, whose salary is accounted for separately. The cost requests for these years are to allow for staff time and advanced training in statistical methods, some of which may involve travel. All the software and supporting documentation that is necessary to begin this work has already been purchased by the GCMRC.

BIO 2.R11.08	
Native Fishes Habitat Data Analysis (FY2007–10)	
	Fiscal year 2008
GCMRC personnel costs (19% burden)	24,323
GCMRC project-related travel/training (19% burden)	—
GCMRC operations/supplies (19% burden)	—
GCMRC equipment purchase/replacement (19% burden)	—
AMP logistical support (19% burden)	—
Outside GCMRC and contract science labor (19% and/or other burden rate)	—
Cooperative/interagency agreements (6.09% GCMRC burden plus cooperator's burden)	—
Project Subtotal	24,323
DOI customer burden (combined 6.09%, 19% and/or other rates)	4,621
Project Total (Gross)	28,944
Percent outsourced (outside of GCMRC; includes 50% of logistics)	0%

BIO 2.R12.08: Trammel Net Effects

BIO 2.R13.08: Remote PIT Tag Reading

BIO 2.R14.08: Test Sonic Tags

Start Date

October 2006

End Date

September 2009

Principal Investigator(s)

Matthew E. Andersen and Lew G. Coggins, Ph.D., Grand Canyon Monitoring and Research Center, with assistance from the Arizona Game and Fish Department and U.S. Fish and Wildlife Service

Geographic Scope

The mainstem Colorado River in Grand Canyon

Project Goals/Tasks

The objective addressed by these projects is the following:

- To determine and refine the most appropriate method(s) for estimating the population size of HBC and other Grand Canyon fishes, including sampling design, gear selection, and development of remote monitoring methods (The method(s) developed and selected should be consistent with the second edition of the Colorado River Endangered Fishes Recovery Goals. The USFWS has scheduled revision of the goals to be initiated in 2007.)

The specific goal of the tasks identified in this project description is to provide evaluations of currently used and potential monitoring techniques. In 2007 and 2008, a study will be conducted to investigate potential improvements in the use of trammel nets, one of the most common gear types in the Colorado River system, but also a gear type that has been implicated in causing stress to fish, a factor of particular importance when handling endangered fishes. This study should also provide quantification of the percentage of native fish populations sampled by trammel nets, an important metric to quantify in order to allow trammel net capture data to contribute to stock assessments. This project also proposes to test two types of monitoring that do not require repeated handling of fishes: (1) remote antennae that can read the PIT tags already implanted in a large fraction of the adult the Grand Canyon HBC population, and (2) sonic tags that once implanted in fish can be read by stationary readers.

Need for Project

A limited number of HBC and other native fishes are present in the modern day Colorado River in Grand Canyon. Nonnative fish species are also present, and are important to study because of their potential to prey on and/or compete with native fishes. Scientists and managers wish to know how many of these species are present and the age-class structures of these populations. Because of the limited numbers, however, scientists and managers wish to know just how effective their gear is in sampling populations; they also wish to obtain population information in the least intrusive manner(s) possible, especially when sampling the endangered HBC. Although more gear types remain to be tested, the four studies described herein begin to investigate gear efficiencies and potentially useful new gear types.

Strategic Science Questions

Primary SSQ addressed:

SSQ 1-8. How can native and nonnative fishes best be monitored while minimizing impacts from capture and handling or sampling?

Links/Relationships to Other Projects

Just which mainstem habitats are most important for native fishes is still a matter of debate among scientists and managers who study the Colorado River in Grand Canyon. The river is deep, wide, and swift in Grand Canyon, making fish sampling challenging. Remote-sensing techniques may provide increased documentation of fish habitat use. This will be especially useful if it turns out that fishes spend a measurable proportion of their time in habitats not susceptible to traditional gear types, such as nets and electroshocking. With increasing knowledge and quantification of fish habitat preferences, scientists and managers can make increasingly specific recommendations for dam releases that favor creation and maintenance of specific riverine habitat types.

Information Needs Addressed

Trammel Net

RIN 2.2.12. What are the impacts of research activities on mortality, recruitment, and the population size of humpback chub?

Trammel nets can be utilized to track the relative abundance of native and nonnative fishes in the Colorado River. If the nets are used in this way they should be deployed so as to be most effective and as safe as possible.

Remote PIT Tag Reading and Sonic Tags

RIN 2.6.5. How are movement patterns for flannelmouth sucker, bluehead sucker, and speckled dace in the Colorado River ecosystem affected by age, natal stream, and dam operations?

CMIN 2.4.1. Determine and track the abundance and distribution of nonnative predatory fish species in the Colorado River ecosystem and their impacts on native fish.

CMIN 2.6.1. Determine and track the abundance and distribution of flannelmouth sucker, bluehead sucker, and speckled dace populations in the Colorado River ecosystem.

General Methods

Beginning in 2007 a graduate student will be partially supported by GCDAMP funds to pursue study of trammel nets. The student will work with faculty at Northern Arizona University (NAU) led by Dr. Alice Gibb. Initial studies will be conducted at an AZGFD hatchery in large rolloff bins with aquaculture-grade liners used to hold water and fish. The expected study animals will be closely related *Gila* species, probably roundtail chub or bonytail.

Experimentation with the use of remote antennae to read PIT tags will be conducted mainly by personnel from the AZGFD. The study area will focus, at least initially, on the LCR.

Experimentation with sonic tags will be led by GCMRC and FWS personnel, working closely with the product manufacturer, who is based in Tucson. Initial efforts will focus on capturing nonnative fish that will be implanted with these tags and released to see if the equipment is effective in the large Colorado River.

Products/Reports

- The preliminary results of the trammel net study are expected by the summer of 2008, and a completed master's thesis on the topic should be completed by the summer of 2009.
- Annual reports, including results and recommendations, will be provided on the use of the three remote-sensing techniques by September 30th of each year. These reports will be used to evaluate whether additional studies are warranted or whether one or more techniques should be abandoned.

Budget

BIO 2.R12.08	
Trammel Net Effects (FY2007–09)	
	Fiscal year 2008
GCMRC personnel costs (19% burden)	—
GCMRC project-related travel/training (19% burden)	—
GCMRC operations/supplies (19% burden)	—
GCMRC equipment purchase/replacement (19% burden)	—
AMP logistical support (19% burden)	—
Outside GCMRC and contract science labor (19% and/or other burden rate)	—
Cooperative/interagency agreements (6.09% GCMRC burden plus cooperator's burden)	36,250
Project Subtotal	36,250
DOI customer burden (combined 6.09%, 19% and/or other rates)	2,208
Project Total (Gross)	38,458
Percent outsourced (outside of GCMRC; includes 50% of logistics)	100%

BIO 2.R13.08	
Remote PIT Tag Reading (FY2007–09)	
	Fiscal year 2008
GCMRC personnel costs (19% burden)	—
GCMRC project-related travel/training (19% burden)	—
GCMRC operations/supplies (19% burden)	—
GCMRC equipment purchase/replacement (19% burden)	—
AMP logistical support (19% burden)	4,600
Outside GCMRC and contract science labor (19% and/or other burden rate)	—
Cooperative/interagency agreements (6.09% GCMRC burden plus cooperator's burden)	27,477
Project Subtotal	32,077
DOI customer burden (combined 6.09%, 19% and/or other rates)	2,547
Project Total (Gross)	34,624
Percent outsourced (outside of GCMRC; includes 50% of logistics)	93%

BIO 2.R14.08	
Test Sonic Tags (FY2007–09)	
	Fiscal year 2008
GCMRC personnel costs (19% burden)	8,873
GCMRC project-related travel/training (19% burden)	—
GCMRC operations/supplies (19% burden)	—
GCMRC equipment purchase/replacement (19% burden)	15,000
AMP logistical support (19% burden)	5,000
Outside GCMRC and contract science labor (19% and/or other burden rate)	—
Cooperative/interagency agreements (6.09% GCMRC burden plus cooperator's burden)	39,595
Project Subtotal	68,468
DOI customer burden (combined 6.09%, 19% and/or other rates)	7,897
Project Total (Gross)	76,365
Percent outsourced (outside of GCMRC; includes 50% of logistics)	61%

GCDAMP Goal 4. Maintain a naturally reproducing population of rainbow trout above the Paria River, to the extent practicable and consistent with the maintenance of viable populations of native fish.

BIO.4.M1.08: Status and Trends of Lees Ferry Trout

Start Date

Ongoing

End Date

Ongoing

Principal Investigator(s)

Arizona Game and Fish Department; Ecometric, Inc.; and Grand Canyon Monitoring and Research Center

Geographic Scope

Colorado River from Glen Canyon Dam to Lees Ferry

Project Goals/Tasks

The objective addressed by this project is the following:

- To monitor the RBT population below GCD to monitor responses to various flows

Operation of the GCD affects the ecology of nonnative RBT and the aquatic food base in the Lees Ferry reach (McKinney and others, 1999). The Lees Ferry fishery was recognized as a resource of concern in the Operation of Glen Canyon Dam Final Environmental Impact Statement (FEIS) (1995): “NPS, AZGFD, Hualapai, and Navajo objectives for the trout fishery are to provide a recreational resource while maintaining and recovering native fish in Grand Canyon. In the Glen Canyon reach, their objective is to encourage natural reproduction, survival, and growth of trout to blue ribbon quality sizes.” This project is designed to monitor the status of the trout fishery to contribute to evaluation of whether this goal from the EIS is being met. Information needs still exist to understand how the trout population, especially regarding reproduction and survival and growth of young fish, responds to modified low fluctuating flow (MLFF) alternative. An additional task has been added to address this need. These protocols will potentially be modified considering the recommendations of the 2007 PEP.

Need for Project

The downstream fish community is an assemblage of native and nonnative fish that occur in the CRE. The status and trends of the fishery are regulated by biotic and abiotic mechanisms that may in turn be affected by the operations of GCD. Monitoring basic population statistics including abundance and distribution of native and nonnative fishes provide information necessary to assess the status of these resources and inform the Adaptive Management Program.

The AZGFD has worked with other fishery cooperators including the GCMRC, FWS, and SWCA Environmental Consultants over the past 5 years to develop consistent, repeatable sampling methods for fishes in both the

mainstem Colorado River and LCR. The overall objective of this proposal is to continue standardized sampling and continue to work to develop a long-term monitoring program for all fish populations. The AZGFD will also assist with other special projects and research needs as appropriate.

Strategic Science Questions

Primary SSQ addressed:

SSQ 3-6. What GCD operations (ramping rates, daily flow range, etc.) maximize trout fishing opportunities and catchability?

Links/Relationships to Other Projects

Understanding the status of the Lees Ferry RBT population is critical to estimating and monitoring the risk that this species may pose to native fishes both in the Lees Ferry reach and further downstream. Following implementation of a 4-year project to remove RBT from the LCR reach of the Colorado River, it will be critical to understand the status and trends of Lees Ferry RBT to help evaluate any repopulation of downstream reaches that may occur.

Information Needs Addressed

Primary information needs addressed:

CMIN 4.1.2. Determine annual proportional stock density of rainbow trout in the Lees Ferry reach.

CMIN 4.1.4. Determine annual standard condition (Kn) and relative weight of rainbow trout in the Lees Ferry reach.

There are a number of RINs that are partially addressed by this project, or which depend, in part, on the results of this project. The primary RIN addressed is the following:

RIN 4.1.1. What is the target proportional stock density (i.e., tradeoff between numbers and size) for rainbow trout in the Lees Ferry reach?

The data collected with the monitoring in this project provide the data on which managers depend to monitor the size and condition of the current RBT population.

General Methods

The fishery is sampled by electrofishing to estimate biological parameters to assess the status and trends of the fishery. Electrofishing provides information on size composition, relative abundance (catch per minute as an index of population size), and condition (length-weight relationships), and samples are collected for whirling disease examination. Samples are collected at 27 stratified random and 9 fixed electrofishing transects three times per year in an augmented, serially alternating sampling design as recommended by the PEP. Present sampling design can detect a 6–10-percent linear change in abundance over a 5-year period. Work is currently underway to assess the statistical power of intra- and interannual comparisons. We are evaluating other methods to estimate abundance, including snorkel surveys (Korman and others, 2006), mark-recapture population estimates similar to those done in 1991 and 1998, and depletion sampling to convert CPUE estimates to population estimates.

Present methods for assessing abundance using a catch rate index may or may not be adequate for addressing management objectives and targets. If managers need an “n” (number of fish), further work needs to be done to find the most cost-effective way to generate reliable population estimates. We are working to evaluate different abundance estimators and discussing management targets with managers (AZGFD) and anglers. We will likely suggest some alternative methods to assess the abundance objective rather than “annual population estimates” as stated in CMIN 4.1, or attempt to clarify the CMIN.

Ongoing analysis of and sampling for trout redds and YoY trout will be conducted. The scope of work for this project is being negotiated and finalized.

Products/Reports

Separate reports will be provided for the mainstem sampling on or before January 1 of the year following the sampling for internal and external review. The revised final deliverable will be submitted on or before March 31 of the year following the sampling.

Budget

BIO 4.M1.08	
Status & Trends of Lees Ferry Trout (Ongoing)	
	Fiscal year 2008
GCMRC personnel costs (19% burden)	—
GCMRC project-related travel/training (19% burden)	—
GCMRC operations/supplies (19% burden)	—
GCMRC equipment purchase/replacement (19% burden)	—
AMP logistical support (19% burden)	2,780
Outside GCMRC and contract science labor (19% and/or other burden rate)	—
Cooperative/interagency agreements (6.09% GCMRC burden plus cooperator's burden)	124,200
Project Subtotal	126,980
DOI customer burden (combined 6.09%, 19% and/or other rates)	8,092
Project Total (Gross)	135,072
Percent outsourced (outside of GCMRC; includes 50% of logistics)	99%

BIO 4.E1.08: Monitoring Rainbow Trout Redds and Larvae

Start Date

October 1, 2007

End Date

September 30, 2008

Principal Investigator(s)

Joshua Korman, Ecometric, Inc.

Geographic Scope

The Lees Ferry reach of the Colorado River

Project Goals/Tasks

The objectives addressed by this project are the following:

- To monitor the responses of RBT redds and larvae in the Lees Ferry reach to dam release patterns
- To complete model development and testing
- To document extent of biases and precision of model parameters under a range of sampling strategies (e.g., how many redd and fry surveys are required)
- To analyze the 2003–07 data within this modeling framework to provide the most reliable estimates of the effects of the 2003 and 2005 experimental flows and other nonexperimental impacts (e.g., 2004 flood, September 1 minimum flow reduction)

Need for Project

Increased flow fluctuations are thought to affect RBT abundance through two mechanisms. Variation in river stage would lead to exposure and temporary desiccation of redds, which in turn could lead to a reduction in egg and alevin survival rates. A number of studies have shown that egg stages are relatively insensitive to short-term desiccation events but that eleutheroembryos and preemergent alevins are very sensitive to desiccation (e.g., Becker and others, 1982). Variation in river stage would also destabilize shoreline habitats, forcing fry and parr to move from protective shoreline cover and increase their predation risks and energetic costs, ultimately leading to reduced survival and growth. McKinney and others (2001) hypothesized that this was the most likely mechanism explaining the increase in the abundance of small RBT observed following reduced flow fluctuations from GCD.

In theory, monitoring the abundance of adults would provide an assessment of the effects of increased flow fluctuations on RBT populations in Glen or Grand Canyons. In practice, this approach is complicated by a number of factors: (1) it may be institutionally difficult to implement a sound experimental design where the treatment is held constant for sufficient time for its effects to become observable in the adult population that is monitored. Recruitment pulses generated from single-year experiments may be swamped by the presence of multiple year-classes in the adult population; and (2) there will be a considerable lag (3+ yrs) between the time that recruitment of juvenile trout is changed and the time it takes for this change to be potentially noticeable in the adult population if a long-term experiment is implemented.

Considering these challenges, it is reasonable to investigate other monitoring alternatives that are more amenable to detecting the effects of short-term (i.e., 1- or 2-year) experiments with minimal lag time. Korman and others (2005) documented the utility of estimating survival rates of incubating life stages and YoY RBT to assess the effects of experimental fluctuating flows in 2003 and 2004. The methodology appears to have considerable

promise to estimate key biological parameters (survival, growth, habitat use, movement) that determine the strength of the age-1 year-class that ultimately determines the size of the adult population. The study continued in 2006 at a reduced level of funding. The year 2006 was the first nontreatment year when ROD flows were implemented throughout the spawning and incubation period. This year therefore represents a control period that can be used to compare incubation and fry survival rates under ROD conditions with those experienced under experimental flows. Data from the 2006 surveys proved very useful to managers in documenting at least a 90 percent decline in spawning intensity in the winter of 2006. In spite of the large decline in spawning activity, the July 2006 fry survey results suggested that densities are $\frac{1}{2}$ to $\frac{3}{4}$ of the levels in 2004. These data suggest that there is a strong compensatory survival response of young trout. This observation is of interest to managers both for assessing the efficacy of reducing trout abundance by focusing on incubation stages, as well as whether to stock fry due to the recent downturn in abundance.

Strategic Science Questions

Primary SSQ addressed:

SSQ 1.1. Does a decrease in the abundance of rainbow trout and other cold- and warmwater nonnatives in Marble and eastern Grand Canyons result in an improvement in the recruitment rate of juvenile humpback chub to the adult population?

Assuming that at least some of the RBT observed in the LCR reach are produced in the Lees Ferry reach, understanding the cohort success of the Lees Ferry RBT will be important to correlate with HBC population dynamics.

Links to Other Projects

Because RBT were hypothesized to have negative impacts on the native fish population, the Adaptive Management Program implemented an experimental nonnative fishes removal project from 2003 through 2006. The primary species removed was RBT, although smaller numbers of other nonnative fish species were also removed. Although the source of the RBT removed has not been determined with certainty, there is evidence that the fish migrated downstream from the Lees Ferry reach. If RBT do have negative impacts on native fishes and these RBT are migrating from the Lees Ferry reach, then the condition of the Lees Ferry RBT population is of great importance to managers as they determine actions to take that benefit the native fish population in Grand Canyon.

Information Needs Addressed

RIN 4.1.1. What is the target proportional stock density (i.e., tradeoff between numbers and size) for rainbow trout in the Lees Ferry reach?

RIN 4.1.2. What is the minimum quantity and quality of spawning substrate necessary for maintaining a wild reproducing rainbow trout population in the Lees Ferry reach?

EIN 4.1.1. How does rainbow trout abundance, proportional stock density, length at age, condition, spawning habitat, natural recruitment, whirling disease and other parasitic infections change in response to an experiment performed under the ROD, unanticipated event, or other management action?

General Methods

Incubation Mortality Model

We have developed a model that integrates the effects of spawn timing, redd hypsometry (proportion of redds by elevation band), and the effects of intergravel water temperature on incubation time and mortality, to predict the relative incubation mortality for daily spawning cohorts. The model consists of five components:

1. Spawn timing—The spawn timing model is used to predict the number of redds created by day over a one-year period (November 1 to October 31).
2. Redd hypsometry—Redds created on each model day are distributed across four elevation bands (<5,000 cfs; 5,000–8,000 cfs; 8,000–12,000 cfs; 12,000–15,000 cfs; and 15,000–20,000 cfs) based on the observed hypsometry.
3. Incubation time—The number of days from spawning to hatch are computed based on the time required to exceed an accumulated thermal unit (ATU) threshold of 329 degree (°C)-days (Jensen and others, 1992) using daily average intergravel temperature measurements. Incubation time varies for each spawning day and elevation band because of differences in temperature histories. The number of days between hatching and emergence was fixed at 30 days, which was determined from the otolith microstructure of a large sample of age-0 rainbow.
4. Temperature-dependent incubation mortality—Daily maximum and minimum intergravel water temperatures over the incubation period are compared to temperature-mortality thresholds. If the daily maximum or minimum temperatures over the incubation period exceed assumed lethal thresholds, redds created on that day are recorded as not producing viable young.
5. Predict hatch timing—The numbers of both viable and total redds are summed across elevation bands for each spawning day. The ratio of the number of nonviable redds to the total number of redds provides a prediction of the daily relative incubation mortality rate. Predictions are shifted from spawning date to hatch date based on the computed incubation time for each spawning date and elevation band.

The predicted hatch date distribution based on the combined spawn timing and incubation mortality models, hereafter referred to as incubation mortality-based predictions, represents the distribution that would result assuming temporal and spatial variation in incubation mortality determined by dewatering frequency. The predicted hatch date distribution based on the spawn timing model alone, hereafter referred to as spawn timing-based predictions, represents the distribution that would result assuming constant incubation mortality. This latter prediction represents the null model as it assumes that mortality is not dependent on dewatering frequency determined by the flow regime.

This model will be applied using count, hypsometry, and temperature data aggregated over the entire Lees Ferry reach. Maximum and minimum daily temperature thresholds during the egg (spawn-to-hatch) and alevin (hatch-emergence) incubation periods were derived from the literature.

Products/Reports

- Document data, model, and results from 2003–07 (4 study years) in four manuscripts to be submitted to peer-reviewed journals by June 2008.
- Provide two brief management reports covering (1) data/methods not described in sufficient detail in the peer-reviewed manuscripts, and (2) results from the sampling strategy analysis.
- Present findings at one TWG or AMWG meeting at the end of the project and to the trout PEP.

Budget

BIO 4.E1.08	
Monitoring Rainbow Trout Redds & Larvae (FY2007)	
	Fiscal year 2008
GCMRC personnel costs (19% burden)	—
GCMRC project-related travel/training (19% burden)	—
GCMRC operations/supplies (19% burden)	—
GCMRC equipment purchase/replacement (19% burden)	—
AMP logistical support (19% burden)	—
Outside GCMRC and contract science labor (19% and/or other burden rate)	—
Cooperative/interagency agreements (6.09% GCMRC burden plus cooperator's burden)	—
Project Subtotal	—
DOI customer burden (combined 6.09%, 19% and/or other rates)	—
Project Total (Gross)	—
Percent outsourced (outside of GCMRC; includes 50% of logistics)	0%

*No funding needed in FY2008

GCDAMP Goal 5: Maintain or attain viable populations of Kanab ambersnail.

BIO 5.R1.08: Monitor Kanab ambersnail (Includes backwater monitoring of small-bodied fishes)

Start Date

April 2007

End Date

September 2010

Principal Investigator(s)

Barbara E. Ralston, Ph.D., Grand Canyon Monitoring and Research Center; and Arizona Game and Fish Department

Geographic Scope

Vaseys Paradise, located 31.5 RM downstream of Lees Ferry; surveys encompass the springs around the pour-off at Vaseys Paradise. The monitoring of KAS is conducted in conjunction with monitoring of backwater habitats for small-bodied fishes.

Project Goals/Tasks

The goals of this project are to determine the extent and kind of vegetation that exists as habitat for the KAS and to track the abundance and distribution of KAS at Vaseys Paradise. The following is a list of tasks required to meet these goals:

- Sample vegetation plots at Vaseys Paradise to determine patch composition and areal extent (spring and fall of each year). Sample for the presence of snails in plots.
- Enter data and conduct quality control on data entry. Provide data to the GCMRC for vegetation analysis.
- Compare previous vegetation composition to previous vegetation/habitat surveys to assess habitat. Provide abundance estimates of snails. Report writing by the GCMRC (winter of each year).
- Provide snail density estimates based on sampling or model estimates.

Need for Project

Knowing the extent of habitat is needed in the event of a high flow to support development of a biological opinion and to help determine snail densities. Changes in snail numbers can be associated with changes in vegetation. By monitoring the vegetation at Vaseys Paradise, the snails are indirectly monitored, based on the assumption that if the preferred habitat is present then snails will also be present. Total habitat can be measured using remote methods, but the composition of the habitat may still require on-the-ground sampling. Sampling at Vaseys Paradise can also provide data for GCDAMP goal 6, which refers to the protection and improvement of riparian and spring communities.

Strategic Science Questions

There are no SSQs that are directly related to the goal of maintaining or attaining viable KAS populations. The specific information needs addressed by the project are indicated below.

Links/Relationships to Other Projects

Riparian vegetation, including vegetation at springs, is a critical interface between aquatic and terrestrial environments around the world. There are multiple components that riparian and spring communities either contribute to or influence (e.g., food base, available habitat). In the CRE, the spring vegetation itself serves as a host for invertebrates, like KAS, provides breeding and foraging habitat for small mammals and birds, provides cover in the heat of the day, and the spring water may be used for ceremonial purposes. Changes in the composition or structure of riparian spring communities like expansion of an exotic species may alter these interactions. Riparian and spring vegetation regulates nutrient exchange between the land and water, and leaf litter is a terrestrial carbon source that may influence in-stream invertebrate production. The relative importance of terrestrial carbon in the aquatic food web is, in part, being addressed through the food base initiative. The linkage could be further defined through studies that focused on terrestrial productivity and processes. Again, changes in abundance or kind of riparian carbon sources may influence aquatic and terrestrial productivity processes.

Information Needs Addressed

Primary information needs addressed:

CMIN 5.1.1. Determine the abundance and distribution of Kanab ambersnails at Vaseys Paradise in the lower (below 100,000 cfs) and upper zone (above 100,000 cfs).

CMIN 5.2.1. Determine and track the size and composition of habitat used by Kanab ambersnail at Vaseys Paradise.

General Methods

Habitat Sampling

Determine percent cover, diversity, and distribution of vegetation that constitutes KAS habitat. Random samples in the habitat record percent cover, plant height of dominant plants, and soil moisture. Survey total habitat and plots using conventional or alternative survey methods. Habitat area is calculated by the GCMRC survey department. Data are analyzed using univariate and multivariate approaches. A minimum of 10 samples will be taken for each patch, if possible.

- Sample snails in each patch. Record number and size of snails encountered.
- Enter and quality check data. Deliver data to the GCMRC for analysis.
- Evaluate current habitat parameters to historic data for comparison.
- Monitor relocated vegetation associated with high flow experimental conservation measures.

Products/Reports

- An annual report for KAS habitat and density estimates is produced by Arizona Game and Fish Department.
- An annual report of the results of backwater seining for small-bodied fishes is produced by the GCMRC.

Budget

BIO 5.R1.08	
Monitor Kanab Ambersnail (FY1995–FY2010)	
	Fiscal year 2008
GCMRC personnel costs (19% burden)	4,003
GCMRC project-related travel/training (19% burden)	—
GCMRC operations/supplies (19% burden)	—
GCMRC equipment purchase/replacement (19% burden)	—
AMP logistical support (19% burden)	10,300
Outside GCMRC and contract science labor (19% and/or other burden rate)	—
Cooperative/interagency agreements (6.09% GCMRC burden plus cooperator's burden)	16,325
Project Subtotal	30,628
DOI customer burden (combined 6.09%, 19% and/or other rates)	3,712
Project Total (Gross)	34,340
Percent outsourced (outside of GCMRC; includes 50% of logistics)	70%

GCDAMP Goal 6: Protect or improve the biotic riparian and spring communities, including threatened and endangered species and their critical habitat.

BIO 6.R1.08: Vegetation Mapping

BIO 6.R2.08: Vegetation Transects

Start Date

October 2006

End Date

September 2010

Principal Investigator(s)

Barbara E. Ralston, Grand Canyon Monitoring and Research Center; and other cooperators (e.g., U.S. Geological Survey, Northern Arizona University) to be determined

Geographic Scope

The riparian zone, including the old high-water zone (OHWZ; >97,000 cfs), in the Colorado River corridor from Glen Canyon Dam to Lake Mead

Project Goals/Tasks

The goals of these projects are to determine the areal extent of vegetation classes among the major habitat zones in the CRE (e.g., new high-water zone (NHWZ), sand beach community, old high-water zone) and how yearly GCD operations affect vegetation cover, richness, diversity, and wetland indicator value by surface elevation. The following tasks are designed to reach these goals:

1. Conduct field surveys to identify community constituents and determine if vegetation overstory species and cover values have changed. Use community analysis—ordination, two-way species analysis—to identify how understory communities may be changing (September and October 2007). Pending PEP recommendations.
2. Use image processing software (e.g., ENVI, ERDAS) to classify imagery into identified vegetation classes (fall/winter 2008). Imagery is from 2009 overflight and compared with the 2002 imagery for the purposes of change detection.
3. Groundtruth accuracy of vegetation classification (September 2008).
4. Compare revised vegetation map to previous vegetation map to determine area change for vegetation classes. Report writing (fall 2008).
5. Conduct field surveys of vegetation transects perpendicular to the river at specific stage elevations (15,000 cfs; 25,000 cfs; 35,000 cfs; 45,000 cfs; and 60,000 cfs) (fall 2008). Pending PEP recommendations.
6. Prepare and analyze data, including entering data, evaluating quality control, and analyzing data for diversity, cover, richness, and wetland score across elevations. Community analysis for marsh plots. Compare with previous years to assess trends (Winter 2008).

7. Report and incorporate results into a yearly report and into SCORE reporting (winter/spring 2008 and each subsequent year).

Need for Project

Riparian vegetation expansion, since operations at GCD began in 1963, has had a pivotal role in the ecology of the postdam river corridor. The reduction in annual flood volumes has allowed vegetation to expand and more permanently occupy land previously subjected to scouring in most years. The expansion has included marsh habitat occurring throughout the CRE, whereas previously, these habitats were restricted to Glen Canyon and the western Grand Canyon (Clover and Jotter, 1944; Turner and Karpiscak, 1980). The plants associated with the expansion include alien species like salt cedar (*Tamarix ramossissima*), camel thorn (*Alhagi maurorum*), and peppergrass (*Lepidium latifolium*), but also native species, arrowweed (*Pluchea sericea*), seepwillow (*Baccharis emoryi*), and coyote willow (*Salix exigua*). The variable operations over the years have resulted in an ebb and flow of vegetation expansion with vegetated area generally increasing over time (Waring 1995; Ralston and others, unpub. data). The increase in terrestrial vegetation contributes to aboveground primary productivity, arthropod densities, and associated food resources for terrestrial and aquatic vertebrates; is a source of culturally important plant species; and also can cause conflicts with recreational activities like available camping area. Because riparian vegetation is linked to multiple resources, knowing how vegetation is changing via monitoring (e.g., which species are expanding or declining and where) is an important source of data when evaluating dam operations.

To address the AMWG needs associated with riparian vegetation requires systemwide assessment of vegetation change at the broad scale (new high-water zone) as well as at the local scale (plot data). While knowing how much vegetation in the river corridor exists is useful, it is equally useful to know how the species that make up the vegetation may be changing. Because riparian vegetation contributes to aquatic productivity (Naiman and others, 2005) and serves as a host to terrestrial invertebrates and higher order vertebrates (e.g., lizards, birds), assessing the quality of these plants can help explain changes observed in higher order vertebrate abundances, including fish species (Nakano and Murakami, 2001). Changes in riparian vegetation are associated with dam operations (Stevens and others, 1995; Kearsley, 2006) and can affect the propagation of exotic species like tamarisk (Porter, 2002). Yearly transects assess year-to-year operations that can detect changes among herbaceous species, including invasives, while remotely sensed data can assess changes in overstory wood species that change more slowly.

The three riparian vegetation studies proposed in the annual work plan are composed of two field-based studies, (1) vegetation dynamics, and (2) vegetation mapping that utilizes corridorwide overflight data scheduled in 2009); and an office-based study, (3) riparian vegetation synthesis Part I). The two field-based projects complement each other rather than replicate efforts. Vegetation dynamics is an annual monitoring effort that records species diversity, richness, and cover at specific stage elevations. The changes in vegetation parameters that this monitoring detects is relevant to perennial and annual herbaceous species like bunch grasses, marsh species and invasive species that can change on an annual basis. Vegetation mapping utilizes the digital overflight imagery (product of the DASA Program) to quantify larger scale area changes (e.g., expansion of arrowweed patches, or extent and type of vegetated shoreline). Imagery is from 2009 overflight and compared with the 2002 imagery for the purposes of change detection. Analysis of change detection in the vegetation mapping project would incorporate the annual transect survey results to help explain patterns of change that may occur over a 5-year timeframe. The two projects complement each other because they provide information about changes in riparian habitat at different ecological scales which may affect other riparian community constituents like invertebrate biomass and riparian bird abundances. Lastly, the vegetation synthesis would use results from both of these studies and previous mapping and monitoring results to test mechanisms that affect riparian vegetation establishment and expansion including rates of change potential colonization sites.

Strategic Science Questions

Primary SSQs addressed:

SSQ 2-1. Do dam-controlled flows affect (increase or decrease) rates of erosion and vegetation growth at archaeological sites and TCP sites, and if so, how?

SSQ 4-2. How important are backwaters and vegetated shoreline habitats to the overall growth and survival of YoY and juvenile native fish? Does the long-term benefit of increasing these habitats outweigh short-term potential costs (displacement and possible mortality of young humpback chub) associated with high flows?

SSQ 5-7. How do warmer releases affect viability and productivity of native/nonnative vegetation?

GCDAMP goal 6 is directed at the protection or improvement of riparian and spring communities. This goal is based on the recognition that the riparian and spring environments are hosts for some endangered species like the southwestern willow flycatcher (*Empidonax traillii extimus*). The protection of these species' critical habitats is part of this goal. Riparian plant communities can be viewed at either a single resource level without ecosystem linkages, or at an integrative level where riparian vegetation is linked to aquatic and terrestrial ecosystem processes (e.g., contributes to secondary production, cover), interacts with cultural resources associated with recreation (e.g., camping sites) and TCPs, or affects aeolian sand transport and possibly archaeological site erosion rates. Understanding how riparian vegetation responds to flows and affects other resources of concern forms a basis for managing critical resources like native fish, archaeological properties, and recreational components.

Links/Relationships to Other Projects

Riparian vegetation is a critical interface between aquatic and terrestrial environments around the world. In the CRE, the vegetation itself serves as a host for invertebrates, provides breeding and foraging habitat for birds, provides cover in the heat of the day, and may be harvested for cultural utility. Changes in the composition or structure of riparian vegetation like expansion of an exotic species may alter these interactions. Riparian vegetation regulates nutrient exchange between the land and water, and leaf litter is a terrestrial carbon source that may influence in-stream invertebrate production. The relative importance of terrestrial carbon in the aquatic food web is, in part, being addressed through the food base initiative. The linkage could be further defined through studies that focus on terrestrial productivity and processes. Again, changes in abundance or kind of riparian carbon sources may influence aquatic productivity processes. The 2005 Knowledge Assessment Workshop (KAW) revealed that there was some certainty about the relationship of marsh community development and flows for the CRE, but that this certainty decreased as one progresses upslope. The outcome of the KAW and the science questions for riparian habitats indicate that, besides knowing the influence of flow on composition and extent of riparian vegetation, an understanding of the integrated role of riparian vegetation with other resources is needed (e.g., aquatic or cultural resources). This understanding would come from a combination of monitoring, synthesis, and field research.

Information Needs Addressed

Parameters and metrics to be measured, and the CMINs that each element addresses.

Determine and track the status and trends of the identified riparian communities (e.g., marsh community, sand beach, nonnative invasive species, etc.) at the appropriate timescale (CMIN 6.1.1, 6.2.1, 6.5.1, 6.6.1). This need will be addressed through the following:

- Semidecadal color infrared (CIR) digital imagery mapping that would quantify (1) area change of dominant overstory species, (2) community composition and possibly changes in understory community composition through groundtruthing associated with mapping, and (3) coarse primary productivity estimates for riparian vegetation.

- Annual vegetation transects/grid surveys that correlate with river stage elevations of 15,000 cfs; 25,000 cfs; 35,000 cfs; 45,000 cfs; and 60,000 cfs. Quantifies cover, richness, and diversity and wetland species scores at each stage elevation. This work would be most informative for herbaceous annuals and perennials, including invasive species. This component would need to incorporate marsh-monitoring needs of tribes.

General Methods

Vegetation Mapping

Community identification will be done using relevé plots in the field that are used to record relative cover. Cover scales use a Daubenmire scale. Data are recorded as categorical data, but plant height of the dominant species is also recorded. Number of samples for each plot is dependent on the abundance of the vegetation type. A minimum of 10 samples will be taken for each community (6 types identified in 2002). These data will be analyzed using multivariate statistics (ordination techniques) to identify the dominant communities along the river corridor.

Vegetation classification will use supervised classification routines that are available in an image-processing software package (ENVI, 2005). Training areas will be selected from previous base map groundtruth. Classes that will likely be used for this effort include tamarisk, baccharis/salix, marsh/wetlands, mesquite/acacia, arrow weed and bare ground. User and producer accuracies will be determined and class aggregation may be required to meet national vegetation mapping standards. The scheduled 2009 overflight would be compared with 2005 and 2002 imagery for vegetation area change detection purposes.

Quantification of changes in riparian communities will be done using a Geographic Information Systems (GIS) platform (ArcMap, ESRI, Inc. 2002).

Vegetation Transects

Data collection involves recording vegetation cover of species within each of four 1-m² plots at each elevation. Transects are located throughout the river corridor and sampled in a rotated panel design so that some plots are sampled every year (n = 20) and others are sampled every 3 years (n = 40). Marsh data will be incorporated for tribal monitoring (August/September 2008 and each year following, pending PEP recommendations).

Sample locations are determined by using the sediment transport and river simulation (STARS) model of Randle and Pemberton (1987), which predicts elevation rise based on river stage in combination with the Colorado River flow, and sediment storage/graphic user interface (CRFSSGUI) model (Ecometric, Inc.), which uses STARS model data and information on channel gradient, width, and roughness to predict the timing and height of the hydrograph at any point along the river.

Vegetation sampling of each transect corresponds to five stage elevations (15,000 cfs; 25,000 cfs; 35,000 cfs; 45,000 cfs; and 60,000 cfs). At each elevation point, a 1-by-1-m sighting frame (per Floyd and Anderson, 1982) with 100 crosshair intersections was placed and leveled with one side along the transect and the riverward corner of the transect side directly over the pin flag. Once a frame was surveyed, the frame is moved upstream or downstream at the same level so that four 1-by-1-m areas are sampled (two frames upstream of the transect and two downstream).

Vegetation data are recorded in the following way. First, all species present in the 1-by-1-m areas are recorded. These data are included in the univariate measures (cover, richness, diversity) but are excluded from the multivariate analyses.

To estimate percent vegetative cover in each frame, the number of sighting points which intercepted each species is counted. If multiple species were present under a single sighting point, all are recorded once so that the total cover of all species can collectively sum to more than 100 percent. Species which are encountered in at least one of the frames, but which are not seen beneath any of the 400 sighting points, are assigned an arbitrary "trace" cover value of 0.001 percent.

Products/Reports

- Annual report for vegetation transect monitoring and a single 5-year report for vegetation mapping change detection.
- Annual progress report will be provided for mapping/change detection project.
- Peer-reviewed articles from vegetation mapping project regarding change detection as well as remote-sensing technology and its utility in mapping vegetation in the arid Southwest.

Budget

BIO 6.R1.08	
Vegetation Mapping (FY2007–10)	
	Fiscal year 2008
GCMRC personnel costs (19% burden)	74,416
GCMRC project-related travel/training (19% burden)	3,000
GCMRC operations/supplies (19% burden)	1,000
GCMRC equipment purchase/replacement (19% burden)	—
AMP logistical support (19% burden)	13,000
Outside GCMRC and contract science labor (19% and/or other burden rate)	—
Cooperative/interagency agreements (6.09% GCMRC burden plus cooperator's burden)	—
Project Subtotal	91,416
DOI customer burden (combined 6.09%, 19% and/or other rates)	17,369
Project Total (Gross)	108,785
Percent outsourced (outside of GCMRC; includes 50% of logistics)	7%

BIO 6.R2.08	
Vegetation Transects (FY2007–10)	
	Fiscal year 2008
GCMRC personnel costs (19% burden)	8,500
GCMRC project-related travel/training (19% burden)	—
GCMRC operations/supplies (19% burden)	1,000
GCMRC equipment purchase/replacement (19% burden)	—
AMP logistical support (19% burden)	13,000
Outside GCMRC and contract science labor (19% and/or other burden rate)	—
Cooperative/interagency agreements (6.09% GCMRC burden plus cooperator's burden)	59,300
Project Subtotal	81,800
DOI customer burden (combined 6.09%, 19% and/or other rates)	7,886
Project Total (Gross)	89,686
Percent outsourced (outside of GCMRC; includes 50% of logistics)	80%

BIO 6.R3.08: Vegetation Synthesis

Start Date

October 2006

End Date

September 2010

Principal Investigator(s)

Barbara E. Ralston, Ph.D., Grand Canyon Monitoring and Research Center; and other cooperators to be determined

Geographic Scope

The riparian zone, including the old high-water zone (>97,000 cfs), in the Colorado River corridor from Glen Canyon Dam to Lake Mead

Project Goals/Tasks

The goal of the project is to utilize existing data from the riparian zones to characterize temporal and spatial responses of riparian vegetation to GCD operations (FY2007–08). The following tasks are designed to meet the goal of this project:

1. Conduct literature and data review of research associated with GCD and data from other rivers to identify appropriate datasets for synthesis at multiple scales (local, reach, systemwide)
2. Topic discussed per scale (local, reach, etc.):
 - Biomass
 - Species diversity
 - Rates of change—community scale
 - Incorporation of physical resource information
 - Determination between site/scale differences
 - Aquatic and terrestrial linkages—preliminary analysis
3. Report results
4. Utilize local and reach-based parameter values to produce a submodel of riparian vegetation response to changes in operations
5. Identify modeling tool for use (e.g., Structural Thinking Experiential Learning Laboratory with Animation [STELLA], GCM)
6. Develop model—parameter definitions and model run (Use intermediate disturbance hypothesis (Huston, 1979; Roxburgh and others, 2004) to test changes in parameters and conceptual model of riparian vegetation response to operations in CRE.)
7. Verify model using published data
8. Report results—incorporate into a yearly report and into SCORE reporting

Incorporate vegetation synthesis results into terrestrial faunal aquatic biology research and cultural program to improve CRE model (FY2009–10).

Need for Project

The GCMRC recognizes that there is a large amount of information in the gray literature associated with riparian vegetation for the Colorado River. The synthesis is intended to utilize the results of these data to construct a synthesis for riparian vegetation. The synthesis would evaluate vegetation change, interactions, and ecosystem function at local, geomorphic, and systemwide scales. The synthesis will incorporate data from other disciplines, most notably the physical science program, as it has completed a synthesis in 2004 (Schmidt and others, 2004). The synthesis should result in several papers that would be submitted for publication in peer-reviewed journals.

The synthesis would consist of two phases with the first phase representing a summary of information and hypotheses generation from review of the material and incorporation of other studies from other rivers. The second phase would be model development to test hypotheses for riparian vegetation change along the river corridor. The model would contribute to our conceptual model of carbon cycling within the CRE.

The identification of mechanisms of change provides loose predictive capabilities regarding the response of riparian vegetation to operations and the associated response in terrestrial and aquatic fauna that are affected by riparian community structure and composition. The compilation and synthesis of sediment and gage data since 1965 and earlier (Topping and others, 2003; Schmidt and others, 2004) provides a rich dataset that forms a basis for study of how discharge and sediment volumes influence community structure within the riparian community.

Strategic Science Questions

Primary SSQs addressed:

SSQ 2-1. Do dam-controlled flows affect (increase or decrease) rates of erosion and vegetation growth at archaeological sites and TCP sites, and if so, how?

SSQ 4-2. How important are backwaters and vegetated shoreline habitats to the overall growth and survival of YoY and juvenile native fish? Does the long-term benefit of increasing these habitats outweigh short-term potential costs (displacement and possible mortality of young humpback chub) associated with high flows?

SSQ 5-7. How do warmer releases affect viability and productivity of native/nonnative vegetation?

GDDAMP goal 6 for terrestrial resources is directed at the protection or improvement of riparian and spring communities. Included in the goal is the recognition that the riparian and spring environments are hosts for some endangered species like the southwestern willow flycatcher (*Empidonax traillii extimus*). The protection of these species' critical habitats is part of this goal. Riparian plant communities can be viewed at either a single-resource level without ecosystem linkages, or at an integrative level where riparian vegetation is linked to aquatic and terrestrial ecosystem processes (e.g., contributes to secondary production, cover), interacts with cultural resources associated with recreation (e.g., camping sites) and TCPs, or affects aeolian sand transport and possibly archaeological site erosion rates. Understanding how riparian vegetation responds to flows and affects other resources of concern forms a basis for managing critical resources like native fish, archaeological properties, and recreational components.

Links/Relationships to Other Projects

The expansion of vegetation along the river corridor affects multiples resources. The increased shoreline vegetation contributes to aquatic drift and may serve as supplemental source of carbon for aquatic food webs in addition to in-stream production. The ecology of human behaviors along the river corridor is affected by riparian vegetation. Exotic species that spread by tributary introductions (e.g., camel thorn) impact campable area by making some beaches unusable. Available campsite area is dependent on amount of open sand, availability of trees and shrubs for shade and wind breaks, and accessibility to the river (i.e., steepness of bank) among other variables (Kearsley and others, 1994; Kaplinski and others, 2005). In a similar vein, culturally important plants and locations have been monitored under the auspices of the adaptive management program since the 1990s (Phillips and Jackson, 1996; Austin and others, 1997; Lomaomvaya and others, 2001). How these data have

changed over time also needs to be incorporated into a synthesis to provide a holistic view of the riparian community.

Information Needs Addressed

The primary information needs addressed by these projects are CMINs 6.1.1., 6.2.1, 6.5.1, and 6.6.1, which are summarized as the following:

- Determine and track the abundance, composition, distribution, and area of terrestrial native and nonnative vegetation species in the CRE.
- Determine parameters and metrics to be measured, and the information needs that address each element.
- Determine how the abundance, composition, and distribution of the OHWZ, NHWZ, and sand beach community have changed since dam closure (1963), high flows (1984), interim flows (1991), and the implementation of ROD operations (RIN 6.2.1, 6.3.1, 6.4.1, 6.5.1, 6.5.2, 6.5.3).
- Compare vegetation patches from the 2002 vegetation base map with previous vegetation maps (Waring, 1995) that were completed for sections of the river for years 1965, 1973, 1984, 1990, and 1991 to determine distribution and abundance information at a gross scale (e.g., NHW, OHW, sand beach, marsh). Area coverage will be provided for different zones.
- Compositional changes are more difficult to determine. Will attempt modeling after assessing local, historic plot data (e.g., Stevens and Ayers, 1993, 1997; Kearsley and Ayers, 1996) and identifying local and reach scale factors that influence community assembly rules. Validation of model using 2005 CIR imagery and ground surveys that coincide with mapping project (FY2008).
- Perform change detection between years to identify change in area and distributional changes for woody exotics (e.g., tamarisk).

General Methods

Landscape change detection using GIS analysis tools to identify area change for vegetation classes or zones of interest between years. Identification of tamarisk in black and white imagery will be done using 2002 and 2005 imagery and comparing imagery characteristics of the vegetation. The scanning project in DASA is orthorectifying historic imagery that will permit retrospective analysis of vegetation change.

Quantification of changes in riparian communities will be done using a GIS platform (ArcMap, ESRI, Inc. 2002).

Vegetation Modeling

1. Incorporate parameters
2. Determine physical parameters (discharge, rates of sediment loss/gain by reach)
3. Precipitation records—decadal
4. Rates of change for vegetation class—obtained from change detection question.
5. Utilize intermediate disturbance hypothesis (Roxburgh and others, 2004; Barnes and others, 2006) to test assumptions of species interactions within the CRE
6. Validate using vegetation transect data and composition data associated with vegetation mapping

Products/Reports

- Semiannual report of progress
- Final report
- Individual reports for segments of synthesis submitted for publication

Budget

BIO 6.R3.08	
Vegetation Synthesis (FY2007–10)	
	Fiscal year 2008
GCMRC personnel costs (19% burden)	31,720
GCMRC project-related travel/training (19% burden)	3,000
GCMRC operations/supplies (19% burden)	5,000
GCMRC equipment purchase/replacement (19% burden)	—
AMP logistical support (19% burden)	—
Outside GCMRC and contract science labor (19% and/or other burden rate)	—
Cooperative/interagency agreements (6.09% GCMRC burden plus cooperator's burden)	20,000
Project Subtotal	59,720
DOI customer burden (combined 6.09%, 19% and/or other rates)	8,765
Project Total (Gross)	68,485
Percent outsourced (outside of GCMRC; includes 50% of logistics)	33%

GCDAMP Goal 7: Establish water temperature, quality, and flow dynamics to achieve the Adaptive Management Program ecosystem goals.

BIO 7.R1.08: Water Quality Monitoring of Lake Powell and the Glen Canyon Dam Tailwater

Start Date

Ongoing (current Interagency Agreement with Reclamation in place through September 30, 2009)

End Date

September 2009

Principal Investigator(s)

William S. Vernieu, Hydrologist, Grand Canyon Monitoring and Research Center

Geographic Scope

Lake Powell and its major tributary arms, inflow tributaries entering Lake Powell, and the tailwater from Glen Canyon Dam to Lees Ferry

Project Goals/Tasks

The objectives addressed by this project are the following:

- To maintain water-quality monitoring program for Lake Powell to predict and track processes in the reservoir that may influence GCD release water quality
- To maintain water-quality monitoring in GCD tailwater to directly evaluate the quality of GCD releases, the effects of GCD operations, and suitability for downstream aquatic resources
- To develop, in conjunction with Reclamation, CE-QUAL-W2 model to predict future changes to the water quality of Lake Powell and GCD releases; simulate the effects of various proposed and hypothetical climate, experimental, and operational scenarios; and guide future monitoring program revisions
- To complete comprehensive database of water-quality information from 40-year monitoring program and publish results as USGS Data Report for further interpretation, synthesis, and analysis
- To revise monitoring program, as needed, in conjunction with development of CE-QUAL-W2 model and historical data analysis, to ensure most efficient means of maintaining cost-effective and reliable monitoring program

Need for Project

Processes within Lake Powell, climate changes in the upper Colorado River Basin, the structure of GCD, and various aspects of dam operation affect the quality of water released from GCD to the CRE in Grand Canyon. Temperature, dissolved oxygen concentrations, nutrient concentrations, biological composition, and other characteristics of GCD releases can have a profound effect on the aquatic ecosystem below the dam.

The 5-year period of below-normal inflows in the upper Colorado River Basin from 2000 to 2004 resulted in a drawdown of Lake Powell of over 140 ft to 3,555 ft, representing 38 percent of total capacity, in 2005. Increasing influence of Lake Powell surface layers on GCD releases can be expected to cause warmer release temperatures, decreased release nutrient concentrations, and increased export of aquatic biota from Lake Powell. The lowering of warm surface layers in relation to the penstock withdrawal elevation has resulted in above-normal late-summer release temperatures since 2003. Release temperatures of 16 °C were recorded in October 2005, representing the warmest releases since 1971. Resuspension of exposed deltaic sediments from reservoir drawdown by 2005 inflow currents resulted in a plume of hypoxic water that appeared at GCD and began to be incorporated in GCD releases in July 2005. This resulted in dam releases containing the lowest concentrations of dissolved oxygen on record, reaching 3.3 mg/L in October 2005. Changes to individual turbine operations at GCD in September and October 2005 were shown to have a significant effect on the reaeration of hypoxic releases.

Differential routing of winter inflow currents can cause longer-term changes to the water quality of Lake Powell and eventual dam releases. For the past 7 years, with the exception of 2006, winter underflow density currents moved along the bottom of the reservoir and refreshed oxygen concentrations in the deepest layers of Lake Powell. In contrast, from 1994 to 1999 and during other periods in Lake Powell's history, winter density currents moved through the reservoir in intermediate layers as an interflow, which caused stagnation and a reduction of dissolved oxygen concentrations in the deepest hypolimnetic water of the reservoir. This interflow pattern again appeared in 2006. Exceptionally cold winter inflows in caused an underflow in January 2007, increasing hypolimnetic density and increasing the likelihood of future interflow conditions, which may cause reductions in hypolimnetic dissolved oxygen in future years.

The GCMRC works in conjunction with Reclamation on development of the CE-QUAL-W2 model by providing monitoring data that is used for model calibration and verification. This monitoring data consists of information describing the quality of water in GCD releases, Lake Powell, and tributary inflows into Lake Powell. In addition, the GCMRC has provided comments on the direction of model development so that a product can be developed that meets the needs of the GCMRC program. Reclamation has had at least two people working on model development and it has been felt that actual model development should rest with Reclamation, rather than have the GCMRC pursue a redundant modeling effort. It is recognized that once a functional model is in place, the different entities involved will have different questions to be addressed by the model. While model development is in progress and a full-functioning model is not yet in place, many components of the water-quality monitoring program can be addressed with results from the model, such as identifying parameters for which the model is more or less sensitive and restructuring monitoring efforts appropriately. Examples are identifying the need for more detailed inflow water-quality monitoring, establishment and maintenance of additional meteorological stations at the reservoir, and modifying sampling methods for biological parameters such as chlorophyll and plankton, in order to refine the model's ability to simulate productivity processes.

Strategic Science Questions

While the recent KA specified many science questions addressing the effects of water quality on various resources (sediment, food base, fisheries, recreation), no SSQs were proposed directly dealing with tracking and predicting changes in water quality in Lake Powell or GCD releases. The following questions are the most critical SSQs related to the effects of water quality on key resources:

SSQ 3-5. How is invertebrate flux affected by water quality (e.g., temperature, nutrient concentrations, turbidity) and dam operations?

SSQ 5-1. How do dam release temperatures, flows (average and fluctuating component), meteorology, canyon orientation and geometry, and reach morphology interact to determine mainstem and nearshore water temperatures throughout the CRE?

SSQ 5-3. To what extent do temperature and fluctuations in flow limit spawning and incubation success for native fish?

Links/Relationships to Other Projects

The quality of dam releases and subsequent in-stream changes can have a profound effect on various aspects of the aquatic ecosystem in Grand Canyon. Temperature affects metabolic rates of various organisms, including bacteria, plants, invertebrates, and vertebrates. It also affects reproductive processes, larval development, and behavior of native and nonnative fishes. Nutrient concentrations in dam releases can influence primary productivity processes in the clear water Lees Ferry reach. Dissolved oxygen is essential to maintaining healthy fish and invertebrate populations throughout Grand Canyon. Temperature and dissolved oxygen have been shown to have the most direct effect on native and nonnative fish populations. Suspended sediment concentrations limit the light available for primary productivity and affect the behavior of various fishes. Tracking status and trends of these water-quality parameters forms a direct link to various food base and fishery studies currently underway in Grand Canyon.

Information Needs Addressed

The following information needs (including supporting information needs [SINs]) (as updated June 23, 2003) relate directly to water-quality monitoring in Lake Powell and the GCD tailwater.

CMIN 7.1.1. Determine the water temperature dynamics in the main channel, tributaries (as appropriate), backwaters, and nearshore areas throughout the Colorado River ecosystem.

CMIN 7.2.1. Determine the seasonal and yearly trends in turbidity, water temperature, conductivity, DO, and pH changes in the main channel throughout the Colorado River ecosystem.

CMIN 7.3.1. What are the status and trends of water quality released from GCD?

SIN 7.2.1. How do the hydrodynamics and stratification of Lake Powell influence the food base or fisheries downstream?

SIN 7.2.2. Which water-quality variables influence food base and fisheries in the Colorado River ecosystem?

RIN 7.3.1. Develop simulation models for Lake Powell and the Colorado River to predict water-quality conditions under various operating scenarios, supplant monitoring efforts and elucidate understanding of the effects of dam operations, climate, and basin hydrology on Colorado River water quality.

7.3.1.a. Determine status and trends of chemical and biological components of water quality in Lake Powell as a function of regional hydrologic conditions and their relation to downstream releases.

7.3.1.b. Determine stratification, convective mixing patterns, and behavior of advective currents in Lake Powell and their relation to GCD operations to predict seasonal patterns and trends in downstream releases.

RIN 7.3.3. How do dam operations affect reservoir limnology?

SIN 7.3.1. Measure appropriate water-quality parameters to determine the influence of these parameters on biological resources in the Colorado River ecosystem.

EIN 7.3.1. How does the water quality of releases from GCD change in response to an experiment performed under the ROD, unanticipated event, or other management action?

Other information needs (as updated June 23, 2003) require supporting information from water-quality monitoring in Lake Powell and the GCD tailwater:

RIN 7.1.1. What are the desired ranges of spatial and temporal patterns of water temperatures for the CRE?

RIN 7.1.2. What are the most likely downstream temperature responses to a variety of scenarios involving a TCD on GCD?

RIN 7.1.3. What are the potential ecological effects of increasing mainstem water temperature?

RIN 7.2.1. Which major ions should be measured? Where and how often?

RIN 7.2.2. Which nutrients should be measured? Where and how often?

RIN 7.2.3. Which metals should be measured? Where and how often?

General Methods

Lake Powell monitoring is conducted monthly in the forebay and quarterly throughout the reservoir. Profiles of physical parameters (temperature, specific conductance, pH, dissolved oxygen, turbidity, redox potential) are collected through the water column at each site in the reservoir. Chemical (major ions and nutrients) and biological samples (chlorophyll and plankton) are collected at selected sites to characterize major strata and advective currents in the reservoir.

GCD tailwater monitoring consists of continuous monitoring (temperature, specific conductance, pH, dissolved oxygen, turbidity) with monthly chemical and biological sample collection. Grand Canyon monitoring consists primarily of collection of temperature and conductance at various locations.

Monitoring parameters include temperature, conductance, pH, dissolved oxygen, redox potential, and turbidity. Chemical analyses include determination of major ionic constituents and nutrient compounds of phosphorus and nitrogen. Plankton analyses include enumeration and identification of species, biomass estimates, and relative abundance calculations. All measurements and laboratory analyses are performed in accordance with standard approved methods.

Reservoir modeling is performed in cooperation between Reclamation and the GCMRC to achieve predictive capabilities and supplant or redirect some aspects of monitoring. Current model development has progressed to include calibrations for dissolved oxygen concentration, algal components, and oxygen demand from deltaic resuspension.

Products/Reports

- An annual report for FY2007 is in development and will be published in FY2008.
- Periodic reports of water-quality conditions will be posted via Internet.
- Updates on water-quality conditions will be provided to AMWG, TWG, and other interested parties through written reports or oral presentations periodically.

Budget

Reclamation provides direct funding to the GCMRC for the Lake Powell water-quality monitoring program. In addition, Reclamation also provides field support for monitoring activities, both from its own staff and through a service agreement with the NPS. It also provides laboratory analyses for nutrients, major ions, and chlorophyll through a service agreement with its Lower Colorado Regional Lab in Boulder City, Nevada. Reclamation is also taking the lead in development and calibration of the CE-QUAL-W2 simulation model for Lake Powell. A table of cost estimates for FY2008 follows.

Bio 7.R1.08	
Water-Quality Monitoring Lake—Powell & Tailwaters (FY2007–09)	
	Fiscal year 2008
GCMRC personnel costs (19% burden)	140,962
GCMRC project-related travel/training (19% burden)	10,720
GCMRC operations/supplies (19% burden)	22,500
GCMRC equipment purchase/replacement (19% burden)	4,500
AMP logistical support (19% burden)	—
Outside GCMRC and contract science labor (19% and/or other burden rate)	—
Cooperative/interagency agreements (6.09% GCMRC burden plus cooperator's burden)	—
Project Subtotal	178,682
DOI customer burden (combined 6.09%, 19% and/or other rates)	33,949
Project Total (Gross)	212,631
Percent outsourced (outside of GCMRC; includes 50% of logistics)	0%

PHY 7.M1.08: Core Monitoring of Downstream Integrated Quality of Water (below Glen Canyon Dam)

The downstream integrated quality-of-water (IQW) project focuses mostly on monitoring but can also support implementation of flow research related to stable flow testing, evaluation of alternative fluctuating flows, tests of beach/habitat-building flows (BHBF) and ongoing development and evaluation of numerical modeling. In some instances, it is difficult to separate these elements from experimental elements because they support each other. For example, monitoring the suspended-sediment budget may be considered core monitoring, but it is also required to assess a trigger for a BHBF such that it could be considered experimental research support. In the section on project goals/tasks, the individual project elements are described along with the associated category(s).

Start Date

October 2007

End Date

September 2008 (This project is intended to provide core-monitoring information to meet the information needs related GCDAMP goals 7 and 8 under an ongoing schedule during FY2008 and beyond).

Principal Investigator

David Topping, Ph.D., U.S. Geological Survey, Grand Canyon Monitoring and Research Center

Geographic Scope

The downstream IQW project is primarily focused on the main channel of the Colorado River from just below GCD (RM -15) downstream to the upper end of Lake Mead (as measured at the gaging station above Diamond Creek at RM 226). In addition, an important component of the project is a combination of monitoring and modeling of tributary sediment inputs such that sediment and flow monitoring activities are also carried out in various tributary watersheds, such as the Paria River at Lees Ferry, the LCR near Cameron, Arizona, another site above the confluence with the mainstem Colorado River, and various lesser tributaries in Glen, Marble, and Grand Canyons.

Project Goals/Tasks

The downstream IQW monitoring project is focused primarily on measurements of surface flow throughout the river ecosystem, as well as quality-of-water parameters such as temperature, specific conductivity, dissolved oxygen, and suspended-sediment transport. The monitoring project directly supports achievement of the following GCDAMP goals:

Goal 7: Establish water temperature, quality, and flow dynamics to achieve GCDAMP ecosystem goals.

Goal 8: Maintain or attain levels of sediment storage within the main channel and along shorelines to achieve GCDAMP ecosystem goals.

Because this monitoring project addresses the physical framework of the ecosystem, which underlies many biological, cultural, and recreational resource objectives, it indirectly supports achievement of almost all other GCDAMP goals, as described below:

Goal 1: Protect or improve the aquatic food base so that it will support viable populations of desired species at higher trophic levels.

The downstream IQW monitoring project supports this goal by providing information on flows, water temperature, and turbidity that aids in food base studies, such as the assessment of primary productivity and allochthonous inputs.

Goal 2: Maintain or attain a viable population of existing native fish, remove jeopardy for HBC and razorback sucker, and prevent adverse modification to their critical habitats.

The downstream IQW monitoring project also supports the native fish program by providing nearshore water temperature data for the assessment of growth rates, sediment concentration data that is used to adjust for catch efficiency in population models, flow and stage data that is important to understanding the effects of nearshore habitat disruption caused by fluctuating flows, and information on sandbars which create backwater habitats that are thought to be important for native fish.

Goal 4: Maintain a wild reproducing population of RBT above the Paria River, to the extent practicable and consistent with the maintenance of viable populations of native fish.

The downstream IQW monitoring project also monitors dam release and Glen Canyon quality of water, which proved critically important in fall 2004 when dissolved oxygen levels were low, requiring modifications to release patterns in order to raise oxygen levels.

Goal 6: Protect or improve the biotic riparian and spring communities within the CRE, including threatened and endangered species and their critical habitat.

The downstream IQW monitoring project also tracks the transport and fate of fine sediment, which provides the substrate for riparian vegetation and marsh communities.

Goal 9: Maintain or improve the quality of recreational experiences for users of the CRE within the framework of GCDAMP ecosystem goals.

The downstream IQW monitoring project also produces monitoring data and supports experimental and modeling research to understand flow dynamics and the size and abundance of sandbars, which are resources that affect the recreational experiences of Colorado River users such as rafters and fishermen.

Goal 11: Preserve, protect, manage, and treat cultural resources for the inspiration and benefit of past, present, and future generations.

The downstream IQW monitoring project also provides monitoring data related to riverine sandbars that provide a source of sediment, through aeolian transport, to high-elevation sand deposits that contain archaeological resources. In addition, the downstream IQW monitoring project has also developed stage modeling capabilities that allow for the assessment of the flow level that inundates a given cultural site.

In August 2004, the AMWG reviewed these goals and identified priority questions. The top five priority questions are as follows:

Priority 1: Why are HBC not thriving, and what can we do about it? How many HBC are there and how are they doing?

Priority 2: Which cultural resources, including TCPs, are within the Area of Potential Effect (APE), which should we treat, and how do we best protect them? What is the status and trends of cultural resources and what are the agents of deterioration?

Priority 3: What is the best flow regime?

Priority 4: What is the impact of sediment loss and what should we do about it?

Priority 5: What will happen when a TCD is tested or implemented? How should it be operated? Are safeguards needed for management?

As with the GCDAMP goals, the downstream IQW monitoring project directly supports some priorities while indirectly supporting others. For example, monitoring and research on flows, sediment transport, and water temperature clearly support priorities 3, 4, and 5 directly, while also indirectly supporting priorities 1 and 2 by providing information on the general physical framework of the riverine environment.

Several project-related tasks, listed below, occur within the downstream IQW monitoring project.

Flow and Stage Monitoring

Continued monitoring of flow and stage at established mainstem locations and major tributaries (RM -15, RM 0, RM 30, RM 61, RM 87, RM 166, RM 226, Paria River at HWY 89 bridge and near Lees Ferry, and two sites on the LCR). Category(s): Core Monitoring. Schedule: Ongoing. Official surface water records are collected at Paria River at the HWY 89 bridge and published by the USGS Utah Water Science Center. Official surface water records are collected and published by the USGS Arizona Water Science Center at the following tributary gage sites: Paria River near Lees Ferry, Ariz.; LCR near Cameron, Ariz.; LCR above the mouth near Desert View, Ariz.; Kanab Creek near Kanab, Utah; Havasu Creek above the mouth near Supai, Ariz.; and at the mainstem gages at RM 0, RM 8, and RM 226, Ariz. The RM -15 flow measurements are reported by Reclamation.

Quality-of-Water Monitoring

Continued monitoring of water temperature at established mainstem locations and major tributaries (RM -15, RM 0, RM 30, RM 61, RM 87, RM 166, RM 226, RM 246, Paria River at Lees Ferry, two sites on the LCR, and Kanab and Havasu Creeks). Continuation of a new nearshore/backwater temperature monitoring program. Continued monitoring of conductivity at established stations (RM -15, RM 0, RM 30, RM 61, RM 87, and RM 226). Continued monitoring of turbidity at established stations (RM 30, RM 61, and RM 226). Category(s): Core Monitoring. Schedule: Ongoing for mainstem temperature, conductivity, and turbidity monitoring; continuation of nearshore/backwater monitoring program in FY2008, then ongoing; monitoring data supports completion of downstream thermal model development during FY2008, applications ongoing.

Suspended-Sediment Flux Monitoring

Continued monitoring of suspended-sediment flux at established mainstem locations and major tributaries (RM 30, RM 61, RM 87, RM 166, RM 226, Paria River at Lees Ferry, and one site along the LCR [near Cameron, Ariz.]). Because BHBF triggers are based on sediment retention within the mainstem, it is insufficient to monitor tributary inputs only. Category(s): Core Monitoring. Schedule: ongoing.

Collaboration with and Support of Aquatic Food Base Program

Integrated research studies with the aquatic food base program, including submerged aquatic vegetation and bed texture classification with acoustics, monitoring algal drift with acoustics, and quantification of tributary inputs of organic material. Category(s): Support for Research and Development. Schedule: ongoing.

Coordination with Other Resource Areas

Regular meetings and interaction with other resource area personnel, particularly at the Program Manager level, in order to facilitate an ecosystem approach to our scientific studies and ensure that the downstream IQW monitoring project is providing useful information regarding the physical environment to the other resource areas. Category(s): Program Management. Schedule: ongoing.

Need for the Project

Information on flow, water quality, and suspended-sediment transport is critical to understanding the physical environment upon which biological and sociocultural resources depend (see details in Section 1 of this project description). In order to understand responses of these resources to dam operations, we must first understand the effects of dam operations on the physical environment. The goal of the downstream IQW project is to provide this information and link dam operations to changes in the physical environment.

Strategic Science Questions

The downstream IQW monitoring project is designed with the goal of providing data that supports answering the two primary physical resources questions identified during the KAW conducted in the summer of 2005, as follows:

SSQ 4-1. Is there a “Flow-Only” operation (i.e., a strategy for dam releases, including managing tributary inputs with BHBFs, without sediment augmentation) that will restore and maintain sandbar habitats over decadal timescales?

SSQ 5-1. How do dam release temperatures, flows (average and fluctuating component), meteorology, canyon orientation and geometry, and reach morphology interact to determine mainstem and nearshore water temperatures throughout the CRE?

Also, as detailed throughout this project description, the downstream IQW monitoring project provides information on the physical environment that is critical to other resource areas and will thus contribute indirectly to answering a variety of other science questions related to other resources.

Links/Relationships to Other Projects

Aquatic Food Web Research

The downstream IQW monitoring project supports new research focused on the food web of the river ecosystem by providing continuous data on surface flow in the main channel and major tributaries, as well as related quality-of-water data, such as water temperature, specific conductivity, dissolved oxygen and suspended-sediment concentrations and grain size for suspended particles in transport.

Fisheries Monitoring and Research

The downstream IQW monitoring project also supports science activities in the fisheries program by providing flow and quality-of-water data that may be used by fisheries biologists in evaluating their fish catch data, as well as growth, movement, and habitat use information.

Information Needs Addressed

The downstream IQW monitoring project directly addresses several of the CMINs and RINs related to GCDAMP goals 7 and 8. A selection of the information needs that are addressed by downstream IQW monitoring project are listed below. The downstream IQW monitoring project addresses many more CMINs, but the ones listed below are considered most relevant to answering the science questions outlined above.

CMIN 7.4.1. Determine and track flow releases from GCD under all operating conditions.

CMIN 7.1.2. Determine and track LCR discharge and temperature near the mouth (below springs).

CMIN 7.1.1. Determine the water temperature dynamics in the mainstem, tributaries, backwaters, and nearshore areas throughout the CRE.

CMIN 8.1.3. Track, as appropriate, the monthly sand and silt/clay volumes and grain-size characteristics, by reach, as measured or estimated at the Paria and LCR [near Cameron, Ariz., and above the confluence] stations, other major tributaries like Kanab and Havasu Creeks, and “lesser” tributaries?

CMIN 8.1.2. What are the monthly sand and silt/clay export volumes and grain-size characteristics, by reach, as measured or estimated at Lees Ferry, Lower Marble Canyon, Grand Canyon, and Diamond Creek Stations?

The monitoring data from the downstream IQW monitoring project not only fulfill the CMINs listed above, but are also intended to feed new information directly into modeling efforts (see PHY 07.R1.08) that will allow sediment-transport modelers the opportunity to address RINs related to GCDAMP goals 7 and 8.

RIN 7.4.1. What is the desired range of seasonal and annual flow dynamics associated with powerplant operations, BHBFs, and habitat maintenance flows, or other flows that meet GCDAMP goals and objectives?

RIN 7.3.1. Develop simulation models for Lake Powell and the Colorado River to predict water-quality conditions under various operating scenarios, supplant monitoring efforts, and elucidate understanding of the effects of dam operations, climate, and basin hydrology on Colorado River water quality.

RIN 8.5.1. What elements of ROD operations (upramp, downramp, maximum and minimum flow, MLFF, high modified flow (HMF), and BHBF) are most/least critical to conserving new fine sediment inputs, and stabilizing sediment deposits above the 25,000 cfs stage?

General Methods

Flow, stage, water temperature, conductivity, turbidity, and suspended-sediment data are collected using standard USGS protocols with Quality Assurance/Quality Control (QA/QC) (described in “Techniques of Water-Resources Investigations of the U.S. Geological Survey,” Book 3, Sections A and C). Suspended-sediment sampling is supplemented through the use of emerging technologies, including acoustics and laser-diffraction (Melis and others, 2003; Topping and others, 2004, 2006, in press). Stage, water temperature, conductivity, turbidity, and suspended-sediment surrogates (i.e., acoustics and laser-diffraction) are monitored with in situ instrumentation recording at 15-min intervals. River flow is measured episodically and used to develop a stage-discharge rating curve, providing 15-min flow records (described in “Techniques of Water-Resources Investigations of the U.S. Geological Survey,” Book 3, Section A). Similarly, suspended-sediment concentration is measured episodically and used to calibrate acoustic and laser diffraction instrumentation, providing 15-min records of concentration (sand, silt/clay, and sand grain size).

Products/Reports

- Streamflow, stage, and tributary sediment data will be published annually in Arizona and Utah Water Resources Data reports (surface water and sediment records published by the USGS Utah and Arizona Water Science Centers) and served through the GCMRC Web page (data delivered on or before February 28, 2009).
- Mainstem sediment transport and water-quality data will be summarized in a biennial data report; data will also be served through the GCMRC Web page (The GCMRC leads in preparing these reports.).
- Conference abstracts and proceedings articles (2–4), journal articles (1–3), and frequent presentations at stakeholder meetings will result from this project.

Budget

PHY 7.M1.08	
Integrated Quality-of-Water Monitoring (Downstream of GCD; FY2007–Ongoing)	
	Fiscal year 2008
GCMRC personnel costs (19% burden)	337,000
GCMRC project-related travel/training (19% burden)	10,000
GCMRC operations/supplies (19% burden)	34,785
GCMRC equipment purchase/replacement (19% burden)	20,000
AMP logistical support (19% burden)	50,000
Outside GCMRC and contract science labor (19% and/or other burden rate)	345,400
Cooperative/interagency agreements (6.09% GCMRC burden plus cooperator's burden)	—
Project Subtotal	797,185
DOI customer burden (combined 6.09%, 19% and/or other rates)	85,839
Project Total (Gross)	883,024
Percent outsourced (outside of GCMRC; includes 50% of logistics)	46%

PHY 7.R1.08: Modeling Support Linked with Downstream Integrated Quality-of-Water Monitoring Project (below Glen Canyon Dam)

The modeling support activity linked with the Downstream Integrated Quality-of-Water (IQW) project focuses on advancement of simulation capabilities needed to predict the fate of flow releases from GCD and associated water-quality constituents such as temperature and suspended sediment. This subelement of the downstream IQW monitoring project is intended to refine existing models that are being developed to predict downstream thermal regimes, as well as the fate of fine sediment inputs that enter the ecosystem from sources such as the Paria and Little Colorado Rivers.

Start Date

October 2006

End Date

Ongoing. This project parallels the downstream IQW monitoring project, and it is expected that support for model development and improvements will continue in parallel to the monitoring program. As new data are collected, existing models can be continuously tested, improved, and applied.

Principal Investigator(s)

Scott A. Wright, U.S. Geological Survey, Water Resources Discipline; Mark Schmeckle, Arizona State University; David M. Rubin, U.S. Geological Survey, Geological Discipline; David J. Topping, U.S. Geological Survey, Biological Resources Discipline.

Geographic Scope

For the most part, the modeling support activities are linked to the IQW project in a spatially parallel way and are, therefore, also focused on the main channel of the CRE, between GCD (RM -15) to Diamond Creek (RM 226). However, an important component of the downstream IQW is a combination of monitoring and modeling of tributary sediment inputs such that research and monitoring activities are carried out in various tributary watersheds as well, such as the Paria and Little Colorado Rivers. Because of this, the proposed modeling activities are also tied to monitoring efforts within these two major tributaries, particularly related to model simulations that predict sand production during runoff events.

Project Goals/Tasks

Ongoing development of models to simulate flow, sediment transport, and downstream water temperature are intended to be closely interfaced with ongoing monitoring activities throughout the science program. As stated in the previous section, the downstream IQW monitoring project (Project PHY 07.M1.08) is focused primarily on measurements of surface flow throughout the river ecosystem as well as quality-of-water parameters such as temperature, specific conductivity, dissolved oxygen, and suspended-sediment transport. As described in the section on Project PHY 07.M1.08, the monitoring project directly supports achievement of the following GCDAMP goals and specific modeling tasks:

Goal 7: Establish water temperature, quality, and flow dynamics to achieve GCDAMP ecosystem goals.

Goal 8: Maintain or attain levels of sediment storage within the main channel and along shorelines to achieve GCDAMP ecosystem goals.

Continued Support for the Development of Nearshore Temperature Models

Under the assumption that the development of nearshore temperature models will continue to be funded through a separate agreement with Reclamation (TCD funds), PI Wright will provide support and guidance to GCMRC

hydrologist Craig Anderson in this endeavor. In particular, in coordination with the second task outlined below, PI Wright will evaluate available multidimensional models, in particular Delft-3D, for use in nearshore temperature modeling. Category(s): Research and Development. Schedule: Ongoing.

Sediment Transport Modeling of Eddy-sandbar Environments.

This is one of the primary recommendations of the external review panel from the Sediment Transport Modeling Review Workshop conducted in Santa Cruz, Calif., on February 15–16, 2007 (Simoes and others, 2007). This is only one of the many recommendations made by the review panel; however, budget constraints dictate that not all recommendations can be addressed in FY2008. There is currently a need to further our understanding of how sediment is exchanged between the main channel and eddy sandbars, as well as how sandbars build and erode under various flows and sediment supply. Reliable predictions of the area and volume of eddy-sandbar deposits requires improvement in this area. Further, improvements in the one-dimensional routing model, as recommended by the Modeling Review Panel, can be facilitated by improving the parameterization of sandbar deposition and erosion mechanics through the use of multidimensional models. Various readily available multidimensional models will be evaluated for use in Grand Canyon. Also, existing datasets will be assessed to determine the need for further field data collection and/or laboratory experiments. Once applicable models are chosen and datasets are selected and/or collected, the datasets and models will be used to further our understanding of how eddy sandbars form and evolve under a variety of flow and sediment supply conditions. The results from these modeling exercises will be used to improve one-dimensional sand routing algorithms as well as to evaluate the effects of various dam operations (e.g., BHBFs, ramping rates) on eddy-sandbar deposits. Category(s): Research and Development. Schedule: Ongoing, through at least FY2008.

Need for Project

Information on flow, water quality, and suspended-sediment transport is critical to understanding the physical environment upon which biological and sociocultural resources depend (see details Project PHY 07.M1.08 description). In order to understand responses of these resources to dam operations, we must first understand the effects of dam operations on the physical environment. The goal of the modeling support activities linked to the downstream IQW monitoring project is to provide increased predictive capabilities (simulations) that can be used as planning tools for linking dam operations to changes in the physical environment, as well as exploring interdisciplinary relationships with biological, cultural, economic, and recreational elements of GCDAMP.

Strategic Science Questions

The downstream IQW modeling activities are designed with the objective of providing predictive capability that supports answering the two primary physical resource questions identified during the KAW conducted in the summer of 2005:

SSQ 4-1. Is there a “Flow-Only” operation (i.e., a strategy for dam releases, including managing tributary inputs with BHBFs, without sediment augmentation) that will restore and maintain sandbar habitats over decadal timescales?

SSQ 5-1. How do dam release temperatures, flows (average and fluctuating component), meteorology, canyon orientation and geometry, and reach morphology interact to determine mainstem and nearshore water temperatures throughout the CRE?

Both of the above questions can be only partially addressed through collection of monitoring data. Likewise, both questions are related to issues that can be at least partially resolved through focused experimental research in combination with ongoing modeling research activities. Following collection of monitoring data in Project PHY 07.M1.08, development and refinement of the models for simulating flow, suspended-sediment transport, and downstream temperature evolution is the next step toward resolving these critical questions in the next phase of monitoring and research.

Links/Relationship to Other Projects

Because ongoing modeling efforts are linked to the downstream IQW monitoring project, it is also intended to address and support elements of the physical framework of the ecosystem, which underlies many biological, cultural, and recreational resource objectives. As a result, the modeling efforts scheduled indirectly support achievement of almost all other GCDAMP goals, as described in the previous section on Project PHY 07.M1.08. The ongoing activities associated with development of simulation capabilities and verification of existing models already in existence can effectively benefit from the collection of monitoring data from the downstream IQW project. These simulation models include flow routing, suspended-sediment transport, sandbar evolution, and downstream thermal simulations throughout the main channel. Having predictive capabilities for physical resources related to dam operations is potentially a valuable support tool in planning future experimental treatments, as well as evaluating proposed management actions in the river ecosystem that generally relate to GCDAMP goal 1, goal 2, goal 4, goal 6, goal 9, and goal 11. In addition, goal 12 is also supported by efforts to advance modeling activities for the ecosystem.

Aquatic Food Web Research

Both the downstream IQW monitoring project and its modeling support link to thermal and suspended-sediment transport can help to support new research focused on the food web of the river ecosystem by providing continuous data on surface flow in the main channel and major tributaries, as well as related quality-of-water data, such as water temperature, specific conductivity, dissolved oxygen, and suspended-sediment concentrations and grain size for suspended particles in transport, but also by providing simulations for predicting downstream boundary conditions that limit in-stream productivity.

Fisheries Monitoring and Research

The downstream IQW modeling activities provide support beyond IQW data by making simulations for physical habitat changes, such as backwaters, available to fishery scientists before future BHBF tests. Such information can assist scientists in planning better integrated studies.

Information Needs Addressed

The modeling support subelement of the downstream IQW directly addresses several of the RINs related to GCDAMP goals 7 and 8:

RIN 7.4.1. What is the desired range of seasonal and annual flow dynamics associated with powerplant operations, BHBFs, and habitat maintenance flows, or other flows that meet GCDAMP goals and objectives?

RIN 7.3.1. Develop simulation models for Lake Powell and the Colorado River to predict water quality conditions under various operating scenarios, supplant monitoring efforts, and elucidate understanding of the effects of dam operations, climate, and basin hydrology on Colorado River water quality.

RIN 8.5.1. What elements of ROD operations (upramp, downramp, maximum and minimum flow, MLFF, HMF, and BHBF) are most/least critical to conserving new fine sediment inputs, and stabilizing sediment deposits above the 25,000 cfs stage?

General Methods

The method used for verification of the existing flow, sediment, and thermal models will vary from one model to another, depending upon how managers and scientists propose to use the models to support planning activities. Generally, historical monitoring data will be used in combination with real or projected boundary conditions for the ecosystem (on a reach-scale basis) to determine how accurately models can recreate conditions measured around specific flow periods or events, such as the fate of Paria River sand inputs, BHBF releases, etc. For downstream temperature simulations, model behavior will be evaluated and compared to measured responses for purposes of testing and calibrating the temperature model. Additional meteorological data (if available) may also be added to the model to further evaluate performance with respect to historical patterns. Projected release

patterns for flow and temperature (from the Lake Powell model) shall also be used to evaluate future conditions of downstream temperature in the main channel and along nearshore habitats.

Products/Reports

- Development and documentation of nearshore water temperature modeling capabilities, including simplified “pond” backwater models and detailed multidimensional models of areas with available bathymetry (by the end of calendar year 2009)
- Development and documentation of multidimensional models of eddy-sandbar environments, including evaluation and summary of available datasets for sediment transport and morphology of eddy-sandbar environments (by the end of calendar year 2009)
- Preparation of conference abstracts and proceedings articles (1+ per year), journal articles (1+ per year), and presentations at GCDAMP meetings (as necessary).

Budget

PHY 7.R1.08	
Modeling Support Linked with Integrated Quality-of-Water Monitoring (FY2007–08)	
	Fiscal year 2008
GCMRC personnel costs (19% burden)	29,319
GCMRC project-related travel/training (19% burden)	5,995
GCMRC operations/supplies (19% burden)	—
GCMRC equipment purchase/replacement (19% burden)	2,000
AMP logistical support (19% burden)	—
Outside GCMRC and contract science labor (19% and/or other burden rate)	67,169
Cooperative/interagency agreements (6.09% GCMRC burden plus cooperator's burden)	5,000
Project Subtotal	109,483
DOI customer burden (combined 6.09%, 19% and/or other rates)	7,394
Project Total (Gross)	116,877
Percent outsourced (outside of GCMRC; includes 50% of logistics)	66%

GCDAMP Goal 8: Maintain or attain levels of sediment storage within the main channel and along shorelines to achieve the Adaptive Management Program ecosystem goals.

PHY 8.M1.08: Core Monitoring for the Sediment Budget and Sandbar Status throughout the CRE Utilizing Direct Topographic/Bathymetric Measurements and Remote Sensing

Start Date

October 2007

End Date

Ongoing

Principal Investigator(s)

Roderic Parnell, Matt Kaplinski, and Joseph E. Hazel, Jr., Northern Arizona University, Department of Geology; David J. Topping, U.S. Geological Survey, Grand Canyon Monitoring and Research Center.; and David M. Rubin, U.S. Geological Survey, Coastal and Marine Team

Geographic Scope

Core monitoring for the sediment budget and sandbar status throughout the CRE utilizing direct topographic/bathymetric measurements and remote sensing (SED TREND) is focused on detecting long-term (i.e., 4-year to multidecadal) trends in the CRE sediment budget for both fine (sand and finer material) and coarse sediment. In addition, this project utilizes a combination of direct topographic measurement and remote sensing to monitor the status of high-elevation (> the stage associated with a discharge of 8,000 cfs) sandbars on an annual to 4-year basis. The annual topographic measurements of high-elevation sandbars are covered under project REC 9.R1.08: Sandbar and Campable Area Monitoring under goal 9 (see below). Note that the remote-sensing aspect of the SED TREND monitoring **will not** occur during FY2008 but is scheduled as part of the 2007–11 Monitoring and Research Plan, with implementation in FY2009–10. The total geographic extent of this monitoring is from GCD to the upper end of Lake Mead (near Separation Canyon). During FY2008, SED TREND monitoring will focus on RM 0 (Lees Ferry) through RM 30; this segment of the CRE is referred to herein as upper Marble Canyon.

Project Goals/Tasks

The SED TREND monitoring directly supports achievement of the following GCDAMP goals:

Goal 8: Maintain or attain levels of sediment storage within the main channel and along shorelines to achieve AMP ecosystem goals.

Goal 9: Maintain or improve the quality of recreational experiences for users of the Colorado River ecosystem within the framework of AMP ecosystem goals. The SED TREND monitoring provides information on the size and abundance of sandbars, which are resources that affect the recreational experiences of Colorado River users.

Goal 11: Preserve, protect, manage, and treat cultural resources for the inspiration and benefit of past, present, and future generations. The SED TREND monitoring is the project that collects information on the sandbars that provide a source of sediment, through aeolian transport, to high-elevation sand deposits that contain archaeological resources.

Because SED TREND monitoring addresses the physical framework of the ecosystem, which underlies many biological resource objectives, it also indirectly supports achievement of the following AMP goals:

Goal 1: Protect or improve the aquatic food base so that it will support viable populations of desired species at higher trophic levels. The SED TREND monitoring supports this goal by providing information on coarse sediment inputs which provide the substrate for parts of the aquatic food base.

Goal 2: Maintain or attain a viable population of existing native fish, remove jeopardy for humpback chub and razorback sucker, and prevent adverse modification to their critical habitats. The SED TREND monitoring supports this goal by providing information on sandbars which create backwater habitats that are thought to be important for native fish.

Goal 6: Protect or improve the biotic riparian and spring communities within the Colorado River ecosystem, including threatened and endangered species and their critical habitat. The SED TREND monitoring monitors the status of the fine sediment deposits which provides the substrate for riparian vegetation and marsh communities.

The primary objective of the goal 8 SED TREND monitoring is to determine magnitudes and trends in fine sediment storage throughout the CRE in the main channel and eddies at all elevations, specifically broken down into bins below the stage associated with a discharge of 8,000 cfs (where over 90 percent of the fine sediment in the CRE is typically stored), between the stages associated with discharges of 8,000 and 25,000 cfs, and above the stage associated with a discharge of 25,000 cfs.

The secondary goals of this project are to determine magnitudes and trends in campsite area and distribution (this supports goal 9), backwater geometry (area plus depths) and distribution (this supports goal 2), and the availability of open dry sand on sandbars that can be transported by the wind upslope into archeological sites thereby helping preserve these resources (this supports goal 11).

These multigoal objectives are proposed to be met through the following three monitoring tasks on an annual to quadrennial basis, in keeping with managers' information needs.

Task 1. Annual Effectiveness Monitoring for Higher Elevation Sand Deposits (subsample of sandbar with emphasis on campsite areas)

Task 1 involves the monitoring of the area and volume of fine sediment above the stage associated with 8,000 cfs for subsets of sandbars and campsites throughout the CRE using conventional ground-based surveying methods. This dataset is commonly referred to as the "NAU sandbar time series" and is the longest running dataset on the state of sandbars currently available (initiated in 1990). This task is conducted in coordination with goal 9 core monitoring and will take place in the fall of each year. Task 1 is covered under project REC 9.R1.08: Sandbar and Campable Area Monitoring under goal 9 (see below).

Task 2: Repeat Systemwide Inventory of Higher Elevation Sand Deposits

Approximately every 4 years (but only in years without BHBFs, see "Schedule by task" section below for details), monitoring of systemwide area and volume of fine sediment (especially open sand) above the stage associated with a discharge of 8,000 cfs (i.e., approximately 10 percent of the fine sediment in the CRE) based on aerial overflight data (light detection and ranging [LIDAR] and orthorectified hyperspectral aerial photography). These remote-sensing data are also used to help monitor the magnitude and trends in campsite area, backwater area and distribution, the availability of open dry sand on sandbars, as well as for other resource areas such as riparian vegetation monitoring. These data will also be used to help quantify the inputs of gravel from tributaries.

These gravel inputs provide important substrate for the aquatic food web. Note that task 2 will not occur during FY2008 but would occur in coordination with the next planned remote-sensing overflight scheduled for FY2009.

Task 3. Annual Repeat Mapping of Lower Elevation Channel Sand Deposits

Annually (but only in years without BHBFs, see “Schedule by task” section for details), monitoring the area and volume of fine sediment at all elevations over long reaches using multibeam bathymetric surveys, ground-based topographic surveys, underwater video transects, and limited underwater microscope data collection for bed grain size. Ideally, this task would be performed on a systemwide basis every 5–10 years in order to estimate fine sediment budgets over timescales for which the goal 7 mass balance sediment budgets likely become inconclusive due to accumulating measurement errors. However, since it is currently logistically impossible to survey the bathymetry of the entire river in any given year, surveys will be completed annually of sections of the river with a different section surveyed each year on a rotating basis. The sections (or reaches) will correspond to the same reaches outlined in the goal 7 mass balance core-monitoring project, as follows: reach 1: RM 0 to RM 30 (upper Marble Canyon); reach 2: RM 30 to RM 61 (lower Marble Canyon); reach 3: RM 61 to RM 87 (eastern Grand Canyon); reach 4: RM 87 to RM 166 (central Grand Canyon); reach 5: RM 166 to RM 226 (western Grand Canyon).

These reach surveys will occur in the late spring and will only be completed in years without BHBFs (see “Schedule by task” section for details); thus, in the absence of BHBFs, each reach would be surveyed every 5 years, or, if BHBFs occurred on average every other year, then each reach would be surveyed on average every 10 years. The 5–10 year interval is considered by sediment scientists to be sufficient to detect long-term trends in the fine sediment budget based on changes in topography and bathymetry. Finally, since some reaches are longer than others, it is possible that some reaches will be too long to survey completely in a single river trip (e.g., reaches 4 and 5); for these reaches, available side-scan sonar data will be used to identify the portions of these reaches that are most likely to store fine sediment. It is also possible that continued technological advancements and improvements in methods will allow for complete surveys of these reaches in the future. In addition to providing key sediment budget information (i.e., the status of the fine sediment “bank account”), these data will provide information on the location and geometries of backwaters thought to be important habitat for native fish. Task 3 is the focus of this narrative proposed for FY2008 SED TREND monitoring.

The schedule for SED TREND monitoring under goal 8 is complicated by the potential for BHBFs, except for task 1 sandbar and campsite surveys which will occur annually in the fall whether or not a BHBF is scheduled (covered under project REC 9.R1.08: Sandbar and Campable Area Monitoring under goal 9). For task 2, remote-sensing missions and task 3 reach surveys, it is advantageous to have these occur in years without BHBFs so that the monitoring data are not dominated by the effects of a single BHBF (BHBF monitoring is described under a separate science plan developed by the GCMRC in 2007). Rather, remote-sensing and reach survey monitoring should represent the integral response of the system to several years of dam operations and tributary inputs. Further, logistical constraints would make it difficult to conduct the remote-sensing and reach survey core monitoring in addition to the BHBF monitoring. Thus, without knowing the exact frequency of BHBFs, it is impossible to outline the exact schedule for goal 8 SED TREND monitoring.

It is possible, though, to outline potential schedules based on assumptions regarding BHBF frequency. Table 3 presents two possible 10-year schedules for illustrative purposes. The first is the schedule in the absence of BHBFs where the exact schedule can be delineated. The second schedule assumes that BHBFs occur every other year, which would be the approximate frequency under previous triggers based on tributary sediment supply. In reality, even if the frequency were every other year on average, there would likely be periods with successive years of BHBFs and successive years without BHBFs such that the core-monitoring schedule for remote-sensing and reach surveys must be flexible.

Table 3. Two possible schedules for the completion of the tasks outlined under project PHY 8.M1.08.

Year	Schedule without BHBFs			With BHBFs every other year		
	Task 1: subsample campsites/sandbars	Task 2: 4-year over flights	Task 3: flux-reach surveys	Task 1: subsample campsites/sandbars	Task 2: 4-year over flights	Task 3: flux-reach surveys
2008	X		X (R1)	X		X (R1)
2009 (BHBF)	X	X	X (R2)	X		
2010	X		X (R3)	X	X	X (R2)
2011 (BHBF)	X		X (R4)	X		
2012	X		X (R5)	X		X (R3)
2013 (BHBF)	X	X	X (R1)	X		
2014	X		X (R2)	X	X	X (R4)
2015 (BHBF)	X		X (R3)	X		
2016	X		X (R4)	X		X (R5)
2017 (BHBF)	X	X	X (R5)	X		

Need for Task 3

Sediment forms the physical template for the CRE downstream from GCD (DOI, 1995; National Research Council, 1996). The endangered and threatened native fishes evolved in a highly turbid river (Gloss and Coggins, 2005), with turbidity predominantly due to suspended silt and clay and, to a lesser degree, suspended sand. Before the closure of GCD, 60 percent of upstream sediment supply from the Colorado River in Glen Canyon was silt and clay (Topping and others, 2000). Closure of GCD reduced the supply of silt and clay by about 96 percent at the upstream boundary of Grand Canyon National Park, with the Paria River now the major supplier of silt and clay at this location (Topping and others, 2000). The postdam Colorado River in Marble and Grand Canyons is much less turbid (with clearer water conditions than ever occurred naturally) and, because the in-channel storage of sand, silt, and clay in the postdam Colorado River is greatly reduced from predam conditions, the Colorado River in the CRE is now turbid only during periods of tributary activity downstream from the dam.

Sandbars and other sandy deposits in and along the Colorado River in Grand Canyon National Park were an integral part of the natural riverscape, and are important for riparian habitat, native fish habitat, protection of archeological sites, and recreation (Rubin and others, 2002; Wright and others, 2005). Recent work has shown that the low-elevation parts of these sandbars (< the stage associated with a discharge of 8,000 cfs) in lateral recirculation eddies contain the bulk of the sand, silt, and clay in storage (Hazel and others, 2006), and the surface grain size of these sandbars is the dominant regulator of sand transport over multiyear timescales (Topping and others, 2008). Thus, the low-elevation parts of sandbars and the channel (as will be shown below) comprise the long-term bank account or reserve for sediment in the CRE. These deposits have eroded substantially following the 1963 closure of GCD that reduced the supply of sand at the upstream boundary of Grand Canyon National Park by about 94 percent (Topping and others, 2000). In response to this reduction in sand supply and the alteration of the natural hydrograph by dam operations (Topping and others, 2003), sandbars in Marble Canyon and the upstream part of Grand Canyon have substantially decreased in size since closure of the dam (Schmidt and others, 2004) and are still in decline under normal powerplant operations at the dam (Wright and others, 2005).

A major outstanding question is whether repeated BHBFs conducted under sediment-enriched conditions (such as those that existed during the 2004 BHBF test) can result in the rebuilding and maintenance of sandbars throughout the CRE. Scour of the low-elevation eddy and channel pool environments during sand-depleted

BHBF tests, such as the 1996 Controlled Flood, is not subsequently offset by deposition of new sand under normal powerplant releases (Schmidt and others, 2004; Topping and others, 2006). Analysis of surveys conducted one to four times per year during the 1990s indicates that sandbars in Marble Canyon and the upstream part of Grand Canyon contained about 25 percent less sand at lower elevations in 2000 than in 1991, and that the lower elevation parts of these sandbars and the adjacent channel bed never fully recovered in sand volume after scouring during the 1996 flood. This net decrease in low-elevation fine sediment volume occurred despite the fact that tributary inputs of sand during this period were well above average. Thus, controlled floods conducted under sediment-depleted conditions, such as those that existed in 1996, cannot be used to sustain sandbar area and volume. In addition, the dominant response (downstream from the upstream half of Marble Canyon) during the 2004 BHBF test was that eddies lost sand (although less than was gained in upper Marble Canyon). By definition, if BHBFs are to be a successful tool for the rebuilding and maintenance of sandbars in the CRE, then the volume of fine sediment stored at lower elevations (i.e., in the long-term fine sediment reserve) must not decrease over longer timescales as a result of the occurrence of repeated BHBFs.

Computing fine sediment budgets for various reaches in the CRE over long (i.e., decadal or longer) timescales is required for evaluating the effects of dam operations, including BHBFs. Over shorter timescales (up to perhaps several years), this is best done by the “mass balance” approach described in the section on goal 7 of the Draft Core Monitoring Report. However, because of the increasing uncertainties over time associated with the mass balance approach, another approach is needed to track the fine sediment budget for the CRE over longer timescales. This other complementary approach (described herein) is required to evaluate whether future dam releases (including BHBFs) continue to mine the sediment reserve or whether the reserve (stored largely at elevations less than the stage associated with a discharge of 8,000 cfs) remains stable or increases under future dam releases. If the amount of sediment in the reserve continues to decrease, then operations will ultimately not be able to sustain the fine sediment resources at higher elevations.

At the 2004 AMWG priority-setting workshop, questions relating specifically to sediment (and tracked by the herein described SED TREND monitoring) were identified under three of the top five priorities of the AMP. These priorities were, in decreasing order of relevance to sediment:

GCDAMP Priority 4: What is the impact of sediment loss and what should we do about it?

GCDAMP Priority 3: What is the best flow regime?

GCDAMP Priority 2: Which cultural resources, including traditional cultural properties, are within the Area of Potential Effect, which should we treat, and how do we best protect them? What is the status and trends of cultural resources and what are the agents of deterioration?

Strategic Science Questions

Several SSQs were identified by scientists and managers during the KAW conducted in the summer of 2005 (Melis and others, 2006). The SED TREND monitoring project provides valuable information to help answer several of the questions related to sediment conservation, and in particular the primary sediment question:

SSQ 4-1. Is there a “Flow Only” operation (i.e., a strategy for dam releases, including managing tributary inputs with BHBFs, without sediment augmentation) that will rebuild and maintain sandbar habitats over decadal timescales?

Links/Relationships to Other Projects

The SED TREND monitoring provides data (i.e., the maps showing the topography and distribution of sediment types over about 30-mile reaches of the river) that are essential to the development and testing of numerical predictive models of discharge, stage, sediment transport, and sandbar morphology. These predictive models can be used to evaluate a wide range of resource responses, such as the fate of sandbar habitats, to various dam release scenarios, such as controlled floods, steady flows, fluctuating flows, etc.

The SED TREND monitoring provides the data used to evaluate the effectiveness of dam operations (including BHBFs) on rebuilding and maintaining sandbars in the CRE. Additionally, the SED TREND monitoring will provide the data showing whether dam operations continue to mine the long-term fine sediment reserve stored at elevations below the stage associated with a discharge of 8,000 cfs (over 90 percent of the fine sediment in the system is currently stored below this elevation). If the amount of sediment in this “bank account” continues to decrease, then operations will ultimately not be able to sustain the fine sediment resources at higher elevations.

The SED TREND monitoring supports the campsite inventories conducted under goal 9 by characterizing the status and trends of the sandbars used as campsites (covered under project REC 9.R1.08: Sandbar and Campable Area Monitoring under goal 9).

The SED TREND monitoring supports goal 11 by characterizing the status of fine sediment at higher elevations in and around cultural sites, and by characterizing the amount of open dry sand available to be transported by the wind into these cultural sites (thereby helping preserve these sites). (This aspect of this project will occur during FY2009 when the next remote-sensing mission is scheduled).

The SED TREND monitoring also supports new research focused on the food web of the river ecosystem by providing data on the input of gravel used as a substrate by the aquatic food web. (This aspect of this project will occur during FY2009 when the next remote-sensing mission is scheduled)

The SED TREND monitoring also provides information on the distribution of the fine sediment deposits that form the substrate for the riparian ecology.

Finally, the SED TREND monitoring supports science activities in the fisheries program by providing the data (as part of the long about 30-mile data collection effort described under task 3) to characterize the locations and geometries of backwaters thought to be important habitat for native fish.

Information Needs Addressed

The 2003 AMP Strategic Plan identified Core Monitoring Information Needs (CMINs) related to sediment storage (goal 8). The CMINS that are addressed by the SED TREND monitoring are listed below. For each, the prioritization ranking applied by the AMP SPG in 2006 is also included. The SED TREND monitoring during FY2008 will directly address the third of the top five goal 8 CMIN priorities; the first two of these five are addressed by the mass balance project described under goal 7.

CMIN 8.1.1. Determine and track the biennial fine sediment volume and grain-size changes in the main channel below 5,000 cfs stage, by reach (third-ranked goal 8 CMIN).

CMIN 8.4.1. Track, as appropriate, the biennial or annual sandbar area, volume, and grain-size changes within eddies between 5,000 and 25,000 cfs stage, by reach (fourth-ranked goal 8 CMIN).

CMIN 8.5.1. Track, as appropriate, the biennial sandbar area, volume, and grain-size changes above 25,000 cfs stage, by reach (fifth-ranked goal 8 CMIN).

During FY2008, the SED TREND monitoring also addresses these unranked goal 8 CMINs:

CMIN 8.2.1. Track, as appropriate, the biennial or annual sandbar area, volume, and grain-size changes outside of eddies between 5,000 and 25,000 cfs stage, by reach.

CMIN 8.3.1. Track, as appropriate, the biennial or annual sandbar area, volume, and grain-size changes within eddies below 5,000 cfs stage, by reach.

During FY2009, the SED TREND monitoring will address this unranked goal 8 CMIN:

CMIN 8.6.1. Track, as appropriate, changes in coarse sediment (> 2 mm) abundance and distribution.

The SED TREND monitoring also directly addresses this top-ranked goal 9 CMIN priority (covered under project REC 9.R1.08: Sandbar and Campable Area Monitoring under goal 9):

CMIN 9.3.1. Determine and track the size frequency, and distribution of camping beaches by reach and stage level in Glen and Grand Canyons (top-ranked goal 9 CMIN).

Developing and testing monitoring protocols for these CMINs was the primary focus of research and development conducted during FY1998–FY2006, as reviewed by SEDS-PEP III.

General Methods

During FY2008, SED TREND monitoring will focus on task 3 described above. Task 3 is conducted using standard ground-based surveying protocols and multibeam-sonar bathymetric surveying protocols (including error analyses) described in Kaplinski and others (2000, 2007, unpub. data). The grain-size data collected under task 3 (recommended by the final PEP, Wohl and others, 2006) are collected and processed using protocols described in Rubin and others (2006, in press) and Rubin (2004).

Products/Reports

Annual updates of the NAU sandbar time series showing trends in the area and volume of the high-elevation parts of sandbars, in addition to providing annual data showing the effectiveness of dam operations on rebuilding and maintaining sandbars (covered under project REC 9.R1.08: Sandbar and Campable Area Monitoring under goal 9) (by the end of calendar year 2009).

Topographic maps of the CRE in the first of five long reaches: upper Marble Canyon, lower Marble Canyon, eastern Grand Canyon, central Grand Canyon, and western Grand Canyon. During FY2008, monitoring will focus on upper Marble Canyon. These maps will be produced one to two times per decade for each reach on average. These maps will characterize the geometries of the backwaters (thought to be important habitat for native fish) in each about 30-mile reach (by the end of calendar year 2009).

Mapping conducted during FY2008 will ultimately result in decadal timescale sediment budgets for these five reaches of the CRE. These data will provide managers information on the long-term status of the fine sediment reserve. These sediment budgets will be compared to the sediment budgets computed for these reaches under the complementary mass balance project described under goal 7. This comparison will help evaluate the uncertainties associated with the SED TREND monitoring and mass balance approaches (by the end of calendar year 2009).

Where possible, data collected in Upper Marble Canyon in FY2008 will be compared with earlier multi-beam-sonar data collected in 2000, 2001, and as part of the 2002–04 FIST project to evaluate volume changes in the fine sediment reserve (2000 vs. 2008) (by the end of calendar year 2009).

Annual peer-reviewed USGS data reports documenting results of the monitoring project. Contribution to other research-related peer-reviewed publications (such as models). Biannual presentations at GCDAMP meetings and GCMRC science symposiums. (By the end of calendar year 2009).

Budget

PHY 8.M1.08	
Long-Term Monitoring of Changes in Sediment Storage (FY2008—Ongoing)	
	Fiscal year 2008
GCMRC personnel costs (19% burden)	17,700
GCMRC project-related travel/training (19% burden)	—
GCMRC operations/supplies (19% burden)	—
GCMRC equipment purchase/replacement (19% burden)	40,000
AMP logistical support (19% burden)	—
Outside GCMRC and contract science labor (19% and/or other burden rate)	—
Cooperative/interagency agreements (6.09% GCMRC burden plus cooperator's burden)	118,447
Project Subtotal	176,147
DOI customer burden (combined 6.09%, 19% and/or other rates)	18,176
Project Total (Gross)	194,323
Percent outsourced (outside of GCMRC; includes 50% of logistics)	67%

NOTE: \$63,394 of this funding is from fiscal year 2007 carry forward. \$130,929 is funded from the fiscal year 2008 GCDAMP power revenue budget.

GCDAMP Goal 9: Maintain or improve the quality of recreational experiences for users of the Colorado River ecosystem, within the framework of GCDAMP ecosystem goals.

REC 9.R1.08/PHYS 8.M2.08: Sandbar and Campable Area Monitoring

Start Date

October 2007 (This monitoring project is a continuation of monitoring efforts that have been occurring annually since 1990 for sandbar area and volume and since 1998 for campable area.)

End Date

Ongoing

Principal Investigator(s)

R. Parnell, M. Kaplinski, and J. Hazel, Northern Arizona University, Geology Department; in cooperation with Grand Canyon Monitoring and Research Center staff scientists

Geographic Scope

Sandbar and campable area monitoring has historically focused on 45 sandbars along the main channel of the Colorado River between GCD (RM -15) and Diamond Creek (RM 226). However, about five additional sites are being proposed for inclusion in this monitoring project below RM 225, downstream to the western boundary of the geographical scope of the GCDAMP program (approximately RM 278). The reach below Diamond Creek has been of increasing interest to managers due to the persistent period of lower reservoir elevations and storage in Lake Mead, and large sandbars that are now exposed along a flowing river reach. This westernmost reach of the study area is frequently used for recreational camping and boating, and additional biological studies are also underway below Diamond Creek (fishery monitoring, etc.).

Project Goals/Tasks

The goal of this project is to track change in sandbar volumes and topography and link these data to changes in campable area using established monitoring protocols while alternative monitoring approaches using remotely sensed data are being explored and tested.

The specific objectives of this study include the following:

- Measuring sandbar area, volume, and campsite area at a series of long-term monitoring sandbar sites annually
- Evaluating changes in campsite area in relation to bar volume and topography
- Evaluating changes in campsite area in relation to past monitoring results at different flow stages

Need for Project

Public concern with the ongoing loss of sandbar “beaches” and recreational capacity in the Colorado River corridor was a key factor leading to the development of the 1995 Glen Canyon Dam Final Environmental Impact Statement and passage of the Grand Canyon Protection Act (GCPA) in 1992. The protection of visitor use values

is specifically identified as a goal of GCPA. This project directly addresses one part of the top-priority core-monitoring information need (change in campsite size) for goal 9 of the GCDAMP Strategic Plan.

This project will also provide data to managers about the status and trend of sandbars throughout the CRE below GCD that have been monitored annually since 1990. Sandbar measurement data (area and volume relative to stage elevations) at these long-term monitoring sites have been reported annually to the AMWG since its formation in 1997, and were also presented to managers annually during the Glen Canyon Environmental Studies (GCES) II era of the EIS. The Strategic Plan of the GCDAMP AMWG identifies conservation of fine sediment as a desired program outcome (GCDAMP goal 8). Recently the GCDAMP (August 2004) identified sediment resources as the program's fourth-priority area of concern and interest, as articulated in the following question:

GCDAMP Priority 4: What is the impact of sediment loss and what should we do about it?

Strategic Science Questions

There is still uncertainty about the future fate of sandbars below GCD under proposed operational strategies intended to promote sand conservation of tributary inputs. The supply of new sand below the dam is estimated to be about 6 percent of the predam supply in Marble Canyon and about 16 percent of the predam supply below the confluence with the LCR (RM 61–278). NAU sandbar monitoring data is extremely useful in addressing specific SSQs and evaluating the ROD operations, as well as alternative operations being considered by managers. Monitoring data pertaining to sandbar volume change address the following SSQ:

SSQ 4-1. Is there a “Flow Only” operation (i.e., a strategy for dam releases, including managing tributary inputs with BHBFs, without sediment augmentation) that will restore and maintain sandbar habitats over decadal timescales?

In terms of questions that are specific to the recreation goal, this project also directly addresses the following SSQ:

SSQ 3-9. How do varying flows positively or negatively affect campsite attributes that are important to visitor experience?

Because campsite size, distribution, and physical attributes are known to affect visitor experience, this project also indirectly addresses two other important science questions related to recreation in the CRE:

SSQ 3-7. How do dam-controlled flows affect visitors' recreational experiences, and what is/are the optimal flows for maintaining a high-quality recreational experience in the CRE?

SSQ 3-8. What are the drivers for recreational experiences in the CRE, and how important are flows relative to other drivers in shaping recreational experience outcomes?

Links/Relationships to Other Projects

This monitoring project incorporates the long-term NAU sandbar survey project that has been underway since the early 1990s and the associated campable area surveys that have occurred annually at a subset of these sandbars since 1998. Although formerly distinguished as separate monitoring projects, one directed at goal 8 and the other at goal 9, both the NAU sandbar survey and campable area monitoring projects are concerned with monitoring sandbar sediment, albeit in different respects. The NAU sandbar survey tracks changes in total area and volume of the sandbars above the 5,000 cfs level, while the campable area monitoring project specifically evaluates changes in campable area at a subset of these sandbar sites. In combination, these two projects provide a holistic assessment of how flows are affecting the sandbar habitats used by recreational boaters for camping.

Campsite Inventory and GIS Atlas

The assessments of campable area throughout the river ecosystem will be evaluated as a subset of sites included in the campsite inventory. Data resulting from this monitoring project will be incorporated into the GIS campsite atlas that is under development in FY2007–08 (REC 9.R3.08).

In addition to recreation resources, sandbars are closely linked with other resources of GCDAMP concern, such as terrestrial and aquatic habitats related to native fish rearing areas (backwaters) and cultural site preservation, as discussed in more detail below.

Changes in Nearshore Habitats (Shoreline Types and Abundance of Backwaters)

At those study sites with well-defined return-current channels, topographic measurements made at the long-term sandbar monitoring sites also incorporate the morphology and size of backwaters. Three-dimensional topography data can therefore be used to analyze local river stage versus depth and area relationships for backwaters at these monitoring sites as one means of addressing what operational ranges of flow are most conducive to backwater size and stability. The sandbar and campable area data will be incorporated into the shoreline habitat study planned for FY2007–08 (DASA 12.D6.08).

Archaeological Sites

Generally, sandbar monitoring tracks changes in higher elevation sand areas and volumes at a subsample of sites throughout the system. The abundance of sand above the active fluctuating-flow operating zone (above 25,000 cfs stage) provides information about whether sand storage in those areas is stable, increasing or decreasing through time in response to normal operations or experimental high flows intended to promote conservation of new sand supplies. The abundance of sand along shorelines that is available for transport by wind to higher elevations where archaeological preservation sites are located is thought to be related to the potential for eroded sites to be reburied by new sand. In the future, additional process studies at such cultural sites may be tied more directly to sandbar monitoring at existing sites, as well as by adding additional monitoring sites over time that are proximal to cultural research sites.

Information Needs Addressed

Sandbar monitoring above the 5,000 cfs level directly addresses information needs specified within the “Fine Sediment” section (GCDAMP goal 8) of the GCDAMP Strategic Plan:

CMIN 8.2.1. Track, as appropriate, the biennial sandbar area, volume, and grain-size changes outside of eddies between 5,000 and 25,000 cfs stage, by reach.

CMIN 8.5.1. Track, as appropriate, the biennial sandbar area, volume, and grain-size changes above 25,000 cfs stage, by reach.

This project also directly addresses one part of the top-priority CMIN for goal 9 (campsite size):

CMIN 9.3.1. Determine and track the size, quality, and distribution of camping beaches by reach and stage level in Glen and Grand Canyons. (This project specifically addresses the part of the CMIN concerned with campsite size.)

This project partially addresses a second campsite CMIN (9.3.2) that is very closely related to the top-priority CMIN for camping beaches:

CMIN 9.3.2. Determine and track the effects of ROD operations on the size, quality, and distribution of camping beaches in the CRE.

This monitoring project will also contribute to tracking one outcome of experimental flows on camping beaches (campable area), as defined by EIN 9.3.1:

EIN 9.3.1. How do the size, quality, and distribution of camping beaches change in response to an experiment performed under the ROD, unanticipated event, or other management action?

General Methods

Repeat surveys of long-term sandbar monitoring sites have been conducted since 1990 using trained field personnel under the joint direction of the GCMRC's survey department staff and scientists from the NAU Department of Geology. Campable area survey protocols have been established and applied consistently by the same team of scientists since the late 1990s (Kaplinski and others, 2005). As described in the State of the Colorado River Ecosystem in Grand Canyon report (Kaplinski and others, 2005, p. 196), campable area surveys are conducted annually in the fall, at the conclusion of the prime river recreation season. Survey crews from NAU Department of Geology survey selected study sites using standard total station survey techniques (U.S. Army Corps of Engineers, 1994). Topographic data are collected and referenced to AZ State-Plane Coordinates generated through the GCMRC's survey control network activities throughout the CRE. Data are reduced and analyzed by the NAU team in cooperation with GCMRC partners and presented in a variety of formats, but most typically are reported as cumulative area and volume totals. The volumes and areas are also assessed relative to flow and stage elevations linked to dam operations. While methods for surveying "offshore" topography within eddies below the 5,000 cfs stage are being evaluated by external peer reviewers in FY2006–07, monitoring data will continue to be collected in shallower portions of the eddies and in the terrestrial portions of the sandbars using the established, conventional ground-survey methods. Once the protocol for measuring sandbar topography in deeper offshore areas is resolved, then information relating to CMIN 8.1.1 shall be collected at these sites (presumably starting in FY2009).

Surveyors follow the criteria of Kearsley (1995) and Kearsley and Quartaroli (1997) to identify campable area. Campable area is defined as "a smooth substrate (preferably sand) with no more than eight degrees of slope with little or no vegetation" (Kaplinski and others, 2005, p.196). Although the goal is to capture the total campable area at each site, camping areas located at considerable distance (>100 m) from the main mooring/cooking areas are generally not included in the totals. In the future, these protocols may be adjusted to measure all campable area with variable slope criteria within the National Park Service (NPS)-defined campsite boundaries using remotely sensed data (see research project description 9.R2.08 in the FY2007 work plan); however, until new protocols are tested and refined, the existing monitoring program will continue.

Products/Reports

Annual report documenting the change in sandbar topography, volume, area, and campable area will be prepared that summarize the implications of annual findings for fine sediment storage throughout the main channel. The data gathered as a result of the project will also be served through the GCMRC Web page. Project findings will also be presented at the biennial GCMRC science symposium.

Budget

REC 9.R1.08/PHY 8.M2.08	
Sandbar and Campable Area Monitoring (FY2007–11)	
	Fiscal year 2008
GCMRC personnel costs (19% burden)	9,390
GCMRC project-related travel/training (19% burden)	1,200
GCMRC operations/supplies (19% burden)	500
GCMRC equipment purchase/replacement (19% burden)	—
AMP logistical support (19% burden)	15,000
Outside GCMRC and contract science labor (19% and/or other burden rate)	14,788
Cooperative/interagency agreements (6.09% GCMRC burden plus cooperator's burden)	92,500
Project Subtotal	133,378
DOI customer burden (combined 6.09%, 19% and/or other rates)	13,400
Project Total (Gross)	146,778
Percent outsourced (outside of GCMRC; includes 50% of logistics)	86%

REC 9.R3.08: Compile Campsite Inventory and GIS Atlas (Year 2)

Start Date

October 2006

End Date

September 2008

Principal Investigator(s)

Project will be jointly funded and jointly implemented by staff from Grand Canyon National Park and Grand Canyon Monitoring and Research Center; Linda Jalbert, Outdoor Recreation Planner, Grand Canyon National Park; and Helen Fairley, Sociocultural Program Manager, Grand Canyon Monitoring and Research Center

Geographic Scope

Entire Colorado River ecosystem, from base of Glen Canyon Dam to Lake Mead (RM 277)

Project Goals/Tasks

The goal of this project is to compile a comprehensive current inventory of campsites in the CRE and document the spatial extent, geographic distribution, and associated attributes of these campsites in a GIS atlas. The atlas will document attributes of current campsites that are important to recreation experience and that have the potential to be affected by flows (e.g., campable area, amount of open sand area, type and amount of vegetation cover, and mooring characteristics under varying flows.) The atlas will also document locations and attributes of past campsites that have disappeared due to loss of sediment and/or vegetation encroachment. The atlas will serve as an electronic repository for all data (e.g., repeat photographs, campable area survey data, vegetation transect data, etc.) that have been collected for each campsite over the past few decades.

This inventory and atlas will serve as the baseline for future monitoring and research projects. It will define the boundaries of current campsites in a GIS environment so that future evaluations that rely on remotely sensed data and statistical samples to quantify change in campsite attributes relative to dam operations have a common spatial basis for evaluating change through time. The atlas will have broad utility for both NPS recreation managers (e.g., Colorado River Management Plan [CRMP] monitoring), as well as for monitoring effects of dam operations on campsites.

Need for Project

Baseline inventories provide the foundation for long-term monitoring programs and research studies. Comprehensive campsite inventories in the CRE conducted initially in 1973 were repeated in 1984 (Weeden and others, 1975; Brian and Thomas, 1984). The last comprehensive campsite inventory was completed 15 years ago in 1991 (Kearsley and Warren, 1993). The 1991 inventory showed a dramatic decline in number and size of campsites compared with previous inventories (Kaplinski and others, 2003). A new comprehensive inventory is needed (Kaplinski and others, 2003, 2005; Loomis and others, 2005) to document the current number, size, and distribution of campsites throughout the CRE and to document the boundaries of the areas that NPS proposes to manage as campsites in the future. This database will serve as a baseline and will document the total pool of sites that will be sampled for various research and monitoring projects in the future. This atlas will also serve as the central repository for all campsite data collected during future inventory and monitoring projects. The 2005 recreation PEP identified this as the highest priority research need under management objective 9.3.

Strategic Science Questions

Primary SSQ directly addressed:

SSQ 3-9. How do varying flows positively or negatively affect campsite attributes that are important to visitor experience?

Indirectly, this project will also provide information that is relevant for addressing a second SSQ about the effects of flows on the quality of recreational experience in the CRE:

SSQ 3-8. What are the drivers for recreational experiences in the CRE, and how important are flows relative to other drivers in shaping recreational experience outcomes?

Links/Relationships to Other Projects

This project is being undertaken in cooperation with staff from Grand Canyon National Park. In addition to meeting GCDAMP needs, data from this project will be used by the NPS as they develop implementation plans and resource monitoring projects tied to the Colorado River Management Plan. Because the NPS has immediate need for some campsite data, \$40,000 in equipment and NPS staff salaries is being contributed by NPS in FY2006 to initiate the project.

The GIS atlas will serve as the definitive source for information on prior and current campsite inventory data. It will provide a foundation and repository for all future research and monitoring projects related to CRE campsites. In addition to documenting the areas used for recreational camping, the GIS campsite layer will document areas of the CRE most heavily impacted by humans. This information will be useful for assessing human impacts rates on near by cultural resources such as archaeological sites and TCPs.

Information Needs Addressed

This project will lay the foundation for future research and monitoring efforts that are designed to address management objective 9.3 and the top-priority CMIN for goal 9:

CMIN 9.3.1. Determine and track the size, quality, and distribution of camping beaches by reach and stage level in Glen and Grand Canyons.

CMIN 9.3.1 is very closely related to a second CMIN under M.O. 9.3

CMIN 9.3.2. Determine and track the effects of ROD operations on the size, quality, and distribution of camping beaches in the CRE.

The current recreation monitoring program is only focused on one aspect of CMIN 9.3.1: campsite size. Note that this project will allow for the tracking of the other key relevant campsite variables, e.g., campsite distribution and quality. This project will also have utility for monitoring effects of experimental flows on camping beaches (campable area), as defined by EIN 9.3.1.

EIN 9.3.1. How do the size, quality, and distribution of camping beaches change in response to an experiment performed under the ROD, unanticipated event, or other management action?

General Methods

- Using existing published sources (e.g., Stevens, 1992; Martin and Whitis, 2004) and the knowledge of experienced river guides, we will identify and map all currently used campsites in the CRE. (NPS task)

- Using existing sources and the knowledge of experienced river guides, the campsite boundaries (as defined by NPS managers) will be documented in a GIS environment. (NPS)
- Campsite boundaries will be field checked and verified. (NPS/GCMRC)
- Campsite attributes that are important to visitor experience (substrate characteristics, mooring characteristics, protection from prevailing winds, proximity to attraction sites) will be identified and documented. (GCMRC/NPS)
- Using published information from prior inventories (e.g., Weeden, 1975; Brian and Thomas, 1984; Kearsley and Warren, 1993) all former campsite locations and associated information will be identified and integrated into the GIS atlas. (GCMRC/NPS)
- Supporting documents and photos will be scanned and linked to GIS/spatial data (document legacy metadata). (NPS/GCMRC)
- Using established slope/area/attribute criteria, current campable areas within the campsite boundaries will be classified to assess current carrying capacity. (NPS/GCMRC)

Products/Reports

A comprehensive inventory of campsites and associated legacy data will be documented and published in an electronic GIS atlas as the final product of this project.

Budget (for Year 2 of the 2-Year Project)

REC 9.R3.08	
Compile Campsite Inventory and GIS Atlas (FY2007–08)	
	Fiscal year 2008
GCMRC personnel costs (19% burden)	19,132
GCMRC project-related travel/training (19% burden)	2,000
GCMRC operations/supplies (19% burden)	2,000
GCMRC equipment purchase/replacement (19% burden)	—
AMP logistical support (19% burden)	15,000
Outside GCMRC and contract science labor (19% and/or other burden rate)	12,000
Cooperative/interagency agreements (6.09% GCMRC burden plus cooperator's burden)	25,000
Project Subtotal	75,132
DOI customer burden (combined 6.09%, 19% and/or other rates)	11,047
Project Total (Gross)	86,179
Percent outsourced (outside of GCMRC; includes 50% of logistics)	59%

*In FY2006, NPS contributed \$40,000 in staff salary, equipment purchases, and supplies in order to get this project started. In FY2007, NPS contributed approximately \$60,000 in staff salaries, equipment purchases, logistics and supplies. These NPS contributions are NOT reflected in the budget shown above.

GCDAMP Goal 10: Maintain power production capacity and energy generation, and increase where feasible and advisable, within the framework of the Adaptive Management ecosystem goals.

HYD 10.M1.08: Monitor Power Generation and Market Values under Current and Future Dam Operations

Start Date

October 2006

End Date

Ongoing

Principal Investigator(s)

Data will be provided by Western Area Power Administration and distributed via the Grand Canyon Monitoring and Research Center Web site

Geographic Scope

Hydropower generation data and market values for the energy generated by Glen Canyon Dam

Project Goals/Tasks

The goal of this core-monitoring project is to monitor and document hourly hydropower generation and potential opportunity (replacement) costs under current and future flow regimes.

Need for Project

Power generated at GCD is marketed mostly in six western states by the Department of Energy's Western Area Power Administration (WAPA). WAPA's primary mission is to sell power from Federal water project powerplants under statutory criteria in the Reclamation Project Act of 1939, the Flood Control Act of 1944, and the Colorado River Storage Project (CRSP) Act of 1956. These criteria include the following:

- Preference in the sale of power must go to municipalities, public corporations, cooperatives, and other nonprofit organizations.
- Power must be marketed at the lowest possible rates consistent with sound business practices.
- Revenues generated from power sales must pay for power generation and all allocated investment costs under the original CRSP Act.
- Projects should generate the greatest amount of power and energy that can be sold at firm power and energy rates, consistent with other project purposes.

Tracking generation (as impacted by operations for other project purposes), power market rates, necessary power purchases, and Basin Fund cash flow provides the means to assess the impact of changes in GCD operations in relation to the four statutory criteria.

Currently, there are no ongoing core-monitoring activities related to goal 10. Although data on GCD hydropower generation and opportunity costs under MLFF operations are currently being gathered by Reclamation and WAPA as routine agency functions, these data are not readily accessible to the GCDAMP. The need for this information in a readily accessible format has been identified as a program need, and this project will help to fill this critical information gap.

Strategic Science Questions

Primary SSQs addressed:

SSQ 3-3. What are the annual hydropower replacement costs of the MLFF since 1996?

SSQ 3-4. What are the projected hydropower costs associated with the various alternative flow regimes being discussed for future experimental science (as defined in the next phase experimental design)?

Links/Relationships to Other Projects

This project is directly linked to the newly proposed adaptive management assessment initiative proposed for goal 12. It also is specifically related to the current overall long-term planning needs of the GCDAMP.

Information Needs Addressed

This project responds to the core-monitoring information need for goal 10, as originally articulated in the 2003 version of the GCDAMP Strategic Plan, and redefined by the SPG:

IN 10.1. Determine and track the impacts to power users from implementation of ROD dam operations and segregate those effects from other causes such as changes in the power market.

CMIN 10.1.1 (as redefined by SPG). Determine and track the marketable capacity and energy produced through dam operations in relation to the various release scenarios (hourly/daily/monthly volumes, daily fluctuation limit, upramp and downramp rates and limits, etc.).

General Methods

WAPA and Reclamation continuously schedule and monitor power generation to meet anticipated and real-time power demand. This information is available on an hourly time step reported daily, weekly, and monthly from System Control and Data Acquisition (SCADA) data. WAPA and its customers track power source, availability, and market changes on an hourly basis in assessing the need, cost, and accessibility for additional power resources to meet contractual obligations or unanticipated demand. Market pricing, resulting cost of purchases, and the impact on Basin Fund cash flow are recorded in the WAPA Energy Tracking Database (ISA). This information is reported monthly and annually and is available through WAPA-CRSP, but not publicly published. Table 4 summarizes the metrics and frequency of data collection for power costs.

Table 4. Metrics and frequency of data collection for power costs.

Objective	Parameters	Methods	Location(s)	Frequency	Accuracy & Precision
Monitor monthly energy generation	MW	SCADA	SCADA Phoenix – Dumped Energy Management System (ISA)	Hourly	N/A
Monitor hourly power market price	\$/MWH	WAPA Energy Tracking Database (ISA)	WAPA – Montrose	Hourly	N/A
Monitor monthly firming power purchases	\$ and MW purchased	WAPA Energy Tracking Database (ISA)	WAPA-Montrose	Monthly	N/A
Monitor monthly Basin Fund Balance	\$	WAPA Energy Tracking Database (ISA)	WAPA-CRSP	Monthly	N/A

Data Sources

Energy generated: The SCADA system that measures generation at GCD is reported to a database that is accessible by the WAPA Phoenix office. Currently, those data are dumped into the CRSP-Montrose office ISA, and from ISA monthly generation is calculated by summing all the hourly values. Hourly generation totals are not currently reported but can be accessed by WAPA-CRSP or WAPA-Montrose. For the purposes of this project, hourly data will be reported.

Hourly market prices: Market prices vary at different purchase points throughout the system. The price that WAPA-Montrose pays for power is pertinent to WAPA and its customers. This value is available only for the hours in which WAPA buys or sells power; therefore, the dataset is incomplete. If complete data is needed by WAPA-Montrose, they may look at the Dow Jones for a representative point of sale and record that data price. These data can be accessed via the Web and reported to an Excel spreadsheet if access is requested and granted by WAPA-Montrose.

Basin fund balance: The financial manager for the CRSP office completes an end-of-month cash balance and Basin Fund balance report found on WAPA’s Web site. The reports are usually completed by the 15th of the month. These data will be for the previous month’s billing on the 2 months previous services.

Monthly firming purchases: These data is found in the WAPA-Montrose TDB database. Purchases made by WAPA for customers are reported by the 10th of the following month, broken out by customer (purchased from). This report is sent to WAPA and can be made available.

Products/Reports

Hourly data will be collected by WAPA and delivered to the GCMRC on a daily basis. These data will be served through the GCMRC Web site. Monthly data will be delivered to the GCMRC at the conclusion of each month.

Budget

HYD 10.M1.08	
Monitor Power Generation and Market Values under Current and Future Dam Operations (FY2007–Ongoing)	
	Fiscal year 2008
GCMRC personnel costs (19% burden)	15,465
GCMRC project-related travel/training (19% burden)	—
GCMRC operations/supplies (19% burden)	500
GCMRC equipment purchase/replacement (19% burden)	—
AMP logistical support (19% burden)	—
Outside GCMRC and contract science labor (19% and/or other burden rate)	—
Cooperative/interagency agreements (6.09% GCMRC burden plus cooperator's burden)	—
Project Subtotal	15,965
DOI customer burden (combined 6.09%, 19% and/or other rates)	3,033
Project Total (Gross)	18,998
Percent outsourced (outside of GCMRC; includes 50% of logistics)	0%

* GCMRC salaries are for setting up Web site and connections to receive and deliver the data.

GCDAMP Goal 11: Preserve, protect, manage, and treat cultural resources for the inspiration and benefit of past, present, and future generations.

CUL 11.R1.08: Research and Development towards Core Monitoring, Phase II

Start Date

October 2007

End Date

September 2010

Principal Investigator(s)

Individual tasks will be accomplished using a combination of Grand Canyon Monitoring and Research Center personnel and outside contractors and/or cooperators. It is anticipated that the National Park Service will assist with the pilot monitoring efforts; other components will involve a combination of university cooperators, U.S. Geological Survey researchers, and independent contractors.

Geographic Scope

Colorado River ecosystem as defined in the Glen Canyon Dam Adaptive Management Program Strategic Plan

Project Goals/Tasks

This cultural monitoring project is part of a phased program of research and development towards implementation of a long-term core-monitoring program. The first phase of this project (FY2006–07) focused on conducting research and development (R&D) for refinement of monitoring protocols. More specifically, the focus of the first 2 years of this project was on completing a comprehensive assessment of the geomorphic and archaeological attributes of sites to aid in the development of the long-term monitoring approach and testing field protocols to be included in a long-term, core-monitoring program for archaeological resources in the CRE.

FY2008 will be the first year of a 3-year monitoring cycle employing the refined protocols developed during the preceding R&D phase. In FY2008 we will continue to build on several R&D activities initiated in FY2006, including (1) continuing to gather data on several short-term, small-scale studies to evaluate the effectiveness, efficiency, and accuracy of various field measurement techniques before implementing them as part of a long-term monitoring program (including weather monitoring, LIDAR mapping, and thalweg survey measurements at a subset of sites); (2) continuing to monitor check dam effectiveness at a subset of sites and expanding this monitoring to additional sites in the CRE; and (3) applying the preliminary results from the first 2 years of the R&D work toward defining and implementing a pilot monitoring project for the next 3 years. The ultimate outcome of this R&D effort will be a final report with specific monitoring protocol recommendations. The program will ultimately be subject to a final review by a PEP in late FY2010 or early in FY2011, with additional refinement of protocols (if necessary) before being implemented as the long-term program.

The scope of this project encompasses the full range of archaeological resources in the Colorado River corridor during the time of human occupation. The actual number of archaeological sites that will be included in the pilot monitoring program will be determined upon completion of the assessment phase of this project.

This project does not address R&D for monitoring of tribally valued resources other than archaeological sites, because in FY2006–07, the six affiliated tribes participating in the GCDAMP are reviewing and defining their monitoring data needs, with the aim of ensuring that the values of importance to each tribe are clearly identified and addressed in future tribal and nontribal monitoring efforts. This initial phase of tribal monitoring program definition is underway through sole-source contracts between Reclamation and the tribes. Integration of these planning efforts into the core-monitoring program will depend on the focus of monitoring projects proposed by the tribes, and will be accomplished after completing the initial research and development phase of these projects, during implementation of the pilot monitoring phase.

Need for Project

The FY2000 cultural PEP recommended redesigning the 1999–2000 programmatic agreement monitoring program to focus more specifically on tracking effects of dam operations and evaluating the efficacy of erosion control efforts (Doelle, 2000). Subsequently, the SPG and Cultural Resources Ad Hoc Group (CRAHG) redefined the primary core-monitoring need for historic properties to track status and trends of site condition and integrity through monitoring rates of erosion, visitor impacts, and other variables or processes known to affect archaeological site condition. This project will explore and test various options for measuring change and achieving these defined monitoring objectives, before implementing a long-term core-monitoring program.

Grand Canyon is one of the classic erosional landscapes of the world, and to some degree erosion of unconsolidated deposits along the Colorado River corridor is inevitable. Yet many cultural resources are being damaged by rapid gully erosion, and recent studies have shown that erosion of the sediment that forms the context of cultural sites has increased in the past few decades (Hereford and others, 1993). Previous research raised several basic questions that are ongoing issues in the river corridor: (1) What are the geomorphic controls and other environmental factors contributing to gully erosion, and what are the ultimate causes of this gully erosion? (2) What is the effectiveness of installed erosion-control measures? (3) Are there accurate, low-impact, cost-efficient monitoring methods that can replace the qualitative assessments and high-impact ground surveys used in the past? Results of recent research by Pederson and others (2003) indicate that the exploration of remote-sensing options for monitoring could potentially be redirected from photogrammetry to high intensity LIDAR. Also, erosion-control efforts—brush check dams in particular—appear to be effective at slowing erosion, but results thus far are from a single-year study, and a longer term assessment is needed to help narrow the focus of future post-treatment monitoring approaches.

Monitoring of the deposition and erosion of sediment at archaeological sites along the Colorado River corridor in Grand Canyon has been done mostly through qualitative observation documented with repeat photography. This approach has been supplemented by total station ground surveys at a select number of sites in the river corridor. Although the total station survey method is highly accurate and precise, it is labor intensive and expensive for long-term, frequent monitoring of multiple sites. Perhaps more importantly for cultural resource management, intensive survey monitoring has its own erosional impacts through significant trampling of cryptobiotic crusts and trailing. Research findings by Pederson and others (2003) showed that erosion is primarily focused at knickpoints and channel heads, and it also indicated that monitoring could be effective with a relatively limited analysis of thalweg and channel cross-section profiles rather than full-terrain total station surveys. These preliminary findings will be tested and evaluated as part of this research effort towards establishing long-term monitoring protocols for archaeological sites.

Since conclusion of the Pederson study, the GCMRC has tested C survey technology for tracking sandbar changes along the Colorado River corridor. This state-of-art technology has potential advantages over photogrammetry or total station surveys of topography by being significantly less labor intensive to produce, having lower technician error or bias, and lower overall impacts to the terrain. Initial indications are that the accuracy of LIDAR data in this setting is at least as good as that of the photogrammetry reported in Pederson and others (2003) (Mike Breedlove, personal communication), but this technology has not been tested for its utility in tracking gully erosion, nor has its repeat accuracy been rigorously determined. Although traditional ground survey will be employed in this project, alternative remote-sensing methods for monitoring treatment effectiveness (and erosion) at archaeological sites in the future will also be explored.

Strategic Science Questions

Primary SSQs addressed:

SSQ 2-1. Do dam-controlled flows affect (increase or decrease) rates of erosion, and vegetation growth, at archaeological sites and TCP sites in the CRE, and if so, how?

SSQ 2-4. How effective are various treatments (e.g., check dams, vegetation management, etc.) in slowing rates of erosion at archaeological sites over the long term?

Links/Relationships to Other Projects

This project is linked to the treatment planning effort that was initiated by Reclamation in FY2006. Specifically, it will extend the site assessment process initiated for treatment planning purposes to include assessments of sites that appear to be stable at this time, but that could be affected by the propagating effects of dam operations in the future. It will also build upon a pilot research project conducted by Utah State University (USU) in FY2006 to assess effectiveness of check dams; this will be accomplished by formalizing and extending the USU study for a second year to assess the utility of monitoring geomorphic change using similar measurement protocols as those being piloted in the FY2006 study.

As noted above, opportunities for integrating the results of this R&D effort with those of the tribal monitoring projects will be explored after completing the initial R&D phase of these projects. This delay in integration is necessary in order for the needs and approaches of the tribal monitoring programs and the Federal agencies to be articulated and appropriate protocols identified. Integration of monitoring efforts, as appropriate, will occur during implementation of the pilot monitoring phase (FY2008–10).

This project builds upon the work of Draut and Rubin (2005, 2006) by incorporating weather monitoring at a small number of sites on a pilot basis as one of the long-term monitoring protocols to be evaluated. This study is also linked to the NPS CRMP implementation efforts, in that monitoring protocols for assessing impacts of human visitation at archaeological sites are being developed cooperatively with NPS to serve the monitoring data needs of both GCDAMP and the CRMP.

Information Needs Addressed

This project is an R&D effort aimed at addressing the highest priority CMIN for historic properties (as revised by the CRAHG and SPG in fall 2005), specifically, the properties known as archaeological sites:

CMIN 11.1.1 (SPG revised). Determine the condition and integrity of prehistoric and historic sites in the CRE through tracking rates of erosion, visitor impacts, and other relevant variables. Determine the condition and integrity of TCPs in the CRE.

It will also directly address CMIN 11.1.2 of the GCDAMP Strategic Plan (renumbered by CRAHG/SPG as EIN 11.1):

EIN 11.1. Determine the efficacy of treatments for mitigation of adverse effects to historic properties.

This project also addresses a RIN (no number) (formerly identified as CMIN 11.1.4 in the GCDAMP Strategic Plan):

How effective is monitoring, what are the appropriate strategies to capture change at an archaeological site—qualitative, quantitative?

General Methods

Task 1. Analyze the assessment data collected in fy2006–07 and cluster data based on geomorphic characteristics and archaeological attributes and values

The assessment of archaeological sites for the development of the long-term monitoring program began in FY2006. Work has consisted of assembling, evaluating, and verifying legacy information (NPS monitoring data) regarding the archaeological resources and updating the information where warranted. In FY2006, this work was coordinated with the site-specific assessments being conducted by USU geomorphologists Dr. Joel Pederson and Mr. Gary O'Brien, and Dr. Jonathan Damp from Zuni Cultural Resources Enterprise for Reclamation's Section 106 Grand Canyon treatment plan development. In FY2007, assessment work continued at the remaining sites not proposed for inclusion in the treatment plan. The goal of the assessment phase was to ensure that accurate, up-to-date, comparable levels of information existed for all of the potentially monitored archaeological sites. Uniform baseline data is critical for selecting a statistically valid sample of the sites to be used in the long-term core-monitoring program.

Variables assessed for each site included the attributes and characteristics that contribute to site significance (elements of integrity as reflected in the nature of artifact assemblage, numbers and types of constructed features, presence and extent of subsurface cultural deposits, specific research values, and association with historical events or people). A concurrent assessment process evaluated the geomorphic context and attributes that affect site stability and/or degree of erosion (e.g., topographic setting, gully catchment characteristics, resilience of the substrate to erosion, degree to which individual gullies are integrated with the river).

In fall 2007, following completion of the final 2007 data collection effort and its integration within a database, the resulting data will be clustered using Gower's coefficient for mixed variables types (Gower, 1971) and Ward's minimum variance clustering algorithm (Ward, 1963), or a similarly suitable method, in order to group sites for defining future monitoring protocols that are relevant to particular groups of sites and for stratifying the site population for future sampling purposes. Clusters will be analyzed in a GIS to detect spatial distributional patterns and potential biases. Samples will then be drawn from each of the clusters for the pilot monitoring effort, which is scheduled to commence in spring 2008. Subtask allocation for compiling, exploring, and analyzing these data; integrating them with the other data collected under task 2 (see below); evaluating all data in a GIS environment; and preparing the final pilot monitoring plan is \$44,600.

Task 2. Implement pilot monitoring protocols for geomorphic change detection and erosion control effectiveness monitoring

In order to test and evaluate quantitative monitoring protocols, approximately six study sites (each study site consisting of two to three archaeological sites located in close proximity to each other) were repeatedly mapped and intensively evaluated in FY2006–07. As of May 2007, these data were being compiled and analyzed. In FY2008, the same study sites evaluated in FY2006–07, plus approximately six additional locations, will be intensively surveyed to quantify rates of gully erosion and topographic surface change occurring throughout the canyon in a variety of geomorphic settings. Specific study sites will be determined based on the results of the geomorphic assessment analysis that will be completed following the final FY2007 field session.

Continue Collecting Field Data on Gully Erosion Rates

Previous research findings by Pederson and others (2003) showed that gully erosion is clearly focused at knickpoints and channel heads, and it also indicated that monitoring could potentially be effective with a relatively limited analysis of thalweg and cross-sectional channel profiles rather than full-terrain surveys. This study is building upon Pederson's prior research for the purpose of testing and evaluating the most appropriate and cost-effective methods to measure geomorphic change at archaeological sites and also to evaluate the effectiveness of erosion control devices that may be installed at various sites in the future.

Consistent with previous years, field data collection will occur two times per year, once before the monsoon season and once after the monsoon in the late fall. Basic geomorphic data related to thalweg evolution will be collected, and repeat photographs of check dam and gully features will be taken to track changes coincident with

hydrologic events or other disturbances. Data collected will include field observations of piping and overland flow features, integrity of check dams, and evidence of past and recent aeolian activity. The effectiveness of installed check dams will continue to be evaluated by comparison of monitoring data from FY2008 to previous thalweg and topographic data collected at the same localities, as well as to unmitigated gullies at nearby sites. Criteria for effectiveness are (1) whether check dams remain competent and in place, (2) whether measurements and/or photographs reveal they successfully trap and store sediment, and (3) whether surveys document that channel widening and knickpoint recession are constrained or do not happen at all during runoff events.

Care will be taken to prevent gully wall failure and disturbance around cultural sites during ground surveys, and gully surveys will be limited to the essential data provided by gully thalweg profiles and topography and at select cross-sectional channel profiles at major knickpoints. This will minimize the amount of trampling of study sites by intensive surveys outside of already-disturbed gully channels. Subtask allocation for collecting and analyzing gully thalweg measurements and preparing a report on findings is \$85,000, not including GCMRC personnel costs. This funding will be provided to USU through a cooperative agreement.

Continue to Monitor Topographic Change and Establish New Baseline Topographic Records

In FY2008, we will continue to develop data pertaining to topographic change at archaeological sites using a combination of conventional total station mapping and ground-based high-density LIDAR data at selected study sites. Both LIDAR-produced digital terrain models and ground-survey data will be georeferenced and provided by the GCMRC. Total station ground surveys will be directed by GCMRC personnel following methods employed by previous GCMRC researchers for capturing topographic changes using high-density data collection methods (e.g., Yeatts, 1996; Hazel and others, 2000; Pederson and others, 2003). LIDAR data will be manually edited and filtered to produce a “bare-earth” terrain model without reflections from vegetation canopy. In FY2008, approximately six additional sites with high potential for topographic change (based on the geomorphic assessment results) will be added to the sample of sites being monitored in this fashion. Subtask budget allocation for LIDAR survey work is \$70,000, not including GCMRC personnel costs. This funding will be provided to USGS Western Coastal Geology and Marine Division through an internal USGS suballotment.

Weather Monitoring

In FY2007, 10 weather stations were established at 8 study sites in the CRE. The study sites include the same ones where gully measurements and LIDAR surveys are occurring, plus two additional sites. In FY2008–10, these stations will continue to monitor precipitation amount and intensity, wind direction and velocity, temperature, humidity, barometric pressure, and sediment transport rates. Because of the spatially isolated nature of monsoon thunderstorms and the significant role that precipitation and wind play in downcutting and backfilling gullies, weather stations and sand traps have been placed in proximity to the study sites where intensive mapping and monitoring of erosion control effectiveness is occurring, so that changes detected from repeat topographic mapping can potentially be related to timing and duration of local or regional weather events. These stations are outfitted with automated data loggers that can store up to 3 months of data. We are also in the process of exploring telemetry and automated sand traps as a means to facilitate data collection. Subtask budget allocation for weather monitoring is \$113,000. This includes equipment replacement and technical maintenance, data processing, quality control, and analysis. The equipment maintenance, data collection, and processing tasks are being managed internally by the GCMRC; data analysis will be handled through an internal USGS suballotment to USGS Western Coastal Geology and Marine Division.

Site Condition Evaluations

Concurrent with the gully thalweg surveys and topographic mapping work, data will be collected on surface indicators of condition using a standardized recording format. These data will document a combination of indicators that reflect both geomorphic and human agents of change affecting site condition in the CRE. The specific recording formats will vary, depending on the type of site being monitored, which be determined from the clustering exercise to be conducted in fall and winter 2007, following completion of the field assessment work in September and October 2007. Subtask budget allocation for the condition indicator monitoring task is \$70,000, not including GCMRC personnel costs. This amount will be allocated to the NPS for participation in the field work activities and compiling data in a database.

Products/Reports

Several peer-reviewed reports will be prepared at the conclusion of this study. Each report will focus on a specific element/task of the R&D project:

Report on analysis of FY2006–07 assessment data and development of site clusters from the archaeological and geomorphic assessment data

Sample selection for the pilot monitoring project

LIDAR maps and gully thalweg measurements from sites surveyed in FY2006–07 and preliminary survey data from FY2008

Summary of first-year pilot project monitoring results (winter 2009)

Budget

CUL 11.R1.08	
Research & Development toward Core Monitoring (FY2007)	
	Fiscal year 2008
GCMRC personnel costs (19% burden)	73,585
GCMRC project-related travel/training (19% burden)	3,700
GCMRC operations/supplies (19% burden)	7,500
GCMRC equipment purchase/replacement (19% burden)	16,000
AMP logistical support (19% burden)	40,000
Outside GCMRC and contract science labor (19% and/or other burden rate)	118,000
Cooperative/interagency agreements (6.09% GCMRC burden plus cooperator's burden)	172,000
Project Subtotal	430,785
DOI customer burden (combined 6.09%, 19% and/or other rates)	37,224
Project Total (Gross)	468,009
Percent outsourced (outside of GCMRC; includes 50% of logistics)	72%

CUL 11.R2.08: Implement Tribal Monitoring Projects

Start Date

To be decided, following TWG approval of the individual tribal monitoring plans

End Date

September 2008

Principal Investigator(s)

To be decided by each tribe

Geographic Scope

Colorado River ecosystem

Project Goals/Tasks

The goal of this project is to collect data on tribally valued resources, including culturally valued elements of the terrestrial ecosystem and TCPs, and evaluate their condition in concordance with the individual tribe's perspectives and value systems.

Need for Project

Tribal stakeholders (i.e., the Hopi Tribe, Hualapai Nation, Kaibab Band of Paiute Indians, Navajo Nation, Paiute Tribe of Utah, and Pueblo of Zuni) have expressed interest in participating in the evaluation of CRE resources in a manner consistent with tribal concerns and value systems. The tribal stakeholders maintain that current monitoring approaches based exclusively on Western science paradigms do not adequately capture tribal interests or concerns. In response to these issues, in FY2006 the GCDAMP funded the tribes to articulate their concerns more explicitly, design monitoring approaches that will more fully meet their needs, and bring their proposed monitoring programs forward to TWG for GCDAMP consideration and formal approval. This FY2008 project has been funded at the request of the TWG, subject to the stipulation that the tribes first complete their current contractual agreements with Reclamation to develop and report on their proposed monitoring programs and present them to TWG for formal approval.

Strategic Science Questions

Primary SSQ addressed:

SSQ 2-7. Are dam-controlled flows affecting TCPs and other tribally valued resources in the CRE, and if so, in what respects are they being affected, and are those effects considered positive or negative by the tribes who value these resources?

Links/Relationships to Other Projects

The tribes' interests in the CRE are broad, encompassing both cultural-historical sites and biological elements. Many archaeological sites in the CRE are also TCPs for individual tribes. Thus, the monitoring program for archaeological sites, which is driven in large measure by Western science interests in historical information preservation, overlaps with the interests of tribes, who share a concern for retaining these cultural landmarks for a variety of different reasons. The archaeological site monitoring project and tribal monitoring projects are currently being developed separately but on parallel tracks with the understanding that once areas of mutual concern have been identified, we will seek ways to reduce monitoring cost and field effort and improve efficiencies for all programs by combining monitoring efforts where feasible.

The tribes also have a long-standing interest in the condition of traditionally valued plants and animal resources. These interests are often place specific, in that the cultural value of biological resources may be enhanced by their association with TCPs. In FY2008, the tribes will be asked to participate in the terrestrial ecosystem PEP and provide an overview of their monitoring approaches and existing terrestrial ecosystem monitoring (TEM)-related data for potential incorporation into long-term TEM protocols. Again, the concept is to identify areas of mutual interest between western scientific approaches and tribal concerns, so that opportunities for reducing monitoring costs and improving program efficiency can be identified, while at the same time, ensuring that information relevant to tribal interests are obtained as part of the long-term core-monitoring program.

Information Needs Addressed

This project is directly responsive to the highest priority core-monitoring information need for cultural resources, as revised by the Cultural Resource Ad Hoc Group and the SPG:

CMIN 11.1.1. Determine the condition and integrity of prehistoric and historic sites in the CRE through tracking rates of erosion, visitor impacts, and other relevant variables. Determine the condition and integrity of TCPs in the CRE.

It is also directly responsive to the second highest priority CMIN for cultural resources:

CMIN 11.2.1. Determine the condition of traditionally important resources and locations using tribal perspectives and values.

General Methods

Monitoring methods are being determined by each tribe in conjunction with completing their current (FY2006-funded) contractual obligation with Reclamation.

Products/Reports

Before this project is initiated, a formal written report and oral presentation will be provided by each tribe to TWG in FY2008 describing each tribe's proposed monitoring approach for FY2008 and beyond.

An annual report documenting the assumptions, methods, annual outcome, and relationship of annual monitoring results to long-term status and trends of tribally valued resources, will be produced by each participating tribe at the conclusion of this study.

Budget (See Reclamation project C.5)

CUL 11.R2.08	
Implement Tribal Monitoring Projects	
	Fiscal year 2008
GCMRC personnel costs (19% burden)	—
GCMRC project-related travel/training (19% burden)	—
GCMRC operations/supplies (19% burden)	—
GCMRC equipment purchase/replacement (19% burden)	—
AMP logistical support (19% burden)	—
Outside GCMRC and contract science labor (19% and/or other burden rate)	—
Cooperative/interagency agreements (6.09% GCMRC burden plus cooperator's burden)	—
Project Subtotal	—
DOI customer burden (combined 6.09%, 19% and/or other rates)	—
Project Total (Gross)	—
Percent outsourced (outside of GCMRC; includes 50% of logistics)	0%
NOTE: See funding in the Reclamation Budget Section.	

*No funding needed in FY2008

GCDAMP Goal 12: Maintain a high-quality monitoring, research, and adaptive management program

DASA 12.D1.08: Preparation for Monitoring Data Acquisition (remote sensing)

Start Date

October 2007

End Date

September 2008 (ongoing annually to support quadrennial, systemwide overflights)

Principal Investigator(s)

Glenn Bennett, Data Acquisition, Storage, and Analysis Program Manager, Grand Canyon Monitoring and Research Center; Thomas Gushue, GIS Coordinator, Grand Canyon Monitoring and Research Center; and Michael Breedlove, Ph.D., Geographer, Utah State University

Geographic Scope

Entire Colorado River ecosystem corridor from forebay of Glen Canyon Dam to upper Lake Mead

Project Goals/Tasks

Revision of current processing protocols, technology comparison to determine applicability and efficient resolutions for area, volume, and classification techniques, FY2009 mission planning, and contract solicitation.

Need for Project

Although no remote-sensing missions are currently planned until FY2009, the DASA program is tasked with preparation for the next scheduled canyonwide overflight.

A primary fiscal objective is to reserve sufficient funding to cover mission costs during implementation in 2009. Additionally, the data collection permit must be reviewed and updated through Grand Canyon National Park to reflect the types of remote-sensing technologies that will be required to help fulfill the core-monitoring and experimental research needs for all GCMRC programs. During FY2008, mission planning and contract solicitation will begin for the next canyonwide data collection effort. Currently, no salaries are funded for this project in FY2008, and so work performed in this realm will be addressed by GIS personnel funded by the GIS General Support Project (DASA 12.D5.08). Due to the dependent nature of remote-sensing and GIS technologies, products described in this project will result from a combination of efforts across other DASA projects.

Additionally, and as time permits, an evaluation of existing remotely sensed data previously collected by the GCMRC will be conducted to determine the appropriateness of different monitoring techniques and required data inputs to achieve desired accuracies for future core-monitoring and research efforts in support of sediment storage, vegetation mapping, habitat classification, and cultural site studies. This undertaking will involve an in-depth simulation testing of data densities (resolutions) and editing requirements of inputs from a variety of sources including multiband imagery, LIDAR, topographic data, hydrographic data and digital surface models. Two key aspects that this project will investigate are as follows:

- A revision of current processing protocols of these datasets that have previously resulted in a massive amount of manual editing before analysis.

- An exploration of remotely sensed data at different resolutions (i.e., density of points) in comparison to final surfaces and classifications to determine the most applicable and efficient resolutions needed to achieve the necessary output with a minimum of error.

Particular attention will be given to datasets collected in conjunction with the 5-year research and development project (2001–06) for monitoring sand storage changes; however, additional canyonwide sand analysis will be conducted and statistical tabulations provided in support of resource monitoring for the cultural and biological programs. Included in this will be an extension of the 2002 sand/campsite analysis up through 2005 with particular emphasis given to effects of experimental flows on camping beaches.

Strategic Science Questions

The airborne data to be collected are multispectral orthorectified images of the CRE. Area and volumetric analysis of these datasets are used to identify and classify elements of interest. Comparison of datasets acquired over time allow for change detection as long as the data continue to be collected. Airborne data is the basis for many of the science questions and research activities conducted in the Grand Canyon. Sandbar habitat change including vegetation encroachment, shoreline location and character at different flow regimes and the distance to cultural sites, backwater existence and changes, and maps used for positioning GCMRC monitoring areas are a few of the applications of airborne data. Some of the resource areas and science questions identified during the 2005 Knowledge Assessment and found within the GCMRC's Strategic Science and Monitoring and Research Plans (see appendix A) that can be addressed with airborne image data include those listed below.

Other Strategic Science Questions

Additional SSQs addressed:

SSQ 4-1. Is there a “Flow-Only” operation (i.e., a strategy for dam releases, including managing tributary inputs with BHBFs, without sediment augmentation) that will restore and maintain sandbar habitats over decadal timescales?

SSQ 5-1. How do dam release temperatures, flows (average and fluctuating component), meteorology, canyon orientation and geometry, and reach morphology interact to determine mainstem and nearshore water temperatures throughout the CRE?

SSQ 1-7. Which tributary and mainstem habitats are most important to native fishes and how can these habitats best be made useable and maintained?

SSQ 2-1. Do dam-controlled flows affect (increase or decrease) rates of erosion and vegetation growth at archaeological sites and TCP sites, and if so, how?

SSQ 2-2. How do flows impact old high-water zone terraces in the CRE (where the majority of archaeological sites occur), and what kinds of important information about the historical ecology and human history of the CRE are being lost due to ongoing erosion of the Holocene sedimentary deposits?

SSQ 3-9. How do varying flows positively or negatively affect campsite attributes that are important to visitor experience?

Links/Relationships to Other Projects

Acquisition of systemwide digital images in this project supports addressing numerous resource questions within other programs, such as abundance and systemwide distribution of both aquatic and terrestrial habitats related to fish, vegetation, and availability and status of campsites along the CRE. The digital products procured by the DASA directly support a varied array of projects within GCDAMP goals 1–11, such as detecting shoreline habitat and changes tied to dam operations and high-flow tests. Additionally, these data are used in terrestrial vegetation and sandbar mapping projects for determining surface texture and land cover classifications within designated study reaches, as well as canyonwide over subsequent years following the overflights (trend analysis).

Information Needs Addressed

Numerous GCDAMP goals and resource area programs that are concerned with remote-sensing analysis are the chief beneficiaries.

IN 12.1. Develop information that can be used by the TWG, in collaboration with the GCMRC, to establish current and target levels for all resources within the GCDAMP as called for in the GCDAMP strategic plan.

CMIN 4.1.6. Determine quantity and quality of spawning habitat for RBT in the Lees Ferry reach as measured at 5-year intervals.

CMIN 6.1.1. Determine and track the abundance, composition, distribution, and area of the marsh community as measured at 5-year or other appropriate intervals based on life cycles of the species and rates of change for the community.

CMIN 6.4.1. Determine and track composition, abundance, and distribution of the sand beach community as measured at 5-year or other appropriate intervals based on life cycles of the species and rates of change for the community.

CMIN 9.3.1. Determine and track the size, quality, and distribution of camping beaches by reach and stage level in Glen and Grand Canyons.

RIN 6.1.1. How has the abundance, composition, distribution, and area of the marsh community changed since dam closure (1963), high flows (1984), interim flows (1991), and the implementation of ROD operations (1996)?

RIN 8.6.1. How do ongoing inputs of coarse-sediment from tributaries influence storage of fine sediment within pools, runs and eddies throughout the CRE?

EIN 4.1.1. How does RBT abundance, proportional stock density, length at age, condition, spawning habitat, natural recruitment, whirling disease and other parasitic infections change in response to an experiment performed under the Record of Decision, unanticipated event, or other management action?

EIN 6.1.1. How do marsh community abundance, composition, distribution, and area change in response to an experiment performed under the Record of Decision, unanticipated event, or other management action?

EIN 6.4.1. How do the abundance, composition, and distribution of the sand beach community change in response to an experiment performed under the Record of Decision, unanticipated event, or other management action?

EIN 9.3.1. How do the size, quality, and distribution of camping beaches change in response to an experiment performed under the Record of Decision, unanticipated event, or other management action?

In total, approximately one-third of the GCDAMP information needs may be directly or indirectly addressed through analysis and use of the systemwide digital imagery.

General Methods

Existing remote-sensing datasets of different resolutions and point densities will be evaluated by DASA staff on the basis of recommendations that were produced during the remote-sensing research and development initiative (2000–05). DASA staff will determine if there are significant changes in volumetric calculations of surfaces and between various resource classifications, based on the density of values within the data.

Products/Reports

- Report on horizontal accuracy assessment of digital imagery datasets (2002 and 2005) using the National Spatial Reference System (NSRS) 2007 adjusted control point database. (End of 3d quarter, FY2008).
- A report comparing and assessing usefulness of remote-sensing technologies used for area and volume calculations of fine-grained sediment sites. (End of 4th quarter, FY2008).

If above products are completed ahead of schedule and as time permits, DASA staff will develop assessments of vertical accuracies achieved through various remote-sensing technologies, including automated photogrammetry, airborne LIDAR, ground-based LIDAR, NAU topography data for areas above 8,000 cfs, and multibeam sonar bathymetry and scanning hydrographic operation airborne **LIDAR** survey (SHOALS) for below 8,000 cfs. This will include developing various scenarios of data collection for the CRE considering lessons learned from previous data collection efforts. Ideas such as determining how to collect data only in areas known or expected to change (e.g., 8,000–25,000 cfs, or 8,000–45,000 cfs) will be explored. Topics covered in this assessment will include accuracies of datasets, geographic coverage/extent and cost for each method of data collection, results and limitations/operational constraints, and a discussion of expected advances in remote-sensing technologies for both those that are listed here as well as any new technologies that may have been developed or are currently being developed by the remote-sensing industry.

Budget

DASA 12.D1.08	
Preparation for Monitoring Data Acquisition (Remote Sensing; FY2007–Ongoing)	
	Fiscal year 2008
GCMRC personnel costs (19% burden)	—
GCMRC project-related travel/training (19% burden)	—
GCMRC operations/supplies (19% burden)	—
GCMRC equipment purchase/replacement (19% burden)	—
AMP logistical support (19% burden)	—
Outside GCMRC and contract science labor (19% and/or other burden rate)	218,487
Cooperative/interagency agreements (6.09% GCMRC burden plus cooperator's burden)	—
Project Subtotal	218,487
DOI customer burden (combined 6.09%, 19% and/or other rates)	41,513
Project Total (Gross)	260,000
Percent outsourced (outside of GCMRC; includes 50% of logistics)	100%

DASA 12.D2.08: Grand Canyon Integrated Oracle Database Management System

Start Date

October 2007

End Date

September 2008, and ongoing annually

Principal Investigator(s)

Glenn Bennett, Data Acquisition, Storage, and Analysis Program Manager, Grand Canyon Monitoring and Research Center; and Paul Alley, Database Administrator, Grand Canyon Monitoring and Research Center

Geographic Scope

Entire Grand Canyon Monitoring and Research Center study area, from the forebay of Lake Powell to upper Lake Mead

Project Goals/Tasks

The goal of the database management system at the GCMRC is to provide an organized, secure, and readily available electronic repository for all scientific data collected in the ongoing research and monitoring activities of the center. The Relational Database Management System (RDBMS) also serves as the electronic storage foundation of the Center's GIS, providing the repository for all aerial photography, survey control, and geographic layers. The program is therefore a vital component of the decision support process and for the adaptive management of the GCD. In support of these goals, the following are tasks will be completed during FY2008:

- Electronically archive all incoming datasets in their original form.
- Error check and import newly collected datasets to the centralized RDBMS.
- Administer database, including backup, recovery, and security.
- Continue to consolidate and import legacy data to the system.
- Continue to support data acquisition, import, and analyses by disciplines such as fish and water sampling in the Colorado River, and survey control.
- Extend database structure to incorporate newly acquired datasets, such as aquatic food base and daily downstream water quality.
- Develop routines to automate the process of error checking and importing datasets.
- Extend Web application architecture to distribute newly collected datasets.
- Provide data analysis support for scientific monitoring and research analyses.

Need for Project

This project establishes the electronic repository and tools necessary to analyze and interpret scientific data collected by the center, thereby providing a fundamental support service to GCMRC scientific investigations and decision support processes.

Strategic Science Questions

This project provides the foundation for all projects concerned with scientific data analysis.

Links/Relationships to Other Projects

Most programs generate datasets that will be archived, served, and analyzed using DASA database services. The best example of the power and utility of the Oracle database is its ability to handle terabytes of data generated of multiple years, such as those data that are associated with systemwide airborne digital imagery.

Information Needs Addressed

Provides access for analysis for all GCMRC datasets.

IN 12.1. Develop information that can be used by the TWG, in collaboration with the GCMRC, to establish current and target levels for all resources within the GCDAMP as called for in the GCDAMP strategic plan.

RIN 12.3.1. As necessary, investigate the most effective methods to integrate and synthesize resource data.

RIN 12.5.4. What is the most effective way to distribute information to our stakeholders and the public in a secure and accessible fashion?

General Methods

Working with data stewards from each scientific program at the GCMRC, the integrated database design will be extended in modular fashion to accommodate both newly collected data, such as with aquatic food base monitoring, and legacy data that have yet to be imported into the RDBMS. This process involves extensive review of existing datasets as well as current data collection protocols, and the information needs of each discipline. As these information needs are fully understood by programming staff, applications will be written that enable users to extract related datasets from the RDBMS and perform appropriate analyses. Generally these applications are written with a Web interface, as this technology provides the greatest flexibility and availability.

Products/Reports

- Fine-grained sediment transport module and Web application (3d quarter FY2008)
- Aquatic food base module and Internet/Intranet application (4th quarter FY2008)
- Downstream water quality and temperature Web application (4th quarter FY2008)

If above products completed ahead of schedule, the following products will be produced as time permits:

- Terrestrial biology database module and Web application
- KAS database module
- Survey control point module and Web application
- Stanton repeat photography Web application
- Extended database design document to include new datasets
- Metadata Web application

Budget

DASA 12.D2.08	
Grand Canyon Integrated Oracle Database Management System (FY2007–Ongoing)	
	Fiscal year 2008
GCMRC personnel costs (19% burden)	97,590
GCMRC project-related travel/training (19% burden)	1,500
GCMRC operations/supplies (19% burden)	2,000
GCMRC equipment purchase/replacement (19% burden)	—
AMP logistical support (19% burden)	—
Outside GCMRC and contract science labor (19% and/or other burden rate)	29,000
Cooperative/interagency agreements (6.09% GCMRC burden plus cooperator's burden)	22,434
Project Subtotal	152,524
DOI customer burden (combined 6.09%, 19% and/or other rates)	26,083
Project Total (Gross)	178,607
Percent outsourced (outside of GCMRC; includes 50% of logistics)	34%

DASA 12.D3.08: Library Operations

Start Date

October 2007

End Date

September 2008, ongoing annually

Principal Investigator(s)

Glenn Bennett, Data Acquisition, Storage, and Analysis Program Manager, Grand Canyon Monitoring and Research Center

Geographic Scope

Entire Grand Canyon Monitoring and Research Center study area—forebay of Glen Canyon Dam and upper Lake Mead

Project Goals/Tasks

Library operations facilitate monitoring and research by providing a centralized repository for hard copy information such as books, reports, maps, photography, and videos.

Need for Project

The GCMRC library acts as the physical repository for reports and data generated by GCMRC scientists as well as materials related to the Colorado River, Grand Canyon and Adaptive Management.

Strategic Science Questions

This project provides a research resource to aid in answering science questions.

General Methods

The library catalogs all new materials that come from staff scientists, contractors, and cooperators as well as items related to Grand Canyon, the Colorado River, and Adaptive Management. Library staff provide support to cooperators, contractors, and staff scientists by researching and obtaining current and legacy articles and reports related to science projects.

Library operations facilitate monitoring and research by providing a centralized repository for hard copy information such as books, reports, maps, photography, and videos.

Links/Relationships to Other Projects

This project supports all other projects.

Information Needs Addressed

The library provides access to current and historical scientific findings of the GCDAMP.

RIN 12.5.4. What is the most effective way to distribute information to our stakeholders and the public in a secure and accessible fashion?

Products/Reports

- Online library catalog which provides access to more than 8,000 publications continually updated
- Catalog records of all materials—continually updated
- Monthly update of new reports received in the library
- Assistance to cooperators, stakeholders, media contacts, and the public by providing access to reports, aerial photos, maps, slides, and photos in hard-copy and digital form
- Research in locating contemporary and legacy materials
- A research facility for researchers, GCMRC employees, cooperators, and the public

Budget

DASA 12.D3.08	
Library Operations (FY2007–Ongoing)	
	Fiscal year 2008
GCMRC personnel costs (19% burden)	29,628
GCMRC project-related travel/training (19% burden)	—
GCMRC operations/supplies (19% burden)	6,200
GCMRC equipment purchase/replacement (19% burden)	—
AMP logistical support (19% burden)	—
Outside GCMRC and contract science labor (19% and/or other burden rate)	—
Cooperative/interagency agreements (6.09% GCMRC burden plus cooperator's burden)	—
Project Subtotal	35,828
DOI customer burden (combined 6.09%, 19% and/or other rates)	6,807
Project Total (Gross)	42,635
Percent outsourced (outside of GCMRC; includes 50% of logistics)	0%

DASA 12.D4.08: Legacy Analog Data Conversion (Analog to Digital—Reports and Imagery)

Start Date

October 2007

End Date

September 2008, and ongoing through 2012

Principal Investigator(s)

Glenn Bennett, Data Acquisition, Storage, and Analysis Program Manager, Grand Canyon Monitoring and Research Center; and Esther Quinn, Computer Assistant, Grand Canyon Monitoring and Research Center

Geographic Scope

Entire Grand Canyon Monitoring and Research Center study area—forebay of Glen Canyon Dam and upper Lake Mead

Project Goals/Tasks

The library has undertaken a project to convert all materials in the library to digital format and make them accessible and searchable on the GCMRC Web site. Having materials available through the Web site will allow multiple users to access data concurrently from remote locations as well as protect unique items from damage or loss. Overflight imagery digitally available for spatial analysis will extend the historical spatial record allowing change detection throughout the CRE.

Need for Project

The conversion project will allow for greater access to and protection of legacy and current materials.

Strategic Science Questions

This project provides a research resource for answering spatially defined science questions and extending the period of record of digitally available overflight imagery.

Links/Relationships to Other Projects

This project supports projects concerned with spatial change over time.

Information Needs Addressed

IN 12.1. Develop information that can be used by the TWG, in collaboration with the GCMRC, to establish current and target levels for all resources within the GCDAMP as called for in the GCDAMP strategic plan.

CMIN 6.1.1. Determine and track the abundance, composition, distribution, and area of the marsh community as measured at 5-year or other appropriate intervals based on life cycles of the species and rates of change for the community.

RIN 6.1.1. How have the abundance, composition, distribution, and area of the marsh community changed since dam closure (1963), high flows (1984), interim flows (1991), and the implementation of Record of Decision operations (1996)?

RIN 6.4.1. How have the abundance, composition, and distribution of the sand beach community changed since dam closure (1963), high flows (1984), interim flows (1991), and the implementation of Record of Decision operations (1996)?

EIN 6.1.1. How do marsh community abundance, composition, distribution, and area change in response to an experiment performed under the Record of Decision, unanticipated event, or other management action?

General Methods

- Scanning and converting paper reports into digital PDF files, making the documents searchable by using optical character recognition software, and then posting the files in the library database on the GCMRC Web site
- Scanning all analog aerial film and photos using the Vexcel Ultrascan 5000, allowing the digital results to be used for 2-D and 3-D change detection
- Digitizing flight line maps to provide a searchable mechanism to locate individual scanned aerial photos
- Converting VHS tapes to DVDs
- Scanning all legacy slides to create digital images using the Nikon SuperCoolScan scanner

Products/Reports

- Access to 17,652 aerial photographs, 9,000 digital aerial images, 8,000 hard-copy reports, 8,000 photos and slides, and 700 videos in broadcast and VHS format. In addition, once the library scanning project is complete, this information will be available in digital format from the library via digital media such as DVD and online via the Web.
- As these conversion products are produced, they are cataloged and made available: see DASA 12.D3.08: Library Operations.

Budget

DASA 12.D4.08	
Legacy Analog Data Conversion (Analog to Digital—Reports & Imagery; FY2007–11)	
	Fiscal year 2008
GCMRC personnel costs (19% burden)	56,232
GCMRC project-related travel/training (19% burden)	—
GCMRC operations/supplies (19% burden)	5,475
GCMRC equipment purchase/replacement (19% burden)	—
AMP logistical support (19% burden)	—
Outside GCMRC and contract science labor (19% and/or other burden rate)	—
Cooperative/interagency agreements (6.09% GCMRC burden plus cooperator's burden)	5,000
Project Subtotal	66,707
DOI customer burden (combined 6.09%, 19% and/or other rates)	12,029
Project Total (Gross)	78,736
Percent outsourced (outside of GCMRC; includes 50% of logistics)	7%

DASA 12.D5.08: GIS General Support for Integrated Analyses and Projects, GIS Lead

Start Date

October 2007

End Date

September 2008, ongoing annually

Principal Investigator(s)

Glenn Bennett, Data Acquisition, Storage, and Analysis Program Manager, Grand Canyon Monitoring and Research Center; and Thomas Gushue, GIS Coordinator, Grand Canyon Monitoring and Research Center

Geographic Scope

Entire Colorado River ecosystem corridor between Glen Canyon Dam and Lake Mead, and the greater Colorado River Basin

Project Goals/Tasks

Specialized maps, advanced spatial analysis, and intuitive data retrieval

Need for Project

The traditional role of the GIS program is inherently service oriented, providing spatial database development and programming and analysis support to the science programs and their cooperators on both a planned and an as-needed basis. To continue functioning in this capacity it is imperative to factor in designated blocks of time to maintain and in some cases improve the level of GIS support. GIS general support benefits core monitoring, experimental programs, and research and development projects alike in the form of GIS and remote-sensing software installation, maintenance and support, creation and maintenance of spatial databases used by science projects, and the development of mapping and analysis tools for use by GCMRC staff and cooperators across all resource programs. There is also a need for a higher level of support for more specific GIS application development and analysis of available spatial data. This higher level of support is often achieved through automation of data processing and manipulation procedures to standardize and streamline repetitive tasks as well as provide a basis for standard operating procedures.

Strategic Science Questions

The spatial aspects of Grand Canyon investigations are addressed in this project.

Links/Relationships to Other Projects

Most GCMRC projects have a spatial component tied to the data being collected in support of the science questions developed for each project. The GIS provides a stable platform upon which all data collected along the CRE are catalogued within a consistent spatial reference system. At the most basic level, this allows for the overlaying and querying of datasets collected from any and all projects within the GCMRC.

Information Needs Addressed

Classification, inventory, and change detection of geomorphic, biological, and cultural areas and volumes.

IN 12.1. Develop information that can be used by the TWG, in collaboration with the GCMRC, to establish current and target levels for all resources within the GCDAMP as called for in the GCDAMP strategic plan.

RIN 12.3.1. As necessary, investigate the most effective methods to integrate and synthesize resource data.

RIN 12.5.4. What is the most effective way to distribute information to our stakeholders and the public in a secure and accessible fashion?

General Methods

The collection of spatial data is achieved through a variety of methods that include, but are not limited to, remote-sensing data collection missions, traditional survey and global positioning system (GPS) operations, field mapping using hard-copy map or pen tablet computers, onscreen digitizing using previously collected remote-sensing data as source information, and through other standard data entry methods. Spatial data are generally stored in one of the standard ESRI file types (shape file, coverage, geodatabase) as well as in ASCII format. Methods used for spatial data processing and analysis will vary depending on the questions that need to be answered.

Products/Reports

As a result of GIS support, a wide range of products will be produced:

- Maps for publications; generation and printing of maps and graphics for posters
- Creation of improved base maps for Lake Powell and Grand Canyon
- Instructional sessions for staff, cooperators, and contractors on GIS layer development, integration and analysis
- Advanced spatial analysis for monitoring projects

Budget

DASA 12.D5.08	
Geographic Information Systems (GIS) Support for Integrated Analyses and Projects, GIS Lead (FY2007–Ongoing)	
	Fiscal year 2008
GCMRC personnel costs (19% burden)	91,721
GCMRC project-related travel/training (19% burden)	3,000
GCMRC operations/supplies (19% burden)	7,317
GCMRC equipment purchase/replacement (19% burden)	—
AMP logistical support (19% burden)	—
Outside GCMRC and contract science labor (19% and/or other burden rate)	—
Cooperative/interagency agreements (6.09% GCMRC burden plus cooperator's burden)	100,000
Project Subtotal	202,038
DOI customer burden (combined 6.09%, 19% and/or other rates)	25,477
Project Total (Gross)	227,515
Percent outsourced (outside of GCMRC; includes 50% of logistics)	49%

DASA 12.D6.08: Integrated Analysis and Modeling—Mapping Shoreline Habitat Changes

Start Date

October 2007

End Date

September 2008, and ongoing through FY2011

Principal Investigator(s)

Glenn Bennett, Data Acquisition, Storage, and Analysis Program Manager, Grand Canyon Monitoring and Research Center; Thomas Gushue, GIS Coordinator, Grand Canyon Monitoring and Research Center; Timothy Andrews, Geographic Information Systems Engineer, Utah State University; and Michael Breedlove, Ph.D., Geographer, Utah State University

Geographic Scope

Entire Colorado River ecosystem corridor between forebay of Glen Canyon Dam and upper Lake Mead

Project Goals/Tasks

Shoreline habitat classification and change detection of shoreline habitat for the following years: 2002, 2004, and 2005.

Task 1: Review, refine and implement edits to Surficial Geomorphic Classes

Task 2: Develop a draft coverage of vegetated and nonvegetated GIS layers for both 2002 and 2005

Task 3: Define methods for backwater delineation and onscreen digitizing of backwater polygons, including a spatial dataset inventory of backwaters for 2002 and 2005 imagery datasets

Task 4: Conduct change detection, statistical analysis, and tabulations

Need for Project

A wealth of remote-sensing data have been collected over the past few years in support of various core-monitoring and experimental programs within the GCMRC. However, the full value of these data have yet to be realized due to a lack of time between consecutive data collection missions to process these data into more usable information. Currently, a need exists to utilize these data to study the shoreline environment along the CRE downstream of GCD. A baseline dataset of shoreline habitat currently exists as a linear classification of six habitat types at 8,000 cfs for the year 2000. Three other remote-sensing datasets exist for the years 2002, 2004, and 2005 which will be used to extend the time series of the shoreline habitat for a 5-year period. Additionally, a need exists to expand this classification into higher stages (above 8,000 cfs up to at least 45,000 cfs) in an effort to better correlate shoreline habitat with fish data and recreation habitat data also collected by the GCMRC and its cooperators. The original classification scheme for the shoreline will be extended to include backwater habitats, providing an update to the existing USU backwater dataset up to the year 2005 (Goeking and others, 2005). In addition to the classification effort, an automated suite of methods could be developed to facilitate shoreline change detection across a range of stages.

Strategic Science Questions

Primary SSQs addressed:

SSQ 3-9. How do varying flows positively or negatively affect campsite attributes that are important to visitor experience?

SSQ 1-4. Is there a “Flow-Only” (nonsediment augmentation) operation that will restore and maintain sandbar habitats over decadal timescales?

SSQ 4-2. How important are backwaters and vegetated shoreline habitats to the overall growth and survival of YoY and juvenile native fish? Does the long-term benefit of increasing these habitats outweigh short-term potential costs (displacement and possibly mortality of young humpback chub) associated with high flows?

Other science questions:

- What is the rate of change in eddy storage (erosion) during time intervals between BHBFs?
- What are the most appropriate methods for detecting change in shoreline habitat along the entire CRE given the available datasets collected using different technologies (scanned analog vs. digital), different platforms (Leica ADS-40/ISTAR vs. DMC/3001, Inc.), and different image resolutions (30 cm, 22 cm, or 18 cm)? What is the most appropriate scale/minimum mapping unit to map the shoreline habitat for all years in order to support related science questions?
- What level of change can be detected in shoreline habitat using remotely sensed data collected in the past 5 years? What changes have occurred to the shoreline habitat across the CRE in the past 5 years?
- Where have the most significant changes taken place in shoreline habitat along the CRE in the past 5 years, and within which shoreline habitat classes are the most noticeable changes? How does the shoreline habitat relate to backwater environments/habitats? What have been the changes in backwater abundance/size/shape over the past 5 years?
- As historical analog overflights become available in digital format, can the timeline be extended back to previous years?

A time-series comparison of shoreline characteristics may prove quite useful for the following SSQ:

SSQ 1-7. Which tributary and mainstem habitats are most important to native fishes and how can these habitats best be made useable and maintained?

A time-series comparison of backwater change in size and existence/nonexistence of habitats may answer questions of species abundance due to changing availability of useable habitat. Other changes in shoreline characteristics may provide insight on nonbackwater habitats utilized in different lifecycle stages.

Links/Relationships to Other Projects

A number of projects in the past few years have used the shoreline habitat data developed from the March 2000 imagery dataset. Shoreline habitat type has been used in conjunction with native and nonnative downstream fish sampling in the mainstem of the Colorado River, and it has also been used as a guide to delineate sampling sites of redds in Glen and Marble Canyons. Similarly, this data is currently being incorporated into the new aquatic food base initiative at the GCMRC. This layer has also been applied to studies of the terrestrial environment including the vegetation mapping project and initial campsite monitoring efforts conducted over the past 2 years. It is expected that new, more recent classifications will be used in similar fashion for future analysis.

Information Needs Addressed

Primary information needs addressed:

IN 12.1. Develop information that can be used by the TWG, in collaboration with the GCMRC, to establish current and target levels for all resources within the GCDAMP as called for in the GCDAMP strategic plan.

CMIN 4.1.6. Determine quantity and quality of spawning habitat for RBT in the Lees Ferry reach as measured at 5-year intervals.

CMIN 8.2.1. Track, as appropriate, the biennial sandbar area, volume, and grain-size changes outside of eddies between 5,000 and 25,000 cfs stage, by reach.

CMIN 8.4.1. Track, as appropriate, the biennial or annual sandbar area, volume, and grain-size changes within eddies between 5,000 and 25,000 cfs stage, by reach.

CMIN 8.5.1. Track, as appropriate, the biennial sandbar area, volume, and grain-size changes above 25,000 cfs stage, by reach.

CMIN 9.3.1. Determine and track the size, quality, and distribution of camping beaches by reach and stage level in Glen and Grand Canyons.

RIN 6.1.1. How has the abundance, composition, distribution, and area of the marsh community changed since dam closure (1963), high flows (1984), interim flows (1991), and the implementation of Record of Decision operations (1996)?

EIN 6.1.1. How do marsh community abundance, composition, distribution, and area change in response to an experiment performed under the Record of Decision, unanticipated event, or other management action?

EIN 6.4.1. How does the abundance, composition, and distribution of the sand beach community change in response to an experiment performed under the Record of Decision, unanticipated event, or other management action?

EIN 9.3.1. How do the size, quality, and distribution of camping beaches change in response to an experiment performed under the Record of Decision, unanticipated event, or other management action?

General Methods

Advanced remote sensing and GIS techniques will be applied to several datasets. Interaction with GCMRC researchers will guide final products in terms of “cutoff” points for certain physical interpretations.

Products/Reports

Spatial databases, spatial analysis results, and associated metadata:

- Surface texture classifications for entire river corridor (GCD to Lake Mead) for years 2002 and 2005. This will include spatial datasets representing coarseness (rough vs. smooth) including sand/no sand data (end of 3d quarter, FY2008). Surface texture classification for 2004 data may possibly have a higher level of noise in the imagery and will likely require different texture algorithms due to nature of the dataset (scanned analog vs. digitally collected) (end of 4th quarter, FY2008 for 2004 dataset).
- First-level normalized difference vegetation index (NDVI) for entire river corridor for 2002 and 2005 imagery datasets using the near-infrared and red bands of the remote-sensing imagery. Resultant GIS layers will show generalized vegetated vs. nonvegetated areas canyonwide for both datasets (end of 3d quarter, FY2008). This may require a review by Barbara Ralston, GCMRC Terrestrial Biologist, and/or comparison to vegetation mapping results in order to ensure that vegetation index thresholds used for this project comply with the vegetation mapping project.

- Preliminary surficial geomorphic classifications (cliff, talus, debris fan, and cobble bar) for entire river corridor as based on existing classification scheme (March 2000 shoreline). Resultant GIS polygons will represent these broad geomorphic classes (end of 4th quarter, FY2008). Completion of final geomorphic classifications is contingent upon the ability of those involved with the project to perform field verification on preliminary classification polygons and will require integrating into another funded river trip, potentially sometime during FY2008.
- Final composite shoreline classification scheme and computer programs for generating surface texture and surficial geomorphic classifications and shoreline habitat attributes. Methodology and procedures will be documented for further application with future remote-sensing data collections (end of 4th quarter, FY2008).

If above products are completed ahead of schedule, the following products will be produced as time permits:

- Update and extend USU backwater time series through year 2005. GIS polygon layer will represent inventory for interpretable backwater areas from 2002, 2004 and 2005 imagery datasets.
- Nearshore habitat classifications and statistical summaries for selected flow regimes in the CRE between Lees Ferry and Diamond Creek. In order to do canyonwide flow regimes, more stage discharge elevation data are needed for Glen Canyon and western Grand Canyon below Diamond Creek. Currently, Hydrologic Engineering Center River Analysis System (HEC-RAS) cross sections developed by Chris Magirl do not exist for these reaches. Also, future analysis of flow regimes will be dependent upon completion of the Magirl HEC-RAS documentation and peer review as well as any need for reprocessing of virtual shorelines for use in statistical summaries of nearshore habitat classifications.

Budget

DASA 12.D6.08	
Integrated Analysis and Modeling—Mapping Shoreline Habitat Changes (FY2007–08)	
	Fiscal year 2008
GCMRC personnel costs (19% burden)	—
GCMRC project-related travel/training (19% burden)	—
GCMRC operations/supplies (19% burden)	—
GCMRC equipment purchase/replacement (19% burden)	—
AMP logistical support (19% burden)	—
Outside GCMRC and contract science labor (19% and/or other burden rate)	—
Cooperative/interagency agreements (6.09% GCMRC burden plus cooperator's burden)	109,236
Project Subtotal	109,236
DOI customer burden (combined 6.09%, 19% and/or other rates)	6,652
Project Total (Gross)	115,888
Percent outsourced (outside of GCMRC; includes 50% of logistics)	100%

Logistics, Support, and Control

SUP 12.S1.08: Logistics Base Costs

Start Date

Ongoing

End Date

Ongoing

Principal Investigator(s)

Carol Fritzing, Logistics and Survey Program Manager, Grand Canyon Monitoring and Research Center

Geographic Scope

Entire Colorado River ecosystem corridor between Glen Canyon Dam and Lake Mead, and the greater Colorado River Basin

Project Goals/Tasks

Provide logistical support for GCMRC projects

Need for Project

The GCMRC will provide complete logistical support for 25 to 40 research, monitoring, and administrative river trips through the Grand Canyon annually. These trips range in length from 7 to 21 days and from 4 to 36 people in size. Trips will utilize a variety of motor- and oar-powered boats operated by contracted boat operators. Projects operating in the Glen Canyon reach of the Colorado River (GCD to Lees Ferry) will be supported by a variety of motor-powered boats operated by GCMRC researchers and contracted boat operators. Additionally, research activities on the LCR and at other locations outside of the Grand Canyon National Park boundaries are supported by helicopter services contracted with Reclamation. Ground-based support for other research activities outside of the river corridor are also coordinated with the use of GCMRC leased vehicles.

Strategic Science Questions

N/A

Links/Relationships to Other Projects

All GCMRC projects which have field data collection components are supported by the GCMRC logistics program.

Information Needs Addressed

N/A

General Methods

The GCMRC will use Government-owned boats and river logistical equipment in conjunction with a contracted vendor who supplies technical and logistical boat operators. Put-in and takeout transportation is provided with the use of General Service Administration (GSA) leased vehicles and contracted shuttle drivers.

Effective communication with principal investigators and sensitivity to and awareness of the challenges they face in implementing their studies enable the GCMRC to offer more customized (and therefore more cost-effective and productive) logistical support than other support strategies utilized previously. Retaining control over the process of supporting trips also facilitates compliance with NPS regulations and allows greater control over issues sensitive to the general public and the “recreational river community.”

Products/Reports

Research projects supported by the GCMRC must obtain permits from Federal, State, tribal, and local agencies to comply with requirements where project activities are conducted. Research activities conducted within Grand Canyon National Park and Glen Canyon National Recreation Area require NPS Research and Collecting Permits and Access Permits for all river launches, back country use, overflights, and media (filming) production. All NPS permits acquired for GCMRC supported projects are processed and submitted by the GCMRC Logistics Coordinator to the NPS Science Center Research Permitting Coordinator.

Budget

SUP 12.S1.08	
Logistics Base Costs (Other costs dispersed throughout projects; Ongoing)	
	Fiscal year 2008
GCMRC personnel costs (19% burden)	86,360
GCMRC project-related travel/training (19% burden)	—
GCMRC operations/supplies (19% burden)	—
GCMRC equipment purchase/replacement (19% burden)	20,103
AMP logistical support (19% burden)	—
Outside GCMRC and contract science labor (19% and/or other burden rate)	—
Cooperative/interagency agreements (6.09% GCMRC burden plus cooperator's burden)	—
Project Subtotal	106,463
DOI customer burden (combined 6.09%, 19% and/or other rates)	20,228
Project Total (Gross)	126,691
Percent outsourced (outside of GCMRC; includes 50% of logistics)	0%

SUP 12.S2.08: Survey Operations

Start Date

Ongoing

End Date

Ongoing

Principal Investigator(s)

Keith Kohl, Grand Canyon Monitoring and Research Center, U.S. Geological Survey

Project Goals/Tasks

- Supply GCMRC principal investigators with the necessary equipment, supplies, and survey knowledge to perform the spatial data collection required by their research
- Provide updated control point coordinates and associated errors to researchers as well as any associated processed survey data required by researchers to complete analysis
- Collect control point descriptions to publish and populate in the control point database
- Publish control point atlas and make it available for all CRE field survey activities
- Translate and rotate historical survey datasets to updated network control coordinates
- Integrate the prioritized historical survey datasets into the CRE database
- Educate principal investigators and researchers regarding the limits of various mapping techniques
- Evaluate innovative mapping techniques supporting research goals
- Publish reports evaluating (1) ground-based LIDAR techniques for monitoring cultural sites, and (2) ground-based LIDAR, oblique photogrammetry, and survey techniques as alternative methods for collecting topographic data.

Need for Project

The GCMRC is charged with providing credible, objective scientific information to the GCDAMP on the downstream resources of the Colorado River using an ecosystemwide approach. This approach requires the integration of physical, biological, and cultural spatial data into a regional GIS database to allow for accurate, long-term change detection analysis of CRE resources and for interrelationship analysis between CRE resources. All spatial data collected under the direction of the GCMRC requires referencing to the geodetic control network established by the National Geodetic Survey (NGS) and the GCMRC, which allows for accurate change detection computations including volumetric and surface area computations. Survey operations support research and monitoring activities by collecting survey data following survey protocols, and by delivering the data in formats consistent with the established data standards. Survey operations maintain survey equipment for field use including conventional total station equipment, static, kinematics and real-time kinematics GPS equipment, digital cameras, hand-held communication devices (radios), echo sounders, acoustic Doppler and bathymetry systems, and field control point atlas.

The GCMRC's survey department supports the research needs of the scientists and includes acquiring topographic data, positioning remotely sensed data, evaluating innovative mapping techniques that support research goals, validating accuracy of topographic and spatial data, and compiling and updating historical data. The survey support offered by the GCMRC allows consistent data collection methods by technically trained personnel familiar with the surveying equipment and the logistical constraints of Grand Canyon fieldwork.

Collection of topographic and spatial data enables comparison and analysis in the GIS database to monitor change. Data storage and database protocols for all survey data are developed and maintained by the survey department to ensure consistency and permit straightforward integration into the GIS database. The GIS database is referenced to the geodetic control network of highly accurate real-world coordinate values for control points from which measurements computing change are calculated. Historical, current, and future data are referenced to the control network, ensuring a common reference system between datasets collected for ecosystem monitoring studies, research, and analysis.

General Methods

The survey department is primarily responsible for assisting researchers to collect and use field measurements for scientific investigations to achieve accurate spatial data with realistic error assessments for reliable data analysis and database integration. This responsibility consists either of collecting field data, whereby the survey operations obtains, maintains, and upgrades all survey equipment required to fulfill project needs; or updating historical legacy data for analysis inclusion into the GIS database. Associated with updating legacy data, the survey department is currently collecting control point information to populate a control point database which concurrently verifies existence and condition of control points along the CRE and includes historical control points. Furthermore, the survey department advises researchers on the appropriate methods of collecting topographic or spatial data to meet the requirements of a scientific study and also evaluates new technology and its applicability to research and monitoring projects.

Survey operations in support of programs within the GCMRC are summarized in the following list:

- Position historical and newly collected spatial data
- Reference historical spatial data to modern control network
- Populate the control point database
- QA/QC of spatial data
- Evaluate innovative mapping techniques
- Maintain and upgrade equipment

Control points are established and spatial data is collected using both GPS and conventional survey methods. Surveys follow protocols developed by the GCMRC with technical support from the NGS, U.S. Army Corps of Engineers, and the Federal Geodetic Data Committee.

All programs within the GCMRC require spatial data measurements. Integration with each program's requirements and the GIS database is imperative to the process of survey data collection, postprocessing, storage, and evaluation. The survey department is available to all GCMRC principal investigators and can often collect data for multiple projects during the same mission.

The funding level requested in the FY2008 budget will allow for more historical datasets to be integrated into the GIS database for accurate change detection. Additional funds will be preserved by attaining accurate positions and elevations of spatial data before integration into GIS database. All survey equipment will be maintained and/or upgraded.

Should a high-flow experiment take place in FY2008, the funding available to support survey activities will be reduced and this reduction will delay progress to some degree on the development of the control point database and updating of legacy data. Under the "nonexperimental" scenario, funds for this area of the program are the same as originally proposed in the February 2007 version of the work plan.

Budget

SUP 12.S2.08	
Survey Operations (Ongoing)	
	Fiscal year 2008
GCMRC personnel costs (19% burden)	52,635
GCMRC project-related travel/training (19% burden)	4,000
GCMRC operations/supplies (19% burden)	5,000
GCMRC equipment purchase/replacement (19% burden)	9,430
AMP logistical support (19% burden)	15,000
Outside GCMRC and contract science labor (19% and/or other burden rate)	—
Cooperative/interagency agreements (6.09% GCMRC burden plus cooperator's burden)	—
Project Subtotal	86,065
DOI customer burden (combined 6.09%, 19% and/or other rates)	16,352
Project Total (Gross)	102,417
Percent outsourced (outside of GCMRC; includes 50% of logistics)	9%

SUP 12.S3.08: Control Network (Ongoing)

Start Date

Ongoing

End Date

Ongoing

Principal Investigator

Keith Kohl, Grand Canyon Monitoring and Research Center

Geographic Scope

Geodetic control now encompasses the entire Colorado River ecosystem corridor between Glen Canyon Dam and Lake Mead, and the greater Colorado River Basin

Project Goals/Tasks

The objective of this project is to develop a high-precision control network throughout the CRE:

- Control monuments will be established at consistent intervals throughout the CRE and at locations required for accurate positions and elevations of past, current, and future datasets.
- Expansion of the control network into the necessary areas before spatial data collection required by GCMRC research and core-monitoring activities.
- Complete stable control monuments and accurate coordinates before spatial data acquisition begins to allow for reduction in the effort required in postprocessing methods, and conservation of both human and funding resources.
- Historical datasets are accurately rectified for integration into the database.

Need for Project

The geodetic control network serves as the foundation for all spatial measurements necessary for long-term monitoring. This control network also serves as the spatial framework for the GIS. The referencing of spatial data must be consistent in order to perform accurate change detection. All spatial data collected within the CRE requires georeferencing to the primary geodetic control network established by the National Spatial Reference System, the GCMRC, and the NGS. While current remote-sensing and long-term monitoring sites have been referenced to this network, additional GCMRC monitoring activities require expanded network control efforts.

It has been shown that horizontal positions can be efficiently attained with the use of GPS techniques. While the vertical component is more problematic, heights referencing the ellipsoid can be effectively calculated throughout much of the CRE. These horizontal and vertical coordinates are required for previously collected datasets before inclusion in the CRE Oracle database. Spatial reference is also required in areas of future data collection to eliminate the need to translate and rotate surveys collected in local or superseded coordinate systems. Substantial project cost savings are achieved when the geodetic control is established within study areas before field data collection in support of monitoring and research projects.

QA/QC is required for all remotely sensed spatial datasets. The Colorado River ecosystem elevation database is designed to give positions and elevations at visible “hard points” along the river corridor. This dataset can be used to check accuracy of LIDAR (Horizons, John Chance) and digital aerial photography (ISTAR, 3001, Inc.) remote-sensing techniques, both on a canyonwide basis and for a local assessment of position and height accuracies of each day’s flight. With the high cost of remote-sensing data collection, QA/QC is critical to

analyzing the usefulness of each data subset. Additionally, this elevation database can also be used to georeference scanned photos from previous missions to study change detection.

Strategic Science Questions

All spatial data aspects of Grand Canyon investigations are addressed in this project.

Links/Relationships to Other Projects

Accurate spatial positioning of scientific data from the cultural, biological, and physical programs is necessary for facilitating change detection methods. Historical data must be adjusted to reliable coordinates before integration into the database and before resource assessments can be made. Often, past surveys that relate to current monitoring efforts have been referenced to local datums. These sites also require accurate positional and height data before the data can be entered into the GIS database for examination and change detection.

Information Needs Addressed

Accurate spatial positioning of scientific data collected within the cultural, biological, and physical programs is necessary for facilitating change detection methods.

General Methods

Control points are established using both GPS and conventional survey methods. GPS techniques utilize relative positioning where antennas and receivers are placed at both known and unknown network positions. Distances are measured between the known and unknown points by time-dependant calculations from GPS satellite data. Conventional survey techniques involve the use of a total station (a survey instrument which combines the horizontal and vertical angle measurement abilities of a transit with electronic distance measurements). Conventional traverse surveys begin at a known reference point, measure through a series of line-of-sight stations, and close at either the point of beginning or another known reference point. Both conventional and GPS measurements will be required for (1) coordinate determinations of positions and elevations throughout the CRE, and (2) realistic error estimates for each network control station.

Products/Reports

- A network of survey control points established in specific research areas and throughout the CRE, referenced to the NSRS 2007 established by the GCMRC and the NGS.
- A published paper on the geodetic control network and efforts can be expected in FY2008. The paper will include methods, results, coordinates, and realistic positional and height accuracy estimates for all network control stations and photo-identifiable fixed points. The paper will also evaluate the ability of Geoid 2003 to model NAVD88 orthometric heights from GPS-derived ellipsoidal heights.
- A separate paper will be written offering standards for all spatial data collection within the GCDAMP. This paper will be presented to the Spatial Positioning Protocol Evaluation Program targeted for fall 2008. The panel will review the document and provide further recommendations for centerwide use and acceptance. This document will provide standards and specifications for both in-house and contractual spatial data collection requirements for all collection including the 2009 remote-sensing mission.
- Coordinate results and index maps showing the locations of the network control stations, as well as QA/QC photo-identifiable fixed points which will be used to georeference past datasets and evaluate horizontal and vertical accuracies of current remotely sensed data.
- GIS layers with control station information.

Budget

SUP 12.S3.08	
Control Network (Ongoing)	
	Fiscal year 2008
GCMRC personnel costs (19% burden)	69,394
GCMRC project-related travel/training (19% burden)	3,000
GCMRC operations/supplies (19% burden)	2,085
GCMRC equipment purchase/replacement (19% burden)	770
AMP logistical support (19% burden)	22,000
Outside GCMRC and contract science labor (19% and/or other burden rate)	—
Cooperative/interagency agreements (6.09% GCMRC burden plus cooperator's burden)	18,000
Project Subtotal	115,249
DOI customer burden (combined 6.09%, 19% and/or other rates)	19,574
Project Total (Gross)	134,823
Percent outsourced (outside of GCMRC; includes 50% of logistics)	25%

PLAN 12.P1.08: Enhancing the Conceptual Ecosystem Model to Identify Critical Ecosystem Interactions and Data Gap (First priority to be funded with any additional FY2008 funds that become available)

Start Date

October 2007

End Date

September 2008

Geographic Scope

Entire Grand Canyon Monitoring and Research Center study area, from the forebay of Lake Powell to upper Lake Mead

Principal Investigator(s)

John Hamill, Chief, Grand Canyon Monitoring and Research Center

Project Goals/Tasks

In FY2007–08, the GCMRC will work with the SAs to identify and incorporate more robust integrated ecosystem science approaches into its overall program effort. The first step will be to evaluate redesign and expansion of the Colorado River CEM. A preliminary list of priority expansions of the CEM model include the following:

- Expanding the fishery elements to address coldwater and warmwater fish predation on HBC, YoY, HBC habitat use, etc.
- Modeling outcomes of nonflow management activities (i.e., operation of a temperature control device, mechanical removal of nonnatives, translocation efforts for HBC, tributary triggers for BHBFs).
- Linking Lake Powell and downstream temperature simulations to fine sediment, food web, and fisheries submodels.
- Expanding the model to provide a broader landscape perspective by incorporating Lake Powell, the Lower Colorado River, and Paria River, and addressing relationships to terrestrial habitats in the CRE.
- Enhancing the use of climatic input data and simulations.
- Recreational use and campsite size/abundance/distribution.
- Cultural site change and protection strategies (archaeological sites, TCPs).
- Financial impact simulations coupled to the flow/dam operations submodels.

The GCMRC proposes to recruit a part-time/visiting ecosystem scientist/ecologist to work with GCMRC staff and cooperators to develop and implement an integrated, interdisciplinary ecosystem science program. The primary focus of the visiting scientist will be to integrate SA recommendations and results of the CEM exercise into the GCMRC science program. The efficacy of this action will be reviewed based on the SA's above proposed evaluation/recommendations related to opportunities for incorporating an ecosystem science approach into the current science program.

Need for Project

Conceptual models summarize our current understanding of ecosystem or community function, or species life history, clarify likely responses to management actions and pressures (i.e., stressors, causes of change) (Atkinson

and others, 2004). In 1998, Walters and others (2000) conducted an Adaptive Environmental Assessment and Management Workshop to assist Grand Canyon scientists and managers in development of a conceptual model of the CRE affected by GCD operations. The model proved to be useful at helping to understand the relationship among various ecosystem components, identify knowledge gaps, and predict the response of some ecosystem components to policy change. However, it was lacking in its ability to predict the effects of policy decisions on several key areas such as long-term sediment storage, fisheries response to habitat restoration, and socioeconomic effects. Expanded design, development, and use of the CEM is needed to increase its utility in ecosystem science planning and management processes, to make it more user friendly to scientists and managers, and to provide information that is relevant to each high-priority GCDAMP goal/question.

Strategic Science Questions

The model will be directed at addressing priority AMWG questions and information needs and related SSQs in an integrated modeling effort.

Link/Relationship to Other Projects

One of the primary purposes of the CEM is to identify the linkages and relationships between various ecosystem components. Information derived from the model will assist in identifying data gaps and critical dependencies between/among science projects and allow for the effective design of an integrated, interdisciplinary science program.

Information Needs Addressed

N/A

General Methods

- The GCMRC will work with the SA and TWG to review the current CEM and identify needed updates and revision (FY2008).
- A RFP will be developed/issued to update the CEM in accordance with the findings and recommendations of the SA (FY2008).
- Two conceptual modeling workshops will be held to revise/update the model to address GCDAMP information needs and to identify data gaps and experiments or R&D projects to fill critical data gaps (FY2008). The workshops will be planned and conducted by the contractor.
- A part-time/visiting ecosystem scientist/ecologist will be recruited to work with GCMRC staff and cooperators to develop and implement an integrated, interdisciplinary ecosystem science program (FY2008–09).

Products/Reports

- SA recommendations for enhancing the CEM and improving integrated ecosystem science in the GCDAMP.
- A revised and fully documented CEM (with metadata).
- Report of workshop activities, results, and recommendations.

Budget

This budget includes \$75,000 for visiting scientist; however, GCMRC costs are covered under program management and DASA support and SA costs covered under SA budget.

Plan 12.P1.08	
Enhancing the Conceptual Ecosystem Model to Identify Critical Ecosystem Interactions and Data Gap (Science Advisor's conduct work in FY2007; Funding in Independent Reviews, ADM 12.A4.07; FY2007–08)	
	Fiscal year 2008
GCMRC personnel costs (19% burden)	—
GCMRC project-related travel/training (19% burden)	—
GCMRC operations/supplies (19% burden)	—
GCMRC equipment purchase/replacement (19% burden)	—
AMP logistical support (19% burden)	—
Outside GCMRC and contract science labor (19% and/or other burden rate)	84,034
Cooperative/interagency agreements (6.09% GCMRC burden plus cooperator's burden)	—
Project Subtotal	84,034
DOI customer burden (combined 6.09%, 19% and/or other rates)	15,966
Project Total (Gross)	100,000
Percent outsourced (outside of GCMRC; includes 50% of logistics)	100%

NOTE: \$100,000 of this funding is from fiscal year 2007 carry forward recommended by the AMWG August 30, 2007, Motion 5.

PLAN 12.P2.08: GCDAMP Effectiveness Workshop Followup

Start Date

November 2007

End Date

October 2008

Principal Investigator(s)

Grand Canyon Monitoring and Research Center will administer the project for the GCDAMP; an GCDAMP project advisory group will help design and oversee the workshop followup activities

Scope

Glen Canyon Dam Adaptive Management Program

Need for Project

In FY2008, an effectiveness workshop will be conducted that focuses on the following recommendations from the Roles Ad Hoc Group Report (2007):

- Establish a common mission/goal for the GCDAMP
- Create incentives for participants to work collaboratively to achieve common goals and desired future resources conditions
- Define desired future resources conditions
- Update or develop a charter and operating procedures for all the elements of the GCDAMP (AMWG, TWG, GCMRC, and Secretary's Designee) to reflect a more collaborative approach and so all parties clearly understand the mission and responsibilities of the group they serve on and the protocols or processes for how business will be conducted
- Utilize facilitation and mediation expertise more broadly throughout the GCDAMP
- Establish a full-time executive coordinator/manager for the program
- Determine if there is adequate time for collaboration to be successful
- Determine if a balanced range of interests are willing to participate

A goal of the FY2008 effectiveness workshop is to develop and complete an action plan to address the above issues and recommendations over the next 5 years. There is a need to provide funding to support continued facilitated discussion related to these topics and the other recommendations included in the roles report for improving the effectiveness of the AMP. Issues that may be examined in followup workshops may include the following:

- What strategies/approaches are most suitable for more effectively (1) addressing the value-based conflicts reflected by the diverse interests in the GCDAMP and (2) integrating the use of scientific information into the GCDAMP process?
- What improvements could be made in GCDAMP structure, procedures, and operations (looking individually at AMWG, TWG, GCMRC, and SAB) to improve efficiency and effectiveness of the overall program?
- Are the respective roles and responsibilities of the GCMRC, AMWG, TWG, and SAB clearly articulated and adhered to?

- Are there clear procedures in place to resolve disagreements between various GCDAMP entities?
- How could the conflict resolution procedures of the GCDAMP be improved?
- What decision support tools are available/appropriate to assist scientists and managers to improve the use of scientific knowledge in the resource management decisionmaking process?
- How can Native American involvement in and input to GCDAMP be improved?

Strategic Science Questions

N/A

Project Goals/Tasks

Results of the 2007 effectiveness workshop and other information (SAs AMP review, roles report, etc.) would be used as the basis for developing followup workshop activities. The followup activities would focus on the following:

- Providing refresher training on concepts and practical application of adaptive management and collaborative resource management. Training will be provided by experts in collaboration, partnerships, Native American involvement, and/or conflict resolution.
- Addressing priority issues and action items from the 2007 AMP effectiveness workshop.
- Addressing other issues from the roles report and AMP effectiveness survey that were not addressed in the 2007 AMP effectiveness workshop.
- Any followup workshops will be held in a centralized location and in conjunction with regularly scheduled AMWG/TWG meetings to minimize travel costs.

Followup workshop findings and recommendations developed will be reviewed by the AMWG and recommendations forwarded to the Secretary of the Interior for consideration and implementation.

Links/Relationships to Other Projects

This project has implications for the overall operation of the GCDAMP. The workshop will also provide information directly relevant to the Decision Support System Feasibility Study planned for FY2008.

Information Needs Addressed

GCDAMP RINs: 12.3.1, 12.3.2, 12.3.3, 12.5.1, 12.5.2, 12.5.4, 12.7.1, 12.8.1, 12.9.2, 12.11.1

Products/Reports

- Summary of workshop findings and recommendations
- An action plan for implementing workshop findings and recommendations

In addition to products, the workshop is designed to produce important outcomes, including (1) increased understanding of the reality of practice of adaptive management in the Glen Canyon Adaptive Management Program, (2) improved relationships among the GCDAMP stakeholders that will result in more effective incorporation of scientific information into management decisions, (3) specific practical recommendations addressing key issues related to the effectiveness and use of science information, and (4) a specific action plan for implementing the recommendations.

Budget

Plan 12.P2.08	
AMP Effectiveness Workshop (FY2007–08)	
	Fiscal year 2008
GCMRC personnel costs (19% burden)	—
GCMRC project-related travel/training (19% burden)	15,000
GCMRC operations/supplies (19% burden)	2,000
GCMRC equipment purchase/replacement (19% burden)	—
AMP logistical support (19% burden)	—
Outside GCMRC and contract science labor (19% and/or other burden rate)	8,000
Cooperative/interagency agreements (6.09% GCMRC burden plus cooperator's burden)	—
Project Subtotal	25,000
DOI customer burden (combined 6.09%, 19% and/or other rates)	4,750
Project Total (Gross)	29,750
Percent outsourced (outside of GCMRC; includes 50% of logistics)	32%

NOTE: Not funded in FY2008 per recommendation of the TWG.

PLAN 12.P3.08 Maintain a high-quality monitoring, research, and adaptive management program

AMWG Requested Project—low steady summer flows—data and research compilation, synopsis, and synthesis

Start Date

January 2008

End Date

January 2010 (conducted in phases with specific end dates)

Principal Investigator(s)

Barbara Ralston will coordinate the effort with cooperators involved in low steady summer flows data collection, Grand Canyon Monitoring and Research Center Data Acquisition Storage and Analysis Group

Geographic Scope

Entire Colorado River ecosystem corridor from forebay of Glen Canyon Dam to upper Lake Mead

Project Goals/Tasks

The overall goal of this project is to develop a synthesis of the effects of the 2000 low steady summer flows (LSSF) experiment on the CRE. The tasks involved in reaching the final goal can be broken into four phases:

- Phase I. Status of reports/data and synopsis. Identify data and products associated with the 2000 LSSF experiment; synopsise the results of the individual projects.
- Phase II. Data evaluation and identification of secondary analyses. Evaluate individual datasets and provide recommendations for further analysis resulting in integration of resource responses to operations.
- Phase III. Synthesis. Use integrated analysis results to develop a synthesis of the effects of the 2000 LSSF Experiment on the CRE.
- Phase IV. Publication. Publication of secondary analysis in a special volume of a journal or USGS circular or other publishing source.

The project outcome is intended to provide managers, and others interested in resource management, with information about how multiple resources respond to a series of flows that varied in duration from several days to several months and in magnitude from 8,000 cfs to 31,000 cfs.

Need for Project

The AMWG identified in August 2007 that there was a need to produce a summary document of the effects of the LSSF experiment (implemented in spring and summer 2000) on resources. The managers requested this summary project so that the results could be used by managers as they implement long-term experiments associated with the environmental impact statement currently under development by Reclamation.

The data collected in association with the 2000 experiment were in the areas of sediment transport and storage, mainstem and shoreline temperature, small-bodied fish sampling, long-term monitoring methods development for mainstem fishes, vegetation change, and recreational aspects of the varied flows. To date several of the data collection efforts have resulted in data reports or journal publications, while other projects remain incomplete,

lacking a final report. The lack of a unifying document regarding the flow experiment may be perceived as an impediment to learning and applying this knowledge in an adaptive management setting. It is for this reason that a summary document is being proposed that synthesizes individual resource response and considers collective resource responses within an ecosystem framework to create a subsequent synthesis.

Strategic Science Questions

Hypothetically, the LSSF experiment affected multiple resources and similarly, there are multiple SSQs that pertain to the flow experiment. The summary project will investigate whether and to what degree these SSQs were addressed by the 2000 LSSF experiment. Those SSQs most pertinent to the LSSF experiment are listed below.

SSQ 4-1. Is there a “Flow-Only” operation (i.e., a strategy for dam releases, including managing tributary inputs with BHBFs, without sediment augmentation) that will restore and maintain sandbar habitats over decadal timescales?

SSQ 5-1. How do dam release temperatures, flows (average and fluctuating component), meteorology, canyon orientation and geometry, and reach morphology interact to determine mainstem and nearshore water temperatures throughout the CRE?

SSQ 4-2. How important are backwaters and vegetated shoreline habitats to the overall growth and survival of YoY and juvenile native fish? Does the long-term benefit of increasing these habitats outweigh short-term potential costs (displacement and possibly mortality of young humpback chub) associated with high flows?

SSQ 1-7. Which tributary and mainstem habitats are most important to native fishes and how can these habitats best be made useable and maintained?

SSQ 2-1. Do dam-controlled flows affect (increase or decrease) rates of erosion and vegetation growth at archaeological sites and TCP sites, and if so, how?

SSQ 3-9. How do varying flows positively or negatively affect campsite attributes that are important to visitor experience?

Links/Relationships to Other Projects

Because much of the biological data collected in 2000, in association with the LSSF, represent a single growing season or single cohort, data from subsequent years could be used to understand the effects of a single year on recruitment signals or species compositions in subsequent surveys. These LSSF data would be linked to monitoring data from fisheries and vegetation collected since 2000, including using retrospective analysis of imagery to assess change through time. The sediment response throughout the duration of the project can be incorporated into the current shoreline study project to understand the relationship of reworking eddy sand supply and available shoreline habitats through remote-sensing analysis. In the same vein, water temperature data collected in 2000 is applicable to current water temperature modeling efforts for shoreline habitats. Lastly, recreational aspects associated with downstream travel and visitation could be interpreted under the current Colorado River Management Plan to determine how similar flows, if they occur in the future, might affect recreational experiences.

Information Needs Addressed

Information needs that pertain to work done during the LSSF are focused on experimental information needs for each resource. Specific information needs that focus on adaptive management and that are pertinent to the proposed project are the following:

IN 12.1. Develop information that can be used by the TWG, in collaboration with the GCMRC, to establish current and target levels for all resources within the GCDAMP as called for in the GCDAMP strategic plan.

RIN 12.3.1. As necessary, investigate the most effective methods to integrate and synthesize resource data.

General Methods

As a part of the 1994 biological opinion associated with the operations of GCD, the USFWS provided reasonable and prudent alternatives (RPAs). One element of the RPAs directed Reclamation to initiate a program of experimental dam releases consisting of high steady spring flows and LSSFs. The intention of these experimental releases was to reduce the risk of further jeopardizing endangered native fishes.

A plan of flows was developed by SWCA Environmental Consultants, Inc. (SWCA, 2000). The plan divides the flows into three time periods: March–May (high flows of 21,000 cfs with a 31,000-cfs spike), June–September (steady flows of 8,000 cfs, ending with a 31,000-cfs spike), and October–February (8,000-cfs flows). The flows that were implemented in spring 2000 were slightly different in that the high flows in the spring were a slightly lower discharge of 17,500 cfs rather than 21,000 cfs, and the duration of the flows was shorter by approximately a month (fig 4).




Figure 4. Hydrograph from March–December 2000 including discharge pattern associated with the LSSF experiment.

Data collected around these flows focused on physical resources (sediment, water temperature), biological resources (aquatic productivity, fisheries, vegetation), and cultural resources (recreation, economics). SWCA (2000) provided some hypotheses regarding the benefits and risks to abiotic and biotic resources relative to each flow period (table 5). It is proposed that these hypotheses form the basis for data consolidation, synopsis, secondary analysis, and subsequent synthesis.

Table 5. Hypothesized effects of flows on physical and biological resources.

Benefits/risks to resources	Period I: March–May	Period II: June–September	Period III: October–February
Benefit to physical resources/ habitat	–Scouring backwaters –May spike flow to mobilize and store sands and sediment	–Storing of sand and sediment in river channel –Expansion of campable beach area –September spike flow –Resuspension, storing of sand from summer tributary inputs	–No significant risks –Increased survival of young native fishes –Maintenance of stable winter conditions to minimize energy expenditure –Maintenance of overwinter autotrophic production in mainstem, shorelines, backwaters
Risks to physical resources/ habitat	–Export of sediment, reduction of campsite areas	–September spike flow, export of sand and sediment instead of storing it	
Benefits to biotic resources	–Ponded tributary inflows as thermal refuges for drifting larvae and young fish –Ponded tributary inflows ease access for spawning native fishes –Destabilizing of habitats to disadvantage nonnatives –Redistribution of nutrients –Resetting of community production –Spike flows to flush nonnative fish from nearshore habitats	–Increased growth and survival of young native fishes –Increased autotrophic algal and macroinvertebrate production –Possible mainstem hatching success –Spike flows to flush nonnatives fish from nearshore habitats	
Risk to biotic resources	–Attraction of nonnative fish predators/competitors to arm-ponded tributaries	–Mainstem reproduction by nonnative fishes –Increased growth and survival of nonnative fishes –Increased infestation of parasites and diseases –Decreased drift of food for fish –Minimized thermal plume at 30-mile may reduce survival of young HBC –Increased water clarity leading to increased predation of native fish by sight predators	–Possible overwinter survival and expansion of nonnative fishes –Possible greater spawning success of downstream populations of trout –Increased predation by sight feeders –Decreased drift of food for fish

Phase I. Status of reports/data and synopsis (6 months)

- Identification of studies in LSSF plan—There were 25 studies identified in the LSSF plan as well as several overflights conducted throughout the period of March through September (table 6). Determine/describe the scope of each study and how many of the proposed studies were executed.
- Determination of location of data and other deliverables—call PIs to determine status of project, location of data, identification of any work that was not done and/or cannot be done and consolidating data.
- Synthesize project results and describe the status of data (metadata report).

Table 6. List of projects identified in LSSF Plan.

Resource Area	Project Title	Principal Investigators
Physical resources		
Sediment storage and transport	Effects of low steady summer flows (LSSF) experiment on the stratification, composition, and hydrodynamics of Lake Powell, and the downstream effects of that limnology	Vernieu (USGS)
	Effect of discharge and flows on temperatures in aquatic habitats	Vernieu (USGS)
	Effect of discharge on shoreline channel and tributary habitat	Protiva (SWI)
	Additional monitoring of effects of flow on suspended sediment and turbidity levels in the main channel of the Colorado River	Rubin, Anima, Topping (USGS, WRD)
	Effects of short-duration high releases and long-duration low steady flows on deposition and erosion of fine sediment at selected eddy complexes utilizing the 34 Northern Arizona University (NAU) long-term eddy complex and sandbar study sites, plus four newly established channel margin monitoring sites	NAU Sandbar Group
	Remote-sensing component to investigate change detection of suspended sediment concentration, turbidity, and river-bottom cover type	Chavez (USGS)
	Additional sediment and streamflow modeling along the Colorado River ecosystem between Glen Canyon Dam (GCD) and Phantom Ranch in support of LSSF testing	Wiele (USGS)
	A collaborative research project before, during, and after the 33,000-cfs fall test flow with integrated and alternative methods to monitor sand storage	NAU, Utah State University, USGS
Biological resources		
Riparian vegetation	Effects of steady vs. fluctuating flows on creation of “vegetated shoreline” for juvenile fish and exotic recruitment into newly available habitat	Porter/ Kearsley
Aquatic productivity	Effect of a 31,000-cfs spike flow and low steady flows on drift and benthic mass and composition in the Lees Ferry reach	Arizona Game and Fish Dept. (AGFD)
	Algal colonization and recolonization response rates during experimental LSSFs	Yard (NAU)
Fishes	Effect of low steady flows on drift and benthic mass and composition in the Lees Ferry reach and downstream	Shannon (NAU)
	Effect of steady flows on growth, relative abundance, and distribution of young-of-year fish along shoreline below the Little Colorado River	SWCA Environmental Consultants
	Monitoring of Colorado River fish community	Grand Canyon Monitoring and Research Center (GCMRC), AGFD, U.S. Fish and Wildlife Service (USFWS), SWCA
	Coupling hydrodynamic and individual-based fish movement models to evaluate the effects of flow and temperature releases from GCD on the accessibility of suitable habitat for humpback chub juveniles in the Colorado River	Korman/Wiele (Ecometric/ USGS)
	Effect of LSSFs on Lees Ferry trout	AGFD
Sociocultural resources		
	Economic impacts to power customers	Palmer (WAPA)
	Whitewater boating safety studies below Lees Ferry	Jalbert (National Park Service)
	Economic impacts to whitewater and angling concessionaires	?

Economic impacts to private whitewater boaters and anglers Effects on recreational river trip characteristics	? Jalbert, Roberts (NPS, NAU)
Changes in campable beach areas	?
Overflight data	
Topographic base mapping of the Colorado River corridor from GCD to Lake Mead—horizons flight in April 2000 Pre-experiment color-infrared (CIR) orthorectified digital photomosaic of the Colorado River corridor from GCD to Lake Mead at 8,000-cfs steady flows and 30-cm resolution. Steady flow (late June) CIR aerial photography of the Colorado River corridor from GCD to Lake Mead at 8,000-cfs steady flows and 10-cm-pixel resolution (RM -15 to 277) Steady flow (late June) thermal infrared imagery of the Colorado River corridor from GCD to approximately Phantom Ranch at 8,000 cfs steady flows and one meter pixel resolution. (RM -15 to 277) Pre-fall-spike black and white stereo digital and hard-copy orthophotography with 60-percent overlap of the first 100 miles of the Colorado River corridor from GCD to Phantom Ranch (river mile [RM] -15 to 85) at 8,000 cfs and 10-cm-pixel resolution Peak fall spike 1:4800 CIR stereo aerial photography of the first 100 miles of the Colorado River corridor from GCD to Phantom Ranch (RM -15 to 85) at 8,000 cfs and 10-cm-pixel resolution Post-fall-spike 1:4800 black and white stereo digital and hard-copy orthophotography with 60-percent overlap of the first 100 miles of the Colorado River corridor from GCD to Phantom Ranch (RM -15 to 85) at 8,000 cfs and 10-cm-pixel resolution High-resolution (25-cm) topography of 20 miles of sandbars yet to be determined within the first 100 miles of the Colorado Canyon corridor up to 300,000 cfs before and after fall spike. This product to be derived from pre- and post-spike orthophotography collected above or high-resolution LIDAR yet to be determined	
High-resolution (25-cm) topography of 20 campable beach sites yet to be determined within the first 100 miles of the Colorado Canyon corridor	

Phase II. Data evaluation and identification of secondary analyses (3 months)

- Evaluation of data compatibility in collaboration with DASA and resource specialists within the GCMRC, SAs, PIs involved in LSSF and other meta-analysis experts. Utilize a workshop setting to initiate discussion and determine the utility of secondary analysis.
- Identification of potential secondary analyses of data including incorporation more recent monitoring and research data to provide a longer term analysis of effects. Utilize proposed senior scientist to help identify potential and critical secondary analyses.
- Identification of PIs available for secondary analysis and collaboration determination of funding needs and timelines.
- Development of statements of work for subsequent secondary analysis and obligate funds.

Phase III. Secondary analysis and synthesis (15 months)

- Execution of secondary analyses incorporating more recent monitoring data and identification of publishing venue for research (e.g., special issue in Ecological Applications, American Geophysical Union?). Collaborators identified in Phase II.
- Incorporation of results into conceptual modeling exercise (e.g., Ecosim/ecopath) for ecosystem response analysis and system-wide synthesis (possibly concluding chapter in volume). Utilize talents of proposed Ecosystem scientist.

- Writing of results and discussion of secondary analyses and conceptual modeling effort to create synthesis document.
- In coordination with editing staff at the GCMRC/SBSC, setup review and complete draft manuscripts.

Phase IV. Publication (3 months)

- In coordination with editing staff at the GCMRC/SBSC complete publication of manuscripts in target journal or circular.

Products/Reports

- Phase I. USGS open-file report providing background information about LSSF, synopses of individual project, metadata, background information about LSSF. Draft submitted by June 2008; Finalized by August 2008.
- Phase II. Evaluation of data, identification of potential secondary analysis through workshop bringing together LSSF PIs, SAs and others familiar with meta- analysis. Workshop anticipated in September 2008 to be led by ecosystem scientist.. Work plans for secondary analysis. Statements of work established for secondary analysis. Draft report submitted by November 2008; Finalized by December 2008.
- Phase III. Initiation of secondary analysis and synthesis. Collation of finalized manuscripts reviewed and ready for submission to target journal or circular for publication. Submitted by March 2010.
- Phase IV. Completed publication of manuscripts. Completed by July 2010.

Budget

Plan 12.P3.08	
Lees Ferry Trout Study FY2008: Low Steady Summer Flows—Data and Research Compilation, Synopsis and Synthesis	
	Fiscal year 2008
GCMRC personnel costs (19% burden)	34,700
GCMRC project-related travel/training (19% burden; Workshop Related)	30,000
GCMRC operations/supplies (19% burden)	1,503
GCMRC equipment purchase/replacement (19% burden)	—
AMP logistical support (19% burden)	—
Outside GCMRC and contract science labor (19% and/or other burden rate)	—
Cooperative/interagency agreements (6.09% GCMRC burden plus cooperator's burden)	20,000
Project Subtotal	86,203
DOI customer burden (combined 6.09%, 19% and/or other rates)	13,797
Project Total (Gross)	100,000
Percent outsourced (outside of GCMRC; includes 50% of logistics)	23%

NOTE: \$100,000 of this funding is from fiscal year 2007 carry forward recommended by the AMWG August 30, 2007, Motion 5.

Budget Detail for Phase I and II

Determining the status of reports/data and synopsis outlined for phase I will require 6 months and involve the cost of two pay periods for the coordinator (Ralston) and personnel costs associated with NAU cooperators (\$20,000). Phase II, data evaluation and identification of secondary analyses (3 months), will involve a data evaluation workshop and identification of secondary analysis, which will require three pay periods for GCMRC DASA personnel and five pay periods of ecosystem ecologist salary. Costs associated with phase III and IV pending results/recommendations of Phase II.

Administrative Operations

ADM 12.A1.08: Administrative Operations

Start Date

Ongoing

End Date

Ongoing

Principal Investigator

John Hamill, Chief, Grand Canyon Monitoring and Research Center

Scope

Grand Canyon Monitoring and Research Center

Need for Project

It is necessary to have smooth running, transparent administrative operations that ensure that the GCMRC scientists can focus on their research rather than on the administrative details involved with the payment of rent and utilities, timekeeping concerns, filing, and various other administrative topics. Administrative operations activities provide the oversight and management of facilities, burden, and overhead; personnel issues; expenditure tracking; processing and financial management of cooperative and interagency agreements; processing of contracts; timekeeping; bank card tracking and reconciliation; travel plans and voucher processing; and liaison activities between the USGS administrative groups (Flagstaff Science Center Administration, Western Region Budget and Fiscal Services and Contracting Offices, Headquarters in Reston, and the Biological Headquarters). In addition, this project is innately involved with the USGS nationwide budget tracking and reporting system known as BASIS+, which is used by the USGS Headquarters and Regional offices to make their annual reports to Congress, as well as to respond to Congressional inquiries with turnaround times as short as 12 hours. (As part of the Glen Canyon Dam Adaptive Management Program, GCMRC administrators have been called upon to provide information of this type from the system on many occasions.)

In FY2008, standard overhead charges including facilities, space, general office supplies, costs for the USGS local network, Flagstaff Science Center support, and USGS regional services including contracting and personnel, as well as the salaries and general travel for the GCMRC secretary and budget analyst, have been moved to the SBSC's overhead account. By doing this, the GCMRC complies with regulations dictated by USGS headquarters and has identified a savings in administrative costs that has been applied to science projects. Only charges directly tied and traceable to the GCMRC continue to be directly charged to the Administrative Operations account. These charges include GSA vehicle lease and maintenance; DOI vehicle gas, maintenance, and replacement costs; funding for editorial support; safety and/or other non-project-specific mandated training; GCMRC non-project-specific personnel support; telecommunications and shipping charges; and others.

Strategic Science Questions

N/A

Project Goals/Tasks

The goals of the project are to provide budgetary oversight and support to the Chief, Program Managers, and all employees of the GCMRC so that they may conduct their responsibilities in the most ethical, professional, and efficient manner possible; to enable the employees to be unburdened, to the largest extent possible, by mundane administrative matters; and to support the USGS and GCMRC missions of conducting unbiased scientific research.

General Methods

General methods will include standard accounting procedures and regulatory and legal standards as required by the USGS and other Federal agencies with legal oversight. Quarterly updates to Program Managers will be provided as well as budgetary and other information provided upon request. The GCMRC will follow USGS guidelines as assigned for personnel, travel, and other processes. Administrative personnel will focus on how to accomplish requests within Federal laws and regulations. The Administrative Officer for SBSC and the Budget Analyst for the GCMRC will report annually to the AMWG/TWG on year-end projections and on the actual expenditures for the previous fiscal year.

Links/Relationships to Other Projects

This project is innately linked to all other projects. All project budgets are impacted by burden charges that are tracked and managed through Administrative Operations, all employees are required to track their time through a USGS personnel system, and many Program Managers use cooperative or interagency agreements that are processed and tracked financially via Administrative Operations. Every project is given an account number and must be entered into and tracked, via its budget and its narrative, through the BASIS+ system. Administrative Operations activities are tied to each project at the project's earliest development.

Information Needs Addressed

N/A

Products/Reports

The Administrative Officer for SBSC and the Budget Analyst for the GCMRC will produce a projection report (usually at the August AMWG meeting) for year end. In addition, they will present a report in actual expenditures for the previous fiscal year that will normally be presented at the March AMWG meeting.

Budget

ADM 12.A1.08	
Administrative Operations (Ongoing)	
	Fiscal year 2008
GCMRC personnel costs (19% burden)	40,000
GCMRC project-related travel/training (19% burden)	5,000
GCMRC operations/supplies (19% burden)	102,675
GCMRC equipment purchase/replacement (19% burden)	5,000
AMP logistical support (19% burden)	—
Outside GCMRC and contract science labor (19% and/or other burden rate)	—
Cooperative/interagency agreements (6.09% GCMRC burden plus cooperator's burden)	44,000
Project Subtotal	196,675
DOI customer burden (combined 6.09%, 19% and/or other rates)	31,688
Project Total (Gross)	228,363
Percent outsourced (outside of GCMRC; includes 50% of logistics)	21%

ADM 12.A2.08: Program Planning and Management

Start Date

Ongoing

End Date

Ongoing

Principal Investigator

John Hamill, Chief, Grand Canyon Monitoring and Research Center

Scope

Grand Canyon Monitoring and Research Center

Need for Project

Successful scientific research and reporting can be enhanced by strong and effective leadership that provides close working relationships between managers and employees. Good managers can apply knowledge as management actions that can enhance scientific research and imagination. In addition to their program management responsibilities, the GCMRC Program Managers are also subject area experts in their respective fields. It is important that GCMRC Program Managers and scientific staff maintain this expertise so they can provide high-quality technical assistance in the form of expert analysis, opinion, and advice to the Chief, TWG, and AMWG, as requested. The Sociocultural Program Manager also functions as the Native American Coordinator. The Program Managers supervise additional technical and support staff, and act as project leads with their cooperators.

Beginning in FY2006, in an effort to simplify distribution of program planning and management salaries and travel, the Program Manager salaries were assigned to this category exclusively. In FY2008, the full salary of the DASA Program Manager's salary is included in this line item. The SBSC's IT Director salary has been removed and has been included in the general overhead account for the Cost Center, as have many of the charges described in the Administrative Operations section. Travel expenses in support of the program, but separate from TWG and AMWG meeting travel, are also included. Salaries and travel costs for Program Managers, the Chief, and Deputy Chief are included in program planning and management budget.

Strategic Science Questions

N/A

Project Goals/Tasks

The GCMRC's goal is to deliver a comprehensive ecosystem science program over the next 5 years that is effective in responding to management needs articulated through the GCDAMP and by DOI. Productive, well-qualified personnel are critical to achieving this goal.

General Methods

In order to provide strong leadership of a quality science program that is responsive to the needs of the GCDAMP, the GCMRC will be administered by a core program management staff that includes the following key positions:

Center Chief

Establishes Center science policies and strategic direction and provides accountability for the GCMRC budget. Interfaces with USGS management, Secretary's GCDAMP Designee, and GCDAMP managers to ensure that quality science is provided in a timely manner on priority issues identified by the GCDAMP leadership.

Deputy Chief

The Deputy Chief will be responsible for day-to-day management and supervision of the Physical Science and Modeling Program and ensure that integrated ecosystem science methods and procedures are utilized in science design and analysis.

Program Managers

Responsible for the timely execution of the science program within their program area; interaction with other program areas to ensure integrated ecosystem approaches, quality control of products and contractors/cooperators; contract/agreement management; management of budget within their program area, and providing reports to GCDAMP work groups as needed. The GCMRC activities now encompass five major program areas:

1. The Physical Science and Modeling Program conducts research and monitoring activities on physical elements of the CRE including studies of sediment storage and transport in the regulated river, and integrated downstream water-quality monitoring and research. The program has been responsible for conducting several experimental high-flow releases from GCD to conserve sediment resources for building beaches and improving habitat for native aquatic species in the Colorado River. More recent tasks have included development of a downstream temperature model for the ecosystem.
2. The Data Acquisition, Storage, and Analysis (DASA) Program provides GIS, data quality control, data management, and library services support to all program areas. In addition, DASA oversees the GCMRC peer-review process.
3. The Biological Program provides scientific information that supports the conservation of native species in the Grand Canyon and the Lees Ferry trout fishery. Elements of the program include assessing the effects of GCD on fishery resources; characterizing the aquatic food base; evaluating terrestrial contributions to the aquatic food base; improving fish community monitoring, developing, and testing of techniques to control nonnative fishes; evaluating terrestrial vegetation changes as a result of dam operations; and water-quality monitoring and modeling in Lake Powell and the Colorado River below GCD.
4. The Cultural and Socioeconomic Program focuses on culturally significant sites and artifacts, and recreation activities based in Grand Canyon. The current focus is on development of comprehensive monitoring programs to assess the condition of the culturally significant sites affected by the operation of GCD.
5. The Logistics Program supports up to 40 river trips per year and coordinates research permit management for the Grand Canyon Monitoring and Research Center. The Logistics Program also provides survey support to various program and activities.

Links/Relationships to Other Projects

This project is linked by nature to all other projects, since each project must be managed by a Program Manager or the Chief.

Information Needs Addressed

N/A

Products/Reports

All products and reports produced by the GCMRC are a result of this project.

Budget

ADM 12.A2.08	
Program Planning & Management (Ongoing)	
	Fiscal year 2008
GCMRC personnel costs (19% burden)	840,904
GCMRC project-related travel/training (19% burden)	38,880
GCMRC operations/supplies (19% burden)	10,500
GCMRC equipment purchase/replacement (19% burden)	—
AMP logistical support (19% burden)	—
Outside GCMRC and contract science labor (19% and/or other burden rate)	—
Cooperative/interagency agreements (6.09% GCMRC burden plus cooperator's burden)	—
Project Subtotal	890,284
DOI customer burden (combined 6.09%, 19% and/or other rates)	169,154
Project Total (Gross)	1,059,438
Percent outsourced (outside of GCMRC; includes 50% of logistics)	0%

ADM 12.A3.08: AMWG/TWG Meeting Travel Funds

Start Date

Ongoing

End Date

Ongoing

Principal Investigator

John Hamill, Chief, Grand Canyon Monitoring and Research Center

Scope

Grand Canyon Monitoring and Research Center

Need for Project

This project is an account to hold funds for travel expenses only for employees who participate in AMWG and TWG meetings. Project-related travel expenses are accounted for by projects, and administrative travel (e.g., general safety and security training) is planned under the Administrative Operations budget.

Strategic Science Questions

N/A

Project Goals/Tasks

To provide travel funds for employees who participate in AMWG and TWG meetings.

General Methods

Methods used are standard USGS travel authorizations and vouchers.

Links/Relationships to Other Projects

N/A

Information Needs Addressed

N/A

Products/Reports

N/A

Budget

ADM 12.A3.08	
AMWG/TWG Meeting Travel Funds (Ongoing)	
	Fiscal year 2008
GCMRC personnel costs (19% burden)	—
GCMRC project-related travel/training (19% burden)	15,191
GCMRC operations/supplies (19% burden)	—
GCMRC equipment purchase/replacement (19% burden)	—
AMP logistical support (19% burden)	—
Outside GCMRC and contract science labor (19% and/or other burden rate)	—
Cooperative/interagency agreements (6.09% GCMRC burden plus cooperator's burden)	—
Project Subtotal	15,191
DOI customer burden (combined 6.09%, 19% and/or other rates)	2,886
Project Total (Gross)	18,077
Percent outsourced (outside of GCMRC; includes 50% of logistics)	0%

ADM 12.A4.08: Independent Reviews

ADM 12.A6.08: Science Symposium

Start Date

Ongoing

End Date

Ongoing

Principal Investigator

John Hamill, Chief, Grand Canyon Monitoring and Research Center

Scope

Grand Canyon Monitoring and Research Center

Need for Project

Independent external review is at the heart of the GCMRC's approach to program management and implementation. Together with the competitive process, independent external peer review ensures the quality and objectivity of the GCMRC's programs. Independent review panels are used to evaluate the GCMRC's plans and activities. All proposals, reports, programs, etc., are subject to independent peer review according to the GCMRC's peer-review protocols. Managing the GCMRC's peer-review process requires 3–6 person-months, but requires no additional salary and is the responsibility of the Librarian/Review Coordinator. The Review Coordinator reports to the Chief directly, but works under the guidance of the DASA Coordinator for all non-review-related activities.

Strategic Science Questions

N/A

Project Goals/Tasks

To increase the efficiency and quality of the science being developed by the GCMRC and used by the AMWG and the Secretary of the Interior, the GCMRC will establish a peer-review process to ensure that all unsolicited, solicited, or in-house proposals and all draft reports received by the GCMRC undergo independent, external peer review.

To ensure program integrity, the SAB will provide independent scientific oversight and technical advice to ensure that GCMRC science programs are efficient, unbiased, objective, and scientifically sound. The SAs individually will be expected upon request, among other things, to review and comment on the following:

Results of ongoing and completed monitoring and research program activities, as well as any synthesis and assessment activities initiated by the GCMRC.

The appropriateness of the GCMRC's RFPs, especially their responsiveness to management objectives

Protocols used in GCMRC-sponsored scientific activities, including a 5-year review of GCMRC monitoring and research protocols

GCMRC's long-term monitoring plan

GCMRC's annual monitoring and research plans

GCMRC's annual budget proposals, to ensure that the science program is efficiently and effectively responding to AMWG goals (i.e., management objectives).

The SAB may also provide other program specific scientific and technical advice it is asked to address by the AMWG, the GCMRC, or the Secretary of the Interior.

General Methods

Peer Review

All of GCMRC's scientific activities undergo an independent, external peer review including all unsolicited, solicited, or in-house proposals. Similarly, all draft reports received by the GCMRC undergo independent, external peer review. The peer-review protocols developed by the GCMRC meet or exceed the standards articulated by the Secretary of the Interior for DOI.

Peer review for proposals received by the GCMRC in response to an RFP is conducted through a panel process, while peer reviews for unsolicited and in-house proposals, as well as project reports, are conducted through correspondence. In all cases, the reviewers are offered anonymity, and the individual and panel reviews, where applicable, are provided to the PIs along with comments from the GCMRC. In addition, the GCMRC conducts PEPs to review and assess GCMRC's projects and methodologies. To date, PEPs have been held for remote-sensing, physical, survey control, terrestrial and aquatic, cultural resource, and the water-quality program.

The GCMRC review process is handled by a Review Coordinator to ensure that the peer-review process is not under the immediate supervision of individual GCMRC Program Managers to guard against any conflicts of interest—real or perceived. Strict conflict-of-interest guidelines are adhered to. GCMRC annually recruits new peer reviewers and maintains a database of almost 500 potential reviewers, organized by area of expertise. GCMRC peer reviewers come from academia; Federal, State, and tribal governmental and nongovernmental organizations; and the private sector. Reviewers are selected on the basis of their record of scientific accomplishment and expertise.

Budget

ADM 12.A4.08	
Independent Reviews (Ongoing)	
	Fiscal year 2008
GCMRC personnel costs (19% burden)	20,883
GCMRC project-related travel/training (19% burden)	12,000
GCMRC operations/supplies (19% burden)	4,800
GCMRC equipment purchase/replacement (19% burden)	—
AMP logistical support (19% burden)	12,000
Outside GCMRC and contract science labor (19% and/or other burden rate)	26,200
Cooperative/interagency agreements (6.09% GCMRC burden plus cooperator's burden)	—
Project Subtotal	75,883
DOI customer burden (combined 6.09%, 19% and/or other rates)	14,418
Project Total (Gross)	90,301
Percent outsourced (outside of GCMRC; includes 50% of logistics)	42%

Science Advisors

The GCMRC works with a group of Science Advisors as one of its independent review panels. The SAB is an advisory group and not a decisionmaking body. It is an interdisciplinary group composed of scientists who are

qualified on the basis of their record of publication in the peer-reviewed literature, or other demonstrable scientific achievements. An Executive Secretary leads the SAs and serves as the liaison officer to the AMWG and the GCMRC.

The SAs, together and individually, will be expected in FY2008 to review and comment to the AMWG and the GCMRC on (1) GCMRC's annual work plan and budget proposal, (2) GCMRC's long-term monitoring and research plan (MRP), (3) the results of GCMRC's completed monitoring and research activities, (4) the results of any synthesis and assessment activities initiated by the GCMRC, and (5) any other activities (i.e., developing a monitoring plan, enhancing opportunities for integrated science, and other program-specific scientific advice) it is asked to address by the GCMRC Chief or the AMWG. Table 7 summarizes SA activities planned for FY2008.

Table 7. Summary of Science Advisors activities for fiscal year (FY) 2008.

Requesting group	Type of activity	Service request	Completion date and months required
TWG	Advisory service	Risk assessment of proposed experimental options and FY2007–11 GCMRC/GCDAMP science program.	7/07; 10
GCMRC	Advisory service	Assist GCMRC in designing and implementing ecosystem science approaches in research and monitoring programs, experimental options, modeling, sampling designs, etc.	10/07; 12

Budget

ADM 12.A4.08	
Executive Director of Science Advisors Review and Coordination; includes Science Advisors' Expenses (Independent Reviews; ongoing)	
	Fiscal year 2008
GCMRC personnel costs (19% burden)	—
GCMRC project-related travel/training (19% burden)	5,000
GCMRC operations/supplies (19% burden)	—
GCMRC equipment purchase/replacement (19% burden)	—
AMP logistical support (19% burden)	—
Outside GCMRC and contract science labor (19% and/or other burden rate)	175,000
Cooperative/interagency agreements (6.09% GCMRC burden plus cooperator's burden)	—
Project Subtotal	180,000
DOI customer burden (combined 6.09%, 19% and/or other rates)	34,200
Project Total (Gross)	214,200
Percent outsourced (outside of GCMRC; includes 50% of logistics)	97%

Science Symposium

In the winter of 2008, the GCMRC will host a science symposium to summarize recent research findings and recommendations since publication of the SCORE report in FY2005. Reports will be provided on all major activities and findings that are relevant to high-priority AMWG questions and related SSQs. The GCMRC will explore the interests in holding its science symposium in conjunction with similar meetings that are held for the Recovery Implementation Program for Endangered Fish Species in the Upper Colorado River Basin, the San Juan River Recovery Implementation Program, and the Lower Colorado River Multi-Species Habitat Conservation Plan to help promote increased technical dialogue and information exchange among these programs. Some of the costs for the Science Symposium will be recovered through a modest registration fee for participants.

Budget

ADM 12.A6.08	
2008 Science Symposium (Ongoing, every other year)	
	Fiscal year 2008
GCMRC personnel costs (19% burden)	—
GCMRC project-related travel/training (19% burden)	11,000
GCMRC operations/supplies (19% burden)	4,000
GCMRC equipment purchase/replacement (19% burden)	—
AMP logistical support (19% burden)	—
Outside GCMRC and contract science labor (19% and/or other burden rate)	10,000
Cooperative/interagency agreements (6.09% GCMRC burden plus cooperator's burden)	—
Project Subtotal	25,000
DOI customer burden (combined 6.09%, 19% and/or other rates)	4,750
Project Total (Gross)	29,750
Percent outsourced (outside of GCMRC; includes 50% of logistics)	40%

Links/Relationships to Other Projects

N/A

Information Needs Addressed

N/A

Products/Reports

- Final products will include final work plans that have undergone peer review (comments maintained on file at GCMRC) and peer-review comments on draft final reports produced related to projects included in the work plan (comments maintained on file at GCMRC).
- Summaries of papers presented at the 2008 Science Symposium will be published by the GCMRC pursuant to USGS Fundamental Science Practices.

ADM 12.A5.08: GCMRC Component of SBSC Systems Administration Support

Start Date

FY2005

End Date

Ongoing

Principal Investigator(s)

John Hamill, Chief, Grand Canyon Monitoring and Research Center

Scope

Grand Canyon Monitoring and Research Center

Need for Project

The IT Department of the SBSC supports a variety of technology needs of the GCMRC's various program areas: computer security, systems administration and procurement of new servers and computers, as well as Web site development and Web page maintenance. These support, development, and maintenance services are cost shared between the GCMRC, the SBSC, and the IT Department, and coordinated by the Center's Deputy Director so as to meet the IT needs of all four research stations.

Strategic Science Questions

N/A

Project Goals/Tasks

It is the IT Department's goal to ensure that GCMRC and all stations within SBSC are able to conduct scientific and administrative functions smoothly and with the least amount of disruption in service as possible. It is the IT Department's task to make IT functions as transparent as possible, to ensure each program has adequate current and future storage, and to provide excellent customer service at all times. IT maintains the security of GCMRC and SBSC networks up to current Federal standards and ensures that all those who access the systems meet Federal security standards in order to protect personal information and scientific research that has not yet been released to the public. At the same time, the IT Department ensures that the public has full and easy access to publicly released data via GCMRC Web sites and works closely with the DASA program to make this possible.

General Methods

The IT Department follows all Federal, DOI, and USGS regulations regarding purchase of, access to, distribution and release of electronic information. Methods also include the following:

- Network environment—Computer interconnectivity is provided using transmission control protocol/internet protocol (TCP/IP) network communication protocol running on a 1000baseT and 100baseT network media. Network traffic is arbitrated by 4 3COM switches and hubs operating at 100 Mbps and 1 Gbps.
- Internet connectivity—The GCMRC computer network is linked to the Internet through the Flagstaff Field Center GEOnet-3 router that provides a DS-3 (45 Mbps) virtual circuit to Menlo Park, where it joins the USGS GEOnet network. Also located in Menlo Park is a network portal to the Internet operated by the USGS and NASA through a peering partnership. GEOnet provides a secure Survey-wide networking environment that interconnects headquarter region, district, and field offices located throughout the United States.

- Intranet Web site—GCMRC’s intranet offers a secure centralized medium for information exchange among GCMRC employees. Among things to be internally shared via the intranet are standard operating procedures, personnel availability and contact info, vehicle and equipment loans, and an IT support system. The GCMRC intranet is served from a Windows 2000 Server utilizing Active Server Pages (ASP).
- Computer security—Network security is provided by firewalls, routers, a system update server (SUS), a systems management server (SMS), and antivirus software. Firewalls and routers are configured and maintained to restrict outside access to authorized systems. Operating systems are updated to minimize vulnerabilities using SUS that automates a central delivery system for patch management. Antivirus updates are downloaded from the Web as released and pushed to all systems the same night.
- Desktop and servers—GCMRC’s computing environment is based upon the PC platform, Microsoft Windows operating system, and Microsoft Office, office automation software. Systems maintenance is performed using a combination of warranty service, service contracts, and in-house service as needed to facilitate quick turnaround, minimize downtime, and reduce costs.
- System backup and disaster recovery—System backup and disaster recovery is accomplished using dual linear tape open (LTO) tape drives in a 30-slot carriage with a capacity of 3 Tbytes. Tapes are stored locally in a fire vault and archival tapes are stored off-site. Server disks are configured to run either a raid-5 array or mirrored for redundancy.
- Troubleshooting and maintenance—Helpdesk support is provided as requested/required. Requests are received via the Web, email, and telephone. Support is tracked in a searchable database with solutions to facilitate prioritization and resolution.
- Assistance with GCMRC’s data storage—Over 7 Tbytes of online disk storage is provided by multiple servers with small computer system interface (SCSI) disk arrays. Server disk arrays are hot swappable to minimize downtime. GCMRC also utilizes networked attached storage (NAS) devices. Integrated Drive Electronics (IDE) drives connected to a SCSI backplane. NAS units are used to provide bulk storage capacity at less expense.

Links/Relationships to Other Projects

All projects are integrated with IT support. Refer to the DASA section for more information on integration with these projects.

Information Needs Addressed

N/A

Products/Reports

The primary products and services of the SBSC Information Technology Department with respect to ongoing support of the GCMRC’s needs are as follows:

- Comprehensive and fully functional Web site development and maintenance, with access to all nonsensitive digital data and information relating to the effects of dam operations on the CRE. Nondigital data and information will be cataloged electronically with instructions on how to obtain it.
- Coordination with GCMRC’s DASA to ensure and support a comprehensive and fully functional library containing all hard copy and digital media containing data and information relating to the effects of dam operations on the CRE cataloged and accessible. Sensitive and nonreleasable data and information will be archived and secured separately from releasable data and information.
- Fully functional and integrated computing environment.
- Web and FTP Services—The GCMRC Web site and file transfer protocol (FTP) site serve to make the mission and findings of GCMRC accessible to the public. The sites offer our updated work plan, descriptions

of our program areas, and various interactive stores of data including our Internet Map Server and our online library.

- Assistance and support of online discussion forums—The GCMRC hosts online discussions forums for the AMWG, GCMRC, and the USGS LIDAR discussion group. These forums provide a widely accessible medium for informal discussions and announcements relating to the respective topics.

Budget

ADM 12.A5.08	
GCMRC Component of SBSC Systems Admin Support (FY2005–Ongoing)	
	Fiscal year 2008
GCMRC personnel costs (19% burden)	—
GCMRC project-related travel/training (19% burden)	—
GCMRC operations/supplies (19% burden)	65,000
GCMRC equipment purchase/replacement (19% burden)	100,000
AMP logistical support (19% burden)	—
Outside GCMRC and contract science labor (19% and/or other burden rate)	5,000
Cooperative/interagency agreements (6.09% GCMRC burden plus cooperator's burden)	—
Project Subtotal	170,000
DOI customer burden (combined 6.09%, 19% and/or other rates)	32,300
Project Total (Gross)	202,300
Percent outsourced (outside of GCMRC; includes 50% of logistics)	3%

References Cited

- Anders, P., M. Bradford, P. Higgins, K.H. Nislow, Rabeni, C., and Tate, C., 2001, Final Report of the Aquatic Protocol Evaluation Program Panel: Grand Canyon Monitoring and Research Center.
- Atkinson, A.J., Trenham, P.C., Fisher, R.N., Hathaway, S.A., Johnson, B.S., Torres, S.G., and Moore, Y.C., 2004, Designing monitoring programs in an adaptive management context for regional multiple species conservation plans: U.S. Geological Survey Technical Report, 69 p.
- Austin, D.E., Osife, C.E., Drye, B., Phillips, A.M., III, Gardner, A., and Suinarich, J., 1997, 1997 Southern Paiute Consortium Colorado River corridor monitoring and education program: Summary report, Bureau of Reclamation, cooperative agreement no. 4-FC-40-15620, 12 p.
- Barnes, B., Sidhu, H.S., and Roxburgh, S.H., 2006, A model integrating patch dynamics, competing species and intermediate disturbance hypothesis: *Ecological Modelling*, v. 194, p. 414–420.
- Becker, C.D., Neitzel, D.A., and Fickeisen, D.H., 1982, Effects of dewatering on Chinook salmon redds: tolerance of four developmental phases to daily dewaterings: *Transactions of the American Fisheries Society*, v. 111, p. 624–637.
- Benenati, P.L., Shannon, J.P., and Blinn, D.W., 1998, Desiccation and recolonization of phyto­benthos in a regulated desert river: Colorado River at Lees Ferry, Arizona, USA: *Regulated Rivers: Research & Management*, v. 14, p. 519–532.
- Blinn, D.W., Shannon, J.P., Stevens, L.E., and Carder, J.P., 1995, Consequences of fluctuating discharge for lotic communities: *Journal of the North American Benthological Society*, v. 14, p. 233–248.
- Brian, N.J., and Thomas, J.R., 1984, 1983 Colorado River sand bar campsite inventory: National Park Service, Division of Resources Management, Grand Canyon National Park, Ariz.
- Chick, J.H., and McIvor, C.C., 1994, Patterns in the abundance and composition of fishes among beds of different macrophytes—viewing a littoral-zone as a landscape: *Canadian Journal of Fisheries and Aquatic Sciences*, v. 51, p. 2873–2882.
- Chick, J.H., and McIvor, C.C., 1997, Habitat selection by three littoral zone fishes: effects of predation pressure, plant density and macrophyte type: *Ecology of Freshwater Fish*, v. 6, p. 27–35.
- Clover, E.U., and Jotter, L., 1944, Floristic studies in the canyon of the Colorado and tributaries: *American Midland Naturalist*, v. 32, p. 591–642.
- Coggins, L.G., Jr., Pine, W.E., Walters, C.J., Van Haverbeke, D.R., Ward, D., and Johnstone, H.C., 2006, Abundance trends and status of the Little Colorado River population of humpback chub: *North American Journal of Fisheries Management*, v. 26, p. 233–245.
- Coggins, L.G., and Van Haverbeke, 2001, Fisheries monitoring activities in the Little Colorado River within Grand Canyon during 2000: A final report submitted to the Grand Canyon Monitoring and Research Center: Flagstaff, Ariz., U.S. Fish Wildlife Service, Arizona Fishery Resources Office, 101 p.
- Coggins, L., Yard, M.D., and Paukert, C., 2002, An operational plan: piscivory by non-native salmonids in the Colorado River and an evaluation of the efficacy of mechanical removal of non-native salmonids: U.S. Geological Survey Grand Canyon Monitoring and Research Center report, Flagstaff, Ariz.
- Doelle, W.H., 2000. Final Report: Cultural Resource Program Assessment: manuscript on file, Grand Canyon Monitoring and Research Center, Flagstaff, Ariz.
- Draut, A.E., and Rubin, D.M., 2005, Measurements of wind, aeolian sand transport, and precipitation in the Colorado River corridor, Grand Canyon, Arizona—November 2003 to December 2004: U.S. Geological Survey Open-File Report 2005-1309.
- Draut, A.E., and Rubin, D.M., 2006, Measurements of wind, aeolian sand transport, and precipitation in the Colorado River corridor, Grand Canyon, Arizona—November 2003 to December 2004: U.S. Geological Survey Open-File Report 2006-1188.

- Ecometric Inc., Colorado River Flow, Stage and Sediment Model (CRFSS) for the Colorado River in the Grand Canyon. For Grand Canyon Monitoring and Research Center, U.S. Geological Survey, Flagstaff, Ariz. Downloadable program, <http://flow.mountainsoft.net/>, accessed May 17, 2008.
- ESRI, Inc., 2005, Environmental Systems Research Institute, Inc., ArcMap Version 9.1. Redlands, CA, USA.
- Floyd, D.A., and Anderson, J.E., 1982, A new point interception frame for estimating cover of vegetation: *Vegetation*, v. 50, p. 185–186.
- Gloss, S.P., and Coggins, L.G., 2005, Fishes of Grand Canyon, in Gloss, S.P., Lovich, J.E., and Melis, T.S., eds., *The state of the Colorado River ecosystem in Grand Canyon: U.S. Geological Survey Circular 1282*, p. 33–68.
- Gloss, S.P., Lovich, J.E., and Melis, T.S., eds., 2005, *The state of the Colorado River ecosystem in Grand Canyon: U.S. Geological Survey Circular 1282*, 220 p.
- Goeking, S.A., Schmidt, J.C., and Webb, M.K., 2003, Spatial and temporal trends in the size and number of backwaters between 1935 and 2000, Marble and Grand Canyons, Arizona: Cooperative Agreement OWRAG0059.
- Gorman, O.T., and Stone, D.M., 1999, Ecology of spawning humpback chub (*Gila cypha*), in the Little Colorado River near Grand Canyon, Arizona: *Environmental Biology of Fishes*, v. 55, p. 11–133.
- Gower, J.C., 1971, A general coefficient of similarity and some of its properties: *Biometrics*, v. 27, p. 857–871.
- Hazel, J.E. Jr., Kaplinski, M., Parnell, R., and M. Manone, 2000, Sand deposition in the Colorado River ecosystem from flooding of the Paria River and the effects of the November 1997, Glen Canyon Dam Test Flow: Final Report to the Grand Canyon Monitoring and Research Center: Flagstaff, Ariz., Northern Arizona University, 37 p.
- Hazel, J.E., Jr., Kaplinski, M., Parnell, R., Manone, M., and Dale, A., 1999, Topographic and bathymetric changes at thirty-three long-term study sites, in Webb, R.H., Schmidt, J.C., Marzolf, G.R., and Valdez, R.A., eds., *The 1996 controlled flood in Grand Canyon: Washington, D.C., American Geophysical Union, Geophysical Monograph 110*, p. 161–184.
- Hazel, J., Jr., Topping, D.J., Schmidt, J.C., and Kaplinski, M., 2006, Influence of a dam on fine-sediment storage in a canyon river: *Journal of Geophysical Research*, v. 111, F01025, 16 p.
- Hereford, R., Fairley, H.C., Thompson, K.S., and Balsom, J.R., 1993, Surficial geology, geomorphology and erosion of archeologic sites along the Colorado River, eastern Grand Canyon, Grand Canyon National Park, Arizona. Grand Canyon National Park in cooperation with the Bureau of Reclamation, Glen Canyon Environmental Studies, Flagstaff, Ariz.: U.S. Geological Survey Open-File Report 93-517.
- Holling, C.S., 1978, *Adaptive environmental assessment and management: Chichester, New York, Wiley*, 785 p.
- Huston, M.A., 1979, A general hypothesis of species diversity: *American Naturalist*, v. 113, p. 81.
- Jensen, J.O.T., McLean, W.E., Rombough, P.J., and Septav T., 1992, Salmonid incubation and rearing programs for IBM-compatible computers: Canadian Technical Report of Fisheries and Aquatic Sciences 1878, 46 p.
- Kaeding, L.R., and Zimmerman, M.A., 1983, Life history and ecology of the humpback chub in the Little Colorado and Colorado Rivers of the Grand Canyon: *Transactions of the American Fisheries Society*, v. 112, p. 577–594.
- Kaplinski, M., Behan, J., Hazel, J.E., Jr., Manone, M., and Parnell, R., 2003, Evaluation of campsite studies in the Colorado River ecosystem: analysis and recommendations for long-term monitoring: Final report for Cooperative Agreement 00PG400255,0001. On file, Grand Canyon Monitoring and Research Center, Flagstaff, Ariz.
- Kaplinski, M., Behan, J., Hazel, J.E., Jr., Parnell, R.A., and Fairley, H.C., 2005, Recreational values and campsites in the Colorado River ecosystem, in Gloss, S.P., Lovich, J.E., and Melis, T.S., eds., *The state of the Colorado River ecosystem in Grand Canyon: U.S. Geological Survey Circular 1282*, p. 193–205.
- Kaplinski, M., Hazel, J.E., Jr., Parnell, R., Breedlove, M., and Schmidt, J.C., 2007, Integrating bathymetric, topographic, and LiDAR surveys of the Colorado River in Grand Canyon to assess the effect of a flow

- experiment from Glen Canyon Dam on the Colorado River ecosystem: Proceedings of the Hydrographic Society of America 2007 Annual Meeting, May 14–17, Norfolk, Va., 22 p.
- Kaplinski, M., Hazel, J.E., Parnell, R., Manone, M., and Gonzales, M., 2000, Evaluation of hydrographic survey techniques used for channel mapping by the Grand Canyon Monitoring and Research Center in the Colorado River ecosystem, Grand Canyon, Arizona: Final Report to the Grand Canyon Monitoring and Research Center, Northern Arizona University, Flagstaff, Ariz., 37 p.
- Kearsley, L.H., 1995, Monitoring the effects of Glen Canyon Dam interim flows on campsite size along the Colorado River in Grand Canyon National Park (final report): National Park Service, Division of Resources Management, Grand Canyon National Park.
- Kearsley, M.J.C., 2006. Vegetation dynamics in Kearsley, M.J.C. ed., Inventory and monitoring of terrestrial riparian resources in the Colorado River corridor of Grand Canyon: an integrative approach: Final Report from Northern Arizona University submitted to Grand Canyon Monitoring and Research Center, U.S. Geological Survey, Flagstaff, Ariz., 262 p.
- Kearsley, M.J.C., and Ayers, T.J., 1996, The effects of interim flows from Glen Canyon Dam on riparian vegetation in the Colorado River corridor, Grand Canyon National Park, Arizona. Final report to Grand Canyon Science Center, Grand Canyon National Park, Grand Canyon, Ariz., 724 p.
- Kearsley, M.J.C., Cobb, N., Yard, H., Lightfoot, D., Brantley, S., Carpenter, G., and Frey, J., 2006, Inventory and monitoring of terrestrial riparian resources in the Colorado River corridor of Grand Canyon: an integrative approach: 2004 draft report to the U.S. Geological Survey Grand Canyon Monitoring and Research Center, cooperative agreement no. 01-WRAG-0044 and 01-WRAG-0034, 126 p.
- Kearsley, L.H., and Quartaroli, R., 1997, Effects of a sand bar/habitat building flow on campsites in Grand Canyon: Final report of Applied Technology Associates for the Glen Canyon Environmental Studies, 18 p.
- Kearsley, L.H., Schmidt, J.C., and Warren, K.D., 1994, Effects of Glen Canyon Dam on Colorado River sand deposits used as campsites in Grand Canyon National Park, USA: Regulated Rivers: Research & Management, v.9, p. 137–149.
- Kearsley, L.H., and Warren, K., 1993, River campsites in Grand Canyon National Park: inventories and effects of discharge on campsite size and availability (final report): National Park Service, Division of Resources Management, Grand Canyon National Park.
- Korman, Josh, Kaplinski, Matthew, Hazel, J.E., III, and Melis, T.S., 2005, Effects of the experimental fluctuating flows from Glen Canyon Dam in 2003 and 2004 on the early life history stages of rainbow trout in the Colorado River: Final report, prepared for the Grand Canyon Monitoring and Research Center, cooperative agreement no. 04WRAG0006, modification no. 002.
- Korman, Josh, Yard, Mike, and Speas, Dave, 2006, An evaluation of the utility of snorkel surveys for estimating population size and tracking trends in relative abundance of rainbow trout in the Lee's Ferry reach of the Colorado river. Final report prepared for the Arizona Game and Fish Department, 43 p.
- Leibfried, W.C., 1988, M.E., 2002, The utilization of Cladophora Glomerata and epiphytic diatoms as a food resource by rainbow trout in the Colorado River below Glen Canyon Dam, Arizona Riparian vegetation responses to contrasting managed flows of the Colorado River in Grand Canyon, Arizona: Flagstaff, Northern Arizona University, M.S. thesis, 33 p.
- Lomaomvaya, M., Ferguson, T.J., and Yeatts, M., 2001, Ongtuvqava Sakwtala: Hopi ethnobotany in the Grand Canyon: Submitted to the Grand Canyon Monitoring and Research Center.
- Loomis, J., Cole, D., Foti, P., Manning, R., Moisey, R. N., Ratcliffe, R., 2005. Final Report of the Protocol Evaluation Panel (PEP) for the Recreation Monitoring Program of the Grand Canyon Monitoring and Research Center (GCMRC): report dated September 28, 2005, on file, Grand Canyon Monitoring and Research Center, Flagstaff, Ariz.
- Maddux, H.R., Kubly, D.M., DeVos, J.C., Jr., Persons, W.R., Staedicke, R., and Wright, R.L., 1987, Effects of varied flow regimes on aquatic resources of Glen and Grand Canyons: Bureau of Reclamation Glen Canyon Environmental Studies report, National Technological Information Series PB88-1 83439/AS.

- Martin, T. and Whitis, D., 2004, Guide to the Colorado River in the Grand Canyon: Lees Ferry to South Cove: Flagstaff, Ariz., Vishnu Temple Press.
- McKinney, T., Rogers, R.S., and Persons, W.R., 1999, Effects of flow reductions on aquatic biota of the Colorado River below Glen Canyon Dam, Arizona: *North American Journal of Fisheries Management*, v. 19, p. 984–991.
- McKinney, Ted, Speas, D.W., Rogers, R.S., and Persons, W.R., 2001, Rainbow trout in a regulated river below Glen Canyon Dam, Arizona, following increased minimum flows and reduced discharge variability: *North American Journal of Fisheries Management*, v. 21, p. 216–222.
- Melis, T.S., Topping, D.J., and Rubin D.M., 2003, Testing laser-based sensors for continuous in situ monitoring of suspended sediment in the Colorado River, Arizona, in Bogen, J., Fergus, T., and Walling, D.E., eds., *Erosion and sediment transport measurement in rivers: technological and methodological advances*: Wallingford, Oxfordshire, United Kingdom, IAHS Press, IAHS Publication 283, p. 21-27.
- Melis, T.S., Wright, S.A., Ralston, B., Fairley, H., Kennedy, T., Coggins, L., and Korman, J., 2006, Knowledge assessment of the effects of Glen Canyon Dam on the Colorado River ecosystem, 77 p.
- Naiman, R.J., Décamps, H., McClain, M.E., eds., 2005, *Riparia: Ecology, Conservation and Management of Streamside Communities*: London, Elsevier Academic Press.
- Nakano, S., and Murakami, M., 2001, Reciprocal subsidies: dynamic interdependence between terrestrial and aquatic food webs: *Proceedings of the National Academy of Sciences*, v. 98, p. 166–170.
- National Research Council, 1996, *River Resource Management in the Grand Canyon*: Washington, D.C., National Academy Press, 226 p.
- Palmer, M., Baron, J., Dale, V., Gunderson, L., Howard, A., Kitchell, J., Robertson, D., Schwartz, D., Watkins, J., and Garrett, D., 2004, A Review of the GCMRC Food Base Science Program by the GCD AMP Science Advisors, 16 p. <http://www.gcmrc.gov/webopac/main?siteid=GCMRC>, accessed May 19, 2008.
- Pederson, J.L., Petersen, P.A., MacFarlane, W.W., Gonzales, M.F., and Kohl, K., 2003, Mitigation, monitoring, and geomorphology related to gully erosion of cultural sites in Grand Canyon. Final report in fulfillment of CA-01-WRAG-0074: on file, Grand Canyon Monitoring and Research Center, Flagstaff, Ariz.
- Pelicice, F.M., Agostinho, A.A., and Thomaz, S.M., 2005, Fish assemblages associated with *Egeria* in a tropical reservoir: investigating the effects of plant biomass and diel period: *Acta Oecologia-International Journal of Ecology*, v. 27, p. 9–16.
- Phillips, A.M., III, and Jackson, L., 1996, Evaluation and mitigation efforts for the March 1996 Colorado River test flow experiment: Final report: Peach Springs, Ariz., Hualapai Tribe Cultural Resources Division, 10 p.
- Porter, M.E., 2002, Riparian vegetation responses to contrasting managed flows of the Colorado River in Grand Canyon, Arizona: Flagstaff, Northern Arizona University, M.S. thesis, 33 p.
- Randle, T.J. and Pemberton, E.L., 1987, Results and analysis of STARS (sediment transport and river simulation model) modeling efforts of the Colorado River in Grand Canyon: Final Report to Glen Canyon Environmental Studies: Salt Lake City, Utah, U.S. Bureau of Reclamation, NTIS PB88-183421, 182 p.
- Rubin, D.M., 2004, A simple autocorrelation algorithm for determining grain size from digital images of sediment: *Journal of Sedimentary Research*, v. 74, p. 160–165.
- Rubin, D.M., Chezar, H., Harney, J.N., Topping, D.J., Melis, T.S., and Sherwood, C.R., 2006, Underwater microscope for measuring spatial and temporal changes in bed-sediment grain size: U.S. Geological Survey Open-File Report 2006-1360, <http://pubs.usgs.gov/of/2006/1360/>, accessed March 30, 2008.
- Rubin, D.M., Chezar, H., Topping, D.J., Melis, D.J., and Harney, J., in press, Two new approaches for measuring spatial and temporal changes in bed-sediment grain size: *Sedimentary Geology*.
- Rubin, D.M., Topping, D.J., Schmidt, J.C., Hazel, J., Kaplinski, K., and Melis, T.S., 2002, Recent sediment studies refute Glen Canyon Dam hypothesis: *EOS, Transactions, American Geophysical Union*, v. 83, n. 25, p. 273, 277–278.

- Roxburgh, S.H., Shea, K., and Wilson, J.B., 2004, The intermediate disturbance hypothesis: patch dynamics and mechanisms of species coexistence: *Ecology*, v. 85, p. 359–371.
- Schmidt, J.C., Topping, D.J., Grams, P.E., and Hazel, J.E., 2004, System-wide changes in the distribution of fine sediment in the Colorado River corridor between Glen Canyon Dam and Bright Angel Creek, Arizona: Final report submitted to Grand Canyon Monitoring and Research Center, Flagstaff, Ariz., 117 p., http://www.gcmrc.gov/library/reports/physical/Fine_Sed/Schmidt2004.pdf, accessed March 30, 2008.
- Shannon, J.P., Blinn, D.W., Haden, G.A., Benenati, E.P., and Wilson, K.P., 2001, Food web implications of delta C-13 and delta N-15 variability over 370 km of the regulated Colorado River, USA: *Isotopes in Environmental and Health Studies*, v. 37, p. 179–191.
- Shaver, M.L., Shannon, J.P., Wilson, K.P., Benenati, P.L., and Blinn, D.W., 1997, Effects of suspended sediment and desiccation on the benthic tailwater community in the Colorado River, USA: *Hydrobiologia*, v. 357, p. 63–72.
- Simoes, F.J., Goodwin, P., Hanes, D.M., Sherwood, C., Schoellhamer, D.H., and Sloff, K., 2007, Final Report of the External Review Panel, Sediment Transport Modeling Review Workshop: U.S. Geological Survey Grand Canyon Monitoring and Research Center Administrative Document, 14 p.
- Stevens, L. E., 1994, *The Colorado River in Grand Canyon: A Comprehensive Guide to Its Natural and Human History* (4th ed.): Flagstaff, Ariz., Red Lake Books
- Stevens, L.E., and Ayers, T.J., 1993, The impacts of Glen Canyon Dam on riparian vegetation and soil stability in the Colorado River corridor, Grand Canyon, Arizona. Final report to The National Park Service, Cooperative Park Studies Unit, Northern Arizona University, Flagstaff, Ariz., 54 p.
- Stevens, L.E., Schmidt, J.C., Ayers, T.J., and Brown, B.T., 1995, Flow regulation, geomorphology, and Colorado River marsh development in the Grand Canyon, Arizona: *Ecological Applications*, v. 5, p. 1025–1039.
- SWCA, Inc., 2000, A program of experimental flows for endangered and native fishes of the Colorado River in Grand Canyon. Final report to Grand Canyon Monitoring and Research Center, U.S. Geological Survey, Flagstaff, Ariz, 57 p.
- Topping, D.J., Melis, T.S., Rubin, D.M., and Wright, S.A., 2004, High-resolution monitoring of suspended-sediment concentration and grain size in the Colorado River in Grand Canyon using a laser-acoustic system, in Hu, C., and Tan, Y, eds., *Proceedings of the Ninth International Symposium on River Sedimentation*, October 18–21, 2004: Yichang, China: People's Republic of China, Tsinghua University Press, p. 2507–2514.
- Topping, D.J., Rubin, D.M., and Schmidt, J.C., 2008, Update on regulation of sand transport in the Colorado River by changes in the surface grain size of eddy sandbars over multiyear timescales: U.S. Geological Survey Scientific Investigations Report 2008-5042, 24 p., <http://pubs.usgs.gov/sir/2008/5042/>, accessed May 19, 2008.
- Topping, D.J., Rubin, D.M., Schmidt, J.C., Hazel, J.E., Jr., Melis, T.S., Wright, S.A., Kaplinski, M., Draut, A.E., and Breedlove, M.J., 2006, Comparison of sediment-transport and bar-response results from the 1996 and 2004 controlled-flood experiments on the Colorado River in Grand Canyon: CD-ROM Proceedings of the 8th Federal Inter-Agency Sedimentation Conference, Reno, Nev., April 2-6, 2006, ISBN 0-9779007-1-1.
- Topping, D.J., Rubin, D.M., and Vierra, L.E., Jr., 2000, Colorado River sediment transport 1. Natural sediment supply limitation and the influence of Glen Canyon Dam: *Water Resources Research*, v. 36, p. 515–542.
- Topping, D.J., Schmidt, J.C., and Vierra, L.E., Jr., 2003, Computation and analysis of the instantaneous-discharge record for the Colorado River at Lees Ferry, Arizona—May 8, 1921, through September 30, 2000: Reston, Va., U.S. Geological Survey Professional Paper 1677, 118 p.
- Topping, D.J., Wright, S.A., Melis, T.S., and Rubin, D.M., 2006, High-resolution monitoring of suspended-sediment concentration and grain size in the Colorado River using laser-diffraction instruments and a three-frequency acoustic system: CD-ROM Proceedings of the 8th Federal Inter-Agency Sedimentation Conference, Reno, Nev., April 2–6, 2006, ISBN 0-9779007-1-1.
- Topping, D.J., Wright, S.A., Melis, T.S., and Rubin, D.M., in press, High-resolution measurements of suspended-sediment concentration and grain size in the Colorado River in Grand Canyon using a multi-frequency acoustic

- system: Proceedings of the Tenth International Symposium on River Sedimentation, August 1–4, 2007, Moscow, Russia.
- Turner, R.M., and Karpiscak, M.M., 1980, Recent vegetation changes along the Colorado River between Glen Canyon Dam and Lake Mead, Arizona: U.S. Geological Survey Professional Paper 1132, 125 p.
- Tyus, H.M., and Saunders, J.F., III, 2000, Nonnative fish control and endangered fish recovery: lessons from the Colorado River: *Fisheries*, 25:17–24.
- U.S. Army Corps of Engineers, 1994, Engineering and design-topographic accuracy standards: EM1110-1-1005, p. 2–1 to 2–12.
- U.S. Department of the Interior, 1995, Operation of Glen Canyon Dam Final Environmental Impact Statement: Bureau of Reclamation, Upper Colorado Region, 337 p. plus appendices, Salt Lake City, Utah.
- Valdez, R.A., and Ryel, R.J., 1995, Life history and ecology of the humpback chub (*Gila cypha*) in the Colorado River, Grand Canyon, Arizona: Final report to the Bureau of Reclamation, Salt Lake City, Utah, contract no. 0-CS-40-09110: Logan, Utah, BIO/WEST, Inc.
- Van Haverbeke, D.R., 2003, Stock assessment and fisheries monitoring activities in the Little Colorado River within Grand Canyon during 2002: final report submitted to the Grand Canyon Monitoring and Research Center: Flagstaff, Ariz., U.S. Fish Wildlife Service, Arizona Fishery Resources Office, 87 p.
- Van Haverbeke, D.R., 2004, Stock assessment and fisheries monitoring activities in the Little Colorado River within Grand Canyon during 2003: final report submitted to the Grand Canyon Monitoring and Research Center: Flagstaff, Ariz., U.S. Fish Wildlife Service, Arizona Fishery Resources Office, 74 p.
- Van Haverbeke, D.R., and Coggins, L.G., Jr., 2003, Stock assessment and fisheries monitoring activities in the Little Colorado River within Grand Canyon during 2001: final report submitted to the Grand Canyon Monitoring and Research Center: Flagstaff, Ariz., U.S. Fish Wildlife Service, Arizona Fishery Resources Office, 87 p.
- Walters, C.J., 1986, Adaptive management of renewable resources: New York, Macmillan, 374 p.
- Walters, C., Korman, J., Stevens, L.E., and Gold, B., 2000, Ecosystem modeling for evaluation of adaptive management policies in the Grand Canyon: *Journal of Conservation Ecology*, v. 4, no.2, <http://www.consecol.org/vol4/iss2/art1>, accessed May 19, 2008.
- Ward, J.H., 1963, Hierarchical Grouping to optimize an objective function: *Journal of American Statistical Association*, v. 58, no. 301, p. 236–244.
- Waring, G.L., 1995, Current and historical riparian vegetation trends in Grand Canyon, using multitemporal remote sensing analyses of GIS sites: Final report, National Park Service, 24 p.
- Weeden, H., Borden, F., Turner, B., Thompson, D., Strauss, C., and Johnson, R., 1975, Grand Canyon National Park campsite inventory, contract no. CX 001-3-0061 with the National Park Service: University Park, Pennsylvania State University.
- Wohl, E., Bennett, J.P., Blum, M.D., Grant, G.E., Hanes, D.M., Howard, A.D., Mueller, D.S., Schoellhamer, D.H., and Simoes, F.J., 2006, Protocols Evaluation Program (PEP-SEDS III): Final report of the physical resources monitoring peer review panel. U.S. Geological Survey Grand Canyon Monitoring and Research Center Administrative Document, 27 p., <http://www.gcmrc.gov/library/reports/PEP/Wohl2006.pdf>, accessed may 19, 2008.
- Wright, S.A., Melis, T.S., Topping, D.J., and Rubin, D.M., 2005, Influence of Glen Canyon Dam operations on downstream sand resources of the Colorado River in Grand Canyon, in Gloss, S.P., Lovich, J.E., and Melis, T.S., eds., *The state of the Colorado River ecosystem in Grand Canyon*: U.S. Geological Survey Circular 1282, p. 17–31.
- Yeatts, M., 1996, High elevation sand retention following the 1996 Spike Flow: An Assessment for Cultural Resources Stabilization, chap. 4 of Balsom, J.R., and Larralde, Signa, eds., *Mitigation and Monitoring of*

Cultural Resources in Response to the Experimental Habitat Building Flow in Glen and Grand Canyons, Spring 1996: Salt Lake City, Utah, Bureau of Reclamation, Upper Colorado River Region.

Glossary

Acronym	Term
ASP	Active server pages
AMPSP	Final Draft GCDAMP Strategic Plan
AMWG	Adaptive Management Work Group
ASMR	Age-structured mark recapture
AZGFD	Arizona Game and Fish Department
BHBF	Beach/habitat-building flows
Bureau of Reclamation	Reclamation
CEM	Conceptual ecosystem model
CMINs	Core monitoring information needs
CRAHG	Cultural Resources Ad Hoc Group
CRE	Colorado River ecosystem
CREDA	Colorado River Energy Distributors Association
CRMP	Colorado River Management Plan
DASA	Data Acquisition, Storage, and Analysis Program
EIS	Environmental impact statement
FEIS	Final Environmental Impact Statement
FIST	Fine-grained integrated sediment transport
GCD	Glen Canyon Dam
GCDAMP	Glen Canyon Dam Adaptive Management Program
GCMRC	Grand Canyon Monitoring and Research Center
GIS	Geographic Information Systems
GCNP	Grand Canyon National Park
HBC	Humpback chub
HEC-RAS	Hydrologic Engineering Center River Analysis System
IDE	Integrated drive electronics
IQW	Integrated quality of water
ISA	WAPA Energy Tracking Database
KA	Knowledge assessment
KAR	Knowledge Assessment Report
KAS	Kanab ambersnail
KAW	Knowledge Assessment Workshop
LCR	Lower Colorado River
LSSF	Low steady summer flows
LTEP	Long-Term Experimental Plan
LTO	Linear tape open
MLFF	Modified low fluctuating flows
MRP	Monitoring and Research Plan
NAS	Networked attached storage
NAU	Northern Arizona University
NGS	National Geodetic Survey
NHPA	National Historic Preservation Act
NPS	National Park Service
OHWZ	Old high water zone
PCMP	Provisional Core Monitoring Plan
PEP	Protocols evaluation panel
PIT	Passive integrated transponder
QA/QC	Quality Assurance/Quality Control
RBT	Rainbow trout
RFP	Request for Proposals
RINs	Research information needs
ROD	Record of Decision
RTK	Real-time kinematics

SA	Science Advisors
SAB	Science Advisors Board
SBSC	Southwest Biological Science Center
SCORE	The State of the Colorado River Ecosystem in Grand Canyon report
SCSI	Small computer system interface
SED TREND	Direct topographic/bathymetric measurements and remote sensing
SEDS-PEP	Sediment transport modeling review-protocols evaluation panel
SHOALS	Scanning hydrographic operation airborne LIDAR survey
SINs	Supporting information needs
SMS	Systems management server
SPG	Science planning group
SSP	Strategic Science Plan
SSQs	Strategic science questions
STARS	Sediment transport and river simulation model
SUS	System update server
SWCA	SWCA Environmental Consultants
TCD	Temperature control device
TCP/IP	Transmission control protocol/internet protocol TCP/IP
TCPs	Traditional cultural properties
TEM	Terrestrial ecosystem monitoring
TWG	Technical Work Group
USFWS	U.S. Fish and Wildlife Service
USGS	U.S. Geological Survey
USU	Utah State University
WAPA	Western Area Power Administration
YoY	Young-of-year

Appendix A. Key Science Questions Addressed in the FY2007–11 Science Program

AMWG Priority 1: Why are the humpback chub not thriving, and what can we do about it? How many humpback chub are there and how are they doing? (GCDAMP goal 2)

Key Strategic Science Questions

1. To what extent are adult populations of native fish controlled by production of young fish from tributaries, spawning and incubation in the mainstem, survival of young-of-year (YoY) and juvenile stages in the mainstem, or by changes in growth and maturation in the adult population as influenced by mainstem conditions? [FY2006–11]
2. Does a decrease in the abundance of rainbow trout (RBT) and other cold- and warmwater nonnatives in Marble and eastern Grand Canyons result in an improvement in the recruitment rate of juvenile humpback chub to the adult population? [FY2006–11]
3. Do RBT immigrate from Glen to Marble and eastern Grand Canyons, and, if so, during what life stages? To what extent do Glen Canyon immigrants support the population in Marble and eastern Grand Canyons? [FY2007–11]
4. Can long-term decreases in abundance of RBT in Marble and eastern Grand Canyons be sustained with a reduced level of effort of mechanical removal or will recolonization from tributaries and from downstream and upstream of the removal reach require that mechanical removal be an ongoing management action? This question also applies to future removal programs targeting other nonnative species. [FY2007–11]
5. What are the important pathways, and the rate of flux among them, that link lower trophic levels with fish and how will they link to dam operations? [FY2006–09]
6. Are trends in the abundance of fish populations, or indicators from fish such as growth, condition, and body composition (e.g., lipids), correlated with patterns in invertebrate flux? [FY2006–09].
7. Which tributary and mainstem habitats are most important to native fishes and how can these habitats best be made useable and maintained? [FY2008–09].
8. How can native and nonnative fishes best be monitored while minimizing impacts from capture and handling or sampling? [FY2007–11].

AMWG Priority 2: Which cultural resources, including traditional cultural properties, are within the Area of Potential Effect, which should we treat, and how do we best protect them? What is the status and trends of cultural resources and what are the agents of deterioration? (GCDAMP goal 11).

Key Strategic Science Questions

1. Do dam-controlled flows affect (increase or decrease) rates of erosion and vegetation growth at archaeological sites and traditional cultural properties (TCP) sites, and if so, how? [FY2007–11]
2. How do flows impact old high-water zone terraces in the Colorado River ecosystem (CRE) (where the majority of archaeological sites occur), and what kinds of important information about the historical ecology and human history of the CRE are being lost due to ongoing erosion of the Holocene sedimentary deposits? [FY2004–11]
3. If dam-controlled flows are contributing to (influencing rates of) archaeological site/TCP erosion, what are the optimal flows for minimizing future impacts to historic properties? [FY2009–11]
4. How effective are various treatments (e.g., check dams, vegetation management, etc.) in slowing rates of erosion at archaeological sites over the long term? [FY2006–11]
5. What are the TCPs in the CRE, and where are they located? [FY2006–11]
6. How can tribal values/data/analyses be appropriately incorporated into a science-driven adaptive management process in order to evaluate the effects of flow operations and management actions on TCPs? [FY2006–08]
7. Are dam-controlled flows affecting TCPs and other tribally valued resources in the CRE, and, if so, in what respects are they being affected, and are those effects considered positive or negative by the tribes who value these resources? [FY2006–11]

AMWG Priority 3: What is the best flow regime? (GCDAMP goals 1–11)

Key Strategic Science Questions

1. Is there a “Flow-Only” operation (i.e., a strategy for dam releases, including managing tributary inputs with BHBFs, without sediment augmentation) that will restore and maintain sandbar habitats over decadal timescales? [FY2008–11]
2. To what extent could predation impacts by nonnative fish be mitigated by higher turbidities or dam-controlled high-flow releases? [FY2007–08]
3. What are the hydropower replacements costs of the modified low fluctuating flow (MLFF) (annually, since 1996)? [FY2007–08]
4. What are the projected hydropower costs associated with the various alternative flow regimes being discussed for future experimental science (as defined in the next phase experimental design)? [FY2006–07]
5. How is invertebrate flux affected by water quality (e.g., temperature, nutrient concentrations, turbidity) and dam operations? [FY2006–08]
6. What Glen Canyon Dam operations (ramping rates, daily flow range, etc.) maximize trout fishing opportunities and catchability? [FY2007–08]
7. How do dam-controlled flows affect visitors’ recreational experiences, and what is/are the optimal flows for maintaining a high-quality recreational experience in the CRE? [FY2007–08]

8. What are the drivers for recreational experiences in the CRE, and how important are flows relative to other drivers in shaping recreational experience outcomes? [FY2007–09]
9. How do varying flows positively or negatively affect campsite attributes that are important to visitor experience? [FY2009–11]
10. How can safety and navigability be reliably measured relative to flows? [FY2007–08]
11. How do varying flows positively or negatively affect visitor safety, health, and navigability of the rapids? [FY2007–09]
12. How do varying flows regimes positively or negatively affect group encounter rates, campsite competition, and other social parameters that are known to be important variables of visitor experience? [FY2007–09]

AMWG Priority 4: What is the impact of sediment loss and what should we do about it? (GCDAMP goal 8)

Key Strategic Science Questions

1. Is there a “Flow-Only” operation (i.e., a strategy for dam releases, including managing tributary inputs with BHBFs, without sediment augmentation) that will restore and maintain sandbar habitats over decadal timescales? (FY2008–11)
2. How important are backwaters and vegetated shoreline habitats to the overall growth and survival of YoY and juvenile native fish? Does the long-term benefit of increasing these habitats outweigh short-term potential costs (displacement and possibly mortality of young humpback chub) associated with high flows? [FY2007–11]

AMWG Priority 5: What will happen when we test or implement the Temperature Control Device (TCD)? How should it be operated? Are safeguards needed for management? (GCDAMP goals 1–4 and 7–10)

Strategic Science Questions

1. How do dam release temperatures, flows (average and fluctuating component), meteorology, canyon orientation and geometry, and reach morphology interact to determine mainstem and nearshore water temperatures throughout the CRE? [FY2006–08]
2. How is invertebrate flux affected by water quality (e.g., temperature, nutrient concentrations, turbidity) and dam operations? [FY2006–08]
3. To what extent do temperature and fluctuations in flow limit spawning and incubation success for native fish? [FY2003–08]
4. What is the relative importance of increased water temperature, shoreline stability, and food availability on the survival and growth of YoY and juvenile native fish? [FY2003–08]
5. Will increased water temperatures increase the incidence of Asian tapeworm in humpback chub or the magnitude of infestation, and if so, what is the impact on survival and growth rates? [FY2003–08]
6. Do the potential benefits of improved rearing habitat (warmer, more stable, more backwater and vegetated shorelines, more food) outweigh negative impacts due to increases in nonnative fish abundance? [FY2007–11]
7. How do warmer releases affect viability and productivity of native/nonnative vegetation? [FY2007–11]

Appendix B. GCDAMP Fiscal Year 2008 Budget Explanatory Material

The draft FY2008 GCDAMP budget, which includes budgets for GCDAMP activities performed by Reclamation and the U.S. Geological Survey (USGS) Grand Canyon Monitoring and Research Center, is attached separately. Table B.1 explains the information found in various columns of the budget document. Following the table is an explanation of USGS policy on cost-recovery accounting and cost share.

Table B.1. Explanation of information found in columns of draft fiscal year 2008 (FY2008) Glen Canyon Dam Adaptive Management Program (GCDAMP) budget.

Column	Title	Key
A	GCMRC Project ID	Column 1–3 Program Area BIO: Biology PHY: Physical Science REC: Recreation HYD: Hydropower CUL: Cultural DASA: Data Acquisition, Storage and Analysis SUP: Support ADM: Administration PLA: Planning Column 4–5 GCDAMP goal number Column 6–7 project number Column 8–9 fiscal year
B	Status	O: Ongoing N: New C: Complete D: Deferred NA: Not applicable
C	Funding emphasis	APM: Administrative program management. Activities/projects that are administrative in nature or are conducted in support of the overall GCMRC science program, including base funding for program managers, logistics staff and permanent DASA staff. COR: Core-monitoring project. Monitoring projects that have been piloted, subjected to initial and secondary protocols evaluation panel (PEP) reviews, documented through a core-monitoring report and formally adopted as a core-monitoring project by the TWG. CRD: Core-monitoring research and development project. Monitoring projects that are currently undergoing research and development, including projects that have been piloted and peer reviewed but which have not yet been formally documented with a core-monitoring report or formally adopted as a core-monitoring project by the TWG. LTE: Long-term experiment. Projects specifically undertaken as part of or in direct support of the Long-Term Experimental Plan. ORD: Other research and development projects. Other research projects or R&D work that is NOT directly tied to the development of core-monitoring projects.
D	Project description	Project title (start date–end date)
F	Estimated FY2008 budget	Estimated FY2008 gross cost of project

Explanation of USGS Policy on Cost Share

In FY2003, the U.S. Geological Survey (USGS) began full-cost recovery accounting and instituted a Department of the Interior (DOI) customer rate of 15 percent against all DOI agency reimbursable funding. In FY2007, the customer rate is estimated at the 15-percent DOI customer rate with an additional 2 percent added to achieve the required additional facilities costs. The DOI customer rate was established by the USGS Bureau Headquarters and determined to be significantly lower than the “full” burden rate that varies annually and includes facilities and the Cost Center and the Bureau-level burdens. In addition to the above rates, a special “pass through” rate of 6 percent was also instated. As a transitional aid to GCMRC, which had received under a previous administration the guarantee that USGS would not charge the power revenue funds any burden, the Bureau allowed the entire GCMRC power revenue budget to be charged only the 6-percent special rate (3 percent was retained by the Cost Center and 3 percent by Headquarters) for FY2003 only.

In FY2004, USGS Headquarters approved the special rate of 6 percent for only a portion of GCMRC’s power revenue funding. This rate was applied to approximately \$2 million of funding that went directly to GCMRC cooperators. The balance of power revenue funds were charged the full DOI customer rate of 15 percent. As a part of the full-cost recovery policy, the USGS established a process referred to as cost share as a means of handling a limited electronic financial system.

Cost share is the funding that “covers” the balance of the full burden rate minus the DOI customer rate. In most cases, reimbursable funding from non-DOI agencies is charged the full burden rate. In FY2004, the full burden rate for GCMRC was approximately 30 percent. The difference between the full rate of 30 percent and the DOI customer rate of 15 percent equals 15 percent (all percentages are approximate). In FY2004 the cost share funding requirement for all DOI agency reimbursable dollars received by GCMRC equaled almost \$1 million. USGS policy requires that cost share funding be from appropriated dollars only, and those funds are also charged the Cost Center burden rate. In essence, the \$1 million appropriation provided by USGS to GCMRC in FY2004 had the effect of not adding funding, but merely filling the holes created by the cost share policy.

In FY2005 and FY2006 the USGS appropriation requested for GCMRC (also \$1 million each fiscal year) was used for cost share funding. However, information has been forwarded to the GCMRC that the required DOI cost share funds will be provided by the USGS Headquarters, so as to continue allowing for the reduced customer rate to apply to the GCMRC science program in FY2007. Per the full-cost accounting policy and the requirement that cost share dollars be appropriated dollars only, the effect of these appropriations is entirely transparent and does not add funding to the GCDAMP. The issue relating to how these cost share funds are derived in the future is a major area of concern for the GCMRC science program.

Appendix C. GCDAMP Fiscal Year 2008 Budget

The oversized budget sheets are provided starting on the following page.

APPENDIX C
Draft GCDAMP FY08 Budget for the USBR and the USGS GCMRC
03/13/2008 VERSION

	A	B	C	D	E	F	G	H	I	J	K	L	M	N	O	P	
			ID	Project Descriptions	Approved BOR FY07 Budget (inc. CPI increase)	BOR Proposed FY08 Budget - Revised 12/06/07		Comments									
1																	
2	Reclamation Administration Power Revenue Under Cap Funded Projects																
3			A	Adaptive Management Work Group													
4			1	Personnel Costs	154,628	158,958											
5			2	AMWG Member Travel Reimbursement	16,197	16,651											
6			3	Reclamation Travel	13,390	13,765											
7			4	Facilitation Contract	25,000	25,700											
8			5	POAHG Expenses	51,500	52,942											
9			6	Other	7,390	7,597											
10				Reclamation AMWG Subtotal	268,105	275,612											
11			B	Technical Work Group													
12			1	Personnel Costs	70,657	72,635											
13			2	TWG Member Travel Reimbursement	22,211	22,833											
14			3	Reclamation Travel	16,375	16,834											
15			4	TWG Chair Reimbursement	22,835	23,474											
16			5	Other	2,112	2,171											
17				Reclamation TWG Subtotal	134,190	137,947											
18			C	Other													
19			1	Compliance Documents	263,622	271,003											
20			2	Administrative Support for NPS Permitting	110,000	113,300											
21			3	Contract Administration	32,413	33,321											
22			4	Experimental Carryover Funds - to be held by BOR	500,000	500,000											
23			5	Integrated Tribal Resources Monitoring	132,500	136,210											
24			6	USFWS HBC Genetics Mgmt Plan	0	-											
25				Other Subtotal	1,038,535	1,053,834											
26				Reclamation Administrative Subtotal	1,440,830	1,467,393											
27			D	Programmatic Agreement Cultural Resources													
28			1	Reclamation Administration	71,892	57,354											
29			2	NPS Support for Archaeological Site Assessment	67,500	-											
30			3	NN & GLCA Treatment Plan and Implementation	-	-											
31			4	Canyon Treatment Plan and Implementation	145,000	300,000											
32				Programmatic Agreement Subtotal	284,392	357,354											
33				Reclamation Power Revenue Under Cap Program Subtotal:	1,725,222	1,824,747											
34																	
35				Reclamation Appropriated Funded Projects													
36			HCA	Development of a LCR Management Plan	-	-											
37				Tribal Consultation													
38			A	Cooperative Agreements with Tribes													
39			1	Hopi Tribe	95,000	95,000											
40			2	Hualapai Tribe	95,000	95,000											
41			3	Navajo Nation	95,000	95,000											
42			4	Pueblo of Zuni	95,000	95,000											
43			5	Southern Paiute	95,000	95,000											
44			6	DOI Handling Fee	-	-											
45				Tribal Consultation Subtotal	475,000	475,000											
46				Reclamation Appropriated Projects Subtotal:	475,000	475,000											
47																	
48				BUREAU OF RECLAMATION TOTAL AMP PROGRAM COSTS:	2,200,222	2,299,747											

APPENDIX C
Draft GCDAMP FY08 Budget for the USBR and the USGS GCMRC
03/13/2008 VERSION

A	B	C	D	E	F	G	H	I	J	K	L	M	N	O	P	
GCMRC Project ID	Status	Funding Emphasis	Project Descriptions	Approved FY07 Budget (inc. CPI Increase)	Proposed FY08 Budget - Gross (inc. Burden) FINAL VERSION	DOI Customer Burden (Combined 6.09%, 19% and/or Other Rate)	Project Subtotal (w/o Burden)	GCMRC Personnel Costs (19% Burden)	GCMRC Project Related Travel / Training (19% Burden)	GCMRC Operations / Supplies (19% Burden)	GCMRC Equipment Purchase / Replacement (19% Burden)	AMP Logistics Support (19% Burden)	Outside GCMRC Contract & Science Labor (19% and/or Other Burden Rate)	Coop & Inter Agency Agmts (6.09% GCMRC Burden plus Cooperator's Burden)	Comments	
51 U.S. Geological Survey - Biological Resource Division - GCMRC - Power Revenues Under Cap Funded Projects																
52																
53 GOAL 1 - FOOD BASE																
54	BIO 1.R1.08	O	CRD	Aquatic Food Base (FY07--FY09)	489,917	513,630	58,575	455,055	114,845	3,000	3,010	6,200	112,000	-	216,000	Coop agmt mod of gross \$216K subject to the Univ of Wyoming est FY08 burden rate of 17%.
55	BIO 1.R4.08	N	CRD	Impacts of Various Flow Regimes on the Aquatic Food Base (FY08-FY09; Note 1)	-	72,700	11,608	61,093	40,000	-	1,500	19,593	-	-	-	Add'l funding added w/FY07 carry forward - Note 1 see table at end of spread sheet.
56	BIO 1.R3.08	C	CRD	Diet, Drift and Predation Data Analysis (FY07)	68,829	-	-	-	-	-	-	-	-	-	-	
57				SUB-TOTAL GOAL 1:	558,746	586,330	70,182	516,148	154,845	3,000	4,510	25,793	112,000	-	216,000	
58 GOAL 2 - NATIVE FISHES																
59	BIO 2.R1.08	O	CRD	LCR HBC Monitoring Lower 15km (HBC Population Est; Ongoing)	388,128	407,680	27,397	380,283	13,328	-	-	-	19,500	-	347,455	USFWS burden rate @ 17%
60	BIO 2.R2.08	O	CRD	LCR HBC Monitoring Lower 1,200m; Ongoing)	37,312	73,088	5,838	67,250	-	-	5,000	-	8,500	-	53,750	AGFD burden rate @ 31% on labor only
61	BIO 2.R3.08	O	CRD	HBC Monitoring Above Chute Falls; Ongoing)	73,504	79,652	6,745	72,907	2,857	-	-	-	15,000	-	55,050	USFWS burden rate @ 17%
62	BIO 2.R4.08	O	ORD	Monitoring Mainstem Fishes (includes Diamond Down; Ongoing)	388,452	518,436	52,551	465,885	63,618	4,967	14,000	12,000	92,700	-	278,600	AGFD burden rate @ 31% on labor only
63	BIO 2.R5.08	O	ORD	Nonnative Control Planning (FY07--FY10)	95,144	109,016	17,406	91,610	89,550	1,500	560	-	-	-	-	
64	BIO 2.R6.08	O	ORD	Nonnative Control Pilot Testing (FY07--FY10)	114,338	121,579	14,044	107,535	15,556	500	-	10,000	32,000	-	49,479	
65	BIO 2.R7.08	O	CRD	Stock Assessment of Native Fish in Grand Canyon (FY07--Ongoing)	35,360	41,392	6,609	34,783	34,033	750	-	-	-	-	-	
66	BIO 2.R8.08	O	CRD	Abundance Estimation Procedures (FY07--Ongoing)	35,360	41,392	6,609	34,783	34,033	750	-	-	-	-	-	
67	BIO 2.R9.08	O	CRD	Bioenergetics Modeling (FY07--FY10)	35,359	41,392	6,609	34,783	34,033	-	750	-	-	-	-	
68	BIO 2.R11.08	O	CRD	Native Fishes Habitat Data Analysis (FY07--FY10)	32,884	28,944	4,621	24,323	24,323	-	-	-	-	-	-	
69	BIO 2.R12.08	O	CRD	Trammel Net Effects (FY07--FY09)	37,789	38,458	2,208	36,250	-	-	-	-	-	-	36,250	
70	BIO 2.R13.08	O	CRD	Remote PIT Tag Reading (FY07--FY09)	61,536	34,624	2,547	32,077	-	-	-	-	4,600	-	27,477	
71	BIO 2.R14.08	O	CRD	Test Sonic Tags (FY07--FY09)	78,434	76,365	7,897	68,468	8,873	-	-	15,000	5,000	-	39,595	
72	BIO 2.R15.08	NA	CRD	Test DIDSON Camera (FY07--FY09)	13,151	-	-	-	-	-	-	-	-	-	-	
73				SUB-TOTAL GOAL 2:	1,426,751	1,612,019	161,082	1,450,937	320,204	8,467	20,310	37,000	177,300	-	887,656	
74 GOAL 3 - EXTIRPATED SPECIES																
75	07.3.00	NA	NA	None Identified	-	-	-	-	-	-	-	-	-	-	-	
76				SUB-TOTAL GOAL 3:	-	-	-	-	-	-	-	-	-	-	-	
77 GOAL 4 - RAINBOW TROUT																
78	BIO 4.M1.08	O	COR	Status & Trends of Lees Ferry Trout (Ongoing)	122,768	135,072	8,092	126,980	-	-	-	-	2,780	-	124,200	
79	BIO 4.E1.08	O	LTE	Monitoring Rainbow Trout Redds & Larvae (FY07)	48,029	-	-	-	-	-	-	-	-	-	-	
80				SUB-TOTAL GOAL 4:	170,797	135,072	8,092	126,980	-	-	-	-	2,780	-	124,200	
81 GOAL 5 - KANAB AMBERSNAIL																
82	BIO 5.R1.08	O	CRD	Monitor Kanab Ambersnail (FY95--FY10)	32,727	34,340	3,712	30,628	4,003	-	-	-	10,300	-	16,325	
83				SUB-TOTAL GOAL 5:	32,727	34,340	3,712	30,628	4,003	-	-	-	10,300	-	16,325	
84 GOAL 6 - SPRINGS / RIPARIAN																
85	BIO 6.R1.08	O	CRD	Vegetation Mapping (FY07--FY10)	112,496	108,785	17,369	91,416	74,416	3,000	1,000	-	13,000	-	-	
86	BIO 6.R2.08	O	COR	Vegetation Transects (FY07--FY10)	95,501	89,686	7,886	81,800	8,500	-	1,000	-	13,000	-	59,300	
87	BIO 6.R3.08	O	CRD	Vegetation Synthesis (FY07--FY10)	66,502	68,485	8,765	59,720	31,720	3,000	5,000	-	-	-	20,000	
88				SUB-TOTAL GOAL 6:	274,499	266,956	34,020	232,936	114,636	6,000	7,000	-	26,000	-	79,300	

APPENDIX C
Draft GCDAMP FY08 Budget for the USBR and the USGS GCMRC
03/13/2008 VERSION

	A	B	C	D	E	F	G	H	I	J	K	L	M	N	O	P
	GCMRC Project ID	Status	Funding Emphasis	Project Descriptions	Approved FY07 Budget (inc. CPI Increase)	Proposed FY08 Budget - Gross (inc. Burden) FINAL VERSION	DOI Customer Burden (Combined 6.09%, 19% and/or Other Rate)	Project Subtotal (w/o Burden)	GCMRC Personnel Costs (19% Burden)	GCMRC Project Related Travel / Training (19% Burden)	GCMRC Operations / Supplies (19% Burden)	GCMRC Equipment Purchase / Replacement (19% Burden)	AMP Logistics Support (19% Burden)	Outside GCMRC Contract & Science Labor (19% and/or Other Burden Rate)	Coop & Inter Agency Agmts (6.09% GCMRC Burden plus Cooperator's Burden)	Comments
50																
51	U.S. Geological Survey - Biological Resource Division - GCMRC - Power Revenues Under Cap Funded Projects															
89	GOAL 7 - QUALITY-OF-WATER															
90	BIO 7.R1.08	O	CRD	Water Quality Monitoring Lake - Powell & Tailwaters (Budget presented below; FY07--FY09)	-	-	-	-	-	-	-	-	-	-	-	-
91	PHY 7.M1.08	O	CRD	Integrated Quality-of-Water Monitoring (Downstream of GCD; FY07--Ongoing)	879,852	883,024	85,839	797,185	337,000	10,000	34,785	20,000	50,000	345,400	-	Suballocations to AZ WRD in the gross amt of \$316K is subject to the AZWRD est FY08 burden rate of 43.7%; the suballocation to UT WRD in the gross amt of \$29.4K is subject to the UT WRD est FY08 burden rate of 47%.
92	PHY 7.R1.08	O	CRD	Modeling Support Linked with Integrated Quality-of-Water Monitoring (FY07--FY08)	76,465	116,877	7,394	109,483	29,319	5,995	-	2,000	-	67,169	5,000	Coop agreement modification in the gross amt of \$5k is subject to the ASU est FY08 burden amt of 17.5% and the USGS/GCMRC "pass-through" burden of 6.09%. Suballocation to CA WRD in the gross amt of \$67,169 is subject to the CA WRD est FY08 burden rate of 70.6%.
93	SUB-TOTAL GOAL 7:				956,317	999,901	93,233	906,668	366,319	15,995	34,785	22,000	50,000	412,569	5,000	
94	GOAL 8 - SEDIMENT															
95	PHY 8.M1.08	N	COR	Longterm Monitoring of Changes in Sediment Storage	-	130,929	14,537	116,392	17,700	-	-	-	40,000	-	58,692	Project fully funded to \$194,323 by FY07 carry forward in spread sheet below.
96	SUB-TOTAL GOAL 8:				-	130,929	14,537	116,392	17,700	-	-	-	40,000	-	58,692	
97	GOAL 9 - RECREATIONAL EXPERIENCE															
98	REC 9.R1.08 / PHY 8.M2.08	O	CRD	Sand Bar and Campable Area Monitoring (FY07--FY11)	130,208	146,778	13,400	133,378	9,390	1,200	500	-	15,000	14,788	92,500	Addresses Goal 8 also.
99	REC 9.R2.08	C	CRD	Evaluate Campable Area Monitoring Results Using Measured Field Data vs. Remotely Sensed Data (FY07)	47,031	-	-	-	-	-	-	-	-	-	-	
100	REC 9.R3.08	O	CRD	Compile Campsite Inventory and GIS Atlas (FY07--FY08)	64,445	86,179	11,047	75,132	19,132	2,000	2,000	-	15,000	12,000	25,000	
101	REC 9.R4.08	D	CRD	Compile and Analyze Existing Safety Data	-	-	-	-	-	-	-	-	-	-	-	Deferred.
102	REC 9.R5.08	D	CRD	Evaluate Relation between Flows and Recreation Experience	-	-	-	-	-	-	-	-	-	-	-	Deferred.
103	SUB-TOTAL GOAL 9:				241,684	232,957	24,448	208,510	28,522	3,200	2,500	-	30,000	26,788	117,500	
104	GOAL 10 - HYDROPOWER															
105	HYD 10.M1.08	O	CRD	Monitor Power Generation and Market Values under Current and Future Dam Operations (FY07--Ongoing)	18,135	18,998	3,033	15,965	15,465	-	500	-	-	-	-	
106	SUB-TOTAL GOAL 10:				18,135	18,998	3,033	15,965	15,465	-	500	-	-	-	-	
107	GOAL 11 - CULTURAL															
108	CUL 11.R1.08	O	CRD	Research & Development toward Core Monitoring (FY07)	316,418	468,009	37,224	430,785	73,585	3,700	7,500	16,000	40,000	118,000	172,000	The sub-allocation of \$118K is subject to the FY08 estimated burden rate of the USGS Coastal & Marine Geology Team of 51%. The cooperative agreement amount to USU is subject to the estimated FY08 USU burden rate of 17%. The interagency agreement funding to NPS is subject to any burden charges allowed by NPS.
109	CUL 11.R2.08	N	CRD	Implement Tribal Monitoring Projects (See funding in BOR section)	-	-	-	-	-	-	-	-	-	-	-	
110	SUB-TOTAL GOAL 11:				316,418	468,009	37,224	430,785	73,585	3,700	7,500	16,000	40,000	118,000	172,000	

APPENDIX C
Draft GCDAMP FY08 Budget for the USBR and the USGS GCMRC
03/13/2008 VERSION

	A	B	C	D	E	F	G	H	I	J	K	L	M	N	O	P
50	GCMRC Project ID	Status	Funding Emphasis	Project Descriptions	Approved FY07 Budget (inc. CPI Increase)	Proposed FY08 Budget - Gross (inc. Burden) FINAL VERSION	DOI Customer Burden (Combined 6.09%, 19% and/or Other Rate)	Project Subtotal (w/o Burden)	GCMRC Personnel Costs (19% Burden)	GCMRC Project Related Travel / Training (19% Burden)	GCMRC Operations / Supplies (19% Burden)	GCMRC Equipment Purchase / Replacement (19% Burden)	AMP Logistics Support (19% Burden)	Outside GCMRC Contract & Science Labor (19% and/or Other Burden Rate)	Coop & Inter Agency Agmts (6.09% GCMRC Burden plus Cooperator's Burden)	Comments
51	U.S. Geological Survey - Biological Resource Division - GCMRC - Power Revenues Under Cap Funded Projects															
111	GOAL 12 - HIGH QUALITY MONITORING, RESEARCH & AEAM															
112	DASA 12.D1.08	O	CRD	Preparation for Monitoring Data Acquisition (Remote Sensing; FY07--Ongoing)	148,400	260,000	41,513	218,487		-	-	-	-	218,487	-	
113	DASA 12.D2.08	O	APM	Grand Canyon Integrated Oracle Database Management System (FY07--Ongoing)	171,445	178,607	26,083	152,524	97,590	1,500	2,000	-	-	29,000	22,434	
114	DASA 12.D3.08	O	APM	Library Operations (FY07--Ongoing)	48,745	42,635	6,807	35,828	29,628	-	6,200	-	-	-	-	
115	DASA 12.D4.08	O	APM	Legacy Analog Data Conversion (Analog to Digital - Reports & Imagery) (FY07--FY11)	104,463	78,736	12,029	66,707	56,232	-	5,475	-	-	-	5,000	
116	DASA 12.D5.08	O	APM	GIS Support for Integrated Analyses and Projects, GIS Lead (FY07--Ongoing)	231,704	227,515	25,477	202,038	91,721	3,000	7,317	-	-	-	100,000	
117	DASA 12.D6.08	O	CRD	Integrated Analysis and Modeling - Mapping Shoreline Habitat Changes (FY07--08)	84,199	115,888	6,652	109,236	-	-	-	-	-	-	109,236	
118	<i>Sub-total Goal 12 DASA Portion:</i>				<i>788,956</i>	<i>903,382</i>	<i>118,562</i>	<i>784,820</i>	<i>275,171</i>	<i>4,500</i>	<i>20,992</i>	<i>-</i>	<i>-</i>	<i>247,487</i>	<i>236,670</i>	
119	SUP 12.S1.08	O	APM	Logistics Base Costs (See BNELA for project related logistics costs; Ongoing)	135,252	126,691	20,228	106,463	86,360	-	-	20,103	-	-	-	
120	SUP 12.S2.08	O	APM	Survey Operations (Ongoing)	112,082	102,417	16,352	86,065	52,635	4,000	5,000	9,430	15,000	-	-	
121	SUP 12.S3.08	O	APM	Control Network (Ongoing)	126,496	134,823	19,574	115,249	69,394	3,000	2,085	770	22,000	-	18,000	
122	<i>Sub-total Goal 12 Support Portion:</i>				<i>373,831</i>	<i>363,931</i>	<i>56,154</i>	<i>307,777</i>	<i>208,389</i>	<i>7,000</i>	<i>7,085</i>	<i>30,303</i>	<i>37,000</i>	<i>-</i>	<i>18,000</i>	
123	PLAN 12.P1.08	N	CRD	Enhancing the Conceptual Ecosystem Model to Identify Critical Ecosystem Interactions and Data Gap (Science Advisor's conduct work in FY07; Funding in Independent Reviews, ADM 12.A4.07; FY07--FY08; Note 1)	-	-	-	-	-	-	-	-	-	-	-	Project added due to FY07 carry over funds availability. See Note1 and table at end of spreadsheet.
124	PLAN 12.P2.08	O	APM	AMP Effectiveness Workshop (FY07-FY08) Will not be conducted in FY08	46,800	-	-	-	-	-	-	-	-	-	-	FY07 funding carried over to FY08; FY08 funding (\$29,750) applied to other projects.
125	PLAN 12.P3.08	N	TBD	Low Steady Summer Flows - Data and Research Compilation, Synopsis and Synthesis (Note 1)	-	-	-	-	-	-	-	-	-	-	-	Project added due to FY07 carry over funds availability. See Note1 and table at end of spreadsheet.
126	<i>Sub-total Goal 12 Planning Portion:</i>				<i>46,800</i>	<i>-</i>	<i>-</i>	<i>-</i>	<i>-</i>	<i>-</i>	<i>-</i>	<i>-</i>	<i>-</i>	<i>-</i>	<i>-</i>	

APPENDIX C
Draft GCDAMP FY08 Budget for the USBR and the USGS GCMRC
03/13/2008 VERSION

	A	B	C	D	E	F	G	H	I	J	K	L	M	N	O	P
	GCMRC Project ID	Status	Funding Emphasis	Project Descriptions	Approved FY07 Budget (inc. CPI Increase)	Proposed FY08 Budget - Gross (inc. Burden) FINAL VERSION	DOI Customer Burden (Combined 6.09%, 19% and/or Other Rate)	Project Subtotal (w/o Burden)	GCMRC Personnel Costs (19% Burden)	GCMRC Project Related Travel / Training (19% Burden)	GCMRC Operations / Supplies (19% Burden)	GCMRC Equipment Purchase / Replacement (19% Burden)	AMP Logistics Support (19% Burden)	Outside GCMRC Contract & Science Labor (19% and/or Other Burden Rate)	Coop & Inter Agency Agmts (6.09% GCMRC Burden plus Cooperator's Burden)	Comments
50																
51	U.S. Geological Survey - Biological Resource Division - GCMRC - Power Revenues Under Cap Funded Projects															
127	ADM 12.A1.08	O	APM	Administrative Operations (Ongoing)	790,942	228,363	31,688	196,675	40,000	5,000	102,675	5,000	-	-	44,000	Reduced funding by moving Secretary and Budget Analyst salaries and Flagstaff Science Center overhead costs to the SBSC overhead account per USGS direction. Made additional cuts decreasing training, travel, supply and equipment expenses.
128	ADM 12.A2.08	O	APM	Program Planning & Management (Ongoing)	908,802	1,059,438	169,154	890,284	840,904	38,880	10,500	-	-	-	-	In FY08 moved DASA Program Mgr salary from DASA projects to this project to correctly reflect his position in planning. This project also maintains the Physical Program Managers' position (not filled as of 6/1/07) as well as the GCMRC Chief, Deputy Chief, Sociocultural, Biology and Survey/Logistics Program Managers' salaries, and travel and training budgets.
129	ADM 12.A3.08	O	APM	AMWG/TWG Meeting Travel Funds (Ongoing)	17,550	18,077	2,886	15,191	-	15,191	-	-	-	-	-	
130	ADM 12.A4.08	O	APM	Independent Reviews and 2008 Science Symposium	275,514	90,301	14,418	75,883	20,883	12,000	4,800	-	12,000	26,200	-	
131	ADM 12.A4.08	O	APM	Executive Director of Science Advisors Review and Coordination; includes Science Advisors' Expenses (Ongoing)	-	214,200	34,200	180,000	-	5,000	-	-	-	175,000	-	
132	ADM 12.A6.08	NA	APM	2008 Science Symposium (Intermittent)	-	29,750	4,750	25,000	-	11,000	4,000	-	-	10,000	-	Science Symposium will be held in the fall (October or later) of 2008 and the venue will be expanded.
133	ADM 12.A5.08	O	APM	GCMRC Component of SBSC Sys Admin Support (FY05-Ongoing)	320,438	202,300	32,300	170,000	-	-	65,000	100,000	-	5,000	-	SBSC covering salaries and travel costs for IT support in basic overhead costs.
134	<i>Sub-total Goal 12 Administrative/Management Portion:</i>				2,313,247	1,842,429	289,396	1,553,033	901,787	87,071	186,975	105,000	12,000	216,200	44,000	
135	SUB-TOTAL GOAL 12:				3,522,834	3,109,742	464,111	2,645,630	1,385,347	98,571	215,052	135,303	49,000	463,687	298,670	
136	GCMRC Power Revenues Under Cap Projects Sub-totals:				7,518,907	7,595,253	913,675	6,681,578	2,480,626	138,933	292,157	236,096	537,380	1,021,044	1,975,343	
137	GCMRC Power Revenue Funded Projects (NOT Capped) and Other Funded Projects															
139	BIO 7.R1.08	O	CRD	Water Quality Monitoring - Lake Powell & Tailwaters (FY07-09)	220,632	212,631	33,949	178,682	140,962	10,720	22,500	4,500	-	-	-	
140	BIO TBD	O	CRD	Environmental Research Agrmt (Temperature Control Device-TCD)	125,000	-	-	-	-	-	-	-	-	-	-	
141	GCMRC Other Agreements Projects Subtotals:				345,632	212,631	33,949	178,682	140,962	10,720	22,500	4,500	-	-	-	
142																
143	GCMRC TOTAL AMP FY2008 PLANNED PROGRAM COSTS:				7,864,539	7,807,884	947,624	6,860,260	2,621,587	149,653	314,657	240,596	537,380	1,021,044	1,975,343	
144																
145	NPS GCDAMP Project Funding Contributions															
146	NA	NA	NA	NPS Treatment Plan Support of Archaeological Resources in FY08 (10 sites)	-	261,180	-	-	-	-	-	-	-	-	-	
147	NA	NA	NA	NPS Cost Share Contributions to Monitoring of Archaeological Resources in FY08 (141 sites)	-	141,410	-	-	-	-	-	-	-	-	-	
148																
149	Other Agencies Projects Subtotals:				-	402,590	-	-	-	-	-	-	-	-	-	
150																
151	ALL AGENCIES TOTAL AMP FY2008 PLANNED PROGRAM COSTS:				7,864,539	8,210,474	947,624	6,860,260	2,621,587	149,653	314,657	240,596	537,380	1,021,044	1,975,343	

APPENDIX C
Draft GCDAMP FY08 Budget for the USBR and the USGS GCMRC
03/13/2008 VERSION

	A	B	C	D	E	F	G	H	I	J	K	L	M	N	O	P		
	GCMRC Project ID	Status	Funding Emphasis	Project Descriptions	Approved FY07 Budget (inc. CPI Increase)	Proposed FY08 Budget - Gross (inc. Burden) FINAL VERSION	DOI Customer Burden (Combined 6.09%, 19% and/or Other Rate)	Project Subtotal (w/o Burden)	GCMRC Personnel Costs (19% Burden)	GCMRC Project Related Travel / Training (19% Burden)	GCMRC Operations / Supplies (19% Burden)	GCMRC Equipment Purchase / Replacement (19% Burden)	AMP Logistics Support (19% Burden)	Outside GCMRC Contract & Science Labor (19% and/or Other Burden Rate)	Coop & Inter Agency Agmts (6.09% GCMRC Burden plus Cooperator's Burden)	Comments		
50																		
51	U.S. Geological Survey - Biological Resource Division - GCMRC - Power Revenues Under Cap Funded Projects																	
152																		
153	PROGRAM COSTS:	BOR Power Revenues Under Cap Program Costs:				FISCAL YEAR 2007	FISCAL YEAR 2008	PROGRAM FUNDING:	BOR Power Revenues Under Cap Program Funding:				FISCAL YEAR 2007	FISCAL YEAR 2008				
154		BOR Power Revenues Under Cap Program Costs (gross)				1,725,222	1,824,747		BOR Power Revenues Under Cap Program Funding:				1,725,222	1,824,747				
155		GCMRC Power Revenues Under Cap Program Costs (gross)				7,438,133	7,595,253		GCMRC Power Revenues Under Cap Program Funding:				7,438,133	7,595,253				
156		Subtotal BOR & GCMRC Power Revenue Under Cap Program Costs				9,163,355	9,420,000		Subtotal BOR & GCMRC Power Revenue Under Cap Funding:				9,163,355	9,420,000				
157																		
158		BOR Appropriated and Other Program Costs :				FISCAL YEAR 2007	FISCAL YEAR 2008		BOR Appropriated and Other Program Funding:				FISCAL YEAR 2007	FISCAL YEAR 2008				
159		BOR Appropriated and Other Program Costs (gross)				475,000	475,000		BOR Tribal Participation Funding:				95,000	95,000				
160		GCMRC Appropriated and Other Program Costs (gross)				345,632	212,631		BOR TCD Experimental Actions Funding:				125,000	0				
161		Subtotal BOR & GCMRC Power Revenue (Non-Capped) and Other Funded Program Costs				820,632	687,631		BOR to GCMRC Lake Powell IWQP (Ops and Maintenance)				220,632	212,631				
162		Subtotal BOR Appropriations and Other Program Funding:				440,632	307,631											
163																		
164	USGS Appropriated Program Costs:				FISCAL YEAR 2007	FISCAL YEAR 2008	USGS Appropriated Program Funding:				FISCAL YEAR 2007	FISCAL YEAR 2008						
165	USGS-GCMRC Estimated Cost Share Expense Required by USGS Policy				1,000,000	1,000,000	USGS Tribal Participation Funding:				95,000	95,000						
166	Subtotal USGS Appropriated Program Costs:				1,000,000	1,000,000	USGS-GCMRC Estimated Cost Share Funding (Per USGS Policy):				1,000,000	1,000,000						
167	Subtotal BOR & GCMRC Power Revenue Under Cap Funding:				1,095,000	1,095,000												
168																		

APPENDIX C
Draft GCDAMP FY08 Budget for the USBR and the USGS GCMRC
03/13/2008 VERSION

	A	B	C	D	E	F	G	H	I	J	K	L	M	N	O	P
	GCMRC Project ID	Status	Funding Emphasis	Project Descriptions	Approved FY07 Budget (inc. CPI Increase)	Proposed FY08 Budget - Gross (inc. Burden) FINAL VERSION	DOI Customer Burden (Combined 6.09%, 19% and/or Other Rate)	Project Subtotal (w/o Burden)	GCMRC Personnel Costs (19% Burden)	GCMRC Project Related Travel / Training (19% Burden)	GCMRC Operations / Supplies (19% Burden)	GCMRC Equipment Purchase / Replacement (19% Burden)	AMP Logistics Support (19% Burden)	Outside GCMRC Contract & Science Labor (19% and/or Other Burden Rate)	Coop & Inter Agency Agmts (6.09% GCMRC Burden plus Cooperator's Burden)	Comments
50																
51	U.S. Geological Survey - Biological Resource Division - GCMRC - Power Revenues Under Cap Funded Projects															
169	Other Agency Appropriated Program Costs:				FISCAL YEAR 2007	FISCAL YEAR 2008	Other Agency Appropriated Funding:				FISCAL YEAR 2007	FISCAL YEAR 2008				
170	NPS/GRCA GCDAMP PROGRAM CONTRIBUTIONS:					402,590	NPS Contribution to Treatment of Archaeological Resources in FY08				0	261,180				
171	Subtotal Other Agency Appropriated Program Costs:					402,590	NPS Cost Share Proposal Funding:				0	141,410				
172							NPS Tribal Participation Funding:				95,000	95,000				
173	FY 2008 GCMRC PROGRAM NEEDS UNMET BY FY2008 FUNDING - FY 2007 CARRY FORWARD NEEDS FROM GCDAMP POWER REVENUES UNDER CAP						FWS Tribal Participation Funding:				95,000	95,000				
174	Refer to Line ### for complete budget details on the projects, below:				FISCAL YEAR 2007	FISCAL YEAR 2008	BIA Tribal Participation Funding:				95,000	95,000				
175	TBD	TBD	TBD	FY07 and FY08: LTEP - EIS Program Work	100,000	65,000	Subtotal Other Agency Appropriated Funding:				285,000	687,590				
176	NA	NA	NA	FY07: Editorial and Technical Writer Support; FY08: CPI Reduction, GCMRC Portion (FY07 CPI reduction of 81K was handled by GCMRC through savings found in FY07 work plan expenditures.)	24,000	16,374										
177	BIO 1.R4.08	N	CRD	FY07: Purchase Sonic Tags / FY08: Impacts of Various Flow Regimes on the Aquatic Food Base (FY08-FY09; Note 1)	20,000	17,500	USGS-GCMRC FY2006/2007 Carry Forward Funding Available for Use (Power Revenues Under Cap):				FISCAL YEAR 2007	FISCAL YEAR 2008				
178	PHY 8.M1.08	N	COR	FY07: Staff Support for Evaluation of Proposal for this project / FY08: Longterm Monitoring of Changes in Sediment Storage (Completion of Funding by FY07 Carry Forward Funds)	23,000	63,394	AMP Effectiveness Workshop Funds carried forward per TWG/AMWG directions				-	46,800				
179	PLAN 12.P1.08	N	CRD	FY07: Facilitation Assistance / FY08: Enhancing the Conceptual Ecosystem Model to Identify Critical Ecosystem Interactions and Data Gap (Science Advisor's conduct work in FY07; Funding in Independent Reviews, ADM 12.A4.07; FY07-FY08; Note 1)	6,000	100,000	Funds saved in Administration in FY07 to assist w/FY08 budget (refer to previous budget version)				-	16,451				
180	PLAN 12.P3.08	N	TBD	FY07: Funded Lees Ferry Trout Study / FY08: Low Steady Summer Flows - Data and Research Compilation, Synopsis and Synthesis (Note 1)	63,000	100,000	Funds carried forward in Admin, misc in FY07				-	16,517				
181	Subtotal FY08 GCMRC Program COSTS Met by GCDAMP FY07 Carry Forward:				236,000	362,268	Funds carried forward from FY07 (refer to Note 1 - Motion 5 at August 2007 AMWG meeting)				-	217,500				
182							LTEP-EIS Funds brought forward from FY06/07				100,000	65,000				
183	PROGRAM COSTS:						PROGRAM FUNDING:									
184							Funds carried forward from misc savings in FY06				136,000	-				
185							Subtotal Other Agency Appropriated Funding:				236,000	362,268				
186					FISCAL YEAR 2007	FISCAL YEAR 2008	FISCAL YEAR 2007				FISCAL YEAR 2008					
187	TOTAL GCDAMP PROGRAM COSTS:				11,219,987	11,872,488	TOTAL GCDAMP PROGRAM FUNDING:				11,219,987	11,872,489				
188																
189																
190																
191																
192																

FUNDING SUMMARY	FY2007	FY2008
DIFFERENCE BETWEEN COSTS AND FUNDING	1	1

APPENDIX C
Draft GCDAMP FY08 Budget for the USBR and the USGS GCMRC
03/13/2008 VERSION

	A	B	C	D	E	F	G	H	I	J	K	L	M	N	O	P
	GCMRC Project ID	Status	Funding Emphasis	Project Descriptions	Approved FY07 Budget (inc. CPI Increase)	Proposed FY08 Budget - Gross (inc. Burden) FINAL VERSION	DOI Customer Burden (Combined 6.09%, 19% and/or Other Rate)	Project Subtotal (w/o Burden)	GCMRC Personnel Costs (19% Burden)	GCMRC Project Related Travel / Training (19% Burden)	GCMRC Operations / Supplies (19% Burden)	GCMRC Equipment Purchase / Replacement (19% Burden)	AMP Logistics Support (19% Burden)	Outside GCMRC Contract & Science Labor (19% and/or Other Burden Rate)	Coop & Inter Agency Agmts (6.09% GCMRC Burden plus Cooperator's Burden)	Comments
50																
51	U.S. Geological Survey - Biological Resource Division - GCMRC - Power Revenues Under Cap Funded Projects															
193	DETAIL BUDGET FOR FY 2008 PROJECTS FUNDED BY FY 2007 CARRY FORWARD FUNDS -															
194	TBD	TBD	TBD	LTEP - EIS Program Work	35,000	65,000	-	-	-	-	-	-	-	-	-	Funds in FY07 sub-allocated for work to be conducted by SA Wright of the CA Water Resources Discipline (35k); FY08 funding to apply to various LTEP-EIS objectives.
195	NA	NA	NA	USGS-GCMRC Portion of CPI Reduction for FY2008	-	16,374	-	-	-	-	-	-	-	-	-	Total CPI Reduction: 3% Estimated; 2.8% Actual; Difference of 0.2%. Difference between \$9,420,000 allowed by BOR and \$9,438,256 used as planning for budget purposes is \$18,256. Portions negotiated are \$1,882 BOR GCDAMP and \$16,376 USGS/GCMRC GCDAMP.
196	BIO 1.R4.08	N	CRD	Impacts of Various Flow Regimes on the Aquatic Food Base (FY08-FY09; Note 1)	-	17,500	-	-	-	-	-	-	-	-	-	Add'l funding added w/FY07 carry forward; Note 1 - Motion 5.
197	PHY 8.M1.08	N	COR	Longterm Monitoring of Changes in Sediment Storage (Completion of Funding by FY07 Carry Forward Funds)	-	63,394	3,639	59,755	-	-	-	-	-	-	59,755	Project fully funded to \$194,323 by FY07 carry forward; See line 95 where partial funding takes place with FY08 funds.
198	PLAN 12.P1.08	N	CRD	Enhancing the Conceptual Ecosystem Model to Identify Critical Ecosystem Interactions and Data Gap (Science Advisor's conduct work in FY07; Funding in Independent Reviews, ADM 12.A4.07; FY07--FY08; Note 1)	-	100,000	11,825	88,175	-	-	-	-	-	50,000	38,175	Project added due to FY07 carry over funds availability; Note 1 - Motion 5.
199	PLAN 12.P3.08	N	TBD	Low Steady Summer Flows - Data and Research Compilation, Synopsis and Synthesis (Note 1)	-	100,000	13,797	86,203	34,700	30,000	1,503	-	-	-	20,000	Project added due to FY07 carry over funds availability; Note 1 - Motion 5.
200																
201	SUB-TOTAL FY07 CARRY FORWARD FUNDSNEEDED:				NA	362,268	29,261	234,133	34,700	30,000	1,503	-	-	50,000	117,930	
202	<div style="border: 1px solid black; padding: 5px;"> <p>NOTE 1: Per motion Number 5, passed at the Adaptive Management Work Group meeting held in Flagstaff August 29-30, 2007, Project PLAN 12.P1.08, entitled "Enhancing the Conceptual Ecosystem Model to Identify Critical Ecosystem Interactions and Data Gap" (Line 123) and Project PLAN 12.P3.08, entitled "Low Steady Summer Flows - Data and Research Compilation, Synopsis and Synthesis" (Line 125) were both added for a total of \$100K each, to be funded by USGS-GCMRC FY2007 carry forward funds. In addition, per this motion, an additional amount of funding in existing Project BIO 1.R4.08, "Impacts of Various Flow Regimes on the Aquatic Food Base" (Line 55) was added from USGS-GCMRC FY2007 carry forward funds to increase the total project funding from \$72K to \$89.5K. The total amount of FY2007 carry forward funding from the previous year under this agreement totals \$280,235 (Line 167) and is added to the FY2008 funding table.</p> </div>															
203																
204																
205																
206																
207																
208																
209																
210																
211																
212																
213																
214																
215																
216																
217																
218																
219																
220																

Explanation of information found in columns of Fiscal Year 2008 Glen Canyon Dam Adaptive Management Program (GCDAMP) Budget.		
Column	Title	Key
A	GCMRC Project ID	Col 1-3 Program Area BIO: Biology PHY: Physical Science REC: Recreation HYD: Hydropower CUL: Cultural DASA: Data Acquisition, Storage and Analysis SUP: Support PLA: Planning ADM: Administration Col 4-5 GCDAMP Goal Number Col 6-7 Project Number Col 8-9 Fiscal Year
B	Status	O: Ongoing N: New C: Complete
C	Category	APM: Admin & Program Mgmt COR: Core Monitoring CRD: Core Monitoring Research & Development ORD: Ongoing Research & Development LTE: Long-Term Experiment NA: Not Applicable
D	Project Description	Project Title (Start Date -- End Date)
F	Est FY08 Costs	Estimated FY08 Costs of Projects

