



## **GEOLOGIC MAP OF THE RIVERSIDE EAST 7.5' QUADRANGLE, RIVERSIDE COUNTY, CALIFORNIA**

By Douglas M. Morton<sup>1</sup> and Brett F. Cox<sup>2</sup>

*Digital preparation by Michael Dawson<sup>1</sup> and Timothy O'Brien<sup>3</sup>*

*Prepared in cooperation with*  
CALIFORNIA DIVISION OF MINES AND GEOLOGY  
*and*  
U.S. AIR FORCE

Open-File Report OF 01-452

2001

This report is preliminary and has not been reviewed for conformity with U.S. Geological Survey editorial standards or with the North American Stratigraphic Code. Any use of trade, firm, or product names in this publication is for descriptive purposes only and does not imply endorsement by the U.S. Government.

U.S. DEPARTMENT OF INTERIOR  
U.S. GEOLOGICAL SURVEY

<sup>1</sup> U.S. Geological Survey, Department of Earth Sciences,  
University of California, Riverside CA 92521

<sup>2</sup> U.S. Geological Survey  
345 Middlefield Rd.  
Menlo Park, CA 94025

<sup>3</sup> Department of Earth Sciences  
University of California  
Riverside, CA 92521

## TABLE OF CONTENTS

- Introduction
  - General
- How to obtain paper plots
- Database contents
  - Data package
  - Plot package
  - Symbols package
  - Other files
- Software utilities
- How to obtain the digital files
  - Digital databases
  - Postscript plot files
  - Portable Document Format (.pdf) files
- How to extract the geologic map database from the tar files
  - Digital database
  - Postscript plot files
- How to convert the ARC/INFO interchange (export) files
- Digital geologic map specifications
  - Digital compilation
  - Base map
  - Spatial resolution
  - Map accuracy standards
  - Faults and landslides
  - Database specifics
    - General
    - Lines
    - Polygons
    - Points
- References

## INTRODUCTION

### General

Open-File Report 01-452 contains a digital geologic map database of the Riverside East 7.5' quadrangle, Riverside County, California that includes:

1. ARC/INFO (Environmental Systems Research Institute, <http://www.esri.com>) version 7.2.1 coverages of the various elements of the geologic map.
2. A Postscript file to plot the geologic map on a topographic base, containing a Correlation of Map Units diagram (CMU), a Description of Map Units (DMU), and an index map.
3. Portable Document Format (.pdf) files of:
  - a. This Readme; includes in Appendix I, data contained in rse\_met.txt
  - b. The same graphic as plotted in 2 above. Test plots have not produced 1:24,000-scale map sheets. Adobe Acrobat page size setting influences map scale.

The Correlation of Map Units and Description of Map Units is in the editorial format of USGS Miscellaneous Investigations Series (I-series) maps but has not been edited to comply with I-map standards. Within the geologic map data package, map units are identified by standard geologic map criteria such as formation-name, age, and lithology. Where known, grain size is indicated on the map by a subscripted letter or letters following the unit symbols as follows: lg, large boulders; b, boulder; g, gravel;

a, arenaceous; s, silt; c, clay; e.g. Qyf<sub>a</sub> is a predominantly young alluvial fan deposit that is arenaceous. Multiple letters are used for more specific identification or for mixed units, e.g., Qfy<sub>sa</sub> is a silty sand. In some cases, mixed units are indicated by a compound symbol; e.g., Qyf<sub>2sc</sub>. Marine deposits are in part overlain by local, mostly alluvial fan, deposits and are labeled Qomf. Grain size follows f.

Even though this is an Open-File Report and includes the standard USGS Open-File disclaimer, the report closely adheres to the stratigraphic nomenclature of the U.S. Geological Survey. Descriptions of units can be obtained by viewing or plotting the .pdf file (3b above) or plotting the postscript file (2 above).

This Readme file describes the digital data, such as types and general contents of files making up the database, and includes information on how to extract and plot the map and accompanying graphic file. Metadata information can be accessed at <http://geo-nsdi.er.usgs.gov/metadata/open-file/01-452> and is included in Appendix I of this Readme.

## HOW TO OBTAIN PAPER PLOTS

For those having access to large-format plotters such as HP650C, HP755C, and HP2500C, plots may be made directly from the included plot file.

## DATABASE CONTENTS

The files constituting the geologic map database of this Open-File Report are listed below along with the interchange files from which they were extracted.

### Data Package

All files listed below are in a compressed tar file named rse.tar.gz (2 Mb); see section below titled, SOFTWARE UTILITIES.

<u>ARC/INFO interchange files</u>	<u>Riverside East coverages</u>	<u>Contains</u>
rse_geo.e00	rse_geo	Contacts, faults, geologic unit labels
rse_ano.e00	rse_ano	Annotation subclasses: GEO (for plotting unit labels)
rse_str.e00	rse_str	Attitudes and their dip values. Dip values plotted as annotation.

The directory, info/, is produced in the process of importing interchange files to ARC coverages in ARC/INFO. The rse (Riverside East) info/ directory contains:

### Feature Attribute Tables

Polygon attribute table	rse_geo.pat
Arc attribute table	rse_geo.aat
	rse_ano.aat
Point attribute table	rse_str.pat

<u>Raster file</u>	<u>Resultant image</u>	<u>Contains</u>
rse.tif	Riverside East base map	Topographic base from 500 dpi scan of USGS Riverside East 7.5' quadrangle, 1967

### **Plot Package**

Postscript plot files of the geologic map and explanation; please see section below titled, SOFTWARE UTILITIES for additional information.

<u>Compressed file</u>	<u>Resultant image</u>	<u>Contains</u>
rse_map.ps.gz	rse_map.ps	PostScript plot file of geologic map and CMU/DMU

The Postscript file is compressed using winzip.

The uncompressed PostScript file rse\_map.ps will plot a 1:24,000 scale, full color geologic map of the Riverside East quadrangle on the topographic base. A detailed CMU diagram, a DMU are included on the sheet. The sheet is in the editorial format of the U.S. Geological Survey's Miscellaneous Investigations (I) map series, and is approximately 46 X 32 inches in size. The map sheet has been successfully plotted on Hewlett-Packard large-format plotters, models HP650C, HP755C, and HP2500C.

### **Symbols Package**

Files in the plot package have been prepared to produce optimum plots using the shade and marker sets and fonts listed below; these symbol sets and fonts are included in a compressed tar file named symbols.tar.gz (0.04 Mb); see section below titled SOFTWARE UTILITIES.

geoSCAMP2.lin	Lineset
geoSCAMP2.mrk	Markerset for points
alc1.shd	Colors
geology2.shd	Pattern fills
fnt026	Font required for geoSCAMP2.lin
fnt037	Font required for geoSCAMP2.mrk
fnt035	Font required for geology2.shd

Special geologic characters used in unit designations are from the Geoage font group and may be obtained at the following web site:

Server:	onyx.wr.usgs.gov
UserID:	anonymous
Password:	Your e-mail address
Directory:	pub/wpg/supplies/geoage

### **Other files**

README.pdf	This document
rse_map.pdf	Postscript plot file of geologic map and CMU/DMU

## **SOFTWARE UTILITIES**

Files which have .gz file extension were compressed using gzip. Gzip utilities are available free of charge via the Internet at the gzip home page, <http://www.gzip.org>. Files with a .zip file extension were compressed using WinZip, available at <http://www.winzip.com>.

The data package and symbols package are additionally bundled into a single tar (tape archive) file. The individual files must be extracted using a tar utility, available free of charge via the Internet through links on the Common Internet File Formats page, <http://www.matisse.net/files/format.html>. One such utility is WinZip, available at <http://www.winzip.com>.

## HOW TO OBTAIN THE DIGITAL FILES

The export files, and subsequently the data and plot files, constituting the geologic map database of this Open-File Map may be obtained in two ways, both over the Internet.

1. The files can be obtained via the Web from Western Region Geologic Information Server. Go to the web page at <http://geopubs.wr.usgs.gov/open-file/of01-452> and follow the directions to download the files.
2. The files can also be obtained by anonymous ftp over the Internet from [wrgis.wr.usgs.gov](http://wrgis.wr.usgs.gov). The files are located in the directory /pub/open-file/. Be sure to use binary transfer mode or ASCII mode for individual .e00 (ARC interchange file format) files.

## HOW TO EXTRACT THE GEOLOGIC MAP DATABASE FROM THE TAR FILE

### Digital database

After downloading the files, they must be uncompressed using a gzip utility such as gzip itself or WinZip. The data files must then be extracted using a tar utility or Winzip.

This process will create a directory, rse/, that will contain the ARC/INFO interchange files and supporting files. The directory should contain the following files:

```
rse/  
  rse_geo.e00  
  rse_str.e00  
  rse_ano.e00  
  
  rse.tif
```

The symbols.tar.gz file is imported using the same methods as for the lkm.tar.gz file. It will create a directory, symbols/ that will contain the following files:

```
geoSCAMP2.lin  
geoSCAMP2.mrk  
alc1.shd  
geology2.shd  
fnt026  
fnt037  
fnt035
```

The following are not included in the database tar file, and are downloaded separately.

```
rse_map.ps.gz  
Readme.pdf  
rse_map.pdf
```

### **Postscript plot files**

Make a 47.3 MB uncompressed file, rse\_map.ps (plot of complete map), by typing gzip -d rse\_map.ps.gz (or use gzip utility of choice).

### **Portable Document Format (.pdf) files**

PDF files are not stored as gzip files. They are accessed using Adobe Acrobat Reader software, available free from the Adobe website <http://www.adobe.com>. Follow instructions at the website to download and install the software. Acrobat Reader contains an on-line manual and tutorial.

## **HOW TO CONVERT THE ARC/INFO INTERCHANGE (EXPORT) FILES**

The ARC interchange (.e00) files are converted to ARC coverages using the ARC command IMPORT.

ARC interchange files can also be read by some other Geographic Information Systems, including ArcView (ESRI) and MapInfo (<http://www.mapinfo.com>), (Environmental Systems Research Institute, Inc., 1998). Please consult your GIS documentation to see if you can use ARC interchange files and the procedure to import them.

## **DIGITAL GEOLOGIC MAP SPECIFICATIONS**

### **Digital compilation**

The geologic map information was hand digitized from a base-stable original (ink on a greenline) of the geologic map at 1:24,000 scale. Digital tics were placed by hand at latitude/longitude intersections. The lines, points, and polygons were edited using standard ARC/INFO commands, and in some places, interactively by hand using graphical user interface ALACARTE (Fitzgibbon, 1991, Fitzgibbon and Wentworth, 1991, Wentworth and Fitzgibbon, 1991). Digitization and editing artifacts significant enough to display at a scale of 1:24,000 were corrected.

### **Base map**

The base map image (rse.tif) was prepared by scanning a scale-stable clear film of the U.S. Geological Survey, 1:24,000 Riverside East 7.5' quadrangle (1967) topographic map. Scanning was done using an Anatech Eagle 4080 monochrome 800 dpi scanner; at a resolution of 500 dpi. The raster scan was converted to a monochromatic image in ARC/INFO, and registered and rectified to the Riverside East 7.5' quadrangle. No elements of the base layer are attributed. The base map is provided for reference only.

### **Spatial resolution**

Use of this digital geologic map database should not violate the spatial resolution of the data. Although the digital form of the data removes the constraint imposed by the scale of a paper map, the detail and accuracy inherent in map scale are also present in the digital data. The fact that this database was edited at a scale of 1:24,000 means that higher resolution information is not generally present in the dataset. Plotting at scales larger than 1:24,000 will not yield greater *real* detail, although it may reveal fine-scale irregularities above the intended resolution of the database. Similarly, although higher resolution data is incorporated at a few places, the resolution of the combined output will be limited by the lower resolution data.

## Map accuracy standards

Until uniform National geologic map standards are developed and adopted, lines and points on SCAMP 1:24,000 scale geologic maps that are located to within 15 meters, relative to accurately located features on the base map, are considered to meet map accuracy standards. Dashed lines, indicated in the database as approximately located or inferred, are generally located within 30 meters, relative to accurately located features on the base map.

## Faults and landslides

This database is sufficiently detailed to identify and characterize many actual and potential geologic hazards represented by faults and landslides, but it is not sufficiently detailed for site-specific determinations. Faults shown do not take the place of fault rupture hazard zones designated by the California State Geologist (see Hart, 1998).

## Database specifics

**General**--The map database consists of ARC/INFO format coverages which are stored in polyconic projection (Table 1), and a series of data tables. Digital tics define a 2.5 minute grid of latitude and longitude in the geologic coverages corresponding to the 2.5 minute tic grid on the topographic base map.

Table 1 --- Map Projection

Projection	Polyconic
Datum	NAD27
Zunits	No
Units	Meters
Spheroid	Clark 1866
X shift	0.000000000
Y shift	0.000000000
Parameters	-117 18 45.000 longitude of central meridian
	33 52 30.00 latitude of projections origin
	0.00000 false easting (meters)
	0.00000 false northing (meters)

The content of the geologic database can be described in terms of feature classes that include lines, points, and areas that compose the map. See the metadata text file (Appendix I) for detailed descriptions.

**Lines** – Lines are recorded as strings of arcs and are described in an arc attribute (.aat) table. Complete lists of the line types (LTYPE) used in the quadrangle are available in Appendix I. They represent contacts and faults, which define the boundaries of map units and map boundaries.

**Polygons** --- Geologic map units (polygons) are described in the polygon attribute (.pat) table (details in Appendix I). For traditional descriptions of the map units, see the Portable Document Format file rse\_map.pdf or the Postscript map plot, rse\_map.ps. A list of all map units in the database is given in Appendix I.

**Points** – Point information (attitudes of planar and linear features) is recorded as coordinate and related information. Complete lists of the point types (PTTYPE) used in the point coverage are available in Appendix I.

## REFERENCES

- Environmental Systems Research Institute, Inc, 1991, ARC/INFO command references 6.0: Proprietary software manual
- Fitzgibbon, T.T., 1991, ALACARTE installation and system manual (version 1.0): U.S. Geological Survey, Open-File Report 91-587B
- Fitzgibbon, T.T., and Wentworth, C.M., 1991, ALACARTE user interface – AML code and demonstration Maps (version 1.0): U.S. Geological Survey, Open-File Report 91-587A
- Wentworth, C.M., and Fitzgibbon, T.T., 1991, ALACARTE user manual (version 1.0): U.S. Geological Survey Open-File Report 91-587C

## Appendix I

(Original metadata text)

Identification\_Information:

Citation:

Citation\_Information:

Originator: Douglas M. Morton

Originator: Brett F. Cox

Publication\_Date: 2001

Title: Geologic Map of the Riverside East 7.5' Quadrangle, Riverside County, California

Edition: (Version 1.0, October 03, 2001)

Geospatial\_Data\_Presentation\_Form: vector digital data

Series\_Information:

Series\_Name: U.S. Geological Survey Open-File Report

Issue\_Identification: USGS OF 01-452

Publication\_Information:

Publication\_Place: Menlo Park, California

Publisher: U.S. Geological Survey

Online\_Linkage: <http://geopubs.wr.usgs.gov/open-file/of01-452>

Description:

Abstract:

This data set maps and describes the geology of the Riverside East 7.5' quadrangle, Riverside County, California. Created using Environmental Systems Research Institute's ARC/INFO software, the data base consists of the following items: (1) a map coverage containing geologic contacts and units, (2) a coverage containing structural data, (3) a coverage containing geologic unit annotation and leaders, and (4) attribute tables for geologic units (polygons), contacts (arcs), and site-specific data (points). In addition, the data set includes the following graphic and text products: (1) a postscript graphic plot-file containing the geologic map, topography, cultural data, a Correlation of Map Units (CMU) diagram, a Description of Map Units (DMU), and a key for point and line symbols, and (2) PDF files of the Readme (including the metadata file as an appendix), and the graphic produced by the Postscript plot file.

The Riverside East quadrangle is located in the northern part of the Peninsular Ranges Province within the central part of the Perris block, a relatively stable, rectangular in plan area located between the Elsinore and San Jacinto fault zones.

The quadrangle is underlain predominantly by Cretaceous plutonic rocks which are part of the composite Peninsular Ranges batholith. Within the quadrangle, the batholithic rocks represent a wide variety of mafic to intermediate composition granitic rocks ranging in composition from monzogranite to gabbro, but tonalite predominates. Most of the granitic rocks are faintly to intensely foliated. Many are heterogeneous and contain varying amounts of meso- and melanocratic discoidal-shaped inclusions. Some



rock is composed almost wholly of inclusion material and some are migmatitic. Included within these granitic rocks are a few septa of Paleozoic(?) biotite schist, marble, and calcsilicate rock of upper amphibolite metamorphic grade.

Metamorphic rocks of Paleozoic(?) age occur primarily in the northwest part of the quadrangle. These rocks include coarse-grained marble bodies that have been quarried in the past. North Hill, at the northwest corner of the quadrangle is the site of the 'Old City' quarry, where tonalite intrudes marble producing pyroxene-hornfels grade garnet-pyroxene skarn. South of Riverside several disconnected bodies of marble, impure quartzite, and calcsilicate rock were quarried at the 'New City' (Victoria Ave) quarry. There, the thicker of the two marble bodies was quarried exposing skarn developed at a contact between the marble and intrusive biotite-hornblende tonalite. The composition of the skarn is highly varied and includes pyroxene-garnet, idocrase, scapolite-pyroxene, and magnetite-pageite skarns.

Biotite-hornblende tonalite of the relatively large Val Verde pluton dominates the quadrangle west of Interstate 215. In most places this tonalite has a northwest oriented crude to well developed planar fabric produced by oriented biotite and hornblende. In the northern part of the pluton northeast striking planar fabric dominates. Schlieren and massive clots of mafic tonalite occur locally. Discoidal-to pancake-shaped mafic inclusions are widespread and are oriented in the plane of the biotite and hornblende. Typically, the planar fabric dips moderately to the northeast, but is locally horizontal to subhorizontal or grades to an isotropic fabric.

Granitic rocks in the northeastern part of the quadrangle are part of the Box Springs plutonic complex. This composite intrusion is an elliptical, flat-floored granitic complex centered on the Box Springs Mountains. The exposed part of the complex is apparently the lower part of a granitic diapir. In the center of the complex massive to indistinctly primarily layered biotite tonalite grades outward into well foliated biotite tonalite. Further outward the rocks are a heterogeneous assemblage of primarily granodioritic plutonic rocks.

The geologic map data base contains original U.S. Geological Survey data generated by detailed field observation recorded on 1:24,000 scale aerial photographs. The map was created by transferring lines from the aerial photographs to a 1:24,000 scale topographic base. The map was digitized and lines, points, and polygons were subsequently edited using standard ARC/INFO commands. Digitizing and editing artifacts significant enough to display at a scale of 1:24,000 were corrected. Within the database, geologic contacts are represented as lines (arcs), geologic units are polygons, and site-specific data as points. Polygon, arc, and point attribute tables (.pat, .aat, and .pat, respectively) uniquely identify each geologic datum.

Purpose: The data set for the Riverside East 7.5' quadrangle was prepared under the U.S. Geological Survey Southern California Areal Mapping Project (SCAMP) as part of an ongoing effort to develop a regional geologic framework of southern California, and to utilize a Geographic Information System (GIS) format to create regional digital geologic databases. These regional databases are being developed as contributions to the National Geologic Map Database of the National Cooperative Geologic Mapping Program of the USGS.

Supplemental\_Information: none

Time\_Period\_of\_Content:

Time\_Period\_Information:

Single\_Date/Time:

Calendar\_Date: 2001

Currentness\_Reference: New data

Status:

Progress: Complete

Maintenance\_and\_Update\_Frequency: As Needed

Spatial\_Domain:

Bounding\_Coordinates:

West\_Bounding\_Coordinate: -117.3750914

East\_Bounding\_Coordinate: -117.2499086

North\_Bounding\_Coordinate: 33.99999995

South\_Bounding\_Coordinate: 33.87498415

Keywords:

Theme:

Theme\_Keyword\_Thesaurus: None  
Theme\_Keyword: geologic map  
Theme\_Keyword: geology  
Theme\_Keyword: bedrock geology  
Theme\_Keyword: alluvial geology

Place:

Place\_Keyword\_Thesaurus: None  
Place\_Keyword: California  
Place\_Keyword: Riverside County  
Place\_Keyword: Riverside East 7.5' quadrangle

Stratum:

Stratum\_Keyword\_Thesaurus: None  
Stratum\_Keyword: Cretaceous tonalite and granodiorite  
Stratum\_Keyword: Quaternary deposits

Temporal:

Temporal\_Keyword\_Thesaurus: None  
Temporal\_Keyword: Cretaceous  
Temporal\_Keyword: Quaternary

Access\_Constraints: None

Use\_Constraints:

The Riverside East 7.5' geologic-map database should be used to evaluate and understand the geologic character of the Riverside East 7.5' quadrangle as a whole. The data should not be used for purposes of site-specific land-use planning or site-specific geologic evaluations. The database is sufficiently detailed to identify and characterize many actual and potential geologic hazards represented by faults and landslides and posed by ground subsidence and earthquake-generated ground shaking. However, it is not sufficiently detailed for site-specific determinations or evaluations of these features. Faults shown do not take the place of fault-rupture hazard zones designated by the California State Geologist (see Hart, 1988).

Use of this digital geologic-map database should not violate the spatial resolution of the data. Although the digital form of the data removes the constraint imposed by the scale of a paper map, the detail and accuracy inherent in map scale are also present in the digital data. The fact that this database was compiled and edited at a scale of 1:24,000 means that higher resolution information may not have been uniformly retained in the dataset. Plotting at scales larger than 1:24,000 will not yield greater real detail, although it may reveal fine-scale irregularities below the intended resolution of the database. Similarly, although higher resolution data is incorporated in most of the map, the resolution of the combined output will be limited by the lower resolution data.

Point\_of\_Contact:

Contact\_Information:

Contact\_Person\_Primary:

Contact\_Person: Douglas M. Morton

Contact\_Organization: U.S. Geological Survey, Western Region, Earth Surface Processes Team

Contact\_Position: Project geologist

Contact\_Address:

Address\_Type: mailing address

Address: U.S. Geological Survey

Address: Department of Earth Sciences

Address: University of California, Riverside

City: Riverside

State\_or\_Province: California

Postal\_Code: 92521

Country: United States of America

Contact\_Voice\_Telephone: (909) 276-6397

Contact\_Facsimile\_Telephone: (909) 276-6295

Contact\_Electronic\_Mail\_Address: scamp@usgs.gov

Data\_Set\_Credit: Geologic mapping and digital preparation of this report were sponsored jointly by (1) the National Cooperative Geologic Mapping Program of the U.S. Geological Survey, (2) the California Division of Mines and Geology, (3) the Southern California Areal Mapping Project (SCAMP), and (4) the U.S. Air Force.

Native\_Data\_Set\_Environment:

SunOS, 5.8, sun4m UNIX

ARC/INFO version 7.2.1

Cross\_Reference:

Citation\_Information:

Originator: Morton, D.M.

Publication\_Date: 1999

Title: Preliminary digital geologic map of the Santa Ana 30'x60' quadrangle, southern California, version 1.0.

Geospatial\_Data\_Presentation\_Form: vector digital data

Series\_Information:

Series\_Name: U.S. Geological Survey Open-File Report

Issue\_Identification: USGS OF 99-172

Publication\_Information:

Publication\_Place: California

Publisher: U.S. Geological Survey

Online\_Linkage: <http://geopubs.wr.usgs.gov/open-file/of99-172>

Data\_Quality\_Information:

Attribute\_Accuracy:

Attribute\_Accuracy\_Report:

Geologic-map units in the Riverside East quadrangle database were described using standard field methods. Consistent with these methods, the database author has assigned standard geologic attributes to geologic lines, points, and polygons identified in the database.

Nation-wide geologic-map accuracy standards have not been developed and adopted by the U.S. Geological Survey and

other earth-science entities. Until such standards are adopted, the SCAMP project has developed internal map-accuracy standards for 1:24,000-scale geologic maps produced by the project.

Geologic lines and points on 1:24,000 scale geologic maps are judged to meet SCAMP's internal map-accuracy standards if they are located to within +/-15 meters, relative to topographic or cultural features on the base map.

On any derivative geologic-map plot, line data that are judged to meet the SCAMP internal map-accuracy standard are denoted by solid lines; line data that may not meet the SCAMP internal map-accuracy standard are denoted by dashed or dotted lines. There is no cartographic device for denoting the map-accuracy for geologic-point data (e.g., symbols representing bedding, foliation, lineations, etc.).

Logical\_Consistency\_Report:

Polygon and chain-node topology present.

The areal extent of the map is represented digitally by an appropriately projected (polyconic projection), mathematically generated box. Consequently, polygons intersecting the lines that comprise the map boundary are closed by that boundary. Polygons internal to the map boundary are completely enclosed by line segments which are themselves a set of sequentially numbered coordinate pairs. Point data are represented by coordinate pairs.

Completeness\_Report: The geologic map database of the Riverside East 7.5' quadrangle contains new data that have been subjected to rigorous review and are a substantially complete representation of the current state of knowledge concerning the geology of the quadrangle.

Positional\_Accuracy:

Horizontal\_Positional\_Accuracy:

Horizontal\_Positional\_Accuracy\_Report: The maximum transformation RMS error acceptable for a 7.5' quadrangle transformation and data input is 0.003 (1.8 meters). Horizontal positional accuracy was checked by visual comparison of hard-copy plots with base-stable source data.

Lineage:

Process\_Step:

Process\_Description: Field mapping and aerial photograph interpretation; iterative process (D.M. Morton).

Process\_Date: 1978, 1991, 1995-96

Process\_Step:

Process\_Description: Field mapping and aerial photograph interpretation; iterative process (B.F. Cox).

Process\_Date: 1980

Process\_Step:

Process\_Description: Digitization of geologic linework and point data from a scale-stable cartographic base of quadrangle. ARC/INFO database established; cleanup of artifacts; polygon, arc, and point attribute tables established. Digitizing and editing artifacts significant enough to display at a scale of 1:24,000 were corrected (R.M. Alvarez and V.M. Diep).

Process\_Date: 1999-2001

Process\_Step:

Process\_Description: Description of map units and correlation of map units (F.K. Miller).

Process\_Date: 2001

Process\_Step:

Process\_Description:

First draft of metadata created by Michael J. Watson using  
FGDCMETA.AML ver. 1.2 05/14/98 on ARC/INFO data set  
/scamp26/mwatson/rse\_ofr/rse\_geo

Process\_Date: 20011003

Spatial\_Data\_Organization\_Information:

Direct\_Spatial\_Reference\_Method: Vector

Point\_and\_Vector\_Object\_Information:

SDTS\_Terms\_Description:

SDTS\_Point\_and\_Vector\_Object\_Type: Point

Point\_and\_Vector\_Object\_Count: 183

SDTS\_Point\_and\_Vector\_Object\_Type: String

Point\_and\_Vector\_Object\_Count: 458

SDTS\_Point\_and\_Vector\_Object\_Type: GT-polygon composed of chains

Point\_and\_Vector\_Object\_Count: 184

Spatial\_Reference\_Information:

Horizontal\_Coordinate\_System\_Definition:

Planar:

Map\_Projection:

Map\_Projection\_Name: Polyconic

Polyconic:

Latitude\_of\_True\_Scale: 33.875

Longitude\_of\_Central\_Meridian: -117.3125

False\_Easting: 0.00000

False\_Northing: 0.00000

Planar\_Coordinate\_Information:

Planar\_Coordinate\_Encoding\_Method: coordinate pair

Coordinate\_Representation:

Abscissa\_Resolution: 1.0

Ordinate\_Resolution: 1.0

Planar\_Distance\_Units: Meters

Geodetic\_Model:

Horizontal\_Datum\_Name: North American Datum of 1927  
Ellipsoid\_Name: Clarke 1866  
Semi-major\_Axis: 6378206.4  
Denominator\_of\_Flattening\_Ratio: 294.98

Entity\_and\_Attribute\_Information:

Overview\_Description:

Entity\_and\_Attribute\_Overview:

Version 1.0 of the Riverside East 7.5' quadrangle comprises three ARC/INFO coverages, of which two contain geologic data, and one contains cartographic features: rse\_geo (geology), rse\_str (structural data), and rse\_ano (annotation and leaders).

Geologic data represented by line entities and the polygons they delineate are contained in the coverage RSE\_GEO. For display purposes, the annotation coverage contains one annotation subclass: anno\_geo contains unit labels.

Geological point data includes site-specific information describing the types and the orientation of bedding, foliation, and lineations. Annotation is respective dip and plunge values associated with individual point data.

>

>RSE\_GEO.PAT:

>

>COLUMN ITEM NAME WIDTH OUTPUT TYPE N.DEC ALTERNATE NAME

> 1	AREA	4	12	F	3	
> 5	PERIMETER	4	12	F	3	
> 9	RSE_GEO#	4	5	B	-	
> 13	RSE_GEO-ID	4	5	B	-	
> 17	PTYPE	35	35	C	-	
> 52	LABL	35	35	C	-	
> 87	PLABL	35	35	C	-	
> 122	SHD	3	3	I	-	
> 125	SHDFIL	3	3	I	-	
> 128	NAME	200	200	C	-	

>

>

>RSE\_GEO.AAT:

>

>COLUMN ITEM NAME WIDTH OUTPUT TYPE N.DEC ALTERNATE NAME

> 1	FNODE#	4	5	B	-	
> 5	TNODE#	4	5	B	-	
> 9	LPOLY#	4	5	B	-	
> 13	RPOLY#	4	5	B	-	
> 17	LENGTH	4	12	F	3	
> 21	RSE_GEO#	4	5	B	-	
> 25	RSE_GEO-ID	4	5	B	-	
> 29	LTYPE	35	35	C	-	
> 64	L-SYMB	3	3	I	-	

>

>

Entity\_and\_Attribute\_Detail\_Citation: none

Detailed\_Description:

Entity\_Type:

Entity\_Type\_Label: rse\_geo.pat

Entity\_Type\_Definition: Geologic units (LABL) and their corresponding names (NAME) identified in the Riverside East 7.5' quadrangle

Attribute:

Attribute\_Label: LABL  
Attribute\_Definition: geologic map unit label, in plain text  
Attribute\_Domain\_Values:  
Enumerated\_Domain:  
Enumerated\_Domain\_Value: Kba  
Enumerated\_Domain\_Value\_Definition: Amphibolitic gabbro of Box Springs plutonic complex  
Enumerated\_Domain:  
Enumerated\_Domain\_Value: Kbfg  
Enumerated\_Domain\_Value\_Definition: Biotite granodiorite and tonalite of Box Springs plutonic complex  
Enumerated\_Domain:  
Enumerated\_Domain\_Value: Kbfgi  
Enumerated\_Domain\_Value\_Definition: Biotite granodiorite and tonalite containing abundant inclusions, Box Springs plutonic complex  
Enumerated\_Domain:  
Enumerated\_Domain\_Value: Kbft  
Enumerated\_Domain\_Value\_Definition: Biotite-hornblende tonalite of Box Springs plutonic complex  
Enumerated\_Domain:  
Enumerated\_Domain\_Value: Kbg  
Enumerated\_Domain\_Value\_Definition: Porphyritic granodiorite of Box Springs plutonic complex  
Enumerated\_Domain:  
Enumerated\_Domain\_Value: Kbg  
Enumerated\_Domain\_Value\_Definition: Heterogeneous granodiorite and tonalite of Box Springs plutonic complex  
Enumerated\_Domain:  
Enumerated\_Domain\_Value: Kbhg  
Enumerated\_Domain\_Value\_Definition: Heterogeneous porphyritic granodiorite of Box Springs plutonic complex  
Enumerated\_Domain:  
Enumerated\_Domain\_Value: Kbhg1  
Enumerated\_Domain\_Value\_Definition: Layered heterogeneous porphyritic granodiorite of Box Springs plutonic complex  
Enumerated\_Domain:  
Enumerated\_Domain\_Value: Kbt  
Enumerated\_Domain\_Value\_Definition: Biotite tonalite of Box Springs plutonic complex  
Enumerated\_Domain:  
Enumerated\_Domain\_Value: Kg  
Enumerated\_Domain\_Value\_Definition: Granitic dikes  
Enumerated\_Domain:  
Enumerated\_Domain\_Value: Kgb  
Enumerated\_Domain\_Value\_Definition: Gabbro  
Enumerated\_Domain:  
Enumerated\_Domain\_Value: Kgd  
Enumerated\_Domain\_Value\_Definition: Granodiorite, undifferentiated  
Enumerated\_Domain:  
Enumerated\_Domain\_Value: KgPz  
Enumerated\_Domain\_Value\_Definition: Intermixed schist and gneiss and granitic rocks  
Enumerated\_Domain:  
Enumerated\_Domain\_Value: Khg  
Enumerated\_Domain\_Value\_Definition: Heterogeneous granitic rocks  
Enumerated\_Domain:  
Enumerated\_Domain\_Value: Kp  
Enumerated\_Domain\_Value\_Definition: Granitic pegmatite dikes  
Enumerated\_Domain:  
Enumerated\_Domain\_Value: Krg

Enumerated\_Domain\_Value\_Definition: Granite of the Riverside area

Enumerated\_Domain:  
Enumerated\_Domain\_Value: Kt  
Enumerated\_Domain\_Value\_Definition: Tonalite, undifferentiated

Enumerated\_Domain:  
Enumerated\_Domain\_Value: Kvt  
Enumerated\_Domain\_Value\_Definition: Val Verde tonalite

Enumerated\_Domain:  
Enumerated\_Domain\_Value: Kvt?  
Enumerated\_Domain\_Value\_Definition: Val Verde tonalite

Enumerated\_Domain:  
Enumerated\_Domain\_Value: Pzc  
Enumerated\_Domain\_Value\_Definition: Calc-silicate rocks

Enumerated\_Domain:  
Enumerated\_Domain\_Value: Pzm  
Enumerated\_Domain\_Value\_Definition: Marble

Enumerated\_Domain:  
Enumerated\_Domain\_Value: Pzms  
Enumerated\_Domain\_Value\_Definition: Marble and schist, undifferentiated

Enumerated\_Domain:  
Enumerated\_Domain\_Value: Pzq  
Enumerated\_Domain\_Value\_Definition: Impure quartzite

Enumerated\_Domain:  
Enumerated\_Domain\_Value: Pzs  
Enumerated\_Domain\_Value\_Definition: Biotite schist

Enumerated\_Domain:  
Enumerated\_Domain\_Value: QTs  
Enumerated\_Domain\_Value\_Definition: Unnamed late Cenozoic sedimentary rocks in Riverside and Corona areas

Enumerated\_Domain:  
Enumerated\_Domain\_Value: Qaf  
Enumerated\_Domain\_Value\_Definition: Artificial fill

Enumerated\_Domain:  
Enumerated\_Domain\_Value: Qofa  
Enumerated\_Domain\_Value\_Definition: Old alluvial fan deposits, arenaceous

Enumerated\_Domain:  
Enumerated\_Domain\_Value: Qvofa  
Enumerated\_Domain\_Value\_Definition: Very old alluvial fan deposits, arenaceous

Enumerated\_Domain:  
Enumerated\_Domain\_Value: Qyaa  
Enumerated\_Domain\_Value\_Definition: Young axial channel deposits, arenaceous

Enumerated\_Domain:  
Enumerated\_Domain\_Value: Qyfa  
Enumerated\_Domain\_Value\_Definition: Young alluvial fan deposits, arenaceous

Attribute:  
Attribute\_Label: PLABL  
Attribute\_Definition: Geological map unit label used to generate plot labels with relevant stratigraphic symbols. The geologic units with LABL designating Mesozoic (Mz) have keystone substitute characters, }, that call their corresponding symbols from the Stratagem Font Group. Geologic map unit labels will plot on derivative map plots with appropriate stratigraphic symbols if PLABL is used as the source for unit labels.

Attribute:  
Attribute\_Label: SHD  
Attribute\_Definition: polygon color (as integer value) from shadeset alc1.shd (included in the data package)

Attribute:

Attribute\_Label: SHDFIL  
 Attribute\_Definition: polygon fill pattern (as integer value) from shadeset geology2.shd (included in the data package)  
 Attribute:  
   Attribute\_Label: NAME  
   Attribute\_Definition: Geologic name of map unit (see list under LABL attribute)  
 Detailed\_Description:  
 Entity\_Type:  
   Entity\_Type\_Label: rse\_geo.aat  
   Entity\_Type\_Definition: Geologic features such as contacts and faults that bound rock-unit polygons  
 Attribute:  
   Attribute\_Label: LTYPE  
   Attribute\_Definition: Description of types of lines on the geologic map (contact, fault).  
   Attribute\_Domain\_Values:  
     Enumerated\_Domain:  
       Enumerated\_Domain\_Value: contact, certain  
       Enumerated\_Domain\_Value: fault, approx. located  
       Enumerated\_Domain\_Value: fault, concealed  
       Enumerated\_Domain\_Value: map boundary  
 Attribute:  
   Attribute\_Label: L-SYMB  
   Attribute\_Definition: stores appropriate line symbol value from the lineset geoscamp2.lin  
 Detailed\_Description:  
 Entity\_Type:  
   Entity\_Type\_Label: rse\_str.pat  
   Entity\_Type\_Definition: Geological point data includes site-specific information describing the types and the orientation of bedding, foliation, and lineations. One annotation subclass is included in the geologic points coverage, RSE\_STR which displays the respective dip and plunge values associated with individual point data.  
 Attribute:  
   Attribute\_Label: PTTYPER  
   Attribute\_Definition: describes type of point data (bedding, horizontal bedding, foliation)  
 Attribute:  
   Attribute\_Label: P-SYMB  
   Attribute\_Definition: Coded integer value that relates point to cartographic point symbol in markerset geoscamp2.mrk  
 Attribute:  
   Attribute\_Label: STRIKE  
   Attribute\_Definition: Azimuthal strike of planar feature  
 Attribute:  
   Attribute\_Label: DIP  
   Attribute\_Definition: Dip of planar feature  
 Detailed\_Description:  
 Entity\_Type:  
   Entity\_Type\_Label: rse\_ano.aat  
   Entity\_Type\_Definition: Annotation leaders  
 Attribute:  
   Attribute\_Label: L-SYMB  
   Attribute\_Definition: Coded integer value (1) that relates arcs to cartographic line symbol in lineset geoscamp2.lin  
 Distribution\_Information:  
 Distributor:  
   Contact\_Information:  
     Contact\_Organization\_Primary:  
       Contact\_Organization: U.S. Geological Survey Information Services



Contact\_Address:

Address\_Type: mailing address  
Address: Box 25286 Denver Federal Center  
City: Denver  
State\_or\_Province: Colorado  
Postal\_Code: 80225  
Country: USA  
Contact\_Voice\_Telephone: (303)202-4700

Contact\_Facsimile\_Telephone: (303)202-4693

Distribution\_Liability:

The U.S. Geological Survey (USGS) provides these geographic data "as is." The USGS makes no guarantee or warranty concerning the accuracy of information contained in the geographic data. The USGS further makes no warranties, either expressed or implied as to any other matter whatsoever, including, without limitation, the condition of the product, or its fitness for use lies entirely with the user. Although these data have been processed successfully on computers at the USGS, no warranty, expressed or implied, is made by the USGS regarding the use of these data on any other system, nor does the fact of distribution constitute or imply any such warranty.

In no event shall the USGS have any liability whatsoever for payment of any consequential, incidental, indirect, special, or tort damages of any kind, including, but not limited to, any loss of profits arising out of use of or reliance on the geographic data or arising out of the delivery, installation, operation, or support by USGS.

This digital geologic map database of the Riverside East 7.5' quadrangle, 1:24,000 map-scale, and any derivative maps thereof, is not meant to be used or displayed at any scale larger than 1:24,000 (e.g., 1:12,000).

Metadata\_Reference\_Information:

Metadata\_Date: 20011003  
Metadata\_Review\_Date: 20011115

Metadata\_Contact:

Contact\_Information:

Contact\_Organization\_Primary:  
Contact\_Organization: U.S. Geological Survey  
Contact\_Person: Rachel M. H. Alvarez  
Contact\_Position: Geologist  
Contact\_Address:  
Address\_Type: mailing address  
Address: U.S. Geological Survey  
Address: Department of Earth Sciences  
Address: University of California, Riverside  
City: Riverside  
State\_or\_Province: California  
Postal\_Code: 92521  
Country: USA  
Contact\_Voice\_Telephone: (909) 276-6397  
Contact\_Facsimile\_Telephone: (909) 276-6295  
Contact\_Electronic\_Mail\_Address: (rhauser@usgs.gov)

Metadata\_Standard\_Name: FGDC Content Standards for Digital Geospatial Metadata

Metadata\_Standard\_Version: Version of June 8, 1994

Metadata\_Access\_Constraints: none

Metadata\_Use\_Constraints: none