

Yellowstone National Park's physical landscape has been and is being created by many geological forces. Here, some of the Earth's most active volcanic, hydrothermal (water + heat), and earthquake systems make this national park a priceless treasure. In fact, Yellowstone was established as the world's first national park primarily because of its extraordinary geysers, hot springs, mudpots and steam vents, and other wonders such as the Grand Canyon of the Yellowstone River.

What Lies Beneath

Yellowstone's geologic story provides examples of how geologic processes work on a planetary scale. The foundation to understanding this story begins with the structure of the Earth and how this structure gives rise to forces that shape the planet's surface.

The Earth is frequently depicted as a ball with a central core surrounded by concentric layers that culminate in the crust or surface layer (*see at right*). The distance from the Earth's surface to its center or core is approximately 4,000 miles. The core may once have been entirely molten, but, as the planet cooled, the inner core (about 1,500 miles thick) solidified. The outer core (about 1,400 miles thick) remains molten and is surrounded by a 1,800 mile thick mantle of dense, mostly solid rock. Above this layer is the relatively thin crust, three to forty-eight miles thick, on which the continents and ocean floors are found.

The Earth's lithosphere (crust and upper mantle; *see illustration next page*) is divided into many plates, which are in constant motion. Where plate edges meet, one plate may slide past another, one plate may be driven beneath another (subduction), or upwelling volcanic material pushes the plates apart at mid-ocean ridges. Continental plates are made of less dense rocks (granites) than oceanic plates (basalts) and thus, "ride" higher than oceanic plates. Many theories have been proposed to explain crustal plate movement. Currently, most evidence supports the theory that con-

YELLOWSTONE'S GEOLOGIC SIGNIFICANCE

- One of the most geologically dynamic areas on Earth due to shallow source of magma and resulting volcanic activity.
 - One of the largest volcanic eruptions known to have occurred in the world, creating one of the largest known calderas.
 - More than 10,000 hydrothermal features, including more than 300 geysers.
 - The largest concentration of active geysers in the world—
- approximately half of the world's total.
- Most of the undisturbed geyser basins left in the world. (Kamchatka Peninsula has the others; the rest have been modified or destroyed by human development.)
 - One of the few places in the world where active travertine terraces are found, at Mammoth Hot Springs.
 - Site of many petrified trees resulting from repeated volcanic eruptions over the ages.

vection currents in the partially molten asthenosphere (the zone of mantle beneath the lithosphere) move the rigid crustal plates above. The volcanism that has so greatly shaped today's Yellowstone is a product of plate movement combined with upwellings of molten rock, as described on the next pages.

3

Ancient Yellowstone

Lamar Canyon contains outcrops of granite and granitic gneiss more than two billion years old. Over time, heat and pressure have altered these rocks and obscured their early history.

Illustrations on pages 55–58 & page 64 adapted with permission from Windows Into the Earth, Dr. Robert Smith and Lee J. Siegel, 2000.

Yellowstone Geologic History (mya=millions of years ago)

570 to 66 mya, area covered by inland seas

50–40 million years ago
—Absaroka Volcanics—

Most of Earth's history (from the beginning to approximately 570 million years ago) is known as the Precambrian time. Rocks of this age are found in northern Yellowstone and in the hearts of the Teton, Beartooth, Wind River, and Gros Ventre ranges.

During the Precambrian and the subsequent Paleozoic and Mesozoic eras (570 to 66 million years ago), the western United States was covered at times by ocean. At other times it was a land of sand dunes, tidal flats, and vast plains. Near the end of the Mesozoic, mountain building processes created the Rocky Mountains.

During the Cenozoic era (approximately the last 66 million years of Earth's history), widespread mountain-building, volcanism, faulting, and glaciation sculpted the Yellowstone area. The Absaroka Range along the park's north and east sides was formed by numerous volcanic eruptions about 50 million years ago. Volcanic debris buried trees that are seen today as fossilized remnants along Specimen Ridge in northern Yellowstone. This period of volcanism is not related to the present Yellowstone volcano.

Approximately 30 million years ago, vast expanses of the West began stretching apart

along an east-west axis. This stretching process increased about 17 million years ago and continues today, creating the modern basin and range topography (north-south mountain ranges interspersed with long north-south valleys) characterizing much of the Western landscape.

About 16.5 million years ago, an intense period of volcanism appeared near the area now marked by the convergence of the Nevada, Oregon, and Idaho state lines. Repeated volcanic eruptions can be traced across southern Idaho into Yellowstone National Park. *This volcanism remains a driving force in Yellowstone today.*


Magma & Hotspots

Magma (molten rock from Earth's mantle) rises to within a few miles of the surface in Yellowstone. This heat fuels the Yellowstone volcano and its associated hydrothermal areas. How it rises, whether or not a hotspot is involved, and the definition of a hotspot all remain the subject of much scientific research and discussion.

One hotspot theory holds that a plume of molten rock rises from Earth's core-mantle boundary to trigger volcanic eruptions at

the surface. Other theories relate the rise of molten rock to areas in Earth's crust weakened by stretching and thinning such as that which is ongoing throughout the interior West. Some of these theories also propose a shallower mantle origin for hotspots. Still other theories define Yellowstone's hotspot

(see below). Upward pressure from the shallow magma chamber caused overlying rocks to break, forming faults and causing earthquakes. Eventually, these faults reached the deep magma chamber. Magma oozed through these cracks, releasing pressure within the chamber and allowing trapped



as the surface manifestation of longlived volcanism.

Regardless of its origins and subsurface behavior, the magma feeding Yellowstone's volcano has been close to the surface for some 16.5 million years, erupting repeatedly and leaving a track of 100 huge calderas across 500 miles from the Nevada-Oregon border northeast up Idaho's Snake River Plain and into central Yellowstone. This trail of evidence was created as the North American plate moved in a southwestern direction over the shallow magma body. Earth's surface bulges above it, which is why the Yellowstone area averages 1,700 feet higher than surrounding regions.

About 2.1 million years ago, the movement of the North American plate brought the Yellowstone area into proximity with the shallow magma body. The heat melted rocks in the crust, creating a magma chamber of partially molten, partially solid rock

gases to expand rapidly. A massive eruption then occurred, spewing volcanic ash and gas high into the atmosphere and causing fast-moving superhot debris (pyroclastic) flows on the ground. As the underground magma chamber emptied, the ground above it sunk, creating a huge caldera known as the Huckleberry Ridge Caldera (also called the Island Park Caldera). Smaller lava flows eventually filled in the caldera over tens to hundreds of thousands of years.

The volume of material ejected during this eruption is estimated to have been 6,000 times the size of the 1980 eruption of Mt. St. Helens in Washington (see illustration next page), and ash has been found as far away as Missouri. Approximately 1.3 million years ago, a second, smaller volcanic eruption occurred on the western edge of the earlier caldera and created the Henry's Fork Caldera. Then 640,000 years ago, the third massive volcanic eruption in central Yellowstone created the Yellowstone

At Yellowstone and some other volcanoes, some scientists theorize that Earth's crust fractures and cracks in a concentric or ring-fracture pattern. When these cracks reach the magma chamber, pressure is released and the volcano explodes. The huge amount of material released causes the volcano to collapse into a huge caldera.

Names for the three major Yellowstone volcano eruptions & the resulting calderas vary, in part, on who is doing the research and what they are studying. In the text, we use names given by the scientists of the Yellowstone Volcano Observatory, but the illustrations may show different names.

*West Thumb
date change*



Caldera, 30 by 45 miles in size. Then approximately 180,000 years ago, a volcanic eruption created a smaller caldera now filled by the West Thumb of Yellowstone Lake. Yellowstone remains atop the shallow magma. The pressure and movement of the underlying heat, magma, and fluids cause the entire caldera floor to inflate and deflate rapidly (compared to more typical geologic processes).

Rising magma has created two large bulges in the Earth called resurgent domes (Sour Creek and Mallard Lake), which we see as large hills.

Sour Creek Dome, east of the Yellowstone River from LeHardys Rapids through Hayden Valley, currently has a net uplift of about 1½ inches per year. This uplift is causing Yellowstone Lake to tilt southward. Larger sandy beaches can now be found on the north shore of the lake, and flooded areas can be found in the southern arms.

From the summit of Mt. Washburn, one can look south into much of the vast caldera. The rim is also visible along the park road system at Gibbon Falls, Lewis Falls, and Lake Butte.

Future Volcanic Activity

Will Yellowstone's volcano erupt again? Over the next thousands to millions of years, probably. In the next few hundred years? Not likely.

The most likely activity would be lava flows, such as those that occurred after the last major eruption. Such a lava flow would ooze slowly over months and years, allowing plenty of time for park managers to evaluate the situation and protect people. No scientific evidence indicates such a lava flow will occur soon.

Geyser Basin Systems

Yellowstone’s hydrothermal features would not exist without the underlying magma body that releases tremendous heat. They also depend on sources of water, such as in the mountains surrounding the Yellowstone Plateau. There, snow and rain slowly percolate through layers of porous rock riddled with cracks and fissures. Some of this cold water meets hot brine directly heated by the shallow magma body. The water’s temperature rises well above the boiling point but the water remains in a liquid state due to the great pressure and weight of the overlying rock and water. The result is superheated water with temperatures exceeding 400°F.

The superheated water is less dense than the colder, heavier water sinking around it. This creates convection currents that allow the

lighter, more buoyant, superheated water to begin its journey back to the surface following the cracks, fissures, and weak areas through rhyolitic lava flows. As hot water travels through this rock, the high water temperatures dissolve some silica in the rhyolite.

While in solution underground, some silica coats the walls of the cracks and fissures to form a nearly pressure-tight seal. This locks in the hot water and creates a natural “plumbing” system that can withstand the great pressure needed to produce a geyser. At the surface, silica precipitates to form either geyserrite or sinter, creating the massive geyser cones, the scalloped edges of hot springs, and the seemingly barren landscape of geyser basins.



Cone geysers, such as Riverside in Upper Geyser Basin (above) erupt in a narrow jet of water, usually from a cone. **Fountain geysers**, such as Echinus in Norris Geyser Basin (right) shoot water in various directions, typically from a pool.

Geysers are hot springs with constrictions in their plumbing, usually near the surface, that prevent water from circulating freely to the surface where heat would escape. The deepest circulating water can exceed the surface boiling point (199°F/93°C). Surrounding pressure also increases with depth, much as it does with depth in the ocean. Increased pressure exerted by the enormous weight of the overlying rock and water prevents the water from boiling. As the water rises, steam forms. Bubbling upward, steam expands as it nears the top of the water column until the bubbles are too large and numerous to pass freely through the tight spots. At a critical point, the confined bubbles actually lift the water above, causing the geyser to splash or overflow. This decreases pressure on the system, and violent boiling results. Tremendous amounts of steam force water out of the vent, and an eruption begins. Water is expelled faster than it can enter the geyser’s plumbing system, and the heat and pressure gradually decrease. The eruption stops when the water reservoir is depleted or when the system cools.



3

Hydrothermal Features

Fumaroles or steam vents, are the hottest hydrothermal features in the park. They have so little water that it all flashes into steam before reaching the surface. At places like Roaring Mountain (right), the result is a loud hissing of steam and gases.



Travertine terraces, found at Mammoth Hot Springs (right), are formed from limestone (calcium carbonate). Thermal waters rise through the limestone, carrying high amounts of dissolved calcium carbonate. At the surface, carbon dioxide is released and calcium carbonate is deposited as travertine, the chalky white rock of the terraces. Due to the rapid rate of deposition, these features constantly and quickly change.




Mudpots such as Fountain Paint Pot (center, right) are acidic hot springs with a limited water supply. Some microorganisms use hydrogen sulfide, which rises from deep within the earth, as an energy source. They help convert the gas to sulfuric acid, which breaks down rock into clay. Various gases escape through the wet clay mud, causing it to bubble. Mudpot consistency and activity vary with the seasons and precipitation.



Hot springs such as this one at West Thumb (right) are the most common hydrothermal features in the park. Their plumbing has no constrictions. Superheated water cools as it reaches the surface, sinks, and is replaced by hotter water from below. This circulation, called convection, prevents water from reaching the temperature needed to set off an eruption.






Beneath Yellowstone Lake

Until the late 1990s, few details were known about the geology beneath Yellowstone Lake. In 1996, researchers saw anomalies on the floor of Bridge Bay as they took depth soundings. They deployed a submersible remotely operated vehicle (ROV) equipped with photographic equipment and sector-scan sonar. Large targets appeared on the sonar image, then suddenly very large, spire-like structures appeared in the photographic field of view (*photo at right*). These structures looked similar to hydrothermal structures found in deep ocean areas, such as the Mid-Atlantic Ridge and the Juan de Fuca Ridge. They also provided habitat for aquatic species such as freshwater sponges and algae.

Lake-bottom Surveys

From 1999 to 2003, scientists from the U.S. Geological Survey and a private company, Eastern Oceanics, surveyed the bottom of Yellowstone Lake using high-resolution, multi-beam swath sonar imaging, seismic reflection profiling, and a ROV. The survey showed the northern half of the lake to be inside the 640,000-year-old Yellowstone Caldera and mapped previously unknown features such as large hydrothermal explosion craters, siliceous spires, hundreds of hydrothermal vents and craters, active fissures, and domal features containing gas pockets and deformed sediments. It also mapped young previously unmapped faults, landslide deposits, and submerged older lake shorelines. These features are part of an undulating landscape shaped by old rhyolitic lava flows that filled the caldera. The southern half of the lake lies outside the

Hydrothermal vents in northern Yellowstone Lake (above) were mapped as part of a five-year project. Scientists also are studying spires from Bridge Bay (below) that were discovered in 1996. They seem to be very old hydrothermal vents.



caldera and has been shaped by glacial and other processes. The floor of the Southeast Arm has many glacial features, similar to the glacial terrain seen on land in Jackson Hole, south of the park.

These new surveys give an accurate picture of the geologic forces shaping Yellowstone Lake and determine geologic influences affecting the present-day aquatic biosphere.

For example, hydrothermal explosions formed craters at Mary Bay and Turbid Lake. Spires may form similarly to black smoker chimneys, which are hydrothermal features associated with oceanic plate boundaries.

Spire Analysis

With the cooperation of the National Park Service, scientists from the University of Wisconsin–Milwaukee collected pieces of spires and a complete small spire for study by several teams. They conducted a CAT scan of the spire, which showed structures seeming to be conduits, perhaps for hydrothermal circulation. When they cut open the spire, they confirmed the presence of

conduits and also saw a layered structure.

Tests by the U.S. Geological Survey show that the spire is about 11,000 years old, which indicates it was formed after the last glaciers retreated. In addition to silica, the spire contains diatom tests (shells) and silica produced by underwater hydrothermal processes. The spire's interior shows


evidence of thermophilic bacteria. Scientists say this suggests that silica precipitated on bacterial filaments, thus playing an important role in building the spire.

Both research projects expanded our understanding of the geological forces at work beneath Yellowstone Lake. Additional study of the spires and other underwater features will continue to contribute to our understanding of the relationship between these features and the aquatic ecosystem.


Illustrations on this page adapted from originals by Dr. Lisa A. Morgan, U.S.G.S. Research Geologist



Initial Spire Growth



Mature Spire



Earthquakes

Yellowstone is the most seismically active area in the Intermountain West. Approximately 2,000 earthquakes occur each year in the Yellowstone area; most are not felt.

Earthquakes occur along fault zones in the crust where forces from crustal plate movement build to a significant level. The rock along these faults becomes so stressed that eventually it slips or breaks. Energy is then released as shock waves (seismic waves) that reverberate throughout the surrounding rock.

Different kinds of seismic waves are released inside the earth during an earthquake. Primary waves (“P-waves”) move quickly in the direction of travel, compressing and stretching the rock. Secondary waves (“S-waves”) move up, down, and sideways through rock in a rolling motion. Once a seismic wave reaches the surface of the Earth, it may be felt. Surface waves affect the ground, which can roll, crack open, or be vertically and/or laterally displaced. Structures are susceptible to earthquake damage because the ground motion is usually horizontal.

Earthquakes in Yellowstone help to maintain hydrothermal activity by keeping the “plumbing” system open. Without periodic disturbance of relatively small earthquakes,

the small fractures and conduits that supply hot water to geysers and hot springs might be sealed by mineral deposition. Some earthquakes generate changes in Yellowstone’s hydrothermal systems. For example, the 1959 Hebgen Lake (*see below*) and 1983 Borah Peak earthquakes caused measurable changes in Old Faithful Geyser and other hydrothermal features.

Sometimes Yellowstone experiences an “earthquake swarm.” Dr. Jake Lowenstern of the Yellowstone Volcano Observatory defines this as “a packet of earthquakes close in time and space but with no large earthquakes relative to the others.” The most active swarm since 1985 began on December 26, 2008 under Yellowstone Lake, and lasted one week. More than 500 earthquakes were recorded; the largest was 3.9 magnitude. No changes in hydrothermal activity were detected.

Earthquakes help us understand the subsurface geology around and beneath Yellowstone. The energy from earthquakes travels through hard and molten rock at different rates. We can “see” the subsurface and make images of the magma chamber and the caldera by “reading” the energy emitted during earthquakes. An extensive geological monitoring system is in place to aid in that interpretation.

... 9 quakes in all that I’ve felt in 24 hours. Lake residents are becoming familiar with the Richter scale. Experience the quake ride at Lake!—Ranger Matt Johnson, December 28, 2008

In 2008, more than 1800 earthquakes occurred in the park. Hundreds of them occurred in the last week of the year.

Real-time data about earthquakes in Yellowstone is at www.seis.utah.edu, maintained by the University of Utah Seismograph Stations.

The Night that Woke Up Yellowstone

Twenty-seven minutes before midnight, August 17, 1959, most visitors in Yellowstone were asleep after their day’s adventures. Then a massive earthquake struck, jolting them awake. People inside the park were lucky—rocks fell and buildings were damaged, no one was seriously hurt. West of the park, though, twenty-eight people died that night—most of them when the side of a mountain slid onto a crowded campground.

Measuring at 7.5 magnitude, this quake is known as the Hebgen Lake Earthquake. The mountain slide dammed the Madison River, creating Quake Lake west of the park. Geysers and hot springs in Yellowstone changed overnight, and at least one new geyser appeared—Earthquake Geyser in the Lower Geyser Basin. It erupted 125 feet high for two weeks then settled to a low, perpetual spouting that continues today.

This summer, the Forest Service is commemorating the 50th anniversary of the Hebgen Lake Earthquake at the Earthquake Lake Visitor Center, located at the site of the mountain slide. For more information, call 406-823-6961 or go to www.fs.fed.us/r1/gallatin/?page=recreation/areas/visitor_centers/quake_lake_vc




Rock slide at Gibbon Falls Overlook

3

Glaciers

The extent of two major glaciations is shown on this map: Bull Lake—orange outline; Pinedale—blue outline



Glaciers

Glaciers result when, for a period of years, more snow falls in an area than melts. Once the snow reaches a certain depth, it turns into ice and begins to move under the force of gravity or the pressure of its own weight. During this movement, rocks are picked up and carried in the ice, and these rocks grind Earth's surface, eroding and carrying material away. Glaciers also deposit materials. Large U-shaped valleys, ridges of debris (moraines), and out-of-place boulders (erratics) are evidence of a glacier's passing.

Yellowstone and much of North America have experienced numerous periods of glaciation during the last two million years. Succeeding periods of glaciation have destroyed most surface evidence of previous glacial periods, but scientists have found evidence of them in sediment cores taken on land and in the ocean. In Yellowstone, a glacial deposit near Tower Fall dates back 1.3 million years. Evidence of such ancient glaciers is rare.

The Bull Lake Period glaciers covered the region about 140,000 years ago. Evidence exists that this glacial episode extended farther south and west of Yellowstone than the

subsequent Pinedale Glaciation (described in the next paragraph), but no surface evidence of it is found to the north and east. This indicates that the Pinedale Glaciation covered or eroded surface evidence of Bull Lake Glaciation in these areas.

The Yellowstone region's last (and most studied) major glaciation, the Pinedale, began as early as 70,000 years ago. It ended 13,000–14,000 years ago. Glaciers advanced and retreated from the Beartooth Mountains, scouring the landscape we know today as the northern range. Glacial dams backed up and released enormous amounts of water, contributing to the formation of Hayden Valley and the Black Canyon of the Yellowstone River. During the peak (16,000–20,000 years ago) nearly all of Yellowstone was covered by a huge ice cap 4,000 feet thick (*see above*). Mount Washburn and Mount Sheridan were both completely covered by ice. This ice cap was not part of the continental ice sheet extending south from Canada. The ice cap occurred here, in part, because the hotspot beneath Yellowstone had pushed up the area to a higher elevation with colder temperatures and more precipitation than the surrounding land.

Scientific understanding of glacier dates, sequence, and extent continues to evolve, and varying information appears in different references (including previous editions of this book). The information here is considered current by Yellowstone's geologist as of January 2009.

Sedimentation & Erosion

Not all the rocks in Yellowstone are of “recent” volcanic origin. Precambrian igneous and metamorphic rock in the northeastern portion of the park and Beartooth Mountains are at least 2.7 billion years old. These rocks are very hard and erode slowly.

Sedimentary sandstones and shales, deposited by seas during the Paleozoic and Mesozoic eras (570 million to 66 million years ago) can be seen in the Gallatin Range and Mount Everts. Sedimentary rocks in Yellowstone tend to erode more easily than the Precambrian rocks.

Weathering breaks down earth materials from large sizes to small particles, and happens in place. The freeze/thaw action of ice is one type of weathering common in Yellowstone. Agents of erosion—wind, water, ice, and waves—move weathered materials from one place to another.

When erosion takes place, sedimentation—the deposition of material—also eventually occurs. Through time, sediments are buried by more sediments and the material hardens into rock. This rock is eventually exposed (through erosion, uplift, and/or faulting), and the cycle repeats itself. Sedimentation and erosion are “reshapers” and “refiners” of the landscape—and they also expose Yellowstone’s past life as seen in fossils like the petrified trees (*see next page*).



Above: The Beartooth Mountains northeast of Yellowstone are actually an uplifted block of Precambrian rock.
Left: Mt. Everts, near Mammoth, exposes sedimentary rock, which erodes easily and often tumbles or slides into Gardner Canyon.

Fossils

Paleobotany

Nearly 150 species of fossil plants (exclusive of fossil pollen specimens) from Yellowstone have been described, including ferns, horse-tail rushes, conifers, and deciduous plants such as sycamores, walnuts, oaks, chestnuts, maples, and hickories. Sequoia is abundant,



In Yellowstone, many petrified trees can be seen (above). Resulting from volcanic eruptions about 50 million years ago, they present questions that scientists continue to ponder: Were the trees petrified in place and thus represent layers of forest? Or were they scattered before and after petrification, which means the number of forests cannot be determined?

and other species such as spruce and fir are also present.

Most petrified wood and other plant fossils come from Eocene deposits about 50 million years old, which occur in many northern parts of the park, including the Gallatin Range, Specimen Creek, Tower, Crescent Hill, Elk Creek, Specimen Ridge, Bison Peak, Barronette Peak, Abiathar Peak, Mount Norris, Cache Creek, and Miller Creek. Petrified wood is also found along streams in areas east of Yellowstone Lake. The most accessible petrified tree site is on Specimen Ridge.

The first fossil plants from Yellowstone were collected by the early Hayden Survey parties. In his 1878 report, Holmes made the first reference to Yellowstone's fossil "forests." The report identified the petrified trees on the north slope of Amethyst Mountain opposite the mouth of Soda Butte Creek, about eight miles southeast of Junction Butte.

Around 1900, F. H. Knowlton identified 147 species of fossil plants from Yellowstone, 81 of them new to science. He also proposed

the theory that the petrified trees on the northwest end of Specimen Ridge were forests petrified in place.

Another theory proposes that the trees were uprooted by volcanic debris flows and transported to lower elevations. The 1980 eruption of Mount St. Helens supported this idea. Mud flows not only transported trees to lower elevations, they also deposited the trees upright.

Cretaceous marine and nonmarine sediments are exposed on Mount Everts. The area is under study; fossil leaves, ferns, clam-like fossils, shark teeth, and several species of vertebrates have been found. In 1994 fossil plants were discovered in Yellowstone during the East Entrance road construction project, which uncovered areas containing fossil sycamore leaves and petrified wood.

Fossil Invertebrates

Fossil invertebrates are abundant in Paleozoic rocks, especially the limestones associated with the Madison Group in the northern and south-central parts of the park. They include corals, bryozoans, brachiopods, trilobites, gastropods, and crinoids. Trace fossils, such as channeling and burrowing of worms, are found in some petrified tree bark.

Fossil Vertebrates

Fossil remains of vertebrates are rare, but perhaps only because of insufficient field research. A one-day survey led by paleontologist Jack Horner, of the Museum of the Rockies, Bozeman, Montana, resulted in the discovery of the skeleton of a Cretaceous vertebrate. Other vertebrate fossils found in Yellowstone include:

- Fish: crushing tooth plate; phosphatized fish bones; fish scales; fish teeth.
- Horse: possible Pleistocene horse, *Equus nebraskensis*, reported in 1939.
- Other mammals: Holocene mammals recovered from Lamar Cave; Titanother (type of rhinoceros) tooth and mandible found on Mt. Hornaday in 1999.

Please remember that collecting any natural resources, including rocks and fossils, is illegal in Yellowstone National Park.

Staff reviewers: Hank Heasler, Geologist; Cheryl Jaworowski, Geologist; Elaine Hale, Archeologist

General

- Anderson, Roger J. and D. Harmon, eds. 2002. *Yellowstone Lake: Hotbed of Chaos or Reservoir of Resilience? Proceedings of the 6th Biennial Scientific Conference on the Greater Yellowstone Ecosystem*. Yellowstone Center for Resources and George Wright Society.
- Ehrlich, Gretel. 1987. *Land of Fire and Ice*. New York: Harper Collins.
- Fritz, William J. 1985. *Roadside Geology of the Yellowstone Country*. Missoula: Mountain Press Publishing Company.
- Good, John M. and Kenneth L. Pierce. 1996. *Interpreting the Landscapes of Grand Teton and Yellowstone National Parks: Recent and Ongoing Geology*. Moose, WY: Grand Teton Natural History Association.
- Hamilton, Wayne L. Geological investigations in Yellowstone National Park, 1976–1981. In *Wyoming Geological Association Guidebook*.
- Keefer, William R. 1976. *The Geologic Story of Yellowstone National Park*. U.S. Geological Survey.
- Raymo, Chet. *The Crust of Our Earth*. Englewood Cliffs, NJ: Prentice-Hall.
- Smith, Robert B. and Robert L. Christiansen. 1980. Yellowstone Park as a window on the earth's interior. *Scientific American*. 242(2): 1004–117. February.
- Smith, Robert B. and Lee J. Siegel. 2000. *Windows Into the Earth: The Geologic Story of Yellowstone and Grand Teton National Parks*. New York: Oxford University Press.
- Tuttle, Sherwood D. 1997. Yellowstone National Park in *Geology of National Parks*. Dubuque, IA: Kendall–Hunt Publishing Company.

Volcanic

- Christiansen, Robert L. 2000. *The Quaternary + Pliocene, Yellowstone Plateau Volcanic Field of Wyoming, Idaho, and Montana*. USGS Professional Paper 729–6.
- Christiansen, Robert L. et al. 2007. Preliminary assessment of volcanic and hydrothermal hazards in Yellowstone National Park and vicinity. USGS Open-file Report 2007-1071; v. 1.1. pubs.usgs.gov/of/2007/1071/.
- Christiansen, Robert L. et al. 2002. Upper-mantle origin of the Yellowstone hotspot. *Geological Society of America Bulletin*. October. 114:10, pgs. 1245–1256.
- Christiansen, Robert L. et al. 1994. *A Field-Trip Guide to Yellowstone National Park, Wyoming, Montana, and Idaho—Volcanic, Hydrothermal, and Glacial Activity in the Region*. U.S. Geological Survey Bulletin 2099.
- Cottrell, Dr. William H. 1987. *Born of Fire: The Volcanic Origin of Yellowstone National Park*. Boulder: Roberts Rinehart.
- Francis, Peter. 1983. Giant Volcanic Calderas. *Scientific American*. June.
- Puskas, C. M. 2000. *Deformation of the Yellowstone Caldera, Hebgen Lake fault zone, and eastern Snake River plain from GPS, seismicity, and moment release*. Masters thesis. U Utah.
- Scientific American. 1982. *Volcanoes and the Earth's Interior*. W.H. Freeman & Co.
- Watt, Fiona. *Usborne Guide: Earthquakes and Volcanoes*. Tulsa: EDC Publishing.

Hydrothermal

- Bryan, T. Scott. 1990. *Geysers: What They Are and How They Work*. Niwot, CO: Roberts Rinehart.
- Bryan, T. Scott. 2008. *The Geysers of Yellowstone*. Boulder: Colorado Associated University Press. Fourth Edition.
- Fouke, B. W. et al. 2000. Depositional facies and aqueous-solid geochemistry of travertine-depositing hot springs (Angel Terrace, Mammoth Hot Springs, Yellowstone National Park, U.S.A.) *J. Sedimentary Res.* 70(3): 565–585.
- Fournier, R. O. 1989. Geochemistry and dynamics of the Yellowstone National Park hydrothermal system. *Ann. Rev. Earth Planet. Sci.* 17:13–53.
- Gallant, Roy A. 1997. *Geysers: When Earth Roared*. Franklin Watts.
- Ingebritsen, S. E. and S. A. Rojstaczer. 1993. Controls on geyser periodicity. *Science*. 262: 889–892.
- Ingebritsen, S. E. and S. A. Rojstaczer. 1996. Geyser periodicity and the response of geysers to deformation. *J. Geophys. Res.* 101(B10): 21,891–21,905.
- Marler, George D. 1973. Inventory of thermal features of the Firehole River geyser basins and other selected areas of Yellowstone National Park. U.S. Department of Commerce, National Technical Information Service, Publication PB221 289.
- Marler, George D. 1969. *The Story of Old Faithful*. Mammoth, WY: Yellowstone Library and Museum Association.
- Rinehart, John S. 1976. *Guide to Geyser Gazing*. Santa Fe: Hyper Dynamics.

Yellowstone Lake

- Cuhel, Russell et al. 2005. The Bridge Bay spires. *Yellowstone Science*. 12(4): 25–40.
- Morgan, Lisa et al. 2003. The floor of Yellowstone Lake is anything but quiet. *Yellowstone Science*. 11(2): 15–30.
- Morgan, Lisa, ed. 2007. *Integrated geoscience studies in the greater Yellowstone area—Volcanic, tectonic, and hydrothermal processes in the Yellowstone geocosystem*. USGS Professional Paper 1717. pubs.usgs.gov/pp/1717/

Earthquakes

- Christopherson, Edmund. 1960, 1962. *The Night the Mountain Fell*. West Yellowstone, MT/Yellowstone Publications.

Glaciers

- Liccardi, J.M., and Pierce, K.L. 2008. Cosmogenic exposure-age chronologies of Pinedale and Bull Lake glaciations in greater Yellowstone and the Teton Range, USA. *Quaternary Science Reviews* 27: 817–831.
- Pierce, Kenneth L. 1979. *History and dynamics of glaciation in the Northern Yellowstone National Park Area*. U.S. Geological Survey Professional Paper 729–F.
- Pierce, Kenneth L. 2004. Pleistocene glaciations of the Rocky Mountains in *Developments in Quaternary Science*, J. Rose (ed). 1: 63–76. Elsevier Press.

Sedimentation/Erosion

- Kharaka, Y.K. and A.S. Maest, eds. 1992. *Proc. of the 7th intern. symp. on water-rock interaction-WRI-7*. Park City, Utah.

Fossils

- Cannon, K.P. et al. 1997. *Results of Archeological and Paleoenvironmental Investigations Along the North Shore of Yellowstone Lake, Yellowstone National Park, Wyoming: 1990–1994*. Lincoln: NPS Midwest Archeological Center.

www.nps.gov/yell.

www.greateryellowstone-science.org/index.html

Yellowstone Science, free from the Yellowstone Center for Resources, in the Yellowstone Research Library, or online at www.nps.gov/yell. Recent articles covered historic sites and park history.

Yellowstone Today, distributed at entrance gates and visitor centers.

Site Bulletins, published as needed, provide more detailed information on park topics such as geology & wildlife. Free; available upon request from visitor centers.

Podcasts and livestream webcam at Old Faithful available at www.nps.gov/yell

3

For More Information

- Hadly, Elizabeth. 1995. *Evolution, ecology, and taphonomy of late-Holocene mammals from Lamar Cave, Yellowstone National Park, Wyoming*. Ph.D. dissertation. UCalfornia, Berkeley.
- Hadly, E. A. 1999. Fidelity of terrestrial vertebrate fossils to a modern ecosystem. *Palaeogeography, Palaeoclimatology, Palaeoecology* 149(1999): 389–409.
- Hadly, E.A. 1990. *Late holocene mammalian fauna of Lamar Cave and its implications for ecosystem dynamics in Yellowstone National Park, Wyoming*. Master's thesis. Northern Arizona University.

Websites

volcanoes.usgs.gov/yvo
www.usgs.gov/corecast
www.seis.utah.edu
Additional information available on numerous other websites.

Videos

The Complete Yellowstone
Yellowstone: Imprints of Time
Yellowstone: A Symphony of Fire and Ice