

Table of Contents

1

2

3

4 **Executive Summary – page 2**

5 **Introduction – page 19**

6 **Chapters:**

7

8 **1. Decision Support for Agricultural Efficiency – page 21**

9 **2. Decision Support for Air Quality – page 34**

10 **3. Decision Support for Assessing Hybrid Renewable Energy**
11 **Systems – page 54**

12 **4. Decision Support for Public Health – page 81**

13 **5. Decision Support for Water Resources – page 93**

14

15 **Appendix A: References by Chapter – page 112**

16 **Appendix B: List of Figures by Chapter – page 145**

17 **Appendix C: Glossary, Acronyms, Symbols &, Abbreviations – page 147**

18 **“Uses and Limitations of Observations, Data, Forecasts, and Other Projections**
19 **in Decision Support for Selected Sectors and Regions”**

20
21 **(Climate Change Science Program, Synthesis and Assessment Product [SAP] 5.1)**

22
23 **Executive Summary**

24
25 Earth information—the diagnostics of Earth’s climate, water, air, land, and other dynamic processes—is essential for
26 our understanding of humankind’s relationship to our natural resources and our environment. Earth information can
27 inform our scientific knowledge, our approach to resource and environmental management and regulation, and our
28 stewardship of the planet for future generations. New data sources, new ancillary and complementary technologies in
29 hardware and software, and ever-increasing modeling and analysis capabilities characterize the current and prospective
30 states of Earth science and are a harbinger of its promise. A host of Earth science data products is enabling a revolution
31 in our ability to understand climate and its anthropogenic and natural variations. Crucial to this relationship, however, is
32 understanding and improving the integration of Earth science information in the activities that support decisions
33 underlying national priorities: ranging from homeland security and public health to air quality and natural resource
34 management.

35
36 Also crucial is the role of Earth information in improving our understanding of the processes and effects of climate as it
37 influences or is influenced by actions taken in response to national priorities. Global change observations, data,
38 forecasts, and projections are integral to informing climate science.

39
40 The Synthesis and Assessment Product (SAP), “Uses and Limitations of Observations, Data, Forecasts, and Other
41 Projections in Decision Support for Selected Sectors and Regions” (SAP 5.1), examines the current and prospective
42 contributions of Earth science information in decision support activities and their relationship to climate change science.
43 The SAP contains a characterization and catalog of observational capabilities in an illustrative set of decision support
44 activities. It also contains a description of the challenges and promise of these capabilities and discusses the interaction

45 between users and producers of information (including the role, measurement, and communication of uncertainty and
46 confidence levels associated with decision support outcomes and their related climate implications).

47

48 **Decision Support Tools and Systems**

49 In 2002, the National Aeronautics and Space Administration (NASA) formulated a conceptual framework in the form of
50 a flow chart (Figure 1) to characterize the link between Earth science data and their potential contribution to resource
51 management and public policy. The framework begins with Earth observations, including measurements made *in situ*
52 and from airborne and space-based instruments. These data are input into Earth system models that simulate the
53 dynamic processes of land, the atmosphere, and the oceans. These models lead in turn to predictions and forecasts to
54 inform decision support tools (DST).

55

56 In this framework, DSTs are typically computer-based models assessing such phenomena as resource supply, the status
57 of real-time events (e.g., forest fires and flooding), or relationships among environmental conditions and other scientific
58 metrics (i.e., water-borne disease vectors and epidemiological data). These tools use data, concepts of relations among
59 data, and analysis functions to allow analysts to build relationships—including spatial, temporal, and process-based—
60 among different types of data, merge layers of data, generate model outcomes, and make predictions or forecasts.

61 Decision support tools are an element of the broader decision making context or Decision Support System (DSS). DSSs
62 include not just computer tools but the institutional, managerial, financial, and other constraints involved in the
63 decision-making process.

64

65 The outcomes in these decision frameworks are intended to enhance our ability to manage resources (management of
66 public lands and measurements for air quality and other environmental regulatory compliance) and evaluate policy
67 alternatives (as promulgated in legislation or regulatory directives) affecting local, state, regional, national, or even
68 international actions. To be exact, for a variety of reasons, many decisions are not based on data or models. In some
69 cases, formal modeling is not appropriate, timely, or feasible for all decisions. But among decisions that are influenced
70 by this information, the flow chart (Figure 1) characterizes a systematic approach for science to be connected to
71 decision processes.

72 For purposes of providing an organizational framework, the CCSP provides additional description of decision support:

73

74 In the context of activities within the CCSP framework, decision-support resources, systems, and
75 activities are climate-related products or processes that directly inform or advise stakeholders in order
76 to help them make decisions. These products or processes include analyses and assessments,
77 interdisciplinary research, analytical methods (including scenarios and alternative analysis
78 methodologies), model and data product development, communication, and operational services that
79 provide timely and useful information to decision makers, including policymakers, resource
80 managers, planners, government officials, and other stakeholders. (*“Our Changing Planet,” CCSP*
81 *FY2007, Chapter 7, p. 155*).

82

83 **Our Approach**

84 Our approach to this SAP has involved two overall tasks. The first task defines and describes an illustrative set of DSTs
85 in areas selected from a number of areas deemed nationally important by NASA and also included in societal benefit
86 areas identified by the intergovernmental Group on Earth Observations (GEO) in leading an international effort to build
87 a Global Earth Observation System of Systems (GOESS) (see Tables 1 and 2).

88

89

90 The areas we have chosen as our case studies are air quality, agricultural efficiency, energy management, water
91 management, and public health. As required by the *SAP 5.1 Prospectus*, in the case studies we:

92

- 93 • explain the observational capabilities that are currently or potentially used in these tools;
- 94 • identify the agencies and organizations responsible for their development, operation, and maintenance;
- 95 • characterize the nature of interaction between users and producers of information in delivering accessing and
96 assimilating information;

- 97 • discuss sources of uncertainty associated with observational capabilities and the decision tools and how they
98 are conveyed in decision support context and to decision makers; and
- 99 • describe relationships between the decision systems and global change information, such as whether the tools
100 at present contribute or in the future could contribute to climate-related predictions or forecasts.

101

102 Because our purpose in this first task is to offer case studies by way of illustration rather than a comprehensive
103 treatment of all DSTs in all national applications, in our second task we have taken steps to catalog other DSTs which
104 use or may use, or which could contribute to, forecasts and projections of climate and global change. The catalog is an
105 exciting first step toward an ever-expanding inventory of existing and emerging DSTs. The catalog can be maintained
106 on-line for community input, expansion, and updating to provide a focal point for information about the status of DSTs
107 and how to access them.

108

109 The information in this report is largely from published literature and interviews with the sponsors and stakeholders of
110 the decision processes, as well as publications by and interviews with the producers of the scientific information used in
111 the tools.

112

113 **Our Case Studies**

114 We illustrate the following DSTs:

- 115 1. The Production Estimate and Crop Assessment Division and its Crop Condition Data Retrieval and Evaluation
116 (PECAD/CADRE) system of the US Department of Agriculture, Foreign Agricultural Service (FAS).
117 PECAD/CADRE is the world's most extensive and longest running (over two decades) operational user of
118 remote sensing for evaluation of worldwide agricultural productivity.
- 119 2. The Community Multiscale Air Quality (CMAQ) modeling system of the US Environmental Protection
120 Agency (EPA). CMAQ is a widely used, US continental/regional/urban-scale air quality decision support tool.
- 121 3. The Hybrid Optimization Model for Electric Renewables (HOMER), a micropower optimization model of the
122 US Department of Energy's National Renewable Energy Laboratory (NREL). HOMER is used around the
123 world to optimize deployment of renewable energy technologies.

124 4. Decision Support System to Prevent Lyme Disease (DDSPL) of the US Centers for Disease Control and
125 Prevention (CDC) and Yale University. DDSPL seeks to prevent the spread of the most common vector-borne
126 disease, Lyme disease, of which there are tens of thousands of cases annually in the US

127 5. RiverWare, developed by the University of Colorado-Boulder's Center for Advanced Decision Support for
128 Water and Environmental Systems (CADSWES) in collaboration with the Bureau of Reclamation, Tennessee
129 Valley Authority, and the Army Corps of Engineers. RiverWare is a hydrologic or river basin modeling
130 system that integrates features of reservoir systems, such as recreation, navigation, flood control, water
131 quality, and water supply, in a basin management tool with power system economics to provide basin
132 managers and electric utilities a method of planning, forecasting, and scheduling reservoir operations.

133

134 Taken together, these DSTs demonstrate a rich variety of applications of observations, data, forecasts, and other
135 predictions. In four of our studies, agricultural efficiency, air quality, water management, and energy management, the
136 DSTs have become well established as a basis for public policy decision making. In the case of public health, our lead
137 author points out reasons why direct applications of Earth observations to public health have tended to lag behind these
138 other applications and thus is a relatively new application area. He also reminds us that management of air quality,
139 agriculture, water, and energy—in and of themselves—have implications for the quality of public health. The DST he
140 selects is a new, emerging tool intended to assist in prevention of the spread of infectious disease.

141

142 Our selection also varies in the geographic breadth of application, illustrating how users of these tools tailor them to
143 relevant regions of analysis and how, in some cases, the geographic coverage of the tools carries over to their
144 requirements for observations. For instance, PECAD/CADRE is used for worldwide study of agricultural productivity
145 and has data requirements of wide geographic scope, HOMER can be used for renewable energy optimization
146 throughout the world, and DDSPL focuses on the eastern, upper Midwest, and West Coast portions of the US. CMAQ is
147 used to predict air quality for the contiguous US as well as regions and urban locales. RiverWare provides basin
148 managers and electric utilities a method of planning, forecasting, and scheduling reservoir operations.

149

150 With the exception of DDSPL, none of the DSTs we considered for potential selection, nor those we discuss in this
151 report, have to date made extensive use of climate change information or been used to study the effect of a changing

152 climate. However, in all cases, the developers and users of these DSTs fully recognize their applicability to climate
153 change science. In the discussion of the five DSTs presented in this SAP, the authors describe how climate data and/or
154 predictions might be used in these DSTs so that long-range decisions and planning might be accomplished.

155

156 **Overview of the Chapters**

157 We next summarize the case studies. For each case study, we describe the DST and its data sources, highlight potential
158 uses as well as limits of the DSTs, note sources of uncertainty in using the tools, and finally, discuss the link between
159 the DST and climate change and variability. After our summary, we offer general observations about similarities and
160 differences among the studies.

161

162 *Agricultural Efficiency*: The Production Estimate and Crop Assessment Division (PECAD) of the US Department of
163 Agriculture, FAS is the world's most extensive and longest running operational user of remote sensing data for
164 evaluation of worldwide agricultural productivity. PECAD supports the FAS mission to collect and analyze global crop
165 intelligence information and provide periodic estimates used to inform official USDA forecasts for the agricultural
166 market, including farmers; agribusiness; commodity traders and researchers; and federal, state, and local agencies.
167 PECAD is often referred to as PECAD/CADRE with one of its major automated components known as the Crop
168 Condition Data Retrieval and Evaluation (CADRE) geospatial database management system. Of all the DSTs we
169 consider in this report, CADRE has the oldest pedigree as the operational outcome of two early, experimental earth
170 observation projects during the 1970s and 1980s: the Large Area Crop Inventory Experiment (LACIE) and the
171 Agriculture and Resources Inventory Surveys through Aerospace Remote Sensing (AgRISTARS).

172

173 Sources of data for CADRE include a large number of weather and other earth observations from US, European,
174 Japanese, and commercial systems. PECAD combines these data with crop models, a variety of GIS tools, and a large
175 amount of contextual information, including official government reports, trade and news sources, and on-the-ground
176 reports from a global network of embassy attaches and regional analysts.

177

178 Potential future developments in PECAD/CADRE could include space-based observations of atmospheric carbon
179 dioxide (CO₂) measurements and measurement of global sea surface salinity to improve understanding of the links

180 between the water cycle, climate, and oceans. Other opportunities for enhancing PECAD/CADRE include
181 improvements in predictive modeling capabilities in weather and climate.

182
183 One of the largest technology gaps in meeting PECAD requirements is the practice of designing earth observation
184 systems for research rather than operational use, limiting the ability of PECAD/CADRE to rely on data sources from
185 non-operational systems. PECAD analysts require dependable inputs, implying use of operational systems that ensure
186 continuous data streams and that minimize vulnerability to component failure through redundancy.

187
188 Sources of uncertainty can arise at each stage of analysis, from the accuracy of data inputs to the assumptions in
189 modeling. PECAD operators have been able to benchmark, validate, verify, and then selectively incorporate additional
190 data sources and automated decision tools by way of detailed engineering reviews. Another aspect of resolving
191 uncertainty in PECAD is the extensive use of a convergence methodology to assimilate information from regional field
192 analysts and other experts. This convergence of evidence analysis seeks to reconcile various independent data sources to
193 achieve a level of agreement to minimize estimate error.

194
195 The relationship between climate and agriculture is complex, as agriculture is influenced not only by a changing
196 climate, but agricultural practices themselves are a contributory factor (e.g., in affecting land use and influencing
197 carbon fluxes. At present, PECAD is not directly used to address these dimensions of the climate-agriculture
198 interaction. However, many of the data inputs for PECAD are climate-related, thereby enabling PECAD to inform
199 understanding of agriculture as a “recipient” of climate-induced changes. For instance, observing spatial and geographic
200 trends in the output measures from PECAD can contribute to understanding how the agricultural sector is responding to
201 a changing climate. Likewise, trends in PECAD’s measures of the composition and production of crops could shed light
202 on the agricultural sector as a “contributor” to climate change (for instance, in terms of greenhouse gas emissions or
203 changes in soil that may affect the potential for agricultural soil carbon sequestration). PECAD may also be influenced
204 by, as well as a barometer of climate-induced changes in land use, such as conversion from food production to biomass
205 fuel production.

206

207 **Air Quality:** The EPA CMAQ modeling system has been designed to approach air quality by including state-of-the-
208 science capabilities for modeling tropospheric ozone, fine particles, toxics, acid deposition, and visibility degradation.
209 CMAQ is used to guide the development of air quality regulations and standards and to create state implementation
210 plans for managing air emissions. CMAQ also can be used to evaluate longer-term as well as short-term transport from
211 localized sources and to perform simulations using downscaled regional climate from global climate change scenarios.

212
213 The CMAQ modeling system contains three types of modeling components: a meteorological modeling system for the
214 description of atmospheric states and motions, emission models for man-made and natural emissions that are injected
215 into the atmosphere, and a chemistry-transport modeling system for simulation of the chemical transformation and fate.
216 Inputs for CMAQ, and their associated regional meteorological model, mesoscale model version 5 (MM5), can include,
217 but are not limited to, the comprehensive output from a general circulation model, anthropogenic and biogenic
218 emissions, description of wildland fires, land use and demographic changes, and meteorological and atmospheric
219 chemical species measurements by *in-situ* and remote sensing platforms, including satellites and aircraft.

220
221 CMAQ can be used to study questions such as: How will present and future emission changes affect attainment of air
222 quality standards? Will present and future emissions and/or climate/meteorological changes affect the frequency and
223 magnitudes of high pollution events? How will land use changes due to urbanization and global warming affect air
224 quality? How does long-range air pollution from other regions affect US air quality? How will changes in the long-
225 range transport due to the climate change affect air quality? How does wildland fire affect air quality and will climate
226 change affect wildland fire and subsequently air quality? How sensitive are the air quality predictions to changes in both
227 anthropogenic and biogenic emissions?

228
229 **Energy Management:** HOMER is a micropower optimization model of the US Department of Energy's NREL.
230 HOMER is able to calculate emission reductions enabled by replacing diesel-generating systems with renewable energy
231 systems in a micro-grid or grid-connected configuration. HOMER helps the user design grid-connected and off-grid
232 renewable energy systems by performing a wide range of design scenarios. HOMER can be used to address questions
233 such as: Which technologies are most cost-effective? What happens to the economics if the project's costs or loads

234 change? Is the renewable energy resource adequate for the different technologies being considered to meet the load?
235 HOMER does this by finding the least-cost combination of components that meet electrical and thermal loads.

236
237 The earth observation information serving as input to HOMER is centered on wind and solar resource assessments
238 derived from a variety of sources. Wind data include surface and upper air station data, satellite-derived ocean and ship
239 wind data, and digital terrain and land cover data. Solar resource data include surface cloud, radiation, aerosol optical
240 depth, and digital terrain and land cover data from both *in-situ* and remote sensing sources.

241
242 All of the input data for HOMER can have a level of uncertainty attached to them. HOMER allows the user to perform
243 sensitivity tests on one or more variables and has graphical capabilities to display these results to inform decision
244 makers. As a general rule, the error in estimating the performance of a renewable energy system over a year is roughly
245 linear to the error in the input resource data.

246
247 One of the largest challenges in HOMER is the absence of direct or *in-situ* solar and wind resource measurements at
248 specific locations to which HOMER is applied. In addition, in many cases, values are not based on direct measurement
249 at all but are approximations based on the use of algorithms to convert a signal into the parameter of interest as is the
250 case with most satellite-derived data products. For example, satellite-derived ocean wind data are not based on direct
251 observation of the wind speed above the ocean surface but from an algorithm that infers wind speed based on wave
252 height observations. Observations of aerosol optical depths (for which considerable research is underway) can be
253 complicated by irregular land-surface features that complicate the application of algorithms for satellite-derived
254 measures.

255
256 For renewable energy resource mapping, improved observations of key weather parameters (for instance, wind speed
257 and direction at various heights above the ground, particularly at the hub height of wind energy turbine systems, and
258 over the open oceans at higher and higher spatial resolutions, and improved ways of differentiating snow cover and
259 bright reflecting surfaces from clouds) will be of value to the renewable energy community. New, more accurate
260 methods of related parameters, such as aerosol optical depth, would also improve the resource data.

261

262 The relationship between HOMER and global change information is largely by way of the dependence of renewable
263 energy resource input measurements on weather and local climate conditions. Although HOMER was not designed to
264 be a climate-related management decision-making tool, by optimizing the mix of hybrid renewable energy technologies
265 for meeting load conditions, HOMER also enables users to respond to climate change and variability in their energy
266 management decisions. HOMER could be used to evaluate how renewable energy systems can be used cost-effectively
267 to displace fossil-fuel-based systems.

268
269 **Public Health:** The DDSPL is operated by the US CDC and Yale University to address questions related to the likely
270 distribution of Lyme disease east of the 100th meridian, where most cases occur. Lyme disease is the most common
271 vector-borne disease in the US, with tens of thousands of cases annually. Most human cases occur in the Eastern and
272 upper Midwest portions of the US, although there is a secondary focus along the West Coast. Vector-borne diseases are
273 those in which parasites are transmitted among people or from wildlife to people by insects or arthropods (as vectors,
274 they do not themselves cause disease). The black-legged tick is typically the carrier of the bacteria causing Lyme
275 disease.

276
277 Early demonstrations during the 1980s showed the utility of earth observations for identifying locations and times that
278 vector-borne diseases were likely to occur, but growth of applications has been comparatively slow. Earth observing
279 instruments have not been designed to monitor disease risk; rather, data gathered from these platforms are “scavenged”
280 for public health risk assessment. DDSPL uses satellite data and derived products, such as land cover together with
281 meteorological data and census data, to characterize statistical predictors of the presence of black-legged ticks. The
282 model is validated by field surveys. The DDSPL is thus a means of setting priorities for the likely geographic extent of
283 the vector; the tool does not at present characterize the risk of disease in the human population.

284
285 Future use of DDSPL partly depends on whether the goal of disease prevention or the goal of treatment drives public
286 health policy decisions. In addition, studies have shown that communication to the public about the risk in regions with
287 Lyme disease often fails to reduce the likelihood of infection. Use of the DDSPL is also limited by restrictions on the
288 dissemination of detailed information on the distribution of human disease. The role of improved Earth science data is
289 unclear in terms of improving the performance of DDSPL because at present the system has a level of accuracy deemed

290 “highly satisfactory.” Future use may instead require a model of sociological/behavioral influences among the
291 population.

292

293 Standard statistical models and in-field validation are used to assess the uncertainty in decision making with DDSPL.

294 The accuracy of clinical diagnoses also influences the ultimate usefulness of DDSPL as an indicator tool to characterize
295 the geographic extent of the vectors.

296

297 The DDSPL is one of the few public health DSTs that has explicitly evaluated the effects of climate variability. Using
298 outputs of a Canadian climate change model, study has shown that with warming global mean temperatures by the year
299 2050 to 2080 the geographic range of the tick vector will decrease at first, with reduced presence in the southern
300 boundary, and then expand into Canada and the central region of North America where it now absent. The range also
301 moves away from population concentrations.

302

303 **Water Management:** RiverWare was developed and is maintained by CADSWES in collaboration with the Bureau of
304 Reclamation, Tennessee Valley Authority, and the Army Corps of Engineers. It is a river basin modeling system that
305 integrates features of reservoir systems, such as recreation, navigation, flood control, water quality, and water supply in
306 a basin management tool, with power system economics to provide basin managers and electric utilities a method of
307 planning, forecasting, and scheduling reservoir operations. RiverWare uses an object-oriented software engineering
308 approach in model development. The object oriented software-modeling strategy allows computational methods for new
309 processes, additional controllers for providing new solution algorithms, and additional objects for modeling new
310 features to be added easily to the modeling system. RiverWare is data intensive in that a specific river/reservoir system
311 and its operating policies must be characterized by the data supplied to the model. This allows the models to be
312 modified as new features are added to the river/reservoir system and/or new operating policies are introduced. The
313 data-intensive feature allows the model to be used for water management in most river basins.

314

315 Riverware is menu driven through a graphical user interface (GUI). The basin topology is developed through the
316 selection of a reservoir, reach, confluence, and other necessary objects and by entering the data associated with each
317 object manually or through importing files. Utilities within RiverWare provide a means to automatically execute many

318 simulations, to access data from external sources, and to export model results. Users also define operating policies
319 through the GUI as system constraints or rules for achieving system management goals (e.g., related to flood control,
320 water supplies, water quality, navigation, recreation, and power generation). The direct use of earth observations in
321 RiverWare is limited. Unlike traditional hydrologic models that track the transformation of precipitation (e.g., rain and
322 snow) into soil moisture and streamflow, RiverWare uses supplies of water to the system as input data. These data are
323 derived from a hydrologic model where direct use of earth observations can be and have been made. Application of
324 RiverWare is limited by the specific implementation defined by the user and by the quality of the input data. It has
325 tremendous flexibility in the kinds of data it can use, but long records of data are required to overcome the issue of data
326 non-stationarity.

327

328 The specific application of RiverWare in the context of mid- or long-range planning for a specific river basin will reflect
329 whether decisions may rely on global change information. For mid-range planning of reservoir operations,
330 characterization and projections of interannual and decadal-scale climate variability (e.g., monitoring, understanding,
331 and predicting interannual climate phenomena such as the El Nino-Southern Oscillation) are important. For long-term
332 planning, global warming has moved from the realm of speculation to general acceptance. The impacts of global
333 warming on water resources, and their implications for management, have been a major focus in the assessments of
334 climate change. The estimates of potential impacts of climate change on precipitation have been mixed, leading to
335 increasing uncertainty about the reliability of future water supplies.

336

337 **General Observations**

338 Application of all of the DSTs involves a variety of input data types, all of which have some degree of uncertainty in
339 terms of their accuracy. The amount of uncertainty associated with resource data can depend heavily on how the data
340 are obtained. Quality *in-situ* measurements of wind and solar data suitable for application in HOMER are can have
341 uncertainties of less than $\pm 3\%$ of true value; however, when estimation methods are required, such as the use of earth
342 observations, modeling, and empirical techniques, uncertainties can be as much as $\pm 10\%$ or more. The DSTs address
343 uncertainty by allowing users to perform sensitivity tests on variables. With the exception of HOMER, a significant
344 amount of additional traditional on-the-ground reports are a critical component. In the case of PECAD/CADRE,
345 uncertainty is resolved in part by extensive use of a convergence methodology to assimilate information from regional

346 field analysts and other experts. This brings a large amount of additional information to PECAD/CADRE forecasts, well
347 beyond the automated outputs of DSTs. In RiverWare, streamflow and other hydrologic variables respond to
348 atmospheric factors such as precipitation, and obtaining quality precipitation estimates is a formidable challenge,
349 especially in the western US where orographic effects produce large spatial variability and where there is a scarcity of
350 real-time precipitation observations and poor radar coverage.

351

352 In terms of their current or prospective use of climate change predictions or forecasts as DST *inputs*, or the
353 contributions of DST *outputs* to understanding, monitoring, and responding to a changing climate, the status is mixed.
354 DDSPL is one of the few public health decision support tools that has explicitly evaluated the potential impact of
355 climate change scenarios on an infectious disease system. None of the other DSTs at present is directly integrated with
356 climate change measurements, but all of them can and may in the future take this step. PECAD/CADRE's assessment of
357 global agricultural production will certainly be influenced by observations and forecasts of climate change and
358 variability as model inputs, just as the response of the agricultural sector to a changing climate will feedback into
359 PECAD/CADRE production estimates. HOMER's renewable energy optimization calculations will be directly affected
360 by climate-related changes in renewable energy resource supplies and will enhance our ability to adapt to climate-
361 induced changes in energy management and forecasting. Air quality will definitely be affected by global climate
362 change. The ability of CMAQ to predict those affects is conditional on acquiring accurate predictions of the
363 meteorology under the climate change conditions that will take place in the US and accurate emission scenarios for the
364 future. Given these inputs to CMAQ, reliable predictions of the air quality and their subsequent health affects can be
365 ascertained. It was noted that there is great difficulty in integrating climate change information into RiverWare and
366 other such water management models. The multiplicity of scenarios and vague attribution of their probability for
367 occurrence, which depends on feedback among social, economic, political, technological, and physical processes,
368 complicates conceptual integration of climate change impacts assessment results in a practical water management
369 context. Furthermore, the century timescales of climate change exceed typical planning and infrastructure design
370 horizons in water management.

371

372 **Audience and Intended Use**

373 The *CCSP SAP 5.1 Prospectus* describes the audience and intended use of this report:

374

375 This synthesis and assessment report is designed to serve decision makers and stakeholder
376 communities interested in using global change information resources in policy, planning, and other
377 practical uses. The goal is to provide useful information on climate change research products that
378 have the capacity to inform decision processes. The report will also be valuable to the climate change
379 science community because it will indicate types of information generated through the processes of
380 observation and research that are particularly valuable for decision support. In addition, the report will
381 be useful for shaping the future development and evaluation of decision-support activities, particularly
382 with regard to improving the interactions with users and potential users.


383

384 There are a number of national and international programs focusing on the use of Earth observations
385 and related prediction capacity to inform decision support tools (see Table 3, “Related National and
386 International Activities”). These programs both inform and are informed by the CCSP and are
387 recognized in the development of this product. (*CCSP Synthesis and Assessment Product 5.1,*
388 *Prospectus for “Uses and Limitations of Observations, Data, Forecasts, and Other Projections in*
389 *Decision Support for Selected Sectors and Regions,” 28 February 2006)*

390

391

392



393

394

395 Figure 1: The flow of information associated with decision support in the context of variability and change in climate
 396 and related systems (Source: CCSP Product 5.1 Prospectus, Appendix D).

397

398 Table 1: List of NASA National Applications Areas (*Appendix B, CCSP SAP 5.1 Prospectus*).

399

400

Nationally Important Applications	Nationally Important Applications
Agricultural Efficiency	Ecological Forecasting
Air Quality	Energy Management
Aviation	Homeland Security
Carbon Management	Invasive Species
Coastal Management	Public Health
Disaster Management	Water Management

401

402

403 Table 2. Societal benefit areas identified by the Group on Earth Observations (GEO) for the Global Earth Observations

404 System of Systems (GEOSS) (http://www.earthobservations.org/about/about_GEO.html) (accessed May 2007)

405

406

GEOSS Socio-Benefit Area Keywords	GEOSS Socio-Benefit Area Descriptions
Health	Understanding environmental factors affecting human health and well-being
Disasters	Reducing loss of life and property from natural and human-induced disasters
Forecasts	Improving weather information, forecasting, and warning
Energy	Improving management of energy resources
Water	Improving water resource management through better understanding of the water cycle
Climate	Understanding, assessing, predicting, mitigating, and adapting to climate variability and change

Agriculture	Supporting sustainable agriculture and combating desertification
Ecology	Improving the management and protection of terrestrial, coastal, and marine ecosystems

407

408

409 Table 3. References to Related National and International Activities (*Source: Appendix C, CCSP SAP 5.1 Prospectus*)

410

Priority	National	International
Climate Change	Climate Change Science Program and Climate Change Technology Program	Intergovernmental Panel on Climate Change and World Climate Research Programme
Global Earth Observations	NSTC CENR U.S. Interagency Working Group on Earth Observations	Group on Earth Observations (GEO)
Weather	U.S. Weather Research Program (USWRP)	World Meteorological Organization
Natural Hazards	NSTC CENR Subcommittee on Disaster Reduction	International Strategy for Disaster Reduction
Sustainability	NSTC CENR Subcommittee on Ecosystems	World Summit on Sustainable Development
E-Government	Geospatial One-Stop and the Federal Geographic Data	World Summit on the Information Society

411

412

413 **“Uses and Limitations of Observations, Data, Forecasts, and Other Projections**
414 **in Decision Support for Selected Sectors and Regions”**

415

416 **(Climate Change Science Program, Synthesis and Assessment Product [SAP] 5.1)**

417

418

419

Introduction

420

421 This Synthesis and Assessment Product (SAP), “Uses and Limitations of Observations, Data, Forecasts, and Other
422 Projections in Decision Support for Selected Sectors and Regions” (SAP 5.1), examines the current and prospective
423 contribution of Earth science information/data in decision support activities and their relationship to climate change
424 science. The SAP contains a characterization and catalog of observational capabilities in an illustrative set of decision
425 support activities. It also contains a description of the challenges and promises of these capabilities and discusses the
426 interaction between users and producers of information, including the role, measurement, and communication of
427 uncertainty and confidence levels associated with decision support outcomes and their related climate implications.

428

429 The organizing basis for the chapters in this SAP is the decision support tools (DST), which are typically computer-
430 based models assessing such phenomena as resource supply, the status of real-time events (e.g., , forest fires and
431 flooding), or relationships among environmental conditions and other scientific metrics (for instance, water-borne
432 disease vectors, and epidemiological data). These tools use data, concepts of relations among data, and analysis
433 functions to allow analysts to build relationships—including spatial, temporal, and process-based—among different
434 types of data, merge layers of data, generate model outcomes, and make predictions or forecasts. DSTs are an element
435 of the broader decision-making context, the decision support system (DSS). DSSs include not just computer tools but
436 also the institutional, managerial, financial, and other constraints involved in decision making.

437

438 Our approach to this SAP is to define and describe an illustrative set of DSTs in areas selected from topics deemed
439 nationally important and included in societal benefit areas identified by the intergovernmental Group on Earth

440 Observations in leading an international effort to build a Global Earth Observation System of Systems. The areas we
441 have chosen as our focus are air quality, agricultural efficiency, energy management, water management, and public
442 health. The DSTs we illustrate are:

- 443 1. The Production Estimate and Crop Assessment Division and its Crop Condition Data Retrieval and Evaluation
444 system (PECAD/CADRE) of the US Department of Agriculture, Foreign Agricultural Service (FAS).
445 PECAD/CADRE is the world's most extensive and longest running (over two decades) operational user of
446 remote sensing for evaluation of worldwide agricultural productivity.
- 447 2. The Community Multiscale Air Quality (CMAQ) modeling system of the US Environmental Protection
448 Agency (EPA). CMAQ is a widely used, US continental/regional/urban-scale air quality decision support tool.
- 449 3. The Hybrid Optimization Model for Electric Renewables (HOMER), a micropower optimization model of the
450 US Department of Energy's National Renewable Energy Laboratory (NREL). HOMER is used around the
451 world to optimize deployment of renewable energy technologies.
- 452 4. The Decision Support System to Prevent Lyme Disease (DDSPL) of the US Centers for Disease Control and
453 Prevention (CDC) and Yale University. DDSPL seeks to prevent the spread of the most common vector-borne
454 disease, Lyme disease, of which there are tens of thousands of reported cases annually in the United States.
- 455 5. RiverWare, developed by the University of Colorado-Boulder's Center for Advanced Decision Support for
456 Water and Environmental Systems (CADSWES) in collaboration with the Bureau of Reclamation, Tennessee
457 Valley Authority, and the Army Corps of Engineers, is a hydrologic or river basin modeling system that
458 integrates features of reservoir systems, such as recreation, navigation, flood control, water quality, and water
459 supply, in a basin management tool with power system economics to provide basin managers and electric
460 utilities a method of planning, forecasting, and scheduling reservoir operations.

461
462 Taken together, these DSTs demonstrate a rich variety of applications of observations, data, forecasts, and other
463 predictions. In four of our studies—agricultural efficiency, air quality, water management, and energy management—
464 the DSTs have become well established as a basis for public policy decision making. In the case of public health, our
465 lead author points out reasons why direct applications of Earth observations to public health have tended to lag behind
466 these other applications and thus is a relatively new applications area. He also reminds us that management of air

467 quality, agriculture, water, and energy—in and of themselves—have implications for the quality of public health. The
468 DST selected for public health is a new and emerging tool intended to assist in prevention of the spread of infectious
469 disease.

470

471 With the exception of DDSPL, none of the DSTs we considered for potential selection, nor those we discuss in this
472 report, have to date made extensive use of climate change information or been used to study the effect of a changing
473 climate. However, in all cases, the developers and users of these DSTs fully recognize their applicability to climate
474 change science. In the discussion of the five DSTs presented in this SAP, the authors describe how climate data and/or
475 predictions might be used in these DSTs so that long-range decisions and planning might be accomplished.

476

Chapter 1

Decision Support for Agricultural Efficiency

Lead Author: Molly K. Macauley

1. Introduction

The efficiency of agriculture has been one of the most daunting challenges confronting mankind in its need to manage natural resources within the constraints of weather, climate, and other environmental conditions. Defined as maximizing output per unit of input, agricultural efficiency reflects a complex relationship among factors of production (including seed, soil, human, and physical capital) and the exogenous influence of nature (such as temperature, sunlight, weather, and climate). The interaction of agricultural activity with the environment creates another source of interdependence, (e.g., the effect on soil and water from applications of pesticides, fungicides, and fertilizer). Agricultural production has long been a large component of international trade and of strategic interest as an indicator of the health and security of nations.

The relationship between climate change and agriculture is complex. A changing climate can influence agricultural practices (e.g., climate-induced changes in patterns of rainfall could lead to changes in these practices). Agriculture is not only influenced by a changing climate, but agricultural practices themselves are a contributory factor through emissions of greenhouse gases and influences on fluxes of carbon through photosynthesis and respiration. In short, agriculture is both a contributor to and a recipient of the effects of a changing climate (Rosen Zweig, 2003; National Assessment Synthesis Team, 2004).

The use of Earth observations by the agricultural sector has a long history. The Large Area Crop Inventory Experiment (LACIE), jointly sponsored by the US National Aeronautics and Space Administration (NASA), the US Department of Agriculture (USDA), and the National Oceanic and Atmospheric Administration (NOAA) conducted from 1974 to 1978 demonstrated the potential for satellite observations to make accurate, extensive, and repeated surveys for global crop forecasts. LACIE used observations from the Landsat series of multi-spectral scanners on sun-synchronous satellites. The Agriculture and Resources Inventory Surveys through Aerospace Remote Sensing

Do Not Cite or Quote

505 (AgRISTARS) followed LACIE and extended the use of satellite observations to include early warning of production
506 changes, inventory and assessment of renewable resources, and other activities (Congressional Research Service, 1983;
507 National Research Council, 2007; Kaupp *et al.*, 2005). Today these data are used by agencies of the federal government,
508 commodity trading companies, farmers, relief agencies, other governments, and essentially anyone with an interest in
509 crop production at a global scale.

510

511 An approach, among others, to increasing agricultural efficiency is to expand and enhance uses of Earth
512 observation data for (1) policy and resource management decision support, (2) monitoring and measuring climate
513 change affects, and (3) providing policy and resource climate change decision support. The foremost example of the
514 application of Earth observations in agriculture is found in the USDA's crop-monitoring decision-support system, the
515 Production Estimates and Crop Assessment Division (PECAD) of the USDA's Foreign Agricultural Service (FAS).
516 (Reorganization at USDA finds the PECAD functionality, but not the name, residing within the USDA's FAS as part of
517 the Office of Global Analysis, Impact Analysis Division, International Production Assessment
518 [USDA/FAS/OGA/IAD/IPA]). PECAD is now the world's most extensive and longest running (over two decades)
519 operational user of remote sensing data for evaluation of worldwide agricultural productivity (NASA, 2001). A
520 Description of the PECAD decision-support system, its functionality, its analysis style, how it deals with making
521 decisions under uncertainty, and its future uses form the basis of this chapter.

522

523 **2. Description of PECAD**

524 The USDA/FAS uses PECAD to analyze global agricultural production and crop conditions affecting planting,
525 harvesting, marketing, commodity export and pricing, drought monitoring, and food assistance. Access to and uses of
526 PECAD are largely by the federal government, rather than state and local governments, as a means of assessing regions
527 of interest in global agricultural production.

528 PECAD uses satellite data, worldwide weather data, and agricultural models in conjunction with FAS overseas
529 post reports, foreign government official reports, and agency travel observations to support decision making. FAS also
530 works closely with the USDA Farm Service Agency and the Risk Management Agency to provide early warning and
531 critical analysis of major crop events in the US. (FAS OnLine Crop Assessment at
532 http://www.fas.usda.gov/pecad2/crop_assmnt.html, accessed April 2007). FAS seeks to promote the security and

533 stability of the US food supply, improve foreign market access for U.S. agricultural products, provide reports on world
534 food security, and advise the US government on international food aid requirements. FAS bears the primary
535 responsibility for USDA's overseas activities: market development, international trade agreements and negotiations, and
536 the collection and analysis of statistics and market information. FAS also administers USDA's export credit guarantee
537 and food aid programs.

538 PECAD's Crop Condition Data Retrieval and Evaluation (CADRE) database management system, the
539 operational outcome of the LACIE and AgRISTARs projects, was one of the first geographic information systems (GIS)
540 designed specifically for global agricultural monitoring (Reynolds, 2001). CADRE is used to maintain a large satellite
541 imagery archive to permit comparative interpretation of incoming imagery with that of past weeks or years. The
542 database contains multi-source weather data and other environmental data that are incorporated as inputs for models to
543 estimate parameters such as soil moisture, crop stage, and yield. These models also indicate the presence and severity of
544 plant stress or injury. The information from these technologies is used by PECAD to produce, in conjunction with the
545 World Agricultural Outlook Board, official USDA foreign crop production estimates. (FAS OnLine Crop Assessment at
546 http://www.fas.usda.gov/pecad2/crop_assmnt.html, accessed April 2007)

547 Figure 1 (Kaupp *et al.*, 2005, p. 5) illustrates the global data sources and decision support tools for PECAD.
548 The left-hand portion of the figure shows sources of data for the CADRE geospatial DBMS. These inputs include
549 station data from the World Meteorological Organization and coarse resolution data from Meteosat, Scanning
550 Multichannel Microwave Radiometer (SSMR), and Geostationary Satellite (GOES). Meteosat, operated by the
551 European Organization for the Exploitation of Meteorological Satellites (EUMETSAT), provides visible and infrared,
552 weather-oriented imaging. The SSMR and its successor, the Special Sensor Microwave/Imager (SSM/I), are microwave
553 radiometric instruments in the US Air Force Defense Meteorological Satellite Program. Additional weather data come
554 from the US GOES program.

555 Medium resolution satellite data include Advanced Very High Resolution Radiometer (AVHRR)/NOAA,
556 Spot-Vegetation, and Terra/Aqua MODIS. AVHRR/NOAA, operated by NOAA, provides cloud cover and land, water,
557 and sea surface temperatures at approximately 1-km spatial resolution. The Systeme Pour L'Observation de la Terre
558 (SPOT) supplies commercial optical Earth imagery at resolutions from 2.5 to 20 meters (m); SPOT-Vegetation is a
559 sensor providing daily coverage at 1 km resolution. The NASA Moderate Resolution Imaging Spectroradiometers
560 (MODIS) on the Terra and Aqua satellites, part of the US Earth Observation System, show rapid biological and

561 meteorological changes at 250 to 1,000 m spatial resolution every two days. NASA's Global Inventory Modeling and
562 Mapping Studies (NASA/GIMMS) group processes data acquired from SPOT and Terra/Aqua MODIS.
563 NASA/GIMMS provides PECAD with cross-calibrated global time series of Normalized Difference Vegetation Index
564 maps from AVHRR and SPOT-Vegetation. Moderate-resolution Earth observation data are also used from the US
565 Landsat program.

566 Sources of high resolution and radar altimeter satellite data include SPOT, IKONOS, Poseidon, and Jason.
567 IKONOS is a commercial Earth imaging satellite providing spatial resolution of 1 and 4 m. Data from Poseidon and its
568 successor, Jason, provide lake and reservoir surface elevation estimates. Poseidon, part of the TOPEX/Poseidon
569 mission, and Jason-1, a follow-on mission, are joint ventures between NASA and the Centre National d'Etudes
570 Spatiales (CNES) using radar altimeters to map ocean surface topography (including sea surface height, wave height,
571 and wind speed above the ocean). These data enable analysts to assess drought or high water-level conditions within
572 some of the world's largest lakes and reservoirs to predict effects on downstream irrigation potential and inform
573 production capacity estimates (Birkett and Doorn, 2004; Kanarek, 2005). The assimilation of these data into PECAD is
574 described in detail in a recent systems engineering report (NASA, 2004b).

575 PECAD combines the satellite and climate data, crop models (along the bottom portion of the figure), a variety
576 of GIS tools, and a large amount of contextual information, including official government reports, trade and new
577 sources, and on-the-ground reports from a global network of embassy attaches and regional analysts. The integration
578 and analysis is attained by "convergence of evidence analysis" (Kaupp *et al.*, 2005). This convergence methodology
579 seeks to reconcile various independent data sources to achieve a level of agreement to minimize estimate error (NASA,
580 2004a).

581 The crop assessment products indicated along the right-hand side of the PECAD architecture in figure 1
582 represent the periodic global estimates used to inform official USDA forecasts. These products are provided to the
583 agricultural market, including farmers; agribusiness; commodity traders and researchers; and federal, state, and local
584 agencies. In addition to CADRE, other automated components include two features providing additional types of
585 information. The FAS Crop Explorer (middle of diagram) is a feature on the FAS Web site since 2002 (Kanarek, 2005).
586 Crop Explorer offers near-real-time global crop condition information based on satellite imagery and weather data from
587 the CADRE database and NASA/GIMMS. Thematic maps of major crop growing regions show vegetation health,
588 precipitation, temperature, and soil moisture. Time-series charts show growing season data for agro-meteorological

589 zones. For major agriculture regions, Crop Explorer provides crop calendars and crop areas. Through Archive Explorer,
590 PECAD provides access to an archive of moderate- to high-resolution data, allowing USDA users (access is controlled
591 by user name and password) to search an image database.

592

593 **3. Potential Future Use and Limits**

594 The most recent enhancements to PECAD/CADRE have included the integration and evaluation of MODIS,
595 Topex/Poseidon, and Jason-1 products (NASA, 2006a). Figure 2 summarizes the Earth system models, Earth
596 observations data, and the CADRE DBMS and characterizes their outputs. Several planned Earth observations missions
597 anticipated when this image was prepared (indicated in italics) show how PECAD/CADRE could incorporate new
598 opportunities, including those with additional land, atmosphere, and ocean observations. These would include space-
599 based observations of atmospheric carbon dioxide (CO₂) from the Orbiting Carbon Observatory (OCO) and
600 measurement of global sea surface salinity (Aquarius) to improve understanding of the links between the water cycle,
601 climate, and the ocean. Other opportunities for enhancing PECAD/CADRE could include improvements in predictive
602 modeling capabilities in weather and climate (National Aeronautics and Space Administration, 2006a).

603 In a recent evaluation report for PECAD, NASA has acknowledged that one of the largest technology gaps in
604 meeting PECAD requirements is the design of NASA systems for research purposes rather than for operational uses
605 (NASA, 2004a). PECAD analysts require dependable inputs, implying the use of operational systems that ensure
606 continuous data streams and that minimize vulnerability to component failure through redundancy. The report also
607 emphasizes that PECAD requires systems that deliver real-time or near-real-time data. Many NASA missions have
608 traded timeliness for experimental research or improvements in other properties of the information delivered.
609 Additionally, the report identifies several potential Earth science data streams that have not yet been addressed,
610 including water balance, the radiation budget (including solar and long wave radiation flux), and elevation, and
611 expresses concern about the potential continuity gap between Landsat 7 and the Landsat Data Continuity Mission.

612 A 2006 workshop convened at the United Nations Food and Agriculture Organization (FAO) by the Integrated
613 Global Observations of Land (IGOL) team identified priorities for agricultural monitoring during the next 5 to 10 years
614 as part of the emerging GEOSS. In summary, the meeting called for several initiatives including the following (United
615 Nations Food and Agriculture Organization, 2006):

616 (1) the need for an international initiative to fill the data gap created by the malfunction of Landsat 7;

- 617 (2) a system to collect cloud-free, high resolution (10 to 20 m) visible, near-infrared, and shortwave infrared
618 observations at 5 to 10-day intervals;
- 619 (3) workshops on global agricultural data coordination and on integrating satellite and *in situ* observations;
- 620 (4) an inventory and evaluation of existing agro-meteorological data sets to identify gaps in terrestrial networks, the
621 availability of data, and validation and quality control in order to offer specific recommendations to the World
622 Meteorological Organization to improve its database;
- 623 (5) funding to support digitizing, archiving, and dissemination of baseline data; and
- 624 (6) an international workshop within the GEOSS framework to develop a strategy for “community of practice” for
625 improved global agricultural monitoring.

626 A recent study by the National Research Council (NRC) of the use of land remote sensing expressed additional
627 concerns about present limits on the usefulness of Earth observations in agricultural assessment) (National Research
628 Council, 2007). These include data integration, communication of results, and capacity to use and interpret data.

629 Specifically, the NRC identified these concerns:

- 630 (1) inadequate integration of spatial data with socioeconomic data (locations and vulnerabilities of human populations
631 and access to infrastructure) to provide information that is effective in generating response strategies to disasters or
632 other factors influencing access to food or impairing agricultural productivity;
- 633 (2) a lack of communication between remote sensing mission planners, scientists and decision makers to ascertain what
634 types of information enable the most effective food resource management; and
- 635 (3) shortcomings in the acquisition, archiving, and access to long-term environmental data and development of capacity
636 to interpret these data, including maintaining continuity of satellite coverage over extended time frames, providing
637 access to affordable data, and improving capacity to interpret data.

638

639 **4. Uncertainty**

640 Two aspects of PECAD provide means of validation and verification of crop assessments. One is the maturity
641 of PECAD as a decision support system. Over the years, it has been able to benchmark, validate, verify, and then
642 selectively incorporate additional data sources and automated decision tools. An example of the systems engineering
643 review associated with a decision to incorporate Poseidon and Jason data, for example, is offered in a detailed NASA
644 study (NASA, 2004b).

645 Another example demonstrates how data product accuracy, delivery, and coverage are tested through
646 validation and verification during the process of assimilating new data sources, as well as to ascertain the extent to
647 which different data sources corroborate model outputs (Kaupp *et al.*, 2005). Essential considerations included
648 enhanced repeatability of results, increased accuracy, and increased throughput speed.

649 Another significant aspect of resolving uncertainty in PECAD is its extensive use of a convergence
650 methodology to assimilate information from regional field analysts and other experts. PECAD seeks to provide accurate
651 and timely estimates of production, yet must accommodate physical and biological influences (e.g., weather or pests),
652 the fluctuations in agricultural markets, and developments in public policy impacting the agricultural sector (Kaupp *et*
653 *al.*, 2005). The methodology brings a large amount of additional information to the PECAD forecasts, well beyond the
654 automated outputs of the decision support tools. This extensive additional analysis may not fully correct for, but
655 certainly mitigates, the uncertainty inherent in the data and modeling at the early stages. Figure 3, a simplified version
656 of Figure 1, shows the step represented by the analyses that take place during this convergence of information in
657 relation to the outputs obtained from the decision support tools and their data inputs. Figure 4 further describes the
658 nature of information included in the convergence methodology in addition to the outputs of the data and automated
659 decision support tools. Official reports, news reports, field travel, and attaché reports are additional inputs at this stage.
660 The process is described as one in which, “while individual analysts reach their conclusions in different ways, giving
661 different weight to various inputs, analysts join experts from the USDA’s Economic Research Service and National
662 Agricultural Statistics Service once a month in a ‘lock-up.’ In this setting, the convergence of evidence approach is fully
663 realized as analysts join together in committee formed by (agricultural) commodity. Final commodity production
664 estimates are achieved by committee consensus” (NASA, 2004a, p. 4).

665 The convergence methodology is at the heart of analysis and the final step prior to official world agricultural
666 production estimates and suggests that uncertainty inherent in data and automated models at earlier stages of the
667 analysis are “scrubbed” in a broader context at this final stage.

668

669 **5. Global change information and PECAD**

670 The relationship between climate and agriculture is complex. Agriculture is not only influenced by a changing
671 climate, but agricultural practices themselves are a contributory factor through emissions of greenhouse gases and

672 influences on fluxes of carbon through photosynthesis and respiration. In short, agriculture is both a contributor to and a
673 recipient of the effects of a changing climate (Rosenzweig, 2003).

674 At present, PECAD is not directly used to address these dimensions of the climate-agriculture interaction.
675 However, many of the data inputs for PECAD are climate-related, thereby enabling PECAD to inform understanding of
676 agriculture as a “recipient” of climate-induced changes in temperature, precipitation, soil moisture, and other variables.
677 If reliable climate change prediction of temperature, precipitation, soil moisture, and other necessary variables become
678 available, then these variables can be used as input to PECAD and the results may be used to provide long-range
679 planning of agricultural practices. In addition, spatial and geographic trends in the output measures from PECAD have
680 the potential to contribute to understanding of how the agricultural sector is responding to a changing climate.

681 The output measures of PECAD also can serve to inform understanding of agriculture as a “contributor” to
682 climate changes. For example, observing trends in PECAD’s measures of production and composition of crops can shed
683 light on the contribution of the agriculture sector to agricultural soil carbon sequestration.

684

685 *The effects of a changing climate on agricultural efficiency as measured by PECAD:*

686

687 PECAD relies on several data sources for agro-meteorological phenomena that affect crop production and the quality of
688 agricultural commodities. These include data that are influenced by climate (e.g., precipitation, temperatures, snow
689 depth, and soil moisture). The productivity measures from PECAD (yield multiplied by area) are also influenced by
690 climate-induced changes in these data.

691 In addition, the productivity measures of PECAD can be indirectly but significantly affected by possible
692 climate-induced changes in land use. Examples of such changes include the reallocation of land from food production to
693 biomass fuel production or from food production to forestry cultivation as a means of carbon sequestration. In all of
694 these cases, Earth observations can contribute to understanding climate-related effects on agricultural efficiency
695 (National Research Council, 2007). Much of the research to integrate Earth observations into climate and agriculture
696 decision support tools is relatively recent; for example, in FY05, NASA, and USDA began climate simulations using
697 GISS GCM ocean temperature data and also completed fieldwork for verification and validation of a climate-based crop
698 yield model (NASA, 2006b). The UN FAO has begun to coordinate similar research on integrating Earth observations

699 and decision support systems to study possible effects of changing climate on food production and distribution (e.g., see
700 United Nations Food and Agriculture Organization, no date).

701


702 *The effects of agricultural practices and efficiency on climate:*

703


704 In addition to consideration of the effects of climate on agriculture, the feedback from agricultural practices to climate
705 has also been a topic of study (e.g., see <http://www.fao.org/NES/1997/971201-e.htm>, accessed April 2007). The crop
706 assessments and estimates from PECAD, by revealing changes in agricultural practices, could play a role as early
707 indicators to inform forecasting future agricultural-induced effects on climate. The Agricultural Research Service within
708 USDA and NASA have undertaken research using Earth observation data to study scale-dependent Earth—atmosphere
709 interactions, suggesting that significant changes in regional land use or agricultural practices could affect local and
710 regional climate (NASA, 2001).

711


712 **Figure 1: The PECAD Decision Support System: Data Sources and Decision Support Tools** (Source:
 713 Kaupp and coauthors, 2005, p. 5).
 714



715 **Figure 2. The PECAD Decision Support System: Earth System Models, Earth Observations, Decision**
 716 **Support Tools, and Outputs** (Source: National Aeronautics and Space Administration, 2006a, p. 32).




717



718

719

720 **Figure 3: The PECAD Decision Support System: The Role of Convergence of Evidence Analysis** (Source: National
 721 Aeronautics and Space Administration, 2004a, p. 8).



From: http://www.fas.usda.gov/pecad/remote/overview/frame_OV.htm

722 **Figure 4: The PECAD Decision Support System: Information Sources for the Convergence of Evidence Analysis**

723 (Source: National Aeronautics and Space Administration, 2004a, p. 5).

Chapter 2

Decision Support for Air Quality

(Use of CMAQ as a Decision Support Tool for Air Quality to Climate Change)

Lead Author: Daewon W. Byun

1. Introduction

Our ability to understand and forecast the quality of the air we breathe, as well as our ability to understand the science of chemical and physical atmospheric interactions, is at the heart of models of air quality. The quality of air is affected by and has implications for the topics presented in our other chapters. Air quality is affected by energy management and agricultural practices, for instance, and is a major factor in public health. Models of air quality also provide a means of evaluating the effectiveness of air pollution and emission control policies and regulations.

While numerous studies examine the potential impact of climate change on forests and vegetation, agriculture, water resources, and human health (examples are found in Brown *et al.*, 2004; Mearns, 2003; Leung and Wigmosta 1999; Kalkstein and Valimont 1987), attempts to project the response of air quality to changes in global and regional climates have long been hampered by the absence of proper tools that can transcend the different spatial and temporal scales involved in climate predictions and air quality assessment and by the uncertainties in climate change predictions and associated air quality changes.

One of the popular modeling tools to study air quality as a whole, including tropospheric ozone, fine particles, toxics, acid deposition, and visibility degradation is the US Environmental Protection Agency's (EPA) Community Multiscale Air Quality (CMAQ) modeling system. CMAQ has as its primary objectives to (1) improve the ability of environmental managers to evaluate the impact of air quality management practices for multiple pollutants at multiple scales, (2) enhance scientific ability to understand and model chemical and physical atmospheric interactions (<http://www.epa.gov/asmdnerl/CMAQ/>), and (3) guide the development of air quality regulations and standards and to create state implementation plans. It has been also used to evaluate longer-term pollutant climatologies as well as short-term transport from localized sources, and it can be used to perform simulations using downscaled regional climate from global climate change scenarios listed in IPCC (2000). Various observations from the ground and from *in-situ*,

751 aircrafts, and satellite platforms can be used at almost at every step of the processing of this Decision Support System
752 (DSS) for air quality.

753 Although there are significant effects of long-range transport, most of the serious air pollution problems are caused
754 by meteorological as well as chemical processes and their changes at regional and local areas, at scales much smaller
755 than those resolved by global climate models (GCM), which are typically applied at a resolution of several hundred
756 kilometers. Current-day regional climate simulations, which typically employ horizontal resolutions of 30 to 60 km, are
757 insufficient to resolve small-scale processes that are important for regional air quality, including low-level jets, land-sea
758 breezes, local wind shears, and urban heat island effects (Leung *et al.*, 2006). In addition, climate simulations place
759 enormous demands on computer storage. As a result, most climate simulations only archive a limited set of
760 meteorological variables, the time interval for the archive is usually 6 to 24 hours (e.g., Liang *et al.*, 2006), and some
761 critical information required for air quality modeling is missing.

762 The interaction and feedback between climate and air chemistry is another issue. Climate and air quality are linked
763 through atmospheric chemical, radiative, and dynamic processes at multiple scales. For instance, aerosols in the
764 atmosphere may modify atmospheric energy fluxes by attenuating, scattering, and absorbing solar and infrared
765 radiation, and may also modify cloud formation by altering the growth and droplet size distribution in the clouds. The
766 changes in energy fluxes and cloud fields may, in turn, alter the concentration and distribution of aerosols and other
767 chemical species. Although a few attempts have been made to address these issues, our understanding of climate
768 change is based largely on modeling studies that have neglected these feedback mechanisms.

769 The impact of climate change on air emissions is also of concern. Changes in temperature, precipitation, soil
770 moisture patterns, and clouds associated with global warming may directly alter emissions, including biogenic
771 emissions (e.g., isoprene and terpenes). Isoprene, an important natural precursor of ozone, is emitted mainly by
772 deciduous tree species. Emission rates are dependent on the availability of solar radiation in visual range and are highly
773 temperature sensitive. Emissions of terpenes (semi-volatile organic species) may induce formation of secondary organic
774 aerosols. The accompanying changes in the soil moisture, atmospheric stability, and flow patterns complicate these
775 effects, and it is difficult to predict whether climatic change will eventually lead to increased degradation of air quality.

776 This chapter discusses how CMAQ is used as the DSS for studying climate change impact on air quality addressing
777 the focus areas required by the *SAP 5.1 Prospectus*: (1) observational capabilities used in the DSS, (2) agencies and
778 organizations responsible, (3) characterization of interactions between users and the DSS information producers, (4)

779 sources of uncertainties with observation and the decision support tools, and (5) description of the relation between the
780 DSS and climate change information.

781

782

783 **2. Description of CMAQ**

784 The US EPA CMAQ modeling system (Byun and Ching, 1999; Byun and Schere, 2006) has the capability to
785 evaluate relationships between emitted precursor species and ozone at urban/regional scales (Appendix W to Part 51 of
786 40CFR: Guideline on Air Quality Models in “[http://www.epa.gov/fedrgstr/EPA-AIR/1995/August/Day-09/pr-
787 912.html](http://www.epa.gov/fedrgstr/EPA-AIR/1995/August/Day-09/pr-912.html)”). CMAQ uses state-of-the-science techniques for simulating all atmospheric and land processes that affect the
788 transport, transformation, and deposition of atmospheric pollutants. The primary modeling components in the CMAQ
789 modeling system include (1) a meteorological modeling system (e.g., The Fifth-Generation NCAR/Penn State
790 Mesoscale Model, MM5) or a Regional Climate Model (RCM) for the description of atmospheric states and motions,
791 (2) inventories of man-made and natural emissions of precursors that are injected into the atmosphere, and (3) the
792 CMAQ Chemistry Transport Modeling (CTM) system for the simulation of the chemical transformation and fate of the
793 emissions. The model can operate on a large range of time scales from minutes to days to weeks as well as on numerous
794 spatial (geographic) scales ranging from local to regional to continental.

795 The base CMAQ system is maintained by the U.S. EPA. The Center for Environmental Modeling for Policy
796 Development (CEMPD), University of North Carolina at Chapel Hill (UNC), is contracted to establish a Community
797 Modeling and Analysis System (CMAS) (<http://www.cmascenter.org/>) for supporting community-based air quality
798 modeling. CMAS helps development, application, and analysis of environmental models and helps distribution of the
799 DSS and related tools to the modeling community. The model performance has been evaluated for various applications
800 (e.g., Zhang *et al.*, 2006; Eder *et al.*, 2006; Tong and Mauzerall, 2006; Yu *et al.*, 2007). Table 1 lists Earth observations
801 (of all types-remote sensing and *in situ*) presently used in the CMAQ DSS.

802 Within this overall DSS structure as shown in Table 1, CMAQ is an emission-based, three-dimensional (3-D) air
803 quality model that does not utilize daily observational data directly for the model simulations. The databases utilized in
804 the system represent typical surface conditions, and demographic distributions. An example is the EPA’s Biogenic
805 Emissions Land Use Database, version 3 (BELD3) database (<http://www.epa.gov/ttn/chief/emch/biogenic/>) that
806 contains land use and land cover as well as the demographic and socioeconomic information. At present the initial

807 conditions are not specified using observed data even for those species routinely measured as part of the controlled
808 criteria species listed in the National Clean Air Act and its Amendments (CAAA) in an urban area using a dense
809 measurement network. This is because of the difficulty in specifying the multi-species conditions that satisfy chemical
810 balance in the system, which is subject to the diurnal evolution of radiative conditions and of the atmospheric boundary
811 layer as well as temporal changes in the emissions that reflect constantly changing human activities.

812 The main output of the CMAQ and its DSS is the concentration and deposition amount of atmospheric trace gases
813 and particulates at the grid resolution of the model, usually at 36 km for the continental United States (CONUS)
814 domain, and 12 km or 4 km for regional or urban scale domains. The end users of the DSS want information on the
815 major scientific uncertainties and our ability to resolve them subject to the information on socioeconomic context and
816 impacts. They seek information on the implications at the national, regional, and local scales and on the baseline and
817 future air quality conditions subject to climate change to assess the effectiveness of current and planned environmental
818 policies. Local air quality managers would want to know if the DSS could help assess methods of attaining current and
819 future ambient air quality standards and evaluate opportunities to mitigate the climate change impacts. Decision makers
820 would ask modelers to simulate the air quality in the future for a few plausible variations in the model inputs that
821 represent plausible climate scenarios of regional implications. Through sensitivity simulations of the DSS with
822 different assumptions on the meteorological and emissions inputs, the effectiveness of such policies and uncertainties in
823 the system can be studied. The results can be also compared with the historic air quality observations with similar
824 ambient conditions to validate predictions of the DSS.

825

826 **3. Potential Future Uses and Limits**

827 Although one of the major strengths of CMAQ is its reliance on the first principles of physics and chemistry, a few
828 modeling components, such as cloud processes, fine scale turbulence, radiative processes, etc., rely on
829 parameterizations or phenomenological concepts to represent intricate and less-well known atmospheric processes. The
830 present limitations in science parameterizations and modeling difficulties will continuously be improved as new
831 understanding of these phenomena are obtained through various measurements and model evaluation/verification. The
832 development of the chemical mechanism, Carbon Bond 05 (CB05), which recently replaced CB-4 is a case in point.
833 The reliability of the CMAQ simulation result is subject to quality of the emission inputs, both at the global and
834 regional scales, which depend heavily on socio-economic conditions. Because such estimates are obtained using

835 projection models in relevant socio-economic disciplinary areas, their accuracy must be scrutinized when used for the
836 decision-making process. The CMAQ DSS users/operators may not always have domain expertise to discern the
837 validity of such results.

838 CMAQ needs to have the ability to utilize available observations to specify more accurately the critical model
839 inputs, although they have been chosen based on best available information and experience currently. A data assimilation
840 approach may be used to improve the system performance at different processing steps.

841

842 Table 1. Input data used for operating the CMAQ-based DSS.

843 <<footnotes: PNNL, UIUC, NCEP, EPA, USGS, NASA>>

844 PNNL: Pacific Northwest National Laboratory

845 UIUC: University of Illinois at Urbana-Champaign

846 NCEP: National Center for Environmental Prediction

847 EPA: Environmental Protection Agency

848 USGS: US Geological Survey

849 NASA: National Aeronautics and Space Agency

850

851 For example, research has been undertaken to use satellite remote sensing data products together with high-
852 resolution land use and land cover (LULC) data to improve the land-surface parameterizations and boundary layer
853 schemes in the RCMs (e.g., Pour-Biazar *et al.*, 2007). Active research in chemical data assimilation (e.g.,
854 Constantinescu *et al.*, 2007a and b) is currently conducted with models such as STEM-II (Carmichael *et al.*, 1991) and
855 Goddard Earth Observing System (GEOS)-Chem (Bey *et al.*, 2001), which utilize both *in situ* and satellite observations
856 (e.g., Sandu *et al.*, 2005; Kopacz *et al.*, 2007; Fu *et al.*, 2007). Because of the coarse spatial and temporal resolutions of
857 the satellite data collected in the 1960s through the 1980s, and gas measurements through the launch of EOS Aura in
858 2004, most research in this area has been performed with global chemistry-transport models. As the horizontal
859 footprints of modern satellite instruments reach the resolution suitable for regional air quality modeling, these data can
860 be used to evaluate and then improve the bottom-up emissions inputs in the regional air quality models. However, they
861 do not provide required vertical information. The exception is occultation instruments, but these do not measure low

862 enough in altitude for air quality applications. In-*situ* and remote sensing measurements from ground and aircraft
863 platforms could be used to augment the satellite data in these data assimilation experiments.

864 Utilization of the column-integrated satellite measurements in a high-resolution 3-D grid model like CMAQ poses
865 serious challenges in distributing the pollutants vertically and separating those within and above the atmospheric
866 boundary layer. Because similar problems exist for the retrieval of meteorological profiles of moisture and temperature,
867 experiences in including these can be adapted for a few well-behaved chemical species. A data assimilation tool can be
868 used to improve the initial and boundary conditions using various in *situ* and satellite measurements of atmospheric
869 constituents. At present, however, an operational assimilation system for CMAQ is not yet available, although prototype
870 assimilation codes have recently been generated (Hakami, *et al.*, 2007; Zhang *et al.*, 2007). Should these data
871 assimilation tools become part of the DSS, various conventional and new satellite products, including Tropospheric
872 Emission Spectrometer (TES) ozone profiles, Geostationary Operational Environmental Satellites (GOES) hourly total
873 ozone column (GhTOC) data, Ozone Monitoring Instrument (OMI) total ozone column (TOC), The Cloud-Aerosol
874 Lidar and Infrared Pathfinder Satellite Observation (CALIPSO) (<http://www-calipso.larc.nasa.gov/>) attenuated
875 backscatter profiles, and OMI aerosol optical thickness (AOT) data can be utilized to improve the urban-to-regional
876 scale air quality predictions.

877 Because of the critical role of the RCM as the driver of CMAQ in climate change studies, RCM results for the long-
878 term simulations must be verified thoroughly. To date, evaluation of the RCM has been performed for the air quality
879 related operations only for relatively short simulation periods. For example, the simulated surface temperature, pressure,
880 and wind speed must be compared to surface observations to determine how well the model captures the mean land-
881 ocean temperature and pressure gradients, the mean sea breeze wind speeds, the average inland penetration of sea
882 breeze, the urban heat island effect, and the seasonal variations of these features. Comparisons with rawinsonde
883 soundings and atmospheric profiler data would determine how well the model reproduces the averaged characteristics of
884 the afternoon mixed layer heights and of the early morning temperature inversion, as well as the speed and the vertical
885 wind shears of the low-level jets. In addition to these mesoscale phenomena, changes in other factors can also alter the
886 air pollution patterns in the future and need to be carefully examined. These factors include the diurnal maximum,
887 minimum, and mean temperature; cloud cover; thunderstorm frequency; surface precipitation and soil moisture patterns;
888 and boundary layer growth and nocturnal inversion strength.

889 As demonstrated in global model applications, satellite measured biomass burning emissions data should be utilized
890 in the regional air quality modeling (e.g., Duncan *et al.*, 2003; Hoelzemann, *et al.*, 2004). Duncan *et al.* (2003)
891 presented a methodology for estimating the seasonal and interannual variation of biomass burning, designed for use in
892 global chemical transport models using fire-count data from the Along Track Scanning Radiometer (ATSR) and the
893 Advanced Very High Resolution Radiometer (AVHRR) World Fire Atlases. The Total Ozone Mapping Spectrometer
894 (TOMS) Aerosol Index (AI) data product was used as a surrogate to estimate interannual variability in biomass burning.
895 Also Spracklen *et al.* (2007) showed that the wildfire contribution to the interannual variability of organic carbon
896 aerosol can be studied using the area-burned data and ecosystem specific fuel loading data. A similar fire emissions
897 data set at the regional scales could be developed for use in a study of climate impact on air quality. For retrospective
898 application, a method similar to that used by the National Oceanic and Atmospheric Administration's (NOAA) Hazard
899 Mapping System (HMS) for Fire and Smoke (<http://www.ssd.noaa.gov/PS/FIRE/hms.html>) may be used to produce a
900 long-term regional scale fire emissions inventory for climate impact analysis.

901

902 **4. Uncertainty**

903 The CMAQ modeling system as currently operated has several sources of uncertainty in addition to those
904 associated with some of the limits described in the previous section. In particular, when CMAQ is used to study the
905 effects of climate change and air quality, improvements in several areas are necessary to reduce uncertainty. First, the
906 regional air quality models employ limited modeling domains and, as such, they are ignorant of air pollution events
907 outside the domains unless proper dynamic boundary conditions are provided. Second, because the pollutant transport
908 and chemical reactions are fundamentally affected by the meteorological conditions, improving both the global climate,
909 regional climate models, and the downscaling methods by evaluating and verifying physical algorithms that have been
910 implemented with observations as necessary in order to improve the system's overall performance. Third, the basic
911 model inputs, including land use/vegetation cover descriptions and emissions inputs must be improved. Fourth, the
912 model representativeness issues, including grid resolution problems, compensating errors among the model components,
913 and incommensurability of the model results compared with the dimensionality of the measurements (i.e., inherent
914 differences in the modeled outputs that represent volume and time averaged quantities to the point or path-integrated

915 measurements), as discussed in Russell and Dennis (2000) and NARSTO (2000), need to be addressed. These factors
916 are the principal cause of simulation/prediction errors.

917 Although the models incorporated in this DSS are first-principle based environmental models, they have difficulties
918 in representing forcing terms in the system, in particular, the influence of the earth's surface, long-range transport, and
919 uncertainties in the model inputs such as daily emissions changes due to anthropogenic and natural events. There is
920 ample opportunity to reduce some uncertainties associated with CMAQ through model evaluation and verification using
921 current and future meteorological and atmospheric chemistry observations. Satellite data products assimilated in the
922 global chemical transport models (GCTM) could provide better dynamic lateral boundary conditions for the regional air
923 quality modeling (e.g., Al-Saddi, *et al.*, 2005). Additional opportunities to reduce the model uncertainty include
924 comparison of model results with observed data at different resolutions, quantification of effects of initial and boundary
925 conditions and chemical mechanisms, application of CMAQ to estimate the uncertainty of input emissions data, and
926 ensemble modeling (using a large pool of simulations among a variety of models) as a means to estimate model
927 uncertainty.

928 A limitation in CMAQ applications, and therefore a source of uncertainty, has been the establishment of initial
929 conditions. The default initial conditions and lateral boundary conditions in CMAQ are provided under the assumption
930 that after spin-up of the model, they no longer play a role, and in time, surface emissions govern the air quality found in
931 the lower troposphere. Song *et al.* (2007) showed that the effects of the lateral boundary conditions differ for different
932 latitudes and altitudes, as well as seasons. In the future, dynamic boundary conditions can be provided by fully
933 integrating the GCTMs as part of the system. Several research groups are actively working on this, but the simulation
934 results are not yet available in open literature. A scientific cooperative forum, the Task Force on Hemispheric Transport
935 of Air Pollution (<http://www.htap.org/index.htm>), is endeavoring to bring together the national and international
936 research efforts at the regional, hemispheric, and global scales to develop a better understanding of air pollution
937 transport in the Northern Hemisphere. This task force is currently preparing its 2007 Interim Report addressing various
938 long-range transport of air pollutant issues (http://www.htap.org/activities/2007_Interim_Report.htm). Although the
939 effort does not directly address climate change issues, many of findings and tools used are very relevant to
940 meteorological and chemical downscaling issues.

941 Ultimately, CMAQ should consider all the uncertainties in the inputs. The system's response may be directly
942 related to the model configuration and algorithms (e.g., structures, resolutions, and chemical and transport algorithms),
943 compensating errors, and the incommensurability of modeling nature, as suggested by Russell and Dennis (2000).

944

945 **5. Global Change Information and CMAQ**

946 CMAQ could be used to help answer several questions about the relationship between air quality and climate
947 change. For instance:

948

949 1) How will global warming affect air quality in a region?

950 2) How will land use change due to climate, urbanization, or intentional management decisions affect air quality?

951 3) How much will climate change alter the frequency, seasonal distribution, and intensity of synoptic weather patterns
952 that influence pollution in a region?

953 4) How sensitive are air quality simulations to uncertainty in wildfire projections and to potential land management
954 scenarios?

955 5) How might the contribution of the local production and long-range transport of pollutants differ due to different
956 climate change scenarios?

957 6) Will future emissions scenarios or climate changes affect the frequency and magnitude of high pollution events?

958 To provide answers to these questions, CMAQ will rely heavily on climate-change-related information. In addition
959 to the influence of greenhouse gases and global warming, other forcing functions include population growth, land use
960 changes, new emission controls being implemented, and new energy sources to be available to replace the existing high-
961 carbon sources. Different scenarios can be chosen either to study potential impacts or to estimate the range of
962 uncertainties of the predictions. The two upstream climate models, GCMs and RCMs, generate the climate change data
963 that drive a GCTM and CMAQ. Both the GCMs and RCMs are expected to represent future climate change conditions
964 while simulating historic climate conditions that can be verified with comprehensive datasets such as the NCEP
965 Reanalysis data provided by the NOAA/OAR/ESRL PSD, Boulder, CO, from their Web site,
966 <http://www.cdc.noaa.gov/cdc/data.ncep.reanalysis.html> The meteorology simulated by the climate models represents

967 conditions in future year scenarios, reflecting changing atmospheric conditions. Furthermore, emissions inputs used for
968 the GCTM and CMAQ must reflect the natural changes and/or anthropogenic developments related to climate change
969 and other factors (e.g., population growth and geographical population shifts due to climate change).

970 In recent years, the EPA Science to Achieve Results (STAR) program has funded several projects on the possible
971 effects of climate change on air quality and on ecosystems. A majority of these projects have adopted CMAQ as the
972 base study tool. Figure 1 provides a general schematic of the potential structure of a CMAQ-based climate change DSS.
973 The figure shows potential uses of CMAQ for climate study; most climate-related CMAQ applications are not yet
974 configured as fully as indicated in the figure.

975 The projects linking CMAQ and climate study have used upstream models and downstream tools, including those
976 identified in Table 2. Related projects that use regional air quality models other than CMAQ are also listed. For the
977 GCMs, NCAR Community Climate Model (CCM) (Kiehl *et al.*, 1996), NASA Goddard Institute for Space Studies
978 (GISS) model (e.g., Hansen *et al.*, 2005), and NOAA Geophysical Fluid Dynamics Laboratory (GFDL) CM2 (Delworth
979 *et al.*, 2006) are the most popular global models for providing meteorological inputs representing climate change events.
980 A recent description for the GISS model can be found in Schmidt *et al.* (2006) (<http://www.giss.nasa.gov/tools/>) and for
981 the CCM in Kiehl *et al.* (1996) (<http://www.cgd.ucar.edu/cms/ccm3/>). A newer version of the CCM was released on
982 May 17, 2002 with a new name—the Community Atmosphere Model (CAM) ([http://www.cesm.ucar.edu/models/atm-](http://www.cesm.ucar.edu/models/atm-cam)
983 [cam](http://www.cesm.ucar.edu/models/atm-cam)). The model is described in Hurrell *et al.* (2006).

984

985 Table 2. An illustrative example of the potential uses of the models and upstream and downstream tools for a CMAQ-
986 based Climate Change Impact Decision Support System.

987 <footnote: WRF-ARW, WRF-NMM, SLEUTH>>

988

989 As shown in Table 2, for climate change studies, CMAQ is linked with upstream models such as a global climate
990 model (GCM), a global tropospheric chemistry model (GTCM), and a regional climate model (RCM) to provide
991 emissions sensitivity analysis, source apportionment, and data assimilation to assist policy and management decision-
992 making activities, including health impact analysis. Certain EPA STAR projects (Hogrefe *et al.*, 2004 and 2005;
993 Knowlton *et al.*, 2004; Civerolo *et al.*, 2007) have utilized the CMAQ-based DSS to assess whether climate change

994 would influence the effectiveness of current and future air pollution policy decisions subject to the potential changes in
995 local and regional meteorological conditions.

996 Other EPA STAR projects employ global climate change information from a GCM. For example, Tagaris *et al.*
997 (2007) and Liao *et al.* (2007a,b) use the results of GCM simulation with the well-mixed greenhouse gases—CO₂, CH₄,
998 N₂O, and halocarbons—updated yearly from observations for 1950 to 2000 (Hansen *et al.*, 2002) and for 2000 to 2052
999 following the A1B SRES scenario from the Intergovernmental Panel on Climate Change (IPCC 2001). The simulation
1000 used ozone and aerosol concentrations in the radiative scheme fixed at present-day climatological value provided in
1001 Mickley, *et al.* (2004).

1002
1003 To resolve the meteorological features affecting air pollution transport and transformation at a regional scale, the
1004 coarse scale meteorological data representing the climate change effects derived from a GCM are downscaled using an
1005 RCM. An RCM is often based on a limited-domain regional mesoscale model, such as MM5, the Regional
1006 Atmospheric Modeling System (RAMS), Eta, and WRF/ARW or WRF/NMM. An alternative method for constructing
1007 regional scale climate change data is through a statistical downscaling, which evaluates observed spatial and temporal
1008 relationships between large-scale (predictors) and local (predictands) climate variables over a specified training period
1009 and domain (Spak *et al.*, 2007). Because of the need to use a meteorological driver that satisfies constraints of dynamic
1010 consistency (i.e., mass and momentum conservations) for regional scale air quality modeling (e.g., Byun, 1999 a and b),
1011 the CMAQ modeling system relies exclusively on the dynamic downscaling method.

1012 Regional chemistry/transport models, like CMAQ, are better suited for regional air quality simulations than a
1013 GCTM because of the acute air pollution problems that are managed and controlled through policy decisions at specific
1014 geographic locations. Difficulty in prescribing proper boundary conditions (BC), especially in the upper troposphere, is
1015 one of the deficiencies of CMAQ simulations of air quality (e.g., Tarasick *et al.*, 2007; Tang *et al.*, 2007). Therefore,
1016 one of the main roles of the global CTM is to provide proper dynamic boundary conditions for CMAQ to represent
1017 temporal variation of chemical conditions that might be affected by the long-range transport of pollution (e.g., particle
1018 from large-scale biomass burnings) from outside the regional domain boundaries (Holloway, *et al.*, 2002; In *et al.*,
1019 2007). The contemporary EPA funded projects on climate change impact on air quality mainly use two 3-D GCTM
1020 models: the NASA/Harvard GEOS-Chem (Bey *et al.*, 2001) and the National Center for Atmospheric Research (NCAR)
1021 Model of Ozone and Related Chemical Tracers (MOZART) (Brasseur *et al.*, 1998; Horowitz *et al.*, 2003).

1022 The GEOS-Chem model (<http://www-as.harvard.edu/chemistry/trop>) is a global model for predicting tropospheric
1023 composition. The model was originally driven by the assimilated meteorological observation data from the GEOS of the
1024 NASA Global Modeling and Assimilation Office (GMAO). GEOS-Chem has been used as community assessment
1025 models for NASA Global Model Initiative, climate change studies with the NASA/GISS GCM, chemical data
1026 assimilation of tropospheric gaseous and aerosol species at NASA GMAO, and regulatory models for air pollution, in
1027 particular providing long-range transport information for regional air quality models. Long-term retrospective studies
1028 are possible with the GEOS data, which are available from 1985 to present at horizontal resolution of 2 degrees
1029 (latitude) by 2.5 degrees (longitude) until the end of 1999 and 1 degree by 1 degree afterward. For climate studies, the
1030 NASA GISS GCM meteorological outputs are used instead. Emission inventories include a satellite-based inventory of
1031 fire emissions (Duncan *et al.*, 2003) with expanded capability for daily temporal resolution (Heald *et al.*, 2003) and the
1032 National Emissions Inventory for 1999 (NEI 1999) for the US with monthly updates in order to achieve adequate
1033 consistency with the CMAQ fields at the GEOS-Chem/CMAQ interface.

1034 MOZART (<http://gctm.acd.ucar.edu/mozart/models/m3/index.shtml>) is built on the framework of the Model of
1035 Atmospheric Transport and Chemistry (MATCH) that can be driven with various meteorological inputs and at different
1036 resolutions such as meteorological reanalysis data from the National Centers for Environmental Prediction (NCEP),
1037 NASA GMAO, and the European Centre for Medium-Range Weather Forecasts (ECMWF). For climate change
1038 applications, meteorological inputs from the NCAR CCM3 are used. MOZART includes a detailed chemistry scheme
1039 for tropospheric ozone, nitrogen oxides, and hydrocarbon chemistry, semi-Lagrangian transport scheme, dry and wet
1040 removal processes, and emissions inputs. Emission inputs include sources from fossil fuel combustion, biofuel and
1041 biomass burning, biogenic and soil emissions, and oceanic emissions. The surface emissions of NO_x, CO, and NMHCs
1042 are based on the inventories described in Horowitz *et al.* (2003), aircraft emissions based on Friedl (1997), and lightning
1043 NO_x emissions that are distributed at the location of convective clouds.

1044 GCTMs are applied to investigate numerous tropospheric chemistry issues, involving gases – CO, CH₄, OH, NO_x,
1045 HCHO, and isoprene– and inorganic (sulfates and nitrates) and organic (elemental carbons, organic carbons)
1046 particulates. Various *in situ*, aircraft, and satellite-based measurements are used to provide the necessary inputs, to
1047 verify the science process algorithms, and to perform general model evaluations. They include vertical profiles from
1048 aircraft observations as compiled by Emmons *et al.* (2000), multiyear analysis of ozonesonde data (Logan, 1999), and
1049 those available at the Community Data Web site managed by the NCAR Earth and Sun Systems Laboratory (ESSL)

1050 Atmospheric Chemistry Division (ACD); and multiyear surface observations of CO reanalysis (Novelli *et al.*, 2003).
1051 Current and previous atmospheric measurement campaigns are listed in Web pages by NOAA Earth Systems Research
1052 Laboratory (ESRL), <http://www.esrl.noaa.gov/>; NASA, Tropospheric Integrated Chemistry Data Center, [http://www-](http://www-air.larc.nasa.gov/)
1053 [air.larc.nasa.gov/](http://www-air.larc.nasa.gov/); and NCAR ESSL (Earth and Sun Systems Laboratory) Atmospheric Chemistry Division (ACD)
1054 Community Data, <http://www.acd.ucar.edu/Data/>. These observations are used to set boundary conditions for the slow
1055 reacting species, including CH₄, N₂O, and CFCs, and to evaluate other modeled species, including CO, NO_x, PAN,
1056 HNO₃, HCHO, acetone, H₂O₂, and non-methane hydrocarbons. In addition, several satellite measurements of CO, NO₂,
1057 and HCHO from the Global Ozone Monitoring Experiment (GOME), The SCanning Imaging Absorption SpectroMeter
1058 for Atmospheric CHartographY (SCIAMACHY), and OMI instruments have been used extensively to verify the
1059 emissions inputs and performance of the GCTM.

1060 The grid resolutions used in the studies discussed above are much coarser than those used in the air quality models
1061 for studying emission control policy issues, such as evaluating state implementation plans (SIP). SIP modeling
1062 typically utilizes over 20 vertical layers at around 4-km horizontal grid spacing to reduce uncertainties in the model
1063 predictions near the ground and around high-emission source areas including urban and industrial centers. Although
1064 Civerolo *et al.*, (2007) applied CMAQ at a higher resolution, the duration of the CMAQ simulation was far too short a
1065 time scale to evaluate the regional climate impacts in detail.

1066 One of the additional key limitations of using the CMAQ for climate change studies is that the linkages between
1067 climate and air quality and from the global scale to regional scale models are only one-way (i.e., no feedback). Jacob
1068 and Gilliland (2005) stated that one-way assessment of the global change scenarios would be less useful for projection
1069 of air pollutant emissions because the evolution of regional air quality policies were not accounted for in these
1070 storylines. Also, to represent the interactions between atmospheric chemistry and meteorology, such as radiation and
1071 cloud/precipitation microphysics, particulates and heterogeneous chemistry, a two-way linkage must be established
1072 between the meteorology and chemistry models. An on-line modeling approach as implemented in WRF-chem is an
1073 example of such a linkage, but still there is a need to develop a link between the global and regional scales. A multi-
1074 resolution modeling system such as demonstrated by Jacobson (2001 a, b) might be necessary to address the true
1075 linkage between air pollution forcing and climate change and to provide the urban-to-global connection.

1076 In addition, there would be significant benefits to linking other multimedia models describing subsoil conditions,
1077 vegetation dynamics, hydrological processes, and ocean dynamics, including the physical/chemical interactions between

1078 the ocean micro-sublayer and atmospheric boundary layer to an air quality model. To generate such a megamodel under
1079 one computer coding structure would require handling of extremely different state variables in each multimedia model
1080 with substantially different data. Furthermore, interactions among the multimedia models need multidirectional data
1081 inputs, quality assurance checkpoints, and the decision-support entries. A more generalized on-line and two-way data
1082 exchange tools currently being developed under the Earth System Modeling Framework (ESMF)
1083 (<http://www.esmf.ucar.edu/>) may be a viable option.

1084 Observations not only represent the real changes in the climate but also provide a fundamental database to verify
1085 various modeling components in the DSS. The meteorological reanalysis data are available both in regional and global
1086 scales, but similar atmospheric chemistry database for air quality is lacking. An ozone database from ozonesonde
1087 system and other *in situ* measurements are useful for global scale studies. But for regional air quality studies, the
1088 availability of such measurements representing urban and local conditions in long-term is limited. Satellite or other
1089 remote sensing platform observations may provide additional data sources to build an atmospheric chemistry reanalysis
1090 database at global and regional scales, but these observations are mainly limited to ozone and aerosols. Such chemical
1091 reanalysis database can be utilized to study long-term air quality trends; to evaluate science process components in the
1092 air quality models, emissions, and other model inputs and configurations; and to improve model predictions through
1093 data assimilation approaches.

1094
1095 Figure 1. Configuration of CMAQ-based Decision Support System for climate change impact study




Figure 1. Configuration of CMAQ-based Decision Support System for climate change impact study

1096

1097 Table 1. Input data used for operating the CMAQ-based DSS.

Data Set	Type of Information	Source	Usage
Regional climate model output	Simulation results from a regional climate model (RCM) used as a driver for CMAQ modeling; processed through meteorology-chemistry interface processor (MCIP)	RCM modeling team; PNNL, UIUC, NCEP, EPA, and universities	Regional climate characterization, driver data for air quality simulations, and emissions processing
Land use, land cover, subsoil category, and topography data; topography for meteorological modeling	Describes land surface conditions and vegetation distribution for surface exchange processes	Various sources from USGS, NASA, NCEP EPA, states, etc.	Usually the data are associated with RCM's land surface module; need to be consistent with vegetation information, such as BELD3 if possible
Biogenic emissions land use database version 3 (BELD3)	Land use and biomass data and vegetation/tree species fractions	EPA	Processing of biogenic emissions; used to provide activity data for county-based emission estimates; now also used for land surface modeling in RCM
Air emissions inventories: national emissions inventories (NEI) and state/special	Amount and type of pollutants into the atmosphere. Includes: - Chemical or physical identity of pollutants - Geographic area covered - Institutional entities	EPA, regional program organizations (RPO), states and local government, and foreign governments	Preparation of model-ready emission inputs; perform speciation for the chemical mechanism used; used to evaluate "top-down" emissions (i.e.,

inventories; often called as “bottom-up” inventories	<ul style="list-style-type: none"> - Time period over which the emissions are estimated - Types of activities that cause emissions 		from inversion of satellite observations though air chemistry models)
Chemical species initial and boundary conditions	Clean species concentration profiles initial input and boundary conditions used for CMAQ simulations; originally from observations from clean background locations	EPA (fixed profiles), GEOS-Chem (Harvard & Univ. Houston), MOZART (NCAR); dynamic concentrations with diurnal variations (daily, monthly or seasonal)	CMAQ simulations; fixed profiles are used for outer domains where no significant emissions sources are located
AQS/AIRNow	Near real-time (AIRNow) and archived datasets (AQS) for ozone, PM, and some toxics species	Joint partnership between EPA and state and local air quality agencies	Measurement data used for model evaluations; report and communicate national air quality conditions for

1098

1099 Table 2. An illustrative example of the potential uses of the models and upstream and downstream tools for a CMAQ-
 1100 based Climate Change Impact Decision Support System.

Component	Functions	Model Name: Owner	Users
Global climate models (GCM)	Performs climate change simulations over the globe for different SRES climate scenarios. Typical resolution for a long-term (50 yearr) simulation is at 4° x 5° latitude and longitude	Community Climate Model (CCM): NCAR Goddard Institute for Space Studies (GISS) GCM: NASA CM2: Geophysical Fluid Dynamics Laboratory (GFDL) of NOAA	Climate research institutes, universities, and government institutions
Global chemistry transport models (GCTM)	Computes global scale chemical states in the atmosphere; uses same resolution as GCM	GEOS-Chem: NASA, Harvard University MOZART: NCAR (ESSL/Atmospheric Chemistry Division)	Global chemistry research organizations, universities, and government institutions
Regional climate models (RCM)	Simulates regional scale climate and meteorological conditions downscaling the GCM output; for US application ~36 km resolution used	MM5-based: NCAR, PNNL, UIUC, and others; the weather research and forecasting (WRF) model - advanced research WRF (WRF-ARW) core based: NCAR, UIUC Eta-based: NCEP (before June, 2006) The WRF- nonhydrostatic mesoscale model (WRF-NMM) core based: NCEP (after June,	Regional climate research groups, universities, and government institutions

		2006)	
Regional air quality models (AQM)	Performs air quality simulations at regional and urban scales at the same resolution as the RCM	Community multiscale air quality (CMAQ): EPA Comprehensive air quality model with extensions (CAMx): Environment WRF-Chem: NOAA/NCAR STEM-II: University of Iowa	Regional, state, and local air quality organizations; universities; private industries; and consulting companies
Downstream tools for decision support	Performs additional computations to help decision support, such as sensitivity and source apportionment studies, exposure studies	CMAQ/DDM: GIT CMAQ/4Dvar: CalTech/VT/UH Stochastic human exposure and dose simulation (SHEDS): EPA Total risk integrated methodology (TRIM): EPA	Universities and consulting companies
Upstream tools for representing climate change impacts on input data	Performs additional computations to generate model inputs that affect simulations	Land surface models SLEUTH: USGS, UC Santa Barbara (captures urban patterns) CLM (community land model): NCAR (used for RCM and biogenic emission estimates after growth)	Universities and consulting companies

1101

1102

1103 **DELIVERABLE ONE: REPORT ON THE DST(S) FOR YOUR NATIONAL APPLICATIONS AREA**

1104

1105

1106

Climate Change Science Program

1107

1108 **Uses and Limitations of Observations, Data, Forecasts, and Other Projections in**

1109

Decision Support for Selected Sectors and Regions

1110

1111 **Chapter 3. “Decision Support System for Assessing Hybrid Renewable Energy Systems”**

1112

1113

David S. Renné

1114

1115

National Renewable Energy Laboratory

1116

1617 Cole Boulevard

1117

Golden, CO 80401-3393

1118

1119 **1. Introduction**

1120

The national application area addressed in this chapter is the deployment of renewable energy technologies.

1121

Renewable energy technologies are being used around the world to meet local energy loads, to supplement grid-wind

1122

electricity supply, to perform mechanical work such as water pumping, to provide fuels for transportation, to provide

1123

hot water for buildings, and to support heating and cooling requirements for building energy design. Numerous

1124

organizations and research institutions around the world have developed a variety of decision-support tools to address

1125

how these technologies might perform in a most cost-effective manner to address specific applications. This chapter

1126

will focus on one specific tool, known as the Micropower Optimization Model, Hybrid Optimization Model for Electric

1127

Renewables (HOMER), that has been under consistent development and improvement at the US Department of

1128

Energy’s National Renewable Energy Laboratory and is used extensively around the world.

1129

1130 HOMER relies heavily on knowledge of the renewable energy resource available to the technologies being analyzed.
1131 Renewable energy resources, particularly for solar and wind technologies, are highly dependent on weather and climate
1132 phenomena, and are also driven by local microclimatic processes. Given the absence of a sufficiently dense ground
1133 network of reliable solar and wind observations, we must rely on validated numerical models, empirical knowledge of
1134 microscale weather characteristics, and collateral (indirect) observations derived from Earth observations, such as
1135 reanalysis data and satellite-borne remote sensors, to develop reliable knowledge of the geospatial characteristics and
1136 extent of these resources. Thus, the Decision Support System (DSS) described in this chapter includes HOMER as an
1137 end-use application and is described in the context of the renewable energy resource information required as input, as
1138 well as some intermediate steps that can be taken to organize these data, using Geographic Information Systems (GIS)
1139 software, to facilitate the application of HOMER.

1140

1141

1142

1143 2a. Description of the HOMER DSS

1144

1145 The HOMER DSS described in this chapter consists of three main components: (1) the renewable energy resource
1146 information required to estimate technology performance and operational characteristics, (2) (optional) organization of
1147 the resource data into a GIS framework so that the data can be easily imported into the decision support tool, and (3)
1148 NREL's Micropower Optimization Model known as HOMER, which ingests the renewable resource data for
1149 determining the optimal mix of power technologies for meeting specified load conditions at specified locations. This
1150 section describes each of these components separately. Although climate-based Earth observational data are primarily
1151 relevant only to the first component, some related Earth observation information could also be associated with the
1152 second and even the third component. Furthermore, it will be apparent that the first component is of major importance
1153 in the successful use of the HOMER DSS.

1154

1155 Although HOMER handles a number of power technologies, we will focus our attention in this chapter on solar and
1156 wind technologies and the resources required to run these technologies.

1157

1158

1159 Solar and Wind Resource Assessments

1160

1161 The first component of the HOMER DSS is properly formatted, reliable renewable energy resource data. The significant
1162 data requirements for this component are time-dependent measurements of wind and solar resources, as well as Earth
1163 observational data and data from numerical models, to provide the necessary spatial information for these resources,
1164 which can vary significantly over relatively small distances due to local microclimatic effects. Because of this natural
1165 variability, it is necessary to examine these energy resources geospatially in order to determine optimal siting of
1166 renewable energy technologies; alternatively, if a renewable energy technology is sited at a specific site in order to meet
1167 a nearby load requirement (such as a solar home system), it is necessary to know what the resource availability is at that
1168 location, since microclimatic variability may make even nearby data sources irrelevant.

1169

1170 Examples of the products derived from the methodologies described below can be found for many areas around the
1171 world. One significant project that has recently been completed is the Solar and Wind Energy Resource Assessment
1172 (SWERA) Project, which provided high-resolution wind and solar resource maps for 13 countries around the world.
1173 SWERA was a project funded by the Global Environment Facility and was cost-shared by several technical
1174 organizations around the world: NREL; the State University of New York at Albany, the NASA's Langley Research
1175 Center, and the USGS/EROS Data Center in the U.S.; Riso National Laboratory in Denmark; the German Aerospace
1176 Institute (DLR); the Energy Resources Institute (New Delhi, India); and the Brazilian Spatial Institute (INPE) in Sao
1177 Jose dos Campos, Brazil. The United Nations Environment Programme (UNEP) managed the project. Besides the
1178 solar and wind resource maps and underlying data sets, a variety of other relevant data products came out of this
1179 program. All of the final products and data can be found on the SWERA archive, hosted at the UNEP/GRID site,
1180 collocated with the USGS/EROS data center in Sioux Falls, South Dakota (<http://swera.unep.net>).

1181

1182 For wind resource assessments, NREL's approach, known as Wind Resource Assessment Mapping System (WRAMS)
1183 relies on mesoscale numerical models such as MM5 or Weather Research and Forecasting (WRF), which can provide
1184 simulations of near-surface wind flow characteristics in complex terrain or where sharp temperature gradients might
1185 exist (such as land-sea contrasts). Typically, these numerical models use available weather data, such as the National

1186 Climatic Data Center’s Integrated Surface Hourly (ISH) data network and National Center for Atmospheric Research-
1187 National Centers for Environmental Protection (NCAR-NCEP) reanalysis data as inputs. In coastal areas or island
1188 situations NREL’s wind resource mapping also relies heavily on SeaWinds data from the Quikscat satellite to obtain
1189 near-shore and near-island wind resources. WRAMS also relies on Global Land Cover Characterization (GLCC) 1-km
1190 and Regional Gap Analysis Program (ReGAP) 200-m land cover data, as well as Moderate Resolution Imaging
1191 Spectroradiometer (MODIS) data from the Aqua and Terra Earth Observation System satellites, to obtain information
1192 such as percent of tree cover and other land use information. This information is used not only to determine roughness
1193 lengths in the numerical mesoscale models but also to screen sites suitable for both wind and solar development in the
1194 second component of the HOMER DSS.

1195
1196 The numerical models are typically run at a 2.5-km resolution. However, wind resource information is often reported at
1197 the highest resolution at which a digital elevation model (DEM) can provide. Globally this has traditionally been 1-km
1198 resolution; however, in some cases in the US 400-m DEM data are available. Furthermore, the Shuttle Radar Topology
1199 Mission has now been able to provide users with a
1200 90-m DEM for much of the world. Thus, additional steps are needed beyond the 2.5-km resolution model output to
1201 depict wind resources at the higher resolutions offered by these DEM’s. This can be accomplished by using a
1202 secondary high-resolution mesoscale model, empirical methods, or both. For example, with NREL’s WRAMS
1203 methodology, GISD-based empirical modeling tools have been developed to modify results from the numerical models
1204 that appear to have provided unreliable results in complex-terrain areas.

1205
1206 The numerical models generally provide outputs at multiple levels above the ground. The WRAMS methodology
1207 provides values at a single specified height above the ground, nominally 50 m, or near the hub-height of modern-day
1208 large wind turbines (although with the recent advent of larger and larger wind turbines, hub heights are approaching 100
1209 m, so this standard height designation is changing). Where measured data are used to assess wind resources, a simple
1210 “power law” relationship is used to extrapolate the measured data to the desired height (Elliott *et al.*, 1987), i.e.

1211
1212
$$V_R/V_a = (Z_R/Z_a)^\alpha \quad (1)$$

1213

1214 where α , the power law coefficient, is normally assumed to be $1/7$, V_R is the wind speed at reference height Z_R
1215 (nominally, 50 m), and V_a is the wind speed at the measurement height Z_a .

1216

1217 The output of the WRAMS methodology is typically a value of wind power density at every grid-cell representative of
1218 an annual average (in order to produce monthly values, the procedure outlined above would have to be repeated for each
1219 month of the year). For mapping purposes, a classification scheme has been set up that relates a “wind power class” to
1220 a range of wind power densities. The classification scheme ranges from 1 to >7 , and applies to a specific height above
1221 ground. Normally, for grid-connected applications, a wind power class of 4 or above is best, while for small wind
1222 turbine applications where machines can operate in lower wind speeds, wind power class of 3 or above is suitable. Of
1223 course, the wind maps are not intended to identify sites at which large wind turbines can be installed, but rather are
1224 intended to provide information to developers on where they might most effectively install wind measurement systems
1225 for further site assessment. The maps also provide a useful tool for policy makers to obtain reliable estimates on the
1226 total wind energy potential for a region.

1227

1228 Other well-known approaches besides NREL’s WRAMS methodology are also used to produce large-area wind
1229 resource mapping. For example, Riso National Laboratory calculates wind speeds within 200 m above the Earth’s
1230 surface using the Karlsruhe Atmospheric Mesoscale Model (KAMM). Although KAMM also uses NCEP/NCAR
1231 reanalysis data, the model is based on large-scale geostrophic winds, and simulations are performed for classes of
1232 different geostrophic wind. The classes are weighted with their frequency to obtain statistics for the simulated winds.
1233 The results can then be treated as similar to real observations to make wind atlas files for the Wind Atlas Analysis and
1234 Application Program (WASP), which are employed to predict local winds at a much higher resolution than KAMM can
1235 provide. WASP calculations are based on wind data measured or simulated at specific locations and includes a complex
1236 terrain flow model, a roughness change model, and a model for sheltering obstacles. More on WASP can be found at
1237 <http://www.wasp.dk/>.

1238

1239 Due to the scarcity of high-quality, ground-based solar resource measurements, large-area solar resource assessments in
1240 the US have historically relied on the analysis of surface National Weather Service cloud cover observations. These
1241 observations are far more ubiquitous than solar measurements, and allowed NREL to develop a 1961 to 1990 National

1242 Solar Radiation Database for 239 surface sites. However, more recently in the US more and more reliance has been
1243 placed on GOES visible channel data to obtain surface reflectance information that can be used to derive high-resolution
1244 (~10-km) site-time specific solar resource data (see for example Perez, *et al.*, 2002). In fact, this approach has become
1245 commonplace in Europe, using Meteosat data. And the NASA Langley Research Center has recently completed a 20-
1246 year worldwide 100-km resolution Surface Solar Energy Data set derived from International Satellite Cloud
1247 Climatology Project data, which is derived from data collected by all of the Earth's geostationary and polar orbiting
1248 satellites (<http://eosweb.larc.nasa.gov/sse>).

1249
1250 The use of satellite imagery for estimating surface solar resource characteristics over large areas has been studied for
1251 some years, and Renné *et al.* (1999) published a summary of approaches developed around the world. These satellite-
1252 derived assessments require good knowledge of the aerosol optical depth over time and space, which can be obtained in
1253 part from MODIS and Advanced Very High Resolution Radiometer (AVHRR) data from polar orbiting environmental
1254 satellites. The assessments provide information both on Global Horizontal Irradiance (GHI), which is useful for
1255 estimating resources available to flat plate collectors such as photovoltaic panels or solar water heating systems, and
1256 Direct Normal Irradiance (DNI), which is needed for determining the resources available to solar concentrators that
1257 track the sun.

1258
1259 Besides NREL and NASA, other organizations perform similar types of high-resolution solar resource data sets. For
1260 example, the German Space Agency (DLR) has been applying similar methods to Meteosat data for developing solar
1261 resource maps and data for Europe and northern Africa. DLR was also involved in the SWERA project and applied
1262 their methodologies to several SWERA countries.

1263

1264

1265 Geospatial Toolkit

1266

1267 Recently, NREL has begun to format the solar and wind resource information into GIS software-compatible formats,
1268 and has incorporated this information, along with other geospatial data relevant to renewable energy development, into
1269 a Geospatial Toolkit (GsT). The GsT is a stand-alone, downloadable, and executable software package that allows the

1270 user to overlay the wind and solar data with other geospatial data sets available for the region, such as transmission
1271 lines, transportation corridors, population (load) centers, locations of power plant facilities and substations, land use and
1272 land form data, terrain data, etc. Not only can the user overlay various data sets of their choosing, there are also simple
1273 queries built into the toolkit, such as the amount of “windy” land (e.g. Class 3 and above) available within a distance of
1274 10-km of all transmission lines (minus specified exclusion areas, such as protected lands). The GsT developed at NREL
1275 makes use of the Environmental Science and Research Institute’s (ESRI) MapObjects software, although other
1276 platforms, including on-line, Web-based platforms, could also be used.

1277
1278 In a sense, the GsT is a DSS, since it allows the user to manipulate resource information with other critical data relevant
1279 to the deployment of renewable energy technologies to assist decision makers in identifying and conducting preliminary
1280 assessments of possible sites for installing these systems and supporting renewable energy policy decisions. However,
1281 up to now NREL has only prepared GsT’s for a few locations: the countries of Sri Lanka, Afghanistan, and Pakistan;
1282 Hebei Province in China; the state of Oaxaca in Mexico; and the state of Nevada in the US. By the time of publication
1283 of this chapter, additional toolkits may also be available. As with the resource data, all toolkits developed by NREL are
1284 available for download from NREL’s Web site. Those toolkits developed under the SWERA project are also available
1285 from the SWERA Web site.

1286

1287 HOMER: NREL’s Micropower Optimization Model

1288

1289 The primary decision support tool that makes up the DSS being described here is HOMER, NREL’s Micropower
1290 Optimization Model. HOMER is a computer model that simplifies the task of evaluating design options for both off-
1291 grid and grid-connected power systems for remote, stand-alone, and distributed generation applications. HOMER’s
1292 optimization and sensitivity analysis algorithms allow the user to evaluate the economic and technical feasibility of a
1293 large number of technology options and to account for variation in technology costs and energy resource availability.
1294 HOMER can also address system component sizing and the adequacy of the available renewable energy resource.
1295 HOMER models both conventional and renewable energy technologies:

1296 **Power sources:**

- 1297 • solar photovoltaic

- 1298 • wind turbine
- 1299 • run-of-river hydropower
- 1300 • Generator: diesel, gasoline, biogas, alternative and custom fuels, co-fired
- 1301 • electric utility grid
- 1302 • microturbine
- 1303 • fuel cell
- 1304 **Storage:**
- 1305 • battery bank
- 1306 • hydrogen
- 1307 **Loads:**
- 1308 • daily profiles with seasonal variation
- 1309 • deferrable (e.g., water pumping and refrigeration)
- 1310 • thermal (e.g., space heating and crop drying)
- 1311 • efficiency measures

1312

1313 In order to find the least cost combination of components that meet electrical and thermal loads, HOMER simulates
1314 thousands of system configurations, optimizes for lifecycle costs, and generates results of sensitivity analyses on most
1315 inputs. HOMER simulates the operation of each technology being examined by making energy balance calculations for
1316 each of the 8,760 hours in a year. For each hour, HOMER compares the electric and thermal load in the hour to the
1317 energy that the system can supply in that hour. For systems that include batteries or fuel-powered generators, HOMER
1318 also decides for each hour how to operate the generators and whether to charge or discharge the batteries. If the system
1319 meets the loads for the entire year, HOMER estimates the lifecycle cost of the system, accounting for the capital,
1320 replacement, operation and maintenance, and fuel and interest costs. The user can obtain screen views of hourly energy
1321 flows for each component as well as annual costs and performance summaries.

1322

1323 This and other information about HOMER are available on NREL's Web site: <http://www.nrel.gov/homer/>. The Web
1324 site also provides extensive examples of how HOMER is used around the world to evaluate optimized hybrid renewable

1325 power systems to meet load requirements in remote villages. Figure 1 shows a typical example of an output graphic
1326 available from HOMER.

1327

1328 In order to accomplish these tasks, HOMER requires information on the hourly renewable energy resources available to
1329 the technologies being studied. However, typically hour-by-hour wind and solar data are not available for most sites.
1330 Thus, the user is requested to provide monthly or average information on solar and wind resources; HOMER then uses
1331 an internal weather generator to provide the best estimate of a simulated hour-by-hour data set, taking into consideration
1332 diurnal variability if the user can provide an indication of what this should be. However, these approximations
1333 represent a source of uncertainty in the model. For those locations where a GsT is available, the GsT offers a
1334 mechanism for the user to easily ingest data from the toolkit into HOMER for the specific location of interest. However,
1335 since the toolkit contains only monthly solar and wind data, the limitations described above still apply. More
1336 information on the weather generator can be found in the HOMER Help files.

1337

1338 The HOMER developers have implemented various methods to facilitate access to reliable resource data that provide
1339 some of the input for simulations. For example, a direct link with the NASA SSE data site enables the user to download
1340 monthly and annual solar data from any location on Earth. The 100-km resolution NASA data have become a
1341 benchmark of solar resource information, due to the high quality of the modeling capability used to generate the data,
1342 the fact that the SSE is validated against numerous ground stations, and the fact that it is global in scope and now covers
1343 a 20-year period. However, the data set is still limited by a somewhat coarse resolution and no validation in areas where
1344 ground data do not exist. The procedures used to generate the SSE also have problems where land-ocean interfaces
1345 occur, and in snow-covered areas.

1346

1347 Linking HOMER to higher-resolution regional solar data sets would likely improve these uncertainties somewhat, but in
1348 general these data sets are also limited to monthly and seasonal values. However, since these methods rely on
1349 geostationary satellite data that provide frequent imagery of the Earth's surface, an opportunity exists to produce hourly
1350 time series data for up to several years at a 10-km resolution. This option will require significant data storage and
1351 retrieval capabilities on a server, but such a possibility now exists for future assessments.

1352

1353 Wind data available to HOMER is also generally limited to annual and at best monthly values. The standard HOMER
1354 interface allows the user to also designate a Weibull “k” value if this information is available. The Weibull k is a
1355 statistical means of defining the frequency distribution of the long-term hourly wind speeds at a location; this value can
1356 vary substantially depending on local terrain and microclimatic conditions. HOMER also has a provision for the user to
1357 designate the diurnal range of wind speeds and the timing when maximum and minimum winds occur. This
1358 information then provides improved simulation of the hour-by-hour wind values. The difficulty is that there may be
1359 applications where even these statistical values are not known to the user and are not available from the standard wind
1360 resource maps produced for a region, but this limitation may not be critical and requires further study to determine the
1361 impact on model output uncertainties.

1362

1363 2b. Access to the HOMER DSS

1364

1365 HOMER was originally developed and has always been maintained by the National Renewable Energy Laboratory.
1366 The model can be downloaded free of charge from NREL’s Web site at <http://www.nrel.gov/homer/default.asp>. The
1367 user is required to register, and registration must be updated every six months. The Web site also contains a variety of
1368 guides for getting started and using the software.

1369

1370 Resource information required as input to HOMER is generally freely available at the Web sites of the institutions
1371 developing the data. These institutions also generally maintain and continuously update the data. For example,
1372 renewable energy resource information can be found in several places on NREL’s Web site, such as <http://rredc.nrel.gov>
1373 or www.nrel.gov/GIS. NASA solar energy data, which can be easily input to HOMER, is available at
1374 <http://eosweb.larc.nasa.gov/sse>. In fact, there is a specific feature built into HOMER that automatically accesses and
1375 inputs the SSE data for the specific location that the model is analyzing. Wind and solar resource data for the 13
1376 SWERA countries can be found at <http://unep.swera.net>. This Web site is currently undergoing expansion and
1377 upgrading by the USGS/EROS Data Center in Sioux Falls, SD, and will eventually become a major clearing house for
1378 resource data from around the world in formats that can be readily ingested into tools such as HOMER.

1379

1380 2c. Definition of HOMER information requirements

1381

1382 The ideal input data format to HOMER is an hourly time series of wind and solar resource data covering a complete
1383 year (8,760 values). In addition, the wind data should be representative of the wind turbine hub height that is being
1384 analyzed within HOMER. Unfortunately data sets such as these are seldom available at the specific locations for which
1385 HOMER is being applied. More typically, the HOMER user will have to identify input data sets from resource maps
1386 (even within the GsT, the resource data are based on what is incorporated into the map, which, in the case of wind, may
1387 represent only a single annual value). Because monthly and annual mean data are more typically available, HOMER
1388 has been designed to take monthly mean wind speeds (in m/s) and monthly mean solar resource values (in kw-h/m²-
1389 day). In the case of wind, HOMER also allows for the specification of other statistical parameters related to wind speed
1390 distributions and diurnal characteristics. Furthermore, if the wind data available for input to HOMER do not represent
1391 the same height above the ground as the wind turbine's hub height being analyzed, HOMER has internal algorithms to
1392 adjust for this. The user must specify the height above the ground for which the data represent, and a power law
1393 conversion adjusts the wind speed value to the hub height of the specific wind turbine being analyzed. HOMER then
1394 utilizes an internal weather generator that takes the input information and creates an hour-by-hour data profile
1395 representing a one-year data file. Then, HOMER calculates turbine energy output by converting each hourly value to
1396 the energy production of the machine using the manufacturer's turbine power curve.

1397

1398 Besides the mean monthly wind speeds, the statistical parameters required by HOMER to generate the hourly data sets
1399 include the following:

1400

- 1401 • The altitude above sea level (to adjust for air density, since turbine performance is typically rated at sea level);
- 1402 • The Weibull k value, which typically ranges from 1.5 to 2.5, depending on terrain type;
- 1403 • An auto-correlation factor, which is a measure of how strongly the wind speed in 1 hour depends (on average)
1404 on the wind speed in the previous hour (these values typically range from 0.85 to 0.90);
- 1405 • A diurnal pattern strength, which is a measure of how strongly the wind speed depends on the time of day
1406 (values are typically 0.0 to 0.4); and
- 1407 • The hour of the peak wind speed (over land areas this is typically 1400 to 1600 local time)

1408

1409 In the US as elsewhere, wind resource maps often depict the resource in terms of wind power density, in units of watts-
1410 m^{-2} rather than in wind speeds. In this case, the wind power density must be converted back to a mean wind speed. The
1411 relationship between wind power density (P) and wind speed (v) is given as follows:

1412

$$1413 \quad P = \frac{1}{2}\rho \sum_i v_i^3, \quad (2)$$

1414

1415 where ρ is the density of the air and i is the individual hourly wind observation. Since the frequency distribution of
1416 wind speed over the period of a year or so follows a Weibull distribution shape, the wind power density can be
1417 converted back to a wind speed if the “k” factor in the Weibull distribution is known, as well as the height above sea
1418 level of the site (to determine the air density).

1419

1420 2d. Access to and use of the HOMER DSS among the federal, state, and local levels

1421

1422 Because of the easy access to HOMER and to the related resource assessment data products, the HOMER DSS is freely
1423 available to all government and private entities in the US and worldwide. Thousands of users from all economic sectors
1424 are using HOMER to evaluate renewable energy technology applications, particularly for off-grid use.

1425

1426 2e. Variation of the HOMER DSS by geographic region or characteristic

1427

1428 A key feature of HOMER is the evaluation of specific renewable energy technologies and related energy systems for
1429 different regions and for different applications. The HOMER model contains information on renewable energy
1430 technology characteristics; however, these characteristics, such as power curves of difference wind turbine models,
1431 generator fuel curves, and other factors are not affected by location. Because of the location-specific dependency of
1432 resource data, use of data that is not representative of the specific region of analysis will introduce additional
1433 uncertainties in the model results. Thus, the user should evaluate the accuracy and relevancy of any default information
1434 that is built into HOMER, or any resource data chosen as input to HOMER before completing the final analyses.

1435

1436 **3. Observations used by the HOMER DSS now and of potential use in the future**

1437

1438 This section focuses on the Earth observations (of all types, from remote sensing and *in situ*) used or of potential use in
1439 the HOMER DSS.

1440

1441 3a. Kinds of observations being used

1442

1443 In the previous section we provided a description of the renewable energy resource assessment related to solar and wind
1444 technologies that are required as input to HOMER when these technologies are being modeled. As noted in that section,
1445 developing this resource information requires the use of a variety of Earth observations. In this section we list out these
1446 observations for each resource category, as well as other types of observations relevant to the HOMER DSS.

1447

1448 Wind Resources

1449

1450 The ideal observational platform for obtaining reliable wind resource data to be input into HOMER would be calibrated
1451 wind speed measurements from a meteorological tower installed at the location of interest. These measurements should
1452 be obtained at the hub height of the wind turbine being modeled, should be of sufficient sampling frequency to provide
1453 hourly measurements, and should be of sufficient quality and duration to result in at least one full year of continuous
1454 measurements. Although measurements of this quality are typically necessary at project sites where significant
1455 investments in large grid-connected wind turbines are anticipate, and where a decision has already been made to
1456 implement a large-scale project, it is extremely rare that this level of observation is available for most HOMER
1457 applications, where the user is examining potential applications for proposed projects. Thus, some indirect means to
1458 establish wind characteristics at a proposed site, such as extrapolating wind resource measurements available from a
1459 nearby location or developing a wind resource map such as described in Section 2, is required. The major global data
1460 sets typically used by NREL for wind resource assessment are summarized in table 1.

1461

1462 More discussion on some of these data sets is provided below.

1463

1464 Surface Station Data

1465

1466 In the US, as well as in most other countries, the main source of routine surface wind observations would be
1467 observations from nearby national weather stations, such as those routinely maintained to support aircraft operations at
1468 airports. These data can be made available to the user from the National Climatic Data Center (NCDC) in the form of
1469 the Integrated Surface Hourly (ISH) data set. This database is composed of worldwide surface weather observations
1470 from about 20,000 stations, collected and stored from sources such as the Automated Weather Network (AWN), the
1471 Global Telecommunications System (GTS), the Automated Surface Observing System (ASOS), and data keyed from
1472 paper forms (see, http://gcmd.nasa.gov/records/GCMD_C00532.html).

1473

1474 Satellite-Derived Ocean Wind Data

1475

1476 Ocean wind data can be obtained from the SeaWinds Scatterometer (see <http://manati.orbit.nesdis.noaa.gov/quikscat/>)
1477 mounted aboard NASA's Quick Scatterometer (QuickSCAT) satellite. QuickSCAT was launched on June 19, 1999 in a
1478 sun-synchronous polar orbit. A longer-term ocean winds data set is available from the Special Sensor
1479 Microwave/Imager (SSM/I) data products as part of NASA's Pathfinder Program. The SSM/I geophysical dataset
1480 consists of data derived from observations collected by SSM/I sensors carried onboard the series of Defense
1481 Meteorological Satellite Program (DMSP) polar orbiting satellites (see
1482 http://www.ssmi.com/ssmi/ssmi_description.html#ssmi). An example of how Scatterometer data were used in support
1483 of a wind resource assessment in Pakistan is provided in figure 2 (see also
1484 http://www.nrel.gov/international/rr_assess_pakistan.html). Airborne or space borne Synthetic Aperture Radar systems
1485 can also provide information on ocean wind data, although these data are not commonly used for this purpose in the US,
1486 since Scatterometer data products are more readily and freely available.

1487

1488 Reanalysis Upper Air Data

1489

1490 The US reanalysis data set was first made available in 1996 to provide gridded global upper air and vertical profiles of
1491 wind data derived from 1,800 radiosonde and pilot balloon observations stations (Kalnay, *et al.* 1997). The reanalysis
1492 data were prepared by NCAR-NCEP and can be found at <http://www.cdc.noaa.gov/cdc/reanalysis/>. An early analysis of

1493 the data set (Schwartz, George, and Elliott, 1999) showed that for wind resource assessments the dataset was a
1494 promising tool for gaining a more complete understanding of vertical wind profiles around the world but that
1495 discrepancies with actual radiosonde observations still existed. Since that time, continuous improvements have been
1496 made to the NCAR-NCEP dataset, and it is has become an ever-increasingly important data source for contributing to
1497 reliable wind resource mapping activities.

1498

1499 Digital Terrain Data

1500

1501 Digital Elevation Models (DEM) have been accessed from the USGS/EROS data center. These models consist of a
1502 raster grid of regularly spaced elevation values that have been derived primarily from the USGS topographic map series.
1503 The USGS no longer offers DEMs, and for the US these can now be accessed from the National Elevation Dataset
1504 (<http://ned.usgs.gov/>). The Shuttle Radar Topographic Mission (SRTM) offers much higher resolution terrain data sets,
1505 which are now beginning to be used in some wind mapping exercises. These are also being distributed by USGS/EROS
1506 under agreement with NASA (<http://srtm.usgs.gov/>).

1507

1508 Digital Land Cover Data

1509

1510 Land cover data are used to estimate roughness length parameters required for the mesoscale meteorological models
1511 used in the wind mapping process. Data from the Global Land Cover Characterization dataset provide this information
1512 at a 1-km resolution (see <http://edcsns17.cr.usgs.gov/glcc/background.html>). The Moderate Imaging Spectroradiometer
1513 (MODIS) is used to obtain global percent tree cover values at a spatial resolution of 0.5 km (Hansen, *et al.*, 2003).
1514 Existing natural vegetation is also being mapped at a 200-m resolution as part of the USGS Regional Gap Analysis
1515 program. Gap analysis is a scientific method for identifying the degree to which native animal species and natural
1516 communities are represented in our present-day mix of conservation lands (Jennings and Scott, 1997).

1517

1518 Solar Resources

1519

1520 As with wind, the ideal solar resource data set for incorporation into HOMER would be data derived from a quality,
1521 calibrated surface solar measurement system consisting of a pyranometer and a pyrliometer that can provide a
1522 continuous stream of hourly data for at least one year. Such data are seldom available at the site for which HOMER is
1523 being applied. Although interpolation to nearby surface radiometer data sets can be accomplished with reasonable
1524 reliability, we usually resort to an estimation scheme to derive an *in-situ* data set. The solar resource assessments that
1525 NREL and others undertake make use of several different observational datasets, such as ground-based cloud cover
1526 measurements, satellite-derived cloud cover measurements, or the use of the visible channel from satellite imagery data.
1527 The major global data sets used for solar resource assessments are summarized in table 2.

1528

1529 More discussion on some of these data products is described below.

1530

1531 World Radiation Data Center

1532

1533 Since the early 1960s the World Radiation Data Center, located at the Main Geophysical Institute in St. Petersburg,
1534 Russia, has served as a clearinghouse for worldwide solar radiation measurements collected at national weather stations.
1535 The WRDC is under the auspices of the World Meteorological Organization. A Web-based data set was developed by
1536 NREL in collaboration with the WRDC and can be accessed at <http://wrdc-mgo.nrel.gov/>. This data archive covers the
1537 period 1964 to 1993. For more recent data, the user should go directly to the WRDC home page at
1538 <http://wrdc.mgo.rssi.ru/>.

1539

1540 Aerosol Optical Depths (AOD)

1541

1542 After clouds, atmospheric aerosols have the greatest impact on the distribution and characteristics of solar resources at
1543 the Earth's surface. However, routine *in-situ* observations of this parameter have only recently begun. Consequently, a
1544 variety of surface-based and satellite-based observations are used to derive the best information possible of the temporal
1545 and spatial characteristics of the atmospheric AOD. The most prominent of the surface data sets is the AERONET
1546 (<http://aeronet.gsfc.nasa.gov/>), a network of automated multiwavelength sun photometers located around the world.
1547 This network also has links to other networks, where the data may be less reliable. AERONET data can be used to

1548 provide ground-truth data for different satellite sensors that have been launched on a variety of sun-synchronous
1549 orbiting platforms since the 1980s, such as the Total Ozone Mapping Spectrometer (TOMS), the Advanced Very High
1550 Resolution Radiometer (AVHRR), the Moderate Resolution Imaging Spectroradiometer (MODIS), and the Multi-Angle
1551 Imaging Spectroradiometer (MISR), the latter two mounted on NASA's Terra satellite. As noted by Gueymard (2003)
1552 determination of AOD from satellite observations is still subject to inaccuracies, particularly over land areas, due to a
1553 variety of problems such as insufficient cloud screening or interference with highly reflective surfaces. The Global
1554 Aerosol Climatology Project (GACP), established in 1998 as part of the NASA Radiation Sciences Program and the
1555 Global Energy and Water Experiment (GEWEX), has as its main objectives to analyze satellite radiance measurements
1556 and field observations in order to infer the global distribution of aerosols, their properties, and their seasonal and
1557 interannual variations and to perform advanced global and regional modeling studies of the aerosol formation,
1558 processing, and transport (<http://gacp.giss.nasa.gov/>).

1559 Other sources of aerosol optical depth data include the Global Ozone Chemistry Aerosol Transport (GOCART) model
1560 (<http://code916.gsfc.nasa.gov/People/Chin/gocartinfo.html>) which is derived from a chemical transport model. An
1561 older dataset, the Global Aerosol Dataset (GADS), which can be found at [http://www.lrz-](http://www.lrz-muenchen.de/~uh234an/www/radaer/gads.html)
1562 [muenchen.de/~uh234an/www/radaer/gads.html](http://www.lrz-muenchen.de/~uh234an/www/radaer/gads.html), is a theoretical data set providing aerosol properties averaged in space
1563 and time on a $5^{\circ} \times 5^{\circ}$ grid. (Koepke, *et al.*, 1997).

1564 Other Renewable Energy Resources

1565 Although the scope of this chapter focuses on wind and solar energy resources, it is evident that many of the Earth
1566 observation data sets listed above can apply to other renewable energy resources as well. For example, hydropower
1567 resources can be determined by analysis of high resolution DEM data, along with knowledge of the rainfall amounts
1568 over specific watersheds and the land use characteristics of these watersheds. Biomass resource assessments can be
1569 enhanced through use of MODIS data as well as other weather-related data, and through evaluation of MODIS and
1570 AVHRR data to determine the Normalized Vegetation Index (NVI).

1571 3b. Limitations on the usefulness of observations

1572

1573 In the absence of direct solar and wind resource measurements at the location for which HOMER is being applied, the
1574 observations described in Section 3a, when used in the wind and solar resource mapping techniques described in
1575 Section 2, will together provide useful approximations of the data required as input to HOMER. However, the
1576 observations all have limitations in that they do not explicitly provide direct observation of the data value required for
1577 the mapping techniques but only approximations based on the use of algorithms to convert a signal into the parameter of
1578 interest. These limitations for some of these data sets can be summarized here:

1579

1580 Surface Station Data: These are generally not available at the specific locations at which HOMER would be applied, so
1581 interpolation is required. Furthermore, they generally do not have actual solar measurements, but rather proxies for
1582 these measurements (i.e., cloud cover). The wind data are generally collected at 10 m above the ground or less, and the
1583 anemometer may not be in a well-exposed condition. When the station observations are derived from human
1584 observations, they represent samples of a few minutes duration every 1 or 3 hours; therefore, many of the observations
1585 are missing. For those stations that have switched from human observations to Automated Surface Observation Stations
1586 (ASOS), the means of observation have changed significantly from the human observations, representing a
1587 discontinuity in long-term records. Occasionally, the location of the station is changed without changing the station ID
1588 number, which can also cause a discontinuity in observations. Similarly, equipment changes can cause a discontinuity in
1589 observations

1590

1591 Satellite-Derived Ocean Wind Data: These data are not based on direct observation of the wind speed at
1592 10 m above the ocean surface, but rather from an algorithm that infers wind speeds based on the wave height
1593 observations provided by the scatterometers or Synthetic Aperture Radar

1594

1595 Satellite-Derived Cloud Cover and Solar Radiation Data: These data sets are derived from observations of the
1596 reflectance of the solar radiation from the Earth-atmosphere system. Although it could be argued that this method does
1597 provide a direct observation of clouds, the solar radiation values are determined from an algorithm that converts
1598 knowledge of the reflectance observation, the incoming solar radiation at the top of the atmosphere, and the
1599 transmissivity characteristics of the atmosphere to develop estimates of solar radiation.

1600

1601 Aerosol Optical Depth: Considerable research is underway to improve the algorithms used to convert multi-spectral
1602 imagery of the Earth's surface to aerosol optical depth. The satellite-derived methods have additional shortcomings
1603 over land surfaces, where irregular land-surface features make application of the algorithms complicated and uncertain.

1604

1605

1606 3c. Reliability of the observations

1607

1608 For those observations that provide inputs to the solar and wind resource data, their reliability can vary from parameter
1609 to parameter. Generally all of the observations used to produce data values required for solar and wind assessments
1610 have undergone rigorous testing, evaluation, and validation. This research has been undertaken by a variety of
1611 institutions, including the institutions gathering the observations (e.g., NASA and NOAA) as well as the institutions
1612 incorporating the observations into resource mapping techniques (e.g., NREL). Many of the satellite-derived
1613 observations of critical parameters will be less reliable than *in-situ* observations; however, satellite-derived observations
1614 must still be used due to the scarcity of *in-situ* measurement stations.

1615

1616 3d. What kinds of observations could be useful in the near future

1617

1618 All of the observations currently available will continue to be of critical value in the near future. For renewable energy
1619 resource mapping, improved observations of key weather parameters (wind speed and direction at various heights above
1620 the ground and over the open oceans at higher and higher spatial resolutions, improved ways of differentiating snow
1621 cover and bright reflecting surfaces from clouds, etc.) will always be of value to the renewable energy community.
1622 New, more accurate methods of related parameters such as aerosol optical depth would result in improvements in the
1623 resource data. All of these steps will lead to improvements in the quality of outputs from renewable energy decision
1624 support tools such as HOMER.

1625

1626 **4. Uncertainty**

1627

1628 Application of the HOMER DSS involves a variety of input data types, all of which can have a level of uncertainty
1629 attached to them. HOMER addresses uncertainties by allowing the user to perform sensitivity analyses for any
1630 particular input variable or combination of variables. HOMER repeats its optimization process for each value of that
1631 variable and provides displays to allow the user to see how results are affected. An input variable for which the user has
1632 specified multiple values is called a sensitivity variable, and users can define as many of these variables as they wish.
1633 In HOMER, a “one-dimensional” sensitivity analysis is done if there is a single sensitivity variable, such as the mean
1634 monthly wind speed. If there are two or more sensitivity variables, the sensitivity analysis is “two” or “multi-
1635 dimensional.” HOMER has powerful graphical capabilities to allow the user to examine the results of sensitivity
1636 analyses of two or more dimensions. This is important for the decision maker, who must factor in the uncertainties of
1637 input variables in order to make a final judgment on the outputs of the model.

1638
1639 The amount of uncertainty associated with resource data is largely dependent on how the data are obtained and on the
1640 nature of the analysis being undertaken. For some types of analyses, very rough estimates of the wind resource would
1641 be sufficient; for others, detailed hourly average data based on surface measurements would be necessary. Quality *in-*
1642 *situ* measurements of wind and solar data in formats suitable for renewable energy applications over a sufficient period
1643 of time (one year or more) can have uncertainties of less than $\pm 3\%$ of the true value. However, when estimation
1644 methods are required, such as the use of Earth observations and modeling and empirical techniques, uncertainties can be
1645 as much as $\pm 10\%$ or more. These uncertainties are highest for shorter-term data sets, and are lower when annual
1646 average values are being used, since throughout the year errors in the estimation methods have a tendency to
1647 compensate among the individual values.

1648
1649 Based on wind turbine and solar technology operating characteristics, it is possible that the error in estimating a
1650 renewable energy system performance over a year is roughly linear to the error in the input resource data. For example,
1651 for wind energy systems, even though the power of the wind available to a wind turbine is a function of the cube of the
1652 wind speed, it turns out that the turbine operating characteristics, where turbines typically do not provide any power at
1653 all until a certain threshold speed is reached, and then the power output increases linearly with wind speed until the
1654 winds are so high that the turbine must shut down. This results in an annual turbine power output that is roughly linear
1655 to the mean annual wind speed for certain mean wind speed rangers. This would mean that, in some cases, an

1656 uncertainty in the annual wind or solar resource of $\pm 10\%$ results in an uncertainty of expected renewable energy
1657 technology output of approximately $\pm 10\%$.

1658

1659

1660 **5. Global change information and the HOMER DSS**

1661

1662 This section expands the discussion of the HOMER DSS to include the relationship of HOMER and its input data
1663 requirements with global change information

1664

1665 5a. Reliance of HOMER DSS global change information

1666

1667 As shown in the previous section, a number of observations that provide information on global change are also used in
1668 either direct or indirect ways as input to HOMER. These observations related primarily to the renewable energy
1669 resource information that is required for HOMER applications. Renewable energy system performance is highly
1670 dependent on the local energy resources available to the technologies. The extent and characteristics of these resources
1671 is driven by weather and local climate conditions, which happens to be the primary area in which Earth observational
1672 systems monitoring climate change are addressing. Thus, as users seek access to observations to support renewable
1673 energy resource assessments, they will invariably be seeking certain global change observational data.

1674

1675 Specifically, users will be seeking global change data related to atmospheric properties that support the assessment of
1676 solar and wind energy resources, such as wind and solar data, and atmospheric parameters important for estimating
1677 these data. For example, major data sets used in solar and wind energy assessments include long-term reanalysis data,
1678 climatological surface weather observations, and a variety of satellite observations from both active and passive
1679 onboard remote sensors.

1680

1681 Key factors in affecting the choice of these observational data are their relevance to conducting reliable solar and wind
1682 energy resource assessment, their ease of access, and low or no cost to the user. The extensive list of observational data

1683 being used in the assessment of renewable energy resources represents strong leveraging of major, taxpayer-supported
1684 observational programs that are geared primarily for global change assessment.

1685

1686 There is also an important consideration regarding the potential influence of long-term climate change on the renewable
1687 energy resources that are used as input into HOMER. Through the Intergovernmental Panel on Climate Change there
1688 has been a significant improvement in the reliability and spatial resolution of General Circulation Models (GCM) used
1689 to estimate the impacts of greenhouse gas emissions on climate change. As weather patterns change under changing
1690 climate conditions, wind and solar energy resources at a specific location can also change over time. The GCM results
1691 indicate that these renewable energy resources can be measurably different 50 to 100 years from now than today in
1692 specific locations and regions. These changes may have a noticeable impact on the results of HOMER simulations in
1693 the future; however, significant uncertainties exist in GCM results. Until these uncertainties are reduced sufficiently,
1694 implementation of GCM results will produce unreliable HOMER simulations.


1695

1696 5b. How the HOMER DSS can support climate-related management decision-making among US government agencies

1697

1698 Although HOMER was not intentionally designed to be a climate-related management decision-making tool, the
1699 HOMER DSS has attributes that can support these decisions. For example, as we explore mechanisms for mitigating
1700 the growth of carbon emissions in the atmosphere, the HOMER DSS can be deployed to evaluate how renewable energy
1701 systems can be used cost-effectively to displace energy systems dependent on fossil fuels. Clearly, the science results
1702 and global change data and information products coming out of our reanalysis and satellite-borne programs are of
1703 critical importance to HOMER for supporting this decision-making process. Given that the pertinent observational data
1704 sets have been developed primarily by federal agencies, these data sets tend to be freely available or available at a
1705 relatively small cost, given the costs involved in making the observations in the first place. However, as we have noted
1706 in previous sections, the use of global change observations as input to the resource assessment data required by
1707 HOMER is not the optimal choice of data; ideally, *in-situ* (site-specific) measurements of wind and solar data relevant
1708 to the technologies being analyzed would be the most useful and accurate data to have for HOMER, if they were
1709 available.

1710 Figure 1: Example of HOMER output graphic. The column on the left provides a diagram showing the load
1711 characteristics and the types of equipment considered to meet the load. The optimal system design graphic shows the
1712 range within specified diesel fuel prices and wind energy resources for which various system types are most economical
1713 (e.g., a wind/diesel/battery system becomes the most optimal configuration to meet the load requirement for wind
1714 speeds greater than 5 m/s and fuel costs at 0.45 to 0.75\$/l.
1715




1716
1717

1718 Figure 2. Example of ocean wind resource assessment output for the offshore regions of Pakistan. These data were
1719 derived from the SeaWinds scatterometer aboard NASA's QuickSCAT satellite. The assessment provides estimated
1720 mean annual wind speeds at 10-m above the ocean surface, averaged over the period 1988 to 2002.

1721

1722



1723

1724 Table 1: Major Global Data Sets Used by NREL for Wind Resource Assessment

1725

Data Set	Type of Information	Source	Period of Record
Surface station data	Surface observations from more than 20,000 stations worldwide	NOAA/NCDC	Variable up to 2006
Upper air station data	Rawinsonde and pibal observations at 1,800 stations	NCAR	1973–2005
Satellite-derived ocean wind data	Wind speeds at 10 m above the ocean surface gridded to 0.25 ⁰	NASA/JPL	1988–2006
Marine climatic atlas of the world	Gridded (1.0 ⁰) statistics of historical ship wind observations	NOAA/NCDC	1854–1969
Reanalysis upper air data	Model-derived gridded (~200-km) upper air data	NCAR-NCEP	1958–2005
Global upper air climatic atlas	Model-derived gridded (2.5 ⁰) upper air statistics	NOAA/NCDC	1980–1991
Digital geographic data	Political, hydrograph, etc.	ESRI	N/A
Digital terrain data	Elevation at 1-km spatial resolution	USGS/EROS	N/A
Digital land cover data	Land use/cover and tree cover density at 0.5-km resolution	NASA/USGS	N/A

1726

1727 Table 2: Major global data sets used for solar resource assessments

1728

1729

Data Set	Type of Information	Source	Period of Record
Surface station data	Surface cloud observations from more than 20,000 stations worldwide	NOAA/NCDC	Variable up to 2006
World Radiation Data Center	Surface radiation observations from over 1,000 stations worldwide	WRDC, St. Petersburg	1964–1993
Satellite imagers	Imagery from the visible channel of geostationary weather satellites, 1-km resolution	NASA/NOAA	1997– present
International Satellite Cloud Climatology Project	Used in the 1 ⁰ global surface solar energy meteorological data set	NASA/SSE	1983–2003
AERONET	Observations of aerosol optical depth from around the world	NASA/Goddard	Variable depending on station
GACP	Aerosol optical depths (generally over oceans) at 1 ⁰ x 1 ⁰ from AVHRR data	NASA	1981–2005
MODIS, MISR, TOMS	Aerosol optical depth	NASA	Variable since

			1980s
GOCART	Aerosol optical depth for turbid areas	NASA	March 30- May 3, 2001
GADS	Aerosol optical depth derived from theoretical calculations and proxies		Compilation of Measurements & Models
Digital geographic data	Political, hydrography, etc.	ESRI	N/A
Digital terrain data	Elevation at 1-km spatial resolution	USGS/EROS	N/A
Digital land cover data	Land use/cover and tree cover density at 0.5-km resolution	NASA/USGS	N/A

1730

Chapter 4

Decision Support for Public Health

Lead Author: Gregory E. Glass

1. Introduction

Public health is an approach to protect and improve the health of community members by preventive medicine, health education, control of communicable diseases, application of sanitary measures, and monitoring of environmental hazards (<http://www.answers.com/topic/public-health?cat=health>). This overall task is achieved by assessing and monitoring populations at risk to identify health problems and establishing priorities, to formulate policies to solve identified problems and to ensure populations have access to appropriate care, including health promotion, disease prevention, and evaluation of care. During the past century, the notable public health achievements as identified by the US Centers for Disease Control and Prevention (CDC) include vaccinations and treatments against infectious diseases, injury prevention strategies, reduced occupational exposures to toxins, improved food and water safety, decreases in childhood and maternal mortality, and safer water sources. Thus, many of the key issues related to public health are incorporated in previous chapters in this report, though they may not be characterized as public health. Regardless, public health may represent a key factor in problem solving under climate change situations. Many of the anticipated public health consequences of climate change are due to the influences of temperature and precipitation patterns, as well as land cover with consequences for the affected human communities. For example, changes in the availability of food resources and the quality of drinking water are anticipated to directly affect nutritional status, the spread of communicable infectious agents, and the impacts of poor air quality on vulnerable populations and in extreme situations the creation of “environmental refugees” (Huntingford *et al.*, 2007).

1758 Because public health is an important outcome component of decision support tools (DST) involving air
1759 quality, water management, energy management and agricultural efficiency issues, it was decided to focus on a
1760 unique public health aspect of DST/DSS by examining infectious disease systems. Infectious diseases remain
1761 a significant burden to populations both globally, as well as within the US. Some of these, such as syphilis and
1762 measles involve a relatively simple dynamic of the human host population and the parasite—be it a virus,
1763 bacterium, or other micro-organism. These diseases, therefore, tend to be influenced by social behavior and
1764 the ability to provide resources and of health education to significantly alter human behavior. However, other
1765 disease systems include additional species for their successful transmission—either wildlife species that
1766 maintain the micro-organism (zoonoses) or there are insect or arthropod vectors that serve to transmit the
1767 parasites either among people or from the wildlife to people (vector-borne diseases).

1768 Some of the most significant diseases globally are vector-borne or zoonotic diseases. Examples include
1769 malaria and dengue. In addition, many newly recognized (i.e., emerging) diseases either are zoonoses, such as
1770 SARS, or appear to have been derived from zoonoses that became established in human populations (e.g.,
1771 HIV). Changes in rates of contact between component populations of these disease systems alter the rates of
1772 infectious disease (Glass 2007). Many of these changes come about through activities involving the movement
1773 of human populations into areas where these pathogen systems normally occur or they can occur because
1774 people introduce materials with infectious agents into areas where they were not known previously (Gubler et
1775 al. 2001). The introduction of West Nile virus from its endemic area in Africa, the Middle East, and Eastern
1776 Europe into North America and its subsequent spread across the continent is a recent example. The impacts of
1777 the virus on wildlife, human, and agricultural production are an excellent example of the economic
1778 consequence of such emergent disease systems.

1779 More recently, attention has focused on the potential impact that climate change could have on
1780 infectious disease systems, especially those with vector or zoonotic components (e.g., Gubler *et al.*, 2001).
1781 Alterations in climate could impact the abundances or interactions of vector and reservoir populations, or the
1782 way in which human populations interact with them (Gubler, 2004). In addition, there is speculation that
1783 climate change will alter the locations where disease systems are established, shifting the human population
1784 that is at risk from these infectious diseases (e.g., Brownstein *et al.*, 2005a; Fox, 2007)

1785 Unlike many of the other applications in this report where Earth observations and modeling are of
1786 growing importance, the use of Earth observations by the public health community has been sporadic and
1787 incomplete. Although early demonstrations showed their utility for identifying locations and times that vector-
1788 borne diseases were likely to occur (e.g., Linthicum *et al.*, 1987; Beck *et al.*, 1997), growth of their application
1789 has been comparatively slow. Details of the barriers to implementation include the need to “scavenge” data
1790 from Earth observation platforms, as none of these are designed for monitoring disease risk. This is not an
1791 insurmountable problem and in fact, only few applications for Earth observations have dedicated sensors.
1792 However, disease monitoring requires a long history of recorded data to provide information concerning the
1793 changes in population distribution and the environmental conditions associated with outbreaks of disease.
1794 Detailed spectral and spatial data need to be of sufficient resolution and the frequency of observations must be
1795 high enough to enable identification of changing conditions (Glass 2007). As a consequence, many DSTs
1796 undergoing development have substantial integration of Earth observations but lack an end-to-end public
1797 health outcome, particularly when focusing on infectious diseases. Therefore, the Decision Support System to
1798 Prevent Lyme Disease (DDSPL) supported by the CDC and Yale University was selected to demonstrate the
1799 potential utility of these systems within the context of climate change science. Lyme disease is a vector-borne,
1800 zoonotic bacterial disease. In the US it is caused by the spirochete, *Borrelia burgdorferi*, and it is the most
1801 common vector-borne disease with tens of thousands of reported cases annually (Piesman and Gern 2004).
1802 Most human cases occur in the Eastern and upper Mid-West portions of the US, although there is a secondary
1803 focus along the West Coast of the country. In the primary focus, the black-legged tick (or deer tick), of the
1804 genus *Ixodes*, is most often found infected with *B. burgdorferi*.

1805

1806

1807 2. Description of DDSPL

1808 The diverse ways in which Lyme disease presents itself in different people has made it a public health
1809 challenge to ensure that proper priorities are established, to formulate policies to solve the problem, and to
1810 ensure that populations have access to appropriate care. The CDC uses DDSPL to address questions related to
1811 the likely distribution of Lyme disease east of the 100th meridian, where most cases occur (Brownstein *et al.*,
1812 2003). This is done by identifying the likely geographic distribution of the primary tick vector (the black-

1813 legged) tick in this region. DDSPL uses field reports of the known distribution of collected tick vectors, as
1814 well as sites with repeated sampling without ticks as the outcome space. DDSPL uses satellite data, and
1815 derived products such as land cover characteristics, and census boundary files and meteorological data files to
1816 identify the best statistical predictor of the presence of black-legged ticks within the region. Land cover is
1817 derived from multi-date Landsat TM imagery and 10-m panchromatic imagery.

1818 DDSPL combines the satellite and climate data with the field survey data of *Ixodes* ticks sampled at
1819 locally sampled sites throughout the region (Brownstein *et al.*, 2003) or from rates of reported cases of Lyme
1820 disease (Brownstein *et al.*, 2005b) in spatially explicit statistical models to generate assessment products of the
1821 distribution of the tick vector or human disease risk, respectively. These models are validated by field surveys
1822 in additional areas and the sensitivity and specificity of the results determined (figure 1). Thus, the DDSPL is
1823 primarily a DST for prioritizing the likely geographic extent of the primary vector of Lyme disease in this
1824 region (figures 1 and 2). It currently stops short of characterizing the risk of disease in the human population
1825 but is intended to delimit the area within which Lyme disease (and other diseases caused by additional
1826 pathogens carried by the ticks) might occur (Figure 2). Researchers at Yale University are responsible for
1827 developing and validating appropriate analytical methods to develop interpretations that can deal with many of
1828 the challenges of spatially structured data, as well as the acquisition of Earth science data that are used for
1829 model DDSPL predictions. The distinction between the presence/abundance of the tick vector and actual
1830 human risk relies on the effects of human population abundance and behavioral heterogeneity (e.g., work or
1831 recreational activity) that can alter the contact rate between the tick vector and susceptible humans. However,
1832 such detailed human studies (especially behavioral heterogeneity) are typically not available (Malouin *et al.*,
1833 2003). In Brownstein *et al.* (2005b) analysis, they found that although the entomological risk (the abundance
1834 of infected ticks) increased with landscape fragmentation, the human incidence of Lyme disease decreased,
1835 thus indicating there is a complex relationship between the landscape, the population of ticks, and the human
1836 response resulting in the health outcome.

1837

1838 3. Potential Future Use and Limits

1839 Future use of DDSPL depends to a great extent on public health policy decisions exterior to the DST. The
1840 perspective of the role that Lyme disease prevention rather than treatment of diseased individuals will play is a

1841 key aspect of the importance that DDSPL will experience. For example, studies have shown that even in Lyme
1842 disease endemic regions, risk communication often fails to reduce the likelihood of infection (Malouin, *et al.*,
1843 2003). In principle, policy makers may decide that it is more cost effective to provide improved treatment
1844 modalities rather than investing in educational programs that fail to reduce disease burden. Alternatively, the
1845 development of vaccines is time consuming, costly, and may have additional risks of unacceptable side effects
1846 that affect the likelihood that this would be a policy choice. Thus, depending on policy decisions and the
1847 effects of alternative interventions, the DDSPL might be used to forecast risk areas for educational
1848 interventions, to inform health care providers in making diagnoses, or to plan mass vaccination campaigns.

1849 Currently, the removal of the licensed Lyme disease vaccine from the general public has eliminated this as
1850 a strategy to reduce the disease burden. The apparent lack of impact of targeted education also makes this a
1851 less likely strategy. Thus, the extent to which treatment modalities rather than prevention of infection will drive
1852 the public health response in the near future will play a major role in the use of DDSPL. However, even if the
1853 decision is made to focus on treatment of potentially infected individuals, DDSPL may still be useful by
1854 identifying regions where disease risk may be low, helping health care workers to focus clinical diagnoses on
1855 alternate causes.

1856 Presuming that the DST continues to be used, the need for alternative/improved Earth science data to
1857 clarify environmental data for DDSPL such as land cover, temperature, and moisture regimes is currently
1858 uncertain. The present system reports a sensitivity of 88 percent and specificity of 89 percent—generally
1859 considered a highly satisfactory result. Sensitivity and specificity are considered the two primary measures of a
1860 method's validity in public health analyses. Sensitivity in the DDSPL model refers to the expected proportion
1861 of times (88 percent) that ticks would be found when field surveys were conducted at sites that the DDSPL
1862 predicted they should occur. Specificity refers to the proportion of times (89 percent) that a survey would not
1863 be able to find ticks at sites where the DDSPL excluded them from occurring. These two measures provide an
1864 estimate of the “confidence” the user can have in the DST prediction (Selvin 1991). These analyses extended
1865 geographically from the East Coast to the 100th meridian and were validated by field sampling for the presence
1866 of *Ixodes* ticks at sites throughout the region.

1867 Typically, patterns of weather regimes appear to have a greater impact on distribution than more
1868 detailed information on land cover patterns. However, some studies indicate that fragmentation of forest cover

1869 and landscape distribution at fairly fine spatial resolution can substantially alter patterns of human disease risk
1870 (Brownstein *et al.*, 2005b). These results also suggest that human incidence of disease may, in some areas of
1871 high transmission, be decoupled from the model constructed for vector abundance, reemphasizing the
1872 distinction between a key component (the vector) and actual human risk. When coupled with the stated
1873 accuracy of the DDSPL in identifying vector distribution, this would suggest that future efforts will probably
1874 require an additional model structure that includes sociological/behavioral factors of the human population that
1875 puts it at varying degrees of risk. An additional limit of the DDSPL is that it does not explicitly incorporate
1876 human health outcomes in its analyses. In part, this reflects a public health infrastructure issue that limits
1877 detailed information on the distribution of human disease to (typically) local and state health agencies. For
1878 example, confidentiality of health records, including detailed locational data, such as home addresses, are often
1879 shielded in the absence of explicit permission. This makes establishing the relationship between monitored
1880 environmental conditions and human health outcomes difficult. One solution is to aggregate data to some
1881 jurisdictional level. However, this produces the well know “ecological fallacy” in establishing relationships
1882 between environmental factors and health outcomes (Selvin 1991). With appropriate planning or the
1883 movement of the technology into local public health agencies, these challenges could be overcome. Some
1884 localized data (e.g., Brownstein *et al.*, 2005b) of human health outcomes have been used to evaluate the utility
1885 of DDSPL and indicate that there is good potential for the DSS to provide important information on local risk
1886 factors.

1887

1888

1889 **4. Uncertainty**

1890 Uncertainty in decision making from DDSPL is based on the results of statistical analyses in which
1891 standard statistical models with spatially explicit components, such as autologistic intercepts of logistic
1892 models, are used to account for spatial autocorrelation in outcomes. The statistical analyses are well-supported
1893 theoretically. Typical calibration approaches involve model construction followed by in-field validation.
1894 Accuracy of classification is then assessed in a sensitivity-specificity paradigm.

1895 However, little attention is paid in the current model to assessing uncertainty in the environmental data
1896 obtained from remotely sensed (or even *in situ*) monitors of the environment. For example, most of the

1897 derivative data, such as land cover, may change with population growth and development. In addition, the use
1898 of average environmental conditions provide an approximate characterization of local edaphic conditions that
1899 may affect the abundance of the the tick vectors.

1900 Whether these are the primary sources of “error” in the sensitivity and specificity results (although these
1901 are considered excellent results) of the DDSPL is not addressed and is an area the public health applications
1902 need to consider in future applications. Alternatively, there are biological reasons for the errors in the model,
1903 including the interaction of climatic factors and tick activity that may be responsible for sites predicted to have
1904 ticks that were not found to have them. To resolve some of the biological/environmental issues, validation is
1905 ongoing.

1906 There also are a number of public health issues that affect the certainty of the DDSPL (and any DST)
1907 that are extrinsic to the system or tool. Accuracy in clinical diagnoses (both false positives and negatives), as
1908 well as reporting accuracy can affect the evaluation of the tool’s utility. Currently, this is an issue of serious
1909 contention and forms part of the rationale for focusing on accurately identifying the distribution of the primary
1910 tick vector, as an integral step in delimiting the distribution of the disease and evaluating needs for the
1911 community.

1912

1913 **5. Global Change Information and DDSPL**

1914 The relationship between climate and public health outcomes is complex. It is affected both by the
1915 direction and strength of the relationship between climatic variability and the component populations that make
1916 up a disease system, as well as the human response to changes in disease risk (Gubler 2004).

1917 The DDSPL is one of the few public health DSTs that has explicitly evaluated the potential impact of
1918 climate change scenarios on this infectious disease system. Assuming that evolutionary responses of the black-
1919 legged tick, *B. burgdoferi* and the reservoir zoonotic species remains little changed under rapid climate change,
1920 Brownstein *et al.*, (2005a) evaluated anticipated changes in the distribution and extent of disease risk.

1921 This analysis used the basic climate-land cover suitability model developed for DDSPL and selected the
1922 Canadian Global Coupled Model (CGCM1) under two historically forced integrations. The first with a 1
1923 percent per year increase in greenhouse gas emissions and the second with greenhouse gas and sulfate aerosol
1924 changes, resulted in a 4.9 and 3.8° Celsius increase in global mean temperature by the year 2080. Near (2020),

1925 mid (2050) and farpoint (2080) outcomes were evaluated (Figure 3). The choice of CGCM1 was based on the
1926 Intergovernmental Panel on Climate Change criteria for vintage, resolution, and validity (Brownstein *et al.*,
1927 2005a).

1928 Extrapolation of the analyses suggest that the tick vector will experience a significant range expansion
1929 into Canada but will also experience a likely loss of habitat range in the current southern portion of its range
1930 (figure 3). This loss of range is thought to be due to impact of increased temperatures causing decreased
1931 survival in ticks when they are off their feeding hosts. It also is anticipated that its range will shift in the central
1932 region of North America – where it is currently absent. When coupled with the anticipated continued human
1933 movement to more southern portions of the country, the numbers of human cases are expected to show an
1934 overall small decrease.

1935 These long-range forecasts disguise a more dynamic process with ranges initially decreasing during near
1936 and mid-term timeframes. This range reduction is later reversed in the long-term producing the overall pattern
1937 described by the authors. The impact in range distribution also produces an overall decrease in human disease
1938 risk as suitable areas move from areas of primary human concentration to areas that are anticipated to be less
1939 well populated.

1940 Thus, DSS similar to those developed for Lyme disease have the potential for providing both near- and
1941 far-term forecasts of potential infectious disease risk that are so important for public health planning. In
1942 addition, detailed studies (e.g. Brownstein *et al.*, 2005b) provide public health agencies with important
1943 information on drivers of human risk that have been difficult to obtain by other means. As a consequence,
1944 DSS using remotely sensed data sources either in part or whole have the potential to significantly improve the
1945 health of communities.

1946 The primary challenges for the Earth science community involve understanding the needs of the public
1947 health community for the appropriate data at the appropriate spatial, temporal, and spectral scales. This will
1948 involve understanding a historically entrenched set of methodologies for interpreting health data and
1949 establishing causal relationships between inputs (environmental data) and outputs (health outcomes). In
1950 addition, there is the challenge of performing these tasks in the presence of limited resources for a community
1951 that has little cultural understanding of both the strengths and limitations of the data derived from these
1952 sources.

1953

1954




Figure 1. Relationship between the occurrence of black-legged tick presence at a site and minimum temperature (top) and evaluation of model (bottom). From Brownstein et al. 2003 Env. Hlth Perspect. **Top Panel:** Log odds plot for relationship between *I. Scapularis* population maintenance and minimum temperature (T). Minimum temperature showed a strong positive association with odds of an established *I. Scapularis* population. According to good-ness of fit testing, the relationship was fit best by a fourth order polynomial regression ($R^2 = 0.97$) $\text{Log odds} = 0.0000067^4 + 0.00027^3 - 0.0027T^2 + 0.0002T - 0.8412$. **Bottom Panel:** ROC Plot describing the accuracy of the auto logistic model. This method graphs sensitivity versus 1-specificity over all possible cutoff probabilities. The AUC is a measure of overall fit, where 0.5 {a 1:1 line} indicates a chance performance {dashed line}. The plot for the auto logistic model significantly outperformed the chance model with an accuracy of 0.95 { $p < 0.00005$ }.

1955





Figure 2. Forecast geographic distribution of the black-legged tick vector east of the 100th meridian in the United States for DSSPL. From Brownstein et al (2003) *Envr. Hlth. Perspect.* 2a. New distribution map for *I. Scapularis* in the United States. To determine whether a given cell can support *I. Scapularis* populations, a probability cutoff point for habitat suitability from the auto logistic model was assessed by sensitivity analysis. A threshold of 21% probability of establishment was selected, giving a sensitivity of 97% and a specificity of 86%. This cutoff was used to reclassify the reported distribution map {Dennis et al. 1998}. The auto logistic model defined 81% of the reported locations {n=427} as established and 14% of the absent areas {n=2,327} as suitable. All other reported and absent areas were considered unsuitable. All areas previously defined as established maintained the same classification.

1956
 1957
 1958
 1959
 1960
 1961
 1962
 1963



1964

Figure 3. Forecast change in black-legged tick distribution in Eastern and Central North America under climate change scenarios using DSSPL. From Brownstein et al (2005a) EcoHealth

1965
1966
1967
1968
1969
1970
1971
1972
1973
1974
1975
1976
1977
1978
1979
1980
1981
1982
1983
1984
1985
1986
1987
1988
1989
1990
1991
1992

Climate Change Science Program

Uses and Limitations of Observations, Data, Forecasts, and Other Projections in Decision Support for Selected Sectors and Regions

Chapter 5. “Decision Support for Water Resources Management”

Holly C. Hartmann

Department of Hydrology and Water Resources, The University of Arizona

1. Introduction

Water resource managers have long been incorporating information related to climate in their decisions. The tremendous, regionally ubiquitous, investments in infrastructure to reduce flooding (e.g., levees and reservoirs) or assure reliable water supplies (e.g., reservoirs, groundwater development, irrigation systems, water allocation, and transfer agreements) reflect societal goals to mitigate the impacts of climate variability at multiple time and space scales. As the financial, political, social, and environmental costs of infrastructure options have become less tractable, water management institutions have undergone comprehensive reform, shifting their focus to optimizing operations of existing projects and managing increasingly diverse, and often conflicting, demands on the services provided by water resources (Bureau of Reclamation [BOR], 1992; Beard, 1993; Congressional Budget Office, 1997; Stakhiv, 2003; National Research Council [NRC], 2004). Governments have also made substantial investments to improve climate information and understanding over the past decades through satellites, *in situ* measuring networks, supercomputers, and research programs. National and international programs have explicitly identified as an important objective ensuring that improved data products, conceptual models, and predictions are useful to the water resources management community (Endreny et al., 2003; Lawford et al., 2005). Although exact accounting is difficult, potential values associated with appropriate use of accurate hydrometeorologic predictions generally range from the millions to the billions of dollars (e.g., National Hydrologic Warning Council, 2002). There are also non-monetary values associated with more efficient, equitable, and environmentally sustainable decisions related to water resources.

1993 Droughts, floods, and increasing demands on available water supplies continue to create concern, and even
1994 crises, for water resources management. Many communities have faced multiple hydrologic events that were earlier
1995 thought to have low probabilities of occurrence (e.g., NRC, 1995), and long-term shifts in streamflows have been
1996 observed (Lettenmaier et al., 1994; Lins and Slack, 1999; Douglas et al., 2000), leading to questions about the relative
1997 impacts of shifts in river hydraulics, land use, and climate conditions.

1998 Until the last two decades, climate was viewed largely as a collection of random processes, and this paradigm
1999 informed much of the water resource management practices developed over the past 50 years that persist today.
2000 However, climate is now recognized as a chaotic process, shifting among distinct regimes with statistically significant
2001 differences in average conditions and variability (Hansen et al., 1997). As instrumental records have grown longer and
2002 extremely long time-series of paleoclimatological indicators have been developed (Ekwurzal, 2005), they increasingly
2003 belie one of the fundamental assumptions behind most extant water resources management—stationarity. Stationary
2004 time series have time-invariant statistical characteristics (e.g., mean or variance), meaning that different parts of the
2005 historical record can be considered equally likely. Within the limits posed by sampling, statistics computed from
2006 stationary time series can be used to define a probability distribution that will also then faithfully represent expectations
2007 for the future (Salas, 1993).

2008
2009 Further, prospects for climate change due to global warming have moved from the realm of speculation to
2010 general acceptance (Intergovernmental Panel on Climate Change [IPCC] 1990, 1995a, 2001a, 2007). The potential
2011 impacts of climate on water resources, and their implications for management, have been central topics of concern in
2012 climate change assessments (e.g., EPA, 1989; IPCC, 1995b, 2001b; National Assessment Synthesis Team, 2000; Gleick
2013 and Adams, 2000; Barnett *et al.*, 2004). These studies are becoming increasingly confident in their conclusions that the
2014 future portends statistically significant changes in hydroclimatic averages and variability.

2015 There has been persistent and broad disappointment in the extent to which improvements in hydroclimatic
2016 science from large-scale research programs have affected resource management practices in general (Pielke, 1995,
2017 2001; NRC, 1998a, 1999a) and water resource management in particular (NRC, 1998b, 1999b,c). For example,
2018 seasonal climate outlooks have been slow to be entered into the water management decision processes, even though
2019 they have improved greatly over the past 20 years (Hartmann et al., 2002a, 2003). Water managers have been even more
2020 resistant to incorporating notions of hydrologic non-stationarity in general and climate change in particular in decision

2021 processes. Until recently, hydrologic analysis techniques have been seen as generally sufficient (e.g., Matalas, 1997;
2022 Lins and Stakhiv, 1998), especially in the context of slow policy and institutional evolution (Stakhiv, 2003). However,
2023 an inescapable message for the water resource management community is the inappropriateness of the stationarity
2024 assumption in the face of climate change.

2025 Several ongoing efforts are leading the way forward to establish more effective ways of incorporating climate
2026 understanding and earth observations into water resources management (Pulwarty, 2002; Office of Global Programs,
2027 2004; NASA, 2005). While diverse in their details, these efforts seek to link hydroclimatological variability, analytical
2028 and predictive technologies, and water management decisions within an end-to-end context extending from
2029 observational data through large-scale analyses and predictions, uncertainty evaluation, impacts assessment,
2030 applications, and evaluations of applications (e.g., Young, 1995; Miles et al., 2000). Some end-to-end efforts focus on
2031 cultivating information and management networks; designing processes for recurrent interaction among research,
2032 operational product generation, management, and constituent communities; and developing adaptive strategies for
2033 accommodating climate variability, uncertainty, and change. Other end-to-end efforts focus on the development of
2034 decision support tools (DST) that embody unique resource management circumstances to enable formal and more
2035 objective linkages between meteorological, hydrologic, and institutional processes. Typically, end-to-end DST
2036 applications are developed for organizations making decisions with high-impact (e.g., state or national agencies) or
2037 high-economic value (e.g., hydropower production) and that possess the technical and managerial abilities to efficiently
2038 exploit research advances (e.g., Georgakakos et al, 1998, 2004, 2005; Georgakakos, 2006). If linked to socioeconomic
2039 models incorporating detailed information about the choices open to decision-makers and their tolerance for risk, these
2040 end-to-end tools could also enable explicit assessment of the impacts of scientific and technological research advances.

2041 This chapter describes a river management DST, RiverWare, which facilitates coordinated efforts among the
2042 research, operational product generation, and water management communities. RiverWare emerged from an early and
2043 sustained effort by several federal agencies to develop generic tools to support the assessment of water resources
2044 management options in river basins with multiple reservoirs and multiple management objectives (Frevert et al., 2006).
2045 RiverWare was selected for use as a case study because it has been used in a variety of settings, by multiple agencies,
2046 over a longer period than many other water management DSTs. Furthermore, RiverWare can explicitly accommodate a
2047 broad range of resource management concerns (e.g., flood control, recreation, navigation, water supply, water quality,
2048 and power production). RiverWare can also consider perspectives ranging from day-to-day scheduling of operations to

2049 long-range planning and can accommodate a variety of climate observations, forecasts, and even climate change
2050 projections. RiverWare can incorporate hydrologic risk, whereby event consequences and their magnitudes are mediated
2051 by their probability of occurrence, in strategic planning applications and design studies, which can offer a way forward
2052 for decision makers reluctant to shift away from use of traditional, stationarity-based, statistical analysis of historical
2053 data (Lee, 1999; Davis and Pangburn, 1999).

2054

2055 **2. Description of RiverWare**

2056 RiverWare is a software framework used to develop detailed models of how water moves and is managed
2057 throughout complex river basin systems. RiverWare applications include physical processes (e.g., streamflow, bank
2058 storage, and solute transport), infrastructure (e.g., reservoirs, hydropower generating turbines, spillways, and diversion
2059 connections), and policies (e.g., minimum instream flow requirements and trades between water users) (Zagona et al.,
2060 2001, 2005). At a minimum, RiverWare applications require streamflow hydrographs as input for multiple locations
2061 throughout a river system. While hydrographs can be generated within the DST, they can also be input from other
2062 sources, with the latter approach being especially important in advanced end-to-end assessments. Detailed discussion of
2063 the role of observations and considerations of global change using RiverWare are discussed in later sections. RiverWare
2064 can be applied to address diverse water management concerns, including real-time operations, strategic planning for
2065 seasonal to interannual variability in water supplies and demands, and examining impacts of hydrologic non-
2066 stationarity. Because infrastructure, management rules, and policies can be easily changed, RiverWare also allows
2067 examination of alternative options for achieving management objectives over short-, medium-, and long-term planning
2068 horizons.

2069 RiverWare was developed by the University of Colorado-Boulder's Center for Advanced Decision Support for
2070 Water and Environmental Systems (CADSWES) in collaboration with the BOR, Tennessee Valley Authority, and the
2071 Army Corps of Engineers (Frevert et al., 2006). CADSWES continues to develop and maintain the RiverWare software,
2072 as well as offer training and support for RiverWare users (see <http://cadswes.colorado.edu>). According to CADSWES,
2073 RiverWare is used by more than 75 federal and state agencies, private sector consultants, universities and research
2074 institutes, and water districts, among others.

2075

2076 *Example Applications*

2077 Consistent with the intent of its original design, the use of RiverWare varies widely, depending on the specific
2078 application. An early application was its use for scheduling reservoir operations by the Tennessee Valley Authority
2079 (Eshenbach et al., 2001). In that application, RiverWare was used to define the physical and economic characteristics of
2080 the multi-reservoir system, including power production economics, to prioritize the policy goals that governed the
2081 reservoir operations and to specify parameters for linear optimization of system objectives. In another application,
2082 RiverWare was used to balance the competing priorities of minimum instream flows and consumptive water use in the
2083 operation of the Flaming Gorge Reservoir in Colorado (Wheeler et al., 2002).

2084 While day-to-day scheduling of reservoir operations is more a function of weather than climate, the use of
2085 seasonal climate forecasts to optimize reservoir operations has long been a goal for water resources management.
2086 RiverWare is being implemented for the Truckee-Carson River basin in Nevada to investigate the impact of
2087 incorporating climate outlooks into an operational water management framework that prioritizes irrigation water
2088 supplies, interbasin diversions, and fish habitat (Grantz et al., 2007). Another example application to the Truckee-
2089 Carson River using a hypothetical operating policy indicated that fish populations could benefit from purchases of water
2090 rights for reservoir releases to mitigate warm summer stream temperatures resulting from low flows and high air
2091 temperatures (Neumann et al., 2006).

2092 RiverWare has also been used to evaluate politically charged management strategies, including water transfers
2093 proposed in California's Quantification Settlement Agreement and the BOR's Inadvertant Overrun Policy, maintaining
2094 instream flows sufficient to restore biodiversity in the Colorado River delta, and conserving riparian habitat while
2095 accommodating future water and power development in the BOR Multiple Species Conservation Program (Wheeler et
2096 al., 2002). RiverWare also played a key role in negotiations by seven western states concerning how the Colorado River
2097 should be managed and the river flow should be distributed among the states during times of drought. The BOR
2098 implemented a special version of the RiverWare model of the Colorado River and its many reservoirs, diversions, and
2099 watersheds (Jerla, 2005). The model was used to provide support to the Basin States Modeling Work Group Committee
2100 over an 18-month period, as they assessed different operational strategies under different hydrologic scenarios,
2101 including extreme drought (U.S. Department of Interior, 2007).

2102

2103 *Implementation*

2104 RiverWare requirements are multi-dimensional. A specific river system and its infrastructure operating policies
2105 are defined by data files supplied to RiverWare. This allows incorporation of new basin features (e.g., reservoirs),
2106 operating policies, and hydroclimatic conditions without users having to write software code. Utilities within RiverWare
2107 enable users to automatically execute many simulations, including accessing external data or exporting results of model
2108 runs. Users can also write new modules that CADSWES can integrate into RiverWare for use in other applications. For
2109 example, in an application for the Pecos River in New Mexico, engineers developed new methods and software code for
2110 realistic downstream routing of summer monsoon-related flood waves (Boroughs and Zagona, 2002). RiverWare is
2111 implemented for use on Windows or Unix Solaris systems, as described in the requirements document
2112 (<http://cadswes.colorado.edu/PDF/RiverWare/RecommendedMinimumSystemsRequirements.pdf>). An extensive
2113 manual is also available (<http://cadswes.colorado.edu/PDF/ReleaseNotes/RiverWareHelp.pdf>).

2114 RiverWare applications can be implemented by any group that can pay for access, both in terms of finances
2115 and educational effort. Development of RiverWare applications requires a site license from CADSWES. Significant
2116 investment is required to learn to use RiverWare as well. CADSWES offers two 3-day RiverWare training courses, an
2117 initial class covering general simulation modeling, managing scenarios, and incorporating policy options through rule-
2118 based simulation, and a second class covering rule-based simulation in more detail, creating basin policies, and
2119 examining water policy options. Costs for the original license, annual renewals, technical support, and training require
2120 several thousand dollars. The costs of licensing and learning RiverWare mean that small communities and civic groups
2121 are unlikely to implement their own applications for assessing water management options. Rather, large agencies with
2122 technical staff or the financial means to fund university research or consultants are the most frequent users of
2123 RiverWare. The agencies then mediate the access of stakeholders to assessments of water management options through
2124 traditional public processes (e.g., U.S. Department of Interior, 2007). Conflicts may arise in having academic research
2125 groups conduct analyses funded by stakeholder groups, with inherent tensions between the open publication of research
2126 required by academia and the limited access to results required by strategic negotiations among interest groups.

2127

2128 **3. Current and Future Use of Observations**

2129 The specific combination of observations used by a RiverWare application depends on both the decision
2130 context and the use of other models and DSTs to provide input to RiverWare that more comprehensively or accurately
2131 describes the character, conditions, and response of the river basin system. Figure 1 illustrates the information flow

2132 linking observations, RiverWare, other models and DSTs, and water management decisions; it shows that RiverWare
2133 has tremendous flexibility in the kinds of observations that could be useful in hydrologic modeling and river system
2134 assessment and management. The types of observations that may ultimately feed into RiverWare applications also
2135 depend on the timescale of the situation.

2136 A detailed discussion of the role of satellite observations in RiverWare applications and selected input models
2137 and DSTs (e.g., the BOR's ET Toolbox and Precipitation Runoff Modeling System [PRMS]) is given by the
2138 "Evaluation Report for AWARDS ET Toolbox and RiverWare Decision Support Tools" (Hydrological Sciences
2139 Branch, 2007). Briefly, RiverWare can use a combination of observations from multiple sources, including satellites,
2140 products derived from land-atmosphere or hydrologic models, and combinations of both. Satellite observations can
2141 assist models in estimating evapotranspiration, precipitation, snow water equivalent, soil moisture, groundwater storage
2142 and aquifer volumes, reservoir storage, and water quality, among other variables. Measurements from sensors aboard a
2143 variety of satellites are being considered for their usefulness within DST contexts and their impacts on reducing water
2144 management uncertainty, including the MODIS (Moderate Resolution Imaging Spectroradiometer) sensor aboard the
2145 Earth Observing System (EOS) Terra and Aqua satellites, Landsat TM data, Advanced Spaceborne Thermal Emission
2146 and Reflection Radiometer (ASTER), Shuttle Radar Topography Mission (SRTM), Advanced Microwave Scanning
2147 Radiometer–EOS (AMSR-E), Gravity Recovery and Climate Experiment (GRACE), and Tropical Rainfall Mapping
2148 Mission (TRMM), among others. Future and planned satellites with hydrologically relevant sensors and measurements
2149 include CloudSAT, the Global Precipitation Mission (GPM), and the National Polar-Orbiting Operational
2150 Environmental Satellite (NPOESS). Use of these observations can be enhanced by assimilating them into land surface
2151 models to produce spatially-distributed estimates of snowpack, soil moisture, evapotranspiration, energy fluxes, and
2152 runoff, which then provide inputs to RiverWare to base a more comprehensive assessment of river basin conditions. The
2153 land surface models include the Community Land Model (CLM), Mosaic, Noah, and VIC, among others, supported by
2154 NASA's Land Data Assimilation System (LDAS) and Land Information System (LIS) (NASA, 2006a).

2155 NASA has several pilot projects specifically focused on assessing the impact of satellite observations in a
2156 variety of hydrologic models and DSTs as they feed into RiverWare applications (NASA, 2005, 2006b, 2007). For
2157 example, one project is comparing Terra and Aqua MODIS snow cover products for the Yakima-Columbia River basins
2158 with land-based snow telemetry measurements, testing their use for LIS simulations that also use the North American
2159 LDAS, connecting assimilated snow data with the Modular Modeling System (MMS) Precipitation-Runoff Modeling

2160 System (PRMS), and then supplying the simulated runoff as inputs to RiverWare. Another project on the Rio Grande
2161 River basin is assessing MODIS and Landsat data to improve evapotranspiration estimates generated by the BOR DST,
2162 the Agricultural Water Resources Decision Support (AWARDS) ET Toolbox, which then provides water demand time
2163 series to RiverWare. While application of specific hydrologic models and observations depend on the specific
2164 RiverWare application, significant processing of both model and observations are required and can be resource
2165 intensive (e.g., calibration and aggregation/disaggregation).

2166 Operational scheduling of reservoir releases depend on orders of water from downstream users (e.g., irrigation
2167 districts) that are largely affected by day-to-day weather conditions as well as seasonally varying demands. In these
2168 cases, the important observations are the near real-time estimates of conditions within the river basin system (e.g., soil
2169 moisture or infiltration capacity), which affect the transformation of precipitation into runoff into the river system,
2170 relative to constraints on system operation (e.g., reservoir storage levels or water temperatures at specific river
2171 locations). Meteorological prospects are mediated by those placing the water orders or through short-term weather
2172 forecasts that may affect operations when the system is near some constraint (e.g., flood flows when reservoir levels are
2173 near peak storage capacity). In these situations, the important observations are recent extreme precipitation events and
2174 their location, which may be provided, separately or in some combination, by *in situ* monitoring networks, radar, or
2175 satellites.

2176 For mid-range applications, such as strategic planning for operations over the next season or year, outlooks of
2177 total seasonal water supplies are routinely used in making commitments for water deliveries, determining industrial and
2178 agricultural water allocation, and carrying out reservoir operations. In these applications, it is also important for water
2179 managers to keep track of the current state of the watershed. Such observations are often used as input to one of the
2180 many independent hydrologic models that can provide input to a specific RiverWare application. In these situations, the
2181 important observations are those that provide boundary or forcing conditions for the independent hydrologic models,
2182 including snowpack moisture storage, soil moisture, precipitation (intensity, duration, and spatial distribution), air
2183 temperature, humidity, winds, and other meteorological conditions.

2184 For long-term planning and design applications, observations are less important because the effects of recent
2185 conditions have less impact on long-term outcomes than future meteorological uncertainty, or even institutions at multi-
2186 decadal time scales. In these applications, accurate representation of anticipated natural hydroclimatological variability
2187 is important. In many western U.S. applications, observed streamflows are adjusted to remove the effects of reservoir

2188 management, interbasin diversions, and water withdrawals. The adjusted flows, termed “naturalized flows.” may be
2189 used as input to RiverWare applications to assess the impact of different management options. Use of naturalized flows
2190 is fraught with problems. A central issue is poor monitoring of actual human impacts, especially withdrawals,
2191 diversions, and return flows (e.g., from irrigation). Alternative approaches include the use of proxy streamflows (e.g.,
2192 from paleoclimatological indicators) or output from hydrologic modeling studies (Hartmann, 2005). For example,
2193 Tarboton (1995) developed hydrologic scenarios for severe sustained drought in the Colorado River basin based on
2194 streamflows reconstructed from centuries of tree-ring records; the scenarios were used in an assessment of management
2195 options using a precursor to the current RiverWare application to the Colorado River system.

2196 The usefulness of the observations used within RiverWare depends on the specific implementation, as well as
2197 the quality of the information itself. For example, one direct use of climate information for long-term planning includes
2198 hydrologic and hydraulic routing of “design storms” of various magnitudes and likelihoods, with the storms based on
2199 analyses of the available instrumental record (Urbanas and Roesner, 1993). However, those instrumental records have
2200 often been too short to adequately express climate variability and resulting impacts, regardless of the specific DSTs
2201 used to do the hydrologic or hydraulic routing. In short- and mid-range forecasting applications, the use of observations
2202 is mediated by the hydrologic model or DST that transforms weather and climate into streamflows, evaporative water
2203 demands, and other hydrologic processes. In these situations, from an operational perspective, the stream of
2204 observational inputs must be dependable, without downtime or large data gaps, and data processing, model simulation,
2205 and creation of forecast products must be fast and efficient. The usefulness of observations may be limited by other
2206 issues as well. The water resources management milieu is complex and diverse, and climate influences are only one
2207 factor among many affecting water management policies and practices. Factors limiting the use of observations or
2208 subsequent hydrologic model input to RiverWare for actual water management include lack of familiarity with the
2209 available information, disconnects between the specific information available (e.g., variables and spatiotemporal scales)
2210 and their relevance to decision makers, skepticism about the quality and applicability of information, conservative
2211 decision preferences due to accountability for poor consequences, and institutional impediments such as the inflexible
2212 nature of many multi-jurisdictional water management agreements (Changnon, 1990; Kenney, 1995; Pulwarty and
2213 Redmond, 1997; Pagano et al., 2001, 2002; Jacobs, 2002; Jacobs and Pulwarty, 2003; Rayner et al., 2005).

2214

2215

2216 **4. Uncertainty**

2217 The reliability of observations for driving hydrologic models that may provide input to RiverWare applications
2218 is the subject of much ongoing research. The hydrologic models, because they incompletely describe the physical
2219 relationships among important watershed components (e.g., vegetation processes that link the atmosphere and different
2220 levels of soil and surface and groundwater interactions), are themselves the subject of much research to determine their
2221 reliability. Streamflow and other hydrologic variables are intimately responsive to atmospheric factors, especially
2222 precipitation, that drive a watershed's behavior; however, errors in precipitation estimates are often amplified in the
2223 hydrologic response (Oudin et al., 2006).

2224 Obtaining quality precipitation estimates is a formidable challenge, especially in the western U.S. where
2225 orographic effects produce large spatial variability and where there is a scarcity of real-time precipitation gage data and
2226 radar beam blockage by mountains. In principal, outputs from atmospheric models can serve as surrogates for
2227 observations, as well as providing forecasts of meteorological variables that can be used to drive hydrologic models.
2228 One issue in integrating atmospheric model output into hydrologic models for small watersheds (<1000 km²) is that the
2229 spatial resolution of atmospheric models is lower than the resolution of hydrologic models. For example, quantitative
2230 precipitation forecasts (QPF) produced by some atmospheric models may cover several thousand square kilometers, but
2231 the hydrologic models used for predicting daily streamflows require precipitation to be downscaled to precipitation
2232 fields for watersheds covering only tens or hundreds of square kilometers. One approach to produce output consistent
2233 with the needs of hydrologic models is to use nested atmospheric models, whereby outputs from large scale but coarse
2234 resolution models are used as boundary conditions for models operating over smaller domains with higher resolution.
2235 However, the error characteristics of atmospheric model products (e.g., bias in precipitation and air temperature) also
2236 can have significant effects on subsequent streamflow forecasts. Bias corrections require knowledge of the
2237 climatologies (i.e., long-term distributions) of both modeled and observed variables.

2238 Although meteorological uncertainty may be high for the periods addressed by streamflow forecasts, accurate
2239 estimates of the state of watershed conditions prior to the forecast period are important because they are used to
2240 initialize hydrologic model states, with significant consequences for forecast results. However, watershed conditions
2241 can be difficult to measure, especially when streamflow forecasts must be made quickly, as in the case of flash flood
2242 forecasts. One option is to continuously update watershed states by running the hydrologic models continuously and by
2243 using inputs from recent meteorological observations and/or atmospheric models. Regardless of the source of inputs,

2244 Westrick et al. (2002) found it essential to obtain observational estimates of initial conditions to keep streamflow
2245 forecasts realistic; storm-by-storm corrections of model biases determined over extended simulation periods were
2246 insufficient. Recent experimental end-to-end forecasts of streamflow produced in a simulated operational setting (Wood
2247 et al., 2001) highlighted the critical role of quality estimates of spring and summer soil moisture used to initialize
2248 hydrologic model states for the eastern U.S.

2249 Where streamflows may be largely comprised of snowmelt runoff, quality estimates of snow conditions are
2250 important. The importance of reducing errors in the timing and magnitude of snowmelt runoff are especially acute in
2251 regions where a large percentage of annual water supplies derive from snowmelt runoff, snowmelt impacts are highly
2252 non-linear with increasing deviation from long-term average supplies, and reservoir storage is smaller than interannual
2253 variation of water supplies. However, resources for on-site monitoring of snow conditions have diminished rather than
2254 grown, relative to the increasing costs of errors in hydrologic forecasts (Davis and Pangburn, 1999). Research activities
2255 of the NWS National Office of Hydrology Remote Sensing Center (NOHRSC) have long been directed at improving
2256 estimates of snowpack conditions through aerial and satellite remote sensing (Carroll, 1985). However, the cost of aerial
2257 flights prohibits routine use (T. Carroll, NOHRSC, personal communication, 1999), while satellite estimates have
2258 qualitative limitations (e.g., not considering fractional snow coverage over large regions) and have not found broad use
2259 operationally.

2260 Multiple techniques exist to more accurately represent the uncertainty inherent in understanding and predicting
2261 potential hydroclimatic variability. Stochastic hydrology techniques use various forms of autoregressive models to
2262 generate multiple synthetic streamflow time series with statistical characteristics matching available observations. For
2263 example, in estimating the risk of low flows for the Sacramento River Basin in California, the BOR (Frevert et al.,
2264 1989) generated 20 one-thousand-year streamflow time series matching selected statistics of observed flows (adjusted to
2265 compensate for water management impacts on natural flows); the non-exceedance probabilities of low flows were
2266 computed by counting the occurrences of low flows within 1- through 10-year intervals for all 20 one-thousand-year
2267 sequences. The U.S. Army Corps of Engineers (1992) used a similar approach to estimate flood magnitudes with return
2268 periods exceeding 1,000 years, using Monte Carlo sampling from within the 95% confidence limits of a Log Pearson III
2269 distribution developed by synthesizing multiple streamflow time series.

2270 The ability to automatically execute many model runs within RiverWare, including accessing data from
2271 external sources and exporting model results, facilitates using stochastic hydrology approaches for representing

2272 uncertainty. For example, Carron et al. (2006) demonstrated RiverWare's capability to identify and quantify significant
2273 sources of uncertainty in projecting river and reservoir conditions, using a first-order, second-moment (FOSM)
2274 algorithm that is computationally more efficient than more traditional Monte Carlo approaches. The FOSM processes
2275 uncertainties in inputs and models to provide estimates of uncertainty in model results that can be used directly within a
2276 risk management decision framework. The case study presented by Carron et al. (2006) evaluated the uncertainties
2277 associated with meeting goals for reservoir water levels beneficial for recovering endangered fish species within the
2278 lower Colorado River.

2279 With regard to RiverWare applications concerned with mid-range planning and use of hydrologic forecasts, at
2280 the core of any forecasting system is the predictive model, whether a simple statistical relationship or a complex
2281 dynamic numerical model. Advances in hydrologic modeling have been notable, especially those associated with the
2282 proper identification of a model's parameters (e.g., Duan et al., 2002) and the development of models that consider the
2283 spatially distributed characteristics of watersheds, rather than treating entire basins as a single point (Grayson and
2284 Bloschl, 2000). Conceptual rainfall-runoff models offer some advantages over statistical techniques in support of long-
2285 range planning for water resources management. These models represent, with varying levels of complexity, the
2286 transformation of precipitation and other meteorological forcing variables (e.g., air temperature and humidity) to
2287 watershed runoff and streamflow, including accounting for hydrologic storage conditions (e.g., snowpack, soil moisture,
2288 and groundwater). These models can be used to assess the impacts and implications of various climate scenarios by
2289 using historic meteorological time series as input, generating hydrologic time series, and then using those hydrologic
2290 scenarios as input to RiverWare. This approach enables consideration of current landscape and river channel conditions,
2291 which may be quite different than recorded in early instrumental records and which can dramatically alter a watershed's
2292 hydrologic behavior (Vorosmarty et al., 2004). Furthermore, the use of multiple input time series, system
2293 parameterizations, or multiple models, enables a probabilistic assessment of an ensemble of scenarios. The Hydrological
2294 Ensemble Prediction Experiment (HEPEX) (Schaake et al., 2007) aims to address the unique challenges of expressing
2295 uncertainty associated with ensemble forecasts for water resources management.

2296 An additional concern for mid- and long-range planning is that, as instrumental records have grown longer,
2297 they often show trends (e.g., Baldwin and Lall, 1999; Olsen et al., 1999; Andreadis and Lettenmaier, 2006) or persistent
2298 regimes (i.e., periods characterized by distinctly different statistics) (e.g., Angel and Huff, 1995; Quinn, 1981, 2002),
2299 with consequences for estimation of hydrologic risk (Olsen et al., 1998). Observed regimes and trends can have multiple

2300 causes, including climatic changes, watershed and river transformations, and management impacts (e.g., irrigation
2301 return flows and trans-basin water diversions). These issues enter into RiverWare applications directly through the use
2302 of naturalized flows, which are notoriously unreliable. For example, in assessments of water management options on the
2303 San Juan River in Colorado and New Mexico, the reliability of naturalized flows was considered to be affected by the
2304 inconsistent accounting of consumptive uses between irrigation and non-irrigation data, use of reservoir evaporation
2305 rates with no year-to-year variation, neglecting time lags in the accounting of return flows from irrigation to the river,
2306 errors in river gage readings that underestimated flows in critical months, and the lack of documentation of diversions
2307 that reduce river flows as well as subsequent adjustments to data used to compute naturalized flows.

2308

2309 **5. Global Change Information and RiverWare**

2310 *Climate Variability*

2311 Decision makers increasingly recognize that climate is an important source of uncertainty and potential
2312 vulnerability in long-term planning for the sustainability of water resources (Hartmann, 2005). With the appropriate
2313 investment in site licenses, training of personnel, implementation for a specific river system, and assessment efforts,
2314 RiverWare is capable of supporting climate-related water resources management decisions by U.S. agencies. However,
2315 technology alone is insufficient to resolve conflicts among competing water uses. Early in the development of
2316 RiverWare, Reitsma et al. (1996) investigated its potential role as a DST within complex negotiations between
2317 hydroelectric, agricultural, and flood control interests. Results indicated that while DSTs can help identify policies that
2318 can satisfy specific management requirements and constraints, as well as expand the range of policy options considered,
2319 they are of limited value in helping decision makers understand interactions within the river system. Furthermore, the
2320 burdens of direct use by decision makers of a DST that embodies a complex system are significant; a more useful
2321 approach is to have specialists support decision makers by making model runs and presenting the results in an iterative
2322 manner. This is the approach used by the Bureau of Reclamation in the application of RiverWare to support interstate
2323 negotiations concerning the sharing of Colorado River water supply shortages during times of drought (Jerla, 2005; U.S.
2324 Department of Interior, 2007).

2325 From the perspective of mid-range water management issues, the use of forecasts within RiverWare
2326 applications constitutes an important pathway for supporting climate-related decision making. Each time a prediction is
2327 made, science has an opportunity to address and communicate the strengths and limitations of current understanding.

2328 Each time a decision is made, managers have an opportunity to confront their understanding of scientific information
2329 and forecast products. Furthermore, each prediction and decision provides opportunities for interaction between
2330 scientists and decision makers and for making clear the importance of investments in scientific research. Perceptions of
2331 poor forecast quality are a significant barrier to more effective use of hydroclimatic forecasts (Changnon, 1990; Pagano
2332 et al., 2001, 2002; Rayner et al., 2005); however, recent advances in modeling and predictive capabilities naturally lead
2333 to speculation that hydroclimatic forecasts can be used to improve the operation of water resource systems.

2334 Great strides have been made in monitoring, understanding, and predicting interannual climate phenomena
2335 such as the El Nino-Southern Oscillation (ENSO). This improved understanding has resulted in long-lead (up to about a
2336 year) climate forecast capabilities that can be exploited in streamflow forecasting. Techniques have been developed to
2337 directly incorporate variable climate states into probabilistic streamflow forecast models based on linear discriminant
2338 analysis (LDA) with various ENSO indicators, (e.g., the Southern Oscillation Index [SOI]) (Peichota and Dracup, 1999;
2339 Piechota et al., 2001). Recent improved understanding of decadal-scale climate variability also has contributed to
2340 improved interannual hydroclimatic forecast capabilities. For example, the Pacific Decadal Oscillation (PDO) (Mantua
2341 et al., 1997) has been shown to modulate ENSO-related climate signals in the West. Experimental streamflow
2342 forecasting systems for the Pacific Northwest have been developed based on long-range forecasts of both PDO and
2343 ENSO (Hamlet and Lettenmaier, 1999). In the U.S., the Pacific Northwest, California, and the Southwest are strong
2344 candidates for the use of long-lead forecasts because ENSO and PDO signals are particularly strong in these regions and
2345 each region's water supplies are closely tied to accumulation of winter snowfall, amplifying the impacts of climatic
2346 variability.

2347 While many current water management decision processes use single-value deterministic approaches,
2348 probabilistic forecasts enable quantitative estimation of the inevitable uncertainties associated with weather and climate
2349 systems. From a decision maker's perspective, probabilistic forecasts are more informative because they explicitly
2350 communicate uncertainty and are more useful because they can be directly incorporated into risk-based calculations.
2351 Probabilistic forecasts of water supplies can be created by overlaying a single prediction with a normal distribution of
2352 estimation error determined at the time of calibration of the forecast equations (Garen, 1992). However, to account for
2353 future meteorological uncertainty, new developments have focused on ensembles, whereby multiple possible futures
2354 (each termed an ensemble trace) are generated; statistical analysis of the ensemble distribution then provides the basis
2355 for a probabilistic forecast.

2356 Changnon (2000), Rayner et al. (2005), and Pagano et al. (2002) found that improved climate prediction
2357 capabilities are initially incorporated into water management decisions informally, using subjective, ad hoc procedures
2358 on the initiative of individual water managers. While improvised, those decisions are not necessarily insignificant. For
2359 example, the Salt River Project, among the largest water management agencies in the Colorado River Basin and primary
2360 supplier to the Phoenix metropolitan area, decided in August 1997 to substitute groundwater withdrawals with reservoir
2361 releases, expecting increased surface runoff during a wet winter related to El Nino. With that decision, they risked
2362 losses exceeding \$4 million in an attempt to realize benefits of \$1 million (Pagano et al., 2002). Because these informal
2363 processes are based in part on confidence in the predictions, overconfidence in forecasts can be even more problematic
2364 than lack of confidence, as a single incorrect forecast that provokes costly shifts in operations can devastate user
2365 confidence in subsequent forecasts (e.g., Glantz, 1982).

2366 The lack of verification of hydroclimatic forecasts is a significant barrier to their application in water
2367 management, but it is not easy to resolve with traditional research efforts, because the level of acceptable skill varies
2368 widely depending on the intended use (Hartmann et al., 2002a; Pagano et al., 2002). Information on forecast
2369 performance has rarely been available to, and framed for, decision makers, although hydrologic forecasts are reviewed
2370 annually by the issuing agencies in the U.S (Hartmann et al., 2002b). Hydrologic forecast verification is an expanding
2371 area of research (Franz et al., 2003; Hartmann et al., 2003; Bradley et al, 2004; Pagano et al., 2004; Kruger et al., 2007),
2372 but much work remains and could benefit from approaches developed within the meteorological community (Welles et
2373 al., 2007). Because uncertainty exists in all phases of the forecast process, forecast systems designed to support risk-
2374 based decision making need to explicitly quantify and communicate uncertainties from the entire forecast system and
2375 from each component source, including model parameterization and initialization, meteorological forecast uncertainty at
2376 the multiple spatial and temporal scales at which they are issued, adjustment of meteorological forecasts (e.g., through
2377 downscaling) to make them usable for hydrologic models, implementation of ensemble techniques, and verification of
2378 hydrologic forecasts.

2379

2380 *Climate Change*

2381 From the perspective of long-range water management issues, the potential impacts of climate change on water
2382 resources, and their implications for management, are central topics of concern. Estimates of prospective impacts of
2383 climate change on precipitation have been mixed, leading, in many cases, to increasing uncertainty about the reliability

2384 of future water supplies. However, where snow provides a large fraction of annual water supplies, prospective
2385 temperature increases dominate hydrologic impacts, leading to stresses on water resources and increased hydrologic
2386 risk. Higher temperatures effectively shift the timing of the release of water stored in the snowpack “reservoir” to earlier
2387 in the year, reducing supplies in summer when demands are greatest, while also increasing the risk of floods due to rain-
2388 on-snow events. While not using RiverWare, several river basin studies have assessed the risks of higher temperatures
2389 on water supplies and management challenges. The near universal analytical approach has been one of sensitivity
2390 analysis (Lettenmaier, 2003):

- 2391 1) downscaling outputs from a dynamic general circulation model of the global land-atmosphere-ocean system to
2392 generate regional- or local-scale meteorological time series over many decades,
- 2393 2) using the meteorological time series as input to rainfall-runoff models to generate hydrologic time series,
- 2394 3) using the hydrologic scenarios as input to water management models, and
- 2395 4) assessing differences among baseline and change scenarios using a variety of metrics.

2396 Early assessments of warming impacts on large river basins generally showed extant water management systems to
2397 be effective for all but the most severe scenarios (Hamlet and Lettenmaier, 1999; Lettenmaier et al., 1999), with a
2398 notable exception being the Great Lakes system where increased lake heat storage was tied to loss of ice cover,
2399 increased winter lake evaporation, lower lake levels, and potential failure to meet Lake Ontario regulation objectives
2400 under extant operating rules (Croley, 1990; Hartmann, 1990; Lee et al., 1994; Lee et al., 1997; Sousounis et al., 2000;
2401 Lofgren et al., 2002).

2402 Extensive detailed studies of the ability of existing reservoir systems and operational regulation rules to meet water
2403 management goals under changed climates are fairly recent (e.g., Saunders and Lewis, 2003; Christensen, et. al, 2004;
2404 Payne et. al, 2004; VanRheenan et. al, 2004; Maurer, 2007). However, there is a rapidly growing literature on broad
2405 considerations of climate change in water resources management (Frederick et al., 1997; Gamble et al., 2003;
2406 Lettenmaier, 2003; Loomis et al., 2003; Snover et al., 2003; Stakhiv, 2003; Ward et al., 2003; Vicuna et al., 2007).
2407 Some (Matalas, 1997) that contend that existing approaches are sufficient for water resource management planning and
2408 risk assessment because they contain safety factors; however, an inescapable message for the water resource
2409 management community is the inappropriateness of the stationarity assumption in the face of climate change. While
2410 precipitation changes may remain too uncertain for consideration in the near term, temperature increases are more
2411 certain and can have strong hydrologic consequences.

2412 Cognitively, climate change information is difficult to integrate into water resources management. First, within
2413 the water resources engineering community, the stationarity assumption is a fundamental element of professional
2414 training. Second, the century timescales of climate change exceed typical planning and infrastructure design horizons
2415 and are remote from human experience. Third, even individuals trying to stay up-to-date can face confusion in
2416 conceptually melding the burgeoning climate change impacts literature. Assessments are often repeated as general
2417 circulation and hydrologic model formulations advance or as new models become available throughout the research
2418 community. Furthermore, assessments can employ a variety of techniques for downscaling. Transposition techniques
2419 (e.g., Croley et al., 1998) are more intuitive than the often mathematically complex statistical and dynamical
2420 downscaling techniques (e.g., Clark et al., 1999; Westrick and Mass, 2001; Wood et al., 2002; Benestad, 2004).

2421 GCMs and their downscaled corollaries provide one unique perspective on long-term trends related to global
2422 change. Another unique perspective is provided by tree-ring reconstructions of paleo-streamflows, which, for example,
2423 indicate that in the U.S. Southwest droughts over the past several hundred years have been more intense, regionally
2424 extensive, and persistent than those reflected in the instrumental record (Woodhouse and Lukas, 2006). Decision makers
2425 have expressed interest in combining the perspectives of paleoclimatological information and GCMs. While some
2426 studies have linked instrumental records to paleoclimatological information (e.g., Prairie, 2006) and others with GCMs
2427 (e.g., Christensen and Lettenmaier, 2006), few link all three (an exception is Smith et al., 2007).

2428 Conceptual integration of climate change impacts assessment results in a practical water management context
2429 is complicated by the multiplicity of scenarios and vague attribution of their prospects for occurrence, which depend so
2430 strongly on feedbacks among social, economic, political, technological, and physical processes. For decision makers, a
2431 critical issue concerns the extent to which the various scenarios reflect the actual uncertainty of the relevant risks versus
2432 the uncertainty due to methodological approaches and biases in underlying models. The difficulties facing decision
2433 makers in reconciling disparate climate change impact assessments are exemplified by the Upper Colorado River Basin,
2434 where reductions in naturalized flow by the mid-21st century have been estimated to range from about 45% by Hoerling
2435 and Eischeid (2007), 10 to 25% by Milly et al (2005), about 18% by Christensen et al. (2004), and about 6% by
2436 Christensen and Lettenmaier (2006). Furthermore, using the difference between precipitation and evapotranspiration as
2437 a proxy for runoff, Seager et al. (2007) suggest an “imminent transition to a more arid climate in southwestern North
2438 America.”

2439 However, in the face of circumstances nearing or exceeding the effectiveness of existing management
2440 paradigms, individuals can become more cognizant of the need to consider climate change. In the U.S. Southwest, over
2441 1999–2004, Lake Powell levels declined faster than previously considered in scenarios of extreme sustained drought
2442 (e.g., Harding et al., 1995; Tarboton, 1995), from full to only 38% capacity in November 2004 (BOR, 2004). Resource
2443 managers, policymakers, and the general public are now actively seeking scientific guidance in exploring how
2444 management practices can be more responsive to the uncertainties associated with a changing climate.

2445




Figure 1: Illustration depicting the flow of information

Appendix A

References by Chapter

Chapter 1 References – Decision Support for Agricultural Efficiency:

Birkett, Charon and Brad Doorn. 2004. "A New Remote Sensing Tool for Water Resources Management, *Earth Observation Magazine*, October 13 (6).

Congressional Research Service, Science Policy Research Division. 1983. "United States Civilian Space Programs, Volume II Applications Satellites," prepared for the Subcommittee on Space Science and Applications of the Committee on Science and Technology, U.S. House of Representatives, May.

Kanarek, Harold. 2005. "The FAS Crop Explorer: A Web Success Story," *FAS Worldwide*, June (<http://www.fas.usda.gov/info/fasworldwide/2005/06-2005/Cropexplorer.htm> (accessed April 2007)).

Kaupp, Verne, Charles Hutchinson, Sam Drake, Tim Haithcoat, Willem van Leeuwen, Vlad Likholetov, David Tralli, Rodney McKellip, and Brad Doorn. 2005. "Benchmarking the USDA Production Estimates and Crop Assessment Division DSS Assimilation," v.3 (01.04.06), report prepared for Production Estimates and Crop Assessment Division, Foreign Agricultural Service, US Department of Agriculture, September.

National Aeronautics and Space Administration, 2001. Aeronautics and Space Report of the President, NASA, Washington DC at <http://history.nasa.gov/presrep01/pages/usda.html> accessed April 2007.

- 2473 National Aeronautics and Space Administration, John C. Stennis Space Center. 2004a. "Decision Support Tools
2474 Evaluation Report for FAS/PECAD," Version 2.0, January.
2475
- 2476 National Aeronautics and Space Administration, John C. Stennis Space Center. 2004b. "PECAD's Global Reservoir and
2477 Lake Monitor: A Systems Engineering Report," Version 1.0, December.
2478
- 2479 National Aeronautics and Space Administration. 2006a. "NASA Science Mission Directorate: Earth-Sun System
2480 Applied Sciences Program Agricultural Efficiency Program Element FY2006-2010 Plan," 30 June at
2481 http://aiwg.gsfc.nasa.gov/esappdocs/Agricultural_Efficiency_FINAL_06.pdf accessed April 2007.
2482
- 2483 National Aeronautics and Space Administration, 2006b. "NASA Science Mission Directorate – Applied Sciences
2484 Program: Agricultural Efficiency – FY 2005 Annual Report at
2485 <http://aiwg.gsfc.nasa.gov/esappdocs/annualreports/> accessed April 2007.
2486
- 2487 National Assessment Synthesis Team. 2004. *Climate Change Impacts on the United States: The Potential
2488 Consequences of Climate Variability* (Boston, MA: Cambridge University Press).
2489
- 2490 National Research Council, Board on Earth Sciences and Resources. 2007. *Contributions of Land Remote Sensing for
2491 Decisions about Food Security and Human Health: Workshop Report* (Washington, DC: National Academies
2492 Press).
2493
- 2494 Reynolds, Curt A. 2001. "CADRE Soil Moisture and Crop Models," at
2495 <http://www.pecad.fas.usda.gov/cropexplorer/datasources.cfm> accessed April 2007.
2496
- 2497 Rosenzweig, Cynthia. 2003. "Climate Change and Agriculture: Mitigation and Adaptation," Testimony before the
2498 Senate Committee on Environment and Public Works, Subcommittee on Clean Air, Climate Change, and
2499 Nuclear Safety, July 8 at http://epw.senate.gov/108th/Rosenzweig_070803.htm accessed April 2007.
2500

- 2501 United Nations Food and Agriculture Organization. 2006. "Agricultural Monitoring Meeting Convened for the
2502 Integrated Global Observations for Land (IGOL) Theme," Rome, Italy (8-11 March 2006), 28 June.
2503
- 2504 United Nations Food and Agriculture Organization. No date. "Agriculture and Climate Change: FAO's Role" at
2505 <http://www.fao.org/News/1997/971201-e.htm> Accessed April 2007.
2506

2507

2508

Chapter 2 References – Decision Support for Air Quality:

2509

2510 Al-Saadi, J., J. Szykman, R. B. Pierce, C. Kittaka, D. Neil, D. A. Chu, L. Remer, L. Gumley, E. Prins, L. Weinstock, C.

2511 MacDonal, R. Wayland, F. Dimmick, and J. Fishman, 2005: Improving national air quality forecast with

2512 satellite aerosol observations. *Bulletin of the American Meteorological Society*, Volume 86, Issue 9, 1249-

2513 1261.

2514

2515 Bey, I., D. J. Jacob, R. M. Yantosca, J. A. Logan, B. D. Field, A. M. Fiore, Q. Li, H. Y. Liu, L. J. Mickley, and M. G.

2516 Schultz (2001), Global modeling of tropospheric chemistry with assimilated meteorology: Model description

2517 and evaluation, *J. Geophys. Res.*, 106, 23,073– 23,095.

2518

2519 Brasseur, G. P., J. T. Kiehl, J.-F. Mueller, T. Schneider, C. Granier, X. X. Tie, and D. Hauglustaine, 1998: Past and

2520 future changes in global tropospheric ozone: Impact on radiative forcing, *Geophys. Res. Lett.*, 25, 3807– 3810.

2521

2522 Brown, T.J., B.L. Hall, and A.L. Westerling, 2004: The impact of twenty-first century climate change on wildland fire

2523 danger in the western United States: An applications perspective. *Climatic Change*, 62, 365–388.

2524

2525 Byun, D.W., 1999a: Dynamically consistent formulations in meteorological and air quality models for multi-scale

2526 atmospheric applications: Part I. Governing equations in generalized coordinate system. *Journal of*

2527 *Atmospheric Science*, Vol 56, 3789-3807.

2528

2529 Byun, D.W., 1999b: Dynamically consistent formulations in meteorological and air quality models for multi-scale

2530 atmospheric applications: Part II. Mass conservation issues. *Journal of Atmospheric Science*, Vol 56, 3808-

2531 3820.

2532

- 2533 Byun, D.W. and Ching, J.K.S. (eds.), 1999: Science algorithms of the EPA Models-3 Community Multiscale Air
2534 Quality Model (CMAQ) modeling system. EPA/600/R-99/030, U. S. Environmental Protection Agency, Office
2535 of Research and Development, Washington, DC 20460.
- 2536
- 2537 Byun, D.W., and K. L. Schere, 2006: Review of the Governing Equations, Computational Algorithms, and Other
2538 Components of the Models-3 Community Multiscale Air Quality (CMAQ) Modeling System . Applied
2539 Mechanics Reviews, Volume 59, Number 2 (March 2006), pp. 51-77.
- 2540
- 2541 Carmichael, G. R., L. K. Peters, and R. D. Saylor, 1991: The STEM-II regional scale acid deposition and photochemical
2542 oxidant model—I. An overview of model development and applications, Atmos. Environ., 25(10), 2077– 2090
- 2543
- 2544 Civerolo, K., C. Hogrefe, B. Lynn, J. Rosenthal, J.-Y. Ku, W. Solecki, J. Cox, C. Small, C. Rosenzweig, R. Goldberg,
2545 K. Knowlton, and P. Kinney, 2007: Estimating the effects of increased urbanization on surface meteorology
2546 and ozone concentrations in the New York City metropolitan region. Atmos. Environ., 41, 1803-1818,
2547 doi:10.1016/j.atmosenv.2006.10.076.
- 2548
- 2549 Constantinescu, E.M., A. Sandu, T. Chai, and G.R. Carmichael, 2007a: Ensemble-based Chemical Data Assimilation I:
2550 General Approach. Quarterly Journal of the Royal Meteorological Society, in print.
- 2551
- 2552 Constantinescu, E.M. , A. Sandu, T. Chai, and G.R. Carmichael, 2007b: Ensemble-based Chemical Data Assimilation
2553 II: Covariance Localization. Quarterly Journal of the Royal Meteorological Society, in print.
- 2554
- 2555 Delworth, T.L., A. J. Broccoli, A. Rosati, R. J. Stouffer, V. Balaji, J. A. Beesley, W. F. Cooke, and 37 co-authors, 2006:
2556 GFDL’s CM2 Global Coupled Climate Models. Part I: Formulation and Simulation Characteristics, Journal of
2557 Climate-Special Section, Vol. 19, 643-674
- 2558

- 2559 Duncan, B.N., R.V. Martin, A.C. Staudt, R. Yevich, J.A. Logan, 2003: Interannual and Seasonal Variability of Biomass
2560 Burning Emissions Constrained by Satellite Observations, *J. Geophys. Res.*, 108(D2), 4040,
2561 doi:10.1029/2002JD002378.
- 2562
- 2563 Eder, B., D. Kang, R. Mathur, S. Yu, K. Schere, 2006: An operational evaluation of the Eta-CMAQ air quality forecast
2564 model, *Atmospheric Environment* 40, 4894-4905
- 2565
- 2566 Emmons, L. K., D. A. Hauglustaine, J.-F. Muller, M. A. Carroll, G. P. Brasseur, D. Brunner, J. Staehelin, V. Thouret,
2567 and A. Marengo, 2000: Data composites of airborne observations of tropospheric ozone and its precursors, *J.*
2568 *Geophys. Res.*, 105, 20,497– 20,538.
- 2569
- 2570 Friedl, R. (ed.), 1997: Atmospheric effects of subsonic aircraft: Interim assessment report of the advanced subsonic
2571 technology program, NASA Ref. Publ. 1400, 143 pp., 1997.
- 2572
- 2573 Fu, T.-M., D. J. Jacob, P. I. Palmer, K. Chance, Y. X. Wang, B. Barletta, D. R. Blake, J. C. Stanton, and M. J. Pilling,
2574 2007: Space-based formaldehyde measurements as constraints on volatile organic compound emissions in east
2575 and south Asia and implications for ozone, *J. Geophys. Res.*, 112, D06312, doi:10.1029/2006JD007853.
- 2576
- 2577 Hakami, A., D.K. Henze, J.H. Seinfeld, K. Singh, A. Sandu, S. Kim, D. Byun, and Q. Li, 2007: The adjoint of CMAQ,
2578 (submitted to *J. Geophys. Res.*)
- 2579
- 2580 Hansen, J., Mki. Sato, L. Nazarenko, R. Ruedy, A. Lacis, D. Koch, I. Tegen, T. Hall, and 20 co-authors, 2002: Climate
2581 forcings in Goddard Institute for Space Studies SI2000 simulations. *J. Geophys. Res.* 107, no. D18, 4347,
2582 doi:10.1029/2001JD001143.
- 2583
- 2584 Hansen, J., Mki. Sato, R. Ruedy, L. Nazarenko, A. Lacis, G.A. Schmidt, G. Russell, and 38 co-authors, 2005: Efficacy
2585 of climate forcings. *J. Geophys. Res.* 110, D18104, doi:10.1029/2005JD005776.
- 2586

- 2587 Heald, Colette L., Daniel J. Jacob, Paul I. Palmer, Mathew J. Evans, Glen W. Sachse, Hanwant B. Singh and Donald R.
2588 Blake, 2003: Biomass burning emission inventory with daily resolution: application to aircraft observations of
2589 Asian outflow, *J. Geophys. Res.*, 108(D4), 8368, doi:10.1029/2002JD002732.
2590
- 2591 Hoelzemann, J.J., M. G. Schultz, G. P. Brasseur, and C. Granier, 2004: Global Wildland Fire Emission Model
2592 (GWEM): Evaluating the use of global area burnt satellite data, *J. Geophys. Res.*, 109, D14S04,
2593 doi:10.1029/2003JD003666.
2594
- 2595 Hogrefe, C., B. Lynn, K. Civerolo, J.-Y. Ku, J. Rosenthal, C. Rosenzweig, R. Goldberg, S. Gaffin, K. Knowlton, and P.
2596 L. Kinney, 2004: Simulating changes in regional air pollution over the eastern United States due to changes in
2597 global and regional climate and emissions, *J. Geophys. Res.*, 109, D22301, doi:10.1029/2004JD004690.
2598
- 2599 Hogrefe C, LR Leung, LJ Mickley, SW Hunt, and DA Winner, 2005: Considering Climate Change in U.S. Air Quality
2600 Management. *EM: Air & Waste Management Association's magazine for environmental managers* October
2601 2005:19-23.
2602
- 2603 Holloway, T., H. Levy II, and G. Carmichael, 2002: Transfer of reactive nitrogen in Asia: development and evaluation
2604 of a source–receptor model. *Atmospheric Environment*, 36(26), 4251-4264.
2605
- 2606 Horowitz, L. W., and Coauthors, 2003: A global simulation of tropospheric ozone and related tracers: Description and
2607 evaluation of MOZART, version 2. *J. Geophys. Res.*, 108, 4784, doi:10.1029/2002JD002853.
2608
- 2609 Hurrell, J.W., J.J. Hack, A.S. Phillips, J. Caron, and J. Yin, 2006: The Dynamical Simulation of the Community
2610 Atmosphere Model Version 3 (CAM3) *Journal of Climate*: Vol. 19, pp 2162-2183.
2611
- 2612 In, H.-J., D. W. Byun, R. J. Park, N.-K. Moon, S. Kim, and S. Zhong, 2007: Impact of transboundary transport of
2613 carbonaceous aerosols on the regional air quality in the United States: A case study of the South American
2614 wildland fire of May 1998, *J. Geophys. Res.*, 112, D07201, doi:10.1029/2006JD007544.

2615
2616 IPCC (Intergovernmental Panel on Climate Change), 2000: Emissions Scenarios, Cambridge University Press,
2617 Cambridge, UK
2618
2619 IPCC (Intergovernmental Panel on Climate Change), 2001: The Scientific Basis. Cambridge University Press,
2620 Cambridge, UK
2621
2622 Jacob, D.J., and A.B. Gilliland, 2005: Modeling the impact of air pollution on global climate change, Environmental
2623 Manager, pp. 24-27, October 2005, Air & Waste Management Association. Pittsburgh, PA.
2624
2625 Jacobson, M. Z., GATOR-GCMM, 2001a: A global through urban scale air pollution and weather forecast model. 1.
2626 Model design and treatment of subgrid soil, vegetation, roads, rooftops, water, sea ice, and snow. J. Geophys.
2627 Res., 106, 5385-5402.
2628
2629 Jacobson, M. Z., 2001b: GATOR-GCMM: 2. A study of day- and nighttime ozone layers aloft, ozone in national parks,
2630 and weather during the SARMAP Field Campaign, J. Geophys. Res., 106, 5403-5420, 2001
2631
2632 Kalkstein, L. S., and K. M. Valimont, 1987: Climate effects on human health. EPA Science and Advisory Committee
2633 Monograph, no. 25389: 122-152. Washington D. C., U. S. EPA.
2634
2635 Kiehl, J.T., J. Hack, G. Bonan, B. Boville, B. Briegleb, D. Williamson, and P. Rasch, 1996: Description of the NCAR
2636 Community Climate Model (CCM3). NCAR Technical Note. NCAR/TN-420+STR, Ntl. Center for Atmos.
2637 Research, Boulder, CO, 152 pp. [Available Ntl. Cen. Atmos. Res., P.O. Box 3000, Boulder, CO, 80305.]
2638
2639 Knowlton, K., Rosenthal, J.E., Hogrefe, C., Lynn, B., Gaffin, S., Goldberg, R., Rosenzweig, C., Civerolo, K., Ku, J.-Y.,
2640 Kinney, P.L., 2004. Assessing ozone-related health impacts under a changing
2641

- 2642 Kopacz, M., D. J. Jacob, D. Henze, C. L. Heald, D. G. Streets, Q. Zhang, 2007: Comparison of adjoint and analytical
2643 Bayesian inversion methods for constraining Asian sources of carbon monoxide using satellite (MOPITT)
2644 measurements of CO columns, Submitted to Journal of Geophysical Research – Atmospheres.
2645
- 2646 Leung, L. R., and M. S. Wigmosta, 1999: Potential climate change impacts on mountain watersheds in the Pacific
2647 Northwest. *J. Amer. Water Resour. Assoc.*, 35(6): 1463-1471.
2648
- 2649 Leung, L. R., S. J. Ghan, Z.-C. Zhao, Y. Luo, W.-C. Wang, and H. Wei, 1999: Intercomparison of regional climate
2650 simulations of the 1991 summer monsoon in East Asia. *J. Geophys. Res.*, 104(D6): 6425-6454.
2651
- 2652 Leung LR, Y Kuo, and J Tribbia. 2006: Research Needs and Directions of Regional Climate Modeling Using WRF and
2653 CCSM." *Bulletin of the American Meteorological Society* 87(12):1747-1751.
2654
- 2655 Liang, X.-Z., J. Pan, J. Zhu, K.E. Kunkel, J.X.L. Wang, and A. Dai, 2006: Regional climate model downscaling of the
2656 U.S. summer climate and future change. *J. Geophys. Res.*, 111, D10108.
2657
- 2658 Liao, K.-J., E. Tagaris, K. Manomaiphiboon, J.-H. Woo, S. He, P. Amar, and A.G. Russell, 2007: Sensitivities of Ozone
2659 and Fine Particulate Matter Formation to Emissions under the Impact of Potential Future Climate Change,
2660 (submitted to Journal of Geophysical Research)
2661
- 2662 Liao, K.-J., E. Tagaris, K. Manomaiphiboon, A.G. Russell, C. Wang, J.-H. Woo, P. Amar, and S. He, 2007: Quantifying
2663 the Uncertainties in Forecasts of Regional Air Quality under Impact of Future Climate Change, (submitted to
2664 Journal of Geophysical Research)
2665
- 2666 Logan, J.A., 1999: An analysis of ozonesonde data for the troposphere: Recommendations 601 for testing 3-D models
2667 and development of a gridded climatology for tropospheric ozone, *J. Geophys. Res.*, 104, D13, 16,115-
2668 16,149.
2669

- 2670 Mearns, L.O., 2003: Issues in the impacts Climate variability and change on agriculture—Applications to the
2671 southeastern United States. *Climate Change*, 60, 1–6.
2672
- 2673 Mickley, L.J., D.J. Jacob, B.D. Field, and D. Rind, 2004: Effects of future climate change on regional air pollution
2674 episodes in the United States, *Geophys. Res. Lett.*, 30, L24103, doi:10.1029/2004GL021216.
2675
- 2676 NARSTO, 2000: An assessment of tropospheric ozone pollution—a North American Perspective. NARSTP Management
2677 Coordinator's Office (Envair). Pasco, Washington, <http://www.narsto.org/>
2678
- 2679 Novelli, P. C., K. A. Masarie, P. M. Lang, B. D. Hall, R. C. Myers, and J.W. Elkins, 2003: Reanalysis of tropospheric
2680 CO trends: Effects of the 1997– 1998 wildfires, *J. Geophys. Res.*, 108(D15), 4464, doi:10.1029/
2681 2002JD003031.
2682
- 2683 Pour-Biazar, A., R.T. McNider, S.J. Roselle, R. Suggs, G. Jedlovex, D.W. Byun, S.T. Kim, C.J. Lin, T.C. Ho, S.
2684 Haines, B. Dornblaser, and R. Cameron, 2007: Correcting photolysis rates on the basis of satellite observed
2685 clouds, *J. Geophys. Res.*, 112, D10302, doi:10.1029/2006JD007422.
2686
2687
- 2688 Russell, A., and R. Dennis, 2000: NARSTO critical review of photochemical models and modeling. *Atmos. Environ.*,
2689 34, 2283-2324.
2690
- 2691 Sandu, A., D. Daescu, G.R. Carmichael, and T. Chai, 2005: Adjoint sensitivity analysis of regional air quality models.
2692 *Journal of Computational Physics*, 204:222–252.
2693
- 2694 Schmidt, G.A., R. Ruedy, J.E. Hansen, I. Aleinov, N. Bell, M. Bauer, S. Bauer, B. Cairns, V. Canuto, Y. Cheng, A. Del
2695 Genio, G. Faluvegi, A.D. Friend, T.M. Hall, Y. Hu, M. Kelley, N.Y. Kiang, D. Koch, A.A. Lacis, J. Lerner,
2696 K.K. Lo, R.L. Miller, L. Nazarenko, V. Oinas, J. Perlwitz, D. Rind, A. Romanou, G.L. Russell, M. Sato, D.T.
2697 Shindell, P.H. Stone, S. Sun, N. Tausnev, D. Thresher, and M.S. Yao, 2006: Present day Atmos. simulations

- 2698 using GISS Model E: Comparison to in-situ, satellite and reanalysis data. *J. Clim.*, 19, 153–192,
2699 doi:10.1175/JCLI3612.1.
- 2700
- 2701 Song, C.-K., D.W. Byun, R.B. Pierce, J.A. Alsaadi, T.K. Schaack, and F. Vukovich, 2007: Downscale linkage of global
2702 model output for regional chemical transport modeling: method and general performance. (Submitted to
2703 *Journal of Geophysical Research*.)
- 2704
- 2705 Spak, S.N., T. Holloway, B. Lynn, R. Goldberg, 2007: A Comparison of Statistical and Dynamical Downscaling for
2706 Surface Temperature in North America, *J. Geophys. Res.*, 112, D08101, doi:10.1029/2005JD006712.
- 2707
- 2708 Spracklen, D.V., J. A. Logan, L. J. Mickley, R. J. Park, R. Yevich, A.L. Westerling, and D. Jaffe, 2007: Wildfires drive
2709 interannual variability of organic carbon aerosol in the western U.S. in summer: implications for trends.
2710 Submitted to *Journal of Geophysical Research*.
- 2711
- 2712 Tang, Y., G. R. Carmichael, N. Thongboonchoo, T. Chai, L. W. Horowitz, R. B. Pierce, J. A., Al-Saadi, G., Pfister, J.
2713 N. Vukovich, M. A. Avery, G. W. Sachse, T. B. Ryerson, J. S. Holloway, E. L. Atlas, F. M. Flocke, R. J.
2714 Weber, L. G. Huey, J. E. Dibb, D. G. Streets, W. H. Brune, 2007: Influence of lateral and top boundary
2715 conditions on regional air quality prediction: A multiscale study coupling regional and global chemical
2716 transport models, *J. Geophys. Res.*, 112, D10S18, doi:10.1029/2006JD007515.
- 2717
- 2718 Tagaris, E., K. Manomaiphiboon, K.-J. Liao, L. R. Leung, J.-H. Woo, S. He, P. Amar, A. G. Russell, 2007: Impacts of
2719 Global Climate Change and Emissions on Regional Ozone and Fine Particulate Matter Concentrations over the
2720 United States, submitted to *Journal of Geophysical Research*.
- 2721
- 2722 Tarasick, D. W. M. D. Moran, A. M. Thompson, T. Carey-Smith, Y. Rochon, V. S. Bouchet, W. Gong, P. A. Makar, C.
2723 Stroud, S. Ménard, L.-P. Crevier, S. Cousineau, J. A. Pudykiewicz, A. Kallaur, R. Moffet, R. Ménard, A.
2724 Robichaud, O. R. Cooper, S. J. Oltmans, J. C. Witte, G. Forbes, B. J. Johnson, J. Merrill, J. L. Moody, G.
2725 Morris, M. J. Newchurch, F. J. Schmidlin, E. Joseph, 2007: Comparison of Canadian air quality forecast

2726 models with tropospheric ozone profile measurements above midlatitude North America during the
2727 IONS/ICARTT campaign: Evidence for stratospheric input, *J. Geophys. Res.*, doi:10.1029/2006JD007782, in
2728 press.

2729

2730 Tong, D.Q. and D.L. Mauzerall, 2006: Spatial Variability of Summertime Tropospheric Ozone over the Continental
2731 United States: Implications of an evaluation of the CMAQ model , *Atmospheric Environment*, 40, 3041-3056.
2732

2733 Yu, S.C., R. Mathur, D. Kang, K. Schere, J. Pleim, and T.L. Otte, 2007: A detailed evaluation of the Eta-CMAQ
2734 forecast model performance for O₃, its related precursors, and meteorological parameters during the 2004
2735 ICARTT study, *J. Geophys. Res.*, 112, D12S14, doi:10.1029/2006JD007715.
2736

2737 Zhang, F., N. Bei, J. W. Nielsen-Gammon, G. Li, R. Zhang, A. Stuart, and A. Aksoy, 2007: Impacts of meteorological
2738 uncertainties on ozone pollution predictability estimated through meteorological and photochemical ensemble
2739 forecasts, *J. Geophys. Res.*, 112, D04304, doi:10.1029/2006JD007429.
2740

2741 Zhang, Y., Liu, P., Pun, B., Seigneur, C., 2006: A comprehensive performance evaluation of MM5-CMAQ for the
2742 summer 1999 southern oxidants study episode, Part I. Evaluation protocols, databases and meteorological
2743 predictions. *Atmospheric Environment*, this issue, doi:10.1016/j.atmosenv. 2005.12.043.
2744
2745

2746 **Chapter 3 References – Decision Support for Assessing Hybrid Renewable Energy Systems:**

2747

2748

2749 Elliott, D. L., C. G. Holladay, W. R. Barchet, H. P. Foote, and W. R. Sandusky, 1987. Wind Energy Resource Atlas of
2750 the United States. Pacific Northwest National Laboratory, Richland, Washington. DOE/CH 10093-4, March,
2751 1987.

2752

2753 Gueymard, C., SWERA Position Paper, 2003: Methodological Issues Related to Aerosol Data. Personal
2754 communication to the National Renewable Energy Laboratory.

2755

2756 Hansen, M.C., R. S. DeFries, J. R. G. Townshend, M. Carroll, C. Dimiceli, and R. A. Sohlberg, 2003. Global Percent
2757 Tree Cover at a Spatial Resolution of 500 Meters: First Results of the MODIS Vegetation Continuous Fields
2758 Algorithm. *Earth Interactions* 7(10):1-15.

2759

2760 Jennings, Michael and J. Michael Scott, 1997: Official Description of the GAP Analysis Program.

2761 [http://gapanalysis.nbi.gov/portal/server.pt/gateway/PTARGS_0_2_1021_200_458_43/http%3B/gapcontent1%](http://gapanalysis.nbi.gov/portal/server.pt/gateway/PTARGS_0_2_1021_200_458_43/http%3B/gapcontent1%3B7087/publishedcontent/publish/public_sections/gap_home_sections/descriptionofficial/highlights_content.html)
2762 [3B7087/publishedcontent/publish/public_sections/gap_home_sections/descriptionofficial/highlights_content.ht](http://gapanalysis.nbi.gov/portal/server.pt/gateway/PTARGS_0_2_1021_200_458_43/http%3B/gapcontent1%3B7087/publishedcontent/publish/public_sections/gap_home_sections/descriptionofficial/highlights_content.html)
2763 [ml.](http://gapanalysis.nbi.gov/portal/server.pt/gateway/PTARGS_0_2_1021_200_458_43/http%3B/gapcontent1%3B7087/publishedcontent/publish/public_sections/gap_home_sections/descriptionofficial/highlights_content.html)

2764

2765 Koepke, P., M. Hess, I. Schult, and E. P. Shettle, 1997. Global Aerosol Data Set. Report No. 243, Max-Planck-Institut
2766 fur Meteorologie, Hamburg, ISSN 0937-1060.

2767

2768 Perez, R., P. Ineichen, K. Moore, M. Kmiecik, C. Chain, R. George, and F. Vignola, 2002: A New Operational
2769 Satellite-to-Irradiance Model. *Solar Energy* 73(5), pp. 307-317.

2770

2771 Renné, David S., Richard Perez, Antoine Zelenka, Charles Whitlock, and Roberta DiPasquale, 1999: Use of Weather
2772 and Climate Research Satellites for Estimating Solar Resources. Chapter 5 in *Advances in Solar Energy*,

- 2773 Volume 13, Edited by D. Yogi Goswami and Karl W. Boer. The American Solar Energy Society, 2400
2774 Central Ave. Suite G1, Boulder, Colorado 80301. Pp. 171-240.
2775
- 2776 Schwartz, M., R. George, and D. Elliott, 1999. The Use of Reanalysis Data for Wind Resource Assessment at the
2777 National Renewable Energy Laboratory. Proceedings, European Wind Energy Conference, Nice, France,
2778 March 1-5, 1999.

Chapter 4 References – Decision Support for Public Health:

- 2779
2780
- 2781 Beck, L.R. M.H Rodriguez, S.W. Dister, A.D. Rodríguez, R.K. Washino, D.R. Roberts and M.A. Spanner 1997:
2782 Assessment of a remote sensing-based model for predicting malaria transmission risk in villages of Chiapas,
2783 Mexico. *American Journal of Tropical Medicine and Hygiene* 56: 99-107.
2784
- 2785 Brownstein, J.S., T.R. Holford and D. Fish. 2003: A climate-based model predicts the spatial distribution of Lyme
2786 disease vector *Ixodes scapularis* in the United States. *Environmental Health Perspectives* 111: 1152- 1157.
2787
- 2788 Brownstein, J.S., T.R. Holford and D. Fish 2005a: Effect of climate change on Lyme disease risk in North America.
2789 *EcoHealth* 2:38-46.
2790
- 2791 Brownstein, J.S., D. K Skelly, T.R. Holford and D. Fish. 2005b: Forest fragmentation predicts local scale heterogeneity
2792 of Lyme disease risk. *Oecologia* 146: 469-475
2793
- 2794 Fox,D. 2007: Back to the no-analog future? *Science* 316:823-825
2795
- 2796 Glass, G.E. 2007: Rainy with a chance of plague: forecasting disease outbreaks from satellites. *Future Virology* 2:225-
2797 229
2798
- 2799 Gubler, D.J. 2004: The changing epidemiology of yellow fever and dengue 1900 to 2003: full circle? *Comparative*
2800 *Immunology Microbiology and Infectious Diseases* 27:319-330.
2801
- 2802 Gubler, D.J., P. Reiter, K.L. Ebi, W. Yap, R. Nasci and J.A. Patz 2001: Climate variability and change in the United
2803 States: potential impacts on vector- and rodent-borne diseases. *Environmental Health Perspectives* 109:223.
2804
- 2805 Huntingford, C., D. Hemming, J.H.C. Gash, N Gedney, and P.A. Nuttall (2007) Impact of climate change on health:
2806 what is required of climate modellers? *Trans. Royal Soc. Trop. Med. Hyg.*, 101, 97-103.

- 2807
- 2808 Linthicum, K.J., C.L. Bailey, F.G. Davies, and C.J. Tucker 1987: Detection of Rift Valley fever viral activity in Kenya
2809 by satellite remote sensing imagery. *Science* 235:1656-1659.
- 2810
- 2811 Malouin, R, P Winch, E Leontsini, G Glass, D Simon, EB Hayes & BS Schwartz. 2003. Longitudinal evaluation of an
2812 educational intervention to prevent tick bites in an area of endemic Lyme disease in Baltimore County,
2813 Maryland. *Am J Epidemiol* 157:1039-1051.
- 2814
- 2815 Piesman, J. and L. Gern 2004: Lyme borreliosis in Europe and North America. *Parasitology* 129:191-220.
- 2816
- 2817 Selvin, S. 1991: *Statistical Analysis of Epidemiologic Data*. Oxford University Press. New York 375

Chapter 5 References – Water Management

- 2818
- 2819
- 2820 Andreadis, K. and D. Lettenmaier, 2006: Trends in 20th century drought over the continental United States.
- 2821 *Geophysical Research Letters* 33, L10403.
- 2822
- 2823 Angel, J.R. and F.A. Huff, 1995: Seasonal distribution of heavy rainfall events in the Midwest. *Journal of Water*
- 2824 *Resources Planning and Management* 121, 110-115.
- 2825
- 2826 Baldwin, C. and U. Lall, 1999: Seasonality of streamflow: the upper Mississippi River, *Water Resources Research*
- 2827 35(4), 1143.
- 2828
- 2829 Barnett, T., R. Malone, W. Pennell, D. Astammer, B. Demter, and W. Washington, 2004: The effects of climate change
- 2830 on water resources in the West: introduction and overview. *Climatic Change* 62, 1-11.
- 2831
- 2832 Beard, D., 1993: *Blueprint for Reform: The Commissioner's Plan for Reinventing Reclamation*. Bureau of Reclamation,
- 2833 Washington, D.C.
- 2834
- 2835 Benestad, R.E., 2004: Empirical-statistical downscaling in climate modeling. *EOS, Transactions, American*
- 2836 *Geophysical Union* 85, 417-422.
- 2837
- 2838 Boroughs, C.B. and E.A. Zagona. 2002: Daily flow routing with the Muskingum-Cunge method in the Pecos River
- 2839 RiverWare Model, *Proceedings of the Second Federal Interagency Hydrologic Modeling Conference*, Las
- 2840 Vegas, NV.
- 2841
- 2842 Bradley, A., S. Schwartz, and T. Hashino, 2004: Distributions-oriented verification of ensemble streamflow predictions,
- 2843 *Journal of Hydrometeorology* 5(3), 532-545.
- 2844

- 2845 Bureau of Reclamation, 1992: *A Long Term Framework for Water Resource Management, Development, and*
2846 *Protection*. U.S. Department of Interior, Washington, DC.
- 2847
- 2848 Carroll, T., 1985: Snow surveying, in *Yearbook of Science and Technology*, pp. 386-388, McGraw-Hill, New York,
2849 N.Y.
- 2850
- 2851 Carroll, T., 1999: personal communication, National Operational Hydrologic Remote Sensing Center, National Weather
2852 Service.
- 2853
- 2854 Carron, J., E. Zagona, and T. Fulp, 2006: Modeling uncertainty in an object-oriented reservoir operations model. *J.*
2855 *Irrig. and Drain. Engrg.*, 132(2), 104-111.
- 2856
- 2857 Changnon, S.A. (1990) The dilemma of climatic and hydrologic forecasting for the Great Lakes. In: *Proceedings of The*
2858 *Great Lakes Water Level Forecast and Statistics Symposium*, H.C. Hartmann and M.J. Donahue (Eds.), Great
2859 Lakes Commission, Ann Arbor, MI, pp. 13-25.
- 2860
- 2861 Changnon, D., 2000: Who used and benefited from the El Nino forecasts? In: *El Nino 1997-1998: The Climate Event of*
2862 *the Century*, S.A. Changnon (Ed.), Oxford University Press, New York, NY, pp. 109-135.
- 2863
- 2864 Christensen, N. and D.P. Lettenmaier, 2006: A multimodel ensemble approach to assessment of climate change impacts
2865 on the hydrology and water resources of the Colorado River basin, *Hydrology and Earth System Sciences*, 3, 1-
2866 44.
- 2867
- 2868 Christensen, N.S., A.W. Wood, N. Voisin, D.P. Lettenmaier, and R.N. Palmer, 2004: Effects of climate change on the
2869 hydrology and water resources of the Colorado River Basin. *Climatic Change* 62, 337-363.
- 2870 Clark, M.P., L.E. Hay, G.J. McCabe, G.H. Leavesley, and R.L. Wilby, 1999: Towards the use of atmospheric forecasts
2871 in hydrologic models, I, Forecast drift and scale dependencies. *EOS Transactions AGU* 80 Fall Meeting
2872 Supplement, Abstract H32G-10, F406-407.

2873

2874 Congressional Budget Office, 1997: *Water Use Conflicts in the West: Implications of Reforming the Bureau of*
2875 *Reclamation's Water Supply Policies*, Congressional Budget Office, Washington, DC.

2876

2877 Croley, T.E., 1990: Laurentian Great Lakes double-CO₂ climate change hydrological impacts. *Climatic Change* 17, 27-
2878 48.

2879 Croley, T., F. Quinn, K. Kunkel, and S. Changnon, 1998: Great Lakes hydrology under a transposed climate. *Climatic*
2880 *Change* 38, 405-433.

2881

2882 Davis, R. E. and T. Pangburn, 1999: Development of new snow products for operational water control and management
2883 in the Kings River Basin, California. *EOS Transactions AGU*, 81, Spring Meeting Supplement, Abstract
2884 H22D-07, S110.

2885

2886 Douglas, E.M., R.M. Vogel, and C.N. Kroll, 2000: Trends in flood and low flows across the U.S. *Journal of Hydrology*
2887 240, 90-105.

2888

2889 Duan, Q., H. V. Gupta, S. Sorooshian, A. N. Rousseau, and R. Turcotte, (eds.) 2002: *Calibration of Watershed Models*,
2890 American Geophysical Union, Washington, D. C.

2891

2892 Endreny, T., B. Felzer, J.W. Shuttleworth, and M. Bonell, 2003: Policy to coordinate watershed hydrological, social,
2893 and ecological needs: the HELP Initiative. In: *Water: Science, Policy, and Management*, R. Lawford, D. Fort,
2894 H. Hartmann, and S. Eden (Eds.), American Geophysical Union, Washington, DC, pp. 395-411.

2895

2896 Environmental Protection Agency, 1989: *The Potential Effects of Global Climate Change on the United States. Report*
2897 *to Congress*. J.B. Smith and D. Tirpak, (Eds), EPA Office of Policy, Planning and Evaluation, Washington,
2898 D.C.

2899

- 2900 Eschenbach, E.A., T. Magee, E. Zagona, M. Goranflo, and R. Shane, 2001: Goal Programming Decision Support
2901 System for Multiobjective Operation of Reservoir Systems. *Journal of Water Resources Planning and*
2902 *Management*, 127, 71-141.
- 2903
- 2904 Ezurkwal, B., 2005: The role and importance of paleohydrology in the study of climate change and variability. In:
2905 *Encyclopedia of Hydrological Sciences*, M.G. Anderson (Ed.), John Wiley and Sons, Ltd., West Sussex, UK.
- 2906
- 2907 Franz, K., H.C. Hartmann, S. Sorooshian, and R. Bales, 2003: An evaluation of National Weather Service ensemble
2908 streamflow predictions for water supply forecasting in the Colorado River Basin. *Journal of Hydrometeorology*
2909 4, 1105-1118.
- 2910
- 2911 Frederick, K., D. Major, and E. Stakhiv, (Eds.) 1997: *Climate Change and Water Resources Planning Criteria*. Kluwer
2912 Academic Publishers, Dordrecht, Netherlands.
- 2913
- 2914 Frevert, D.K., M.S. Cowan, and W.L. Lane, 1989: Use of stochastic hydrology in reservoir operation. *Journal of*
2915 *Irrigation and Drainage Engineering* 115, 334-343.
- 2916
- 2917 Frevert, D., T. Fulp, E. Zagona, G. Leavesley, and H. Lins, 2006: Watershed and River Systems Management Program:
2918 Overview of Capabilities. *J. Irrig. and Drain. Engrg.* 132(2), 92-97.
- 2919
- 2920 Gamble, J.L., J. Furlow, A.K. Snover, A.F. Hamlet, B.J. Morehouse, H. Hartmann, and T. Pagano, 2003: Assessing the
2921 impact of climate variability and change on regional water resources: the implications for stakeholders. In:
2922 *Water: Science, Policy, and Management*, R. Lawford, D. Fort, H. Hartmann, and S. Eden (Eds.), American
2923 Geophysical Union, Washington, DC, pp. 341-368.
- 2924
- 2925 Garen, D.C., 1992: Improved techniques in regression-based streamflow volume forecasting, *J. Water Resour. Planning*
2926 *and Manag.*, 118, 654-670.
- 2927

- 2928 Georgakako, A., 2006: Decision support systems for integrated water resources management with an application to the
2929 Nile Basin. In: *Topics on System Analysis and Integrated Water Resources Management*, A. Castelletti and R.
2930 Soncini-Sessa (Eds.), Elsevier, New York, NY.
- 2931
- 2932 Georgakakos, A., H. Yao, M. Mullusky, and K. Georgakakos, 1998: Impacts of climate variability on the operational
2933 forecast and management of the Upper Des Moines River Basin, *Water Resour. Res.*, 34, 799-821.
- 2934
- 2935 Georgakakos, K., E. Shamir, S. Taylor, T. Carpenter, and N. Graham, 2004: *Integrated Forecast and Reservoir*
2936 *Management INFORM - A Demonstration for Northern California Phase 1 Progress Report*. HRC Limited
2937 Distribution Rept. No. 17, Hydrologic Research Center, San Diego, CA.
- 2938
- 2939 Georgakako, K., N. Graham, T. Carpenter, A. Georgakakos, and H. Yao, 2005: Integrating climate-hydrology forecasts
2940 and multi-objective reservoir management for northern California. *EOS Transactions* 86, 122, 127.
- 2941
- 2942 Gilmore, A., T. Magee, T. Fulp, and K. Strezepek, 2000: Multiobjective optimization of the Colorado River.
2943 Proceedings of the ASCE 2000 Joint Conference on Water Resources Engineering and Water Resources
2944 Planning and Management, Minneapolis, MN.
- 2945
- 2946 Glantz, M.H., 1982: Consequences and responsibilities in drought forecasting- the case of Yakima, 1977, *Water Resour.*
2947 *Res.*, 18, 3-13.
- 2948
- 2949 Gleick, P.H. and D.B. Adams, 2000: *Water: The Potential Consequences of Climate Variability and Change for Water*
2950 *Resources of the United States*. Pacific Institute, Oakland, CA.
- 2951
- 2952 Grantz, K., B. Rajagopalan, E. Zagona, and M. Clark, 2007: Water management applications of climate-based
2953 hydrologic forecasts: case study of the Truckee-Carson River basin, Nevada. *Journal of Water Resources*
2954 *Planning and Management*.
- 2955

- 2956 Grayson, R., and G. Bloschl, 2000: *Spatial Patterns in Catchment Hydrology: Observations and Modelling*, Cambridge
2957 University Press, Cambridge, U. K.
2958
- 2959 Hamlet, A. F., and D. P. Lettenmaier, 1999: Columbia River streamflow forecasting based on ENSO and PDO climate
2960 signals, *J. Water Resour. Planning and Manag.*, 125, 333-341.
2961
- 2962 Hansen, J., M. Sato, R. Ruedy, A. Lacis, K. Asamoah, K. Beckford, S. Borenstein, E. Brown, B. Cairns, B. Carlson, B.
2963 Curran, S. de Castro, L. Druyan, P. Etwarrow, T. Ferede, M. Fox, D. Gaffen, J. Glascoe, H. Gordon, S.
2964 Hollandsworth, X. Jiang, C. Johnson, N. Lawrence, J. Lean, J. Lerner, K. Lo, J. Logan, A. Lueckert, M. P.
2965 McCormick, R. McPeters, R. Miller, P. Minnis, I. Ramberran, G. Russell, P. Russell, P. Stone, I. Tegen, S.
2966 Thomas, L. Thomason, A. Thompson, J. Wilder, R. Willson, and J. Zawodny, 1997: Forcings and chaos in
2967 interannual to decadal climate change, *J. Geophys. Res.*, 102, 25679-25720.
2968
- 2969 Harding, B.J., T.B. Sangoyomi, and E.A. Payton, 1995: Impacts of severe sustained drought on Colorado River water
2970 resources. *Water Resources Bulletin* 31, 815-824.
2971
- 2972 Hartmann, H.C., 1990: Impacts on Laurentian Great Lakes levels. *Climatic Change* 17, 49-68.
2973
- 2974 Hartmann, H.C., T.C. Pagano, S. Sorooshian, and R. Bales, (2002a): Confidence builders: evaluating seasonal climate
2975 forecasts from user perspectives. *Bulletin of the American Meteorological Society* 83(5), 683-698.
2976
- 2977 Hartmann, H.C., R. Bales, and S. Sorooshian, (2002b): Weather, climate, and hydrologic forecasting for the U.S.
2978 Southwest: a survey. *Climate Research* 21, 239-258.
2979
- 2980 Hartmann, H., A. Bradley, and A. Hamlet, (2003): Advanced hydrologic prediction for improving water management.
2981 In: Lawford, R., Fort, D., Hartmann, H., and S. Eden (Editors), *Water: Science, Policy, and Management*.
2982 Water Resources Monograph 16, American Geophysical Union, Washington, DC, pp.285-307.
2983

- 2984 Hartmann, H.C., 2005: Use of climate information in water resources management. In: *Encyclopedia of Hydrological*
2985 *Sciences*, M.G. Anderson (Ed.), John Wiley and Sons Ltd., West Sussex, UK, Chapter 202.
2986
- 2987 Hoerling, M. and J. Eischeid, 2007: Past peak water in the Southwest, *Southwest Hydrology* 6(10),18-19, 35.
2988
- 2989 Hydrological Sciences Branch, 2007: Evaluation Report for AWARDS ET Toolbox and RiverWare Decision Support
2990 Tools. NASA Goddard Space Flight Center, Greenbelt, MD, 28 pp. (URL:
2991 http://wmp.gsfc.nasa.gov/projects/project_RiverWare.php)
2992
- 2993 IPCC, 1990: *Scientific Assessment of Climate Change: Report of Working Group I to the First Assessment Report of the*
2994 *IPCC*. Cambridge University Press, Cambridge.
2995
- 2996 IPCC, 1995a: *Climate Change 1995: IPCC Second Assessment*. Cambridge University Press, Cambridge.
2997
- 2998 IPCC, 1995b: *Impacts, Adaptations and Mitigations: Contributions of Working Group II to the Second Assessment*
2999 *Report of the IPCC*. Cambridge University Press, Cambridge.
3000
- 3001 IPCC, 2001a: *Climate Change 2001: Synthesis Report*. Third Assessment Report of the IPCC. Cambridge University
3002 Press, Cambridge.
3003
- 3004 IPCC, 2001b: *Impacts, Adaptations, and Vulnerability: Contribution of Working Group II to the Third Assessment*
3005 *Report of the IPCC*. Cambridge University Press, Cambridge.
3006
- 3007 IPCC, 2007: *Climate Change 2007: Climate Change Impacts, Adaptation and Vulnerability*. Working Group II
3008 Contribution to the IPCC Fourth Assessment Report (<http://www.ipcc.ch/SPM6avr07.pdf>).
3009

- 3010 Jacobs, K., 2002: *Connecting Science, Policy, and Decision-Making: A Handbook for Researchers and Science*
3011 *Agencies*. Office of Global Programs, National Oceanic and Atmospheric Administration, Silver Spring, MD.
3012
- 3013 Jacobs, K. and R. Pulwarty, 2003: Water resource management: science, planning and decision-making. In: *Water:*
3014 *Science, Policy, and Management*, R. Lawford, D. Fort, H. Hartmann, and S. Eden (Eds.), American
3015 Geophysical Union, Washington, DC, pp. 177-204.
3016
- 3017 Jerla, C., 2005: *An Analysis of Coordinated Operation of Lakes Powell and Mead under Low Reservoir Conditions*.
3018 M.S. Thesis, University of Colorado-Boulder, Boulder, CO, 187 pp.
3019
- 3020 Kenney, D., 1995: Institutional options for the Colorado River. *Water Resources Bulletin* 31(5), 837-850.
3021
- 3022 Kruger, A., S. Khandelwal, and A. Bradley, 2007: AHPSVER: A web-based system for hydrologic forecast verification.
3023 *Computers and Geosciences* 33(6), 739-748.
3024
- 3025 Lawford, R., R. Try, and S. Eden, 2005: International research programs in global hydroclimatology. In: *Encyclopedia*
3026 *of Hydrological Sciences*, M.G. Anderson (Ed.), John Wiley and Sons, Ltd., West Sussex, UK.
3027
- 3028 Lee, D.H., Quinn, F.H., D. Sparks, and J.C. Rassam, 1994: Modification of Great Lakes regulation plans for simulation
3029 of maximum Lake Ontario outflows. *Journal of Great Lakes Research* 20, 569-582
3030
- 3031 Lee, D.H., T.E. Croley, II, and F.H. Quinn, 1997: Lake Ontario regulation under transposed climates. *Journal of the*
3032 *American Water Resources Association* 33, 55-69
3033
- 3034 Lee, D.H., 1999: Institutional and technical barriers to implementing risk-based water resources management: a case
3035 study. *Journal of Water Resources Planning and Management* 125, 186-193.

- 3036 Lettenmaier, D.P., 2003: The role of climate in water resources planning and management. In: *Water: Science, Policy,*
3037 *and Management*, R. Lawford, D. Fort, H. Hartmann, and S. Eden (Eds.), American Geophysical Union,
3038 Washington, DC, pp. 247-266.
- 3039
- 3040 Lettenmaier, D.P., E.F. Wood, and J.R. Wallis, 1994: Hydro-climatological trends in the continental United States,
3041 1948-88. *Journal of Climate* 7, 586-607.
- 3042
- 3043 Lettenmaier, D., A. Wood, R. Palmer, E. Wood, and E. Stakhiv, 1999: Water resources implications of global warming:
3044 a U.S. regional perspective. *Climatic Change* 43, 537-579.
- 3045
- 3046 Lins, H.F. and J.R. Slack, 1999: Streamflow trends in the United States. *Geophysical Research Letters* 26, 227-230
- 3047
- 3048 Lins, H.F. and E.Z. Stakhiv, 1998: Managing the nation's water in a changing climate. *Journal of the American Water*
3049 *Resources Association* 34, 1255-1264.
- 3050
- 3051 Lofgren, B.M., F.H. Quinn, A.H. Clites, R.A. Assel, A.J. Eberhardt, and C.L. Luukkonen, 2002: Evaluation of potential
3052 impacts on Great Lakes water Resources based on climate scenarios of two GCMs. *Journal of Great Lakes*
3053 *Research*. 28, 537-554.
- 3054
- 3055 Loomis, J., J. Koteen, and B. Hurd, 2003: Economic and institutional strategies for adapting to water-resource effects of
3056 climate change. In: *Water and Climate in the Western United States*. W. Lewis (Ed.), University Press of
3057 Colorado, Boulder, CO, pp. 235-249.
- 3058
- 3059 Mantua, N., S. Hare, Y. Zhang, J. M. Wallace, and R. Francis, 1997: A Pacific interdecadal climate oscillation with
3060 impacts on salmon production, *Bull. Amer. Meteor. Soc.*, 78, 1069-1079.
- 3061
- 3062 Matalas, N.C., 1997: Stochastic hydrology in the context of climate change. *Climatic Change* 37, 89-101.
- 3063

- 3064 Maurer, E., 2007: Uncertainty in hydrologic impacts of climate change in the Sierra Nevada, California, under two
3065 emissions scenarios. *Climatic Change* 82, 309-325.
3066
- 3067 Miles, E. L., A.K. Snover, A.F. Hamlet, B. Callahan, and D. Fluharty, 2000: Pacific northwest regional assessment: the
3068 impacts of climate variability and change on the water resources of the Columbia river basin. *Journal of the*
3069 *American Water Resources Association* 36, 399-420.
3070
- 3071 Milly, P., K. Dunne, and A. Vecchia, 2005: Global patterns of trends in streamflow and water availability in a changing
3072 climate, *Nature* 438, 347-350.
3073
- 3074 NRC, 1995) *Flood Risk Management and the American River Basin: An Evaluation*. National Academy Press,
3075 Washington, DC.
3076
- 3077 NRC, 1998a: *GCIP: A Review of Progress and Opportunities*. National Academy Press, Washington, DC.
3078
- 3079 NRC, 1998b: *Hydrologic Sciences: Taking Stock and Looking Ahead*. National Academy Press, Washington, DC.
3080
- 3081 NRC, 1999a: *Making Climate Forecasts Matter*. National Academy Press, Washington, DC.
3082
- 3083 NRC, 1999b: *A Vision for the National Weather Service: Road Map for the Future*. National Academy Press,
3084 Washington, DC.
3085
- 3086 NRC, 1999c: *Hydrologic Science Priorities for the U.S. Global Change Research Program: An Initial Assessment*,
3087 National Academy Press, Washington, DC.
3088
- 3089 NRC, 2004: *Analytical Methods and Approaches for Water Resources Project Planning*, National Academies Press,
3090 Washington, DC, 151 pp.

- 3091 NASA, 2005a: *Earth-Sun System Applied Sciences Program Water Management Program Element FY2005-2009 Plan*,
3092 Washington, DC.
- 3093
- 3094 NASA, 2005b: *Water Management Annual Report*, Goddard Space Flight Center, December.
- 3095
- 3096 NASA, 2006: Applied sciences program, *The Subcommittee on Hydrology Newsletter* 1, 12-14.
- 3097
- 3098 NASA, 2007: *Water Management Progress Report Jan-Mar 2007*, Goddard Space Flight Center, 27 April.
- 3099
- 3100 National Assessment Synthesis Team (2000) *Climate Change Impacts on the United States: The Potential*
3101 *Consequences of Climate Variability and Change*. U.S. Global Change Research Program, Washington, DC.
- 3102
- 3103 National Hydrologic Warning Council (2002) *Use and Benefits of the National Weather Service River and Flood*
3104 *Forecasts*. National Weather Service Office of Hydrologic Development, Silver Spring, Md.
- 3105
- 3106 Neumann, D., E. Zagona, and B. Rajagopalan, 2006: A decision support system to manage summer stream
3107 temperatures. *Journal of the American Water Resources Association* 42, 1275-1284.
- 3108
- 3109 Olsen, J.R., J.H. Lambert, and Y.Y. Haimes, 1998: Risk of extreme events under nonstationary conditions. *Risk*
3110 *Analysis* 18, 497-510.
- 3111
- 3112 Olsen, J.R., J.R. Stedinger, N.C. Matalas, and E.Z. Stakhiv, 1999: Climate variability and flood frequency estimation
3113 for the upper Mississippi and lower Missouri rivers. *Journal of the American Water Resources Association* 35,
3114 1509-1523.
- 3115
- 3116 Office of Global Programs, 2004: *Regional Integrated Sciences and Assessments*. National Oceanic and Atmospheric
3117 Administration, <http://www.risa.ogp.noaa.gov>, 17 March 2004.
- 3118

- 3119 Oudin, L., C. Perrin, T. Mathevet, V. Andreassian, and C. Michel, 2006: Impact of biased and randomly corrupted
3120 inputs on the efficiency and the parameters of watershed models. *Journal of Hydrology* pp. 1-2, 62-83.
3121
- 3122 Pagano, T.C., H.C. Hartmann, and S. Sorooshian, 2001: Using climate forecasts for water management: Arizona and the
3123 1997-98 El Nino, *Journal of the American Water Resources Association* 37, 1139-1153.
3124
- 3125 Pagano, T.C., H.C. Hartmann, and S. Sorooshian, 2002: Use of climate forecasts for water management in Arizona: a
3126 case study of the 1997-98 El Niño. *Climate Research* 21, 59-269.
3127
- 3128 Pagano, T., D. Garen, and S. Sorooshian, 2004: Evaluation of official Western U.S. seasonal water supply outlooks,
3129 1922-2002. *Journal of Hydrometeorology* 5(5), 896-909.
3130
- 3131 Payne, J.T., A.W. Wood, A.F. Hamlet, R.N. Palmer, and D.P. Lettenmaier, 2004: Mitigating the effects of climate
3132 change on the water resources of the Columbia River Basin. *Climatic Change* 62, 233-256.
3133
- 3134 Piechota, T.C., and J.A. Dracup, 1999: Long range streamflow forecasting using El Niño-Southern Oscillation
3135 indicators, *J. Hydrol. Engineer.*, 4, 144-151.
3136
- 3137 Piechota, T.C., F.H.S. Chiew, J.A. Dracup, and T.A. McMahon, 2001: Development of an exceedance probability
3138 streamflow forecast using the El Niño-Southern Oscillation, *J. Hydrol. Engineer.*, 4, 20-28.
3139
- 3140 Pielke, R.A., Jr., 1995: Usable information for policy: an appraisal of the U.S. global change research program. *Policy*
3141 *Sciences* 38, 39-77.
3142
- 3143 Pielke, R.A., Jr., 2001: *The Development of the U.S. Global Change Research Program: 1987 to 1994*. Policy Case
3144 Study, National Center for Atmospheric Research, Boulder, CO.

- 3145 Prairie, J.R., 2006: *Stochastic nonparametric framework for basin wide streamflow and salinity modeling: application*
3146 *for the Colorado River basin*. Civil Environmental and Architectural Engineering Ph.D. Dissertation,
3147 University of Colorado, Boulder, Colorado.
- 3148
- 3149 Pulwarty, R.S., 2002: *Regional Integrated Sciences and Assessment Program*. Office of Global Programs, National
3150 Oceanic and Atmospheric Administration, Silver Spring, MD.
- 3151
- 3152 Pulwarty, R.S. and K.T. Redmond, 1997: Climate and salmon restoration in the Columbia River basin: the role and
3153 usability of seasonal forecasts. *Bulletin of the American Meteorological Society* 78, 381-397.
- 3154
- 3155 Quinn, F.H., 1981: Secular changes in annual and seasonal Great Lakes precipitation, 1854-1979, and their implications
3156 for Great Lakes water resources studies. *Water Resources Research* 17, 1619-1624.
- 3157
- 3158 Quinn, F.H., 2002: Secular changes in Great Lakes water level changes. *Journal of Great Lakes Research* 28, 451-465.
- 3159
- 3160 Rayner, S., D. Lach, and H. Ingram, 2005: Weather forecasts are for wimps: why water resource managers do not use
3161 climate forecasts. *Climatic Change* 69, 197-227.
- 3162
- 3163 Reitsma, R.F., 1996: Structure and support of water resources management and decision making, *Journal of Hydrology*,
3164 177(1), 253-268.
- 3165
- 3166 Reitsma, R., I. Zigurs, C. Lewis, V. Wilson, and A. Sloane, 1996: Experiment with simulation models in water
3167 resources negotiations, *Journal of Water Resources Management and Planning*, ASCE 122, 64-70.
- 3168
- 3169 Salas, J.D., 1993: Analysis and modeling of hydrologic time series. In: *Handbook of Hydrology*, D.R. Maidment (Ed.),
3170 McGraw-Hill, Inc., New York, NY, Chapter 19.
- 3171

- 3172 Saunders, J.F., III and W.M. Lewis, Jr., 2003: Implications of climatic variability for regulatory low flows in the South
3173 Platte River Basin, Colorado. *Journal of the American Water Resources Association* 39, 33-45.
- 3174
- 3175 Seager, R., M. Ting, I. Held, Y. Kushnir, J. Lu, G. Vecchi, H.P. Huang, N. Harnik, A. Leetma, N.C. Lau, C. Li, J.
3176 Velez, and N. Naik, 2007: Model projections of an imminent transition to a more arid climate in Southwestern
3177 North America, *Science* 316(5828), 1181-1184.
- 3178
- 3179 Schaake, J., T. Hamill, R. Buizza, and M. Clark, 2007: HEPEX, the Hydrological Ensemble Prediction Experiment.
3180 *Bulletin of the American Meteorological Society*, in press.
- 3181
- 3182 Smith, J. B., K.C. Hallet, J. Henderson, and K.M. Strzepek, 2007: Expanding the tool kit for water management in an
3183 uncertain climate. *Southwest Hydrology*, 6, 24-35, 36.
- 3184
- 3185 Snover, A.K., A.F. Hamlet, and D.P. Lettenmaier, 2003: Climate change scenarios for water planning studies: pilot
3186 applications in the Pacific Northwest. *Bulletin of the American Meteorological Society* 84, 1513-1518.
- 3187
- 3188 Sousounis, P., G. Albercook, D. Allen, J. Andresen, A. Brooks, D. Brown, H.H. Cheng, M. Davis, J. Lehman, J.
3189 Lindeberg, J. Root, K. Kunkel, B. Lofgren, F. Quinn, J. Price, T.D. Stead, J. Winkler, and M. Wilson, 2000:
3190 *Preparing for a Changing Climate: The Potential Consequences of Climate Variability and Change for the*
3191 *Great Lakes*. U.S. Global Change Research Program, Washington, DC.
- 3192
- 3193 Stakhiv, E., 2003: What can water managers do about climate variability and change? In: *Water and Climate in the*
3194 *Western United States*. W. Lewis (Ed.), University Press of Colorado, Boulder, CO, pp. 131-142.
- 3195
- 3196 Tarboton, D., 1995: Hydrologic scenarios for severe sustained drought in the Southwestern United States. *Water*
3197 *Resources Bulletin* 31(5), 803-813.
- 3198

- 3199 Urbanas, B.R. and L.A. Roesner, 1993: Hydrologic design for urban drainage and flood control. In: *Handbook of*
3200 *Hydrology*, D.R. Maidment (Ed.), McGraw-Hill, Inc., New York, NY, Chapter 28.
- 3201 U.S. Army Corps of Engineers, 1992: *Guidelines for Risk and Uncertainty Analysis in Water Resources Planning,*
3202 *Volumes I and II.* IWR Report 92-R-1, 92-R-2. Institute for Water Resources, Fort Belvoir, VA.
- 3203
- 3204 U.S. Department of Interior, 2007: *Colorado River Interim Guidelines for Lower Basin Shortages and Coordinated*
3205 *Operations for Lake Powell and Lake Mead, Draft Environmental Impact Statement, Volume 1,* Bureau of
3206 Reclamation, Boulder City, NV (URL: [http://www.usbr.gov/lc/](http://www.usbr.gov/lc/region/programs/strategies/draftEIS/index.html)
3207 [region/programs/strategies/draftEIS/index.html](http://www.usbr.gov/lc/region/programs/strategies/draftEIS/index.html))
- 3208
- 3209 VanRheenen, N., A.W. Wood, R.N. Palmer, and D.P. Lettenmaier, 2004: Potential implications of PCM climate change
3210 scenarios for Sacramento-San Joaquin River basin hydrology and water resources. *Climatic Change* 62, 257-
3211 281.
- 3212
- 3213 Vicuna, S., E. Maurer, B. Joyce, J. Dracup, and D. Purkey, 2007: The sensitivity of California water resources to
3214 climate change scenarios. *Journal of the American Water Resources Association* 43, 482-498.
- 3215
- 3216 Vorosmarty, C., D. Lettenmaier, C. Leveque, M. Meybeck, C. Pahl-Wostl, J. Alcamo, W. Cosgrove, H. Grassl, H. Hoff,
3217 P. Kabat, F. Lansigan, R. Lawford, R. Naiman, 2004: Humans transforming the global water system. *EOS,*
3218 *Transactions, American Geophysical Union* 85, 509-514.
- 3219
- 3220 Walker, A. E., and B. E. Goodison, 1993: Discrimination of wet snow cover using passive microwaver satellite data,
3221 *Annals of Glaciology*, 17, 307-311.
- 3222
- 3223 Ward, R.C., R. Pielke Sr., and J. Salas, (Eds.) 2003: Special Issue: Is Global Climate Change Research Relevant to Day-
3224 to-day Water Resources Managers? *Water Resources Update* 124.
- 3225

- 3226 Welles, E., S. Sorooshian, G. Carter, and B. Olsen, 2007: Hydrologic verification: a call for action and collaboration.
3227 *Bulletin of the American Meteorological Society* 88(4), 503-511.
3228
- 3229 Westrick, K. J. and C. F. Mass, 2001: An evaluation of a high resolution hydrometeorological modeling system for
3230 prediction of a cool-season flood event in a coastal mountainous watershed. *Journal of Hydrometeorology* 2,
3231 161-180.
3232
- 3233 Westrick, K. J., P. Storck, and C. F. Mass, 2002: Description and evaluation of a hydrometeorological forecast system
3234 for mountainous watersheds, *Wea. and Forecasting*, 17, 250-262.
3235
- 3236 Wheeler, K., T. Magee, T. Fulp, and E. Zagona, 2002: Alternative policies on the Colorado River. *Proceedings of*
3237 *Natural Resources Law Center Allocating and Managing Water for a Sustainable Future: Lessons From*
3238 *Around the World*, Natural Resources Law Center, University of Colorado, Boulder, CO.
3239
- 3240 Wood, E., E.P. Maurer, A. Kumar, and D.P. Lettenmaier, 2002: Long-range experimental hydrologic forecasting for the
3241 eastern United States. *Journal of Geophysical Research, Atmospheres* 107, 4423-4429.
3242
- 3243 Wood, A.W., A. Hamlet, D.P. Lettenmaier, and A. Kumar, 2001: Experimental real-time seasonal hydrologic
3244 forecasting for the Columbia River Basin, *Proc., 26th Annual Climate Diagnostics and Prediction Workshop*,
3245 National Weather Service, PB92-167378, National Technical Information Service, Springfield, VA.
3246
- 3247 Woodhouse, C., and J.J. Lukas, 2006: Multi-Century Tree-Ring Reconstruction of Colorado Streamflow for water
3248 resource planning. *Climatic Change*, (78), 293-315.
3249
- 3250 Young, R.A., (Ed.) 1995: Special Issue: Managing the Colorado River in a severe sustained drought. *Water Resources*
3251 *Bulletin* 35, 779-944.
3252

- 3253 Zagona, E.A., T.J. Fulp, H. Goranflo, and R. Shane, 1998: RiverWare: a general river and reservoir modeling
3254 environment. *Proceedings of the First Federal Interagency Hydrologic Modeling Conference*, Las Vegas, NV,
3255 19-23 April, 5,113-120.
- 3256
- 3257 Zagona, E., T.J. Fulp, R. Shane, T. Magee, and H. Morgan Goranflo, 2001: RiverWare: a generalized tool for complex
3258 reservoir systems modeling. *Journal of the American Water Resources Association* 37, 913-929.
- 3259
- 3260 Zagona, E., T. Magee, D. Frevert, T. Fulp, M. Goranflo, and J. Cotter, 2005: RiverWare. In: *Watershed Models*, V.
3261 Singh & D. Frevert (Eds.), Taylor & Francis/CRC Press: Boca Raton, FL.
- 3262
- 3263
- 3264

APPENDIX B

List of Figures by Chapter

3265

3266

3267

3268

3269

Chapter 1:

3271

3272 **Figure 2:** The PECAD Decision Support System: Data Sources and Decision Support Tools

3273 **Figure 2:** The PECAD Decision Support System: Earth System Models, Earth Observations, Decision Support
3274 Tools, and Outputs

3275 **Figure 3:** The PECAD Decision Support System: The Role of Convergence of Evidence Analysis

3276 **Figure 4:** The PECAD Decision Support System: Information Sources for the Convergence of Evidence
3277 Analysis

3278

3279

Chapter 2:

3281

3282 **Figure 1:** Configuration of CMAQ-based Decision Support System for climate change impact study

3283

Chapter 3:

3285

3286 **Figure 1:** Example of HOMER output graphic.

3287 **Figure 2:** Example of ocean wind resource assessment output for the offshore regions of Pakistan.

3288

3289

3290

3291

3292 **Chapter 4:**

3293

3294 **Figure 1:** Relationship between the occurrence of black-legged tick presence at a site and minimum
3295 temperature (top) and evaluation of model (bottom)..

3296 **Top Panel:** Log odds plot for relationship between *I. Scapularis* population maintenance and minimum
3297 temperature (T).

3298 **Bottom Panel:** ROC Plot describing the accuracy of the autologistic model.

3299 **Figure 2:** Forecast geographic distribution of the black-legged tick vector east of the
3300 100th meridian in the United States for DSSPL.

3301 **Figure 3:** Forecast change in black-legged tick distribution in Eastern and Central North America under
3302 climate change scenarios using DSSPL

3303

3304 **Chapter 5:**

3305

3306 **Figure 1:** Illustration depicting the flow of information

3307

3308

3309

Appendix C

3310

Glossary, Acronyms, Symbols & Abbreviations

3-D	Three-dimensional
ACD	Atmospheric Chemistry Division
AERONET	Aerosol RObotic NETwork
AgRISTARS	Agriculture and Resources Inventory Surveys through Aerospace Remote Sensing
AI	Aerosol Index
AOD	Aerosol Optical Depths
ASOS	Automated Surface Observation Stations
ATSR	Along Track Scanning Radiometer
AVHRR	Advanced Very High Resolution Radiometer
AWN	Automated Weather Network
BC	Boundary Conditions
BELD3	Biogenic Emissions Land Use Database version 3
CAAA	Clean Air Act and its Amendments
CAM	Community Atmosphere Model
CAMx	Comprehensive Air quality Model with Extensions
CBv	Carbon Bond V
CCSP	Climate Change Science Program
CDC	Disease Control and Prevention
CEMPI	Center for Environmental Modeling for Policy Development
CENR	Committee on Environment and Natural Resources Research
CH4	Methane
CM2	Climate Model 2
CMAQ	Community Multiscale Air Quality

CMAS	Community Modeling and Analysis System
CNES	Centre National d'Etudes Spatiales
CO₂	Carbon Dioxide
CONUS	Continental United States
CTM	Chemistry Transport Modeling
DDSPL	Decision Support System to Prevent Lyme disease
DEM	Digital Elevation Models
DG	Distributed generation
DLR	German Aerospace Center (DLR) (German: Deutsches Zentrum für Luft- und Raumfahrt e.V.)
DMSP	Defense Meteorological Satellite Program
DSSs	Decision Support Systems
DSTs	Decision Support Tools
ECMWF	European Centre for Medium-Range Weather Forecasts
EPA	Environmental Protection Agency
ESMP	Earth System Modeling Framework
ESRI	Environmental Science and Research Institute
ESRL	Earth Systems Research Laboratory
ESSL	Earth and Sun Systems Laboratory
EUTMETSAT	European Organization for the Exploitation of Meteorological Satellites
FAS	Foreign Agricultural Service
GACP	Global Aerosol Climatology Project
GADS	Global Aerosol Dataset
GCM	Global Climate Model
GCTM	Global Chemistry Transport Models
GEO	Group on Earth Observations
GEOS	Goddard Earth Observing System
GEWEX	Global Energy and Water Experiment
GFDL	Geophysical Fluid Dynamics Laboratory

GhTOC	HourlyTotal Ozone Column
GIS	geographic information system
GISS	Goddard Institute for Space Studies
GLCC	Global Land Cover Characterization
GMAO	Global Modeling and Assimilation Office
GOCAT	Global Ozone Chemistry Aerosol Transport
GOES	Geostationary Environmental Operational Satellite
GOME	Global Ozone Monitoring Experiment
GsT	Geospatial Toolkit
GTCM	Global Tropospheric Chemistry Model
GTS	Global Telecommunications System
HMS	Hazard Mapping System
HOMER	Hybrid Optimization Model for Electric Renewables
IGOL	Integrated Global Observations of Land
INPE	Brazilian Spatial Institute
IPCC	Intergovernmental Panel on Climate Change
ISH	Integrated Surface Hourly
KAMM	Karlsruhe Atmospheric Mesoscale Model
kw	Kilowatt
h	Hour
m	Meter
LACIE	Large Area Crop Inventory Experiment
LULC	Land Use and Land Cover
MATCH	Model of Atmospheric Transport and Chemistry
MCIP	Metorology-Chemistry Interface Processor
MIST	Multi-Angle Imaging Spectroradiometer
MM5	Mesoscale Model Version 5
MODIS	Moderate Resolution Imaging Spectroradiometer

MOZART	Model of Ozone and Related Chemical Tracers
N₂O	Nitrous oxide
NASA	National Aeronautics and Space Administration
JPL	Jet Propulsion Laboratory
SSE	Surface meteorology and Solar Energy
USGS	US Geological Survey
NCAR	National Center for Atmospheric Research
NCAR- NCEP	National Center for Atmospheric Research-National Centers for Environmental Protection
NCDC	National Climatic Data Center
NCEP	National Centers for Environmental Prediction
NOAA	National Oceanic and Atmospheric Administration
NRC	National Research Council
NREL	National Renewable Energy Laboratory
NSTC	National Science and Technology Council
OCO	Orbiting Carbon Observatory
OMI	Ozone Monitoring Instrument
AOT	aerosol optical thickness
TOC	Total Ozone Content
P	Wind Power Density
PECAD	Production Estimate and Crop Assessment Division
CADRE	Crop Condition Data Retrieval and Evaluation system
PNNL	Pacific Northwest National Laboratory
PV	Photovoltaic
RCM	Regional Climate Model
ReGAP	Regional Gap Analysis Program
RPOs	Regional Program Organizations
SAP	Synthesis and Assessment Product
SHEDS	Stochastic Human Exposure and Dose Simulation

SIP	State Implementation Plans
SPOT	Systeme Pour L'Observation de la Terre
SRTM	Shuttle Radar Topographic Mission
SSE	Surface meteorology and Solar Energy
SSM/I	Special Sensor Microwave/Imager
SSMR	Scanning Multichannel Microwave Radiometer
STAR	Science to Achieve Results
SWERA	Solar and Wind Energy Resource Assessment
TES	Tropospheric Emission Spectrometer
TOMS	Total Ozone Mapping Spectrometer
TOMS	Total Ozone Mapping Spectrometer
TRI,	Total Risk Integrated Methodology
U.S.	United States
UIUC	Unknown Sent email to Daewon
UNC	University of North Carolina at Chapel Hill
UNEP	United Nations Environment Programme
USDA	Department of Agriculture
EROS	Earth Resources Observation Systems
USWRP	United States Weather Research Program
V	Wind Speed
WAsP	Wind Atlas Analysis and Application Program
WRAMS	Wind Resource Assessment Mapping System
WRDC	World Radiation Data Centre
WRF	Weather Research and Forecasting

3311

3312

3313