

A Dendroclimatic Record of Paleoclimate of the Last 10,000 Years, Glacier Bay National Park and Preserve 2005 Progress Report

Studies Conducted As Part of Research Project: Long-term tidewater and terrestrial glacier dynamics, glacier hydrology, and Holocene and historic glacier activity and climate change in Glacier Bay National Park and Preserve

Daniel Lawson¹, Greg Wiles², Laura Conkey³ and David Finnegan¹

¹ CRREL, 72 Lyme Road, Hanover, NH 03755

² Department of Geology, College of Wooster, Wooster, OH 44691

³ Department of Geography, Dartmouth College, Hanover, NH 03755

Abstract

Global climate is changing, and humans may have a significant role in affecting those changes. Our knowledge of how the climate system works is hampered by a lack of long-term climatic data, which are needed to demonstrate the full range of natural variability of the climate system on both short and long time scales. In addition, as global warming progresses, we expect to see some of the greatest effects in the higher latitudes. It is with this need in mind that we are conducting research on the paleoclimate of Glacier Bay, a climatically-sensitive region of the North Pacific region. Our research involves analysis of the climatically-driven glacial cycle during the Holocene, and linking this record to a unique 10,000-year chronology of high-latitude climatic events using tree rings from ancient wood of trees overridden and killed by successive ice advances. We began collecting cores of modern trees to compare the modern chronologies with instrumented climatic data and assess the strength and power of tree growth to climate.

Our continuing primary objective is to collect sections of ancient trees overridden by the glaciers during the Holocene excursions across Glacier Bay before they are lost to erosion and decay. The recovery and processing of these samples is the primary resource in building a tree ring record of the last 10,000 years. Presently we are engaged in showing that this effort is possible and that the ring-width data we are collecting from archived and samples taken in 2005 reflect paleoclimate by crossdating the Glacier Bay records with other records from The Gulf of Alaska..

Introduction

Normally heavy snowfall in the high mountains surrounding Glacier Bay feeds one of the larger active glacier complexes in North America, a part of the fourth largest glaciated region in the world (Meier 1984) and recent changes in Glacier Bay alone have had a

significant effect on global sea level (Arendt et al., 2002; Larsen et al. 2005). With the exception of some lowlands at the southeastern and southwestern margins, Glacier Bay was covered by ice as recently as 250 years ago during the Little Ice Age. Glacial retreat since that time is one of the best documented in the world, with ice margins retreating distances as far as 90 km at some of the highest rates ever recorded. During this retreat, forests that were overridden by ice advance were uncovered, and are now being exposed by erosion, uplift of shoreline areas and continued retreat of terrestrial glaciers. These interstadial forests reveal that in addition to the advance during the Little Ice Age, ice apparently advanced into Glacier Bay several other times beginning about 12,500 years ago (Lawson et al, 2006).

Numerous global warming scenarios (IPCC 2001) indicate that Arctic and Subarctic regions are particularly sensitive to current and predicted climatic changes, but our knowledge is hampered by relatively short-term climatic records from instrumented sources. The large repository of interstadial wood within Glacier Bay could provide detailed data on long-term changes in the climate of the North Pacific region, a region thought to be particularly sensitive to annual, decadal and longer periods of climatic change. Paleoclimate data from our analysis will provide critical parameters that are now lacking but required to calibrate Global Climate Models (GCMs) and better predict future changes in climate. Other scientists studying modern and exhumed wood from areas in the western Gulf of Alaska have found that the wood samples crossdate and correlate with climate (Barclay et al. 1999, Wiles et al. 1999). However, these studies are limited in their expanse of time, spanning only the last 1000 years or so; the former forests in Glacier Bay provide the only known Subarctic North American repository of wood that may continuously span the last 10,000 years.

As part of our research in Glacier Bay in 2005, we initiated the dendrochronological analysis of new wood sections and crossdating of samples from the period 1400 years BP, a period of time with few tree ring records in any Subarctic region. With such data, various parameters such as long-term annual and seasonal temperatures and trends in precipitation can be assessed. We also obtained cores of living trees to examine the relationship between modern measurements of climate and tree-ring width, and to link these modern (last 100 – 200 years) tree-ring records to those of the first millennia for which we acquired additional sections of interstadial trees.

2005 Objectives

Our studies of the paleoclimate of Glacier Bay in 2005 specifically focused on the following:

- 1) Collect tree cores from extant trees growing at locations that we now know were free of ice during the last glacial advance to examine the relationship of tree-ring widths to local climate records and define our ability to crossdate modern tree ring records;
- 2) Obtain sections of interstadial wood that grew within the last 1000 to 1200 years;
- 3) Examine selected groups of existing radiocarbon-dated cross sections and initiate cross-dating with tree ring records from the western Gulf of Alaska.

Methodology

We collected cores of modern trees to meet two requirements. First, there is the need to assess the suitability of the two major tree species in the Park, Sitka spruce (*Picea sitchensis* (Bong.) Carr.) and western hemlock (*Tsuga heterophylla* (Raf.) Sarg.), as sources of tree-ring records that reflect climate. Dendroclimatic studies involve the statistical comparison of tree growth (such as measured ring widths) to important climatic factors such as regional temperatures; this is most often accomplished with instrumented climate records (Fritts 1976). We must show that modern trees have tree-ring widths that statistically vary with climate, thus providing the basis for future analyses of paleoclimate in the 10,000-year chronology we plan to establish. Climate data now being collected at sites across Glacier Bay (Finnegan et al 2006) will provide data on how climate varies regionally across the Park and allow us to determine how representative historical records from climate stations outside Glacier Bay reflect the climate within the bay.

Secondly, the radiocarbon dates from the exhumed wood are at their youngest 250 to 500 years old, which puts the older wood outside the range (~100 years) of instrumented climatic data for this region. In order to know the exact year of growth on each of these older specimens, we need to connect them to the present day, which can be done with the oldest still-living trees that we have begun to sample. We know of at least three locations where trees of up to 700 years in age still live in the Park; it is these trees we have begun to sample in order to develop a long-term chronology of exact known ages using both living trees and the older cross-sections.

Tree-ring records of the interstadial trees are developed from sections cut from in situ stumps and logs in glacial sediments. Small pieces of wood from each stump sample are radiocarbon dated using the high-resolution AMS technique. We must examine multiple sections to insure reproducibility of the records and account for missing rings; rings may be lost due to various stresses in the environment, as well as to account for sub-regional differences that may be present in the climate. Each sample is located by GPS, photographed and various parameters recorded about the wood, such as dimensions, position of each sample section relative to the roots, and species. We base each year's sampling on the results of dating and off-season lab work.

The tree-ring analyses are conducted by students and PI's in the tree-ring laboratories at Dartmouth College, the College of Wooster and CRREL.

2005 Results

We focused our efforts on obtaining cores of living trees for subsequently calibrating modern climate signals to the tree-ring record, and obtaining select cross sections of ancient trees overridden by glaciers at different times during the Holocene to develop the tree-ring record of paleoclimate.

We cored living trees from five stands of old and young growth spruce and hemlock, obtaining sections from Dundas Bay and Bartlett Cove. Several of the sites that we sampled in Dundas Bay have trees that are at least 600-700 years old, which will allow for some age overlap with the youngest ancient cross-sections previously sampled in lower Glacier Bay. These cores are critical to filling gaps in the paleo-record and linking our ancient wood data to that of the present, thus making the annual and seasonal reconstructions of climate from the longer chronology. Core samples of the older trees in Dundas Bay will allow us to explore the statistical relationships between tree growth and climate in the period of roughly 100 years for which we have modern climatic data. The shorter (approx. 200 year-old) chronologies sampled previously at Bartlett Cove, Geikie Inlet and a higher elevation site above Tlingit Point will provide depth in the more recent time period, along with good spatial and altitudinal coverage of the Bay.

The initial analysis of the tree cores is underway in the Tree-Ring Lab at Dartmouth College. We have prepared the cores for analysis and completed dating of those from Bartlett Cove area. We are now at work on the longer record samples from Dundas Bay. Some of these cores however have proven unuseable due primarily to gaps in the record and additional core sampling is planned for 2006. Data from these cores will go a long way towards bridging the gap between the present day (since the trees have known ring dates) and the youngest cross sections of interstadial wood for which we have approximate (radiocarbon) dates, and ultimately allow for exact calendar-dating of many of the cross sections we have sampled.

We also obtained 20 cross-sections from newly exposed, in situ interstadial stumps and logs across Glacier Bay, focusing our efforts on areas that we believed would fill gaps in the longer chronology. Sections were obtained from the head of Geikie Inlet where radiocarbon dating had previously shown the ancient trees were growing during the critical time period of ~1800 to 1100 years BP. We also obtained 4 sections from younger stumps on Lester Island, where a single previous radiocarbon date indicated an age of about 500 years BP. Both groups of sections are critical to linking modern and ancient tree-ring records. The interstadial cross sections have been processed at CRREL and we have initiated their analysis at the Wooster Tree Ring Lab of the sections of ~ 1000 yrs BP interval. Selected 1000 yr BP samples from previous years were also examined to determine whether they crossdated with tree-ring records from Columbia Bay, Prince William Sound in the central Gulf of Alaska region. Although limited samples were available, these results are extremely encouraging (Figure 1). The tree-ring record of ten logs from Glacier Bay was compiled into a ring-width series and then dated with the master chronology from Columbia Bay (Wiles, unpublished). The dating, although preliminary, is encouraging and we plan to increase the sample size through this important interval to strengthen the common signal in the record. This work is significantly extending the Gulf of Alaska tree-ring record by almost 400 years (Figure 1) and is extremely encouraging for the future development of dendroclimatology in the North Pacific region.




Figure 1. Tree-ring dating of the Glacier Bay samples. The green curve is a ring-width record composite from logs in Columbia Bay, Prince William Sound. The inset shows graphically the matching ring-width variations from the Glacier Bay chronology and the Prince William Sound record.

Continuing Work

Based on our initial results, we anticipate being able to show that the modern trees growing in Glacier Bay exhibit variations in tree-ring widths that relate to climatic changes, mainly temperature and precipitation. Similar species of trees in other parts of the Gulf of Alaska have been shown to vary climatically (Barclay et al. 1999, Wiles et al. 1999). We will continue laboratory analyses of the ancient sections of wood of the last millennia over the next 6 months, as well as to complete the analysis of the modern core records. During field work in 2006, we will continue to obtain crucial cores and sections to complete the record of the last millennia and allow us to verify what our limited sampling has shown to date – a strong crossdating with the western Gulf tree-ring records and thus a clear link to ancient changes in climate.

Literature Cited

Arendt, A. A., K.A. Echelmeyer, W.D. Harrison, C.S. Lingle, and V.B. Valentine, 2002. Rapid wastage of Alaska glaciers and their contribution to rising sea level, *Science*, 297, 382-386.

Barclay, D.J., Wiles, G.C. and Calkin, P.E., 1999. A 1119-year tree-ring-width chronology from western Prince William Sound, southern Alaska. *Holocene* 9(1):79-84.

Finnegan, D.F., Lawson, D.E. and Kopczynski, S.E. 2006. Assessing contemporary and Holocene glacial and glacial-marine environments. Proceedings Glacier Bay Science Symposium, Juneau, AK, Oct. 2004. In Press.

Fritts, H.C. 1976. *Tree Rings and Climate*. New York: Academic Press.

IPCC (Intergovernmental Panel on Climate Change) 2001. *Climate Change 2000: The Science of Climate Change*. Cambridge and New York: Cambridge University Press.

Larsen, C.F., Motyka, R.J., Freymuller, J.T., Echelmeyer, K.A., and Ivins, E.R., 2004. Rapid uplift of southern Alaska caused by recent ice loss. *Geophys. Int. Jour.*, 158(3). 1118-1133.

Lawson, D.E., Finnegan, D.C., Kopczynski, S.E., and Bigl, S.R. 2004. Long-term studies of tidewater and terrestrial glacier dynamics, glacier hydrology, and Holocene and historic climate activity, Glacier Bay National Park and Preserve, Alaska. 2004 Progress Report to Glacier Bay National Park and Preserve. Unpublished, Cold Regions Research and Engineering Laboratory, Hanover, NH.

Lawson, D.E., Finnegan, D.F., Kopczynski, S.E. and Bigl, S.B. 2006. Early to mid-Holocene glacier fluctuations in Glacier Bay, Alaska. Proceedings Glacier Bay Science Symposium, Juneau, AK, Oct. 2004. In Press.

Meier, M, 1984. Contribution of Small Glaciers to Global Sea Level. *Science* 226(4681) 1418-1421

Wiles, G.C., Barclay D.J., and Calkin P.E. 1999. Tree-ring-dated "Little Ice Age" histories of maritime glaciers from western Prince William Sound, Alaska. *Holocene* 9(2):163-173.