

# Polymorphisms in the novel gene acyloxyacyl hydroxylase (AOAH) are associated with asthma and associated phenotypes

Kathleen C. Barnes, PhD,<sup>a,b,c</sup> Audrey Grant, MS,<sup>a</sup> Peisong Gao, MD, PhD,<sup>a</sup> Daniela Baltadjieva, MD, PhD,<sup>a</sup> Tiina Berg, BS,<sup>a</sup> Peter Chi, MHS,<sup>a</sup> Shu Zhang, MA,<sup>a</sup> April Zambelli-Weiner, PhD,<sup>a</sup> Eva Ehrlich, MS,<sup>a</sup> Omeed Zardkoobi, MD,<sup>d</sup> Mary E. Brummet, MS,<sup>a</sup> Maria Stockton, MA,<sup>a</sup> Tonya Watkins, BS,<sup>a</sup> Li Gao, MD, PhD,<sup>a</sup> Marquita Gittens, PhD,<sup>e</sup> Marsha Wills-Karp, PhD,<sup>f</sup> Christopher Cheadle, PhD,<sup>a</sup> Lisa A. Beck, MD,<sup>a,g</sup> Terri H. Beaty, PhD,<sup>b</sup> Kevin G. Becker, PhD,<sup>h</sup> Joe G. N. Garcia, MD,<sup>c</sup> and Rasika A. Mathias, ScD<sup>b,i</sup> Baltimore, Md, Barbados, West Indies, and Cincinnati, Ohio

**Background:** The gene encoding acyloxyacyl hydroxylase (AOAH), an enzyme that hydrolyzes secondary fatty acyl chains of LPS, is localized on chromosome 7p14-p12, where evidence for linkage to total IgE (tIgE) concentrations and asthma has been previously reported.

**Objective:** We hypothesized that variants in AOAH are associated with asthma and related phenotypes. Because both AOAH and soluble CD14 respond to LPS, we tested for gene-gene interaction.

**Methods:** We investigated the association between 28 single nucleotide polymorphisms throughout the AOAH gene and asthma, concentrations of tIgE, the ratio of IL-13/IFN- $\gamma$ , and soluble CD14 levels among 125 African Caribbean, multiplex asthmatic pedigrees (n = 834). Real-time PCR was used to assess whether AOAH cDNA expression differed with AOAH genotype.

**Results:** Significant effects were observed for all 4 phenotypes and AOAH markers in 3 distinct regions (promoter, introns 1-6, and the intron 12/exon 13 boundary/intron 13 region) by means of single-marker and haplotype analyses, with the strongest evidence for a 2-single-nucleotide-polymorphism haplotype and log[tIgE] ( $P = .006$ ). There was no difference in AOAH expression levels by AOAH genotype for any of the markers. Comparing genotypic distributions at both the AOAH marker rs2727831 and *CD14*(-260)C>T raises the possibility of gene-gene interaction ( $P = .006-.036$ ).

**Conclusion:** Our results indicate that polymorphisms in markers within the AOAH gene are associated with risk of asthma and associated quantitative traits (IgE and cytokine levels) among asthmatic subjects and their families in Barbados, and there is an interactive effect on tIgE and asthma concentrations between an AOAH marker and the functional *CD14*(-260)C>T polymorphism.

**Clinical implications:** AOAH is a novel innate immunity candidate gene associated with asthma and related phenotypes in an African ancestry population. (*J Allergy Clin Immunol* 2006;118:70-7.)

**Key words:** *CD14*, acyloxyacyl hydroxylase, association, asthma, total IgE, soluble *CD14*, family-based association test

From <sup>a</sup>the Division of Allergy and Clinical Immunology, Department of Medicine, The Johns Hopkins University, Baltimore; <sup>b</sup>the Department of Epidemiology, Bloomberg School of Public Health, Johns Hopkins University, Baltimore; <sup>c</sup>the Division of Pulmonary and Critical Care Medicine, Department of Medicine, Johns Hopkins School of Medicine, Baltimore; <sup>d</sup>Johns Hopkins School of Medicine, The Johns Hopkins University, Baltimore; <sup>e</sup>the School of Clinical Medicine and Research, University of the West Indies, Barbados; <sup>f</sup>the Division of Immunobiology, Children's Hospital Medical Center, Cincinnati; <sup>g</sup>the Department of Dermatology, Johns Hopkins University, Baltimore; <sup>h</sup>DNA Array Unit, National Institute of Aging, Baltimore; and <sup>i</sup>the Genometrics Section, Inherited Disease Research Branch, National Human Genome Research Institute, National Institutes of Health, Baltimore.

Supported by National Institutes of Health (NIH) grants U01 HL66615, AI50024-03, and HL67736 and an Asthma and Allergy Foundation of America New Investigator Award Grant (KCB). K. C. B. was supported in part by the Mary Beryl Patch Turnbull Scholar Program, and R. A. M. was supported by the Intramural Research Program of the National Human Genome Research Institute, National Institutes of Health.

Disclosure of potential conflict of interest: R. A. Mathias is employed by the Inherited Disease Research Branch, National Human Genome Research Institute, and the National Institutes of Health. The rest of the authors have declared that they have no conflict of interest.

Received for publication February 18, 2005; revised March 20, 2006; accepted for publication March 27, 2006.

Available online May 28, 2006.

Reprint requests: Kathleen C. Barnes, PhD, The Johns Hopkins Asthma and Allergy Center, 5501 Hopkins Bayview Circle, Rm 3A.62, Baltimore, MD 21224. E-mail: kbarnes@jhmi.edu.

0091-6749

doi:10.1016/j.jaci.2006.03.036

#### Abbreviations used

AOAH: Acyloxyacyl hydroxylase  
FBAT: Family-based association test  
LD: Linkage disequilibrium  
OR: Odds ratio  
sCD14: Soluble CD14  
SNP: Single nucleotide polymorphism  
tIgE: Total IgE

marker and asthma.<sup>11-13</sup> LeVan et al<sup>14</sup> showed this polymorphism is functional, resulting in higher soluble CD14 (sCD14) concentrations. To date, the *CD14*(-260)C>T polymorphism is one of the most reproducible associations for asthma and its associated traits.

Another interesting candidate, but one for which relatively little is known, is acyloxyacyl hydroxylase (AOAH), a highly conserved,<sup>15</sup> eukaryotic lipase that releases secondary acyl chains from the LPS found on cell walls of gram-negative bacteria.<sup>16-18</sup> A unique leukocyte enzyme, AOAH is potentially an important host defense molecule because deacylated LPS is at least 100-fold less potent than intact LPS.<sup>19</sup> *In vitro* assays show deacylated LPS is an effective LPS antagonist.<sup>20,21</sup> Recently, it was demonstrated that AOAH also appears to modulate B-cell proliferation and polyclonal antibody production in response to gram-negative bacterial infection.<sup>22</sup>

AOAH is located on chromosome 7p14-p12,<sup>23</sup> a region where evidence for linkage to asthma, bronchial hyper-reactivity, and tIgE concentrations has been reported.<sup>24-26</sup> Collectively, these observations suggest that AOAH is a candidate gene in the LPS signaling pathway for tIgE levels in particular and possibly for other allergic diseases, including asthma.

## METHODS

### Study subjects

Nuclear and extended asthmatic families were recruited in Barbados as a part of an ongoing asthma genetics study (described elsewhere<sup>27,28</sup>). Patients with a positive family history of asthma ( $\geq 1$  asthmatic siblings) were referred by physicians cooperating with coinvestigators at the University of the West Indies. Blood samples were collected from all subjects for serum extraction of DNA for genotyping, and total RNA was collected from a subset ( $n = 30$ ) for expression experiments.

Asthma and tIgE concentrations were measured, as described previously.<sup>27,29</sup> IgE values were log transformed to minimize skewness and then adjusted for age and sex. Soluble CD14 levels were measured with a commercially available ELISA kit supplied by Biosource (Europe S.A.), and concentration was determined by means of extrapolation from a standard curve estimated from a panel of standards with known concentrations. The minimum detectable concentration was estimated to be 1 ng/mL and defined as the sCD14 concentration corresponding to the average OD of 20 replicates of the zero standard +2 SD. Serum IL-13 and IFN- $\gamma$  levels were measured by means of ELISA with specific anti-IL-13 and anti-IFN- $\gamma$  mAbs obtained from BD Biosciences (San Diego,

Calif), according to the manufacturer's instructions. Serum IL-13 and IFN- $\gamma$  levels were log transformed to approximate a normal distribution. Family-based association tests (FBATs) on the qualitative phenotype asthma and the quantitative traits were performed on 834 persons from 125 pedigrees (both nuclear and extended).

All subjects provided verbal and written consent, as approved by the Johns Hopkins Bayview Hospital Institutional Review Board and the Ministry of Health in Barbados.

### Genotyping

A total of 29 single nucleotide polymorphisms (SNPs) from the gene encoding AOAH (accession no. NT\_007819.13) were genotyped in these data. Of these, 26 were validated SNPs selected from Applied Biosystems (Foster City, Calif) by using SNPBrowser software 3.0 ([www.allsnps.com/snpbrowser](http://www.allsnps.com/snpbrowser)), a freely available Applied Biosystems tool. Of these, one nonsynonymous variant (Gly/Ala) was not polymorphic in this African Caribbean population. Two additional SNPs that were part of an initial pilot study on AOAH were also included. The final panel comprised 28 SNPs, all of which were intronic, with the exception of four 5' UTR SNPs; this, however, reflects the lack of reported SNPs in exons in this gene ( $\sim 1\%$  exonic).

DNA was extracted by using standard protocols. When necessary, whole-genome amplification of DNA was prepared by OmniPlex Technology (Rubicon Genomics Inc, Ann Arbor, Mich). Genetic screening for the *CD14*(-260) variant (rs2569190) was performed as previously described<sup>29</sup> through PCR amplification and restriction digest (described elsewhere<sup>4</sup>). Genotyping of the AOAH SNPs was performed by using the TaqManR probe-based, 5' nuclease assay with minor groove binder chemistry (Applied Biosystems). Genotyping quality was high, with an average completion rate of 97%, with no discordances on repeat genotyping of a random 10% of the sample and a low rate of Mendelian inconsistencies.

### Quantitative RT-PCR analysis

Reverse transcription was performed by using total RNA isolated from human PBMCs from subjects selected on the basis of their rs2727831 (AOAH IVS12-355C>T) genotype (10 CCs, 10 CTs, and 10 TTs) and processed with the Applied Biosystems High-Capacity cDNA Archive kit first-strand synthesis system for RT-PCR, according to the manufacturer's protocol. Quantitative RT-PCR was performed by using the TaqMan assay system from Applied Biosystems. All PCR amplifications were carried out on an Applied Biosystems Prism 7300 Sequence Detection System by using a fluorogenic 5' nuclease assay (TaqMan probes). Probes and primers were designed and synthesized by Applied Biosystems. Relative gene expressions were calculated by using the  $2^{-Ct}$  method, in which Ct indicates cycle threshold, the fractional cycle number at which the fluorescent signal reaches a detection threshold.<sup>30</sup> The normalized  $\Delta Ct$  value of each sample was calculated by using a total of 3 endogenous human control genes (*GAPDH*, *ACTB*, and *PGK1*). Fold change values are presented as  $Averagefoldchange = 2^{-(AverageCt)}$  for genes in treated samples relative to control samples. Error bars represent the SEM for multiple biologic replicates (10 for each group).

### Statistical analyses

Clinical characteristics of the study population, including mean age, proportion by sex, proportion of asthmatic subjects, and mean values with SDs for serum measures and quantitative traits considered (log[tIgE], sIL-13, sIFN- $\gamma$ , the IL-13/IFN- $\gamma$  ratio, and sCD14), were calculated for all subjects with available data who were genotyped for AOAH, for founder members of the population, and for asthmatic and nonasthmatic subjects by using STATA 8.2 (StataCorp, College Station, Tex). Equality of means was tested between asthmatic and nonasthmatic subjects for serum measures and quantitative traits by

using linear regression, and proportions by *CD14*(-260) genotype were also calculated for these subsets of the study population.

Gene-gene interaction was investigated by using logistic regression models for asthma with interaction terms according to genetic transmission model, applying the robust “sandwich” estimator of variance in STATA 8.2.<sup>31</sup> Both regression coefficients and within-family correlation were estimated by using generalized estimating equations in regression models for quantitative traits.<sup>32</sup>

Mendelian inconsistencies were identified with Sib-Pair (version 99.9). Departures from Hardy-Weinberg equilibrium proportions at each marker were tested among founders. Pairwise linkage disequilibrium (LD) was measured with HaploView (<http://www-genome.wi.mit.edu/personal/jcbarret/haplo>). Association between markers in *CD14* and AOA and asthma, log[tIgE], sCD14, log[IL-13]/[IFN- $\gamma$ ], and haplotype associations were tested by using the FBAT (version 1.5.5).<sup>33,34</sup> For haplotype analyses, an expectation-maximization (EM) algorithm that maximizes the likelihood of phased haplotype frequencies based on all observed genotypes in the nuclear family was used. All analyses with FBAT were first conducted under a general genotype model and subsequently under an additive or recessive model, as suggested by the general model.

Multiple comparisons resulting from sliding-window haplotype tests is of concern, and the significance of haplotype tests was evaluated by using permutations to deal with this issue. Specifically, empiric *P* values are calculated through permutation analysis by using Monte Carlo methods under the null hypothesis of complete independence between haplotypes and the observed phenotypes. The FBAT sampling procedure stops when at least 100 Monte Carlo-based test statistic values are generated that are less than the observed value or after 100,000 replicate samples are generated, whichever comes first.

## RESULTS

Clinical characteristics of subjects and distribution of *CD14*(-260)C>T genotypes are presented in Table E1 (available in the Online Repository at [www.jacionline.org](http://www.jacionline.org)). We previously reported a significant association between the TT genotype at this *CD14* marker and lower asthma severity scores ( $P = .002$ ) in these families from Barbados.<sup>29</sup> The frequency of CC homozygotes among African Caribbean families (46%) is nearly twice that reported in non-African populations,<sup>4,5,7,8,35,36</sup> and the frequency of TT homozygotes (8%) is substantially lower.

Distributions of tIgE, serum IL-13, serum IFN- $\gamma$ , and sCD14 concentrations were similar to those described before.<sup>29,37</sup> Age and sex showed no effect on log[serum IL-13] or log[serum IFN- $\gamma$ ] levels. There was no significant difference in IL-13 levels between asthmatic subjects ( $728 \pm 1035$  pg/mL) and nonasthmatic subjects ( $604 \pm 1406$  pg/mL,  $P = .149$ ), nor were there significant differences in IFN- $\gamma$  levels when comparing asthmatic subjects ( $418 \pm 1002$  pg/mL) and nonasthmatic subjects ( $371 \pm 1244$  pg/mL,  $P = .614$ ). Neither log[serum IL-13] nor log[serum IFN- $\gamma$ ] levels were correlated with log[tIgE] levels. Serum IL-13 and IFN- $\gamma$  concentrations were highly correlated with each other ( $r = 0.6037$ ,  $P < .0001$ ). When IL-13 and IFN- $\gamma$  levels were analyzed as the log[IL-13]/log[IFN- $\gamma$ ] ratio, there was no difference in means by asthma status, and values were not correlated with log[tIgE] levels.

## Single-marker analyses

The 28 AOA SNPs spanning 313,068 bp are summarized in Table E2 (available in the Online Repository at [www.jacionline.org](http://www.jacionline.org)). All 28 SNPs were in Hardy-Weinberg equilibrium among founders ( $n = 269$ ). Fig 1 shows overall there was low LD in this gene, reflecting its large size and an average of 11.6 kb between neighboring SNPs. However, we did observe high LD for 13 of 27 pairings of contiguous SNPs ( $D'$  ranging from 0.84-1.0). This was reflected in three 2-SNP blocks, according to the definition of Gabriel et al,<sup>38</sup> comprised of markers rs3935953 and rs10275462 (block 1), rs4720210 and rs11770435 (block 2), and rs2727833 and rs2727831 (block 3), with a  $D'$  0.96, 1.0, and 0.94, respectively ( $r^2 = 0.69, 0.25, \text{ and } 0.37$ , respectively).


All single SNP analyses were first performed by using the robust genotype model (data not shown), and the strongest association signals suggested a recessive model was most likely; therefore all results are presented using a recessive model. All results of the 2-point tests for linkage and association between 28 AOA markers and asthma and log[tIgE], log[IL-13]/log[IFN- $\gamma$ ], and sCD14 concentrations with FBAT statistics are presented in Table E3 (available at [www.jacionline.org](http://www.jacionline.org)).

Eight SNPs provide evidence for significant association across the 4 phenotypes, 5 of which showed evidence for association with more than one phenotype. The most compelling associations were for the 2 markers in LD block 3 in intron 12: (1) rs2727833, with a strong negative association (ie, minor allele associated with lower levels) for log[tIgE] and log[IL-13]/log[IFN- $\gamma$ ] ( $P = .008$  and  $P = .008$ , respectively), and (2) rs2727831, with a similar negative association for log[tIgE], sCD14, and log[IL-13]/log[IFN- $\gamma$ ] ( $P = .006$ ,  $P = .041$ , and  $P = .023$ , respectively). Additional regions of the AOA gene highlighted by single-marker analyses included (1) the promoter region (rs1592497,  $P = .032$  for sCD14), (2) introns 3 and 4 (rs11978395,  $P = .014$  for sCD14; rs4720210,  $P = .022$  for log[tIgE] and  $P = .023$  for asthma; rs11979472,  $P = .012$  for log[IL-13]/log[IFN- $\gamma$ ]), and (3) introns 18 and 19 at the 3' end of the gene (rs1986468,  $P = .027$  for log[tIgE] and  $P = .002$  for log[IL-13]/log[IFN- $\gamma$ ]; rs3801297,  $P = .017$  for asthma and  $P = .010$  for log[IL-13]/log[IFN- $\gamma$ ]).

## Haplotype analyses

Restricting haplotype tests within observed LD blocks would have disregarded most of the data, and therefore a systematic sliding window approach was implemented, considering windows of 2 to 4 SNPs per window beginning with the first (5') marker and working across the gene one marker at a time. Haplotype tests revealed multiple association signals in 3 specific regions (defined as signals of association across contiguous SNPs with little to no LD between signals) that overlapped with the single-SNP results described above.

Significant evidence for linkage and association was observed for all 4 phenotypes in the promoter region



**FIG 1.** Genomic structure and LD estimates for 28 SNPs in the AOA gene. Illustration of gene structure indicating exons (pink), introns (light gray), and the flanking region (dark gray) spanning 211 kb and location of 28 SNPs drawn to scale with gene transcribed from left to right. Pairwise LD in 269 founders is represented as red squares for strong LD, blue squares for nonsignificant LD, and white squares for little or no LD. LD blocks are identified as noted.

(Fig 2). The strongest signals for each phenotype were as follows: 4-, 3-, and 2-SNP haplotypes, all beginning with marker rs2006882, for asthma ( $P = .016$ ), log[tIgE] ( $P = .002$ ), and log[IL-13]/log[IFN- $\gamma$ ] ( $P = .045$ ), respectively. The only association in this region for sCD14 occurred with the marker rs1592497 ( $P = .032$ ), as described above.

Significant associations for all 4 phenotypes were observed in the second region (rs10275462-rs2727822). Associations for asthma, log[tIgE], and log[IL-13]/log[IFN- $\gamma$ ] were centered between markers rs4720210 to rs11979472; 3- and 4-SNP windows provided the strongest signals for these 3 traits ( $P = .020$ ,  $P = .029$ , and  $P = .044$ , respectively). The 4-SNP window providing the strongest signal for sCD14 ( $P = .014$ ) was 5' to this marker but overlapped with signals for the other 3 phenotypes.

The third and most compelling region was the 3' end of AOA and included marker rs2727831. All 4 phenotypes demonstrated significant associations in this region flanked by markers rs2727833 and rs10951482. The most consistent signal was for log[tIgE], with 7 of 15 haplotype tests ( $P < .05$ ), including the 2-SNP haplotype of rs2727833 and rs2727831 ( $P = .006$ ). Modest associations were observed for asthma, log[IL-13]/log[IFN- $\gamma$ ], and sCD14 ( $P = .022$ ,  $P = .012$ , and  $P = .038$ , respectively).


Analyses performed using the additive model (data not shown) supported these results from a recessive model without revealing any additional signals.

### AOAH expression according to AOAH genotype

RNA samples from PBMCs were collected from 30 individuals stratified by the marker providing the most compelling evidence for association (rs2727831) to determine whether the C>T variant affected transcriptional levels of AOA. Subjects were also stratified according to 4 other AOA markers showing significant associations to more than one phenotype (rs4720210, rs2727833, rs1986468, and rs3801297). Comparison of cDNA expression of AOA according to the 3 genotypes at each AOA marker was performed for each of the 5 markers individually. There was no difference in AOA expression levels by genotype at any of the AOA markers tested.

### Evidence for gene-gene interaction between CD14(-260)C>T and AOAH marker rs2727831

We evaluated evidence for gene-gene interaction between AOA marker rs2727831 and the *CD14*(C-260) C>T polymorphism because both variants were associated with lower tIgE concentrations in this Barbados population. On the basis of the above results, we used a recessive model for both SNPs and accounted for dependence within families. The CC and CT genotypes were combined into a comparison group for both polymorphisms (ie, subjects with CC/CT at AOAH rs2727831 and CC/CT at *CD14*[-260]C>T were the reference



**FIG 2.** FBAT results showing global  $-\log(P)$  against chromosomal location for asthma (A),  $\log[\text{tIgE}]$  concentrations (B),  $\log[\text{IL-13}]/\log[\text{IFN-}\gamma]$  (C), and sCD14 (D). Two-point and haplotype results for 2-, 3-, and 4-marker windows across the AOA gene are shown. Black vertical lines represent all individual SNP tests ( $n = 28$ ), and colored horizontal lines represent haplotype tests.

group). Fig 3 depicts odds ratios (ORs) and  $P$  values for asthma, contrasting various genotypic groups with this group, and suggests that the presence of the TT genotype at either loci (but not both) reduced the risk for asthma (ORs of  $<1.0$ ) and conferred lower tIgE levels (data not shown) compared with those in the reference group, but the presence of the TT genotype at both loci increased risk (ORs  $>1$ ), suggesting the presence of a qualitative interaction. When modeling interactions, the interaction coefficient was significant for both asthma (OR for interaction = 5.28,  $P = .036$ ) and tIgE ( $P = .006$ ). When sCD14 or  $\log[\text{IL-13}]/\log[\text{IFN-}\gamma]$  was examined, there was no significant evidence of interaction.

## DISCUSSION

This is the first report of association between variants in AOA and human allergic disease. We have presented evidence for linkage and association among 8 markers in AOA and the phenotypes of asthma,  $\log[\text{tIgE}]$ , sCD14, and  $\log[\text{IL-13}]/\log[\text{IFN-}\gamma]$ . The most compelling signal was for 2 SNPs flanking exon 13 and low tIgE concentrations.


The precise relationship between AOA and the  $T_H2/T_H1$  lymphocyte-cytokine milieu in the context of asthma and allergic disease remains unclear. The first evidence for

*in vivo* function for AOA was recently presented by Lu et al,<sup>22</sup> who showed that deacylation of LPS affected longer-term immune responses, specifically B-cell proliferation and polyclonal antibody production in response to gram-negative infection in mice. AOA is an attractive candidate gene for asthma because it is localized on chromosome 7p14-p12, where evidence for linkage to asthma, bronchial hyperreactivity, and tIgE concentrations has been reported from separate genome screens.<sup>24-26</sup> Interestingly, linkage has also been observed at this locus for other inflammatory diseases, including multiple sclerosis,<sup>39</sup> autoimmune thyroiditis,<sup>40</sup> inflammatory bowel disease,<sup>41</sup> and systemic lupus erythematosus.<sup>42</sup> In unpublished gene-profiling studies to test for the effects of the *CD14* ( $-260$ )C>T genotype on AOA expression, we compared cDNA expression from PBMCs of asthmatic subjects stratified by genotype (CC, CT, or TT) at *CD14* ( $-260$ ). AOA was consistently differentially expressed, and by using real-time RT-PCR, we validated these findings and determined that the expression of AOA was significantly higher among asthmatic subjects with *CD14* ( $-260$ ) CC and CT genotypes compared with those with the TT genotype (1.46- to 1.48-fold increase, 1-sided  $P$  value = .045-.014; unpublished data). These findings suggested that AOA might play an important role in the *CD14*-MD-2-TLR4 signaling pathway.

LPS recognition is essential for effective host defense, and the primary mechanism is through the CD14–MD-2–TLR4 signaling complex, which in turn initiates an inflammatory response by producing  $T_H1$  cytokines, chemokines, and other mediators.<sup>43</sup> Soluble CD14 further enhances LPS-stimulated responses by mediating the response of cells lacking membrane-bound CD14.<sup>44</sup> The agonistic effect of sCD14 on endothelial cells<sup>45</sup> has been attributed to overactivation of nonspecific immune response to LPS, which underlies conditions such as septic shock.<sup>46,47</sup> Alternatively, sCD14 is recognized as being important in neutralizing LPS and minimizing the damaging effects of endotoxin-induced cell activation.<sup>48-50</sup> Deacylation of LPS by AOAHS diminishes stimulation of cells through this signaling complex,<sup>19</sup> and *in vitro* murine studies in renal cortical tubule cells suggest that deacylation by AOAHS limits inflammatory reactions to gram-negative bacteria.<sup>51</sup> Both sCD14 and AOAHS seem to function for a common end point: to deactivate LPS and regulate the inflammatory cascade associated with endotoxin-mediated host response.

Because the *CD14*(-260C>T) promoter substitution results in decreased affinity of Sp protein binding and enhanced transcriptional activity,<sup>14</sup> there has been a great deal of focus on the effects of this variant on high sCD14 levels.<sup>4,6,52-55</sup> Soluble CD14 has been shown to be correlated with development of recurrent wheezing in the first year of life and other markers of atopy, such as IFN- $\gamma$ .<sup>56</sup> Previously, we did not observe any association between sCD14 concentrations and *CD14*(-260)C>T<sup>29</sup>; however, when we tested for association between variants in AOAHS and sCD14 concentration, we found several AOAHS variants/haplotypes were associated with low sCD14 levels. We also observed a significant association between a marker of the  $T_H2$  phenotype (IL-13/IFN- $\gamma$  ratio) and several AOAHS markers, suggesting that the AOAHS gene plays a role in the  $T_H2$  milieu in addition to its effect on sCD14 levels. We speculate that AOAHS activities are downstream of CD14 in the LPS signaling pathway, given that the conversion of LPS to deacylated LPS (resulting in approximately 0.2% to 1% as much activity) might take many hours or even days,<sup>22,57-59</sup> whereas CD14 initiates a rapid response that normally begins within minutes of LPS exposure. Thus it is possible that abnormalities in AOAHS expression might contribute to persistent inflammation in diseases such as asthma.

Taken together, these studies suggest an interaction between *CD14* and AOAHS. An evaluation of distributions of AOAHS and *CD14* genotypes among asthmatic subjects raises the possibility of gene-gene interaction or epistasis (Fig 3). These observed *CD14* and AOAHS markers might combine to control sCD14 levels, which in turn would affect the  $T_H1/T_H2$  environment (eg, tIgE and cytokine concentrations). Consequently, increased concentrations of sCD14 associated with *CD14*(-260) TT status could cause lower AOAHS expression, either through reducing LPS-induced activation through competitive binding between sCD14 and membrane-bound CD14<sup>60,61</sup> or by enhancing the rate of LPS efflux (eg, transferring cell-



**FIG 3.** Interaction of *CD14*(C-260T) and AOAHS rs2727831(C/T) genotypes for asthma. Bars indicate the ORs between the different combinations of genotypes for asthma. Under the recessive model, the CC and CT genotypes were collapsed, and the CC/CT group at both genes was used as the baseline comparison group.

bound LPS to plasma proteins).<sup>50</sup> In this family sample from Barbados, both the AOAHS CT haplotype and the single marker rs2727831 variant influenced sCD14, suggesting that high sCD14 concentration is conferred, in part, by a wild-type AOAHS allele. Alternatively, an epistatic mechanism for these 2 candidate genes might be through their effect on tIgE concentration. Homozygotes for the variant at both *CD14*(-260) and rs2727831 had lower tIgE concentrations than carriers of wild-type alleles. Alterations in AOAHS might affect deacylation and subsequently result in sustained concentrations of circulating serum LPS, which in turn could modulate IgE regulation by favoring  $T_H1$  differentiation, suppressing  $T_H2$ -dependent IgE responses, or both. Direct effects of sCD14 on tIgE concentrations have also been described in which sCD14 inhibits IgE production by interfering with CD40 signaling in B cells, increasing CD40 ligand expression on T cells, and inhibiting IL-2, IL-4, and IL-6.<sup>62,63</sup> Our sample is unique in that very high levels of circulating sCD14 were observed among predominantly asthmatic subjects (mean, 6710  $\pm$  3243 ng/mL), but there was no inverse relationship between sCD14 and tIgE concentrations ( $r = -0.0201$ ,  $P = .7988$ ). It is possible that this sample lacked sufficient power to detect any association between this variant and sCD14 because of low frequency (8.2%). Another explanation is environmental: we previously reported exceptionally high household concentrations of endotoxin in these Caribbean families enriched for allergic disease.<sup>29</sup> Because increased sCD14 concentrations have been reported for *in vitro* models of chronic exposure to trace amounts of endotoxin,<sup>64</sup> domestic exposure could account for the high sCD14 concentrations seen here. Similarly, AOAHS activity can be induced after LPS exposure in murine dendritic cells<sup>59</sup> and macrophages.<sup>65</sup> Perhaps only subjects with either or both of these variants would have increased tIgE concentrations.

In the current study we examined 28 SNPs throughout the AOA gene, including 4 SNPs within approximately 5 kb of each other in intron 12, flanking exon 13. It is possible that one or more causal variants lie within or flank the TA/CT haplotype for markers rs2727833 and rs2727831 ( $P = .006$ ), which showed the strongest association with serum tIgE levels (Fig 2). AOA is a highly polymorphic gene (211 kb), with large introns ( $n = 20$ ; mean size, 10.5 kb), relatively small exons ( $n = 21$ ; mean size, 107 bp) and very few SNPs in coding regions (5 synonymous and 5 nonsynonymous), raising the possibility of splicing variants playing a major role. These 28 AOA markers averaged more than 11 kb apart, distances not typically found with strong LD. Conservative estimates suggest more than 150 haplotype-tagging AOA SNPs would be necessary to fully cover AOA in a population of African descent. Given the limited coverage of markers across AOA, plus the fact that none of these variants are known to be functional, it is premature to draw conclusions about the functional relevance of the associations reported here. Analysis of AOA expression from RNA in subjects of different AOA genotypes did not support the hypothesis that these AOA variants affected transcriptional levels of AOA in peripheral blood, despite our previous unpublished studies showing significant differences in AOA expression between TT and C genotypes at the functional *CD14*(-260) variant. After fine mapping of AOA in this dataset and refinement of the linkage and association loci, appropriate future functional studies might be conducted.

In summary, *CD14* and AOA are 2 innate immunity candidate genes that might control risk to allergic diseases and quantitative traits associated with asthma in an interactive manner. Replication of these findings in other populations of African descent, as well as in non-African samples, will be needed to confirm these associations. With more extensive genotyping, causal variants might ultimately be identified.

We thank all the families and volunteers who graciously agreed to participate in this study. We are especially grateful to Mrs Pissamai Maul, Mr Trevor Maul, Dr Kathryn Held, Ms Sharon Patterson, Caitlin Dorer, and Kimberly Donnelly for field work and to Drs Harold Watson, Paul Levett, and Raana Naidu for facilitating the Barbados study. We also thank Robert Munford for helpful discussions and Roxann Ashworth, Gordon Spangler, Pat Oldewurtel, and Lianhe Shao for technical assistance.

## REFERENCES

- Braun-Fahrlander C, Riedler J, Herz U, Eder W, Waser M, Grize L, et al. Environmental exposure to endotoxin and its relation to asthma in school-age children. *N Engl J Med* 2002;347:869-77.
- Liu AH. Endotoxin exposure in allergy and asthma: reconciling a paradox. *J Allergy Clin Immunol* 2002;109:379-92.
- Wright SD. CD14 and innate recognition of bacteria. *J Immunol* 1995; 155:6-8.
- Baldini M, Lohman IC, Halonen M, Erickson RP, Holt PG, Martinez FD. A polymorphism\* in the 5' flanking region of the CD14 gene is associated with circulating soluble CD14 levels and with total serum immunoglobulin E. *Am J Respir Cell Mol Biol* 1999;20:976-83.
- Gao PS, Mao XQ, Baldini M, Roberts MH, Adra CN, Shirakawa T, et al. Serum total IgE levels and CD14 on chromosome 5q31. *Clin Genet* 1999;56:164-5.
- Leung TF, Tang NL, Sung YM, Li AM, Wong GW, Chan IH, et al. The C-159T polymorphism in the CD14 promoter is associated with serum total IgE concentration in atopic Chinese children. *Pediatr Allergy Immunol* 2003;14:255-60.
- Koppelman GH, Reijmerink NE, Colin Stine O, Howard TD, Whittaker PA, Meyers DA, et al. Association of a promoter polymorphism of the CD14 gene and atopy. *Am J Respir Crit Care Med* 2001;163:965-9.
- Buckova D, Holla LI, Schuller M, Znojil V, Vacha J. Two CD14 promoter polymorphisms and atopic phenotypes in Czech patients with IgE-mediated allergy. *Allergy* 2003;58:1023-6.
- Choudhry S, Avila PC, Nazario S, Ung N, Kho J, Rodriguez-Santana JR, et al. CD14 tobacco gene-environment interaction modifies asthma severity and immunoglobulin E levels in Latinos with asthma. *Am J Respir Crit Care Med* 2005;172:173-82.
- Sackesen C, Karaaslan C, Keskin O, Tokol N, Tahan F, Civelek E, et al. The effect of polymorphisms at the CD14 promoter and the TLR4 gene on asthma phenotypes in Turkish children with asthma. *Allergy* 2005;60: 1485-92.
- Woo JG, Assa'ad A, Heizer AB, Bernstein JA, Hershey GK. The -159 C>T polymorphism of CD14 is associated with nonatopic asthma and food allergy. *J Allergy Clin Immunol* 2003;112:438-44.
- Sharma M, Batra J, Mabalirajan U, Goswami S, Ganguly D, Lahkar B, et al. Suggestive evidence of association of C-159T functional polymorphism of the CD14 gene with atopic asthma in northern and northwestern Indian populations. *Immunogenetics* 2004;56:544-7.
- Martin AC, Laing IA, Khoo SK, Zhang G, Rueter K, Teoh L, et al. Acute asthma in children: relationship between CD14 and CC16 genotype, plasma levels and severity. *Am J Respir Crit Care Med* 2005;15:617-22.
- LeVan TD, Bloom JW, Bailey TJ, Karp CL, Halonen M, Martinez FD, et al. A common single nucleotide polymorphism in the CD14 promoter decreases the affinity of Sp protein binding and enhances transcriptional activity. *J Immunol* 2001;167:5838-44.
- Staab JF, Fosmire S, Zhang M, Varley AW, Munford RS. Distinctive structural features are shared by human, lapine, and murine acyloxyacyl hydrolases. *J Endotoxin Res* 1999;5:205-8.
- Hall CL, Munford RS. Enzymatic deacylation of the lipid A moiety of *Salmonella typhimurium* lipopolysaccharides by human neutrophils. *Proc Natl Acad Sci U S A* 1983;80:6671-5.
- Munford RS, Hall CL. Purification of acyloxyacyl hydrolase, a leukocyte enzyme that removes secondary acyl chains from bacterial lipopolysaccharides. *J Biol Chem* 1989;264:15613-9.
- Hagen FS, Grant FJ, Kuijper JL, Slaughter CA, Moomaw CR, Orth K, et al. Expression and characterization of recombinant human acyloxyacyl hydrolase, a leukocyte enzyme that deacylates bacterial lipopolysaccharides. *Biochemistry* 1991;30:8415-23.
- Munford RS, Hall CL. Detoxification of bacterial lipopolysaccharides (endotoxins) by a human neutrophil enzyme. *Science* 1986;234:203-5.
- Pohlman TH, Munford RS, Harlan JM. Deacylated lipopolysaccharide inhibits neutrophil adherence to endothelium induced by lipopolysaccharide in vitro. *J Exp Med* 1987;165:1393-402.
- Kitchens RL, Ulevitch RJ, Munford RS. Lipopolysaccharide (LPS) partial structures inhibit responses to LPS in a human macrophage cell line without inhibiting LPS uptake by a CD14-mediated pathway. *J Exp Med* 1992;176:485-94.
- Lu M, Zhang M, Takashima A, Weiss J, Apicella MA, Li XH, et al. Lipopolysaccharide deacylation by an endogenous lipase controls innate antibody responses to Gram-negative bacteria. *Nat Immunol* 2005;6: 989-94.
- Whitmore TE, Mathewes SL, O'Hara PJ, Durnam DM. Chromosomal localization of the acyloxyacyl hydrolase (AOAH) gene to 7p14-p12 using fluorescence in situ hybridization. *Genomics* 1994;21:457-8.
- Daniels SE, Bhattacharya S, James A, Leaves NI, Young A, Hill MR, et al. A genome-wide search for quantitative trait loci underlying asthma. *Nature* 1996;383:247-50.
- Dizier MH, Besse-Schmittler C, Guilloud-Bataille M, Annesi-Maesano I, Boussaha M, Bousquet J, et al. Genome screen for asthma and related phenotypes in the French EGEA study. *Am J Respir Crit Care Med* 2000;162:1812-8.

26. Laitinen T, Daly MJ, Rioux JD, Kauppi P, Laprise C, Petays T, et al. A susceptibility locus for asthma-related traits on chromosome 7 revealed by genome-wide scan in a founder population. *Nat Genet* 2001; 28:87-91.
27. Barnes KC, Neely JD, Duffy DL, Freidhoff LR, Breazeale DR, Schou C, et al. Linkage of asthma and total serum IgE concentration to markers on chromosome 12q: evidence from Afro-Caribbean and Caucasian populations. *Genomics* 1996;37:41-50.
28. Barnes KC, Freidhoff LR, Nickel R, Chiu Y-F, Juo S-H, Hizawa N, et al. Dense mapping of chromosome 12q13.12-q23.3 and linkage to asthma and atopy. *J Allergy Clin Immunol* 1999;104:485-91.
29. Zambelli-Weiner A, Ehrlich A, Stockton ML, Grant AV, Zhang S, Levett PN, et al. Evaluation of the CD14/-260 polymorphism and house dust endotoxin exposure in the Barbados asthma genetics study. *J Allergy Clin Immunol* 2005;115:1203-9.
30. Livak KJ, Schmittgen TD. Analysis of relative gene expression data using real-time quantitative PCR and the 2(-Delta Delta C(T)) Method. *Methods* 2001;25:402-8.
31. Zeger SL, Liang KY. Longitudinal data analysis for discrete and continuous outcomes. *Biometrics* 1986;42:121-30.
32. Williams RL. A note on robust variance estimation for cluster-correlated data. *Biometrics* 2000;56:645-6.
33. Horvath S, Wei E, Xu X, Palmer LJ, Bauer M. Family-based association test method: age of onset traits and covariates. *Genet Epidemiol* 2001; 21(suppl 1):S403-8.
34. Rabinowitz D, Laird N. A unified approach to adjusting association tests for population admixture with arbitrary pedigree structure and arbitrary missing marker information. *Hum Hered* 2000;50:211-23.
35. Heinzmann A, Dietrich H, Jerkic SP, Kurz T, Deichmann KA. Promoter polymorphisms of the CD14 gene are not associated with bronchial asthma in Caucasian children. *Eur J Immunogenet* 2003;30:345-8.
36. Sengler C, Haider A, Sommerfeld C, Lau S, Baldini M, Martinez F, et al. Evaluation of the CD14 C-159 T polymorphism in the German Multicenter Allergy Study cohort. *Clin Exp Allergy* 2003;33:166-9.
37. Barnes KC, Grant AV, Baltadzhieva D, Zhang S, Berg T, Shao L, et al. Variants in the gene encoding C3 are associated with asthma and related phenotypes among African Caribbean families. *Genes Immun* 2006;7: 27-35.
38. Gabriel SB, Schaffner SF, Nguyen H, Moore JM, Roy J, Blumenstiel B, et al. The structure of haplotype blocks in the human genome. *Science* 2002;296:2225-9.
39. Sawcer S, Jones HB, Feakes R, Gray J, Smaldon N, Chataway J, et al. A genome screen in multiple sclerosis reveals susceptibility loci on chromosome 6p21 and 17q22. *Nat Genet* 1996;13:464-8.
40. Allen EM, Hsueh WC, Sabra MM, Pollin TI, Ladenson PW, Silver KD, et al. A genome-wide scan for autoimmune thyroiditis in the Old Order Amish: replication of genetic linkage on chromosome 5q11.2-q14.3. *J Clin Endocrinol Metab* 2003;88:1292-6.
41. Satsangi J, Parkes M, Louis E, Hashimoto L, Kato N, Welsh K, et al. Two stage genome-wide search in inflammatory bowel disease provides evidence for susceptibility loci on chromosomes 3, 7 and 12. *Nat Genet* 1996;14:199-202.
42. Gaffney PM, Ortmann WA, Selby SA, Shark KB, Ockenden TC, Rohlf KE, et al. Genome screening in human systemic lupus erythematosus: results from a second Minnesota cohort and combined analyses of 187 sib-pair families. *Am J Hum Genet* 2000;66:547-56.
43. Medzhitov R, Janeway C Jr. Innate immunity. *N Engl J Med* 2000;343: 338-44.
44. Frey EA, Miller DS, Jahr TG, Sundan A, Bazil V, Espevik T, et al. Soluble CD14 participates in the response of cells to lipopolysaccharide. *J Exp Med* 1992;176:1665-71.
45. Arditi M, Zhou J, Dorio R, Rong GW, Goyert SM, Kim KS. Endotoxin-mediated endothelial cell injury and activation: role of soluble CD14. *Infect Immun* 1993;61:3149-56.
46. Kruger C, Schutt C, Obertacke U, Joka T, Muller FE, Knoller J, et al. Serum CD14 levels in polytraumatized and severely burned patients. *Clin Exp Immunol* 1991;85:297-301.
47. Landmann R, Reber AM, Sansano S, Zimmerli W. Function of soluble CD14 in serum from patients with septic shock. *J Infect Dis* 1996;173: 661-8.
48. Grunwald U, Kruger C, Schutt C. Endotoxin-neutralizing capacity of soluble CD14 is a highly conserved specific function. *Circ Shock* 1993;39: 220-5.
49. Wurfel MM, Hailman E, Wright SD. Soluble CD14 acts as a shuttle in the neutralization of lipopolysaccharide (LPS) by LPS-binding protein and reconstituted high density lipoprotein. *J Exp Med* 1995;181:1743-54.
50. Kitchens RL, Thompson PA, Viriyakosol S, O'Keefe GE, Munford RS. Plasma CD14 decreases monocyte responses to LPS by transferring cell-bound LPS to plasma lipoproteins. *J Clin Invest* 2001;108:485-93.
51. Feulner JA, Lu M, Shelton JM, Zhang M, Richardson JA, Munford RS. Identification of acyloxyacyl hydrolase, a lipopolysaccharide-detoxifying enzyme, in the murine urinary tract. *Infect Immun* 2004;72:3171-8.
52. Eng HL, Wang CH, Chen CH, Chou MH, Cheng CT, Lin TM. A CD14 promoter polymorphism is associated with CD14 expression and *Chlamydia*-stimulated TNF alpha production. *Genes Immun* 2004;5:426-30.
53. Kabesch M, Hasemann K, Schickinger V, Tzotcheva I, Bohnert A, Carr D, et al. A promoter polymorphism in the CD14 gene is associated with elevated levels of soluble CD14 but not with IgE or atopic diseases. *Allergy* 2004;59:520-5.
54. Amar J, Ruidavets JB, Bal dit Sollier C, Bongard V, Boccalon H, Chamontin B, et al. CD14 C(-260)T gene polymorphism, circulating soluble CD14 levels and arteriosclerosis. *J Hypertens* 2004;22:1523-8.
55. Shimada K, Miyauchi K, Mokuno H, Watanabe Y, Iwama Y, Shigekiyo M, et al. Promoter polymorphism in the CD14 gene and concentration of soluble CD14 in patients with in-stent restenosis after elective coronary stenting. *Int J Cardiol* 2004;94:87-92.
56. Guerra S, Lohman IC, Halonen M, Martinez FD, Wright AL. Reduced interferon gamma production and soluble CD14 levels in early life predict recurrent wheezing by 1 year of age. *Am J Respir Crit Care Med* 2004;169:70-6.
57. Luchi M, Munford RS. Binding, internalization, and deacylation of bacterial lipopolysaccharide by human neutrophils. *J Immunol* 1993;151:959-69.
58. Katz SS, Weinrauch Y, Munford RS, Elsbach P, Weiss J. Deacylation of lipopolysaccharide in whole *Escherichia coli* during destruction by cellular and extracellular components of a rabbit peritoneal inflammatory exudate. *J Biol Chem* 1999;274:36579-84.
59. Lu M, Zhang M, Kitchens RL, Fosmire S, Takashima A, Munford RS. Stimulus-dependent deacylation of bacterial lipopolysaccharide by dendritic cells. *J Exp Med* 2003;197:1745-54.
60. Schutt C, Schilling T, Grunwald U, Schonfeld W, Kruger C. Endotoxin-neutralizing capacity of soluble CD14. *Res Immunol* 1992;143:71-8.
61. Haziot A, Rong GW, Lin XY, Silver J, Goyert SM. Recombinant soluble CD14 prevents mortality in mice treated with endotoxin (lipopolysaccharide). *J Immunol* 1995;154:6529-32.
62. Rey Nores JE, Bensusan A, Vita N, Stelter F, Arias MA, Jones M, et al. Soluble CD14 acts as a negative regulator of human T cell activation and function. *Eur J Immunol* 1999;29:265-76.
63. Arias MA, Rey Nores JE, Vita N, Stelter F, Borysiewicz LK, Ferrara P, et al. Cutting edge: human B cell function is regulated by interaction with soluble CD14: opposite effects on IgG1 and IgE production. *J Immunol* 2000;164:3480-5.
64. Nockher WA, Scherberich JE. Monocyte cell-surface CD14 expression and soluble CD14 antigen in hemodialysis: evidence for chronic exposure to LPS. *Kidney Int* 1995;48:1469-76.
65. Cody MJ, Salkowski CA, Henricson BE, Detore GR, Munford RS, Vogel SN. Effect of inflammatory and anti-inflammatory stimuli on acyloxyacyl hydrolase gene expression and enzymatic activity in murine macrophages. *J Endotoxin Res* 1997;4:371-9.