

UNITED STATES DEPARTMENT OF THE INTERIOR, Oscar L. Chapman, *Secretary*
FISH AND WILDLIFE SERVICE, Albert M. Day, *Director*

STUDIES OF GEORGES BANK HADDOCK

Part I: Landings by Pounds, Numbers, and Sizes of Fish

BY HOWARD A. SCHUCK



FISHERY BULLETIN 66
From Fishery Bulletin of the Fish and Wildlife Service
VOLUME 52

UNITED STATES GOVERNMENT PRINTING OFFICE • WASHINGTON : 1951

For sale by the Superintendent of Documents, U. S. Government Printing Office, Washington 25, D. C.
Price 20 cents

ABSTRACT

The haddock has been New England's most valuable fishery resource for nearly three decades. After 1929, production declined markedly and as a consequence a study of the resources was begun to determine what caused the decline, what could be done to maintain or increase production, and what prediction of future landings might be possible. Presented in this first paper of a series reporting the results of this study is information on pounds, numbers, average weights, and sizes of haddock landed from Georges Bank for the years 1931 to 1948, and information on trends and seasonal cycles in the landings.

CONTENTS

| | Page |
|--|------|
| Introduction..... | 151 |
| Fishing banks and areas studied..... | 152 |
| Organization of study..... | 153 |
| Development of data..... | 153 |
| Ports of landing..... | 153 |
| Categories of fish..... | 156 |
| Seasons..... | 157 |
| Segregating landings by subareas..... | 157 |
| Pounds of haddock landed..... | 157 |
| Average weights of haddock landed..... | 159 |
| Numbers of haddock landed..... | 162 |
| Size compositions of haddock landed..... | 162 |
| Scrod haddock..... | 164 |
| Large haddock..... | 164 |
| Total haddock..... | 164 |
| Undersized haddock..... | 166 |
| Scrod versus large haddock..... | 166 |
| Discussion and summary..... | 174 |
| Literature cited..... | 176 |

STUDIES OF GEORGES BANK HADDOCK

Part I: Landings by Pounds, Numbers, and Sizes of Fish

By HOWARD A. SCHUCK, *Fishery Research Biologist*

The haddock, *Melanogrammus aeglefinus*, has been New England's most valuable fishery resource, and one of the most important in the United States, for nearly three decades. In the early days, this fish was little sought and the annual New England catch was small—only about 40-odd million pounds until well into the 1900's. With the development of filleting and freezing methods the market for haddock grew, and during the 1920's New England landings increased greatly. They reached a peak of about 250 million pounds in 1929, but after that production declined rapidly.

From Georges Bank, source of most United States haddock, production dropped from about 223 million pounds in 1929 to 115 million pounds in 1931. In addition, an index of abundance indicates that the size of the stock on Georges Bank declined greatly over these years.

The decline of haddock landings and abundance aroused concern in the fishing industry, and in 1930 funds were made available to the United States Bureau of Fisheries (now the Fish and Wildlife Service) to study the haddock and the haddock fishery. The general purposes of the investigation were to determine (1) what caused the decline of the fishery in waters fished by United States fishermen, (2) what could be done to increase abundance and production, or at least to prevent them from decreasing further, and (3) what predictions of future production were possible.

During the years 1931-48, a large quantity of data was collected, partly at sea but mostly at the important haddock ports (Boston, Gloucester, and New Bedford, Mass., and Portland, Maine) where collectors and interviewers have worked systematically since 1931. These data, the basis of this and other papers, were obtained with the cooperation of fishermen at sea and of boat owners, dealers, and fish handlers—especially those on the Boston Fish Pier (fig. 1).

William C. Herrington, in charge of the Haddock Investigation from 1931 to 1947, planned the collection of these data obtained in various years during the period 1931-48 by many employees of the Fish and Wildlife Service. Among these were H. M. Bearse, F. E. Firth, D. F. Hammack, J. J. Miggins, J. M. Shuval, and J. R. Webster. Assisting in tabulating and summarizing data at various times during the years 1945-49 were E. L. Arnold, Jr., F. A. Dreyer, Dorothy B. Monahan, Elizabeth V. Nugent, E. S. Phillips, S. L. Cogswell, and L. D. Stringer.

At sea, data were collected on commercial fishing vessels; on the *Atlantis*, a research vessel leased from the Woods Hole Oceanographic Institution; and on the fishery-research vessels *Albatross II* (1931 and 1932) and *Albatross III* (beginning in 1948). Most of these data were collected to determine how to protect small haddock, destroyed in large numbers by the otter-trawl (fig. 2) fleet. Line trawlers (fig. 3) were used in the early days of the haddock fishery, but now only two are operating out of Boston, Mass., the major haddock port. Results of these studies on the small haddock situation were reported by Herrington (1933, 1935, 1936, 1941).¹ In addition, a small amount of tagging was done to determine migrations and interdependence of populations. Most of this work remains unreported, but one publication refers to phases of it (Rounsefell 1942). And since the commissioning of the *Albatross III* in 1948, further experiments on mesh sizes, studies of survival of young haddock that escape through larger mesh, some tagging, and a census of the population of all ages of haddock have been undertaken.

At the important haddock ports considerable quantities of data were obtained. These data are largely unreported, although contributions of Herrington (1944, 1948) and Schuck (1949) have presented segments of them and certain conclu-

¹ Publications referred to parenthetically by date are listed in the Literature Cited, p. 176.




FIGURE 1.—Part of the Boston Fish Pier, where most of the United States production of haddock is landed.




FIGURE 2.—Modern otter trawler: predominant type of vessel in the present-day New England haddock fishery.




FIGURE 3.—Line trawler: predominant type of vessel in the early years of the New England haddock fishery.

sions regarding the fishery. At the ports, since 1931, the following data have been collected: (1) Almost complete records of the poundages landed from various banks, with records of depths and locations from which the fish were taken, the gear used, and the days actually spent fishing; (2) randomized samples of the lengths of fish in the landings; (3) selected samples of scales; and (4) length-weight data.

FISHING BANKS AND AREAS STUDIED

The United States haddock fishery has depended upon Georges Bank and the Nova Scotian banks. To the north of these banks, haddock are found, but are little fished by United States fishermen. To the south, haddock are not found, except for stragglers.

Georges Bank is the most important area for the United States haddock fishery, with about 67 percent of the total United States landings coming from this area during recent years (1931 to 1948).

The haddock on Georges Bank are apparently a relatively distinct and homogeneous stock. Present knowledge indicates that the Fundian Channel, which separates Georges Bank from the Nova Scotian banks, is a natural barrier to the intermigration of bottom-living stages of haddock. Evidence of this comes from studies of size compositions, growth rates, tagging, and vertebral counts. The size composition of the stock and the sizes of haddock of various ages on Georges Bank are decidedly different from those on Browns Bank across the Fundian Channel (Needler 1930, Schuck and Arnold in press). Although the number of tagged haddock is not large, there is no evidence from the returns that any of them crossed this channel (Needler 1930, Schroeder 1942, United States Fish and Wildlife Service unpublished data). There is a seasonal migration in the spring from Georges Bank north along the coast of Massachusetts and Maine as far as the Bay of Fundy and a return to Georges Bank in the fall, but very few haddock are caught on this northward migration.

Because, first, the Georges Bank area was the most important for the United States haddock fishery and, second, the haddock on Georges Bank formed a relatively distinct population and, third, haddock production from this bank had declined more seriously than production from the Nova Scotian banks, we decided to study first the

Georges Bank haddock—before the Nova Scotian haddock.

The Georges Bank region comprises most of International Area XXII, shown in figure 4. International Area XXII was established by the North American Council on Fishery Investigation when the western North Atlantic Ocean was divided along natural, political, and ecological lines. By Georges Bank we mean specifically the following subareas (fig. 5) of Area XXII:

| | <i>International subarea</i> |
|--|----------------------------------|
| 1. Northern Edge and Northeast Peak..... | J |
| 2. Southeast Part of Georges..... | M |
| 3. Southwest Georges..... | N |
| 4. South Channel and Nantucket Shoals..... | G, H, O ¹ |

¹ Data include very small quantities from subareas Q, R, and S.

The manner by which these subareas were established is described by Rounsefell (1948).

ORGANIZATION OF STUDY

Russell (1942) has expressed the dynamics of a fish population by the equation

$$S_1 + (G + R) - (C + N) = S_2$$

where

S_1 = size of population at the beginning of the year,

G = additions to the population during the year by growth,

R = additions to the population by recruitment of young fish,

C = deductions from the population during the year by fishery,

N = deductions from the population during the year due to natural mortality,

S_2 = size of population at the end of the year.

The main problems, as we see them, are (1) to obtain accurate measures of the various quantities expressed in this equation for each year, (2) to determine what effect variations of catch, natural mortality, growth, and recruitment have had on the size of the stock, (3) to determine what effect variations in the size of the stock have had upon each of these factors, and (4) to show what effect other factors in the environment (hydrographic conditions and stocks of other species of competing fishes) have had upon (a) the size of the stock and (b) the four factors—catch, growth, recruitment, and natural mortality.

With this information at hand, if the relative effects of the fishery and of the environment on

the stock are sufficiently clear, it should be possible (1) to predict the abundance and production of haddock, and (2) to determine what measures, if any, would maintain or increase the catch of haddock from the important populations.

Most of the material in this series is devoted to solving these problems. The purpose of the remainder of the present paper is restricted to determining the total landings of Georges Bank haddock for each season and year, 1931 to 1948, in terms of pounds, numbers, average weights, and numbers of each size.

Obtaining "total" values implies adding together not only those portions of the landings of the various ports that originated on Georges Bank, but adding together also data for two artificial market categories, the limits of which vary from season to season, from year to year, and among different areas of the bank.

Where we refer to totals we refer, of course, to our best estimate of such values. All such values are subject to a certain amount of error due to limitations in collecting and assembling statistics and to sampling error.

The values developed in this paper represent landings but not catches because the smallest sizes of haddock are discarded at sea as they lack sufficient marketable value to be brought to port.

DEVELOPMENT OF DATA

Ports of landing

Haddock are caught in North American waters by fishermen from New England, New York, Canada, Newfoundland, and various European countries.

Canadian and Newfoundland landings were excluded from this study, as no records could be found to indicate that any of their haddock were caught in the Georges Bank area. McKenzie (1946) has shown that all Canadian haddock landings for the years 1938 to 1940 came from banks to the north and east of Georges Bank. Herrington (unpublished manuscript) lists all Canadian landings for the years 1918 to 1940 as having originated from banks other than Georges.

European fishermen, mainly interested in cod, frequent the Newfoundland banks and the most easterly of the Nova Scotian banks. Records show that Europeans fished on Georges Bank during early years, but not during the years covered in this summary.




FIGURE 4.—International statistical areas off the Atlantic coast of North America.




Figure 5.—Subareas currently used in International Areas XXI and XXII.

Thus United States fishermen were the only ones to land haddock from Georges Bank. However, we could not use the total of all United States landings of haddock for this study because United States fishermen took varying quantities of haddock from other banks as well as from Georges.

Inasmuch as Georges Bank lies at a considerable distance off shore, it is exploited mainly by large vessels. These vessels land at only a few ports where, for the most part, accurate records have been kept on the origin of haddock landings. Thus for Boston and Gloucester we were able to determine the quantities of haddock landed from Georges Bank each year. We included also in our tabulations the quantities of Georges Bank haddock landed at Portland, Maine, during the years 1931 to 1946. And beginning in 1942, landings of haddock at the port of New Bedford became quite large, so the New Bedford landings of Georges Bank haddock for the years 1942 to 1948 were included. As almost all haddock landed at New York City are taken from the Georges area, the total of that port's landings for all years also were included. We included also the total landings for Groton, Conn. for 1931 and 1932—landings at this port were negligible after 1932. To these quantities, we added the entire amount of haddock landed on Cape Cod, which lies next to Georges Bank. This is the only area where small boats land Georges Bank haddock, and almost all landings there are from Georges.

The sums of these quantities we have accepted as the total poundages² of haddock originating in the Georges area that were landed and sold.

Categories of fish

Immediately after capture at sea, haddock are separated into two market categories, scrod and large. This division of the catch makes it necessary to collect complete data on each market category and later to combine the data to obtain total statistics for the species haddock.

As defined by the New England Fish Exchange, scrod haddock (scrod) are those weighing from 1½

to 2½ pounds (gutted weight), and large haddock are those weighing more than 2½ pounds. These definitions are only approximate owing to variations in culling and to a practice of marketing, as scrod, many fish weighing less than 1½ pounds.

We have tabulated records of the landings for both market categories, large and scrod, for all years. Small amounts of "mixed" haddock were added to scrod in New Bedford. When OPA price control regulations were in effect (which allowed a higher price for "large" haddock), New Bedford landings showed an artificial scarcity of scrod and an overabundance of large. For the period July 1943 to June 1946, therefore, we used the percentage that scrod made up of the monthly total of scrod and large for the ports of Boston, Gloucester, and Portland, from any subarea in any month, to estimate the proportion of scrod in the New Bedford landings from these same subareas in that month.

Where we refer to "undersized" haddock we mean those less than 1½ pounds, the lower limit of the market category of scrod, although at present there is no State or Federal regulation that classifies such fish as undersized. When we refer to "total haddock" or merely "haddock", we mean the total of all haddock regardless of market category.

Most haddock are landed as drawn or gutted fish, but some are landed in the "round". Where poundages of fish in the round were obtained, they were reduced by 15 percent. Thus all poundages are in terms of gutted fish.

Landings of large haddock in the round were negligible but landings of round scrod were more numerous and were of two types, (1) regular-sized scrod that were left ungutted because of rough weather or gluts of fish on deck, and (2) unusually small-sized scrod, or baby scrod. Landings of baby scrod became unusually large in the winter of 1940, owing to a scarcity of large haddock and a high abundance of baby haddock (year class 1939).

The landings of baby scrod from the winter of 1940 to the summer of 1943 were considered to be so large that in the initial steps of the analysis they were treated separately from scrod or large haddock. These landings of baby scrod amounted to approximately the following:

² Sources of data are the former U. S. Bureau of Fisheries and the present U. S. Fish and Wildlife Service publications, "Current Fishery Statistics" for all years, and unpublished records of various fish companies assembled by William C. Herrington.

| | <i>Thousands of pounds</i> |
|-------------|--------------------------------|
| Year 1940: | |
| Fall..... | 33 |
| Winter..... | 1, 097 |
| Year 1941: | |
| Spring..... | 3, 153 |
| Summer..... | 1, 683 |
| Fall..... | 913 |
| Winter..... | 339 |
| Year 1942: | |
| Spring..... | 239 |
| Summer..... | 380 |
| Fall..... | 275 |
| Winter..... | 362 |
| Year 1943: | |
| Spring..... | 2, 212 |
| Summer..... | 429 |
| Fall..... | 25 |

Seasons

A "haddock year" is the summation of spring, summer, fall, and winter seasons, and differs from a calendar year by one month. These seasons are as follows:

| | <i>Months</i> |
|-------------|--|
| Spring..... | February, March, April. |
| Summer..... | May, June, July. |
| Fall..... | August, September, October. |
| Winter..... | November, December, January (of following year). |

These seasons agree with the Georges Bank haddock life-cycle better than any other 3-month grouping, for the months of February, March, and April constitute the spawning period. During these months the size and age composition of the catch is considerably different from that of each of the other seasons.

All data were collected initially on a monthly basis, then assembled into seasons, and then into haddock years.

Segregating landings by subareas

Inasmuch as different sizes of haddock are caught on various parts of Georges Bank, we wished in the initial steps of development of the data to segregate the landings by subareas. For the ports of Boston, Gloucester, New Bedford, and Portland, accurate information was obtained on the amounts of haddock landed from each subarea. These ports received the bulk of the total landings (88 percent for all years), thus we allotted the remainder of the landings to subareas

on the basis of the subarea contribution at these ports.

The subareas shown in figure 5 were in use from 1939 through 1948. In the years before 1939, there were several different systems of naming and segregating the various sections of Georges Bank. The data from earlier years, therefore, were arranged to conform, as much as possible, to the modern subareas. One exception should be noted, however. During the years 1931 through 1935, published statistics furnished a breakdown by only (1) South Channel and Nantucket Shoals, and (2) the rest of Georges Bank proper—roughly J, M, and N of the modern terminology.

In all tables showing pounds and numbers of fish, values were rounded off to the nearest thousand. Total as well as individual values were rounded off. Thus, individual values do not add up exactly to the totals in some cases.

POUNDS OF HADDOCK LANDED

Table 1 shows the pounds of scrod and large haddock landed from the four subareas of Georges Bank by seasons and years, from 1931 through 1948. Whether particular subareas of Georges Bank contributed more or less haddock in recent years can be studied through this table. Their importance, relative to one another, is shown in table 2 (percent contribution by years, 1936-48 only). The landings are summarized, by seasons, for scrod in table 3, for large in table 4, and for total haddock in table 5. Landings by years only are shown also in tables 3, 4, and 5, and in figure 6.




FIGURE 6.—Pounds of scrod, large, and total haddock landed from Georges Bank, 1931 to 1948.

TABLE 3.—*Scrod haddock landed, by seasons and years*
[In thousands of pounds]

| Year | Spring | Summer | Fall | Winter | Total |
|---------|--------|---------|---------|--------|---------|
| 1931 | 894 | 1,255 | 5,266 | 10,086 | 17,501 |
| 1932 | 6,955 | 8,737 | 11,799 | 4,022 | 31,513 |
| 1933 | 4,116 | 4,520 | 8,631 | 1,531 | 18,798 |
| 1934 | 1,605 | 3,488 | 6,038 | 1,845 | 12,976 |
| 1935 | 963 | 5,717 | 10,554 | 8,303 | 25,537 |
| 1936 | 3,872 | 9,604 | 12,933 | 3,541 | 29,950 |
| 1937 | 5,514 | 8,423 | 14,665 | 2,482 | 31,084 |
| 1938 | 4,307 | 7,982 | 20,414 | 7,204 | 39,907 |
| 1939 | 7,524 | 11,743 | 17,716 | 6,142 | 43,125 |
| 1940 | 6,614 | 9,393 | 9,507 | 4,086 | 29,600 |
| 1941 | 11,614 | 13,578 | 21,066 | 7,506 | 53,764 |
| 1942 | 11,484 | 17,026 | 13,757 | 7,213 | 49,480 |
| 1943 | 11,618 | 13,907 | 10,032 | 1,950 | 37,507 |
| 1944 | 3,978 | 5,485 | 4,822 | 963 | 15,248 |
| 1945 | 1,040 | 2,983 | 5,441 | 2,210 | 11,674 |
| 1946 | 1,009 | 7,215 | 9,147 | 3,749 | 21,120 |
| 1947 | 5,637 | 8,337 | 20,873 | 6,058 | 40,905 |
| 1948 | 6,176 | 12,669 | 15,982 | 9,729 | 44,556 |
| Total | 94,920 | 152,062 | 218,643 | 88,620 | 554,245 |
| Average | 5,273 | 8,448 | 12,147 | 4,923 | 30,791 |

AVERAGE WEIGHTS OF HADDOCK
LANDED

Average weights of fish landed, in each season, year, subarea, and market category, were computed by combining length samples of haddock landed with seasonal length-weight relations. This procedure is described in the following paragraphs.

At the Boston Fish Pier, lengths of representative samples of the haddock landed were obtained from 1931 through 1948. In general, 50 scrod and 100 large haddock were measured from a "trip" when a vessel had fished in only one subarea of Georges Bank, and as many vessels were sampled as time permitted.

Each fish was measured from the tip of the snout to the fork of the tail. Lengths were recorded by centimeter groups, that is, fish measuring from 40.0 centimeters to and including 40.9 centimeters were recorded as 40 centimeters, fish from 41.0 centimeters to and including 41.9 centimeters as 41 centimeters, and so on. No distinction as to sex was possible as most haddock, when landed, are already dressed.

The numbers of Georges Bank haddock that were measured, by years, seasons, and market categories are shown in table 6. In all, measurements of 627,996 haddock from Georges Bank were utilized in this analysis.

Table 7 illustrates the general method used to compute the average weight of haddock landed. The steps of this method are as follows: (1) The number of fish of each centimeter size group in the total sample for the season was entered in column II; (2) the length-weight relation was available by seasons (table 8 and figure 7) and the average weights for each centimeter size group were listed in column III, the total weight of all fish measured of each centimeter size group was computed in column IV, and the total weight of *all sizes* in the season's sample was entered at the bottom of column IV; and finally (3) the total weight of the sample was divided by the number of fish in the sample to give the average weight of the fish in the sample. We used this same general method for each season, year, subarea, and market category.

Summaries of average weights are given in table 9 and figure 8; to save space, values for the various subareas are not shown.

TABLE 4.—*Large haddock landed, by seasons and years*
[In thousands of pounds]

| Year | Spring | Summer | Fall | Winter | Total |
|---------|---------|---------|---------|---------|-----------|
| 1931 | 29,611 | 33,610 | 23,827 | 10,491 | 97,539 |
| 1932 | 18,136 | 19,534 | 23,303 | 12,934 | 73,907 |
| 1933 | 18,367 | 20,715 | 19,495 | 4,265 | 62,842 |
| 1934 | 7,261 | 13,953 | 12,780 | 2,914 | 36,908 |
| 1935 | 5,166 | 18,623 | 18,041 | 11,814 | 53,644 |
| 1936 | 13,828 | 17,218 | 16,359 | 6,663 | 54,068 |
| 1937 | 19,705 | 17,431 | 17,647 | 9,588 | 64,371 |
| 1938 | 15,283 | 15,637 | 12,834 | 8,726 | 52,480 |
| 1939 | 15,811 | 18,118 | 17,376 | 10,105 | 61,410 |
| 1940 | 15,763 | 22,204 | 16,490 | 8,588 | 63,045 |
| 1941 | 19,674 | 23,808 | 15,961 | 8,519 | 67,962 |
| 1942 | 16,870 | 20,300 | 12,916 | 7,525 | 57,611 |
| 1943 | 14,202 | 17,779 | 15,385 | 4,711 | 52,077 |
| 1944 | 16,310 | 27,942 | 24,648 | 11,844 | 80,744 |
| 1945 | 14,643 | 20,319 | 19,260 | 12,375 | 66,597 |
| 1946 | 13,049 | 27,825 | 28,603 | 13,289 | 82,766 |
| 1947 | 19,693 | 19,190 | 17,668 | 7,809 | 64,360 |
| 1948 | 12,810 | 12,798 | 14,147 | 9,212 | 48,967 |
| Total | 286,182 | 367,004 | 326,740 | 161,372 | 1,141,298 |
| Average | 15,899 | 20,389 | 18,152 | 8,965 | 63,405 |

TABLE 5.—*Total haddock landed, by seasons and years*
[In thousands of pounds]

| Year | Spring | Summer | Fall | Winter | Total |
|---------|---------|---------|---------|---------|-----------|
| 1931 | 30,505 | 34,865 | 29,093 | 20,577 | 115,040 |
| 1932 | 25,091 | 28,271 | 35,102 | 16,956 | 105,420 |
| 1933 | 22,483 | 25,235 | 28,126 | 5,796 | 81,640 |
| 1934 | 8,866 | 17,441 | 18,818 | 4,759 | 49,884 |
| 1935 | 6,129 | 24,340 | 28,595 | 20,117 | 79,181 |
| 1936 | 17,700 | 26,822 | 29,242 | 10,204 | 84,018 |
| 1937 | 25,219 | 25,864 | 32,312 | 12,070 | 95,455 |
| 1938 | 19,680 | 23,619 | 35,248 | 15,930 | 92,387 |
| 1939 | 23,335 | 29,861 | 35,092 | 16,247 | 104,535 |
| 1940 | 22,377 | 31,597 | 25,997 | 12,674 | 92,645 |
| 1941 | 31,288 | 37,386 | 37,027 | 16,025 | 121,726 |
| 1942 | 28,354 | 37,326 | 26,673 | 14,738 | 107,091 |
| 1943 | 25,820 | 31,686 | 25,417 | 6,661 | 89,584 |
| 1944 | 20,288 | 33,427 | 29,470 | 12,807 | 95,992 |
| 1945 | 15,683 | 23,302 | 24,701 | 14,585 | 78,271 |
| 1946 | 14,058 | 35,040 | 37,750 | 17,038 | 103,886 |
| 1947 | 25,330 | 27,527 | 38,541 | 13,867 | 105,265 |
| 1948 | 18,986 | 25,467 | 30,129 | 18,941 | 93,523 |
| Total | 381,102 | 519,066 | 545,383 | 249,992 | 1,695,543 |
| Average | 21,172 | 28,837 | 30,299 | 13,888 | 94,196 |




FIGURE 7.—Relation between length and weight for Georges Bank haddock, by seasons.




FIGURE 8.—Average weights of scrod, large, and total haddock landed from Georges Bank, by years.

TABLE 6.—Numbers of haddock measured for length, by seasons and years

| Season | Scrod | Large | Total |
|-------------------|---------------|---------------|---------------|
| Year 1931: | | | |
| Spring..... | 513 | 5,042 | 5,555 |
| Summer..... | 1,194 | 4,054 | 5,248 |
| Fall..... | 3,285 | 4,577 | 7,862 |
| Winter..... | 4,102 | 2,582 | 6,684 |
| Total..... | 9,094 | 16,235 | 25,329 |
| Year 1932: | | | |
| Spring..... | 2,913 | 3,484 | 6,397 |
| Summer..... | 2,445 | 6,245 | 8,690 |
| Fall..... | 4,849 | 8,558 | 13,407 |
| Winter..... | 3,741 | 3,662 | 7,403 |
| Total..... | 13,948 | 21,949 | 35,897 |
| Year 1933: | | | |
| Spring..... | 3,082 | 3,834 | 6,914 |
| Summer..... | 1,702 | 3,775 | 5,477 |
| Fall..... | 2,455 | 5,349 | 7,804 |
| Winter..... | 911 | 2,157 | 3,068 |
| Total..... | 18,150 | 15,115 | 23,265 |
| Year 1934: | | | |
| Spring..... | 675 | 3,326 | 4,001 |
| Summer..... | 2,014 | 3,341 | 5,355 |
| Fall..... | 2,588 | 3,924 | 6,512 |
| Winter..... | 2,691 | 1,831 | 4,522 |
| Total..... | 7,968 | 12,422 | 20,390 |
| Year 1935: | | | |
| Spring..... | 1,440 | 3,398 | 4,838 |
| Summer..... | 4,582 | 7,357 | 11,939 |
| Fall..... | 7,199 | 6,462 | 13,661 |
| Winter..... | 3,318 | 2,981 | 6,299 |
| Total..... | 16,539 | 20,198 | 36,737 |
| Year 1936: | | | |
| Spring..... | 3,643 | 6,914 | 10,557 |
| Summer..... | 9,533 | 11,089 | 20,622 |
| Fall..... | 9,740 | 9,997 | 19,737 |
| Winter..... | 3,849 | 5,595 | 9,444 |
| Total..... | 26,765 | 33,595 | 60,360 |
| Year 1937: | | | |
| Spring..... | 3,383 | 8,781 | 12,164 |
| Summer..... | 5,394 | 8,777 | 14,171 |
| Fall..... | 5,129 | 5,296 | 10,425 |
| Winter..... | 4,055 | 5,387 | 9,442 |
| Total..... | 17,961 | 28,241 | 46,202 |
| Year 1938: | | | |
| Spring..... | 4,419 | 7,574 | 11,993 |
| Summer..... | 4,592 | 6,520 | 11,112 |
| Fall..... | 5,250 | 4,668 | 9,918 |
| Winter..... | 3,860 | 3,716 | 7,576 |
| Total..... | 18,121 | 22,478 | 40,599 |
| Year 1939: | | | |
| Spring..... | 2,540 | 4,002 | 6,542 |
| Summer..... | 5,244 | 6,835 | 12,079 |
| Fall..... | 4,448 | 7,712 | 12,160 |
| Winter..... | 3,043 | 4,141 | 7,184 |
| Total..... | 15,275 | 22,690 | 37,965 |

TABLE 6.—Numbers of haddock measured for length, by seasons and years—Continued

| Season | Scrod | Large | Total |
|-------------|----------|----------|----------|
| Year 1940: | | | |
| Spring..... | 4, 219 | 9, 324 | 13, 543 |
| Summer..... | 4, 085 | 8, 588 | 12, 674 |
| Fall..... | 3, 356 | 4, 784 | 8, 140 |
| Winter..... | 4, 501 | 4, 379 | 8, 880 |
| Total..... | 16, 162 | 27, 075 | 43, 237 |
| Year 1941: | | | |
| Spring..... | 6, 080 | 8, 145 | 14, 225 |
| Summer..... | 5, 287 | 6, 069 | 11, 356 |
| Fall..... | 8, 167 | 6, 179 | 14, 346 |
| Winter..... | 4, 853 | 3, 334 | 8, 187 |
| Total..... | 24, 387 | 23, 727 | 48, 114 |
| Year 1942: | | | |
| Spring..... | 4, 516 | 6, 380 | 10, 896 |
| Summer..... | 7, 163 | 8, 453 | 15, 616 |
| Fall..... | 6, 247 | 6, 186 | 12, 433 |
| Winter..... | 3, 933 | 4, 345 | 8, 278 |
| Total..... | 21, 559 | 25, 364 | 47, 223 |
| Year 1943: | | | |
| Spring..... | 6, 082 | 6, 644 | 12, 726 |
| Summer..... | 4, 796 | 4, 834 | 9, 630 |
| Fall..... | 3, 237 | 6, 420 | 9, 657 |
| Winter..... | 644 | 2, 304 | 2, 948 |
| Total..... | 14, 759 | 20, 202 | 34, 961 |
| Year 1944: | | | |
| Spring..... | 1, 471 | 3, 295 | 4, 766 |
| Summer..... | 1, 532 | 5, 183 | 6, 715 |
| Fall..... | 1, 984 | 5, 262 | 7, 246 |
| Winter..... | 200 | 1, 890 | 2, 090 |
| Total..... | 5, 187 | 15, 630 | 20, 817 |
| Year 1945: | | | |
| Spring..... | 250 | 1, 644 | 1, 894 |
| Summer..... | 649 | 1, 797 | 2, 446 |
| Fall..... | 950 | 3, 150 | 4, 100 |
| Winter..... | 699 | 3, 266 | 3, 965 |
| Total..... | 2, 548 | 9, 857 | 12, 405 |
| Year 1946: | | | |
| Spring..... | 750 | 2, 800 | 3, 550 |
| Summer..... | 2, 600 | 6, 147 | 8, 747 |
| Fall..... | 3, 250 | 6, 660 | 9, 910 |
| Winter..... | 2, 234 | 3, 387 | 5, 621 |
| Total..... | 8, 834 | 18, 994 | 27, 828 |
| Year 1947: | | | |
| Spring..... | 2, 230 | 3, 651 | 5, 881 |
| Summer..... | 2, 037 | 2, 870 | 4, 907 |
| Fall..... | 3, 776 | 7, 861 | 11, 637 |
| Winter..... | 3, 205 | 4, 468 | 7, 673 |
| Total..... | 11, 248 | 18, 850 | 30, 098 |
| Year 1948: | | | |
| Spring..... | 3, 507 | 4, 181 | 7, 688 |
| Summer..... | 3, 480 | 2, 217 | 5, 697 |
| Fall..... | 7, 101 | 7, 417 | 14, 518 |
| Winter..... | 4, 763 | 3, 903 | 8, 666 |
| Total..... | 18, 851 | 17, 718 | 36, 569 |
| All years: | | | |
| Spring..... | 51, 713 | 92, 419 | 144, 132 |
| Summer..... | 68, 330 | 104, 151 | 172, 481 |
| Fall..... | 83, 011 | 110, 462 | 193, 473 |
| Winter..... | 54, 602 | 63, 308 | 117, 910 |
| Total..... | 257, 656 | 370, 340 | 627, 996 |

TABLE 7.—Method used to compute average weight of haddock

Example used: 1948, Spring, Southeast Part, Scrod

| Length group ¹ | Number in sample | Average weight | Total weight of sample |
|---------------------------|------------------|----------------|------------------------|
| (I) | (II) | (III) | (IV) |
| | | Pounds | Pounds |
| 29 cm..... | 1 | 0.58 | 0.58 |
| 30 cm..... | 5 | .64 | 3.20 |
| 31 cm..... | 11 | .70 | 7.70 |
| 32 cm..... | 17 | .76 | 12.92 |
| 33 cm..... | 29 | .83 | 24.07 |
| 34 cm..... | 36 | .90 | 32.40 |
| 35 cm..... | 40 | .98 | 39.20 |
| 36 cm..... | 44 | 1.06 | 46.64 |
| 37 cm..... | 45 | 1.14 | 51.30 |
| 38 cm..... | 41 | 1.23 | 50.43 |
| 39 cm..... | 31 | 1.32 | 40.92 |
| 40 cm..... | 53 | 1.4 | 74.2 |
| 41 cm..... | 54 | 1.5 | 81.0 |
| 42 cm..... | 82 | 1.6 | 131.2 |
| 43 cm..... | 133 | 1.7 | 226.1 |
| 44 cm..... | 142 | 1.8 | 255.6 |
| 45 cm..... | 188 | 2.0 | 376.0 |
| 46 cm..... | 188 | 2.1 | 394.8 |
| 47 cm..... | 183 | 2.2 | 402.6 |
| 48 cm..... | 160 | 2.4 | 384.0 |
| 49 cm..... | 160 | 2.5 | 400.0 |
| 50 cm..... | 93 | 2.6 | 241.8 |
| 51 cm..... | 62 | 2.8 | 173.6 |
| 52 cm..... | 38 | 2.9 | 110.2 |
| 53 cm..... | 17 | 3.1 | 52.7 |
| 54 cm..... | 11 | 3.2 | 35.2 |
| 55 cm..... | 6 | 3.4 | 20.4 |
| 56 cm..... | 2 | 3.6 | 7.2 |
| 57 cm..... | | 3.8 | |
| 58 cm..... | | 4.0 | |
| 59 cm..... | 1 | 4.2 | 4.2 |
| Total..... | 1,873 | 1,965 | 3,680.16 |

¹ By 1-cm. intervals.

² 3,680.16 pounds
1,873 fish = 1.965 pounds.

TABLE 8.—Length-weight relation by seasons, in terms of centimeter size groups and drawn weight in pounds

| Length ¹ | Drawn weight in pounds | | | |
|---------------------|------------------------|--------|------|--------|
| | Spring | Summer | Fall | Winter |
| 18 cm..... | 0.15 | 0.12 | 0.15 | 0.14 |
| 19 cm..... | .17 | .14 | .17 | .16 |
| 20 cm..... | .20 | .17 | .20 | .19 |
| 21 cm..... | .23 | .20 | .23 | .21 |
| 22 cm..... | .27 | .23 | .26 | .25 |
| 23 cm..... | .30 | .26 | .30 | .28 |
| 24 cm..... | .34 | .29 | .33 | .32 |
| 25 cm..... | .38 | .33 | .38 | .36 |
| 26 cm..... | .43 | .36 | .42 | .40 |
| 27 cm..... | .47 | .41 | .47 | .45 |
| 28 cm..... | .52 | .45 | .52 | .50 |
| 29 cm..... | .58 | .50 | .57 | .55 |
| 30 cm..... | .64 | .55 | .63 | .61 |
| 31 cm..... | .70 | .60 | .69 | .67 |
| 32 cm..... | .76 | .66 | .75 | .73 |
| 33 cm..... | .83 | .72 | .82 | .80 |
| 34 cm..... | .90 | .79 | .89 | .88 |
| 35 cm..... | .98 | .85 | .96 | .95 |
| 36 cm..... | 1.06 | .92 | 1.05 | 1.04 |
| 37 cm..... | 1.14 | 1.00 | 1.13 | 1.12 |
| 38 cm..... | 1.23 | 1.08 | 1.22 | 1.21 |
| 39 cm..... | 1.32 | 1.16 | 1.31 | 1.31 |
| 40 cm..... | 1.4 | 1.2 | 1.4 | 1.4 |
| 41 cm..... | 1.5 | 1.3 | 1.5 | 1.5 |

See footnote at end of table.

TABLE 8.—Length-weight relation by seasons, in terms of centimeter size groups and drawn weight in pounds—Con.

| Length ¹ | Drawn weight in pounds | | | |
|---------------------|------------------------|--------|------|--------|
| | Spring | Summer | Fall | Winter |
| 42 cm | 1.6 | 1.4 | 1.6 | 1.6 |
| 43 cm | 1.7 | 1.5 | 1.7 | 1.7 |
| 44 cm | 1.8 | 1.6 | 1.8 | 1.8 |
| 45 cm | 2.0 | 1.7 | 2.0 | 2.0 |
| 46 cm | 2.1 | 1.8 | 2.1 | 2.1 |
| 47 cm | 2.2 | 2.0 | 2.2 | 2.2 |
| 48 cm | 2.4 | 2.1 | 2.3 | 2.4 |
| 49 cm | 2.5 | 2.2 | 2.5 | 2.5 |
| 50 cm | 2.6 | 2.3 | 2.6 | 2.7 |
| 51 cm | 2.8 | 2.5 | 2.8 | 2.8 |
| 52 cm | 2.9 | 2.6 | 2.9 | 3.0 |
| 53 cm | 3.1 | 2.8 | 3.1 | 3.2 |
| 54 cm | 3.2 | 2.9 | 3.2 | 3.4 |
| 55 cm | 3.4 | 3.1 | 3.4 | 3.5 |
| 56 cm | 3.6 | 3.2 | 3.6 | 3.7 |
| 57 cm | 3.8 | 3.4 | 3.8 | 3.9 |
| 58 cm | 4.0 | 3.6 | 4.0 | 4.1 |
| 59 cm | 4.2 | 3.8 | 4.1 | 4.3 |
| 60 cm | 4.4 | 3.9 | 4.3 | 4.5 |
| 61 cm | 4.6 | 4.1 | 4.5 | 4.8 |
| 62 cm | 4.8 | 4.3 | 4.8 | 5.0 |
| 63 cm | 5.0 | 4.5 | 5.0 | 5.2 |
| 64 cm | 5.2 | 4.7 | 5.2 | 5.5 |
| 65 cm | 5.4 | 4.9 | 5.4 | 5.7 |
| 66 cm | 5.7 | 5.1 | 5.6 | 6.0 |

See footnote at end of table.

TABLE 8.—Length-weight relation by seasons, in terms of centimeter size groups and drawn weight in pounds—Con.

| Length ¹ | Drawn weight in pounds | | | |
|---------------------|------------------------|--------|------|--------|
| | Spring | Summer | Fall | Winter |
| 67 cm | 5.9 | 5.4 | 5.9 | 6.2 |
| 68 cm | 6.2 | 5.6 | 6.1 | 6.5 |
| 69 cm | 6.4 | 5.8 | 6.4 | 6.8 |
| 70 cm | 6.7 | 6.1 | 6.7 | 7.1 |
| 71 cm | 7.0 | 6.3 | 6.9 | 7.4 |
| 72 cm | 7.2 | 6.6 | 7.2 | 7.7 |
| 73 cm | 7.5 | 6.8 | 7.5 | 8.0 |
| 74 cm | 7.8 | 7.1 | 7.8 | 8.3 |
| 75 cm | 8.1 | 7.4 | 8.1 | 8.7 |
| 76 cm | 8.4 | 7.7 | 8.4 | 9.0 |
| 77 cm | 8.7 | 7.9 | 8.7 | 9.4 |
| 78 cm | 9.0 | 8.2 | 9.0 | 9.7 |
| 79 cm | 9.3 | 8.6 | 9.3 | 10.1 |
| 80 cm | 9.7 | 8.9 | 9.6 | 10.4 |
| 81 cm | 10.0 | 9.2 | 10.0 | 10.8 |
| 82 cm | 10.3 | 9.5 | 10.3 | 11.1 |
| 83 cm | 10.6 | 9.8 | 10.6 | 11.5 |
| 84 cm | 10.9 | 10.1 | 10.9 | 11.8 |
| 85 cm | 11.4 | 10.3 | 11.4 | 12.3 |
| 86 cm | 11.7 | 10.7 | 11.7 | 12.7 |
| 87 cm | 12.2 | 11.2 | 12.2 | 13.1 |
| 88 cm | 12.6 | 11.5 | 12.6 | 13.6 |
| 89 cm | 12.9 | 11.8 | 12.9 | 14.1 |
| 90 cm | 13.5 | 12.3 | 13.5 | 14.6 |

¹ Size groups by 1-cm. intervals.

TABLE 9.—Average weights in pounds of scrod, large, and total haddock, by seasons and years

| Year | Scrod | | | | | Large | | | | | Total | | | | |
|------------------|--------|--------|-------|--------|-------|--------|--------|-------|--------|-------|--------|--------|-------|--------|-------|
| | Spring | Summer | Fall | Winter | Total | Spring | Summer | Fall | Winter | Total | Spring | Summer | Fall | Winter | Total |
| 1931 | 1.517 | 1.540 | 1.653 | 1.541 | 1.585 | 3.648 | 3.112 | 3.866 | 3.490 | 3.473 | 3.543 | 3.079 | 3.112 | 2.154 | 2.940 |
| 1932 | 1.654 | 1.679 | 1.942 | 1.921 | 1.793 | 3.732 | 3.350 | 3.154 | 3.322 | 3.374 | 2.769 | 2.562 | 2.621 | 2.832 | 2.670 |
| 1933 | 1.938 | 1.248 | 1.714 | 1.636 | 1.604 | 3.607 | 3.062 | 3.171 | 3.639 | 3.277 | 3.116 | 2.429 | 2.515 | 2.750 | 2.643 |
| 1934 | 1.890 | 1.648 | 1.614 | 1.402 | 1.617 | 3.580 | 3.126 | 3.271 | 3.195 | 3.263 | 3.082 | 2.650 | 2.460 | 2.136 | 2.580 |
| 1935 | 1.574 | 1.594 | 1.705 | 1.624 | 1.658 | 3.706 | 3.014 | 3.044 | 3.476 | 3.174 | 3.212 | 2.492 | 2.360 | 2.363 | 2.451 |
| 1936 | 1.905 | 1.456 | 1.710 | 1.586 | 1.626 | 3.602 | 3.009 | 3.025 | 3.343 | 3.187 | 3.014 | 2.177 | 2.267 | 2.415 | 2.374 |
| 1937 | 1.950 | 1.528 | 1.820 | 1.793 | 1.748 | 3.580 | 3.160 | 3.289 | 4.051 | 3.432 | 3.027 | 2.344 | 2.407 | 3.218 | 2.613 |
| 1938 | 1.972 | 1.510 | 1.709 | 1.656 | 1.679 | 4.902 | 3.199 | 3.348 | 4.051 | 3.716 | 3.692 | 2.322 | 2.107 | 2.401 | 2.438 |
| 1939 | 1.890 | 1.633 | 1.718 | 1.674 | 1.715 | 3.955 | 3.083 | 2.933 | 3.492 | 3.285 | 2.925 | 2.285 | 2.161 | 2.475 | 2.384 |
| 1940 | 1.989 | 1.748 | 1.867 | 1.558 | 1.803 | 3.434 | 3.218 | 3.357 | 3.998 | 3.399 | 2.827 | 2.575 | 2.598 | 2.656 | 2.650 |
| 1941 | 1.688 | 1.541 | 1.681 | 1.823 | 1.662 | 3.991 | 3.330 | 3.377 | 4.030 | 3.592 | 2.650 | 2.342 | 2.145 | 2.572 | 2.375 |
| 1942 | 2.012 | 1.690 | 1.701 | 1.742 | 1.766 | 3.644 | 3.195 | 3.121 | 3.536 | 3.340 | 2.743 | 2.272 | 2.182 | 2.351 | 2.366 |
| 1943 | 1.924 | 1.586 | 1.837 | 1.809 | 1.757 | 3.495 | 3.138 | 3.306 | 2.774 | 3.239 | 2.555 | 2.195 | 2.513 | 2.400 | 2.393 |
| 1944 | 1.926 | 1.617 | 1.800 | 2.049 | 1.772 | 3.464 | 3.031 | 3.231 | 3.965 | 3.290 | 2.995 | 2.651 | 2.859 | 3.705 | 2.896 |
| 1945 | 1.940 | 1.296 | 1.644 | 1.736 | 1.573 | 3.678 | 3.157 | 3.548 | 3.766 | 3.481 | 3.472 | 2.667 | 2.827 | 3.199 | 2.948 |
| 1946 | 1.665 | 1.449 | 1.770 | 1.778 | 1.642 | 3.630 | 3.077 | 3.406 | 3.824 | 3.377 | 3.346 | 2.499 | 2.782 | 3.052 | 2.780 |
| 1947 | 1.876 | 1.481 | 1.580 | 2.291 | 1.670 | 3.725 | 3.635 | 3.622 | 4.194 | 3.719 | 3.055 | 2.524 | 2.130 | 3.077 | 2.519 |
| 1948 | 1.842 | 1.493 | 1.651 | 1.592 | 1.623 | 3.959 | 3.251 | 3.472 | 3.743 | 3.572 | 2.882 | 2.050 | 2.218 | 2.209 | 2.272 |
| Weighted average | 1.871 | 1.556 | 1.719 | 1.697 | 1.691 | 3.718 | 3.163 | 3.306 | 3.661 | 3.398 | 2.984 | 2.430 | 2.413 | 2.596 | 2.554 |

NUMBERS OF HADDOCK LANDED

Dividing poundage by average weight gave the number of fish landed—for each season, subarea, market category, and year. Excepting subarea values, all of these numbers are shown in the following tables.

Tables 10, 11, and 12 show the numbers of scrod, large, and total haddock landed, by seasons and years. Relative contributions of scrod and large haddock to the total, by seasons, are shown in figure 9. Figure 10 shows the yearly trends, and here it can be seen that much of the variation in total landings by years is due to variations in scrod landings. The importance of

these small-sized haddock to the present fishery is thus evident.

SIZE COMPOSITIONS OF HADDOCK LANDED

Now having available the number of haddock that were landed (in each season, year, subarea, and market category), and having also the lengths of samples of haddock (in each similar subdivision), we estimated how many haddock of each size were landed. This was accomplished by multiplying the number of fish measured in each centimeter size group by the proportion of the number landed to the number measured. This

calculation assumes that the fish measured were representative samples of the landings. Precautions had been taken to avoid bias in sampling, and many uniformity trials showed that the samples could be considered as representative of the landing.




FIGURE 9.—Numbers of scrod, large, and total haddock landed from Georges Bank in the average year, by seasons.




FIGURE 10.—Numbers of scrod, large, and total haddock landed from Georges Bank, by years.

TABLE 10.—Numbers of scrod haddock landed, by seasons and years

[In thousands of fish]

| Year | Spring | Summer | Fall | Winter | Total |
|---------|--------|--------|---------|--------|---------|
| 1931 | 492 | 816 | 3,186 | 6,547 | 11,041 |
| 1932 | 4,204 | 5,206 | 6,075 | 2,094 | 17,579 |
| 1933 | 2,124 | 3,623 | 5,035 | 936 | 11,718 |
| 1934 | 849 | 2,117 | 3,742 | 1,316 | 8,024 |
| 1935 | 514 | 3,587 | 6,190 | 5,113 | 15,404 |
| 1936 | 2,033 | 6,598 | 7,561 | 2,232 | 18,424 |
| 1937 | 2,828 | 5,512 | 8,056 | 1,384 | 17,780 |
| 1938 | 2,193 | 5,285 | 11,945 | 4,350 | 23,773 |
| 1939 | 3,980 | 7,190 | 10,313 | 3,670 | 25,153 |
| 1940 | 3,325 | 5,373 | 5,093 | 2,623 | 16,414 |
| 1941 | 6,879 | 8,811 | 12,535 | 4,117 | 32,342 |
| 1942 | 5,708 | 10,077 | 8,088 | 4,140 | 28,013 |
| 1943 | 6,040 | 8,771 | 5,460 | 1,078 | 21,349 |
| 1944 | 2,065 | 3,393 | 2,679 | 470 | 8,607 |
| 1945 | 536 | 2,301 | 3,310 | 1,273 | 7,420 |
| 1946 | 606 | 4,978 | 5,169 | 2,108 | 12,861 |
| 1947 | 3,004 | 5,628 | 13,213 | 2,644 | 24,489 |
| 1948 | 3,352 | 8,484 | 9,510 | 6,113 | 27,459 |
| Total | 50,732 | 97,750 | 127,160 | 52,208 | 327,850 |
| Average | 2,818 | 5,431 | 7,065 | 2,900 | 18,214 |

TABLE 11.—Numbers of large haddock landed, by seasons and years

[In thousands of fish]

| Year | Spring | Summer | Fall | Winter | Total |
|---------|--------|---------|--------|--------|---------|
| 1931 | 8,117 | 10,799 | 6,164 | 3,006 | 28,086 |
| 1932 | 4,859 | 5,831 | 7,318 | 3,894 | 21,902 |
| 1933 | 5,092 | 6,765 | 6,147 | 1,172 | 19,176 |
| 1934 | 2,028 | 4,464 | 3,907 | 912 | 11,311 |
| 1935 | 1,394 | 6,179 | 5,927 | 3,309 | 16,809 |
| 1936 | 3,839 | 5,723 | 5,408 | 1,993 | 16,963 |
| 1937 | 5,504 | 5,517 | 5,366 | 2,367 | 18,754 |
| 1938 | 3,118 | 4,888 | 3,833 | 2,285 | 14,124 |
| 1939 | 3,998 | 5,876 | 5,924 | 2,894 | 18,692 |
| 1940 | 4,590 | 6,899 | 4,912 | 2,148 | 18,540 |
| 1941 | 4,930 | 7,150 | 4,726 | 2,114 | 18,920 |
| 1942 | 4,630 | 6,353 | 4,138 | 2,128 | 17,249 |
| 1943 | 4,064 | 5,665 | 4,653 | 1,698 | 16,080 |
| 1944 | 4,708 | 9,218 | 7,629 | 2,987 | 24,542 |
| 1945 | 3,981 | 6,436 | 5,428 | 3,287 | 19,132 |
| 1946 | 3,595 | 9,043 | 8,399 | 3,475 | 24,512 |
| 1947 | 5,287 | 5,279 | 4,878 | 1,862 | 17,306 |
| 1948 | 3,236 | 3,937 | 4,075 | 2,461 | 13,709 |
| Total | 76,970 | 116,022 | 98,832 | 44,082 | 335,906 |
| Average | 4,276 | 6,445 | 5,491 | 2,449 | 18,661 |

TABLE 12.—Numbers of total haddock landed, by seasons and years

[In thousands of fish]

| Year | Spring | Summer | Fall | Winter | Total |
|---------|---------|---------|---------|--------|---------|
| 1931 | 8,609 | 11,615 | 9,350 | 9,553 | 39,127 |
| 1932 | 9,063 | 11,037 | 13,393 | 5,988 | 39,481 |
| 1933 | 7,216 | 10,388 | 11,182 | 2,108 | 30,894 |
| 1934 | 2,877 | 6,581 | 7,649 | 2,228 | 19,335 |
| 1935 | 1,908 | 9,766 | 12,117 | 8,512 | 32,303 |
| 1936 | 5,872 | 12,321 | 12,969 | 4,225 | 35,387 |
| 1937 | 8,332 | 11,029 | 13,422 | 3,751 | 36,534 |
| 1938 | 5,311 | 10,173 | 15,778 | 6,635 | 37,897 |
| 1939 | 7,978 | 13,066 | 16,237 | 6,564 | 43,845 |
| 1940 | 7,915 | 12,272 | 10,005 | 4,771 | 34,963 |
| 1941 | 11,809 | 15,961 | 17,261 | 6,231 | 51,262 |
| 1942 | 10,338 | 16,430 | 12,226 | 6,268 | 45,262 |
| 1943 | 10,104 | 14,436 | 10,113 | 2,776 | 37,429 |
| 1944 | 6,773 | 12,611 | 10,308 | 3,457 | 33,149 |
| 1945 | 4,517 | 8,737 | 8,738 | 4,560 | 26,552 |
| 1946 | 4,201 | 14,021 | 13,568 | 5,583 | 37,373 |
| 1947 | 8,291 | 10,907 | 18,091 | 4,506 | 41,795 |
| 1948 | 6,588 | 12,421 | 13,585 | 8,574 | 41,168 |
| Total | 127,702 | 213,772 | 225,992 | 96,290 | 663,756 |
| Average | 7,095 | 11,876 | 12,555 | 5,349 | 36,875 |

The size compositions for subareas were combined, and thus we obtained a size composition representing all of Georges Bank, for each season, year, and market category. A certain amount of irregularity in these curves was due to sampling variations, inasmuch as only a limited sample from a very large population of fish had been obtained. To eliminate some of this irregularity we smoothed each distribution by a moving average of three.

Scrod haddock

Tables 13, 14, 15, and 16 show the size compositions³ of the landings of scrod, in each of the 72 seasons, from 1931 through 1948. Table 17 shows the size compositions of scrod by years. Table 18 and figure 11 show the average size compositions of scrod for each season in all of the 18 years, and table 19 shows the size composition of scrod that were landed in the average year, and also the percentage size composition.

Large haddock

Tables 20, 21, 22, and 23 show the size compositions of large haddock in each of the 72 seasons over the 18-year period. Table 24 shows the size composition of large haddock by years. Table 25 and figure 11 show, by seasons, the average size

³ For convenience in handling the large mass of data, we grouped all length frequencies by 3-centimeter groups: Fish of the 29-, 30-, and 31-centimeter groups were recorded as 30 centimeters, fish of the 32-, 33-, and 34-centimeter groups as 33 centimeters, and so on. In graphs and tables where centimeters are shown, they are shown as 30, 33, and 36 rather than 30.5, 33.5, and 36.5 (the true midpoints of the groups) inasmuch as the original centimeter measurements were recorded as 29 when the midpoint was 29.5, 30 instead of 30.5, 31 instead of 31.5, etc. Where inches are shown in graphs, they represent actual values: The inch equivalents opposite 30.5 rather than 30, opposite 31.5 rather than 31, and so on.

The sizes in inches corresponding to the true midpoints of the 3-centimeter groups are as follows:

| 3-centimeter groups: | Inches |
|----------------------|--------|
| 18 cm..... | 7.3 |
| 21 cm..... | 8.5 |
| 24 cm..... | 9.6 |
| 27 cm..... | 10.8 |
| 30 cm..... | 12.0 |
| 33 cm..... | 13.2 |
| 36 cm..... | 14.4 |
| 39 cm..... | 15.6 |
| 42 cm..... | 16.7 |
| 45 cm..... | 17.9 |
| 48 cm..... | 19.1 |
| 51 cm..... | 20.3 |
| 54 cm..... | 21.6 |
| 57 cm..... | 22.6 |
| 60 cm..... | 23.8 |
| 63 cm..... | 25.0 |
| 66 cm..... | 26.2 |
| 69 cm..... | 27.4 |
| 72 cm..... | 28.5 |
| 75 cm..... | 29.7 |
| 78 cm..... | 30.9 |
| 81 cm..... | 32.1 |
| 84 cm..... | 33.3 |
| 87 cm..... | 34.4 |

composition of large haddock that were landed in all 18 years, and table 26 shows the size composition of large haddock that were landed in the average year, and also the percentage size composition.




FIGURE 11.—Size compositions of scrod, large, and total haddock landed from Georges Bank in the average year, by seasons.

Total haddock

Tables 27, 28, 29, and 30, and figures 12a, 12b, and 12c show the size compositions of total haddock (scrod and large combined) in each of the 72 seasons over the 18-year period.

The presence of modes (figures 12a, 12b, and 12c), at slightly increasing sizes of fish in succeeding seasons, suggests that each series of modes may be composed largely of the same year class of haddock. In some instances these year classes (if they are year classes) apparently were the chief source of supply of the fishery for several succeeding seasons, and even for succeeding years.

These modes are more obvious if one season (spring, for example) in a particular year is com-

pared with the average of that season for all years. Figures 13a, 13b, and 13c show such contrasts in terms of deviations from seasonal means.




FIGURE 12a.—Size compositions of total haddock landings from Georges Bank, by seasons and years, 1931 to 1936.




FIGURE 12b.—Size compositions of total haddock landings from Georges Bank, by seasons and years, 1937 to 1942.

Table 31 and figure 14 show the yearly size compositions for total haddock. Table 32 shows the four seasonal size compositions for the average of all 18 years. These values are shown also in figure 11.

In figure 14, it can be seen that there was considerable variation in the relative numbers of various sizes in different years. To study these differences more readily, we plotted (fig. 15) devia-




FIGURE 12c.—Size compositions of total haddock landings from Georges Bank, by seasons and years, 1943 to 1948.




FIGURE 13a.—Deviations from the average size compositions, by seasons, 1931 to 1936.

tions from the average year. Here, it can be seen that a scarcity of small-sized fish characterized some years such as 1931, 1940, 1944, 1945, and 1946. In other years, such as 1943 and 1948, a scarcity of large-sized fish occurred. In still others, an abundance of either small-sized or large-sized haddock occurred, or a scarcity or an abundance of both—the scarce years of 1933, 1934, and 1935, and the abundant year of 1941 demonstrate this. In other years, such as 1937, all sizes were taken in approximately average numbers.

The differences in size composition help to explain how different average weights (shown in table 9) occurred. As one example, the years 1936




FIGURE 13b.—Deviations from the average size compositions, by seasons, 1937 to 1942.




FIGURE 13c.—Deviations from the average size compositions, by seasons, 1943 to 1948.

and 1941 had an identical, low average weight of 2.37 pounds. In 1936, this low average weight was associated with a slight abundance of small-sized and a scarcity of large-sized haddock, while in 1941 it was associated with factors entirely different—an abundance of all sizes, but with small haddock much more abundant than large-sized haddock.

It is obvious that average weight is dependent upon the relative numbers of the various sizes and not upon the actual numbers of fish of various sizes.

In table 33 are shown the size composition of the average year and the percent size composition.

Undersized haddock

The New England Fish Exchange defines scrod haddock as 1½ to 2½ pounds. The average length




FIGURE 14.—Size compositions of total haddock landings from Georges Bank, by years.

of 1½ pound haddock is about 41 centimeters. Thus, most fish up to and including the 39-centimeter size group could be considered as undersized. From table 33, we see that in the average year about 4,974,000 undersized fish were landed, or 13.5 percent of the total. In all years the total number of undersized haddock landed was about 89,513,000. The numbers of undersized haddock that were landed in each year are shown in table 34.

Scrod versus large haddock

Table 35 shows the percentages of each size group that were scrod and large haddock; figure 16 shows the actual size compositions of scrod and large haddock.

The dividing line between scrod and large haddock for the average of the 18-year period was about 48 centimeters. Below 48 centimeters most fish landed were classified as scrod; above 48 most were classified as large haddock.

This dividing line has varied from year to year, owing to differences in relative abundance of fish of difference sizes and to market conditions. Such





FIGURE 15.—Deviations from average size compositions, by years.



FIGURES 16.—Size compositions of scrod and large haddock landings from Georges Bank in average year.

variation made it necessary to measure samples of each category in every year for which we desired an accurate measurement of size composition of the total haddock landings.

The amount of overlap in length between the two market categories has been considerable. For instance, haddock as long as 63 centimeters were occasionally landed as scrod, and fish as small as 36 centimeters were landed as large haddock. This was due to difficulties and mistakes in sorting haddock into two arbitrary categories at sea under varying conditions of weather, haste, and so on.

TABLE 13.—Size compositions of scrod haddock, spring seasons

[In thousands of fish]

| Length ¹ | 1931 | 1932 | 1933 | 1934 | 1935 | 1936 | 1937 | 1938 | 1939 | 1940 | 1941 | 1942 | 1943 | 1944 | 1945 | 1946 | 1947 | 1948 |
|---------------------|------|-------|-------|------|------|-------|-------|-------|-------|-------|-------|-------|-------|-------|------|------|-------|-------|
| 21 cm. | | | | | | | 1 | | | | | | | | | | | |
| 24 cm. | | | | | | | 1 | | | | | | | | | | | |
| 27 cm. | | | | | | | | 2 | | | | | | | | | | |
| 30 cm. | 1 | 7 | | | | 1 | 9 | 30 | | | | 13 | | | | | | |
| 33 cm. | | 31 | 2 | 2 | 1 | 74 | 40 | 113 | 9 | 16 | 108 | 5 | 12 | | | | | |
| 36 cm. | 15 | 292 | 44 | 14 | 13 | 98 | 105 | 173 | 162 | 101 | 1,211 | 125 | 208 | 62 | 2 | 38 | 10 | 2 |
| 39 cm. | 56 | 927 | 172 | 62 | 68 | 152 | 278 | 185 | 460 | 295 | 1,121 | 385 | 774 | 250 | 30 | 97 | 382 | 531 |
| 42 cm. | 160 | 1,464 | 491 | 242 | 144 | 370 | 554 | 314 | 962 | 654 | 819 | 1,117 | 1,536 | 480 | 144 | 198 | 892 | 1,036 |
| 45 cm. | 175 | 1,111 | 828 | 348 | 163 | 650 | 866 | 520 | 1,231 | 989 | 1,184 | 1,948 | 1,667 | 614 | 230 | 188 | 899 | 925 |
| 48 cm. | 68 | 331 | 478 | 156 | 94 | 520 | 726 | 542 | 851 | 911 | 1,249 | 1,530 | 1,165 | 481 | 112 | 57 | 549 | 574 |
| 51 cm. | 14 | 39 | 102 | 24 | 28 | 117 | 215 | 253 | 234 | 311 | 533 | 493 | 441 | 145 | 13 | 5 | 171 | 173 |
| 54 cm. | | 2 | | | 3 | 17 | 29 | 45 | 31 | 43 | 78 | 71 | 116 | 17 | 3 | | 26 | 24 |
| 57 cm. | | | | | | 6 | 4 | 11 | | 5 | 4 | | 36 | 3 | | | | |
| 60 cm. | | | | | | | | 4 | | | | | 18 | | | | | |
| 63 cm. | | | | | | | | 1 | | | | | 8 | | | | | |
| 66 cm. | | | | | | | | | | | | | 3 | | | | | |
| 69 cm. | | | | | | | | | | | | | 1 | | | | | |
| 72 cm. | | | | | | | | | | | | | 1 | | | | | |
| Total | 492 | 4,204 | 2,124 | 849 | 514 | 2,033 | 2,828 | 2,193 | 3,980 | 3,325 | 6,879 | 5,708 | 6,040 | 2,065 | 536 | 606 | 3,004 | 3,352 |

¹ Size groups by 3-cm. intervals.

TABLE 14.—Size compositions of scrod haddock, summer seasons

[In thousands of fish]

| Length ¹ | 1931 | 1932 | 1933 | 1934 | 1935 | 1936 | 1937 | 1938 | 1939 | 1940 | 1941 | 1942 | 1943 | 1944 | 1945 | 1946 | 1947 | 1948 |
|---------------------|------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|--------|-------|-------|-------|-------|-------|-------|
| 21 cm | | | | | | 1 | | | | | | | | | | | | |
| 24 cm | | | | | | 2 | | | | | | | | | | | | |
| 27 cm | | | 14 | | | 5 | | 1 | | | | 2 | | | | | | |
| 30 cm | | | 152 | | | 19 | | 3 | | | | 4 | | | | | | |
| 33 cm | | 1 | 290 | 2 | 1 | 17 | | 16 | 1 | | | 15 | 10 | 24 | 4 | | 29 | |
| 36 cm | | 16 | 373 | 12 | 13 | 128 | 17 | 62 | 25 | 163 | 106 | 214 | 10 | 29 | 210 | 17 | 52 | |
| 39 cm | | 8 | 114 | 524 | 103 | 189 | 635 | 265 | 366 | 345 | 134 | 949 | 531 | 655 | 85 | 333 | 555 | 696 |
| 42 cm | | 74 | 659 | 636 | 225 | 623 | 1,550 | 1,119 | 1,275 | 1,059 | 587 | 2,046 | 1,230 | 1,276 | 442 | 868 | 940 | 1,366 |
| 45 cm | 240 | 1,594 | 592 | 472 | 899 | 1,702 | 1,722 | 1,477 | 1,768 | 1,176 | 1,925 | 1,758 | 2,166 | 996 | 688 | 1,033 | 1,515 | 2,420 |
| 48 cm | 324 | 1,807 | 593 | 750 | 1,028 | 1,365 | 1,343 | 980 | 2,048 | 1,427 | 1,402 | 2,724 | 2,254 | 1,100 | 261 | 1,372 | 1,249 | 2,341 |
| 51 cm | 152 | 842 | 347 | 463 | 656 | 900 | 718 | 769 | 1,502 | 1,416 | 1,491 | 2,706 | 1,575 | 636 | 104 | 714 | 732 | 1,068 |
| 54 cm | 17 | 159 | 92 | 83 | 154 | 258 | 218 | 267 | 405 | 564 | 693 | 885 | 539 | 112 | 17 | 104 | 177 | 135 |
| 57 cm | 1 | 14 | 9 | 5 | 20 | 29 | 54 | 33 | 31 | 56 | 107 | 116 | 57 | 8 | 1 | 8 | 16 | 12 |
| 60 cm | | | | 1 | 4 | 3 | 36 | 5 | 6 | 4 | 14 | 10 | 8 | | | | | |
| 63 cm | | | | 1 | 1 | 1 | 18 | 1 | | | | | 2 | | | | | 2 |
| Total | 816 | 5,206 | 3,623 | 2,117 | 3,587 | 6,598 | 5,512 | 5,285 | 7,190 | 5,373 | 8,811 | 10,077 | 8,771 | 3,393 | 2,301 | 4,978 | 5,628 | 8,484 |

¹ Size groups by 3-cm. intervals.

TABLE 15.—Size compositions of scrod haddock, fall seasons

[In thousands of fish]

| Length ¹ | 1931 | 1932 | 1933 | 1934 | 1935 | 1936 | 1937 | 1938 | 1939 | 1940 | 1941 | 1942 | 1943 | 1944 | 1945 | 1946 | 1947 | 1948 |
|---------------------|-------|-------|-------|-------|-------|-------|-------|--------|--------|-------|--------|-------|-------|-------|-------|-------|--------|-------|
| 24 cm | | | | | | | 2 | | | | 1 | | | | | | | |
| 27 cm | | | 1 | | | 17 | 166 | | | 6 | 10 | 1 | | | 1 | 1 | | |
| 30 cm | 1 | | 17 | 2 | 1 | 4 | 30 | | | 59 | 5 | 3 | | 3 | 6 | 8 | 4 | 4 |
| 33 cm | 7 | 1 | 44 | 41 | 24 | 103 | 383 | 40 | 43 | 183 | 136 | 58 | 41 | 50 | 12 | 36 | 54 | 104 |
| 36 cm | 127 | 29 | 254 | 362 | 256 | 345 | 385 | 370 | 354 | 242 | 664 | 648 | 337 | 139 | 174 | 267 | 1,122 | 1,076 |
| 39 cm | 736 | 318 | 1,228 | 1,094 | 1,310 | 1,298 | 1,050 | 2,291 | 1,591 | 414 | 2,519 | 1,901 | 836 | 315 | 812 | 884 | 4,250 | 2,158 |
| 42 cm | 1,299 | 1,547 | 1,943 | 1,096 | 2,173 | 2,690 | 2,444 | 4,587 | 3,262 | 1,191 | 4,546 | 2,294 | 1,221 | 732 | 1,244 | 1,496 | 4,482 | 2,389 |
| 45 cm | 777 | 2,592 | 1,121 | 693 | 1,648 | 2,172 | 2,395 | 3,359 | 3,178 | 1,535 | 3,180 | 1,785 | 1,553 | 929 | 798 | 1,504 | 2,205 | 2,262 |
| 48 cm | 203 | 1,362 | 373 | 385 | 660 | 776 | 994 | 1,092 | 1,546 | 1,076 | 1,106 | 1,095 | 1,162 | 418 | 213 | 798 | 921 | 1,241 |
| 51 cm | 28 | 203 | 53 | 67 | 111 | 140 | 188 | 177 | 308 | 347 | 277 | 264 | 281 | 77 | 32 | 164 | 147 | 236 |
| 54 cm | 7 | 21 | 1 | 1 | 7 | 14 | 19 | 25 | 27 | 47 | 35 | 34 | 24 | 14 | 16 | 11 | 24 | 32 |
| 57 cm | 1 | 2 | | 1 | | 2 | | 3 | 3 | 3 | 2 | 3 | 2 | 2 | 2 | | 4 | 6 |
| 60 cm | | | | | | | | | 1 | | | | | | | | | 2 |
| Total | 3,186 | 6,075 | 5,035 | 3,742 | 6,190 | 7,561 | 8,056 | 11,945 | 10,313 | 5,093 | 12,535 | 8,088 | 5,460 | 2,679 | 3,310 | 5,169 | 13,213 | 9,510 |

¹ Size groups by 3-cm. intervals.

TABLE 16.—Size compositions of scrod haddock, winter seasons

[In thousands of fish]

| Length ¹ | 1931 | 1932 | 1933 | 1934 | 1935 | 1936 | 1937 | 1938 | 1939 | 1940 | 1941 | 1942 | 1943 | 1944 | 1945 | 1946 | 1947 | 1948 |
|---------------------|-------|-------|------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|------|-------|-------|-------|-------|
| 24 cm | | | | | | 2 | | | | | | | | | | | | |
| 27 cm | | | | | 16 | 11 | 5 | 2 | 4 | 2 | 7 | 1 | | | | | | |
| 30 cm | 1 | | | | 80 | 140 | 37 | 18 | 67 | 28 | 63 | 55 | 10 | | | 3 | | 4 |
| 33 cm | 34 | 1 | 14 | 109 | 485 | 168 | 74 | 243 | 135 | 321 | 114 | 49 | 5 | 5 | 7 | 24 | 32 | 105 |
| 36 cm | 694 | 19 | 68 | 176 | 632 | 343 | 149 | 415 | 434 | 566 | 169 | 204 | 63 | 28 | 37 | 107 | 254 | 893 |
| 39 cm | 2,054 | 152 | 189 | 292 | 643 | 362 | 155 | 644 | 719 | 419 | 403 | 779 | 199 | 52 | 192 | 330 | 813 | 1,712 |
| 42 cm | 2,264 | 567 | 362 | 389 | 989 | 460 | 210 | 1,199 | 826 | 288 | 1,076 | 1,351 | 274 | 79 | 448 | 661 | 878 | 1,537 |
| 45 cm | 1,205 | 545 | 231 | 204 | 1,345 | 565 | 395 | 1,176 | 887 | 450 | 1,412 | 1,076 | 281 | 191 | 457 | 595 | 444 | 1,132 |
| 48 cm | 274 | 445 | 63 | 44 | 718 | 250 | 292 | 481 | 510 | 384 | 698 | 523 | 208 | 107 | 120 | 296 | 177 | 576 |
| 51 cm | 19 | 59 | 8 | 6 | 136 | 36 | 79 | 105 | 116 | 113 | 156 | 132 | 40 | 8 | 11 | 74 | 42 | 134 |
| 54 cm | 1 | 6 | 1 | | 11 | 5 | 9 | 11 | 12 | 17 | 26 | 14 | 7 | | 1 | 14 | 3 | 17 |
| 57 cm | 1 | | | | 1 | 1 | 1 | 2 | 1 | 1 | 1 | 1 | 1 | | | 3 | 1 | 3 |
| 60 cm | | | | | | | | | | | | | | | | | | |
| Total | 6,547 | 2,094 | 936 | 1,316 | 5,113 | 2,232 | 1,384 | 4,350 | 3,670 | 2,623 | 4,117 | 4,140 | 1,078 | 470 | 1,273 | 2,108 | 2,644 | 6,113 |

¹ Size groups by 3-cm. intervals.

TABLE 17.—Size composition, scrod haddock, in each of the 18 years
[In thousands of fish]

| Length ¹ | 1931 | 1932 | 1933 | 1934 | 1935 | 1936 | 1937 | 1938 | 1939 | 1940 | 1941 | 1942 | 1943 | 1944 | 1945 | 1946 | 1947 | 1948 |
|---------------------|--------|--------|--------|-------|--------|--------|--------|--------|--------|--------|--------|--------|--------|-------|-------|--------|--------|--------|
| 21 cm. | | | | | | 1 | 1 | | | | 1 | | | | | | | |
| 24 cm. | | | 14 | | 2 | | 2 | 1 | | | 6 | | | | | | | |
| 27 cm. | | | 153 | 16 | 11 | 15 | 32 | 9 | 2 | 7 | 34 | 3 | 2 | | 1 | 12 | 4 | |
| 30 cm. | 3 | 8 | 307 | 84 | 142 | 101 | 193 | 114 | 39 | 111 | 237 | 30 | 39 | 9 | 48 | 48 | 4 | 8 |
| 33 cm. | 42 | 49 | 433 | 164 | 523 | 473 | 514 | 458 | 240 | 529 | 968 | 246 | 323 | 76 | 48 | 285 | 113 | 263 |
| 36 cm. | 844 | 454 | 890 | 655 | 1,090 | 1,421 | 904 | 1,357 | 1,295 | 1,043 | 2,993 | 1,508 | 1,263 | 314 | 546 | 967 | 2,007 | 2,752 |
| 39 cm. | 2,920 | 2,056 | 2,225 | 1,673 | 2,644 | 3,362 | 2,602 | 4,395 | 3,829 | 1,715 | 6,089 | 4,295 | 3,085 | 1,059 | 1,902 | 2,251 | 6,811 | 6,159 |
| 42 cm. | 3,963 | 5,172 | 3,388 | 2,199 | 4,205 | 5,222 | 4,930 | 7,577 | 6,818 | 3,309 | 8,366 | 6,520 | 5,197 | 2,287 | 2,524 | 3,388 | 7,767 | 7,382 |
| 45 cm. | 2,481 | 6,355 | 2,773 | 1,995 | 4,184 | 4,752 | 4,999 | 6,035 | 7,344 | 4,401 | 7,178 | 7,533 | 5,755 | 2,834 | 1,746 | 3,659 | 4,797 | 6,660 |
| 48 cm. | 697 | 2,980 | 1,261 | 1,048 | 2,128 | 2,446 | 2,730 | 2,884 | 4,409 | 3,787 | 4,544 | 5,854 | 4,100 | 1,642 | 549 | 1,865 | 2,379 | 3,459 |
| 51 cm. | 78 | 460 | 255 | 180 | 429 | 551 | 700 | 802 | 1,063 | 1,335 | 1,659 | 1,774 | 1,301 | 342 | 73 | 347 | 537 | 678 |
| 54 cm. | 11 | 43 | 18 | 7 | 41 | 65 | 111 | 114 | 101 | 163 | 246 | 235 | 204 | 39 | 21 | 33 | 69 | 85 |
| 57 cm. | 2 | 2 | 1 | 2 | 5 | 12 | 41 | 21 | 12 | 13 | 21 | 15 | 47 | 5 | 4 | 3 | 5 | 9 |
| 60 cm. | | | | 1 | | 1 | 18 | 5 | 1 | 1 | | | 20 | | | 2 | | 4 |
| 63 cm. | | | | | | | 2 | 1 | | | | | 8 | | | 1 | | |
| 66 cm. | | | | | | | | | | | | | 3 | | | | | |
| 69 cm. | | | | | | | | | | | | | 1 | | | | | |
| 72 cm. | | | | | | | | | | | | | | | | | | |
| Total | 11,041 | 17,579 | 11,718 | 8,024 | 15,404 | 18,424 | 17,780 | 23,773 | 25,153 | 16,414 | 32,342 | 28,013 | 21,349 | 8,607 | 7,420 | 12,861 | 24,489 | 27,459 |

¹ Size groups by 3-cm. intervals.

TABLE 18.—Average size composition of scrod haddock, in each of the seasons
[In thousands of fish]

| Length ¹ | Spring | Summer | Fall | Winter |
|---------------------|--------|--------|-------|--------|
| 24 cm. | | 1 | | |
| 27 cm. | 1 | 10 | 3 | 3 |
| 30 cm. | 12 | 23 | 19 | 28 |
| 33 cm. | 56 | 81 | 76 | 107 |
| 36 cm. | 157 | 393 | 397 | 292 |
| 39 cm. | 346 | 985 | 1,389 | 562 |
| 42 cm. | 643 | 1,341 | 2,258 | 770 |
| 45 cm. | 808 | 1,354 | 1,871 | 716 |
| 48 cm. | 577 | 933 | 857 | 342 |
| 51 cm. | 184 | 271 | 172 | 71 |
| 54 cm. | 29 | 32 | 20 | 9 |
| 57 cm. | 4 | 5 | 2 | 1 |
| 60 cm. | 1 | 1 | | |
| 63 cm. | 1 | | | |
| Total | 2,819 | 5,430 | 7,064 | 2,901 |

¹ Size groups by 3-cm. intervals.

TABLE 19.—Size composition of scrod haddock in the average year
[In thousands of fish]

| Length ¹ | Average number | Percent of total |
|---------------------|----------------|------------------|
| 24 cm. | 1 | 0.1 |
| 27 cm. | 17 | .4 |
| 30 cm. | 82 | 1.8 |
| 33 cm. | 320 | 6.8 |
| 36 cm. | 1,240 | 18.0 |
| 39 cm. | 3,281 | 27.5 |
| 42 cm. | 5,012 | 26.1 |
| 45 cm. | 4,747 | 14.9 |
| 48 cm. | 2,710 | 3.8 |
| 51 cm. | 699 | .5 |
| 54 cm. | 89 | .1 |
| 57 cm. | 12 | |
| 60 cm. | 3 | |
| 63 cm. | 1 | |
| Total | 18,214 | 100.0 |

¹ Size groups by 3-cm. intervals.

TABLE 20.—Size compositions of large haddock, spring seasons
[In thousands of fish]

| Length ¹ | 1931 | 1932 | 1933 | 1934 | 1935 | 1936 | 1937 | 1938 | 1939 | 1940 | 1941 | 1942 | 1943 | 1944 | 1945 | 1946 | 1947 | 1948 |
|---------------------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|
| 30 cm. | | | | | | | | | | | | | | | | 1 | | |
| 33 cm. | | | | | | | | | | | | | | | | | | |
| 36 cm. | | | | | | | | 1 | | | | | | 1 | | 4 | 1 | |
| 39 cm. | | | | | | | | 2 | 4 | | | | | 7 | 3 | 20 | 3 | |
| 42 cm. | 6 | 1 | 16 | 5 | 4 | 1 | 8 | 7 | 2 | 1 | 3 | 1 | 4 | 36 | 25 | 118 | 16 | 11 |
| 45 cm. | 36 | 40 | 16 | 5 | 4 | 89 | 118 | 38 | 36 | 112 | 51 | 75 | 73 | 148 | 112 | 444 | 96 | 45 |
| 48 cm. | 256 | 279 | 181 | 67 | 44 | 276 | 132 | 450 | 282 | 647 | 395 | 462 | 416 | 596 | 395 | 663 | 461 | 206 |
| 51 cm. | 825 | 548 | 637 | 276 | 132 | 450 | 636 | 192 | 282 | 647 | 395 | 462 | 416 | 596 | 395 | 663 | 461 | 206 |
| 54 cm. | 1,398 | 684 | 1,074 | 410 | 240 | 777 | 1,225 | 526 | 652 | 1,171 | 1,025 | 945 | 958 | 1,068 | 673 | 608 | 946 | 626 |
| 57 cm. | 1,567 | 753 | 897 | 394 | 289 | 798 | 1,171 | 658 | 778 | 1,018 | 1,169 | 992 | 978 | 1,045 | 791 | 551 | 1,139 | 627 |
| 60 cm. | 1,537 | 834 | 753 | 317 | 233 | 668 | 897 | 599 | 715 | 683 | 940 | 824 | 686 | 732 | 728 | 453 | 871 | 590 |
| 63 cm. | 1,185 | 714 | 632 | 231 | 178 | 435 | 608 | 444 | 560 | 402 | 604 | 586 | 416 | 477 | 585 | 322 | 641 | 410 |
| 66 cm. | 742 | 497 | 450 | 153 | 140 | 306 | 412 | 306 | 416 | 241 | 363 | 382 | 252 | 283 | 350 | 228 | 545 | 346 |
| 69 cm. | 364 | 300 | 282 | 99 | 76 | 174 | 232 | 183 | 266 | 180 | 198 | 167 | 162 | 172 | 185 | 109 | 275 | 228 |
| 72 cm. | 132 | 148 | 118 | 47 | 38 | 87 | 122 | 92 | 167 | 88 | 101 | 94 | 67 | 84 | 87 | 46 | 122 | 162 |
| 75 cm. | 51 | 43 | 41 | 18 | 17 | 34 | 54 | 44 | 78 | 42 | 52 | 44 | 25 | 39 | 32 | 19 | 63 | 64 |
| 78 cm. | 18 | 16 | 7 | 6 | 3 | 10 | 17 | 19 | 26 | 10 | 23 | 14 | 8 | 16 | 9 | 7 | 19 | 19 |
| 81 cm. | | 2 | 4 | 4 | | 1 | 2 | 5 | 2 | 5 | 5 | 4 | 4 | | | 1 | 1 | 1 |
| 84 cm. | | | | | | | | | 1 | | | | | | | | | |
| Total | 8,117 | 4,859 | 5,092 | 2,028 | 1,394 | 3,839 | 5,504 | 3,118 | 3,998 | 4,590 | 4,930 | 4,630 | 4,064 | 4,708 | 3,981 | 3,595 | 5,287 | 3,236 |

¹ Size groups by 3-cm. intervals.

TABLE 21.—Size compositions of large haddock, summer seasons

[In thousands of fish]

| Length ¹ | 1931 | 1932 | 1933 | 1934 | 1935 | 1936 | 1937 | 1938 | 1939 | 1940 | 1941 | 1942 | 1943 | 1944 | 1945 | 1946 | 1947 | 1948 |
|---------------------|--------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|
| 33 cm | | | | | | 1 | | 1 | | | | | | 1 | 11 | 15 | | |
| 36 cm | | | | | | | | | | | | | | | 19 | 49 | 49 | 1 |
| 39 cm | 1 | 2 | 14 | 3 | 2 | 11 | 4 | 6 | 5 | 2 | 6 | 2 | 6 | 19 | 49 | 140 | | |
| 42 cm | 46 | 27 | 44 | 23 | 18 | 33 | 19 | 20 | 23 | 12 | 14 | 12 | 16 | 95 | 119 | 140 | 11 | 19 |
| 45 cm | 291 | 318 | 280 | 143 | 210 | 315 | 127 | 85 | 129 | 97 | 56 | 105 | 65 | 457 | 362 | 604 | 104 | 147 |
| 48 cm | 1,191 | 876 | 1,065 | 629 | 938 | 1,215 | 650 | 486 | 848 | 676 | 461 | 767 | 550 | 1,444 | 782 | 1,480 | 394 | 675 |
| 51 cm | 2,099 | 936 | 1,555 | 971 | 1,457 | 1,458 | 1,352 | 1,054 | 1,562 | 1,589 | 1,378 | 1,517 | 1,370 | 2,043 | 1,049 | 1,624 | 739 | 905 |
| 54 cm | 2,138 | 789 | 1,374 | 869 | 1,351 | 1,090 | 1,227 | 1,107 | 1,644 | 1,726 | 1,349 | 1,448 | 1,448 | 1,862 | 1,203 | 1,411 | 901 | 813 |
| 57 cm | 1,784 | 758 | 932 | 689 | 983 | 698 | 859 | 868 | 798 | 1,176 | 1,458 | 1,036 | 993 | 1,306 | 1,264 | 1,332 | 906 | 521 |
| 60 cm | 1,427 | 773 | 628 | 542 | 607 | 409 | 550 | 583 | 520 | 927 | 669 | 560 | 913 | 806 | 1,107 | 686 | 354 | |
| 63 cm | 910 | 630 | 424 | 327 | 344 | 257 | 316 | 354 | 380 | 451 | 552 | 439 | 332 | 521 | 411 | 651 | 672 | 238 |
| 66 cm | 507 | 389 | 242 | 160 | 175 | 146 | 190 | 179 | 202 | 273 | 291 | 249 | 187 | 291 | 216 | 359 | 439 | 117 |
| 69 cm | 255 | 209 | 128 | 74 | 66 | 60 | 116 | 90 | 112 | 149 | 153 | 111 | 87 | 172 | 108 | 181 | 260 | 52 |
| 72 cm | 103 | 82 | 56 | 25 | 22 | 24 | 67 | 39 | 54 | 65 | 86 | 59 | 38 | 60 | 37 | 66 | 114 | 75 |
| 75 cm | 36 | 32 | 13 | 9 | 5 | 5 | 28 | 12 | 13 | 28 | 30 | 27 | 10 | 22 | 14 | 14 | 34 | 12 |
| 78 cm | 7 | 7 | 8 | | 1 | 1 | 11 | 4 | 5 | 13 | 10 | 10 | 2 | 10 | 5 | 8 | 17 | 8 |
| 81 cm | 2 | 2 | 2 | | | | | | 2 | 2 | 2 | 1 | 1 | 2 | | 1 | 2 | |
| 84 cm | 1 | 1 | | | | | | | | | | | | | | | | |
| 87 cm | 1 | | | | | | | | | | | | | | | | | |
| Total | 10,799 | 5,831 | 6,765 | 4,464 | 6,179 | 5,723 | 5,517 | 4,888 | 5,876 | 6,899 | 7,150 | 6,353 | 5,665 | 9,218 | 6,436 | 9,043 | 5,279 | 3,937 |

¹ Size groups by 3-cm. intervals.

TABLE 22.—Size compositions of large haddock, fall seasons

[In thousands of fish]

| Length ¹ | 1931 | 1932 | 1933 | 1934 | 1935 | 1936 | 1937 | 1938 | 1939 | 1940 | 1941 | 1942 | 1943 | 1944 | 1945 | 1946 | 1947 | 1948 |
|---------------------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|
| 36 cm | | 1 | | | | 1 | | | | | | 1 | | | 6 | 3 | | |
| 39 cm | 14 | 4 | 6 | 4 | 1 | 12 | 4 | 2 | 5 | 1 | 6 | 4 | 3 | 6 | 49 | 22 | 3 | 2 |
| 42 cm | 59 | 43 | 48 | 44 | 19 | 44 | 21 | 19 | 26 | 10 | 30 | 11 | 16 | 62 | 181 | 74 | 23 | 14 |
| 45 cm | 233 | 545 | 401 | 213 | 324 | 337 | 203 | 199 | 362 | 90 | 156 | 129 | 109 | 436 | 406 | 346 | 126 | 132 |
| 48 cm | 463 | 1,726 | 1,173 | 617 | 1,320 | 1,081 | 913 | 729 | 1,451 | 640 | 901 | 825 | 684 | 1,248 | 554 | 1,316 | 559 | 694 |
| 51 cm | 852 | 1,802 | 1,585 | 944 | 1,770 | 1,605 | 1,327 | 1,043 | 1,978 | 1,326 | 1,145 | 1,361 | 1,267 | 1,771 | 820 | 2,047 | 1,005 | 950 |
| 54 cm | 1,029 | 1,158 | 1,277 | 777 | 1,174 | 1,182 | 1,179 | 805 | 1,196 | 1,242 | 1,122 | 883 | 1,109 | 1,671 | 984 | 1,521 | 1,023 | 729 |
| 57 cm | 1,089 | 716 | 762 | 584 | 712 | 600 | 788 | 486 | 505 | 776 | 752 | 438 | 713 | 1,217 | 936 | 797 | 531 | |
| 60 cm | 956 | 514 | 405 | 372 | 336 | 274 | 468 | 282 | 195 | 413 | 449 | 229 | 356 | 671 | 706 | 878 | 494 | 413 |
| 63 cm | 722 | 392 | 223 | 210 | 166 | 138 | 228 | 149 | 116 | 210 | 221 | 124 | 193 | 327 | 387 | 613 | 360 | 272 |
| 66 cm | 410 | 244 | 156 | 90 | 63 | 78 | 121 | 64 | 55 | 113 | 143 | 71 | 115 | 128 | 104 | 343 | 255 | 156 |
| 69 cm | 171 | 110 | 67 | 37 | 24 | 38 | 70 | 28 | 19 | 53 | 54 | 42 | 50 | 55 | 118 | 164 | 131 | 93 |
| 72 cm | 103 | 48 | 28 | 9 | 11 | 12 | 27 | 17 | 6 | 25 | 24 | 14 | 20 | 25 | 60 | 52 | 80 | 52 |
| 75 cm | 49 | 12 | 10 | 5 | 6 | 5 | 11 | 7 | 8 | 10 | 16 | 5 | 12 | 9 | 23 | 20 | 16 | 20 |
| 78 cm | 11 | 2 | 2 | 1 | 1 | 1 | 5 | 2 | 2 | 3 | 6 | 1 | 4 | 3 | 2 | 2 | 4 | 7 |
| 81 cm | 2 | 1 | 3 | | | | 1 | 1 | | | 1 | | 2 | | 1 | 2 | | 1 |
| 84 cm | 1 | | 1 | | | | | | | | | | | | 1 | | | |
| Total | 6,164 | 7,318 | 6,147 | 3,907 | 5,927 | 5,408 | 5,366 | 3,833 | 5,924 | 4,912 | 4,726 | 4,138 | 4,653 | 7,629 | 5,428 | 8,399 | 4,878 | 4,075 |

¹ Size groups by 3-cm. intervals.

TABLE 23.—Size compositions of large haddock, winter seasons

[In thousands of fish]

| Length ¹ | 1931 | 1932 | 1933 | 1934 | 1935 | 1936 | 1937 | 1938 | 1939 | 1940 | 1941 | 1942 | 1943 | 1944 | 1945 | 1946 | 1947 | 1948 |
|---------------------|-------|-------|-------|------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|
| 30 cm | | | | | | | | | | | | | | | 2 | | | |
| 33 cm | | 1 | | | | | | | | | | | | | 4 | | | 1 |
| 36 cm | | 1 | | | | | | | | | | | | | 25 | | | 3 |
| 39 cm | 6 | 2 | 1 | 1 | 1 | 1 | 1 | 4 | 2 | 18 | 1 | 1 | 2 | 7 | 48 | | | 7 |
| 42 cm | 96 | 36 | 4 | 10 | 12 | 13 | 7 | 16 | 16 | 41 | 2 | 10 | 14 | 29 | 138 | 38 | 5 | |
| 45 cm | 410 | 276 | 39 | 87 | 216 | 119 | 49 | 90 | 124 | 64 | 36 | 55 | 68 | 90 | 314 | 137 | 51 | 94 |
| 48 cm | 404 | 837 | 157 | 191 | 734 | 384 | 220 | 315 | 474 | 232 | 148 | 307 | 289 | 193 | 446 | 394 | 145 | 420 |
| 51 cm | 420 | 911 | 257 | 229 | 748 | 431 | 355 | 451 | 734 | 448 | 356 | 536 | 432 | 459 | 322 | 598 | 257 | 486 |
| 54 cm | 459 | 557 | 240 | 161 | 526 | 364 | 439 | 416 | 564 | 470 | 439 | 470 | 346 | 561 | 380 | 624 | 318 | 389 |
| 57 cm | 422 | 396 | 179 | 98 | 376 | 265 | 406 | 318 | 374 | 310 | 371 | 298 | 219 | 569 | 461 | 528 | 310 | 326 |
| 60 cm | 332 | 341 | 118 | 64 | 276 | 170 | 322 | 249 | 229 | 220 | 303 | 197 | 145 | 470 | 451 | 444 | 258 | 256 |
| 63 cm | 229 | 258 | 86 | 36 | 222 | 116 | 249 | 182 | 165 | 146 | 201 | 118 | 86 | 279 | 327 | 354 | 198 | 196 |
| 66 cm | 121 | 167 | 50 | 18 | 160 | 71 | 155 | 124 | 111 | 104 | 122 | 69 | 46 | 174 | 200 | 188 | 148 | 136 |
| 69 cm | 67 | 67 | 26 | 10 | 82 | 37 | 90 | 71 | 57 | 51 | 70 | 36 | 30 | 99 | 101 | 100 | 101 | 81 |
| 72 cm | 32 | 32 | 9 | 3 | 36 | 16 | 48 | 35 | 31 | 18 | 42 | 17 | 12 | 43 | 40 | 44 | 47 | 44 |
| 75 cm | 8 | 9 | 5 | 4 | 8 | 5 | 20 | 12 | 10 | 12 | 18 | 11 | 6 | 12 | 19 | 16 | 17 | 15 |
| 78 cm | | 2 | 1 | | 2 | 1 | 5 | 2 | 2 | 6 | 3 | 2 | 1 | 1 | 8 | 4 | 6 | 4 |
| 81 cm | | 1 | | | | | 1 | | 1 | 2 | 2 | 1 | 2 | | 1 | 1 | 1 | 1 |
| 84 cm | | | | | | | | | | 1 | | | | | | | | |
| Total | 3,006 | 3,894 | 1,172 | 912 | 3,399 | 1,993 | 2,367 | 2,285 | 2,894 | 2,148 | 2,114 | 2,128 | 1,698 | 2,987 | 3,287 | 3,475 | 1,862 | 2,461 |

¹ Size groups by 3-cm. intervals.

TABLE 24.—Size composition of large haddock, in each of the 18 years
[In thousands of fish]

| Length ¹ | 1931 | 1932 | 1933 | 1934 | 1935 | 1936 | 1937 | 1938 | 1939 | 1940 | 1941 | 1942 | 1943 | 1944 | 1945 | 1946 | 1947 | 1948 |
|---------------------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|
| 30 cm | | | | | | | | | | | | | | | 2 | 1 | | |
| 33 cm | | 1 | | | | | | | | | | | | | 4 | 1 | | |
| 36 cm | | 2 | | | | | | | | | | | | | 1 | | | |
| 39 cm | 27 | 9 | 21 | 9 | 4 | 2 | 11 | 16 | 14 | 22 | 13 | 8 | 15 | 3 | 42 | 22 | 1 | 1 |
| 42 cm | 237 | 146 | 112 | 82 | 53 | 99 | 55 | 62 | 72 | 72 | 49 | 43 | 61 | 222 | 149 | 96 | 6 | 6 |
| 45 cm | 1,190 | 1,418 | 901 | 510 | 794 | 860 | 497 | 412 | 651 | 363 | 299 | 364 | 315 | 1,131 | 1,194 | 1,531 | 377 | 51 |
| 48 cm | 2,883 | 3,987 | 3,032 | 1,713 | 3,124 | 3,130 | 2,419 | 1,722 | 3,055 | 2,185 | 1,605 | 2,361 | 1,939 | 3,451 | 2,177 | 3,853 | 1,559 | 1,995 |
| 51 cm | 4,789 | 4,333 | 4,471 | 2,554 | 4,215 | 4,271 | 4,259 | 3,074 | 4,962 | 4,534 | 3,904 | 4,359 | 4,027 | 5,341 | 2,864 | 4,877 | 2,947 | 2,876 |
| 54 cm | 5,193 | 3,257 | 3,788 | 2,201 | 3,340 | 3,434 | 4,016 | 2,986 | 3,761 | 4,374 | 4,456 | 3,094 | 3,881 | 5,139 | 3,358 | 4,107 | 3,361 | 2,558 |
| 57 cm | 4,832 | 2,704 | 2,626 | 1,688 | 2,304 | 2,231 | 2,950 | 2,271 | 2,392 | 2,945 | 3,521 | 2,596 | 2,611 | 3,824 | 3,389 | 3,309 | 2,984 | 1,968 |
| 60 cm | 3,900 | 2,342 | 1,783 | 1,209 | 1,397 | 1,288 | 1,948 | 1,558 | 1,504 | 1,757 | 2,283 | 1,681 | 1,477 | 2,545 | 2,751 | 2,079 | 1,433 | |
| 63 cm | 2,603 | 1,777 | 1,183 | 726 | 872 | 817 | 1,205 | 991 | 1,077 | 1,048 | 1,337 | 1,063 | 863 | 1,410 | 1,475 | 1,846 | 1,775 | 1,054 |
| 66 cm | 1,402 | 1,100 | 780 | 367 | 474 | 469 | 698 | 550 | 634 | 650 | 754 | 586 | 510 | 765 | 795 | 999 | 1,117 | 637 |
| 69 cm | 625 | 534 | 339 | 168 | 210 | 222 | 398 | 281 | 355 | 341 | 378 | 283 | 234 | 410 | 414 | 491 | 614 | 388 |
| 72 cm | 289 | 205 | 134 | 55 | 86 | 86 | 196 | 135 | 169 | 150 | 204 | 134 | 95 | 167 | 169 | 181 | 304 | 235 |
| 75 cm | 111 | 69 | 35 | 24 | 22 | 25 | 76 | 50 | 57 | 60 | 87 | 57 | 36 | 59 | 71 | 59 | 74 | 66 |
| 78 cm | 4 | 4 | 5 | 5 | 4 | 4 | 23 | 13 | 19 | 27 | 24 | 17 | 11 | 18 | 15 | 14 | 28 | 20 |
| 81 cm | 2 | 1 | | | | | 3 | 1 | 5 | 5 | 6 | 2 | 5 | 2 | 4 | 5 | | 3 |
| 84 cm | 1 | | | | | | | | | | | | | | | | | |
| 87 cm | | | | | | | | | | | | | | | | | | |
| Total | 28,086 | 21,902 | 19,176 | 11,311 | 16,899 | 16,963 | 18,754 | 14,124 | 18,692 | 18,549 | 18,920 | 17,249 | 16,080 | 24,542 | 19,132 | 24,512 | 17,306 | 13,709 |

¹ Size groups by 3-cm. intervals.

TABLE 25.—Average size composition of large haddock, in each of the seasons

[In thousands of fish]

| Length ¹ | Spring | Summer | Fall | Winter |
|---------------------|--------|--------|-------|--------|
| 36 cm | | 2 | 1 | 2 |
| 39 cm | 21 | 10 | 8 | 6 |
| 42 cm | 37 | 38 | 41 | 27 |
| 45 cm | 126 | 216 | 264 | 129 |
| 48 cm | 457 | 840 | 922 | 349 |
| 51 cm | 828 | 1,370 | 1,967 | 468 |
| 54 cm | 868 | 1,307 | 1,114 | 429 |
| 57 cm | 731 | 1,020 | 744 | 346 |
| 60 cm | 524 | 710 | 467 | 269 |
| 63 cm | 356 | 456 | 281 | 192 |
| 66 cm | 203 | 256 | 156 | 120 |
| 69 cm | 100 | 132 | 74 | 65 |
| 72 cm | 42 | 60 | 34 | 31 |
| 75 cm | 14 | 19 | 14 | 12 |
| 78 cm | 3 | 7 | 3 | 3 |
| 81 cm | | 1 | 1 | 1 |
| Total | 4,276 | 6,444 | 5,491 | 2,449 |

¹ Size groups by 3-cm. intervals.

TABLE 26.—Size composition of large haddock in the average year

[In thousands of fish]

| Length ¹ | Average number | Percent |
|---------------------|----------------|---------|
| 36 cm | 4 | |
| 39 cm | 27 | 0.1 |
| 42 cm | 128 | 0.7 |
| 45 cm | 735 | 3.9 |
| 48 cm | 2,569 | 13.8 |
| 51 cm | 4,032 | 21.7 |
| 54 cm | 3,718 | 19.9 |
| 57 cm | 2,841 | 15.2 |
| 60 cm | 1,970 | 10.6 |
| 63 cm | 1,285 | 6.9 |
| 66 cm | 736 | 3.9 |
| 69 cm | 371 | 2.0 |
| 72 cm | 167 | 0.9 |
| 75 cm | 59 | 0.3 |
| 78 cm | 16 | 0.1 |
| 81 cm | 3 | |
| Total | 18,661 | 100.0 |

¹ Size groups by 3-cm. intervals.

TABLE 27.—Size compositions of total haddock, spring seasons

[In thousands of fish]

| Length ¹ | 1931 | 1932 | 1933 | 1934 | 1935 | 1936 | 1937 | 1938 | 1939 | 1940 | 1941 | 1942 | 1943 | 1944 | 1945 | 1946 | 1947 | 1948 |
|---------------------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|--------|--------|--------|-------|-------|-------|-------|-------|
| 21 cm | | | | | | | | | | | | | | | | | | |
| 24 cm | | | | | | | | | | | | | | | | | | |
| 27 cm | | | | | | | | | | | | | | | | | | |
| 30 cm | 1 | 7 | | | | 28 | 9 | 2 | 9 | | 13 | | | | | | | |
| 33 cm | | 31 | 2 | 2 | 1 | 74 | 40 | 113 | 37 | 16 | 555 | 33 | 63 | 11 | 15 | 10 | 2 | |
| 36 cm | 15 | 292 | 44 | 14 | 13 | 98 | 105 | 174 | 162 | 101 | 1,211 | 125 | 208 | 63 | 2 | 42 | 76 | 87 |
| 39 cm | 62 | 928 | 172 | 63 | 68 | 153 | 280 | 189 | 462 | 296 | 1,121 | 386 | 778 | 257 | 33 | 117 | 385 | 531 |
| 42 cm | 196 | 1,504 | 507 | 247 | 148 | 379 | 562 | 321 | 969 | 663 | 822 | 1,127 | 1,551 | 516 | 169 | 316 | 908 | 1,047 |
| 45 cm | 431 | 1,390 | 1,009 | 415 | 207 | 739 | 984 | 558 | 1,267 | 1,101 | 1,235 | 2,023 | 1,740 | 762 | 342 | 632 | 995 | 970 |
| 48 cm | 893 | 879 | 1,115 | 432 | 226 | 970 | 1,362 | 734 | 1,133 | 1,558 | 1,644 | 1,992 | 1,571 | 1,077 | 507 | 720 | 1,010 | 780 |
| 51 cm | 1,412 | 723 | 1,176 | 434 | 268 | 894 | 1,440 | 779 | 886 | 1,482 | 1,558 | 1,438 | 1,399 | 1,213 | 686 | 613 | 1,117 | 699 |
| 54 cm | 1,569 | 755 | 904 | 395 | 292 | 815 | 1,200 | 703 | 809 | 1,061 | 1,247 | 1,063 | 1,094 | 1,062 | 794 | 551 | 1,165 | 651 |
| 57 cm | 1,537 | 834 | 753 | 317 | 233 | 674 | 901 | 610 | 717 | 688 | 944 | 825 | 722 | 735 | 730 | 453 | 971 | 590 |
| 60 cm | 1,185 | 714 | 632 | 231 | 178 | 435 | 608 | 448 | 561 | 402 | 604 | 586 | 434 | 477 | 585 | 322 | 641 | 410 |
| 63 cm | 742 | 497 | 450 | 153 | 140 | 306 | 412 | 307 | 416 | 241 | 363 | 382 | 260 | 283 | 350 | 228 | 545 | 346 |
| 66 cm | 364 | 300 | 282 | 99 | 76 | 174 | 232 | 183 | 266 | 160 | 198 | 197 | 165 | 172 | 185 | 109 | 275 | 228 |
| 69 cm | 132 | 148 | 118 | 47 | 38 | 87 | 122 | 92 | 167 | 88 | 101 | 94 | 68 | 84 | 87 | 46 | 122 | 162 |
| 72 cm | 51 | 43 | 41 | 18 | 17 | 34 | 54 | 44 | 78 | 42 | 52 | 44 | 26 | 39 | 32 | 19 | 63 | 64 |
| 75 cm | 18 | 16 | 7 | 6 | 3 | 10 | 17 | 19 | 26 | 10 | 23 | 14 | 8 | 16 | 15 | 9 | 7 | 19 |
| 78 cm | | 2 | 4 | 4 | | | 1 | 2 | 5 | 5 | 5 | 4 | 4 | | | | 1 | 1 |
| 81 cm | | | | | | | | | | | | | | | | | | |
| 84 cm | | | | | | | | | | | | | | | | | | |
| Total | 8,609 | 9,063 | 7,216 | 2,877 | 1,908 | 5,872 | 8,332 | 5,311 | 7,978 | 7,915 | 11,809 | 10,338 | 10,104 | 6,773 | 4,517 | 4,201 | 8,291 | 6,588 |

¹ Size groups by 3-cm. intervals.

TABLE 28.—Size compositions of total haddock, summer seasons

[In thousands of fish]

| Length ¹ | 1931 | 1932 | 1933 | 1934 | 1935 | 1936 | 1937 | 1938 | 1939 | 1940 | 1941 | 1942 | 1943 | 1944 | 1945 | 1946 | 1947 | 1948 |
|---------------------|--------|--------|--------|-------|-------|--------|--------|--------|--------|--------|--------|--------|--------|--------|-------|--------|--------|--------|
| 21 cm. | | | | | | 1 | | | | | | | | | | | | |
| 24 cm. | | | 14 | | | 2 | | 1 | | | 2 | | | | | | | |
| 27 cm. | | | 152 | | | 5 | | 3 | | | 4 | 1 | 1 | | | 11 | | |
| 30 cm. | | 1 | 290 | 2 | 1 | 19 | | 16 | 1 | | 15 | 10 | 24 | 4 | | 29 | | |
| 33 cm. | | 16 | 373 | 12 | 13 | 128 | 17 | 62 | 25 | 9 | 163 | 106 | 214 | 10 | 29 | 211 | 17 | 52 |
| 36 cm. | 8 | 114 | 524 | 103 | 189 | 636 | 265 | 397 | 345 | 134 | 949 | 531 | 655 | 86 | 344 | 570 | 556 | 696 |
| 39 cm. | 75 | 661 | 650 | 228 | 625 | 1,561 | 1,123 | 1,281 | 1,064 | 589 | 2,052 | 1,232 | 1,282 | 461 | 917 | 989 | 1,366 | 1,759 |
| 42 cm. | 286 | 1,621 | 636 | 495 | 917 | 1,735 | 1,741 | 1,497 | 1,791 | 1,188 | 1,939 | 1,770 | 2,182 | 1,091 | 807 | 1,173 | 1,526 | 2,439 |
| 45 cm. | 615 | 2,125 | 873 | 893 | 1,238 | 1,680 | 1,470 | 1,065 | 2,177 | 1,524 | 1,458 | 2,829 | 2,319 | 1,557 | 623 | 1,976 | 1,353 | 2,488 |
| 48 cm. | 1,343 | 1,718 | 1,412 | 1,092 | 1,594 | 2,115 | 1,368 | 1,255 | 2,350 | 2,092 | 1,952 | 3,473 | 2,125 | 2,080 | 886 | 2,194 | 1,126 | 1,743 |
| 51 cm. | 2,116 | 1,095 | 1,647 | 854 | 1,611 | 1,716 | 1,570 | 1,321 | 1,967 | 2,153 | 2,071 | 2,402 | 1,909 | 2,155 | 1,066 | 1,728 | 916 | 1,040 |
| 54 cm. | 2,139 | 803 | 1,383 | 874 | 1,371 | 1,119 | 1,281 | 1,140 | 1,254 | 1,700 | 1,833 | 1,465 | 1,505 | 1,870 | 1,204 | 1,419 | 917 | 825 |
| 57 cm. | 1,784 | 758 | 933 | 690 | 987 | 701 | 895 | 873 | 804 | 1,180 | 1,472 | 1,046 | 1,001 | 1,306 | 1,264 | 1,332 | 906 | 521 |
| 60 cm. | 1,427 | 773 | 628 | 543 | 607 | 410 | 568 | 584 | 520 | 722 | 927 | 669 | 562 | 913 | 806 | 1,108 | 686 | 356 |
| 63 cm. | 910 | 630 | 424 | 327 | 344 | 257 | 318 | 354 | 380 | 451 | 552 | 439 | 332 | 521 | 411 | 652 | 672 | 238 |
| 66 cm. | 507 | 389 | 242 | 160 | 175 | 146 | 190 | 179 | 202 | 273 | 291 | 249 | 187 | 291 | 216 | 359 | 439 | 117 |
| 69 cm. | 255 | 209 | 128 | 74 | 66 | 60 | 116 | 90 | 112 | 149 | 153 | 111 | 87 | 172 | 108 | 181 | 260 | 52 |
| 72 cm. | 103 | 82 | 56 | 25 | 22 | 24 | 67 | 39 | 54 | 65 | 86 | 59 | 38 | 60 | 37 | 66 | 114 | 75 |
| 75 cm. | 36 | 32 | 13 | 9 | 5 | 5 | 28 | 12 | 13 | 28 | 30 | 27 | 10 | 22 | 14 | 14 | 34 | 12 |
| 78 cm. | 7 | 7 | 8 | 1 | 1 | 1 | 11 | 4 | 5 | 13 | 10 | 10 | 2 | 10 | 5 | 8 | 17 | 8 |
| 81 cm. | 2 | 2 | 2 | | | | 1 | | 2 | 2 | 2 | 1 | 1 | 2 | | 1 | 2 | |
| 84 cm. | 1 | 1 | | | | | | | | | | | | | | | | |
| 87 cm. | 1 | | | | | | | | | | | | | | | | | |
| Total..... | 11,615 | 11,037 | 10,388 | 6,381 | 9,766 | 12,321 | 11,029 | 10,173 | 13,066 | 12,272 | 15,961 | 16,430 | 14,436 | 12,611 | 8,737 | 14,021 | 10,907 | 12,421 |

¹ Size groups by 3-cm. intervals.

TABLE 29.—Size compositions of total haddock, fall seasons

[In thousands of fish]

| Length ¹ | 1931 | 1932 | 1933 | 1934 | 1935 | 1936 | 1937 | 1938 | 1939 | 1940 | 1941 | 1942 | 1943 | 1944 | 1945 | 1946 | 1947 | 1948 |
|---------------------|-------|--------|--------|-------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|-------|--------|--------|--------|
| 24 cm. | | | | | | 4 | 30 | | | | 1 | | | | | | | |
| 27 cm. | | | 1 | | | 17 | 166 | 1 | | 6 | 10 | 1 | | | 1 | 1 | | |
| 30 cm. | 1 | 7 | 17 | 2 | 1 | 17 | 383 | 40 | 43 | 48 | 59 | 5 | 3 | 3 | 6 | 8 | 4 | 4 |
| 33 cm. | 7 | 1 | 44 | 41 | 24 | 103 | 383 | 370 | 354 | 183 | 136 | 58 | 41 | 50 | 12 | 36 | 54 | 104 |
| 36 cm. | 127 | 30 | 254 | 362 | 256 | 346 | 385 | 370 | 354 | 242 | 664 | 649 | 337 | 139 | 180 | 270 | 1,122 | 1,076 |
| 39 cm. | 750 | 322 | 1,234 | 1,098 | 1,311 | 1,310 | 1,054 | 2,293 | 1,596 | 415 | 2,525 | 1,905 | 839 | 321 | 861 | 906 | 4,253 | 2,160 |
| 42 cm. | 1,358 | 1,590 | 1,991 | 1,140 | 2,192 | 2,734 | 2,465 | 4,606 | 3,288 | 1,201 | 4,576 | 2,305 | 1,237 | 794 | 1,425 | 1,570 | 4,505 | 2,403 |
| 45 cm. | 1,010 | 3,137 | 1,522 | 906 | 1,972 | 2,509 | 2,598 | 3,558 | 3,540 | 1,625 | 3,336 | 1,914 | 1,662 | 1,365 | 1,204 | 1,850 | 2,331 | 2,394 |
| 48 cm. | 666 | 3,088 | 1,546 | 1,002 | 1,980 | 1,857 | 1,907 | 1,821 | 2,997 | 1,716 | 1,707 | 1,920 | 1,846 | 1,666 | 767 | 2,114 | 1,480 | 1,635 |
| 51 cm. | 880 | 2,005 | 1,638 | 1,011 | 1,881 | 1,745 | 1,515 | 1,220 | 2,286 | 1,673 | 1,422 | 1,625 | 1,548 | 1,848 | 852 | 2,211 | 1,152 | 1,195 |
| 54 cm. | 1,036 | 1,179 | 1,278 | 778 | 1,181 | 1,196 | 1,198 | 830 | 1,223 | 1,289 | 1,157 | 917 | 1,133 | 1,685 | 1,000 | 1,532 | 1,047 | 761 |
| 57 cm. | 1,090 | 718 | 762 | 585 | 712 | 602 | 788 | 489 | 608 | 779 | 754 | 441 | 715 | 1,219 | 938 | 996 | 901 | 537 |
| 60 cm. | 956 | 514 | 405 | 372 | 336 | 274 | 468 | 282 | 195 | 414 | 449 | 229 | 356 | 671 | 706 | 878 | 494 | 415 |
| 63 cm. | 722 | 392 | 223 | 210 | 166 | 138 | 228 | 149 | 116 | 210 | 221 | 124 | 193 | 327 | 387 | 613 | 360 | 272 |
| 66 cm. | 410 | 244 | 156 | 90 | 63 | 78 | 121 | 64 | 55 | 113 | 143 | 71 | 115 | 128 | 194 | 343 | 255 | 156 |
| 69 cm. | 171 | 110 | 67 | 37 | 24 | 38 | 70 | 28 | 19 | 53 | 54 | 42 | 50 | 55 | 118 | 164 | 131 | 93 |
| 72 cm. | 103 | 48 | 28 | 9 | 11 | 12 | 27 | 17 | 6 | 25 | 24 | 14 | 20 | 25 | 60 | 52 | 80 | 52 |
| 75 cm. | 49 | 12 | 10 | 5 | 6 | 5 | 11 | 7 | 8 | 10 | 16 | 5 | 12 | 9 | 23 | 20 | 16 | 20 |
| 78 cm. | 11 | 2 | 2 | 1 | 1 | 1 | 5 | 2 | 2 | 3 | 6 | 1 | 4 | 3 | 2 | 2 | 4 | 7 |
| 81 cm. | 2 | 1 | 3 | | | | 1 | 1 | | | 1 | | 2 | | 1 | 2 | 2 | 1 |
| 84 cm. | 1 | | 1 | | | | | | | | | | | | 1 | | | |
| Total..... | 9,350 | 13,393 | 11,182 | 7,649 | 12,117 | 12,969 | 13,422 | 15,778 | 16,237 | 10,005 | 17,261 | 12,226 | 10,113 | 10,308 | 8,738 | 13,568 | 18,091 | 13,585 |

¹ Size groups by 3-cm. intervals.

TABLE 30.—Size compositions of total haddock, winter seasons

[In thousands of fish]

| Length ¹ | 1931 | 1932 | 1933 | 1934 | 1935 | 1936 | 1937 | 1938 | 1939 | 1940 | 1941 | 1942 | 1943 | 1944 | 1945 | 1946 | 1947 | 1948 |
|---------------------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|
| 24 cm. | | | | | 2 | | | | | | | | | | | | | |
| 27 cm. | | | | 16 | 11 | 5 | 2 | 4 | 2 | 1 | 7 | 1 | | | | | | |
| 30 cm. | 1 | | | 80 | 140 | 37 | 18 | 67 | 28 | 63 | 55 | 10 | | | 2 | 3 | | 4 |
| 33 cm. | 34 | 2 | 14 | 109 | 485 | 168 | 74 | 243 | 135 | 321 | 114 | 49 | 5 | 5 | 11 | 24 | 32 | 105 |
| 36 cm. | 694 | 20 | 68 | 176 | 632 | 343 | 149 | 418 | 434 | 571 | 169 | 204 | 63 | 29 | 62 | 107 | 254 | 894 |
| 39 cm. | 2,060 | 154 | 190 | 293 | 644 | 363 | 156 | 648 | 721 | 437 | 404 | 780 | 201 | 59 | 240 | 335 | 813 | 1,715 |
| 42 cm. | 2,360 | 603 | 366 | 399 | 1,001 | 473 | 217 | 1,215 | 842 | 329 | 1,078 | 1,361 | 288 | 108 | 586 | 699 | 883 | 1,544 |
| 45 cm. | 1,615 | 1,121 | 270 | 291 | 1,561 | 684 | 444 | 1,266 | 1,011 | 514 | 1,448 | 1,131 | 349 | 281 | 771 | 732 | 495 | 1,226 |
| 48 cm. | 678 | 1,282 | 220 | 235 | 1,452 | 634 | 512 | 796 | 984 | 616 | 846 | 830 | 497 | 300 | 566 | 690 | 322 | 996 |
| 51 cm. | 439 | 970 | 265 | 235 | 884 | 467 | 434 | 556 | 850 | 561 | 512 | 668 | 472 | 467 | 333 | 672 | 299 | 620 |
| 54 cm. | 460 | 563 | 241 | 161 | 537 | 369 | 448 | 427 | 576 | 487 | 465 | 484 | 353 | 561 | 381 | 638 | 321 | 406 |
| 57 cm. | 423 | 396 | 179 | 98 | 377 | 266 | 407 | 320 | 375 | 311 | 372 | 299 | 220 | 569 | 461 | 531 | 311 | 329 |
| 60 cm. | 332 | 341 | 118 | 64 | 276 | 170 | 322 | 249 | 229 | 220 | 303 | 197 | 145 | 470 | 451 | 445 | 258 | 256 |
| 63 cm. | 229 | 258 | 86 | 36 | 222 | 116 | 249 | 182 | 165 | 146 | 201 | 118 | 86 | 279 | 327 | 354 | 198 | 198 |
| 66 cm. | 121 | 167 | 50 | 18 | 160 | 71 | 155 | 124 | 111 | 104 | 122 | 69 | 46 | 174 | 200 | 188 | 148 | 136 |
| 69 cm. | 67 | 67 | 26 | 10 | 82 | 37 | 90 | 71 | 57 | 51 | 70 | 36 | 30 | 99 | 101 | 100 | 101 | 81 |
| 72 cm. | 32 | 32 | 9 | 3 | 36 | 16 | 48 | 35 | 31 | 18 | 42 | 17 | 12 | 43 | 40 | 44 | 47 | 44 |
| 75 cm. | 8 | 9 | 5 | 4 | 8 | 5 | 20 | 12 | 10 | 12 | 18 | 11 | 6 | 12 | 19 | 16 | 17 | 15 |
| 78 cm. | 2 | 1 | | | 2 | 1 | 5 | 2 | 2 | 6 | 3 | 2 | 1 | 1 | 8 | 4 | 6 | 4 |
| 81 cm. | | 1 | | | | | 1 | | 1 | 2 | 2 | 1 | 2 | | 1 | 1 | 1 | 1 |
| 84 cm. | | | | | | | | | | 1 | | | | | | | | |
| Total..... | 9,553 | 5,988 | 2,108 | 2,228 | 8,512 | 4,225 | 3,751 | 6,635 | 6,564 | 4,771 | 6,231 | 6,268 | 2,776 | 3,457 | 4,560 | 5,553 | 4,506 | 8,574 |

¹ Size groups by 3-cm. intervals.

TABLE 31.—Size composition of landings of total haddock, in each of the 18 years

[In thousands of fish]

| Length ¹ | 1931 | 1932 | 1933 | 1934 | 1935 | 1936 | 1937 | 1938 | 1939 | 1940 | 1941 | 1942 | 1943 | 1944 | 1945 | 1946 | 1947 | 1948 |
|---------------------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|
| 21 cm. | | | | | | 1 | 1 | | | | 1 | | | | | | | |
| 24 cm. | | | 14 | | 2 | 2 | 3 | 1 | | | 6 | | | | | | | |
| 27 cm. | | | 153 | 16 | 11 | 15 | 32 | 9 | 2 | 7 | 34 | 3 | 2 | | 1 | 12 | | |
| 30 cm. | 3 | 8 | 307 | 84 | 142 | 101 | 193 | 114 | 39 | 111 | 237 | 30 | 39 | 9 | 8 | 49 | | 8 |
| 33 cm. | 42 | 50 | 433 | 164 | 523 | 473 | 514 | 458 | 240 | 529 | 968 | 246 | 323 | 76 | 52 | 286 | 113 | 263 |
| 36 cm. | 844 | 456 | 890 | 655 | 1,090 | 1,423 | 904 | 1,359 | 1,295 | 1,048 | 2,993 | 1,509 | 1,263 | 317 | 588 | 989 | 2,008 | 2,753 |
| 39 cm. | 2,947 | 2,065 | 2,246 | 1,682 | 2,648 | 3,387 | 2,613 | 4,411 | 3,843 | 1,737 | 6,102 | 4,303 | 3,100 | 1,098 | 2,051 | 2,347 | 6,817 | 6,165 |
| 42 cm. | 4,200 | 5,318 | 3,500 | 2,281 | 4,258 | 5,321 | 4,985 | 7,639 | 6,890 | 3,381 | 8,415 | 6,583 | 5,258 | 2,509 | 2,987 | 3,758 | 7,822 | 7,433 |
| 45 cm. | 3,671 | 7,773 | 3,674 | 2,505 | 4,978 | 5,612 | 5,496 | 6,447 | 7,995 | 4,764 | 7,477 | 7,897 | 6,070 | 3,965 | 2,940 | 5,190 | 5,174 | 7,078 |
| 48 cm. | 3,580 | 6,967 | 4,293 | 2,761 | 5,252 | 5,576 | 5,149 | 4,606 | 7,464 | 5,982 | 6,149 | 8,125 | 6,039 | 5,123 | 2,726 | 5,718 | 3,938 | 5,454 |
| 51 cm. | 4,847 | 4,793 | 4,726 | 2,734 | 4,644 | 4,822 | 4,959 | 3,876 | 5,989 | 5,869 | 5,563 | 6,133 | 5,328 | 5,683 | 2,937 | 5,224 | 3,484 | 3,554 |
| 54 cm. | 5,204 | 3,300 | 3,806 | 2,208 | 3,381 | 3,499 | 4,127 | 3,100 | 3,862 | 4,537 | 4,702 | 3,929 | 4,085 | 5,178 | 3,379 | 4,140 | 3,450 | 2,643 |
| 57 cm. | 4,834 | 2,706 | 2,627 | 1,690 | 2,309 | 2,243 | 2,991 | 2,292 | 2,404 | 2,958 | 3,542 | 2,611 | 2,656 | 3,829 | 3,393 | 3,312 | 2,989 | 1,977 |
| 60 cm. | 3,900 | 2,342 | 1,783 | 1,210 | 1,397 | 1,289 | 1,966 | 1,583 | 1,505 | 1,758 | 2,283 | 1,681 | 1,497 | 2,531 | 2,548 | 2,753 | 2,079 | 1,437 |
| 63 cm. | 2,603 | 1,777 | 1,153 | 726 | 872 | 817 | 1,207 | 992 | 1,077 | 1,048 | 1,337 | 1,063 | 871 | 1,410 | 1,475 | 1,847 | 1,775 | 1,054 |
| 66 cm. | 1,402 | 1,100 | 730 | 367 | 474 | 469 | 698 | 550 | 634 | 650 | 754 | 586 | 513 | 765 | 795 | 999 | 1,117 | 637 |
| 69 cm. | 625 | 534 | 339 | 168 | 210 | 222 | 398 | 281 | 355 | 341 | 378 | 283 | 235 | 410 | 414 | 491 | 614 | 388 |
| 72 cm. | 289 | 205 | 134 | 55 | 86 | 86 | 196 | 135 | 169 | 150 | 204 | 134 | 96 | 167 | 169 | 181 | 304 | 235 |
| 75 cm. | 111 | 69 | 35 | 24 | 22 | 25 | 76 | 50 | 57 | 60 | 87 | 57 | 36 | 59 | 71 | 59 | 74 | 66 |
| 78 cm. | 18 | 13 | 15 | 5 | 4 | 4 | 23 | 13 | 19 | 27 | 24 | 17 | 11 | 18 | 15 | 14 | 28 | 20 |
| 81 cm. | 4 | 4 | 5 | | | | 3 | 1 | 5 | 5 | 6 | 2 | 5 | 2 | 2 | 4 | 5 | 3 |
| 84 cm. | 2 | 1 | | | | | | | 1 | | | | | | 1 | | | |
| 87 cm. | 1 | | | | | | | | | | | | | | | | | |
| Total..... | 39,127 | 39,481 | 30,894 | 19,335 | 32,303 | 35,387 | 36,534 | 37,897 | 43,845 | 34,963 | 51,262 | 45,262 | 37,429 | 33,149 | 26,552 | 37,373 | 41,795 | 41,168 |

¹ Size groups by 3-cm. intervals.

TABLE 32.—Average size composition of total ¹ haddock, in each of the seasons
[In thousands of fish]

| Length ² | Spring | Summer | Fall | Winter |
|---------------------|--------|--------|--------|--------|
| 24 cm..... | | 1 | | |
| 27 cm..... | 1 | 10 | 3 | 3 |
| 30 cm..... | 12 | 23 | 19 | 28 |
| 33 cm..... | 56 | 81 | 76 | 107 |
| 36 cm..... | 157 | 395 | 398 | 294 |
| 39 cm..... | 349 | 995 | 1,397 | 567 |
| 42 cm..... | 664 | 1,380 | 2,299 | 797 |
| 45 cm..... | 933 | 1,570 | 2,135 | 844 |
| 48 cm..... | 1,035 | 1,773 | 1,779 | 692 |
| 51 cm..... | 1,012 | 1,641 | 1,539 | 539 |
| 54 cm..... | 896 | 1,339 | 1,134 | 438 |
| 57 cm..... | 735 | 1,025 | 746 | 347 |
| 60 cm..... | 525 | 712 | 467 | 269 |
| 63 cm..... | 357 | 456 | 281 | 192 |
| 66 cm..... | 204 | 256 | 156 | 120 |
| 69 cm..... | 100 | 132 | 74 | 65 |
| 72 cm..... | 42 | 60 | 34 | 31 |
| 75 cm..... | 14 | 19 | 14 | 12 |
| 78 cm..... | 3 | 7 | 3 | 3 |
| 81 cm..... | | 1 | 1 | 1 |
| Total..... | 7,095 | 11,876 | 12,555 | 5,349 |

¹ All values calculated by dividing 18-year total for total haddock by 18 rather than by summing 18-year averages of scrod plus large.
² Size groups by 3-cm. intervals.

TABLE 33.—Size composition of total haddock in the average year
[In thousands of fish]

| Length ¹ | Average number | Percent |
|---------------------|----------------|---------|
| 24 cm..... | 1 | |
| 27 cm..... | 17 | 0.1 |
| 30 cm..... | 82 | .2 |
| 33 cm..... | 320 | .9 |
| 36 cm..... | 1,244 | 3.4 |
| 39 cm..... | 3,308 | 9.0 |
| 42 cm..... | 5,140 | 13.9 |
| 45 cm..... | 5,482 | 14.9 |
| 48 cm..... | 5,279 | 14.3 |
| 51 cm..... | 4,731 | 12.8 |
| 54 cm..... | 3,807 | 10.3 |
| 57 cm..... | 2,853 | 7.7 |
| 60 cm..... | 1,973 | 5.4 |
| 63 cm..... | 1,286 | 3.5 |
| 66 cm..... | 736 | 2.0 |
| 69 cm..... | 371 | 1.0 |
| 72 cm..... | 167 | .4 |
| 75 cm..... | 59 | .2 |
| 78 cm..... | 16 | |
| 81 cm..... | 3 | |
| Total..... | 36,875 | 100.0 |

¹ Size groups by 3-cm. intervals.

TABLE 34.—Undersized haddock landed, by years
[In thousands of fish]

| Year | Number of fish |
|--------------|----------------|
| 1931..... | 3,836 |
| 1932..... | 2,579 |
| 1933..... | 4,043 |
| 1934..... | 2,601 |
| 1935..... | 4,416 |
| 1936..... | 5,402 |
| 1937..... | 4,260 |
| 1938..... | 6,352 |
| 1939..... | 5,419 |
| 1940..... | 3,432 |
| 1941..... | 10,341 |
| 1942..... | 6,091 |
| 1943..... | 4,727 |
| 1944..... | 1,500 |
| 1945..... | 2,700 |
| 1946..... | 3,683 |
| 1947..... | 8,942 |
| 1948..... | 9,189 |
| Total..... | 89,513 |
| Average..... | 4,974 |

TABLE 35.—Division of landings for each size

| Length ¹ | Percent of landings | |
|-----------------------|---------------------|-------|
| | Scrod | Large |
| 33 cm. and under..... | 100.0 | |
| 36 cm..... | 99.7 | 0.3 |
| 39 cm..... | 99.2 | .8 |
| 42 cm..... | 97.5 | 2.5 |
| 45 cm..... | 86.6 | 13.4 |
| 48 cm..... | 51.3 | 48.7 |
| 51 cm..... | 14.8 | 85.2 |
| 54 cm..... | 2.3 | 97.7 |
| 57 cm..... | .4 | 99.6 |
| 60 cm..... | .2 | 99.8 |
| 63 cm..... | .1 | 99.9 |
| 66 cm. and over..... | | 100.0 |
| All sizes..... | 49.4 | 50.6 |

¹ Size groups by 3-cm. intervals.

DISCUSSION AND SUMMARY

1. Presented in this paper is an outline of a study of Georges Bank haddock and also details of landings for the years of 1931 to 1948. Pounds, numbers, and average weights of fish, and size compositions of landings are given for scrod, for large, and for total haddock. While these data are presented primarily as background for further studies, the averages and ranges are informative. The values presented, in our opinion, are as nearly complete a record of the quantities of Georges Bank haddock that were landed and sold as can be readily assembled. They are more nearly complete than values previously given (Schuck 1949), which represent only Georges Bank haddock landed at the ports of Boston, Gloucester, and New Bedford, Mass., and Portland, Maine.

2. The industry is most affected, not by the average or ordinary condition of the fishery, but by deviations from the normal, be it in terms of pounds of fish, of numbers of fish, of numbers of certain sizes as compared with previous years, or of a change in the seasonal cycle of the above. But, in order to measure deviations, it is first necessary to determine the norm from which they deviate. We can define the average year as follows: In the average year (during the period 1931-1948) there were 94,196,000 pounds of haddock (30,791,000 pounds of scrod and 63,405,000 pounds of large) landed from Georges Bank. The average weight of these fish was 2.55 pounds (1.69 for scrod, 3.40 for large) and 36,875,000 individual fish (18,214,000 scrod and 18,661,000 large) were landed. Of these numbers landed, there were practically none less than 27 centimeters (9.6 inches), and none more than 81 centimeters (32.1 inches) in length. The 45-centimeter (17.9-inch)

group contained the most fish and over 66 percent of all haddock landed were between the 42-centimeter (16.2-inch) group and the 54-centimeter (22.1-inch) group in length.

Also in the average year about 4,974,000 fish or 13.5 percent of the total number landed were smaller than the established minimum market size of 1½ pounds.

3. So far as subareas of Georges Bank are concerned, in the average year (1936 to 1948 only) the Northern Edge, though not the largest area, has been the largest producer, with 35 percent of the total poundage.

Percentages for scrod, large, and total haddock from the four areas are as follows:

| | Scrod | Large | Total haddock |
|---------------------|-------|-------|---------------|
| Northern Edge..... | 39.5 | 32.9 | 35.2 |
| Southeast Part..... | 26.3 | 23.3 | 24.4 |
| South Channel..... | 28.6 | 36.4 | 33.6 |
| Southwest Part..... | 5.6 | 7.4 | 6.8 |
| | 100.0 | 100.0 | 100.0 |

4. The seasonal landings, for the average year, are shown in table 36 by pounds, numbers, and average weights.

TABLE 36.—Seasonal average weights and quantities landed

| | Pounds of fish (thousands) | Number of fish (thousands) | Average weight per fish (pounds) |
|----------------|----------------------------|----------------------------|----------------------------------|
| Spring: | | | |
| Scrod..... | 5,273 | 2,819 | 1.871 |
| Large..... | 15,899 | 4,276 | 3.718 |
| Total..... | 21,172 | 7,095 | 2.984 |
| Summer: | | | |
| Scrod..... | 8,448 | 5,430 | 1.556 |
| Large..... | 20,359 | 6,444 | 3.163 |
| Total..... | 28,837 | 11,876 | 2.430 |
| Fall: | | | |
| Scrod..... | 12,147 | 7,064 | 1.719 |
| Large..... | 18,152 | 5,491 | 3.306 |
| Total..... | 30,299 | 12,555 | 2.413 |
| Winter: | | | |
| Scrod..... | 4,923 | 2,901 | 1.697 |
| Large..... | 8,965 | 2,449 | 3.661 |
| Total..... | 13,888 | 5,349 | 2.596 |
| Year: | | | |
| Scrod..... | 30,791 | 18,214 | 1.691 |
| Large..... | 63,405 | 18,660 | 3.398 |
| Total..... | 94,196 | 36,875 | 2.554 |

From table 36, we have computed the percent by weight and the percent by number for scrod, large, and total haddock of the year's landings. They are as follows:

| | By weight | By number |
|-----------------------|-----------|-----------|
| Scrod: | | |
| Spring..... | 17.1 | 15.5 |
| Summer..... | 27.4 | 29.8 |
| Fall..... | 39.5 | 38.8 |
| Winter..... | 16.0 | 15.9 |
| Total year..... | 100.0 | 100.0 |
| Large: | | |
| Spring..... | 25.1 | 22.9 |
| Summer..... | 32.2 | 34.6 |
| Fall..... | 28.6 | 29.4 |
| Winter..... | 14.1 | 13.1 |
| Total year..... | 100.0 | 100.0 |
| Total haddock: | | |
| Spring..... | 22.5 | 19.2 |
| Summer..... | 30.6 | 32.2 |
| Fall..... | 32.2 | 34.1 |
| Winter..... | 14.7 | 14.5 |
| Total year..... | 100.0 | 100.0 |

Landings of undersized haddock were greatest in the fall season, when 38 percent of the yearly average landings of undersized fish occurred. The summer season accounted for 30 percent, the winter season for 20 percent, and the spring season for the least quantity, 12 percent. Considering each season separately, the percentages of haddock landed that were undersized are as follows:

| | Percent undersized |
|-----------------|--------------------|
| Spring..... | 8.1 |
| Summer..... | 12.7 |
| Fall..... | 15.1 |
| Winter..... | 18.7 |
| Total year..... | 13.5 |

5. Having thus developed average values of important characteristics of the landings, each individual year can be evaluated by comparing it with these norms. For instance, considering 1934 (the poorest year of haddock production), we see that only 12,976,000 pounds of scrod as compared with the average of 30,791,000 pounds were landed; only 36,908,000 pounds of large haddock as compared with the average of 63,405,000; and only 49,884,000 pounds of all haddock as compared with the average of 94,196,000. Average weights for 1934 as compared to the average year were:

| | 1934 | Average year |
|--------------------|------|--------------|
| Scrod..... | 1.62 | 1.69 |
| Large..... | 3.26 | 3.40 |
| Total haddock..... | 2.58 | 2.55 |

The numbers of fish landed in 1934 as compared with 18-year averages were: scrod 8,024,000 (18,214,000), large 11,311,000 (18,661,000), total haddock 19,335,000 (36,875,000).

In addition to such yearly deviations, seasonal deviations for 1934 can be compared with average seasonal values, and subarea contributions can be evaluated in terms of average subarea contributions.

6. For a rapid evaluation of how each of the 18 years deviate in the more important characteristics from the average year, table 37 has been prepared. Shown are the percentages that the individual years are above or below the 18-year average;

pounds, numbers, and average weights are treated, for large, scrod and total haddock.

7. The data in this paper serve (1) as a record of the total landings of haddock from Georges Bank in terms of pounds, average weights, numbers and sizes of scrod, large, and total haddock, by seasons and years over the 18-year period, 1931 to 1948; and (2) as a basis for developing other data, among which will be the age composition of the landings; the size of various ages; year class contributions; and estimates of the relative size of the stock on the banks, of rates of decline of year classes, and of mortality rates.

TABLE 37.—Percentage deviations of quantities and average weights from the average year

| | 1931 | 1932 | 1933 | 1934 | 1935 | 1936 | 1937 | 1938 | 1939 | 1940 | 1941 | 1942 | 1943 | 1944 | 1945 | 1946 | 1947 | 1948 |
|----------------------------------|-------|------|-------|-------|-------|-------|------|-------|------|------|------|------|-------|-------|-------|-------|------|-------|
| Pounds: | | | | | | | | | | | | | | | | | | |
| Scrod..... | -43.2 | 2.3 | -38.9 | -57.9 | -17.1 | -2.7 | 1.0 | 29.6 | 40.1 | -3.9 | 74.6 | 60.7 | 21.8 | -50.5 | -62.1 | -31.4 | 32.8 | 44.7 |
| Large..... | 53.8 | 16.6 | -.9 | -41.8 | -15.4 | -14.7 | 1.5 | -17.2 | -3.2 | -.6 | 7.2 | -9.1 | -17.9 | 27.4 | 5.0 | 30.5 | 1.5 | -22.8 |
| Total..... | 22.1 | 11.9 | -13.3 | -47.0 | -15.9 | -10.8 | 1.3 | -1.9 | 11.0 | -1.6 | 29.2 | 13.7 | -4.9 | 1.9 | -16.9 | 10.3 | 11.8 | -.7 |
| Numbers: | | | | | | | | | | | | | | | | | | |
| Scrod..... | -39.4 | -3.5 | -35.7 | -55.9 | -15.4 | 1.2 | -2.4 | 30.5 | 38.1 | -9.9 | 77.6 | 53.8 | 17.2 | -52.8 | -59.3 | -29.4 | 34.4 | 50.8 |
| Large..... | 50.5 | 17.4 | 2.8 | -39.4 | -9.4 | -9.1 | .5 | -24.3 | .2 | -.6 | 1.4 | -7.6 | -13.8 | 31.5 | 2.5 | 31.4 | -7.3 | -26.5 |
| Total..... | 6.1 | 7.1 | -16.2 | -47.6 | -12.4 | -4.0 | -.9 | 2.8 | 18.9 | -5.2 | 39.0 | 22.7 | 1.5 | -10.1 | -28.0 | 1.4 | 13.3 | 11.6 |
| Average weights (pounds): | | | | | | | | | | | | | | | | | | |
| Scrod..... | -6.2 | 6.0 | -5.1 | -4.3 | -1.9 | -3.8 | 3.4 | -.7 | 1.4 | 6.7 | -1.7 | 4.5 | 3.9 | 4.8 | -6.9 | -2.9 | -1.2 | -4.0 |
| Large..... | 2.2 | -.7 | -3.6 | -4.0 | -6.6 | -6.2 | 1.0 | 9.4 | -3.3 | 0 | 5.7 | -1.7 | -4.7 | -3.2 | 2.5 | -.6 | 9.4 | 5.1 |
| Total..... | 15.1 | 4.5 | 3.4 | 1.0 | -4.0 | -7.1 | 2.3 | -4.5 | -5.7 | 3.7 | -7.0 | -7.4 | -6.3 | 13.4 | 15.4 | 8.8 | -1.4 | -11.1 |

LITERATURE CITED

HERRINGTON, WILLIAM C.

1933. Conservation of immature fish in otter trawling. *Trans. Amer. Fish. Soc.*, vol. 62 (1932), pp. 57-63.

1935. Modifications in gear to curtail the destruction of undersized fish in otter trawling. Bureau of Fisheries, U. S. Dept. Commerce, Investigational Report No. 24, 48 pp.

1936. Decline in haddock abundance on Georges Bank and a practical remedy. Bureau of Fisheries, U. S. Dept. Commerce, Fishery Circular No. 23, 22 pp.

1941. A crisis in the haddock fishery. Fish and Wildlife Service, U. S. Dept. Interior, Fishery Circular No. 4, 14 pp.

1944. Factors controlling population size. *Trans. Ninth North Amer. Wildlife Conf.*, 1944, pp. 250-263.

1948. Limiting factors in fish production. Some theories and an example. *Bulletin, Bingham Oceanographic Collection*, vol. 11, Article 4, pp. 229-283.

McKENZIE, R. A.

1946. The haddock fishery of grounds fished by Canadians. *Bull. 69, Fish. Res. Bd. Can.*, 1946, 30 pp.

NEEDLER, A. W. H.

1930. The migrations of haddock and the interrelationships of haddock populations in North American waters. *Contr. Canad. Biol. Fish., N. S.*, vol. 6, No. 10, 1930, pp. 243-313.

ROUNSEFELL, GEORGE A.

1942. Field experiments in selecting the most efficient tag for use in haddock studies. *Trans. Amer. Fish. Soc.*, vol. 71, 1941, pp. 228-235.

1948. Development of fishery statistics in the North Atlantic. U. S. Fish and Wildlife Service, Spec. Sci. Report No. 47, 18 pp.

RUSSELL, E. S.

1942. *The overfishing problem.* Cambridge Univ. Press, London, 1942, 130 pp.

SCHROEDER, WILLIAM C.

1942. Results of haddock tagging in the Gulf of Maine from 1923 to 1932. *Sears Foundation: Jour. Marine Res.*, vol. 5, No. 1, June 1942, pp. 1-19.

SCHUCK, HOWARD A.

1949. Relationship of catch to changes in population size of New England haddock. *Amer. Stat. Assn.: Biometrics*, vol. 5, No. 3, Sept. 1949, pp. 213-231.

SCHUCK, HOWARD A., and EDGAR L. ARNOLD, JR.

1951. Comparison of haddock from Georges and Browns Banks. U. S. Fish and Wildlife Service, *Fishery Bulletin* 67, vol. 52, pp. 177-185.