

STUDY PLAN FOR AVIAN EGG INJECTION STUDY

HUDSON RIVER NATURAL RESOURCE DAMAGE ASSESSMENT

HUDSON RIVER NATURAL RESOURCE TRUSTEES

STATE OF NEW YORK

U.S. DEPARTMENT OF COMMERCE

U.S. DEPARTMENT OF THE INTERIOR

PUBLIC RELEASE VERSION*

FINAL

MAY 12, 2006

Available from:

U.S. Department of Commerce

National Oceanic and Atmospheric Administration

Hudson River NRDA, Lead Administrative Trustee

Damage Assessment Center, N/ORR31

1305 East-West Highway, Rm 10219

Silver Spring, MD 20910-3281

**Names of certain individuals and affiliations have been removed to maintain confidentiality*



EXECUTIVE SUMMARY

Past and continuing discharges of polychlorinated biphenyls (PCBs) have contaminated the natural resources of the Hudson River. The Hudson River Natural Resource Trustees - New York State, the U.S. Department of Commerce, and the U.S. Department of the Interior - are conducting a natural resource damage assessment (NRDA) to assess and restore those natural resources injured by PCBs.

In 2002, the Trustees conducted an avian egg exposure preliminary investigation for the Hudson River. That preliminary investigation revealed that of the eleven avian species tested, the highest PCB levels were found in belted kingfisher and spotted sandpiper.

Based on the results of avian investigations conducted by the Trustees, and considering factors such as the life histories of various Hudson River avian species, avian toxicology, and the goals of the NRDA, the Trustees determined that it was appropriate to conduct further investigations focused on avian species. Pursuant to that determination and to the Hudson River NRDA Plan, the Trustees conducted a study of belted kingfisher, spotted sandpiper and tree swallow in 2004 and 2005. The Trustees further proposed conducting an avian egg injection study.

A Draft Study Plan for the egg injection work was peer reviewed and made available to the public for review and comment. All comments received on the Draft Study Plan, as part of the peer and public review process, have been considered. The Trustees evaluated peer and public comments and, where warranted, incorporated these comments in the Draft Study Plan to produce the Final Study Plan. In the remaining instances, public comments on the Draft Study Plan have been addressed by letters to the commenters, acknowledging receipt of comments and providing an initial response and noting that a more detailed Responsiveness Summary will be provided by the Trustees in the near future.

The Trustees will conduct an avian egg injection study of tree swallow and American kestrel in 2006 to evaluate whether specific avian species in the vicinity of the Hudson River are injured due to exposure to PCBs. Additionally, egg injection experiments will be conducted using chicken eggs to provide a point of reference for impacts observed in other species and in relation to effects levels identified in the toxicology literature for PCBs, dioxins, and other chemicals. Work on the species studied in 2006 or other species may continue in 2007.

The objective of the investigation is to evaluate the toxicity and adverse effects of embryonic exposure of multiple avian species to dose ranges of PCB 126 or a PCB mixture. This study will be used to evaluate whether avian reproduction and/or development is affected as a result of exposure to PCBs from the Hudson River.

Pursuant to the Hudson River NRDA Plan, the results of the work conducted pursuant to this Study Plan will be peer reviewed upon completion of the study, and the results then released to the public.

TABLE OF CONTENTS

1.0 BACKGROUND.	1
2.0 INTRODUCTION.	3
3.0 PURPOSE AND OBJECTIVE.	3
4.0 METHODS.	4
4.1 YEAR 1 (2006) AVIAN EGG INJECTION STUDY OF TREE SWALLOW, AMERICAN KESTREL, AND CHICKEN	4
4.2 YEAR 2 (2007) AVIAN EGG INJECTION STUDY	4
5.0 QUALITY ASSURANCE/QUALITY CONTROL.	5
6.0 SPECIAL PROVISIONS.	5
7.0 LITERATURE CITED.	6

APPENDIX A: FINAL WORK PLAN FOR TREE SWALLOW, AMERICAN KESTREL, AND CHICKEN EGG INJECTION STUDIES

APPENDIX B: EGG INJECTION DOSING MIXTURE

1.0 BACKGROUND

Past and continuing discharges of polychlorinated biphenyls (PCBs) have contaminated the natural resources of the Hudson River. The Hudson River Natural Resource Trustees - New York State, the U.S. Department of Commerce, and the U.S. Department of the Interior - are conducting a natural resource damage assessment (NRDA) to assess and restore those natural resources injured by PCBs (Hudson River Natural Resource Trustees 2002).

The Hudson River and surrounding area support more than 150 species of birds, including waterfowl, wading birds, shorebirds, songbirds, and rare species such as the bald eagle, peregrine falcon, and osprey (Andrle and Carroll, 1988). Birds are an integral part of the ecosystem and provide a number of important ecosystem services such as seed distribution, plant pollination, and insect control. Birds are also an important source of prey to other species. Birds may be exposed to PCBs through direct ingestion of contaminated water, sediment, and soil. A more important likely exposure pathway is their consumption of food items that contain PCBs derived from the Hudson River and its floodplain. PCB-contaminated food items linked to the river may include fish, amphibians, benthic invertebrates, adult insects that develop from aquatic larvae, plants growing in or near the river, and mammals that forage in the floodplain.

In 2002, the Trustees conducted an avian egg exposure preliminary investigation for the Hudson River. The investigation entailed collection of eggs, and subsequent analysis for PCBs, from six primary species (belted kingfisher (*Ceryle alcyon*), American robin (*Turdus migratorius*), Eastern phoebe (*Sayornis phoebe*), spotted sandpiper (*Actitis macularia*), red winged blackbird (*Agelaius phoeniceus*), and American woodcock (*Scolopax minor*)) and from five additional species (Eastern screech owl (*Otus asio*), common grackle (*Quiscalus quiscula*), northern rough-winged swallow (*Stelgidopteryx serripennis*), barn swallow (*Hirundo rustica*), and Eastern bluebird (*Sialia sialis*)) based on the opportunities for survey team members to locate the nests of these species. The geographic scope of the 2002 avian egg investigation was the Hudson River and its floodplains, from Hudson Falls to Lower Schodack Island, New York.

That preliminary investigation was undertaken by the Trustees to assist in determining the extent to which avian species in the Hudson River are contaminated with PCBs, to determine if additional pathway and injury assessment studies focused on avian species should be conducted as part of the Hudson River NRDA, and for potential use in the design of future studies to assess the health of Hudson River birds. The Trustees noted in the Hudson River NRDA Plan that, based on the results of the bird egg study, the Trustees would determine whether injury determination and quantification studies were warranted.

That preliminary investigation revealed that of the eleven avian species tested, the highest PCB levels were found in belted kingfisher and spotted sandpiper (Hudson River Natural Resource Trustees, 2003). Spotted sandpiper eggs contained a mean of 15 parts per million (ppm) PCBs (as total homologues, fresh weight basis). Of the eleven species tested, spotted sandpiper eggs exhibited the highest individual egg concentration of PCBs (56 ppm) as well as the highest average PCB concentration (15 ppm). Of the eleven species tested, belted kingfisher eggs exhibited the second highest individual egg concentration of PCBs (43 ppm).

Based on the results of avian investigations conducted by the Trustees, including the tree swallow (*Tachycineta bicolor*) work (McCarty and Secord 1999a, 1999b, Secord *et al.* 1999) and the 2002 avian egg preliminary investigation (Hudson River Natural Resource Trustees 2003), and input from a panel of avian experts, and considering factors such as the life histories of various Hudson River avian species, avian toxicology, and the goals of the NRDA, the Trustees determined that it was appropriate to conduct further investigations focused on avian species, and initially on belted kingfisher, spotted sandpiper, and tree swallow, to be started in the year 2004.

Pursuant to that determination and to the Hudson River NRDA Plan, the Trustees released in 2004 a "Study Plan for Year 2004 Avian Investigations for the Hudson River - Final, Public Release Version," dated June 15, 2004 (Hudson River Natural Resource Trustees, 2004). That Avian Injury Study Plan described the activities that constituted the Trustees' planned approach to conducting investigations of injury to avian species as part of the Hudson River NRDA.

As noted in the Avian Injury Study Plan, the Trustees planned to assess the following potential injuries to birds: reduced avian reproduction and overt external malformations. The Trustees planned to: (1) assess the relationship between contaminant concentrations in nest sample eggs and parameters of nest reproduction by application of appropriate statistical analysis of data to determine whether reproductive success of spotted sandpipers, tree swallows and belted kingfishers nesting on the Hudson River is negatively affected by PCB exposure; (2) assess the incidence of gross deformities in embryos or hatchlings; (3) assess organic contaminant accumulation rates in belted kingfisher chicks on the Hudson River; and (4) initiate an avian egg injection pilot study in 2004.

Pursuant to the Final Avian Injury Study Plan and a May 4, 2005 Modification to that Study Plan (Hudson River Natural Resource Trustees, 2005a), the U.S. Geological Survey (USGS) conducted a study of belted kingfisher, spotted sandpiper and tree swallow in 2004 and 2005. The USGS study was directed at items (1), (2) and (3) above. Trustee review of the data and results from the USGS study is ongoing.

Regarding item (4) above, the Trustees' Final Avian Injury Study Plan proposed a "pilot" study, a preliminary investigation focused on incubation of eggs of Hudson River avian species in 2004 with injection of PCBs into eggs of avian species of interest in a subsequent year. The Trustees subsequently determined that, considering preliminary work done by the Trustees and the literature on avian egg injection studies, it was not necessary to conduct a separate incubation-focused pilot study prior to initiating an avian egg injection study.

A Draft Study Plan for an avian egg injection experiment (Hudson River Natural Resource Trustees 2006) was developed, and that Draft Study Plan was peer reviewed and made available to the public for review and comment. All comments received on the Draft Study Plan, as part of the peer and public review process, have been considered. The Trustees evaluated peer and public comments and, where warranted, incorporated these comments in the Draft Study Plan to produce this Final Study Plan. In the remaining instances, public comments on the Draft Study Plan have been addressed by letters to the commenters, acknowledging receipt of comments and providing an initial response and noting that a more detailed Responsiveness Summary will be provided by the Trustees in the near future.

2.0 INTRODUCTION

Avian egg injection is a well-established technique to assess the effects of contaminants on a developing avian embryo (Allred and Strange 1977, Blankenship et al. 2003, Blomqvist et al. 2006, Boily et al. 2003, Bruggeman et al. 2003, Brunström 1986, Brunström et al. 1999, Brunström 1990, Brunström 1988, Brunström and Andersson 1988, de Roode et al. 2000, DeWitt et al. 2005a, DeWitt et al. 2005b, Drake et al. 2006, Fernie et al. 2005, Fox and Grasman 1999, Goff et al. 2005, Gould et al. 1997, Grasman and Whitacre 2001, Halldin et al. 2002, Heinz et al. 2006, Hoffman et al. 1998, Ivnitski et al. 2001, Janz and Bellward 1996, Jin et al. 2001, Katynski et al. 2004, Lim et al. 2005, Meneely and Wuttenbach 1989, Murvoll and Skaare 2005, Nosek et al. 1993, Ottinger et al. 2005, Powell et al. 1996a, Powell et al. 1996b, Powell et al. 1997a, Powell et al. 1997b, Powell et al. 1998, Stanton et al. 2003, U.S. Environmental Protection Agency 2001, Walker and Catron 2000, Walker et al. 1997, Wilhelms et al. 2006, Zhang et al. 2002, and Zhao et al. 1997).

To conduct an avian egg injection experiment, eggs are collected and brought into a laboratory where they are injected with the substance being tested. In avian egg injection experiments, various doses of a contaminant of concern (for example, PCBs in a vehicle or carrier solution) are typically injected into the yolk sac (for example, Hoffman et al. 1998), air cell (for example, Brunström and Andersson 1988, Fox and Grasman 1999), or albumen (for example, Nosek et al. 1993) of eggs. The eggs are then incubated in a laboratory and their development monitored. Measurement endpoints may include embryomortality, malformations, and hatching success. Measurement endpoints may also extend to hatchlings, for which chick growth and development, for example, may be measured.

Results reported in the literature of injecting contaminants, such as PCBs, into avian eggs include embryomortality and malformation. Death, including embryomortality, for example, and physical deformation, such as external malformation, skeletal deformities, and organ and soft tissue malformation, are injuries pursuant to the regulations written by the U.S. Department of the Interior contained in Title 43 of the Code of Federal Regulations Part 11, Natural Resource Damage Assessment (the "DOI NRDA Regulations"), and would be relevant to determining injury as part of the NRDA.

3.0 PURPOSE AND OBJECTIVE

The Trustees will conduct an avian egg injection study of tree swallow and American kestrel in 2006 to evaluate whether specific avian species in the vicinity of the Hudson River are injured due to exposure to PCBs. Additionally, egg injection experiments will be conducted using chicken eggs to provide a point of reference for impacts observed in other species and in relation to effects levels identified in the toxicology literature for PCBs, dioxins, and other chemicals. Work on the species studied in 2006 or other species may continue in 2007.

The objective of the investigation is to evaluate the toxicity and adverse effects of embryonic exposure of multiple avian species to dose ranges of PCB 126 or a PCB mixture. The PCB mixture is made up of individual PCB congeners and fits a similar profile to the mixture of PCBs occurring in the eggs of birds nesting in the Upper Hudson River. This study will be used to evaluate whether avian reproduction and/or development is affected as a result of exposure to PCBs from the Hudson River. The work will inform the Trustees regarding injury to avian resources and guide their future efforts to identify pathway and specific injuries to birds from PCBs, determine causation, and scale restoration, as defined in the DOI NRDA Regulations. The work will be used to identify and evaluate the type(s) of injury(ies), if any, that PCBs are causing to Hudson River birds at the embryonic level. This work will also be used to help determine whether future studies will be performed, and if so, to help in their design.

4.0 METHODS

4.1 YEAR 1 (2006) AVIAN EGG INJECTION STUDY OF TREE SWALLOW, AMERICAN KESTREL, AND CHICKEN

The attached work plan entitled, "Final Work Plan for Tree Swallow, American Kestrel, and Chicken Egg Injection Studies" (Appendix A) describes the avian egg injection investigation that the Trustees will implement to evaluate whether specific avian species in the vicinity of the Hudson River are injured due to exposure to PCBs. The attached work plan includes information regarding the experimental design, Quality Assurance/Quality Control, and Standard Operating Procedures that will be used in the study.

In Year 1 (2006), the Trustees will focus on injection of test PCBs and incubation methods for eggs from chosen species. If injection and incubation methods are successful, tissues will be collected for analysis, providing initial datasets for these species. Year 1 (2006) work will focus on those species with eggs that are more easily obtainable than others.

In 2006, work will be conducted on tree swallow and American kestrel (*Falco sparverius*). These species have been selected because they represent different positions in the ecosystem, breed in the Hudson River basin, and may be sensitive to PCB exposure. Eggs of American kestrel will be obtained from Patuxent Wildlife Research Center, Maryland. Eggs of tree swallows will be obtained from Patuxent National Wildlife Refuge, Maryland, from a breeding colony on Great Sacandaga Lake, New York, and from the Hudson River, New York. Additionally, egg injection experiments will be conducted using chicken (*Gallus domesticus*) eggs to provide a point of reference for impacts observed in other species and in relation to effects levels identified in the toxicology literature for PCBs, dioxins and other chemicals. Work will begin in Spring 2006.

Trials will be conducted using select PCBs administered early in embryonic development. As described in the "Final Work Plan for Tree Swallow, American Kestrel, and Chicken Egg Injection Studies," eggs will be injected with PCBs (PCB 126 or a PCB mixture reflective of chemical exposures in the Hudson River region). A vehicle control and an untreated group of eggs will be included. The PCBs to be injected into the eggs have been selected by the Trustees based on existing contaminants data from Hudson River biota and other relevant factors. Appendix B provides information on the PCB congener mixture to be used in the egg injections.

There will be a separate experiment conducted for each species. The timing of each experiment will depend on the availability of eggs for that species. In the case of chickens, for example, eggs will be available over much of the year, so those experiments will be scheduled around the work with the other species.

The "Final Work Plan for Tree Swallow, American Kestrel, and Chicken Egg Injection Studies" (Appendix A) notes the endpoints to be assessed and the methods that will be used.

4.2 YEAR 2 (2007) AVIAN EGG INJECTION STUDY

Year 1 work focuses on injection of test PCBs and incubation methods for eggs from tree swallow, American kestrel and chicken. These studies are projected to continue into a second year to allow further development of injection and incubation protocols for eggs from wild species and, in some cases, to produce larger sample sizes. Work on the species from 2006 or other species may be conducted in 2007. Work in 2007 will be conducted pursuant to a Study Plan Amendment for Year 2007.

5.0 QUALITY ASSURANCE/QUALITY CONTROL

This study is being conducted in accordance with the Quality Assurance Management Plan for the Hudson River NRDA (Hudson River Natural Resources Trustees, 2005b).

Strict chain-of-custody procedures will be used throughout the study. All samples collected under this Study Plan will be maintained under chain-of-custody upon collection, and through processing, storage and shipment to the testing laboratory, analytical laboratory or archive facility.

Analysis will be by appropriate methods approved by the Trustees. Analytes may include congener-specific PCBs, including the non-*ortho* congeners, polychlorinated dibenzo-p-dioxins (PCDDs), polychlorinated dibenzofurans (PCDFs), polybrominated diphenyl ethers (PBDEs), organochlorine pesticides, and metals, as determined appropriate by the Trustees.

In order to minimize analytical costs, and reduce the overall cost associated with the project, the Trustees may conduct the chemical analyses in stages, using initial work to inform subsequent decisions regarding which analyses to conduct on which samples.

The laboratories performing analytical work will be contracted to follow the Trustees' Analytical Quality Assurance Plan for the Hudson River NRDA (Hudson River Natural Resource Trustees 2005b). Laboratories will provide fully documented data packages which will enable data validation to be performed based on the criteria provided in the Analytical Quality Assurance Plan for the Hudson River NRDA, applicable laboratory Standard Operating Procedures, and the U.S. Environmental Protection Agency guidelines (1999).

6.0 SPECIAL PROVISIONS

Any necessary collection permits, such as those from New York State or Maryland where eggs will be collected, or from the U.S. Fish and Wildlife Service, will be obtained.

7.0 LITERATURE CITED

- Allred, P. M. and J. R. Strange. 1977. The effects of 2,4,5-trichlorophenoxyacetic acid and 2,3,7,8-tetrachlorodibenzo-p-dioxin on developing chicken embryos. *Arch. Environ. Contam. Toxicol.* 6: 483-489.
- Andrle, R.F. and J.R. Carroll. 1988. *The Atlas of Breeding Birds in New York State*. Cornell Univ. Press. 548 pp.
- Blankenship, A. L., K. Hilscherova, M. Nie, K. K. Coady, S. A. Villalobos, K. Kannan, D. C. Powell, S. J. Bursian, and J. P. Giesy. 2003. Mechanisms of TCDD-induced abnormalities and embryo lethality in white leghorn chickens. *Comp. Biochem. Physiol., C: Comp. Pharmacol. Toxicol.* 136: 47-62.
- Blomqvist, A., C. Berg, L. Holm, I. Brandt, Y. Ridderstråle, and B. Brunström. 2006. Defective reproductive organ morphology and function in domestic rooster embryonically exposed to o,p'-DDT or ethynylestradiol. *Biol. Reprod.* 74: 481-486.
- Boily, M. H., A. Ndayibagira, and P. A. Spear. 2003. Retinoids, LRAT and REH activities in eggs of Japanese quail following maternal and *in ovo* exposures to 3,3',4,4'-tetrachlorobiphenyl. *Ecotoxicology* 12: 9-21.
- Bruggeman, V., Q. Swennen, B. De Ketelaere, O. Onagbesan, K. Tona, and E. Decuyper. 2003. Embryonic exposure to 2,3,7,8-tetrachlorodibenzo-p-dioxin in chickens: effects of dose and embryonic stage on hatchability and growth. *Comparative Biochemistry and Physiology, C: Comparative Pharmacology and Toxicology* 136(1): 17-28.
- Brunström, B. 1986. Differences in sensitivity of some avian species to the embryotoxicity of a PCB, 3,3',4,4'-tetrachlorobiphenyl, injected into the eggs. *Environ. Pollut. (Series A)* 42: 37-45.
- Brunström, B. and L. Andersson. 1988. Toxicity and 7-ethoxyresorufin-O-deethylase-inducing potency of coplanar polychlorinated biphenyls (PCBs) in chick embryos. *Arch. Toxicol.* 62: 263-266.
- Brunström, B., D. Broman, and C. Naf. 1999. Embryotoxicity of polycyclic aromatic hydrocarbons (PAHs) in three domestic avian species, and of PAHs and coplanar polychlorinated biphenyls in the common eider. *Environ. Pollut.* 67: 133-143.
- Brunström, B. 1990. Mono-ortho-chlorinated chlorobiphenyls: toxicity and induction of 7-ethoxyresorufin O-deethylase (EROD) activity in chick embryos. *Arch. Toxicol.* 64: 188-192.
- Brunström, B. 1988. Sensitivity of embryos from duck, goose, herring gull and various chicken breeds to 3,3',4,4'-tetrachlorobiphenyl. *Poult. Sci.* 67: 52-57.
- de Roode, D. F., L. Balk, J. H. Koeman, and A. T. C. Bosveld. 2000. Development of a bioassay to test the possible role of thiamine disturbance as a mechanism behind pollution-induced reproductive failure in birds. *Arch. Environ. Contam. Toxicol.* 39: 386-391.
- DeWitt, J. C., E. B. Meyer, and D. S. Henshel. 2005a. Environmental toxicity studies using chickens as surrogates for wildlife: effects of vehicle volume. *Arch. Environ. Contam. Toxicol.* 48: 260-269.

- DeWitt, J. C., E. B. Meyer, and D. S. Henshel. 2005b. Environmental toxicity studies using chickens as surrogates for wildlife: effects of injection day. *Arch. Environ. Contam. Toxicol.* 48: 270-277.
- Drake, V. J., S. L. Koprowski, J. W. Lough, and S. M. Smith. 2006. Gastrulating chick embryo as a model for evaluating teratogenicity: a comparison of three approaches. *Birth Defects Research. Part A, Clinical and Molecular Teratology* 76: 66-71.
- Fernie, K. J., J. L. Shutt, G. Mayne, D. Hoffman, R. J. Letcher, K. G. Drouillard, and I. J. Ritchie. 2005. Exposure to polybrominated diphenyl ethers (PBDEs): changes in thyroid, vitamin A, glutathione homeostasis, and oxidative stress in American kestrels (*Falco sparverius*). *Toxicological Sciences* 88: 375-383.
- Fox, L. L. and K. A. Grasman. 1999. Effects of PCB 126 on primary immune organ development in chicken embryos. *Journal of Toxicology and Environmental Health, Part A* 58: 233-244.
- Goff, K. F., B. E. Hull, and K. A. Grasman. 2005. Effects of PCB 126 on primary immune organs and thymocyte apoptosis in chicken embryos. *Journal of Toxicology and Environmental Health, Part A* 68: 485-500.
- Gould, J. C., K. R. Cooper, and C. G. Scanes. 1997. Effects of three polychlorinated biphenyl mixtures and three specific congeners on growth and circulating growth-related hormones. *Journal of Comparative Endocrinology* 106: 221-230.
- Grasman, K. A. and L. L. Whitacre. 2001. Effects of PCB 126 on thymocyte surface marker expression and immune organ development in chicken embryos. *Journal of Toxicology and Environmental Health, Part A* 62: 191-206.
- Halldin, K., L. Holm, Y. Ridderstråle, and B. Brunström. 2002. Reproductive impairment in Japanese quail (*Coturnix japonica*) after in ovo exposure to o,p'-DDT. *Arch. Toxicol.* 77: 116-122.
- Heinz, G. H., D. J. Hoffman, S. L. Kondrad, and C. A. Erwin. 2006. Factors affecting the toxicity of methylmercury injected into eggs. *Arch. Environ. Contam. Toxicol.* 50: 264-279.
- Hoffman, D. J., M. J. Melancon, P. N. Klein, J. D. Eisemann, and J. W. Spann. 1998. Comparative developmental toxicity of planar polychlorinated biphenyl congeners in chickens, American kestrels, and common terns. *Environ. Toxicol. Chem.* 17: 747-757.
- Hudson River Natural Resource Trustees. 2002. Hudson River Natural Resource Damage Assessment Plan. September 2002. U.S. Department of Commerce, Silver Spring, MD.
- Hudson River Natural Resource Trustees. 2003. Avian Egg Database. Version 2.1. U.S. Department of Commerce, Silver Spring, MD.
- Hudson River Natural Resource Trustees. 2004. Study Plan for Year 2004 Avian Investigations for the Hudson River. Final. Public Release Version. June 15, 2004. U.S. Department of Commerce, Silver Spring MD.
- Hudson River Natural Resource Trustees. 2005a. Modification to Study Plan for Avian Investigations for the Hudson River. USGS Study Plan Amendment for 2005. Final. Public Release Version. May 4, 2005. U.S. Department of Commerce, Silver Spring, MD.
- Hudson River Natural Resource Trustees. 2005b. Analytical Quality Assurance Plan for the Hudson River Natural Resource Damage Assessment. Final. Public Release Version. September 1, 2005. Version 2.0. U.S. Department of Commerce, Silver Spring, MD.

- Hudson River Natural Resource Trustees. 2006. Study Plan for Avian Egg Injection Study. Hudson River Natural Resource Damage Assessment. Draft for Public Review and Comment. February 14, 2006. U.S. Department of Commerce, Silver Spring, MD.
- Ivnitski, I., R. Elmaoued, and M. K. Walker. 2001. 2,3,7,8-tetrachlorodibenzo-p-dioxin (TCDD) inhibition of coronary development is preceded by a decrease in myocyte proliferation and an increase in cardiac apoptosis. *Teratology* 64: 201-212.
- Janz, D. M. and G. D. Bellward. 1996. In ovo 2,3,7,8-tetrachlorodibenzo-p-dioxin exposure in three avian species. 1. Effects on thyroid hormones and growth during the perinatal period. *Toxicol. Appl. Pharmacol.* 139: 281-291.
- Jin, X., S. W. Kennedy, T. DiMuccio, and T. W. Moon. 2001. Role of oxidative stress and antioxidant defense in 3,3',4,4',5-pentachlorobiphenyl-induced toxicity and species-differential sensitivity in chicken and duck embryos. *Toxicol. Appl. Pharmacol.* 172: 241-248.
- Katynski, A. L., M. M. Vijayan, S. W. Kennedy, and T. W. Moon. 2004. 3,3',4,4',5-pentachlorobiphenyl (PCB 126) impacts hepatic lipid peroxidation, membrane fluidity, and beta-androreceptor kinetics in chick embryos. *Comp. Biochem. Physiol., C: Comp. Pharmacol. Toxicol.* 137: 81-83.
- Lim, J., J. DeWitt, J. B. I. Watkins, and D. S. Henshel. 2005. Analysis of TCDD-induced oxidative stress in chicken liver during development. Society of Environmental Toxicology and Chemistry, 26th Annual Meeting, November 13-17, 2005, Baltimore, Maryland, USA.
- McCarty, J.P. and A.L. Secord. 1999a. Reproductive Ecology of Tree Swallows (*Tachycineta bicolor*) With High Levels of Polychlorinated Biphenyl Contamination. *Environ. Toxicol. Chem.* 18:1433-1439.
- McCarty, J.P. and A.L. Secord. 1999b. Nest-Building Behavior in PCB-Contaminated Tree Swallows. *The Auk* 116:55-63.
- Meneely, G. A. and C. R. Wuttenbach. 1989. Effects of the organophosphate insecticides diazinon and parathion on bobwhite quail embryos: skeletal defects and acetylcholinesterase activity. *J. Exp. Zool.* 252: 60-70.
- Murvoll, K. M., B. M. Jenssen, and J. U. Skaare. 2005. Effects of polybrominated diphenyl ether (PBDE-99) on vitamin status in domestic ducks (*Anas platyrhynchos*) hatchlings. *Journal of Toxicology and Environmental Health, Part A* 68: 515-533.
- Nosek, J. A., J. R. Sullivan, S. R. Craven, A. Gendron-Fitzpatrick, and R. E. Peterson. 1993. Embryotoxicity of 2,3,7,8-tetrachlorodibenzo-p-dioxin in the ring-necked pheasant. *Environ. Toxicol. Chem.* 12: 1215-1222.
- Ottinger, M. A., J. M. Wu, J. L. Hazelton, M. Abdelnabi, N. Thompson, M. J. Quinn, Jr., D. Donogue, F. Schenck, M. Ruscio, J. Beavers, and M. Jaber. 2005. Assessing the consequences of the pesticide methoxychlor: neuroendocrine and behavioral measures as indicators of biological impact of an estrogenic environmental chemical. *Brain Res. Bull.* 65: 199-209.
- Powell, D. C., R. J. Aulerich, J. C. Meadows, D. E. Tillitt, J. P. Giesy, K. L. Stromborg, and S. J. Bursian. 1996a. Effects of 3,3',4,4',5-pentachlorobiphenyl (PCB 126) and 2,3,7,8-tetrachlorodibenzo-p-dioxin (TCDD) injected into the yolks of chicken (*Gallus domesticus*) eggs prior to incubation. *Arch. Environ. Contam. Toxicol.* 31: 404-409.
- Powell, D. C., R. J. Aulerich, K. L. Stromborg, and S. J. Bursian. 1996b. Effects of 3,3',4,4'-tetrachlorobiphenyl, 2,3,3',4,4'-pentachlorobiphenyl, and 3,3',4,4',5-pentachlorobiphenyl on the developing chicken embryo when injected prior to incubation. *J. Toxicol. Environ. Health* 49: 319-338.

- Powell, D. C., R. J. Aulerich, J. C. Meadows, D. E. Tillitt, J. F. Powell, J. C. Restum, K. L. Stromborg, J. P. Giesy, and S. J. Bursian. 1997a. Effects of 3,3',4,4',5-pentachlorobiphenyl (PCB 126), 2,3,7,8-tetrachlorodibenzo-p-dioxin (TCDD), or an extract derived from field-collected cormorant eggs injected into double-crested cormorant (*Phalacrocorax auritus*) eggs. Environ. Toxicol. Chem. 16: 1450-1455.
- Powell, D. C., R. J. Aulerich, J. C. Meadows, D. E. Tillitt, K. L. Stromborg, T. J. Kubiak, J. P. Giesy, and S. J. Bursian. 1997b. Organochlorine contaminants in double-crested cormorant eggs from Green Bay, Wisconsin: II. Effects of an extract derived from cormorant eggs on the chicken embryo. Arch. Environ. Contam. Toxicol. 32: 316-322.
- Powell, D. C., R. J. Aulerich, J. C. Meadows, D. E. Tillitt, M. E. Kelly, K. L. Stromborg, M. J. Melancon, S. D. Fitzgerald, and S. J. Bursian. 1998. Effects of 3,3',4,4',5-pentachlorobiphenyl (PCB 126), 2,3,7,8-tetrachlorodibenzo-p-dioxin (TCDD) injected into the yolks of double-crested cormorant (*Phalacrocorax auritus*) eggs prior to incubation. Environ. Toxicol. Chem. 17: 2035-2040.
- Secord, A.L., J.P. McCarty, K.R. Echols, J.C. Meadows, R.W. Gale, and D.E. Tillitt. 1999. Polychlorinated Biphenyls and 2,3,7,8-tetrachlorodibenzo-p-dioxin equivalents in tree swallows from the Upper Hudson River, New York State, USA. Environ. Toxicol. Chem. 18:2519-2525.
- Stanton, B., J. DeWitt, D. Henshel, S. Watkins, and B. Lasley. 2003. Fatty acid metabolism in neonatal chickens (*Gallus domesticus*) treated with 2,3,7,8-tetrachlorodibenzo-p-dioxin (TCDD) or 3,3',4,4',5-pentachlorobiphenyl (PCB 126) in ovo. Comp. Biochem. Physiol., C: Comp. Pharmacol. Toxicol. 136: 73-84.
- U.S. Environmental Protection Agency. 1999. USEPA Contract Laboratory Program National Functional Guidelines for Organic Data Review. Office of Emergency and Remedial Response, Washington, D.C. EPA540/R-99/008. 118 pp.
- U.S. Environmental Protection Agency. 2001. Analyses of Laboratory and Field Studies of Reproductive Toxicity in Birds Exposed to Dioxin-Like Compounds for Use in Ecological Risk Assessment. Office of Research and Development, National Center for Environmental Assessment, Cincinnati, OH. EPA/600/R-03/114F. 39 pp plus appendices.
- Walker, M. K. and T. F. Catron. 2000. Characterization of cardiotoxicity induced by 2,3,7,8-tetrachlorodibenzo-p-dioxin and related chemicals during early chick embryo development. Toxicol. Appl. Pharmacol. 167: 210-221.
- Walker, M. K., R. S. Pollenz, and S. M. Smith. 1997. Expression of the aryl hydrocarbon (AhR) and AhR nuclear translocator during chick cardiogenesis is consistent with 2,3,7,8-tetrachlorodibenzo-p-dioxin-induced heart defects. Toxicol. Appl. Pharmacol. 143: 407-419.
- Wilhelms, K. W., K. F. Fitzpatrick, C. G. Scanes, and L. L. Anderson. 2006. In ovo exposure to a triazine herbicide: effects of atrazine on circulating reproductive hormones and gonadal histology in young Japanese quail. Arch. Environ. Contam. Toxicol. 51: 117-122.
- Zhang, C., C. Fang, L. Lui, G. Xia, and H. Qiao. 2002. Disrupting effects of polychlorinated biphenyls on gonadal development and reproductive functions in chickens. J. Environ. Sci. Health, Part A 37: 509-519.
- Zhao, F., K. Mayura, N. Kocurek, J. F. Edwards, L. F. Kubena, S. H. Safe, and T. D. Phillips. 1997. Inhibition of 3,3',4,4',5-pentachlorobiphenyl-induced chicken embryotoxicity by 2,2',4,4',5,5'-hexachlorobiphenyl. Fundam. Appl. Toxicol. 35: 1-8.

APPENDIX A

FINAL WORK PLAN FOR TREE SWALLOW, AMERICAN KESTREL, AND CHICKEN EGG INJECTION STUDIES

FINAL WORK PLAN
FOR
TREE SWALLOW,
AMERICAN KESTREL,
AND CHICKEN
EGG INJECTION STUDIES

HUDSON RIVER NATURAL RESOURCE DAMAGE ASSESSMENT

12th May 2006

Principal Investigator

Co-Principal Investigator

Quality Assurance Coordinator

INVESTIGATION TEAM ACKNOWLEDGEMENT OF WORK PLAN REVIEW AND COMPLIANCE

By my signature, I acknowledge that I have read this Work Plan and understand it, and will comply with it in performing this work.

Name (printed): _____ Name (printed): _____

Signature: _____ Signature: _____

Date: _____ Date: _____

Title: _____ Title: _____

Name (printed): _____ Name (printed): _____

Signature: _____ Signature: _____

Date: _____ Date: _____

Title: _____ Title: _____

Name (printed): _____ Name (printed): _____

Signature: _____ Signature: _____

Date: _____ Date: _____

Title: _____ Title: _____

Name (printed): _____ Name (printed): _____

Signature: _____ Signature: _____

Date: _____ Date: _____

Title: _____ Title: _____

Name (printed): _____ Name (printed): _____

Signature: _____ Signature: _____

Date: _____ Date: _____

Title: _____ Title: _____

ABBREVIATIONS

ED	Embryonic Day
EIA	Enzyme Immuno Assay
EROD	Ethoxyresorufin-o-dealkylase
GnRH-I	Gonadotropin Releasing Hormone-I
HPLC	High Pressure Liquid Chromatography
PCB	Polychlorinated Biphenyl
RIA	Radioimmunoassay
ELISA	Enzyme Linked ImmunoSorbent Assay

TABLE OF CONTENTS

1.	INTRODUCTION	6
2.	WORK PLAN.....	6
2.1	Animals.....	6
2.2	Treatments.....	7
2.3	End Points.....	7
2.4	Experimental Design.....	7
2.4.1	Egg Collection	7
2.4.2	Egg Incubation and Injection.....	8
2.4.3	Dosing Solutions.....	9
2.4.4	Egg Hatching and Tissue Sampling.....	9
2.4.5	Biological Tissue Analysis	10
2.4.6	Statistical Analyses.....	10
3	QUALITY ASSURANCE/QUALITY CONTROL.....	14
3.1	Data Quality Objectives, Indicators, and Assessment.....	14
3.1.1	Overview.....	14
3.1.2	Project Management	15
3.2	Data Generation and Acquisition.....	16
3.2.1	Data Quality Objectives.....	16
3.2.2	Study Documentation.....	16
3.2.3	Sample Identification Procedures	17
3.3	Assessment and Oversight.....	17
3.4	Data Validation and Usability.....	18
4	PRINCIPAL INVESTIGATORS	20
5.	LITERATURE CONSULTED.....	20
6.	STANDARD OPERATING PROCEDURES.....	21
6.1	Recording and Handling Data for Avian Egg Injection Study	21
6.2	Field Collection of Tree Swallow Eggs for Avian Egg Injection Study, Hudson River NRDA	22
6.3	Removal of Avian Egg Contents for Contaminants Analysis	25
6.4	Egg Injection and Incubation Procedure for Tree Swallow (<i>Tachycineta bicolor</i>) Eggs ..	30
6.5	Egg Injection and Incubation Procedure for American Kestrel (<i>Falco sparverius</i>).....	37
6.6	Egg Injection Procedure for Chicken (<i>Gallus domesticus</i>) Eggs.....	40
6.7	Necropsy of Hatchling Birds	41
6.8	Histological Analysis of Avian Embryo Tissue: Gonads and Bursa of Fabricius.....	44
6.9	Ethoxyresorufin-o-dealkylase (EROD) Assay For CYP450 Activity In Liver Samples..	47
6.10	High Pressure Liquid Chromatography (HPLC):Monoamine Analysis.....	51
6.11	GnRH-I ELISA	55
6.12	Protein Assay	58
6.13	Extraction Of Steroid Hormones From Avian Feces.....	59
6.14	Radioimmunoassay of Androgens In Avian Serum or Fecal Extract.....	62
6.15	Radioimmunoassay of Estradiol In Avian Serum or Fecal Extract.....	68
6.16	Digestion Of Thyroid Gland And Radioimmunoassay of Thyroid Hormones.....	72
6.17	Analysis of Aromatase Activity in Hypothalamic Tissue.....	74

6.19 Tissue Storage, RNA Isolation And Microarray Analysis..... 77

6.20 Field Collection of Tree Swallow Eggs from Great Sacandaga Lake, New York for
Avian Egg Injection Study, Hudson River NRDA 78

6.21 Field Collection of Tree Swallow Eggs from Upper Hudson River, New York for Injury
Assessment Hudson River NRDA 81

1. INTRODUCTION

This work plan is for assessment of the biological impacts of selected chemicals of concern for the Hudson River assessment area, using an embryo bioassay on sentinel avian species for this region.

These experiments are projected over two years to allow development of injection and incubation protocols for eggs from wild species and to produce statistically useful sample sizes. This work plan defines work to be done in year one with three species for which we have identified egg sources: American kestrel (*Falco sparverius*), tree swallow (*Tachycineta bicolor*) and domestic chicken (*Gallus domesticus*). In addition to ready egg sources being available, the two wild species have been chosen because they represent a range of positions in the ecosystem and individuals from each species can be found for some or all of the year in the Hudson River basin. An experiment with chickens will be conducted to round out a species sensitivity comparison.

Our objective is to use egg injections as a method for evaluating the toxicity of PCB 126 and an environmentally realistic mixture of PCB congeners on the embryos of kestrels, tree swallows and chickens.

Based on data from other studies on the consequences of endocrine disrupting chemical exposure in birds, we expect sensitive endpoints to include 1) viability—embryo survival, pipping, hatching success, gross abnormalities, 2) gonadal and thyroid gland morphology, 3) endocrine measures—including hormone levels, 4) status of neural systems important in metabolic and reproductive axes function—hypothalamic neurotransmitters, aromatase enzyme, and gonadotropin releasing hormone (GnRH-I), 5) Cytochrome P450 enzyme activity and 6) Endocrine Disrupting Chemical (EDC) induced abnormalities, such as gonadal, biochemical and hormonal end points that are evident in the hatchling. Therefore, we will include these measurement end points in an embryo testing protocol that we have developed in other studies. These trials will be conducted by administering the PCB solutions early in embryonic development (before gonadal differentiation). Samples will be taken at hatch for measurement of selected end points known to be steroid responsive and sexually dimorphic. It is expected that endocrine active substances will affect these end points, thereby making treated individuals less sexually dimorphic.

2. WORK PLAN

2.1 Animals

We will receive eggs from two species in the 2006 breeding season: American kestrel (*Falco sparverius*) and tree swallow (*Tachycineta bicolor*). Kestrel eggs will be from the captive colony maintained at Patuxent National Wildlife Refuge (NWR)/Research Center (PWRC). Eggs from tree swallows will be collected from nest boxes located on the Patuxent National Wildlife Refuge and from Great Sacandaga Lake in New York state. In addition, chicken eggs

will be acquired in the Summer and (or) Fall from a commercial supplier such as CBT Farms, MD. Protocols for egg incubation and tissue sampling will be reviewed by appropriate Animal Care and Use Committees.

2.2 Treatments

This experiment will evaluate the effects of exposure to a dose range of PCB 126 in tree swallows and a PCB mixture in kestrels and chickens. The PCBs will be injected into eggs early in incubation, and tissues will be sampled from hatchlings. The time of egg injection will be at approximately 18% of incubation.

2.3 End Points

- Embryo survival/embryo mortality
- Gross morphology—including abnormalities
- Organ weights for liver, heart, brain, thyroid, thymus, bursa, spleen, and gonads
- Gonad histology and morphology of the reproductive tract
- Biochemical analyses
 - Liver CYP4501A
 - Brain monoamines/aromatase/GnRH-I (gonadotropin releasing hormone-I)
 - Thyroid hormones: T3 and T4 in thyroid gland; serum samples will be archived for T3 and T4 analysis if deemed necessary
 - Estradiol and androgen concentrations in fecal samples
- Bursal morphology
- Cardiac histology*
- Microarray analysis of RNA from liver and brain samples

* Hearts will be dissected and preserved for histological analysis.

Tissues will be collected for all biochemistries and histology but analyses will be made as deemed appropriate based on sample sizes.

2.4 Experimental Design

2.4.1 Egg Collection

Eggs will be collected under permits from Kathryn Jahn, USFWS. Eggs will be assigned to treatment groups: untreated, vehicle injected or PCB injected on the day of injection. Assignment to treatment group will be made under guidance of our statistical consultant. Since we can only estimate the number of eggs available for the study, the statistician will provide guidance on treatment group assignment with consideration of sample size, statistical power, sampling day, eggs per breeding pair, and estimated median lethal dose. The goal will be to maximize the number of eggs per independent parent within each treatment group and to attain data demonstrating a normal distribution. We will prioritize treatment groups based on data from other avian studies and we will evaluate which treatment groups can be eliminated to maximize sample size and dose response. Since we expect to obtain hundreds of eggs from tree

swallows, but only two dozen kestrel eggs we shall achieve a broader number of treatments for tree swallows and only a control and two or three PCB exposed groups for kestrels. A subset of tree swallow eggs will be assigned to contaminants analysis to verify the eggs are suitable as a 'clean' source. At a minimum, 2 eggs from each of three nests but potentially as many as 2 eggs from each of 10 nests will be kept for contaminants analyses. Because the number of eggs collected from PWRC may be limited, additional eggs will be assigned to contaminant analysis from eggs that fail to develop from the un-injected control group. Since the sources of kestrel eggs are birds that are fed a controlled diet, and because the number of eggs we will attain is limited, we will not submit fresh eggs for contaminants analysis but will retain yolks from hatchlings for analyses of experimental exposure (and background contamination from untreated eggs). Furthermore, we will keep infertile eggs and eggs containing dead embryos for contaminant analyses from the tree swallows and kestrels; this will substantially increase the amount of eggs available for analyses. There are also historical data available for Patuxent tree swallow contaminant burden available for consideration.

Tree swallow eggs will be collected from nests that will be monitored daily. We will follow the collection practices of Dr. Chris Custer, United States Geological Survey (USGS), in which the monitored nests will be observed daily for eggs, which are laid at one day intervals. When the fifth egg is laid, then two eggs are collected; if another egg is laid on the following day, it is possible to also collect that egg. In this way, the female should not abandon the nest.

We will also sample tissues from fifteen or more tree swallow embryos collected from the Upper Hudson River area.

The kestrels are housed in pairs at Patuxent Wildlife Center; pairs are monitored daily. Egg collection will be conducted according to husbandry practices for these colonies.

Standard Operating Procedure (SOP): "Field Collection of Avian Eggs for Avian Egg Injection Study Hudson River NRDA", "Removal of Avian Egg Contents for Contaminants Analysis Hudson River NRDA", "Field Collection of Tree Swallow Eggs from Great Sacandaga Lake, New York for Avian Egg Injection Study", "Field Collection of Tree Swallow Eggs from Upper Hudson River, New York for Injury Assessment Hudson River NRDA".

Data Sheet: "Avian Egg Collection Data Sheet", "Avian Egg Processing Data Sheet"

2.4.2 Egg Incubation and Injection

Eggs will be injected at a time point approximately 18% of incubation and is specifically defined in the SOP. Eggs will be candled during incubation at time points approximately equivalent to 18% (i.e., before injection), 24% (one day post-injection), 41%, and 82% of incubation and specifically defined in the SOPs. At time of candling, any dead or infertile eggs will be removed and any dead embryos evaluated for stage of development and deformities.

SOP: "Egg Injection and Incubation Procedure for Tree Swallow (*Tachycineta bicolor*) Eggs", "Egg Injection and Incubation Procedure for American Kestrel (*Falco sparverius*) Eggs".

Data Sheets: "Incubator Record", "Log of checking of eggs", "Egg Moisture Loss".

2.4.3 Dosing Solutions

PCB 126 solutions were prepared in corn oil, Sigma C8267, and provided by the USGS Columbia Environmental Research Center (CERC), Columbia, Missouri. Concentrations of the solutions provided by CERC are 0 (corn oil only), 5.9 ng/uL, 60 ng/uL, 110 ng/uL, 500 ng/uL, 5,000 ng/uL, and 10,000 ng/uL.

The PCB mixture solutions prepared in corn oil (Sigma-Fisher Scientific) will be provided by CERC at concentrations of 0 (corn oil), 8 µg/uL, 15 µg/uL, 31 µg/uL, 62 µg/uL, 123 µg/uL and 246 µg/uL if one microliter were injected per gram of egg, but since we are injecting 0.4 µL/gram of egg our actual egg doses will be 3 µg/g, 6 µg/g, 12 µg/g, 25 µg/g, 49 µg/g, and 98 µg/g.

SOP: “Egg Injection and Incubation Procedure for Tree Swallow (*Tachycineta bicolor*) Eggs”, “Egg Injection and Incubation Procedure for American Kestrel (*Falco sparverius*) Eggs”.

Data Sheets: “Egg treatment and incubation log”, “Log of checking of eggs”

2.4.4 Egg Hatching and Tissue Sampling

Initially, embryos will be sampled at 90% of incubation to maximize the number of viable embryos for tissue collection. If viability at 90% of hatchability is high we may monitor hatchability in subsequent sets of eggs. Any eggs that fail to hatch will be opened and condition of the embryo noted. Deformities will be scored for presence or absence of crossed bill, shortened upper bill, unusually small or large eyes, edema of the neck and head area, incomplete ossification of skull (brain not enclosed in skull), gastroschisis in post stage 45 embryos (if incubation is taken to hatch), malformed or clubbed feet, asymmetrical body form, mal-position in the egg, and any other abnormal appearances shall be noted shall be noted on the data sheet.

Data Sheet: “Deformity Score Sheet”

Samples from each bird will be identified by a unique code encompassing the species, egg code and year, e.g. TRES-1-2006 for a tree swallow collected in 2006. The four letter codes of AMKE and TRES will be used for American Kestrel and tree swallow samples respectively. CKN will be used for chicken samples. Series of numbers 1-199 for Patuxent Research Refuge, 200-799 for Great Sacandaga Lake and 800-899 for Hudson River will be used for tree swallow eggs collected at these sites. For American kestrels collected from Patuxent Research Colony, the unique code will reflect the nest number and clutch order number followed by year, e.g., AMKE-634-6-2006, for the sixth egg laid in nest 634. For chicken a series of numbers 1 and higher will be used, e.g. CKN-1-2006. Sampling of hatchlings will include body weight, organ weights, and collection of tissue.

SOP: “Necropsy of Hatchling Birds”

Data Sheet: “Hatchling Sampling Data Sheet”

2.4.5 Biological Tissue Analysis

Gonads and bursa of Fabricius will be analyzed for all birds by histological assessment. Slides will be labeled and well organized for retrieval and review. Liver tissue will be analyzed for EROD activity as a biomarker of planar PCB exposure. Because the amount of tissue will be limited, we will prioritize the analyses for the hypothalami and will perform aromatase enzyme in all individuals, then monoamines and GnRH-I when possible, i.e., when tissue samples are large enough. Please note that the monoamines and GnRH-I are analyzed in the same extract. Hypothalamic monoamines, GnRH-I and aromatase activity will be measured from the posterior hypothalamus, which is the area of storage and release of GnRH-I. Monoamines are measured by HPLC with electrochemical detection. GnRH-I is measured by an EIA (enzyme immunoassay), using an antibody specific for GnRH-I. Aromatase will be measured by radiological enzymatic activity assay. Estradiol and androgens in fecal matter and thyroid hormone content of thyroid gland will be measured by RIA (radioimmunoassay), according to techniques used in our laboratory. RIAs will be validated for tree swallows and American kestrels. Additional tissue will be collected from hatchlings and stored for microarray analysis from any or all of the following: hypothalamus, gonad, bursa, thyroid, heart and liver. Lab notebooks and records from these assays will be dated and initialed.

SOPs:

- “Histological Analysis Of Avian Tissue: Gonads and Bursa of Fabricius”
- “EROD Assay For CYP450 Activity In Liver Samples”
- “High Pressure Liquid Chromatography: Monoamine Analysis”
- “GnRH-I ELISA”
- “Protein Assay”
- “Extraction of Steroid Hormones from Avian Feces”
- “Radioimmunoassay of Androgens In Avian Serum or Fecal Extract”
- “Radioimmunoassay of Estradiol In Avian Serum or Fecal Extract”
- “Digestion Of Thyroid Gland And Radioimmunoassay of Thyroid Hormones”.
- “Analysis of Aromatase Activity in Hypothalamic Tissue”
- “Tissue Storage, RNA Isolation and Microarray Analysis”

2.4.6 Statistical Analyses

Data will be analyzed by testing for normality and proceeding with parametric ANOVAs or non-parametric tests, and regressions as appropriate. Mortality data will be analyzed with Fisher Exact Probability test and probit analysis for determining median lethal doses or effects. When necessary, further analyses would be used to understand the significance of dose-responses and non-monotonic trends. If the predictions warrant the use of one-tailed tests, these tests will be used with consultation with our statistician. Additional tests may include bootstrap techniques if data are not normally distributed and sample sizes are low. A biostatistician has agreed to serve as a statistical consultant. These statistical tests may be revised, including not performed, by the Principal Investors (PI) based on data collection. Further, the PIs may conduct additional statistical tests not noted above.

The PIs plan to conduct the following comparisons. Null (HO) and alternative (HA) hypotheses are presented below. "PCB" refers to PCB 126 for work with tree swallows and PCB mixture for work with chickens and American kestrels.

2.4.6.1 Compare the embryo survival or hatchability of eggs injected with the PCB with eggs that are un-injected (control) or are injected with only the vehicle solution for the PCB (vehicle control).

- General Hypotheses
HO: Hatchability of eggs injected with the PCB is equal to the hatchability of control eggs
HA: Hatchability of eggs injected with the PCB is less than the hatchability of control eggs in a dose response manner
- Statistical tests
Fisher Exact probability tests and probit analysis will be used for determining significant decreases in survival or hatchability and for determining median lethal doses.

2.4.6.2 Compare occurrence and severity of deformities between PCB exposed embryos and unexposed embryos.

- General Hypotheses
HO: The occurrence and severity of deformities are equal in control and PCB exposed embryos
HA: The occurrence and severity of deformities are increased in PCB exposed embryos compared to controls
- Statistical tests

Fisher Exact probability tests and probit analysis will be used for determining significant increases in deformities and for determining median effect concentrations.

2.4.6.3 Compare organ weights of PCB exposed embryos and unexposed embryos.

- General Hypotheses
HO: Organ weights in PCB exposed embryos are not different than controls
HA: Organ weights in PCB exposed embryos are different compared to controls and are proportionally related to the dose of treatment

For liver weight, HA: PCB exposure will enlarge liver.

- Statistical tests

All data will be tested for normality and homogeneity of variances. If assumptions of normality and homogeneity of variances are met, organ weights will be compared across treatment groups by treatment and sex, and possible interactions examined, using 2-way ANOVA with appropriate post hoc comparisons. Alternately, appropriate non-parametric tests shall be used. Furthermore, regression analyses with estimates of confidence intervals or dose-response statistics such as the Jonckheere test shall be used to evaluate dose related effects.

2.6.4.4 Compare histology of gonad, bursa and heart of PCB exposed embryos to unexposed embryos.

- General Hypotheses
 HO: Gonad, bursa, and heart morphology in PCB exposed embryos are not different than controls
 HA: Gonad, bursa, and heart morphology PCB exposed embryos are different compared to controls and are proportionally related to the dose of treatment
- Statistical tests
 All data will be tested for normality and homogeneity of variances. If assumptions of normality and homogeneity of variances are met, histological indices of morphology will be compared across treatment groups by treatment and sex, and possible interactions examined, using 2-way ANOVA with appropriate post hoc comparisons. Alternately, appropriate non-parametric tests shall be used. Furthermore, regression analyses with estimates of confidence intervals or dose-response statistics such as the Jonckheere test shall be used to evaluate dose related effects.

2.4.6.5 Compare cytochrome P450 activity in liver microsomes from PCB exposed embryos to that of unexposed embryos.

- General Hypotheses
 HO: Cytochrome P450 activity in PCB exposed embryos is not different than controls
 HA: Cytochrome P450 activity in PCB exposed embryos is higher than controls and is proportionally related to the dose of treatment
- Statistical tests
 All data will be tested for normality and homogeneity of variances. If assumptions of normality and homogeneity of variances are met, cytochrome P450 activity will be compared across treatment groups by treatment and sex, and possible interactions examined, using 2-way ANOVA with appropriate post hoc comparisons. Alternately, appropriate non-parametric tests shall be used. Furthermore, regression analyses with estimates of confidence intervals

or dose-response statistics such as the Jonckheere test shall be used to evaluate dose related effects.

2.4.6.6 Compare thyroid hormone (T3 and T4) content of thyroid glands from PCB exposed embryos to that of unexposed embryos.

- General Hypotheses
HO: Thyroid hormone (T3 and T4) contents of thyroid glands in PCB exposed embryos are not different than controls
HA: Thyroid hormone (T3 and T4) contents of thyroid glands in PCB exposed embryos are lower than controls and are proportionally related to the dose of treatment
- Statistical tests
All data will be tested for normality and homogeneity of variances. If assumptions of normality and homogeneity of variances are met, thyroid hormone concentrations will be compared across treatment groups by treatment and sex, and possible interactions examined, using 2-way ANOVA with appropriate post hoc comparisons. Alternately, appropriate non-parametric tests shall be used. Furthermore, regression analyses with estimates of confidence intervals or dose-response statistics such as the Jonckheere test shall be used to evaluate dose related effects.

2.4.6.7 Compare estradiol and androgen content in fecal matter from PCB exposed embryos to that of unexposed embryos.

- General Hypotheses
HO: Estradiol and androgen content in PCB exposed embryos are not different than controls
HA: Estradiol and androgen content in PCB exposed embryos are different than controls and are proportionally related to the dose of treatment
- Statistical tests
All data will be tested for normality and homogeneity of variances. If assumptions of normality and homogeneity of variances are met, steroid hormone concentrations will be compared across treatment groups by treatment and sex, and possible interactions examined, using 2-way ANOVA with appropriate post hoc comparisons. Alternately, appropriate non-parametric tests shall be used. Furthermore, regression analyses with estimates of confidence intervals or dose-response statistics such as the Jonckheere test shall be used to evaluate dose related effects.

2.4.6.8 Compare hypothalamic aromatase activity, catecholamine content and GnRH-I concentration in PCB exposed embryos to unexposed embryos.

- General Hypotheses

HO: Hypothalamic aromatase activity, catecholamine content and GnRH-I concentration in PCB exposed embryos do not differ from controls

HA: Hypothalamic aromatase activity, catecholamine content and GnRH-I concentration in PCB exposed embryos are different than controls and are proportionally related to the dose of treatment

- **Statistical tests**
All data will be tested for normality and homogeneity of variances. If assumptions of normality and homogeneity of variances are met, hypothalamic aromatase activity, catecholamine content and GnRH-I concentration will be compared across treatment groups by treatment and sex, and possible interactions examined, using 2-way ANOVA with appropriate post hoc comparisons. Alternately, appropriate non-parametric tests shall be used. Furthermore, regression analyses with estimates of confidence intervals or dose-response statistics such as the Jonckheere test shall be used to evaluate dose related effects.

These hypotheses and statistical tests may be revised, or not performed by the PI based on data collected. Further, the PI may test other hypotheses and conduct additional statistical tests not noted above.

3. QUALITY ASSURANCE/QUALITY CONTROL

3.1 *Data Quality Objectives, Indicators, and Assessment*

3.1.1 Overview

This study is being conducted in accordance with the Quality Assurance Management Plan for the Trustees' Hudson River NRDA. As described in the plan, four general elements of quality assurance/quality control (QA/QC) must be addressed for each data collection effort:

- Project Management
- Data Generation and Acquisition
- Assessment and Oversight
- Data Validation and Usability

This section describes the Quality Assurance Plan (QAP) for the avian egg injection study, based on these four general elements. The objectives of the study are outlined in Section 1 of this Work Plan. To achieve these objectives, the following requirements must be met:

- \$ All samples, from the initial eggs through embryos, hatchlings, dead or infertile eggs, necropsy samples, and egg products must be identified and stored following documented procedures to insure proper identification and handling.

- \$ All procedures for assessment of biological impacts, including egg injections, observation and measurement of live birds, necropsy, and biological tissue analyses, must be performed following documented procedures to ensure consistent, comparable data.
- \$ PCB mixture preparation and egg contaminant levels: The laboratories performing chemical contaminant testing will follow the requirements of the Hudson River NRDA Analytical QA Plan. This effort is not part of the current work plan and will be funded separately.

3.1.2 Project Management

The study team is organized based on tasks and levels of responsibility to ensure good communication between all personnel. The Assessment Manager (Kathryn Jahn, USFWS) has overall project oversight responsibility and provides direction to the Quality Assurance Coordinator. The Assessment Manager also provides direction to the Principal Investigator and Co-Principal Investigator via the Project Coordinator. The Project Coordinator is responsible for ensuring that adequate coordination and communication occurs amongst the Assessment Manager, Quality Assurance Coordinator, Principal Investigator and Co-Principal Investigator. The Principal Investigator and Co-Principal Investigator are responsible for the project's design and implementation and provide guidance and technical expertise as needed to the study team and statistician. They will also work with the Project Coordinators and Quality Assurance Coordinator to ensure that the study is consistent with the overall QA objectives of the NRDA.

The work plan was developed to provide detailed and explicit instructions for the research staff to follow in collecting the study data. The plan has been reviewed, commented on, and approved by key parties to the study before the beginning of egg injection. Reliance on a detailed, explicit, and fully reviewed plan ensures that:

- Study objectives, methods, procedures, and details are documented before sampling.
- Data are collected in a systematic and consistent way throughout the study.
- Each member of the study team adheres to the requirements of the plan. In particular, the Principal Investigator and Co-Principal Investigator must ensure that their research staff adheres to the plan. Each team member is required to sign a statement that they have read the plan and understand it.

Events may arise during this study that require changes to the procedures documented in the work plan. Deviations from the work plan will be documented in writing, with a detailed explanation of the reasons for these deviations. Predetermined deviations from the plan will be conducted only after the approval of the Principal Investigator and/or Co-Principal Investigator.

3.2 Data Generation and Acquisition

3.2.1 Data Quality Objectives

Data developed in this study must meet standards of precision, accuracy, completeness, and comparability, and be consistent with sound scientific methodology appropriate to the data quality objectives.

Precision is the degree of mutual agreement among individual measurements of the same property under similar prescribed conditions, such as replicate measurements of the same sample. Precision is concerned with the “closeness” of the results. For this study, repeated independent measurements will be performed to assess the precision of several biological assays. Precision will be expressed as the relative standard deviation (RSD) between these replicate measurements on a single sample, and for the hormone assays, will be expressed as Coefficient of Variation.

Accuracy is the degree of agreement of a measurement with an accepted reference value and may be expressed as the difference between the two measured values or as a percentage of the reference value. For this study, evaluation of accuracy may be performed using a positive control sample or reference standard for most biological assays. Another measure of accuracy will be the closeness of duplicate determinations of the same sample in an assay.

Completeness is defined for this study as the percentage of the planned data collections compared to data actually collected within the work plan specifications. Because there is uncertainty due to the variables in number and viability of available eggs and hatchlings, the assessment of completeness achieved will be assessed in two ways. First, completeness will be assessed by comparing planned sampling versus samples collected at the end of the study. Secondly, the DQO for completeness of data analysis is 95%, which pertains to no more than 5% of the data points collected are to be rejected as unreliable.

Comparability is defined as the measure of confidence with which results from this study may be compared to another similar data set. For this study, evaluation of comparability will be performed using external reference standards or an internal standard prepared from a serum pool extract or a standard prepared within our laboratory, aliquoted and frozen into individual units for utilization within each assay as an internal quality control measure. These comparisons will also take into consideration inter-assay variability due to reagent differences. For example, antibodies used in hormone assays may differ in the forms of their cross reactivity with closely related hormones thereby providing differing absolute concentrations.

3.2.2 Study Documentation

All study procedures and results will be documented on data sheets, which will be placed in binders and retained for review. To the extent possible, information will be recorded on pre-formatted data sheets. The use of pre-formatted data sheets is a QA/QC measure designed to:

- ensure that all necessary and relevant information is recorded for each sample and each sampling activity
- serve as checklists for the Principal Investigator, Co-Principal Investigator and their staff to help ensure completeness of the data collection effort
- assist the research staff by making data recording more efficient
- minimize the problem of illegible or hard-to-follow notebook entries

The researcher performing each procedure will be responsible for recording information on data forms.

Data entries will be made in waterproof ink, and corrections will be made with a single line through the error accompanied by the correction date and corrector's initials. Each completed data sheet will be reviewed, corrected (if necessary), and initialed by the Principal Investigator, Co-Principal Investigator, or their designee. Following completion of the study, data sheet originals will be retained.

3.2.3 Sample Identification Procedures

Strict sample identification procedures will be used throughout the study. The sample identification procedure will begin when an egg is collected. Each egg will be identified by a unique sample code including links to the year of collection, species, and egg code (see section 2.4.4).

The sample identification described in section 2.4.4 will be recorded on all data sheets used to document all procedures. This identification along with tissue type will be transferred to all other sample types originating from the egg, including hatchlings (live and sacrificed), and necropsy samples.

The sample ID will be used to uniquely identify all samples, either on a label or written directly on the container. The code will be recorded using a waterproof marker. If applicable, the label should also include the type of sample and date of collection and researcher's initials.

3.3 Assessment and Oversight

The QA management plan specifies that studies that generate data will be audited to ensure that the project-specific plans are being properly implemented. Several mechanisms for internal audits of the data generation process will be used for the avian egg injection study. These mechanisms include:

- A project management structure that defines clear lines of responsibility and ensures communication between researchers and trustees. Clear responsibilities and communication can serve as a means of providing internal audits of the study as it

proceeds.

- A requirement that laboratory notebooks and data forms be completed daily and be reviewed weekly by the Principal Investigator or Co-Principal Investigator.
- The use of pre-formatted data sheets that serve as a checklist for study procedures and assay results.
- The study will not begin until approval is received from the Quality Assurance Coordinator or their delegate. The Quality Assurance Coordinator will conduct an audit of the procedures and documentation of the study.

3.4 Data Validation and Usability

This study employs documented, repeatable procedures to perform the experiments and assays required to generate the data for this study. The work plan has been reviewed for the adequacy of the design and proposed methodology. The original data sheets and other study records will be maintained and archived for a minimum of eight years. Disposal of these records will require the approval of the Assessment Manager. Findings from this study can be reviewed against the data sheets to ensure that the data presented in the reports represent complete and accurate information. Chemistry contaminant data will be validated as specified in the Analytical QA Plan.

The Principal Investigator or Co-Principal Investigator will perform oversight of all egg injections and data collection for measurement endpoints. They will validate that Project Scientists and Technicians are correctly following the standard operating procedures and correctly documenting the results.

Data analysis will be performed using JMP IN version 5, release 5.1, SAS Institute Inc and SAS programming but not be limited to these statistical programs. All numeric data presented in reports will contain basic statistical properties and uncertainty. The robustness of each parameter studied will be presented.

3.5 Chain of Custody Procedures

Chain of Custody (COC) procedures will be used during the field sample collection and transfer to the laboratories for incubation or analysis. The purpose of COC is to assure the integrity of each sample and be able to clearly identify who was responsible for the sample at each step. The COC procedure will begin when an egg is collected from the nest. That collection is documented on field data forms (Avian Egg Collection Data Sheets), which clearly identify the team member(s) responsible, as well as the date and time. The egg collection forms will clearly identify to whom the sample was delivered for further processing, and will also include the date and time.

The immediate team members are personally responsible for the care and custody of the samples that are in their possession. A sample is in custody of the immediate team member if any of the following occur:

- The sample is in the individual's physical possession;
- The sample is within view after being in possession;
- The sample is in a locked or sealed container that prevents tampering after being in possession; or,
- The sample is in a designated secure area.

When the samples are packed in coolers or other containers for shipment to the laboratory or storage facility, completed COC records will accompany the samples. The COC form will contain the following information:

- Project name;
- Sample identification (unique for each sample);
- Sample matrix (e.g., egg contents, liver) which may be part of the sample ID;
- Name and signature of individual relinquishing custody;
- Name and signature of individual accepting custody;
- Sample shipping date and mode.

Other information such as date of sample collection, collection location, and jar sizes may be on the COC form or on accompanying documentation.

An original COC record for the samples in that cooler will accompany each shipping container. All sections of the COC form will be completed. Indication of the number of coolers per shipment (e.g., 1 of 3) will be listed on the form if more than 1 container is shipped. Once the form is completely filled out, it will be placed securely inside the cooler (in a plastic sealable bag to keep it dry). Field personnel will maintain a copy of the COC to keep with the airbill. The cooler will be sealed with custody seals or the containers inside the cooler may be sealed with custody seals. Custody seals are used to detect unauthorized tampering with samples after sample collection until the time of use or analysis. Signed and dated gummed paper seals may be used for this purpose. The seals will be attached so that they must be broken to open the shipping container. Each cooler will be sturdy and well sealed with strapping or other tape. All samples will be kept in locked locations or with custody seals at all times until shipped.

An air bill, Federal Express shipping label, etc. can be used to document the transfer of a sample from the field team to an intermediate storage location, the analytical laboratory, or archive freezer.

Coolers or other containers containing samples will be opened at the analytical laboratories or archiving facility only by a person authorized to receive the samples. The containers will first be inspected for integrity of the chain of custody seals or other signs of tampering. The receipt of each sample in the coolers or containers will be verified on the COC forms. The signed COC forms will be photocopied, and the photocopy will be mailed to the sending party. Samples will be stored in a secure area according to procedures documented for each analytical facility.

4. PRINCIPAL INVESTIGATORS

Principal Investigator

The PI is a neuroendocrinologist with twenty five years of experience studying avian neuroendocrinology and reproduction. The PI will oversee all aspects of the studies.

Co-Principal Investigator

The Co-PI is an avian toxicologist with experience in egg injection studies and immune and endocrine disruption studies in birds. The Co-PI will plan the logistics of all aspects of the study and participate in assays, data collection and data analysis.

Research Technicians with expertise in endpoints required as part of the study will conduct assays and analyze data as needed. The lead technician will devote 50% time to this work.

Other staff will assist with animal care, sampling and assays as needed.

The full names, contact information, written signature and written initials of all individuals working on this project shall be maintained in the project file.

5. LITERATURE CONSULTED

Brunstrom, B., Halldin, K. 1998 EROD induction by environmental contaminants in avian embryo livers. *Comparative Biochemistry and Physiology Part C* 121: 213-219.

6. STANDARD OPERATING PROCEDURES

6.1 Recording and Handling Data for Avian Egg Injection Study

This protocol describes procedures for recording and handling data in this laboratory.

Procedure

- 1) Blank data sheets are available in electronic format on the lab server in the "Lab Protocols" folder.
- 2) Data entry:
 - Entries will be made in ink.
 - All blank cells in the sheets should be filled with data, or marked with "NA". Large areas left blank (such as the bottom part of a partially-filled sheet) should be crossed out.
 - Any changes will be made by crossing through the error with a single line, and initialing and dating the change.
- 3) After hard copies of data sheets are filled out they must be reviewed by the PI or Co-PI then stored in the project notebook in the Co-PI's office in a locked filing cabinet.
- 4) Data should be input as soon as possible, after collection, into electronic files (Excel or JMP) and files stored on the PI's or Co-PI's computers. Data entry must be 100% verified against the hard copy by someone other than the person who performed the initial data entry.
- 5) Back-up copies should be made to a CD after any additions or changes to files are made. A back-up copy of data on CD will be made weekly and will be stored at the homes of the PI or Co-PI.
- 6) Any deviations from the protocols will be written out in detail by the Principal Investigator and added to the project notebook.

6.2 Field Collection of Tree Swallow Eggs for Avian Egg Injection Study, Hudson River NRDA

INTRODUCTION

Avian egg injection is a well-established technique to assess the effects of contaminants on a developing avian embryo. To conduct an avian egg injection experiment, eggs from Patuxent Wildlife Research Refuge will be collected and brought into a laboratory where they are injected with the substance being tested. Proper handling of the eggs during collection and transit to the laboratory is essential to maintain viability in eggs that will subsequently be injected with contaminants and incubated. A subsample of eggs will be selected for contaminants analysis.

MATERIALS AND EQUIPMENT

FIELD:

- Scientific collecting permits
- Field notebook, writing instruments (pencils/pens/permanent markers)
- Padded egg collection boxes (hard-sided container, e.g., Tupperware or tackle box, with foam padding)
- Labels
- Styrofoam or other type of cooler or ice chest
- Frozen cooler pack(s)
- Avian Egg Collection Data Sheets

PROCEDURES

FIELD:

- Collected eggs should be whole and not cracked.
- For tree swallows, the following approach should be used: Incubation of tree swallow eggs doesn't start until the clutch is complete. Eggs are generally laid at one day intervals. Monitor the laying of eggs, waiting till the fifth egg is laid. On the day the fifth egg is laid collect 2 eggs from that nest. Continue monitoring the nest. If another egg shows up in that nest the next day (so the total clutch would have been 6), a third egg can be collected (leaving three).
- For each egg collected, complete the appropriate information on the Avian Egg Collection Datasheet.
-
- For egg that are going to be incubated: Using a soft pencil, mark the Egg Code on the egg.
- Transport to lab in hard container with sufficient padding.
- Complete chain of custody transfer of samples from field collection crew to laboratory crew on Egg Collection Data Sheet.
- For eggs to be incubated: Follow Work Plan procedures.
- For eggs that are going to be analyzed for contaminants and not incubated: Refrigerate

eggs until opened, no longer than 48 hrs. Processing of eggs for contaminants analysis will be completed on a daily basis as much as practical. Follow Standard Operating Procedure for Removal of Avian Egg Contents for Contaminants Analysis, Hudson River NRDA.

Collector _____ Data Recorder _____
 Name Signature Name Signature

Egg Code ¹	Location ²	Date Collected ³	Time Collected ⁴	Clutch Size	Eggs Warm Yes or No	Comments

¹ Egg Code: Numeric code beginning at 1; ² Site, species and nest number, such as Patuxent TS 2 where TS=tree swallow and AK=American kestrel; ³ In MM/DD/YEAR format, such as 04/30/2006 for April 30, 2006; ⁴ In 24-hour format, such as 1300 for 1PM

Custody of samples listed above transferred from field collection crew to laboratory crew as follows:

Relinquished by: _____
 Signature Print Name Company/Title Date Time
 Received by: _____
 Signature Print Name Company/Title Date Time

Data Sheet checked by: _____ Date: _____
 Name/Initials

6.3 Removal of Avian Egg Contents for Contaminants Analysis Hudson River NRDA

INTRODUCTION

Avian eggs are a common sample for contaminants analysis. An accurate analysis depends upon getting the egg contents from the shell to a clean sample jar without introducing other sources of contamination. This protocol, which has been developed and refined by many researchers over the decades, was written for those who have minimal experience. Your first egg should be a practice egg. *It is suggested that all personnel practice on several quail eggs to improve technique. Chicken eggs may be used if quail eggs are not available.*

LABORATORY MATERIALS AND EQUIPMENT

- Avian Egg Processing Data Sheets
- paper or other towels
- green scrubby or sponge
- Acculab V-200 balance, weighs to nearest 0.01 gm
- calipers
- Chemically-clean jars, 1 per sample
 - ✓ Make sure they are cleaned for the contaminants you are sampling, e.g., I-Chem pesticide/PCBs Series 200 or 300.
 - ✓ Size: 4 oz.
- chemically-clean stainless steel scalpel blades (No. 21 or No. 22 with No. 4 handles work well)
- chemically-clean forceps
- aluminum foil sheets (approximately 30 x 30 cm square), 1 per egg
- sharps container for used blades or disposable scalpels

LABORATORY PROCEDURES

- Fill out the Avian Egg Processing Data Sheet; use one data sheet per egg.
- If debris is present, rinse egg in cool water while gently scrubbing with green scrubby or sponge. Do not soak the egg.
- Dry and weigh whole egg to the nearest .01 g
- Take three measurements each of egg length and maximum egg width with calipers. Compute average of three measurements for final width and length measurements.
- Transfer egg contents to chemically-clean jar using the following procedure:
 1. Use nitrile gloves for this part of the procedure. Avoid letting contents run over your hands into the sample jar.
 2. Create a catch basin out of the aluminum foil by turning edges up and securing the corners. This will catch egg contents in case they spill over the edge of the jar. Use a separate piece of foil for each sample. The foil also is a clean place to place your instruments when they are not in use.
 3. Weigh the clean empty jar with lid on, and note this tare weight on data sheet.

4. Place jar in center of aluminum foil, and loosen the lid.
5. Score equator with serrated blade or scalpel blade. Use a new, chemically-clean scalpel blade for each egg. This part takes practice. Cradle the egg in one hand (don't squeeze too tightly!) and gently score while rotating the egg. Many light strokes are preferable to a fewer deeper strokes, increasing the evenness of the score and decreasing the possibility of eggshells not separating cleanly or of punching through the shell. Continue to work on your score until you see the membrane, which usually appears gray underneath the white of the eggshell. When you see the first bit of membrane, remove the lid from the jar so that it will be ready as soon as you need it. Avoid getting shell dust, or anything else besides the egg contents, in the jar. Try to expose the membrane evenly around the entire egg. Often the score line can be used to help pick the egg shell apart using forceps.
6. Place the egg over the jar and cut through membranes with the scalpel. For large eggs a new scalpel blade may be used at this point to reduce the potential for cross contamination and since the blade may become dull during the cutting process. The scalpel can also be used to finish scoring down to the membranes. Pour contents into jar, or use the scalpel to gently scrape if that is necessary. Small stainless steel scoops may also be used to help remove the contents. Use forceps to remove any shell fragments from the jar. Cover the jar.
7. For swallows, hold the egg vertically with air cell end up. Using scissors cut the top of the eggshell off above the air cell if possible. Pour contents into the jar, or use the scalpel to gently scrape if that is necessary. Use forceps to remove any shell fragments from the jar. Avoid getting shell dust, or anything else besides the egg contents, in the jar.
8. The target for the minimum weight of egg tissue is 4 grams for analysis. It may be possible to analyze smaller samples ranging from 1 – 2 grams. Analysis of these samples may result in a lower ability to detect contaminants due to the lack of mass. An effort must be made to maximize the amount of each sample that is usable. The weight of each sample should be made in the laboratory during egg processing using the following procedure:
 - a. Place a small jar on a balance that reads to at least 1 milligram and that has been appropriately calibrated.
 - b. Tare the jar or record the jar weight if the balance cannot be tared.
 - c. Open the egg, according to the procedures referenced above and empty the contents into the jar.
 - d. Record the weight, to the nearest .01g, of the egg contents if the balance was tared. If the balance was not tared, then record the weight for the egg contents and the jar, then subtract the previously recorded weight of the jar. Record the weight of the egg contents in the field notebook and on the jar label.
 - e. If egg is developed, estimate age of embryo. Wet weight conversion will be made based on the weight and egg measurements. A photographic record of the contents of each egg will be made. Documentation of embryo development is very limited (Powell *et al.* 1998; Bird *et al.* 1984), therefore, documenting this phase of the egg processing is important. Note

amount of decay or anything else pertinent to your study, and examine for deformities, particularly bill deformities such as crossed bills or lack of jaws, but also lack of skull bones, club feet, rotated ankles, or dwarfed appendages (Gilbertson *et al.* 1991).

- f. Repeat these procedures for any other eggs that need to be added to the sample jar. Using these procedures, the weight of each egg's contents will be measured, even for eggs whose contents are combined into a single jar.
- ✓ Do not touch or move the jar between steps b. and d. above. It is preferable to add the egg contents to the jar while the jar is still on the balance, immediately after taring the jar.
- Place label on jar. Place clear tape over the label to keep it from getting wet.
- Prepare Chain of Custody records and maintain egg samples under chain of custody.
- Freeze samples. Ship under Chain of Custody overnight on dry ice to the sample archive or analytical laboratory.

Literature Consulted

Bird, D.M, J. Gautier, and V. Montpetit. 1984. Embryonic growth of American kestrels. *Auk* 101:392-396.

Gilbertson, M., T. Kubiak, J. Ludwig, G. Fox. 1991. Great Lakes embryo mortality, edema, and deformities syndrome (GLEMEDS) in colonial fish-eating birds: similarity to chick-edema disease. *J. Toxicol. Environ. Health* 33:455-520.

Powell, D.C., R.J. Aulerich, R.J. Balander, K.L. Stromborg, and S.J. Bursian. 1998. A photographic guide to the development of double-crested cormorant embryos. *Colonial Waterbirds* 21(3):348-355.

These egg-processing guidelines were developed by the U.S. Fish and Wildlife Service and modified for the project based on consultation with the author of these guidelines and on conversations with the Quality Assurance Coordinator for this project.

Appendix A: Chemically-Clean Instruments for Collecting Contaminants Samples

To minimize cross-contamination when collecting biological samples for contaminants analysis, a primary requirement is use of chemically-clean instruments. These are made of appropriate materials (stainless steel or teflon) and rinsed with alcohol and solvents to remove contamination and organics. Once rinsed, the instruments should be treated as sterile instruments, e.g. not placed on unclean surfaces.

Because every laboratory situation is different, this document tells you what to do, but not how to do it. The chemicals used for rinsing are hazardous, so you should follow proper safety and laboratory protocols when using them. This includes proper personal protective equipment (lab coats, gloves specific to the chemical, eye protection), proper laboratory equipment and procedures (use of hood, proper storage and disposal methods), and knowledge of chemical hazards such as flammability, reactivity, and toxicity (MSDS required). If this is all new to you, enlist the help of a chemist to help you make the proper decisions and reduce your risks of exposure and accident.

For organics, rinse with a reagent grade isopropyl alcohol, air-dry, rinse with reagent-grade hexanes, and air-dry.

Rinsing should be done using glass pipettes or wash bottles (made of appropriate material for the rinsing agent). Glass funnels, wide enough to accommodate your instruments and foil sheets, are invaluable in directing the flow of used chemicals into disposal containers or waste jars. Use disposal containers that are the same as your source chemical containers (e.g. brown glass). Never rinse into or pour unused chemicals back into your source chemical bottle.

Avian Egg Processing Data Sheet

Processor(s): Name _____ Name _____

Signature _____ Signature _____

Date Processed: _____

Egg Code: _____

Sample ID: _____

Egg Length (three measurements, mm): _____ , _____ , _____ Average _____

Egg Width (three measurements, mm): _____ , _____ , _____ Average _____

Jar lot number _____ Balance within limits? Yes OR No

Whole Egg Weight (g): _____

Contents weight:

Weight of jar (g) : _____

Weight of jar + contents (g): _____

Weight of contents (g): _____

Membrane location: ___ with embryo OR ___ with eggshell

Contents condition (embryo development ¹, state of decay, etc.) and other comments:

Other comments: _____

Contaminants disposition (catalog number and date submitted, etc):

¹ None, 1/4, 1/2, 3/4 , full term

Data Sheet checked by: _____ Date: _____

Name/Initials

6.4 Egg Injection and Incubation Procedure for Tree Swallow (*Tachycineta bicolor*) Eggs

This protocol outlines procedures for incubating eggs and injecting chemicals into the eggs of tree swallows. The purpose of which is to mimic maternal deposition of chemicals into the egg and determine toxicity toward the embryo.

Robertson et al. (1992) in *Birds of North America* describe the eggs, their incubation and hatching of tree swallow chicks and is summarized herein:

Eggs are 18.7 x 13.2 mm and 1.4 to 2.6 g in size with an average weight of 1.9 g.

Approximately 14% of mass is lost between laying and the end of incubation. Egg color translucent and rosy pink at time of laying turning to pure white (without any markings) around the fourth day of incubation. Eggs become more glossy during incubation. Incubation length for tree swallow eggs averages 14-15 days but ranges from 11 to 19 days. Female incubation rhythms have been reported as 11 minutes on the nest and 9 minutes off the nest. An embryo takes one to two hours to hatch from start of pipping and clutches hatch over a one to two day period, occasionally over three days. Hatchlings weigh 1.5 to 1.7 g, eyes are closed, skin is uniform pink and the gape is yellow. Hatchling is able to raise head to beg and position itself with the dorsal side up.

Incubation Procedures

- 1) Upon receipt of the eggs, examine them noting any evidence of damage or embryonic development (by candling) and assign each egg a number. Note on the coding sheet the source, nest number, egg number for the clutch etc. Write the egg number on the egg in soft pencil on the pointed end of the egg.
- 2) Assign eggs to treatment groups with consideration of number of eggs available, number of eggs from the same clutch (consult the biostatistician).
- 3) Weigh eggs and note weight.
- 4) Place eggs in an incubator in an egg rack of suitable size on their sides. Incubation will be done at 99.5°F and humidity adjusted as needed to ensure correct moisture loss (see no. 5). Eggs will be turned hourly by automatic rotation in the incubator for a total of 60° every two hours. In addition turn eggs 180° by hand or using a scoopula twice per day (before 10 am and after 4 pm). Mark the eggs with an X and an O on opposite side to confirm eggs are turned.
- 5) Check moisture loss by weighing a subset of eggs every other day and adjust the humidity appropriately to ensure correct moisture loss with egg mass loss averaging 14% over entire incubation period. For an average 1.9 g tree swallow egg over a 15 day incubation, moisture loss should average approximately 1% or 0.02 g per day.
On days four, seven, and thirteen of incubation, candle the eggs and remove infertile and dead eggs. Retain infertile and dead eggs for contaminants analysis; follow SOP "Removal of Avian Egg Contents for Contaminants Analysis, Hudson River NRDA".
- 6) Two days before estimated hatch date, (~90% of incubation) open the eggs and sample tissues from the embryos.

Injection Procedures

- 1) Two and half days into incubation (expected equivalent age to a 3 day old quail embryo, i.e. 18% of incubation based on a 15 day incubation for a tree swallow egg), candle eggs, note center of air cell with a dot of sharpie and remove infertile or dead eggs. Retain any infertile and dead eggs for contaminant analysis if warranted.
- 2) Assign the eggs to treatment groups and weigh each egg to the nearest one hundredth of a gram. Calculate and record the volume of dosing solution to be added to each egg. Round the volume to the nearest 0.01 μL .
- 3) Make injections into the egg as follows, allowing the eggs to be outside the incubator for not more than 30 minutes:
 - a. Wipe the blunt (air cell) end of the egg with 70% ethanol.
 - b. Gently make a hole in the egg with Dremel drill with a fine drill bit.
 - c. Inject the vehicle or PCB 126 solution, 0.4 $\mu\text{L/g}$ egg into the air cell, with a micro-pipettor and extended tip or Hamilton syringe.
 - d. Seal the hole with paraffin.
 - e. Allow the egg to sit pointed end down until the end of the 30 minute period.
- 4) Place eggs back into the incubator on their sides in an egg rack of suitable size. Randomly place treatment groups in the egg racks. Avoid placing eggs in the very top, very bottom, very back and very front of the incubator.

Equipment Needed

Incubators: Natureform NMC2000 or GQF Sportsman 1502

Egg trays

Light for candling

Ethanol and tissue or alcohol wipes

22 gauge needles or dremel drill

Hamilton syringes: one per treatment

Paraffin and tool to apply it to eggs

Heating block

Scales (510 - 0.001 g) Mettler Toledo PG503-S

Rainin Pipettman with extended tips: one tip per egg

Data Sheets

“Avian Egg Collection Data Sheet”, “Incubator Record”, “Egg Treatment and Incubation Log”, “Log of checking eggs”, “Moisture Loss Data Sheet”, “Deformity Score Sheet”.

Literature Consulted

Robertson, R. J., Stutchbury, B. J., and R. R. Cohen. 1992. Tree Swallow. *In* The Birds of North America, No. 11 (A. Poole, P. Stettenheim, and F. Gill, Eds.). Philadelphia: The Academy of Natural Sciences; Washington, DC: The American Ornithologists' Union.

Incubator Record

				% Humidity or	Eggs Turned (O/X)	List any adjustments	Initials
Incubator	Date	Time	Temperature	Wet Bulb Temp			

Reviewed by _____ Date _____

Egg Treatment and Incubation Log

Chemical:

Species:

Source:

Treatment (ug/g)	Egg ID	Egg Mass (g)	Dosing Concentration ug/uL	uL injected	Date of Injection (ED2.5)	Date Death Detected	Stage at death	Initials

Make any additional notes on reverse.

Reviewed by: _____

May 2006

Log of checking
of eggs

Species: _____

Egg Code
Range: _____

Date & Time Eggs
Set: _____

Date
Injected: _____

Date	Time	Investigator	Embryonic Day	# infertile	#dead	# remaining	Initials

Make any additional notes on reverse.

Reviewed
by: _____ Date: _____

Egg Moisture Loss

Species:

Source:

Date	Time	Egg ID	Embryonic Day	Weight (g)	Comments	Initials

Reviewed by: _____

Date: _____

Deformity Score Sheet

		Egg	Date Death		Cross	Short	Abnormal	Neck/head	Incomplete	Clubbed	Mal-	Gastroschisis		
Date	Species	Code	Detected	Stage*	Bill**	Upper Bill	Eye Size	Edema	Skull	Feet	position	(post stage 45)	Other	Initials

Reviewed by: _____ Date: _____

* If embryo is not old enough to detect a structure, or is too decomposed note "NS" for not scored under the deformity type.

** Note 'Y' (yes) or 'N' (no) to note presence or absence of the deformity.

6.5 Egg Injection and Incubation Procedure for American Kestrel (*Falco sparverius*)

This protocol outlines procedures for incubating eggs and injecting chemicals into the eggs of American kestrels, the purpose of which is to mimic maternal deposition of chemicals into the egg and determine toxicity toward the embryo.

Smallwood and Bird (2002) in *Birds of North America* describe the eggs, their incubation and hatching of American kestrel chicks and is summarized herein:

Eggs are approximately 34 x 28 mm and 10 to 18 g in weight. Egg color is variable from white to cream to yellowish to light red-brown with blotches and mottling of varying shades but especially brown shades. Eggs are generally not glossy. Incubation length for American kestrel eggs averages 27 to 29 days in captivity but approximately 30 days in the wild. “Apparently relatively cold-hardy...Captive-produced eggs hatched successfully in an incubator that shut down twice due to power failures to the point of ice forming on added water.” Kestrels are considered semi-altricial. An embryo takes approximately 48-52 hours to hatch from start of pipping, the female assists the chick out of the shell. Hatchling’s skin is pinkish and covered in sparse white down, bill, cere and talons are white-pink and legs and yellowish. Belly is prominently protruding and nearly naked. Hatchling is able to raise head, open its bill and ‘peep’.

Egg Collection from Patuxent NWRC Kestrel Colony

- 1) Collect eggs between 8 and 9:30 am (during feeding to minimize disturbance).
- 2) Have rubber gloves on to collect the eggs. Have a pair of leather gloves on hand in case you have to push aside a female kestrel.
- 3) Label eggs in pencil at the pointed end. If a nest box is #660, then label the egg 660-1, or 660-2 etc depending on egg order.
- 4) Place the eggs in a cushioned container for transport back to the laboratory.

Incubation Procedures

- 1) Upon receipt of the eggs at the laboratory, examine them noting any evidence of damage or embryonic development (by candling). Note on the coding sheet the source, nest number, egg number for the clutch etc. Wash eggs in a 40°C 1% betadine solution, and then rinse in 40°C water. Submerge eggs for less than 5 seconds in the betadine solution, and lightly scrub the cuticle off with hands and dry with a paper towel (wash one egg at a time as a pencil label often rubs off).
- 2) Re-label the egg with its number if it has washed off.
- 3) Weigh eggs to the nearest one hundredth of a gram and note weight. Eggs collected from the Patuxent colony in thus far in 2006 weigh between 13 and 18 grams (personal communication with Moira McKernan).
- 4) Hold eggs in cold storage (13°C) for not more than 4 days (Pisenti et al. 2001).

- 5) Warm eggs by leaving them at room temperature for one hour.
- 6) Place eggs on their sides in the Kuhl incubator in Kuhl brand pheasant egg racks. Incubate the eggs at 99.5°F and 55-65% humidity (84°F wet bulb). In addition to the hourly turning (60°) of the eggs done automatically by the incubator, turn the eggs 180° twice a day at 9 am and 5 pm +/- 1 hr. Draw an O and an X on opposite sides of the egg. At the morning time point turn the egg so that the O is showing and at the afternoon time point turn the egg so that the X is showing. This step provides additional turning that may be necessary for wild bird eggs.
- 7) Check moisture loss by weighing the eggs on Monday, Wednesday and Friday of each week and adjust the humidity appropriately to ensure correct moisture loss with egg mass loss averaging between 9 and 14% over entire incubation period. For an average 14 g kestrel egg over a 28 day incubation, moisture loss should average 2 grams or approximately 0.5% or 0.07 g per day to achieve a 14% weight loss over 28 days.

Steps number 6 and 7 are adapted from the methods of Pisenti et al. (2001) and personal communication with Dr. Gary Heinz, Patuxent NWRC. Pisenti et al. (2001) described a 9% egg mass loss for embryos surviving to hatch and Dr. Heinz adjusted humidity as needed to attain a 14% mass loss with good hatchability using a Kuhl brand incubator.

- 8) On approximately days seven, twelve and twenty-four of incubation, candle the eggs and remove infertile and dead eggs. Open eggs containing dead embryos and stage the embryo based on the guide in Pisenti et al. (2001). Note the stage of the embryo and any deformities on the egg treatment log.
- 9) On embryonic day 24, transfer eggs to a 'hatcher' incubator (99.5°F and 70-75% humidity) or separate tray in the same incubator with each egg placed in its own compartment fashioned from plastic mesh.
- 10) Necropsy within 24 hours after hatch.

Injection Procedures

- 1) Five days into incubation (expected equivalent age to a 3 day old quail embryo, i.e. 18% of incubation based on a 28 day incubation for a kestrel egg), candle eggs and remove infertile or dead eggs if possible to see through the egg shell. Retain any infertile and dead eggs for contaminant analysis if warranted.
- 2) Assign the eggs to treatment groups with consideration of number of eggs available, number of eggs from the same clutch, and optimal number of treatment groups (consult the biostatistician). Weigh each egg to the nearest one hundredth of a gram. Calculate and record the volume of dosing solution to be added to each egg. Round the volume to the nearest 0.01 µL.
- 3) Make injections into the egg as follows, allowing the eggs to be outside the incubator for not more than 30 minutes:
 - a. Wipe the blunt (air cell) end of the egg with 70% ethanol.
 - b. Gently make a hole in the egg with the Dremel drill.
 - c. Inject the vehicle or PCB mixture solution, 0.4 µL/g egg into the air cell, with a micro-pipettor and extended tip.
 - d. Seal the hole with paraffin.

- 4) Place eggs back into the incubator on their sides. Randomly place treatment groups in the egg racks. Avoid placing eggs in the very top, very bottom, very back and very front of the incubator.

Equipment Needed

Rubber and leather gloves

Foam filled case for egg transport

Betadine

Incubators: Natureform NMC2000 or GQF Sportsman 1502 or Kuhl

Pheasant egg trays

Light for candling

Ethanol and tissue or alcohol wipes

Dremel drill with fine point attachment

Paraffin and tool to apply it to eggs

Heating block

Scales (510 - 0.001 g) Mettler Toledo PG503-S

Rainin Pipettman with extended tips: one tip per egg

Data Sheets

“Avian Egg Collection Data Sheet”, “Incubator Record”, “Egg Treatment and Incubation Log”, “Log of checking eggs”, “Moisture Loss Data Sheet”, “Deformity Score Sheet”.

Literature Consulted

Smallwood, J. A., and D. M. Bird. 2002. American Kestrel (*Falco sparverius*). In *The Birds of North America*, No. 602 (A. Poole and F. Gill, eds.). The Birds of North America, Inc., Philadelphia, PA.

Pisenti JM, Santolo GM, Yamamoto JT, Morzenti AA. 2001 Embryonic development of the American Kestrel (*Falco sparverius*): External criteria for staging. *J. Raptor Res.* 35:194-206.

6.6 Egg Injection Procedure for chicken (*Gallus domesticus*) Eggs

This protocol outlines procedures for incubating eggs and injecting chemicals into the eggs of domestic chickens. The purpose of these procedures is to mimic maternal deposition of chemicals into the egg for determination of toxicity to the embryo or chick.

- 1) Upon receipt from supplies, store the eggs at 15°C until needed (but no more than 5 days).
- 2) Place eggs in an incubator in an egg rack of suitable size with the pointed end down. Incubate the eggs at 99.5°F and 55-65% humidity (84°F wet bulb) with hourly turning of the eggs.
- 3) On embryonic day four, candle the eggs and remove those that are infertile or which contain dead embryos.
- 4) Assign the eggs to treatment groups, mark each egg with unique code in pencil and weigh each egg to the nearest one tenth of a gram. Calculate and record the volume of dosing solution to be added to each egg. Round the volume to the nearest 0.01 μL .
- 5) Make injections into the egg as follows, allowing the eggs to be outside the incubator for not more than 15 minutes:
 - a. Wipe the blunt (air cell) end of the egg with 70% ethanol.
 - b. Gently make a hole in the egg with a 20 gauge needle.
 - c. Inject the vehicle or PCB mixture, 0.4 $\mu\text{L/g}$ egg into the air cell, with a micro-pipettor and extended tip or Hamilton syringe.
 - d. Seal the hole with paraffin.
- 6) Return eggs to the incubator.
- 7) On days five, eleven, and eighteen of incubation, candle the eggs and remove infertile and dead eggs.
- 8) On ED 18, transfer eggs to a 'hatcher' incubator, with each egg in an individual compartment.
- 9) Necropsy at approximately 24 hours after hatch.

Equipment Needed

Scales (510 - 0.001 g) Mettler Toledo PG503-S

Incubators: Natureform NMC2000

Egg trays

Light for candling

Ethanol and tissue or alcohol wipes

20 gauge needles

Rainin Pipettman with extended tips: one tip per egg
or Hamilton Syringe

Paraffin and tool to apply it to egg

Heating plate

Data Sheets: "Avian Egg Collection Data Sheet", "Incubator Record", "Egg Treatment and Incubation Log", "Log of checking eggs", "Moisture Loss Data Sheet", "Deformity Score Sheet".

6.7 Necropsy of Hatchling Birds

Hatchling birds are maintained in the incubator in which they hatch for 18-24 hours before necropsy to allow complete drying of feathers. Birds are sampled before 24 hours after hatch. This protocol outlines appropriate dissection techniques and sample storage conditions for several tissues including:

- Blood for serum
- Brain for hypothalamic monoamines analysis
- Liver for CYP450
- Fecal matter from lower intestine for estradiol and androgen measurements
- Thymus for mass and Bursa for mass and histology
- Thyroid for thyroid hormone radioimmunoassay
- Gonads for histology
- Tissues for RNA isolation**

- 1) Bring ten to twenty hatchlings at a time to the necropsy room and keep warm in a table top incubator.
- 2) Record time necropsy is initiated and completed. Record all data on appropriate data sheet.
- 3) Weigh the hatchling.
- 4) Kill the hatchling by cervical dislocation and decapitate with scissors. Immediately collect trunk blood into a 12x75 mm glass tube. Set tube aside allowing blood to clot for serum collection.

** For tissues requiring microarray analysis, transfer a liver sample or hypothalamus/brain sample to a microcentrifuge tube and snap freeze in liquid nitrogen . Store at -80°C.

- 5) Immediately remove the brain from the head, intact, and drop it directly into dry ice powder in an ice bucket. Make brain dissection as follows: cut along the sides of the skull on 3 sides including anterior; pull skull up; remove dura mater, break nerves and exteriorize the brain. After at least one minute on dry ice, fold the brain into a cold piece of aluminum foil and keep temporarily on dry ice.
- 6) By pinching around the cloaca, remove a fecal sample to a pre-tared 1.5 mL micro-centrifuge tube. Weigh and record the weight of the fecal sample. Store on dry ice.
- 7) Record the condition of the umbilicus and internal yolk sac, e.g. has it retracted properly into the abdomen? Is it adhered to the internal umbilicus? Tare a chemically clean jar on the scale and collect the yolk and transfer it to the jar, noting the weight.
- 8) Dissect the liver, remove the gall bladder and weigh the liver. Place the liver in a cryovial, or if it is too big for one vial, mince it and divide the tissue between multiple vials. Flash freeze the liver in liquid nitrogen for CYP450 analysis.
- 9) Dissect out the spleen, weigh it, then discard it.
- 10) Dissect out the heart, weigh it and preserve in appropriate fixative: Trim the heart of blood vessels in a standard manner from sample to sample, being careful not to remove any heart muscle. Weigh and then preserve the heart in 10% buffered formalin for 24-48hours. For storage longer than 48 hours, replace the formalin with 70% ethanol.
- 11) Remove both right and left thyroid at the same time. The thyroid is located at the caudal point of the thymus just anterior to the heart. Thyroid is within the thorax, ventral to and

bound by fascia to the carotid artery. Weigh both thyroids separately but store together in a 1.5 mL micro-centrifuge tube. Weigh the thyroids on plastic, weighing paper will cause desiccation. Freeze thyroids on dry ice.

- 12) Remove the bursa, weigh it and place it in a 1.5 mL microcentrifuge tube in appropriate fixative.
- 13) Identify the gonads to determine gender. Males have two kidney shaped testicles. Females have one left ovary. Leave the gonads intact and remove them on a portion of the back of the carcass. Fix in 10% neutral buffered formalin or other appropriate fixative.
- 14) Place the remaining carcass in freezer bags by treatment and place in the freezer.

Long term storage:

Brains, serum, liver, thyroids, feces: -80°C

Yolk: -20°C

Fixed tissue: room temperature

Tissue for RNA analysis: -80°C

Equipment Needed

Box

Heating plate

Scales sensitive to 0.00001 grams

(Mettler MT5)

Scales (510 - 0.001 g) Mettler Toledo PG503-S

Scissors for decapitation

Dissecting scissors and forceps

12x75 mm glass borosilicate tubes

Dry Ice Powder

Aluminum foil

Cryovials

1.5 mL microcentrifuge tubes

Bouin's Fixative or Davidson's Fixative

10% buffered formalin

Liquid nitrogen

RNA Later if needed

Labor: ideally a minimum of 4 people participate to ensure rapid dissection and storage of tissues

Data Sheets

“Hatchling Sampling Data Sheet

6.8 *Histological Analysis Of Avian Embryo Tissue: Gonads and Bursa of Fabricius*

This protocol describes the process of dehydrating, embedding, sectioning and staining tissue for histological study under light microscopy.

Procedure

1. Fix tissue with 10% buffered formalin
2. Dehydration:
 - a. 70% EtOH 1 hour
 - b. 85% EtOH 1 hour
 - c. 95% EtOH 40 minutes
 - d. 95% EtOH 40 minutes
 - e. 100% EtOH 40 minutes
 - f. 100% EtOH 1 hour
 - g. AMERICLEAR 40 minutes
 - h. AMERICLEAR 1 hour
 - i. Paraffin 40 minutes @ 56°C
 - j. Paraffin 1 hour @ 56°C
 - k. Paraffin 1 hour @ 56°C
3. Embed tissue in paraffin with desired orientation (longitudinal orientation for bursas; ****this is very important****). This step must happen quickly in order for the paraffin to solidify in one block. See note below for gonad orientation (##).
4. Section tissue embedded in paraffin into 5-10 μm sections. Make three slides. Place 3-5 sections on one slide. Use plus-treated slides, as the tissue will stay on the slides while they are taken through the washing and staining process.
5. Place slides in an oven or on a hotplate at 60°C for 30-60 minutes. This helps with keeping the sections bonded to the slide.
6. Stain the tissue by washing the slides as follows:
 - a. Xylenes 2 minutes
 - b. Xylenes 2 minutes
 - c. 100% EtOH 1 minute
 - d. 100% EtOH 1 minute
 - e. 95% EtOH 1 minute
 - f. 95% EtOH 1 minute
 - g. Tap water (non-running) 10 minute
 - h. Mayer's hematoxylin 15 minutes
 - i. Lukewarm running tap water 20 minutes
 - j. Eosin 2 minutes

k. Non-running tap water	1 minute
l. 95% EtOH	2 minutes
m. 95% EtOH	2 minutes
n. 100% EtOH	2 minutes
o. 100% EtOH	2 minutes
p. 100% EtOH	2 minutes
q. Xylenes	2 minutes
r. Xylenes	2 minutes
s. Xylenes	2 minutes

7. Set slides out to dry.

8. Mount cover slip with mounting medium after the slides are dry. Try not to use too much mounting medium.

Notes for gonads:

Cut tissue in 3-5 μM sections. Place multiple sections on each of at least 5 slides for every embedded block of tissue. For testes cut transverse. For ovaries, cut to acquire both end and middle regions, i.e., several sections of each of three regions per ovary. For ease cut the organ before dehydration and embedding.

Alternative staining: For immature gonads stain with Toluidine Blue. For mature ovaries stain with Toluidine Blue. For mature testes stain with Berg's stain to distinguish spermatozoa.

Endpoints:

Bursa: For bursa analysis, measure number of follicles per section, number of vacuoles, thickness of epithelial layer, and follicle size. Other qualitative aspects to also consider with each section is arrangement of bursal buds and arrangement of epithelial layers.

- a. Male gonad:
 - i. Differentiate cortex and medulla
 - ii. Area of seminiferous tubules
 - iii. Number of spermatozoa
 - iv. Abnormalities in Sertoli or Leydig cells and outer cortex
 - v. Presence of lipid filled vacuoles indicating steroidogenic activity
- b. Female gonad:
 - i. Number of primordial follicles and oocytes
 - ii. Differentiate cortex and medulla
 - iii. Relative area of cortex and medulla
 - iv. Abnormalities

Record a digital image of each section used.

Equipment Needed:

Tissue Tek(VIP1000) for tissue dehydration and embedding

Microtome for tissue sectioning

Warm water bath for mounting tissue onto slides

Hot plate for warming slides prior to washing

Additional Supplies:

Paraplast

10-20 baths for washing fluids

Plus-treated slides

Cover slips

Mounting medium

Hematoxylin

Eosin

Ethanol

Xylenes

Literature Consulted

Reaves, M.E. 2002. Effects of organochlorines on reproductive structures of wild fish-eating birds and domestic chickens. PhD Dissertation, Wright State University, Dayton, OH.

6.9 Ethoxyresorufin-o-dealkylase (EROD) Assay For CYP450 Activity In Liver Samples

This protocol outlines a method for preparing microsomes from hatchling avian liver tissue by differential centrifugation and measuring enzyme activity of cytochrome P450-1A (a monooxygenase or mixed function oxidase) in those microsomes. The expression/activity of CYP450 is elevated by exposure to xenobiotics. Halogenated aromatic hydrocarbons (HAHs), especially planar dioxins, furans and polychlorinated biphenyls increase CYP450 expression via interaction with the aryl hydrocarbon receptor.

Expression of CYP450 is a biomarker of exposure to HAHs. CYP450 is a phase I metabolic enzyme that has dealkylase activity. The enzyme activity is measured by assaying EROD (ethoxyresorufin-o-dealkylase) activity. 7-ethoxyresorufin serves as a substrate for this enzyme which yields resorufin as a fluorescent product. NADPH is the cofactor which donates electrons. The reaction is $\text{NADPH}_2 + \text{substrate} + \text{O}_2 \rightarrow \text{substrate-O} + \text{H}_2\text{O} + \text{NADP}$. Enzyme activity is expressed as nmol or pmol of resorufin per mg protein per minute. This protocol is adapted from Melancon (1997) and Brunstrom and Halldin (1998).

Reagents

Homogenization buffer: Na/K Phosphate pH 7.4

Stock A 0.2M KH_2PO_4 = 27.22 g/L Potassium Phosphate-Monobasic, using distilled water

Stock B 0.2 M Na_2PO_4 = 28.40 g/L Sodium Phosphate-Dibasic, using distilled water

The homogenizing buffer, at pH 7.4, is prepared by mixing approximately 1 part stock A to 4 parts stock B until a pH 7.4 is reached. (Stock A will decrease pH while stock B will increase it)

Microsomal resuspension buffer: 0.05M Na/K Phosphate containing 10^{-3} M Disodium Ethylenediamine Tetraacetate (EDTA), pH 7.6

Stock C = 6.80g KH_2PO_4 + 0.372g EDTA/L using distilled water

Stock D = 7.10g Na_2PO_4 + 0.372g EDTA/L using distilled water

The resuspension buffer is prepared by starting with Stock C and raising pH to 7.6 using Stock D.

Tris Buffer

0.066 M Tris-HCl, pH 7.4

Start with 7.99 g of Trizma base and lower pH to 7.4 or start with 10.4 g of Trizma HCl and raise pH with NaOH to 7.4.

For avian sample:

Stock Solutions::

1.00 mM solution of 7-ethoxyresorufin(Sigma E3763 “Resorufin ethyl ether”) in methanol (HPLC-grade)

2.0mM solution of Resorufin sodium salt in methanol (HPLC grade)

Working Solutions:

2.5 μ M 7-ethoxyresorufin in Tris-HCl

1 x 10⁻⁶ M resorufin in Tris-HCl

0.25 mM solution of NADPH in Tris-HCl

Procedure

Keep all samples on cool on wet ice between each step

- 1) Weigh excised liver (without gall bladder). Place the liver in a cryovial and flash freeze in liquid nitrogen if storage is necessary before microsomal preparation.
- 2) Thaw liver sample on crushed ice.
- 3) Using Melancon’s approach for small liver samples, homogenize the liver in **4X** of ice cold **homogenizing** buffer per **weight** of tissue. Keep the tube in a beaker of crushed ice while homogenizing. Homogenizer: motor driven stainless steel and teflon pestle in a glass homogenizing vessel is suggested. Use **no less than 1 mL** of buffer! Divide the homogenate into two. Retain one half of the homogenate for potential additional endpoints and continue the procedure with the second half of the homogenate.
- 4) Centrifuge homogenate for 20 min at 11,000g in the **pre-cooled**, 4°C, Eppendorf 5402 centrifug (~11500 rpm).
- 5) Collect **supernatant** from step 4 and centrifuge it at 100,000 xg for 60 min at 4°C to obtain microsomes. Use the Sorvall MC120 centrifuge, rotor RP80-T at 46,000 rpm. If extra volume in the tubes is needed, it is permissible to add more homogenizing buffer.
- 6) Resuspend microsomes in **resuspension** buffer. If necessary, aliquot samples into 2 or more tubes to allow one tube to be kept in the ultracold without all microsomes undergoing freeze thaw cycle. Store the microsomes in aliquots at -80°C and upon thawing vortex vigorously before proceeding. Multiple freeze/thaw cycles can affect activity. Refer to literature for guidance on microsome concentration for each species, or if unavailable, literature for a species with similar size liver to the species you are analyzing.

The quantity of microsomes used should be that which gives a linear response over the time of the assay, within the range of the standard curve. If enzyme activity is highly induced, one may need to re-run the assay with a smaller number of microsomes. In one microtiter plate run 7 triplicate samples, four resorufin standard curves of 10⁻⁸, 10⁻⁹, 10⁻¹², and 10⁻¹⁵ dilutions and 4 wells of non-induced reference microsomes, e.g. from Japanese quail. A total volume of 260 uL per well is optimal.

***As of August 2005, samples (S1, S2, etc.) were made in 1x(100%), 1:2(50%) and 1:4(25%) dilutions.

1:2 = 100ul of tris + 100ul of microsomes = 200ul, 50ul of this mixture per well and 50ul left for protein

1:4 = 150ul of tris + 50ul of microsomes = 200ul, 50ul of this mixture per well and 50ul left for protein

Example Plate Layout:

	1	2	3	4	5	6	7	8	9	10	11	12
A	250 ul 10 ⁻⁸	250 ul 10 ⁻⁹	250 ul 10 ⁻¹²	250 ul 10 ⁻¹⁵	S1 1x	S1 1x	S1 1x	S3 1:4	S3 1:4	S3 1:4	S6 1:2	S6 1:4
B	100ul 10⁻⁸ 100ul tris	100ul 10⁻⁹ 100ul tris	100ul 10⁻¹² 100ul tris	100ul 10⁻¹⁵ 100ul tris	S1 1:2	S1 1:2	S1 1:2	S4 1x	S4 1x	S4 1x	S6 1:2	S6 1:4
C	100ul tris 100ul B1	100ul tris 100ul B2	100ul tris 100ul B3	100ul tris 100ul B4	S1 1:4	S1 1:4	S1 1:4	S4 1:2	S4 1:2	S4 1:2	S6 1:2	S6 1:4
D	100ul tris 100ul C1	100ul tris 100ul C2	100ul tris 100ul C3	100ul tris 100ul C4	S2 1x	S2 1x	S2 1x	S4 1:4	S4 1:4	S4 1:4	S7 1x	S7 1:4
E	100ul tris 100ul D1	100ul tris 100ul D2	100ul tris 100ul D3	100ul tris 100ul D4	S2 1:2	S2 1:2	S2 1:2	S5 1x	S5 1x	S5 1x	S7 1x	S7 1:4
F	100ul tris 100ul E1	100ul tris 100ul E2	100ul tris 100ul E3	100ul tris 100ul E4	S2 1:4	S2 1:4	S2 1:4	S5 1:2	S5 1:2	S5 1:2	S7 1x	S7 1:4
G	100ul tris 100ul F1	100ul tris 100ul F2	100ul tris 100ul F3	100ul tris 100ul F4	S3 1x	S3 1x	S3 1x	S5 1:4	S5 1:4	S5 1:4	S7 1:2	S7 1:2
H	100ul tris 100ul G1	100ul tris 100ul G2	100ul tris 100ul G3	100ul tris 100ul G4	S3 1:2	S3 1:2	S3 1:2	S6 1x	S6 1x	S6 1x	S7 1:2	250ul Tris

- 7) Using a round bottom, 96 well plate, suitable for fluorescence assays: To each well add 200 uL of Tris buffer containing ethoxyresorufin and 50 uL of Tris buffer containing microsomes.
- 8) Preincubate in the dark at 37°C for 10 mins.
- 9) Add 10 uL of Tris buffer containing NADPH (0.25 mM) to each well and place the plate in the fluorescent reader. Take an initial reading then seven additional readings at 90 sec intervals (75 sec for each plate reading), excitation 544 nm, emission 590 nm. Calculate the results from the initial reading. The additional multiple readings serve only to determine that there is a stable increase in product over time. This serves as a quality control to ensure nothing unusual occurred.
- 10) Determine protein concentrations at the dilution used for the assay.
- 11) Calculate results using standard curve and protein concentration to nmol (or pmol) product per mg protein per minute.

Equipment Needed

Homogenizer

Centrifuge (11,000 xg)

Ultra centrifuge (100,000 xg)

Fluorescence microplate reader, excitation 544 nm, emission 590 nm

Literature Consulted

Melancon, M. J., 1997. Development of Cytochromes P450 in Avian Species as a Biomarker for Environmental Contaminant Exposure and Effect: Procedures and Baseline Values. Environmental Toxicology and Risk Assessment: Biomarkers and Risk Assessment (5th. Volume), ASTM STP 1306, David A. Bengston and Diane S. Henshel, Eds., American Society for Testing and Materials, Philadelphia.

Brunstrom, B, Halldin, K. 1998. EROD induction by environmental contaminants in avian embryo livers. Comparative Biochemistry and Physiology Part C 121: 213-219.

Personal Communication with Mark Melancon provided some detail.

6.10 High Pressure Liquid Chromatography (HPLC): Monoamine Analysis

Background:

HPLC analysis allows determination of monoamines in brain homogenates. This SOP covers machine set-up, standard preparation and typical tissue analysis used in our lab. The Bioanalytical Systems (BAS, West Lafayette, IN; also provides analysis software package) manual provides further information regarding machine set-up. Heat and light degrade catecholamines, so acidified samples are stored in the dark at -80°C . There is also freeze/thaw degradation, so standards and samples are prepared at assay concentrations the same day they are to be analyzed.

Brain sample monoamines are expressed per mg protein. Protein measurement protocol is listed under Protein Measurement.

These samples may be further analyzed for GnRH-I.

Materials:

Use all HPLC grade chemicals and deionized water (diH_2O).

We use a Bioanalytical Systems (BAS) HPLC with the following BAS parts and subunits:

Electrochemical detector unit (2 of these) CC-5

Temperature controller LC 22C

Amperometric Detector LC 4C

Solvent Delivery System PM 80

Data Acquisition System DA 5

Injection loop: 10 μl using a Hamilton 25 μl syringe (Fisher # 80275, 1702N, 22s/ 2²/3)

BAS software upgraded 12/03 to Epsilon LC, ver. 2.34B (epsilon-web.net)

Column: BAS # MF 6213, Phase II, ODS, 3 μm , 100x3.2 mm

Working electrode: BAS # F1000 glassy carbon

Reference electrode RE6 (pack of 3): BAS # MF2078

Some parts are available through BAS while others (e.g., stainless tubing pre-cut lengths) are available from Upchurch. BAS can recommend other suppliers.

Procedure:

Three steps are required to analyze samples with the HPLC:

I- Preparation of the standard, mobile phase and samples.

II- Setting the machine.

III- Data analysis: using the BAS software

I. Standard Preparation:

In order to develop a standard curve for each chemical, we inject (analyze) at least 2 concentrations for each compound. A concentration of 2ng/10 μl is diluted serially (1:2) to 1ng and 0.5 ng as follow:

Stock A- to be kept frozen –80°C

Compound name	abbr.	molecular weight of compound	molecular weight of parent compound	equivalent weight of parent compound	amount of compound weighed (gm)
L-Arterenol bitartrate	NE	319.3	169.2	25mg	0.0472
5-Hydroxytryptamine creatine sulfate complex	5HT	387.4	176.2	25mg	0.0549
5-Hydroxyindole-3-acetic acid	5HIAA	191.2	191.2	25mg	0.0250
3-Hydroxytryptamine	DA	189.6	153.2	25mg	0.0310

Weigh out 25.0 mg or equivalent weight from the previous table (last column) in separate volumetric flasks (5 mL.).

Dissolve in 5 ml. HCL 0.1 M.

To prepare 250 ml. of HCl 0.1M, add 2.05 ml. Conc. acid to 250 diH₂O.

Stock B (to be kept frozen in small brown glass vials)

Take 10 ul from each chemical of stock A and add to 25 ml of HCl 0.1M in separate 25 ml volumetric flasks

Stock C (prepared at the time it is to be used- keep on wet ice and in the dark)

Take 100 ul of each of the 8 standards from stock B and add 200 ul HCl 0.1 M, so the total volume is 1000 uL. This stock now is ready for injection. If an individual standard compound is to be injected, use 100 uL of that B stock standard and 900 ul HCl. If there will be a 4 compound standard (eg NE, 5HIAA, DA, and 5HT) use 100 ul of each of those B stocks and 600 ul HCl.

To inject, vortex prepared standard and draw up into a syringe (this will overfill the 10 ul loop). Follow the procedure for the BAS Epsilon software to inject/begin the data acquisition.

NOTE:

stock A- 25 mg is dissolved in 5 ml, so 10 ul contains $0.010 \times 5/25 = .050\text{mg} = 50 \text{ug}$

stock B- 50 ug is dissolved in 25 ml, so 100 ul contains $0.100 \times 50/25 = .200 \text{ug} = 200 \text{ng}$

stock C- 200 ng is dissolved in 1000 ul, so 10 ul (injection volume) contains $10 \times 200/1000 = 2 \text{ng}/10 \text{ul}$.

The maximum capacity or volume of the stainless steel loop going from the injector to HPLC is 10 ul. The loop is overfilled to be sure that it fills to its 10 ul capacity.

Mobile Phase: to prepare 1 Liter

Chemicals:

EDTA (0.67 mM): 0.250 g (MW 372.2)

Monochloroacetic Acid (MCAA, 0.13 M) : 12.3 g (MW 94.4)

Sodium Hydroxide (NaOH 0.1N): 4.5 g (MW 40)

Procedure

This acidic mobile phase is stable under refrigeration and can be prepared (4 L / time). Note that harmful chemicals are used in the preparation of this mobile phase. Dispose of waste chemicals using approved procedures.

Measured chemicals are added with enough diH₂O to bring the volume to 1L.

Adjust Ph to 3.1 using NaOH to raise pH or MCAA to lower it.

Add: Octyl Sodium Sulfate (SOS 0.86mM) : 0.200 g (MW 232.3)

Store in the refrigerator until use.

When ready to use the mobile phase:

Filter solution through a 0.2 um filter using a water vacuum system.

Allow the solution to degas 20-45 min.(stir under vacuum)

Prior to beginning a cycle of analysis and then periodically during that series, one may need to follow the BAS suggested procedure for cleaning the column using 100% acetonitrile followed by 40:60 acetonitrile:diH₂O. Inspect the BAS manual troubleshooting guide for indications of the latter condition.

Degas the required volume of the mobile phase (0.5L- 1.0L).

Add acetonitrile (HPLC-Grade; 2%) and Tetrahydrofuran, THF (HPLC-Grade; 0.1%). For each liter we need 20 mL acetonitrile and 1 mL THF. Measure this mobile phase + additions in a volumetric flask for consistency's sake.

Begin the process of establishing a quiet baseline and standard runs with good peak separation. This may take a few days.

Tissues sample preparation:

Add 500 ul of 0.1M HCl to tissue samples in 12 x 75 borosilicate tubes.

Using a small bore probe and an electric tissue grinder (PowerGen125, Fisher Scientific), homogenize to completely and evenly grind tissue. Avian brain punches will be homogenized in 5 sec or less (sample hypothalamus and median eminence with a 2mm biopsy scoop, Roboz Co. #6384). If a sample is larger and needs longer homogenizing, do not overheat; homogenize no more than 10 seconds and then return the sample to the ice bucket. Keep samples in refrigerator or on ice/ in dark except during the homogenization. Punched, non-homogenized samples may be stored at -80°C in 0.1 M HCl.

Rinse homogenizer in distilled water after after each sample, and blot dry. Change the rinse water with each sample.

Centrifuge homogenized samples at 4C for 20 min at 2000 RPM (or long enough to get a firm pellet) in the Sorvall RT6000-D centrifuge.

Pour the supernatant into labeled spin-filters, Costar Spin-x #8161. Centrifuge the spin-filters at 4°C for 40 min at 9000 rpm in an Eppendorf model 5402 centrifuge.

Keep the spin-filtered sample on ice in the dark together with its standard set until the entire sample group (for that day) has been processed (injected). After processing, the remainders of the standard and sample groups are stored in the dark at -80°C .

Comment

If samples are to be analyzed for GnRH-I, a minimum of 100 ul of supernatant is required for each assay replicate.

II. Setting the Machine:

The HPLC machine should display the following settings (turning off the power to the unit will return the settings to default values):

Potential level : +

Manual offset : LEFT toggle:OFF & RIGHT toggle:100

Temperature : 30°C keep the detector cell cabinet shut during operation to better stabilize the temperature of the column

Output : as close to 0 as possible; toggle the autozero lever to zero when needed to keep the reading near 0

AppE potential : 0.800

E gen : -0.200 V

I gen : .000 uA

Flow rate : 1.0 mL/ min this is during active operation. Reduce flow rate to .3 ml/min during non-test/ off use periods

Local/ remote : local

Detector Range : W1, the top detector in our arrangement, 5 nA: W2, the bottom detector in our arrangement, 2 nA. Remember that 5 nA is the LESS sensitive, and 2 nA is the MORE sensitive

Typical flow pressure for our mobile phase is 3000-3400 psi. The pressure will be lower for 100% acetonitrile and 40:60 acetonitrile:water solutions. The machine is set to shut down at pressures greater than 4000 psi. The electrochemical detector is left ON at all times except during servicing at which time it is in STANDBY. During off/non test periods (e.g., overnight) the flow rate is reduced and the temperature controller is turned off.

Adjusting the range depends on the concentration of the chemical on the tissue. The lower the value the higher the sensitivity.

Reference:

C.S. Duchala, M.A. Ottinger, and E. Russek. 1984 The Developmental Distribution of Monoamines in the Brain of Male Japanese Quail (*Coturnix coturnix japonica*). Poultry Science 63:1052-1060.

6.11 GnRH-I ELISA

This assay measures GnRH-I in tissue extracts or serum.

Procedure

1. Coat plates, 50 uL/ well, goat anti rabbit IgG (5ug/ml in coating buffer) for 1 hour @37⁰ C or overnight @ 4⁰C
2. Wash plate 3 times with wash buffer, 250 uL/ well, then blot plate
3. Pipet PEG (2%) blocker, 200 uL/ well, then incubate for 1 hour @ 37⁰ C
4. Wash plate 3 times with wash buffer, 250 uL/ well, then blot plate
5. Pipet GnRH (1:200 of purified Ab in standard diluent), 50 uL/ well, then incubate for 1.5 hour @ 37⁰ C
6. Wash plate 3 times with wash buffer, 250 uL/ well, then blot plate
7. Pipet standards or samples (prepared in standard diluent, SEE **Notes**.) 100 uL/ well, then incubate for 1.5 hour @ 37⁰ C
8. DO NOT WASH PLATE BEFORE THE NEXT STEP
9. Pipet GnRH conjugate (1:2500 in standard diluent), 50 uL/ well, then incubate for 1.25 hour @ 37⁰ C
10. Wash plate 5 times with wash buffer, 250 uL/ well, allowing the last wash to remain on the plate, room temperature, 10 minutes, then blot plate
11. Pipet SAP (1:2000 in enzyme diluent), 50 uL/ well, then incubate for 0.5 hour @ 37⁰ C
12. Wash plate 5 times with wash buffer, 250 uL/ well; pipet 200 uL/ well PBS, allowing the buffer to remain on the plate, room temperature, 10 minutes, then blot plate
13. Pipet PNPP substrate (1 mg/ ml in substrate buffer), 100 uL/ well, then incubate for at least 1 hour @ 37⁰ C while periodically checking color development
14. Read the plate @ 405 nm and plot using a cubic-spline vs. log-linear method.

Standards: pipet in duplicate with column placement for best space efficiency. Use 7 serial 1:10 dilutions of Avian LhRH with 1 ug/ml as the highest concentration and an 8th well for a standard diluent blank.

The standards will have these concentrations:

1. 1 ug/ ml
2. .1 ug/ ml
3. .01 ug/ ml
4. .001 ug/ ml
5. .0001 ug/ ml
6. .00001 ug/ ml
7. .000001 ug/ ml

Plate sealers may be reused as long as they are completely sealing the rim of the plate to prevent dehydration of the wells.

Plates may be coated in advance and stored; plates sealed and in a plastic bag, @ -200 C. follow the method through step 3. Do not remove the PEG before freezing.

PNPP is light sensitive. Prepare it at the beginning of step 11 to allow it time to dissolve. If there is a pronounced yellow color, the PNPP is unusable.

Follow good lab techniques in handling and pipetting ELISA plates. Touching the underside of the plate or splashing reagents out of a well will distort readings. Examine the %CV for any duplicates; only less than 10% is acceptable. See the HPLC method for preparing tissue extracts. Serum is used without any treatment. There must be 100 μ L of any standard or sample. Any volume less than that for a sample can be corrected using standard diluent and the dilution of the sample noted for later mathematical correction.

Coating Buffer (pH 9.6)

1. Weigh 15.9 g Na_2CO_3 .
2. Weigh 29.3 g NaHCO_3 .
3. Add all into 1000 mL Distilled water.
4. Adjust pH to 9.6.
5. Store in refrigerator.

Standard Diluent- Stock Solution (1% BSA, 1% NaN_3 solution in 10X PBS)

1. Weigh 0.5 g BSA (bovine serum albumin).
2. Weigh 0.5 g NaN_3 .
3. Add both to 50 mL 10X PB
4. Mix well.

*** Put 5 mL of solution in 50 mL plastic tubes and store at 0 degrees C (or put 1.5 mL in 15 mL tubes)***

Standard Diluent – Working Solution

1. Weigh 0.844 g NaHCO_3 .
2. Weigh 1.255 g Na_2CO_3 .
3. Weigh 0.203 g MgCl_2 .
4. Add all to 1000 mL Distilled water and mix well.

Wash Buffer

1. Measure out 8 mL of Tween-20.
2. Weigh 4 g of NaN_3 .
3. Measure out 400 mL of PBS (10X).

Add all to 3600 mL of Distilled Water and mix well.

Equipment Needed

Platwasher

Pipettors, repeater pipetors, and multi-tip pipettors

pH meter

96 well plates with disposable sealers

warming oven

refrigerator

-70⁰ C freezer

centrifuge

Homogenizer

Literature Consulted

Q. Li, L. Tamarkin, P. Levantine, M.A. Ottinger. 1994 Estradiol and Androgen Modulate Chicken Luteinizing Hormone-Releasing Hormone-I Release in Vitro. *Biology of Reproduction* 51: 896-903.

6.12 Protein Assay

This protocol describes a procedure for measuring protein content using a commercially available kit, it is used with the GnRH-I assay and HPLC analysis of hypothalamic monoamines to generate protein values. BCA Protein Assay Reagent Kit (Pierce 23225X)

Procedure:

Making Standards:

Always run a test curve with a few samples to see if they fall in the middle range of the curve. If the samples do not fit the curve, dilute the samples appropriately.

1. Standard #1 is the bovine serum albumin provided in the kit (2mg/mL)
2. Standard #2 is 0.5mL of standard #1 mixed with 0.5mL 1X phosphate buffered saline (PBS).
3. Standards #3-7 are serial dilutions of standard #2, made with 0.5mL of standard and 0.5mL 1X PBS.
4. Standard #8 is 1X PBS (a negative control).

Preparing Working Solution of Assay Reagent:

1. Mix 50 parts solution A with 1 part solution B.
2. If using the test tube method, mix 50mL of solution A with 1 ml solution B for one plate.
3. If using the microtiter plate method mix 25mL solution A with 0.5mL solution B for one plate.

Assay Using Microtiter Plates:

1. Measure samples in duplicates at a minimum.
2. Add 10 μ L of standard and samples into wells.
3. Quickly add 200 μ L of working assay reagent into each well.
4. Mix by gentle agitation and incubate 30 minutes at 37°C
5. Remove plate from incubation and allow to equilibrate at room temperature for 15 minutes.
6. Read at 570nm on Bio-Rad plate reader.

6.13 Extraction Of Steroid Hormones From Avian Feces

For samples from HATCHLINGS: Weigh feces WHEN THEY ARE COLLECTED - you will extract the entire sample. If feces were not weighed at collection they must be weighed before extraction which can result in sample loss. The extraction should be done in glass.

Add 1 mL of distilled water and 9 mL of 100% ethanol (ETOH) to each tube. If running an extraction efficiency the samples can be spiked with radiolabeled hormone (i.e. probably 20 uL of hot estradiol and 20 uL of hot testosterone¹). The labeled/unlabeled hormone should be added first along with the water and then vortexed for one minute. After vortexing the ethanol should be added and vortexed again for one minute. Remember everything is radioactive from now on so proceed accordingly.

Cap tubes with purple/green lids and parafilm on top of the caps.

Secure tubes horizontally to a shaker/rocker (room 3144) for end-to-end mixing **OVERNIGHT** (12 hr minimum).

The next day centrifuge samples at ~3000 RPM for 25minutes at 4°C using the table top Sorvall centrifuge in the RIA room.

Pour off supernatant into a clean 16 x 125 test tube. **SAVE THE SUPERNATANT** and discard old sample tube (*into radiation waste if necessary).

Place tubes with supernatant into water bath heated to 60 °C. (Any hotter and the supernatant might pop/boil resulting in lost sample and possible radioactive contamination). Place the water bath under the multi-tube manifold apparatus in the hood (room 3144) and gently blow off the ethanol using nitrogen gas. The combination of heat from the water bath and nitrogen gas will cause the ethanol to evaporate off much quicker.

Once the tubes are **completely dry**, resuspend the samples in 3 mL of 100% ETOH. Sonicate and vortex for one minute each.

This next step is designed for analysis of small samples, i.e., hatchlings, and is necessary for measuring estradiol and androgens from the same extract with minimal dilution of the hormone because we use two different diluents for the two assays.

¹ Before you add tracer to the extraction you need to check expiration date for the estradiol and cpm's for the testosterone - see androgen RIA SOP for the appropriate CPM range for the testosterone and check it on the scintillation counter and run the efficiency from the same batches of tracer. Write down everything you do clearly.

Remove 1.5 mL (i.e. one half) of the ethanol/hormone extract to a new tube. Pipet accurately since ethanol is volatile and tends to leak out of the pipette tip if not transferred quickly. Label one extract tube 'E' for estradiol and one extract tube 'A' for androgens.

Return samples to water bath/nitrogen gas setup and evaporate off ETOH for the final time. The hormone has now been extracted from the fecal sample and should be resuspended in RIA standard diluent depending on RIA needs:

Resuspend the 'E' tube extract in MP Biomedicals E₂ RIA kit - zero pg/mL standard. Since the estradiol assay requires 50 uL of extract per tube, should be done in duplicate and you need to have enough extract to measure hormone recovery, resuspend the dried 'E' extract in 150 uL of zero standard - that should yield 2x 50 uL for assay, 1x 20 uL for recovery and extra that will be stuck in the tube and not removable.

Resuspend the 'A' tube extract in androgen assay standard diluent. Remember there are two recipes for this diluent - one from the old fecal extract protocol and one from the RIA protocol. It probably doesn't matter which one you use, just be consistent although I would say use the recipe from the RIA SOP. Since the androgen assay requires 100 uL of extract per tube, should be done in duplicate and you need to have enough extract to measure hormone recovery, resuspend the dried 'A' extract in 250 uL of zero standard - that should yield 2x 100 uL for assay, 1x 20 uL for recovery and extra that will be stuck in the tube and not removable.

Sonicate and vortex the reconstituted extracts and allow equilibration for 30 minutes, i.e., let stand at room temperature.

Securely cap the extracts and freeze at -80°C until analysis. They are radioactive - label and store them appropriately!!

Analyze extracts for androgens and estradiol using SOPs: "Radioimmunoassay of Androgens In Avian Serum or Fecal Extract" and "Radioimmunoassay of Estradiol In Avian Serum or Fecal Extract"

This will ensure that the sample hormone will not be overdiluted and still in the range of the standard curve for the RIA. IF after all this, the hormone is not detectable on the standard curve, you will probably have to reduce the amount of diluent and only run the samples in singlets, not duplicates.

For extraction efficiencies (these are done at the time of RIA):

Estradiol: For each sample place 20 uL of extract into a tube for counts for recovery. Prepare a 'total count' tube containing 20 uL of the hot estradiol, i.e. exactly what you spiked the fecal sample with. Count on the gamma counter. Cpms of extract *2 then divided by cpms of total count and the result multiplied by 100 will give the extraction efficiency as a percent. The sample cpms are multiplied by two because we divided the sample into two after extraction.

Androgens: For each sample, place 20 uL of extract into a scintillation vial. Add 5 mL of scintillation cocktail. Prepare 'total count' vials with 20 uL of hot androgen that you originally

spiked the sample with and add 5 mL scintillation cocktail. Measure cps on the scintillation counter. As for estradiol:

$\frac{\text{CPM extract} * 2}{\text{CPM total counts}}$ and multiply result by 100 to express % extraction efficiency

6.14 Radioimmunoassay of Androgens In Avian Serum or Fecal Extract

The purpose of this protocol is to measure androgens in serum from blood of birds. The method and reagents described here have been validated for hatchling and adult Japanese quail and adult bobwhite quail. To use this method with any other species or age requires validation. The radioimmunoassay is a competitive binding assay that includes use of a tritiated testosterone and requires training in radiation safety for any personnel working on the assay. The method was validated for quail by serum extraction of steroids and parallelism experiments.

Procedure

Steroid extraction with spiking for recovery

- 1) Calculate the amount of serum required for the assay. This is dependent upon results of the validation which should be done whenever a series of assays is started, whenever a procedure is adapted (different sex, age or species) or when a different kit is tried. Serum may need to be concentrated or diluted depending on androgen concentrations.
- 2) For serum volume less than 1 mL, double extract with 2 mL ethyl ether:
 - a. Add serum to a 16x100 mm borosilicate glass tube labelled 'A' and with the sample number.
 - b. Spike sample with 10 μL ^3H testosterone ('hot') to estimate extraction efficiency
 - c. Under the fume hood add 2 mL of ether and vortex vigorously.
 - d. Wait for the serum and ether layers to separate (~1 min).
 - e. Freeze the sample either in a ethanol and dry ice bath in a beaker or covered with foil in a -80° freezer if working with many samples. The serum will freeze, the ether fraction will not.
 - f. Pour off the ether fraction into tube 'B'.
 - g. Allow the serum fraction in tube A to thaw and repeat steps c to f.
 - h. Dry down the sample by evaporation of the ether leaving the steroid residue on the glass tube either by leaving the tubes in a fume hood overnight or by forcing nitrogen gas over the sample in a fume hood for rapid drying.
- 3) Assay requires the following volumes (samples should be assayed in duplicate unless samples are from hatchlings and volume is limiting):

Japanese Quail:

Hatchling (male) serum assayed in singlets (most samples have to be pooled): 120 μL , assayed in duplicates, 220 μL . 100 μL per assay tube, 10 μL for recovery, 10 μL lost on tube. Serum reconstituted 1:1.

Adult serum assayed in duplicate: 250 μL , reconstituted 1:1 for males and females yielding 250 μL : 100 μL per assay tube and 20 μL for recovery = 220 μL .

Bobwhite Quail:

Adult serum assayed in duplicate: 150 μL reconstituted 1:2 to yield 300 μL serum for assay: 100 μL per assay tube and 20 μL for recovery = 220 μL .

Validation/Parallelism

- 1) Acquire a sample of pooled serum. Volume is dependent upon number of serial dilutions or repeated measurements to be done. An example is as follows:

To validate a 2X concentrate and 1:1 through 1:16 serial dilution in duplicate requires 600 μL of serum. Double extract the 600 μL and reconstitute in 300 μL of standard diluent. This yields a 2x concentrated sample. Remove 150 μL of this 2X concentration to a new tube, dilute it 1:1 with zero standard and label the tube "X" for the 1:1 reconstitution. From the "X" tube remove 150 μL to a tube labelled "1:2" and add 150 μL standard diluent. From that "1:2" dilution remove 150 μL to another tube, labelled "1:4" and add 150 μL standard diluent to yield a 1:4 dilution. Continue as such to create a 1:8 and 1:16 dilution. From these serial dilution tubes take 50 μL of reconstituted sample for each assay tube (assay in duplicate). Follow the methods for the radioimmunoassay below. Compare the resulting data to the standard curve. The serially diluted samples' curve should parallel that of the standard and final concentrations should be close to one half of the higher dilution for each sample. The non-linear portion of the standard curve is usually the upper and lower 5%. Unknowns in these high and low areas of the curve are more likely to have some error. Therefore, samples are run at a dilution predicted to be in the working range of the assay. Precision is determined by repeated (5-8 samples) assay of a sample. The CV should be no more than 10%. Sample data in Appendix A.

Radioimmunoassay

- Run a standard before you begin to check that the assay and all reagents are acting normally.

1. Label 12x75mm borosilicate glass tubes in duplicate with Total, NSB (non-specific binding), Bo and S1-S10 and then number tubes of unknowns in duplicate.
2. Prepare standards:

Take 10 μL of testosterone standard solution C and dilute it in 990 μL standard diluent yielding 1 mL of 10 ng/mL, this is solution D and is not used in the assay.

Prepare 10 standards by serial dilution solution D to be used in the assay and designated S1 to S10.

S10 = 250 μL of solution D plus 250 μL of standard diluent and is 5 ng/mL.

S9 = 250 μL of S10 plus 250 μL standard diluent and is 2.5 ng/mL.

S8 = 250 μL of S9 plus 250 μL standard diluent and is 1.25 ng/mL.

S7 = 250 μL of S8 plus 250 μL standard diluent and is 0.625 ng/mL.

S6 = 250 μL of S7 plus 250 μL standard diluent and is 0.3125 ng/mL.

S5 = 250 μL of S6 plus 250 μL standard diluent and is 0.156 ng/mL.

S4 = 250 μL of S5 plus 250 μL standard diluent and is 0.08 ng/mL.

S3 = 250 μL of S4 plus 250 μL standard diluent and is 0.04 ng/mL.

S2 = 250 μL of S3 plus 250 μL standard diluent and is 0.02 ng/mL.

S1 = 250 μL of S2 plus 250 μL standard diluent and is 0.01 ng/mL.

3. Prepare anti-androgen antibody: 100 μ L antibody to 9.9 mL standard diluent.
4. Add 100 μ L standard diluent to Total, NSB and Bo tubes.
5. Add 100 μ L of standards to each respective tube.
6. Add 100 μ L of unknowns and control sera to each respective tube.
7. Add 100 μ L of antibody solution to all tubes EXCEPT total and NSB.
8. Add 150 μ L of 3H-testosterone to all tubes.
9. Cover, vortex and incubate at room temperature for 1.5-2 hours.
10. Place tubes in ice bath for 5 minutes to stop the reaction.
11. Add 400 μ L per tube of dextran coated charcoal EXCEPT total tubes, add 400 μ L phosphate buffer to total tubes.
12. Vortex and incubate on ice for 15 minutes.
13. Centrifuge 3000 rpm on Sorvall table top centrifuge for 15 minutes at 2-8°C.
14. Remove 0.5 mL supernatant, carefully without disturbing the charcoal, and add to a scintillation vial.
15. Add 5 mL of scintillation cocktail and vortex vigorously.
16. Count on scintillation counter for four minutes per sample.

Guidelines:

NSB should be less than 10% of the total count.

Bo should be 20-25% of the total count.

CV of duplicate samples should be less than 5% ideally.

Waste Disposal

A and B tubes from the extraction, assay tubes must be disposed of in containers provided by radiation safety for ^3H waste (cardboard containers). Scintillation tube waste must be placed in the appropriate drum. Amount of radiation must be calculated and listed on waste containers.

All tips used with radioactive samples must be disposed of in a plastic bag and the bag disposed of in the dry waste container.

Conduct wipe tests of lab and equipment on a monthly basis to check for radioactive contamination.

Calculations

Using Graphpad Prism, convert counts per minute to androgen concentration as follows:

1. Create new project
2. Format data for duplicate 'Y' values.
3. Type standard curve concentrations into X values column
4. Type standard curve counts per minute (cpm) and unknown cpms into A-Y values column
5. Type NSB cpms into B-Y values column
6. "Analyze" data - remove baseline and column math
Baseline values = B, D, F
Calculate = difference: value-baseline

- Replicates = calculate the mean of the replicates
7. "Analyze" data - transforms
 $X = \log(X)$
 $Y = Y/K$ (where $K=B_0$ (baseline subtracted)/100)
 8. "Analyze" data - nonlinear regression (curve fit)
 Choose an equation = sigmoidal dose response (variable slope)
 Also calculate = unknowns from standard curve
 9. Pull down: interpolate X values
 10. "Analyze" data - transforms $X = 10^X$
 11. "Graph" data using $\log(X)$ vs. % binding

Equipment Needed

Ethyl ether
 Borosilicate glass tubes, 16x100 and 12x75 mm, tube racks
 Pipettors
 Ice
 Vortex
 Radioactive waste disposal containers
 Scintillation Counter

Reagents

0.1 M Phosphate Buffer, pH 7.6

K_2HPO_4	13.93g
NaH_2PO_4	3.12g
NaCl	8.76g
NaN_3	1.0g
DDW	to 1000 mL

Adjust pH to 7.6 and store at 4°C

Standard Diluent, pH 7.4

0.1 M phosphate buffer	100 mL
DDW	900 mL
NaCl	7.6g

Bovine Serum Albumin, fraction V, Sigma A-7888, suitable for insulin RIA 10g
 NaN_3 1.0g
 Adjust to pH 7.4 and store at 4°C

Dextran Coated Charcoal

Activated charcoal	1.5g
Dextran	0.15g
0.1 M phosphate buffer	300 mL

Mix and store at 4°C (keep away from the light)

Testosterone Standard


1. Weigh 10 mg powdered testosterone and dissolve in 10 mL EtOH to make a 1000 $\mu\text{g/mL}$ solution and label this 'A'.
2. In a new tube dilute 100 μL of A with 900 μL EtOH to make solution B (100 $\mu\text{g/mL}$).
3. Prepare solution C with 10 μL of B into 990 μL of standard diluent (1000 ng/mL).
4. Make 20 μL aliquots of solution C and store at -80°C .

Testosterone from Amersham Biosciences, UK.

1,2,6,7- ^3H -testosterone, 250 μCi , 250 μL , catalog # TRK402, store at -20°C .
 "Tracer" 3-5 μL of tritiated testosterone in 100 mL standard diluent and stir well.
 Count 20 μL with scintillation cocktail. Want cpm between 1000 and 2000.

Anti-androgen from Esoterix (previously Endocrine Sciences, Inc) Calabasas Hills, CA

Store at -80°C , rehydrate per directions on vial (1.2 mL water) and prepare 110 μL aliquots, refreeze at -80°C . Working solution (prepare shortly before use) = 100 μL of frozen stock in 9.9 mL of standard diluent (1:100 dilution).

AppendicesA. Sample validation/parallelism data:**Transform of Baseline-corrected of Data 1:Transformed data**

On graph:

A-B is standard

C-D is serial dilution of unknown

On table:

A-B is percent binding (Y axis)

Std Conc are unknowns from greatest dilution to 2X concentrate (X axis shown log transformed).

Literature Consulted

Bales, KL, Abdelnabi, M, Cushing, BS, Ottinger, MA, Carter CS. 2004 Effects of neonatal oxytocin manipulations on male reproductive potential in prairie voles. *Physiology and Behavior* 81: 519-526.

Hahn, DC, Hatfield, JS, Abdelnabi, M, Wu, J, Igl, LD, Ottinger, MA. In press. Inter-species variation in yolk steroid levels in a cowbird host comparison. *Journal of Avian Biology*.

Ottinger, MA, Pitts, S, Abdelnabi, MA. 2001 Steroid hormones during embryonic development in Japanese Quail: Plasma, gonadal and adrenal levels. *Poultry Science* 80:795-799.

Ottinger, MA and Brinkley, HJ. 1979. Testosterone and sex related physical characteristics during the maturation of the male Japanese quail. *Biology of Reproduction* 20:905-909.

6.15 Radioimmunoassay of Estradiol In Avian Serum or Fecal Extract

The purpose of this protocol is to measure estradiol in serum from blood of birds. The method and reagents described here have been validated for hatchling and adult Japanese quail and adult bobwhite quail. To use this method with any other species or age requires validation. The radioimmunoassay is a competitive binding assay that includes use of a iodinated estradiol and requires training in radiation safety for any personnel working on the assay. Currently we use a double-antibody kit from MP Biomedicals designed for human serum. The kit was validated for quail by serum extraction of steroids and parallelism and precision experiments. Steroids are first extracted from the serum sample by ether partition which removes interfering aqueous compounds.

Procedure

Steroid extraction with spiking for recovery

- 1) Calculate the amount of serum required for the assay. This is dependent upon results of the validation which should be done whenever a series of assays is started, whenever a procedure is adapted (different sex, age or species) or when a different source of reagents is tried. Serum may need to be concentrated or diluted depending on estradiol concentrations.
- 2) For serum volume less than 1 mL, double extract with 2 mL ethyl ether:
 - a. Add serum to a 16x100 mm borosilicate glass tube labelled 'A' and with the sample number.
 - b. Spike sample with 10 μL ^{125}I estradiol ('hot') to estimate extraction recovery (efficiency) and incubate for at least 30 minutes at room temperature to equilibrate.
 - c. Under the fume hood add 2 mL of ethyl ether and vortex vigorously.
 - d. Wait for the serum and ether layers to separate (~1 min).
 - e. Freeze the sample either in a ethanol and dry ice bath in a beaker or covered with foil in a -80° freezer if working with many samples. The serum (bottom, aqueous layer) will freeze, the ether fraction will not (top layer including steroid hormones)
 - f. Pour off the ether fraction into corresponding tube 'B'.
 - g. Allow the serum fraction in tube A to thaw and repeat steps c to f.
 - h. Dry down the sample in tube B by evaporation of the ether leaving the steroid residue on the glass tube either by leaving the tubes in a fume hood overnight or by forcing nitrogen gas over the sample in a fume hood for rapid drying.
- 3) Recent assays have required the following volumes (samples should be assayed in duplicate unless samples are from hatchlings and volume is limiting):

Japanese Quail:

Hatchling (female) serum assayed in singlets (some samples have to be pooled):
 80 μL , assayed in duplicates, 130 μL . 50 μL per assay tube, 20 μL for recovery, 10 μL lost on tube. If assayed in singlets. Serum reconstituted 1:1.

80µL of serum is extracted as described above then reconstituted with the same amount of buffer: 50µL for the assay and 20 µL for calculating the recovery. Adult serum assayed in duplicate: 100 µL plasma extracted then reconstituted (1:4) with buffer for females yielding 400 µL and 1:3 for males yielding 300 µL: 50 µL per assay tube and 50 µL for recovery = 150 µL.

Bobwhite Quail:

Adult serum assayed same as adult Japanese quail serum.

Validation/Parallelism

1. Acquire a sample of pooled serum from each new species. Volume is dependent upon number of serial dilutions or repeated measurements to be done. An example follows: To validate a 2X concentrate and 1:1 through 1:16 serial dilution in duplicate requires 600 µL of serum. Double extract the 600 µL and reconstitute in 300 µL of standard diluent solution. This yields a 2x concentrated sample. Remove 150 µL of this 2X concentration to a new tube, dilute it 1:1 with zero standard and label the tube "X" for the 1:1 reconstitution. From the "X" tube remove 150 µL to a tube labelled "1:2" and add 150 µL zero standard. From that "1:2" dilution remove 150 µL to another tube, labelled "1:4" and add 150 µL zero standard to yield a 1:4 dilution. Continue as such to create a 1:8 and 1:16 dilution. From these serial dilution tubes take 50 µL of reconstituted sample for each assay tube (assay in duplicate). Follow the methods for the radioimmunoassay below. Compare the resulting data to the standard curve. The serially diluted samples' curve should parallel that of the standard and final concentrations should be close to one half of the higher dilution for each sample. The non-linear portion of the standard curve is usually the upper and lower 5%. Unknowns in these high and low areas of the curve are more likely to have some error. Therefore, samples are run at a dilution predicted to be in the working range of the assay. Precision is determined by repeated (5-8 samples) assay of a sample. The CV should be no more than 10%. Sample data in Appendix A.

Radioimmunoassay

1. Follow the directions given in the MP Biomedicals protocol. A description follows:
2. Bring all reagents to room temperature. Reconstitute extracted samples with zero standard, vortex and allow to stand at least 30 minutes.
3. Label 12x 75 mm tubes 1-16 for standards and 17+ for unknowns. 1-2 is non-specific binding (NSB), 3-4 is zero with 15-16 being 3000 pg/mL.
4. Add 500 µL of the steroid diluent into tubes 1-2.
5. Add 50 µL of zero standard into tubes 1-4 and 50µL from each standard to its respective tube. (Known amount in duplicate 5-16). Add 50 µL of each unknown and control sera to each appropriate tube.
6. Vortex each sample and add 50 µL of each into its appropriate tubes.
7. Add 500 µL of ¹²⁵I estradiol (blue) to all tubes.
8. Add 500 µL of anti-estradiol antibody (yellow) to all tubes EXCEPT 1-2.
9. Vortex, cover tubes with foil, label as radioactive, and incubate for 90 minutes in a 37°C waterbath.

10. Add 500 μL of precipitant solution (red) to all tubes and vortex.
11. Centrifuge all samples at 2300 rpm ($\sim 1000 \text{ xg}$) for 25 minutes* at 2-8°C on the Sorvall table top centrifuge. (*This is a slight increase in time from the MP Biomedicals protocol and reduces slippage of pellets when decanting).
12. Decant supernatants into a radioactive waste container and blot each tube on absorbent bench paper. If pellets start to slip, centrifuge the samples again for 5 minutes.
13. Count the precipitate on a gamma counter calibrated for ^{125}I for 1 minute. Also count two or more tubes of 10 μL tracer each for Total Counts for recovery calculations.

Waste Disposal

A and B tubes from the extraction, assay tubes must be disposed of in containers provided by radiation safety for ^{125}I waste (blue plastic barrels). Liquid ^{125}I waste must be poured into plastic carbuoy supplied by radiation safety. Amount of radiation must be calculated and listed on waste containers. All tips used with radioactive samples must be disposed of in a plastic bag and the bag disposed of in the dry waste container. Conduct wipe tests of lab and equipment on a monthly basis to check for radioactive contamination.

Calculations

Using Graphpad Prism, convert counts per minute to estradiol concentration as follows:

1. Create new project
2. Format data for duplicate 'Y' values.
3. Type standard curve concentrations into X values column
4. Type standard curve and unknowns cpms into A-Y values column
5. Type NSB cpms into B-Y values column
6. "Analyze" data - remove baseline and column math
 Baseline values = B, D, F
 Calculate = difference: value-baseline
 Replicates = calculate the mean of the replicates
7. "Analyze" data - transforms
 $X = \log(X)$
 $Y = Y/K$ (where $K = B_0$ (baseline subtracted)/100)
8. "Analyze" data - nonlinear regression (curve fit)
 Choose an equation = sigmoidal dose response (variable slope)
 Also calculate = unknowns from standard curve
9. Pull down: interpolate X values
10. "Analyze" data - transforms $X = 10^X$
11. "Graph" data using $\log(X)$ vs. % binding

Equipment Needed


17 β estradiol double antibody kit
 # 07-138-102 MP Biomedicals
 Ethyl ether
 Borosilicate glass tubes, 16x100 and 12x75 mm, tube racks
 Pipettors

Extra 0 pg/mL standard (“zero standard”)
 # 07138171 for 50 mL MP Biomedicals
 Radioactive waste disposal containers
 Gamma Counter

Appendices

A. Sample validation/parallelism data:

Transform of Baseline-corrected of Data 1:Transformed data



On graph:
 A-B is standard
 C-D is serial dilution of unknown
 On table:
 A-B is percent binding (Y axis)
 pg/mL are unknowns from greatest dilution to 2X concentrate (X axis shown log transformed).

Literature Consulted

This method and MP Biomedicals reagents have recently been validated for Japanese and bobwhite quail. Prior work was similar but with different sources for reagents.

Ottinger, MA, Pitts, S, Abdelnabi, MA. 2001 Steroid hormones during embryonic development in Japanese Quail: Plasma, gonadal and adrenal levels. *Poultry Science* 80:795-799.

Watson, JT, Abdelnabi, M, Wersinger, X, Ottinger, MA, Adkins-Regan, E. 1990 Circulating Estradiol and activation of male and female copulatory behavior in Japanese quail (*Coturnix japonica*). *General Comparative Endocrinology* 77:229-238.

6.16 Digestion Of Thyroid Gland And Radioimmunoassay of Thyroid Hormones

The thyroid gland is site of thyroid hormone synthesis and storage. While measuring circulating thyroid hormone in the blood is a potentially useful endpoint, measurement of thyroid hormone stores in the thyroid gland is a more sensitive indicator of thyroid function (McNabb et al. 2004). The thyroid gland must maintain a store of hormone for times of growth or stress. This protocol outlines methods for digesting thyroid glands to extract hormone and radioimmunoassay for quantification of T₃ (triiodothyronine) and T₄ (thyroxine) content.

Procedure

Digestion of thyroid gland

This procedure was supplied by Anne McNabb, Virginia Polytechnic University, Blacksburg VA in August 2004.

Reagents:

Digestion Mix (100 mL)

1.21g TRIS

153.65mg L-Glutathione reduced, Sigma G4251

42.55mg PTU – light sensitive (6-n-propyl-2-thiouracil Cat# P-3755 from Sigma)

1ml Triton X

100ml dd H₂O

1. Weigh all dry components in the following order: a)TRIS, b)Glutathione, c)PTU
2. Combine 100 mL ddH₂O and dry components in an Erlenmeyer beaker and stir on stir-plate at low speed.
3. As the solution is stirring, add the Triton X. (To ease the Triton in and out of the pipette tip, cut about 1cm off the pipette tip before putting it on the pipette.) Cover the beaker with aluminum foil as it stirs.
4. Pour the solution into a amber bottle (it is light sensitive) and store at 2-8°C.

Pronase Solution

Pronase: Calbiochem Cat#53702

1. The average weight of thyroid glands (TG) to be digested is approximately 5mg.
2. Calculate the amount of Pronase needed. E.g. For 68 thyroid glands and 25 mg Pronase per gland, $68 \times 25 = 1700$ mg (or 1.7 g) Pronase for entire group. Use a volume of 350 μ L of digestion mixture per gland, $0.35 \text{ mL} \times 68 \text{ glands} = 23.8 \text{ mL}$. Mix the 1.7 g

of Pronase into 23.8 mL of digestion mix by swirling it around until all the Pronase has gone into solution (do not use a stirring rod; Pronase tends to stick to anything that touches it). Make your actual calculations for a few more glands than you have to ensure you have enough solution since some is lost in the container or when pipetting.

3. Pour a small amount of the mixed Pronase/digestion mixture into the weigh boat that was used to weigh the Pronase and swirl (to remove any Pronase that is stuck to the sides of the boat).

100% EtOH (ice cold)

1. Pipette 350 μ L of pronase solution in each 1.5 mL tube that holds the gland tissue. Vortex, and incubate in a 37°C water bath for 24 hours.
2. Place about 100 mL of 100% EtOH into the freezer for about 20 minutes. Remove tubes from the water bath and add 1 ml of the ice cold 100% EtOH (for a total of 1.35ml).
3. Vortex and store tubes at -20°C for 24 hours.
4. Remove tubes from the freezer and centrifuge at 14,000 RPM and -4°C for 8 minutes.
5. Decant supernatant into a clean, labeled, 1.5ml microcentrifuge tube and discard the tube with the pellet. Store the extract at -20°C until analysis.

Radioimmunoassay

Measure T3 and T4 using DPC coat-a-count kits for total canine T3 and T4. Follow manufacturers directions. Include an interassay control if running samples over more than one assay. Prepare standards for appropriate hormone ranges from scratch in 75% ethanol as needed.

Equipment Needed

Stir plate Water bath 37°C Gamma Counter
DPC coat-a-count kits for total canine T3 and T4 (Catalog #s TKC3 and TKC4)

Literature Consulted

McNabb, FMA, Cheng, MF. 1985. Thyroid development in altricial ring doves, *Streptopelia risoria*. *General and Comparative Endocrinology* 58: 243-251.

McNabb, FMA, Larsen, CT, Pooler, PS. 2004 Ammonium perchlorate effects on thyroid function and growth in bobwhite quail chicks. *Environmental Toxicology and Chemistry* 23(4):997-1003.

6.17 Analysis of Aromatase Activity in Hypothalamic Tissue

Background:

Aromatase is responsible for the biosynthesis of estradiol. This assay measures amounts of aromatase. Testosterone is converted to estradiol in the brain by aromatase.

Materials:

- Charcoal Pellets
- K-phosphate buffer
- NADPH/buffer solution
- Ice
- 16 x 100 test tubes
- microcentrifuge tubes
- androstenedione
- scintillation vials

Procedures:

*Directions for 25 samples

*Allow 2 days for assay completion

Day 1

Prepare:

- a.) Charcoal Pellets
- b.) K-phosphate buffer
- c.) Prepare NADPH/buffer solution

A.) Charcoal Pellet Preparation

0.5 ml of a 5% aqueous charcoal solution is placed into a 1.5 ml Eppendorf snap top centrifuge tube. Centrifuge (3000 RPM in Sorval for 10 min) and draw off supernatant which is discarded. Tubes must air dry before use, so let sit with caps open overnight – 2 days. Store in a cool dry place with the lids closed.

B.) Prepare K-phosphate buffer - into deionized water add:

- 1.) K-phosphate monobasic (0.1M)
 $x(\text{g})/136.09\text{g/mol} = 0.1\text{M (mol/L)}$
 $x = 13.69 \text{ g (for 1 L)}$
 $x = 6.8045 \text{ g (for 0.5 L)}$
 $x = 2.738 \text{ g (for 200 ml)}$
- 2.) K-phosphate dibasic, anhydrous (0.1M)

$$x/174.18\text{g/mol} = 0.1\text{M}$$

$$x = 17.418 \text{ g (for 1 L)}$$

$$x = 8.709 \text{ g (for 0.5 L)}$$

$$x = 3.483 \text{ g (for 200ml)}$$

- 3.) Sucrose (0.25M)
- $$x/342.30 = 0.25$$
- $$x = 85.575\text{g (for 1 L)}$$
- $$x = 42.7875\text{g (for 0.5 L)}$$
- $$x = 17.115 \text{ g (for 200ml)}$$

Bring solution to pH 7.2 (pH 7.4 for poikilotherms) and store in refrigerator until use

C.) NADPH -***keep on ice***

1. Dissolve 0.01775g NADPH in 1mL K-phosphate buffer.
2. Store at -80°C with samples in microcentrifuge tube.

Day 2

Preparation:

1. Turn on water bath to 37° C and double check temperature (for quail; for other poikilotherms, water bath must be at the temperature animals were acclimated to).
2. Set shake to 2.5
3. Label (16x100) test tubes: 2 for each sample, 2 blanks
4. Obtain ice.

Experiment:

1. Defrost samples but keep on ice in microcentrifuge tubes.
2. Add K-phosphate buffer and homogenize for 30 seconds (100 µl for adult dissections or 50ul for hatchling dissections).
3. Dissolve 72 µL androstenedione in 1mL propylene glycol.
Do this under hood or radioactive labeled countertop.
4. To each test tube that will hold sample add:
 - 233 µL K-phosphate buffer
 - 30 µL dissolved NADPH
 - 12 µL dissolved androstenedione
5. To each test tube that will hold blanks add:
 - 258 µL K-phosphate buffer
 - 30 µL dissolved NADPH
 - 12 µL dissolved androstenedione
6. Add 25 µL sample homogenate to each appropriately labeled sample test tube.
7. Place all tubes in gently shaking water bath for 3 hours.

Check temperature of bath regularly to insure it is the correct temperature.

8. Remove tubes from bath, place under hood in room 3144 and add 4mL methylene chloride to stop the reaction.
9. Add 1.7 mL deionized water to each tube and vortex GENTLY.
10. Centrifuge in bucket centrifuge for 5 minutes @ 1000 rpm.
Ensure buckets are balanced exactly by using balance located next to centrifuge.
11. Remove 1mL aqueous portion (top layer) and add to charcoal pellets in microcentrifuge tubes.
12. Let incubate for 20 minutes at 4°C in radioactive fridge vortex every 5 minutes.
13. Centrifuge for 5 minutes @ 14,000g or 6,750 rpm (using centrifuge in room 3120).
14. Add 200 µL supernatant to scintillation vial, .
300 µL MilleQ water to scintillation vial.
5 mL scintillation cocktail to scintillation vial.
15. Vortex until white precipitate is swirled.
16. Place vials in scintillation counter.

Notes: order information

Sigma: *B* NADPH tetra Na salt N1630, buy in lots of 250 mg or more

Perkin Elmer/ NEN : Androst-4-ene 3,17- dione [1 *B* – ³H(N)] NET926250UC, buy as 250 uCi or more

Round bottom snap top tubes

Campus Chemistry store for methylene chloride

SAMPLE Calculation: **Androst-ene dione** calculation- for 200 samples, 12 ul x 200= 2,400 ul or 2.4 ml... so need 216 ul ³H dissolved in 3 ml propylene glycol. **NADPH** calculation- .01775 g in 1 ml K-phosphate & 30 ul /sample x 200 = 6 ml or 0.1065gx0.01775g NADPH so would need to order 2 x 250 mg

6.19 Tissue Storage, RNA Isolation And Microarray Analysis

1. Collect tissue and make cuts using a sterile razor blade. Place tissue sample in a clean microcentrifuge tube and snap freeze in liquid nitrogen. Store at -80°C.
2. Isolate RNA using Qiagen RNeasy kits. Use appropriate kit depending upon type and size of tissue as described in RNeasy handbook.
3. Submit RNA samples for microarray analysis.
 - Tissues will be collected and prepared as in previous experiments and microarrays have been validated for several avian species. Because these microarrays are constructed with normalized cDNAs, they offer higher cross species utility than those produced with oligos, especially with species having a high homology to galliforms. Our previous study has used the following procedure.
 1. Hypothalamic tissue samples from were frozen in liquid nitrogen and stored at -80°C. Tissue samples ranged from 115 to 238 mg in weight.
 2. RNA was isolated with RNeasy MAXI kit per manufacturers instructions.
 3. Between 75 and 190 µg of RNA was obtained from each tissue sample.
 4. 260/280 ratio for isolated RNA was between 1.5 and 1.6.
 5. RNA was run on a gel to check for degradation; degradation was minimal.
 6. 25 µg RNA from each sample was submitted for microarray analysis on slides developed for chickens.
 7. The RNA successfully hybridized with the chicken oligonucleotides on the chicken microarray.

Detailed descriptions of the microarrays, including the characterization and analytical approaches can be found in:

Laura E. Ellestad, Wilfrid Carre, Michael Muchow, Sultan A. Jenkins, Xiaofei Wang, Larry A. Coghurn, Tom E. Porter. Gene expression profiling during cellular differentiation *in the embryonic pituitary* gland using cDNA microarrays. *In* "Functional Avian Endocrinology", Allistair Dawson and Peter Sharp, *Editors*.

Wilfrid Carre, Xiaofei Wang, Tom E. Porter, Yves Nys, Jianshan Tang, Erin Bernberg, Robin Morgan, Joan Burnside, Samuel E. Aggrey, Jean Simon and Larry A. Coghurn. 2006 Chicken Genomics Resource: Sequencing and Annotation of 35,407 ESTs from Single and Multiple Tissue cDNA Libraries and CAP3 Assembly of a Chicken Gene Index. *Physiological Genomics*, Articles in Press. (March 21, 2006).

6.20 *Field Collection of Tree Swallow Eggs from Great Sacandaga Lake, New York for Avian Egg Injection Study, Hudson River NRDA*

INTRODUCTION

Avian egg injection is a well-established technique to assess the effects of contaminants on a developing avian embryo. To conduct an avian egg injection experiment, eggs are collected from the field and brought into a laboratory where they are injected with the substance being tested. Proper handling of the eggs during collection and transit to the laboratory is essential to maintain viability in eggs that will subsequently be injected with contaminants and incubated. A subsample of eggs will be selected for contaminants analysis.

MATERIALS AND EQUIPMENT

FIELD:

- ❑ Scientific collecting permits
- ❑ Field notebook, writing instruments (pencils/pens/permanent markers)
- ❑ Padded egg collection boxes (hard-sided container, e.g., Tupperware or tackle box, with padding such as sawdust or holofill)
- ❑ Avian Egg Collection Data Sheets

PROCEDURES

FIELD:

- ❑ Collected eggs should be whole and not cracked.
- ❑ For tree swallows, the following approach should be used: Incubation of tree swallow eggs doesn't start until the clutch is complete. Eggs are generally laid at one day intervals. Monitor the laying of eggs daily. Tree swallows generally lay eggs at one day intervals with a maximum clutch size of about 5-7 eggs. When the number of eggs in the nest has not increased since the day before the clutch should be considered complete and should be collected promptly. Collect all eggs in the clutch.
- ❑ For each egg collected, complete the appropriate information on the Avian Egg Collection Datasheet.
- ❑ Place eggs in individually numbered compartments (one for each egg or eggs from each clutch). A list of the nest box associated with each compartment will be placed inside the

container. A fishing tackle box with compartments lined with sawdust or holofill is ideal – all eggs should be treated the same. Place this box in a hard-sided container with sufficient padding. Transport to the New York State Department of Environmental Conservation (NYSDEC) laboratory in a hard container avoiding temperature extremes and jostling.

- At the NYSDEC laboratory, eggs which are going to be incubated should be separated from eggs which are going to undergo contaminants analysis.
- For eggs that are going to be analyzed for contaminants and not incubated: Randomly select 10 nests from among those from which eggs were collected and from each of those 10 nests, randomly select 2 eggs from each nest. These 2 eggs will each form a composite sample for chemical analysis. Refrigerate eggs until opened, no longer than 48 hours. Processing of eggs for contaminants analysis will be completed on a daily basis as much as practical. Follow Standard Operating Procedure for Removal of Avian Egg Contents for Contaminants Analysis, Hudson River NRDA, compositing the 2 eggs from each nest in one jar. Note that the eggs were taken for contaminant analysis on the Egg Collection Data Sheet. Freeze samples at NYSDEC laboratory. Ship under Chain of Custody overnight on dry ice to the sample archive or analytical laboratory, as directed by the Trustees.
- For eggs that are going to be incubated: Transport promptly to the laboratory. Injection of tree swallow eggs at day ED2.5-3 will be done in the laboratory; as eggs will have already undergone potentially one day of incubation in the nest in the field, prompt transport under appropriate conditions is essential. Use of a “Koolatron” to maintain a proper temperature of eggs during transport is recommended. Maintain a temperature of about 90 to 100 degrees F, unless the transport time is going to be 8 hours or more, in which case a temperature as close as possible to 99.5 degrees should be maintained. Complete chain of custody transfer of samples from NYSDEC field collection crew to the laboratory crew on Egg Collection Data Sheet.

Avian Egg Collection Data Sheet – Hudson River Avian Egg Injection Study – Great Sacandaga Lake, New York

Collector: _____ Data Recorder: _____
 Print Name Signature Print Name Signature

Species: Tree Swallow

Compartment #	Egg Code ¹	Location ²	Date Collected ³	Time Collected ⁴	Clutch Size	Eggs Warm Yes/No	Prior Day Clutch Size	Prior Day Time	Comments	Fresh Weight (g) ⁵

¹ Egg Code: Numeric code between 200 and 799; ² Site and nest number, such as Gr Sac Box 2 where Gr Sac = Great Sacandaga Lake Box Number 2; ³ In MM/DD/YEAR format, such as 04/30/2006 for April 30, 2006; ⁴ In 24-hour format, such as 1300 for 1PM; ⁵ Record at the laboratory

Custody of samples listed above transferred from field collection crew to laboratory crew as follows:

Relinquished by: _____
 Signature Print Name Company/Title Date Time

Received by: _____
 Signature Print Name Company/Title Date Time

Data Sheet checked by: _____ Date: _____

6.21 Field Collection of Tree Swallow Eggs from Upper Hudson River, New York for Injury Assessment Hudson River NRDA

INTRODUCTION

Tree swallow eggs from a PCB-contaminated location will be collected late in incubation and incubated to hatching. A subsample of eggs from the PCB-contaminated location will be selected for contaminants analysis.

MATERIALS AND EQUIPMENT

FIELD:

- ❑ Scientific collecting permits
- ❑ Field notebook, writing instruments (pencils/pens/permanent markers)
- ❑ Padded egg collection boxes (hard-sided container, e.g., Tupperware or tackle box, with padding such as sawdust or holofill)
- ❑ Avian Egg Collection Data Sheets

PROCEDURES

FIELD:

- ❑ Collected eggs should be whole and not cracked.
- ❑ For tree swallows, the following approach should be used: Incubation of tree swallow eggs doesn't start until the clutch is complete. Monitor nests every two to three days. Tree swallows generally lay eggs at one day intervals with a maximum clutch size of about 5-7 eggs. When a nest is 2-5 days pre-hatch (based on when the clutch was completed and incubation began), collect three eggs from that nest -- one egg will be incubated at the laboratory and the other 2 sibling eggs will be subject to contents collection for contaminants analysis. In order to facilitate transport of eggs to the laboratory, eggs for contaminant analysis can be collected independently of those to be transported to the laboratory. Eggs should be collected from a total of 20 nests (60 eggs total).
- ❑ For each egg collected, complete the appropriate information on the Avian Egg Collection Datasheet. Maintain separate Avian Egg Collection Datasheets for eggs to be transported to the laboratory and for eggs to be analyzed for contaminants.
- ❑ Place eggs in individually numbered compartments (one for each egg or eggs from each clutch). A list of the nest box associated with each compartment will be placed inside the container. A fishing tackle box with compartments lined with sawdust or holofill is ideal – all eggs should be treated the same. Place this box in a hard-sided container with sufficient padding. Transport to the processing laboratory in a hard container avoiding temperature extremes and jostling.
- ❑ For eggs that are going to be analyzed for contaminants and not incubated: Refrigerate eggs until opened, no longer than 48 hours. Processing of eggs for contaminants analysis will be completed on a daily basis as much as practical. Follow Standard Operating

Procedure for Removal of Avian Egg Contents for Contaminants Analysis, Hudson River NRDA, compositing the 2 eggs from each nest in one jar. Archive samples at NYSDEC laboratory within two weeks of collection.

- For eggs that are going to be incubated: Transport promptly to the laboratory. Prompt transport under appropriate conditions is essential. Use of a “Koolatron” to maintain a proper temperature of eggs during transport is recommended. A hot water bottle can be substituted if a Koolatron is not practical or malfunctions. Maintain a temperature of about 90 to 95 degrees F, unless the transport time is going to be 8 hours or more, in which case a temperature as close as possible to 99.5 degrees should be maintained. Complete chain of custody transfer of samples from field collection crew to the laboratory crew on Egg Collection Data Sheet.

Avian Egg Collection Data Sheet – Hudson River Avian Egg Injection Study – Upper Hudson River

Collector: _____ Data Recorder: _____
 Print Name Signature Print Name Signature

Species: Tree Swallow

Compartment #	Egg Code ¹	Location ²	Date Collected ³	Time Collected ⁴	Clutch Size	Eggs Warm Yes/No	Embryonic day of incubation	Comments	Fresh Weight (g) ⁵

¹ Egg Code: Numeric code between 800 and 899; ² Site followed by box number, such as UHR313 where UHR = Upper Hudson River, Box Number 313; ³ In MM/DD/YEAR format, such as 04/30/2006 for April 30, 2006; ⁴ In 24-hour format, such as 1300 for 1PM; ⁵ Weighed at the laboratory
 Custody of samples listed above transferred from field collection crew to laboratory crew as follows:

Relinquished by: _____
 Signature Print Name Company/Title Date Time

Received by: _____
 Signature Print Name Company/Title Date Time

Data Sheet checked by: _____ Date: _____

[This page intentionally left blank]

APPENDIX B

EGG INJECTION DOSING MIXTURE

Table B1. Final selected PCB congeners, total mass and percentage of total PCB in the stock 50mL aliquant of the original PCB mixture solution.

PCB Congener	PCB congener mass (ug) in the 50 mL aliquant of the original stock solution received from AccuStandard.	Relative Amount (%)
PCB 28	40880	10.46%
PCB 66	36660	9.38%
PCB 74	26840	6.87%
PCB 118	25530	6.53%
PCB 47	14420	3.69%
PCB 48	9405	2.41%
PCB 75	802	0.21%
PCB 138	17000	4.35%
PCB 163	3980	1.02%
PCB 164	1607	0.41%
PCB 101	19980	5.11%
PCB 89	600	0.15%
PCB 52	20100	5.14%
PCB 49	15820	4.05%
PCB 43	1816	0.46%
PCB 153	16215	4.15%
PCB 99	15935	4.08%
PCB 70	14515	3.72%
PCB 105	12280	3.14%
PCB 31	11800	3.02%
PCB 56	5780	1.48%
PCB 60	5760	1.47%
PCB 41	1998	0.51%
PCB 71	1614	0.41%
PCB 64	6780	1.74%
PCB 110	7220	1.85%
PCB 85	6940	1.78%
PCB 87	4818	1.23%
PCB 115	398.4	0.10%
PCB 128	4026	1.03%
PCB 149	3380	0.87%
PCB 139	68	0.02%
PCB 92	3413.5	0.87%
PCB 180	2802	0.72%
PCB 158	2822	0.72%
PCB 146	2583.5	0.66%
PCB 97	2584	0.66%
PCB 156	2412	0.62%
PCB 95	2398	0.61%
PCB 187	2196.5	0.56%
PCB 170	1814	0.46%
PCB 190	400.8	0.10%
PCB 117	1992	0.51%
PCB 141	1788	0.46%
PCB 130	1390	0.36%
PCB 109	1206	0.31%
PCB 137	1198.5	0.31%
PCB 42	795	0.20%
PCB 59	197.2	0.05%
PCB 114	1008	0.26%
PCB 167	796	0.20%
PCB 123	599	0.15%
PCB 157	594	0.15%
PCB 77	400	0.10%
PCB 81	200.4	0.05%
PCB 126	84.35	0.02%
PCB 189	70.2	0.02%
PCB 169	0.72	0.0002%
Total	390713	100.0%

Table B2 notes the target and nominal concentrations of total PCB in each dosing solution. The "target concentration" is the concentration that the laboratory hoped to achieve when making the solutions. The "nominal concentration" is the concentration that the laboratory believes has been achieved based on the weights and measures taken during the preparation process. The laboratory will be determining "measured concentrations," which are concentrations based on analyses of the dosing solutions.

Table B2. Target and nominal concentrations of total PCB in each dosing solution.

Dose ID	Vial & Cap #	Solution Description	Target Concentration (ug/uL) ¹	Nominal Concentration (ug/uL) ²
7	3-7	Stock 58-congener PCB mixture	246	244
6	3-8	2-fold dilution	123	122
5	3-9	4-fold dilution	62	61
4	3-10	8-fold dilution	31	30.5
3	3-11	16-fold dilution	15	15
2	3-12	32-fold dilution	8	7.5
1	3-4	Isooctane blank	0	0

¹Target concentrations for the dosing solutions of the custom 58-congener PCB mixture.
²Nominal total PCB concentrations are based on the sum of the certified analyte concentrations from AccuStandard (see attached), a volume of 50 mL of the 250 mL of original custom 58-congener PCB mixture, a stock PCB emulsion volume of 1.6 mL, and the appropriate serial dilution for each dose.



AccuStandard Inc.

125 Market Street
New Haven, CT 06513
USA

CERTIFICATE OF ANALYSIS



Ph: 203-788-5290
Fax: 203-786-5287
E-mail: usa@accustandard.com
www.accustandard.com

CATALOG NO: S-13907-250ML

DESCRIPTION: Custom PCB Congener Standard

LOT: B5110052

SOLVENT: Isooctane

EXPIRATION: Nov 9, 2015

See reverse for additional certification information.

This product is guaranteed accurate to + 0.5% of the Certified Analyte concen
Expiration Date on the Label.

Component	CAS #	Purity % (GC/MS)	Prepared Concentration ¹ (µg/mL)	Certified Analyte Concentration ² (µg/mL)
2,4,4'-Trichlorobiphenyl	7012-37-5	100	817.6	± 32.70 817.6
2,3',4,4'-Tetrachlorobiphenyl	32598-10-0	100	733.2	± 29.33 733.2
2,4,4',5-Tetrachlorobiphenyl	32690-93-0	100	536.8	± 21.47 536.8
2,3',4,4',5-Pentachlorobiphenyl 113	31508-00-6	99.5	513.2	± 20.53 510.6
2,2',4,4'-Tetrachlorobiphenyl	2437-79-8	100	288.4	± 11.54 288.4
2,2',4,5-Tetrachlorobiphenyl 48	70362-47-9	99.9	188.3	± 7.53 188.1
2,4,4',6-Tetrachlorobiphenyl	32598-12-2	99.5	16.12	± 0.64 16.04
2,2',3,4,4',5'-Hexachlorobiphenyl 138	35065-28-2	100	340.0	± 13.60 340.0
2,3,3',4',5,6-Hexachlorobiphenyl 163	74472-44-9	99.0	80.40	± 3.22 79.60
2,3,3',4',5,6-Hexachlorobiphenyl 164	74472-45-0	99.7	32.24	± 1.29 32.14
2,2',4,5,5'-Pentachlorobiphenyl 101	37680-73-2	99.4	402.0	± 16.08 399.6
2,2',3,4,6'-Pentachlorobiphenyl 84	73555-57-2	99.9	12.01	± 0.48 12.00
2,2',5,5'-Tetrachlorobiphenyl 52	35693-99-3	100	402.0	± 16.08 402.0
2,2',4,5'-Tetrachlorobiphenyl 49	41464-40-8	100	316.4	± 12.66 316.4
2,2',3,5-Tetrachlorobiphenyl 43	70362-46-8	99.9	36.36	± 1.45 36.32
2,2',4,4',5,5'-Hexachlorobiphenyl 153	35065-27-1	99.6	325.6	± 13.02 324.3
2,2',4,4',5-Pentachlorobiphenyl 99	38380-01-7	99.6	320.0	± 12.80 318.7
2,3',4,5-Tetrachlorobiphenyl 70	32598-11-1	99.0	293.2	± 11.73 290.3
2,3,3',4,4'-Pentachlorobiphenyl 105	32598-14-4	100	245.6	± 9.82 245.6
2,4',5-Trichlorobiphenyl 31	16606-02-3	100	236.0	± 9.44 236.0
2,3,3',4-Tetrachlorobiphenyl 56	41464-43-1	99.6	116.1	± 4.64 115.6
2,3,4,4'-Tetrachlorobiphenyl 60	33025-41-1	99.0	116.4	± 4.66 115.2
2,2',3,4-Tetrachlorobiphenyl 41	52663-59-9	W.3	40.24	± 1.61 39.96
2,3',4,6-Tetrachlorobiphenyl 71	41464-46-4	100	32.28	± 1.29 32.28
2,3,4,6-Tetrachlorobiphenyl 64	52663-58-8	99.0	137.0	± 5.43 135.6
2,3,3',4,6-Pentachlorobiphenyl 110	38380-03-9	99.7	144.8	± 5.79 144.4
2,2',3,4,4'-Pentachlorobiphenyl 85	65510-45-4	99.0	140.2	± 5.61 138.8
2,2',3,4,5'-Pentachlorobiphenyl 87	38380-02-8	W.5	96.84	± 3.87 96.36
2,3,4,4',6-Pentachlorobiphenyl	74472-38-1	99.5	8.008	± 0.32 7.968
2,2',3,3',4,4'-Hexachlorobiphenyl 125	38380-07-3	99.7	80.76	± 3.23 80.52
2,2',3,4',5,6-Hexachlorobiphenyl 149	38380-04-0	99.0	68.28	± 2.73 67.60
2,2',3,4,4',6-Hexachlorobiphenyl 139	56030-56-9	99.4	1.368	± 0.05 1.360
2,2',3,5,5'-Pentachlorobiphenyl 92	52663-61-3	99.7	68.48	± 2.74 68.27
2,2',3,4,4',5,5'-Heptachlorobiphenyl 180	35065-29-3	100	56.04	± 2.24 56.04
2,3,3',4,4',6-Hexachlorobiphenyl 158	74472-42-7	100	56.44	± 2.26 56.44
2,2',3,4',5,5'-Hexachlorobiphenyl 146	51908-16-8	98.9	52.24	± 2.09 51.67
2,2',3',4,5-Pentachlorobiphenyl 97	41464-51-1	99.0	52.20	± 2.09 51.68

1. All weights are traceable through NIST. Test No 822/270236-04
 2. Certified Analyte Concentration = Purity x Prepared Concentration. The Uncertainty calculated for this product is the Combined Uncertainty $u_c(y)$. It represents an estimated standard deviation equal to the positive square root of the total variance of the uncertainty of components. The Expanded Uncertainty is U which is $U_c(y) \times K$ where K is the coverage factor at the 95% confidence level ($K=2$). Values reported above are Expanded Combined Uncertainty
 3. A product with a suffix (-1A, -2B, etc.) on its lot# has had its expiration date extended and is identical to the same lot# without the suffix

R. Cooper



AccuStandard Inc.

125 Market Street
New Haven, CT 06513
USA



Ph: 203-786-5290
Fax: 203-786-5287

E-mail: usa@accustandard.com
www.accustandard.com

CERTIFICATE OF ANALYSIS

CATALOG NO: S-13907-250ML

EXPIRATION: Nov 9, 2015

DESCRIPTION: Custom PCB Congener Standard

LOT: B5110052

See reverse for additional certification information.

SOLVENT: Isooctane

This product is guaranteed accurate to +0.5% of the Certified Analyte concentration through the Expiration Date on the Label.

Component	CAS #	Purity % (GC/MS)	Prepared Concentration ¹ (µg/mL)	Certified Analyte Concentration ² (µg/mL)
2,3,3',4,4',5-Hexachlorobiphenyl 156	38380-08-4	100	48.24	± 1.93 48.24
2,2',3,5',6-Pentachlorobiphenyl 155	38379-99-6	99.1	48.40	± 1.94 47.96
2,2',3,4',5,5',6-Heptachlorobiphenyl	52663-68-0	99.4	44.20	± 1.77 43.93
2,2',3,3',4,4',5-Heptachlorobiphenyl 170	35065-30-6	100	36.28	± 1.45 36.28
2,3,3',4,4',5,6-Heptachlorobiphenyl 190	41411-64-7	1 0	8.016	± 0.32 8.016
2,3,4',5,6-Pentachlorobiphenyl 168	68194-11-6	99.0	40.24	± 1.61 39.84
2,2',3,4,5,5'-Hexachlorobiphenyl 191	52712-04-6	99.0	36.12	± 1.44 35.76
2,2',3,3',4,5'-Hexachlorobiphenyl 130	52663-66-8	99.3	28.00	± 1.12 27.80
2,3,3',4,6-Pentachlorobiphenyl	74472-35-8	100	24.12	± 0.96 24.12
2,2',3,4,4',5-Hexachlorobiphenyl 134	35694-06-3	99.7	24.04	± 0.96 23.97
2,2',3,4'-Tetrachlorobiphenyl 42	36559-22-5	99.4	16.00	± 0.64 15.90
2,3,3',6-Tetrachlorobiphenyl 59	74472-33-6	98.5	4.004	± 0.16 3.944
2,3,4,4',5-Pentachlorobiphenyl 114	74472-37-0	100	20.16	± 0.81 20.16
2,3',4,4',5,5'-Hexachlorobiphenyl 167	52663-72-6	99.0	16.08	± 0.64 15.92
2',3,4,4',5-Pentachlorobiphenyl 123	65510-44-3	99.3	12.00	± 0.48 11.98
2,3,3',4,4',5'-Hexachlorobiphenyl 177	69782-90-7	99.3	12.00	± 0.48 11.88
3,3',4,4'-Tetrachlorobiphenyl 77	32598-13-3	100	8.000	± 0.32 8.000
3,4,4',5-Tetrachlorobiphenyl 81	70362-50-4	100	4.008	± 0.16 4.008
3,3',4,4',5-Pentachlorobiphenyl 1	57465-28-8	100	1.687	± 0.07 1.687
2,3,3',4,4',5,5'-Heptachlorobiphenyl 189	39635-31-9	100	1.404	± 0.06 1.404
3,3',4,4',5,5'-Hexachlorobiphenyl 169	32774-16-6	100	0.0144	± 0.00 0.0144

58 Components

1. All weights are traceable through NIST, Test No 822/270236-04

2. Certified Analyte Concentration = Purity x Prepared Concentration. The Uncertainty calculated for this product is the Combined Uncertainty $u_c(y)$. It represents an estimated standard deviation equal to the positive square root of the total variance of the uncertainty of components. The Expanded Uncertainty is U which is $U_c(y) * K$ where K is the coverage factor at the 95% confidence level ($K=2$). Values reported above are Expanded Combined Uncertainty

3. A product with a suffix (-1A, 2B, etc.) on its lot# has had its expiration date extended and is identical to the same lot# without the suffix.

Certified by: _____

R. Cooper

Page 2 of 2

This product was manufactured to meet the quality system requirements of ISO 9001

QR-ORC11W01002

CERTIFICATION REPORT

- 1. Intended Use:** The product covered by this Certificate is designed for Calibration or for use in Quality Control procedures for the specified chemical compounds listed on the reverse side. This product can be used for Identification and/or Quantification. This product can also be used as a Reference Material to validate analytical procedures, subject to the conditions under Section 8.
- 2. Raw Materials:** Reference standards are prepared from the highest quality starting materials with defined purities. All analytes and solvents are obtained from pre-qualified vendors and then analyzed or evaluated prior to use according to ISO9001 requirements.
- 3. Manufacturing:** AccuStandard, Inc. manufactures its products under an ISO 9001 certified quality system. Balances used in the manufacturing process are calibrated regularly. All weights are traceable through the National Institute of Standards and Technology (NIST), Test No. 822/254480.
- 4. Homogeneity Assessment:** Homogeneity of the finished product is assessed by analyzing sample batches or by other methods consistent with the intended use of the product and by procedures that comply with the ISO 9001' Quality System.
- 5. Stability Assessment:** AccuStandard, Inc. guarantees the stability of this solution through the expiration date stated on the label, when handled and stored according to the conditions stated on the label. To ensure a uniform solution, mix the contents of the sealed container thoroughly prior to use. Care should be taken not to contaminate the contents of the original container.
- 6. Analytical Quality Control:** Products are tested by validated analytical methods covered under the company's ISO 9001 Quality System.
- 7. Uncertainty Statistics and Confidence Limits:** The maximum Uncertainty stated on the face of this certificate has been calculated in accordance with the EURACHEM/CITAC Guide - Quantifying Uncertainty in Analytical Measurement - Second Edition. The Uncertainty given is the Expanded Combined Uncertainty and represents an estimated Standard Deviation equal to the positive square root of the total variance of the uncertainty of components. The Expanded Uncertainty is U which is $U_c(y) * K$, where K is the coverage factor at the 95% confidence level ($K=2$). The Expanded Uncertainty is based on the combination of uncertainties associated with each individual operation involved in the preparation of the product.
- 8. Legal Notice and Limit of Liability:** This product is for research use only. No warranty for any particular application is expressed or implied. Due to their hazardous nature, they should be handled by trained personnel. The company's liability will be limited to replacement of product or refund of purchase price. Notice of claims must be made within thirty (30) days from date of delivery.



AccuStandard[®], Inc.

Chemical Reference Standards • The Standard for Excellence

WARRANTIES:

Manufacture]: (AccuStandard[®], Inc.) warrants that its products shall conform to the description of such products as provided in its catalog or on the specific products' label. This warranty is exclusive, and AccuStandard, Inc. makes no other Warranty, express or implied, including any implied warranty of merchantability or fitness for any particular purpose.

PRODUCT STABILITY:

AccuStandard's products are monitored regularly to ensure they meet Catalog Specifications (on-going stability studies). The integrity of these products is dependent upon proper handling and storage by the end-user.

AccuStandard recommends the following storage conditions:

Volatiles	-10 to -20 °C
Semi-Volatiles	4 °C

Exceptions: Highly concentrated solutions (e.g. Z-014J) should be stored at room temperature.

Note: Allow ampules to equilibrate to 20 °C prior to opening.

LIABILITY:

AccuStandard, Inc. products are for research use only. Due to their hazardous nature, they should be handled by trained personnel. AccuStandard's liability will be limited to replacement of products or refund of purchase price. Failure to give notice of claim within thirty (30) days from date of delivery will constitute a waiver by buyer of any and all claims.



AccuStandard Inc.

125 Market Street
New Haven, CT 06513
USA



Ph: 203-786-5290
Fax: 203-786-5287
E-mail: usa@accustandard.com
www.accustandard.com

CERTIFICATE OF ANALYSIS

CATALOG NO. S-13907-BLK-250ML

DESCRIPTION: Isooctane Control Blank

EXPIRATION: Nov 9, 2006

LOT: B5110053

SOLVENT: N/A

See reverse for additional certification information.

This product is guaranteed accurate to + 0.5% of the Certified Analyte concentration through the Expiration Date on the Label.

Component	CAS #	Purity % MFG	Prepared Concentration ¹	Certified Analyte Concentration ²
Isooctane	540-84-1	99.9	N/A ±0	N/A

Please note: AccuStandard follows the U.S. conventions in reporting numerical values, on both certificates and labels.

A comma (,) is used to separate units of one-thousand or greater.
A period (.) is used as a decimal place marker.

1. All weights are traceable through National Institute of Standards & Technology, Test No.
2. Certified Analyte Concentration = Purity x Prepared Concentration
3. A product with a suffix (-1A, -2B, etc.) on its lot# has had its expiration date extended and is identical to the same lot# without the suffix.

Certified by: _____

R. Cooper

This product was manufactured to meet the quality system requirements of ISO 9001

QR-ORG/INO-001
Rev. 11/02

CERTIFICATION REPORT

- 1. Intended Use:** The product covered by this Certificate is designed for **Calibration** or for use in Quality Control procedures for the specified chemical compounds listed on the reverse side. This product **can be** used for Identification **and/or** Quantification. This product **can also be** used as a Reference Material to validate analytical procedures, subject to the conditions under Section 8.
- 2. Raw Materials:** Reference Standards are prepared from the highest quality starting materials with defined purities. All **analytes** and solvents are obtained from **pre-qualified** vendors and then analyzed or evaluated prior to use according to ISO9001 requirements.
- 3. Manufacturing:** AccuStandard, Inc. manufactures its products under an ISO 9001 certified quality system. Balances used in the manufacturing process are calibrated regularly. All weights are traceable **through** the National Institute of Standards and Technology (NIST), Test No, 822/254480.
- 4. Homogeneity Assessment:** Homogeneity of the finished product is assessed by analyzing sample batches or by other methods consistent with the intended use of the product and by **procedures** that comply with the ISO 9001 Quality System.
- 5. Stability Assessment:** AccuStandard, Inc. guarantees the stability of this solution through the expiration **date** stated on the label, when handled and stored **according** to the conditions stated on the label. To ensure a uniform solution, **mix** the contents of the **sealed** container thoroughly prior to use. Care should be **taken** not to contaminate the contents of the **original container**.
- 6. Analytical Quality Control:** Products are tested by validated analytical methods **covered** under the company's ISO 9001 Quality System.
- 7. Uncertainty Statistics and Confidence Limits:** The maximum Uncertainty stated on the face of this certificate has been calculated in accordance with the EURACHEM/CITAC Guide –Quantifying Uncertainty in Analytical Measurement - Second Edition. The Uncertainty given is the Expanded Combined Uncertainty and represents an estimated Standard Deviation equal to the positive square root of the total variance of the uncertainty of components. The Expanded Uncertainty is **U which is $U_c(y) * K$** , where K is the coverage factor at **the 95% confidence level (K=2)**. The Expanded Uncertainty is **based on the combination** of uncertainties associated with each individual operation involved in the **preparation** of the product.
- 8. Legal Notice and Limit of Liability:** This product is for research use only. No warranty for **any** particular application is expressed or implied. Due to their hazardous nature, they should be handled by trained personnel. The company's liability will be limited to replacement of **product** or **refund** of purchase price. Notice of claims **must** be made within thirty (30) **days** from date of delivery.

