

77/09/02 TABLE 290A. DEATHS FOR 60 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1960-67

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	TOTAL	UNDER 1 YEAR	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS	65-74 YEARS	75-84 YEARS	85+ YEARS	N.S.
ALL CAUSES													
BOTH SEXES													
1967	1,851,323	79,028	13,506	16,893	37,706	34,786	74,039	165,041	294,162	437,919	469,669	227,987	587
1966	1,863,149	85,516	15,089	17,133	36,326	34,398	74,836	165,611	293,362	443,835	471,022	225,388	633
1965	1,828,136	92,866	15,396	16,675	33,093	33,581	74,821	162,848	287,465	435,392	458,302	217,147	550
1964	1,798,051	99,783	15,976	16,750	31,477	33,639	74,884	160,649	283,049	430,893	443,128	207,212	611
1963	1,813,549	103,390	16,571	16,524	29,321	32,879	74,277	160,429	282,960	440,362	445,667	210,541	628
1962	1,756,720	105,479	16,254	16,410	27,345	32,295	72,837	156,940	272,873	428,038	426,467	201,203	579
1961	1,701,522	107,956	16,629	16,248	25,504	32,241	71,081	152,744	265,470	415,906	407,864	189,275	604
1960	1,711,982	110,873	17,682	16,537	25,533	33,414	72,103	154,876	270,197	420,312	405,209	184,526	720
MALE													
1967	1,045,945	45,442	7,651	10,361	27,185	22,918	45,745	105,363	192,486	260,882	236,663	90,906	343
1966	1,052,827	49,173	8,385	10,477	26,176	22,254	46,045	105,957	192,431	264,071	237,439	90,015	404
1965	1,035,200	53,419	8,528	10,204	23,754	21,601	45,684	103,961	187,464	260,039	232,319	87,882	345
1964	1,017,778	57,348	8,878	10,236	22,398	21,546	45,815	102,389	183,854	256,774	224,428	83,749	363
1963	1,027,686	59,734	9,140	9,955	20,680	20,841	45,053	102,905	183,050	263,231	226,255	86,472	370
1962	994,789	60,939	8,836	9,918	19,274	20,279	44,197	100,809	176,129	254,923	215,923	83,228	334
1961	966,827	62,120	9,184	9,881	18,111	20,246	43,395	98,061	171,542	247,871	207,421	78,638	357
1960	975,648	63,936	9,844	10,038	18,106	21,010	43,828	100,140	174,059	251,434	206,063	76,753	437
FEMALE													
1967	805,378	33,586	5,855	6,532	10,521	11,868	28,294	59,678	101,676	177,037	233,006	137,081	244
1966	810,322	36,343	6,704	6,656	10,150	12,144	28,791	59,654	100,931	179,764	233,583	135,373	229
1965	792,936	39,447	6,868	6,471	9,339	11,980	29,137	58,887	100,001	175,353	225,983	129,265	205
1964	780,273	42,435	7,098	6,514	9,079	12,093	29,069	58,260	99,195	174,119	218,700	123,463	248
1963	785,863	43,656	7,431	6,569	8,641	12,038	29,224	57,524	99,910	177,131	219,412	124,069	258
1962	761,931	44,540	7,418	6,492	8,071	12,016	28,640	56,131	96,744	173,115	210,544	117,975	245
1961	734,695	45,836	7,445	6,367	7,393	11,995	27,686	54,683	93,928	168,035	200,443	110,637	247
1960	736,334	46,937	7,838	6,499	7,427	12,404	28,275	54,736	96,138	168,878	199,146	107,773	283

Note: Figures by race exclude data for residents of New Jersey for 1962 and 1963.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290A for 1960-67 and separate unpublished tables for the black population and Accidents.

77/09/02 TABLE 290A. DEATHS FOR 60 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1960-67

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	TOTAL	UNDER 1 YEAR	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS	65-74 YEARS	75-84 YEARS	85+ YEARS	N.S.
ALL CAUSES													
WHITE, BOTH SEXES													
1967	1,627,151	57,533	9,993	13,616	30,536	25,130	56,258	136,205	253,274	389,463	439,386	215,403	354
1966	1,634,461	61,749	11,089	13,800	29,682	25,141	56,858	136,578	251,930	394,450	440,131	212,666	387
1965	1,605,498	67,198	11,348	13,420	27,135	24,684	57,190	134,653	246,392	388,605	429,101	205,419	353
1964	1,578,674	72,728	11,811	13,502	25,933	24,855	57,466	132,489	241,867	386,077	415,364	196,203	379
1963	1,533,015	73,727	11,795	12,904	23,402	23,567	55,139	127,193	231,845	377,753	402,593	192,697	400
1962	1,488,113	75,812	11,784	12,927	21,757	23,385	54,437	124,926	223,886	368,469	386,217	184,121	392
1961	1,498,332	80,781	12,577	13,388	20,785	24,012	55,249	126,748	227,415	374,432	383,540	179,028	377
1960	1,505,335	82,479	13,200	13,488	20,880	24,895	56,157	127,927	231,480	378,847	380,982	174,563	437
WHITE, MALE													
1967	919,514	33,565	5,687	8,401	22,187	16,825	35,350	88,474	168,860	233,968	220,794	85,191	212
1966	924,342	36,010	6,224	8,481	21,600	16,527	35,693	89,009	168,384	236,569	221,423	84,171	251
1965	910,511	39,098	6,370	8,293	19,694	16,211	35,704	87,596	164,046	233,806	216,948	82,523	222
1964	895,518	42,211	6,629	8,288	18,689	16,303	35,912	86,385	160,599	231,645	209,853	78,780	224
1963	871,994	42,969	6,583	7,889	16,721	15,432	34,411	83,709	153,679	227,632	204,083	78,656	230
1962	846,458	44,270	6,454	7,884	15,609	15,103	34,036	82,197	148,108	221,351	195,320	75,895	231
1961	854,569	46,989	7,036	8,179	15,041	15,526	34,672	83,448	150,511	224,454	194,677	73,804	232
1960	860,857	48,063	7,409	8,255	15,063	16,221	35,139	84,963	152,430	227,995	193,078	71,974	267
WHITE, FEMALE													
1967	707,637	23,968	4,306	5,215	8,349	8,305	20,908	47,731	84,414	155,495	218,592	130,212	142
1966	710,119	25,739	4,865	5,319	8,082	8,614	21,165	47,569	83,546	157,881	218,708	128,495	136
1965	694,987	28,100	4,978	5,127	7,441	8,473	21,486	47,057	82,346	154,799	212,153	122,896	131
1964	683,156	30,517	5,182	5,214	7,244	8,552	21,554	46,104	81,268	154,432	205,511	117,423	155
1963	661,021	30,758	5,212	5,015	6,681	8,135	20,728	43,484	78,166	150,121	198,510	114,041	170
1962	641,655	31,542	5,330	5,043	6,148	8,282	20,401	42,729	75,778	147,118	190,897	108,226	161
1961	643,763	33,792	5,541	5,209	5,744	8,486	20,577	43,300	76,904	149,978	188,863	105,224	145
1960	644,478	34,416	5,791	5,233	5,817	8,674	21,018	42,964	79,050	150,852	187,904	102,589	170

Note: Figures by race exclude data for residents of New Jersey for 1962 and 1963.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290A for 1960-67 and separate unpublished tables for the black population and Accidents.

77/09/02 TABLE 290A. DEATHS FOR 60 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1960-67

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	TOTAL	UNDER 1 YEAR	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS	65-74 YEARS	75-84 YEARS	85+ YEARS	N.S.
ALL CAUSE													
ALL OTHER, BOTH SEXES													
1967	224,172	21,495	3,513	3,277	7,170	9,656	17,781	28,836	40,888	48,456	30,283	12,584	233
1966	228,688	23,767	4,000	3,333	6,644	9,257	17,978	29,033	41,432	49,385	30,891	12,722	246
1965	222,638	25,668	4,048	3,255	5,958	8,897	17,631	28,195	41,073	46,787	29,201	11,728	197
1964	219,377	27,055	4,165	3,248	5,544	8,784	17,418	28,160	41,182	44,816	27,764	11,009	232
1963	216,662	26,511	4,335	3,153	5,188	8,274	16,466	27,136	40,464	45,749	27,813	11,364	209
1962	206,583	26,535	4,019	3,032	4,906	7,965	15,894	26,165	38,454	43,005	25,777	10,654	177
1961	203,190	27,175	4,052	2,860	4,719	8,229	15,832	25,996	38,055	41,474	24,324	10,247	227
1960	206,647	28,394	4,482	3,049	4,653	8,519	15,946	26,949	38,717	41,465	24,227	9,963	283
ALL OTHER, MALE													
1967	126,431	11,877	1,964	1,960	4,998	6,093	10,395	16,889	23,626	26,914	15,869	5,715	131
1966	128,485	13,163	2,161	1,996	4,576	5,727	10,352	16,948	24,047	27,502	16,016	5,844	153
1965	124,689	14,321	2,158	1,911	4,060	5,390	9,980	16,365	23,418	26,233	15,371	5,359	123
1964	122,260	15,137	2,249	1,948	3,709	5,243	9,903	16,004	23,255	25,129	14,575	4,969	139
1963	120,406	14,907	2,318	1,791	3,461	4,775	9,035	15,393	22,592	25,879	14,796	5,332	127
1962	114,262	14,828	2,155	1,762	3,192	4,603	8,696	14,913	21,307	24,116	13,602	4,992	96
1961	112,258	15,131	2,148	1,702	3,070	4,720	8,723	14,613	21,031	23,417	12,744	4,834	125
1960	114,791	15,873	2,435	1,783	3,043	4,789	8,689	15,177	21,629	23,439	12,985	4,779	170
ALL OTHER, FEMALE													
1967	97,741	9,618	1,549	1,317	2,172	3,563	7,386	11,947	17,262	21,542	14,414	6,869	102
1966	100,203	10,604	1,839	1,337	2,068	3,530	7,626	12,085	17,385	21,883	14,875	6,878	93
1965	97,949	11,347	1,890	1,344	1,898	3,507	7,651	11,830	17,655	20,554	13,830	6,369	74
1964	97,117	11,918	1,916	1,300	1,835	3,541	7,515	12,156	17,927	19,687	13,189	6,040	93
1963	96,256	11,604	2,017	1,362	1,727	3,499	7,431	11,743	17,872	19,870	13,017	6,032	82
1962	92,321	11,707	1,864	1,270	1,714	3,362	7,198	11,252	17,147	18,889	12,175	5,662	81
1961	90,932	12,044	1,904	1,158	1,649	3,509	7,109	11,383	17,024	18,057	11,580	5,413	102
1960	91,856	12,521	2,047	1,266	1,610	3,730	7,257	11,772	17,088	18,026	11,242	5,184	113

Note: Figures by race exclude data for residents of New Jersey for 1962 and 1963.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290A for 1960-67 and separate unpublished tables for the black population and Accidents.

77/09/02 TABLE 290A. DEATHS FOR 60 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1960-67

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	TOTAL	UNDER 1 YEAR	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS	65-74 YEARS	75-84 YEARS	85+ YEARS	N.S.
Tuberculosis, all forms (001-019)													
BOTH SEXES													
1967	6,901	18	37	23	55	192	595	1,125	1,550	1,716	1,224	364	2
1966	7,625	12	62	28	69	222	711	1,226	1,736	1,821	1,342	388	8
1965	7,934	28	53	28	96	235	750	1,257	1,797	1,917	1,363	402	8
1964	8,303	18	61	36	109	303	816	1,353	1,844	2,007	1,386	366	4
1963	9,311	34	81	30	97	316	909	1,543	2,089	2,307	1,517	383	5
1962	9,506	43	91	51	90	401	986	1,621	2,129	2,292	1,459	342	1
1961	9,938	23	100	37	107	414	1,119	1,681	2,202	2,427	1,456	366	6
1960	10,866	37	91	45	137	545	1,239	1,913	2,416	2,547	1,521	368	7
MALE													
1967	5,002	7	20	10	25	105	372	800	1,196	1,359	880	228	-
1966	5,602	6	37	16	26	116	451	896	1,398	1,446	949	255	6
1965	5,789	14	18	14	46	131	429	909	1,447	1,543	988	243	7
1964	6,055	10	33	16	48	160	492	986	1,524	1,583	983	216	4
1963	6,835	17	39	16	53	156	546	1,122	1,698	1,835	1,101	249	3
1962	6,942	25	49	31	37	200	580	1,174	1,724	1,851	1,059	211	1
1961	7,254	15	54	20	43	208	667	1,251	1,795	1,939	1,048	211	3
1960	7,887	18	47	19	54	262	723	1,449	2,006	2,012	1,056	235	6
FEMALE													
1967	1,899	11	17	13	30	87	223	325	354	357	344	136	2
1966	2,023	6	25	12	43	106	260	330	338	375	393	133	2
1965	2,145	14	35	14	50	104	321	348	350	374	375	159	1
1964	2,248	8	28	20	61	143	324	367	320	424	403	150	-
1963	2,476	17	42	14	44	160	363	421	391	472	416	134	2
1962	2,564	18	42	20	53	201	406	447	405	441	400	131	-
1961	2,684	8	46	17	64	206	452	430	407	488	408	155	3
1960	2,979	19	44	26	83	283	516	464	410	535	465	133	1

- Quantity zero

Note: Figures by race exclude data for residents of New Jersey for 1962 and 1963.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290A for 1960-67 and separate unpublished tables for the black population and Accidents.

77/09/02 TABLE 290A. DEATHS FOR 60 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1960-67

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	TOTAL	UNDER 1 YEAR	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS	65-74 YEARS	75-84 YEARS	85+ YEARS	N.S.
Tuberculosis, all forms (001-019)													
WHITE, BOTH SEXES													
1967	4,883	10	20	10	25	76	269	687	1,138	1,308	1,038	300	2
1966	5,486	4	29	15	30	83	357	783	1,327	1,395	1,133	324	6
1965	5,797	13	23	14	41	88	391	831	1,351	1,534	1,165	343	3
1964	6,092	7	38	18	46	124	443	917	1,396	1,591	1,201	310	1
1963	6,586	17	39	17	41	120	472	1,009	1,526	1,777	1,266	298	4
1962	6,732	19	53	33	35	174	520	1,086	1,529	1,789	1,214	279	1
1961	7,391	11	62	25	48	166	605	1,180	1,675	2,029	1,274	315	1
1960	8,163	16	47	26	49	269	706	1,381	1,883	2,146	1,326	308	6
WHITE, MALE													
1967	3,574	5	9	4	11	47	160	489	884	1,036	745	184	-
1966	4,096	1	13	8	15	40	219	572	1,084	1,119	809	211	5
1965	4,323	9	10	6	23	44	240	603	1,100	1,238	843	204	3
1964	4,528	4	22	11	24	65	273	665	1,168	1,259	859	177	1
1963	4,923	9	15	11	24	67	285	736	1,252	1,422	910	189	3
1962	5,028	11	33	21	15	85	308	790	1,265	1,457	871	171	1
1961	5,517	7	32	14	20	84	360	876	1,391	1,641	910	181	1
1960	6,050	8	22	12	25	136	399	1,055	1,582	1,703	914	188	6
WHITE, FEMALE													
1967	1,309	5	11	6	14	29	109	198	254	272	293	116	2
1966	1,390	3	16	7	15	43	138	211	243	276	324	113	1
1965	1,474	4	13	8	18	44	151	228	251	296	322	139	-
1964	1,564	3	16	7	22	59	170	252	228	332	342	133	-
1963	1,663	8	24	6	17	53	187	273	274	355	356	109	1
1962	1,704	8	20	12	20	89	212	296	264	332	343	108	-
1961	1,874	4	30	11	28	82	245	304	284	388	364	134	-
1960	2,113	8	25	14	24	133	307	326	301	443	412	120	-

- Quantity zero

Note: Figures by race exclude data for residents of New Jersey for 1962 and 1963.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290A for 1960-67 and separate unpublished tables for the black population and Accidents.

77/09/02 TABLE 290A. DEATHS FOR 60 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1960-67

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	TOTAL	UNDER 1 YEAR	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS	65-74 YEARS	75-84 YEARS	85+ YEARS	N.S.
Tuberculosis, all forms (001-019)													
ALL OTHER, BOTH SEXES													
1967	2,018	8	17	13	30	116	326	438	412	408	186	64	-
1966	2,139	8	33	13	39	139	354	443	409	426	209	64	2
1965	2,137	15	30	14	55	147	359	426	446	383	198	59	5
1964	2,211	11	23	18	63	179	373	436	448	416	185	56	3
1963	2,362	17	38	12	50	173	398	469	482	448	201	73	1
1962	2,427	21	35	15	52	202	426	473	509	427	212	55	-
1961	2,547	12	38	12	59	248	514	501	527	398	182	51	5
1960	2,703	21	44	19	88	276	533	532	533	401	195	60	1
ALL OTHER, MALE													
1967	1,428	2	11	6	14	58	212	311	312	323	135	44	-
1966	1,506	5	24	8	11	76	232	324	314	327	140	44	1
1965	1,466	5	8	8	23	87	189	306	347	305	145	39	4
1964	1,527	6	11	5	24	95	219	321	356	324	124	39	3
1963	1,645	8	20	4	26	78	236	343	380	346	154	50	-
1962	1,660	14	14	9	22	101	248	336	389	330	161	36	-
1961	1,737	8	22	6	23	124	307	375	404	298	138	30	2
1960	1,837	10	25	7	29	126	324	394	424	309	142	47	-
ALL OTHER, FEMALE													
1967	590	6	6	7	16	58	114	127	100	85	51	20	-
1966	633	3	9	5	28	63	122	119	95	99	69	20	1
1965	671	10	22	6	32	60	170	120	99	78	53	20	1
1964	684	5	12	13	39	84	154	115	92	92	61	17	-
1963	717	9	18	8	24	95	162	126	102	102	47	23	1
1962	767	7	21	6	30	101	178	137	120	97	51	19	-
1961	810	4	16	6	36	124	207	126	123	100	44	21	3
1960	866	11	19	12	59	150	209	138	109	92	53	13	1

- Quantity zero

Note: Figures by race exclude data for residents of New Jersey for 1962 and 1963.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290A for 1960-67 and separate unpublished tables for the black population and Accidents.

77/077/09/02 TABLE 290A. DEATHS FOR 60 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1960-67

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	TOTAL	UNDER 1 YEAR	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS	65-74 YEARS	75-84 YEARS	85+ YEARS	N.S.
Tuberculosis of respiratory system (001-008)													
BOTH SEXES													
1967	6,351	8	10	11	34	147	530	1,041	1,459	1,607	1,153	349	2
1966	7,044	8	23	10	46	187	643	1,144	1,636	1,721	1,249	370	7
1965	7,358	13	16	12	63	206	689	1,162	1,706	1,812	1,291	380	8
1964	7,646	6	16	13	76	255	737	1,247	1,729	1,906	1,315	342	4
1963	8,608	15	25	11	66	273	839	1,446	1,949	2,175	1,437	367	5
1962	8,792	16	28	13	61	358	913	1,533	2,007	2,143	1,394	325	1
1961	9,191	10	30	5	81	359	1,018	1,573	2,074	2,302	1,387	347	5
1960	10,089	18	27	19	105	487	1,143	1,800	2,282	2,403	1,450	348	7
MALE													
1967	4,683	5	5	3	16	86	337	755	1,141	1,282	833	220	-
1966	5,260	3	10	5	19	100	412	854	1,338	1,383	885	245	6
1965	5,446	7	5	5	28	116	398	850	1,389	1,468	943	230	7
1964	5,663	3	8	3	34	135	451	914	1,441	1,518	952	200	4
1963	6,427	9	13	4	36	133	509	1,063	1,613	1,754	1,050	240	3
1962	6,531	9	15	7	25	178	542	1,127	1,650	1,750	1,027	200	1
1961	6,843	8	16	2	31	181	610	1,198	1,712	1,869	1,007	207	2
1960	7,440	11	15	10	40	236	671	1,375	1,914	1,915	1,022	225	6
FEMALE													
1967	1,668	3	5	8	18	61	193	286	318	325	320	129	2
1966	1,784	5	13	5	27	87	231	290	298	338	364	125	1
1965	1,912	6	11	7	35	90	291	312	317	344	348	150	1
1964	1,983	3	8	10	42	120	286	333	288	388	363	142	-
1963	2,181	6	12	7	30	140	330	383	336	421	387	127	2
1962	2,261	7	13	6	36	180	371	406	357	393	367	125	-
1961	2,348	2	14	3	50	178	408	375	362	433	380	140	3
1960	2,649	7	12	9	65	251	472	425	368	488	428	123	1

- Quantity zero

Note: Figures by race exclude data for residents of New Jersey for 1962 and 1963.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290A for 1960-67 and separate unpublished tables for the black population and Accidents.

77/09/02 TABLE 290A. DEATHS FOR 60 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1960-67

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	TOTAL	UNDER 1 YEAR	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS	65-74 YEARS	75-84 YEARS	85+ YEARS	N.S.
Tuberculosis of respiratory system (001-008)													
WHITE, BOTH SEXES													
1967	4,596	5	6	4	14	58	249	651	1,086	1,245	988	288	2
1966	5,159	2	14	6	20	70	331	745	1,264	1,328	1,062	311	6
1965	5,464	4	4	5	26	79	366	793	1,294	1,456	1,108	326	3
1964	5,711	3	9	4	34	105	409	856	1,325	1,525	1,147	293	1
1963	6,199	7	11	8	30	110	446	957	1,448	1,687	1,203	288	4
1962	6,344	7	16	10	25	166	493	1,042	1,462	1,690	1,166	266	1
1961	6,946	5	17	3	36	149	562	1,116	1,595	1,941	1,218	303	1
1960	7,685	8	7	12	36	241	656	1,313	1,798	2,043	1,273	292	6
WHITE, MALE													
1967	3,408	4	3	1	7	40	151	470	851	990	712	179	-
1966	3,894	-	5	3	13	32	203	550	1,046	1,074	760	203	5
1965	4,115	3	2	2	14	39	226	578	1,063	1,183	808	194	3
1964	4,292	1	7	2	17	56	255	622	1,113	1,220	832	166	1
1963	4,689	4	4	4	18	60	272	702	1,202	1,366	871	183	3
1962	4,792	4	10	6	11	79	292	766	1,219	1,396	847	161	1
1961	5,269	4	8	2	15	79	334	841	1,338	1,590	878	179	1
1960	5,777	5	5	8	21	120	374	1,005	1,525	1,635	891	182	6
WHITE, FEMALE													
1967	1,188	1	3	3	7	18	98	181	235	255	276	109	2
1966	1,265	2	9	3	7	38	128	195	218	254	302	108	1
1965	1,349	1	2	3	12	40	140	215	231	273	300	132	-
1964	1,419	2	2	2	17	49	154	234	212	305	315	127	-
1963	1,510	3	7	4	12	50	174	255	246	321	332	105	1
1962	1,552	3	6	4	14	87	201	276	243	294	319	105	-
1961	1,677	1	9	1	21	70	228	275	257	351	340	124	-
1960	1,908	3	2	4	15	121	282	308	273	408	382	110	-

- Quantity zero

Note: Figures by race exclude data for residents of New Jersey for 1962 and 1963.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290A for 1960-67 and separate unpublished tables for the black population and Accidents.

77/09/02 TABLE 290A. DEATHS FOR 60 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1960-67

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	TOTAL	UNDER 1 YEAR	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS	65-74 YEARS	75-84 YEARS	85+ YEARS	N.S.
Tuberculosis of respiratory system (001-008)													
ALL OTHER, BOTH SEXES													
1967	1,755	3	4	7	20	89	281	390	373	362	165	61	-
1966	1,885	6	9	4	26	117	312	399	372	393	187	59	1
1965	1,894	9	12	7	37	127	323	369	412	356	183	54	5
1964	1,935	3	7	9	42	150	328	391	404	381	168	49	3
1963	2,078	8	14	3	34	145	355	427	427	409	188	67	1
1962	2,139	7	12	3	34	170	388	432	461	383	196	53	-
1961	2,245	5	13	2	45	210	456	457	479	361	169	44	4
1960	2,404	10	20	7	69	246	487	487	484	360	177	56	1
ALL OTHER, MALE													
1967	1,275	1	2	2	9	46	186	285	290	292	121	41	-
1966	1,366	3	5	2	6	68	209	304	292	309	125	42	1
1965	1,331	4	3	3	14	77	172	272	326	285	135	36	4
1964	1,371	2	1	1	17	79	196	292	328	298	120	34	3
1963	1,488	5	9	-	17	63	212	319	349	322	145	47	-
1962	1,504	5	5	1	14	86	231	313	364	296	154	35	-
1961	1,574	4	8	-	16	102	276	357	374	279	129	28	1
1960	1,663	6	10	2	19	116	297	370	389	280	131	43	-
ALL OTHER, FEMALE													
1967	480	2	2	5	11	43	95	105	83	70	44	20	-
1966	519	3	4	2	20	49	103	95	80	84	62	17	-
1965	563	5	9	4	23	50	151	97	86	71	48	18	1
1964	564	1	6	8	25	71	132	99	76	83	48	15	-
1963	590	3	5	3	17	82	143	108	78	87	43	20	1
1962	635	2	7	2	20	84	157	119	97	87	42	18	-
1961	671	1	5	2	29	108	180	100	105	82	40	16	3
1960	741	4	10	5	50	130	190	117	95	80	46	13	1

- Quantity zero

Note: Figures by race exclude data for residents of New Jersey for 1962 and 1963.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290A for 1960-67 and separate unpublished tables for the black population and Accidents.

77/09/02 TABLE 290A. DEATHS FOR 60 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1960-67

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	TOTAL	UNDER 1 YEAR	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS	65-74 YEARS	75-84 YEARS	85+ YEARS	N.S.
Tuberculosis, other forms (010-019)													
BOTH SEXES													
1967	550	10	27	12	21	45	65	84	91	109	71	15	-
1966	581	4	39	18	23	35	68	82	100	100	93	18	1
1965	576	15	37	16	33	29	61	95	91	105	72	22	-
1964	657	12	45	23	33	48	79	106	115	101	71	24	-
1963	703	19	56	19	31	43	70	97	140	132	80	16	-
1962	714	27	63	38	29	43	73	88	122	149	65	17	-
1961	747	13	70	32	26	55	101	108	128	125	69	19	1
1960	777	19	64	26	32	58	96	113	134	144	71	20	-
MALE													
1967	319	2	15	7	9	19	35	45	55	77	47	8	-
1966	342	3	27	11	7	16	39	42	60	63	64	10	-
1965	343	7	13	9	18	15	31	59	58	75	45	13	-
1964	392	7	25	13	14	25	41	72	83	65	31	16	-
1963	408	8	26	12	17	23	37	59	85	81	51	9	-
1962	411	16	34	24	12	22	38	47	74	101	32	11	-
1961	411	7	38	18	12	27	57	53	83	70	41	4	1
1960	447	7	32	9	14	26	52	74	92	97	34	10	-
FEMALE													
1967	231	8	12	5	12	26	30	39	36	32	24	7	-
1966	239	1	12	7	16	19	29	40	40	37	29	8	1
1965	233	8	24	7	15	14	30	36	33	30	27	9	-
1964	265	5	20	10	19	23	38	34	32	36	40	8	-
1963	295	11	30	7	14	20	33	38	55	51	29	7	-
1962	303	11	29	14	17	21	35	41	48	48	33	6	-
1961	336	6	32	14	14	28	44	55	45	55	28	15	-
1960	330	12	32	17	18	32	44	39	42	47	37	10	-

- Quantity zero

Note: Figures by race exclude data for residents of New Jersey for 1962 and 1963.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290A for 1960-67 and separate unpublished tables for the black population and Accidents.

77/09/02 TABLE 290A. DEATHS FOR 60 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1960-67

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	TOTAL	UNDER 1 YEAR	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS	65-74 YEARS	75-84 YEARS	85+ YEARS	N.S.
Tuberculosis, other forms (010-019)													
WHITE, BOTH SEXES													
1967	287	5	14	6	11	18	20	36	52	63	50	12	-
1966	327	2	15	9	10	13	26	38	63	67	71	13	-
1965	333	9	19	9	15	9	25	38	57	78	57	17	-
1964	381	4	29	14	12	19	34	61	71	66	54	17	-
1963	387	10	28	9	11	10	26	52	78	90	63	10	-
1962	388	12	37	23	10	8	27	44	67	99	48	13	-
1961	445	6	45	22	12	17	43	64	80	88	56	12	-
1960	478	8	40	14	13	28	50	68	85	103	53	16	-
WHITE, MALE													
1967	166	1	6	3	4	7	9	19	33	46	33	5	-
1966	202	1	8	5	2	8	16	22	38	45	49	8	-
1965	208	6	8	4	9	5	14	25	37	55	35	10	-
1964	236	3	15	9	7	9	18	43	55	39	27	11	-
1963	234	5	11	7	6	7	13	34	50	56	39	6	-
1962	236	7	23	15	4	6	16	24	46	61	24	10	-
1961	248	3	24	12	5	5	26	35	53	51	32	2	-
1960	273	3	17	4	4	16	25	50	57	68	23	6	-
WHITE, FEMALE													
1967	121	4	8	3	7	11	11	17	19	17	17	7	-
1966	125	1	7	4	8	5	10	16	25	22	22	5	-
1965	125	3	11	5	6	4	11	13	20	23	22	7	-
1964	145	1	14	5	5	10	16	18	16	27	27	6	-
1963	153	5	17	2	5	3	13	18	28	34	24	4	-
1962	152	5	14	8	6	2	11	20	21	38	24	3	-
1961	197	3	21	10	7	12	17	29	27	37	24	10	-
1960	205	5	23	10	9	12	25	18	28	35	30	10	-

- Quantity zero

Note: Figures by race exclude data for residents of New Jersey for 1962 and 1963.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290A for 1960-67 and separate unpublished tables for the black population and Accidents.

77/09/02 TABLE 290A. DEATHS FOR 60 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1960-67

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	TOTAL	UNDER 1 YEAR	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS	65-74 YEARS	75-84 YEARS	85+ YEARS	N.S.
Tuberculosis, other forms (010-019)													
ALL OTHER, BOTH SEXES													
1967	263	5	13	6	10	27	45	48	39	46	21	3	-
1966	254	2	24	9	13	22	42	44	37	33	22	5	1
1965	243	6	18	7	18	20	36	57	34	27	15	5	-
1964	276	8	16	9	21	29	45	45	44	35	17	7	-
1963	284	9	24	9	16	28	43	42	55	39	13	6	-
1962	288	14	23	12	18	32	38	41	48	44	16	2	-
1961	302	7	25	10	14	38	58	44	48	37	13	7	1
1960	299	11	24	12	19	30	46	45	49	41	18	4	-
ALL OTHER, MALE													
1967	153	1	9	4	5	12	26	26	22	31	14	3	-
1966	140	2	19	6	5	8	23	20	22	18	15	2	-
1965	135	1	5	5	9	10	17	34	21	20	10	3	-
1964	156	4	10	4	7	16	23	29	28	26	4	5	-
1963	157	3	11	4	9	15	24	24	31	24	9	3	-
1962	156	9	9	8	8	15	17	23	25	34	7	1	-
1961	163	4	14	6	7	22	31	18	30	19	9	2	1
1960	174	4	15	5	10	10	27	24	35	29	11	4	-
ALL OTHER, FEMALE													
1967	110	4	4	2	5	15	19	22	17	15	7	-	-
1966	114	-	5	3	8	14	19	24	15	15	7	3	1
1965	108	5	13	2	9	10	19	23	13	7	5	2	-
1964	120	4	6	5	14	13	22	16	16	9	13	2	-
1963	127	6	13	5	7	13	19	18	24	15	4	3	-
1962	132	5	14	4	10	17	21	18	23	10	9	1	-
1961	139	3	11	4	7	16	27	26	18	18	4	5	-
1960	125	7	9	7	9	20	19	21	14	12	7	-	-

- Quantity zero

Note: Figures by race exclude data for residents of New Jersey for 1962 and 1963.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290A for 1960-67 and separate unpublished tables for the black population and Accidents.

77/09/02 TABLE 290A. DEATHS FOR 60 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1960-67

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	TOTAL	UNDER 1 YEAR	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS	65-74 YEARS	75-84 YEARS	85+ YEARS	N.S.
Syphilis and its sequelae (020-029)													
BOTH SEXES													
1967	2,381	15	-	2	11	18	62	176	563	841	559	130	4
1966	2,193	25	-	-	8	10	45	170	540	791	496	106	2
1965	2,434	25	-	4	7	26	53	240	587	902	514	76	-
1964	2,619	20	1	1	7	30	85	276	692	859	533	115	-
1963	2,666	19	1	-	7	21	91	294	770	879	494	90	-
1962	2,811	29	-	2	10	18	99	332	782	939	492	107	1
1961	2,850	20	1	1	8	37	115	382	838	941	414	90	3
1960	2,945	30	-	2	9	35	111	389	872	1,003	402	88	4
MALE													
1967	1,684	7	-	2	9	10	43	112	412	645	373	69	2
1966	1,558	15	-	-	6	8	24	118	404	597	330	54	2
1965	1,732	14	-	3	4	17	35	177	422	684	340	36	-
1964	1,829	14	1	1	6	21	53	182	515	642	333	61	-
1963	1,904	9	-	-	6	16	65	198	571	678	329	32	-
1962	1,998	20	-	1	8	9	56	232	583	718	316	55	-
1961	2,032	14	-	1	5	23	78	263	630	701	263	51	3
1960	2,145	21	-	2	7	21	75	275	651	778	266	46	3
FEMALE													
1967	697	8	-	-	2	8	19	64	151	196	186	61	2
1966	635	10	-	-	2	2	21	52	136	194	166	52	-
1965	702	11	-	1	3	9	18	63	165	218	174	40	-
1964	790	6	-	-	1	9	32	94	177	217	200	54	-
1963	762	10	1	-	1	5	26	96	199	201	165	58	-
1962	813	9	-	1	2	9	43	100	199	221	176	52	1
1961	818	6	1	-	3	14	37	119	208	240	151	39	-
1960	800	9	-	-	2	14	36	114	221	225	136	42	1

- Quantity zero

Note: Figures by race exclude data for residents of New Jersey for 1962 and 1963.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290A for 1960-67 and separate unpublished tables for the black population and Accidents.

77/09/02 TABLE 290A. DEATHS FOR 60 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1960-67

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	TOTAL	UNDER 1 YEAR	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS	65-74 YEARS	75-84 YEARS	85+ YEARS	N.S.
Syphilis and its sequelae (020-029)													
WHITE, BOTH SEXES													
1967	1,855	6	-	2	9	13	40	96	406	675	490	117	1
1966	1,670	8	-	-	5	7	21	95	391	620	429	94	-
1965	1,789	11	-	3	4	11	23	127	396	707	442	65	-
1964	1,889	8	1	-	6	16	42	139	465	652	459	101	-
1963	1,803	4	-	-	2	8	32	156	482	639	408	72	-
1962	1,918	7	-	2	6	13	42	169	507	680	405	86	1
1961	1,892	8	-	1	7	25	48	166	527	698	338	73	1
1960	2,019	15	-	1	6	16	52	192	570	762	333	70	2
WHITE, MALE													
1967	1,347	3	-	2	7	6	30	70	315	529	322	62	1
1966	1,224	5	-	-	4	7	13	76	299	483	288	49	-
1965	1,288	7	-	2	3	6	16	95	293	545	293	28	-
1964	1,344	5	1	-	5	13	26	99	362	498	283	52	-
1963	1,307	2	-	-	2	7	24	111	367	496	269	29	-
1962	1,391	4	-	1	6	8	28	125	390	528	257	44	-
1961	1,405	6	-	1	5	18	36	121	427	533	219	38	1
1960	1,492	11	-	1	5	12	37	142	437	593	219	33	2
WHITE, FEMALE													
1967	508	3	-	-	2	7	10	26	91	146	168	55	-
1966	446	3	-	-	1	-	8	19	92	137	141	45	-
1965	501	4	-	1	1	5	7	32	103	162	149	37	-
1964	545	3	-	-	1	3	16	40	103	154	176	49	-
1963	496	2	-	-	-	1	8	45	115	143	139	43	-
1962	527	3	-	1	-	5	14	44	117	152	148	42	1
1961	487	2	-	-	2	7	12	45	100	165	119	35	-
1960	527	4	-	-	1	4	15	50	133	169	114	37	-

- Quantity zero

Note: Figures by race exclude data for residents of New Jersey for 1962 and 1963.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290A for 1960-67 and separate unpublished tables for the black population and Accidents.

77/09/02 TABLE 290A. DEATHS FOR 60 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1960-67

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	TOTAL	UNDER 1 YEAR	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS	65-74 YEARS	75-84 YEARS	85+ YEARS	N.S.
Syphilis and its sequelae (020-029)													
ALL OTHER, BOTH SEXES													
1967	526	9	-	-	2	5	22	80	157	166	69	13	3
1966	523	17	-	-	3	3	24	75	149	171	67	12	2
1965	645	14	-	1	3	15	30	113	191	195	72	11	-
1964	730	12	-	1	1	14	43	137	227	207	74	14	-
1963	798	14	1	-	5	13	56	127	274	225	68	15	-
1962	828	21	-	-	4	4	54	159	254	237	76	19	-
1961	958	12	1	-	1	12	67	216	311	243	76	17	2
1960	926	15	-	1	3	19	59	197	302	241	69	18	2
ALL OTHER, MALE													
1967	337	4	-	-	2	4	13	42	97	116	51	7	1
1966	334	10	-	-	2	1	11	42	105	114	42	5	2
1965	444	7	-	1	1	11	19	82	129	139	47	8	-
1964	485	9	-	1	1	8	27	83	153	144	50	9	-
1963	548	7	-	-	4	9	39	77	192	171	46	3	-
1962	564	15	-	-	2	-	27	105	180	176	49	10	-
1961	627	8	-	-	-	5	42	142	203	168	44	13	2
1960	653	10	-	1	2	9	38	133	214	185	47	13	1
ALL OTHER, FEMALE													
1967	189	5	-	-	-	1	9	38	60	50	18	6	2
1966	189	7	-	-	1	2	13	33	44	57	25	7	-
1965	201	7	-	-	2	4	11	31	62	56	25	3	-
1964	245	3	-	-	-	6	16	54	74	63	24	5	-
1963	250	7	1	-	1	4	17	50	82	54	22	12	-
1962	264	6	-	-	2	4	27	54	74	61	27	9	-
1961	331	4	1	-	1	7	25	74	108	75	32	4	-
1960	273	5	-	-	1	10	21	64	88	56	22	5	1

- Quantity zero

Note: Figures by race exclude data for residents of New Jersey for 1962 and 1963.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290A for 1960-67 and separate unpublished tables for the black population and Accidents.

77/09/02 TABLE 290A. DEATHS FOR 60 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1960-67

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	TOTAL	UNDER 1 YEAR	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS	65-74 YEARS	75-84 YEARS	85+ YEARS	N.S.
Dysentery, all forms (045-048)													
BOTH SEXES													
1967	157	32	28	17	5	2	10	15	11	19	11	6	1
1966	172	44	31	8	5	2	7	11	17	21	18	8	-
1965	243	85	35	14	6	1	11	15	18	27	20	11	-
1964	283	108	47	20	1	4	9	13	14	28	31	8	-
1963	294	94	43	31	6	12	11	15	31	15	20	16	-
1962	323	115	46	25	5	8	8	20	28	27	27	14	-
1961	300	102	44	24	3	7	10	22	26	21	29	12	-
1960	355	140	56	23	7	6	11	17	23	31	32	9	-
MALE													
1967	96	18	13	12	3	2	7	13	9	13	6	-	-
1966	102	26	16	6	2	1	4	8	13	11	12	3	-
1965	141	44	21	8	5	1	6	10	13	15	13	5	-
1964	158	59	28	10	1	3	5	10	8	13	18	3	-
1963	171	46	28	17	4	8	9	8	20	13	12	6	-
1962	183	67	23	16	3	5	4	14	19	12	15	5	-
1961	166	54	26	14	1	5	6	15	13	10	17	5	-
1960	189	83	33	9	5	2	7	8	12	13	13	4	-
FEMALE													
1967	61	14	15	5	2	-	3	2	2	6	5	6	1
1966	70	18	15	2	3	1	3	3	4	10	6	5	-
1965	102	41	14	6	1	-	5	5	5	12	7	6	-
1964	125	49	19	10	-	1	4	3	6	15	13	5	-
1963	123	48	15	14	2	4	2	7	11	2	8	10	-
1962	140	48	23	9	2	3	4	6	9	15	12	9	-
1961	134	48	18	10	2	2	4	7	13	11	12	7	-
1960	166	57	23	14	2	4	4	9	11	18	19	5	-

- Quantity zero

Note: Figures by race exclude data for residents of New Jersey for 1962 and 1963.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290A for 1960-67 and separate unpublished tables for the black population and Accidents.

77/09/02 TABLE 290A. DEATHS FOR 60 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1960-67

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	TOTAL	UNDER 1 YEAR	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS	65-74 YEARS	75-84 YEARS	85+ YEARS	N.S.
Dysentery, all forms (045-048)													
WHITE, BOTH SEXES													
1967	97	11	13	14	4	2	6	10	6	17	9	4	1
1966	100	20	11	5	3	1	5	10	11	14	13	7	-
1965	131	32	18	7	5	1	5	11	13	17	14	8	-
1964	156	44	19	14	1	1	7	13	8	18	24	7	-
1963	160	38	19	22	4	5	7	12	18	8	15	12	-
1962	193	52	23	18	3	7	6	17	20	16	20	11	-
1961	184	48	26	20	3	4	6	15	17	15	21	9	-
1960	203	60	30	16	5	5	9	12	13	23	23	7	-
WHITE, MALE													
1967	62	5	6	10	2	2	5	9	5	12	6	-	-
1966	65	11	7	5	2	1	3	7	8	8	10	3	-
1965	83	16	12	5	5	1	3	7	9	11	10	4	-
1964	93	25	13	7	1	1	4	10	5	10	15	2	-
1963	98	19	15	11	4	4	6	8	12	7	8	4	-
1962	114	28	15	12	2	5	4	12	15	7	10	4	-
1961	108	28	15	12	1	3	4	11	9	9	12	4	-
1960	109	38	18	8	4	1	7	5	6	9	11	2	-
WHITE, FEMALE													
1967	35	6	7	4	2	-	1	1	1	5	3	4	1
1966	35	9	4	-	1	-	2	3	3	6	3	4	-
1965	48	16	6	2	-	-	2	4	4	6	4	4	-
1964	63	19	6	7	-	-	3	3	3	8	9	5	-
1963	62	19	4	11	-	1	1	4	6	1	7	8	-
1962	79	24	8	6	1	2	2	5	5	9	10	7	-
1961	76	20	11	8	2	1	2	4	8	6	9	5	-
1960	94	22	12	8	1	4	2	7	7	14	12	5	-

- Quantity zero

Note: Figures by race exclude data for residents of New Jersey for 1962 and 1963.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290A for 1960-67 and separate unpublished tables for the black population and Accidents.

77/09/02 TABLE 290A. DEATHS FOR 60 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1960-67

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	TOTAL	UNDER 1 YEAR	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS	65-74 YEARS	75-84 YEARS	85+ YEARS	N.S.
Dysentery, all forms (045-048)													
ALL OTHER, BOTH SEXES													
1967	60	21	15	3	1	-	4	5	5	2	2	2	-
1966	72	24	20	3	2	1	2	1	6	7	5	1	-
1965	112	53	17	7	1	-	6	4	5	10	6	3	-
1964	127	64	28	6	-	3	2	-	6	10	7	1	-
1963	130	55	24	9	2	6	4	3	11	7	5	4	-
1962	129	63	23	7	2	1	2	2	8	11	7	3	-
1961	116	54	18	4	-	3	4	7	9	6	8	3	-
1960	152	80	26	7	2	1	2	5	10	8	9	2	-
ALL OTHER, MALE													
1967	34	13	7	2	1	-	2	4	4	1	-	-	-
1966	37	15	9	1	-	-	1	1	5	3	2	-	-
1965	58	28	9	3	-	-	3	3	4	4	3	1	-
1964	65	34	15	3	-	2	1	-	3	3	3	1	-
1963	70	26	13	6	-	3	3	-	7	6	4	2	-
1962	68	39	8	4	1	-	-	1	4	5	5	1	-
1961	58	26	11	2	-	2	2	4	4	1	5	1	-
1960	80	45	15	1	1	1	-	3	6	4	2	2	-
ALL OTHER, FEMALE													
1967	26	8	8	1	-	-	2	1	1	1	2	2	-
1966	35	9	11	2	2	1	1	-	1	4	3	1	-
1965	54	25	8	4	1	-	3	1	1	6	3	2	-
1964	62	30	13	3	-	1	1	-	3	7	4	-	-
1963	60	29	11	3	2	3	1	3	4	1	1	2	-
1962	61	24	15	3	1	1	2	1	4	6	2	2	-
1961	58	28	7	2	-	1	2	3	5	5	3	2	-
1960	72	35	11	6	1	-	2	2	4	4	7	-	-

- Quantity zero

Note: Figures by race exclude data for residents of New Jersey for 1962 and 1963.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290A for 1960-67 and separate unpublished tables for the black population and Accidents.

77/09/02 TABLE 290A. DEATHS FOR 60 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1960-67

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	TOTAL	UNDER 1 YEAR	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS	65-74 YEARS	75-84 YEARS	85+ YEARS	N.S.
Scarlet fever and streptococcal sore throat (050,051)													
BOTH SEXES													
1967	47	4	8	9	1	1	1	8	8	1	5	1	-
1966	65	5	7	12	3	5	2	7	8	4	8	4	-
1965	63	3	7	9	7	4	5	5	7	7	6	3	-
1964	95	9	12	11	5	2	8	7	7	7	16	11	-
1963	102	10	14	21	9	2	4	13	8	7	8	6	-
1962	102	16	11	9	11	5	7	4	9	9	13	8	-
1961	130	16	27	18	11	7	7	10	6	7	13	8	-
1960	108	9	17	19	6	11	7	8	9	10	10	2	-
MALE													
1967	24	3	4	4	-	1	-	5	5	-	1	1	-
1966	29	1	3	6	1	4	-	4	4	2	2	2	-
1965	31	2	5	4	3	1	4	1	3	2	5	1	-
1964	49	7	7	7	1	1	5	2	4	2	7	6	-
1963	50	3	7	12	5	1	3	10	1	3	3	2	-
1962	51	8	5	2	5	2	5	2	8	5	6	3	-
1961	66	11	11	10	4	2	3	9	3	3	6	4	-
1960	54	3	12	11	2	3	4	6	3	5	5	-	-
FEMALE													
1967	23	1	4	5	1	-	1	3	3	1	4	-	-
1966	36	4	4	6	2	1	2	3	4	2	6	2	-
1965	32	1	2	5	4	3	1	4	4	5	1	2	-
1964	46	2	5	4	4	1	3	5	3	5	9	5	-
1963	52	7	7	9	4	1	1	3	7	4	5	4	-
1962	51	8	6	7	6	3	2	2	1	4	7	5	-
1961	64	5	16	8	7	5	4	1	3	4	7	4	-
1960	54	6	5	8	4	8	3	2	6	5	5	2	-

- Quantity zero

Note: Figures by race exclude data for residents of New Jersey for 1962 and 1963.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290A for 1960-67 and separate unpublished tables for the black population and Accidents.

77/09/02 TABLE 290A. DEATHS FOR 60 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1960-67

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	TOTAL	UNDER 1 YEAR	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS	65-74 YEARS	75-84 YEARS	85+ YEARS	N.S.
Scarlet fever and streptococcal sore throat (050,051)													
WHITE, BOTH SEXES													
1967	34	2	7	8	1	1	-	5	5	1	3	1	-
1966	53	3	6	10	2	3	1	6	8	4	6	4	-
1965	46	2	7	6	5	3	3	4	4	6	4	2	-
1964	83	6	11	9	4	2	7	4	6	7	16	11	-
1963	67	4	9	10	6	2	1	9	6	6	8	6	-
1962	75	13	8	6	9	4	3	3	4	6	11	8	-
1961	114	11	21	18	10	7	4	9	6	7	13	8	-
1960	95	8	14	16	4	11	5	6	9	10	10	2	-
WHITE, MALE													
1967	19	2	3	4	-	1	-	3	4	-	1	1	-
1966	24	1	3	5	-	2	-	4	4	2	1	2	-
1965	24	1	5	2	2	1	3	1	3	1	4	1	-
1964	42	5	6	5	1	1	5	1	3	2	7	6	-
1963	32	1	4	5	5	1	1	7	1	2	3	2	-
1962	38	7	5	2	3	2	2	2	4	2	6	3	-
1961	56	7	7	10	3	2	3	8	3	3	6	4	-
1960	49	3	10	10	1	3	4	5	3	5	5	-	-
WHITE, FEMALE													
1967	15	-	4	4	1	-	-	2	1	1	2	-	-
1966	29	2	3	5	2	1	1	2	4	2	5	2	-
1965	22	1	2	4	3	2	-	3	1	5	-	1	-
1964	41	1	5	4	3	1	2	3	3	5	9	5	-
1963	35	3	5	5	1	1	-	2	5	4	5	4	-
1962	37	6	3	4	6	2	1	1	-	4	5	5	-
1961	58	4	14	8	7	5	1	1	3	4	7	4	-
1960	46	5	4	6	3	8	1	1	6	5	5	2	-

- Quantity zero

Note: Figures by race exclude data for residents of New Jersey for 1962 and 1963.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290A for 1960-67 and separate unpublished tables for the black population and Accidents.

77/09/02 TABLE 290A. DEATHS FOR 60 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1960-67

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	TOTAL	UNDER 1 YEAR	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS	65-74 YEARS	75-84 YEARS	85+ YEARS	N.S.
Scarlet fever and streptococcal sore throat (050,051)													
ALL OTHER, BOTH SEXES													
1967	13	2	1	1	-	-	1	3	3	-	2	-	-
1966	12	2	1	2	1	2	1	1	-	-	2	-	-
1965	17	1	-	3	2	1	2	1	3	1	2	1	-
1964	12	3	1	2	1	-	1	3	1	-	-	-	-
1963	30	6	4	11	3	-	3	3	-	-	-	-	-
1962	24	3	3	3	1	1	4	1	4	3	1	-	-
1961	16	5	6	-	1	-	3	1	-	-	-	-	-
1960	13	1	3	3	2	-	2	2	-	-	-	-	-
ALL OTHER, MALE													
1967	5	1	1	-	-	-	-	2	1	-	-	-	-
1966	5	-	-	1	1	2	-	-	-	-	1	-	-
1965	7	1	-	2	1	-	1	-	-	1	1	-	-
1964	7	2	1	2	-	-	-	1	1	-	-	-	-
1963	15	2	2	7	-	-	2	2	-	-	-	-	-
1962	11	1	-	-	1	-	3	-	3	3	-	-	-
1961	10	4	4	-	1	-	-	1	-	-	-	-	-
1960	5	-	2	1	1	-	-	1	-	-	-	-	-
ALL OTHER, FEMALE													
1967	8	1	-	1	-	-	1	1	2	-	2	-	-
1966	7	2	1	1	-	-	1	1	-	-	1	-	-
1965	10	-	-	1	1	1	1	1	3	-	1	1	-
1964	5	1	-	-	1	-	1	2	-	-	-	-	-
1963	15	4	2	4	3	-	1	1	-	-	-	-	-
1962	13	2	3	3	-	1	1	1	1	-	1	-	-
1961	6	1	2	-	-	-	3	-	-	-	-	-	-
1960	8	1	1	2	1	-	2	1	-	-	-	-	-

- Quantity zero

Note: Figures by race exclude data for residents of New Jersey for 1962 and 1963.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290A for 1960-67 and separate unpublished tables for the black population and Accidents.

77/09/02 TABLE 290A. DEATHS FOR 60 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1960-67

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	TOTAL	UNDER 1 YEAR	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS	65-74 YEARS	75-84 YEARS	85+ YEARS	N.S.
Diphtheria (055)													
BOTH SEXES													
1967	32	1	11	14	1	-	2	2	-	-	1	-	-
1966	20	-	8	10	-	-	-	-	-	2	-	-	-
1965	18	-	7	9	-	-	-	1	1	-	-	-	-
1964	42	2	16	13	1	1	2	4	1	1	1	-	-
1963	45	3	16	18	2	-	-	1	4	1	-	-	-
1962	41	-	18	15	1	2	1	2	1	1	-	-	-
1961	68	2	30	29	1	-	1	2	2	1	-	-	-
1960	69	2	38	24	1	1	1	1	-	1	-	-	-
MALE													
1967	14	-	8	4	-	-	-	1	-	-	1	-	-
1966	9	-	4	3	-	-	-	-	-	2	-	-	-
1965	10	-	5	5	-	-	-	-	-	-	-	-	-
1964	17	1	8	4	-	-	-	2	1	-	1	-	-
1963	25	2	11	8	1	-	-	1	1	1	-	-	-
1962	23	-	12	8	-	1	1	-	1	-	-	-	-
1961	38	1	19	15	-	-	-	2	1	-	-	-	-
1960	32	1	16	13	1	1	-	-	-	-	-	-	-
FEMALE													
1967	18	1	3	10	1	-	2	1	-	-	-	-	-
1966	11	-	4	7	-	-	-	-	-	-	-	-	-
1965	8	-	2	4	-	-	-	1	1	-	-	-	-
1964	25	1	8	9	1	1	2	2	-	1	-	-	-
1963	20	1	5	10	1	-	-	-	3	-	-	-	-
1962	18	-	6	7	1	1	-	2	-	1	-	-	-
1961	30	1	11	14	1	-	1	-	1	1	-	-	-
1960	37	1	22	11	-	-	1	1	-	1	-	-	-

- Quantity zero

Note: Figures by race exclude data for residents of New Jersey for 1962 and 1963.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290A for 1960-67 and separate unpublished tables for the black population and Accidents.

77/09/02 TABLE 290A. DEATHS FOR 60 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1960-67

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	TOTAL	UNDER 1 YEAR	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS	65-74 YEARS	75-84 YEARS	85+ YEARS	N.S.
Diphtheria (055)													
WHITE, BOTH SEXES													
1967	16	-	5	9	-	-	-	1	-	-	1	-	-
1966	14	-	4	8	-	-	-	-	-	2	-	-	-
1965	10	-	5	3	-	-	-	1	1	-	-	-	-
1964	30	-	11	10	1	-	2	3	1	1	1	-	-
1963	21	1	5	10	-	-	-	1	3	1	-	-	-
1962	22	-	9	9	-	1	1	1	1	-	-	-	-
1961	45	2	16	21	1	-	1	2	1	1	-	-	-
1960	44	2	21	16	1	1	1	1	-	1	-	-	-
WHITE, MALE													
1967	8	-	4	3	-	-	-	-	-	-	1	-	-
1966	5	-	2	1	-	-	-	-	-	2	-	-	-
1965	5	-	3	2	-	-	-	-	-	-	-	-	-
1964	14	-	7	3	-	-	-	2	1	-	1	-	-
1963	15	-	5	7	-	-	-	1	1	1	-	-	-
1962	11	-	5	4	-	-	1	-	1	-	-	-	-
1961	24	1	10	11	-	-	-	2	-	-	-	-	-
1960	23	1	9	11	1	1	-	-	-	-	-	-	-
WHITE, FEMALE													
1967	8	-	1	6	-	-	-	1	-	-	-	-	-
1966	9	-	2	7	-	-	-	-	-	-	-	-	-
1965	5	-	2	1	-	-	-	1	1	-	-	-	-
1964	16	-	4	7	1	-	2	1	-	1	-	-	-
1963	6	1	-	3	-	-	-	-	2	-	-	-	-
1962	11	-	4	5	-	1	-	1	-	-	-	-	-
1961	21	1	6	10	1	-	1	-	1	1	-	-	-
1960	21	1	12	5	-	-	1	1	-	1	-	-	-

- Quantity zero

Note: Figures by race exclude data for residents of New Jersey for 1962 and 1963.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290A for 1960-67 and separate unpublished tables for the black population and Accidents.

77/09/02 TABLE 290A. DEATHS FOR 60 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1960-67

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	TOTAL	UNDER 1 YEAR	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS	65-74 YEARS	75-84 YEARS	85+ YEARS	N.S.
Diphtheria (055)													
ALL OTHER, BOTH SEXES													
1967	16	1	6	5	1	-	2	1	-	-	-	-	-
1966	6	-	4	2	-	-	-	-	-	-	-	-	-
1965	8	-	2	6	-	-	-	-	-	-	-	-	-
1964	12	2	5	3	-	1	-	1	-	-	-	-	-
1963	22	2	11	7	1	-	-	-	1	-	-	-	-
1962	18	-	9	6	-	1	-	1	-	1	-	-	-
1961	23	-	14	8	-	-	-	-	1	-	-	-	-
1960	25	-	17	8	-	-	-	-	-	-	-	-	-
ALL OTHER, MALE													
1967	6	-	4	1	-	-	-	1	-	-	-	-	-
1966	4	-	2	2	-	-	-	-	-	-	-	-	-
1965	5	-	2	3	-	-	-	-	-	-	-	-	-
1964	3	1	1	1	-	-	-	-	-	-	-	-	-
1963	9	2	6	-	1	-	-	-	-	-	-	-	-
1962	12	-	7	4	-	1	-	-	-	-	-	-	-
1961	14	-	9	4	-	-	-	-	1	-	-	-	-
1960	9	-	7	2	-	-	-	-	-	-	-	-	-
ALL OTHER, FEMALE													
1967	10	1	2	4	1	-	2	-	-	-	-	-	-
1966	2	-	2	-	-	-	-	-	-	-	-	-	-
1965	3	-	-	3	-	-	-	-	-	-	-	-	-
1964	9	1	4	2	-	1	-	1	-	-	-	-	-
1963	13	-	5	7	-	-	-	-	1	-	-	-	-
1962	6	-	2	2	-	-	-	1	-	1	-	-	-
1961	9	-	5	4	-	-	-	-	-	-	-	-	-
1960	16	-	10	6	-	-	-	-	-	-	-	-	-

- Quantity zero

Note: Figures by race exclude data for residents of New Jersey for 1962 and 1963.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290A for 1960-67 and separate unpublished tables for the black population and Accidents.

77/09/02 TABLE 290A. DEATHS FOR 60 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1960-67

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	TOTAL	UNDER 1 YEAR	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS	65-74 YEARS	75-84 YEARS	85+ YEARS	N.S.
Whooping cough (056)													
BOTH SEXES													
1967	37	29	6	1	-	-	-	-	-	1	-	-	-
1966	49	33	15	-	-	-	-	-	-	-	1	-	-
1965	55	42	13	-	-	-	-	-	-	-	-	-	-
1964	93	66	21	4	-	2	-	-	-	-	-	-	-
1963	115	88	26	1	-	-	-	-	-	-	-	-	-
1962	83	66	14	1	1	1	-	-	-	-	-	-	-
1961	76	54	20	1	-	-	-	-	-	1	-	-	-
1960	118	73	40	2	1	-	2	-	-	-	-	-	-
MALE													
1967	19	15	2	1	-	-	-	-	-	1	-	-	-
1966	23	18	5	-	-	-	-	-	-	-	-	-	-
1965	34	26	8	-	-	-	-	-	-	-	-	-	-
1964	44	33	8	3	-	-	-	-	-	-	-	-	-
1963	52	42	9	1	-	-	-	-	-	-	-	-	-
1962	45	37	6	1	-	1	-	-	-	-	-	-	-
1961	33	22	11	-	-	-	-	-	-	-	-	-	-
1960	56	44	11	1	-	-	-	-	-	-	-	-	-
FEMALE													
1967	18	14	4	-	-	-	-	-	-	-	-	-	-
1966	26	15	10	-	-	-	-	-	-	-	1	-	-
1965	21	16	5	-	-	-	-	-	-	-	-	-	-
1964	49	33	13	1	-	2	-	-	-	-	-	-	-
1963	63	46	17	-	-	-	-	-	-	-	-	-	-
1962	38	29	8	-	1	-	-	-	-	-	-	-	-
1961	43	32	9	1	-	-	-	-	-	1	-	-	-
1960	62	29	29	1	1	-	2	-	-	-	-	-	-

- Quantity zero

Note: Figures by race exclude data for residents of New Jersey for 1962 and 1963.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290A for 1960-67 and separate unpublished tables for the black population and Accidents.

77/09/02 TABLE 290A. DEATHS FOR 60 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1960-67

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	TOTAL	UNDER 1 YEAR	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS	65-74 YEARS	75-84 YEARS	85+ YEARS	N.S.
Whooping cough (056)													
WHITE, BOTH SEXES													
1967	14	9	4	1	-	-	-	-	-	-	-	-	-
1966	27	17	9	-	-	-	-	-	-	-	1	-	-
1965	23	20	3	-	-	-	-	-	-	-	-	-	-
1964	28	18	6	3	-	1	-	-	-	-	-	-	-
1963	47	41	6	-	-	-	-	-	-	-	-	-	-
1962	34	26	6	-	1	1	-	-	-	-	-	-	-
1961	26	18	7	-	-	-	-	-	-	1	-	-	-
1960	45	24	17	1	1	-	2	-	-	-	-	-	-
WHITE, MALE													
1967	6	4	1	1	-	-	-	-	-	-	-	-	-
1966	13	10	3	-	-	-	-	-	-	-	-	-	-
1965	14	13	1	-	-	-	-	-	-	-	-	-	-
1964	11	8	1	2	-	-	-	-	-	-	-	-	-
1963	20	19	1	-	-	-	-	-	-	-	-	-	-
1962	21	18	2	-	-	1	-	-	-	-	-	-	-
1961	12	8	4	-	-	-	-	-	-	-	-	-	-
1960	19	15	3	1	-	-	-	-	-	-	-	-	-
WHITE, FEMALE													
1967	8	5	3	-	-	-	-	-	-	-	-	-	-
1966	14	7	6	-	-	-	-	-	-	-	1	-	-
1965	9	7	2	-	-	-	-	-	-	-	-	-	-
1964	17	10	5	1	-	1	-	-	-	-	-	-	-
1963	27	22	5	-	-	-	-	-	-	-	-	-	-
1962	13	8	4	-	1	-	-	-	-	-	-	-	-
1961	14	10	3	-	-	-	-	-	-	1	-	-	-
1960	26	9	14	-	1	-	2	-	-	-	-	-	-

- Quantity zero

Note: Figures by race exclude data for residents of New Jersey for 1962 and 1963.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290A for 1960-67 and separate unpublished tables for the black population and Accidents.

77/09/02 TABLE 290A. DEATHS FOR 60 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1960-67

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	TOTAL	UNDER 1 YEAR	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS	65-74 YEARS	75-84 YEARS	85+ YEARS	N.S.
Whooping cough (056)													
ALL OTHER, BOTH SEXES													
1967	23	20	2	-	-	-	-	-	-	1	-	-	-
1966	22	16	6	-	-	-	-	-	-	-	-	-	-
1965	32	22	10	-	-	-	-	-	-	-	-	-	-
1964	65	48	15	1	-	1	-	-	-	-	-	-	-
1963	67	46	20	1	-	-	-	-	-	-	-	-	-
1962	49	40	8	1	-	-	-	-	-	-	-	-	-
1961	50	36	13	1	-	-	-	-	-	-	-	-	-
1960	73	49	23	1	-	-	-	-	-	-	-	-	-
ALL OTHER, MALE													
1967	13	11	1	-	-	-	-	-	-	1	-	-	-
1966	10	8	2	-	-	-	-	-	-	-	-	-	-
1965	20	13	7	-	-	-	-	-	-	-	-	-	-
1964	33	25	7	1	-	-	-	-	-	-	-	-	-
1963	31	22	8	1	-	-	-	-	-	-	-	-	-
1962	24	19	4	1	-	-	-	-	-	-	-	-	-
1961	21	14	7	-	-	-	-	-	-	-	-	-	-
1960	37	29	8	-	-	-	-	-	-	-	-	-	-
ALL OTHER, FEMALE													
1967	10	9	1	-	-	-	-	-	-	-	-	-	-
1966	12	8	4	-	-	-	-	-	-	-	-	-	-
1965	12	9	3	-	-	-	-	-	-	-	-	-	-
1964	32	23	8	-	-	1	-	-	-	-	-	-	-
1963	36	24	12	-	-	-	-	-	-	-	-	-	-
1962	25	21	4	-	-	-	-	-	-	-	-	-	-
1961	29	22	6	1	-	-	-	-	-	-	-	-	-
1960	36	20	15	1	-	-	-	-	-	-	-	-	-

- Quantity zero

Note: Figures by race exclude data for residents of New Jersey for 1962 and 1963.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290A for 1960-67 and separate unpublished tables for the black population and Accidents.

77/09/02 TABLE 290A. DEATHS FOR 60 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1960-67

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	TOTAL	UNDER 1 YEAR	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS	65-74 YEARS	75-84 YEARS	85+ YEARS	N.S.
Meningococcal infections (057)													
BOTH SEXES													
1967	635	167	182	82	69	12	21	34	36	20	9	3	-
1966	876	212	287	100	99	16	28	48	46	24	13	3	-
1965	890	274	292	110	65	21	26	35	35	21	9	2	-
1964	750	204	223	100	84	24	29	31	27	19	8	1	-
1963	733	203	263	81	69	15	19	31	22	15	13	2	-
1962	649	189	229	85	47	12	15	21	26	14	7	4	-
1961	586	163	226	71	35	9	12	26	15	19	10	-	-
1960	644	196	235	74	34	11	17	33	21	16	5	2	-
MALE													
1967	355	105	107	44	35	7	12	17	14	9	4	1	-
1966	482	115	167	51	66	11	8	26	22	10	4	2	-
1965	483	156	158	65	40	10	13	16	16	5	3	1	-
1964	456	122	147	56	60	15	15	17	15	7	2	-	-
1963	396	108	139	45	52	4	12	14	12	6	4	-	-
1962	366	99	137	44	30	8	8	9	17	9	3	2	-
1961	324	88	138	34	22	2	5	10	9	9	7	-	-
1960	372	113	144	40	22	4	11	17	12	6	3	-	-
FEMALE													
1967	280	62	75	38	34	5	9	17	22	11	5	2	-
1966	394	97	120	49	33	5	20	22	24	14	9	1	-
1965	407	118	134	45	25	11	13	19	19	16	6	1	-
1964	294	82	76	44	24	9	14	14	12	12	6	1	-
1963	337	95	124	36	17	11	7	17	10	9	9	2	-
1962	283	90	92	41	17	4	7	12	9	5	4	2	-
1961	262	75	88	37	13	7	7	16	6	10	3	-	-
1960	272	83	91	34	12	7	6	16	9	10	2	2	-

- Quantity zero

Note: Figures by race exclude data for residents of New Jersey for 1962 and 1963.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290A for 1960-67 and separate unpublished tables for the black population and Accidents.

77/09/02 TABLE 290A. DEATHS FOR 60 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1960-67

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	TOTAL	UNDER 1 YEAR	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS	65-74 YEARS	75-84 YEARS	85+ YEARS	N.S.
Meningococcal infections (057)													
WHITE, BOTH SEXES													
1967	481	144	123	58	54	8	13	27	28	16	8	2	-
1966	688	180	223	68	76	9	20	38	39	20	12	3	-
1965	714	223	242	76	52	13	20	30	30	18	8	2	-
1964	583	164	181	73	66	16	18	20	19	18	7	1	-
1963	582	164	207	58	57	11	13	25	19	14	13	1	-
1962	526	154	190	65	39	8	12	16	21	12	5	4	-
1961	486	142	188	56	27	5	9	21	11	17	10	-	-
1960	507	161	188	54	24	7	14	25	15	13	4	2	-
WHITE, MALE													
1967	272	91	72	33	27	4	6	13	13	8	4	1	-
1966	374	96	133	35	46	6	4	22	19	8	3	2	-
1965	394	133	133	48	31	5	10	14	12	4	3	1	-
1964	353	98	121	40	47	10	7	10	11	7	2	-	-
1963	319	89	108	35	43	4	8	12	10	6	4	-	-
1962	294	77	114	31	26	6	6	7	15	8	2	2	-
1961	267	77	113	27	16	2	3	7	6	9	7	-	-
1960	290	95	115	28	15	3	8	12	7	5	2	-	-
WHITE, FEMALE													
1967	209	53	51	25	27	4	7	14	15	8	4	1	-
1966	314	84	90	33	30	3	16	16	20	12	9	1	-
1965	320	90	109	28	21	8	10	16	18	14	5	1	-
1964	230	66	60	33	19	6	11	10	8	11	5	1	-
1963	263	75	99	23	14	7	5	13	9	8	9	1	-
1962	232	77	76	34	13	2	6	9	6	4	3	2	-
1961	219	65	75	29	11	3	6	14	5	8	3	-	-
1960	217	66	73	26	9	4	6	13	8	8	2	2	-

- Quantity zero

Note: Figures by race exclude data for residents of New Jersey for 1962 and 1963.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290A for 1960-67 and separate unpublished tables for the black population and Accidents.

77/09/02 TABLE 290A. DEATHS FOR 60 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1960-67

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	TOTAL	UNDER 1 YEAR	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS	65-74 YEARS	75-84 YEARS	85+ YEARS	N.S.
Meningococcal infections (057)													
ALL OTHER, BOTH SEXES													
1967	154	23	59	24	15	4	8	7	8	4	1	1	-
1966	188	32	64	32	23	7	8	10	7	4	1	-	-
1965	176	51	50	34	13	8	6	5	5	3	1	-	-
1964	167	40	42	27	18	8	11	11	8	1	1	-	-
1963	136	36	49	21	12	3	4	6	3	1	-	1	-
1962	93	26	31	16	7	4	1	4	2	2	-	-	-
1961	100	21	38	15	8	4	3	5	4	2	-	-	-
1960	137	35	47	20	10	4	3	8	6	3	1	-	-
ALL OTHER, MALE													
1967	83	14	35	11	8	3	6	4	1	1	-	-	-
1966	108	19	34	16	20	5	4	4	3	2	1	-	-
1965	89	23	25	17	9	5	3	2	4	1	-	-	-
1964	103	24	26	16	13	5	8	7	4	-	-	-	-
1963	67	17	25	10	9	-	2	2	2	-	-	-	-
1962	57	17	19	11	4	2	-	1	2	1	-	-	-
1961	57	11	25	7	6	-	2	3	3	-	-	-	-
1960	82	18	29	12	7	1	3	5	5	1	1	-	-
ALL OTHER, FEMALE													
1967	71	9	24	13	7	1	2	3	7	3	1	1	-
1966	80	13	30	16	3	2	4	6	4	2	-	-	-
1965	87	28	25	17	4	3	3	3	1	2	1	-	-
1964	64	16	16	11	5	3	3	4	4	1	1	-	-
1963	69	19	24	11	3	3	2	4	1	1	-	1	-
1962	36	9	12	5	3	2	1	3	-	1	-	-	-
1961	43	10	13	8	2	4	1	2	1	2	-	-	-
1960	55	17	18	8	3	3	-	3	1	2	-	-	-

- Quantity zero

Note: Figures by race exclude data for residents of New Jersey for 1962 and 1963.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290A for 1960-67 and separate unpublished tables for the black population and Accidents.

77/09/02 TABLE 290A. DEATHS FOR 60 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1960-67

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	TOTAL	UNDER 1 YEAR	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS	65-74 YEARS	75-84 YEARS	85+ YEARS	N.S.
Acute poliomyelitis (080)													
BOTH SEXES													
1967	16	1	2	2	2	4	2	1	2	-	-	-	-
1966	9	2	1	2	1	-	1	2	-	-	-	-	-
1965	16	2	2	2	-	6	3	-	1	-	-	-	-
1964	17	2	1	6	1	1	5	-	1	-	-	-	-
1963	41	6	6	9	3	10	4	2	1	-	-	-	-
1962	60	4	15	16	7	8	7	1	1	1	-	-	-
1961	90	1	9	25	12	24	13	3	3	-	-	-	-
1960	230	10	34	55	43	52	23	12	-	1	-	-	-
MALE													
1967	7	-	-	1	2	3	-	-	1	-	-	-	-
1966	4	1	-	-	1	-	1	1	-	-	-	-	-
1965	7	-	1	-	-	5	1	-	-	-	-	-	-
1964	12	1	1	4	1	1	3	-	1	-	-	-	-
1963	27	5	5	3	2	6	3	2	1	-	-	-	-
1962	37	3	7	11	4	7	3	1	1	-	-	-	-
1961	50	1	4	15	6	17	4	2	1	-	-	-	-
1960	140	5	25	35	19	32	15	9	-	-	-	-	-
FEMALE													
1967	9	1	2	1	-	1	2	1	1	-	-	-	-
1966	5	1	1	2	-	-	-	1	-	-	-	-	-
1965	9	2	1	2	-	1	2	-	1	-	-	-	-
1964	5	1	-	2	-	-	2	-	-	-	-	-	-
1963	14	1	1	6	1	4	1	-	-	-	-	-	-
1962	23	1	8	5	3	1	4	-	-	1	-	-	-
1961	40	-	5	10	6	7	9	1	2	-	-	-	-
1960	90	5	9	20	24	20	8	3	-	1	-	-	-

- Quantity zero

Note: Figures by race exclude data for residents of New Jersey for 1962 and 1963.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290A for 1960-67 and separate unpublished tables for the black population and Accidents.

77/09/02 TABLE 290A. DEATHS FOR 60 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1960-67

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	TOTAL	UNDER 1 YEAR	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS	65-74 YEARS	75-84 YEARS	85+ YEARS	N.S.
Acute poliomyelitis (080)													
WHITE, BOTH SEXES													
1967	12	1	1	1	1	3	2	1	2	-	-	-	-
1966	7	2	1	1	-	-	1	2	-	-	-	-	-
1965	13	1	1	2	-	5	3	-	1	-	-	-	-
1964	17	2	1	6	1	1	5	-	1	-	-	-	-
1963	33	6	3	9	3	6	3	2	1	-	-	-	-
1962	46	4	9	12	6	7	6	1	-	1	-	-	-
1961	80	1	8	20	11	24	10	3	3	-	-	-	-
1960	194	8	25	44	39	48	20	9	-	1	-	-	-
WHITE, MALE													
1967	5	-	-	1	1	2	-	-	1	-	-	-	-
1966	3	1	-	-	-	-	1	1	-	-	-	-	-
1965	5	-	-	-	-	4	1	-	-	-	-	-	-
1964	12	1	1	4	1	1	3	-	1	-	-	-	-
1963	21	5	3	3	2	3	2	2	1	-	-	-	-
1962	29	3	4	8	4	6	3	1	-	-	-	-	-
1961	45	1	3	12	5	17	4	2	1	-	-	-	-
1960	115	4	19	25	17	31	12	7	-	-	-	-	-
WHITE, FEMALE													
1967	7	1	1	-	-	1	2	1	1	-	-	-	-
1966	4	1	1	1	-	-	-	1	-	-	-	-	-
1965	8	1	1	2	-	1	2	-	1	-	-	-	-
1964	5	1	-	2	-	-	2	-	-	-	-	-	-
1963	12	1	-	6	1	3	1	-	-	-	-	-	-
1962	17	1	5	4	2	1	3	-	-	1	-	-	-
1961	35	-	5	8	6	7	6	1	2	-	-	-	-
1960	79	4	6	19	22	17	8	2	-	1	-	-	-

- Quantity zero

Note: Figures by race exclude data for residents of New Jersey for 1962 and 1963.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290A for 1960-67 and separate unpublished tables for the black population and Accidents.

77/09/02 TABLE 290A. DEATHS FOR 60 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1960-67

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	TOTAL	UNDER 1 YEAR	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS	65-74 YEARS	75-84 YEARS	85+ YEARS	N.S.
Acute poliomyelitis (080)													
ALL OTHER, BOTH SEXES													
1967	4	-	1	1	1	1	-	-	-	-	-	-	-
1966	2	-	-	1	1	-	-	-	-	-	-	-	-
1965	3	1	1	-	-	1	-	-	-	-	-	-	-
1964	-	-	-	-	-	-	-	-	-	-	-	-	-
1963	7	-	3	-	-	4	-	-	-	-	-	-	-
1962	12	-	5	4	1	-	1	-	1	-	-	-	-
1961	10	-	1	5	1	-	3	-	-	-	-	-	-
1960	36	2	9	11	4	4	3	3	-	-	-	-	-
ALL OTHER, MALE													
1967	2	-	-	-	1	1	-	-	-	-	-	-	-
1966	1	-	-	-	1	-	-	-	-	-	-	-	-
1965	2	-	1	-	-	1	-	-	-	-	-	-	-
1964	-	-	-	-	-	-	-	-	-	-	-	-	-
1963	5	-	2	-	-	3	-	-	-	-	-	-	-
1962	7	-	3	3	-	-	-	-	1	-	-	-	-
1961	5	-	1	3	1	-	-	-	-	-	-	-	-
1960	25	1	6	10	2	1	3	2	-	-	-	-	-
ALL OTHER, FEMALE													
1967	2	-	1	1	-	-	-	-	-	-	-	-	-
1966	1	-	-	1	-	-	-	-	-	-	-	-	-
1965	1	1	-	-	-	-	-	-	-	-	-	-	-
1964	-	-	-	-	-	-	-	-	-	-	-	-	-
1963	2	-	1	-	-	1	-	-	-	-	-	-	-
1962	5	-	2	1	1	-	1	-	-	-	-	-	-
1961	5	-	-	2	-	-	3	-	-	-	-	-	-
1960	11	1	3	1	2	3	-	1	-	-	-	-	-

- Quantity zero

Note: Figures by race exclude data for residents of New Jersey for 1962 and 1963.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290A for 1960-67 and separate unpublished tables for the black population and Accidents.

77/09/02 TABLE 290A. DEATHS FOR 60 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1960-67

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	TOTAL	UNDER 1 YEAR	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS	65-74 YEARS	75-84 YEARS	85+ YEARS	N.S.
Measles (085)													
BOTH SEXES													
1967	81	16	36	21	1	2	-	1	1	1	1	1	-
1966	261	54	134	60	6	4	1	1	-	1	-	-	-
1965	276	68	126	73	4	1	1	1	1	-	-	1	-
1964	421	88	201	112	10	1	2	1	1	2	3	-	-
1963	364	78	164	103	9	3	3	2	-	1	1	-	-
1962	408	93	204	87	8	5	2	1	2	3	1	2	-
1961	434	70	234	104	10	8	3	1	2	1	1	-	-
1960	380	85	199	80	4	8	2	-	-	-	2	-	-
MALE													
1967	41	10	22	5	-	1	-	1	-	1	1	-	-
1966	146	31	78	29	5	2	-	-	-	1	-	-	-
1965	129	36	66	24	1	-	1	1	-	-	-	-	-
1964	195	46	94	50	3	-	-	-	-	2	-	-	-
1963	171	34	84	47	2	1	2	1	-	-	-	-	-
1962	180	37	97	37	3	1	-	1	1	1	1	1	-
1961	212	35	116	49	5	2	2	1	1	1	-	-	-
1960	198	46	104	41	2	3	1	-	-	-	1	-	-
FEMALE													
1967	40	6	14	16	1	1	-	-	1	-	-	1	-
1966	115	23	56	31	1	2	1	1	-	-	-	-	-
1965	147	32	60	49	3	1	-	-	1	-	-	1	-
1964	226	42	107	62	7	1	2	1	1	-	3	-	-
1963	193	44	80	56	7	2	1	1	-	1	1	-	-
1962	228	56	107	50	5	4	2	-	1	2	-	1	-
1961	222	35	118	55	5	6	1	-	1	-	1	-	-
1960	182	39	95	39	2	5	1	-	-	-	1	-	-

- Quantity zero

Note: Figures by race exclude data for residents of New Jersey for 1962 and 1963.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290A for 1960-67 and separate unpublished tables for the black population and Accidents.

77/09/02 TABLE 290A. DEATHS FOR 60 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1960-67

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	TOTAL	UNDER 1 YEAR	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS	65-74 YEARS	75-84 YEARS	85+ YEARS	N.S.
Measles (085)													
WHITE, BOTH SEXES													
1967	56	12	24	12	1	2	-	1	1	1	1	1	-
1966	184	36	86	50	5	4	1	1	-	1	-	-	-
1965	192	43	82	58	4	1	1	1	1	-	-	1	-
1964	321	56	153	94	10	-	2	-	1	2	3	-	-
1963	269	47	117	90	7	2	3	1	-	1	1	-	-
1962	279	59	129	70	6	5	2	1	2	2	1	2	-
1961	325	43	167	91	9	7	3	1	2	1	1	-	-
1960	280	59	137	69	4	7	2	-	-	-	2	-	-
WHITE, MALE													
1967	30	8	16	2	-	1	-	1	-	1	1	-	-
1966	106	21	51	26	5	2	-	-	-	1	-	-	-
1965	91	25	46	17	1	-	1	1	-	-	-	-	-
1964	146	28	71	42	3	-	-	-	-	2	-	-	-
1963	127	24	57	40	2	1	2	1	-	-	-	-	-
1962	126	23	65	29	3	1	-	1	1	1	1	1	-
1961	156	23	80	41	5	2	2	1	1	1	-	-	-
1960	150	28	76	39	2	3	1	-	-	-	1	-	-
WHITE, FEMALE													
1967	26	4	8	10	1	1	-	-	1	-	-	1	-
1966	78	15	35	24	-	2	1	1	-	-	-	-	-
1965	101	18	36	41	3	1	-	-	1	-	-	1	-
1964	175	28	82	52	7	-	2	-	1	-	3	-	-
1963	142	23	60	50	5	1	1	-	-	1	1	-	-
1962	153	36	64	41	3	4	2	-	1	1	-	1	-
1961	169	20	87	50	4	5	1	-	1	-	1	-	-
1960	130	31	61	30	2	4	1	-	-	-	1	-	-

- Quantity zero

Note: Figures by race exclude data for residents of New Jersey for 1962 and 1963.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290A for 1960-67 and separate unpublished tables for the black population and Accidents.

77/09/02 TABLE 290A. DEATHS FOR 60 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1960-67

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	TOTAL	UNDER 1 YEAR	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS	65-74 YEARS	75-84 YEARS	85+ YEARS	N.S.
Measles (085)													
ALL OTHER, BOTH SEXES													
1967	25	4	12	9	-	-	-	-	-	-	-	-	-
1966	77	18	48	10	1	-	-	-	-	-	-	-	-
1965	84	25	44	15	-	-	-	-	-	-	-	-	-
1964	100	32	48	18	-	1	-	1	-	-	-	-	-
1963	90	30	45	11	2	1	-	1	-	-	-	-	-
1962	120	33	71	13	2	-	-	-	-	1	-	-	-
1961	109	27	67	13	1	1	-	-	-	-	-	-	-
1960	100	26	62	11	-	1	-	-	-	-	-	-	-
ALL OTHER, MALE													
1967	11	2	6	3	-	-	-	-	-	-	-	-	-
1966	40	10	27	3	-	-	-	-	-	-	-	-	-
1965	38	11	20	7	-	-	-	-	-	-	-	-	-
1964	49	18	23	8	-	-	-	-	-	-	-	-	-
1963	42	9	27	6	-	-	-	-	-	-	-	-	-
1962	51	14	32	5	-	-	-	-	-	-	-	-	-
1961	56	12	36	8	-	-	-	-	-	-	-	-	-
1960	48	18	28	2	-	-	-	-	-	-	-	-	-
ALL OTHER, FEMALE													
1967	14	2	6	6	-	-	-	-	-	-	-	-	-
1966	37	8	21	7	1	-	-	-	-	-	-	-	-
1965	46	14	24	8	-	-	-	-	-	-	-	-	-
1964	51	14	25	10	-	1	-	1	-	-	-	-	-
1963	48	21	18	5	2	1	-	1	-	-	-	-	-
1962	69	19	39	8	2	-	-	-	-	1	-	-	-
1961	53	15	31	5	1	1	-	-	-	-	-	-	-
1960	52	8	34	9	-	1	-	-	-	-	-	-	-

- Quantity zero

Note: Figures by race exclude data for residents of New Jersey for 1962 and 1963.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290A for 1960-67 and separate unpublished tables for the black population and Accidents.

77/09/02 TABLE 290A. DEATHS FOR 60 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1960-67

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	TOTAL	UNDER 1 YEAR	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS	65-74 YEARS	75-84 YEARS	85+ YEARS	N.S.
Other infective and parasitic diseases (030-044,049,052-054,058-074,081-084,086-138)													
BOTH SEXES													
1967	5,959	648	368	338	296	300	401	672	852	971	795	315	3
1966	6,044	692	406	347	287	297	465	618	832	960	810	328	2
1965	5,904	742	428	338	264	286	462	601	821	904	782	275	1
1964	5,889	807	451	375	245	288	430	624	777	926	715	246	5
1963	6,062	790	442	357	275	304	461	654	848	964	701	263	3
1962	5,791	834	434	378	283	275	477	626	746	835	639	264	-
1961	5,606	853	456	334	256	287	428	591	747	806	644	204	-
1960	5,672	861	488	334	258	311	441	597	757	839	571	213	2
MALE													
1967	3,090	341	201	181	141	132	193	341	500	541	378	138	3
1966	3,204	396	229	162	136	139	216	331	480	549	426	139	1
1965	3,113	426	239	179	123	132	209	331	454	514	415	91	-
1964	3,176	463	241	200	128	129	215	320	437	549	373	117	4
1963	3,293	449	238	191	151	148	221	360	491	565	360	119	-
1962	3,013	455	216	184	128	126	217	334	414	483	329	127	-
1961	3,024	510	245	174	129	128	210	309	440	465	335	79	-
1960	3,065	526	263	168	121	119	226	339	420	494	301	87	1
FEMALE													
1967	2,869	307	167	157	155	168	208	331	352	430	417	177	-
1966	2,840	296	177	185	151	158	249	287	352	411	384	189	1
1965	2,791	316	189	159	141	154	253	270	367	390	367	184	1
1964	2,713	344	210	175	117	159	215	304	340	377	342	129	1
1963	2,769	341	204	166	124	156	240	294	357	399	341	144	3
1962	2,778	379	218	194	155	149	260	292	332	352	310	137	-
1961	2,582	343	211	160	127	159	218	282	307	341	309	125	-
1960	2,607	335	225	166	137	192	215	258	337	345	270	126	1

- Quantity zero

Note: Figures by race exclude data for residents of New Jersey for 1962 and 1963.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290A for 1960-67 and separate unpublished tables for the black population and Accidents.

77/09/02 TABLE 290A. DEATHS FOR 60 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1960-67

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	TOTAL	UNDER 1 YEAR	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS	65-74 YEARS	75-84 YEARS	85+ YEARS	N.S.
Other infective and parasitic diseases (030-044,049,052-054,058-074,081-084,086-138)													
WHITE, BOTH SEXES													
1967	4,688	432	271	289	232	194	267	505	676	815	717	289	1
1966	4,750	467	324	291	212	200	327	445	657	791	730	306	-
1965	4,620	496	312	273	188	193	321	457	662	777	690	250	1
1964	4,702	554	347	327	199	188	311	462	609	815	661	226	3
1963	4,696	530	337	280	198	205	321	491	659	816	618	239	2
1962	4,440	540	315	291	209	194	334	472	577	705	559	244	-
1961	4,466	577	328	277	196	209	312	475	624	693	589	186	-
1960	4,499	575	363	277	202	225	313	455	623	734	528	203	1
WHITE, MALE													
1967	2,428	241	144	148	112	86	137	254	395	455	333	122	1
1966	2,506	264	177	136	97	92	155	253	382	444	381	125	-
1965	2,444	285	173	148	87	91	151	252	376	441	363	77	-
1964	2,544	317	182	176	104	80	161	239	348	486	342	107	2
1963	2,568	306	182	156	116	108	149	274	377	479	315	106	-
1962	2,303	300	154	142	98	88	155	251	317	399	281	118	-
1961	2,420	351	180	143	98	104	157	249	376	390	303	69	-
1960	2,421	348	200	139	98	84	161	257	337	432	282	82	1
WHITE, FEMALE													
1967	2,260	191	127	141	120	108	130	251	281	360	384	167	-
1966	2,244	203	147	155	115	108	172	192	275	347	349	181	-
1965	2,176	211	139	125	101	102	170	205	286	336	327	173	1
1964	2,158	237	165	151	95	108	150	223	261	329	319	119	1
1963	2,128	224	155	124	82	97	172	217	282	337	303	133	2
1962	2,137	240	161	149	111	106	179	221	260	306	278	126	-
1961	2,046	226	148	134	98	105	155	226	248	303	286	117	-
1960	2,078	227	163	138	104	141	152	198	286	302	246	121	-

- Quantity zero

Note: Figures by race exclude data for residents of New Jersey for 1962 and 1963.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290A for 1960-67 and separate unpublished tables for the black population and Accidents.

77/09/02 TABLE 290A. DEATHS FOR 60 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1960-67

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	TOTAL	UNDER 1 YEAR	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS	65-74 YEARS	75-84 YEARS	85+ YEARS	N.S.
Other infective and parasitic diseases (030-044,049,052-054,058-074,081-084,086-138)													
ALL OTHER, BOTH SEXES													
1967	1,271	216	97	49	64	106	134	167	176	156	78	26	2
1966	1,294	225	82	56	75	97	138	173	175	169	80	22	2
1965	1,284	246	116	65	76	93	141	144	159	127	92	25	-
1964	1,187	253	104	48	46	100	119	162	168	111	54	20	2
1963	1,207	250	97	72	65	89	119	152	154	128	62	18	1
1962	1,191	276	111	71	64	71	133	134	148	103	64	16	-
1961	1,140	276	128	57	60	78	116	116	123	113	55	18	-
1960	1,173	286	125	57	56	86	128	142	134	105	43	10	1
ALL OTHER, MALE													
1967	662	100	57	33	29	46	56	87	105	86	45	16	2
1966	698	132	52	26	39	47	61	78	98	105	45	14	1
1965	669	141	66	31	36	41	58	79	78	73	52	14	-
1964	632	146	59	24	24	49	54	81	89	63	31	10	2
1963	636	136	52	32	30	34	57	81	94	76	34	10	-
1962	628	148	58	33	26	31	61	72	86	69	36	8	-
1961	604	159	65	31	31	24	53	60	64	75	32	10	-
1960	644	178	63	29	23	35	65	82	83	62	19	5	-
ALL OTHER, FEMALE													
1967	609	116	40	16	35	60	78	80	71	70	33	10	-
1966	596	93	30	30	36	50	77	95	77	64	35	8	1
1965	615	105	50	34	40	52	83	65	81	54	40	11	-
1964	555	107	45	24	22	51	65	81	79	48	23	10	-
1963	571	114	45	40	35	55	62	71	60	52	28	8	1
1962	563	128	53	38	38	40	72	62	62	34	28	8	-
1961	536	117	63	26	29	54	63	56	59	38	23	8	-
1960	529	108	62	28	33	51	63	60	51	43	24	5	1

- Quantity zero

Note: Figures by race exclude data for residents of New Jersey for 1962 and 1963.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290A for 1960-67 and separate unpublished tables for the black population and Accidents.

77/09/02 TABLE 290A. DEATHS FOR 60 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1960-67

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	TOTAL	UNDER 1 YEAR	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS	65-74 YEARS	75-84 YEARS	85+ YEARS	N.S.
ICD-7 Nos. 140-205 Malignant neoplasms													
BOTH SEXES													
1967	310,983	196	1,283	2,711	2,663	4,002	14,384	40,527	72,464	89,786	65,518	17,409	40
1966	303,736	205	1,342	2,587	2,628	4,092	14,722	40,154	70,202	87,640	63,330	16,789	45
1965	297,588	226	1,431	2,565	2,485	4,200	14,780	39,229	68,978	86,409	61,195	16,053	37
1964	289,577	233	1,525	2,539	2,423	4,297	14,910	38,740	66,930	84,105	58,613	15,212	50
1963	285,362	260	1,672	2,616	2,334	4,165	14,582	37,919	65,521	83,922	57,030	15,283	58
1962	278,562	261	1,610	2,597	2,218	4,176	14,409	37,449	63,769	82,058	55,245	14,717	53
1961	273,502	293	1,705	2,617	2,071	4,359	14,421	36,660	62,247	80,765	53,887	14,405	72
1960	267,627	298	1,762	2,415	2,000	4,441	14,374	36,264	61,795	78,504	52,239	13,474	61
MALE													
1967	169,164	106	691	1,546	1,604	1,941	6,071	19,572	41,321	52,266	35,854	8,175	17
1966	164,975	98	729	1,499	1,574	1,941	6,267	19,472	40,063	50,973	34,539	7,793	27
1965	161,422	118	769	1,450	1,488	1,948	6,220	18,832	39,313	50,343	33,314	7,606	21
1964	156,913	128	879	1,435	1,406	2,046	6,235	18,565	38,121	49,035	31,916	7,122	25
1963	154,288	155	948	1,454	1,371	1,979	5,947	18,278	36,965	48,972	30,929	7,253	37
1962	150,009	124	879	1,426	1,335	1,930	5,837	17,944	36,009	47,543	29,922	7,030	30
1961	147,157	146	938	1,490	1,249	2,036	5,752	17,422	34,984	46,873	29,360	6,866	41
1960	143,543	161	1,020	1,370	1,214	2,099	5,747	17,243	34,659	45,561	28,129	6,308	32
FEMALE													
1967	141,819	90	592	1,165	1,059	2,061	8,313	20,955	31,143	37,520	29,664	9,234	23
1966	138,761	107	613	1,088	1,054	2,151	8,455	20,682	30,139	36,667	28,791	8,996	18
1965	136,166	108	662	1,115	997	2,252	8,560	20,397	29,665	36,066	27,881	8,447	16
1964	132,664	105	646	1,104	1,017	2,251	8,675	20,175	28,809	35,070	26,697	8,090	25
1963	131,074	105	724	1,162	963	2,186	8,635	19,641	28,556	34,950	26,101	8,030	21
1962	128,553	137	731	1,171	883	2,246	8,572	19,505	27,760	34,515	25,323	7,687	23
1961	126,345	147	767	1,127	822	2,323	8,669	19,238	27,263	33,892	24,527	7,539	31
1960	124,084	137	742	1,045	786	2,342	8,627	19,021	27,136	32,943	24,110	7,166	29

Note: Figures by race exclude data for residents of New Jersey for 1962 and 1963.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290A for 1960-67 and separate unpublished tables for the black population and Accidents.

77/09/02 TABLE 290A. DEATHS FOR 60 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1960-67

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	TOTAL	UNDER 1 YEAR	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS	65-74 YEARS	75-84 YEARS	85+ YEARS	N.S.
ICD-7 Nos. 140-205 Malignant neoplasms													
WHITE, BOTH SEXES													
1967	279,502	165	1,151	2,406	2,367	3,375	12,152	34,914	64,008	81,208	61,415	16,316	25
1966	273,169	181	1,199	2,298	2,332	3,444	12,357	34,602	62,155	79,432	59,367	15,774	28
1965	268,542	197	1,299	2,300	2,188	3,584	12,484	34,028	61,102	78,646	57,585	15,095	34
1964	261,390	196	1,340	2,230	2,162	3,700	12,610	33,457	59,257	76,862	55,160	14,379	37
1963	246,823	219	1,423	2,250	2,028	3,399	11,706	31,269	55,437	73,327	51,761	13,971	33
1962	241,499	229	1,393	2,203	1,909	3,447	11,707	31,021	54,095	71,815	50,192	13,445	43
1961	247,697	264	1,545	2,373	1,838	3,764	12,190	31,756	55,183	74,180	50,935	13,630	39
1960	242,712	258	1,584	2,186	1,767	3,791	12,216	31,415	54,988	72,232	49,434	12,801	40
WHITE, MALE													
1967	151,451	89	623	1,366	1,427	1,698	5,105	16,654	36,390	47,103	33,404	7,580	12
1966	147,901	87	658	1,348	1,424	1,697	5,262	16,580	35,388	45,988	32,237	7,217	15
1965	145,374	100	699	1,303	1,332	1,741	5,280	16,235	34,791	45,616	31,149	7,110	18
1964	141,333	108	768	1,261	1,259	1,788	5,279	15,934	33,773	44,639	29,840	6,668	16
1963	133,202	134	814	1,264	1,195	1,679	4,816	14,990	31,304	42,539	27,882	6,566	19
1962	130,050	110	776	1,222	1,164	1,665	4,794	14,771	30,665	41,431	27,033	6,396	23
1961	133,219	131	857	1,372	1,113	1,810	4,906	15,069	30,999	42,906	27,601	6,434	21
1960	130,183	141	924	1,254	1,083	1,873	4,893	14,957	30,886	41,723	26,501	5,928	20
WHITE, FEMALE													
1967	128,051	76	528	1,040	940	1,677	7,047	18,260	27,618	34,105	28,011	8,736	13
1966	125,268	94	541	950	908	1,747	7,095	18,022	26,767	33,444	27,130	8,557	13
1965	123,168	97	600	997	856	1,843	7,204	17,793	26,311	33,030	26,436	7,985	16
1964	120,057	88	572	969	903	1,912	7,331	17,523	25,484	32,223	25,320	7,711	21
1963	113,621	85	609	986	833	1,720	6,890	16,279	24,133	30,788	23,879	7,405	14
1962	111,449	119	617	981	745	1,782	6,913	16,250	23,430	30,384	23,159	7,049	20
1961	114,478	133	688	1,001	725	1,954	7,284	16,687	24,184	31,274	23,334	7,196	18
1960	112,529	117	660	932	684	1,918	7,323	16,458	24,102	30,509	22,933	6,873	20

Note: Figures by race exclude data for residents of New Jersey for 1962 and 1963.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290A for 1960-67 and separate unpublished tables for the black population and Accidents.

77/09/02 TABLE 290A. DEATHS FOR 60 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1960-67

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	TOTAL	UNDER 1 YEAR	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS	65-74 YEARS	75-84 YEARS	85+ YEARS	N.S.
ICD-7 Nos. 140-205 Malignant neoplasms													
ALL OTHER, BOTH SEXES													
1967	31,481	31	132	305	296	627	2,232	5,613	8,456	8,578	4,103	1,093	15
1966	30,567	24	143	289	296	648	2,365	5,552	8,047	8,208	3,963	1,015	17
1965	29,046	29	132	265	297	616	2,296	5,201	7,876	7,763	3,610	958	3
1964	28,187	37	185	309	261	597	2,300	5,283	7,673	7,243	3,453	833	13
1963	26,932	36	195	273	248	617	2,254	5,018	7,305	7,062	3,116	786	22
1962	25,740	29	160	313	237	588	2,157	4,832	6,950	6,713	3,023	728	10
1961	25,805	29	160	244	233	595	2,231	4,904	7,064	6,585	2,952	775	33
1960	24,915	40	178	229	233	650	2,158	4,849	6,807	6,272	2,805	673	21
ALL OTHER, MALE													
1967	17,713	17	68	180	177	243	966	2,918	4,931	5,163	2,450	595	5
1966	17,074	11	71	151	150	244	1,005	2,892	4,675	4,985	2,302	576	12
1965	16,048	18	70	147	156	207	940	2,597	4,522	4,727	2,165	496	3
1964	15,580	20	111	174	147	258	956	2,631	4,348	4,396	2,076	454	9
1963	14,875	18	104	136	140	229	889	2,527	4,100	4,396	1,883	438	15
1962	13,884	13	77	157	127	204	822	2,415	3,804	4,071	1,790	397	7
1961	13,938	15	81	118	136	226	846	2,353	3,985	3,967	1,759	432	20
1960	13,360	20	96	116	131	226	854	2,286	3,773	3,838	1,628	380	12
ALL OTHER, FEMALE													
1967	13,768	14	64	125	119	384	1,266	2,695	3,525	3,415	1,653	498	10
1966	13,493	13	72	138	146	404	1,360	2,660	3,372	3,223	1,661	439	5
1965	12,998	11	62	118	141	409	1,356	2,604	3,354	3,036	1,445	462	-
1964	12,607	17	74	135	114	339	1,344	2,652	3,325	2,847	1,377	379	4
1963	12,057	18	91	137	108	388	1,365	2,491	3,205	2,666	1,233	348	7
1962	11,856	16	83	156	110	384	1,335	2,417	3,146	2,642	1,233	331	3
1961	11,867	14	79	126	97	369	1,385	2,551	3,079	2,618	1,193	343	13
1960	11,555	20	82	113	102	424	1,304	2,563	3,034	2,434	1,177	293	9

- Quantity zero

Note: Figures by race exclude data for residents of New Jersey for 1962 and 1963.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290A for 1960-67 and separate unpublished tables for the black population and Accidents.

77/09/02 TABLE 290A. DEATHS FOR 60 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1960-67

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	TOTAL	UNDER 1 YEAR	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS	65-74 YEARS	75-84 YEARS	85+ YEARS	N.S.
Malignant neoplasm of buccal cavity and pharynx (140-148)													
BOTH SEXES													
1967	6,718	1	1	15	30	46	251	1,214	2,011	1,737	1,057	354	1
1966	6,800	-	7	21	35	34	288	1,190	1,935	1,748	1,162	378	2
1965	6,501	-	4	17	33	41	276	1,118	1,898	1,719	1,053	338	4
1964	6,571	1	5	15	21	51	266	1,122	1,861	1,710	1,125	391	3
1963	6,508	-	8	17	29	41	275	1,162	1,806	1,725	1,084	358	3
1962	6,481	1	9	19	34	43	287	1,079	1,694	1,766	1,150	396	3
1961	6,322	1	5	15	28	57	251	1,064	1,638	1,766	1,135	361	1
1960	6,175	-	3	18	28	55	266	1,093	1,541	1,719	1,129	323	-
MALE													
1967	5,013	-	1	10	20	29	172	903	1,596	1,318	749	214	1
1966	5,047	-	5	12	21	24	190	842	1,545	1,364	824	219	1
1965	4,924	-	1	10	21	25	195	833	1,523	1,346	754	212	4
1964	4,977	1	3	11	11	32	184	840	1,454	1,367	834	237	3
1963	4,935	-	4	10	19	29	177	882	1,462	1,343	784	222	3
1962	4,920	-	5	10	21	25	187	851	1,346	1,392	826	256	1
1961	4,823	-	1	9	20	41	168	798	1,336	1,389	827	233	1
1960	4,668	-	1	13	18	32	175	801	1,232	1,356	820	220	-
FEMALE													
1967	1,705	1	-	5	10	17	79	311	415	419	308	140	-
1966	1,753	-	2	9	14	10	98	348	390	384	338	159	1
1965	1,577	-	3	7	12	16	81	285	375	373	299	126	-
1964	1,594	-	2	4	10	19	82	282	407	343	291	154	-
1963	1,573	-	4	7	10	12	98	280	344	382	300	136	-
1962	1,561	1	4	9	13	18	100	228	348	374	324	140	2
1961	1,499	1	4	6	8	16	83	266	302	377	308	128	-
1960	1,507	-	2	5	10	23	91	292	309	363	309	103	-

- Quantity zero

Note: Figures by race exclude data for residents of New Jersey for 1962 and 1963.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290A for 1960-67 and separate unpublished tables for the black population and Accidents.

77/09/02 TABLE 290A. DEATHS FOR 60 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1960-67

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	TOTAL	UNDER 1 YEAR	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS	65-74 YEARS	75-84 YEARS	85+ YEARS	N.S.
Malignant neoplasm of buccal cavity and pharynx (140-148)													
WHITE, BOTH SEXES													
1967	5,915	1	1	13	18	31	164	974	1,797	1,568	1,004	343	1
1966	6,080	-	4	17	23	28	194	1,010	1,737	1,606	1,097	362	2
1965	5,802	-	4	16	24	35	183	945	1,672	1,590	1,010	319	4
1964	5,849	-	5	13	17	39	204	911	1,673	1,553	1,060	371	3
1963	5,643	-	7	13	22	30	208	955	1,552	1,532	998	325	1
1962	5,603	1	8	16	26	32	209	879	1,445	1,570	1,046	368	3
1961	5,755	-	5	15	24	50	197	908	1,468	1,643	1,095	349	1
1960	5,629	-	3	17	24	39	206	937	1,404	1,610	1,080	309	-
WHITE, MALE													
1967	4,409	-	1	8	10	20	111	713	1,431	1,195	711	208	1
1966	4,514	-	3	9	15	18	135	708	1,386	1,251	779	209	1
1965	4,400	-	1	9	16	21	132	702	1,343	1,246	727	199	4
1964	4,422	-	3	9	10	23	146	667	1,306	1,247	787	221	3
1963	4,273	-	3	7	13	19	136	725	1,255	1,197	716	201	1
1962	4,253	-	4	9	16	22	133	692	1,147	1,237	752	240	1
1961	4,388	-	1	9	19	35	132	682	1,199	1,289	797	224	1
1960	4,274	-	1	13	15	22	131	683	1,131	1,277	791	210	-
WHITE, FEMALE													
1967	1,506	1	-	5	8	11	53	261	366	373	293	135	-
1966	1,566	-	1	8	8	10	59	302	351	355	318	153	1
1965	1,402	-	3	7	8	14	51	243	329	344	283	120	-
1964	1,427	-	2	4	7	16	58	244	367	306	273	150	-
1963	1,370	-	4	6	9	11	72	230	297	335	282	124	-
1962	1,350	1	4	7	10	10	76	187	298	333	294	128	2
1961	1,367	-	4	6	5	15	65	226	269	354	298	125	-
1960	1,355	-	2	4	9	17	75	254	273	333	289	99	-

- Quantity zero

Note: Figures by race exclude data for residents of New Jersey for 1962 and 1963.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290A for 1960-67 and separate unpublished tables for the black population and Accidents.

77/09/02 TABLE 290A. DEATHS FOR 60 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1960-67

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	TOTAL	UNDER 1 YEAR	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS	65-74 YEARS	75-84 YEARS	85+ YEARS	N.S.
Malignant neoplasm of buccal cavity and pharynx (140-148)													
ALL OTHER, BOTH SEXES													
1967	803	-	-	2	12	15	87	240	214	169	53	11	-
1966	720	-	3	4	12	6	94	180	198	142	65	16	-
1965	699	-	-	1	9	6	93	173	226	129	43	19	-
1964	722	1	-	2	4	12	62	211	188	157	65	20	-
1963	625	-	1	4	6	10	60	165	192	128	45	13	1
1962	600	-	-	3	6	9	65	155	172	121	56	13	-
1961	567	1	-	-	4	7	54	156	170	123	40	12	-
1960	546	-	-	1	4	16	60	156	137	109	49	14	-
ALL OTHER, MALE													
1967	604	-	-	2	10	9	61	190	165	123	38	6	-
1966	533	-	2	3	6	6	55	134	159	113	45	10	-
1965	524	-	-	1	5	4	63	131	180	100	27	13	-
1964	555	1	-	2	1	9	38	173	148	120	47	16	-
1963	473	-	1	3	5	9	40	122	156	94	34	8	1
1962	455	-	-	1	4	3	46	124	138	92	40	7	-
1961	435	-	-	-	1	6	36	116	137	100	30	9	-
1960	394	-	-	-	3	10	44	118	101	79	29	10	-
ALL OTHER, FEMALE													
1967	199	-	-	-	2	6	26	50	49	46	15	5	-
1966	187	-	1	1	6	-	39	46	39	29	20	6	-
1965	175	-	-	-	4	2	30	42	46	29	16	6	-
1964	167	-	-	-	3	3	24	38	40	37	18	4	-
1963	152	-	-	1	1	1	20	43	36	34	11	5	-
1962	145	-	-	2	2	6	19	31	34	29	16	6	-
1961	132	1	-	-	3	1	18	40	33	23	10	3	-
1960	152	-	-	1	1	6	16	38	36	30	20	4	-

- Quantity zero

Note: Figures by race exclude data for residents of New Jersey for 1962 and 1963.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290A for 1960-67 and separate unpublished tables for the black population and Accidents.

77/09/02 TABLE 290A. DEATHS FOR 60 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1960-67

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	TOTAL	UNDER 1 YEAR	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS	65-74 YEARS	75-84 YEARS	85+ YEARS	N.S.
Malignant neoplasm of digestive organs and peritoneum, not specified as secondary (150-156A,157-159)													
BOTH SEXES													
1967	95,320	16	29	48	134	553	2,566	9,337	20,312	30,053	25,126	7,130	16
1966	95,079	14	39	52	149	569	2,732	9,502	20,063	30,115	24,886	6,945	13
1965	94,305	16	31	34	137	533	2,682	9,217	20,323	30,082	24,515	6,719	16
1964	93,158	30	32	35	158	580	2,816	9,390	19,980	30,059	23,679	6,382	17
1963	92,914	16	39	42	162	533	2,757	9,204	19,745	30,258	23,532	6,609	17
1962	92,047	21	50	52	127	550	2,814	9,380	19,678	30,135	22,872	6,352	16
1961	91,727	23	47	30	123	582	2,892	9,346	19,502	30,050	22,885	6,223	24
1960	91,035	29	45	40	109	604	2,796	9,411	19,800	29,953	22,362	5,861	25
MALE													
1967	51,262	9	17	28	77	306	1,388	5,284	12,052	16,699	12,431	2,965	6
1966	51,186	6	23	35	91	309	1,511	5,443	11,698	16,729	12,534	2,798	9
1965	51,253	6	21	23	73	302	1,452	5,223	12,034	16,860	12,383	2,867	9
1964	50,694	14	21	23	95	307	1,552	5,294	11,824	16,953	11,956	2,647	8
1963	50,361	8	24	20	87	283	1,462	5,248	11,519	17,117	11,806	2,775	12
1962	49,936	13	24	30	76	297	1,507	5,284	11,505	16,887	11,577	2,726	10
1961	50,173	12	22	11	73	322	1,480	5,306	11,349	17,178	11,786	2,621	13
1960	49,792	8	26	28	65	323	1,464	5,366	11,585	17,135	11,328	2,451	13
FEMALE													
1967	44,058	7	12	20	57	247	1,178	4,053	8,260	13,354	12,695	4,165	10
1966	43,893	8	16	17	58	260	1,221	4,059	8,365	13,386	12,352	4,147	4
1965	43,052	10	10	11	64	231	1,230	3,994	8,289	13,222	12,132	3,852	7
1964	42,464	16	11	12	63	273	1,264	4,096	8,156	13,106	11,723	3,735	9
1963	42,553	8	15	22	75	250	1,295	3,956	8,226	13,141	11,726	3,834	5
1962	42,111	8	26	22	51	253	1,307	4,096	8,173	13,248	11,295	3,626	6
1961	41,554	11	25	19	50	260	1,412	4,040	8,153	12,872	11,099	3,602	11
1960	41,243	21	19	12	44	281	1,332	4,045	8,215	12,818	11,034	3,410	12

Note: Figures by race exclude data for residents of New Jersey for 1962 and 1963.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290A for 1960-67 and separate unpublished tables for the black population and Accidents.

77/09/02 TABLE 290A. DEATHS FOR 60 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1960-67

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	TOTAL	UNDER 1 YEAR	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS	65-74 YEARS	75-84 YEARS	85+ YEARS	N.S.
Malignant neoplasm of digestive organs and peritoneum, not specified as secondary (150-156A,157-159)													
WHITE, BOTH SEXES													
1967	85,370	13	29	39	107	431	2,105	7,769	17,560	27,043	23,566	6,698	10
1966	84,981	12	32	41	118	437	2,198	7,868	17,317	27,115	23,290	6,545	8
1965	84,653	12	30	29	105	428	2,214	7,699	17,595	27,122	23,077	6,327	15
1964	83,731	29	25	24	127	467	2,255	7,823	17,336	27,272	22,306	6,056	11
1963	79,908	13	36	33	128	423	2,143	7,298	16,305	26,185	21,300	6,037	7
1962	79,438	17	42	41	92	432	2,178	7,476	16,362	26,245	20,737	5,804	12
1961	82,881	18	43	28	96	464	2,335	7,840	16,996	27,467	21,679	5,906	9
1960	82,463	26	37	32	81	499	2,296	7,937	17,307	27,439	21,193	5,598	18
WHITE, MALE													
1967	45,368	7	17	20	58	240	1,112	4,326	10,347	14,921	11,565	2,750	5
1966	45,145	5	20	29	71	234	1,206	4,427	9,984	14,909	11,677	2,579	4
1965	45,474	5	21	21	56	249	1,189	4,311	10,326	15,053	11,561	2,674	8
1964	45,065	13	17	16	76	247	1,239	4,360	10,218	15,240	11,151	2,484	4
1963	42,786	7	24	17	68	223	1,123	4,112	9,436	14,669	10,600	2,502	5
1962	42,709	11	21	23	55	249	1,150	4,149	9,564	14,621	10,384	2,476	6
1961	44,887	10	21	11	58	249	1,186	4,405	9,796	15,590	11,103	2,454	4
1960	44,669	8	23	22	47	268	1,168	4,489	10,084	15,547	10,688	2,316	9
WHITE, FEMALE													
1967	40,002	6	12	19	49	191	993	3,443	7,213	12,122	12,001	3,948	5
1966	39,836	7	12	12	47	203	992	3,441	7,333	12,206	11,613	3,966	4
1965	39,179	7	9	8	49	179	1,025	3,388	7,269	12,069	11,516	3,653	7
1964	38,666	16	8	8	51	220	1,016	3,463	7,118	12,032	11,155	3,572	7
1963	37,122	6	12	16	60	200	1,020	3,186	6,869	11,516	10,700	3,535	2
1962	36,729	6	21	18	37	183	1,028	3,327	6,798	11,624	10,353	3,328	6
1961	37,994	8	22	17	38	215	1,149	3,435	7,200	11,877	10,576	3,452	5
1960	37,794	18	14	10	34	231	1,128	3,448	7,223	11,892	10,505	3,282	9

Note: Figures by race exclude data for residents of New Jersey for 1962 and 1963.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290A for 1960-67 and separate unpublished tables for the black population and Accidents.

77/09/02 TABLE 290A. DEATHS FOR 60 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1960-67

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	TOTAL	UNDER 1 YEAR	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS	65-74 YEARS	75-84 YEARS	85+ YEARS	N.S.
Malignant neoplasm of digestive organs and peritoneum, not specified as secondary (150-156A,157-159)													
ALL OTHER, BOTH SEXES													
1967	9,950	3	-	9	27	122	461	1,568	2,752	3,010	1,560	432	6
1966	10,098	2	7	11	31	132	534	1,634	2,746	3,000	1,596	400	5
1965	9,652	4	1	5	32	105	468	1,518	2,728	2,960	1,438	392	1
1964	9,427	1	7	11	31	113	561	1,567	2,644	2,787	1,373	326	6
1963	8,896	3	1	6	33	93	500	1,492	2,522	2,649	1,253	334	10
1962	8,545	3	7	8	30	97	515	1,457	2,409	2,505	1,224	286	4
1961	8,846	5	4	2	27	118	557	1,506	2,506	2,583	1,206	317	15
1960	8,572	3	8	8	28	105	500	1,474	2,493	2,514	1,169	263	7
ALL OTHER, MALE													
1967	5,894	2	-	8	19	66	276	958	1,705	1,778	866	215	1
1966	6,041	1	3	6	20	75	305	1,016	1,714	1,820	857	219	5
1965	5,779	1	-	2	17	53	263	912	1,708	1,807	822	193	1
1964	5,629	1	4	7	19	60	313	934	1,606	1,713	805	163	4
1963	5,337	1	-	2	19	50	269	904	1,535	1,652	718	180	7
1962	5,026	2	2	5	19	39	293	887	1,406	1,503	720	146	4
1961	5,286	2	1	-	15	73	294	901	1,553	1,588	683	167	9
1960	5,123	-	3	6	18	55	296	877	1,501	1,588	640	135	4
ALL OTHER, FEMALE													
1967	4,056	1	-	1	8	56	185	610	1,047	1,232	694	217	5
1966	4,057	1	4	5	11	57	229	618	1,032	1,180	739	181	-
1965	3,873	3	1	3	15	52	205	606	1,020	1,153	616	199	-
1964	3,798	-	3	4	12	53	248	633	1,038	1,074	568	163	2
1963	3,559	2	1	4	14	43	231	588	987	997	535	154	3
1962	3,519	1	5	3	11	58	222	570	1,003	1,002	504	140	-
1961	3,560	3	3	2	12	45	263	605	953	995	523	150	6
1960	3,449	3	5	2	10	50	204	597	992	926	529	128	3

- Quantity zero

Note: Figures by race exclude data for residents of New Jersey for 1962 and 1963.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290A for 1960-67 and separate unpublished tables for the black population and Accidents.

77/09/02 TABLE 290A. DEATHS FOR 60 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1960-67

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	TOTAL	UNDER 1 YEAR	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS	65-74 YEARS	75-84 YEARS	85+ YEARS	N.S.
Malignant neoplasm of respiratory system, not specified as secondary (160-164)													
BOTH SEXES													
1967	58,086	5	9	15	42	231	2,377	9,170	17,734	18,605	8,694	1,201	3
1966	54,934	4	2	10	49	264	2,344	8,644	16,996	17,660	7,830	1,122	9
1965	52,039	7	5	12	64	231	2,256	8,375	15,967	16,984	7,086	1,051	1
1964	49,187	3	16	21	41	245	2,189	7,810	15,453	15,843	6,650	908	8
1963	46,973	10	5	12	44	276	2,093	7,545	14,551	15,402	6,109	917	9
1962	44,644	2	5	25	45	257	1,965	7,276	14,068	14,487	5,679	827	8
1961	42,211	8	12	13	45	264	1,868	6,803	13,325	13,834	5,191	836	12
1960	39,730	7	9	16	31	261	1,756	6,562	12,691	12,887	4,767	735	8
MALE													
1967	48,411	3	6	6	27	165	1,757	7,159	15,150	16,159	7,145	833	1
1966	46,070	-	2	3	35	194	1,713	6,902	14,688	15,420	6,381	723	9
1965	43,798	5	3	6	41	159	1,674	6,699	13,862	14,833	5,806	709	1
1964	41,687	3	6	11	29	166	1,622	6,386	13,547	13,928	5,366	616	7
1963	39,817	7	3	9	28	197	1,537	6,159	12,760	13,533	4,958	618	8
1962	38,050	2	3	12	28	193	1,481	6,052	12,399	12,769	4,553	551	7
1961	35,954	3	5	6	33	198	1,411	5,674	11,787	12,150	4,116	562	9
1960	34,047	3	3	7	25	188	1,335	5,517	11,322	11,344	3,816	479	8
FEMALE													
1967	9,675	2	3	9	15	66	620	2,011	2,584	2,446	1,549	368	2
1966	8,864	4	-	7	14	70	631	1,742	2,308	2,240	1,449	399	-
1965	8,241	2	2	6	23	72	582	1,676	2,105	2,151	1,280	342	-
1964	7,500	-	10	10	12	79	567	1,424	1,906	1,915	1,284	292	1
1963	7,156	3	2	3	16	79	556	1,386	1,791	1,869	1,151	299	1
1962	6,594	-	2	13	17	64	484	1,224	1,669	1,718	1,126	276	1
1961	6,257	5	7	7	12	66	457	1,129	1,538	1,684	1,075	274	3
1960	5,683	4	6	9	6	73	421	1,045	1,369	1,543	951	256	-

- Quantity zero

Note: Figures by race exclude data for residents of New Jersey for 1962 and 1963.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290A for 1960-67 and separate unpublished tables for the black population and Accidents.

77/09/02 TABLE 290A. DEATHS FOR 60 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1960-67

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	TOTAL	UNDER 1 YEAR	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS	65-74 YEARS	75-84 YEARS	85+ YEARS	N.S.
Malignant neoplasm of respiratory system, not specified as secondary (160-164)													
WHITE, BOTH SEXES													
1967	52,427	4	8	12	37	185	1,938	7,908	15,895	17,142	8,192	1,104	2
1966	49,534	4	1	7	36	205	1,883	7,411	15,296	16,228	7,407	1,049	7
1965	47,287	7	2	8	54	197	1,863	7,318	14,470	15,669	6,714	984	1
1964	44,772	2	15	16	36	206	1,807	6,815	13,947	14,755	6,319	848	6
1963	40,765	6	5	9	38	213	1,631	6,260	12,591	13,655	5,522	829	6
1962	38,977	2	4	17	38	201	1,596	6,116	12,211	12,874	5,153	758	7
1961	38,562	7	11	13	40	219	1,578	5,962	12,077	12,908	4,946	792	9
1960	36,182	6	7	12	29	220	1,473	5,744	11,490	12,026	4,489	681	5
WHITE, MALE													
1967	43,633	3	6	5	25	132	1,413	6,105	13,552	14,901	6,731	760	-
1966	41,516	-	1	2	26	150	1,355	5,884	13,203	14,188	6,028	672	7
1965	39,793	5	2	3	34	138	1,369	5,830	12,574	13,685	5,493	659	1
1964	37,954	2	5	8	25	137	1,320	5,548	12,248	12,981	5,100	575	5
1963	34,562	4	3	7	24	152	1,190	5,074	11,064	12,015	4,470	554	5
1962	33,193	2	3	9	24	150	1,196	5,070	10,790	11,325	4,119	499	6
1961	32,838	3	5	6	30	167	1,176	4,966	10,701	11,346	3,905	527	6
1960	31,012	2	2	6	23	157	1,099	4,828	10,264	10,585	3,598	443	5
WHITE, FEMALE													
1967	8,794	1	2	7	12	53	525	1,803	2,343	2,241	1,461	344	2
1966	8,018	4	-	5	10	55	528	1,527	2,093	2,040	1,379	377	-
1965	7,494	2	-	5	20	59	494	1,488	1,896	1,984	1,221	325	-
1964	6,818	-	10	8	11	69	487	1,267	1,699	1,774	1,219	273	1
1963	6,203	2	2	2	14	61	441	1,186	1,527	1,640	1,052	275	1
1962	5,784	-	1	8	14	51	400	1,046	1,421	1,549	1,034	259	1
1961	5,724	4	6	7	10	52	402	996	1,376	1,562	1,041	265	3
1960	5,170	4	5	6	6	63	374	916	1,226	1,441	891	238	-

- Quantity zero

Note: Figures by race exclude data for residents of New Jersey for 1962 and 1963.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290A for 1960-67 and separate unpublished tables for the black population and Accidents.

77/09/02 TABLE 290A. DEATHS FOR 60 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1960-67

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	TOTAL	UNDER 1 YEAR	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS	65-74 YEARS	75-84 YEARS	85+ YEARS	N.S.
Malignant neoplasm of respiratory system, not specified as secondary (160-164)													
ALL OTHER, BOTH SEXES													
1967	5,659	1	1	3	5	46	439	1,262	1,839	1,463	502	97	1
1966	5,400	-	1	3	13	59	461	1,233	1,700	1,432	423	73	2
1965	4,752	-	3	4	10	34	393	1,057	1,497	1,315	372	67	-
1964	4,415	1	1	5	5	39	382	995	1,506	1,088	331	60	2
1963	4,268	3	-	2	5	51	386	968	1,341	1,108	348	53	3
1962	3,756	-	1	8	6	45	300	857	1,248	955	293	42	1
1961	3,649	1	1	-	5	45	290	841	1,248	926	245	44	3
1960	3,548	1	2	4	2	41	283	818	1,201	861	278	54	3
ALL OTHER, MALE													
1967	4,778	-	-	1	2	33	344	1,054	1,598	1,258	414	73	1
1966	4,554	-	1	1	9	44	358	1,018	1,485	1,232	353	51	2
1965	4,005	-	1	3	7	21	305	869	1,288	1,148	313	50	-
1964	3,733	1	1	3	4	29	302	838	1,299	947	266	41	2
1963	3,599	2	-	1	4	35	291	826	1,161	954	284	38	3
1962	3,191	-	-	3	3	35	230	724	1,073	850	240	32	1
1961	3,116	-	-	-	3	31	235	708	1,086	804	211	35	3
1960	3,035	1	1	1	2	31	236	689	1,058	759	218	36	3
ALL OTHER, FEMALE													
1967	881	1	1	2	3	13	95	208	241	205	88	24	-
1966	846	-	-	2	4	15	103	215	215	200	70	22	-
1965	747	-	2	1	3	13	88	188	209	167	59	17	-
1964	682	-	-	2	1	10	80	157	207	141	65	19	-
1963	669	1	-	1	1	16	95	142	180	154	64	15	-
1962	565	-	1	5	3	10	70	133	175	105	53	10	-
1961	533	1	1	-	2	14	55	133	162	122	34	9	-
1960	513	-	1	3	-	10	47	129	143	102	60	18	-

- Quantity zero

Note: Figures by race exclude data for residents of New Jersey for 1962 and 1963.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290A for 1960-67 and separate unpublished tables for the black population and Accidents.

77/09/02 TABLE 290A. DEATHS FOR 60 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1960-67

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	TOTAL	UNDER 1 YEAR	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS	65-74 YEARS	75-84 YEARS	85+ YEARS	N.S.
Malignant neoplasm of breast (170)													
BOTH SEXES													
1967	28,217	-	1	-	22	437	2,495	6,159	7,279	6,187	4,239	1,394	4
1966	27,533	-	-	-	25	428	2,525	6,011	7,079	6,068	4,107	1,284	6
1965	27,048	-	-	1	16	437	2,590	5,897	6,762	5,997	4,091	1,253	4
1964	26,162	-	-	-	29	485	2,538	5,916	6,361	5,838	3,749	1,241	5
1963	25,401	2	-	2	21	407	2,518	5,626	6,275	5,799	3,581	1,167	3
1962	24,958	2	-	2	15	461	2,438	5,558	5,991	5,699	3,636	1,150	6
1961	24,542	1	-	1	12	458	2,520	5,408	5,773	5,560	3,633	1,168	8
1960	23,970	5	-	1	16	447	2,501	5,359	5,767	5,351	3,424	1,097	2
MALE													
1967	232	-	-	-	-	-	8	29	68	70	44	13	-
1966	229	-	-	-	-	4	10	29	52	66	57	11	-
1965	243	-	-	-	-	2	11	40	53	76	48	12	1
1964	226	-	-	-	-	1	7	35	53	69	50	11	-
1963	262	-	-	-	1	4	13	34	59	83	49	18	1
1962	225	-	-	-	-	1	8	38	57	69	46	6	-
1961	231	-	-	-	-	3	10	36	51	74	42	14	1
1960	215	2	-	-	1	-	12	21	75	60	35	9	-
FEMALE													
1967	27,985	-	1	-	22	437	2,487	6,130	7,211	6,117	4,195	1,381	4
1966	27,304	-	-	-	25	424	2,515	5,982	7,027	6,002	4,050	1,273	6
1965	26,805	-	-	1	16	435	2,579	5,857	6,709	5,921	4,043	1,241	3
1964	25,936	-	-	-	29	484	2,531	5,881	6,308	5,769	3,699	1,230	5
1963	25,139	2	-	2	20	403	2,505	5,592	6,216	5,716	3,532	1,149	2
1962	24,733	2	-	2	15	460	2,430	5,520	5,934	5,630	3,590	1,144	6
1961	24,311	1	-	1	12	455	2,510	5,372	5,722	5,486	3,591	1,154	7
1960	23,755	3	-	1	15	447	2,489	5,338	5,692	5,291	3,389	1,088	2

- Quantity zero

Note: Figures by race exclude data for residents of New Jersey for 1962 and 1963.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290A for 1960-67 and separate unpublished tables for the black population and Accidents.

77/09/02 TABLE 290A. DEATHS FOR 60 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1960-67

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	TOTAL	UNDER 1 YEAR	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS	65-74 YEARS	75-84 YEARS	85+ YEARS	N.S.
Malignant neoplasm of breast (170)													
WHITE, BOTH SEXES													
1967	25,798	-	1	-	18	354	2,184	5,492	6,633	5,750	4,038	1,326	2
1966	25,308	-	-	-	18	342	2,218	5,414	6,493	5,670	3,923	1,227	3
1965	24,848	-	-	1	13	351	2,242	5,313	6,206	5,633	3,902	1,183	4
1964	24,028	-	-	-	26	405	2,232	5,327	5,813	5,466	3,571	1,185	3
1963	22,206	1	-	-	18	316	2,048	4,833	5,449	5,202	3,269	1,067	3
1962	21,921	-	-	1	10	369	2,007	4,805	5,247	5,110	3,328	1,039	5
1961	22,592	1	-	1	9	392	2,168	4,901	5,303	5,232	3,461	1,118	6
1960	22,053	2	-	1	11	363	2,178	4,812	5,323	5,028	3,274	1,060	1
WHITE, MALE													
1967	209	-	-	-	-	-	8	24	59	64	42	12	-
1966	197	-	-	-	-	4	9	23	43	58	51	9	-
1965	221	-	-	-	-	2	8	36	47	73	43	11	1
1964	195	-	-	-	-	1	6	31	41	60	46	10	-
1963	221	-	-	-	1	4	11	28	47	71	42	16	1
1962	189	-	-	-	-	1	8	30	47	57	41	5	-
1961	212	-	-	-	-	3	10	32	47	68	38	13	1
1960	186	1	-	-	1	-	8	19	64	54	31	8	-
WHITE, FEMALE													
1967	25,589	-	1	-	18	354	2,176	5,468	6,574	5,686	3,996	1,314	2
1966	25,111	-	-	-	18	338	2,209	5,391	6,450	5,612	3,872	1,218	3
1965	24,627	-	-	1	13	349	2,234	5,277	6,159	5,560	3,859	1,172	3
1964	23,833	-	-	-	26	404	2,226	5,296	5,772	5,406	3,525	1,175	3
1963	21,985	1	-	-	17	312	2,037	4,805	5,402	5,131	3,227	1,051	2
1962	21,732	-	-	1	10	368	1,999	4,775	5,200	5,053	3,287	1,034	5
1961	22,380	1	-	1	9	389	2,158	4,869	5,256	5,164	3,423	1,105	5
1960	21,867	1	-	1	10	363	2,170	4,793	5,259	4,974	3,243	1,052	1

- Quantity zero

Note: Figures by race exclude data for residents of New Jersey for 1962 and 1963.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290A for 1960-67 and separate unpublished tables for the black population and Accidents.

77/09/02 TABLE 290A. DEATHS FOR 60 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1960-67

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	TOTAL	UNDER 1 YEAR	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS	65-74 YEARS	75-84 YEARS	85+ YEARS	N.S.
Malignant neoplasm of breast (170)													
ALL OTHER, BOTH SEXES													
1967	2,419	-	-	-	4	83	311	667	646	437	201	68	2
1966	2,225	-	-	-	7	86	307	597	586	398	184	57	3
1965	2,200	-	-	-	3	86	348	584	556	364	189	70	-
1964	2,134	-	-	-	3	80	306	589	548	372	178	56	2
1963	2,047	1	-	2	2	77	343	539	518	348	164	53	-
1962	1,917	2	-	1	3	73	322	499	458	341	165	52	1
1961	1,950	-	-	-	3	66	352	507	470	328	172	50	2
1960	1,917	3	-	-	5	84	323	547	444	323	150	37	1
ALL OTHER, MALE													
1967	23	-	-	-	-	-	-	5	9	6	2	1	-
1966	32	-	-	-	-	-	1	6	9	8	6	2	-
1965	22	-	-	-	-	-	3	4	6	3	5	1	-
1964	31	-	-	-	-	-	1	4	12	9	4	1	-
1963	27	-	-	-	-	-	2	5	8	7	4	1	-
1962	22	-	-	-	-	-	-	5	7	8	2	-	-
1961	19	-	-	-	-	-	-	4	4	6	4	1	-
1960	29	1	-	-	-	-	4	2	11	6	4	1	-
ALL OTHER, FEMALE													
1967	2,396	-	-	-	4	83	311	662	637	431	199	67	2
1966	2,193	-	-	-	7	86	306	591	577	390	178	55	3
1965	2,178	-	-	-	3	86	345	580	550	361	184	69	-
1964	2,103	-	-	-	3	80	305	585	536	363	174	55	2
1963	2,020	1	-	2	2	77	341	534	510	341	160	52	-
1962	1,895	2	-	1	3	73	322	494	451	333	163	52	1
1961	1,931	-	-	-	3	66	352	503	466	322	168	49	2
1960	1,888	2	-	-	5	84	319	545	433	317	146	36	1

- Quantity zero

Note: Figures by race exclude data for residents of New Jersey for 1962 and 1963.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290A for 1960-67 and separate unpublished tables for the black population and Accidents.

77/09/02 TABLE 290A. DEATHS FOR 60 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1960-67

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	TOTAL	UNDER 1 YEAR	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS	65-74 YEARS	75-84 YEARS	85+ YEARS	N.S.
Malignant neoplasm of genital organs (171-179)													
BOTH SEXES													
1967	40,408	-	22	41	282	632	2,092	4,618	7,675	11,555	10,395	3,089	7
1966	40,378	5	26	40	265	708	2,198	4,841	7,516	11,502	10,273	2,994	10
1965	40,454	1	19	48	256	775	2,331	4,955	7,691	11,431	10,050	2,893	4
1964	39,950	2	23	46	238	716	2,365	4,959	7,652	11,378	9,834	2,730	7
1963	39,983	2	24	57	242	750	2,444	4,917	7,737	11,435	9,622	2,746	7
1962	39,590	1	20	47	209	773	2,497	4,990	7,696	11,266	9,409	2,676	6
1961	39,256	2	29	42	185	742	2,513	5,151	7,595	11,235	9,126	2,628	8
1960	38,669	2	20	35	207	795	2,583	5,064	7,638	10,980	8,895	2,441	9
MALE													
1967	17,292	-	15	8	155	200	146	374	1,901	5,481	6,822	2,188	2
1966	16,954	2	15	6	149	195	189	398	1,891	5,403	6,627	2,074	5
1965	16,880	-	12	13	137	191	209	418	1,935	5,344	6,577	2,043	1
1964	16,552	1	16	9	117	199	197	408	1,859	5,455	6,363	1,925	3
1963	16,409	-	15	11	128	181	195	386	1,824	5,509	6,216	1,941	3
1962	16,111	-	12	9	112	190	175	387	1,830	5,357	6,144	1,891	4
1961	15,911	-	20	11	101	192	204	389	1,806	5,261	6,023	1,900	4
1960	15,391	1	15	7	103	188	193	405	1,769	5,223	5,788	1,696	3
FEMALE													
1967	23,116	-	7	33	127	432	1,946	4,244	5,774	6,074	3,573	901	5
1966	23,424	3	11	34	116	513	2,009	4,443	5,625	6,099	3,646	920	5
1965	23,574	1	7	35	119	584	2,122	4,537	5,756	6,087	3,473	850	3
1964	23,398	1	7	37	121	517	2,168	4,551	5,793	5,923	3,471	805	4
1963	23,574	2	9	46	114	569	2,249	4,531	5,913	5,926	3,406	805	4
1962	23,479	1	8	38	97	583	2,322	4,603	5,866	5,909	3,265	785	2
1961	23,345	2	9	31	84	550	2,309	4,762	5,789	5,974	3,103	728	4
1960	23,278	1	5	28	104	607	2,390	4,659	5,869	5,757	3,107	745	6

- Quantity zero

Note: Figures by race exclude data for residents of New Jersey for 1962 and 1963.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290A for 1960-67 and separate unpublished tables for the black population and Accidents.

77/09/02 TABLE 290A. DEATHS FOR 60 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1960-67

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	TOTAL	UNDER 1 YEAR	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS	65-74 YEARS	75-84 YEARS	85+ YEARS	N.S.
Malignant neoplasm of genital organs (171-179)													
WHITE, BOTH SEXES													
1967	34,861	-	21	32	249	506	1,669	3,865	6,453	9,828	9,435	2,800	3
1966	34,917	5	23	35	233	564	1,749	4,043	6,284	9,914	9,344	2,719	4
1965	35,080	-	19	36	230	611	1,877	4,145	6,447	9,890	9,163	2,658	4
1964	34,584	2	21	39	207	597	1,901	4,073	6,411	9,892	8,951	2,486	4
1963	33,379	2	21	48	212	566	1,854	3,918	6,166	9,557	8,571	2,460	4
1962	33,001	1	19	39	186	605	1,926	3,981	6,153	9,347	8,348	2,391	5
1961	34,007	2	29	36	160	593	2,030	4,237	6,315	9,835	8,353	2,413	4
1960	33,649	1	16	31	176	616	2,072	4,186	6,433	9,655	8,212	2,247	4
WHITE, MALE													
1967	14,882	-	15	7	146	191	138	289	1,490	4,493	6,130	1,982	1
1966	14,753	2	13	6	148	189	176	304	1,491	4,554	5,988	1,882	-
1965	14,720	-	12	10	134	185	196	321	1,523	4,498	5,957	1,883	1
1964	14,381	1	14	9	113	193	175	298	1,454	4,632	5,739	1,753	-
1963	13,836	-	15	11	122	171	172	287	1,363	4,449	5,511	1,733	2
1962	13,505	-	11	9	105	180	151	282	1,328	4,295	5,450	1,691	3
1961	13,909	-	20	10	95	185	185	290	1,377	4,510	5,482	1,753	2
1960	13,491	1	11	7	97	184	166	306	1,391	4,468	5,299	1,560	1
WHITE, FEMALE													
1967	19,979	-	6	25	103	315	1,531	3,576	4,963	5,335	3,305	818	2
1966	20,164	3	10	29	85	375	1,573	3,739	4,793	5,360	3,356	837	4
1965	20,360	-	7	26	96	426	1,681	3,824	4,924	5,392	3,206	775	3
1964	20,203	1	7	30	94	404	1,726	3,775	4,957	5,260	3,212	733	4
1963	19,543	2	6	37	90	395	1,682	3,631	4,803	5,108	3,060	727	2
1962	19,496	1	8	30	81	425	1,775	3,699	4,825	5,052	2,898	700	2
1961	20,098	2	9	26	65	408	1,845	3,947	4,938	5,325	2,871	660	2
1960	20,158	-	5	24	79	432	1,906	3,880	5,042	5,187	2,913	687	3

- Quantity zero

Note: Figures by race exclude data for residents of New Jersey for 1962 and 1963.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290A for 1960-67 and separate unpublished tables for the black population and Accidents.

77/09/02 TABLE 290A. DEATHS FOR 60 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1960-67

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	TOTAL	UNDER 1 YEAR	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS	65-74 YEARS	75-84 YEARS	85+ YEARS	N.S.
Malignant neoplasm of genital organs (171-179)													
ALL OTHER, BOTH SEXES													
1967	5,547	-	1	9	33	126	423	753	1,222	1,727	960	289	4
1966	5,461	-	3	5	32	144	449	798	1,232	1,588	929	275	6
1965	5,374	1	-	12	26	164	454	810	1,244	1,541	887	235	-
1964	5,366	-	2	7	31	119	464	886	1,241	1,486	883	244	3
1963	5,162	-	2	6	20	160	476	800	1,278	1,467	742	208	3
1962	5,236	-	1	6	16	149	496	812	1,268	1,506	762	219	1
1961	5,249	-	-	6	25	149	483	914	1,280	1,400	773	215	4
1960	5,020	1	4	4	31	179	511	878	1,205	1,325	683	194	5
ALL OTHER, MALE													
1967	2,410	-	-	1	9	9	8	85	411	988	692	206	1
1966	2,201	-	2	-	1	6	13	94	400	849	639	192	5
1965	2,160	-	-	3	3	6	13	97	412	846	620	160	-
1964	2,171	-	2	-	4	6	22	110	405	823	624	172	3
1963	2,045	-	-	-	2	4	18	86	400	869	517	148	1
1962	2,113	-	1	-	3	6	20	86	439	892	513	152	1
1961	2,002	-	-	1	6	7	19	99	429	751	541	147	2
1960	1,900	-	4	-	6	4	27	99	378	755	489	136	2
ALL OTHER, FEMALE													
1967	3,137	-	1	8	24	117	415	668	811	739	268	83	3
1966	3,260	-	1	5	31	138	436	704	832	739	290	83	1
1965	3,214	1	-	9	23	158	441	713	832	695	267	75	-
1964	3,195	-	-	7	27	113	442	776	836	663	259	72	-
1963	3,117	-	2	6	18	156	458	714	878	598	225	60	2
1962	3,123	-	-	6	13	143	476	726	829	614	249	67	-
1961	3,247	-	-	5	19	142	464	815	851	649	232	68	2
1960	3,120	1	-	4	25	175	484	779	827	570	194	58	3

- Quantity zero

Note: Figures by race exclude data for residents of New Jersey for 1962 and 1963.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290A for 1960-67 and separate unpublished tables for the black population and Accidents.

77/09/02 TABLE 290A. DEATHS FOR 60 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1960-67

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	TOTAL	UNDER 1 YEAR	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS	65-74 YEARS	75-84 YEARS	85+ YEARS	N.S.
Malignant neoplasm of urinary organs (180,181)													
BOTH SEXES													
1967	14,656	13	99	104	27	51	325	1,297	2,987	4,730	3,915	1,105	3
1966	14,166	21	92	87	34	60	352	1,319	2,871	4,480	3,795	1,055	-
1965	13,901	16	119	106	34	57	351	1,266	2,918	4,484	3,574	975	1
1964	13,515	9	122	111	35	62	352	1,305	2,799	4,321	3,457	938	4
1963	13,583	16	117	104	25	60	308	1,334	2,789	4,452	3,414	959	5
1962	13,299	21	127	129	19	62	340	1,279	2,678	4,481	3,220	941	2
1961	13,040	16	161	112	21	74	353	1,241	2,710	4,274	3,175	898	5
1960	12,804	27	159	100	23	56	338	1,199	2,702	4,197	3,183	812	8
MALE													
1967	9,712	5	41	38	15	30	214	920	2,145	3,254	2,473	574	3
1966	9,469	8	41	40	20	33	229	912	2,065	3,079	2,426	616	-
1965	9,094	11	62	48	10	30	227	855	2,091	3,093	2,175	492	-
1964	8,992	5	69	48	17	44	208	905	1,962	3,036	2,184	513	1
1963	9,122	9	70	51	10	35	198	918	2,022	3,113	2,187	508	1
1962	8,818	12	60	57	13	33	214	899	1,898	3,108	2,014	508	2
1961	8,678	8	65	50	13	39	232	866	1,956	2,989	1,984	471	5
1960	8,585	14	84	50	12	35	234	838	1,953	2,928	1,997	436	4
FEMALE													
1967	4,944	8	58	66	12	21	111	377	842	1,476	1,442	531	-
1966	4,697	13	51	47	14	27	123	407	806	1,401	1,369	439	-
1965	4,807	5	57	58	24	27	124	411	827	1,391	1,399	483	1
1964	4,523	4	53	63	18	18	144	400	837	1,285	1,273	425	3
1963	4,461	7	47	53	15	25	110	416	767	1,339	1,227	451	4
1962	4,481	9	67	72	6	29	126	380	780	1,373	1,206	433	-
1961	4,362	8	96	62	8	35	121	375	754	1,285	1,191	427	-
1960	4,219	13	75	50	11	21	104	361	749	1,269	1,186	376	4

- Quantity zero

Note: Figures by race exclude data for residents of New Jersey for 1962 and 1963.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290A for 1960-67 and separate unpublished tables for the black population and Accidents.

77/09/02 TABLE 290A. DEATHS FOR 60 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1960-67

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	TOTAL	UNDER 1 YEAR	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS	65-74 YEARS	75-84 YEARS	85+ YEARS	N.S.
Malignant neoplasm of urinary organs (180,181)													
WHITE, BOTH SEXES													
1967	13,427	11	76	84	22	45	287	1,116	2,650	4,339	3,739	1,055	3
1966	13,077	19	80	81	26	52	309	1,156	2,585	4,149	3,615	1,005	-
1965	12,832	13	97	93	23	40	309	1,115	2,618	4,171	3,408	944	1
1964	12,485	8	102	88	27	50	294	1,133	2,496	4,050	3,323	910	4
1963	11,998	14	94	84	21	52	239	1,096	2,397	3,980	3,130	888	3
1962	11,704	20	106	100	13	48	266	1,050	2,279	3,983	2,971	866	2
1961	12,071	15	137	99	16	57	283	1,070	2,456	3,998	3,069	868	3
1960	11,851	24	137	86	20	48	280	1,015	2,439	3,939	3,076	781	6
WHITE, MALE													
1967	8,975	5	31	29	11	25	194	812	1,928	3,019	2,366	552	3
1966	8,799	8	34	38	16	30	203	809	1,881	2,869	2,327	584	-
1965	8,453	9	50	42	8	21	199	769	1,896	2,900	2,082	477	-
1964	8,365	4	54	39	13	36	180	788	1,777	2,860	2,112	501	1
1963	8,096	9	55	43	6	32	153	766	1,760	2,800	2,003	469	-
1962	7,843	11	54	45	9	26	172	751	1,656	2,782	1,864	471	2
1961	8,074	8	51	42	12	32	187	766	1,790	2,810	1,920	453	3
1960	8,026	13	74	45	10	31	204	730	1,794	2,767	1,935	420	3
WHITE, FEMALE													
1967	4,452	6	45	55	11	20	93	304	722	1,320	1,373	503	-
1966	4,278	11	46	43	10	22	106	347	704	1,280	1,288	421	-
1965	4,379	4	47	51	15	19	110	346	722	1,271	1,326	467	1
1964	4,120	4	48	49	14	14	114	345	719	1,190	1,211	409	3
1963	3,902	5	39	41	15	20	86	330	637	1,180	1,127	419	3
1962	3,861	9	52	55	4	22	94	299	623	1,201	1,107	395	-
1961	3,997	7	86	57	4	25	96	304	666	1,188	1,149	415	-
1960	3,825	11	63	41	10	17	76	285	645	1,172	1,141	361	3

- Quantity zero

Note: Figures by race exclude data for residents of New Jersey for 1962 and 1963.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290A for 1960-67 and separate unpublished tables for the black population and Accidents.

77/09/02 TABLE 290A. DEATHS FOR 60 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1960-67

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	TOTAL	UNDER 1 YEAR	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS	65-74 YEARS	75-84 YEARS	85+ YEARS	N.S.
Malignant neoplasm of urinary organs (180,181)													
ALL OTHER, BOTH SEXES													
1967	1,229	2	23	20	5	6	38	181	337	391	176	50	-
1966	1,089	2	12	6	8	8	43	163	286	331	180	50	-
1965	1,069	3	22	13	11	17	42	151	300	313	166	31	-
1964	1,030	1	20	23	8	12	58	172	303	271	134	28	-
1963	968	2	14	15	4	7	52	179	271	266	126	30	2
1962	1,013	1	20	25	6	10	60	195	267	283	119	27	-
1961	969	1	24	13	5	17	70	171	254	276	106	30	2
1960	953	3	22	14	3	8	58	184	263	258	107	31	2
ALL OTHER, MALE													
1967	737	-	10	9	4	5	20	108	217	235	107	22	-
1966	670	-	7	2	4	3	26	103	184	210	99	32	-
1965	641	2	12	6	2	9	28	86	195	193	93	15	-
1964	627	1	15	9	4	8	28	117	185	176	72	12	-
1963	605	-	10	5	4	3	35	114	165	175	76	17	1
1962	586	1	6	9	4	5	31	123	147	179	64	17	-
1961	604	-	14	8	1	7	45	100	166	179	64	18	2
1960	559	1	10	5	2	4	30	108	159	161	62	16	1
ALL OTHER, FEMALE													
1967	492	2	13	11	1	1	18	73	120	156	69	28	-
1966	419	2	5	4	4	5	17	60	102	121	81	18	-
1965	428	1	10	7	9	8	14	65	105	120	73	16	-
1964	403	-	5	14	4	4	30	55	118	95	62	16	-
1963	363	2	4	10	-	4	17	65	106	91	50	13	1
1962	427	-	14	16	2	5	29	72	120	104	55	10	-
1961	365	1	10	5	4	10	25	71	88	97	42	12	-
1960	394	2	12	9	1	4	28	76	104	97	45	15	1

- Quantity zero

Note: Figures by race exclude data for residents of New Jersey for 1962 and 1963.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290A for 1960-67 and separate unpublished tables for the black population and Accidents.

77/09/02 TABLE 290A. DEATHS FOR 60 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1960-67

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	TOTAL	UNDER 1 YEAR	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS	65-74 YEARS	75-84 YEARS	85+ YEARS	N.S.
Malignant neoplasm of other and unspecified sites (156B,165,190-199)													
BOTH SEXES													
1967	36,761	71	440	886	855	942	2,352	5,232	8,643	9,126	6,376	1,834	4
1966	35,032	75	449	889	879	877	2,313	5,078	8,108	8,667	5,889	1,806	2
1965	34,644	78	474	902	787	972	2,369	4,983	7,911	8,609	5,769	1,787	3
1964	32,835	81	521	868	758	985	2,302	4,826	7,482	7,942	5,430	1,636	4
1963	32,086	82	516	860	737	922	2,233	4,808	7,277	7,908	5,160	1,576	7
1962	30,791	91	495	836	750	938	2,195	4,629	6,898	7,432	5,003	1,518	6
1961	29,977	98	505	874	665	1,000	2,145	4,447	6,615	7,382	4,753	1,487	6
1960	29,561	90	525	815	645	985	2,209	4,401	6,598	7,107	4,702	1,477	7
MALE													
1967	19,802	37	249	491	503	533	1,262	2,899	4,969	4,999	3,122	736	2
1966	19,010	46	228	480	523	487	1,262	2,846	4,786	4,750	2,862	740	-
1965	18,797	39	240	492	491	545	1,281	2,772	4,590	4,714	2,857	774	2
1964	17,738	48	296	467	452	585	1,251	2,718	4,313	4,291	2,628	687	2
1963	17,332	49	276	475	429	534	1,222	2,738	4,176	4,264	2,495	669	5
1962	16,560	40	277	431	453	509	1,161	2,491	4,026	4,051	2,454	664	3
1961	15,948	51	283	457	388	550	1,128	2,470	3,705	3,943	2,329	641	3
1960	15,895	57	296	421	392	566	1,200	2,403	3,774	3,836	2,313	634	3
FEMALE													
1967	16,959	34	191	395	352	409	1,090	2,333	3,674	4,127	3,254	1,098	2
1966	16,022	29	221	409	356	390	1,051	2,232	3,322	3,917	3,027	1,066	2
1965	15,847	39	234	410	296	427	1,088	2,211	3,321	3,895	2,912	1,013	1
1964	15,097	33	225	401	306	400	1,051	2,108	3,169	3,651	2,802	949	2
1963	14,754	33	240	385	308	388	1,011	2,070	3,101	3,644	2,665	907	2
1962	14,231	51	218	405	297	429	1,034	2,138	2,872	3,381	2,549	854	3
1961	14,029	47	222	417	277	450	1,017	1,977	2,910	3,439	2,424	846	3
1960	13,666	33	229	394	253	419	1,009	1,998	2,824	3,271	2,389	843	4

- Quantity zero

Note: Figures by race exclude data for residents of New Jersey for 1962 and 1963.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290A for 1960-67 and separate unpublished tables for the black population and Accidents.

77/09/02 TABLE 290A. DEATHS FOR 60 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1960-67

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	TOTAL	UNDER 1 YEAR	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS	65-74 YEARS	75-84 YEARS	85+ YEARS	N.S.
Malignant neoplasm of other and unspecified sites (156B,165,190-199)													
WHITE, BOTH SEXES													
1967	33,285	59	387	772	778	843	2,087	4,656	7,715	8,259	5,983	1,743	3
1966	31,615	66	399	764	798	786	2,037	4,455	7,261	7,841	5,502	1,705	1
1965	31,474	72	427	798	703	881	2,073	4,424	7,058	7,901	5,452	1,682	3
1964	29,846	65	443	765	675	889	2,037	4,284	6,727	7,280	5,106	1,571	4
1963	28,141	69	433	734	646	789	1,925	4,063	6,309	6,954	4,741	1,474	4
1962	27,071	86	424	702	662	819	1,894	3,944	5,934	6,567	4,623	1,412	4
1961	27,327	95	441	771	600	909	1,930	3,957	5,934	6,798	4,478	1,410	4
1960	26,929	77	453	725	566	886	1,983	3,920	5,911	6,525	4,458	1,421	4
WHITE, MALE													
1967	17,957	31	217	424	457	491	1,123	2,598	4,453	4,526	2,936	700	1
1966	17,191	40	204	413	474	444	1,122	2,502	4,321	4,311	2,665	695	-
1965	17,135	35	216	437	444	498	1,139	2,473	4,118	4,347	2,697	729	2
1964	16,172	37	252	414	400	523	1,121	2,441	3,908	3,948	2,469	657	2
1963	15,245	40	234	410	382	463	1,066	2,347	3,650	3,749	2,282	620	2
1962	14,657	39	244	364	406	450	1,014	2,139	3,519	3,601	2,270	609	2
1961	14,645	48	253	415	346	507	1,035	2,217	3,381	3,653	2,187	602	1
1960	14,570	47	257	383	347	512	1,093	2,175	3,429	3,525	2,199	602	1
WHITE, FEMALE													
1967	15,328	28	170	348	321	352	964	2,058	3,262	3,733	3,047	1,043	2
1966	14,424	26	195	351	324	342	915	1,953	2,940	3,530	2,837	1,010	1
1965	14,339	37	211	361	259	383	934	1,951	2,940	3,554	2,755	953	1
1964	13,674	28	191	351	275	366	916	1,843	2,819	3,332	2,637	914	2
1963	12,896	29	199	324	264	326	859	1,716	2,659	3,205	2,459	854	2
1962	12,414	47	180	338	256	369	880	1,805	2,415	2,966	2,353	803	2
1961	12,682	47	188	356	254	402	895	1,740	2,553	3,145	2,291	808	3
1960	12,359	30	196	342	219	374	890	1,745	2,482	3,000	2,259	819	3

- Quantity zero

Note: Figures by race exclude data for residents of New Jersey for 1962 and 1963.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290A for 1960-67 and separate unpublished tables for the black population and Accidents.

77/09/02 TABLE 290A. DEATHS FOR 60 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1960-67

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	TOTAL	UNDER 1 YEAR	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS	65-74 YEARS	75-84 YEARS	85+ YEARS	N.S.
Malignant neoplasm of other and unspecified sites (156B,165,190-199)													
ALL OTHER, BOTH SEXES													
1967	3,476	12	53	114	77	99	265	576	928	867	393	91	1
1966	3,417	9	50	125	81	91	276	623	847	826	387	101	1
1965	3,170	6	47	104	84	91	296	559	853	708	317	105	-
1964	2,989	16	78	103	83	96	265	542	755	662	324	65	-
1963	2,822	11	66	101	77	103	226	538	696	668	272	63	1
1962	2,686	5	49	108	74	92	223	513	707	602	249	62	2
1961	2,650	3	64	103	65	91	215	490	681	584	275	77	2
1960	2,632	13	72	90	79	99	226	481	687	582	244	56	3
ALL OTHER, MALE													
1967	1,845	6	32	67	46	42	139	301	516	473	186	36	1
1966	1,819	6	24	67	49	43	140	344	465	439	197	45	-
1965	1,662	4	24	55	47	47	142	299	472	367	160	45	-
1964	1,566	11	44	53	52	62	130	277	405	343	159	30	-
1963	1,490	8	34	50	40	58	107	289	369	360	144	30	1
1962	1,337	1	25	50	37	44	106	263	359	301	122	28	1
1961	1,303	3	30	42	42	43	93	253	324	290	142	39	2
1960	1,325	10	39	38	45	54	107	228	345	311	114	32	2
ALL OTHER, FEMALE													
1967	1,631	6	21	47	31	57	126	275	412	394	207	55	-
1966	1,598	3	26	58	32	48	136	279	382	387	190	56	1
1965	1,508	2	23	49	37	44	154	260	381	341	157	60	-
1964	1,423	5	34	50	31	34	135	265	350	319	165	35	-
1963	1,332	3	32	51	37	45	119	249	327	308	128	33	-
1962	1,349	4	24	58	37	48	117	250	348	301	127	34	1
1961	1,347	-	34	61	23	48	122	237	357	294	133	38	-
1960	1,307	3	33	52	34	45	119	253	342	271	130	24	1

- Quantity zero

Note: Figures by race exclude data for residents of New Jersey for 1962 and 1963.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290A for 1960-67 and separate unpublished tables for the black population and Accidents.

77/09/02 TABLE 290A. DEATHS FOR 60 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1960-67

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	TOTAL	UNDER 1 YEAR	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS	65-74 YEARS	75-84 YEARS	85+ YEARS	N.S.
Leukemia and aleukemia (204)													
BOTH SEXES													
1967	14,336	77	607	1,279	675	455	750	1,314	2,158	3,319	2,899	801	2
1966	14,012	69	655	1,168	650	450	781	1,281	2,216	3,189	2,778	773	2
1965	13,628	83	696	1,157	606	449	723	1,246	2,145	3,183	2,683	653	4
1964	13,453	86	709	1,149	568	479	779	1,311	2,050	3,178	2,533	609	2
1963	13,507	104	854	1,213	557	499	789	1,234	2,096	3,141	2,438	580	2
1962	12,997	93	807	1,171	549	415	729	1,202	2,030	3,078	2,353	565	5
1961	12,873	108	826	1,206	534	492	770	1,232	1,959	2,978	2,226	539	3
1960	12,725	111	894	1,149	501	523	776	1,269	2,062	2,932	2,071	437	-
MALE													
1967	8,222	46	317	744	430	265	404	741	1,309	1,924	1,636	404	2
1966	8,060	28	370	704	410	262	425	717	1,336	1,891	1,518	397	2
1965	7,840	46	378	655	357	255	416	695	1,295	1,917	1,497	326	3
1964	7,746	47	408	651	335	282	413	730	1,218	1,902	1,452	307	1
1963	7,807	65	484	647	359	284	412	694	1,274	1,877	1,383	327	1
1962	7,493	43	430	647	342	240	407	704	1,196	1,867	1,330	285	2
1961	7,569	52	467	698	334	265	436	705	1,182	1,820	1,306	301	3
1960	7,371	59	520	673	307	320	438	722	1,211	1,742	1,144	235	-
FEMALE													
1967	6,114	31	290	535	245	190	346	573	849	1,395	1,263	397	-
1966	5,952	41	285	464	240	188	356	564	880	1,298	1,260	376	-
1965	5,788	37	318	502	249	194	307	551	850	1,266	1,186	327	1
1964	5,707	39	301	498	233	197	366	581	832	1,276	1,081	302	1
1963	5,700	39	370	566	198	215	377	540	822	1,264	1,055	253	1
1962	5,504	50	377	524	207	175	322	498	834	1,211	1,023	280	3
1961	5,304	56	359	508	200	227	334	527	777	1,158	920	238	-
1960	5,354	52	374	476	194	203	338	547	851	1,190	927	202	-

- Quantity zero

Note: Figures by race exclude data for residents of New Jersey for 1962 and 1963.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290A for 1960-67 and separate unpublished tables for the black population and Accidents.

77/09/02 TABLE 290A. DEATHS FOR 60 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1960-67

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	TOTAL	UNDER 1 YEAR	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS	65-74 YEARS	75-84 YEARS	85+ YEARS	N.S.
Leukemia and aleukemia (204)													
WHITE, BOTH SEXES													
1967	13,282	65	561	1,181	594	390	679	1,170	1,986	3,101	2,775	779	1
1966	13,054	61	600	1,070	584	409	683	1,158	2,039	3,000	2,694	754	2
1965	12,677	71	641	1,066	535	392	641	1,134	1,969	3,001	2,593	632	2
1964	12,502	72	642	1,034	518	418	709	1,186	1,873	3,002	2,453	593	2
1963	12,043	90	740	1,061	481	417	660	1,064	1,841	2,850	2,293	546	-
1962	11,638	78	706	1,025	483	353	645	1,037	1,791	2,786	2,194	536	4
1961	12,039	97	775	1,120	482	442	677	1,131	1,800	2,830	2,161	521	3
1960	11,922	97	835	1,064	451	465	704	1,153	1,907	2,802	2,019	425	-
WHITE, MALE													
1967	7,599	37	296	686	377	228	369	658	1,198	1,797	1,565	387	1
1966	7,546	25	342	655	372	240	376	652	1,245	1,773	1,476	388	2
1965	7,274	37	347	599	309	223	368	637	1,199	1,803	1,439	312	1
1964	7,186	42	368	582	301	245	369	667	1,116	1,796	1,400	299	1
1963	6,954	58	423	568	301	236	337	606	1,121	1,707	1,291	306	-
1962	6,757	37	381	568	303	204	365	624	1,074	1,685	1,242	272	2
1961	7,086	46	440	657	297	236	386	661	1,089	1,718	1,264	289	3
1960	6,929	52	487	622	273	284	398	668	1,134	1,668	1,115	228	-
WHITE, FEMALE													
1967	5,683	28	265	495	217	162	310	512	788	1,304	1,210	392	-
1966	5,508	36	258	415	212	169	307	506	794	1,227	1,218	366	-
1965	5,403	34	294	467	226	169	273	497	770	1,198	1,154	320	1
1964	5,316	30	274	452	217	173	340	519	757	1,206	1,053	294	1
1963	5,089	32	317	493	180	181	323	458	720	1,143	1,002	240	-
1962	4,881	41	325	457	180	149	280	413	717	1,101	952	264	2
1961	4,953	51	335	463	185	206	291	470	711	1,112	897	232	-
1960	4,993	45	348	442	178	181	306	485	773	1,134	904	197	-

- Quantity zero

Note: Figures by race exclude data for residents of New Jersey for 1962 and 1963.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290A for 1960-67 and separate unpublished tables for the black population and Accidents.

77/09/02 TABLE 290A. DEATHS FOR 60 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1960-67

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	TOTAL	UNDER 1 YEAR	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS	65-74 YEARS	75-84 YEARS	85+ YEARS	N.S.
Leukemia and aleukemia (204)													
ALL OTHER, BOTH SEXES													
1967	1,054	12	46	98	81	65	71	144	172	218	124	22	1
1966	958	8	55	98	66	41	98	123	177	189	84	19	-
1965	951	12	55	91	71	57	82	112	176	182	90	21	2
1964	951	14	67	115	50	61	70	125	177	176	80	16	-
1963	999	12	92	102	61	58	94	121	187	177	78	15	2
1962	893	14	74	111	50	50	60	118	161	165	76	13	1
1961	834	11	51	86	52	50	93	101	159	148	65	18	-
1960	803	14	59	85	50	58	72	116	155	130	52	12	-
ALL OTHER, MALE													
1967	623	9	21	58	53	37	35	83	111	127	71	17	1
1966	514	3	28	49	38	22	49	65	91	118	42	9	-
1965	566	9	31	56	48	32	48	58	96	114	58	14	2
1964	560	5	40	69	34	37	44	63	102	106	52	8	-
1963	565	6	46	50	46	33	49	60	109	107	49	9	1
1962	491	6	38	62	29	29	27	56	87	105	45	7	-
1961	483	6	27	41	37	29	50	44	93	102	42	12	-
1960	442	7	33	51	34	36	40	54	77	74	29	7	-
ALL OTHER, FEMALE													
1967	431	3	25	40	28	28	36	61	61	91	53	5	-
1966	444	5	27	49	28	19	49	58	86	71	42	10	-
1965	385	3	24	35	23	25	34	54	80	68	32	7	-
1964	391	9	27	46	16	24	26	62	75	70	28	8	-
1963	434	6	46	52	15	25	45	61	78	70	29	6	1
1962	402	8	36	49	21	21	33	62	74	60	31	6	1
1961	351	5	24	45	15	21	43	57	66	46	23	6	-
1960	361	7	26	34	16	22	32	62	78	56	23	5	-

- Quantity zero

Note: Figures by race exclude data for residents of New Jersey for 1962 and 1963.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290A for 1960-67 and separate unpublished tables for the black population and Accidents.

77/09/02 TABLE 290A. DEATHS FOR 60 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1960-67

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	TOTAL	UNDER 1 YEAR	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS	65-74 YEARS	75-84 YEARS	85+ YEARS	N.S.
Lymphosarcoma and other neoplasms of lymphatic and hematopoietic tissues (200-203,205)													
BOTH SEXES													
1967	16,481	13	75	323	596	655	1,176	2,186	3,665	4,474	2,817	501	-
1966	15,802	17	72	320	542	702	1,189	2,288	3,418	4,211	2,610	432	1
1965	15,068	25	83	288	552	705	1,202	2,172	3,363	3,920	2,374	384	-
1964	14,746	21	97	294	575	694	1,303	2,101	3,292	3,836	2,156	377	-
1963	14,407	28	109	309	517	677	1,165	2,089	3,245	3,802	2,090	371	5
1962	13,755	29	97	316	470	677	1,144	2,056	3,036	3,714	1,923	292	1
1961	13,554	36	120	324	458	690	1,109	1,968	3,130	3,686	1,763	265	5
1960	12,958	27	107	241	440	715	1,149	1,906	2,996	3,378	1,706	291	2
MALE													
1967	9,218	6	45	221	377	413	720	1,263	2,131	2,362	1,432	248	-
1966	8,950	8	45	219	325	433	738	1,383	2,002	2,271	1,310	215	1
1965	8,593	11	52	203	358	439	755	1,297	1,930	2,160	1,217	171	-
1964	8,301	9	60	215	350	430	801	1,249	1,891	2,034	1,083	179	-
1963	8,243	17	72	231	310	432	731	1,219	1,869	2,133	1,051	175	3
1962	7,896	14	68	230	290	442	697	1,238	1,752	2,043	978	143	1
1961	7,870	20	75	248	287	426	683	1,178	1,812	2,069	947	123	2
1960	7,579	17	75	171	291	447	696	1,170	1,738	1,937	888	148	1
FEMALE													
1967	7,263	7	30	102	219	242	456	923	1,534	2,112	1,385	253	-
1966	6,852	9	27	101	217	269	451	905	1,416	1,940	1,300	217	-
1965	6,475	14	31	85	194	266	447	875	1,433	1,760	1,157	213	-
1964	6,445	12	37	79	225	264	502	852	1,401	1,802	1,073	198	-
1963	6,164	11	37	78	207	245	434	870	1,376	1,669	1,039	196	2
1962	5,859	15	29	86	180	235	447	818	1,284	1,671	945	149	-
1961	5,684	16	45	76	171	264	426	790	1,318	1,617	816	142	3
1960	5,379	10	32	70	149	268	453	736	1,258	1,441	818	143	1

- Quantity zero

Note: Figures by race exclude data for residents of New Jersey for 1962 and 1963.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290A for 1960-67 and separate unpublished tables for the black population and Accidents.

77/09/02 TABLE 290A. DEATHS FOR 60 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1960-67

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	TOTAL	UNDER 1 YEAR	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS	65-74 YEARS	75-84 YEARS	85+ YEARS	N.S.
Lymphosarcoma and other neoplasms of lymphatic and hematopoietic tissues (200-203,205)													
WHITE, BOTH SEXES													
1967	15,137	12	67	273	544	590	1,039	1,964	3,319	4,178	2,683	468	-
1966	14,603	14	60	283	496	621	1,086	2,087	3,143	3,909	2,495	408	1
1965	13,889	22	79	253	501	649	1,082	1,935	3,067	3,669	2,266	366	-
1964	13,593	18	87	251	529	629	1,171	1,905	2,981	3,592	2,071	359	-
1963	12,740	24	87	268	462	593	998	1,782	2,827	3,412	1,937	345	5
1962	12,146	24	84	262	399	588	986	1,733	2,673	3,333	1,792	271	1
1961	12,463	29	104	290	411	638	992	1,750	2,834	3,469	1,693	253	-
1960	12,034	25	96	218	409	655	1,024	1,711	2,774	3,208	1,633	279	2
WHITE, MALE													
1967	8,419	6	40	187	343	371	637	1,129	1,932	2,187	1,358	229	-
1966	8,240	7	41	196	302	388	680	1,271	1,834	2,075	1,246	199	1
1965	7,904	9	50	182	331	404	680	1,156	1,765	2,011	1,150	166	-
1964	7,593	9	55	184	321	383	723	1,134	1,705	1,875	1,036	168	-
1963	7,229	16	57	201	278	379	628	1,045	1,608	1,882	967	165	3
1962	6,944	10	58	195	246	383	605	1,034	1,540	1,828	911	133	1
1961	7,180	16	66	222	256	396	609	1,050	1,619	1,922	905	119	-
1960	7,026	17	69	156	270	415	626	1,059	1,595	1,832	845	141	1
WHITE, FEMALE													
1967	6,718	6	27	86	201	219	402	835	1,387	1,991	1,325	239	-
1966	6,363	7	19	87	194	233	406	816	1,309	1,834	1,249	209	-
1965	5,985	13	29	71	170	245	402	779	1,302	1,658	1,116	200	-
1964	6,000	9	32	67	208	246	448	771	1,276	1,717	1,035	191	-
1963	5,511	8	30	67	184	214	370	737	1,219	1,530	970	180	2
1962	5,202	14	26	67	153	205	381	699	1,133	1,505	881	138	-
1961	5,283	13	38	68	155	242	383	700	1,215	1,547	788	134	-
1960	5,008	8	27	62	139	240	398	652	1,179	1,376	788	138	1

- Quantity zero

Note: Figures by race exclude data for residents of New Jersey for 1962 and 1963.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290A for 1960-67 and separate unpublished tables for the black population and Accidents.

77/09/02 TABLE 290A. DEATHS FOR 60 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1960-67

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	TOTAL	UNDER 1 YEAR	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS	65-74 YEARS	75-84 YEARS	85+ YEARS	N.S.
Lymphosarcoma and other neoplasms of lymphatic and hematopoietic tissues (200-203,205)													
ALL OTHER, BOTH SEXES													
1967	1,344	1	8	50	52	65	137	222	346	296	134	33	-
1966	1,199	3	12	37	46	81	103	201	275	302	115	24	-
1965	1,179	3	4	35	51	56	120	237	296	251	108	18	-
1964	1,153	3	10	43	46	65	132	196	311	244	85	18	-
1963	1,145	4	19	35	40	58	117	216	300	251	88	17	-
1962	1,094	4	8	43	46	63	116	226	260	235	79	14	-
1961	1,091	7	16	34	47	52	117	218	296	217	70	12	5
1960	924	2	11	23	31	60	125	195	222	170	73	12	-
ALL OTHER, MALE													
1967	799	-	5	34	34	42	83	134	199	175	74	19	-
1966	710	1	4	23	23	45	58	112	168	196	64	16	-
1965	689	2	2	21	27	35	75	141	165	149	67	5	-
1964	708	-	5	31	29	47	78	115	186	159	47	11	-
1963	734	1	13	25	20	37	78	121	197	178	57	7	-
1962	663	3	5	27	28	43	69	147	148	141	44	8	-
1961	690	4	9	26	31	30	74	128	193	147	42	4	2
1960	553	-	6	15	21	32	70	111	143	105	43	7	-
ALL OTHER, FEMALE													
1967	545	1	3	16	18	23	54	88	147	121	60	14	-
1966	489	2	8	14	23	36	45	89	107	106	51	8	-
1965	490	1	2	14	24	21	45	96	131	102	41	13	-
1964	445	3	5	12	17	18	54	81	125	85	38	7	-
1963	411	3	6	10	20	21	39	95	103	73	31	10	-
1962	431	1	3	16	18	20	47	79	112	94	35	6	-
1961	401	3	7	8	16	22	43	90	103	70	28	8	3
1960	371	2	5	8	10	28	55	84	79	65	30	5	-

- Quantity zero

Note: Figures by race exclude data for residents of New Jersey for 1962 and 1963.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290A for 1960-67 and separate unpublished tables for the black population and Accidents.

77/09/02 TABLE 290A. DEATHS FOR 60 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1960-67

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	TOTAL	UNDER 1 YEAR	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS	65-74 YEARS	75-84 YEARS	85+ YEARS	N.S.
Benign neoplasms and neoplasms of unspecified nature (210-239)													
BOTH SEXES													
1967	5,013	97	105	230	169	215	425	831	996	996	686	263	-
1966	4,923	85	91	185	189	229	441	758	988	1,029	683	245	-
1965	5,058	105	113	219	211	213	491	818	1,015	977	677	219	-
1964	4,930	95	104	221	168	216	498	819	989	949	653	215	3
1963	4,905	125	91	212	207	233	529	809	957	919	622	200	1
1962	4,681	123	97	236	165	256	572	777	887	854	540	173	1
1961	4,751	124	119	221	176	282	559	811	920	846	519	173	1
1960	4,893	138	103	232	174	264	614	850	940	867	514	197	-
MALE													
1967	2,450	53	62	108	81	99	173	394	516	550	315	99	-
1966	2,298	42	46	96	96	93	168	296	496	541	334	90	-
1965	2,406	46	64	116	112	92	188	344	540	506	317	81	-
1964	2,331	51	66	131	82	92	207	326	518	478	297	81	2
1963	2,280	73	48	117	111	107	182	308	490	472	295	76	1
1962	2,137	58	42	132	74	94	205	311	461	436	264	59	1
1961	2,121	74	48	106	90	114	177	332	469	412	239	60	-
1960	2,227	68	49	131	78	94	236	381	467	434	225	64	-
FEMALE													
1967	2,563	44	43	122	88	116	252	437	480	446	371	164	-
1966	2,625	43	45	89	93	136	273	462	492	488	349	155	-
1965	2,652	59	49	103	99	121	303	474	475	471	360	138	-
1964	2,599	44	38	90	86	124	291	493	471	471	356	134	1
1963	2,625	52	43	95	96	126	347	501	467	447	327	124	-
1962	2,544	65	55	104	91	162	367	466	426	418	276	114	-
1961	2,630	50	71	115	86	168	382	479	451	434	280	113	1
1960	2,666	70	54	101	96	170	378	469	473	433	289	133	-

- Quantity zero

Note: Figures by race exclude data for residents of New Jersey for 1962 and 1963.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290A for 1960-67 and separate unpublished tables for the black population and Accidents.

77/09/02 TABLE 290A. DEATHS FOR 60 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1960-67

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	TOTAL	UNDER 1 YEAR	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS	65-74 YEARS	75-84 YEARS	85+ YEARS	N.S.
Benign neoplasms and neoplasms of unspecified nature (210-239)													
WHITE, BOTH SEXES													
1967	4,290	79	90	175	142	173	341	671	866	880	630	243	-
1966	4,259	73	74	149	158	197	344	619	860	908	643	234	-
1965	4,362	88	91	186	180	172	388	673	882	861	634	207	-
1964	4,227	81	92	191	137	169	374	677	854	847	595	208	2
1963	4,058	105	73	171	163	176	380	642	809	798	557	183	1
1962	3,867	107	80	181	141	193	422	613	735	744	489	161	1
1961	4,056	110	99	177	149	216	410	677	798	772	486	162	-
1960	4,167	119	83	189	141	198	478	700	804	787	482	186	-
WHITE, MALE													
1967	2,138	44	55	82	68	81	150	327	457	494	289	91	-
1966	2,037	38	39	77	86	83	143	251	436	487	314	83	-
1965	2,109	36	52	99	95	77	155	292	473	451	303	76	-
1964	2,049	42	60	117	68	75	166	283	451	428	279	78	2
1963	1,962	61	37	97	88	86	151	252	431	421	267	70	1
1962	1,816	49	30	100	66	74	171	263	393	375	240	54	1
1961	1,872	67	38	86	76	92	148	284	423	377	225	56	-
1960	1,963	59	42	107	65	79	200	327	410	401	213	60	-
WHITE, FEMALE													
1967	2,152	35	35	93	74	92	191	344	409	386	341	152	-
1966	2,222	35	35	72	72	114	201	368	424	421	329	151	-
1965	2,253	52	39	87	85	95	233	381	409	410	331	131	-
1964	2,178	39	32	74	69	94	208	394	403	419	316	130	-
1963	2,096	44	36	74	75	90	229	390	378	377	290	113	-
1962	2,051	58	50	81	75	119	251	350	342	369	249	107	-
1961	2,184	43	61	91	73	124	262	393	375	395	261	106	-
1960	2,204	60	41	82	76	119	278	373	394	386	269	126	-

- Quantity zero

Note: Figures by race exclude data for residents of New Jersey for 1962 and 1963.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290A for 1960-67 and separate unpublished tables for the black population and Accidents.

77/09/02 TABLE 290A. DEATHS FOR 60 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1960-67

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	TOTAL	UNDER 1 YEAR	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS	65-74 YEARS	75-84 YEARS	85+ YEARS	N.S.
Benign neoplasms and neoplasms of unspecified nature (210-239)													
ALL OTHER, BOTH SEXES													
1967	723	18	15	55	27	42	84	160	130	116	56	20	-
1966	664	12	17	36	31	32	97	139	128	121	40	11	-
1965	696	17	22	33	31	41	103	145	133	116	43	12	-
1964	703	14	12	30	31	47	124	142	135	102	58	7	1
1963	685	14	17	38	34	50	129	133	118	92	47	13	-
1962	662	11	15	45	16	54	128	139	127	88	34	5	-
1961	695	14	20	44	27	66	149	134	122	74	33	11	1
1960	726	19	20	43	33	66	136	150	136	80	32	11	-
ALL OTHER, MALE													
1967	312	9	7	26	13	18	23	67	59	56	26	8	-
1966	261	4	7	19	10	10	25	45	60	54	20	7	-
1965	297	10	12	17	17	15	33	52	67	55	14	5	-
1964	282	9	6	14	14	17	41	43	67	50	18	3	-
1963	247	9	11	18	16	16	26	45	41	37	23	5	-
1962	253	6	10	26	2	15	31	41	54	48	18	2	-
1961	249	7	10	20	14	22	29	48	46	35	14	4	-
1960	264	9	7	24	13	15	36	54	57	33	12	4	-
ALL OTHER, FEMALE													
1967	411	9	8	29	14	24	61	93	71	60	30	12	-
1966	403	8	10	17	21	22	72	94	68	67	20	4	-
1965	399	7	10	16	14	26	70	93	66	61	29	7	-
1964	421	5	6	16	17	30	83	99	68	52	40	4	1
1963	438	5	6	20	18	34	103	88	77	55	24	8	-
1962	409	5	5	19	14	39	97	98	73	40	16	3	-
1961	446	7	10	24	13	44	120	86	76	39	19	7	1
1960	462	10	13	19	20	51	100	96	79	47	20	7	-

- Quantity zero

Note: Figures by race exclude data for residents of New Jersey for 1962 and 1963.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290A for 1960-67 and separate unpublished tables for the black population and Accidents.

77/09/02 TABLE 290A. DEATHS FOR 60 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1960-67

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	TOTAL	UNDER 1 YEAR	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS	65-74 YEARS	75-84 YEARS	85+ YEARS	N.S.
Asthma (241)													
BOTH SEXES													
1967	4,137	19	48	92	123	149	344	632	951	1,000	636	143	-
1966	4,324	18	38	108	123	153	315	642	1,009	1,117	665	135	1
1965	4,520	19	51	95	130	165	334	657	1,045	1,149	718	157	-
1964	4,441	16	55	77	115	135	353	678	984	1,151	732	145	-
1963	5,058	25	64	93	109	161	333	698	1,223	1,391	803	158	-
1962	4,896	22	53	93	112	166	356	725	1,085	1,360	762	161	1
1961	4,910	40	58	67	94	178	343	686	1,109	1,338	817	180	-
1960	5,394	24	57	70	94	170	426	764	1,222	1,519	878	169	1
MALE													
1967	2,286	11	18	47	54	54	131	285	576	622	420	68	-
1966	2,498	10	24	65	42	54	114	309	621	751	435	73	-
1965	2,649	11	19	54	49	59	143	330	632	799	462	91	-
1964	2,624	15	33	45	53	55	136	339	626	780	472	70	-
1963	3,160	13	34	54	46	50	123	379	817	1,007	549	88	-
1962	3,102	13	31	56	35	63	151	396	750	990	521	95	1
1961	3,120	20	36	45	29	69	145	381	770	964	563	98	-
1960	3,459	10	30	34	32	61	171	443	870	1,142	579	87	-
FEMALE													
1967	1,851	8	30	45	69	95	213	347	375	378	216	75	-
1966	1,826	8	14	43	81	99	201	333	388	366	230	62	1
1965	1,871	8	32	41	81	106	191	327	413	350	256	66	-
1964	1,817	1	22	32	62	80	217	339	358	371	260	75	-
1963	1,898	12	30	39	63	111	210	319	406	384	254	70	-
1962	1,794	9	22	37	77	103	205	329	335	370	241	66	-
1961	1,790	20	22	22	65	109	198	305	339	374	254	82	-
1960	1,935	14	27	36	62	109	255	321	352	377	299	82	1

- Quantity zero

Note: Figures by race exclude data for residents of New Jersey for 1962 and 1963.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290A for 1960-67 and separate unpublished tables for the black population and Accidents.

77/09/02 TABLE 290A. DEATHS FOR 60 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1960-67

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	TOTAL	UNDER 1 YEAR	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS	65-74 YEARS	75-84 YEARS	85+ YEARS	N.S.
Asthma (241)													
WHITE, BOTH SEXES													
1967	3,399	13	29	60	77	87	236	493	788	896	589	131	-
1966	3,598	11	23	68	72	96	204	488	859	1,016	633	127	1
1965	3,770	10	29	68	78	95	212	505	891	1,059	679	144	-
1964	3,731	13	31	54	81	74	233	543	819	1,056	689	138	-
1963	4,178	15	36	60	62	92	216	532	1,036	1,239	742	148	-
1962	4,087	16	30	64	64	78	231	556	924	1,251	720	152	1
1961	4,186	24	33	42	51	100	232	558	957	1,241	783	165	-
1960	4,655	14	34	43	59	96	294	631	1,080	1,409	835	159	1
WHITE, MALE													
1967	1,909	9	9	27	36	31	82	223	483	557	388	64	-
1966	2,093	7	13	42	20	28	70	230	522	682	411	68	-
1965	2,226	5	9	37	27	31	81	243	533	738	438	84	-
1964	2,220	12	15	29	35	28	86	265	524	708	450	68	-
1963	2,677	8	20	35	22	27	77	298	700	900	506	84	-
1962	2,662	9	15	42	20	31	92	303	647	917	494	91	1
1961	2,716	13	23	29	13	35	98	306	677	894	540	88	-
1960	3,067	7	19	22	17	33	119	372	776	1,066	553	83	-
WHITE, FEMALE													
1967	1,490	4	20	33	41	56	154	270	305	339	201	67	-
1966	1,505	4	10	26	52	68	134	258	337	334	222	59	1
1965	1,544	5	20	31	51	64	131	262	358	321	241	60	-
1964	1,511	1	16	25	46	46	147	278	295	348	239	70	-
1963	1,501	7	16	25	40	65	139	234	336	339	236	64	-
1962	1,425	7	15	22	44	47	139	253	277	334	226	61	-
1961	1,470	11	10	13	38	65	134	252	280	347	243	77	-
1960	1,588	7	15	21	42	63	175	259	304	343	282	76	1

- Quantity zero

Note: Figures by race exclude data for residents of New Jersey for 1962 and 1963.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290A for 1960-67 and separate unpublished tables for the black population and Accidents.

77/09/02 TABLE 290A. DEATHS FOR 60 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1960-67

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	TOTAL	UNDER 1 YEAR	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS	65-74 YEARS	75-84 YEARS	85+ YEARS	N.S.
Asthma (241)													
ALL OTHER, BOTH SEXES													
1967	738	6	19	32	46	62	108	139	163	104	47	12	-
1966	726	7	15	40	51	57	111	154	150	101	32	8	-
1965	750	9	22	27	52	70	122	152	154	90	39	13	-
1964	710	3	24	23	34	61	120	135	165	95	43	7	-
1963	758	10	27	32	44	67	104	149	152	118	47	8	-
1962	692	6	21	26	45	80	117	143	136	87	26	5	-
1961	724	16	25	25	43	78	111	128	152	97	34	15	-
1960	739	10	23	27	35	74	132	133	142	110	43	10	-
ALL OTHER, MALE													
1967	377	2	9	20	18	23	49	62	93	65	32	4	-
1966	405	3	11	23	22	26	44	79	99	69	24	5	-
1965	423	6	10	17	22	28	62	87	99	61	24	7	-
1964	404	3	18	16	18	27	50	74	102	72	22	2	-
1963	403	5	13	18	22	22	39	71	92	83	34	4	-
1962	376	4	15	12	15	28	57	80	87	58	18	2	-
1961	404	7	13	16	16	34	47	75	93	70	23	10	-
1960	392	3	11	12	15	28	52	71	94	76	26	4	-
ALL OTHER, FEMALE													
1967	361	4	10	12	28	39	59	77	70	39	15	8	-
1966	321	4	4	17	29	31	67	75	51	32	8	3	-
1965	327	3	12	10	30	42	60	65	55	29	15	6	-
1964	306	-	6	7	16	34	70	61	63	23	21	5	-
1963	355	5	14	14	22	45	65	78	60	35	13	4	-
1962	316	2	6	14	30	52	60	63	49	29	8	3	-
1961	320	9	12	9	27	44	64	53	59	27	11	5	-
1960	347	7	12	15	20	46	80	62	48	34	17	6	-

- Quantity zero

Note: Figures by race exclude data for residents of New Jersey for 1962 and 1963.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290A for 1960-67 and separate unpublished tables for the black population and Accidents.

77/09/02 TABLE 290A. DEATHS FOR 60 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1960-67

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	TOTAL	UNDER 1 YEAR	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS	65-74 YEARS	75-84 YEARS	85+ YEARS	N.S.
ICD-7 Nos. 260 Diabetes													
BOTH SEXES													
1967	35,049	14	20	97	213	575	1,135	2,789	6,368	11,002	10,069	2,756	11
1966	34,597	17	32	113	225	570	1,196	2,693	6,158	11,068	9,885	2,635	5
1965	33,174	20	31	118	207	581	1,144	2,624	6,130	10,636	9,307	2,365	11
1964	32,279	23	40	109	218	575	1,176	2,614	6,215	10,423	8,756	2,121	9
1963	32,465	15	29	117	244	576	1,214	2,595	6,140	10,774	8,679	2,070	12
1962	31,222	23	33	112	215	545	1,122	2,476	5,950	10,596	8,109	2,032	9
1961	30,098	13	31	121	184	610	1,149	2,361	5,685	10,213	7,943	1,779	9
1960	29,971	17	44	132	216	530	1,092	2,477	5,909	10,267	7,584	1,688	15
MALE													
1967	14,421	6	8	38	88	323	661	1,369	2,895	4,356	3,709	963	5
1966	14,163	9	18	62	98	325	657	1,337	2,682	4,361	3,663	950	1
1965	13,588	10	18	46	95	326	637	1,329	2,651	4,157	3,461	857	1
1964	13,233	13	20	42	108	338	667	1,296	2,710	3,973	3,321	744	1
1963	13,183	7	12	37	94	328	657	1,242	2,624	4,140	3,248	789	5
1962	12,881	13	14	50	92	300	696	1,195	2,550	4,078	3,087	803	3
1961	12,114	3	15	46	87	329	657	1,150	2,387	3,854	2,942	641	3
1960	12,147	11	22	48	97	298	601	1,223	2,401	3,891	2,928	618	9
FEMALE													
1967	20,628	8	12	59	125	252	474	1,420	3,473	6,646	6,360	1,793	6
1966	20,434	8	14	51	127	245	539	1,356	3,476	6,707	6,222	1,685	4
1965	19,586	10	13	72	112	255	507	1,295	3,479	6,479	5,846	1,508	10
1964	19,046	10	20	67	110	237	509	1,318	3,505	6,450	5,435	1,377	8
1963	19,282	8	17	80	150	248	557	1,353	3,516	6,634	5,431	1,281	7
1962	18,341	10	19	62	123	245	426	1,281	3,400	6,518	5,022	1,229	6
1961	17,984	10	16	75	97	281	492	1,211	3,298	6,359	5,001	1,138	6
1960	17,824	6	22	84	119	232	491	1,254	3,508	6,376	4,656	1,070	6

Note: Figures by race exclude data for residents of New Jersey for 1962 and 1963.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290A for 1960-67 and separate unpublished tables for the black population and Accidents.

77/09/02 TABLE 290A. DEATHS FOR 60 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1960-67

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	TOTAL	UNDER 1 YEAR	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS	65-74 YEARS	75-84 YEARS	85+ YEARS	N.S.
ICD-7 Nos. 260 Diabetes													
WHITE, BOTH SEXES													
1967	30,077	14	19	88	165	453	846	2,053	5,014	9,472	9,339	2,607	7
1966	29,640	12	26	99	170	463	863	1,938	4,846	9,568	9,182	2,471	2
1965	28,532	17	28	97	153	481	852	1,909	4,825	9,261	8,687	2,215	7
1964	27,731	14	37	94	171	480	888	1,885	4,846	9,171	8,157	1,985	3
1963	26,836	9	21	99	181	444	864	1,841	4,640	9,063	7,791	1,874	9
1962	25,986	18	30	95	163	448	820	1,711	4,550	9,001	7,289	1,856	5
1961	26,256	10	26	109	143	498	874	1,747	4,564	9,117	7,477	1,686	5
1960	26,121	13	37	111	165	439	813	1,814	4,745	9,253	7,139	1,585	7
WHITE, MALE													
1967	12,686	6	8	34	65	262	531	1,095	2,482	3,817	3,467	915	4
1966	12,451	8	14	54	77	269	516	1,087	2,254	3,873	3,414	884	1
1965	11,922	9	18	38	68	279	506	1,066	2,221	3,682	3,233	802	-
1964	11,632	6	18	35	78	293	535	1,047	2,272	3,554	3,089	705	-
1963	11,203	5	9	30	72	258	516	991	2,134	3,541	2,939	703	5
1962	11,003	11	12	40	68	246	551	949	2,063	3,523	2,793	746	1
1961	10,806	2	12	39	72	288	527	960	2,057	3,447	2,796	605	1
1960	10,761	8	19	36	69	255	493	996	2,018	3,532	2,754	578	3
WHITE, FEMALE													
1967	17,391	8	11	54	100	191	315	958	2,532	5,655	5,872	1,692	3
1966	17,189	4	12	45	93	194	347	851	2,592	5,695	5,768	1,587	1
1965	16,610	8	10	59	85	202	346	843	2,604	5,579	5,454	1,413	7
1964	16,099	8	19	59	93	187	353	838	2,574	5,617	5,068	1,280	3
1963	15,633	4	12	69	109	186	348	850	2,506	5,522	4,852	1,171	4
1962	14,983	7	18	55	95	202	269	762	2,487	5,478	4,496	1,110	4
1961	15,450	8	14	70	71	210	347	787	2,507	5,670	4,681	1,081	4
1960	15,360	5	18	75	96	184	320	818	2,727	5,721	4,385	1,007	4

- Quantity zero

Note: Figures by race exclude data for residents of New Jersey for 1962 and 1963.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290A for 1960-67 and separate unpublished tables for the black population and Accidents.

77/09/02 TABLE 290A. DEATHS FOR 60 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1960-67

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	TOTAL	UNDER 1 YEAR	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS	65-74 YEARS	75-84 YEARS	85+ YEARS	N.S.
ICD-7 Nos. 260 Diabetes													
ALL OTHER, BOTH SEXES													
1967	4,972	-	1	9	48	122	289	736	1,354	1,530	730	149	4
1966	4,957	5	6	14	55	107	333	755	1,312	1,500	703	164	3
1965	4,642	3	3	21	54	100	292	715	1,305	1,375	620	150	4
1964	4,548	9	3	15	47	95	288	729	1,369	1,252	599	136	6
1963	4,266	6	7	15	52	109	311	668	1,249	1,202	527	118	2
1962	3,974	5	2	15	44	79	262	683	1,136	1,118	515	112	3
1961	3,842	3	5	12	41	112	275	614	1,121	1,096	466	93	4
1960	3,850	4	7	21	51	91	279	663	1,164	1,014	445	103	8
ALL OTHER, MALE													
1967	1,735	-	-	4	23	61	130	274	413	539	242	48	1
1966	1,712	1	4	8	21	56	141	250	428	488	249	66	-
1965	1,666	1	-	8	27	47	131	263	430	475	228	55	1
1964	1,601	7	2	7	30	45	132	249	438	419	232	39	1
1963	1,455	2	3	7	17	56	120	210	379	419	188	54	-
1962	1,403	2	2	8	20	45	118	201	377	387	200	42	1
1961	1,308	1	3	7	15	41	130	190	330	407	146	36	2
1960	1,386	3	3	12	28	43	108	227	383	359	174	40	6
ALL OTHER, FEMALE													
1967	3,237	-	1	5	25	61	159	462	941	991	488	101	3
1966	3,245	4	2	6	34	51	192	505	884	1,012	454	98	3
1965	2,976	2	3	13	27	53	161	452	875	900	392	95	3
1964	2,947	2	1	8	17	50	156	480	931	833	367	97	5
1963	2,811	4	4	8	35	53	191	458	870	783	339	64	2
1962	2,571	3	-	7	24	34	144	482	759	731	315	70	2
1961	2,534	2	2	5	26	71	145	424	791	689	320	57	2
1960	2,464	1	4	9	23	48	171	436	781	655	271	63	2

- Quantity zero

Note: Figures by race exclude data for residents of New Jersey for 1962 and 1963.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290A for 1960-67 and separate unpublished tables for the black population and Accidents.

77/09/02 TABLE 290A. DEATHS FOR 60 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1960-67

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	TOTAL	UNDER 1 YEAR	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS	65-74 YEARS	75-84 YEARS	85+ YEARS	N.S.
Anemias (290-293)													
BOTH SEXES													
1967	3,460	85	134	150	181	110	162	199	433	710	911	382	3
1966	3,452	81	128	156	141	111	152	233	400	722	920	408	-
1965	3,539	105	177	157	159	118	161	219	423	722	897	401	-
1964	3,516	89	169	163	157	116	133	221	394	760	891	423	-
1963	3,409	122	175	164	144	89	130	213	357	761	855	398	1
1962	3,398	105	177	170	142	95	141	228	388	712	879	360	1
1961	3,280	107	200	173	123	122	132	191	345	727	813	347	-
1960	3,330	111	199	174	140	147	122	198	382	714	814	326	3
MALE													
1967	1,738	50	63	65	97	50	81	99	238	376	451	165	3
1966	1,659	32	65	72	74	51	75	104	215	383	437	151	-
1965	1,743	58	83	66	87	63	73	97	212	377	466	161	-
1964	1,684	50	82	82	73	52	52	104	217	370	413	189	-
1963	1,634	60	103	64	80	43	64	85	178	407	388	162	-
1962	1,620	56	86	86	74	35	68	110	184	369	416	136	-
1961	1,587	52	101	71	51	50	55	93	183	389	395	147	-
1960	1,567	67	90	69	74	54	45	80	207	387	369	123	2
FEMALE													
1967	1,722	35	71	85	84	60	81	100	195	334	460	217	-
1966	1,793	49	63	84	67	60	77	129	185	339	483	257	-
1965	1,796	47	94	91	72	55	88	122	211	345	431	240	-
1964	1,832	39	87	81	84	64	81	117	177	390	478	234	-
1963	1,775	62	72	100	64	46	66	128	179	354	467	236	1
1962	1,778	49	91	84	68	60	73	118	204	343	463	224	1
1961	1,693	55	99	102	72	72	77	98	162	338	418	200	-
1960	1,763	44	109	105	66	93	77	118	175	327	445	203	1

- Quantity zero

Note: Figures by race exclude data for residents of New Jersey for 1962 and 1963.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290A for 1960-67 and separate unpublished tables for the black population and Accidents.

77/09/02 TABLE 290A. DEATHS FOR 60 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1960-67

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	TOTAL	UNDER 1 YEAR	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS	65-74 YEARS	75-84 YEARS	85+ YEARS	N.S.
Anemias (290-293)													
WHITE, BOTH SEXES													
1967	2,763	33	50	80	83	33	102	149	374	648	839	370	2
1966	2,776	41	45	97	76	39	76	174	325	639	873	391	-
1965	2,829	59	80	87	79	44	77	155	346	661	851	390	-
1964	2,834	46	70	105	80	49	70	160	325	678	844	407	-
1963	2,619	51	65	90	68	29	71	150	284	646	792	372	1
1962	2,699	48	84	115	77	40	80	165	320	628	803	339	-
1961	2,678	53	98	128	54	53	72	146	291	677	775	331	-
1960	2,727	53	107	117	80	62	75	147	331	650	790	312	3
WHITE, MALE													
1967	1,384	15	19	38	43	12	50	71	209	345	419	161	2
1966	1,336	19	22	46	43	19	31	76	182	336	416	146	-
1965	1,396	27	36	32	44	22	37	70	179	349	441	159	-
1964	1,349	26	32	50	41	22	25	77	180	330	386	180	-
1963	1,243	20	38	33	37	17	37	62	147	346	357	149	-
1962	1,294	27	32	55	43	13	37	82	162	335	379	129	-
1961	1,286	27	40	46	26	17	32	74	155	357	375	137	-
1960	1,279	30	43	38	42	21	23	61	184	360	357	118	2
WHITE, FEMALE													
1967	1,379	18	31	42	40	21	52	78	165	303	420	209	-
1966	1,440	22	23	51	33	20	45	98	143	303	457	245	-
1965	1,433	32	44	55	35	22	40	85	167	312	410	231	-
1964	1,485	20	38	55	39	27	45	83	145	348	458	227	-
1963	1,376	31	27	57	31	12	34	88	137	300	435	223	1
1962	1,405	21	52	60	34	27	43	83	158	293	424	210	-
1961	1,392	26	58	82	28	36	40	72	136	320	400	194	-
1960	1,448	23	64	79	38	41	52	86	147	290	433	194	1

- Quantity zero

Note: Figures by race exclude data for residents of New Jersey for 1962 and 1963.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290A for 1960-67 and separate unpublished tables for the black population and Accidents.

77/09/02 TABLE 290A. DEATHS FOR 60 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1960-67

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	TOTAL	UNDER 1 YEAR	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS	65-74 YEARS	75-84 YEARS	85+ YEARS	N.S.
Anemias (290-293)													
ALL OTHER, BOTH SEXES													
1967	697	52	84	70	98	77	60	50	59	62	72	12	1
1966	676	40	83	59	65	72	76	59	75	83	47	17	-
1965	710	46	97	70	80	74	84	64	77	61	46	11	-
1964	682	43	99	58	77	67	63	61	69	82	47	16	-
1963	667	67	101	70	69	58	51	53	66	75	41	16	-
1962	573	54	88	49	62	49	56	52	51	60	39	12	1
1961	602	54	102	45	69	69	60	45	54	50	38	16	-
1960	603	58	92	57	60	85	47	51	51	64	24	14	-
ALL OTHER, MALE													
1967	354	35	44	27	54	38	31	28	29	31	32	4	1
1966	323	13	43	26	31	32	44	28	33	47	21	5	-
1965	347	31	47	34	43	41	36	27	33	28	25	2	-
1964	335	24	50	32	32	30	27	27	37	40	27	9	-
1963	330	38	60	30	37	26	25	18	28	39	20	9	-
1962	271	27	51	27	29	20	27	23	16	24	22	5	-
1961	301	25	61	25	25	33	23	19	28	32	20	10	-
1960	288	37	47	31	32	33	22	19	23	27	12	5	-
ALL OTHER, FEMALE													
1967	343	17	40	43	44	39	29	22	30	31	40	8	-
1966	353	27	40	33	34	40	32	31	42	36	26	12	-
1965	363	15	50	36	37	33	48	37	44	33	21	9	-
1964	347	19	49	26	45	37	36	34	32	42	20	7	-
1963	337	29	41	40	32	32	26	35	38	36	21	7	-
1962	302	27	37	22	33	29	29	29	35	36	17	7	1
1961	301	29	41	20	44	36	37	26	26	18	18	6	-
1960	315	21	45	26	28	52	25	32	28	37	12	9	-

- Quantity zero

Note: Figures by race exclude data for residents of New Jersey for 1962 and 1963.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290A for 1960-67 and separate unpublished tables for the black population and Accidents.

77/09/02 TABLE 290A. DEATHS FOR 60 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1960-67

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	TOTAL	UNDER 1 YEAR	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS	65-74 YEARS	75-84 YEARS	85+ YEARS	N.S.
Meningitis, except meningococcal and tuberculous (340)													
BOTH SEXES													
1967	2,046	679	311	124	68	45	103	150	230	194	111	30	1
1966	2,324	802	343	135	67	65	123	185	216	206	147	34	1
1965	2,363	844	433	125	67	72	101	168	194	200	123	35	1
1964	2,464	953	439	134	67	59	105	176	204	199	100	27	1
1963	2,343	866	459	137	58	68	113	139	192	170	113	26	2
1962	2,322	907	390	151	58	62	102	180	184	189	77	22	-
1961	2,263	876	428	157	58	63	90	138	154	181	99	18	1
1960	2,316	872	447	147	57	69	105	173	165	168	91	21	1
MALE													
1967	1,197	407	184	68	39	27	65	97	141	106	48	15	-
1966	1,298	456	184	76	37	36	80	110	130	108	63	17	1
1965	1,313	467	239	62	36	49	58	106	116	108	55	16	1
1964	1,406	547	245	73	37	31	68	111	125	105	48	15	1
1963	1,343	483	275	76	41	38	69	91	112	95	54	8	1
1962	1,315	510	222	84	34	37	64	107	108	98	37	14	-
1961	1,309	507	249	88	31	40	52	91	88	105	48	9	1
1960	1,358	524	269	89	33	40	62	105	99	78	49	10	-
FEMALE													
1967	849	272	127	56	29	18	38	53	89	88	63	15	1
1966	1,026	346	159	59	30	29	43	75	86	98	84	17	-
1965	1,050	377	194	63	31	23	43	62	78	92	68	19	-
1964	1,058	406	194	61	30	28	37	65	79	94	52	12	-
1963	1,000	383	184	61	17	30	44	48	80	75	59	18	1
1962	1,007	397	168	67	24	25	38	73	76	91	40	8	-
1961	954	369	179	69	27	23	38	47	66	76	51	9	-
1960	958	348	178	58	24	29	43	68	66	90	42	11	1

- Quantity zero

Note: Figures by race exclude data for residents of New Jersey for 1962 and 1963.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290A for 1960-67 and separate unpublished tables for the black population and Accidents.

77/09/02 TABLE 290A. DEATHS FOR 60 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1960-67

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	TOTAL	UNDER 1 YEAR	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS	65-74 YEARS	75-84 YEARS	85+ YEARS	N.S.
Meningitis, except meningococcal and tuberculous (340)													
WHITE, BOTH SEXES													
1967	1,431	455	224	97	47	28	57	96	165	142	94	26	-
1966	1,585	509	233	92	47	40	79	135	147	157	120	26	-
1965	1,626	538	306	99	46	44	61	102	139	156	104	31	-
1964	1,688	595	316	111	47	38	69	110	141	150	85	25	1
1963	1,599	576	303	99	33	47	67	98	141	125	87	23	-
1962	1,552	549	278	109	38	41	67	121	127	141	61	20	-
1961	1,597	558	328	116	39	39	55	99	108	146	93	16	-
1960	1,608	570	315	116	37	44	66	120	114	132	77	17	-
WHITE, MALE													
1967	835	266	129	52	26	17	38	66	106	79	43	13	-
1966	895	289	134	54	26	24	52	78	94	81	49	14	-
1965	920	312	180	50	22	26	37	60	83	88	47	15	-
1964	974	350	182	59	24	24	46	73	83	78	41	13	1
1963	938	334	178	59	24	29	41	69	81	72	43	8	-
1962	878	311	151	55	24	25	42	76	80	71	30	13	-
1961	936	327	196	65	21	24	35	65	59	88	47	9	-
1960	959	352	201	67	24	25	39	73	68	62	39	9	-
WHITE, FEMALE													
1967	596	189	95	45	21	11	19	30	59	63	51	13	-
1966	690	220	99	38	21	16	27	57	53	76	71	12	-
1965	706	226	126	49	24	18	24	42	56	68	57	16	-
1964	714	245	134	52	23	14	23	37	58	72	44	12	-
1963	661	242	125	40	9	18	26	29	60	53	44	15	-
1962	674	238	127	54	14	16	25	45	47	70	31	7	-
1961	661	231	132	51	18	15	20	34	49	58	46	7	-
1960	649	218	114	49	13	19	27	47	46	70	38	8	-

- Quantity zero

Note: Figures by race exclude data for residents of New Jersey for 1962 and 1963.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290A for 1960-67 and separate unpublished tables for the black population and Accidents.

77/09/02 TABLE 290A. DEATHS FOR 60 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1960-67

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	TOTAL	UNDER 1 YEAR	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS	65-74 YEARS	75-84 YEARS	85+ YEARS	N.S.
Meningitis, except meningococcal and tuberculous (340)													
ALL OTHER, BOTH SEXES													
1967	615	224	87	27	21	17	46	54	65	52	17	4	1
1966	739	293	110	43	20	25	44	50	69	49	27	8	1
1965	737	306	127	26	21	28	40	66	55	44	19	4	1
1964	776	358	123	23	20	21	36	66	63	49	15	2	-
1963	670	270	136	35	24	16	43	35	47	38	21	3	2
1962	701	334	101	40	19	19	30	50	51	42	13	2	-
1961	666	318	100	41	19	24	35	39	46	35	6	2	1
1960	708	302	132	31	20	25	39	53	51	36	14	4	1
ALL OTHER, MALE													
1967	362	141	55	16	13	10	27	31	35	27	5	2	-
1966	403	167	50	22	11	12	28	32	36	27	14	3	1
1965	393	155	59	12	14	23	21	46	33	20	8	1	1
1964	432	197	63	14	13	7	22	38	42	27	7	2	-
1963	359	138	81	17	16	5	26	17	29	20	9	-	1
1962	394	184	64	27	9	10	19	26	23	24	7	1	-
1961	373	180	53	23	10	16	17	26	29	17	1	-	1
1960	399	172	68	22	9	15	23	32	31	16	10	1	-
ALL OTHER, FEMALE													
1967	253	83	32	11	8	7	19	23	30	25	12	2	1
1966	336	126	60	21	9	13	16	18	33	22	13	5	-
1965	344	151	68	14	7	5	19	20	22	24	11	3	-
1964	344	161	60	9	7	14	14	28	21	22	8	-	-
1963	311	132	55	18	8	11	17	18	18	18	12	3	1
1962	307	150	37	13	10	9	11	24	28	18	6	1	-
1961	293	138	47	18	9	8	18	13	17	18	5	2	-
1960	309	130	64	9	11	10	16	21	20	20	4	3	1

- Quantity zero

Note: Figures by race exclude data for residents of New Jersey for 1962 and 1963.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290A for 1960-67 and separate unpublished tables for the black population and Accidents.

77/09/02 TABLE 290A. DEATHS FOR 60 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1960-67

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	TOTAL	UNDER 1 YEAR	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS	65-74 YEARS	75-84 YEARS	85+ YEARS	N.S.
Major cardiovascular-renal diseases (300-334,400-468,592-594)													
BOTH SEXES													
1967	1,012,047	563	304	793	2,082	5,079	23,416	71,800	151,767	262,462	321,665	171,848	268
1966	1,021,188	559	364	851	2,188	5,360	23,761	72,308	153,328	268,477	324,237	169,456	299
1965	1,000,787	594	405	828	2,140	5,428	24,109	72,056	150,811	263,532	316,526	164,079	279
1964	984,261	578	447	802	2,210	5,398	24,517	71,704	150,544	263,550	307,438	156,773	300
1963	994,747	524	390	873	2,104	5,612	24,751	72,195	150,887	269,549	309,713	157,829	320
1962	968,809	488	369	898	2,023	5,575	24,458	71,538	147,164	264,744	298,853	152,407	292
1961	936,218	508	378	845	2,071	5,694	23,947	70,326	144,078	258,286	286,194	143,585	306
1960	935,668	501	398	942	2,023	5,883	24,280	70,661	146,973	261,035	283,744	138,878	350
MALE													
1967	554,242	319	176	408	1,044	2,830	15,868	51,605	104,932	155,365	155,840	65,713	142
1966	560,431	307	199	452	1,148	3,024	16,188	52,015	106,047	158,596	157,242	65,049	164
1965	551,694	345	211	402	1,104	3,103	16,452	51,675	103,451	156,220	154,644	63,922	165
1964	542,947	322	243	380	1,141	3,101	16,792	51,067	102,432	155,891	150,417	61,001	160
1963	550,674	289	209	470	1,083	3,266	16,785	51,710	102,057	159,806	152,360	62,472	167
1962	536,486	280	196	442	1,060	3,220	16,659	51,295	99,152	156,547	146,644	60,837	154
1961	521,229	289	203	421	1,132	3,331	16,446	50,221	97,188	152,893	141,199	57,739	167
1960	522,638	304	233	501	1,072	3,391	16,526	50,645	98,487	155,053	140,121	56,121	184
FEMALE													
1967	457,805	244	128	385	1,038	2,249	7,548	20,195	46,835	107,097	165,825	106,135	126
1966	460,757	252	165	399	1,040	2,336	7,573	20,293	47,281	109,881	166,995	104,407	135
1965	449,093	249	194	426	1,036	2,325	7,657	20,381	47,360	107,312	161,882	100,157	114
1964	441,314	256	204	422	1,069	2,297	7,725	20,637	48,112	107,659	157,021	95,772	140
1963	444,073	235	181	403	1,021	2,346	7,966	20,485	48,830	109,743	157,353	95,357	153
1962	432,323	208	173	456	963	2,355	7,799	20,243	48,012	108,197	152,209	91,570	138
1961	414,989	219	175	424	939	2,363	7,501	20,105	46,890	105,393	144,995	85,846	139
1960	413,030	197	165	441	951	2,492	7,754	20,016	48,486	105,982	143,623	82,757	166

Note: Figures by race exclude data for residents of New Jersey for 1962 and 1963.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290A for 1960-67 and separate unpublished tables for the black population and Accidents.

77/09/02 TABLE 290A. DEATHS FOR 60 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1960-67

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	TOTAL	UNDER 1 YEAR	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS	65-74 YEARS	75-84 YEARS	85+ YEARS	N.S.
Major cardiovascular-renal diseases (300-334,400-468,592-594)													
WHITE, BOTH SEXES													
1967	909,993	410	218	601	1,508	3,299	17,412	58,854	130,032	233,032	301,627	162,845	155
1966	916,441	374	243	651	1,604	3,565	17,659	59,288	130,733	238,133	303,621	160,396	174
1965	898,597	424	275	609	1,581	3,671	17,975	58,903	128,113	234,495	296,763	155,624	164
1964	883,834	402	314	585	1,652	3,662	18,420	58,490	127,213	235,417	288,650	148,854	175
1963	858,900	372	248	646	1,489	3,714	17,874	56,342	122,458	230,700	280,083	144,764	210
1962	837,303	335	264	654	1,441	3,727	17,677	56,012	119,402	227,109	270,883	139,613	186
1961	842,521	368	271	642	1,570	3,989	18,230	57,556	122,186	231,784	269,493	136,246	186
1960	842,273	360	261	702	1,523	4,133	18,548	57,739	124,685	234,823	267,422	131,872	205
WHITE, MALE													
1967	500,732	240	127	308	783	1,961	12,619	44,256	92,762	139,856	145,908	61,829	83
1966	505,301	205	133	350	874	2,107	12,854	44,556	93,337	142,566	147,114	61,112	93
1965	497,949	245	148	301	832	2,218	13,098	44,192	91,014	140,725	144,848	60,233	95
1964	490,510	224	172	273	923	2,277	13,501	43,746	89,732	140,908	141,093	57,568	93
1963	478,756	211	134	353	787	2,306	13,023	42,410	86,063	138,527	137,826	57,010	106
1962	467,179	191	142	322	802	2,298	12,917	42,153	83,606	136,134	132,959	55,552	103
1961	472,452	203	143	318	893	2,508	13,406	43,280	85,673	138,601	132,874	54,447	106
1960	473,640	228	153	385	836	2,576	13,569	43,608	86,556	140,980	131,770	52,873	106
WHITE, FEMALE													
1967	409,261	170	91	293	725	1,338	4,793	14,598	37,270	93,176	155,719	101,016	72
1966	411,140	169	110	301	730	1,458	4,805	14,732	37,396	95,567	156,507	99,284	81
1965	400,648	179	127	308	749	1,453	4,877	14,711	37,099	93,770	151,915	95,391	69
1964	393,324	178	142	312	729	1,385	4,919	14,744	37,481	94,509	147,557	91,286	82
1963	380,144	161	114	293	702	1,408	4,851	13,932	36,395	92,173	142,257	87,754	104
1962	370,124	144	122	332	639	1,429	4,760	13,859	35,796	90,975	137,924	84,061	83
1961	370,069	165	128	324	677	1,481	4,824	14,276	36,513	93,183	136,619	81,799	80
1960	368,633	132	108	317	687	1,557	4,979	14,131	38,129	93,843	135,652	78,999	99

Note: Figures by race exclude data for residents of New Jersey for 1962 and 1963.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290A for 1960-67 and separate unpublished tables for the black population and Accidents.

77/09/02 TABLE 290A. DEATHS FOR 60 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1960-67

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	TOTAL	UNDER 1 YEAR	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS	65-74 YEARS	75-84 YEARS	85+ YEARS	N.S.
Major cardiovascular-renal diseases (300-334,400-468,592-594)													
ALL OTHER, BOTH SEXES													
1967	102,054	153	86	192	574	1,780	6,004	12,946	21,735	29,430	20,038	9,003	113
1966	104,747	185	121	200	584	1,795	6,102	13,020	22,595	30,344	20,616	9,060	125
1965	102,190	170	130	219	559	1,757	6,134	13,153	22,698	29,037	19,763	8,455	115
1964	100,427	176	133	217	558	1,736	6,097	13,214	23,331	28,133	18,788	7,919	125
1963	99,494	142	130	197	551	1,673	5,888	12,973	22,691	28,390	18,810	7,950	99
1962	96,036	143	99	221	509	1,648	5,851	12,769	22,044	27,293	17,625	7,734	100
1961	93,697	140	107	203	501	1,705	5,717	12,770	21,892	26,502	16,701	7,339	120
1960	93,395	141	137	240	500	1,750	5,732	12,922	22,288	26,212	16,322	7,006	145
ALL OTHER, MALE													
1967	53,510	79	49	100	261	869	3,249	7,349	12,170	15,509	9,932	3,884	59
1966	55,130	102	66	102	274	917	3,334	7,459	12,710	16,030	10,128	3,937	71
1965	53,745	100	63	101	272	885	3,354	7,483	12,437	15,495	9,796	3,689	70
1964	52,437	98	71	107	218	824	3,291	7,321	12,700	14,983	9,324	3,433	67
1963	52,234	73	70	98	266	826	3,075	7,228	12,147	15,299	9,509	3,589	54
1962	50,240	84	51	110	212	789	3,113	7,154	11,732	14,582	8,864	3,502	47
1961	48,777	86	60	103	239	823	3,040	6,941	11,515	14,292	8,325	3,292	61
1960	48,998	76	80	116	236	815	2,957	7,037	11,931	14,073	8,351	3,248	78
ALL OTHER, FEMALE													
1967	48,544	74	37	92	313	911	2,755	5,597	9,565	13,921	10,106	5,119	54
1966	49,617	83	55	98	310	878	2,768	5,561	9,885	14,314	10,488	5,123	54
1965	48,445	70	67	118	287	872	2,780	5,670	10,261	13,542	9,967	4,766	45
1964	47,990	78	62	110	340	912	2,806	5,893	10,631	13,150	9,464	4,486	58
1963	47,260	69	60	99	285	847	2,813	5,745	10,544	13,091	9,301	4,361	45
1962	45,796	59	48	111	297	859	2,738	5,615	10,312	12,711	8,761	4,232	53
1961	44,920	54	47	100	262	882	2,677	5,829	10,377	12,210	8,376	4,047	59
1960	44,397	65	57	124	264	935	2,775	5,885	10,357	12,139	7,971	3,758	67

Note: Figures by race exclude data for residents of New Jersey for 1962 and 1963.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290A for 1960-67 and separate unpublished tables for the black population and Accidents.

77/09/02 TABLE 290A. DEATHS FOR 60 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1960-67

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	TOTAL	UNDER 1 YEAR	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS	65-74 YEARS	75-84 YEARS	85+ YEARS	N.S.
Diseases of cardiovascular system (330-334,400-468)													
BOTH SEXES													
1967	1,002,111	541	278	666	1,700	4,661	22,595	70,473	150,063	260,393	319,650	170,828	263
1966	1,010,812	530	348	684	1,773	4,850	22,921	70,982	151,596	266,323	322,090	168,419	296
1965	990,192	563	364	670	1,735	4,920	23,263	70,651	149,063	261,373	314,326	162,987	277
1964	973,182	553	408	645	1,779	4,865	23,602	70,296	148,699	261,322	305,071	155,644	298
1963	983,504	503	350	694	1,695	5,036	23,858	70,715	149,030	267,181	307,408	156,720	314
1962	957,437	441	338	724	1,625	4,971	23,471	70,161	145,383	262,370	296,421	151,245	287
1961	924,622	484	353	675	1,635	5,047	22,984	68,907	142,229	255,874	283,748	142,384	302
1960	923,635	479	361	770	1,616	5,235	23,261	69,155	145,087	258,406	281,273	137,651	341
MALE													
1967	548,726	306	163	347	826	2,594	15,401	50,837	103,903	154,235	154,744	65,231	139
1966	554,633	291	190	369	894	2,726	15,704	51,235	105,026	157,365	156,133	64,538	162
1965	545,801	329	191	335	866	2,795	15,957	50,812	102,472	155,014	153,474	63,393	163
1964	536,841	306	221	303	894	2,804	16,264	50,235	101,359	154,669	149,158	60,468	160
1963	544,452	281	190	382	854	2,913	16,276	50,840	100,981	158,516	151,122	61,933	164
1962	530,155	254	176	359	820	2,868	16,085	50,485	98,124	155,187	145,395	60,250	152
1961	514,831	277	184	337	871	2,942	15,881	49,380	96,143	151,553	139,908	57,192	163
1960	516,016	290	210	409	817	2,999	15,930	49,783	97,412	153,632	138,819	55,535	180
FEMALE													
1967	453,385	235	115	319	874	2,067	7,194	19,636	46,160	106,158	164,906	105,597	124
1966	456,179	239	158	315	879	2,124	7,217	19,747	46,570	108,958	165,957	103,881	134
1965	444,391	234	173	335	869	2,125	7,306	19,839	46,591	106,359	160,852	99,594	114
1964	436,341	247	187	342	885	2,061	7,338	20,061	47,340	106,653	155,913	95,176	138
1963	439,052	222	160	312	841	2,123	7,582	19,875	48,049	108,665	156,286	94,787	150
1962	427,282	187	162	365	805	2,103	7,386	19,676	47,259	107,183	151,026	90,995	135
1961	409,791	207	169	338	764	2,105	7,103	19,527	46,086	104,321	143,840	85,192	139
1960	407,619	189	151	361	799	2,236	7,331	19,372	47,675	104,774	142,454	82,116	161

Note: Figures by race exclude data for residents of New Jersey for 1962 and 1963.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290A for 1960-67 and separate unpublished tables for the black population and Accidents.

77/09/02 TABLE 290A. DEATHS FOR 60 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1960-67

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	TOTAL	UNDER 1 YEAR	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS	65-74 YEARS	75-84 YEARS	85+ YEARS	N.S.
Diseases of cardiovascular system (330-334,400-468)													
WHITE, BOTH SEXES													
1967	902,483	393	198	494	1,213	3,032	16,903	57,980	128,839	231,420	299,894	161,964	153
1966	908,521	351	232	512	1,273	3,204	17,122	58,409	129,517	236,444	301,785	159,499	173
1965	890,438	402	247	488	1,244	3,328	17,417	57,916	126,854	232,828	294,883	154,669	162
1964	875,281	383	283	460	1,308	3,288	17,800	57,540	125,919	233,665	286,607	147,854	174
1963	850,503	356	224	503	1,177	3,315	17,290	55,377	121,145	228,938	278,153	143,817	208
1962	828,734	304	242	524	1,116	3,284	17,029	55,104	118,167	225,314	268,826	138,640	184
1961	833,368	350	253	506	1,215	3,496	17,545	56,557	120,828	229,857	267,379	135,197	185
1960	832,888	343	241	565	1,182	3,631	17,838	56,705	123,315	232,756	265,302	130,807	203
WHITE, MALE													
1967	496,486	229	118	260	610	1,788	12,309	43,718	91,997	138,967	144,989	61,419	82
1966	500,843	194	128	286	671	1,889	12,515	44,020	92,613	141,578	146,180	60,677	92
1965	493,365	236	135	249	635	1,995	12,761	43,566	90,271	139,778	143,863	59,783	93
1964	485,735	212	155	215	719	2,054	13,119	43,161	88,945	139,943	140,022	57,097	93
1963	474,056	205	124	284	616	2,048	12,666	41,822	85,270	137,562	136,801	56,552	106
1962	462,325	173	128	264	604	2,022	12,523	41,583	82,856	135,095	131,906	55,069	102
1961	467,365	196	131	251	675	2,195	12,975	42,656	84,889	137,555	131,764	53,973	105
1960	468,420	217	141	313	619	2,252	13,141	42,966	85,753	139,875	130,671	52,367	105
WHITE, FEMALE													
1967	405,997	164	80	234	603	1,244	4,594	14,262	36,842	92,453	154,905	100,545	71
1966	407,678	157	104	226	602	1,315	4,607	14,389	36,904	94,866	155,605	98,822	81
1965	397,073	166	112	239	609	1,333	4,656	14,350	36,583	93,050	151,020	94,886	69
1964	389,546	171	128	245	589	1,234	4,681	14,379	36,974	93,722	146,585	90,757	81
1963	376,447	151	100	219	561	1,267	4,624	13,555	35,875	91,376	141,352	87,265	102
1962	366,409	131	114	260	512	1,262	4,506	13,521	35,311	90,219	136,920	83,571	82
1961	366,003	154	122	255	540	1,301	4,570	13,901	35,939	92,302	135,615	81,224	80
1960	364,468	126	100	252	563	1,379	4,697	13,739	37,562	92,881	134,631	78,440	98

Note: Figures by race exclude data for residents of New Jersey for 1962 and 1963.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290A for 1960-67 and separate unpublished tables for the black population and Accidents.

77/09/02 TABLE 290A. DEATHS FOR 60 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1960-67

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	TOTAL	UNDER 1 YEAR	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS	65-74 YEARS	75-84 YEARS	85+ YEARS	N.S.
Diseases of cardiovascular system (330-334,400-468)													
ALL OTHER, BOTH SEXES													
1967	99,628	148	80	172	487	1,629	5,692	12,493	21,224	28,973	19,756	8,864	110
1966	102,291	179	116	172	500	1,646	5,799	12,573	22,079	29,879	20,305	8,920	123
1965	99,754	161	117	182	491	1,592	5,846	12,735	22,209	28,545	19,443	8,318	115
1964	97,901	170	125	185	471	1,577	5,802	12,756	22,780	27,657	18,464	7,790	124
1963	97,025	137	117	165	468	1,519	5,603	12,516	22,220	27,867	18,501	7,817	95
1962	93,585	128	90	182	454	1,504	5,550	12,339	21,565	26,786	17,313	7,577	97
1961	91,254	134	100	169	420	1,551	5,439	12,350	21,401	26,017	16,369	7,187	117
1960	90,747	136	120	205	434	1,604	5,423	12,450	21,772	25,650	15,971	6,844	138
ALL OTHER, MALE													
1967	52,240	77	45	87	216	806	3,092	7,119	11,906	15,268	9,755	3,812	57
1966	53,790	97	62	83	223	837	3,189	7,215	12,413	15,787	9,953	3,861	70
1965	52,436	93	56	86	231	800	3,196	7,246	12,201	15,236	9,611	3,610	70
1964	51,106	94	66	88	175	750	3,145	7,074	12,414	14,726	9,136	3,371	67
1963	50,896	71	63	81	213	743	2,936	6,978	11,902	15,014	9,328	3,516	51
1962	48,948	77	45	87	181	722	2,954	6,935	11,493	14,299	8,699	3,410	46
1961	47,466	81	53	86	196	747	2,906	6,724	11,254	13,998	8,144	3,219	58
1960	47,596	73	69	96	198	747	2,789	6,817	11,659	13,757	8,148	3,168	75
ALL OTHER, FEMALE													
1967	47,388	71	35	85	271	823	2,600	5,374	9,318	13,705	10,001	5,052	53
1966	48,501	82	54	89	277	809	2,610	5,358	9,666	14,092	10,352	5,059	53
1965	47,318	68	61	96	260	792	2,650	5,489	10,008	13,309	9,832	4,708	45
1964	46,795	76	59	97	296	827	2,657	5,682	10,366	12,931	9,328	4,419	57
1963	46,129	66	54	84	255	776	2,667	5,538	10,318	12,853	9,173	4,301	44
1962	44,637	51	45	95	273	782	2,596	5,404	10,072	12,487	8,614	4,167	51
1961	43,788	53	47	83	224	804	2,533	5,626	10,147	12,019	8,225	3,968	59
1960	43,151	63	51	109	236	857	2,634	5,633	10,113	11,893	7,823	3,676	63

Note: Figures by race exclude data for residents of New Jersey for 1962 and 1963.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290A for 1960-67 and separate unpublished tables for the black population and Accidents.

77/09/02 TABLE 290A. DEATHS FOR 60 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1960-67

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	TOTAL	UNDER 1 YEAR	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS	65-74 YEARS	75-84 YEARS	85+ YEARS	N.S.
ICD-7 Nos. 330-334 Cerebrovascular diseases													
BOTH SEXES													
1967	202,184	107	86	252	461	1,110	3,666	9,831	21,248	48,183	75,011	42,176	53
1966	204,841	101	116	289	492	1,068	3,747	9,668	21,417	49,970	76,067	41,841	65
1965	201,057	155	120	260	449	1,065	3,745	9,917	21,700	49,396	73,966	40,224	60
1964	198,209	125	140	231	486	1,087	3,711	10,066	22,163	49,566	72,479	38,071	84
1963	201,166	142	124	240	440	1,070	3,713	9,942	22,666	51,050	73,388	38,317	74
1962	197,451	128	136	280	420	1,043	3,747	9,962	22,325	50,775	71,092	37,478	65
1961	192,951	144	144	242	427	1,074	3,573	9,789	21,875	50,461	69,326	35,830	66
1960	193,588	169	131	246	427	1,077	3,549	10,078	22,936	51,599	69,099	34,201	76
MALE													
1967	93,071	53	57	135	216	497	1,735	5,050	11,829	25,047	33,310	15,117	25
1966	94,583	51	65	164	261	492	1,834	5,116	12,090	25,725	33,772	14,985	28
1965	93,520	101	61	129	234	483	1,822	5,168	12,176	25,590	33,024	14,701	31
1964	92,511	75	71	118	268	530	1,755	5,171	12,271	25,494	32,642	14,081	35
1963	94,239	80	70	143	229	523	1,770	5,169	12,576	26,356	32,951	14,338	34
1962	92,595	74	76	144	205	476	1,777	5,158	12,035	26,314	32,208	14,105	23
1961	91,025	83	80	124	218	534	1,748	5,064	11,849	26,158	31,488	13,649	30
1960	92,263	105	78	135	230	501	1,721	5,264	12,348	27,151	31,500	13,198	32
FEMALE													
1967	109,113	54	29	117	245	613	1,931	4,781	9,419	23,136	41,701	27,059	28
1966	110,258	50	51	125	231	576	1,913	4,552	9,327	24,245	42,295	26,856	37
1965	107,537	54	59	131	215	582	1,923	4,749	9,524	23,806	40,942	25,523	29
1964	105,698	50	69	113	218	557	1,956	4,895	9,892	24,072	39,837	23,990	49
1963	106,927	62	54	97	211	547	1,943	4,773	10,090	24,694	40,437	23,979	40
1962	104,856	54	60	136	215	567	1,970	4,804	10,290	24,461	38,884	23,373	42
1961	101,926	61	64	118	209	540	1,825	4,725	10,026	24,303	37,838	22,181	36
1960	101,325	64	53	111	197	576	1,828	4,814	10,588	24,448	37,599	21,003	44

Note: Figures by race exclude data for residents of New Jersey for 1962 and 1963.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290A for 1960-67 and separate unpublished tables for the black population and Accidents.

77/09/02 TABLE 290A. DEATHS FOR 60 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1960-67

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	TOTAL	UNDER 1 YEAR	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS	65-74 YEARS	75-84 YEARS	85+ YEARS	N.S.
ICD-7 Nos. 330-334 Cerebrovascular diseases													
WHITE, BOTH SEXES													
1967	176,761	80	75	206	362	729	2,349	6,843	16,146	40,428	69,678	39,844	21
1966	178,487	77	87	232	364	688	2,359	6,669	16,214	41,890	70,463	39,411	33
1965	174,591	120	84	204	343	715	2,368	6,753	16,079	41,282	68,545	38,066	32
1964	172,598	87	109	179	391	716	2,368	6,919	16,348	42,078	67,287	36,074	42
1963	169,922	100	94	181	337	718	2,267	6,585	16,319	41,742	66,197	35,342	40
1962	167,098	92	105	223	311	659	2,313	6,652	15,967	41,795	64,411	34,533	37
1961	168,737	109	114	196	333	721	2,282	6,739	16,233	43,234	64,835	33,907	34
1960	169,053	120	93	201	322	695	2,301	6,897	17,069	44,364	64,605	32,351	35
WHITE, MALE													
1967	80,932	40	51	112	171	334	1,144	3,598	9,267	21,192	30,848	14,164	11
1966	81,912	36	52	134	204	329	1,188	3,602	9,424	21,766	31,143	14,020	14
1965	80,823	79	43	100	187	338	1,158	3,616	9,370	21,604	30,464	13,851	13
1964	80,295	55	58	88	228	373	1,164	3,670	9,421	21,795	30,177	13,247	19
1963	79,458	61	54	105	173	371	1,151	3,539	9,467	21,728	29,655	13,138	16
1962	78,380	54	60	112	157	327	1,143	3,546	8,926	21,898	29,168	12,973	16
1961	79,638	62	64	102	180	391	1,154	3,638	9,193	22,653	29,389	12,795	17
1960	80,486	76	57	116	174	351	1,192	3,731	9,525	23,558	29,334	12,359	13
WHITE, FEMALE													
1967	95,829	40	24	94	191	395	1,205	3,245	6,879	19,236	38,830	25,680	10
1966	96,575	41	35	98	160	359	1,171	3,067	6,790	20,124	39,320	25,391	19
1965	93,768	41	41	104	156	377	1,210	3,137	6,709	19,678	38,081	24,215	19
1964	92,303	32	51	91	163	343	1,204	3,249	6,927	20,283	37,110	22,827	23
1963	90,464	39	40	76	164	347	1,116	3,046	6,852	20,014	36,542	22,204	24
1962	88,718	38	45	111	154	332	1,170	3,106	7,041	19,897	35,243	21,560	21
1961	89,099	47	50	94	153	330	1,128	3,101	7,040	20,581	35,446	21,112	17
1960	88,567	44	36	85	148	344	1,109	3,166	7,544	20,806	35,271	19,992	22

Note: Figures by race exclude data for residents of New Jersey for 1962 and 1963.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290A for 1960-67 and separate unpublished tables for the black population and Accidents.

77/09/02 TABLE 290A. DEATHS FOR 60 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1960-67

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	TOTAL	UNDER 1 YEAR	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS	65-74 YEARS	75-84 YEARS	85+ YEARS	N.S.
ICD-7 Nos. 330-334 Cerebrovascular diseases													
ALL OTHER, BOTH SEXES													
1967	25,423	27	11	46	99	381	1,317	2,988	5,102	7,755	5,333	2,332	32
1966	26,354	24	29	57	128	380	1,388	2,999	5,203	8,080	5,604	2,430	32
1965	26,466	35	36	56	106	350	1,377	3,164	5,621	8,114	5,421	2,158	28
1964	25,611	38	31	52	95	371	1,343	3,147	5,815	7,488	5,192	1,997	42
1963	25,515	38	24	46	88	325	1,309	3,043	5,697	7,674	5,226	2,013	32
1962	24,895	34	28	53	92	351	1,330	3,007	5,704	7,406	4,850	2,013	27
1961	24,214	35	30	46	94	353	1,291	3,050	5,642	7,227	4,491	1,923	32
1960	24,535	49	38	45	105	382	1,248	3,181	5,867	7,235	4,494	1,850	41
ALL OTHER, MALE													
1967	12,139	13	6	23	45	163	591	1,452	2,562	3,855	2,462	953	14
1966	12,671	15	13	30	57	163	646	1,514	2,666	3,959	2,629	965	14
1965	12,697	22	18	29	47	145	664	1,552	2,806	3,986	2,560	850	18
1964	12,216	20	13	30	40	157	591	1,501	2,850	3,699	2,465	834	16
1963	12,186	17	14	29	49	139	554	1,463	2,759	3,805	2,472	868	17
1962	11,815	19	15	29	40	133	584	1,466	2,764	3,646	2,266	847	6
1961	11,387	21	16	22	38	143	594	1,426	2,656	3,505	2,099	854	13
1960	11,777	29	21	19	56	150	529	1,533	2,823	3,593	2,166	839	19
ALL OTHER, FEMALE													
1967	13,284	14	5	23	54	218	726	1,536	2,540	3,900	2,871	1,379	18
1966	13,683	9	16	27	71	217	742	1,485	2,537	4,121	2,975	1,465	18
1965	13,769	13	18	27	59	205	713	1,612	2,815	4,128	2,861	1,308	10
1964	13,395	18	18	22	55	214	752	1,646	2,965	3,789	2,727	1,163	26
1963	13,329	21	10	17	39	186	755	1,580	2,938	3,869	2,754	1,145	15
1962	13,080	15	13	24	52	218	746	1,541	2,940	3,760	2,584	1,166	21
1961	12,827	14	14	24	56	210	697	1,624	2,986	3,722	2,392	1,069	19
1960	12,758	20	17	26	49	232	719	1,648	3,044	3,642	2,328	1,011	22

Note: Figures by race exclude data for residents of New Jersey for 1962 and 1963.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290A for 1960-67 and separate unpublished tables for the black population and Accidents.

77/09/02 TABLE 290A. DEATHS FOR 60 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1960-67

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	TOTAL	UNDER 1 YEAR	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS	65-74 YEARS	75-84 YEARS	85+ YEARS	N.S.
ICD-7 Nos. 400-402, 410-416 Diseases of heart													
BOTH SEXES													
1967	721,268	405	168	341	840	2,833	17,083	56,936	120,797	194,969	218,219	108,483	194
1966	727,002	387	203	333	918	3,078	17,359	57,670	121,995	198,900	219,322	106,625	212
1965	712,087	370	213	346	933	3,118	17,734	56,990	119,484	195,032	214,470	103,205	192
1964	699,861	382	239	344	969	3,137	18,151	56,617	118,644	195,112	207,315	98,750	201
1963	707,830	311	203	391	993	3,288	18,433	57,255	118,709	199,296	208,902	99,840	209
1962	688,009	281	178	384	946	3,297	18,027	56,708	115,817	195,545	201,209	95,422	195
1961	663,391	292	188	370	982	3,341	17,854	55,780	113,321	190,032	191,747	89,274	210
1960	661,712	272	207	453	951	3,566	17,961	55,678	114,912	191,401	189,483	86,586	242
MALE													
1967	415,851	234	89	179	484	1,817	12,785	43,552	86,810	118,832	108,497	42,465	107
1966	420,061	208	110	183	521	1,958	12,988	43,955	87,669	121,148	109,297	41,900	124
1965	412,831	204	113	178	508	2,004	13,243	43,383	85,211	119,063	107,761	41,042	121
1964	405,892	205	132	163	523	2,015	13,618	42,884	83,919	119,178	104,098	39,036	121
1963	411,989	174	103	219	533	2,082	13,629	43,504	83,529	121,998	105,814	40,287	117
1962	400,672	155	88	193	527	2,119	13,412	43,139	81,433	119,128	101,407	38,951	120
1961	388,563	167	93	191	566	2,124	13,289	42,291	79,723	116,184	97,202	36,615	118
1960	388,205	163	116	245	505	2,246	13,247	42,432	80,406	117,230	96,011	35,463	141
FEMALE													
1967	305,417	171	79	162	356	1,016	4,298	13,384	33,987	76,137	109,722	66,018	87
1966	306,941	179	93	150	397	1,120	4,371	13,715	34,326	77,752	110,025	64,725	88
1965	299,256	166	100	168	425	1,114	4,491	13,607	34,273	75,969	106,709	62,163	71
1964	293,969	177	107	181	446	1,122	4,533	13,733	34,725	75,934	103,217	59,714	80
1963	295,841	137	100	172	460	1,206	4,804	13,751	35,180	77,298	103,088	59,553	92
1962	287,337	126	90	191	419	1,178	4,615	13,569	34,384	76,417	99,802	56,471	75
1961	274,828	125	95	179	416	1,217	4,565	13,489	33,598	73,848	94,545	52,659	92
1960	273,507	109	91	208	446	1,320	4,714	13,246	34,506	74,171	93,472	51,123	101

Note: Figures by race exclude data for residents of New Jersey for 1962 and 1963.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290A for 1960-67 and separate unpublished tables for the black population and Accidents.

77/09/02 TABLE 290A. DEATHS FOR 60 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1960-67

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	TOTAL	UNDER 1 YEAR	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS	65-74 YEARS	75-84 YEARS	85+ YEARS	N.S.
ICD-7 Nos. 400-402, 410-416 Diseases of heart													
WHITE, BOTH SEXES													
1967	655,386	292	105	238	579	1,884	13,459	48,403	106,063	175,868	205,448	102,928	119
1966	659,554	245	121	225	657	2,105	13,665	49,054	106,709	179,281	206,274	101,090	128
1965	647,070	259	141	233	646	2,180	13,990	48,469	104,323	176,649	202,110	97,952	118
1964	635,672	259	151	225	694	2,211	14,403	48,040	103,157	176,934	195,642	93,832	124
1963	616,272	226	115	272	663	2,224	14,048	46,344	98,862	172,893	189,184	91,297	144
1962	599,503	192	121	257	635	2,272	13,796	46,090	96,528	169,872	182,466	87,143	131
1961	603,735	205	122	258	741	2,422	14,386	47,432	98,868	173,105	181,268	84,794	134
1960	602,876	196	129	311	682	2,586	14,501	47,343	100,398	174,809	179,346	82,423	152
WHITE, MALE													
1967	379,955	173	52	127	353	1,293	10,639	38,428	78,235	108,534	102,039	40,017	65
1966	383,251	135	62	132	387	1,413	10,778	38,767	78,781	110,484	102,805	39,433	74
1965	377,420	144	77	127	356	1,481	11,069	38,295	76,614	108,941	101,555	38,686	75
1964	371,170	136	82	109	411	1,532	11,414	37,866	75,197	109,231	98,243	36,877	72
1963	361,705	127	59	163	384	1,502	11,010	36,731	71,874	107,119	95,970	36,684	82
1962	352,160	104	60	132	388	1,543	10,883	36,466	70,155	104,859	92,023	35,464	83
1961	356,360	114	60	133	444	1,642	11,332	37,507	71,810	106,743	91,885	34,611	79
1960	356,267	123	71	172	381	1,749	11,354	37,663	72,338	108,056	90,730	33,541	89
WHITE, FEMALE													
1967	275,431	119	53	111	226	591	2,820	9,975	27,828	67,334	103,409	62,911	54
1966	276,303	110	59	93	270	692	2,887	10,287	27,928	68,797	103,469	61,657	54
1965	269,650	115	64	106	290	699	2,921	10,174	27,709	67,708	100,555	59,266	43
1964	264,502	123	69	116	283	679	2,989	10,174	27,960	67,703	97,399	56,955	52
1963	254,567	99	56	109	279	722	3,038	9,613	26,988	65,774	93,214	54,613	62
1962	247,343	88	61	125	247	729	2,913	9,624	26,373	65,013	90,443	51,679	48
1961	247,375	91	62	125	297	780	3,054	9,925	27,058	66,362	89,383	50,183	55
1960	246,609	73	58	139	301	837	3,147	9,680	28,060	66,753	88,616	48,882	63

Note: Figures by race exclude data for residents of New Jersey for 1962 and 1963.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290A for 1960-67 and separate unpublished tables for the black population and Accidents.

77/09/02 TABLE 290A. DEATHS FOR 60 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1960-67

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	TOTAL	UNDER 1 YEAR	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS	65-74 YEARS	75-84 YEARS	85+ YEARS	N.S.
ICD-7 Nos. 400-402, 410-416 Diseases of heart													
ALL OTHER, BOTH SEXES													
1967	65,882	113	63	103	261	949	3,624	8,533	14,734	19,101	12,771	5,555	75
1966	67,448	142	82	108	261	973	3,694	8,616	15,286	19,619	13,048	5,535	84
1965	65,017	111	72	113	287	938	3,744	8,521	15,161	18,383	12,360	5,253	74
1964	64,189	123	88	119	275	926	3,748	8,577	15,487	18,178	11,673	4,918	77
1963	63,535	80	86	108	298	910	3,608	8,511	15,091	18,190	11,660	4,937	56
1962	61,082	83	54	116	284	895	3,514	8,344	14,542	17,505	11,007	4,679	59
1961	59,656	87	66	112	241	919	3,468	8,348	14,453	16,927	10,479	4,480	76
1960	58,836	76	78	142	269	980	3,460	8,335	14,514	16,592	10,137	4,163	90
ALL OTHER, MALE													
1967	35,896	61	37	52	131	524	2,146	5,124	8,575	10,298	6,458	2,448	42
1966	36,810	73	48	51	134	545	2,210	5,188	8,888	10,664	6,492	2,467	50
1965	35,411	60	36	51	152	523	2,174	5,088	8,597	10,122	6,206	2,356	46
1964	34,722	69	50	54	112	483	2,204	5,018	8,722	9,947	5,855	2,159	49
1963	34,502	44	43	49	133	481	2,035	4,969	8,376	10,071	6,014	2,258	29
1962	33,144	49	26	56	120	479	2,006	4,930	8,013	9,577	5,694	2,160	34
1961	32,203	53	33	58	122	482	1,957	4,784	7,913	9,441	5,317	2,004	39
1960	31,938	40	45	73	124	497	1,893	4,769	8,068	9,174	5,281	1,922	52
ALL OTHER, FEMALE													
1967	29,986	52	26	51	130	425	1,478	3,409	6,159	8,803	6,313	3,107	33
1966	30,638	69	34	57	127	428	1,484	3,428	6,398	8,955	6,556	3,068	34
1965	29,606	51	36	62	135	415	1,570	3,433	6,564	8,261	6,154	2,897	28
1964	29,467	54	38	65	163	443	1,544	3,559	6,765	8,231	5,818	2,759	28
1963	29,033	36	43	59	165	429	1,573	3,542	6,715	8,119	5,646	2,679	27
1962	27,938	34	28	60	164	416	1,508	3,414	6,529	7,928	5,313	2,519	25
1961	27,453	34	33	54	119	437	1,511	3,564	6,540	7,486	5,162	2,476	37
1960	26,898	36	33	69	145	483	1,567	3,566	6,446	7,418	4,856	2,241	38

Note: Figures by race exclude data for residents of New Jersey for 1962 and 1963.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290A for 1960-67 and separate unpublished tables for the black population and Accidents.

77/09/02 TABLE 290A. DEATHS FOR 60 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1960-67

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	TOTAL	UNDER 1 YEAR	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS	65-74 YEARS	75-84 YEARS	85+ YEARS	N.S.
Rheumatic fever and chronic rheumatic heart disease (400-402,410-416)													
BOTH SEXES													
1967	14,176	2	12	109	265	546	1,421	2,752	3,823	3,216	1,634	393	3
1966	15,012	4	21	122	273	672	1,643	3,017	3,989	3,205	1,657	409	-
1965	15,471	11	14	133	350	716	1,758	3,199	4,101	3,189	1,597	399	4
1964	15,923	4	31	133	352	738	1,985	3,467	4,233	3,133	1,470	376	1
1963	16,619	3	27	183	405	809	2,101	3,625	4,237	3,214	1,592	421	2
1962	17,751	8	23	166	445	887	2,288	3,901	4,482	3,401	1,701	449	-
1961	17,872	6	26	197	460	999	2,293	3,926	4,361	3,470	1,672	462	-
1960	18,411	3	32	232	474	1,085	2,458	4,036	4,470	3,420	1,702	496	3
MALE													
1967	5,988	1	7	52	124	281	643	1,265	1,647	1,296	558	113	1
1966	6,451	2	10	56	144	315	788	1,353	1,754	1,336	580	113	-
1965	6,770	6	7	64	165	317	845	1,450	1,830	1,363	596	125	2
1964	7,100	2	16	60	155	352	987	1,628	1,899	1,328	554	119	-
1963	7,723	1	14	99	196	400	1,027	1,768	2,016	1,456	610	135	1
1962	8,166	5	11	82	213	421	1,154	1,865	2,094	1,497	681	143	-
1961	8,361	5	13	86	229	489	1,115	1,929	2,069	1,606	681	139	-
1960	8,783	1	16	118	221	529	1,234	2,005	2,189	1,588	713	167	2
FEMALE													
1967	8,188	1	5	57	141	265	778	1,487	2,176	1,920	1,076	280	2
1966	8,561	2	11	66	129	357	855	1,664	2,235	1,869	1,077	296	-
1965	8,701	5	7	69	185	399	913	1,749	2,271	1,826	1,001	274	2
1964	8,823	2	15	73	197	386	998	1,839	2,334	1,805	916	257	1
1963	8,896	2	13	84	209	409	1,074	1,857	2,221	1,758	982	286	1
1962	9,585	3	12	84	232	466	1,134	2,036	2,388	1,904	1,020	306	-
1961	9,511	1	13	111	231	510	1,178	1,997	2,292	1,864	991	323	-
1960	9,628	2	16	114	253	556	1,224	2,031	2,281	1,832	989	329	1

- Quantity zero

Note: Figures by race exclude data for residents of New Jersey for 1962 and 1963.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290A for 1960-67 and separate unpublished tables for the black population and Accidents.

77/09/02 TABLE 290A. DEATHS FOR 60 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1960-67

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	TOTAL	UNDER 1 YEAR	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS	65-74 YEARS	75-84 YEARS	85+ YEARS	N.S.
Rheumatic fever and chronic rheumatic heart disease (400-402,410-416)													
WHITE, BOTH SEXES													
1967	12,909	2	6	55	181	421	1,174	2,475	3,592	3,051	1,582	367	3
1966	13,712	2	10	63	186	527	1,394	2,744	3,750	3,059	1,591	386	-
1965	14,124	8	6	72	224	548	1,507	2,925	3,870	3,030	1,542	390	2
1964	14,564	2	17	78	245	573	1,710	3,200	3,973	3,002	1,407	356	1
1963	14,525	1	12	109	255	622	1,739	3,134	3,811	2,957	1,482	402	1
1962	15,514	3	17	95	309	679	1,942	3,397	3,991	3,083	1,580	418	-
1961	16,381	3	12	123	353	815	2,031	3,613	4,091	3,298	1,607	435	-
1960	16,890	3	19	136	350	877	2,189	3,764	4,228	3,242	1,621	459	2
WHITE, MALE													
1967	5,426	1	3	28	91	223	537	1,136	1,547	1,225	533	101	1
1966	5,859	1	5	32	101	250	672	1,222	1,638	1,275	555	108	-
1965	6,163	3	3	41	101	251	737	1,321	1,715	1,291	577	122	1
1964	6,480	-	8	34	122	300	850	1,504	1,773	1,258	518	113	-
1963	6,766	-	6	63	132	325	868	1,540	1,810	1,334	563	125	-
1962	7,187	1	9	52	160	327	1,000	1,633	1,870	1,363	640	132	-
1961	7,659	3	6	52	181	411	1,002	1,772	1,938	1,512	651	131	-
1960	8,038	1	10	70	170	436	1,109	1,868	2,066	1,491	666	150	1
WHITE, FEMALE													
1967	7,483	1	3	27	90	198	637	1,339	2,045	1,826	1,049	266	2
1966	7,853	1	5	31	85	277	722	1,522	2,112	1,784	1,036	278	-
1965	7,961	5	3	31	123	297	770	1,604	2,155	1,739	965	268	1
1964	8,084	2	9	44	123	273	860	1,696	2,200	1,744	889	243	1
1963	7,759	1	6	46	123	297	871	1,594	2,001	1,623	919	277	1
1962	8,327	2	8	43	149	352	942	1,764	2,121	1,720	940	286	-
1961	8,722	-	6	71	172	404	1,029	1,841	2,153	1,786	956	304	-
1960	8,852	2	9	66	180	441	1,080	1,896	2,162	1,751	955	309	1

- Quantity zero

Note: Figures by race exclude data for residents of New Jersey for 1962 and 1963.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290A for 1960-67 and separate unpublished tables for the black population and Accidents.

77/09/02 TABLE 290A. DEATHS FOR 60 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1960-67

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	TOTAL	UNDER 1 YEAR	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS	65-74 YEARS	75-84 YEARS	85+ YEARS	N.S.
Rheumatic fever and chronic rheumatic heart disease (400-402,410-416)													
ALL OTHER, BOTH SEXES													
1967	1,267	-	6	54	84	125	247	277	231	165	52	26	-
1966	1,300	2	11	59	87	145	249	273	239	146	66	23	-
1965	1,347	3	8	61	126	168	251	274	231	159	55	9	2
1964	1,359	2	14	55	107	165	275	267	260	131	63	20	-
1963	1,369	2	14	68	134	151	265	281	232	139	66	16	1
1962	1,404	4	6	68	125	175	236	287	258	173	57	15	-
1961	1,491	3	14	74	107	184	262	313	270	172	65	27	-
1960	1,521	-	13	96	124	208	269	272	242	178	81	37	1
ALL OTHER, MALE													
1967	562	-	4	24	33	58	106	129	100	71	25	12	-
1966	592	1	5	24	43	65	116	131	116	61	25	5	-
1965	607	3	4	23	64	66	108	129	115	72	19	3	1
1964	620	2	8	26	33	52	137	124	126	70	36	6	-
1963	647	1	8	31	55	59	119	138	121	75	30	9	1
1962	635	4	2	29	45	72	99	136	125	93	23	7	-
1961	702	2	7	34	48	78	113	157	131	94	30	8	-
1960	745	-	6	48	51	93	125	137	123	97	47	17	1
ALL OTHER, FEMALE													
1967	705	-	2	30	51	67	141	148	131	94	27	14	-
1966	708	1	6	35	44	80	133	142	123	85	41	18	-
1965	740	-	4	38	62	102	143	145	116	87	36	6	1
1964	739	-	6	29	74	113	138	143	134	61	27	14	-
1963	722	1	6	37	79	92	146	143	111	64	36	7	-
1962	769	-	4	39	80	103	137	151	133	80	34	8	-
1961	789	1	7	40	59	106	149	156	139	78	35	19	-
1960	776	-	7	48	73	115	144	135	119	81	34	20	-

- Quantity zero

Note: Figures by race exclude data for residents of New Jersey for 1962 and 1963.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290A for 1960-67 and separate unpublished tables for the black population and Accidents.

77/09/02 TABLE 290A. DEATHS FOR 60 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1960-67

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	TOTAL	UNDER 1 YEAR	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS	65-74 YEARS	75-84 YEARS	85+ YEARS	N.S.
Arteriosclerotic heart disease, including coronary disease (420)													
BOTH SEXES													
1967	573,153	12	4	27	170	1,420	12,513	45,621	98,934	159,909	173,011	81,379	153
1966	573,191	18	8	25	199	1,469	12,522	45,997	99,647	162,555	171,737	78,854	160
1965	559,293	20	12	23	173	1,562	12,754	45,464	97,353	158,820	167,080	75,891	141
1964	545,499	12	13	28	213	1,540	13,039	44,793	96,198	158,004	160,129	71,376	154
1963	546,813	9	6	28	177	1,594	13,024	45,005	95,632	160,189	159,575	71,415	159
1962	527,561	14	13	19	132	1,582	12,502	44,079	92,915	156,066	152,274	67,833	132
1961	502,351	15	5	21	156	1,474	12,266	42,840	89,847	150,244	143,083	62,253	147
1960	494,282	12	9	20	143	1,548	12,148	42,557	90,467	149,124	138,913	59,180	161
MALE													
1967	345,154	9	3	16	113	1,058	10,284	36,923	74,186	100,689	88,898	32,888	87
1966	346,855	10	5	17	136	1,140	10,286	37,277	74,729	102,401	88,600	32,158	96
1965	340,077	12	5	16	109	1,235	10,529	36,825	72,549	100,455	87,094	31,156	92
1964	332,676	8	3	10	146	1,198	10,855	36,238	71,364	100,012	83,474	29,267	101
1963	334,974	6	5	19	119	1,208	10,707	36,621	70,643	101,842	83,855	29,853	96
1962	324,183	8	7	11	89	1,242	10,385	35,919	68,662	99,041	79,960	28,773	86
1961	311,189	9	3	10	114	1,175	10,310	35,005	66,577	95,856	75,506	26,536	88
1960	307,377	7	6	14	91	1,222	10,132	35,036	66,711	95,326	73,624	25,108	100
FEMALE													
1967	227,999	3	1	11	57	362	2,229	8,698	24,748	59,220	84,113	48,491	66
1966	226,336	8	3	8	63	329	2,236	8,720	24,918	60,154	83,137	46,696	64
1965	219,216	8	7	7	64	327	2,225	8,639	24,804	58,365	79,986	44,735	49
1964	212,823	4	10	18	67	342	2,184	8,555	24,834	57,992	76,655	42,109	53
1963	211,839	3	1	9	58	386	2,317	8,384	24,989	58,347	75,720	41,562	63
1962	203,378	6	6	8	43	340	2,117	8,160	24,253	57,025	72,314	39,060	46
1961	191,162	6	2	11	42	299	1,956	7,835	23,270	54,388	67,577	35,717	59
1960	186,905	5	3	6	52	326	2,016	7,521	23,756	53,798	65,289	34,072	61

Note: Figures by race exclude data for residents of New Jersey for 1962 and 1963.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290A for 1960-67 and separate unpublished tables for the black population and Accidents.

77/09/02 TABLE 290A. DEATHS FOR 60 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1960-67

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	TOTAL	UNDER 1 YEAR	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS	65-74 YEARS	75-84 YEARS	85+ YEARS	N.S.
Arteriosclerotic heart disease, including coronary disease (420)													
WHITE, BOTH SEXES													
1967	531,322	8	4	19	117	1,014	10,535	40,406	89,404	147,382	164,592	77,734	107
1966	530,782	9	6	22	151	1,101	10,578	40,843	89,872	149,649	163,197	75,247	107
1965	518,737	15	7	15	132	1,175	10,744	40,305	87,773	146,920	159,070	72,487	94
1964	506,356	10	8	18	164	1,163	11,072	39,679	86,561	146,500	152,786	68,293	102
1963	487,193	6	4	21	127	1,153	10,689	38,240	82,617	142,243	146,195	65,779	119
1962	470,711	8	9	10	89	1,187	10,276	37,591	80,275	138,969	139,726	62,469	102
1961	468,311	11	4	17	119	1,146	10,636	38,309	81,375	140,189	136,859	59,541	105
1960	461,419	9	5	19	102	1,220	10,483	37,924	82,080	139,517	133,124	56,820	116
WHITE, MALE													
1967	321,241	5	3	10	78	808	8,992	33,587	68,329	93,648	84,469	31,251	61
1966	322,528	5	5	15	102	907	9,004	33,968	68,741	95,033	84,189	30,495	64
1965	316,770	8	1	11	83	970	9,207	33,515	66,857	93,575	82,929	29,552	62
1964	310,244	8	2	7	119	960	9,552	32,994	65,580	93,451	79,655	27,851	65
1963	300,246	4	3	17	90	924	9,108	31,989	62,460	91,230	76,932	27,416	73
1962	291,009	4	7	5	66	975	8,875	31,427	60,717	89,013	73,392	26,461	67
1961	291,614	7	2	8	87	964	9,256	32,165	61,610	89,952	72,222	25,275	66
1960	288,376	6	4	13	69	1,028	9,086	32,130	61,744	89,780	70,455	23,987	74
WHITE, FEMALE													
1967	210,081	3	1	9	39	206	1,543	6,819	21,075	53,734	80,123	46,483	46
1966	208,254	4	1	7	49	194	1,574	6,875	21,131	54,616	79,008	44,752	43
1965	201,967	7	6	4	49	205	1,537	6,790	20,916	53,345	76,141	42,935	32
1964	196,112	2	6	11	45	203	1,520	6,685	20,981	53,049	73,131	40,442	37
1963	186,947	2	1	4	37	229	1,581	6,251	20,157	51,013	69,263	38,363	46
1962	179,702	4	2	5	23	212	1,401	6,164	19,558	49,956	66,334	36,008	35
1961	176,697	4	2	9	32	182	1,380	6,144	19,765	50,237	64,637	34,266	39
1960	173,043	3	1	6	33	192	1,397	5,794	20,336	49,737	62,669	32,833	42

Note: Figures by race exclude data for residents of New Jersey for 1962 and 1963.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290A for 1960-67 and separate unpublished tables for the black population and Accidents.

77/09/02 TABLE 290A. DEATHS FOR 60 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1960-67

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	TOTAL	UNDER 1 YEAR	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS	65-74 YEARS	75-84 YEARS	85+ YEARS	N.S.
Arteriosclerotic heart disease, including coronary disease (420)													
ALL OTHER, BOTH SEXES													
1967	41,831	4	-	8	53	406	1,978	5,215	9,530	12,527	8,419	3,645	46
1966	42,409	9	2	3	48	368	1,944	5,154	9,775	12,906	8,540	3,607	53
1965	40,556	5	5	8	41	387	2,010	5,159	9,580	11,900	8,010	3,404	47
1964	39,143	2	5	10	49	377	1,967	5,114	9,637	11,504	7,343	3,083	52
1963	37,703	2	2	6	45	366	1,783	4,877	9,143	11,251	7,175	3,020	33
1962	35,725	6	4	9	37	326	1,725	4,742	8,763	10,549	6,714	2,824	26
1961	34,040	4	1	4	37	328	1,630	4,531	8,472	10,055	6,224	2,712	42
1960	32,863	3	4	1	41	328	1,665	4,633	8,387	9,607	5,789	2,360	45
ALL OTHER, MALE													
1967	23,913	4	-	6	35	250	1,292	3,336	5,857	7,041	4,429	1,637	26
1966	24,327	5	-	2	34	233	1,282	3,309	5,988	7,368	4,411	1,663	32
1965	23,307	4	4	5	26	265	1,322	3,310	5,692	6,880	4,165	1,604	30
1964	22,432	-	1	3	27	238	1,303	3,244	5,784	6,561	3,819	1,416	36
1963	21,774	1	2	2	26	227	1,134	3,109	5,385	6,539	3,890	1,441	18
1962	20,655	4	-	6	18	208	1,113	3,046	5,142	6,105	3,663	1,333	17
1961	19,575	2	1	2	27	211	1,054	2,840	4,967	5,904	3,284	1,261	22
1960	19,001	1	2	1	22	194	1,046	2,906	4,967	5,546	3,169	1,121	26
ALL OTHER, FEMALE													
1967	17,918	-	-	2	18	156	686	1,879	3,673	5,486	3,990	2,008	20
1966	18,082	4	2	1	14	135	662	1,845	3,787	5,538	4,129	1,944	21
1965	17,249	1	1	3	15	122	688	1,849	3,888	5,020	3,845	1,800	17
1964	16,711	2	4	7	22	139	664	1,870	3,853	4,943	3,524	1,667	16
1963	15,929	1	-	4	19	139	649	1,768	3,758	4,712	3,285	1,579	15
1962	15,070	2	4	3	19	118	612	1,696	3,621	4,444	3,051	1,491	9
1961	14,465	2	-	2	10	117	576	1,691	3,505	4,151	2,940	1,451	20
1960	13,862	2	2	-	19	134	619	1,727	3,420	4,061	2,620	1,239	19

- Quantity zero

Note: Figures by race exclude data for residents of New Jersey for 1962 and 1963.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290A for 1960-67 and separate unpublished tables for the black population and Accidents.

77/09/02 TABLE 290A. DEATHS FOR 60 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1960-67

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	TOTAL	UNDER 1 YEAR	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS	65-74 YEARS	75-84 YEARS	85+ YEARS	N.S.
Nonrheumatic chronic endocarditis and other myocardial degeneration (421,422)													
BOTH SEXES													
1967	52,697	39	26	38	104	216	705	2,213	5,453	11,279	18,867	13,748	9
1966	53,581	41	35	36	103	199	674	2,110	5,129	11,354	19,749	14,131	20
1965	53,178	34	37	44	93	167	670	1,969	5,074	11,193	19,743	14,142	12
1964	53,255	45	40	31	84	169	602	1,944	5,141	11,564	19,596	14,023	16
1963	55,978	24	33	43	91	182	638	1,872	5,165	12,287	20,809	14,814	20
1962	54,645	35	35	39	88	169	604	1,954	5,020	12,216	20,258	14,203	24
1961	54,632	44	33	33	80	178	611	2,063	5,205	12,294	20,049	14,019	23
1960	57,029	31	39	49	86	195	616	1,997	5,330	13,152	21,122	14,383	29
MALE													
1967	25,887	25	13	22	75	135	507	1,553	3,567	6,340	8,613	5,032	5
1966	26,122	18	17	23	67	119	468	1,431	3,373	6,345	9,113	5,139	9
1965	26,050	17	22	26	60	101	440	1,394	3,290	6,233	9,166	5,296	5
1964	26,112	21	19	17	58	106	400	1,306	3,287	6,469	9,241	5,180	8
1963	27,617	14	17	27	56	97	441	1,248	3,282	6,847	9,923	5,655	10
1962	26,915	22	16	20	61	107	402	1,322	3,152	6,776	9,531	5,492	14
1961	27,122	21	17	24	54	104	381	1,412	3,271	6,841	9,548	5,435	14
1960	28,295	19	21	28	55	122	388	1,316	3,353	7,361	9,949	5,666	17
FEMALE													
1967	26,810	14	13	16	29	81	198	660	1,886	4,939	10,254	8,716	4
1966	27,459	23	18	13	36	80	206	679	1,756	5,009	10,636	8,992	11
1965	27,128	17	15	18	33	66	230	575	1,784	4,960	10,577	8,846	7
1964	27,143	24	21	14	26	63	202	638	1,854	5,095	10,355	8,843	8
1963	28,361	10	16	16	35	85	197	624	1,883	5,440	10,886	9,159	10
1962	27,730	13	19	19	27	62	202	632	1,868	5,440	10,727	8,711	10
1961	27,510	23	16	9	26	74	230	651	1,934	5,453	10,501	8,584	9
1960	28,734	12	18	21	31	73	228	681	1,977	5,791	11,173	8,717	12

Note: Figures by race exclude data for residents of New Jersey for 1962 and 1963.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290A for 1960-67 and separate unpublished tables for the black population and Accidents.

77/09/02 TABLE 290A. DEATHS FOR 60 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1960-67

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	TOTAL	UNDER 1 YEAR	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS	65-74 YEARS	75-84 YEARS	85+ YEARS	N.S.
Nonrheumatic chronic endocarditis and other myocardial degeneration (421,422)													
WHITE, BOTH SEXES													
1967	46,886	25	16	30	76	127	499	1,749	4,409	9,537	17,429	12,988	1
1966	47,846	24	23	26	71	128	479	1,621	4,127	9,657	18,288	13,393	9
1965	47,648	21	24	38	72	112	489	1,514	4,056	9,577	18,328	13,411	6
1964	47,561	32	22	26	66	110	421	1,490	4,033	9,856	18,195	13,302	8
1963	47,388	17	19	38	57	113	384	1,287	3,699	9,924	18,409	13,431	10
1962	46,065	20	24	33	63	102	377	1,332	3,634	9,785	17,864	12,820	11
1961	48,771	34	25	27	60	109	418	1,528	4,011	10,465	18,746	13,335	13
1960	51,069	22	18	39	60	115	436	1,484	4,096	11,381	19,749	13,653	16
WHITE, MALE													
1967	22,886	19	9	17	58	81	367	1,278	2,994	5,444	7,908	4,710	1
1966	23,113	10	10	15	50	79	346	1,151	2,800	5,428	8,423	4,799	2
1965	23,202	11	14	21	46	72	340	1,112	2,738	5,386	8,458	5,002	2
1964	23,208	16	11	15	50	66	303	1,054	2,693	5,570	8,544	4,883	3
1963	23,196	10	11	26	37	64	280	881	2,447	5,591	8,759	5,086	4
1962	22,570	11	8	18	48	66	261	937	2,383	5,520	8,396	4,915	7
1961	24,002	13	13	20	41	68	275	1,077	2,620	5,833	8,905	5,128	9
1960	25,189	15	7	22	42	81	296	1,047	2,687	6,399	9,244	5,340	9
WHITE, FEMALE													
1967	24,000	6	7	13	18	46	132	471	1,415	4,093	9,521	8,278	-
1966	24,733	14	13	11	21	49	133	470	1,327	4,229	9,865	8,594	7
1965	24,446	10	10	17	26	40	149	402	1,318	4,191	9,870	8,409	4
1964	24,353	16	11	11	16	44	118	436	1,340	4,286	9,651	8,419	5
1963	24,192	7	8	12	20	49	104	406	1,252	4,333	9,650	8,345	6
1962	23,495	9	16	15	15	36	116	395	1,251	4,265	9,468	7,905	4
1961	24,769	21	12	7	19	41	143	451	1,391	4,632	9,841	8,207	4
1960	25,880	7	11	17	18	34	140	437	1,409	4,982	10,505	8,313	7

- Quantity zero

Note: Figures by race exclude data for residents of New Jersey for 1962 and 1963.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290A for 1960-67 and separate unpublished tables for the black population and Accidents.

77/09/02 TABLE 290A. DEATHS FOR 60 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1960-67

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	TOTAL	UNDER 1 YEAR	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS	65-74 YEARS	75-84 YEARS	85+ YEARS	N.S.
Nonrheumatic chronic endocarditis and other myocardial degeneration (421,422)													
ALL OTHER, BOTH SEXES													
1967	5,811	14	10	8	28	89	206	464	1,044	1,742	1,438	760	8
1966	5,735	17	12	10	32	71	195	489	1,002	1,697	1,461	738	11
1965	5,530	13	13	6	21	55	181	455	1,018	1,616	1,415	731	6
1964	5,694	13	18	5	18	59	181	454	1,108	1,708	1,401	721	8
1963	6,009	7	14	5	32	50	189	466	1,225	1,774	1,465	774	8
1962	5,825	15	11	5	19	58	176	482	1,111	1,775	1,417	743	13
1961	5,861	10	8	6	20	69	193	535	1,194	1,829	1,303	684	10
1960	5,960	9	21	10	26	80	180	513	1,234	1,771	1,373	730	13
ALL OTHER, MALE													
1967	3,001	6	4	5	17	54	140	275	573	896	705	322	4
1966	3,009	8	7	8	17	40	122	280	573	917	690	340	7
1965	2,848	6	8	5	14	29	100	282	552	847	708	294	3
1964	2,904	5	8	2	8	40	97	252	594	899	697	297	5
1963	3,136	4	6	1	18	22	113	282	681	945	715	344	5
1962	2,981	11	8	1	9	33	104	278	596	910	686	338	7
1961	3,120	8	4	4	13	36	106	335	651	1,008	643	307	5
1960	3,106	4	14	6	13	41	92	269	666	962	705	326	8
ALL OTHER, FEMALE													
1967	2,810	8	6	3	11	35	66	189	471	846	733	438	4
1966	2,726	9	5	2	15	31	73	209	429	780	771	398	4
1965	2,682	7	5	1	7	26	81	173	466	769	707	437	3
1964	2,790	8	10	3	10	19	84	202	514	809	704	424	3
1963	2,873	3	8	4	14	28	76	184	544	829	750	430	3
1962	2,844	4	3	4	10	25	72	204	515	865	731	405	6
1961	2,741	2	4	2	7	33	87	200	543	821	660	377	5
1960	2,854	5	7	4	13	39	88	244	568	809	668	404	5

Note: Figures by race exclude data for residents of New Jersey for 1962 and 1963.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290A for 1960-67 and separate unpublished tables for the black population and Accidents.

77/09/02 TABLE 290A. DEATHS FOR 60 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1960-67

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	TOTAL	UNDER 1 YEAR	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS	65-74 YEARS	75-84 YEARS	85+ YEARS	N.S.
Other diseases of heart (430-434)													
BOTH SEXES													
1967	31,267	351	126	163	274	433	1,252	2,644	5,059	7,326	8,691	4,936	12
1966	31,042	323	137	145	311	504	1,231	2,678	4,886	7,312	8,660	4,845	10
1965	29,177	305	150	143	284	445	1,186	2,477	4,567	7,058	8,166	4,383	13
1964	27,741	321	155	143	286	446	1,191	2,386	4,304	6,735	7,603	4,157	14
1963	27,323	275	137	132	279	440	1,199	2,516	4,372	6,660	7,405	3,898	10
1962	25,876	224	107	151	243	387	1,163	2,421	4,132	6,430	6,981	3,621	16
1961	25,254	226	123	116	260	430	1,222	2,492	4,130	6,166	6,737	3,344	8
1960	25,682	225	126	146	198	451	1,194	2,444	4,147	6,472	6,884	3,381	14
MALE													
1967	17,280	199	66	88	156	230	729	1,734	3,327	4,333	4,372	2,039	7
1966	17,154	178	77	85	159	258	727	1,743	3,247	4,333	4,395	1,947	5
1965	16,197	169	79	71	155	234	690	1,633	3,016	4,169	4,126	1,843	12
1964	15,449	174	94	72	146	227	680	1,586	2,836	4,044	3,838	1,744	8
1963	15,343	153	67	71	147	228	687	1,666	2,831	4,029	3,840	1,621	3
1962	14,754	120	54	79	143	204	733	1,663	2,728	3,901	3,579	1,540	10
1961	14,578	132	59	70	155	230	749	1,690	2,783	3,704	3,520	1,483	3
1960	14,893	135	73	82	112	228	746	1,667	2,788	4,038	3,602	1,413	9
FEMALE													
1967	13,987	152	60	75	118	203	523	910	1,732	2,993	4,319	2,897	5
1966	13,888	145	60	60	152	246	504	935	1,639	2,979	4,265	2,898	5
1965	12,980	136	71	72	129	211	496	844	1,551	2,889	4,040	2,540	1
1964	12,292	147	61	71	140	219	511	800	1,468	2,691	3,765	2,413	6
1963	11,980	122	70	61	132	212	512	850	1,541	2,631	3,565	2,277	7
1962	11,122	104	53	72	100	183	430	758	1,404	2,529	3,402	2,081	6
1961	10,676	94	64	46	105	200	473	802	1,347	2,462	3,217	1,861	5
1960	10,789	90	53	64	86	223	448	777	1,359	2,434	3,282	1,968	5

Note: Figures by race exclude data for residents of New Jersey for 1962 and 1963.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290A for 1960-67 and separate unpublished tables for the black population and Accidents.

77/09/02 TABLE 290A. DEATHS FOR 60 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1960-67

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	TOTAL	UNDER 1 YEAR	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS	65-74 YEARS	75-84 YEARS	85+ YEARS	N.S.
Other diseases of heart (430-434)													
WHITE, BOTH SEXES													
1967	26,060	256	79	130	190	255	850	1,936	3,897	6,019	7,867	4,579	2
1966	25,856	209	81	111	232	285	814	1,962	3,776	6,031	7,853	4,497	5
1965	24,350	215	104	106	207	280	782	1,800	3,457	5,871	7,456	4,064	8
1964	22,975	215	104	98	207	286	741	1,690	3,201	5,612	6,912	3,903	6
1963	21,990	202	80	100	207	267	758	1,736	3,214	5,344	6,529	3,547	6
1962	20,837	161	71	110	160	230	739	1,686	3,011	5,204	6,159	3,297	9
1961	20,924	157	80	90	196	265	807	1,721	3,185	5,181	6,123	3,118	1
1960	21,506	162	87	113	145	287	832	1,767	3,242	5,447	6,277	3,141	6
WHITE, MALE													
1967	14,408	148	37	71	113	144	509	1,320	2,631	3,611	3,952	1,871	1
1966	14,304	119	42	69	126	143	502	1,315	2,591	3,641	3,969	1,784	3
1965	13,515	122	59	53	120	147	491	1,250	2,369	3,479	3,734	1,684	7
1964	12,792	112	61	50	114	154	449	1,170	2,180	3,396	3,484	1,619	3
1963	12,429	113	39	54	117	144	465	1,206	2,187	3,279	3,373	1,450	2
1962	11,939	88	36	56	102	131	501	1,207	2,095	3,188	3,141	1,389	5
1961	12,162	91	38	53	128	152	525	1,236	2,236	3,133	3,192	1,378	-
1960	12,498	101	50	65	87	152	545	1,256	2,258	3,418	3,266	1,297	3
WHITE, FEMALE													
1967	11,652	108	42	59	77	111	341	616	1,266	2,408	3,915	2,708	1
1966	11,552	90	39	42	106	142	312	647	1,185	2,390	3,884	2,713	2
1965	10,835	93	45	53	87	133	291	550	1,088	2,392	3,722	2,380	1
1964	10,183	103	43	48	93	132	292	520	1,021	2,216	3,428	2,284	3
1963	9,561	89	41	46	90	123	293	530	1,027	2,065	3,156	2,097	4
1962	8,898	73	35	54	58	99	238	479	916	2,016	3,018	1,908	4
1961	8,762	66	42	37	68	113	282	485	949	2,048	2,931	1,740	1
1960	9,008	61	37	48	58	135	287	511	984	2,029	3,011	1,844	3

- Quantity zero

Note: Figures by race exclude data for residents of New Jersey for 1962 and 1963.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290A for 1960-67 and separate unpublished tables for the black population and Accidents.

77/09/02 TABLE 290A. DEATHS FOR 60 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1960-67

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	TOTAL	UNDER 1 YEAR	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS	65-74 YEARS	75-84 YEARS	85+ YEARS	N.S.
Other diseases of heart (430-434)													
ALL OTHER, BOTH SEXES													
1967	5,207	95	47	33	84	178	402	708	1,162	1,307	824	357	10
1966	5,186	114	56	34	79	219	417	716	1,110	1,281	807	348	5
1965	4,827	90	46	37	77	165	404	677	1,110	1,187	710	319	5
1964	4,766	106	51	45	79	160	450	696	1,103	1,123	691	254	8
1963	4,659	69	56	28	65	161	422	725	1,050	1,128	679	272	4
1962	4,442	58	33	34	80	148	402	687	1,043	1,061	644	245	7
1961	4,330	69	43	26	64	165	415	771	945	985	614	226	7
1960	4,176	63	39	33	53	164	362	677	905	1,025	607	240	8
ALL OTHER, MALE													
1967	2,872	51	29	17	43	86	220	414	696	722	420	168	6
1966	2,850	59	35	16	33	115	225	428	656	692	426	163	2
1965	2,682	47	20	18	35	87	199	383	647	690	392	159	5
1964	2,657	62	33	22	32	73	231	416	656	648	354	125	5
1963	2,571	38	27	15	27	76	214	426	580	646	380	141	1
1962	2,514	30	16	20	40	70	219	424	586	621	357	126	5
1961	2,416	41	21	17	27	78	224	454	547	571	328	105	3
1960	2,395	34	23	17	25	76	201	411	530	620	336	116	6
ALL OTHER, FEMALE													
1967	2,335	44	18	16	41	92	182	294	466	585	404	189	4
1966	2,336	55	21	18	46	104	192	288	454	589	381	185	3
1965	2,145	43	26	19	42	78	205	294	463	497	318	160	-
1964	2,109	44	18	23	47	87	219	280	447	475	337	129	3
1963	2,088	31	29	13	38	85	208	299	470	482	299	131	3
1962	1,928	28	17	14	40	78	183	263	457	440	287	119	2
1961	1,914	28	22	9	37	87	191	317	398	414	286	121	4
1960	1,781	29	16	16	28	88	161	266	375	405	271	124	2

- Quantity zero

Note: Figures by race exclude data for residents of New Jersey for 1962 and 1963.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290A for 1960-67 and separate unpublished tables for the black population and Accidents.

77/09/02 TABLE 290A. DEATHS FOR 60 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1960-67

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	TOTAL	UNDER 1 YEAR	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS	65-74 YEARS	75-84 YEARS	85+ YEARS	N.S.
Hypertensive heart disease (440-443)													
BOTH SEXES													
1967	49,975	1	-	4	27	218	1,192	3,706	7,528	13,239	16,016	8,027	17
1966	54,176	1	2	5	32	234	1,289	3,868	8,344	14,474	17,519	8,386	22
1965	54,968	-	-	3	33	228	1,366	3,881	8,389	14,772	17,884	8,390	22
1964	57,443	-	-	9	34	244	1,334	4,027	8,768	15,676	18,517	8,818	16
1963	61,097	-	-	5	41	263	1,471	4,237	9,303	16,946	19,521	9,292	18
1962	62,176	-	-	9	38	272	1,470	4,353	9,268	17,432	19,995	9,316	23
1961	63,282	1	1	3	26	260	1,462	4,459	9,778	17,858	20,206	9,196	32
1960	66,308	1	1	6	50	287	1,545	4,644	10,498	19,233	20,862	9,146	35
MALE													
1967	21,542	-	-	1	16	113	622	2,077	4,083	6,174	6,056	2,393	7
1966	23,479	-	1	2	15	126	719	2,151	4,566	6,733	6,609	2,543	14
1965	23,737	-	-	1	19	117	739	2,081	4,526	6,843	6,779	2,622	10
1964	24,555	-	-	4	18	132	696	2,126	4,533	7,325	6,991	2,726	4
1963	26,332	-	-	3	15	149	767	2,201	4,757	7,824	7,586	3,023	7
1962	26,654	-	-	1	21	145	738	2,370	4,797	7,913	7,656	3,003	10
1961	27,313	-	1	1	14	126	734	2,255	5,023	8,177	7,947	3,022	13
1960	28,857	1	-	3	26	145	747	2,408	5,365	8,917	8,123	3,109	13
FEMALE													
1967	28,433	1	-	3	11	105	570	1,629	3,445	7,065	9,960	5,634	10
1966	30,697	1	1	3	17	108	570	1,717	3,778	7,741	10,910	5,843	8
1965	31,231	-	-	2	14	111	627	1,800	3,863	7,929	11,105	5,768	12
1964	32,888	-	-	5	16	112	638	1,901	4,235	8,351	11,526	6,092	12
1963	34,765	-	-	2	26	114	704	2,036	4,546	9,122	11,935	6,269	11
1962	35,522	-	-	8	17	127	732	1,983	4,471	9,519	12,339	6,313	13
1961	35,969	1	-	2	12	134	728	2,204	4,755	9,681	12,259	6,174	19
1960	37,451	-	1	3	24	142	798	2,236	5,133	10,316	12,739	6,037	22

- Quantity zero

Note: Figures by race exclude data for residents of New Jersey for 1962 and 1963.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290A for 1960-67 and separate unpublished tables for the black population and Accidents.

77/09/02 TABLE 290A. DEATHS FOR 60 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1960-67

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	TOTAL	UNDER 1 YEAR	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS	65-74 YEARS	75-84 YEARS	85+ YEARS	N.S.
Hypertensive heart disease (440-443)													
WHITE, BOTH SEXES													
1967	38,209	1	-	4	15	67	401	1,837	4,761	9,879	13,978	7,260	6
1966	41,358	1	1	3	17	64	400	1,884	5,184	10,885	15,345	7,567	7
1965	42,211	-	-	2	11	65	468	1,925	5,167	11,251	15,714	7,600	8
1964	44,216	-	-	5	12	79	459	1,981	5,389	11,964	16,342	7,978	7
1963	45,176	-	-	4	17	69	478	1,947	5,521	12,425	16,569	8,138	8
1962	46,376	-	-	9	14	74	462	2,084	5,617	12,831	17,137	8,139	9
1961	49,348	-	1	1	13	87	494	2,261	6,206	13,972	17,933	8,365	15
1960	51,992	-	-	4	25	87	561	2,404	6,752	15,222	18,575	8,350	12
WHITE, MALE													
1967	15,994	-	-	1	13	37	234	1,107	2,734	4,606	5,177	2,084	1
1966	17,447	-	-	1	8	34	254	1,111	3,011	5,107	5,669	2,247	5
1965	17,770	-	-	1	6	41	294	1,097	2,935	5,210	5,857	2,326	3
1964	18,446	-	-	3	6	52	260	1,144	2,971	5,556	6,042	2,411	1
1963	19,068	-	-	3	8	45	289	1,115	2,970	5,685	6,343	2,607	3
1962	19,455	-	-	1	12	44	246	1,262	3,090	5,775	6,454	2,567	4
1961	20,923	-	1	-	7	47	274	1,257	3,406	6,313	6,915	2,699	4
1960	22,166	-	-	2	13	52	318	1,362	3,583	6,968	7,099	2,767	2
WHITE, FEMALE													
1967	22,215	1	-	3	2	30	167	730	2,027	5,273	8,801	5,176	5
1966	23,911	1	1	2	9	30	146	773	2,173	5,778	9,676	5,320	2
1965	24,441	-	-	1	5	24	174	828	2,232	6,041	9,857	5,274	5
1964	25,770	-	-	2	6	27	199	837	2,418	6,408	10,300	5,567	6
1963	26,108	-	-	1	9	24	189	832	2,551	6,740	10,226	5,531	5
1962	26,921	-	-	8	2	30	216	822	2,527	7,056	10,683	5,572	5
1961	28,425	-	-	1	6	40	220	1,004	2,800	7,659	11,018	5,666	11
1960	29,826	-	-	2	12	35	243	1,042	3,169	8,254	11,476	5,583	10

- Quantity zero

Note: Figures by race exclude data for residents of New Jersey for 1962 and 1963.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290A for 1960-67 and separate unpublished tables for the black population and Accidents.

77/09/02 TABLE 290A. DEATHS FOR 60 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1960-67

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	TOTAL	UNDER 1 YEAR	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS	65-74 YEARS	75-84 YEARS	85+ YEARS	N.S.
Hypertensive heart disease (440-443)													
ALL OTHER, BOTH SEXES													
1967	11,766	-	-	-	12	151	791	1,869	2,767	3,360	2,038	767	11
1966	12,818	-	1	2	15	170	889	1,984	3,160	3,589	2,174	819	15
1965	12,757	-	-	1	22	163	898	1,956	3,222	3,521	2,170	790	14
1964	13,227	-	-	4	22	165	875	2,046	3,379	3,712	2,175	840	9
1963	13,795	-	-	1	22	182	949	2,162	3,441	3,898	2,275	855	10
1962	13,686	-	-	-	23	188	975	2,146	3,367	3,947	2,175	852	13
1961	13,934	1	-	2	13	173	968	2,198	3,572	3,886	2,273	831	17
1960	14,316	1	1	2	25	200	984	2,240	3,746	4,011	2,287	796	23
ALL OTHER, MALE													
1967	5,548	-	-	-	3	76	388	970	1,349	1,568	879	309	6
1966	6,032	-	1	1	7	92	465	1,040	1,555	1,626	940	296	9
1965	5,967	-	-	-	13	76	445	984	1,591	1,633	922	296	7
1964	6,109	-	-	1	12	80	436	982	1,562	1,769	949	315	3
1963	6,374	-	-	-	7	97	455	1,014	1,609	1,866	999	323	4
1962	6,359	-	-	-	8	96	471	1,046	1,564	1,848	965	356	5
1961	6,390	-	-	1	7	79	460	998	1,617	1,864	1,032	323	9
1960	6,691	1	-	1	13	93	429	1,046	1,782	1,949	1,024	342	11
ALL OTHER, FEMALE													
1967	6,218	-	-	-	9	75	403	899	1,418	1,792	1,159	458	5
1966	6,786	-	-	1	8	78	424	944	1,605	1,963	1,234	523	6
1965	6,790	-	-	1	9	87	453	972	1,631	1,888	1,248	494	7
1964	7,118	-	-	3	10	85	439	1,064	1,817	1,943	1,226	525	6
1963	7,421	-	-	1	15	85	494	1,148	1,832	2,032	1,276	532	6
1962	7,327	-	-	-	15	92	504	1,100	1,803	2,099	1,210	496	8
1961	7,544	1	-	1	6	94	508	1,200	1,955	2,022	1,241	508	8
1960	7,625	-	1	1	12	107	555	1,194	1,964	2,062	1,263	454	12

- Quantity zero

Note: Figures by race exclude data for residents of New Jersey for 1962 and 1963.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290A for 1960-67 and separate unpublished tables for the black population and Accidents.

77/09/02 TABLE 290A. DEATHS FOR 60 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1960-67

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	TOTAL	UNDER 1 YEAR	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS	65-74 YEARS	75-84 YEARS	85+ YEARS	N.S.
Other hypertensive disease (444-447)													
BOTH SEXES													
1967	11,151	1	-	1	34	171	585	849	1,225	2,337	3,671	2,277	-
1966	11,380	-	-	6	33	174	549	801	1,301	2,454	3,785	2,275	2
1965	11,667	-	-	5	31	207	546	954	1,326	2,515	3,794	2,284	5
1964	12,191	-	1	1	25	179	615	1,000	1,384	2,700	3,972	2,313	1
1963	12,694	2	2	5	32	218	631	960	1,544	2,915	4,081	2,299	5
1962	12,497	-	1	1	41	213	689	1,074	1,410	2,773	4,020	2,270	5
1961	12,185	-	-	8	32	215	666	1,038	1,507	2,789	3,772	2,156	2
1960	12,662	-	2	4	28	207	773	1,168	1,558	2,984	3,791	2,142	5
MALE													
1967	5,583	1	-	-	13	87	312	501	726	1,239	1,762	942	-
1966	5,731	-	-	2	17	98	315	486	753	1,327	1,775	957	1
1965	5,942	-	-	2	17	123	315	570	753	1,354	1,819	988	1
1964	6,160	-	1	-	10	94	340	565	798	1,430	1,945	977	-
1963	6,383	1	2	-	17	129	353	567	834	1,543	1,951	983	3
1962	6,296	-	-	1	18	112	391	634	788	1,527	1,910	913	2
1961	6,056	-	-	1	20	117	369	598	847	1,423	1,765	914	2
1960	6,385	-	2	1	19	107	454	668	823	1,567	1,824	918	2
FEMALE													
1967	5,568	-	-	1	21	84	273	348	499	1,098	1,909	1,335	-
1966	5,649	-	-	4	16	76	234	315	548	1,127	2,010	1,318	1
1965	5,725	-	-	3	14	84	231	384	573	1,161	1,975	1,296	4
1964	6,031	-	-	1	15	85	275	435	586	1,270	2,027	1,336	1
1963	6,311	1	-	5	15	89	278	393	710	1,372	2,130	1,316	2
1962	6,201	-	1	-	23	101	298	440	622	1,246	2,110	1,357	3
1961	6,129	-	-	7	12	98	297	440	660	1,366	2,007	1,242	-
1960	6,277	-	-	3	9	100	319	500	735	1,417	1,967	1,224	3

- Quantity zero

Note: Figures by race exclude data for residents of New Jersey for 1962 and 1963.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290A for 1960-67 and separate unpublished tables for the black population and Accidents.

77/09/02 TABLE 290A. DEATHS FOR 60 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1960-67

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	TOTAL	UNDER 1 YEAR	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS	65-74 YEARS	75-84 YEARS	85+ YEARS	N.S.
Other hypertensive disease (444-447)													
WHITE, BOTH SEXES													
1967	8,935	1	-	1	20	52	197	435	828	1,916	3,365	2,120	-
1966	9,090	-	-	5	14	47	183	417	853	1,950	3,482	2,139	-
1965	9,296	-	-	4	14	60	183	474	909	2,020	3,488	2,142	2
1964	9,727	-	1	1	15	49	212	521	912	2,213	3,639	2,163	1
1963	9,844	1	2	5	20	76	213	458	1,022	2,313	3,631	2,100	3
1962	9,573	-	1	1	20	62	224	528	897	2,195	3,579	2,063	3
1961	9,659	-	-	7	8	65	228	550	1,051	2,281	3,452	2,016	1
1960	10,089	-	2	4	17	78	317	664	1,089	2,446	3,476	1,995	1
WHITE, MALE													
1967	4,428	1	-	-	6	24	103	258	512	1,035	1,605	884	-
1966	4,542	-	-	1	10	30	111	265	522	1,073	1,638	892	-
1965	4,624	-	-	1	6	39	112	275	526	1,093	1,653	919	-
1964	4,863	-	1	-	7	29	125	308	537	1,167	1,782	907	-
1963	4,852	-	2	-	7	51	120	276	557	1,217	1,728	893	1
1962	4,758	-	-	1	10	37	128	337	519	1,206	1,684	836	-
1961	4,749	-	-	-	6	43	133	341	622	1,162	1,597	844	1
1960	5,054	-	2	1	12	49	205	388	594	1,297	1,663	843	-
WHITE, FEMALE													
1967	4,507	-	-	1	14	28	94	177	316	881	1,760	1,236	-
1966	4,548	-	-	4	4	17	72	152	331	877	1,844	1,247	-
1965	4,672	-	-	3	8	21	71	199	383	927	1,835	1,223	2
1964	4,864	-	-	1	8	20	87	213	375	1,046	1,857	1,256	1
1963	4,992	1	-	5	13	25	93	182	465	1,096	1,903	1,207	2
1962	4,815	-	1	-	10	25	96	191	378	989	1,895	1,227	3
1961	4,910	-	-	7	2	22	95	209	429	1,119	1,855	1,172	-
1960	5,035	-	-	3	5	29	112	276	495	1,149	1,813	1,152	1

- Quantity zero

Note: Figures by race exclude data for residents of New Jersey for 1962 and 1963.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290A for 1960-67 and separate unpublished tables for the black population and Accidents.

77/09/02 TABLE 290A. DEATHS FOR 60 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1960-67

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	TOTAL	UNDER 1 YEAR	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS	65-74 YEARS	75-84 YEARS	85+ YEARS	N.S.
Other hypertensive disease (444-447)													
ALL OTHER, BOTH SEXES													
1967	2,216	-	-	-	14	119	388	414	397	421	306	157	-
1966	2,290	-	-	1	19	127	366	384	448	504	303	136	2
1965	2,371	-	-	1	17	147	363	480	417	495	306	142	3
1964	2,464	-	-	-	10	130	403	479	472	487	333	150	-
1963	2,469	1	-	-	11	131	401	471	473	526	325	128	2
1962	2,547	-	-	-	18	146	436	516	471	486	324	148	2
1961	2,526	-	-	1	24	150	438	488	456	508	320	140	1
1960	2,573	-	-	-	11	129	456	504	469	538	315	147	4
ALL OTHER, MALE													
1967	1,155	-	-	-	7	63	209	243	214	204	157	58	-
1966	1,189	-	-	1	7	68	204	221	231	254	137	65	1
1965	1,318	-	-	1	11	84	203	295	227	261	166	69	1
1964	1,297	-	-	-	3	65	215	257	261	263	163	70	-
1963	1,333	1	-	-	9	70	222	272	243	289	169	56	2
1962	1,354	-	-	-	5	72	249	283	242	270	171	60	2
1961	1,307	-	-	1	14	74	236	257	225	261	168	70	1
1960	1,331	-	-	-	7	58	249	280	229	270	161	75	2
ALL OTHER, FEMALE													
1967	1,061	-	-	-	7	56	179	171	183	217	149	99	-
1966	1,101	-	-	-	12	59	162	163	217	250	166	71	1
1965	1,053	-	-	-	6	63	160	185	190	234	140	73	2
1964	1,167	-	-	-	7	65	188	222	211	224	170	80	-
1963	1,136	-	-	-	2	61	179	199	230	237	156	72	-
1962	1,193	-	-	-	13	74	187	233	229	216	153	88	-
1961	1,219	-	-	-	10	76	202	231	231	247	152	70	-
1960	1,242	-	-	-	4	71	207	224	240	268	154	72	2

- Quantity zero

Note: Figures by race exclude data for residents of New Jersey for 1962 and 1963.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290A for 1960-67 and separate unpublished tables for the black population and Accidents.

77/09/02 TABLE 290A. DEATHS FOR 60 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1960-67

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	TOTAL	UNDER 1 YEAR	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS	65-74 YEARS	75-84 YEARS	85+ YEARS	N.S.
ICD-7 Nos. 450 Atherosclerosis													
BOTH SEXES													
1967	37,564	-	-	-	2	6	57	300	1,452	5,654	14,837	15,244	12
1966	38,907	1	-	-	4	8	60	356	1,604	6,141	15,450	15,275	8
1965	38,102	-	1	-	3	16	64	356	1,531	5,982	15,129	15,010	10
1964	37,176	-	-	-	4	6	52	348	1,636	5,918	14,734	14,472	6
1963	37,429	1	-	-	1	12	48	369	1,580	6,152	14,961	14,287	18
1962	36,848	1	1	2	2	7	55	405	1,500	5,939	14,668	14,252	16
1961	35,388	1	-	1	2	13	56	324	1,589	5,951	13,886	13,548	17
1960	35,876	-	-	-	-	15	68	360	1,695	6,254	14,242	13,226	16
MALE													
1967	16,589	-	-	-	1	4	43	193	875	3,083	6,796	5,588	6
1966	17,470	1	-	-	2	4	38	214	990	3,328	7,214	5,674	5
1965	17,299	-	1	-	1	9	47	242	951	3,308	7,027	5,707	6
1964	16,707	-	-	-	2	3	31	221	1,003	3,170	6,796	5,479	2
1963	17,152	1	-	-	-	8	30	231	971	3,379	7,011	5,518	3
1962	16,855	-	-	2	-	6	33	277	940	3,206	6,884	5,501	6
1961	16,520	1	-	-	2	4	43	207	1,009	3,320	6,611	5,315	8
1960	17,073	-	-	-	-	8	46	254	1,075	3,521	6,901	5,265	3
FEMALE													
1967	20,975	-	-	-	1	2	14	107	577	2,571	8,041	9,656	6
1966	21,437	-	-	-	2	4	22	142	614	2,813	8,236	9,601	3
1965	20,803	-	-	-	2	7	17	114	580	2,674	8,102	9,303	4
1964	20,469	-	-	-	2	3	21	127	633	2,748	7,938	8,993	4
1963	20,277	-	-	-	1	4	18	138	609	2,773	7,950	8,769	15
1962	19,993	1	1	-	2	1	22	128	560	2,733	7,784	8,751	10
1961	18,868	-	-	1	-	9	13	117	580	2,631	7,275	8,233	9
1960	18,803	-	-	-	-	7	22	106	620	2,733	7,341	7,961	13

- Quantity zero

Note: Figures by race exclude data for residents of New Jersey for 1962 and 1963.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290A for 1960-67 and separate unpublished tables for the black population and Accidents.

77/09/02 TABLE 290A. DEATHS FOR 60 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1960-67

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	TOTAL	UNDER 1 YEAR	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS	65-74 YEARS	75-84 YEARS	85+ YEARS	N.S.
ICD-7 Nos. 450 Atherosclerosis													
WHITE, BOTH SEXES													
1967	34,953	-	-	-	1	5	46	243	1,183	4,887	13,975	14,603	10
1966	36,041	1	-	-	2	6	45	272	1,255	5,314	14,532	14,608	6
1965	35,292	-	1	-	3	15	42	266	1,208	5,195	14,192	14,365	5
1964	34,414	-	-	-	4	6	40	259	1,299	5,117	13,824	13,862	3
1963	33,580	1	-	-	1	8	34	278	1,189	5,156	13,622	13,274	17
1962	33,092	1	1	2	2	3	37	294	1,160	4,995	13,374	13,215	8
1961	32,827	1	-	1	1	10	43	250	1,258	5,177	13,086	12,989	11
1960	33,213	-	-	-	-	11	44	283	1,326	5,454	13,458	12,623	14
WHITE, MALE													
1967	15,307	-	-	-	-	3	34	162	726	2,682	6,379	5,316	5
1966	16,069	1	-	-	1	3	28	172	800	2,933	6,746	5,382	3
1965	15,907	-	1	-	1	8	35	188	778	2,897	6,569	5,426	4
1964	15,356	-	-	-	2	3	23	172	816	2,766	6,345	5,228	1
1963	15,294	1	-	-	-	5	21	183	762	2,866	6,354	5,099	3
1962	15,018	-	-	2	-	3	23	213	743	2,684	6,274	5,074	2
1961	15,196	1	-	-	1	4	34	169	825	2,892	6,199	5,066	5
1960	15,675	-	-	-	-	6	31	209	858	3,100	6,498	4,971	2
WHITE, FEMALE													
1967	19,646	-	-	-	1	2	12	81	457	2,205	7,596	9,287	5
1966	19,972	-	-	-	1	3	17	100	455	2,381	7,786	9,226	3
1965	19,385	-	-	-	2	7	7	78	430	2,298	7,623	8,939	1
1964	19,058	-	-	-	2	3	17	87	483	2,351	7,479	8,634	2
1963	18,286	-	-	-	1	3	13	95	427	2,290	7,268	8,175	14
1962	18,074	1	1	-	2	-	14	81	417	2,311	7,100	8,141	6
1961	17,631	-	-	1	-	6	9	81	433	2,285	6,887	7,923	6
1960	17,538	-	-	-	-	5	13	74	468	2,354	6,960	7,652	12

- Quantity zero

Note: Figures by race exclude data for residents of New Jersey for 1962 and 1963.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290A for 1960-67 and separate unpublished tables for the black population and Accidents.

77/09/02 TABLE 290A. DEATHS FOR 60 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1960-67

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	TOTAL	UNDER 1 YEAR	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS	65-74 YEARS	75-84 YEARS	85+ YEARS	N.S.
ICD-7 Nos. 450 Atherosclerosis													
ALL OTHER, BOTH SEXES													
1967	2,611	-	-	-	1	1	11	57	269	767	862	641	2
1966	2,866	-	-	-	2	2	15	84	349	827	918	667	2
1965	2,810	-	-	-	-	1	22	90	323	787	937	645	5
1964	2,762	-	-	-	-	-	12	89	337	801	910	610	3
1963	2,784	-	-	-	-	4	14	83	344	805	912	621	1
1962	2,724	-	-	-	-	4	17	93	298	791	868	645	8
1961	2,561	-	-	-	1	3	13	74	331	774	800	559	6
1960	2,663	-	-	-	-	4	24	77	369	800	784	603	2
ALL OTHER, MALE													
1967	1,282	-	-	-	1	1	9	31	149	401	417	272	1
1966	1,401	-	-	-	1	1	10	42	190	395	468	292	2
1965	1,392	-	-	-	-	1	12	54	173	411	458	281	2
1964	1,351	-	-	-	-	-	8	49	187	404	451	251	1
1963	1,400	-	-	-	-	3	9	43	176	419	467	283	-
1962	1,399	-	-	-	-	3	9	51	169	435	429	299	4
1961	1,324	-	-	-	1	-	9	38	184	428	412	249	3
1960	1,398	-	-	-	-	2	15	45	217	421	403	294	1
ALL OTHER, FEMALE													
1967	1,329	-	-	-	-	-	2	26	120	366	445	369	1
1966	1,465	-	-	-	1	1	5	42	159	432	450	375	-
1965	1,418	-	-	-	-	-	10	36	150	376	479	364	3
1964	1,411	-	-	-	-	-	4	40	150	397	459	359	2
1963	1,384	-	-	-	-	1	5	40	168	386	445	338	1
1962	1,325	-	-	-	-	1	8	42	129	356	439	346	4
1961	1,237	-	-	-	-	3	4	36	147	346	388	310	3
1960	1,265	-	-	-	-	2	9	32	152	379	381	309	1

- Quantity zero

Note: Figures by race exclude data for residents of New Jersey for 1962 and 1963.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290A for 1960-67 and separate unpublished tables for the black population and Accidents.

77/09/02 TABLE 290A. DEATHS FOR 60 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1960-67

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	TOTAL	UNDER 1 YEAR	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS	65-74 YEARS	75-84 YEARS	85+ YEARS	N.S.
Other diseases of circulatory system (451-468)													
BOTH SEXES													
1967	29,944	28	24	72	363	541	1,204	2,557	5,341	9,250	7,912	2,648	4
1966	28,682	41	29	56	326	522	1,206	2,487	5,279	8,858	7,466	2,403	9
1965	27,279	38	30	59	319	514	1,174	2,434	5,022	8,448	6,967	2,264	10
1964	25,745	46	28	69	295	456	1,073	2,265	4,872	8,026	6,571	2,038	6
1963	24,385	47	21	58	229	448	1,033	2,189	4,531	7,768	6,076	1,977	8
1962	22,632	31	22	57	216	411	953	2,012	4,331	7,338	5,432	1,823	6
1961	20,707	47	21	54	192	404	835	1,976	3,937	6,641	5,017	1,576	7
1960	19,797	38	21	67	210	370	910	1,871	3,986	6,168	4,658	1,496	2
MALE													
1967	17,632	18	17	33	112	189	526	1,541	3,663	6,034	4,379	1,119	1
1966	16,788	31	15	20	93	174	529	1,464	3,524	5,837	4,075	1,022	4
1965	16,209	24	16	26	106	176	530	1,449	3,381	5,699	3,843	955	4
1964	15,571	26	17	22	91	162	520	1,394	3,368	5,397	3,677	895	2
1963	14,689	25	15	20	75	171	494	1,369	3,071	5,240	3,395	807	7
1962	13,737	25	12	19	70	155	472	1,277	2,928	5,012	2,986	780	1
1961	12,667	26	11	21	65	163	432	1,220	2,715	4,468	2,842	699	5
1960	12,090	22	14	28	63	137	462	1,165	2,760	4,163	2,583	691	2
FEMALE													
1967	12,312	10	7	39	251	352	678	1,016	1,678	3,216	3,533	1,529	3
1966	11,894	10	14	36	233	348	677	1,023	1,755	3,021	3,391	1,381	5
1965	11,070	14	14	33	213	338	644	985	1,641	2,749	3,124	1,309	6
1964	10,174	20	11	47	204	294	553	871	1,504	2,629	2,894	1,143	4
1963	9,696	22	6	38	154	277	539	820	1,460	2,528	2,681	1,170	1
1962	8,895	6	10	38	146	256	481	735	1,403	2,326	2,446	1,043	5
1961	8,040	21	10	33	127	241	403	756	1,222	2,173	2,175	877	2
1960	7,707	16	7	39	147	233	448	706	1,226	2,005	2,075	805	-

- Quantity zero

Note: Figures by race exclude data for residents of New Jersey for 1962 and 1963.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290A for 1960-67 and separate unpublished tables for the black population and Accidents.

77/09/02 TABLE 290A. DEATHS FOR 60 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1960-67

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	TOTAL	UNDER 1 YEAR	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS	65-74 YEARS	75-84 YEARS	85+ YEARS	N.S.
Other diseases of circulatory system (451-468)													
WHITE, BOTH SEXES													
1967	26,448	20	18	49	251	362	852	2,056	4,619	8,321	7,428	2,469	3
1966	25,349	28	24	50	236	358	870	1,997	4,486	8,009	7,034	2,251	6
1965	24,189	23	21	47	238	358	834	1,954	4,335	7,682	6,548	2,144	5
1964	22,870	37	22	55	204	306	777	1,801	4,203	7,323	6,215	1,923	4
1963	20,885	28	13	45	156	289	728	1,712	3,753	6,834	5,519	1,804	4
1962	19,468	19	14	41	148	288	659	1,540	3,615	6,457	4,996	1,686	5
1961	18,410	35	17	44	132	278	606	1,586	3,418	6,060	4,738	1,491	5
1960	17,657	27	17	49	161	261	675	1,518	3,433	5,683	4,417	1,415	1
WHITE, MALE													
1967	15,864	15	15	21	80	134	389	1,272	3,257	5,524	4,118	1,038	1
1966	15,069	22	14	19	69	114	410	1,214	3,086	5,322	3,848	950	1
1965	14,591	13	14	21	85	129	387	1,192	2,983	5,243	3,622	901	1
1964	14,051	21	14	18	71	117	393	1,145	2,974	4,984	3,475	838	1
1963	12,747	16	9	16	52	119	364	1,093	2,610	4,632	3,094	738	4
1962	12,009	15	8	17	49	112	346	1,021	2,513	4,448	2,757	722	1
1961	11,422	19	7	16	44	115	322	1,001	2,439	4,105	2,694	657	3
1960	10,938	18	11	24	52	97	359	975	2,438	3,864	2,446	653	1
WHITE, FEMALE													
1967	10,584	5	3	28	171	228	463	784	1,362	2,797	3,310	1,431	2
1966	10,280	6	10	31	167	244	460	783	1,400	2,687	3,186	1,301	5
1965	9,598	10	7	26	153	229	447	762	1,352	2,439	2,926	1,243	4
1964	8,819	16	8	37	133	189	384	656	1,229	2,339	2,740	1,085	3
1963	8,138	12	4	29	104	170	364	619	1,143	2,202	2,425	1,066	-
1962	7,459	4	6	24	99	176	313	519	1,102	2,009	2,239	964	4
1961	6,988	16	10	28	88	163	284	585	979	1,955	2,044	834	2
1960	6,719	9	6	25	109	164	316	543	995	1,819	1,971	762	-

- Quantity zero

Note: Figures by race exclude data for residents of New Jersey for 1962 and 1963.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290A for 1960-67 and separate unpublished tables for the black population and Accidents.

77/09/02 TABLE 290A. DEATHS FOR 60 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1960-67

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	TOTAL	UNDER 1 YEAR	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS	65-74 YEARS	75-84 YEARS	85+ YEARS	N.S.
Other diseases of circulatory system (451-468)													
ALL OTHER, BOTH SEXES													
1967	3,496	8	6	23	112	179	352	501	722	929	484	179	1
1966	3,333	13	5	6	90	164	336	490	793	849	432	152	3
1965	3,090	15	9	12	81	156	340	480	687	766	419	120	5
1964	2,875	9	6	14	91	150	296	464	669	703	356	115	2
1963	2,722	18	7	11	71	149	271	408	615	672	378	118	4
1962	2,337	11	8	13	60	108	253	379	550	598	264	92	1
1961	2,297	12	4	10	60	126	229	390	519	581	279	85	2
1960	2,140	11	4	18	49	109	235	353	553	485	241	81	1
ALL OTHER, MALE													
1967	1,768	3	2	12	32	55	137	269	406	510	261	81	-
1966	1,719	9	1	1	24	60	119	250	438	515	227	72	3
1965	1,618	11	2	5	21	47	143	257	398	456	221	54	3
1964	1,520	5	3	4	20	45	127	249	394	413	202	57	1
1963	1,475	9	6	3	22	50	116	231	348	430	206	51	3
1962	1,236	9	4	2	16	35	106	205	305	371	139	44	-
1961	1,245	7	4	5	21	48	110	219	276	363	148	42	2
1960	1,152	4	3	4	11	40	103	190	322	299	137	38	1
ALL OTHER, FEMALE													
1967	1,728	5	4	11	80	124	215	232	316	419	223	98	1
1966	1,614	4	4	5	66	104	217	240	355	334	205	80	-
1965	1,472	4	7	7	60	109	197	223	289	310	198	66	2
1964	1,355	4	3	10	71	105	169	215	275	290	154	58	1
1963	1,247	9	1	8	49	99	155	177	267	242	172	67	1
1962	1,101	2	4	11	44	73	147	174	245	227	125	48	1
1961	1,052	5	-	5	39	78	119	171	243	218	131	43	-
1960	988	7	1	14	38	69	132	163	231	186	104	43	-

- Quantity zero

Note: Figures by race exclude data for residents of New Jersey for 1962 and 1963.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290A for 1960-67 and separate unpublished tables for the black population and Accidents.

77/09/02 TABLE 290A. DEATHS FOR 60 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1960-67

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	TOTAL	UNDER 1 YEAR	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS	65-74 YEARS	75-84 YEARS	85+ YEARS	N.S.
Chronic and unspecified nephritis and other renal sclerosis (592-594)													
BOTH SEXES													
1967	9,936	22	26	127	382	418	821	1,327	1,704	2,069	2,015	1,020	5
1966	10,376	29	16	167	415	510	840	1,326	1,732	2,154	2,147	1,037	3
1965	10,595	31	41	158	405	508	846	1,405	1,748	2,159	2,200	1,092	2
1964	11,079	25	39	157	431	533	915	1,408	1,845	2,228	2,367	1,129	2
1963	11,243	21	40	179	409	576	893	1,480	1,857	2,368	2,305	1,109	6
1962	11,372	47	31	174	398	604	987	1,377	1,781	2,374	2,432	1,162	5
1961	11,596	24	25	170	436	647	963	1,419	1,849	2,412	2,446	1,201	4
1960	12,033	22	37	172	407	648	1,019	1,506	1,886	2,629	2,471	1,227	9
MALE													
1967	5,516	13	13	61	218	236	467	768	1,029	1,130	1,096	482	3
1966	5,798	16	9	83	254	298	484	780	1,021	1,231	1,109	511	2
1965	5,893	16	20	67	238	308	495	863	979	1,206	1,170	529	2
1964	6,106	16	22	77	247	297	528	832	1,073	1,222	1,259	533	-
1963	6,222	8	19	88	229	353	509	870	1,076	1,290	1,238	539	3
1962	6,331	26	20	83	240	352	574	810	1,028	1,360	1,249	587	2
1961	6,398	12	19	84	261	389	565	841	1,045	1,340	1,291	547	4
1960	6,622	14	23	92	255	392	596	862	1,075	1,421	1,302	586	4
FEMALE													
1967	4,420	9	13	66	164	182	354	559	675	939	919	538	2
1966	4,578	13	7	84	161	212	356	546	711	923	1,038	526	1
1965	4,702	15	21	91	167	200	351	542	769	953	1,030	563	-
1964	4,973	9	17	80	184	236	387	576	772	1,006	1,108	596	2
1963	5,021	13	21	91	180	223	384	610	781	1,078	1,067	570	3
1962	5,041	21	11	91	158	252	413	567	753	1,014	1,183	575	3
1961	5,198	12	6	86	175	258	398	578	804	1,072	1,155	654	-
1960	5,411	8	14	80	152	256	423	644	811	1,208	1,169	641	5

- Quantity zero

Note: Figures by race exclude data for residents of New Jersey for 1962 and 1963.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290A for 1960-67 and separate unpublished tables for the black population and Accidents.

77/09/02 TABLE 290A. DEATHS FOR 60 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1960-67

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	TOTAL	UNDER 1 YEAR	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS	65-74 YEARS	75-84 YEARS	85+ YEARS	N.S.
Chronic and unspecified nephritis and other renal sclerosis (592-594)													
WHITE, BOTH SEXES													
1967	7,510	17	20	107	295	267	509	874	1,193	1,612	1,733	881	2
1966	7,920	23	11	139	331	361	537	879	1,216	1,689	1,836	897	1
1965	8,159	22	28	121	337	343	558	987	1,259	1,667	1,880	955	2
1964	8,553	19	31	125	344	374	620	950	1,294	1,752	2,043	1,000	1
1963	8,397	16	24	143	312	399	584	965	1,313	1,762	1,930	947	2
1962	8,569	31	22	130	325	443	648	908	1,235	1,795	2,057	973	2
1961	9,153	18	18	136	355	493	685	999	1,358	1,927	2,114	1,049	1
1960	9,385	17	20	137	341	502	710	1,034	1,370	2,067	2,120	1,065	2
WHITE, MALE													
1967	4,246	11	9	48	173	173	310	538	765	889	919	410	1
1966	4,458	11	5	64	203	218	339	536	724	988	934	435	1
1965	4,584	9	13	52	197	223	337	626	743	947	985	450	2
1964	4,775	12	17	58	204	223	382	585	787	965	1,071	471	-
1963	4,700	6	10	69	171	258	357	588	793	965	1,025	458	-
1962	4,854	18	14	58	198	276	394	570	750	1,039	1,053	483	1
1961	5,087	7	12	67	218	313	431	624	784	1,046	1,110	474	1
1960	5,220	11	12	72	217	324	428	642	803	1,105	1,099	506	1
WHITE, FEMALE													
1967	3,264	6	11	59	122	94	199	336	428	723	814	471	1
1966	3,462	12	6	75	128	143	198	343	492	701	902	462	-
1965	3,575	13	15	69	140	120	221	361	516	720	895	505	-
1964	3,778	7	14	67	140	151	238	365	507	787	972	529	1
1963	3,697	10	14	74	141	141	227	377	520	797	905	489	2
1962	3,715	13	8	72	127	167	254	338	485	756	1,004	490	1
1961	4,066	11	6	69	137	180	254	375	574	881	1,004	575	-
1960	4,165	6	8	65	124	178	282	392	567	962	1,021	559	1

- Quantity zero

Note: Figures by race exclude data for residents of New Jersey for 1962 and 1963.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290A for 1960-67 and separate unpublished tables for the black population and Accidents.

77/09/02 TABLE 290A. DEATHS FOR 60 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1960-67

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	TOTAL	UNDER 1 YEAR	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS	65-74 YEARS	75-84 YEARS	85+ YEARS	N.S.
Chronic and unspecified nephritis and other renal sclerosis (592-594)													
ALL OTHER, BOTH SEXES													
1967	2,426	5	6	20	87	151	312	453	511	457	282	139	3
1966	2,456	6	5	28	84	149	303	447	516	465	311	140	2
1965	2,436	9	13	37	68	165	288	418	489	492	320	137	-
1964	2,526	6	8	32	87	159	295	458	551	476	324	129	1
1963	2,469	5	13	32	83	154	285	457	471	523	309	133	4
1962	2,451	15	9	39	55	144	301	430	479	507	312	157	3
1961	2,443	6	7	34	81	154	278	420	491	485	332	152	3
1960	2,648	5	17	35	66	146	309	472	516	562	351	162	7
ALL OTHER, MALE													
1967	1,270	2	4	13	45	63	157	230	264	241	177	72	2
1966	1,340	5	4	19	51	80	145	244	297	243	175	76	1
1965	1,309	7	7	15	41	85	158	237	236	259	185	79	-
1964	1,331	4	5	19	43	74	146	247	286	257	188	62	-
1963	1,338	2	7	17	53	83	139	250	245	285	181	73	3
1962	1,292	7	6	23	31	67	159	219	239	283	165	92	1
1961	1,311	5	7	17	43	76	134	217	261	294	181	73	3
1960	1,402	3	11	20	38	68	168	220	272	316	203	80	3
ALL OTHER, FEMALE													
1967	1,156	3	2	7	42	88	155	223	247	216	105	67	1
1966	1,116	1	1	9	33	69	158	203	219	222	136	64	1
1965	1,127	2	6	22	27	80	130	181	253	233	135	58	-
1964	1,195	2	3	13	44	85	149	211	265	219	136	67	1
1963	1,131	3	6	15	30	71	146	207	226	238	128	60	1
1962	1,159	8	3	16	24	77	142	211	240	224	147	65	2
1961	1,132	1	-	17	38	78	144	203	230	191	151	79	-
1960	1,246	2	6	15	28	78	141	252	244	246	148	82	4

- Quantity zero

Note: Figures by race exclude data for residents of New Jersey for 1962 and 1963.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290A for 1960-67 and separate unpublished tables for the black population and Accidents.

77/09/02 TABLE 290A. DEATHS FOR 60 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1960-67

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	TOTAL	UNDER 1 YEAR	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS	65-74 YEARS	75-84 YEARS	85+ YEARS	N.S.
ICD-7 Nos. 480-493 Pneumonia and influenza													
BOTH SEXES													
1967	56,892	5,716	1,443	699	735	738	1,702	3,239	5,197	9,946	16,054	11,401	22
1966	63,615	6,961	1,772	790	766	791	1,996	3,635	6,027	11,296	17,276	12,269	36
1965	61,903	8,036	1,889	823	652	803	1,905	3,329	5,644	10,866	16,506	11,423	27
1964	59,451	8,325	2,047	742	625	793	1,734	3,268	5,378	10,346	15,506	10,658	29
1963	70,761	8,914	2,395	882	714	926	2,080	3,947	6,779	13,119	18,313	12,659	33
1962	59,995	8,985	2,271	862	671	775	1,866	3,203	5,325	10,615	15,034	10,368	20
1961	55,175	9,202	2,147	760	549	729	1,744	2,905	4,898	9,638	13,385	9,204	14
1960	66,803	9,828	2,628	929	724	1,091	2,234	4,024	6,725	12,327	15,782	10,469	42
MALE													
1967	31,904	3,244	820	371	435	398	1,043	2,161	3,492	6,379	8,648	4,900	13
1966	35,565	3,988	952	403	442	439	1,204	2,311	4,047	7,164	9,275	5,315	25
1965	34,973	4,604	959	456	367	444	1,113	2,168	3,766	6,968	9,101	5,010	17
1964	33,576	4,745	1,087	376	347	429	1,038	2,178	3,585	6,654	8,416	4,698	23
1963	39,569	5,086	1,261	450	380	455	1,222	2,601	4,461	8,292	9,783	5,562	16
1962	33,517	5,128	1,217	442	367	410	1,068	2,091	3,501	6,658	8,108	4,515	12
1961	31,175	5,142	1,153	408	321	401	1,048	1,949	3,282	6,194	7,242	4,026	9
1960	38,126	5,594	1,388	490	400	566	1,357	2,709	4,538	7,913	8,544	4,601	26
FEMALE													
1967	24,988	2,472	623	328	300	340	659	1,078	1,705	3,567	7,406	6,501	9
1966	28,050	2,973	820	387	324	352	792	1,324	1,980	4,132	8,001	6,954	11
1965	26,930	3,432	930	367	285	359	792	1,161	1,878	3,898	7,405	6,413	10
1964	25,875	3,580	960	366	278	364	696	1,090	1,793	3,692	7,090	5,960	6
1963	31,192	3,828	1,134	432	334	471	858	1,346	2,318	4,827	8,530	7,097	17
1962	26,478	3,857	1,054	420	304	365	798	1,112	1,824	3,957	6,926	5,853	8
1961	24,000	4,060	994	352	228	328	696	956	1,616	3,444	6,143	5,178	5
1960	28,677	4,234	1,240	439	324	525	877	1,315	2,187	4,414	7,238	5,868	16

Note: Figures by race exclude data for residents of New Jersey for 1962 and 1963.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290A for 1960-67 and separate unpublished tables for the black population and Accidents.

77/09/02 TABLE 290A. DEATHS FOR 60 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1960-67

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	TOTAL	UNDER 1 YEAR	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS	65-74 YEARS	75-84 YEARS	85+ YEARS	N.S.
ICD-7 Nos. 480-493 Pneumonia and influenza													
WHITE, BOTH SEXES													
1967	47,759	3,374	883	530	564	455	1,075	2,359	4,215	8,529	14,938	10,823	14
1966	52,779	3,988	1,087	600	600	467	1,249	2,728	4,791	9,650	16,005	11,592	22
1965	51,311	4,739	1,169	604	510	500	1,214	2,535	4,479	9,337	15,336	10,872	16
1964	49,120	5,056	1,261	553	483	455	1,058	2,416	4,305	8,978	14,424	10,114	17
1963	55,834	5,144	1,430	640	530	568	1,295	2,694	4,972	10,656	16,331	11,556	18
1962	47,795	5,299	1,365	633	511	470	1,189	2,242	4,068	8,828	13,567	9,609	14
1961	45,407	5,809	1,394	596	411	450	1,147	2,132	3,874	8,354	12,538	8,697	5
1960	54,091	5,835	1,716	711	557	707	1,448	2,899	5,221	10,530	14,575	9,867	25
WHITE, MALE													
1967	26,444	1,976	489	290	332	240	655	1,550	2,849	5,458	7,977	4,620	8
1966	29,215	2,377	595	319	361	245	757	1,696	3,251	6,104	8,530	4,966	14
1965	28,848	2,831	596	345	291	268	704	1,623	3,025	6,001	8,404	4,750	10
1964	27,545	2,962	668	270	276	237	621	1,630	2,884	5,794	7,766	4,425	12
1963	31,106	2,994	747	327	287	286	752	1,814	3,350	6,808	8,664	5,068	9
1962	26,618	3,112	736	317	291	247	704	1,463	2,712	5,565	7,301	4,161	9
1961	25,471	3,318	768	319	251	236	681	1,421	2,622	5,357	6,733	3,763	2
1960	30,721	3,417	879	380	315	384	897	1,965	3,575	6,770	7,811	4,312	16
WHITE, FEMALE													
1967	21,315	1,398	394	240	232	215	420	809	1,366	3,071	6,961	6,203	6
1966	23,564	1,611	492	281	239	222	492	1,032	1,540	3,546	7,475	6,626	8
1965	22,463	1,908	573	259	219	232	510	912	1,454	3,336	6,932	6,122	6
1964	21,575	2,094	593	283	207	218	437	786	1,421	3,184	6,658	5,689	5
1963	24,728	2,150	683	313	243	282	543	880	1,622	3,848	7,667	6,488	9
1962	21,177	2,187	629	316	220	223	485	779	1,356	3,263	6,266	5,448	5
1961	19,936	2,491	626	277	160	214	466	711	1,252	2,997	5,805	4,934	3
1960	23,370	2,418	837	331	242	323	551	934	1,646	3,760	6,764	5,555	9

Note: Figures by race exclude data for residents of New Jersey for 1962 and 1963.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290A for 1960-67 and separate unpublished tables for the black population and Accidents.

77/09/02 TABLE 290A. DEATHS FOR 60 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1960-67

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	TOTAL	UNDER 1 YEAR	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS	65-74 YEARS	75-84 YEARS	85+ YEARS	N.S.
ICD-7 Nos. 480-493 Pneumonia and influenza													
ALL OTHER, BOTH SEXES													
1967	9,133	2,342	560	169	171	283	627	880	982	1,417	1,116	578	8
1966	10,836	2,973	685	190	166	324	747	907	1,236	1,646	1,271	677	14
1965	10,592	3,297	720	219	142	303	691	794	1,165	1,529	1,170	551	11
1964	10,331	3,269	786	189	142	338	676	852	1,073	1,368	1,082	544	12
1963	13,019	3,553	911	205	162	329	725	1,119	1,624	2,086	1,481	809	15
1962	10,472	3,432	832	196	133	276	608	863	1,091	1,454	1,052	529	6
1961	9,768	3,393	753	164	138	279	597	773	1,024	1,284	847	507	9
1960	12,712	3,993	912	218	167	384	786	1,125	1,504	1,797	1,207	602	17
ALL OTHER, MALE													
1967	5,460	1,268	331	81	103	158	388	611	643	921	671	280	5
1966	6,350	1,611	357	84	81	194	447	615	796	1,060	745	349	11
1965	6,125	1,773	363	111	76	176	409	545	741	967	697	260	7
1964	6,031	1,783	419	106	71	192	417	548	701	860	650	273	11
1963	7,371	1,961	481	107	81	156	430	701	992	1,247	838	370	7
1962	5,930	1,868	445	105	63	154	322	557	673	889	597	254	3
1961	5,704	1,824	385	89	70	165	367	528	660	837	509	263	7
1960	7,405	2,177	509	110	85	182	460	744	963	1,143	733	289	10
ALL OTHER, FEMALE													
1967	3,673	1,074	229	88	68	125	239	269	339	496	445	298	3
1966	4,486	1,362	328	106	85	130	300	292	440	586	526	328	3
1965	4,467	1,524	357	108	66	127	282	249	424	562	473	291	4
1964	4,300	1,486	367	83	71	146	259	304	372	508	432	271	1
1963	5,648	1,592	430	98	81	173	295	418	632	839	643	439	8
1962	4,542	1,564	387	91	70	122	286	306	418	565	455	275	3
1961	4,064	1,569	368	75	68	114	230	245	364	447	338	244	2
1960	5,307	1,816	403	108	82	202	326	381	541	654	474	313	7

Note: Figures by race exclude data for residents of New Jersey for 1962 and 1963.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290A for 1960-67 and separate unpublished tables for the black population and Accidents.

77/09/02 TABLE 290A. DEATHS FOR 60 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1960-67

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	TOTAL	UNDER 1 YEAR	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS	65-74 YEARS	75-84 YEARS	85+ YEARS	N.S.
Influenza (480-493)													
BOTH SEXES													
1967	1,475	79	30	23	16	13	32	58	110	257	471	385	1
1966	2,830	129	53	34	33	30	71	122	236	502	860	759	1
1965	2,295	182	57	20	27	30	47	92	191	391	690	566	2
1964	1,687	137	53	17	16	11	26	75	136	294	488	433	1
1963	7,083	356	159	72	104	82	196	386	741	1,373	2,120	1,490	4
1962	3,431	284	123	77	63	48	101	168	258	598	940	770	1
1961	2,137	254	83	39	34	26	68	96	189	331	592	424	1
1960	7,872	538	189	106	142	154	257	519	846	1,604	2,019	1,488	10
MALE													
1967	648	36	18	14	12	9	14	36	61	123	194	130	1
1966	1,344	71	27	13	21	11	32	60	132	286	375	315	1
1965	1,064	96	31	12	13	17	22	42	111	202	310	207	1
1964	835	79	34	6	6	6	11	43	73	163	228	186	-
1963	3,584	194	77	41	55	42	86	218	438	739	1,038	654	2
1962	1,664	162	64	37	30	22	48	90	122	303	461	324	1
1961	1,031	142	42	24	17	13	37	61	105	162	270	158	-
1960	4,096	293	80	45	69	79	132	314	501	942	1,006	630	5
FEMALE													
1967	827	43	12	9	4	4	18	22	49	134	277	255	-
1966	1,486	58	26	21	12	19	39	62	104	216	485	444	-
1965	1,231	86	26	8	14	13	25	50	80	189	380	359	1
1964	852	58	19	11	10	5	15	32	63	131	260	247	1
1963	3,499	162	82	31	49	40	110	168	303	634	1,082	836	2
1962	1,767	122	59	40	33	26	53	78	136	295	479	446	-
1961	1,106	112	41	15	17	13	31	35	84	169	322	266	1
1960	3,776	245	109	61	73	75	125	205	345	662	1,013	858	5

- Quantity zero

Note: Figures by race exclude data for residents of New Jersey for 1962 and 1963.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290A for 1960-67 and separate unpublished tables for the black population and Accidents.

77/09/02 TABLE 290A. DEATHS FOR 60 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1960-67

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	TOTAL	UNDER 1 YEAR	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS	65-74 YEARS	75-84 YEARS	85+ YEARS	N.S.
Influenza (480-493)													
WHITE, BOTH SEXES													
1967	1,244	30	15	19	12	11	26	45	86	208	432	359	1
1966	2,457	50	32	27	27	21	59	93	192	431	807	718	-
1965	1,902	65	26	13	22	21	38	75	151	312	642	536	1
1964	1,406	50	33	14	13	7	16	51	104	250	458	409	1
1963	5,350	167	89	50	70	51	131	241	464	997	1,805	1,284	1
1962	2,769	141	83	56	51	34	74	106	187	478	844	714	1
1961	1,700	122	47	26	24	19	50	71	142	267	542	389	1
1960	6,231	252	131	84	111	110	175	344	590	1,251	1,806	1,374	3
WHITE, MALE													
1967	549	12	9	10	9	7	12	29	53	104	180	123	1
1966	1,172	36	14	11	17	7	26	46	113	254	351	297	-
1965	881	38	12	9	9	13	21	35	90	169	289	195	1
1964	693	30	22	5	6	4	7	31	54	142	215	177	-
1963	2,730	89	43	29	39	26	62	142	295	552	885	567	1
1962	1,347	84	46	26	24	17	39	53	88	251	419	299	1
1961	783	61	24	13	13	9	27	48	77	127	245	139	-
1960	3,235	139	57	34	58	63	92	214	359	739	892	586	2
WHITE, FEMALE													
1967	695	18	6	9	3	4	14	16	33	104	252	236	-
1966	1,285	14	18	16	10	14	33	47	79	177	456	421	-
1965	1,021	27	14	4	13	8	17	40	61	143	353	341	-
1964	713	20	11	9	7	3	9	20	50	108	243	232	1
1963	2,620	78	46	21	31	25	69	99	169	445	920	717	-
1962	1,422	57	37	30	27	17	35	53	99	227	425	415	-
1961	917	61	23	13	11	10	23	23	65	140	297	250	1
1960	2,996	113	74	50	53	47	83	130	231	512	914	788	1

- Quantity zero

Note: Figures by race exclude data for residents of New Jersey for 1962 and 1963.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290A for 1960-67 and separate unpublished tables for the black population and Accidents.

77/09/02 TABLE 290A. DEATHS FOR 60 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1960-67

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	TOTAL	UNDER 1 YEAR	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS	65-74 YEARS	75-84 YEARS	85+ YEARS	N.S.
Influenza (480-493)													
ALL OTHER, BOTH SEXES													
1967	231	49	15	4	4	2	6	13	24	49	39	26	-
1966	373	79	21	7	6	9	12	29	44	71	53	41	1
1965	393	117	31	7	5	9	9	17	40	79	48	30	1
1964	281	87	20	3	3	4	10	24	32	44	30	24	-
1963	1,667	186	70	19	33	30	59	139	268	366	300	194	3
1962	637	142	38	20	11	14	26	61	69	113	91	52	-
1961	437	132	36	13	10	7	18	25	47	64	50	35	-
1960	1,641	286	58	22	31	44	82	175	256	353	213	114	7
ALL OTHER, MALE													
1967	99	24	9	4	3	2	2	7	8	19	14	7	-
1966	172	35	13	2	4	4	6	14	19	32	24	18	1
1965	183	58	19	3	4	4	1	7	21	33	21	12	-
1964	142	49	12	1	-	2	4	12	19	21	13	9	-
1963	818	102	34	12	15	16	20	73	139	182	141	83	1
1962	300	77	17	10	5	5	9	36	33	46	39	23	-
1961	248	81	18	11	4	4	10	13	28	35	25	19	-
1960	861	154	23	11	11	16	40	100	142	203	114	44	3
ALL OTHER, FEMALE													
1967	132	25	6	-	1	-	4	6	16	30	25	19	-
1966	201	44	8	5	2	5	6	15	25	39	29	23	-
1965	210	59	12	4	1	5	8	10	19	46	27	18	1
1964	139	38	8	2	3	2	6	12	13	23	17	15	-
1963	849	84	36	7	18	14	39	66	129	184	159	111	2
1962	337	65	21	10	6	9	17	25	36	67	52	29	-
1961	189	51	18	2	6	3	8	12	19	29	25	16	-
1960	780	132	35	11	20	28	42	75	114	150	99	70	4

- Quantity zero

Note: Figures by race exclude data for residents of New Jersey for 1962 and 1963.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290A for 1960-67 and separate unpublished tables for the black population and Accidents.

77/09/02 TABLE 290A. DEATHS FOR 60 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1960-67

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	TOTAL	UNDER 1 YEAR	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS	65-74 YEARS	75-84 YEARS	85+ YEARS	N.S.
Pneumonia, except pneumonia of newborn (490-493)													
BOTH SEXES													
1967	55,417	5,637	1,413	676	719	725	1,670	3,181	5,087	9,689	15,583	11,016	21
1966	60,785	6,832	1,719	756	733	761	1,925	3,513	5,791	10,794	16,416	11,510	35
1965	59,608	7,854	1,832	803	625	773	1,858	3,237	5,453	10,475	15,816	10,857	25
1964	57,764	8,188	1,994	725	609	782	1,708	3,193	5,242	10,052	15,018	10,225	28
1963	63,678	8,558	2,236	810	610	844	1,884	3,561	6,038	11,746	16,193	11,169	29
1962	56,564	8,701	2,148	785	608	727	1,765	3,035	5,067	10,017	14,094	9,598	19
1961	53,038	8,948	2,064	721	515	703	1,676	2,809	4,709	9,307	12,793	8,780	13
1960	58,931	9,290	2,439	823	582	937	1,977	3,505	5,879	10,723	13,763	8,981	32
MALE													
1967	31,256	3,208	802	357	423	389	1,029	2,125	3,431	6,256	8,454	4,770	12
1966	34,221	3,917	925	390	421	428	1,172	2,251	3,915	6,878	8,900	5,000	24
1965	33,909	4,508	928	444	354	427	1,091	2,126	3,655	6,766	8,791	4,803	16
1964	32,741	4,666	1,053	370	341	423	1,027	2,135	3,512	6,491	8,188	4,512	23
1963	35,985	4,892	1,184	409	325	413	1,136	2,383	4,023	7,553	8,745	4,908	14
1962	31,853	4,966	1,153	405	337	388	1,020	2,001	3,379	6,355	7,647	4,191	11
1961	30,144	5,000	1,111	384	304	388	1,011	1,888	3,177	6,032	6,972	3,868	9
1960	34,030	5,301	1,308	445	331	487	1,225	2,395	4,037	6,971	7,538	3,971	21
FEMALE													
1967	24,161	2,429	611	319	296	336	641	1,056	1,656	3,433	7,129	6,246	9
1966	26,564	2,915	794	366	312	333	753	1,262	1,876	3,916	7,516	6,510	11
1965	25,699	3,346	904	359	271	346	767	1,111	1,798	3,709	7,025	6,054	9
1964	25,023	3,522	941	355	268	359	681	1,058	1,730	3,561	6,830	5,713	5
1963	27,693	3,666	1,052	401	285	431	748	1,178	2,015	4,193	7,448	6,261	15
1962	24,711	3,735	995	380	271	339	745	1,034	1,688	3,662	6,447	5,407	8
1961	22,894	3,948	953	337	211	315	665	921	1,532	3,275	5,821	4,912	4
1960	24,901	3,989	1,131	378	251	450	752	1,110	1,842	3,752	6,225	5,010	11

Note: Figures by race exclude data for residents of New Jersey for 1962 and 1963.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290A for 1960-67 and separate unpublished tables for the black population and Accidents.

77/09/02 TABLE 290A. DEATHS FOR 60 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1960-67

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	TOTAL	UNDER 1 YEAR	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS	65-74 YEARS	75-84 YEARS	85+ YEARS	N.S.
Pneumonia, except pneumonia of newborn (490-493)													
WHITE, BOTH SEXES													
1967	46,515	3,344	868	511	552	444	1,049	2,314	4,129	8,321	14,506	10,464	13
1966	50,322	3,938	1,055	573	573	446	1,190	2,635	4,599	9,219	15,198	10,874	22
1965	49,409	4,674	1,143	591	488	479	1,176	2,460	4,328	9,025	14,694	10,336	15
1964	47,714	5,006	1,228	539	470	448	1,042	2,365	4,201	8,728	13,966	9,705	16
1963	50,484	4,977	1,341	590	460	517	1,164	2,453	4,508	9,659	14,526	10,272	17
1962	45,026	5,158	1,282	577	460	436	1,115	2,136	3,881	8,350	12,723	8,895	13
1961	43,707	5,687	1,347	570	387	431	1,097	2,061	3,732	8,087	11,996	8,308	4
1960	47,860	5,583	1,585	627	446	597	1,273	2,555	4,631	9,279	12,769	8,493	22
WHITE, MALE													
1967	25,895	1,964	480	280	323	233	643	1,521	2,796	5,354	7,797	4,497	7
1966	28,043	2,341	581	308	344	238	731	1,650	3,138	5,850	8,179	4,669	14
1965	27,967	2,793	584	336	282	255	683	1,588	2,935	5,832	8,115	4,555	9
1964	26,852	2,932	646	265	270	233	614	1,599	2,830	5,652	7,551	4,248	12
1963	28,376	2,905	704	298	248	260	690	1,672	3,055	6,256	7,779	4,501	8
1962	25,271	3,028	690	291	267	230	665	1,410	2,624	5,314	6,882	3,862	8
1961	24,688	3,257	744	306	238	227	654	1,373	2,545	5,230	6,488	3,624	2
1960	27,486	3,278	822	346	257	321	805	1,751	3,216	6,031	6,919	3,726	14
WHITE, FEMALE													
1967	20,620	1,380	388	231	229	211	406	793	1,333	2,967	6,709	5,967	6
1966	22,279	1,597	474	265	229	208	459	985	1,461	3,369	7,019	6,205	8
1965	21,442	1,881	559	255	206	224	493	872	1,393	3,193	6,579	5,781	6
1964	20,862	2,074	582	274	200	215	428	766	1,371	3,076	6,415	5,457	4
1963	22,108	2,072	637	292	212	257	474	781	1,453	3,403	6,747	5,771	9
1962	19,755	2,130	592	286	193	206	450	726	1,257	3,036	5,841	5,033	5
1961	19,019	2,430	603	264	149	204	443	688	1,187	2,857	5,508	4,684	2
1960	20,374	2,305	763	281	189	276	468	804	1,415	3,248	5,850	4,767	8

Note: Figures by race exclude data for residents of New Jersey for 1962 and 1963.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290A for 1960-67 and separate unpublished tables for the black population and Accidents.

77/09/02 TABLE 290A. DEATHS FOR 60 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1960-67

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	TOTAL	UNDER 1 YEAR	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS	65-74 YEARS	75-84 YEARS	85+ YEARS	N.S.
Pneumonia, except pneumonia of newborn (490-493)													
ALL OTHER, BOTH SEXES													
1967	8,902	2,293	545	165	167	281	621	867	958	1,368	1,077	552	8
1966	10,463	2,894	664	183	160	315	735	878	1,192	1,575	1,218	636	13
1965	10,199	3,180	689	212	137	294	682	777	1,125	1,450	1,122	521	10
1964	10,050	3,182	766	186	139	334	666	828	1,041	1,324	1,052	520	12
1963	11,352	3,367	841	186	129	299	666	980	1,356	1,720	1,181	615	12
1962	9,835	3,290	794	176	122	262	582	802	1,022	1,341	961	477	6
1961	9,331	3,261	717	151	128	272	579	748	977	1,220	797	472	9
1960	11,071	3,707	854	196	136	340	704	950	1,248	1,444	994	488	10
ALL OTHER, MALE													
1967	5,361	1,244	322	77	100	156	386	604	635	902	657	273	5
1966	6,178	1,576	344	82	77	190	441	601	777	1,028	721	331	10
1965	5,942	1,715	344	108	72	172	408	538	720	934	676	248	7
1964	5,889	1,734	407	105	71	190	413	536	682	839	637	264	11
1963	6,553	1,859	447	95	66	140	410	628	853	1,065	697	287	6
1962	5,630	1,791	428	95	58	149	313	521	640	843	558	231	3
1961	5,456	1,743	367	78	66	161	357	515	632	802	484	244	7
1960	6,544	2,023	486	99	74	166	420	644	821	940	619	245	7
ALL OTHER, FEMALE													
1967	3,541	1,049	223	88	67	125	235	263	323	466	420	279	3
1966	4,285	1,318	320	101	83	125	294	277	415	547	497	305	3
1965	4,257	1,465	345	104	65	122	274	239	405	516	446	273	3
1964	4,161	1,448	359	81	68	144	253	292	359	485	415	256	1
1963	4,799	1,508	394	91	63	159	256	352	503	655	484	328	6
1962	4,205	1,499	366	81	64	113	269	281	382	498	403	246	3
1961	3,875	1,518	350	73	62	111	222	233	345	418	313	228	2
1960	4,527	1,684	368	97	62	174	284	306	427	504	375	243	3

Note: Figures by race exclude data for residents of New Jersey for 1962 and 1963.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290A for 1960-67 and separate unpublished tables for the black population and Accidents.

77/09/02 TABLE 290A. DEATHS FOR 60 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1960-67

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	TOTAL	UNDER 1 YEAR	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS	65-74 YEARS	75-84 YEARS	85+ YEARS	N.S.
Bronchitis (500-502)													
BOTH SEXES													
1967	6,264	314	214	64	46	45	140	436	1,166	1,814	1,520	503	2
1966	6,151	352	184	62	38	38	140	462	1,107	1,813	1,471	484	-
1965	5,772	470	187	68	32	39	133	415	1,044	1,681	1,278	424	1
1964	5,410	494	244	82	26	38	141	388	911	1,524	1,154	408	-
1963	5,462	547	236	74	43	47	134	349	872	1,575	1,140	444	1
1962	4,665	579	242	74	36	40	113	323	751	1,231	899	375	2
1961	4,064	599	266	70	29	33	104	266	661	1,038	732	265	1
1960	4,343	656	346	69	36	30	103	310	656	1,088	744	304	1
MALE													
1967	4,598	197	121	33	27	22	78	308	901	1,468	1,164	278	1
1966	4,523	203	108	37	21	19	76	316	850	1,476	1,126	291	-
1965	4,238	273	103	46	19	18	78	283	823	1,387	949	259	-
1964	3,941	278	152	47	14	14	75	267	738	1,265	845	246	-
1963	3,947	331	137	37	19	20	63	254	741	1,309	806	229	1
1962	3,321	342	147	45	25	16	64	229	603	1,017	627	205	1
1961	2,868	351	167	32	10	21	66	195	516	846	522	141	1
1960	3,036	404	194	40	11	18	62	228	523	887	517	151	1
FEMALE													
1967	1,666	117	93	31	19	23	62	128	265	346	356	225	1
1966	1,628	149	76	25	17	19	64	146	257	337	345	193	-
1965	1,534	197	84	22	13	21	55	132	221	294	329	165	1
1964	1,469	216	92	35	12	24	66	121	173	259	309	162	-
1963	1,515	216	99	37	24	27	71	95	131	266	334	215	-
1962	1,344	237	95	29	11	24	49	94	148	214	272	170	1
1961	1,196	248	99	38	19	12	38	71	145	192	210	124	-
1960	1,307	252	152	29	25	12	41	82	133	201	227	153	-

- Quantity zero

Note: Figures by race exclude data for residents of New Jersey for 1962 and 1963.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290A for 1960-67 and separate unpublished tables for the black population and Accidents.

77/09/02 TABLE 290A. DEATHS FOR 60 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1960-67

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	TOTAL	UNDER 1 YEAR	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS	65-74 YEARS	75-84 YEARS	85+ YEARS	N.S.
Bronchitis (500-502)													
WHITE, BOTH SEXES													
1967	5,740	204	174	52	36	31	106	385	1,085	1,714	1,466	486	1
1966	5,653	232	148	54	32	29	98	407	1,033	1,723	1,424	473	-
1965	5,252	319	145	48	26	28	105	374	973	1,590	1,235	408	1
1964	4,935	364	196	73	18	27	103	347	845	1,458	1,110	394	-
1963	4,780	400	176	58	38	33	102	293	788	1,417	1,058	416	1
1962	4,033	409	178	65	27	24	90	278	684	1,109	824	343	2
1961	3,635	465	213	60	23	26	78	236	603	974	706	250	1
1960	3,889	513	287	64	28	24	84	270	598	1,018	714	289	-
WHITE, MALE													
1967	4,243	141	102	29	22	15	56	265	837	1,390	1,119	267	-
1966	4,190	137	85	32	18	14	52	276	793	1,406	1,089	288	-
1965	3,883	192	79	33	15	15	64	248	768	1,311	911	247	-
1964	3,637	214	123	44	10	9	56	239	687	1,209	809	237	-
1963	3,492	252	102	30	17	13	49	209	673	1,181	753	212	1
1962	2,899	246	109	42	18	7	50	196	553	920	572	185	1
1961	2,592	280	128	26	10	15	52	170	474	795	505	136	1
1960	2,758	333	165	38	9	15	49	197	476	833	499	144	-
WHITE, FEMALE													
1967	1,497	63	72	23	14	16	50	120	248	324	347	219	1
1966	1,463	95	63	22	14	15	46	131	240	317	335	185	-
1965	1,369	127	66	15	11	13	41	126	205	279	324	161	1
1964	1,298	150	73	29	8	18	47	108	158	249	301	157	-
1963	1,288	148	74	28	21	20	53	84	115	236	305	204	-
1962	1,134	163	69	23	9	17	40	82	131	189	252	158	1
1961	1,043	185	85	34	13	11	26	66	129	179	201	114	-
1960	1,131	180	122	26	19	9	35	73	122	185	215	145	-

- Quantity zero

Note: Figures by race exclude data for residents of New Jersey for 1962 and 1963.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290A for 1960-67 and separate unpublished tables for the black population and Accidents.

77/09/02 TABLE 290A. DEATHS FOR 60 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1960-67

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	TOTAL	UNDER 1 YEAR	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS	65-74 YEARS	75-84 YEARS	85+ YEARS	N.S.
Bronchitis (500-502)													
ALL OTHER, BOTH SEXES													
1967	524	110	40	12	10	14	34	51	81	100	54	17	1
1966	498	120	36	8	6	9	42	55	74	90	47	11	-
1965	520	151	42	20	6	11	28	41	71	91	43	16	-
1964	475	130	48	9	8	11	38	41	66	66	44	14	-
1963	512	138	55	12	5	12	28	41	59	101	39	22	-
1962	471	152	59	8	9	15	17	38	45	74	38	16	-
1961	429	134	53	10	6	7	26	30	58	64	26	15	-
1960	454	143	59	5	8	6	19	40	58	70	30	15	1
ALL OTHER, MALE													
1967	355	56	19	4	5	7	22	43	64	78	45	11	1
1966	333	66	23	5	3	5	24	40	57	70	37	3	-
1965	355	81	24	13	4	3	14	35	55	76	38	12	-
1964	304	64	29	3	4	5	19	28	51	56	36	9	-
1963	328	74	33	5	2	5	13	33	47	80	23	13	-
1962	305	85	36	2	7	9	10	27	33	57	28	11	-
1961	276	71	39	6	-	6	14	25	42	51	17	5	-
1960	278	71	29	2	2	3	13	31	47	54	18	7	1
ALL OTHER, FEMALE													
1967	169	54	21	8	5	7	12	8	17	22	9	6	-
1966	165	54	13	3	3	4	18	15	17	20	10	8	-
1965	165	70	18	7	2	8	14	6	16	15	5	4	-
1964	171	66	19	6	4	6	19	13	15	10	8	5	-
1963	184	64	22	7	3	7	15	8	12	21	16	9	-
1962	166	67	23	6	2	6	7	11	12	17	10	5	-
1961	153	63	14	4	6	1	12	5	16	13	9	10	-
1960	176	72	30	3	6	3	6	9	11	16	12	8	-

- Quantity zero

Note: Figures by race exclude data for residents of New Jersey for 1962 and 1963.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290A for 1960-67 and separate unpublished tables for the black population and Accidents.

77/09/02 TABLE 290A. DEATHS FOR 60 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1960-67

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	TOTAL	UNDER 1 YEAR	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS	65-74 YEARS	75-84 YEARS	85+ YEARS	N.S.
Other bronchopulmonic diseases (525-527)													
BOTH SEXES													
1967	29,360	1,198	136	80	82	122	511	2,053	6,255	10,117	7,149	1,653	4
1966	28,371	1,355	177	54	82	134	551	2,045	6,083	9,714	6,671	1,501	4
1965	26,518	1,366	161	56	72	110	530	2,048	5,579	9,091	6,100	1,402	3
1964	23,427	1,268	159	68	79	154	509	1,851	5,026	7,997	5,148	1,163	5
1963	23,205	1,369	185	53	71	154	510	1,893	5,015	7,958	4,828	1,161	8
1962	20,072	1,307	179	72	78	143	506	1,627	4,330	6,704	4,165	954	7
1961	17,832	1,246	185	74	73	134	455	1,401	3,964	5,859	3,577	857	7
1960	17,196	1,219	193	80	76	140	494	1,508	3,785	5,704	3,258	735	4
MALE													
1967	23,556	668	73	41	46	63	327	1,486	5,173	8,691	5,831	1,154	3
1966	22,798	789	96	27	43	70	338	1,486	5,128	8,280	5,504	1,033	4
1965	21,155	783	83	20	40	60	318	1,526	4,684	7,820	4,869	950	2
1964	18,719	735	84	31	51	76	315	1,419	4,245	6,834	4,128	796	5
1963	18,517	784	99	32	37	79	332	1,418	4,265	6,835	3,835	793	8
1962	15,838	765	108	44	41	77	314	1,203	3,684	5,780	3,187	631	4
1961	13,992	728	100	27	36	70	301	1,035	3,403	5,000	2,742	545	5
1960	13,497	740	112	39	43	81	307	1,174	3,215	4,836	2,482	465	3
FEMALE													
1967	5,804	530	63	39	36	59	184	567	1,082	1,426	1,318	499	1
1966	5,573	566	81	27	39	64	213	559	955	1,434	1,167	468	-
1965	5,363	583	78	36	32	50	212	522	895	1,271	1,231	452	1
1964	4,708	533	75	37	28	78	194	432	781	1,163	1,020	367	-
1963	4,688	585	86	21	34	75	178	475	750	1,123	993	368	-
1962	4,234	542	71	28	37	66	192	424	646	924	978	323	3
1961	3,840	518	85	47	37	64	154	366	561	859	835	312	2
1960	3,699	479	81	41	33	59	187	334	570	868	776	270	1

- Quantity zero

Note: Figures by race exclude data for residents of New Jersey for 1962 and 1963.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290A for 1960-67 and separate unpublished tables for the black population and Accidents.

77/09/02 TABLE 290A. DEATHS FOR 60 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1960-67

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	TOTAL	UNDER 1 YEAR	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS	65-74 YEARS	75-84 YEARS	85+ YEARS	N.S.
Other bronchopulmonic diseases (525-527)													
WHITE, BOTH SEXES													
1967	27,142	766	86	62	66	88	400	1,791	5,819	9,618	6,879	1,563	4
1966	26,220	851	123	40	63	103	433	1,774	5,696	9,250	6,447	1,436	4
1965	24,530	898	117	47	58	87	435	1,767	5,184	8,716	5,881	1,337	3
1964	21,662	851	106	60	62	118	407	1,641	4,675	7,642	4,977	1,119	4
1963	20,828	938	124	39	52	117	390	1,613	4,551	7,370	4,541	1,087	6
1962	17,998	885	119	56	56	92	398	1,416	3,927	6,221	3,926	895	7
1961	16,358	860	123	61	52	102	366	1,244	3,695	5,590	3,440	819	6
1960	15,813	846	139	68	61	111	401	1,324	3,545	5,448	3,163	704	3
WHITE, MALE													
1967	21,925	450	45	31	36	42	257	1,281	4,821	8,263	5,609	1,087	3
1966	21,248	526	64	20	32	56	267	1,285	4,803	7,891	5,315	985	4
1965	19,694	528	58	19	31	47	256	1,315	4,362	7,496	4,683	897	2
1964	17,448	520	59	27	41	56	250	1,266	3,948	6,527	3,984	766	4
1963	16,759	545	71	22	27	63	260	1,208	3,878	6,330	3,610	739	6
1962	14,335	538	72	34	31	49	249	1,058	3,343	5,362	3,002	593	4
1961	12,928	510	63	19	25	52	236	910	3,177	4,775	2,641	516	4
1960	12,523	543	85	34	36	66	242	1,028	3,014	4,621	2,404	448	2
WHITE, FEMALE													
1967	5,217	316	41	31	30	46	143	510	998	1,355	1,270	476	1
1966	4,972	325	59	20	31	47	166	489	893	1,359	1,132	451	-
1965	4,836	370	59	28	27	40	179	452	822	1,220	1,198	440	1
1964	4,214	331	47	33	21	62	157	375	727	1,115	993	353	-
1963	4,069	393	53	17	25	54	130	405	673	1,040	931	348	-
1962	3,663	347	47	22	25	43	149	358	584	859	924	302	3
1961	3,430	350	60	42	27	50	130	334	518	815	799	303	2
1960	3,290	303	54	34	25	45	159	296	531	827	759	256	1

- Quantity zero

Note: Figures by race exclude data for residents of New Jersey for 1962 and 1963.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290A for 1960-67 and separate unpublished tables for the black population and Accidents.

77/09/02 TABLE 290A. DEATHS FOR 60 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1960-67

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	TOTAL	UNDER 1 YEAR	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS	65-74 YEARS	75-84 YEARS	85+ YEARS	N.S.
Other bronchopulmonic diseases (525-527)													
ALL OTHER, BOTH SEXES													
1967	2,218	432	50	18	16	34	111	262	436	499	270	90	-
1966	2,151	504	54	14	19	31	118	271	387	464	224	65	-
1965	1,988	468	44	9	14	23	95	281	395	375	219	65	-
1964	1,765	417	53	8	17	36	102	210	351	355	171	44	1
1963	1,806	415	58	10	15	30	105	238	334	364	180	56	1
1962	1,564	408	58	13	20	45	95	173	281	304	128	39	-
1961	1,474	386	62	13	21	32	89	157	269	269	137	38	1
1960	1,383	373	54	12	15	29	93	184	240	256	95	31	1
ALL OTHER, MALE													
1967	1,631	218	28	10	10	21	70	205	352	428	222	67	-
1966	1,550	263	32	7	11	14	71	201	325	389	189	48	-
1965	1,461	255	25	1	9	13	62	211	322	324	186	53	-
1964	1,271	215	25	4	10	20	65	153	297	307	144	30	1
1963	1,281	227	26	6	8	13	60	180	272	308	139	41	1
1962	1,089	217	34	8	10	25	57	120	235	262	95	26	-
1961	1,064	218	37	8	11	18	65	125	226	225	101	29	1
1960	974	197	27	5	7	15	65	146	201	215	78	17	1
ALL OTHER, FEMALE													
1967	587	214	22	8	6	13	41	57	84	71	48	23	-
1966	601	241	22	7	8	17	47	70	62	75	35	17	-
1965	527	213	19	8	5	10	33	70	73	51	33	12	-
1964	494	202	28	4	7	16	37	57	54	48	27	14	-
1963	525	188	32	4	7	17	45	58	62	56	41	15	-
1962	475	191	24	5	10	20	38	53	46	42	33	13	-
1961	410	168	25	5	10	14	24	32	43	44	36	9	-
1960	409	176	27	7	8	14	28	38	39	41	17	14	-

- Quantity zero

Note: Figures by race exclude data for residents of New Jersey for 1962 and 1963.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290A for 1960-67 and separate unpublished tables for the black population and Accidents.

77/09/02 TABLE 290A. DEATHS FOR 60 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1960-67

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	TOTAL	UNDER 1 YEAR	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS	65-74 YEARS	75-84 YEARS	85+ YEARS	N.S.
Ulcer of stomach and duodenum (540,541)													
BOTH SEXES													
1967	9,825	56	14	14	57	113	430	1,101	2,028	2,600	2,468	942	2
1966	10,321	56	7	14	54	119	463	1,180	2,060	2,854	2,621	891	2
1965	10,424	75	12	11	51	117	460	1,238	2,148	2,841	2,588	879	4
1964	10,969	64	10	14	43	129	538	1,271	2,271	3,034	2,708	886	1
1963	12,161	70	19	16	44	173	608	1,511	2,434	3,426	2,924	932	4
1962	12,228	65	23	19	48	166	632	1,473	2,487	3,508	2,877	929	1
1961	11,540	56	12	11	47	189	585	1,538	2,390	3,312	2,600	798	2
1960	11,360	78	18	15	29	167	601	1,501	2,448	3,290	2,485	724	4
MALE													
1967	6,783	32	9	10	40	79	304	798	1,566	1,900	1,550	493	2
1966	7,187	33	5	5	34	79	338	857	1,565	2,074	1,727	469	1
1965	7,361	43	7	6	36	80	327	914	1,670	2,107	1,687	480	4
1964	7,809	39	7	10	29	87	375	929	1,802	2,269	1,759	502	1
1963	8,779	49	14	8	27	129	433	1,154	1,922	2,576	1,954	509	4
1962	8,836	45	14	14	31	129	462	1,116	1,946	2,645	1,922	512	-
1961	8,459	36	6	3	34	147	438	1,166	1,896	2,520	1,753	459	1
1960	8,507	57	9	5	21	122	455	1,173	1,997	2,530	1,720	414	4
FEMALE													
1967	3,042	24	5	4	17	34	126	303	462	700	918	449	-
1966	3,134	23	2	9	20	40	125	323	495	780	894	422	1
1965	3,063	32	5	5	15	37	133	324	478	734	901	399	-
1964	3,160	25	3	4	14	42	163	342	469	765	949	384	-
1963	3,382	21	5	8	17	44	175	357	512	850	970	423	-
1962	3,392	20	9	5	17	37	170	357	541	863	955	417	1
1961	3,081	20	6	8	13	42	147	372	494	792	847	339	1
1960	2,853	21	9	10	8	45	146	328	451	760	765	310	-

- Quantity zero

Note: Figures by race exclude data for residents of New Jersey for 1962 and 1963.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290A for 1960-67 and separate unpublished tables for the black population and Accidents.

77/09/02 TABLE 290A. DEATHS FOR 60 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1960-67

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	TOTAL	UNDER 1 YEAR	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS	65-74 YEARS	75-84 YEARS	85+ YEARS	N.S.
Ulcer of stomach and duodenum (540,541)													
WHITE, BOTH SEXES													
1967	8,934	36	10	8	42	84	329	938	1,827	2,428	2,333	898	1
1966	9,362	31	5	10	45	95	358	990	1,837	2,629	2,507	854	1
1965	9,536	38	11	8	39	89	371	1,060	1,944	2,655	2,463	854	4
1964	9,990	31	10	14	37	94	426	1,073	2,045	2,820	2,584	856	-
1963	10,748	39	15	12	31	127	483	1,262	2,080	3,083	2,731	884	1
1962	10,800	39	16	17	35	116	497	1,236	2,128	3,157	2,683	875	1
1961	10,576	33	8	11	36	138	493	1,336	2,169	3,092	2,494	765	1
1960	10,447	43	14	11	26	128	491	1,311	2,238	3,080	2,400	702	3
WHITE, MALE													
1967	6,165	23	7	7	32	60	235	671	1,422	1,776	1,462	469	1
1966	6,505	20	3	3	29	61	263	731	1,393	1,912	1,642	447	1
1965	6,734	22	7	4	24	60	265	782	1,524	1,972	1,603	467	4
1964	7,129	21	7	10	24	68	306	795	1,624	2,111	1,676	487	-
1963	7,757	26	11	6	18	97	343	974	1,654	2,329	1,810	488	1
1962	7,778	27	10	12	21	86	365	932	1,679	2,381	1,783	482	-
1961	7,769	20	5	3	26	106	369	1,019	1,742	2,356	1,680	442	1
1960	7,824	33	7	2	19	95	376	1,026	1,834	2,370	1,659	400	3
WHITE, FEMALE													
1967	2,769	13	3	1	10	24	94	267	405	652	871	429	-
1966	2,857	11	2	7	16	34	95	259	444	717	865	407	-
1965	2,802	16	4	4	15	29	106	278	420	683	860	387	-
1964	2,861	10	3	4	13	26	120	278	421	709	908	369	-
1963	2,991	13	4	6	13	30	140	288	426	754	921	396	-
1962	3,022	12	6	5	14	30	132	304	449	776	900	393	1
1961	2,807	13	3	8	10	32	124	317	427	736	814	323	-
1960	2,623	10	7	9	7	33	115	285	404	710	741	302	-

- Quantity zero

Note: Figures by race exclude data for residents of New Jersey for 1962 and 1963.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290A for 1960-67 and separate unpublished tables for the black population and Accidents.

77/09/02 TABLE 290A. DEATHS FOR 60 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1960-67

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	TOTAL	UNDER 1 YEAR	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS	65-74 YEARS	75-84 YEARS	85+ YEARS	N.S.
Ulcer of stomach and duodenum (540,541)													
ALL OTHER, BOTH SEXES													
1967	891	20	4	6	15	29	101	163	201	172	135	44	1
1966	959	25	2	4	9	24	105	190	223	225	114	37	1
1965	888	37	1	3	12	28	89	178	204	186	125	25	-
1964	979	33	-	-	6	35	112	198	226	214	124	30	1
1963	982	27	4	4	12	40	98	189	244	232	105	24	3
1962	969	21	3	2	9	43	113	173	252	229	95	29	-
1961	964	23	4	-	11	51	92	202	221	220	106	33	1
1960	913	35	4	4	3	39	110	190	210	210	85	22	1
ALL OTHER, MALE													
1967	618	9	2	3	8	19	69	127	144	124	88	24	1
1966	682	13	2	2	5	18	75	126	172	162	85	22	-
1965	627	21	-	2	12	20	62	132	146	135	84	13	-
1964	680	18	-	-	5	19	69	134	178	158	83	15	1
1963	704	19	3	2	8	27	72	137	178	165	79	11	3
1962	719	15	2	2	6	38	81	135	182	175	66	17	-
1961	690	16	1	-	8	41	69	147	154	164	73	17	-
1960	683	24	2	3	2	27	79	147	163	160	61	14	1
ALL OTHER, FEMALE													
1967	273	11	2	3	7	10	32	36	57	48	47	20	-
1966	277	12	-	2	4	6	30	64	51	63	29	15	1
1965	261	16	1	1	-	8	27	46	58	51	41	12	-
1964	299	15	-	-	1	16	43	64	48	56	41	15	-
1963	278	8	1	2	4	13	26	52	66	67	26	13	-
1962	250	6	1	-	3	5	32	38	70	54	29	12	-
1961	274	7	3	-	3	10	23	55	67	56	33	16	1
1960	230	11	2	1	1	12	31	43	47	50	24	8	-

- Quantity zero

Note: Figures by race exclude data for residents of New Jersey for 1962 and 1963.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290A for 1960-67 and separate unpublished tables for the black population and Accidents.

77/09/02 TABLE 290A. DEATHS FOR 60 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1960-67

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	TOTAL	UNDER 1 YEAR	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS	65-74 YEARS	75-84 YEARS	85+ YEARS	N.S.
Appendicitis (550-553)													
BOTH SEXES													
1967	1,526	11	27	72	57	47	95	173	253	324	367	100	-
1966	1,627	19	36	113	75	63	94	162	263	335	371	96	-
1965	1,694	15	40	118	76	43	89	178	277	386	355	117	-
1964	1,783	14	32	131	75	64	110	194	279	401	375	108	-
1963	1,749	22	41	110	68	65	116	168	297	396	353	112	1
1962	1,800	21	43	144	86	62	110	197	281	405	346	105	-
1961	1,819	19	58	125	82	69	111	208	307	387	332	121	-
1960	1,871	13	50	143	76	60	123	198	340	443	331	92	2
MALE													
1967	928	8	15	40	27	30	66	113	178	193	211	47	-
1966	965	12	21	64	43	32	53	113	171	211	204	41	-
1965	1,072	10	25	78	48	20	64	121	176	253	226	51	-
1964	1,109	8	19	84	41	41	61	136	199	257	226	37	-
1963	1,066	14	24	64	36	33	74	107	191	259	208	55	1
1962	1,095	10	24	82	50	42	70	139	189	251	192	46	-
1961	1,106	9	37	69	49	37	76	140	219	238	180	52	-
1960	1,172	7	30	74	47	39	68	147	238	282	198	42	-
FEMALE													
1967	598	3	12	32	30	17	29	60	75	131	156	53	-
1966	662	7	15	49	32	31	41	49	92	124	167	55	-
1965	622	5	15	40	28	23	25	57	101	133	129	66	-
1964	674	6	13	47	34	23	49	58	80	144	149	71	-
1963	683	8	17	46	32	32	42	61	106	137	145	57	-
1962	705	11	19	62	36	20	40	58	92	154	154	59	-
1961	713	10	21	56	33	32	35	68	88	149	152	69	-
1960	699	6	20	69	29	21	55	51	102	161	133	50	2

- Quantity zero

Note: Figures by race exclude data for residents of New Jersey for 1962 and 1963.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290A for 1960-67 and separate unpublished tables for the black population and Accidents.

77/09/02 TABLE 290A. DEATHS FOR 60 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1960-67

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	TOTAL	UNDER 1 YEAR	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS	65-74 YEARS	75-84 YEARS	85+ YEARS	N.S.
Appendicitis (550-553)													
WHITE, BOTH SEXES													
1967	1,270	7	20	57	35	40	69	131	203	278	334	96	-
1966	1,355	15	28	85	62	50	61	126	208	290	341	89	-
1965	1,414	10	30	93	56	32	70	132	231	328	328	104	-
1964	1,479	11	28	101	58	46	76	137	230	349	340	103	-
1963	1,404	18	33	81	47	46	81	125	218	345	308	102	-
1962	1,454	17	33	109	67	46	75	149	216	334	310	98	-
1961	1,497	17	44	95	68	46	68	152	243	341	307	116	-
1960	1,562	12	40	109	62	42	90	156	270	385	310	85	1
WHITE, MALE													
1967	768	6	10	31	18	26	44	85	148	166	188	46	-
1966	814	9	14	49	38	26	41	91	139	184	186	37	-
1965	895	8	19	62	37	13	53	90	152	210	204	47	-
1964	923	5	17	64	31	34	44	99	165	226	203	35	-
1963	862	10	21	53	25	26	53	81	140	224	181	48	-
1962	904	8	17	66	38	33	52	111	149	214	174	42	-
1961	916	8	28	53	42	25	51	107	177	212	166	47	-
1960	988	7	25	54	40	28	56	120	191	242	187	38	-
WHITE, FEMALE													
1967	502	1	10	26	17	14	25	46	55	112	146	50	-
1966	541	6	14	36	24	24	20	35	69	106	155	52	-
1965	519	2	11	31	19	19	17	42	79	118	124	57	-
1964	556	6	11	37	27	12	32	38	65	123	137	68	-
1963	542	8	12	28	22	20	28	44	78	121	127	54	-
1962	550	9	16	43	29	13	23	38	67	120	136	56	-
1961	581	9	16	42	26	21	17	45	66	129	141	69	-
1960	574	5	15	55	22	14	34	36	79	143	123	47	1

- Quantity zero

Note: Figures by race exclude data for residents of New Jersey for 1962 and 1963.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290A for 1960-67 and separate unpublished tables for the black population and Accidents.

77/09/02 TABLE 290A. DEATHS FOR 60 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1960-67

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	TOTAL	UNDER 1 YEAR	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS	65-74 YEARS	75-84 YEARS	85+ YEARS	N.S.
Appendicitis (550-553)													
ALL OTHER, BOTH SEXES													
1967	256	4	7	15	22	7	26	42	50	46	33	4	-
1966	272	4	8	28	13	13	33	36	55	45	30	7	-
1965	280	5	10	25	20	11	19	46	46	58	27	13	-
1964	304	3	4	30	17	18	34	57	49	52	35	5	-
1963	290	4	7	28	18	15	27	36	71	39	37	7	1
1962	279	4	9	34	16	14	30	38	52	56	19	7	-
1961	322	2	14	30	14	23	43	56	64	46	25	5	-
1960	309	1	10	34	14	18	33	42	70	58	21	7	1
ALL OTHER, MALE													
1967	160	2	5	9	9	4	22	28	30	27	23	1	-
1966	151	3	7	15	5	6	12	22	32	27	18	4	-
1965	177	2	6	16	11	7	11	31	24	43	22	4	-
1964	186	3	2	20	10	7	17	37	34	31	23	2	-
1963	172	4	3	10	9	5	17	23	46	25	24	5	1
1962	149	2	6	16	9	7	14	21	30	29	11	4	-
1961	190	1	9	16	7	12	25	33	42	26	14	5	-
1960	184	-	5	20	7	11	12	27	47	40	11	4	-
ALL OTHER, FEMALE													
1967	96	2	2	6	13	3	4	14	20	19	10	3	-
1966	121	1	1	13	8	7	21	14	23	18	12	3	-
1965	103	3	4	9	9	4	8	15	22	15	5	9	-
1964	118	-	2	10	7	11	17	20	15	21	12	3	-
1963	118	-	4	18	9	10	10	13	25	14	13	2	-
1962	130	2	3	18	7	7	16	17	22	27	8	3	-
1961	132	1	5	14	7	11	18	23	22	20	11	-	-
1960	125	1	5	14	7	7	21	15	23	18	10	3	1

- Quantity zero

Note: Figures by race exclude data for residents of New Jersey for 1962 and 1963.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290A for 1960-67 and separate unpublished tables for the black population and Accidents.

77/09/02 TABLE 290A. DEATHS FOR 60 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1960-67

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	TOTAL	UNDER 1 YEAR	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS	65-74 YEARS	75-84 YEARS	85+ YEARS	N.S.
Hernia and intestinal obstruction (560,561,570)													
BOTH SEXES													
1967	9,814	685	61	54	68	81	263	629	1,296	2,377	2,975	1,320	5
1966	10,078	716	70	40	65	105	255	691	1,299	2,527	2,967	1,342	1
1965	10,003	852	70	50	60	108	299	694	1,401	2,378	2,806	1,281	4
1964	9,944	860	71	45	76	90	262	657	1,279	2,440	2,931	1,232	1
1963	9,917	910	80	43	57	107	292	688	1,371	2,408	2,778	1,179	4
1962	9,723	889	79	47	39	120	321	704	1,321	2,378	2,692	1,130	3
1961	9,177	931	82	65	61	116	277	640	1,275	2,206	2,479	1,041	4
1960	9,081	922	69	49	49	108	325	716	1,301	2,212	2,308	1,016	6
MALE													
1967	4,462	402	46	33	42	38	122	294	670	1,126	1,234	451	4
1966	4,673	430	41	20	30	56	121	318	699	1,241	1,251	466	-
1965	4,668	492	47	37	27	50	125	318	718	1,197	1,191	463	3
1964	4,658	524	40	30	43	39	115	319	646	1,243	1,207	451	1
1963	4,724	574	58	29	29	43	121	311	721	1,223	1,202	411	2
1962	4,715	521	49	31	15	53	131	351	688	1,229	1,179	466	2
1961	4,513	559	54	44	34	49	125	297	671	1,124	1,121	433	2
1960	4,466	542	47	27	23	53	148	357	702	1,147	1,038	380	2
FEMALE													
1967	5,352	283	15	21	26	43	141	335	626	1,251	1,741	869	1
1966	5,405	286	29	20	35	49	134	373	600	1,286	1,716	876	1
1965	5,335	360	23	13	33	58	174	376	683	1,181	1,615	818	1
1964	5,286	336	31	15	33	51	147	338	633	1,197	1,724	781	-
1963	5,193	336	22	14	28	64	171	377	650	1,185	1,576	768	2
1962	5,008	368	30	16	24	67	190	353	633	1,149	1,513	664	1
1961	4,664	372	28	21	27	67	152	343	604	1,082	1,358	608	2
1960	4,615	380	22	22	26	55	177	359	599	1,065	1,270	636	4

- Quantity zero

Note: Figures by race exclude data for residents of New Jersey for 1962 and 1963.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290A for 1960-67 and separate unpublished tables for the black population and Accidents.

77/09/02 TABLE 290A. DEATHS FOR 60 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1960-67

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	TOTAL	UNDER 1 YEAR	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS	65-74 YEARS	75-84 YEARS	85+ YEARS	N.S.
Hernia and intestinal obstruction (560,561,570)													
WHITE, BOTH SEXES													
1967	8,689	567	48	41	45	55	185	487	1,076	2,120	2,800	1,262	3
1966	8,967	578	49	32	50	80	201	553	1,104	2,253	2,786	1,281	-
1965	8,847	710	48	38	49	80	205	551	1,176	2,106	2,661	1,221	2
1964	8,801	704	53	35	53	63	190	503	1,081	2,192	2,752	1,175	-
1963	8,456	729	59	32	39	64	202	522	1,092	2,073	2,547	1,097	-
1962	8,283	734	62	36	27	86	218	540	1,049	2,052	2,433	1,046	-
1961	8,089	785	62	51	49	75	182	498	1,057	1,997	2,341	989	3
1960	7,975	771	53	33	37	72	234	562	1,097	1,978	2,170	966	2
WHITE, MALE													
1967	3,856	327	38	26	28	24	79	222	555	988	1,142	425	2
1966	4,079	342	29	16	20	44	94	254	593	1,093	1,156	438	-
1965	4,084	409	31	28	23	41	86	259	597	1,052	1,122	435	1
1964	4,042	422	29	23	31	28	78	241	551	1,094	1,125	420	-
1963	3,974	451	43	21	21	23	87	244	571	1,058	1,083	372	-
1962	3,958	431	39	23	10	37	94	264	544	1,047	1,044	425	-
1961	3,928	461	39	35	27	33	82	241	555	1,012	1,039	402	2
1960	3,874	450	35	18	16	35	110	278	603	1,011	957	360	1
WHITE, FEMALE													
1967	4,833	240	10	15	17	31	106	265	521	1,132	1,658	837	1
1966	4,888	236	20	16	30	36	107	299	511	1,160	1,630	843	-
1965	4,763	301	17	10	26	39	119	292	579	1,054	1,539	786	1
1964	4,759	282	24	12	22	35	112	262	530	1,098	1,627	755	-
1963	4,482	278	16	11	18	41	115	278	521	1,015	1,464	725	-
1962	4,325	303	23	13	17	49	124	276	505	1,005	1,389	621	-
1961	4,161	324	23	16	22	42	100	257	502	985	1,302	587	1
1960	4,101	321	18	15	21	37	124	284	494	967	1,213	606	1

- Quantity zero

Note: Figures by race exclude data for residents of New Jersey for 1962 and 1963.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290A for 1960-67 and separate unpublished tables for the black population and Accidents.

77/09/02 TABLE 290A. DEATHS FOR 60 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1960-67

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	TOTAL	UNDER 1 YEAR	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS	65-74 YEARS	75-84 YEARS	85+ YEARS	N.S.
Hernia and intestinal obstruction (560,561,570)													
ALL OTHER, BOTH SEXES													
1967	1,125	118	13	13	23	26	78	142	220	257	175	58	2
1966	1,111	138	21	8	15	25	54	138	195	274	181	61	1
1965	1,156	142	22	12	11	28	94	143	225	272	145	60	2
1964	1,143	156	18	10	23	27	72	154	198	248	179	57	1
1963	1,142	156	17	9	13	35	75	144	235	256	148	50	4
1962	1,090	125	16	10	10	30	95	134	215	235	163	54	3
1961	1,088	146	20	14	12	41	95	142	218	209	138	52	1
1960	1,106	151	16	16	12	36	91	154	204	234	138	50	4
ALL OTHER, MALE													
1967	606	75	8	7	14	14	43	72	115	138	92	26	2
1966	594	88	12	4	10	12	27	64	106	148	95	28	-
1965	584	83	16	9	4	9	39	59	121	145	69	28	2
1964	616	102	11	7	12	11	37	78	95	149	82	31	1
1963	599	107	11	7	7	16	27	61	124	129	80	28	2
1962	584	74	9	8	5	14	35	69	111	136	90	31	2
1961	585	98	15	9	7	16	43	56	116	112	82	31	-
1960	592	92	12	9	7	18	38	79	99	136	81	20	1
ALL OTHER, FEMALE													
1967	519	43	5	6	9	12	35	70	105	119	83	32	-
1966	517	50	9	4	5	13	27	74	89	126	86	33	1
1965	572	59	6	3	7	19	55	84	104	127	76	32	-
1964	527	54	7	3	11	16	35	76	103	99	97	26	-
1963	543	49	6	2	6	19	48	83	111	127	68	22	2
1962	506	51	7	2	5	16	60	65	104	99	73	23	1
1961	503	48	5	5	5	25	52	86	102	97	56	21	1
1960	514	59	4	7	5	18	53	75	105	98	57	30	3

- Quantity zero

Note: Figures by race exclude data for residents of New Jersey for 1962 and 1963.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290A for 1960-67 and separate unpublished tables for the black population and Accidents.

77/09/02 TABLE 290A. DEATHS FOR 60 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1960-67

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	TOTAL	UNDER 1 YEAR	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS	65-74 YEARS	75-84 YEARS	85+ YEARS	N.S.
Gastritis, duodenitis, enteritis, and colitis, except diarrhea of newborn (543,571,572)													
BOTH SEXES													
1967	7,504	1,031	295	81	109	133	256	526	952	1,499	1,746	876	-
1966	7,552	1,352	342	72	91	133	232	555	909	1,481	1,617	768	-
1965	7,899	1,709	401	82	96	150	290	535	911	1,492	1,541	692	-
1964	8,178	2,024	463	91	135	146	291	517	878	1,436	1,499	698	-
1963	8,246	2,093	506	96	106	140	283	564	873	1,409	1,503	672	1
1962	8,194	2,342	503	133	119	137	253	522	861	1,327	1,390	606	1
1961	7,820	2,296	543	97	114	134	252	507	780	1,273	1,327	495	2
1960	7,932	2,622	528	87	100	155	267	484	742	1,257	1,176	512	2
MALE													
1967	3,489	578	168	42	55	66	137	286	537	670	669	281	-
1966	3,559	753	192	35	48	65	128	281	498	684	633	242	-
1965	3,802	957	236	46	48	76	152	294	470	721	570	232	-
1964	3,989	1,119	232	57	72	76	157	286	479	709	584	218	-
1963	4,093	1,196	260	53	52	74	167	309	481	689	605	207	-
1962	4,063	1,302	268	69	60	63	122	301	473	689	521	194	1
1961	4,016	1,289	307	50	54	72	146	276	438	658	528	197	1
1960	4,087	1,457	316	48	43	82	133	270	414	627	501	196	-
FEMALE													
1967	4,015	453	127	39	54	67	119	240	415	829	1,077	595	-
1966	3,993	599	150	37	43	68	104	274	411	797	984	526	-
1965	4,097	752	165	36	48	74	138	241	441	771	971	460	-
1964	4,189	905	231	34	63	70	134	231	399	727	915	480	-
1963	4,153	897	246	43	54	66	116	255	392	720	898	465	1
1962	4,131	1,040	235	64	59	74	131	221	388	638	869	412	-
1961	3,804	1,007	236	47	60	62	106	231	342	615	799	298	1
1960	3,845	1,165	212	39	57	73	134	214	328	630	675	316	2

- Quantity zero

Note: Figures by race exclude data for residents of New Jersey for 1962 and 1963.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290A for 1960-67 and separate unpublished tables for the black population and Accidents.

77/09/02 TABLE 290A. DEATHS FOR 60 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1960-67

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	TOTAL	UNDER 1 YEAR	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS	65-74 YEARS	75-84 YEARS	85+ YEARS	N.S.
Gastritis, duodenitis, enteritis, and colitis, except diarrhea of newborn (543,571,572)													
WHITE, BOTH SEXES													
1967	6,390	504	198	63	93	112	206	473	855	1,389	1,662	835	-
1966	6,218	597	230	53	83	112	205	490	827	1,368	1,529	724	-
1965	6,351	748	267	63	87	128	242	463	820	1,375	1,482	676	-
1964	6,451	934	291	71	119	123	244	446	786	1,328	1,435	674	-
1963	6,282	964	328	71	85	117	240	477	750	1,254	1,385	610	1
1962	6,104	1,106	342	99	101	112	209	430	711	1,164	1,269	561	-
1961	6,054	1,106	378	79	102	115	217	455	686	1,174	1,270	470	2
1960	5,995	1,270	353	70	88	134	223	423	667	1,159	1,127	480	1
WHITE, MALE													
1967	2,881	270	114	28	47	55	110	253	494	621	630	259	-
1966	2,884	334	135	27	46	57	116	255	458	632	599	225	-
1965	2,993	429	159	33	45	64	127	262	423	676	552	223	-
1964	3,062	506	144	45	67	65	131	246	439	659	550	210	-
1963	3,067	558	176	42	40	63	146	267	425	608	556	186	-
1962	2,978	616	185	49	53	56	99	256	403	607	476	178	-
1961	3,063	625	217	39	45	63	131	251	397	610	504	180	1
1960	3,054	728	210	37	37	73	113	242	376	577	478	183	-
WHITE, FEMALE													
1967	3,509	234	84	35	46	57	96	220	361	768	1,032	576	-
1966	3,334	263	95	26	37	55	89	235	369	736	930	499	-
1965	3,358	319	108	30	42	64	115	201	397	699	930	453	-
1964	3,389	428	147	26	52	58	113	200	347	669	885	464	-
1963	3,215	406	152	29	45	54	94	210	325	646	829	424	1
1962	3,126	490	157	50	48	56	110	174	308	557	793	383	-
1961	2,991	481	161	40	57	52	86	204	289	564	766	290	1
1960	2,941	542	143	33	51	61	110	181	291	582	649	297	1

- Quantity zero

Note: Figures by race exclude data for residents of New Jersey for 1962 and 1963.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290A for 1960-67 and separate unpublished tables for the black population and Accidents.

77/09/02 TABLE 290A. DEATHS FOR 60 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1960-67

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	TOTAL	UNDER 1 YEAR	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS	65-74 YEARS	75-84 YEARS	85+ YEARS	N.S.
Gastritis, duodenitis, enteritis, and colitis, except diarrhea of newborn (543,571,572)													
ALL OTHER, BOTH SEXES													
1967	1,114	527	97	18	16	21	50	53	97	110	84	41	-
1966	1,334	755	112	19	8	21	27	65	82	113	88	44	-
1965	1,548	961	134	19	9	22	48	72	91	117	59	16	-
1964	1,727	1,090	172	20	16	23	47	71	92	108	64	24	-
1963	1,702	1,088	168	23	18	20	33	64	81	104	64	39	-
1962	1,789	1,185	144	31	15	20	32	72	103	102	63	21	1
1961	1,766	1,190	165	18	12	19	35	52	94	99	57	25	-
1960	1,937	1,352	175	17	12	21	44	61	75	98	49	32	1
ALL OTHER, MALE													
1967	608	308	54	14	8	11	27	33	43	49	39	22	-
1966	675	419	57	8	2	8	12	26	40	52	34	17	-
1965	809	528	77	13	3	12	25	32	47	45	18	9	-
1964	927	613	88	12	5	11	26	40	40	50	34	8	-
1963	906	615	79	10	11	9	20	28	37	57	26	14	-
1962	935	652	74	18	6	6	15	38	42	51	23	9	1
1961	953	664	90	11	9	9	15	25	41	48	24	17	-
1960	1,033	729	106	11	6	9	20	28	38	50	23	13	-
ALL OTHER, FEMALE													
1967	506	219	43	4	8	10	23	20	54	61	45	19	-
1966	659	336	55	11	6	13	15	39	42	61	54	27	-
1965	739	433	57	6	6	10	23	40	44	72	41	7	-
1964	800	477	84	8	11	12	21	31	52	58	30	16	-
1963	796	473	89	13	7	11	13	36	44	47	38	25	-
1962	854	533	70	13	9	14	17	34	61	51	40	12	-
1961	813	526	75	7	3	10	20	27	53	51	33	8	-
1960	904	623	69	6	6	12	24	33	37	48	26	19	1

- Quantity zero

Note: Figures by race exclude data for residents of New Jersey for 1962 and 1963.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290A for 1960-67 and separate unpublished tables for the black population and Accidents.

77/09/02 TABLE 290A. DEATHS FOR 60 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1960-67

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	TOTAL	UNDER 1 YEAR	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS	65-74 YEARS	75-84 YEARS	85+ YEARS	N.S.
ICD-7 Nos. 581 Chronic liver disease and cirrhosis													
BOTH SEXES													
1967	27,816	35	18	45	118	918	4,096	7,858	8,026	4,795	1,630	268	9
1966	26,692	25	27	50	115	850	4,005	7,539	7,469	4,609	1,717	273	13
1965	24,715	29	28	37	105	778	3,649	7,015	6,800	4,364	1,674	233	3
1964	23,164	30	21	41	94	735	3,401	6,418	6,367	4,191	1,595	263	8
1963	22,456	37	22	38	79	725	3,107	6,409	5,971	4,255	1,546	262	5
1962	21,824	25	27	35	92	705	3,181	6,119	5,702	4,183	1,490	258	7
1961	20,737	29	24	43	86	699	2,987	5,744	5,419	3,983	1,467	250	6
1960	20,296	34	25	50	62	661	2,849	5,644	5,094	4,117	1,488	268	4
MALE													
1967	17,930	22	6	18	47	525	2,561	4,944	5,427	3,245	987	140	8
1966	17,378	9	16	20	54	493	2,447	4,782	5,203	3,139	1,050	155	10
1965	15,984	16	13	12	41	429	2,178	4,432	4,706	2,955	1,073	126	3
1964	15,004	14	8	16	35	395	1,994	4,116	4,370	2,916	1,005	131	4
1963	14,511	20	9	14	32	392	1,828	4,010	4,135	2,979	958	133	1
1962	14,329	14	15	17	41	367	1,871	3,964	4,022	2,926	941	144	7
1961	13,639	17	14	23	37	393	1,750	3,704	3,785	2,841	948	122	5
1960	13,483	16	16	20	25	347	1,698	3,711	3,654	2,935	913	145	3
FEMALE													
1967	9,886	13	12	27	71	393	1,535	2,914	2,599	1,550	643	128	1
1966	9,314	16	11	30	61	357	1,558	2,757	2,266	1,470	667	118	3
1965	8,731	13	15	25	64	349	1,471	2,583	2,094	1,409	601	107	-
1964	8,160	16	13	25	59	340	1,407	2,302	1,997	1,275	590	132	4
1963	7,945	17	13	24	47	333	1,279	2,399	1,836	1,276	588	129	4
1962	7,495	11	12	18	51	338	1,310	2,155	1,680	1,257	549	114	-
1961	7,098	12	10	20	49	306	1,237	2,040	1,634	1,142	519	128	1
1960	6,813	18	9	30	37	314	1,151	1,933	1,440	1,182	575	123	1

- Quantity zero

Note: Figures by race exclude data for residents of New Jersey for 1962 and 1963.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290A for 1960-67 and separate unpublished tables for the black population and Accidents.

77/09/02 TABLE 290A. DEATHS FOR 60 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1960-67

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	TOTAL	UNDER 1 YEAR	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS	65-74 YEARS	75-84 YEARS	85+ YEARS	N.S.
ICD-7 Nos. 581 Chronic liver disease and cirrhosis													
WHITE, BOTH SEXES													
1967	23,859	24	13	38	56	472	2,959	6,741	7,283	4,452	1,560	256	5
1966	22,837	21	21	38	64	416	2,901	6,416	6,755	4,294	1,643	257	11
1965	21,340	22	23	30	61	414	2,640	6,055	6,156	4,112	1,604	220	3
1964	20,344	22	16	31	59	416	2,565	5,631	5,849	3,971	1,521	256	7
1963	18,951	24	18	24	58	391	2,296	5,429	5,192	3,844	1,426	246	3
1962	18,359	15	20	29	53	395	2,360	5,186	4,951	3,733	1,371	240	6
1961	18,458	18	18	34	58	410	2,369	5,114	4,995	3,778	1,417	241	6
1960	18,212	26	21	45	44	417	2,321	5,024	4,688	3,937	1,431	255	3
WHITE, MALE													
1967	15,493	15	3	17	13	293	1,874	4,264	4,950	2,999	933	128	4
1966	15,024	8	13	17	25	240	1,814	4,101	4,741	2,911	1,002	144	8
1965	13,931	12	11	10	24	239	1,609	3,822	4,276	2,782	1,025	118	3
1964	13,290	11	8	12	22	244	1,529	3,611	4,006	2,762	953	128	4
1963	12,406	13	6	10	26	237	1,356	3,443	3,611	2,701	875	127	1
1962	12,242	8	12	16	25	230	1,400	3,423	3,510	2,616	864	132	6
1961	12,277	10	8	18	25	245	1,420	3,321	3,497	2,695	914	119	5
1960	12,231	12	15	17	18	232	1,415	3,328	3,369	2,807	878	138	2
WHITE, FEMALE													
1967	8,366	9	10	21	43	179	1,085	2,477	2,333	1,453	627	128	1
1966	7,813	13	8	21	39	176	1,087	2,315	2,014	1,383	641	113	3
1965	7,409	10	12	20	37	175	1,031	2,233	1,880	1,330	579	102	-
1964	7,054	11	8	19	37	172	1,036	2,020	1,843	1,209	568	128	3
1963	6,545	11	12	14	32	154	940	1,986	1,581	1,143	551	119	2
1962	6,117	7	8	13	28	165	960	1,763	1,441	1,117	507	108	-
1961	6,181	8	10	16	33	165	949	1,793	1,498	1,083	503	122	1
1960	5,981	14	6	28	26	185	906	1,696	1,319	1,130	553	117	1

- Quantity zero

Note: Figures by race exclude data for residents of New Jersey for 1962 and 1963.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290A for 1960-67 and separate unpublished tables for the black population and Accidents.

77/09/02 TABLE 290A. DEATHS FOR 60 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1960-67

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	TOTAL	UNDER 1 YEAR	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS	65-74 YEARS	75-84 YEARS	85+ YEARS	N.S.
ICD-7 Nos. 581 Chronic liver disease and cirrhosis													
ALL OTHER, BOTH SEXES													
1967	3,957	11	5	7	62	446	1,137	1,117	743	343	70	12	4
1966	3,855	4	6	12	51	434	1,104	1,123	714	315	74	16	2
1965	3,375	7	5	7	44	364	1,009	960	644	252	70	13	-
1964	2,820	8	5	10	35	319	836	787	518	220	74	7	1
1963	2,477	12	4	12	18	302	665	677	494	223	58	11	1
1962	2,424	9	7	4	38	280	658	657	467	229	62	12	1
1961	2,279	11	6	9	28	289	618	630	424	205	50	9	-
1960	2,084	8	4	5	18	244	528	620	406	180	57	13	1
ALL OTHER, MALE													
1967	2,437	7	3	1	34	232	687	680	477	246	54	12	4
1966	2,354	1	3	3	29	253	633	681	462	228	48	11	2
1965	2,053	4	2	2	17	190	569	610	430	173	48	8	-
1964	1,714	3	-	4	13	151	465	505	364	154	52	3	-
1963	1,450	6	3	4	6	134	380	387	333	151	41	5	-
1962	1,414	5	3	1	16	123	373	376	315	151	41	9	1
1961	1,362	7	6	5	12	148	330	383	288	146	34	3	-
1960	1,252	4	1	3	7	115	283	383	285	128	35	7	1
ALL OTHER, FEMALE													
1967	1,520	4	2	6	28	214	450	437	266	97	16	-	-
1966	1,501	3	3	9	22	181	471	442	252	87	26	5	-
1965	1,322	3	3	5	27	174	440	350	214	79	22	5	-
1964	1,106	5	5	6	22	168	371	282	154	66	22	4	1
1963	1,027	6	1	8	12	168	285	290	161	72	17	6	1
1962	1,010	4	4	3	22	157	285	281	152	78	21	3	-
1961	917	4	-	4	16	141	288	247	136	59	16	6	-
1960	832	4	3	2	11	129	245	237	121	52	22	6	-

- Quantity zero

Note: Figures by race exclude data for residents of New Jersey for 1962 and 1963.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290A for 1960-67 and separate unpublished tables for the black population and Accidents.

77/09/02 TABLE 290A. DEATHS FOR 60 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1960-67

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	TOTAL	UNDER 1 YEAR	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS	65-74 YEARS	75-84 YEARS	85+ YEARS	N.S.
Cholelithiasis, cholecystitis, and cholangitis (584,585)													
BOTH SEXES													
1967	4,383	1	1	1	21	48	76	248	552	1,231	1,532	672	-
1966	4,592	-	-	-	12	29	113	280	601	1,312	1,560	684	1
1965	4,666	3	-	-	17	38	114	274	586	1,342	1,639	652	1
1964	4,617	2	-	-	10	51	114	291	657	1,333	1,485	673	1
1963	4,676	1	-	-	20	44	130	264	676	1,380	1,528	632	1
1962	4,831	5	-	2	17	49	122	310	660	1,457	1,585	624	-
1961	4,707	1	1	2	16	45	147	308	689	1,440	1,424	634	-
1960	4,619	1	1	-	17	50	148	324	655	1,425	1,449	548	1
MALE													
1967	1,967	1	1	1	4	22	28	105	264	604	690	247	-
1966	1,991	-	-	-	1	11	47	126	292	608	673	232	1
1965	1,960	1	-	-	1	12	38	115	262	605	707	219	-
1964	1,915	1	-	-	2	14	42	120	291	602	612	231	-
1963	1,904	-	-	-	2	10	44	120	292	608	612	216	-
1962	2,014	3	-	2	4	15	51	140	298	664	631	206	-
1961	1,957	-	1	1	2	6	54	122	297	661	582	231	-
1960	1,858	1	-	-	3	10	49	138	267	608	605	177	-
FEMALE													
1967	2,416	-	-	-	17	26	48	143	288	627	842	425	-
1966	2,601	-	-	-	11	18	66	154	309	704	887	452	-
1965	2,706	2	-	-	16	26	76	159	324	737	932	433	1
1964	2,702	1	-	-	8	37	72	171	366	731	873	442	1
1963	2,772	1	-	-	18	34	86	144	384	772	916	416	1
1962	2,817	2	-	-	13	34	71	170	362	793	954	418	-
1961	2,750	1	-	1	14	39	93	186	392	779	842	403	-
1960	2,761	-	1	-	14	40	99	186	388	817	844	371	1

- Quantity zero

Note: Figures by race exclude data for residents of New Jersey for 1962 and 1963.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290A for 1960-67 and separate unpublished tables for the black population and Accidents.

77/09/02 TABLE 290A. DEATHS FOR 60 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1960-67

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	TOTAL	UNDER 1 YEAR	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS	65-74 YEARS	75-84 YEARS	85+ YEARS	N.S.
Cholelithiasis, cholecystitis, and cholangitis (584,585)													
WHITE, BOTH SEXES													
1967	4,104	1	1	1	17	38	65	212	503	1,149	1,471	646	-
1966	4,318	-	-	-	10	24	103	246	548	1,213	1,507	666	1
1965	4,416	2	-	-	13	35	94	247	534	1,269	1,583	639	-
1964	4,386	1	-	-	10	43	92	257	609	1,267	1,453	653	1
1963	4,246	1	-	-	17	32	100	223	599	1,237	1,429	607	1
1962	4,381	4	-	1	14	37	93	262	578	1,332	1,477	583	-
1961	4,450	-	1	-	16	39	126	280	629	1,370	1,372	617	-
1960	4,375	1	-	-	14	42	128	288	608	1,356	1,405	532	1
WHITE, MALE													
1967	1,878	1	1	1	4	19	27	98	250	577	664	236	-
1966	1,882	-	-	-	1	10	44	114	265	571	650	226	1
1965	1,883	1	-	-	1	11	36	110	242	579	689	214	-
1964	1,834	1	-	-	2	12	36	109	274	572	602	226	-
1963	1,767	-	-	-	1	8	37	103	276	557	578	207	-
1962	1,843	2	-	1	3	11	39	125	267	618	586	191	-
1961	1,861	-	1	-	2	6	46	111	275	631	564	225	-
1960	1,774	1	-	-	2	9	42	126	250	588	586	170	-
WHITE, FEMALE													
1967	2,226	-	-	-	13	19	38	114	253	572	807	410	-
1966	2,436	-	-	-	9	14	59	132	283	642	857	440	-
1965	2,533	1	-	-	12	24	58	137	292	690	894	425	-
1964	2,552	-	-	-	8	31	56	148	335	695	851	427	1
1963	2,479	1	-	-	16	24	63	120	323	680	851	400	1
1962	2,538	2	-	-	11	26	54	137	311	714	891	392	-
1961	2,589	-	-	-	14	33	80	169	354	739	808	392	-
1960	2,601	-	-	-	12	33	86	162	358	768	819	362	1

- Quantity zero

Note: Figures by race exclude data for residents of New Jersey for 1962 and 1963.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290A for 1960-67 and separate unpublished tables for the black population and Accidents.

77/09/02 TABLE 290A. DEATHS FOR 60 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1960-67

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	TOTAL	UNDER 1 YEAR	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS	65-74 YEARS	75-84 YEARS	85+ YEARS	N.S.
Cholelithiasis, cholecystitis, and cholangitis (584,585)													
ALL OTHER, BOTH SEXES													
1967	279	-	-	-	4	10	11	36	49	82	61	26	-
1966	274	-	-	-	2	5	10	34	53	99	53	18	-
1965	250	1	-	-	4	3	20	27	52	73	56	13	1
1964	231	1	-	-	-	8	22	34	48	66	32	20	-
1963	258	-	-	-	3	11	23	26	52	86	43	14	-
1962	285	1	-	1	3	7	22	35	61	72	62	21	-
1961	257	1	-	2	-	6	21	28	60	70	52	17	-
1960	244	-	1	-	3	8	20	36	47	69	44	16	-
ALL OTHER, MALE													
1967	89	-	-	-	-	3	1	7	14	27	26	11	-
1966	109	-	-	-	-	1	3	12	27	37	23	6	-
1965	77	-	-	-	-	1	2	5	20	26	18	5	-
1964	81	-	-	-	-	2	6	11	17	30	10	5	-
1963	82	-	-	-	1	2	6	11	10	28	18	6	-
1962	100	1	-	1	1	3	8	10	20	25	26	5	-
1961	96	-	-	1	-	-	8	11	22	30	18	6	-
1960	84	-	-	-	1	1	7	12	17	20	19	7	-
ALL OTHER, FEMALE													
1967	190	-	-	-	4	7	10	29	35	55	35	15	-
1966	165	-	-	-	2	4	7	22	26	62	30	12	-
1965	173	1	-	-	4	2	18	22	32	47	38	8	1
1964	150	1	-	-	-	6	16	23	31	36	22	15	-
1963	176	-	-	-	2	9	17	15	42	58	25	8	-
1962	185	-	-	-	2	4	14	25	41	47	36	16	-
1961	161	1	-	1	-	6	13	17	38	40	34	11	-
1960	160	-	1	-	2	7	13	24	30	49	25	9	-

- Quantity zero

Note: Figures by race exclude data for residents of New Jersey for 1962 and 1963.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290A for 1960-67 and separate unpublished tables for the black population and Accidents.

77/09/02 TABLE 290A. DEATHS FOR 60 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1960-67

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	TOTAL	UNDER 1 YEAR	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS	65-74 YEARS	75-84 YEARS	85+ YEARS	N.S.
Acute nephritis, and nephritis with edema including nephrosis (590,591)													
BOTH SEXES													
1967	1,005	19	36	62	66	50	84	132	164	174	136	81	1
1966	1,164	21	45	52	72	65	113	126	174	213	201	82	-
1965	1,325	21	66	77	89	75	104	177	196	243	205	72	-
1964	1,335	31	53	76	89	86	105	165	206	231	199	94	-
1963	1,434	17	61	100	89	81	138	165	230	267	204	82	-
1962	1,573	42	79	112	99	94	135	179	230	267	241	95	-
1961	1,584	34	86	114	82	81	133	186	232	288	241	107	-
1960	1,651	44	80	116	93	102	131	210	265	303	215	91	1
MALE													
1967	564	12	18	36	39	32	41	72	97	98	81	38	-
1966	656	12	30	27	45	34	57	74	105	122	108	42	-
1965	730	12	37	43	54	40	47	105	117	129	108	38	-
1964	734	18	27	46	52	45	54	94	118	129	113	38	-
1963	805	12	33	51	40	46	75	103	142	150	108	45	-
1962	901	20	49	61	60	50	76	104	134	162	132	53	-
1961	894	20	50	53	53	48	75	118	126	178	133	40	-
1960	921	29	36	67	56	63	73	117	147	169	119	44	1
FEMALE													
1967	441	7	18	26	27	18	43	60	67	76	55	43	1
1966	508	9	15	25	27	31	56	52	69	91	93	40	-
1965	595	9	29	34	35	35	57	72	79	114	97	34	-
1964	601	13	26	30	37	41	51	71	88	102	86	56	-
1963	629	5	28	49	49	35	63	62	88	117	96	37	-
1962	672	22	30	51	39	44	59	75	96	105	109	42	-
1961	690	14	36	61	29	33	58	68	106	110	108	67	-
1960	730	15	44	49	37	39	58	93	118	134	96	47	-

- Quantity zero

Note: Figures by race exclude data for residents of New Jersey for 1962 and 1963.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290A for 1960-67 and separate unpublished tables for the black population and Accidents.

77/09/02 TABLE 290A. DEATHS FOR 60 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1960-67

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	TOTAL	UNDER 1 YEAR	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS	65-74 YEARS	75-84 YEARS	85+ YEARS	N.S.
Acute nephritis, and nephritis with edema including nephrosis (590,591)													
WHITE, BOTH SEXES													
1967	775	14	24	42	50	29	54	93	129	155	118	67	-
1966	907	15	36	37	55	40	64	95	140	173	181	71	-
1965	1,017	20	47	60	69	41	77	115	150	208	175	55	-
1964	1,053	24	35	52	71	49	78	125	172	191	180	76	-
1963	1,043	12	31	69	61	57	87	118	176	194	165	73	-
1962	1,201	30	53	86	78	65	85	120	175	223	203	83	-
1961	1,216	26	57	84	62	55	81	140	175	237	207	92	-
1960	1,280	33	58	89	71	71	87	147	204	253	184	83	-
WHITE, MALE													
1967	452	9	11	26	29	19	33	53	76	89	76	31	-
1966	514	6	24	20	33	18	36	59	85	99	98	36	-
1965	559	12	24	33	43	26	35	67	88	113	91	27	-
1964	591	13	18	32	43	26	39	78	99	108	103	32	-
1963	586	8	19	34	31	38	45	73	107	104	88	39	-
1962	697	14	32	44	51	36	55	71	100	134	113	47	-
1961	694	14	35	39	41	35	46	90	101	145	116	32	-
1960	707	19	25	47	46	46	46	80	116	138	103	41	-
WHITE, FEMALE													
1967	323	5	13	16	21	10	21	40	53	66	42	36	-
1966	393	9	12	17	22	22	28	36	55	74	83	35	-
1965	458	8	23	27	26	15	42	48	62	95	84	28	-
1964	462	11	17	20	28	23	39	47	73	83	77	44	-
1963	457	4	12	35	30	19	42	45	69	90	77	34	-
1962	504	16	21	42	27	29	30	49	75	89	90	36	-
1961	522	12	22	45	21	20	35	50	74	92	91	60	-
1960	573	14	33	42	25	25	41	67	88	115	81	42	-

- Quantity zero

Note: Figures by race exclude data for residents of New Jersey for 1962 and 1963.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290A for 1960-67 and separate unpublished tables for the black population and Accidents.

77/09/02 TABLE 290A. DEATHS FOR 60 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1960-67

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	TOTAL	UNDER 1 YEAR	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS	65-74 YEARS	75-84 YEARS	85+ YEARS	N.S.
Acute nephritis, and nephritis with edema including nephrosis (590,591)													
ALL OTHER, BOTH SEXES													
1967	230	5	12	20	16	21	30	39	35	19	18	14	1
1966	257	6	9	15	17	25	49	31	34	40	20	11	-
1965	308	1	19	17	20	34	27	62	46	35	30	17	-
1964	282	7	18	24	18	37	27	40	34	40	19	18	-
1963	345	5	29	29	27	23	46	38	49	62	28	9	-
1962	324	9	23	20	19	23	42	52	50	38	37	11	-
1961	368	8	29	30	20	26	52	46	57	51	34	15	-
1960	371	11	22	27	22	31	44	63	61	50	31	8	1
ALL OTHER, MALE													
1967	112	3	7	10	10	13	8	19	21	9	5	7	-
1966	142	6	6	7	12	16	21	15	20	23	10	6	-
1965	171	-	13	10	11	14	12	38	29	16	17	11	-
1964	143	5	9	14	9	19	15	16	19	21	10	6	-
1963	190	4	13	17	9	7	26	24	30	38	16	6	-
1962	182	3	15	14	8	13	18	31	32	25	18	5	-
1961	200	6	15	14	12	13	29	28	25	33	17	8	-
1960	214	10	11	20	10	17	27	37	31	31	16	3	1
ALL OTHER, FEMALE													
1967	118	2	5	10	6	8	22	20	14	10	13	7	1
1966	115	-	3	8	5	9	28	16	14	17	10	5	-
1965	137	1	6	7	9	20	15	24	17	19	13	6	-
1964	139	2	9	10	9	18	12	24	15	19	9	12	-
1963	155	1	16	12	18	16	20	14	19	24	12	3	-
1962	142	6	8	6	11	10	24	21	18	13	19	6	-
1961	168	2	14	16	8	13	23	18	32	18	17	7	-
1960	157	1	11	7	12	14	17	26	30	19	15	5	-

- Quantity zero

Note: Figures by race exclude data for residents of New Jersey for 1962 and 1963.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290A for 1960-67 and separate unpublished tables for the black population and Accidents.

77/09/02 TABLE 290A. DEATHS FOR 60 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1960-67

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	TOTAL	UNDER 1 YEAR	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS	65-74 YEARS	75-84 YEARS	85+ YEARS	N.S.
Infections of kidney (600)													
BOTH SEXES													
1967	9,006	14	14	34	62	138	334	709	1,285	2,278	2,748	1,388	2
1966	9,498	22	15	39	91	137	348	789	1,361	2,472	2,842	1,381	1
1965	9,813	40	15	34	106	160	450	862	1,496	2,505	2,754	1,386	5
1964	9,889	48	20	46	97	204	446	920	1,463	2,526	2,812	1,301	6
1963	8,952	36	28	46	87	207	436	838	1,449	2,327	2,403	1,090	5
1962	8,744	41	20	40	83	173	448	798	1,446	2,281	2,317	1,093	4
1961	8,095	41	15	38	81	193	465	795	1,336	2,138	2,108	883	2
1960	7,645	70	14	45	69	197	485	838	1,306	1,916	1,860	843	2
MALE													
1967	4,179	11	7	7	30	55	111	271	578	1,090	1,369	649	1
1966	4,520	13	6	13	49	43	125	327	603	1,272	1,439	630	-
1965	4,652	26	6	12	41	55	158	345	709	1,275	1,426	598	1
1964	4,780	32	12	21	47	73	155	368	694	1,320	1,453	602	3
1963	4,325	28	14	16	35	77	167	338	668	1,233	1,223	524	2
1962	4,306	30	6	22	40	55	166	326	687	1,205	1,214	555	-
1961	4,082	23	5	12	31	65	180	350	663	1,184	1,142	427	-
1960	3,861	45	5	12	37	73	190	383	641	1,082	1,001	391	1
FEMALE													
1967	4,827	3	7	27	32	83	223	438	707	1,188	1,379	739	1
1966	4,978	9	9	26	42	94	223	462	758	1,200	1,403	751	1
1965	5,161	14	9	22	65	105	292	517	787	1,230	1,328	788	4
1964	5,109	16	8	25	50	131	291	552	769	1,206	1,359	699	3
1963	4,627	8	14	30	52	130	269	500	781	1,094	1,180	566	3
1962	4,438	11	14	18	43	118	282	472	759	1,076	1,103	538	4
1961	4,013	18	10	26	50	128	285	445	673	954	966	456	2
1960	3,784	25	9	33	32	124	295	455	665	834	859	452	1

- Quantity zero

Note: Figures by race exclude data for residents of New Jersey for 1962 and 1963.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290A for 1960-67 and separate unpublished tables for the black population and Accidents.

77/09/02 TABLE 290A. DEATHS FOR 60 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1960-67

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	TOTAL	UNDER 1 YEAR	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS	65-74 YEARS	75-84 YEARS	85+ YEARS	N.S.
Infections of kidney (600)													
WHITE, BOTH SEXES													
1967	7,376	7	14	30	53	81	185	482	904	1,808	2,518	1,293	1
1966	7,744	15	11	31	75	79	210	512	955	2,000	2,572	1,284	-
1965	7,979	19	13	25	82	98	270	555	1,045	2,068	2,514	1,287	3
1964	8,093	33	15	37	77	132	276	601	1,065	2,105	2,557	1,194	1
1963	7,000	25	22	37	68	136	248	545	1,015	1,830	2,086	984	4
1962	6,858	23	15	38	63	116	277	510	1,009	1,795	2,010	998	4
1961	6,578	28	12	31	62	144	293	509	961	1,786	1,928	822	2
1960	6,117	50	9	36	47	126	309	543	922	1,584	1,695	795	1
WHITE, MALE													
1967	3,489	6	7	7	27	37	72	203	417	879	1,238	595	1
1966	3,657	9	3	9	39	28	79	207	415	1,007	1,289	572	-
1965	3,775	10	5	8	34	32	109	227	509	1,018	1,279	543	1
1964	3,906	23	8	16	42	53	93	242	514	1,080	1,293	542	-
1963	3,361	18	11	13	28	53	97	228	458	941	1,048	465	1
1962	3,368	15	5	22	30	44	100	207	479	936	1,023	507	-
1961	3,331	14	3	12	24	51	127	222	469	980	1,036	393	-
1960	3,117	31	-	9	27	48	128	261	461	891	901	360	-
WHITE, FEMALE													
1967	3,887	1	7	23	26	44	113	279	487	929	1,280	698	-
1966	4,087	6	8	22	36	51	131	305	540	993	1,283	712	-
1965	4,204	9	8	17	48	66	161	328	536	1,050	1,235	744	2
1964	4,187	10	7	21	35	79	183	359	551	1,025	1,264	652	1
1963	3,639	7	11	24	40	83	151	317	557	889	1,038	519	3
1962	3,490	8	10	16	33	72	177	303	530	859	987	491	4
1961	3,247	14	9	19	38	93	166	287	492	806	892	429	2
1960	3,000	19	9	27	20	78	181	282	461	693	794	435	1

- Quantity zero

Note: Figures by race exclude data for residents of New Jersey for 1962 and 1963.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290A for 1960-67 and separate unpublished tables for the black population and Accidents.

77/09/02 TABLE 290A. DEATHS FOR 60 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1960-67

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	TOTAL	UNDER 1 YEAR	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS	65-74 YEARS	75-84 YEARS	85+ YEARS	N.S.
Infections of kidney (600)													
ALL OTHER, BOTH SEXES													
1967	1,630	7	-	4	9	57	149	227	381	470	230	95	1
1966	1,754	7	4	8	16	58	138	277	406	472	270	97	1
1965	1,834	21	2	9	24	62	180	307	451	437	240	99	2
1964	1,796	15	5	9	20	72	170	319	398	421	255	107	5
1963	1,677	10	5	7	19	62	173	265	392	421	236	86	1
1962	1,615	17	5	1	18	50	157	265	386	406	235	75	-
1961	1,517	13	3	7	19	49	172	286	375	352	180	61	-
1960	1,528	20	5	9	22	71	176	295	384	332	165	48	1
ALL OTHER, MALE													
1967	690	5	-	-	3	18	39	68	161	211	131	54	-
1966	863	4	3	4	10	15	46	120	188	265	150	58	-
1965	877	16	1	4	7	23	49	118	200	257	147	55	-
1964	874	9	4	5	5	20	62	126	180	240	160	60	3
1963	837	9	2	3	7	18	65	101	192	251	140	48	1
1962	824	14	1	-	10	10	62	108	188	236	157	38	-
1961	751	9	2	-	7	14	53	128	194	204	106	34	-
1960	744	14	5	3	10	25	62	122	180	191	100	31	1
ALL OTHER, FEMALE													
1967	940	2	-	4	6	39	110	159	220	259	99	41	1
1966	891	3	1	4	6	43	92	157	218	207	120	39	1
1965	957	5	1	5	17	39	131	189	251	180	93	44	2
1964	922	6	1	4	15	52	108	193	218	181	95	47	2
1963	840	1	3	4	12	44	108	164	200	170	96	38	-
1962	791	3	4	1	8	40	95	157	198	170	78	37	-
1961	766	4	1	7	12	35	119	158	181	148	74	27	-
1960	784	6	-	6	12	46	114	173	204	141	65	17	-

- Quantity zero

Note: Figures by race exclude data for residents of New Jersey for 1962 and 1963.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290A for 1960-67 and separate unpublished tables for the black population and Accidents.

77/09/02 TABLE 290A. DEATHS FOR 60 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1960-67

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	TOTAL	UNDER 1 YEAR	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS	65-74 YEARS	75-84 YEARS	85+ YEARS	N.S.
Hyperplasia of prostate (610)													
BOTH SEXES													
1967	3,136	-	-	-	-	-	3	26	178	796	1,332	801	-
1966	3,217	-	-	-	-	-	1	20	199	857	1,349	791	-
1965	3,413	-	-	-	-	3	2	34	253	896	1,478	747	-
1964	3,847	-	-	-	1	1	4	29	294	993	1,634	889	2
1963	4,248	-	-	-	-	-	3	34	309	1,135	1,836	929	2
1962	4,283	-	-	-	-	1	2	40	336	1,160	1,804	939	1
1961	4,158	-	-	-	-	-	7	42	304	1,120	1,757	927	1
1960	4,449	-	-	-	-	1	8	50	317	1,244	1,868	957	4
MALE													
1967	3,136	-	-	-	-	-	3	26	178	796	1,332	801	-
1966	3,217	-	-	-	-	-	1	20	199	857	1,349	791	-
1965	3,413	-	-	-	-	3	2	34	253	896	1,478	747	-
1964	3,847	-	-	-	1	1	4	29	294	993	1,634	889	2
1963	4,248	-	-	-	-	-	3	34	309	1,135	1,836	929	2
1962	4,283	-	-	-	-	1	2	40	336	1,160	1,804	939	1
1961	4,158	-	-	-	-	-	7	42	304	1,120	1,757	927	1
1960	4,449	-	-	-	-	1	8	50	317	1,244	1,868	957	4
FEMALE													
1967
1966
1965
1964
1963
1962
1961
1960

- Quantity zero

... Not applicable.

Note: Figures by race exclude data for residents of New Jersey for 1962 and 1963.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290A for 1960-67 and separate unpublished tables for the black population and Accidents.

77/09/02 TABLE 290A. DEATHS FOR 60 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1960-67

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	TOTAL	UNDER 1 YEAR	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS	65-74 YEARS	75-84 YEARS	85+ YEARS	N.S.
Hyperplasia of prostate (610)													
WHITE, BOTH SEXES													
1967	2,736	-	-	-	-	-	2	17	124	642	1,200	751	-
1966	2,800	-	-	-	-	-	-	11	136	696	1,221	736	-
1965	2,953	-	-	-	-	3	2	23	174	729	1,334	688	-
1964	3,317	-	-	-	1	1	3	16	210	818	1,459	809	-
1963	3,523	-	-	-	-	-	-	21	202	870	1,588	840	2
1962	3,519	-	-	-	-	1	1	16	209	891	1,571	829	1
1961	3,608	-	-	-	-	-	2	21	224	912	1,586	863	-
1960	3,864	-	-	-	-	1	5	27	226	1,027	1,701	876	1
WHITE, MALE													
1967	2,736	-	-	-	-	-	2	17	124	642	1,200	751	-
1966	2,800	-	-	-	-	-	-	11	136	696	1,221	736	-
1965	2,953	-	-	-	-	3	2	23	174	729	1,334	688	-
1964	3,317	-	-	-	1	1	3	16	210	818	1,459	809	-
1963	3,523	-	-	-	-	-	-	21	202	870	1,588	840	2
1962	3,519	-	-	-	-	1	1	16	209	891	1,571	829	1
1961	3,608	-	-	-	-	-	2	21	224	912	1,586	863	-
1960	3,864	-	-	-	-	1	5	27	226	1,027	1,701	876	1
WHITE, FEMALE													
1967
1966
1965
1964
1963
1962
1961
1960

- Quantity zero

... Not applicable.

Note: Figures by race exclude data for residents of New Jersey for 1962 and 1963.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290A for 1960-67 and separate unpublished tables for the black population and Accidents.

77/09/02 TABLE 290A. DEATHS FOR 60 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1960-67

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	TOTAL	UNDER 1 YEAR	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS	65-74 YEARS	75-84 YEARS	85+ YEARS	N.S.
Hyperplasia of prostate (610)													
ALL OTHER, BOTH SEXES													
1967	400	-	-	-	-	-	1	9	54	154	132	50	-
1966	417	-	-	-	-	-	1	9	63	161	128	55	-
1965	460	-	-	-	-	-	-	11	79	167	144	59	-
1964	530	-	-	-	-	-	1	13	84	175	175	80	2
1963	594	-	-	-	-	-	3	12	95	218	198	68	-
1962	622	-	-	-	-	-	1	24	114	223	181	79	-
1961	550	-	-	-	-	-	5	21	80	208	171	64	1
1960	585	-	-	-	-	-	3	23	91	217	167	81	3
ALL OTHER, MALE													
1967	400	-	-	-	-	-	1	9	54	154	132	50	-
1966	417	-	-	-	-	-	1	9	63	161	128	55	-
1965	460	-	-	-	-	-	-	11	79	167	144	59	-
1964	530	-	-	-	-	-	1	13	84	175	175	80	2
1963	594	-	-	-	-	-	3	12	95	218	198	68	-
1962	622	-	-	-	-	-	1	24	114	223	181	79	-
1961	550	-	-	-	-	-	5	21	80	208	171	64	1
1960	585	-	-	-	-	-	3	23	91	217	167	81	3
ALL OTHER, FEMALE													
1967
1966
1965
1964
1963
1962
1961
1960

- Quantity zero

... Not applicable.

Note: Figures by race exclude data for residents of New Jersey for 1962 and 1963.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290A for 1960-67 and separate unpublished tables for the black population and Accidents.

77/09/02 TABLE 290A. DEATHS FOR 60 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1960-67

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	TOTAL	UNDER 1 YEAR	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS	65-74 YEARS	75-84 YEARS	85+ YEARS	N.S.
Deliveries and complications of pregnancy, childbirth, and the puerperium (640-689)													
BOTH SEXES													
1967	987	-	-	4	347	370	253	13	-	-	-	-	-
1966	1,049	-	-	7	333	423	274	11	-	-	-	-	1
1965	1,189	-	-	3	350	493	336	7	-	-	-	-	-
1964	1,343	-	-	3	369	554	399	18	-	-	-	-	-
1963	1,466	-	-	7	405	610	434	10	-	-	-	-	-
1962	1,465	-	-	3	381	681	393	7	-	-	-	-	-
1961	1,573	-	-	12	385	709	451	16	-	-	-	-	-
1960	1,579	-	-	7	424	672	461	15	-	-	-	-	-
MALE													
1967
1966
1965
1964
1963
1962
1961
1960
FEMALE													
1967	987	-	-	4	347	370	253	13	-	-	-	-	-
1966	1,049	-	-	7	333	423	274	11	-	-	-	-	1
1965	1,189	-	-	3	350	493	336	7	-	-	-	-	-
1964	1,343	-	-	3	369	554	399	18	-	-	-	-	-
1963	1,466	-	-	7	405	610	434	10	-	-	-	-	-
1962	1,465	-	-	3	381	681	393	7	-	-	-	-	-
1961	1,573	-	-	12	385	709	451	16	-	-	-	-	-
1960	1,579	-	-	7	424	672	461	15	-	-	-	-	-

- Quantity zero

... Not applicable.

Note: Figures by race exclude data for residents of New Jersey for 1962 and 1963.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290A for 1960-67 and separate unpublished tables for the black population and Accidents.

77/09/02 TABLE 290A. DEATHS FOR 60 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1960-67

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	TOTAL	UNDER 1 YEAR	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS	65-74 YEARS	75-84 YEARS	85+ YEARS	N.S.
Deliveries and complications of pregnancy, childbirth, and the puerperium (640-689)													
WHITE, BOTH SEXES													
1967	571	-	-	-	189	226	149	7	-	-	-	-	-
1966	605	-	-	1	187	252	157	8	-	-	-	-	-
1965	656	-	-	-	204	252	196	4	-	-	-	-	-
1964	751	-	-	-	198	309	230	14	-	-	-	-	-
1963	797	-	-	-	229	319	242	7	-	-	-	-	-
1962	807	-	-	-	203	382	217	5	-	-	-	-	-
1961	897	-	-	3	213	398	275	8	-	-	-	-	-
1960	936	-	-	2	253	393	281	7	-	-	-	-	-
WHITE, MALE													
1967
1966
1965
1964
1963
1962
1961
1960
WHITE, FEMALE													
1967	571	-	-	-	189	226	149	7	-	-	-	-	-
1966	605	-	-	1	187	252	157	8	-	-	-	-	-
1965	656	-	-	-	204	252	196	4	-	-	-	-	-
1964	751	-	-	-	198	309	230	14	-	-	-	-	-
1963	797	-	-	-	229	319	242	7	-	-	-	-	-
1962	807	-	-	-	203	382	217	5	-	-	-	-	-
1961	897	-	-	3	213	398	275	8	-	-	-	-	-
1960	936	-	-	2	253	393	281	7	-	-	-	-	-

- Quantity zero

... Not applicable.

Note: Figures by race exclude data for residents of New Jersey for 1962 and 1963.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290A for 1960-67 and separate unpublished tables for the black population and Accidents.

77/09/02 TABLE 290A. DEATHS FOR 60 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1960-67

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	TOTAL	UNDER 1 YEAR	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS	65-74 YEARS	75-84 YEARS	85+ YEARS	N.S.
Deliveries and complications of pregnancy, childbirth, and the puerperium (640-689)													
ALL OTHER, BOTH SEXES													
1967	416	-	-	4	158	144	104	6	-	-	-	-	-
1966	444	-	-	6	146	171	117	3	-	-	-	-	1
1965	533	-	-	3	146	241	140	3	-	-	-	-	-
1964	592	-	-	3	171	245	169	4	-	-	-	-	-
1963	619	-	-	7	166	261	182	3	-	-	-	-	-
1962	615	-	-	3	172	273	165	2	-	-	-	-	-
1961	676	-	-	9	172	311	176	8	-	-	-	-	-
1960	643	-	-	5	171	279	180	8	-	-	-	-	-
ALL OTHER, MALE													
1967
1966
1965
1964
1963
1962
1961
1960
ALL OTHER, FEMALE													
1967	416	-	-	4	158	144	104	6	-	-	-	-	-
1966	444	-	-	6	146	171	117	3	-	-	-	-	1
1965	533	-	-	3	146	241	140	3	-	-	-	-	-
1964	592	-	-	3	171	245	169	4	-	-	-	-	-
1963	619	-	-	7	166	261	182	3	-	-	-	-	-
1962	615	-	-	3	172	273	165	2	-	-	-	-	-
1961	676	-	-	9	172	311	176	8	-	-	-	-	-
1960	643	-	-	5	171	279	180	8	-	-	-	-	-

- Quantity zero

... Not applicable.

Note: Figures by race exclude data for residents of New Jersey for 1962 and 1963.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290A for 1960-67 and separate unpublished tables for the black population and Accidents.

77/09/02 TABLE 290A. DEATHS FOR 60 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1960-67

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	TOTAL	UNDER 1 YEAR	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS	65-74 YEARS	75-84 YEARS	85+ YEARS	N.S.
Abortion (650-652)													
BOTH SEXES													
1967	160	-	-	-	64	62	33	1	-	-	-	-	-
1966	189	-	-	2	69	79	39	-	-	-	-	-	-
1965	235	-	-	-	85	109	41	-	-	-	-	-	-
1964	247	-	-	1	90	98	55	3	-	-	-	-	-
1963	280	-	-	2	74	137	66	1	-	-	-	-	-
1962	305	-	-	1	93	153	57	1	-	-	-	-	-
1961	324	-	-	1	92	167	63	1	-	-	-	-	-
1960	289	-	-	2	79	160	45	3	-	-	-	-	-
MALE													
1967
1966
1965
1964
1963
1962
1961
1960
FEMALE													
1967	160	-	-	-	64	62	33	1	-	-	-	-	-
1966	189	-	-	2	69	79	39	-	-	-	-	-	-
1965	235	-	-	-	85	109	41	-	-	-	-	-	-
1964	247	-	-	1	90	98	55	3	-	-	-	-	-
1963	280	-	-	2	74	137	66	1	-	-	-	-	-
1962	305	-	-	1	93	153	57	1	-	-	-	-	-
1961	324	-	-	1	92	167	63	1	-	-	-	-	-
1960	289	-	-	2	79	160	45	3	-	-	-	-	-

- Quantity zero

... Not applicable.

Note: Figures by race exclude data for residents of New Jersey for 1962 and 1963.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290A for 1960-67 and separate unpublished tables for the black population and Accidents.

77/09/02 TABLE 290A. DEATHS FOR 60 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1960-67

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	TOTAL	UNDER 1 YEAR	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS	65-74 YEARS	75-84 YEARS	85+ YEARS	N.S.
Abortion (650-652)													
WHITE, BOTH SEXES													
1967	76	-	-	-	31	26	19	-	-	-	-	-	-
1966	96	-	-	-	34	41	21	-	-	-	-	-	-
1965	106	-	-	-	38	48	20	-	-	-	-	-	-
1964	117	-	-	-	41	44	30	2	-	-	-	-	-
1963	161	-	-	-	44	79	37	1	-	-	-	-	-
1962	149	-	-	-	38	75	35	1	-	-	-	-	-
1961	163	-	-	-	38	82	42	1	-	-	-	-	-
1960	147	-	-	1	34	83	27	2	-	-	-	-	-
WHITE, MALE													
1967
1966
1965
1964
1963
1962
1961
1960
WHITE, FEMALE													
1967	76	-	-	-	31	26	19	-	-	-	-	-	-
1966	96	-	-	-	34	41	21	-	-	-	-	-	-
1965	106	-	-	-	38	48	20	-	-	-	-	-	-
1964	117	-	-	-	41	44	30	2	-	-	-	-	-
1963	161	-	-	-	44	79	37	1	-	-	-	-	-
1962	149	-	-	-	38	75	35	1	-	-	-	-	-
1961	163	-	-	-	38	82	42	1	-	-	-	-	-
1960	147	-	-	1	34	83	27	2	-	-	-	-	-

- Quantity zero

... Not applicable.

Note: Figures by race exclude data for residents of New Jersey for 1962 and 1963.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290A for 1960-67 and separate unpublished tables for the black population and Accidents.

77/09/02 TABLE 290A. DEATHS FOR 60 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1960-67

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	TOTAL	UNDER 1 YEAR	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS	65-74 YEARS	75-84 YEARS	85+ YEARS	N.S.
Abortion (650-652)													
ALL OTHER, BOTH SEXES													
1967	84	-	-	-	33	36	14	1	-	-	-	-	-
1966	93	-	-	2	35	38	18	-	-	-	-	-	-
1965	129	-	-	-	47	61	21	-	-	-	-	-	-
1964	130	-	-	1	49	54	25	1	-	-	-	-	-
1963	107	-	-	2	28	50	27	-	-	-	-	-	-
1962	148	-	-	1	53	73	21	-	-	-	-	-	-
1961	161	-	-	1	54	85	21	-	-	-	-	-	-
1960	142	-	-	1	45	77	18	1	-	-	-	-	-
ALL OTHER, MALE													
1967
1966
1965
1964
1963
1962
1961
1960
ALL OTHER, FEMALE													
1967	84	-	-	-	33	36	14	1	-	-	-	-	-
1966	93	-	-	2	35	38	18	-	-	-	-	-	-
1965	129	-	-	-	47	61	21	-	-	-	-	-	-
1964	130	-	-	1	49	54	25	1	-	-	-	-	-
1963	107	-	-	2	28	50	27	-	-	-	-	-	-
1962	148	-	-	1	53	73	21	-	-	-	-	-	-
1961	161	-	-	1	54	85	21	-	-	-	-	-	-
1960	142	-	-	1	45	77	18	1	-	-	-	-	-

- Quantity zero

... Not applicable.

Note: Figures by race exclude data for residents of New Jersey for 1962 and 1963.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290A for 1960-67 and separate unpublished tables for the black population and Accidents.

77/09/02 TABLE 290A. DEATHS FOR 60 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1960-67

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	TOTAL	UNDER 1 YEAR	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS	65-74 YEARS	75-84 YEARS	85+ YEARS	N.S.
Other complications of pregnancy, childbirth, and the puerperium (640-649,660-689)													
BOTH SEXES													
1967	827	-	-	4	283	308	220	12	-	-	-	-	-
1966	860	-	-	5	264	344	235	11	-	-	-	-	1
1965	954	-	-	3	265	384	295	7	-	-	-	-	-
1964	1,096	-	-	2	279	456	344	15	-	-	-	-	-
1963	1,186	-	-	5	331	473	368	9	-	-	-	-	-
1962	1,160	-	-	2	288	528	336	6	-	-	-	-	-
1961	1,249	-	-	11	293	542	388	15	-	-	-	-	-
1960	1,290	-	-	5	345	512	416	12	-	-	-	-	-
MALE													
1967
1966
1965
1964
1963
1962
1961
1960
FEMALE													
1967	827	-	-	4	283	308	220	12	-	-	-	-	-
1966	860	-	-	5	264	344	235	11	-	-	-	-	1
1965	954	-	-	3	265	384	295	7	-	-	-	-	-
1964	1,096	-	-	2	279	456	344	15	-	-	-	-	-
1963	1,186	-	-	5	331	473	368	9	-	-	-	-	-
1962	1,160	-	-	2	288	528	336	6	-	-	-	-	-
1961	1,249	-	-	11	293	542	388	15	-	-	-	-	-
1960	1,290	-	-	5	345	512	416	12	-	-	-	-	-

- Quantity zero

... Not applicable.

Note: Figures by race exclude data for residents of New Jersey for 1962 and 1963.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290A for 1960-67 and separate unpublished tables for the black population and Accidents.

77/09/02 TABLE 290A. DEATHS FOR 60 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1960-67

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	TOTAL	UNDER 1 YEAR	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS	65-74 YEARS	75-84 YEARS	85+ YEARS	N.S.
Other complications of pregnancy, childbirth, and the puerperium (640-649,660-689)													
WHITE, BOTH SEXES													
1967	495	-	-	-	158	200	130	7	-	-	-	-	-
1966	509	-	-	1	153	211	136	8	-	-	-	-	-
1965	550	-	-	-	166	204	176	4	-	-	-	-	-
1964	634	-	-	-	157	265	200	12	-	-	-	-	-
1963	636	-	-	-	185	240	205	6	-	-	-	-	-
1962	658	-	-	-	165	307	182	4	-	-	-	-	-
1961	734	-	-	3	175	316	233	7	-	-	-	-	-
1960	789	-	-	1	219	310	254	5	-	-	-	-	-
WHITE, MALE													
1967
1966
1965
1964
1963
1962
1961
1960
WHITE, FEMALE													
1967	495	-	-	-	158	200	130	7	-	-	-	-	-
1966	509	-	-	1	153	211	136	8	-	-	-	-	-
1965	550	-	-	-	166	204	176	4	-	-	-	-	-
1964	634	-	-	-	157	265	200	12	-	-	-	-	-
1963	636	-	-	-	185	240	205	6	-	-	-	-	-
1962	658	-	-	-	165	307	182	4	-	-	-	-	-
1961	734	-	-	3	175	316	233	7	-	-	-	-	-
1960	789	-	-	1	219	310	254	5	-	-	-	-	-

- Quantity zero

... Not applicable.

Note: Figures by race exclude data for residents of New Jersey for 1962 and 1963.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290A for 1960-67 and separate unpublished tables for the black population and Accidents.

77/09/02 TABLE 290A. DEATHS FOR 60 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1960-67

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	TOTAL	UNDER 1 YEAR	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS	65-74 YEARS	75-84 YEARS	85+ YEARS	N.S.
Other complications of pregnancy, childbirth, and the puerperium (640-649,660-689)													
ALL OTHER, BOTH SEXES													
1967	332	-	-	4	125	108	90	5	-	-	-	-	-
1966	351	-	-	4	111	133	99	3	-	-	-	-	1
1965	404	-	-	3	99	180	119	3	-	-	-	-	-
1964	462	-	-	2	122	191	144	3	-	-	-	-	-
1963	512	-	-	5	138	211	155	3	-	-	-	-	-
1962	467	-	-	2	119	200	144	2	-	-	-	-	-
1961	515	-	-	8	118	226	155	8	-	-	-	-	-
1960	501	-	-	4	126	202	162	7	-	-	-	-	-
ALL OTHER, MALE													
1967
1966
1965
1964
1963
1962
1961
1960
ALL OTHER, FEMALE													
1967	332	-	-	4	125	108	90	5	-	-	-	-	-
1966	351	-	-	4	111	133	99	3	-	-	-	-	1
1965	404	-	-	3	99	180	119	3	-	-	-	-	-
1964	462	-	-	2	122	191	144	3	-	-	-	-	-
1963	512	-	-	5	138	211	155	3	-	-	-	-	-
1962	467	-	-	2	119	200	144	2	-	-	-	-	-
1961	515	-	-	8	118	226	155	8	-	-	-	-	-
1960	501	-	-	4	126	202	162	7	-	-	-	-	-

- Quantity zero

... Not applicable.

Note: Figures by race exclude data for residents of New Jersey for 1962 and 1963.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290A for 1960-67 and separate unpublished tables for the black population and Accidents.

77/09/02 TABLE 290A. DEATHS FOR 60 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1960-67

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	TOTAL	UNDER 1 YEAR	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS	65-74 YEARS	75-84 YEARS	85+ YEARS	N.S.
ICD-7 Nos. 750-759 Congenital anomalies													
BOTH SEXES													
1967	17,328	11,632	1,516	977	700	368	439	604	514	332	192	46	8
1966	18,158	12,200	1,703	1,056	718	429	458	603	513	293	138	34	13
1965	19,512	13,443	1,683	1,122	737	412	507	622	474	331	147	30	4
1964	20,288	14,197	1,709	1,111	741	374	532	610	503	333	136	38	4
1963	20,817	14,581	1,803	1,063	708	445	548	664	489	331	140	42	3
1962	21,192	14,832	1,880	1,187	725	453	561	633	444	317	125	34	1
1961	21,922	15,515	1,945	1,280	708	409	549	582	465	299	131	37	2
1960	21,860	15,389	2,086	1,287	647	465	527	547	447	303	128	30	4
MALE													
1967	9,346	6,307	797	505	418	195	244	331	265	155	108	17	4
1966	9,721	6,625	857	556	397	234	237	292	275	154	70	17	7
1965	10,375	7,178	861	608	417	212	278	315	246	168	75	14	3
1964	10,840	7,616	874	582	420	223	289	323	252	177	68	14	2
1963	11,209	7,943	880	588	407	234	305	342	240	170	76	22	2
1962	11,562	8,192	937	631	430	232	311	338	231	173	72	15	-
1961	11,919	8,542	971	691	389	217	306	317	227	172	65	22	-
1960	11,965	8,505	1,059	685	377	245	295	302	225	189	60	20	3
FEMALE													
1967	7,982	5,325	719	472	282	173	195	273	249	177	84	29	4
1966	8,437	5,575	846	500	321	195	221	311	238	139	68	17	6
1965	9,137	6,265	822	514	320	200	229	307	228	163	72	16	1
1964	9,448	6,581	835	529	321	151	243	287	251	156	68	24	2
1963	9,608	6,638	923	475	301	211	243	322	249	161	64	20	1
1962	9,630	6,640	943	556	295	221	250	295	213	144	53	19	1
1961	10,003	6,973	974	589	319	192	243	265	238	127	66	15	2
1960	9,895	6,884	1,027	602	270	220	232	245	222	114	68	10	1

- Quantity zero

Note: Figures by race exclude data for residents of New Jersey for 1962 and 1963.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290A for 1960-67 and separate unpublished tables for the black population and Accidents.

77/09/02 TABLE 290A. DEATHS FOR 60 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1960-67

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	TOTAL	UNDER 1 YEAR	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS	65-74 YEARS	75-84 YEARS	85+ YEARS	N.S.
ICD-7 Nos. 750-759 Congenital anomalies													
WHITE, BOTH SEXES													
1967	14,742	9,852	1,239	825	598	319	382	529	454	309	184	44	7
1966	15,464	10,398	1,370	886	628	365	392	537	449	269	126	34	10
1965	16,647	11,420	1,365	959	647	354	437	559	437	303	139	25	2
1964	17,483	12,229	1,399	944	656	318	467	541	464	303	126	33	3
1963	17,352	12,173	1,407	881	593	373	459	558	433	303	131	40	1
1962	17,812	12,453	1,498	1,010	609	382	472	561	393	286	113	34	1
1961	19,080	13,473	1,618	1,113	621	367	488	531	435	275	121	37	1
1960	19,056	13,426	1,719	1,136	589	399	462	480	411	287	116	27	4
WHITE, MALE													
1967	7,959	5,345	656	431	361	173	214	284	232	142	101	17	3
1966	8,265	5,636	692	460	349	197	205	259	241	143	61	17	5
1965	8,825	6,062	699	525	368	182	242	289	227	149	71	10	1
1964	9,308	6,537	703	495	377	188	260	289	234	154	59	10	2
1963	9,344	6,603	706	492	343	195	262	289	208	154	70	22	-
1962	9,671	6,835	733	540	369	199	264	301	201	151	63	15	-
1961	10,389	7,439	814	592	348	191	264	295	208	157	59	22	-
1960	10,392	7,383	866	608	341	216	262	265	200	177	54	17	3
WHITE, FEMALE													
1967	6,783	4,507	583	394	237	146	168	245	222	167	83	27	4
1966	7,199	4,762	678	426	279	168	187	278	208	126	65	17	5
1965	7,822	5,358	666	434	279	172	195	270	210	154	68	15	1
1964	8,175	5,692	696	449	279	130	207	252	230	149	67	23	1
1963	8,008	5,570	701	389	250	178	197	269	225	149	61	18	1
1962	8,141	5,618	765	470	240	183	208	260	192	135	50	19	1
1961	8,691	6,034	804	521	273	176	224	236	227	118	62	15	1
1960	8,664	6,043	853	528	248	183	200	215	211	110	62	10	1

- Quantity zero

Note: Figures by race exclude data for residents of New Jersey for 1962 and 1963.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290A for 1960-67 and separate unpublished tables for the black population and Accidents.

77/09/02 TABLE 290A. DEATHS FOR 60 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1960-67

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	TOTAL	UNDER 1 YEAR	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS	65-74 YEARS	75-84 YEARS	85+ YEARS	N.S.
ICD-7 Nos. 750-759 Congenital anomalies													
ALL OTHER, BOTH SEXES													
1967	2,586	1,780	277	152	102	49	57	75	60	23	8	2	1
1966	2,694	1,802	333	170	90	64	66	66	64	24	12	-	3
1965	2,865	2,023	318	163	90	58	70	63	37	28	8	5	2
1964	2,805	1,968	310	167	85	56	65	69	39	30	10	5	1
1963	2,792	1,933	342	148	91	61	67	80	46	17	4	1	2
1962	2,711	1,925	321	136	91	59	70	49	38	16	6	-	-
1961	2,842	2,042	327	167	87	42	61	51	30	24	10	-	1
1960	2,804	1,963	367	151	58	66	65	67	36	16	12	3	-
ALL OTHER, MALE													
1967	1,387	962	141	74	57	22	30	47	33	13	7	-	1
1966	1,456	989	165	96	48	37	32	33	34	11	9	-	2
1965	1,550	1,116	162	83	49	30	36	26	19	19	4	4	2
1964	1,532	1,079	171	87	43	35	29	34	18	23	9	4	-
1963	1,490	1,067	157	79	46	33	32	36	25	11	2	-	2
1962	1,519	1,096	172	71	51	26	37	26	22	14	4	-	-
1961	1,530	1,103	157	99	41	26	42	22	19	15	6	-	-
1960	1,573	1,122	193	77	36	29	33	37	25	12	6	3	-
ALL OTHER, FEMALE													
1967	1,199	818	136	78	45	27	27	28	27	10	1	2	-
1966	1,238	813	168	74	42	27	34	33	30	13	3	-	1
1965	1,315	907	156	80	41	28	34	37	18	9	4	1	-
1964	1,273	889	139	80	42	21	36	35	21	7	1	1	1
1963	1,302	866	185	69	45	28	35	44	21	6	2	1	-
1962	1,192	829	149	65	40	33	33	23	16	2	2	-	-
1961	1,312	939	170	68	46	16	19	29	11	9	4	-	1
1960	1,231	841	174	74	22	37	32	30	11	4	6	-	-

- Quantity zero

Note: Figures by race exclude data for residents of New Jersey for 1962 and 1963.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290A for 1960-67 and separate unpublished tables for the black population and Accidents.

77/09/02 TABLE 290A. DEATHS FOR 60 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1960-67

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	TOTAL	UNDER 1 YEAR	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS	65-74 YEARS	75-84 YEARS	85+ YEARS	N.S.
ICD-7 Nos. 760-776 Certain conditions originating in the perinatal period													
BOTH SEXES													
1967	48,314	48,271	22	6	4	1	-	-	-	-	-	-	10
1966	51,644	51,608	19	7	4	1	-	-	1	-	-	-	4
1965	55,398	55,375	10	9	-	-	-	-	-	-	-	-	4
1964	60,322	60,298	6	12	1	1	-	-	-	-	1	-	3
1963	62,688	62,664	12	9	3	-	-	-	-	-	-	-	-
1962	64,205	64,185	10	10	-	-	-	-	-	-	-	-	-
1961	65,679	65,662	10	6	-	1	-	-	-	-	-	-	-
1960	67,094	67,074	12	7	1	-	-	-	-	-	-	-	-
MALE													
1967	28,261	28,233	14	4	2	1	-	-	-	-	-	-	7
1966	30,226	30,209	8	2	4	-	-	-	-	-	-	-	3
1965	32,504	32,491	4	5	-	-	-	-	-	-	-	-	4
1964	35,322	35,310	3	6	-	1	-	-	-	-	-	-	2
1963	36,902	36,887	11	3	1	-	-	-	-	-	-	-	-
1962	37,741	37,732	6	3	-	-	-	-	-	-	-	-	-
1961	38,388	38,380	5	3	-	-	-	-	-	-	-	-	-
1960	39,103	39,091	8	4	-	-	-	-	-	-	-	-	-
FEMALE													
1967	20,053	20,038	8	2	2	-	-	-	-	-	-	-	3
1966	21,418	21,399	11	5	-	1	-	-	1	-	-	-	1
1965	22,894	22,884	6	4	-	-	-	-	-	-	-	-	-
1964	25,000	24,988	3	6	1	-	-	-	-	-	1	-	1
1963	25,786	25,777	1	6	2	-	-	-	-	-	-	-	-
1962	26,464	26,453	4	7	-	-	-	-	-	-	-	-	-
1961	27,291	27,282	5	3	-	1	-	-	-	-	-	-	-
1960	27,991	27,983	4	3	1	-	-	-	-	-	-	-	-

- Quantity zero

Note: Figures by race exclude data for residents of New Jersey for 1962 and 1963.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290A for 1960-67 and separate unpublished tables for the black population and Accidents.

77/09/02 TABLE 290A. DEATHS FOR 60 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1960-67

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	TOTAL	UNDER 1 YEAR	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS	65-74 YEARS	75-84 YEARS	85+ YEARS	N.S.
ICD-7 Nos. 760-776 Certain conditions originating in the perinatal period													
WHITE, BOTH SEXES													
1967	35,703	35,668	19	5	4	-	-	-	-	-	-	-	7
1966	38,099	38,069	15	6	4	1	-	-	1	-	-	-	3
1965	40,866	40,846	8	9	-	-	-	-	-	-	-	-	3
1964	44,618	44,599	4	10	1	1	-	-	-	-	1	-	2
1963	45,606	45,588	6	9	3	-	-	-	-	-	-	-	-
1962	47,117	47,100	8	9	-	-	-	-	-	-	-	-	-
1961	49,988	49,974	8	5	-	1	-	-	-	-	-	-	-
1960	51,276	51,260	9	6	1	-	-	-	-	-	-	-	-
WHITE, MALE													
1967	21,231	21,206	13	4	2	-	-	-	-	-	-	-	6
1966	22,637	22,624	5	2	4	-	-	-	-	-	-	-	2
1965	24,279	24,268	3	5	-	-	-	-	-	-	-	-	3
1964	26,470	26,459	3	5	-	1	-	-	-	-	-	-	2
1963	27,112	27,102	6	3	1	-	-	-	-	-	-	-	-
1962	28,058	28,050	5	3	-	-	-	-	-	-	-	-	-
1961	29,579	29,573	4	2	-	-	-	-	-	-	-	-	-
1960	30,211	30,201	6	4	-	-	-	-	-	-	-	-	-
WHITE, FEMALE													
1967	14,472	14,462	6	1	2	-	-	-	-	-	-	-	1
1966	15,462	15,445	10	4	-	1	-	-	1	-	-	-	1
1965	16,587	16,578	5	4	-	-	-	-	-	-	-	-	-
1964	18,148	18,140	1	5	1	-	-	-	-	-	1	-	-
1963	18,494	18,486	-	6	2	-	-	-	-	-	-	-	-
1962	19,059	19,050	3	6	-	-	-	-	-	-	-	-	-
1961	20,409	20,401	4	3	-	1	-	-	-	-	-	-	-
1960	21,065	21,059	3	2	1	-	-	-	-	-	-	-	-

- Quantity zero

Note: Figures by race exclude data for residents of New Jersey for 1962 and 1963.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290A for 1960-67 and separate unpublished tables for the black population and Accidents.

77/09/02 TABLE 290A. DEATHS FOR 60 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1960-67

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	TOTAL	UNDER 1 YEAR	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS	65-74 YEARS	75-84 YEARS	85+ YEARS	N.S.
ICD-7 Nos. 760-776 Certain conditions originating in the perinatal period													
All other, both sexes													
1967	12,611	12,603	3	1	-	1	-	-	-	-	-	-	3
1966	13,545	13,539	4	1	-	-	-	-	-	-	-	-	1
1965	14,532	14,529	2	-	-	-	-	-	-	-	-	-	1
1964	15,704	15,699	2	2	-	-	-	-	-	-	-	-	1
1963	14,969	14,964	5	-	-	-	-	-	-	-	-	-	-
1962	15,051	15,048	2	1	-	-	-	-	-	-	-	-	-
1961	15,691	15,688	2	1	-	-	-	-	-	-	-	-	-
1960	15,818	15,814	3	1	-	-	-	-	-	-	-	-	-
ALL OTHER, MALE													
1967	7,030	7,027	1	-	-	1	-	-	-	-	-	-	1
1966	7,589	7,585	3	-	-	-	-	-	-	-	-	-	1
1965	8,225	8,223	1	-	-	-	-	-	-	-	-	-	1
1964	8,852	8,851	-	1	-	-	-	-	-	-	-	-	-
1963	8,535	8,531	4	-	-	-	-	-	-	-	-	-	-
1962	8,478	8,477	1	-	-	-	-	-	-	-	-	-	-
1961	8,809	8,807	1	1	-	-	-	-	-	-	-	-	-
1960	8,892	8,890	2	-	-	-	-	-	-	-	-	-	-
ALL OTHER, FEMALE													
1967	5,581	5,576	2	1	-	-	-	-	-	-	-	-	2
1966	5,956	5,954	1	1	-	-	-	-	-	-	-	-	-
1965	6,307	6,306	1	-	-	-	-	-	-	-	-	-	-
1964	6,852	6,848	2	1	-	-	-	-	-	-	-	-	1
1963	6,434	6,433	1	-	-	-	-	-	-	-	-	-	-
1962	6,573	6,571	1	1	-	-	-	-	-	-	-	-	-
1961	6,882	6,881	1	-	-	-	-	-	-	-	-	-	-
1960	6,926	6,924	1	1	-	-	-	-	-	-	-	-	-

- Quantity zero

Note: Figures by race exclude data for residents of New Jersey for 1962 and 1963.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290A for 1960-67 and separate unpublished tables for the black population and Accidents.

77/09/02 TABLE 290A. DEATHS FOR 60 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1960-67

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	TOTAL	UNDER 1 YEAR	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS	65-74 YEARS	75-84 YEARS	85+ YEARS	N.S.
Birth injuries, postnatal asphyxia, and atelectasis (760-762)													
BOTH SEXES													
1967	19,354	19,344	3	-	-	1	-	-	-	-	-	-	6
1966	21,050	21,044	1	1	-	-	-	-	1	-	-	-	3
1965	23,149	23,147	-	1	-	-	-	-	-	-	-	-	1
1964	25,245	25,242	1	1	-	-	-	-	-	-	1	-	-
1963	26,889	26,888	-	1	-	-	-	-	-	-	-	-	-
1962	28,199	28,198	1	-	-	-	-	-	-	-	-	-	-
1961	28,945	28,945	-	-	-	-	-	-	-	-	-	-	-
1960	29,697	29,697	-	-	-	-	-	-	-	-	-	-	-
MALE													
1967	11,551	11,541	3	-	-	1	-	-	-	-	-	-	6
1966	12,467	12,463	1	-	-	-	-	-	-	-	-	-	3
1965	13,869	13,868	-	-	-	-	-	-	-	-	-	-	1
1964	15,026	15,025	-	1	-	-	-	-	-	-	-	-	-
1963	16,122	16,121	-	1	-	-	-	-	-	-	-	-	-
1962	16,832	16,831	1	-	-	-	-	-	-	-	-	-	-
1961	17,261	17,261	-	-	-	-	-	-	-	-	-	-	-
1960	17,660	17,660	-	-	-	-	-	-	-	-	-	-	-
FEMALE													
1967	7,803	7,803	-	-	-	-	-	-	-	-	-	-	-
1966	8,583	8,581	-	1	-	-	-	-	1	-	-	-	-
1965	9,280	9,279	-	1	-	-	-	-	-	-	-	-	-
1964	10,219	10,217	1	-	-	-	-	-	-	-	1	-	-
1963	10,767	10,767	-	-	-	-	-	-	-	-	-	-	-
1962	11,367	11,367	-	-	-	-	-	-	-	-	-	-	-
1961	11,684	11,684	-	-	-	-	-	-	-	-	-	-	-
1960	12,037	12,037	-	-	-	-	-	-	-	-	-	-	-

- Quantity zero

Note: Figures by race exclude data for residents of New Jersey for 1962 and 1963.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290A for 1960-67 and separate unpublished tables for the black population and Accidents.

77/09/02 TABLE 290A. DEATHS FOR 60 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1960-67

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	TOTAL	UNDER 1 YEAR	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS	65-74 YEARS	75-84 YEARS	85+ YEARS	N.S.
Birth injuries, postnatal asphyxia, and atelectasis (760-762)													
WHITE, BOTH SEXES													
1967	14,731	14,723	3	-	-	-	-	-	-	-	-	-	5
1966	16,031	16,027	1	-	-	-	-	-	1	-	-	-	2
1965	17,723	17,721	-	1	-	-	-	-	-	-	-	-	1
1964	19,584	19,581	1	1	-	-	-	-	-	-	1	-	-
1963	20,398	20,397	-	1	-	-	-	-	-	-	-	-	-
1962	21,510	21,509	1	-	-	-	-	-	-	-	-	-	-
1961	22,968	22,968	-	-	-	-	-	-	-	-	-	-	-
1960	23,588	23,588	-	-	-	-	-	-	-	-	-	-	-
WHITE, MALE													
1967	8,942	8,934	3	-	-	-	-	-	-	-	-	-	5
1966	9,665	9,662	1	-	-	-	-	-	-	-	-	-	2
1965	10,780	10,779	-	-	-	-	-	-	-	-	-	-	1
1964	11,782	11,781	-	1	-	-	-	-	-	-	-	-	-
1963	12,382	12,381	-	1	-	-	-	-	-	-	-	-	-
1962	12,974	12,973	1	-	-	-	-	-	-	-	-	-	-
1961	13,838	13,838	-	-	-	-	-	-	-	-	-	-	-
1960	14,216	14,216	-	-	-	-	-	-	-	-	-	-	-
WHITE, FEMALE													
1967	5,789	5,789	-	-	-	-	-	-	-	-	-	-	-
1966	6,366	6,365	-	-	-	-	-	-	1	-	-	-	-
1965	6,943	6,942	-	1	-	-	-	-	-	-	-	-	-
1964	7,802	7,800	1	-	-	-	-	-	-	-	1	-	-
1963	8,016	8,016	-	-	-	-	-	-	-	-	-	-	-
1962	8,536	8,536	-	-	-	-	-	-	-	-	-	-	-
1961	9,130	9,130	-	-	-	-	-	-	-	-	-	-	-
1960	9,372	9,372	-	-	-	-	-	-	-	-	-	-	-

- Quantity zero

Note: Figures by race exclude data for residents of New Jersey for 1962 and 1963.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290A for 1960-67 and separate unpublished tables for the black population and Accidents.

77/09/02 TABLE 290A. DEATHS FOR 60 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1960-67

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	TOTAL	UNDER 1 YEAR	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS	65-74 YEARS	75-84 YEARS	85+ YEARS	N.S.
Birth injuries, postnatal asphyxia, and atelectasis (760-762)													
ALL OTHER, BOTH SEXES													
1967	4,623	4,621	-	-	-	1	-	-	-	-	-	-	1
1966	5,019	5,017	-	1	-	-	-	-	-	-	-	-	1
1965	5,426	5,426	-	-	-	-	-	-	-	-	-	-	-
1964	5,661	5,661	-	-	-	-	-	-	-	-	-	-	-
1963	5,391	5,391	-	-	-	-	-	-	-	-	-	-	-
1962	5,655	5,655	-	-	-	-	-	-	-	-	-	-	-
1961	5,977	5,977	-	-	-	-	-	-	-	-	-	-	-
1960	6,109	6,109	-	-	-	-	-	-	-	-	-	-	-
ALL OTHER, MALE													
1967	2,609	2,607	-	-	-	1	-	-	-	-	-	-	1
1966	2,802	2,801	-	-	-	-	-	-	-	-	-	-	1
1965	3,089	3,089	-	-	-	-	-	-	-	-	-	-	-
1964	3,244	3,244	-	-	-	-	-	-	-	-	-	-	-
1963	3,078	3,078	-	-	-	-	-	-	-	-	-	-	-
1962	3,239	3,239	-	-	-	-	-	-	-	-	-	-	-
1961	3,423	3,423	-	-	-	-	-	-	-	-	-	-	-
1960	3,444	3,444	-	-	-	-	-	-	-	-	-	-	-
ALL OTHER, FEMALE													
1967	2,014	2,014	-	-	-	-	-	-	-	-	-	-	-
1966	2,217	2,216	-	1	-	-	-	-	-	-	-	-	-
1965	2,337	2,337	-	-	-	-	-	-	-	-	-	-	-
1964	2,417	2,417	-	-	-	-	-	-	-	-	-	-	-
1963	2,313	2,313	-	-	-	-	-	-	-	-	-	-	-
1962	2,416	2,416	-	-	-	-	-	-	-	-	-	-	-
1961	2,554	2,554	-	-	-	-	-	-	-	-	-	-	-
1960	2,665	2,665	-	-	-	-	-	-	-	-	-	-	-

- Quantity zero

Note: Figures by race exclude data for residents of New Jersey for 1962 and 1963.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290A for 1960-67 and separate unpublished tables for the black population and Accidents.

77/09/02 TABLE 290A. DEATHS FOR 60 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1960-67

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	TOTAL	UNDER 1 YEAR	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS	65-74 YEARS	75-84 YEARS	85+ YEARS	N.S.
Infections of newborn (763-768)													
BOTH SEXES													
1967	3,269	3,269	-	-	-	-	-	-	-	-	-	-	-
1966	3,522	3,521	1	-	-	-	-	-	-	-	-	-	-
1965	3,926	3,926	-	-	-	-	-	-	-	-	-	-	-
1964	4,356	4,356	-	-	-	-	-	-	-	-	-	-	-
1963	4,408	4,408	-	-	-	-	-	-	-	-	-	-	-
1962	4,551	4,551	-	-	-	-	-	-	-	-	-	-	-
1961	4,706	4,706	-	-	-	-	-	-	-	-	-	-	-
1960	4,786	4,786	-	-	-	-	-	-	-	-	-	-	-
MALE													
1967	1,873	1,873	-	-	-	-	-	-	-	-	-	-	-
1966	1,996	1,995	1	-	-	-	-	-	-	-	-	-	-
1965	2,310	2,310	-	-	-	-	-	-	-	-	-	-	-
1964	2,497	2,497	-	-	-	-	-	-	-	-	-	-	-
1963	2,573	2,573	-	-	-	-	-	-	-	-	-	-	-
1962	2,682	2,682	-	-	-	-	-	-	-	-	-	-	-
1961	2,783	2,783	-	-	-	-	-	-	-	-	-	-	-
1960	2,799	2,799	-	-	-	-	-	-	-	-	-	-	-
FEMALE													
1967	1,396	1,396	-	-	-	-	-	-	-	-	-	-	-
1966	1,526	1,526	-	-	-	-	-	-	-	-	-	-	-
1965	1,616	1,616	-	-	-	-	-	-	-	-	-	-	-
1964	1,859	1,859	-	-	-	-	-	-	-	-	-	-	-
1963	1,835	1,835	-	-	-	-	-	-	-	-	-	-	-
1962	1,869	1,869	-	-	-	-	-	-	-	-	-	-	-
1961	1,923	1,923	-	-	-	-	-	-	-	-	-	-	-
1960	1,987	1,987	-	-	-	-	-	-	-	-	-	-	-

- Quantity zero

Note: Figures by race exclude data for residents of New Jersey for 1962 and 1963.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290A for 1960-67 and separate unpublished tables for the black population and Accidents.

77/09/02 TABLE 290A. DEATHS FOR 60 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1960-67

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	TOTAL	UNDER 1 YEAR	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS	65-74 YEARS	75-84 YEARS	85+ YEARS	N.S.
Infections of newborn (763-768)													
WHITE, BOTH SEXES													
1967	2,152	2,152	-	-	-	-	-	-	-	-	-	-	-
1966	2,209	2,209	-	-	-	-	-	-	-	-	-	-	-
1965	2,413	2,413	-	-	-	-	-	-	-	-	-	-	-
1964	2,642	2,642	-	-	-	-	-	-	-	-	-	-	-
1963	2,729	2,729	-	-	-	-	-	-	-	-	-	-	-
1962	2,790	2,790	-	-	-	-	-	-	-	-	-	-	-
1961	2,978	2,978	-	-	-	-	-	-	-	-	-	-	-
1960	3,124	3,124	-	-	-	-	-	-	-	-	-	-	-
WHITE, MALE													
1967	1,263	1,263	-	-	-	-	-	-	-	-	-	-	-
1966	1,269	1,269	-	-	-	-	-	-	-	-	-	-	-
1965	1,439	1,439	-	-	-	-	-	-	-	-	-	-	-
1964	1,537	1,537	-	-	-	-	-	-	-	-	-	-	-
1963	1,591	1,591	-	-	-	-	-	-	-	-	-	-	-
1962	1,678	1,678	-	-	-	-	-	-	-	-	-	-	-
1961	1,790	1,790	-	-	-	-	-	-	-	-	-	-	-
1960	1,854	1,854	-	-	-	-	-	-	-	-	-	-	-
WHITE, FEMALE													
1967	889	889	-	-	-	-	-	-	-	-	-	-	-
1966	940	940	-	-	-	-	-	-	-	-	-	-	-
1965	974	974	-	-	-	-	-	-	-	-	-	-	-
1964	1,105	1,105	-	-	-	-	-	-	-	-	-	-	-
1963	1,138	1,138	-	-	-	-	-	-	-	-	-	-	-
1962	1,112	1,112	-	-	-	-	-	-	-	-	-	-	-
1961	1,188	1,188	-	-	-	-	-	-	-	-	-	-	-
1960	1,270	1,270	-	-	-	-	-	-	-	-	-	-	-

- Quantity zero

Note: Figures by race exclude data for residents of New Jersey for 1962 and 1963.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290A for 1960-67 and separate unpublished tables for the black population and Accidents.

77/09/02 TABLE 290A. DEATHS FOR 60 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1960-67

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	TOTAL	UNDER 1 YEAR	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS	65-74 YEARS	75-84 YEARS	85+ YEARS	N.S.
Infections of newborn (763-768)													
ALL OTHER, BOTH SEXES													
1967	1,117	1,117	-	-	-	-	-	-	-	-	-	-	-
1966	1,313	1,312	1	-	-	-	-	-	-	-	-	-	-
1965	1,513	1,513	-	-	-	-	-	-	-	-	-	-	-
1964	1,714	1,714	-	-	-	-	-	-	-	-	-	-	-
1963	1,554	1,554	-	-	-	-	-	-	-	-	-	-	-
1962	1,654	1,654	-	-	-	-	-	-	-	-	-	-	-
1961	1,728	1,728	-	-	-	-	-	-	-	-	-	-	-
1960	1,662	1,662	-	-	-	-	-	-	-	-	-	-	-
ALL OTHER, MALE													
1967	610	610	-	-	-	-	-	-	-	-	-	-	-
1966	727	726	1	-	-	-	-	-	-	-	-	-	-
1965	871	871	-	-	-	-	-	-	-	-	-	-	-
1964	960	960	-	-	-	-	-	-	-	-	-	-	-
1963	903	903	-	-	-	-	-	-	-	-	-	-	-
1962	936	936	-	-	-	-	-	-	-	-	-	-	-
1961	993	993	-	-	-	-	-	-	-	-	-	-	-
1960	945	945	-	-	-	-	-	-	-	-	-	-	-
ALL OTHER, FEMALE													
1967	507	507	-	-	-	-	-	-	-	-	-	-	-
1966	586	586	-	-	-	-	-	-	-	-	-	-	-
1965	642	642	-	-	-	-	-	-	-	-	-	-	-
1964	754	754	-	-	-	-	-	-	-	-	-	-	-
1963	651	651	-	-	-	-	-	-	-	-	-	-	-
1962	718	718	-	-	-	-	-	-	-	-	-	-	-
1961	735	735	-	-	-	-	-	-	-	-	-	-	-
1960	717	717	-	-	-	-	-	-	-	-	-	-	-

- Quantity zero

Note: Figures by race exclude data for residents of New Jersey for 1962 and 1963.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290A for 1960-67 and separate unpublished tables for the black population and Accidents.

77/09/02 TABLE 290A. DEATHS FOR 60 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1960-67

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	TOTAL	UNDER 1 YEAR	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS	65-74 YEARS	75-84 YEARS	85+ YEARS	N.S.
Other diseases peculiar to early infancy, and immaturity, unqualified (769-776)													
BOTH SEXES													
1967	25,691	25,658	19	6	4	-	-	-	-	-	-	-	4
1966	27,072	27,043	17	6	4	1	-	-	-	-	-	-	1
1965	28,323	28,302	10	8	-	-	-	-	-	-	-	-	3
1964	30,721	30,700	5	11	1	1	-	-	-	-	-	-	3
1963	31,391	31,368	12	8	3	-	-	-	-	-	-	-	-
1962	31,455	31,436	9	10	-	-	-	-	-	-	-	-	-
1961	32,028	32,011	10	6	-	1	-	-	-	-	-	-	-
1960	32,611	32,591	12	7	1	-	-	-	-	-	-	-	-
MALE													
1967	14,837	14,819	11	4	2	-	-	-	-	-	-	-	1
1966	15,763	15,751	6	2	4	-	-	-	-	-	-	-	-
1965	16,325	16,313	4	5	-	-	-	-	-	-	-	-	3
1964	17,799	17,788	3	5	-	1	-	-	-	-	-	-	2
1963	18,207	18,193	11	2	1	-	-	-	-	-	-	-	-
1962	18,227	18,219	5	3	-	-	-	-	-	-	-	-	-
1961	18,344	18,336	5	3	-	-	-	-	-	-	-	-	-
1960	18,644	18,632	8	4	-	-	-	-	-	-	-	-	-
FEMALE													
1967	10,854	10,839	8	2	2	-	-	-	-	-	-	-	3
1966	11,309	11,292	11	4	-	1	-	-	-	-	-	-	1
1965	11,998	11,989	6	3	-	-	-	-	-	-	-	-	-
1964	12,922	12,912	2	6	1	-	-	-	-	-	-	-	1
1963	13,184	13,175	1	6	2	-	-	-	-	-	-	-	-
1962	13,228	13,217	4	7	-	-	-	-	-	-	-	-	-
1961	13,684	13,675	5	3	-	1	-	-	-	-	-	-	-
1960	13,967	13,959	4	3	1	-	-	-	-	-	-	-	-

- Quantity zero

Note: Figures by race exclude data for residents of New Jersey for 1962 and 1963.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290A for 1960-67 and separate unpublished tables for the black population and Accidents.

77/09/02 TABLE 290A. DEATHS FOR 60 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1960-67

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	TOTAL	UNDER 1 YEAR	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS	65-74 YEARS	75-84 YEARS	85+ YEARS	N.S.
Other diseases peculiar to early infancy, and immaturity, unqualified (769-776)													
WHITE, BOTH SEXES													
1967	18,820	18,793	16	5	4	-	-	-	-	-	-	-	2
1966	19,859	19,833	14	6	4	1	-	-	-	-	-	-	1
1965	20,730	20,712	8	8	-	-	-	-	-	-	-	-	2
1964	22,392	22,376	3	9	1	1	-	-	-	-	-	-	2
1963	22,479	22,462	6	8	3	-	-	-	-	-	-	-	-
1962	22,817	22,801	7	9	-	-	-	-	-	-	-	-	-
1961	24,042	24,028	8	5	-	1	-	-	-	-	-	-	-
1960	24,564	24,548	9	6	1	-	-	-	-	-	-	-	-
WHITE, MALE													
1967	11,026	11,009	10	4	2	-	-	-	-	-	-	-	1
1966	11,703	11,693	4	2	4	-	-	-	-	-	-	-	-
1965	12,060	12,050	3	5	-	-	-	-	-	-	-	-	2
1964	13,151	13,141	3	4	-	1	-	-	-	-	-	-	2
1963	13,139	13,130	6	2	1	-	-	-	-	-	-	-	-
1962	13,406	13,399	4	3	-	-	-	-	-	-	-	-	-
1961	13,951	13,945	4	2	-	-	-	-	-	-	-	-	-
1960	14,141	14,131	6	4	-	-	-	-	-	-	-	-	-
WHITE, FEMALE													
1967	7,794	7,784	6	1	2	-	-	-	-	-	-	-	1
1966	8,156	8,140	10	4	-	1	-	-	-	-	-	-	1
1965	8,670	8,662	5	3	-	-	-	-	-	-	-	-	-
1964	9,241	9,235	-	5	1	-	-	-	-	-	-	-	-
1963	9,340	9,332	-	6	2	-	-	-	-	-	-	-	-
1962	9,411	9,402	3	6	-	-	-	-	-	-	-	-	-
1961	10,091	10,083	4	3	-	1	-	-	-	-	-	-	-
1960	10,423	10,417	3	2	1	-	-	-	-	-	-	-	-

- Quantity zero

Note: Figures by race exclude data for residents of New Jersey for 1962 and 1963.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290A for 1960-67 and separate unpublished tables for the black population and Accidents.

77/09/02 TABLE 290A. DEATHS FOR 60 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1960-67

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	TOTAL	UNDER 1 YEAR	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS	65-74 YEARS	75-84 YEARS	85+ YEARS	N.S.
Other diseases peculiar to early infancy, and immaturity, unqualified (769-776)													
ALL OTHER, BOTH SEXES													
1967	6,871	6,865	3	1	-	-	-	-	-	-	-	-	2
1966	7,213	7,210	3	-	-	-	-	-	-	-	-	-	-
1965	7,593	7,590	2	-	-	-	-	-	-	-	-	-	1
1964	8,329	8,324	2	2	-	-	-	-	-	-	-	-	1
1963	8,024	8,019	5	-	-	-	-	-	-	-	-	-	-
1962	7,742	7,739	2	1	-	-	-	-	-	-	-	-	-
1961	7,986	7,983	2	1	-	-	-	-	-	-	-	-	-
1960	8,047	8,043	3	1	-	-	-	-	-	-	-	-	-
ALL OTHER, MALE													
1967	3,811	3,810	1	-	-	-	-	-	-	-	-	-	-
1966	4,060	4,058	2	-	-	-	-	-	-	-	-	-	-
1965	4,265	4,263	1	-	-	-	-	-	-	-	-	-	1
1964	4,648	4,647	-	1	-	-	-	-	-	-	-	-	-
1963	4,554	4,550	4	-	-	-	-	-	-	-	-	-	-
1962	4,303	4,302	1	-	-	-	-	-	-	-	-	-	-
1961	4,393	4,391	1	1	-	-	-	-	-	-	-	-	-
1960	4,503	4,501	2	-	-	-	-	-	-	-	-	-	-
ALL OTHER, FEMALE													
1967	3,060	3,055	2	1	-	-	-	-	-	-	-	-	2
1966	3,153	3,152	1	-	-	-	-	-	-	-	-	-	-
1965	3,328	3,327	1	-	-	-	-	-	-	-	-	-	-
1964	3,681	3,677	2	1	-	-	-	-	-	-	-	-	1
1963	3,470	3,469	1	-	-	-	-	-	-	-	-	-	-
1962	3,439	3,437	1	1	-	-	-	-	-	-	-	-	-
1961	3,593	3,592	1	-	-	-	-	-	-	-	-	-	-
1960	3,544	3,542	1	1	-	-	-	-	-	-	-	-	-

- Quantity zero

Note: Figures by race exclude data for residents of New Jersey for 1962 and 1963.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290A for 1960-67 and separate unpublished tables for the black population and Accidents.

77/09/02 TABLE 290A. DEATHS FOR 60 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1960-67

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	TOTAL	UNDER 1 YEAR	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS	65-74 YEARS	75-84 YEARS	85+ YEARS	N.S.
Symptoms, senility, and ill-defined conditions (780-795)													
BOTH SEXES													
1967	24,098	2,620	381	198	569	772	1,624	2,753	3,958	4,413	4,012	2,746	52
1966	23,960	2,644	448	172	431	716	1,554	2,654	3,786	4,548	4,014	2,932	61
1965	23,414	2,369	396	163	403	656	1,482	2,701	3,745	4,471	4,143	2,849	36
1964	25,259	2,588	479	299	777	1,040	1,858	2,941	3,918	4,462	4,031	2,820	46
1963	22,420	2,533	399	216	412	642	1,411	2,378	3,341	4,140	3,911	2,975	62
1962	19,730	2,266	351	174	361	638	1,235	2,039	2,859	3,457	3,497	2,797	56
1961	19,094	2,395	369	143	313	520	1,057	1,922	2,664	3,415	3,457	2,798	41
1960	20,385	2,518	455	187	307	584	1,120	2,052	2,799	3,659	3,781	2,874	49
MALE													
1967	14,394	1,538	212	109	344	432	951	1,874	2,705	2,812	2,225	1,158	34
1966	14,120	1,488	248	102	259	406	903	1,784	2,570	2,897	2,206	1,210	47
1965	13,929	1,383	215	90	242	351	896	1,788	2,583	2,877	2,289	1,190	25
1964	15,077	1,498	275	186	520	647	1,106	1,888	2,667	2,874	2,224	1,163	29
1963	13,131	1,457	220	110	254	352	819	1,574	2,252	2,642	2,139	1,265	47
1962	11,468	1,266	204	102	212	335	691	1,365	1,937	2,261	1,883	1,182	30
1961	11,122	1,365	199	79	173	289	633	1,310	1,830	2,170	1,869	1,178	27
1960	12,012	1,415	263	111	183	324	675	1,399	1,908	2,364	2,158	1,179	33
FEMALE													
1967	9,704	1,082	169	89	225	340	673	879	1,253	1,601	1,787	1,588	18
1966	9,840	1,156	200	70	172	310	651	870	1,216	1,651	1,808	1,722	14
1965	9,485	986	181	73	161	305	586	913	1,162	1,594	1,854	1,659	11
1964	10,182	1,090	204	113	257	393	752	1,053	1,251	1,588	1,807	1,657	17
1963	9,289	1,076	179	106	158	290	592	804	1,089	1,498	1,772	1,710	15
1962	8,262	1,000	147	72	149	303	544	674	922	1,196	1,614	1,615	26
1961	7,972	1,030	170	64	140	231	424	612	834	1,245	1,588	1,620	14
1960	8,373	1,103	192	76	124	260	445	653	891	1,295	1,623	1,695	16

Note: Figures by race exclude data for residents of New Jersey for 1962 and 1963.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290A for 1960-67 and separate unpublished tables for the black population and Accidents.

77/09/02 TABLE 290A. DEATHS FOR 60 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1960-67

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	TOTAL	UNDER 1 YEAR	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS	65-74 YEARS	75-84 YEARS	85+ YEARS	N.S.
Symptoms, senility, and ill-defined conditions (780-795)													
WHITE, BOTH SEXES													
1967	15,955	1,286	193	121	373	421	943	1,814	2,720	2,909	2,984	2,163	28
1966	15,748	1,252	231	103	290	368	904	1,695	2,523	3,052	2,991	2,307	32
1965	15,383	1,077	179	88	270	335	856	1,712	2,409	3,017	3,103	2,317	20
1964	16,734	1,159	256	190	556	610	1,105	1,874	2,583	3,011	3,080	2,279	31
1963	14,364	1,026	161	134	276	362	851	1,519	2,113	2,685	2,898	2,299	40
1962	12,795	872	174	106	242	366	738	1,299	1,811	2,305	2,653	2,191	38
1961	12,427	975	182	88	191	285	665	1,253	1,700	2,271	2,602	2,189	26
1960	13,109	972	216	108	195	326	682	1,289	1,805	2,439	2,809	2,242	26
WHITE, MALE													
1967	9,732	806	109	76	229	234	547	1,271	1,967	1,922	1,677	872	22
1966	9,509	766	121	63	181	205	539	1,202	1,809	2,018	1,652	926	27
1965	9,404	648	95	47	164	188	535	1,213	1,784	2,046	1,719	948	17
1964	10,237	686	159	121	378	395	673	1,281	1,877	2,012	1,700	934	21
1963	8,608	596	90	68	176	192	519	1,061	1,538	1,810	1,577	950	31
1962	7,616	493	96	68	149	186	443	912	1,326	1,582	1,434	905	22
1961	7,454	576	97	51	104	168	419	908	1,271	1,528	1,411	903	18
1960	7,928	560	130	61	123	182	438	934	1,323	1,656	1,606	896	19
WHITE, FEMALE													
1967	6,223	480	84	45	144	187	396	543	753	987	1,307	1,291	6
1966	6,239	486	110	40	109	163	365	493	714	1,034	1,339	1,381	5
1965	5,979	429	84	41	106	147	321	499	625	971	1,384	1,369	3
1964	6,497	473	97	69	178	215	432	593	706	999	1,380	1,345	10
1963	5,756	430	71	66	100	170	332	458	575	875	1,321	1,349	9
1962	5,179	379	78	38	93	180	295	387	485	723	1,219	1,286	16
1961	4,973	399	85	37	87	117	246	345	429	743	1,191	1,286	8
1960	5,181	412	86	47	72	144	244	355	482	783	1,203	1,346	7

Note: Figures by race exclude data for residents of New Jersey for 1962 and 1963.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290A for 1960-67 and separate unpublished tables for the black population and Accidents.

77/09/02 TABLE 290A. DEATHS FOR 60 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1960-67

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	TOTAL	UNDER 1 YEAR	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS	65-74 YEARS	75-84 YEARS	85+ YEARS	N.S.
Symptoms, senility, and ill-defined conditions (780-795)													
ALL OTHER, BOTH SEXES													
1967	8,143	1,334	188	77	196	351	681	939	1,238	1,504	1,028	583	24
1966	8,212	1,392	217	69	141	348	650	959	1,263	1,496	1,023	625	29
1965	8,031	1,292	217	75	133	321	626	989	1,336	1,454	1,040	532	16
1964	8,525	1,429	223	109	221	430	753	1,067	1,335	1,451	951	541	15
1963	7,816	1,495	233	76	122	257	535	833	1,198	1,420	979	647	21
1962	6,788	1,379	174	66	118	261	484	723	1,031	1,134	825	575	18
1961	6,667	1,420	187	55	122	235	392	669	964	1,144	855	609	15
1960	7,276	1,546	239	79	112	258	438	763	994	1,220	972	632	23
ALL OTHER, MALE													
1967	4,662	732	103	33	115	198	404	603	738	890	548	286	12
1966	4,611	722	127	39	78	201	364	582	761	879	554	284	20
1965	4,525	735	120	43	78	163	361	575	799	831	570	242	8
1964	4,840	812	116	65	142	252	433	607	790	862	524	229	8
1963	4,391	855	128	41	68	145	288	496	695	810	545	305	15
1962	3,791	765	107	32	63	145	243	442	603	672	441	270	8
1961	3,668	789	102	28	69	121	214	402	559	642	458	275	9
1960	4,084	855	133	50	60	142	237	465	585	708	552	283	14
ALL OTHER, FEMALE													
1967	3,481	602	85	44	81	153	277	336	500	614	480	297	12
1966	3,601	670	90	30	63	147	286	377	502	617	469	341	9
1965	3,506	557	97	32	55	158	265	414	537	623	470	290	8
1964	3,685	617	107	44	79	178	320	460	545	589	427	312	7
1963	3,425	640	105	35	54	112	247	337	503	610	434	342	6
1962	2,997	614	67	34	55	116	241	281	428	462	384	305	10
1961	2,999	631	85	27	53	114	178	267	405	502	397	334	6
1960	3,192	691	106	29	52	116	201	298	409	512	420	349	9

Note: Figures by race exclude data for residents of New Jersey for 1962 and 1963.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290A for 1960-67 and separate unpublished tables for the black population and Accidents.

77/09/02 TABLE 290A. DEATHS FOR 60 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1960-67

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	TOTAL	UNDER 1 YEAR	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS	65-74 YEARS	75-84 YEARS	85+ YEARS	N.S.
All other diseases (Residual)													
BOTH SEXES													
1967	57,165	1,864	1,197	1,515	1,913	2,135	4,414	7,079	9,592	12,052	10,891	4,491	22
1966	57,152	2,032	1,355	1,579	1,903	2,137	4,368	7,227	9,524	12,029	10,608	4,366	24
1965	55,393	2,352	1,375	1,584	1,769	1,970	4,259	6,742	9,229	11,761	10,257	4,077	18
1964	54,600	2,604	1,465	1,632	1,659	2,036	4,233	6,599	8,910	11,524	9,915	3,999	24
1963	54,150	2,617	1,488	1,617	1,577	2,044	4,280	6,540	8,970	11,499	9,491	4,014	13
1962	52,173	2,710	1,553	1,510	1,499	2,074	4,142	6,377	8,408	11,117	8,974	3,795	14
1961	51,620	2,890	1,610	1,572	1,554	2,097	4,158	6,265	8,268	10,699	8,859	3,622	26
1960	51,921	2,971	1,734	1,594	1,501	2,153	4,138	6,378	8,303	10,977	8,589	3,559	24
MALE													
1967	31,316	1,057	633	823	1,171	1,195	2,423	4,050	5,709	6,783	5,596	1,865	11
1966	31,156	1,165	735	843	1,130	1,126	2,378	4,189	5,584	6,802	5,420	1,768	16
1965	30,400	1,367	740	864	1,077	1,053	2,267	3,845	5,459	6,678	5,322	1,716	12
1964	29,901	1,505	802	882	980	1,080	2,327	3,755	5,238	6,548	5,051	1,719	14
1963	29,634	1,505	805	830	943	1,067	2,271	3,866	5,279	6,575	4,818	1,670	5
1962	28,569	1,622	818	821	873	1,070	2,244	3,681	4,887	6,392	4,517	1,635	9
1961	28,560	1,677	901	850	918	1,059	2,337	3,661	4,914	6,210	4,470	1,553	10
1960	28,874	1,776	978	856	873	1,143	2,201	3,766	4,960	6,375	4,372	1,559	15
FEMALE													
1967	25,849	807	564	692	742	940	1,991	3,029	3,883	5,269	5,295	2,626	11
1966	25,996	867	620	736	773	1,011	1,990	3,038	3,940	5,227	5,188	2,598	8
1965	24,993	985	635	720	692	917	1,992	2,897	3,770	5,083	4,935	2,361	6
1964	24,699	1,099	663	750	679	956	1,906	2,844	3,672	4,976	4,864	2,280	10
1963	24,516	1,112	683	787	634	977	2,009	2,674	3,691	4,924	4,673	2,344	8
1962	23,604	1,088	735	689	626	1,004	1,898	2,696	3,521	4,725	4,457	2,160	5
1961	23,060	1,213	709	722	636	1,038	1,821	2,604	3,354	4,489	4,389	2,069	16
1960	23,047	1,195	756	738	628	1,010	1,937	2,612	3,343	4,602	4,217	2,000	9

Note: Figures by race exclude data for residents of New Jersey for 1962 and 1963.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290A for 1960-67 and separate unpublished tables for the black population and Accidents.

77/09/02 TABLE 290A. DEATHS FOR 60 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1960-67

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	TOTAL	UNDER 1 YEAR	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS	65-74 YEARS	75-84 YEARS	85+ YEARS	N.S.
All other diseases (Residual)													
WHITE, BOTH SEXES													
1967	48,712	1,406	968	1,280	1,503	1,395	3,114	5,662	8,177	10,782	10,190	4,223	12
1966	48,643	1,518	1,074	1,318	1,493	1,447	3,072	5,837	8,143	10,757	9,877	4,096	11
1965	47,492	1,745	1,101	1,313	1,401	1,334	3,065	5,469	8,018	10,567	9,627	3,839	13
1964	46,814	1,964	1,160	1,351	1,320	1,366	3,032	5,312	7,711	10,454	9,371	3,760	13
1963	45,056	1,924	1,170	1,310	1,199	1,352	3,026	5,168	7,476	10,033	8,682	3,706	10
1962	43,524	2,023	1,192	1,231	1,116	1,377	3,019	5,060	7,028	9,773	8,201	3,497	7
1961	44,534	2,265	1,323	1,351	1,188	1,424	3,098	5,167	7,196	9,737	8,368	3,399	18
1960	44,651	2,306	1,417	1,359	1,178	1,491	3,112	5,207	7,225	9,959	8,046	3,337	14
WHITE, MALE													
1967	26,535	801	522	709	941	769	1,707	3,220	4,858	6,057	5,208	1,739	4
1966	26,330	878	573	700	912	748	1,650	3,381	4,794	6,051	4,996	1,639	8
1965	25,896	1,013	600	725	875	714	1,634	3,086	4,734	5,960	4,948	1,599	8
1964	25,551	1,141	635	735	796	724	1,676	3,027	4,551	5,901	4,745	1,612	8
1963	24,658	1,113	633	680	731	707	1,647	3,108	4,418	5,722	4,380	1,517	2
1962	23,831	1,206	616	686	674	709	1,653	2,940	4,097	5,641	4,109	1,497	3
1961	24,540	1,323	737	718	717	723	1,735	3,041	4,279	5,622	4,204	1,433	8
1960	24,781	1,373	797	733	703	821	1,675	3,096	4,329	5,744	4,063	1,438	9
WHITE, FEMALE													
1967	22,177	605	446	571	562	626	1,407	2,442	3,319	4,725	4,982	2,484	8
1966	22,313	640	501	618	581	699	1,422	2,456	3,349	4,706	4,881	2,457	3
1965	21,596	732	501	588	526	620	1,431	2,383	3,284	4,607	4,679	2,240	5
1964	21,263	823	525	616	524	642	1,356	2,285	3,160	4,553	4,626	2,148	5
1963	20,398	811	537	630	468	645	1,379	2,060	3,058	4,311	4,302	2,189	8
1962	19,693	817	576	545	442	668	1,366	2,120	2,931	4,132	4,092	2,000	4
1961	19,994	942	586	633	471	701	1,363	2,126	2,917	4,115	4,164	1,966	10
1960	19,870	933	620	626	475	670	1,437	2,111	2,896	4,215	3,983	1,899	5

Note: Figures by race exclude data for residents of New Jersey for 1962 and 1963.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290A for 1960-67 and separate unpublished tables for the black population and Accidents.

77/09/02 TABLE 290A. DEATHS FOR 60 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1960-67

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	TOTAL	UNDER 1 YEAR	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS	65-74 YEARS	75-84 YEARS	85+ YEARS	N.S.
All other diseases (Residual)													
ALL OTHER, BOTH SEXES													
1967	8,453	458	229	235	410	740	1,300	1,417	1,415	1,270	701	268	10
1966	8,509	514	281	261	410	690	1,296	1,390	1,381	1,272	731	270	13
1965	7,901	607	274	271	368	636	1,194	1,273	1,211	1,194	630	238	5
1964	7,786	640	305	281	339	670	1,201	1,287	1,199	1,070	544	239	11
1963	7,449	630	259	262	329	618	1,101	1,163	1,188	1,083	580	233	3
1962	7,143	621	309	240	329	634	994	1,094	1,140	999	547	229	7
1961	7,086	625	287	221	366	673	1,060	1,098	1,072	962	491	223	8
1960	7,270	665	317	235	323	662	1,026	1,171	1,078	1,018	543	222	10
ALL OTHER, MALE													
1967	4,781	256	111	114	230	426	716	830	851	726	388	126	7
1966	4,826	287	162	143	218	378	728	808	790	751	424	129	8
1965	4,504	354	140	139	202	339	633	759	725	718	374	117	4
1964	4,350	364	167	147	184	356	651	728	687	647	306	107	6
1963	4,075	356	143	121	179	313	538	640	684	644	326	128	3
1962	3,918	374	177	117	164	324	526	604	645	563	302	116	6
1961	4,020	354	164	132	201	336	602	620	635	588	266	120	2
1960	4,093	403	181	123	170	322	526	670	631	631	309	121	6
ALL OTHER, FEMALE													
1967	3,672	202	118	121	180	314	584	587	564	544	313	142	3
1966	3,683	227	119	118	192	312	568	582	591	521	307	141	5
1965	3,397	253	134	132	166	297	561	514	486	476	256	121	1
1964	3,436	276	138	134	155	314	550	559	512	423	238	132	5
1963	3,374	274	116	141	150	305	563	523	504	439	254	105	-
1962	3,225	247	132	123	165	310	468	490	495	436	245	113	1
1961	3,066	271	123	89	165	337	458	478	437	374	225	103	6
1960	3,177	262	136	112	153	340	500	501	447	387	234	101	4

- Quantity zero

Note: Figures by race exclude data for residents of New Jersey for 1962 and 1963.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290A for 1960-67 and separate unpublished tables for the black population and Accidents.

77/09/02 TABLE 290A. DEATHS FOR 60 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1960-67

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	TOTAL	UNDER 1 YEAR	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS	65-74 YEARS	75-84 YEARS	85+ YEARS	N.S.
ICD-7 Nos. E800-E962 Accidents													
BOTH SEXES													
1967	113,169	2,751	5,074	7,874	21,645	11,856	11,399	12,231	11,595	10,645	11,280	6,736	83
1966	113,563	3,096	5,411	7,958	21,030	11,591	11,543	12,402	11,620	10,706	11,480	6,653	73
1965	108,004	3,316	5,270	7,391	18,688	10,913	11,315	11,968	10,932	10,430	11,188	6,516	77
1964	103,843	3,406	5,184	7,306	17,190	10,739	11,062	11,336	10,536	10,288	10,692	6,025	79
1963	100,669	3,514	5,174	6,962	15,889	10,002	10,527	11,025	10,237	10,194	10,726	6,362	57
1962	97,139	3,653	5,052	6,751	14,557	9,667	10,288	10,599	9,736	10,149	10,419	6,193	75
1961	92,249	3,581	5,041	6,717	13,431	9,518	9,755	9,986	9,148	9,452	9,742	5,801	77
1960	93,806	3,831	5,119	6,836	13,457	9,802	9,858	10,201	9,184	9,689	9,938	5,805	86
MALE													
1967	77,879	1,552	3,046	5,498	17,290	9,605	8,752	9,092	8,330	6,585	5,557	2,504	68
1966	77,946	1,763	3,171	5,497	16,881	9,350	8,839	9,251	8,370	6,572	5,715	2,476	61
1965	74,062	1,887	3,166	5,144	15,069	8,831	8,655	8,890	7,858	6,508	5,551	2,445	58
1964	70,969	1,912	3,015	5,108	13,839	8,646	8,502	8,523	7,494	6,426	5,314	2,134	56
1963	68,639	1,949	3,026	4,847	12,801	8,118	8,117	8,305	7,391	6,355	5,263	2,417	50
1962	66,074	2,026	2,877	4,727	11,809	7,902	7,885	8,014	7,006	6,324	5,122	2,322	60
1961	63,337	2,037	2,915	4,760	11,033	7,751	7,601	7,641	6,650	5,943	4,796	2,147	63
1960	64,346	2,153	2,954	4,804	11,069	8,045	7,740	7,837	6,651	6,128	4,785	2,108	72
FEMALE													
1967	35,290	1,199	2,028	2,376	4,355	2,251	2,647	3,139	3,265	4,060	5,723	4,232	15
1966	35,617	1,333	2,240	2,461	4,149	2,241	2,704	3,151	3,250	4,134	5,765	4,177	12
1965	33,942	1,429	2,104	2,247	3,619	2,082	2,660	3,078	3,074	3,922	5,637	4,071	19
1964	32,874	1,494	2,169	2,198	3,351	2,093	2,560	2,813	3,042	3,862	5,378	3,891	23
1963	32,030	1,565	2,148	2,115	3,088	1,884	2,410	2,720	2,846	3,839	5,463	3,945	7
1962	31,065	1,627	2,175	2,024	2,748	1,765	2,403	2,585	2,730	3,825	5,297	3,871	15
1961	28,912	1,544	2,126	1,957	2,398	1,767	2,154	2,345	2,498	3,509	4,946	3,654	14
1960	29,460	1,678	2,165	2,032	2,388	1,757	2,118	2,364	2,533	3,561	5,153	3,697	14

Note: Figures by race exclude data for residents of New Jersey for 1962 and 1963.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290A for 1960-67 and separate unpublished tables for the black population and Accidents.

77/09/02 TABLE 290A. DEATHS FOR 60 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1960-67

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	TOTAL	UNDER 1 YEAR	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS	65-74 YEARS	75-84 YEARS	85+ YEARS	N.S.
ICD-7 Nos. E800-E962 Accidents													
WHITE, BOTH SEXES													
1967	96,775	1,870	3,738	6,268	18,921	9,708	9,320	10,351	10,025	9,497	10,574	6,446	57
1966	96,813	2,079	4,009	6,336	18,406	9,528	9,446	10,400	10,049	9,448	10,731	6,324	57
1965	92,534	2,267	3,898	5,893	16,413	9,003	9,241	10,145	9,478	9,333	10,581	6,226	56
1964	89,128	2,395	3,869	5,808	15,165	8,869	9,104	9,627	9,158	9,235	10,069	5,768	61
1963	84,001	2,385	3,765	5,361	13,616	8,028	8,516	9,153	8,621	8,809	9,775	5,936	36
1962	81,110	2,483	3,699	5,225	12,477	7,848	8,321	8,797	8,159	8,763	9,543	5,742	53
1961	79,294	2,569	3,797	5,400	11,768	7,905	8,051	8,494	7,908	8,553	9,211	5,581	57
1960	80,268	2,672	3,809	5,451	11,754	8,156	8,153	8,592	7,936	8,755	9,383	5,547	60
WHITE, MALE													
1967	66,033	1,071	2,275	4,404	15,082	7,870	7,141	7,641	7,142	5,814	5,161	2,386	46
1966	66,047	1,193	2,407	4,381	14,762	7,734	7,222	7,741	7,179	5,755	5,297	2,328	48
1965	63,024	1,343	2,389	4,156	13,229	7,297	7,058	7,482	6,761	5,790	5,185	2,291	43
1964	60,513	1,352	2,296	4,100	12,183	7,172	6,965	7,164	6,481	5,724	5,001	2,032	43
1963	57,090	1,362	2,244	3,790	10,947	6,581	6,533	6,887	6,212	5,481	4,778	2,243	32
1962	55,036	1,415	2,180	3,717	10,096	6,434	6,395	6,612	5,850	5,454	4,676	2,164	43
1961	54,292	1,467	2,276	3,889	9,699	6,504	6,292	6,466	5,742	5,363	4,507	2,039	48
1960	54,883	1,528	2,249	3,890	9,663	6,730	6,424	6,586	5,765	5,537	4,475	1,987	49
WHITE, FEMALE													
1967	30,742	799	1,463	1,864	3,839	1,838	2,179	2,710	2,883	3,683	5,413	4,060	11
1966	30,766	886	1,602	1,955	3,644	1,794	2,224	2,659	2,870	3,693	5,434	3,996	9
1965	29,510	924	1,509	1,737	3,184	1,706	2,183	2,663	2,717	3,543	5,396	3,935	13
1964	28,615	1,043	1,573	1,708	2,982	1,697	2,139	2,463	2,677	3,511	5,068	3,736	18
1963	26,911	1,023	1,521	1,571	2,669	1,447	1,983	2,266	2,409	3,328	4,997	3,693	4
1962	26,074	1,068	1,519	1,508	2,381	1,414	1,926	2,185	2,309	3,309	4,867	3,578	10
1961	25,002	1,102	1,521	1,511	2,069	1,401	1,759	2,028	2,166	3,190	4,704	3,542	9
1960	25,385	1,144	1,560	1,561	2,091	1,426	1,729	2,006	2,171	3,218	4,908	3,560	11

Note: Figures by race exclude data for residents of New Jersey for 1962 and 1963.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290A for 1960-67 and separate unpublished tables for the black population and Accidents.

77/09/02 TABLE 290A. DEATHS FOR 60 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1960-67

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	TOTAL	UNDER 1 YEAR	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS	65-74 YEARS	75-84 YEARS	85+ YEARS	N.S.
ICD-7 Nos. E800-E962 Accidents													
ALL OTHER, BOTH SEXES													
1967	16,394	881	1,336	1,606	2,724	2,148	2,079	1,880	1,570	1,148	706	290	26
1966	16,750	1,017	1,402	1,622	2,624	2,063	2,097	2,002	1,571	1,258	749	329	16
1965	15,470	1,049	1,372	1,498	2,275	1,910	2,074	1,823	1,454	1,097	607	290	21
1964	14,715	1,011	1,315	1,498	2,025	1,870	1,958	1,709	1,378	1,053	623	257	18
1963	14,086	1,023	1,288	1,431	1,936	1,742	1,754	1,606	1,335	1,055	623	273	20
1962	13,528	1,073	1,239	1,367	1,771	1,617	1,718	1,540	1,283	1,071	572	256	21
1961	12,955	1,012	1,244	1,317	1,663	1,613	1,704	1,492	1,240	899	531	220	20
1960	13,538	1,159	1,310	1,385	1,703	1,646	1,705	1,609	1,248	934	555	258	26
ALL OTHER, MALE													
1967	11,846	481	771	1,094	2,208	1,735	1,611	1,451	1,188	771	396	118	22
1966	11,899	570	764	1,116	2,119	1,616	1,617	1,510	1,191	817	418	148	13
1965	11,038	544	777	988	1,840	1,534	1,597	1,408	1,097	718	366	154	15
1964	10,456	560	719	1,008	1,656	1,474	1,537	1,359	1,013	702	313	102	13
1963	9,871	530	713	940	1,589	1,355	1,377	1,231	985	680	342	112	17
1962	9,429	559	638	897	1,458	1,297	1,317	1,203	951	677	309	106	17
1961	9,045	570	639	871	1,334	1,247	1,309	1,175	908	580	289	108	15
1960	9,463	625	705	914	1,406	1,315	1,316	1,251	886	591	310	121	23
ALL OTHER, FEMALE													
1967	4,548	400	565	512	516	413	468	429	382	377	310	172	4
1966	4,851	447	638	506	505	447	480	492	380	441	331	181	3
1965	4,432	505	595	510	435	376	477	415	357	379	241	136	6
1964	4,259	451	596	490	369	396	421	350	365	351	310	155	5
1963	4,215	493	575	491	347	387	377	375	350	375	281	161	3
1962	4,099	514	601	470	313	320	401	337	332	394	263	150	4
1961	3,910	442	605	446	329	366	395	317	332	319	242	112	5
1960	4,075	534	605	471	297	331	389	358	362	343	245	137	3

Note: Figures by race exclude data for residents of New Jersey for 1962 and 1963.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290A for 1960-67 and separate unpublished tables for the black population and Accidents.

Deaths for accidents by age and sex for black population: United States, 1960-61, 1964-67

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	TOTAL	UNDER 1 YEAR	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS	65-74 YEARS	75-84 YEARS	85+ YEARS	N.S.
ICD-7 Nos. E800-E962 Accidents													
STATISTICAL WORKSHEET													
BLACK, BOTH SEXES													
1967	14,828	828	1,222	1,497	2,379	1,887	1,901	1,734	1,426	1,047	621	262	24
1966	15,246	951	1,299	1,512	2,318	1,810	1,924	1,849	1,439	1,153	688	291	12
1965	14,047	994	1,278	1,395	2,015	1,675	1,879	1,663	1,327	1,007	546	248	20
1964	13,446	956	1,210	1,393	1,807	1,648	1,778	1,581	1,281	961	578	235	18
1963	---	---	---	---	---	---	---	---	---	---	---	---	---
1962	---	---	---	---	---	---	---	---	---	---	---	---	---
1961	11,762	948	1,157	1,221	1,422	1,419	1,552	1,365	1,134	843	491	191	19
1960	12,348	1,093	1,211	1,289	1,482	1,470	1,572	1,480	1,154	858	490	226	23
BLACK, MALE													
1967	10,720	454	704	1,025	1,950	1,531	1,490	1,337	1,073	694	334	107	21
1966	10,804	528	716	1,050	1,880	1,426	1,484	1,391	1,084	736	373	127	9
1965	9,993	515	721	921	1,642	1,352	1,460	1,289	992	646	318	122	15
1964	9,527	527	658	944	1,482	1,307	1,395	1,261	936	633	279	92	13
1963	---	---	---	---	---	---	---	---	---	---	---	---	---
1962	---	---	---	---	---	---	---	---	---	---	---	---	---
1961	8,175	543	589	808	1,147	1,101	1,200	1,065	815	541	262	90	14
1960	8,565	589	650	845	1,223	1,171	1,215	1,150	811	530	264	96	21
BLACK, FEMALE													
1967	4,108	374	518	472	429	356	411	397	353	353	287	155	3
1966	4,442	423	583	462	438	384	440	458	355	417	315	164	3
1965	4,054	479	557	474	373	323	419	374	335	361	228	126	5
1964	3,919	429	552	449	325	341	383	320	345	328	299	143	5
1963	---	---	---	---	---	---	---	---	---	---	---	---	---
1962	---	---	---	---	---	---	---	---	---	---	---	---	---
1961	3,587	405	568	413	275	318	352	300	319	302	229	101	5
1960	3,783	504	561	444	259	299	357	330	343	328	226	130	2

--- Data not available

Note: Figures by race exclude data for residents of New Jersey for 1962 and 1963.

Source: Unpublished trend table 290A for 1960-67 and separate unpublished tables for the black population and Accidents.

77/09/02 TABLE 290A. DEATHS FOR 60 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1960-67

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	TOTAL	UNDER 1 YEAR	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS	65-74 YEARS	75-84 YEARS	85+ YEARS	N.S.
Motor vehicle accidents (E810-E835)													
BOTH SEXES													
1967	52,924	321	1,746	3,845	15,646	7,217	5,770	5,819	5,083	4,285	2,698	471	23
1966	53,041	355	1,827	3,869	15,298	7,268	6,014	5,899	5,152	4,217	2,635	489	18
1965	49,163	326	1,733	3,526	13,395	6,610	5,985	5,592	4,917	4,077	2,547	433	22
1964	46,930	375	1,709	3,373	12,191	6,513	5,783	5,338	4,700	4,094	2,438	401	15
1963	43,564	321	1,670	3,063	11,123	5,980	5,376	5,073	4,433	3,786	2,327	399	13
1962	40,804	336	1,567	3,028	10,157	5,588	5,113	4,615	4,197	3,696	2,114	379	14
1961	38,091	307	1,584	2,802	9,088	5,472	4,740	4,393	3,874	3,467	1,995	351	18
1960	38,137	332	1,621	2,814	9,117	5,548	4,641	4,388	3,906	3,457	1,938	352	23
MALE													
1967	38,133	159	1,027	2,483	12,169	5,689	4,257	4,144	3,467	2,702	1,704	313	19
1966	38,316	191	1,043	2,551	11,916	5,708	4,462	4,204	3,497	2,650	1,742	337	15
1965	35,584	182	1,009	2,330	10,502	5,235	4,410	3,972	3,350	2,610	1,680	291	13
1964	33,794	188	988	2,253	9,501	5,143	4,238	3,814	3,152	2,668	1,598	243	8
1963	31,715	157	978	2,045	8,726	4,759	4,030	3,642	3,055	2,467	1,568	277	11
1962	29,639	178	877	2,032	8,028	4,466	3,753	3,316	2,874	2,428	1,414	261	12
1961	27,983	153	927	1,887	7,277	4,371	3,561	3,181	2,660	2,349	1,346	254	17
1960	28,084	180	949	1,874	7,290	4,481	3,514	3,187	2,686	2,342	1,337	227	17
FEMALE													
1967	14,791	162	719	1,362	3,477	1,528	1,513	1,675	1,616	1,583	994	158	4
1966	14,725	164	784	1,318	3,382	1,560	1,552	1,695	1,655	1,567	893	152	3
1965	13,579	144	724	1,196	2,893	1,375	1,575	1,620	1,567	1,467	867	142	9
1964	13,136	187	721	1,120	2,690	1,370	1,545	1,524	1,548	1,426	840	158	7
1963	11,849	164	692	1,018	2,397	1,221	1,346	1,431	1,378	1,319	759	122	2
1962	11,165	158	690	996	2,129	1,122	1,360	1,299	1,323	1,268	700	118	2
1961	10,108	154	657	915	1,811	1,101	1,179	1,212	1,214	1,118	649	97	1
1960	10,053	152	672	940	1,827	1,067	1,127	1,201	1,220	1,115	601	125	6

Note: Figures by race exclude data for residents of New Jersey for 1962 and 1963.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290A for 1960-67 and separate unpublished tables for the black population and Accidents.

77/09/02 TABLE 290A. DEATHS FOR 60 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1960-67

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	TOTAL	UNDER 1 YEAR	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS	65-74 YEARS	75-84 YEARS	85+ YEARS	N.S.
Motor vehicle accidents (E810-E835)													
WHITE, BOTH SEXES													
1967	46,284	265	1,394	3,149	14,065	6,114	4,909	5,037	4,479	3,881	2,532	446	13
1966	46,461	292	1,484	3,256	13,752	6,156	5,088	5,077	4,592	3,828	2,469	454	13
1965	43,202	286	1,387	2,974	12,077	5,587	5,104	4,868	4,356	3,716	2,425	406	16
1964	41,428	328	1,415	2,829	11,107	5,581	4,924	4,639	4,179	3,727	2,307	379	13
1963	37,549	281	1,356	2,522	9,856	4,981	4,501	4,332	3,834	3,364	2,149	363	10
1962	35,189	279	1,284	2,507	8,989	4,696	4,264	3,924	3,633	3,278	1,968	358	9
1961	33,612	271	1,338	2,373	8,205	4,678	4,067	3,854	3,430	3,178	1,873	330	15
1960	33,645	285	1,357	2,421	8,225	4,759	3,981	3,776	3,477	3,189	1,835	322	18
WHITE, MALE													
1967	33,203	135	807	2,047	10,927	4,814	3,599	3,572	3,013	2,393	1,590	295	11
1966	33,428	153	842	2,149	10,728	4,840	3,757	3,598	3,062	2,361	1,615	313	10
1965	31,144	163	791	1,978	9,480	4,422	3,746	3,432	2,914	2,352	1,585	270	11
1964	29,637	163	797	1,887	8,642	4,410	3,576	3,271	2,763	2,392	1,503	226	7
1963	27,186	137	778	1,680	7,707	3,973	3,334	3,102	2,614	2,162	1,443	248	8
1962	25,421	149	711	1,694	7,094	3,733	3,118	2,790	2,447	2,121	1,308	248	8
1961	24,618	133	779	1,605	6,589	3,747	3,045	2,762	2,322	2,127	1,261	234	14
1960	24,656	157	795	1,620	6,568	3,840	3,000	2,707	2,359	2,139	1,253	205	13
WHITE, FEMALE													
1967	13,081	130	587	1,102	3,138	1,300	1,310	1,465	1,466	1,488	942	151	2
1966	13,033	139	642	1,107	3,024	1,316	1,331	1,479	1,530	1,467	854	141	3
1965	12,058	123	596	996	2,597	1,165	1,358	1,436	1,442	1,364	840	136	5
1964	11,791	165	618	942	2,465	1,171	1,348	1,368	1,416	1,335	804	153	6
1963	10,363	144	578	842	2,149	1,008	1,167	1,230	1,220	1,202	706	115	2
1962	9,768	130	573	813	1,895	963	1,146	1,134	1,186	1,157	660	110	1
1961	8,994	138	559	768	1,616	931	1,022	1,092	1,108	1,051	612	96	1
1960	8,989	128	562	801	1,657	919	981	1,069	1,118	1,050	582	117	5

Note: Figures by race exclude data for residents of New Jersey for 1962 and 1963.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290A for 1960-67 and separate unpublished tables for the black population and Accidents.

77/09/02 TABLE 290A. DEATHS FOR 60 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1960-67

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	TOTAL	UNDER 1 YEAR	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS	65-74 YEARS	75-84 YEARS	85+ YEARS	N.S.
Motor vehicle accidents (E810-E835)													
ALL OTHER, BOTH SEXES													
1967	6,640	56	352	696	1,581	1,103	861	782	604	404	166	25	10
1966	6,580	63	343	613	1,546	1,112	926	822	560	389	166	35	5
1965	5,961	40	346	552	1,318	1,023	881	724	561	361	122	27	6
1964	5,502	47	294	544	1,084	932	859	699	521	367	131	22	2
1963	5,042	35	287	474	1,045	878	744	636	483	312	119	26	3
1962	4,711	52	260	455	946	776	732	591	459	322	103	10	5
1961	4,479	36	246	429	883	794	673	539	444	289	122	21	3
1960	4,492	47	264	393	892	789	660	612	429	268	103	30	5
ALL OTHER, MALE													
1967	4,930	24	220	436	1,242	875	658	572	454	309	114	18	8
1966	4,888	38	201	402	1,188	868	705	606	435	289	127	24	5
1965	4,440	19	218	352	1,022	813	664	540	436	258	95	21	2
1964	4,157	25	191	366	859	733	662	543	389	276	95	17	1
1963	3,826	18	181	318	850	687	590	472	364	235	87	21	3
1962	3,542	28	150	291	753	637	550	445	356	244	76	8	4
1961	3,365	20	148	282	688	624	516	419	338	222	85	20	3
1960	3,428	23	154	254	722	641	514	480	327	203	84	22	4
ALL OTHER, FEMALE													
1967	1,710	32	132	260	339	228	203	210	150	95	52	7	2
1966	1,692	25	142	211	358	244	221	216	125	100	39	11	-
1965	1,521	21	128	200	296	210	217	184	125	103	27	6	4
1964	1,345	22	103	178	225	199	197	156	132	91	36	5	1
1963	1,216	17	106	156	195	191	154	164	119	77	32	5	-
1962	1,169	24	110	164	193	139	182	146	103	78	27	2	1
1961	1,114	16	98	147	195	170	157	120	106	67	37	1	-
1960	1,064	24	110	139	170	148	146	132	102	65	19	8	1

- Quantity zero

Note: Figures by race exclude data for residents of New Jersey for 1962 and 1963.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290A for 1960-67 and separate unpublished tables for the black population and Accidents.

77/09/02 TABLE 290A. DEATHS FOR 60 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1960-67

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	TOTAL	UNDER 1 YEAR	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS	65-74 YEARS	75-84 YEARS	85+ YEARS	N.S.
Other accidents (E800-E802,E840-E962)													
BOTH SEXES													
1967	60,245	2,430	3,328	4,029	5,999	4,639	5,629	6,412	6,512	6,360	8,582	6,265	60
1966	60,522	2,741	3,584	4,089	5,732	4,323	5,529	6,503	6,468	6,489	8,845	6,164	55
1965	58,841	2,990	3,537	3,865	5,293	4,303	5,330	6,376	6,015	6,353	8,641	6,083	55
1964	56,913	3,031	3,475	3,933	4,999	4,226	5,279	5,998	5,836	6,194	8,254	5,624	64
1963	57,105	3,193	3,504	3,899	4,766	4,022	5,151	5,952	5,804	6,408	8,399	5,963	44
1962	56,335	3,317	3,485	3,723	4,400	4,079	5,175	5,984	5,539	6,453	8,305	5,814	61
1961	54,158	3,274	3,457	3,915	4,343	4,046	5,015	5,593	5,274	5,985	7,747	5,450	59
1960	55,669	3,499	3,498	4,022	4,340	4,254	5,217	5,813	5,278	6,232	8,000	5,453	63
MALE													
1967	39,746	1,393	2,019	3,015	5,121	3,916	4,495	4,948	4,863	3,883	3,853	2,191	49
1966	39,630	1,572	2,128	2,946	4,965	3,642	4,377	5,047	4,873	3,922	3,973	2,139	46
1965	38,478	1,705	2,157	2,814	4,567	3,596	4,245	4,918	4,508	3,898	3,871	2,154	45
1964	37,175	1,724	2,027	2,855	4,338	3,503	4,264	4,709	4,342	3,758	3,716	1,891	48
1963	36,924	1,792	2,048	2,802	4,075	3,359	4,087	4,663	4,336	3,888	3,695	2,140	39
1962	36,435	1,848	2,000	2,695	3,781	3,436	4,132	4,698	4,132	3,896	3,708	2,061	48
1961	35,354	1,884	1,988	2,873	3,756	3,380	4,040	4,460	3,990	3,594	3,450	1,893	46
1960	36,262	1,973	2,005	2,930	3,779	3,564	4,226	4,650	3,965	3,786	3,448	1,881	55
FEMALE													
1967	20,499	1,037	1,309	1,014	878	723	1,134	1,464	1,649	2,477	4,729	4,074	11
1966	20,892	1,169	1,456	1,143	767	681	1,152	1,456	1,595	2,567	4,872	4,025	9
1965	20,363	1,285	1,380	1,051	726	707	1,085	1,458	1,507	2,455	4,770	3,929	10
1964	19,738	1,307	1,448	1,078	661	723	1,015	1,289	1,494	2,436	4,538	3,733	16
1963	20,181	1,401	1,456	1,097	691	663	1,064	1,289	1,468	2,520	4,704	3,823	5
1962	19,900	1,469	1,485	1,028	619	643	1,043	1,286	1,407	2,557	4,597	3,753	13
1961	18,804	1,390	1,469	1,042	587	666	975	1,133	1,284	2,391	4,297	3,557	13
1960	19,407	1,526	1,493	1,092	561	690	991	1,163	1,313	2,446	4,552	3,572	8

Note: Figures by race exclude data for residents of New Jersey for 1962 and 1963.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290A for 1960-67 and separate unpublished tables for the black population and Accidents.

77/09/02 TABLE 290A. DEATHS FOR 60 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1960-67

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	TOTAL	UNDER 1 YEAR	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS	65-74 YEARS	75-84 YEARS	85+ YEARS	N.S.
Other accidents (E800-E802,E840-E962)													
WHITE, BOTH SEXES													
1967	50,491	1,605	2,344	3,119	4,856	3,594	4,411	5,314	5,546	5,616	8,042	6,000	44
1966	50,352	1,787	2,525	3,080	4,654	3,372	4,358	5,323	5,457	5,620	8,262	5,870	44
1965	49,332	1,981	2,511	2,919	4,336	3,416	4,137	5,277	5,122	5,617	8,156	5,820	40
1964	47,700	2,067	2,454	2,979	4,058	3,288	4,180	4,988	4,979	5,508	7,762	5,389	48
1963	46,452	2,104	2,409	2,839	3,760	3,047	4,015	4,821	4,787	5,445	7,626	5,573	26
1962	45,921	2,204	2,415	2,718	3,488	3,152	4,057	4,873	4,526	5,485	7,575	5,384	44
1961	45,682	2,298	2,459	3,027	3,563	3,227	3,984	4,640	4,478	5,375	7,338	5,251	42
1960	46,623	2,387	2,452	3,030	3,529	3,397	4,172	4,816	4,459	5,566	7,548	5,225	42
WHITE, MALE													
1967	32,830	936	1,468	2,357	4,155	3,056	3,542	4,069	4,129	3,421	3,571	2,091	35
1966	32,619	1,040	1,565	2,232	4,034	2,894	3,465	4,143	4,117	3,394	3,682	2,015	38
1965	31,880	1,180	1,598	2,178	3,749	2,875	3,312	4,050	3,847	3,438	3,600	2,021	32
1964	30,876	1,189	1,499	2,213	3,541	2,762	3,389	3,893	3,718	3,332	3,498	1,806	36
1963	29,904	1,225	1,466	2,110	3,240	2,608	3,199	3,785	3,598	3,319	3,335	1,995	24
1962	29,615	1,266	1,469	2,023	3,002	2,701	3,277	3,822	3,403	3,333	3,368	1,916	35
1961	29,674	1,334	1,497	2,284	3,110	2,757	3,247	3,704	3,420	3,236	3,246	1,805	34
1960	30,227	1,371	1,454	2,270	3,095	2,890	3,424	3,879	3,406	3,398	3,222	1,782	36
WHITE, FEMALE													
1967	17,661	669	876	762	701	538	869	1,245	1,417	2,195	4,471	3,909	9
1966	17,733	747	960	848	620	478	893	1,180	1,340	2,226	4,580	3,855	6
1965	17,452	801	913	741	587	541	825	1,227	1,275	2,179	4,556	3,799	8
1964	16,824	878	955	766	517	526	791	1,095	1,261	2,176	4,264	3,583	12
1963	16,548	879	943	729	520	439	816	1,036	1,189	2,126	4,291	3,578	2
1962	16,306	938	946	695	486	451	780	1,051	1,123	2,152	4,207	3,468	9
1961	16,008	964	962	743	453	470	737	936	1,058	2,139	4,092	3,446	8
1960	16,396	1,016	998	760	434	507	748	937	1,053	2,168	4,326	3,443	6

Note: Figures by race exclude data for residents of New Jersey for 1962 and 1963.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290A for 1960-67 and separate unpublished tables for the black population and Accidents.

77/09/02 TABLE 290A. DEATHS FOR 60 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1960-67

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	TOTAL	UNDER 1 YEAR	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS	65-74 YEARS	75-84 YEARS	85+ YEARS	N.S.
Other accidents (E800-E802,E840-E962)													
ALL OTHER, BOTH SEXES													
1967	9,754	825	984	910	1,143	1,045	1,218	1,098	966	744	540	265	16
1966	10,170	954	1,059	1,009	1,078	951	1,171	1,180	1,011	869	583	294	11
1965	9,509	1,009	1,026	946	957	887	1,193	1,099	893	736	485	263	15
1964	9,213	964	1,021	954	941	938	1,099	1,010	857	686	492	235	16
1963	9,044	988	1,001	957	891	864	1,010	970	852	743	504	247	17
1962	8,817	1,021	979	912	825	841	986	949	824	749	469	246	16
1961	8,476	976	998	888	780	819	1,031	953	796	610	409	199	17
1960	9,046	1,112	1,046	992	811	857	1,045	997	819	666	452	228	21
ALL OTHER, MALE													
1967	6,916	457	551	658	966	860	953	879	734	462	282	100	14
1966	7,011	532	563	714	931	748	912	904	756	528	291	124	8
1965	6,598	525	559	636	818	721	933	868	661	460	271	133	13
1964	6,299	535	528	642	797	741	875	816	624	426	218	85	12
1963	6,045	512	532	622	739	668	787	759	621	445	255	91	14
1962	5,887	531	488	606	705	660	767	758	595	433	233	98	13
1961	5,680	550	491	589	646	623	793	756	570	358	204	88	12
1960	6,035	602	551	660	684	674	802	771	559	388	226	99	19
ALL OTHER, FEMALE													
1967	2,838	368	433	252	177	185	265	219	232	282	258	165	2
1966	3,159	422	496	295	147	203	259	276	255	341	292	170	3
1965	2,911	484	467	310	139	166	260	231	232	276	214	130	2
1964	2,914	429	493	312	144	197	224	194	233	260	274	150	4
1963	2,999	476	469	335	152	196	223	211	231	298	249	156	3
1962	2,930	490	491	306	120	181	219	191	229	316	236	148	3
1961	2,796	426	507	299	134	196	238	197	226	252	205	111	5
1960	3,011	510	495	332	127	183	243	226	260	278	226	129	2

Note: Figures by race exclude data for residents of New Jersey for 1962 and 1963.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290A for 1960-67 and separate unpublished tables for the black population and Accidents.

77/09/02 TABLE 290A. DEATHS FOR 60 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1960-67

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	TOTAL	UNDER 1 YEAR	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS	65-74 YEARS	75-84 YEARS	85+ YEARS	N.S.
ICD-7 Nos. E963-E979 Suicide													
BOTH SEXES													
1967	21,325	121	2,244	2,819	3,961	4,412	3,935	2,315	1,246	267	5
1966	21,281	116	2,010	2,757	3,812	4,463	3,955	2,494	1,400	267	7
1965	21,507	104	1,876	2,721	4,064	4,554	4,040	2,525	1,353	262	8
1964	20,588	92	1,736	2,623	3,806	4,466	3,779	2,519	1,297	262	8
1963	20,825	104	1,663	2,621	3,924	4,530	3,864	2,534	1,329	246	10
1962	20,207	103	1,502	2,509	3,670	4,448	3,820	2,504	1,378	262	11
1961	18,999	76	1,258	2,306	3,489	4,232	3,665	2,458	1,267	241	7
1960	19,041	93	1,239	2,284	3,416	4,250	3,690	2,524	1,293	242	10
MALE													
1967	15,187	94	1,671	1,930	2,654	3,007	2,878	1,707	1,016	227	3
1966	15,416	94	1,517	1,895	2,531	3,087	2,986	1,900	1,162	238	6
1965	15,490	89	1,425	1,884	2,677	3,123	3,033	1,928	1,107	217	7
1964	15,092	82	1,326	1,841	2,536	3,172	2,891	1,905	1,097	236	6
1963	15,276	87	1,237	1,809	2,675	3,227	2,941	1,971	1,108	215	6
1962	15,062	90	1,115	1,736	2,565	3,206	2,951	2,003	1,170	218	8
1961	14,460	60	972	1,640	2,517	3,175	2,866	1,944	1,068	212	6
1960	14,539	77	976	1,643	2,472	3,192	2,871	2,028	1,062	208	10
FEMALE													
1967	6,138	27	573	889	1,307	1,405	1,057	608	230	40	2
1966	5,865	22	493	862	1,281	1,376	969	594	238	29	1
1965	6,017	15	451	837	1,387	1,431	1,007	597	246	45	1
1964	5,496	10	410	782	1,270	1,294	888	614	200	26	2
1963	5,549	17	426	812	1,249	1,303	923	563	221	31	4
1962	5,145	13	387	773	1,105	1,242	869	501	208	44	3
1961	4,539	16	286	666	972	1,057	799	514	199	29	1
1960	4,502	16	263	641	944	1,058	819	496	231	34	-

... Not applicable.

Note: Figures by race exclude data for residents of New Jersey for 1962 and 1963.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290A for 1960-67 and separate unpublished tables for the black population and Accidents.

77/09/02 TABLE 290A. DEATHS FOR 60 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1960-67

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	TOTAL	UNDER 1 YEAR	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS	65-74 YEARS	75-84 YEARS	85+ YEARS	N.S.
ICD-7 Nos. E963-E979 Suicide													
WHITE, BOTH SEXES													
1967	20,116	112	1,993	2,509	3,720	4,248	3,805	2,248	1,215	261	5
1966	20,100	108	1,784	2,444	3,628	4,276	3,817	2,418	1,364	254	7
1965	20,342	96	1,659	2,460	3,819	4,368	3,912	2,450	1,315	255	8
1964	19,545	87	1,560	2,358	3,591	4,308	3,657	2,449	1,268	261	6
1963	19,168	91	1,455	2,289	3,597	4,236	3,639	2,361	1,260	231	9
1962	18,677	98	1,307	2,235	3,368	4,170	3,564	2,366	1,309	249	11
1961	18,012	70	1,114	2,056	3,297	4,065	3,537	2,394	1,237	235	7
1960	18,121	90	1,140	2,074	3,206	4,092	3,550	2,455	1,269	235	10
WHITE, MALE													
1967	14,307	89	1,505	1,705	2,474	2,885	2,776	1,656	991	223	3
1966	14,527	90	1,350	1,671	2,391	2,941	2,877	1,842	1,132	227	6
1965	14,624	83	1,267	1,708	2,492	2,984	2,927	1,865	1,077	214	7
1964	14,300	78	1,186	1,642	2,379	3,052	2,800	1,850	1,074	235	4
1963	14,051	78	1,085	1,575	2,430	3,010	2,772	1,840	1,052	203	6
1962	13,933	86	974	1,543	2,347	3,012	2,752	1,890	1,113	208	8
1961	13,677	58	860	1,440	2,378	3,036	2,761	1,889	1,041	208	6
1960	13,825	75	900	1,483	2,311	3,067	2,755	1,976	1,045	203	10
WHITE, FEMALE													
1967	5,809	23	488	804	1,246	1,363	1,029	592	224	38	2
1966	5,573	18	434	773	1,237	1,335	940	576	232	27	1
1965	5,718	13	392	752	1,327	1,384	985	585	238	41	1
1964	5,245	9	374	716	1,212	1,256	857	599	194	26	2
1963	5,117	13	370	714	1,167	1,226	867	521	208	28	3
1962	4,744	12	333	692	1,021	1,158	812	476	196	41	3
1961	4,335	12	254	616	919	1,029	776	505	196	27	1
1960	4,296	15	240	591	895	1,025	795	479	224	32	-

... Not applicable.

- Quantity zero

Note: Figures by race exclude data for residents of New Jersey for 1962 and 1963.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290A for 1960-67 and separate unpublished tables for the black population and Accidents.

77/09/02 TABLE 290A. DEATHS FOR 60 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1960-67

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	TOTAL	UNDER 1 YEAR	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS	65-74 YEARS	75-84 YEARS	85+ YEARS	N.S.
ICD-7 Nos. E963-E979 Suicide													
ALL OTHER, BOTH SEXES													
1967	1,209	9	251	310	241	164	130	67	31	6	-
1966	1,181	8	226	313	184	187	138	76	36	13	-
1965	1,165	8	217	261	245	186	128	75	38	7	-
1964	1,043	5	176	265	215	158	122	70	29	1	2
1963	1,073	13	163	257	231	163	107	96	36	6	1
1962	981	4	162	215	203	156	130	77	29	5	-
1961	987	6	144	250	192	167	128	64	30	6	-
1960	920	3	99	210	210	158	140	69	24	7	-
ALL OTHER, MALE													
1967	880	5	166	225	180	122	102	51	25	4	-
1966	889	4	167	224	140	146	109	58	30	11	-
1965	866	6	158	176	185	139	106	63	30	3	-
1964	792	4	140	199	157	120	91	55	23	1	2
1963	829	9	120	189	177	134	90	73	33	4	-
1962	738	3	114	152	151	121	102	68	23	4	-
1961	783	2	112	200	139	139	105	55	27	4	-
1960	714	2	76	160	161	125	116	52	17	5	-
ALL OTHER, FEMALE													
1967	329	4	85	85	61	42	28	16	6	2	-
1966	292	4	59	89	44	41	29	18	6	2	-
1965	299	2	59	85	60	47	22	12	8	4	-
1964	251	1	36	66	58	38	31	15	6	-	-
1963	244	4	43	68	54	29	17	23	3	2	1
1962	243	1	48	63	52	35	28	9	6	1	-
1961	204	4	32	50	53	28	23	9	3	2	-
1960	206	1	23	50	49	33	24	17	7	2	-

... Not applicable.

- Quantity zero

Note: Figures by race exclude data for residents of New Jersey for 1962 and 1963.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290A for 1960-67 and separate unpublished tables for the black population and Accidents.

77/09/02 TABLE 290A. DEATHS FOR 60 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1960-67

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	TOTAL	UNDER 1 YEAR	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS	65-74 YEARS	75-84 YEARS	85+ YEARS	N.S.
ICD-7 Nos. E963-E979 Suicide													
BLACK, BOTH SEXES													
1967	982	6	200	255	207	138	95	58	20	3	-
1966	956	7	177	266	153	163	104	58	21	7	-
1965	958	7	182	218	201	165	101	55	25	4	-
1964	880	5	150	219	188	139	102	58	17	1	1
1963	---	---	---	---	---	---	---	---	---	---	---	---	---
1962	---	---	---	---	---	---	---	---	---	---	---	---	---
1961	781	5	115	196	153	148	101	45	16	2	-
1960	741	3	72	178	176	126	120	53	11	2	-
BLACK, MALE													
1967	712	2	127	186	154	103	74	47	17	2	-
1966	731	3	134	194	119	128	82	46	19	6	-
1965	732	5	134	151	161	121	86	50	21	3	-
1964	669	4	118	165	137	105	76	47	15	1	1
1963	---	---	---	---	---	---	---	---	---	---	---	---	---
1962	---	---	---	---	---	---	---	---	---	---	---	---	---
1961	622	1	92	157	112	123	83	38	15	1	-
1960	584	2	54	139	139	96	100	43	9	2	-
BLACK, FEMALE													
1967	270	4	73	69	53	35	21	11	3	1	-
1966	225	4	43	72	34	35	22	12	2	1	-
1965	226	2	48	67	40	44	15	5	4	1	-
1964	211	1	32	54	51	34	26	11	2	-	-
1963	---	---	---	---	---	---	---	---	---	---	---	---	---
1962	---	---	---	---	---	---	---	---	---	---	---	---	---
1961	159	4	23	39	41	25	18	7	1	1	-
1960	157	1	18	39	37	30	20	10	2	-	-

... Not applicable.

--- Data not available

- Quantity zero

Note: Figures by race exclude data for residents of New Jersey for 1962 and 1963.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290A for 1960-67 and separate unpublished tables for the black population and Accidents.

Deaths for suicides by age and sex for black population: United States, 1960-61, 1964-67

77/09/02 TABLE 290A. DEATHS FOR 60 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1960-67

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	TOTAL	UNDER 1 YEAR	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS	65-74 YEARS	75-84 YEARS	85+ YEARS	N.S.
ICD-7 Nos. E964-E985 Homicide													
BOTH SEXES													
1967	13,425	226	174	286	2,923	3,326	2,896	1,857	984	491	190	45	27
1966	11,606	211	189	250	2,397	2,744	2,546	1,711	931	399	164	35	29
1965	10,712	213	189	249	2,066	2,635	2,402	1,529	846	386	153	26	18
1964	9,814	219	200	236	1,833	2,329	2,261	1,449	770	329	134	32	22
1963	9,225	203	186	225	1,609	2,259	2,162	1,329	733	334	145	24	16
1962	9,013	214	151	206	1,556	2,198	2,097	1,341	725	344	131	33	17
1961	8,578	194	169	198	1,415	2,158	2,006	1,310	626	321	140	27	14
1960	8,464	199	116	173	1,422	2,208	1,948	1,269	656	312	109	22	30
MALE													
1967	10,236	122	86	152	2,275	2,645	2,223	1,434	782	370	114	21	12
1966	8,729	118	95	137	1,866	2,097	1,969	1,316	711	287	91	21	21
1965	8,148	123	102	150	1,609	2,046	1,842	1,187	661	294	107	16	11
1964	7,367	112	105	133	1,431	1,773	1,725	1,140	597	223	91	19	18
1963	6,921	104	100	124	1,246	1,752	1,649	1,023	581	230	89	14	9
1962	6,707	116	75	104	1,179	1,687	1,550	1,076	574	244	81	12	9
1961	6,346	103	85	117	1,081	1,625	1,476	1,011	493	250	81	16	8
1960	6,269	99	61	98	1,086	1,671	1,447	964	497	236	75	13	22
FEMALE													
1967	3,189	104	88	134	648	681	673	423	202	121	76	24	15
1966	2,877	93	94	113	531	647	577	395	220	112	73	14	8
1965	2,564	90	87	99	457	589	560	342	185	92	46	10	7
1964	2,447	107	95	103	402	556	536	309	173	106	43	13	4
1963	2,304	99	86	101	363	507	513	306	152	104	56	10	7
1962	2,306	98	76	102	377	511	547	265	151	100	50	21	8
1961	2,232	91	84	81	334	533	530	299	133	71	59	11	6
1960	2,195	100	55	75	336	537	501	305	159	76	34	9	8

Note: Figures by race exclude data for residents of New Jersey for 1962 and 1963.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290A for 1960-67 and separate unpublished tables for the black population and Accidents.

77/09/02 TABLE 290A. DEATHS FOR 60 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1960-67

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	TOTAL	UNDER 1 YEAR	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS	65-74 YEARS	75-84 YEARS	85+ YEARS	N.S.
---	-------	-----------------	--------------	---------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	--------------	------

ICD-7 Nos. E964-E985 Homicide

WHITE, BOTH SEXES

1967	6,212	151	116	170	1,182	1,320	1,252	922	561	325	160	38	15
1966	5,382	152	116	160	959	1,090	1,061	853	560	267	116	31	17
1965	5,039	144	126	158	887	1,005	1,035	782	515	244	114	19	10
1964	4,634	145	144	161	765	936	918	740	461	221	105	27	11
1963	4,299	138	139	144	663	896	894	650	409	229	110	20	7
1962	4,228	144	109	152	624	847	880	684	412	237	102	28	9
1961	4,174	130	116	140	595	870	892	702	375	218	107	19	10
1960	3,986	128	77	116	628	829	826	639	409	221	79	17	17

WHITE, MALE

1967	4,501	84	60	80	871	1,033	910	680	436	237	87	17	6
1966	3,785	82	59	86	681	796	800	612	403	174	61	18	13
1965	3,660	87	70	87	651	757	778	583	386	168	76	11	6
1964	3,261	79	80	97	565	670	656	549	341	135	64	16	9
1963	3,060	71	77	81	494	678	657	465	305	155	63	10	4
1962	2,937	75	52	72	432	641	615	512	310	154	60	10	4
1961	2,903	72	60	80	428	627	620	503	283	159	56	9	6
1960	2,832	68	42	65	464	621	585	460	297	159	50	9	12

WHITE, FEMALE

1967	1,711	67	56	90	311	287	342	242	125	88	73	21	9
1966	1,597	70	57	74	278	294	261	241	157	93	55	13	4
1965	1,379	57	56	71	236	248	257	199	129	76	38	8	4
1964	1,373	66	64	64	200	266	262	191	120	86	41	11	2
1963	1,239	67	62	63	169	218	237	185	104	74	47	10	3
1962	1,291	69	57	80	192	206	265	172	102	83	42	18	5
1961	1,271	58	56	60	167	243	272	199	92	59	51	10	4
1960	1,154	60	35	51	164	208	241	179	112	62	29	8	5

Note: Figures by race exclude data for residents of New Jersey for 1962 and 1963.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290A for 1960-67 and separate unpublished tables for the black population and Accidents.

77/09/02 TABLE 290A. DEATHS FOR 60 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1960-67

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	TOTAL	UNDER 1 YEAR	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS	65-74 YEARS	75-84 YEARS	85+ YEARS	N.S.
ICD-7 Nos. E964-E985 Homicide													
ALL OTHER, BOTH SEXES													
1967	7,213	75	58	116	1,741	2,006	1,644	935	423	166	30	7	12
1966	6,224	59	73	90	1,438	1,654	1,485	858	371	132	48	4	12
1965	5,673	69	63	91	1,179	1,630	1,367	747	331	142	39	7	8
1964	5,180	74	56	75	1,068	1,393	1,343	709	309	108	29	5	11
1963	4,733	57	45	73	909	1,320	1,231	649	307	100	29	4	9
1962	4,593	61	36	50	908	1,302	1,168	635	294	100	28	5	6
1961	4,404	64	53	58	820	1,288	1,114	608	251	103	33	8	4
1960	4,478	71	39	57	794	1,379	1,122	630	247	91	30	5	13
ALL OTHER, MALE													
1967	5,735	38	26	72	1,404	1,612	1,313	754	346	133	27	4	6
1966	4,944	36	36	51	1,185	1,301	1,169	704	308	113	30	3	8
1965	4,488	36	32	63	958	1,289	1,064	604	275	126	31	5	5
1964	4,106	33	25	36	866	1,103	1,069	591	256	88	27	3	9
1963	3,730	30	22	40	726	1,041	965	537	266	72	22	4	5
1962	3,633	34	20	30	731	1,011	900	546	251	85	20	2	3
1961	3,443	31	25	37	653	998	856	508	210	91	25	7	2
1960	3,437	31	19	33	622	1,050	862	504	200	77	25	4	10
ALL OTHER, FEMALE													
1967	1,478	37	32	44	337	394	331	181	77	33	3	3	6
1966	1,280	23	37	39	253	353	316	154	63	19	18	1	4
1965	1,185	33	31	28	221	341	303	143	56	16	8	2	3
1964	1,074	41	31	39	202	290	274	118	53	20	2	2	2
1963	1,003	27	23	33	183	279	266	112	41	28	7	-	4
1962	960	27	16	20	177	291	268	89	43	15	8	3	3
1961	961	33	28	21	167	290	258	100	41	12	8	1	2
1960	1,041	40	20	24	172	329	260	126	47	14	5	1	3

Note: Figures by race exclude data for residents of New Jersey for 1962 and 1963.

- Quantity zero

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290A for 1960-67 and separate unpublished tables for the black population and Accidents.

77/09/02 TABLE 290A. DEATHS FOR 60 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1960-67

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	TOTAL	UNDER 1 YEAR	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS	65-74 YEARS	75-84 YEARS	85+ YEARS	N.S.
---	-------	-----------------	--------------	---------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	--------------	------

ICD-7 Nos. E964-E985 Homicide

BLACK, BOTH SEXES

1967	7,037	73	53	113	1,696	1,955	1,610	924	410	155	29	7	12
1966	6,091	58	69	90	1,420	1,615	1,461	833	358	125	46	4	12
1965	5,523	65	61	85	1,155	1,589	1,336	731	315	135	36	7	8
1964	5,074	73	53	75	1,051	1,353	1,327	691	302	105	28	5	11
1963	---	---	---	---	---	---	---	---	---	---	---	---	---
1962	---	---	---	---	---	---	---	---	---	---	---	---	---
1961	4,308	63	52	56	805	1,264	1,094	589	242	100	32	7	4
1960	4,358	68	36	52	772	1,354	1,094	612	241	87	26	4	12

BLACK, MALE

1967	5,611	38	24	70	1,368	1,579	1,293	745	334	124	26	4	6
1966	4,847	36	34	51	1,172	1,275	1,151	686	296	106	29	3	8
1965	4,376	32	30	58	939	1,257	1,045	594	263	120	28	5	5
1964	4,026	32	25	36	853	1,069	1,058	577	251	87	26	3	9
1963	---	---	---	---	---	---	---	---	---	---	---	---	---
1962	---	---	---	---	---	---	---	---	---	---	---	---	---
1961	3,370	30	25	36	643	983	838	492	202	89	24	6	2
1960	3,345	29	18	30	605	1,030	842	488	196	73	22	3	9

BLACK, FEMALE

1967	1,426	35	29	43	328	376	317	179	76	31	3	3	6
1966	1,244	22	35	39	248	340	310	147	62	19	17	1	4
1965	1,147	33	31	27	216	332	291	137	52	15	8	2	3
1964	1,048	41	28	39	198	284	269	114	51	18	2	2	2
1963	---	---	---	---	---	---	---	---	---	---	---	---	---
1962	---	---	---	---	---	---	---	---	---	---	---	---	---
1961	938	33	27	20	162	281	256	97	40	11	8	1	2
1960	1,013	39	18	22	167	324	252	124	45	14	4	1	3

--- Data not available

Note: Figures by race exclude data for residents of New Jersey for 1962 and 1963.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290A for 1960-67 and separate unpublished tables for the black population and Accidents.

Deaths for homicides by age and sex for black population: United States, 1960-61, 1964-67