

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|---------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| ALL CAUSES | | | | | | | | | | | | |
| BOTH SEXES | | | | | | | | | | | | |
| 1978 | 868.0 | 1,381.4 | 67.9 | 33.2 | 115.0 | 131.7 | 238.7 | 610.4 | 1,387.1 | 3,015.9 | 6,709.7 | 15,479.6 |
| 1977 | 864.4 | 1,432.6 | 67.6 | 34.0 | 114.8 | 132.8 | 247.3 | 621.2 | 1,408.5 | 3,044.1 | 6,755.7 | 15,369.0 |
| 1976 | 877.6 | 1,549.4 | 68.8 | 34.2 | 111.7 | 133.2 | 253.9 | 635.3 | 1,452.3 | 3,118.1 | 6,952.0 | 16,058.6 |
| 1975 | 878.5 | 1,603.0 | 69.9 | 35.2 | 117.3 | 140.6 | 266.7 | 649.9 | 1,475.3 | 3,179.5 | 7,034.5 | 15,655.0 |
| 1974 | 906.7 | 1,721.9 | 73.3 | 37.8 | 120.3 | 144.5 | 278.6 | 675.3 | 1,533.0 | 3,318.6 | 7,377.1 | 16,901.1 |
| 1973 | 933.5 | 1,776.9 | 79.0 | 40.7 | 127.1 | 151.9 | 295.9 | 697.6 | 1,598.6 | 3,432.2 | 7,708.5 | 17,678.9 |
| 1972 | 938.4 | 1,820.4 | 80.4 | 40.6 | 126.9 | 152.6 | 302.0 | 710.5 | 1,621.7 | 3,521.2 | 7,805.0 | 17,543.1 |
| 1971 | 932.0 | 1,887.8 | 82.3 | 40.9 | 126.1 | 155.7 | 307.4 | 710.1 | 1,617.4 | 3,482.9 | 7,774.3 | 17,567.0 |
| 1970 | 945.3 | 2,142.4 | 84.5 | 41.3 | 127.7 | 157.4 | 314.5 | 730.0 | 1,658.8 | 3,582.7 | 8,004.4 | 16,344.9 |
| 1969 | 954.4 | 2,199.6 | 88.0 | 42.6 | 129.2 | 158.7 | 321.4 | 732.5 | 1,661.3 | 3,632.7 | 8,097.5 | 18,821.8 |
| 1968 | 967.9 | 2,265.7 | 89.6 | 43.0 | 123.7 | 157.2 | 319.8 | 751.3 | 1,704.4 | 3,724.0 | 8,293.5 | 19,582.7 |
| MALE | | | | | | | | | | | | |
| 1978 | 977.5 | 1,532.8 | 76.7 | 40.5 | 169.4 | 186.8 | 311.6 | 802.1 | 1,884.5 | 4,181.5 | 8,835.6 | 18,211.8 |
| 1977 | 978.9 | 1,599.7 | 75.2 | 41.8 | 169.2 | 188.5 | 321.2 | 818.1 | 1,915.7 | 4,224.3 | 8,898.1 | 18,071.6 |
| 1976 | 993.8 | 1,715.0 | 76.9 | 41.9 | 164.8 | 187.0 | 329.8 | 837.7 | 1,981.6 | 4,331.4 | 9,077.8 | 18,666.5 |
| 1975 | 1,002.0 | 1,786.2 | 76.7 | 43.3 | 174.1 | 198.9 | 345.4 | 859.3 | 2,018.9 | 4,409.4 | 9,154.4 | 18,134.7 |
| 1974 | 1,031.5 | 1,931.5 | 81.5 | 46.4 | 178.9 | 201.8 | 357.4 | 893.1 | 2,099.6 | 4,580.2 | 9,484.0 | 19,276.6 |
| 1973 | 1,065.2 | 2,001.6 | 88.2 | 49.6 | 187.8 | 211.8 | 379.4 | 918.5 | 2,196.2 | 4,727.4 | 9,867.9 | 20,067.1 |
| 1972 | 1,074.9 | 2,055.1 | 89.4 | 48.6 | 186.7 | 209.9 | 385.9 | 941.4 | 2,237.9 | 4,814.0 | 9,949.3 | 19,821.1 |
| 1971 | 1,069.4 | 2,121.4 | 90.6 | 49.3 | 185.4 | 213.0 | 392.0 | 932.1 | 2,221.2 | 4,758.8 | 9,820.3 | 19,901.4 |
| 1970 | 1,090.3 | 2,410.0 | 93.2 | 50.5 | 188.5 | 215.3 | 402.6 | 958.5 | 2,282.7 | 4,873.8 | 10,010.2 | 17,821.5 |
| 1969 | 1,103.0 | 2,484.5 | 95.5 | 51.5 | 191.6 | 217.2 | 410.7 | 967.9 | 2,291.0 | 4,934.9 | 10,068.3 | 20,875.6 |
| 1968 | 1,119.4 | 2,562.6 | 98.2 | 52.3 | 182.8 | 214.0 | 406.0 | 988.7 | 2,344.1 | 5,049.1 | 10,215.2 | 21,732.0 |

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| ALL CAUSES | | | | | | | | | | | | |
| FEMALE | | | | | | | | | | | | |
| 1978 | 764.5 | 1,222.7 | 58.8 | 25.6 | 59.6 | 77.5 | 168.6 | 431.7 | 948.2 | 2,125.3 | 5,444.1 | 14,245.0 |
| 1977 | 756.0 | 1,257.0 | 59.7 | 25.8 | 59.6 | 78.2 | 176.4 | 437.9 | 959.8 | 2,141.1 | 5,473.5 | 14,127.6 |
| 1976 | 767.6 | 1,375.2 | 60.4 | 26.1 | 57.7 | 80.5 | 181.1 | 447.1 | 983.8 | 2,188.8 | 5,668.4 | 14,833.6 |
| 1975 | 761.4 | 1,410.9 | 62.7 | 26.8 | 59.8 | 83.6 | 191.4 | 455.2 | 993.7 | 2,237.3 | 5,743.1 | 14,454.5 |
| 1974 | 788.3 | 1,502.1 | 64.6 | 28.8 | 61.1 | 88.7 | 203.2 | 472.9 | 1,030.6 | 2,351.2 | 6,079.1 | 15,724.7 |
| 1973 | 808.3 | 1,542.2 | 69.4 | 31.3 | 65.9 | 93.3 | 216.1 | 492.4 | 1,067.6 | 2,437.6 | 6,359.4 | 16,477.0 |
| 1972 | 808.7 | 1,575.3 | 71.0 | 32.2 | 66.9 | 96.9 | 222.0 | 496.1 | 1,072.5 | 2,524.0 | 6,438.1 | 16,377.3 |
| 1971 | 801.4 | 1,642.9 | 73.8 | 32.3 | 67.2 | 100.3 | 226.9 | 503.7 | 1,077.5 | 2,494.8 | 6,447.3 | 16,348.5 |
| 1970 | 807.8 | 1,863.7 | 75.4 | 31.8 | 68.1 | 101.6 | 231.1 | 517.2 | 1,098.9 | 2,579.7 | 6,677.6 | 15,518.0 |
| 1969 | 813.5 | 1,903.8 | 80.2 | 33.4 | 68.4 | 102.6 | 237.0 | 513.0 | 1,094.6 | 2,615.2 | 6,775.4 | 17,710.1 |
| 1968 | 824.1 | 1,956.1 | 80.7 | 33.4 | 66.4 | 102.8 | 238.2 | 528.9 | 1,126.2 | 2,680.7 | 6,978.2 | 18,425.0 |
| WHITE, BOTH SEXES | | | | | | | | | | | | |
| 1978 | 880.2 | 1,169.2 | 61.2 | 31.5 | 111.6 | 114.9 | 206.3 | 559.2 | 1,318.5 | 2,945.8 | 6,734.8 | 15,761.1 |
| 1977 | 874.6 | 1,221.4 | 61.1 | 32.4 | 110.8 | 114.9 | 213.6 | 567.6 | 1,338.2 | 2,969.5 | 6,790.6 | 15,642.8 |
| 1976 | 887.7 | 1,320.1 | 62.8 | 32.7 | 106.0 | 113.7 | 218.7 | 580.6 | 1,383.5 | 3,039.8 | 7,007.5 | 16,372.5 |
| 1975 | 886.9 | 1,377.0 | 63.3 | 33.4 | 110.0 | 119.2 | 229.4 | 596.8 | 1,405.4 | 3,094.7 | 7,096.2 | 15,970.0 |
| 1974 | 913.8 | 1,498.2 | 66.4 | 36.1 | 111.7 | 121.2 | 237.9 | 617.3 | 1,457.9 | 3,219.7 | 7,457.4 | 17,251.1 |
| 1973 | 937.7 | 1,545.9 | 70.9 | 38.4 | 117.5 | 126.9 | 252.2 | 633.4 | 1,518.3 | 3,325.2 | 7,778.9 | 18,047.8 |
| 1972 | 941.7 | 1,572.4 | 72.9 | 38.9 | 115.5 | 125.6 | 255.2 | 646.1 | 1,542.0 | 3,421.8 | 7,878.0 | 17,952.6 |
| 1971 | 933.9 | 1,643.3 | 74.0 | 38.6 | 113.3 | 127.8 | 261.1 | 648.8 | 1,538.9 | 3,387.7 | 7,833.0 | 18,004.0 |
| 1970 | 946.3 | 1,869.7 | 75.1 | 39.1 | 115.8 | 129.9 | 267.0 | 666.2 | 1,577.1 | 3,490.1 | 8,043.3 | 16,889.7 |
| 1969 | 953.1 | 1,900.9 | 77.6 | 40.0 | 117.7 | 130.8 | 272.4 | 667.1 | 1,572.8 | 3,535.3 | 8,135.7 | 19,345.5 |
| 1968 | 964.9 | 1,968.4 | 78.3 | 40.3 | 113.9 | 128.5 | 270.4 | 684.9 | 1,608.5 | 3,620.4 | 8,331.1 | 20,083.5 |

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|---------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| ALL CAUSES | | | | | | | | | | | | |
| WHITE, MALE | | | | | | | | | | | | |
| 1978 | 982.7 | 1,302.9 | 69.9 | 38.3 | 164.8 | 162.3 | 267.0 | 735.2 | 1,800.9 | 4,120.8 | 8,923.3 | 18,587.9 |
| 1977 | 983.0 | 1,377.4 | 68.1 | 39.7 | 164.0 | 162.0 | 275.7 | 748.9 | 1,831.8 | 4,165.2 | 9,006.7 | 18,420.8 |
| 1976 | 997.3 | 1,473.1 | 70.5 | 40.0 | 156.8 | 158.5 | 282.8 | 767.0 | 1,900.2 | 4,266.5 | 9,208.3 | 19,108.2 |
| 1975 | 1,004.1 | 1,551.9 | 70.1 | 40.9 | 163.6 | 166.4 | 295.1 | 791.0 | 1,940.9 | 4,343.0 | 9,274.7 | 18,562.2 |
| 1974 | 1,032.4 | 1,695.3 | 74.5 | 44.2 | 166.6 | 168.2 | 304.8 | 821.1 | 2,016.0 | 4,501.2 | 9,621.5 | 19,733.4 |
| 1973 | 1,063.8 | 1,756.5 | 79.2 | 46.6 | 174.5 | 175.6 | 324.0 | 840.3 | 2,110.1 | 4,645.8 | 9,992.0 | 20,561.2 |
| 1972 | 1,072.0 | 1,796.6 | 81.9 | 46.7 | 169.5 | 171.2 | 325.1 | 861.7 | 2,154.4 | 4,739.2 | 10,073.9 | 20,351.6 |
| 1971 | 1,066.5 | 1,860.8 | 81.6 | 46.2 | 166.4 | 173.6 | 334.7 | 860.6 | 2,141.1 | 4,691.8 | 9,940.1 | 20,508.6 |
| 1970 | 1,086.7 | 2,113.2 | 83.6 | 48.0 | 170.8 | 176.6 | 343.5 | 882.9 | 2,202.6 | 4,810.1 | 10,098.8 | 18,551.7 |
| 1969 | 1,097.1 | 2,167.0 | 84.0 | 48.2 | 174.8 | 178.9 | 349.3 | 891.3 | 2,204.7 | 4,861.7 | 10,147.2 | 21,593.7 |
| 1968 | 1,112.7 | 2,251.4 | 85.9 | 48.8 | 168.5 | 175.4 | 346.3 | 912.1 | 2,249.5 | 4,970.8 | 10,300.3 | 22,371.2 |
| WHITE, FEMALE | | | | | | | | | | | | |
| 1978 | 782.7 | 1,027.5 | 52.0 | 24.4 | 57.0 | 67.5 | 146.7 | 392.8 | 890.0 | 2,049.3 | 5,447.1 | 14,501.5 |
| 1977 | 771.3 | 1,056.7 | 53.7 | 24.8 | 56.3 | 67.8 | 152.8 | 396.4 | 899.0 | 2,057.2 | 5,482.8 | 14,384.9 |
| 1976 | 783.1 | 1,159.0 | 54.8 | 25.0 | 54.0 | 68.9 | 156.2 | 405.1 | 923.4 | 2,102.9 | 5,697.0 | 15,121.7 |
| 1975 | 775.1 | 1,191.7 | 56.2 | 25.5 | 55.3 | 72.1 | 165.3 | 414.0 | 928.6 | 2,142.0 | 5,785.4 | 14,749.7 |
| 1974 | 800.5 | 1,290.4 | 57.8 | 27.6 | 55.7 | 74.5 | 172.8 | 425.8 | 960.7 | 2,242.1 | 6,135.9 | 16,040.4 |
| 1973 | 817.3 | 1,324.8 | 62.1 | 29.9 | 59.6 | 78.5 | 182.3 | 439.4 | 990.4 | 2,315.8 | 6,408.3 | 16,820.7 |
| 1972 | 817.3 | 1,337.1 | 63.4 | 30.7 | 60.8 | 80.4 | 187.4 | 444.0 | 994.3 | 2,410.3 | 6,492.7 | 16,745.6 |
| 1971 | 807.4 | 1,414.2 | 66.0 | 30.6 | 60.1 | 82.7 | 189.9 | 450.2 | 999.2 | 2,382.9 | 6,484.1 | 16,721.1 |
| 1970 | 812.6 | 1,614.6 | 66.1 | 29.9 | 61.6 | 84.1 | 193.3 | 462.9 | 1,014.9 | 2,470.7 | 6,698.7 | 15,980.2 |
| 1969 | 816.0 | 1,623.1 | 71.0 | 31.4 | 61.7 | 84.0 | 198.5 | 456.5 | 1,002.9 | 2,503.4 | 6,797.6 | 18,178.5 |
| 1968 | 824.0 | 1,671.5 | 70.4 | 31.3 | 60.5 | 82.8 | 197.6 | 470.5 | 1,028.5 | 2,562.8 | 6,996.4 | 18,875.5 |

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|---------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| ALL CAUSES | | | | | | | | | | | | |
| ALL OTHER, BOTH SEXES | | | | | | | | | | | | |
| 1978 | 790.0 | 2,405.1 | 99.9 | 41.5 | 133.6 | 239.4 | 458.3 | 1,004.5 | 2,005.4 | 3,670.7 | 6,452.1 | 12,250.0 |
| 1977 | 798.6 | 2,455.6 | 98.5 | 41.6 | 137.6 | 250.9 | 474.5 | 1,044.1 | 2,050.3 | 3,751.1 | 6,396.6 | 12,234.4 |
| 1976 | 812.1 | 2,670.8 | 97.5 | 41.8 | 144.2 | 264.3 | 492.8 | 1,076.5 | 2,087.3 | 3,868.6 | 6,380.9 | 12,375.8 |
| 1975 | 823.1 | 2,733.7 | 101.4 | 45.1 | 159.6 | 287.2 | 523.6 | 1,087.0 | 2,121.9 | 4,004.4 | 6,374.5 | 12,040.4 |
| 1974 | 859.1 | 2,831.8 | 107.7 | 47.0 | 171.3 | 307.0 | 563.2 | 1,160.1 | 2,237.0 | 4,287.3 | 6,500.2 | 12,828.1 |
| 1973 | 904.5 | 2,916.7 | 120.9 | 52.8 | 184.9 | 328.2 | 605.6 | 1,240.8 | 2,359.6 | 4,489.1 | 6,900.4 | 13,380.3 |
| 1972 | 915.5 | 3,090.9 | 120.0 | 49.7 | 197.4 | 344.7 | 639.2 | 1,264.5 | 2,381.0 | 4,505.2 | 6,956.8 | 12,817.9 |
| 1971 | 918.4 | 3,193.5 | 126.8 | 54.2 | 208.0 | 352.4 | 647.3 | 1,245.2 | 2,367.5 | 4,422.3 | 7,062.6 | 12,590.0 |
| 1970 | 938.4 | 3,597.1 | 134.1 | 53.7 | 203.6 | 348.7 | 664.3 | 1,290.9 | 2,431.1 | 4,488.4 | 7,511.1 | 10,750.3 |
| 1969 | 963.3 | 3,884.2 | 142.7 | 57.7 | 205.1 | 358.1 | 693.8 | 1,320.2 | 2,515.0 | 4,602.7 | 7,619.6 | 12,889.0 |
| 1968 | 989.6 | 3,870.9 | 150.2 | 59.1 | 190.2 | 363.0 | 700.0 | 1,352.2 | 2,634.5 | 4,761.9 | 7,805.9 | 13,990.2 |
| ALL OTHER, MALE | | | | | | | | | | | | |
| 1978 | 943.2 | 2,652.2 | 109.2 | 51.8 | 195.8 | 357.3 | 639.5 | 1,357.3 | 2,667.3 | 4,749.5 | 7,978.2 | 14,360.3 |
| 1977 | 951.7 | 2,692.3 | 109.2 | 52.4 | 199.5 | 377.3 | 653.8 | 1,402.2 | 2,710.2 | 4,773.4 | 7,875.1 | 14,510.7 |
| 1976 | 969.8 | 2,921.8 | 108.3 | 51.8 | 212.1 | 393.3 | 676.4 | 1,445.1 | 2,756.7 | 4,934.5 | 7,877.8 | 14,241.8 |
| 1975 | 987.6 | 2,967.3 | 109.4 | 56.2 | 237.6 | 438.4 | 720.9 | 1,453.8 | 2,762.7 | 5,040.0 | 7,964.9 | 13,859.3 |
| 1974 | 1,025.1 | 3,125.9 | 117.3 | 58.3 | 254.3 | 454.2 | 756.8 | 1,527.9 | 2,901.9 | 5,341.7 | 8,069.9 | 14,572.0 |
| 1973 | 1,075.3 | 3,222.6 | 135.4 | 66.5 | 271.3 | 487.5 | 805.5 | 1,613.5 | 3,033.6 | 5,506.8 | 8,525.4 | 15,022.9 |
| 1972 | 1,095.2 | 3,402.2 | 129.7 | 59.2 | 296.6 | 506.0 | 860.2 | 1,656.4 | 3,047.8 | 5,531.5 | 8,610.1 | 14,459.6 |
| 1971 | 1,090.5 | 3,534.5 | 139.3 | 66.5 | 310.3 | 511.2 | 848.7 | 1,581.8 | 2,998.7 | 5,398.1 | 8,534.8 | 13,919.1 |
| 1970 | 1,115.9 | 4,020.0 | 144.7 | 65.0 | 304.6 | 504.1 | 873.5 | 1,646.1 | 3,046.6 | 5,474.4 | 8,981.0 | 11,405.2 |
| 1969 | 1,145.5 | 4,310.9 | 157.5 | 70.8 | 305.7 | 511.9 | 914.5 | 1,680.7 | 3,128.8 | 5,630.8 | 9,123.4 | 14,102.3 |
| 1968 | 1,169.1 | 4,260.8 | 165.4 | 72.6 | 281.7 | 512.6 | 899.5 | 1,704.9 | 3,265.0 | 5,809.6 | 9,194.6 | 15,512.2 |
| ALL OTHER, FEMALE | | | | | | | | | | | | |
| 1978 | 651.0 | 2,151.8 | 90.4 | 31.2 | 73.7 | 137.6 | 306.2 | 713.2 | 1,456.0 | 2,835.5 | 5,425.2 | 11,239.4 |
| 1977 | 659.6 | 2,213.0 | 87.6 | 30.7 | 78.0 | 142.0 | 324.3 | 747.2 | 1,499.6 | 2,948.0 | 5,372.3 | 11,008.7 |
| 1976 | 668.6 | 2,425.3 | 86.5 | 31.8 | 78.8 | 153.2 | 339.1 | 768.0 | 1,526.9 | 3,025.4 | 5,357.7 | 11,328.6 |
| 1975 | 673.1 | 2,493.8 | 93.3 | 33.8 | 84.9 | 157.0 | 358.3 | 776.9 | 1,580.7 | 3,180.8 | 5,269.2 | 10,854.8 |
| 1974 | 707.6 | 2,541.9 | 97.9 | 35.6 | 92.0 | 180.5 | 401.2 | 848.5 | 1,671.2 | 3,442.8 | 5,417.5 | 11,802.4 |
| 1973 | 748.4 | 2,603.8 | 106.4 | 39.0 | 102.7 | 191.3 | 438.8 | 921.9 | 1,785.2 | 3,667.4 | 5,777.4 | 12,382.3 |
| 1972 | 750.9 | 2,773.9 | 110.3 | 40.2 | 103.0 | 206.1 | 454.2 | 927.3 | 1,805.9 | 3,678.9 | 5,788.0 | 11,802.6 |
| 1971 | 760.7 | 2,845.2 | 114.2 | 41.8 | 111.4 | 216.6 | 479.8 | 952.3 | 1,816.0 | 3,624.3 | 5,974.3 | 11,734.2 |
| 1970 | 775.3 | 3,169.4 | 123.3 | 42.3 | 108.8 | 215.7 | 490.5 | 979.4 | 1,886.9 | 3,675.6 | 6,392.6 | 10,288.9 |
| 1969 | 796.4 | 3,454.3 | 127.9 | 44.5 | 111.6 | 227.7 | 510.2 | 1,004.3 | 1,970.9 | 3,756.5 | 6,444.1 | 12,067.7 |
| 1968 | 824.8 | 3,461.8 | 134.9 | 45.5 | 105.1 | 236.7 | 532.3 | 1,041.3 | 2,068.5 | 3,892.9 | 6,757.7 | 12,967.2 |

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|---------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| ALL CAUSES | | | | | | | | | | | | |
| BLACK, BOTH SEXES | | | | | | | | | | | | |
| 1978 | 855.1 | 2,596.1 | 104.6 | 43.2 | 136.1 | 262.5 | 508.2 | 1,105.3 | 2,159.9 | 3,876.0 | 6,569.2 | 12,550.7 |
| 1977 | 864.0 | 2,652.2 | 103.2 | 43.1 | 139.8 | 275.5 | 526.4 | 1,146.7 | 2,200.3 | 3,970.1 | 6,497.5 | 12,566.4 |
| 1976 | 875.0 | 2,888.4 | 103.0 | 43.0 | 147.6 | 289.0 | 541.4 | 1,176.6 | 2,228.7 | 4,090.3 | 6,482.3 | 12,752.9 |
| 1975 | 882.5 | 2,940.6 | 106.2 | 45.7 | 164.6 | 313.9 | 574.3 | 1,181.3 | 2,252.7 | 4,219.7 | 6,456.0 | 12,355.4 |
| 1974 | 916.5 | 3,038.9 | 112.1 | 48.3 | 177.5 | 335.3 | 614.6 | 1,251.4 | 2,362.0 | 4,514.2 | 6,586.5 | 13,168.6 |
| 1973 | 960.6 | 3,126.0 | 126.2 | 54.8 | 189.8 | 354.2 | 659.1 | 1,333.0 | 2,488.0 | 4,719.3 | 6,979.8 | 13,680.7 |
| 1972 | 969.2 | 3,299.4 | 122.9 | 50.7 | 203.4 | 370.6 | 696.7 | 1,354.4 | 2,498.3 | 4,727.1 | 7,092.0 | 13,154.5 |
| 1971 | 968.5 | 3,392.3 | 130.5 | 55.4 | 215.6 | 379.6 | 699.3 | 1,325.8 | 2,481.7 | 4,619.1 | 7,220.4 | 12,898.1 |
| 1970 | 999.3 | 3,835.6 | 140.0 | 55.5 | 212.4 | 381.2 | 724.9 | 1,383.8 | 2,570.6 | 4,719.4 | 7,860.7 | 11,300.5 |
| 1969 | 1,011.3 | 4,131.7 | 146.1 | 58.7 | 211.4 | 378.8 | 743.7 | 1,395.4 | 2,635.7 | 4,789.3 | 7,892.6 | 13,336.1 |
| 1968 | 1,035.7 | 4,063.2 | 151.7 | 60.4 | 195.2 | 384.2 | 749.1 | 1,423.8 | 2,757.1 | 4,948.3 | 8,037.0 | 14,289.1 |
| BLACK, MALE | | | | | | | | | | | | |
| 1978 | 1,016.8 | 2,862.5 | 114.0 | 53.9 | 199.4 | 392.5 | 713.5 | 1,499.5 | 2,888.7 | 5,076.8 | 8,072.0 | 14,500.0 |
| 1977 | 1,026.0 | 2,914.7 | 114.8 | 54.4 | 203.1 | 416.3 | 730.8 | 1,543.8 | 2,929.7 | 5,117.0 | 8,024.7 | 14,730.6 |
| 1976 | 1,041.6 | 3,168.1 | 113.6 | 53.4 | 217.9 | 431.8 | 752.4 | 1,583.3 | 2,961.9 | 5,280.3 | 7,979.1 | 14,744.7 |
| 1975 | 1,055.4 | 3,197.0 | 115.2 | 57.1 | 246.2 | 484.0 | 795.4 | 1,584.5 | 2,951.1 | 5,378.1 | 8,034.7 | 14,345.7 |
| 1974 | 1,090.0 | 3,350.4 | 122.7 | 60.1 | 263.9 | 499.1 | 830.7 | 1,652.0 | 3,091.1 | 5,684.7 | 8,183.7 | 15,058.1 |
| 1973 | 1,137.7 | 3,451.9 | 141.4 | 69.9 | 279.8 | 529.1 | 880.9 | 1,735.7 | 3,229.0 | 5,840.6 | 8,609.8 | 15,639.0 |
| 1972 | 1,155.9 | 3,614.1 | 133.8 | 60.0 | 307.7 | 547.2 | 942.4 | 1,781.6 | 3,234.7 | 5,858.8 | 8,740.1 | 15,115.0 |
| 1971 | 1,146.8 | 3,739.2 | 143.8 | 68.0 | 322.5 | 558.4 | 922.6 | 1,690.6 | 3,172.2 | 5,674.5 | 8,761.3 | 14,432.5 |
| 1970 | 1,186.6 | 4,298.9 | 150.5 | 67.1 | 320.6 | 559.5 | 956.6 | 1,777.5 | 3,256.9 | 5,803.2 | 9,454.9 | 12,222.3 |
| 1969 | 1,197.4 | 4,594.8 | 160.5 | 72.1 | 315.6 | 544.6 | 980.6 | 1,785.0 | 3,313.6 | 5,888.6 | 9,452.1 | 14,959.5 |
| 1968 | 1,219.4 | 4,470.6 | 167.2 | 74.4 | 289.8 | 548.2 | 965.5 | 1,805.7 | 3,450.3 | 6,063.8 | 9,520.6 | 16,225.7 |
| BLACK, FEMALE | | | | | | | | | | | | |
| 1978 | 709.5 | 2,324.3 | 95.0 | 32.4 | 75.7 | 150.0 | 337.9 | 780.9 | 1,564.8 | 2,980.3 | 5,565.9 | 11,546.5 |
| 1977 | 718.0 | 2,384.2 | 91.3 | 31.6 | 79.3 | 154.0 | 356.6 | 817.9 | 1,605.1 | 3,109.1 | 5,502.0 | 11,438.3 |
| 1976 | 724.5 | 2,604.9 | 92.2 | 32.5 | 80.4 | 165.8 | 366.5 | 838.8 | 1,622.1 | 3,185.7 | 5,478.8 | 11,701.1 |
| 1975 | 726.1 | 2,679.6 | 97.0 | 34.3 | 87.0 | 167.6 | 390.8 | 843.5 | 1,673.4 | 3,341.2 | 5,387.5 | 11,265.5 |
| 1974 | 759.4 | 2,720.4 | 101.3 | 36.6 | 95.4 | 194.7 | 434.4 | 915.2 | 1,757.3 | 3,623.3 | 5,506.1 | 12,126.9 |
| 1973 | 800.0 | 2,780.8 | 110.8 | 39.5 | 104.6 | 204.8 | 474.2 | 990.5 | 1,871.5 | 3,852.5 | 5,880.1 | 12,580.8 |
| 1972 | 799.4 | 2,965.4 | 111.9 | 41.4 | 104.6 | 220.3 | 492.7 | 989.9 | 1,880.4 | 3,853.2 | 5,907.1 | 12,034.3 |
| 1971 | 806.4 | 3,038.4 | 116.9 | 42.7 | 115.0 | 228.1 | 514.5 | 1,012.2 | 1,897.3 | 3,780.1 | 6,113.9 | 11,982.1 |
| 1970 | 829.2 | 3,368.8 | 129.4 | 43.8 | 111.9 | 231.0 | 533.0 | 1,043.9 | 1,986.2 | 3,860.9 | 6,691.5 | 10,706.6 |
| 1969 | 842.2 | 3,666.7 | 131.7 | 45.2 | 115.0 | 240.1 | 547.0 | 1,058.1 | 2,058.0 | 3,914.9 | 6,710.6 | 12,335.0 |
| 1968 | 868.4 | 3,650.6 | 136.1 | 46.4 | 107.7 | 247.3 | 567.8 | 1,091.5 | 2,157.6 | 4,041.9 | 6,934.2 | 13,333.9 |

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| BACILLARY DYSENTERY AND AMEBIASIS (004,006) | | | | | | | | | | | | |
| BOTH SEXES | | | | | | | | | | | | |
| 1978 | 0.0 | * | * | * | * | * | * | * | * | * | * | * |
| 1977 | 0.0 | * | * | * | * | * | * | * | * | * | * | * |
| 1976 | 0.0 | * | * | * | * | * | * | * | * | * | * | * |
| 1975 | 0.0 | * | * | * | * | * | * | * | * | * | * | * |
| 1974 | 0.0 | * | * | * | * | * | * | * | * | * | * | * |
| 1973 | 0.0 | * | * | * | * | * | * | * | * | * | * | * |
| 1972 | 0.0 | * | * | * | * | * | * | * | * | * | * | * |
| 1971 | 0.0 | * | * | * | * | * | * | * | * | * | * | * |
| 1970 | 0.0 | * | * | * | * | * | * | * | * | * | * | * |
| 1969 | 0.0 | 0.6 | * | * | * | * | * | * | * | * | * | * |
| 1968 | 0.1 | * | * | * | * | * | * | * | 0.1 | * | * | * |
| MALE | | | | | | | | | | | | |
| 1978 | 0.0 | * | * | * | * | * | * | * | * | * | * | * |
| 1977 | 0.0 | * | * | * | * | * | * | * | * | * | * | * |
| 1976 | 0.0 | * | * | * | * | * | * | * | * | * | * | * |
| 1975 | 0.0 | * | * | * | * | * | * | * | * | * | * | * |
| 1974 | 0.0 | * | * | * | * | * | * | * | * | * | * | * |
| 1973 | 0.0 | * | * | * | * | * | * | * | * | * | * | * |
| 1972 | 0.1 | * | * | * | * | * | * | * | * | * | * | * |
| 1971 | 0.1 | * | * | * | * | * | * | * | * | * | * | * |
| 1970 | 0.0 | * | * | * | * | * | * | * | * | * | * | * |
| 1969 | 0.1 | * | * | * | * | * | * | * | * | * | * | * |
| 1968 | 0.1 | * | * | * | * | * | * | * | * | * | * | * |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| BACILLARY DYSENTERY AND AMEBIASIS (004,006) | | | | | | | | | | | | |
| FEMALE | | | | | | | | | | | | |
| 1978 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1977 | 0.0 | * | * | * | * | * | * | * | * | * | * | * |
| 1976 | 0.0 | * | * | * | * | * | * | * | * | * | * | * |
| 1975 | 0.0 | * | * | * | * | * | * | * | * | * | * | * |
| 1974 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1973 | 0.0 | * | * | * | * | * | * | * | * | * | * | * |
| 1972 | 0.0 | * | * | * | * | * | * | * | * | * | * | * |
| 1971 | 0.0 | * | * | * | * | * | * | * | * | * | * | * |
| 1970 | 0.0 | * | * | * | * | * | * | * | * | * | * | * |
| 1969 | 0.0 | * | * | * | * | * | * | * | * | * | * | * |
| 1968 | 0.0 | * | * | * | * | * | * | * | * | * | * | * |
| WHITE, BOTH SEXES | | | | | | | | | | | | |
| 1978 | 0.0 | * | * | * | * | * | * | * | * | * | * | * |
| 1977 | 0.0 | * | * | * | * | * | * | * | * | * | * | * |
| 1976 | 0.0 | * | * | * | * | * | * | * | * | * | * | * |
| 1975 | 0.0 | * | * | * | * | * | * | * | * | * | * | * |
| 1974 | 0.0 | * | * | * | * | * | * | * | * | * | * | * |
| 1973 | 0.0 | * | * | * | * | * | * | * | * | * | * | * |
| 1972 | 0.0 | * | * | * | * | * | * | * | * | * | * | * |
| 1971 | 0.0 | * | * | * | * | * | * | * | * | * | * | * |
| 1970 | 0.0 | * | * | * | * | * | * | * | * | * | * | * |
| 1969 | 0.0 | * | * | * | * | * | * | * | * | * | * | * |
| 1968 | 0.0 | * | * | * | * | * | * | * | * | * | * | * |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| BACILLARY DYSENTERY AND AMEBIASIS (004,006) | | | | | | | | | | | | |
| WHITE, MALE | | | | | | | | | | | | |
| 1978 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1977 | 0.0 | * | * | * | * | * | * | * | * | * | * | * |
| 1976 | 0.0 | * | * | * | * | * | * | * | * | * | * | * |
| 1975 | 0.0 | * | * | * | * | * | * | * | * | * | * | * |
| 1974 | 0.0 | * | * | * | * | * | * | * | * | * | * | * |
| 1973 | 0.0 | * | * | * | * | * | * | * | * | * | * | * |
| 1972 | 0.0 | * | * | * | * | * | * | * | * | * | * | * |
| 1971 | 0.0 | * | * | * | * | * | * | * | * | * | * | * |
| 1970 | 0.0 | * | * | * | * | * | * | * | * | * | * | * |
| 1969 | 0.0 | * | * | * | * | * | * | * | * | * | * | * |
| 1968 | 0.1 | * | * | * | * | * | * | * | * | * | * | * |
| WHITE, FEMALE | | | | | | | | | | | | |
| 1978 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1977 | 0.0 | * | * | * | * | * | * | * | * | * | * | * |
| 1976 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1975 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1974 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1973 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1972 | 0.0 | * | * | * | * | * | * | * | * | * | * | * |
| 1971 | 0.0 | * | * | * | * | * | * | * | * | * | * | * |
| 1970 | 0.0 | * | * | * | * | * | * | * | * | * | * | * |
| 1969 | 0.0 | * | * | * | * | * | * | * | * | * | * | * |
| 1968 | 0.0 | * | * | * | * | * | * | * | * | * | * | * |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| BACILLARY DYSENTERY AND AMEBIASIS (004,006) | | | | | | | | | | | | |
| ALL OTHER, BOTH SEXES | | | | | | | | | | | | |
| 1978 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1977 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1976 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1975 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1974 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1973 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1972 | 0.1 | * | * | * | * | * | * | * | * | * | * | * |
| 1971 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1970 | 0.1 | * | * | * | * | * | * | * | * | * | * | * |
| 1969 | 0.1 | * | * | * | * | * | * | * | * | * | * | * |
| 1968 | 0.2 | * | * | * | * | * | * | * | * | * | * | * |
| ALL OTHER, MALE | | | | | | | | | | | | |
| 1978 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1977 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1976 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1975 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1974 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1973 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1972 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1971 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1970 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1969 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1968 | 0.2 | * | * | * | * | * | * | * | * | * | * | * |
| ALL OTHER, FEMALE | | | | | | | | | | | | |
| 1978 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1977 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1976 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1975 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1974 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1973 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1972 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1971 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1970 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1969 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1968 | * | * | * | * | * | * | * | * | * | * | * | * |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| BACILLARY DYSENTERY AND AMEBIASIS (004,006) | | | | | | | | | | | | |
| BLACK, BOTH SEXES | | | | | | | | | | | | |
| 1978 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1977 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1976 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1975 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1974 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1973 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1972 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1971 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1970 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1969 | 0.1 | * | * | * | * | * | * | * | * | * | * | * |
| 1968 | 0.1 | * | * | * | * | * | * | * | * | * | * | * |
| BLACK, MALE | | | | | | | | | | | | |
| 1978 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1977 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1976 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1975 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1974 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1973 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1972 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1971 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1970 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1969 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1968 | * | * | * | * | * | * | * | * | * | * | * | * |
| BLACK, FEMALE | | | | | | | | | | | | |
| 1978 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1977 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1976 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1975 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1974 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1973 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1972 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1971 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1970 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1969 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1968 | * | * | * | * | * | * | * | * | * | * | * | * |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|--|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| ENTERITIS AND OTHER DIARRHEAL DISEASES (008,009) | | | | | | | | | | | | |
| BOTH SEXES | | | | | | | | | | | | |
| 1978 | 0.8 | 22.0 | 0.4 | * | * | 0.1 | 0.1 | 0.2 | 0.5 | 1.5 | 4.2 | 13.6 |
| 1977 | 0.8 | 19.7 | 0.6 | 0.1 | 0.1 | 0.1 | 0.1 | 0.3 | 0.6 | 1.4 | 4.2 | 15.8 |
| 1976 | 0.9 | 23.0 | 0.7 | 0.1 | * | 0.1 | 0.1 | 0.3 | 0.6 | 1.5 | 5.0 | 15.2 |
| 1975 | 0.9 | 22.6 | 0.7 | 0.1 | 0.1 | 0.1 | 0.1 | 0.4 | 0.6 | 1.8 | 5.2 | 15.2 |
| 1974 | 1.1 | 22.3 | 0.8 | 0.1 | 0.1 | 0.1 | 0.3 | 0.4 | 0.8 | 2.1 | 6.7 | 18.6 |
| 1973 | 1.1 | 22.1 | 1.0 | 0.1 | 0.1 | 0.1 | 0.2 | 0.4 | 0.9 | 2.1 | 7.1 | 19.7 |
| 1972 | 1.1 | 21.3 | 0.8 | 0.1 | 0.1 | 0.1 | 0.3 | 0.5 | 0.9 | 2.2 | 6.6 | 19.2 |
| 1971 | 1.2 | 21.4 | 1.1 | 0.1 | 0.1 | 0.1 | 0.2 | 0.5 | 1.1 | 2.6 | 6.9 | 20.0 |
| 1970 | 1.3 | 24.4 | 1.4 | 0.2 | 0.1 | 0.2 | 0.2 | 0.5 | 1.0 | 2.5 | 7.0 | 21.0 |
| 1969 | 1.3 | 26.8 | 1.4 | 0.2 | 0.1 | 0.1 | 0.3 | 0.5 | 1.0 | 2.5 | 7.1 | 21.2 |
| 1968 | 1.5 | 30.7 | 1.7 | 0.2 | 0.1 | 0.2 | 0.2 | 0.5 | 1.2 | 2.7 | 8.1 | 24.1 |
| MALE | | | | | | | | | | | | |
| 1978 | 0.8 | 24.3 | 0.5 | * | * | * | * | 0.2 | 0.6 | 1.7 | 4.6 | 13.5 |
| 1977 | 0.8 | 21.8 | 0.6 | * | * | * | * | 0.3 | 0.6 | 1.3 | 4.3 | 15.2 |
| 1976 | 0.8 | 24.5 | 0.6 | * | * | * | * | 0.3 | 0.5 | 1.5 | 4.9 | 13.5 |
| 1975 | 0.9 | 24.7 | 0.8 | * | * | 0.1 | * | 0.4 | 0.7 | 1.8 | 5.6 | 13.6 |
| 1974 | 1.0 | 24.9 | 0.8 | 0.1 | * | * | 0.2 | 0.3 | 0.8 | 2.3 | 6.5 | 17.9 |
| 1973 | 1.0 | 24.6 | 1.2 | * | 0.1 | * | 0.2 | 0.4 | 0.8 | 2.1 | 7.0 | 19.0 |
| 1972 | 1.0 | 23.3 | 1.1 | * | 0.1 | * | 0.3 | 0.5 | 0.9 | 2.3 | 6.0 | 19.2 |
| 1971 | 1.2 | 23.1 | 1.2 | 0.1 | * | 0.2 | * | 0.5 | 1.2 | 2.9 | 6.4 | 21.8 |
| 1970 | 1.3 | 28.2 | 1.3 | 0.2 | * | * | 0.2 | 0.5 | 1.1 | 2.6 | 7.1 | 17.9 |
| 1969 | 1.3 | 30.1 | 1.4 | 0.2 | * | * | 0.2 | 0.4 | 1.2 | 2.4 | 6.9 | 20.0 |
| 1968 | 1.5 | 32.9 | 1.8 | 0.2 | 0.1 | 0.2 | 0.2 | 0.5 | 1.4 | 2.9 | 7.8 | 23.6 |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|--|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| ENTERITIS AND OTHER DIARRHEAL DISEASES (008,009) | | | | | | | | | | | | |
| FEMALE | | | | | | | | | | | | |
| 1978 | 0.8 | 19.7 | 0.3 | * | * | * | * | 0.2 | 0.5 | 1.3 | 4.0 | 13.6 |
| 1977 | 0.9 | 17.4 | 0.6 | * | * | * | 0.2 | 0.3 | 0.7 | 1.6 | 4.1 | 16.0 |
| 1976 | 1.0 | 21.5 | 0.7 | * | * | * | * | 0.2 | 0.6 | 1.4 | 5.1 | 16.0 |
| 1975 | 1.0 | 20.5 | 0.6 | * | * | 0.1 | * | 0.3 | 0.5 | 1.8 | 5.0 | 16.0 |
| 1974 | 1.1 | 19.5 | 0.8 | * | * | * | 0.3 | 0.4 | 0.9 | 2.0 | 6.7 | 18.9 |
| 1973 | 1.2 | 19.4 | 0.8 | 0.1 | 0.1 | * | 0.2 | 0.4 | 1.0 | 2.1 | 7.2 | 20.1 |
| 1972 | 1.1 | 19.2 | 0.6 | * | * | * | 0.3 | 0.6 | 0.9 | 2.2 | 6.9 | 19.2 |
| 1971 | 1.2 | 19.7 | 1.1 | 0.1 | * | * | 0.3 | 0.4 | 1.1 | 2.3 | 7.1 | 19.0 |
| 1970 | 1.3 | 20.4 | 1.4 | 0.1 | * | 0.2 | 0.2 | 0.4 | 0.8 | 2.4 | 7.0 | 22.7 |
| 1969 | 1.3 | 23.5 | 1.5 | 0.1 | * | * | 0.3 | 0.6 | 0.9 | 2.5 | 7.2 | 21.8 |
| 1968 | 1.5 | 28.3 | 1.5 | 0.2 | 0.1 | * | 0.2 | 0.5 | 1.0 | 2.6 | 8.3 | 24.4 |
| WHITE, BOTH SEXES | | | | | | | | | | | | |
| 1978 | 0.7 | 13.7 | 0.3 | * | * | 0.1 | 0.1 | 0.2 | 0.5 | 1.5 | 4.1 | 13.8 |
| 1977 | 0.7 | 12.8 | 0.6 | * | 0.1 | 0.1 | 0.1 | 0.3 | 0.6 | 1.4 | 4.2 | 15.8 |
| 1976 | 0.8 | 14.9 | 0.6 | 0.1 | * | * | * | 0.2 | 0.6 | 1.4 | 5.0 | 15.5 |
| 1975 | 0.8 | 15.1 | 0.6 | 0.1 | * | 0.1 | 0.1 | 0.4 | 0.5 | 1.7 | 5.4 | 15.0 |
| 1974 | 1.0 | 15.5 | 0.7 | 0.1 | 0.1 | 0.1 | 0.2 | 0.3 | 0.8 | 2.0 | 6.7 | 19.5 |
| 1973 | 1.0 | 14.9 | 0.8 | 0.1 | 0.1 | 0.1 | 0.2 | 0.4 | 0.9 | 2.0 | 7.1 | 19.7 |
| 1972 | 0.9 | 12.2 | 0.6 | 0.1 | 0.1 | 0.1 | 0.2 | 0.5 | 1.0 | 2.1 | 6.4 | 19.5 |
| 1971 | 1.0 | 12.7 | 1.0 | 0.1 | * | 0.1 | 0.2 | 0.4 | 1.0 | 2.5 | 6.8 | 19.4 |
| 1970 | 1.1 | 14.9 | 1.2 | 0.1 | 0.1 | 0.2 | 0.2 | 0.4 | 0.9 | 2.4 | 7.0 | 21.4 |
| 1969 | 1.0 | 14.8 | 1.2 | 0.1 | * | * | 0.2 | 0.4 | 0.9 | 2.2 | 7.0 | 20.9 |
| 1968 | 1.2 | 17.2 | 1.3 | 0.2 | 0.1 | 0.1 | 0.2 | 0.5 | 1.1 | 2.6 | 8.1 | 24.4 |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|--|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| ENTERITIS AND OTHER DIARRHEAL DISEASES (008,009) | | | | | | | | | | | | |
| WHITE, MALE | | | | | | | | | | | | |
| 1978 | 0.7 | 15.1 | 0.4 | * | * | * | * | * | 0.5 | 1.7 | 4.5 | 14.1 |
| 1977 | 0.7 | 14.4 | 0.5 | * | * | * | * | 0.3 | 0.5 | 1.4 | 4.2 | 15.2 |
| 1976 | 0.7 | 16.2 | 0.5 | * | * | * | * | 0.3 | 0.6 | 1.5 | 4.8 | 14.3 |
| 1975 | 0.8 | 15.8 | 0.7 | * | * | * | * | 0.4 | 0.6 | 1.6 | 5.8 | 12.8 |
| 1974 | 0.9 | 17.7 | 0.8 | * | * | * | * | 0.3 | 0.7 | 2.2 | 6.6 | 18.8 |
| 1973 | 0.9 | 17.0 | 1.0 | * | * | * | * | 0.3 | 0.8 | 2.1 | 6.7 | 19.6 |
| 1972 | 0.9 | 14.1 | 0.8 | * | * | * | 0.2 | 0.4 | 1.0 | 2.2 | 5.9 | 19.8 |
| 1971 | 1.0 | 13.9 | 1.0 | 0.1 | * | * | * | 0.4 | 1.0 | 2.9 | 6.3 | 20.5 |
| 1970 | 1.0 | 16.9 | 1.1 | 0.2 | * | * | * | 0.4 | 1.0 | 2.5 | 7.0 | 18.5 |
| 1969 | 1.0 | 15.9 | 1.2 | 0.2 | * | * | 0.2 | 0.4 | 1.1 | 2.2 | 7.0 | 18.3 |
| 1968 | 1.2 | 18.2 | 1.4 | 0.2 | 0.2 | * | * | 0.5 | 1.3 | 2.9 | 7.8 | 24.1 |
| WHITE, FEMALE | | | | | | | | | | | | |
| 1978 | 0.7 | 12.2 | * | * | * | * | * | * | 0.5 | 1.3 | 3.9 | 13.6 |
| 1977 | 0.8 | 11.0 | 0.6 | * | * | * | 0.2 | 0.3 | 0.7 | 1.5 | 4.1 | 16.1 |
| 1976 | 0.8 | 13.6 | 0.6 | * | * | * | * | 0.2 | 0.5 | 1.4 | 5.1 | 16.1 |
| 1975 | 0.9 | 14.3 | 0.5 | * | * | * | * | 0.3 | 0.4 | 1.7 | 5.1 | 16.1 |
| 1974 | 1.0 | 13.2 | 0.6 | * | * | * | 0.3 | 0.3 | 0.8 | 1.9 | 6.8 | 19.9 |
| 1973 | 1.1 | 12.7 | 0.7 | 0.1 | * | * | * | 0.4 | 0.9 | 2.0 | 7.3 | 19.8 |
| 1972 | 1.0 | 10.2 | 0.4 | * | * | * | 0.2 | 0.6 | 1.0 | 2.1 | 6.7 | 19.3 |
| 1971 | 1.1 | 11.4 | 1.0 | * | * | * | 0.3 | 0.4 | 1.0 | 2.2 | 7.2 | 18.8 |
| 1970 | 1.1 | 12.7 | 1.2 | 0.1 | * | * | 0.2 | 0.4 | 0.8 | 2.3 | 7.0 | 22.9 |
| 1969 | 1.1 | 13.6 | 1.2 | * | * | * | 0.3 | 0.4 | 0.8 | 2.2 | 7.0 | 22.2 |
| 1968 | 1.2 | 16.2 | 1.2 | 0.2 | * | * | 0.2 | 0.5 | 0.9 | 2.4 | 8.4 | 24.6 |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|--|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| ENTERITIS AND OTHER DIARRHEAL DISEASES (008,009) | | | | | | | | | | | | |
| ALL OTHER, BOTH SEXES | | | | | | | | | | | | |
| 1978 | 1.7 | 62.2 | 1.0 | * | * | * | * | * | 1.2 | 1.8 | 5.5 | * |
| 1977 | 1.5 | 53.1 | * | * | * | * | * | * | * | 1.6 | 4.4 | 15.0 |
| 1976 | 1.7 | 62.9 | 1.2 | * | * | * | * | * | * | 2.2 | 5.5 | * |
| 1975 | 1.7 | 60.4 | 1.3 | * | * | * | * | * | 1.2 | 2.6 | 3.7 | 17.1 |
| 1974 | 1.7 | 55.8 | 1.4 | * | * | * | * | * | 1.5 | 2.9 | 6.1 | * |
| 1973 | 1.9 | 57.6 | 1.7 | * | * | * | * | 0.9 | 1.1 | 3.1 | 7.2 | 19.7 |
| 1972 | 2.2 | 67.7 | 1.9 | * | * | * | * | * | * | 2.9 | 8.1 | 16.3 |
| 1971 | 2.4 | 68.3 | 1.8 | * | * | * | * | 0.9 | 2.0 | 2.8 | 7.2 | 26.7 |
| 1970 | 2.5 | 75.2 | 2.4 | * | * | * | * | * | 1.7 | 3.1 | 7.4 | 17.2 |
| 1969 | 3.2 | 94.7 | 2.9 | * | * | * | * | 1.1 | 2.2 | 5.1 | 8.3 | 24.8 |
| 1968 | 3.4 | 103.2 | 3.8 | 0.3 | * | * | * | * | 1.8 | 4.1 | 7.7 | 20.6 |
| ALL OTHER, MALE | | | | | | | | | | | | |
| 1978 | 1.9 | 68.9 | * | * | * | * | * | * | * | * | * | * |
| 1977 | 1.6 | 58.1 | * | * | * | * | * | * | * | * | * | * |
| 1976 | 1.8 | 66.2 | * | * | * | * | * | * | * | * | * | * |
| 1975 | 2.0 | 69.5 | * | * | * | * | * | * | * | * | * | * |
| 1974 | 1.8 | 61.0 | * | * | * | * | * | * | * | * | * | * |
| 1973 | 2.1 | 62.8 | 2.1 | * | * | * | * | * | * | * | 9.8 | * |
| 1972 | 2.3 | 71.1 | 2.3 | * | * | * | * | * | * | * | * | * |
| 1971 | 2.7 | 72.8 | 1.9 | * | * | * | * | * | 2.6 | * | * | * |
| 1970 | 3.0 | 89.3 | 2.5 | * | * | * | * | * | * | * | * | * |
| 1969 | 3.5 | 111.6 | 2.6 | * | * | * | * | * | * | 4.3 | * | * |
| 1968 | 3.8 | 114.0 | 4.4 | * | * | * | * | * | * | * | * | * |
| ALL OTHER, FEMALE | | | | | | | | | | | | |
| 1978 | 1.5 | 55.3 | * | * | * | * | * | * | * | * | 5.4 | * |
| 1977 | 1.4 | 48.0 | * | * | * | * | * | * | * | * | * | * |
| 1976 | 1.6 | 59.8 | * | * | * | * | * | * | * | * | 5.3 | * |
| 1975 | 1.4 | 51.0 | * | * | * | * | * | * | * | * | * | * |
| 1974 | 1.7 | 50.6 | * | * | * | * | * | * | * | * | 6.0 | * |
| 1973 | 1.8 | 52.3 | * | * | * | * | * | * | * | 3.7 | * | * |
| 1972 | 2.0 | 64.2 | * | * | * | * | * | * | * | * | 9.3 | * |
| 1971 | 2.1 | 63.7 | * | * | * | * | * | * | * | * | * | * |
| 1970 | 2.1 | 61.1 | 2.4 | * | * | * | * | * | * | * | * | * |
| 1969 | 2.9 | 77.7 | 3.2 | * | * | * | * | 1.6 | * | 5.7 | 9.7 | * |
| 1968 | 3.0 | 92.0 | 3.3 | * | * | * | * | * | * | 4.8 | * | * |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|--|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| ENTERITIS AND OTHER DIARRHEAL DISEASES (008,009) | | | | | | | | | | | | |
| BLACK, BOTH SEXES | | | | | | | | | | | | |
| 1978 | 1.8 | 67.2 | * | * | * | * | * | * | 1.2 | 1.9 | 5.4 | * |
| 1977 | 1.6 | 57.3 | * | * | * | * | * | * | * | 1.6 | 3.7 | * |
| 1976 | 1.8 | 67.0 | 1.2 | * | * | * | * | * | * | 2.2 | 5.1 | * |
| 1975 | 1.8 | 62.7 | 1.0 | * | * | * | * | * | 1.3 | 2.8 | * | 19.2 |
| 1974 | 1.8 | 57.0 | 1.4 | * | * | * | * | * | 1.5 | 3.0 | 5.6 | * |
| 1973 | 2.0 | 61.8 | 1.9 | * | * | * | * | 0.9 | 1.2 | 2.8 | 7.0 | 19.3 |
| 1972 | 2.2 | 68.8 | 2.0 | * | * | * | * | * | * | 2.8 | 8.0 | 18.2 |
| 1971 | 2.5 | 70.9 | 1.7 | * | * | * | * | 1.0 | 2.0 | 3.0 | 7.0 | 26.2 |
| 1970 | 2.5 | 75.9 | 2.4 | * | * | * | * | * | 1.6 | 3.3 | 6.5 | * |
| 1969 | 3.2 | 96.5 | 2.9 | * | * | * | * | 1.1 | 2.3 | 4.9 | 7.9 | 24.7 |
| 1968 | 3.3 | 98.5 | 3.7 | * | * | * | * | * | 1.9 | 4.1 | 8.0 | * |
| BLACK, MALE | | | | | | | | | | | | |
| 1978 | 2.0 | 73.0 | * | * | * | * | * | * | * | * | * | * |
| 1977 | 1.7 | 63.7 | * | * | * | * | * | * | * | * | * | * |
| 1976 | 1.8 | 69.9 | * | * | * | * | * | * | * | * | * | * |
| 1975 | 2.0 | 70.9 | * | * | * | * | * | * | * | * | * | * |
| 1974 | 1.8 | 62.4 | * | * | * | * | * | * | * | * | * | * |
| 1973 | 2.2 | 66.1 | 2.4 | * | * | * | * | * | * | * | * | * |
| 1972 | 2.3 | 69.7 | 2.4 | * | * | * | * | * | * | * | * | * |
| 1971 | 2.7 | 73.7 | * | * | * | * | * | * | 2.9 | * | * | * |
| 1970 | 3.1 | 92.0 | 2.3 | * | * | * | * | * | * | * | * | * |
| 1969 | 3.6 | 116.2 | 2.6 | * | * | * | * | * | * | * | * | * |
| 1968 | 3.7 | 108.4 | 4.3 | * | * | * | * | * | * | * | * | * |
| BLACK, FEMALE | | | | | | | | | | | | |
| 1978 | 1.6 | 61.3 | * | * | * | * | * | * | * | * | * | * |
| 1977 | 1.4 | 50.8 | * | * | * | * | * | * | * | * | * | * |
| 1976 | 1.7 | 64.2 | * | * | * | * | * | * | * | * | * | * |
| 1975 | 1.5 | 54.4 | * | * | * | * | * | * | * | * | * | * |
| 1974 | 1.7 | 51.6 | * | * | * | * | * | * | * | * | * | * |
| 1973 | 1.9 | 57.2 | * | * | * | * | * | * | * | 3.4 | * | * |
| 1972 | 2.1 | 67.5 | * | * | * | * | * | * | * | * | 9.3 | * |
| 1971 | 2.2 | 68.0 | * | * | * | * | * | * | * | * | * | * |
| 1970 | 2.1 | 59.7 | 2.5 | * | * | * | * | * | * | * | * | * |
| 1969 | 2.8 | 76.8 | 3.2 | * | * | * | * | * | * | 5.4 | 9.3 | * |
| 1968 | 2.9 | 88.5 | 3.2 | * | * | * | * | * | * | 5.0 | * | * |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| TUBERCULOSIS, ALL FORMS (010-019) | | | | | | | | | | | | |
| BOTH SEXES | | | | | | | | | | | | |
| 1978 | 1.3 | * | * | * | 0.1 | 0.2 | 0.5 | 1.5 | 2.7 | 5.0 | 8.8 | 16.2 |
| 1977 | 1.4 | * | * | * | 0.1 | 0.2 | 0.8 | 1.5 | 2.9 | 5.2 | 8.8 | 14.8 |
| 1976 | 1.4 | * | * | * | 0.1 | 0.2 | 0.9 | 1.6 | 3.1 | 5.7 | 10.0 | 14.3 |
| 1975 | 1.5 | * | * | * | 0.1 | 0.3 | 0.9 | 1.8 | 3.6 | 6.0 | 10.1 | 15.2 |
| 1974 | 1.6 | * | * | * | 0.1 | 0.4 | 1.0 | 2.1 | 3.9 | 6.6 | 10.0 | 15.8 |
| 1973 | 1.8 | * | * | * | 0.1 | 0.3 | 1.2 | 2.5 | 4.4 | 7.3 | 11.2 | 15.4 |
| 1972 | 2.1 | * | 0.1 | * | 0.1 | 0.5 | 1.3 | 2.8 | 5.0 | 8.8 | 12.3 | 19.8 |
| 1971 | 2.2 | * | * | * | 0.1 | 0.5 | 1.6 | 3.1 | 5.1 | 8.6 | 13.0 | 20.2 |
| 1970 | 2.6 | * | 0.3 | * | 0.1 | 0.7 | 1.8 | 3.5 | 6.2 | 10.1 | 16.4 | 20.9 |
| 1969 | 2.8 | * | 0.2 | * | 0.2 | 0.7 | 1.9 | 4.0 | 6.6 | 11.0 | 15.8 | 28.8 |
| 1968 | 3.2 | * | 0.3 | 0.0 | 0.2 | 0.8 | 2.2 | 4.4 | 8.0 | 12.8 | 18.3 | 28.3 |
| MALE | | | | | | | | | | | | |
| 1978 | 1.8 | * | * | * | * | 0.3 | 0.7 | 2.1 | 4.3 | 7.9 | 15.0 | 27.1 |
| 1977 | 1.9 | * | * | * | * | 0.3 | 1.1 | 2.2 | 4.4 | 8.8 | 15.3 | 24.7 |
| 1976 | 2.0 | * | * | * | * | 0.3 | 1.2 | 2.6 | 4.7 | 9.1 | 17.4 | 25.7 |
| 1975 | 2.2 | * | * | * | * | 0.4 | 1.3 | 2.8 | 5.3 | 9.6 | 18.0 | 27.1 |
| 1974 | 2.3 | * | * | * | * | 0.5 | 1.4 | 3.2 | 6.2 | 10.8 | 17.2 | 27.6 |
| 1973 | 2.6 | * | * | * | * | 0.4 | 1.7 | 3.8 | 7.1 | 12.2 | 19.4 | 27.0 |
| 1972 | 3.1 | * | * | * | 0.1 | 0.4 | 1.8 | 4.4 | 8.0 | 14.9 | 22.8 | 36.0 |
| 1971 | 3.2 | * | * | * | 0.1 | 0.6 | 2.3 | 4.6 | 8.2 | 14.8 | 23.2 | 35.5 |
| 1970 | 3.8 | * | 0.3 | * | 0.2 | 0.8 | 2.3 | 5.1 | 10.0 | 16.8 | 29.4 | 36.9 |
| 1969 | 4.1 | * | * | * | 0.2 | 0.8 | 2.4 | 6.0 | 10.9 | 19.2 | 28.6 | 47.5 |
| 1968 | 4.6 | * | 0.4 | * | 0.2 | 0.9 | 2.8 | 6.3 | 13.1 | 22.4 | 31.8 | 46.1 |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| TUBERCULOSIS, ALL FORMS (010-019) | | | | | | | | | | | | |
| FEMALE | | | | | | | | | | | | |
| 1978 | 0.9 | * | * | * | * | 0.1 | 0.4 | 0.9 | 1.4 | 2.7 | 5.1 | 11.2 |
| 1977 | 0.8 | * | * | * | * | 0.1 | 0.4 | 0.8 | 1.6 | 2.5 | 5.0 | 10.3 |
| 1976 | 0.9 | * | * | * | * | 0.2 | 0.6 | 0.7 | 1.6 | 3.1 | 5.5 | 9.0 |
| 1975 | 1.0 | * | * | * | * | 0.3 | 0.6 | 1.0 | 2.1 | 3.2 | 5.2 | 9.4 |
| 1974 | 1.0 | * | * | * | 0.1 | 0.3 | 0.6 | 1.0 | 1.8 | 3.4 | 5.6 | 10.0 |
| 1973 | 1.1 | * | * | * | * | 0.3 | 0.8 | 1.4 | 2.1 | 3.5 | 6.1 | 9.6 |
| 1972 | 1.2 | * | * | * | * | 0.5 | 0.9 | 1.4 | 2.2 | 4.0 | 5.6 | 11.6 |
| 1971 | 1.2 | * | * | * | 0.1 | 0.5 | 1.0 | 1.7 | 2.3 | 3.8 | 6.4 | 12.3 |
| 1970 | 1.4 | * | * | * | * | 0.6 | 1.3 | 1.9 | 2.7 | 4.9 | 7.7 | 12.0 |
| 1969 | 1.5 | * | * | * | 0.1 | 0.7 | 1.4 | 2.1 | 2.8 | 4.6 | 7.2 | 18.6 |
| 1968 | 1.8 | * | * | * | 0.2 | 0.8 | 1.5 | 2.6 | 3.5 | 5.2 | 9.1 | 18.5 |
| WHITE, BOTH SEXES | | | | | | | | | | | | |
| 1978 | 1.1 | * | * | * | * | 0.1 | 0.2 | 0.9 | 2.0 | 4.2 | 7.9 | 14.6 |
| 1977 | 1.1 | * | * | * | * | 0.1 | 0.3 | 0.9 | 2.2 | 4.1 | 7.7 | 13.4 |
| 1976 | 1.1 | * | * | * | * | 0.1 | 0.4 | 0.9 | 2.4 | 4.7 | 8.4 | 13.1 |
| 1975 | 1.2 | * | * | * | * | 0.1 | 0.4 | 1.1 | 2.7 | 4.9 | 9.1 | 13.7 |
| 1974 | 1.3 | * | * | * | * | 0.1 | 0.4 | 1.4 | 2.8 | 5.5 | 8.8 | 14.2 |
| 1973 | 1.4 | * | * | * | * | 0.1 | 0.5 | 1.5 | 3.5 | 6.0 | 10.2 | 14.1 |
| 1972 | 1.6 | * | * | * | 0.1 | 0.2 | 0.6 | 1.8 | 3.9 | 7.0 | 11.2 | 17.2 |
| 1971 | 1.7 | * | * | * | * | 0.2 | 0.7 | 2.0 | 4.1 | 7.1 | 11.6 | 18.0 |
| 1970 | 2.1 | * | * | * | 0.1 | 0.3 | 0.9 | 2.3 | 5.0 | 8.5 | 14.9 | 19.5 |
| 1969 | 2.2 | * | * | * | 0.1 | 0.3 | 0.9 | 2.7 | 5.2 | 9.4 | 14.3 | 26.7 |
| 1968 | 2.5 | * | 0.2 | * | 0.1 | 0.3 | 1.0 | 3.1 | 6.4 | 10.6 | 16.7 | 24.8 |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| TUBERCULOSIS, ALL FORMS (010-019) | | | | | | | | | | | | |
| WHITE, MALE | | | | | | | | | | | | |
| 1978 | 1.4 | * | * | * | * | * | 0.2 | 1.3 | 3.0 | 6.6 | 13.3 | 24.4 |
| 1977 | 1.4 | * | * | * | * | * | 0.5 | 1.2 | 3.4 | 6.9 | 13.1 | 21.7 |
| 1976 | 1.6 | * | * | * | * | * | 0.5 | 1.5 | 3.5 | 7.4 | 14.6 | 23.6 |
| 1975 | 1.7 | * | * | * | * | * | 0.6 | 1.7 | 4.1 | 7.7 | 16.8 | 24.6 |
| 1974 | 1.8 | * | * | * | * | * | 0.6 | 2.1 | 4.6 | 8.9 | 15.3 | 24.1 |
| 1973 | 2.1 | * | * | * | * | * | 0.7 | 2.1 | 5.6 | 10.0 | 17.6 | 25.1 |
| 1972 | 2.4 | * | * | * | * | * | 0.7 | 2.7 | 6.4 | 11.8 | 21.0 | 32.0 |
| 1971 | 2.5 | * | * | * | * | 0.2 | 1.1 | 2.9 | 6.3 | 12.0 | 20.7 | 31.3 |
| 1970 | 3.1 | * | * | * | * | 0.4 | 1.1 | 3.4 | 8.2 | 14.2 | 26.8 | 34.9 |
| 1969 | 3.3 | * | * | * | * | 0.3 | 1.2 | 4.2 | 8.5 | 16.3 | 26.2 | 45.1 |
| 1968 | 3.7 | * | * | * | * | 0.3 | 1.2 | 4.4 | 10.4 | 18.9 | 28.8 | 40.9 |
| WHITE, FEMALE | | | | | | | | | | | | |
| 1978 | 0.7 | * | * | * | * | * | 0.2 | 0.5 | 1.1 | 2.3 | 4.7 | 10.2 |
| 1977 | 0.7 | * | * | * | * | * | * | 0.5 | 1.2 | 2.0 | 4.5 | 9.6 |
| 1976 | 0.7 | * | * | * | * | * | 0.3 | 0.4 | 1.3 | 2.7 | 4.7 | 8.2 |
| 1975 | 0.8 | * | * | * | * | 0.2 | 0.2 | 0.6 | 1.5 | 2.7 | 4.5 | 8.6 |
| 1974 | 0.8 | * | * | * | * | * | 0.3 | 0.7 | 1.3 | 3.0 | 4.8 | 9.4 |
| 1973 | 0.9 | * | * | * | * | * | 0.4 | 0.9 | 1.6 | 2.9 | 5.6 | 8.6 |
| 1972 | 0.9 | * | * | * | * | 0.2 | 0.6 | 0.9 | 1.7 | 3.2 | 5.0 | 9.7 |
| 1971 | 1.0 | * | * | * | * | 0.2 | 0.4 | 1.1 | 2.0 | 3.2 | 5.8 | 11.2 |
| 1970 | 1.1 | * | * | * | * | 0.2 | 0.7 | 1.3 | 2.1 | 4.1 | 7.1 | 11.1 |
| 1969 | 1.2 | * | * | * | * | 0.3 | 0.6 | 1.3 | 2.2 | 4.0 | 6.4 | 17.0 |
| 1968 | 1.3 | * | * | * | * | 0.3 | 0.7 | 1.8 | 2.7 | 4.0 | 8.5 | 16.1 |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| TUBERCULOSIS, ALL FORMS (010-019) | | | | | | | | | | | | |
| ALL OTHER, BOTH SEXES | | | | | | | | | | | | |
| 1978 | 2.9 | * | * | * | * | 1.0 | 2.5 | 6.0 | 9.8 | 12.5 | 18.8 | 34.5 |
| 1977 | 3.2 | * | * | * | * | 1.4 | 3.8 | 6.6 | 9.0 | 15.4 | 20.2 | 31.3 |
| 1976 | 3.4 | * | * | * | * | 1.2 | 4.2 | 6.9 | 9.8 | 15.3 | 26.4 | 28.8 |
| 1975 | 3.6 | * | * | * | * | 1.5 | 4.3 | 7.6 | 11.5 | 16.7 | 20.2 | 31.5 |
| 1974 | 4.1 | * | * | * | 0.3 | 2.3 | 4.9 | 7.9 | 13.6 | 17.2 | 23.7 | 34.8 |
| 1973 | 4.5 | * | * | * | * | 1.9 | 6.3 | 11.2 | 13.5 | 19.8 | 23.0 | 31.5 |
| 1972 | 5.1 | * | * | * | * | 2.5 | 6.5 | 11.8 | 14.7 | 26.6 | 25.0 | 50.4 |
| 1971 | 5.5 | * | * | * | 0.6 | 2.7 | 8.0 | 12.9 | 15.1 | 23.6 | 30.2 | 45.8 |
| 1970 | 6.0 | * | 1.0 | * | 0.5 | 3.1 | 8.3 | 13.8 | 17.3 | 25.6 | 34.9 | 35.1 |
| 1969 | 6.7 | * | * | * | 0.8 | 4.0 | 9.4 | 15.4 | 20.5 | 27.2 | 35.5 | 51.4 |
| 1968 | 7.9 | * | * | * | 0.8 | 4.7 | 11.4 | 16.3 | 24.2 | 35.1 | 39.2 | 67.6 |
| ALL OTHER, MALE | | | | | | | | | | | | |
| 1978 | 4.3 | * | * | * | * | 1.5 | 3.6 | 8.8 | 16.7 | 20.5 | 31.2 | 55.2 |
| 1977 | 4.9 | * | * | * | * | 2.2 | 5.9 | 10.8 | 14.4 | 26.0 | 35.2 | 55.4 |
| 1976 | 5.1 | * | * | * | * | 1.6 | 6.2 | 11.9 | 16.4 | 25.6 | 44.4 | 47.3 |
| 1975 | 5.1 | * | * | * | * | 2.2 | 6.2 | 12.0 | 16.5 | 28.0 | 30.2 | 51.9 |
| 1974 | 6.0 | * | * | * | * | 3.3 | 7.2 | 13.0 | 21.4 | 29.2 | 36.7 | 64.0 |
| 1973 | 6.6 | * | * | * | * | 2.4 | 9.3 | 18.4 | 21.9 | 32.8 | 38.4 | 45.8 |
| 1972 | 7.7 | * | * | * | * | 2.4 | 10.3 | 19.6 | 23.9 | 44.8 | 42.4 | 76.6 |
| 1971 | 8.2 | * | * | * | * | 3.2 | 11.6 | 20.2 | 26.7 | 41.1 | 51.7 | 76.6 |
| 1970 | 8.6 | * | * | * | * | 3.8 | 12.0 | 21.0 | 26.8 | 40.8 | 59.6 | 54.1 |
| 1969 | 9.6 | * | * | * | * | 4.7 | 12.0 | 23.1 | 33.5 | 46.5 | 57.4 | 70.5 |
| 1968 | 11.5 | * | * | * | * | 5.4 | 16.2 | 23.8 | 39.6 | 56.7 | 67.7 | 97.6 |
| ALL OTHER, FEMALE | | | | | | | | | | | | |
| 1978 | 1.7 | * | * | * | * | * | 1.5 | 3.6 | 4.0 | 6.2 | 10.3 | 23.9 |
| 1977 | 1.7 | * | * | * | * | * | 2.1 | 3.1 | 4.6 | 7.0 | 9.9 | * |
| 1976 | 1.9 | * | * | * | * | * | 2.6 | 2.8 | 4.3 | 7.2 | 14.1 | * |
| 1975 | 2.2 | * | * | * | * | * | 2.6 | 3.9 | 7.3 | 7.7 | 13.3 | * |
| 1974 | 2.3 | * | * | * | * | 1.5 | 2.9 | 3.6 | 6.9 | 7.6 | 14.8 | * |
| 1973 | 2.5 | * | * | * | * | 1.5 | 3.7 | 5.0 | 6.3 | 9.3 | 12.3 | * |
| 1972 | 2.8 | * | * | * | * | 2.6 | 3.3 | 5.1 | 6.7 | 11.9 | 12.6 | 34.2 |
| 1971 | 2.9 | * | * | * | * | 2.3 | 4.9 | 6.5 | 5.0 | 9.2 | 14.3 | * |
| 1970 | 3.6 | * | * | * | * | 2.6 | 5.2 | 7.4 | 8.9 | 13.1 | 16.2 | * |
| 1969 | 4.1 | * | * | * | * | 3.5 | 7.3 | 8.6 | 9.1 | 11.2 | 18.6 | 38.5 |
| 1968 | 4.6 | * | * | * | * | 4.1 | 7.4 | 9.6 | 10.3 | 17.2 | 17.2 | 47.5 |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| TUBERCULOSIS, ALL FORMS (010-019) | | | | | | | | | | | | |
| BLACK, BOTH SEXES | | | | | | | | | | | | |
| 1978 | 3.1 | * | * | * | * | 1.1 | 2.8 | 6.4 | 10.3 | 12.5 | 17.3 | 32.7 |
| 1977 | 3.4 | * | * | * | * | 1.6 | 4.4 | 7.2 | 9.6 | 15.4 | 19.2 | 29.4 |
| 1976 | 3.6 | * | * | * | * | 1.4 | 4.7 | 7.5 | 10.2 | 15.2 | 25.4 | 27.9 |
| 1975 | 3.7 | * | * | * | * | 1.5 | 4.8 | 8.3 | 11.7 | 16.7 | 17.6 | 24.6 |
| 1974 | 4.2 | * | * | * | * | 2.6 | 5.2 | 8.4 | 14.1 | 17.4 | 22.6 | 31.4 |
| 1973 | 4.6 | * | * | * | * | 2.2 | 6.9 | 12.1 | 14.1 | 18.8 | 21.4 | 28.9 |
| 1972 | 5.2 | * | * | * | * | 2.7 | 6.9 | 12.4 | 14.8 | 27.1 | 22.7 | 49.1 |
| 1971 | 5.6 | * | * | * | 0.5 | 2.9 | 8.6 | 13.7 | 15.5 | 24.1 | 27.6 | 43.9 |
| 1970 | 6.2 | * | * | * | 0.5 | 3.4 | 9.2 | 14.5 | 18.0 | 26.1 | 33.4 | 31.5 |
| 1969 | 6.8 | * | * | * | 0.8 | 4.3 | 10.0 | 15.9 | 20.4 | 27.4 | 33.8 | 48.5 |
| 1968 | 8.0 | * | * | * | 0.9 | 4.9 | 12.2 | 16.8 | 24.7 | 35.3 | 37.5 | 53.3 |
| BLACK, MALE | | | | | | | | | | | | |
| 1978 | 4.5 | * | * | * | * | 1.7 | 4.2 | 9.6 | 18.1 | 21.2 | 29.3 | 51.0 |
| 1977 | 5.3 | * | * | * | * | 2.6 | 6.8 | 12.2 | 16.1 | 27.2 | 34.0 | 49.0 |
| 1976 | 5.4 | * | * | * | * | 1.8 | 6.9 | 12.9 | 17.6 | 25.5 | 42.3 | 46.8 |
| 1975 | 5.2 | * | * | * | * | 2.3 | 7.2 | 12.8 | 17.1 | 28.5 | 26.1 | * |
| 1974 | 6.3 | * | * | * | * | 3.6 | 7.8 | 13.9 | 23.0 | 30.5 | 35.4 | 58.1 |
| 1973 | 6.9 | * | * | * | * | 2.7 | 10.5 | 20.2 | 23.1 | 31.3 | 35.3 | * |
| 1972 | 8.1 | * | * | * | * | 2.7 | 11.2 | 21.2 | 24.9 | 47.8 | 38.6 | 80.0 |
| 1971 | 8.6 | * | * | * | * | 3.5 | 12.7 | 21.6 | 28.5 | 42.4 | 48.9 | 70.0 |
| 1970 | 9.0 | * | * | * | * | 4.3 | 13.6 | 22.2 | 28.0 | 42.5 | 56.9 | 54.4 |
| 1969 | 9.9 | * | * | * | * | 5.0 | 13.0 | 24.4 | 34.6 | 47.9 | 55.1 | 73.0 |
| 1968 | 11.8 | * | * | * | * | 5.8 | 17.8 | 24.7 | 41.5 | 58.7 | 65.5 | 74.3 |
| BLACK, FEMALE | | | | | | | | | | | | |
| 1978 | 1.7 | * | * | * | * | * | 1.7 | 3.8 | 3.8 | 5.9 | 9.2 | 23.2 |
| 1977 | 1.7 | * | * | * | * | * | 2.3 | 3.2 | 4.3 | 6.4 | 9.3 | * |
| 1976 | 2.0 | * | * | * | * | * | 2.9 | 3.1 | 4.1 | 7.4 | 14.0 | * |
| 1975 | 2.2 | * | * | * | * | * | 2.8 | 4.4 | 7.3 | 7.7 | 11.9 | * |
| 1974 | 2.3 | * | * | * | * | 1.7 | 3.1 | 3.7 | 6.6 | 7.3 | 13.9 | * |
| 1973 | 2.6 | * | * | * | * | 1.7 | 3.9 | 5.1 | 6.7 | 9.2 | 11.8 | * |
| 1972 | 2.7 | * | * | * | * | 2.7 | 3.3 | 4.9 | 6.4 | 11.0 | 11.4 | 31.4 |
| 1971 | 3.0 | * | * | * | * | 2.5 | 5.2 | 6.9 | 4.6 | 9.6 | 12.4 | * |
| 1970 | 3.7 | * | * | * | * | 2.7 | 5.7 | 7.9 | 9.6 | 13.1 | 16.1 | * |
| 1969 | 4.0 | * | * | * | * | 3.7 | 7.6 | 8.5 | 8.3 | 11.0 | 18.1 | 33.3 |
| 1968 | 4.6 | * | * | * | * | 4.1 | 7.6 | 9.9 | 10.3 | 16.3 | 16.7 | 41.1 |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|--|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| TUBERCULOSIS OF RESPIRATORY SYSTEM (010-012) | | | | | | | | | | | | |
| BOTH SEXES | | | | | | | | | | | | |
| 1978 | 1.0 | * | * | * | 0.0 | 0.1 | 0.4 | 1.2 | 2.2 | 3.8 | 6.7 | 13.7 |
| 1977 | 1.1 | * | * | * | * | 0.2 | 0.6 | 1.2 | 2.3 | 4.1 | 6.7 | 12.9 |
| 1976 | 1.1 | * | * | * | * | 0.2 | 0.7 | 1.2 | 2.4 | 4.2 | 7.9 | 12.5 |
| 1975 | 1.2 | * | * | * | * | 0.2 | 0.7 | 1.5 | 2.8 | 4.7 | 7.7 | 12.0 |
| 1974 | 1.3 | * | * | * | 0.1 | 0.3 | 0.8 | 1.6 | 3.0 | 5.2 | 8.0 | 13.4 |
| 1973 | 1.4 | * | * | * | * | 0.2 | 1.0 | 2.0 | 3.5 | 5.6 | 8.7 | 12.4 |
| 1972 | 1.7 | * | * | * | 0.1 | 0.4 | 1.1 | 2.4 | 3.9 | 6.8 | 9.6 | 16.2 |
| 1971 | 1.7 | * | * | * | 0.1 | 0.4 | 1.4 | 2.5 | 4.1 | 6.7 | 10.6 | 17.1 |
| 1970 | 2.0 | * | * | * | 0.1 | 0.5 | 1.5 | 2.7 | 5.0 | 7.9 | 13.4 | 17.9 |
| 1969 | 2.2 | * | * | * | 0.1 | 0.6 | 1.5 | 3.2 | 5.3 | 9.0 | 13.0 | 24.5 |
| 1968 | 2.5 | * | * | * | 0.1 | 0.6 | 1.8 | 3.5 | 6.4 | 10.0 | 15.2 | 23.3 |
| MALE | | | | | | | | | | | | |
| 1978 | 1.4 | * | * | * | * | 0.2 | 0.6 | 1.7 | 3.6 | 6.1 | 11.9 | 23.3 |
| 1977 | 1.5 | * | * | * | * | 0.3 | 0.9 | 1.8 | 3.7 | 7.2 | 12.1 | 22.0 |
| 1976 | 1.6 | * | * | * | * | 0.2 | 1.0 | 2.1 | 3.8 | 7.2 | 13.9 | 22.4 |
| 1975 | 1.7 | * | * | * | * | 0.3 | 1.1 | 2.3 | 4.3 | 7.6 | 13.9 | 22.2 |
| 1974 | 1.9 | * | * | * | * | 0.4 | 1.1 | 2.6 | 5.0 | 8.8 | 14.2 | 23.7 |
| 1973 | 2.1 | * | * | * | * | 0.3 | 1.4 | 3.1 | 5.9 | 9.7 | 15.0 | 22.3 |
| 1972 | 2.5 | * | * | * | * | 0.3 | 1.6 | 3.8 | 6.7 | 11.8 | 17.7 | 29.5 |
| 1971 | 2.6 | * | * | * | * | 0.4 | 1.9 | 3.9 | 6.8 | 11.8 | 19.1 | 30.0 |
| 1970 | 3.1 | * | * | * | * | 0.6 | 2.0 | 4.2 | 8.3 | 13.8 | 24.7 | 32.3 |
| 1969 | 3.4 | * | * | * | 0.1 | 0.5 | 2.0 | 5.1 | 9.0 | 16.2 | 23.6 | 39.9 |
| 1968 | 3.8 | * | * | * | * | 0.8 | 2.4 | 5.2 | 10.9 | 17.9 | 26.7 | 38.9 |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|--|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| TUBERCULOSIS OF RESPIRATORY SYSTEM (010-012) | | | | | | | | | | | | |
| FEMALE | | | | | | | | | | | | |
| 1978 | 0.6 | * | * | * | * | * | 0.3 | 0.6 | 0.9 | 2.0 | 3.6 | 9.3 |
| 1977 | 0.6 | * | * | * | * | * | 0.3 | 0.6 | 1.0 | 1.7 | 3.5 | 8.6 |
| 1976 | 0.6 | * | * | * | * | 0.1 | 0.4 | 0.5 | 1.1 | 2.0 | 4.2 | 7.8 |
| 1975 | 0.7 | * | * | * | * | 0.1 | 0.4 | 0.7 | 1.5 | 2.4 | 3.9 | 7.0 |
| 1974 | 0.7 | * | * | * | * | 0.3 | 0.5 | 0.7 | 1.3 | 2.4 | 4.2 | 8.2 |
| 1973 | 0.8 | * | * | * | * | 0.2 | 0.6 | 1.0 | 1.4 | 2.4 | 4.8 | 7.5 |
| 1972 | 0.9 | * | * | * | * | 0.4 | 0.7 | 1.1 | 1.4 | 3.0 | 4.5 | 9.4 |
| 1971 | 0.9 | * | * | * | * | 0.3 | 0.8 | 1.2 | 1.6 | 2.7 | 5.1 | 10.4 |
| 1970 | 1.1 | * | * | * | * | 0.4 | 1.0 | 1.4 | 2.0 | 3.4 | 5.9 | 9.9 |
| 1969 | 1.1 | * | * | * | * | 0.6 | 1.1 | 1.5 | 1.9 | 3.4 | 5.8 | 16.2 |
| 1968 | 1.3 | * | * | * | * | 0.5 | 1.3 | 1.8 | 2.3 | 3.8 | 7.4 | 14.8 |
| WHITE, BOTH SEXES | | | | | | | | | | | | |
| 1978 | 0.8 | * | * | * | * | * | 0.2 | 0.7 | 1.5 | 3.1 | 6.1 | 12.6 |
| 1977 | 0.8 | * | * | * | * | * | 0.3 | 0.7 | 1.7 | 3.3 | 5.9 | 11.7 |
| 1976 | 0.9 | * | * | * | * | * | 0.3 | 0.7 | 1.8 | 3.5 | 6.5 | 11.4 |
| 1975 | 1.0 | * | * | * | * | 0.1 | 0.3 | 0.9 | 2.1 | 3.8 | 7.0 | 11.2 |
| 1974 | 1.0 | * | * | * | * | 0.1 | 0.4 | 1.0 | 2.2 | 4.4 | 7.0 | 12.0 |
| 1973 | 1.1 | * | * | * | * | 0.1 | 0.4 | 1.1 | 2.7 | 4.6 | 7.9 | 11.1 |
| 1972 | 1.3 | * | * | * | * | 0.1 | 0.5 | 1.5 | 3.1 | 5.4 | 8.7 | 14.4 |
| 1971 | 1.3 | * | * | * | * | 0.1 | 0.6 | 1.6 | 3.2 | 5.4 | 9.5 | 14.9 |
| 1970 | 1.6 | * | * | * | * | 0.2 | 0.7 | 1.8 | 4.0 | 6.5 | 12.1 | 16.7 |
| 1969 | 1.8 | * | * | * | * | 0.2 | 0.8 | 2.2 | 4.1 | 7.6 | 11.7 | 22.8 |
| 1968 | 2.0 | * | * | * | * | 0.2 | 0.8 | 2.4 | 5.0 | 8.2 | 13.9 | 20.4 |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|--|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| TUBERCULOSIS OF RESPIRATORY SYSTEM (010-012) | | | | | | | | | | | | |
| WHITE, MALE | | | | | | | | | | | | |
| 1978 | 1.1 | * | * | * | * | * | 0.2 | 1.1 | 2.5 | 4.9 | 10.6 | 21.2 |
| 1977 | 1.2 | * | * | * | * | * | 0.4 | 1.0 | 2.8 | 5.7 | 10.5 | 19.3 |
| 1976 | 1.2 | * | * | * | * | * | 0.4 | 1.2 | 2.8 | 5.8 | 11.6 | 20.5 |
| 1975 | 1.4 | * | * | * | * | * | 0.5 | 1.5 | 3.2 | 6.1 | 13.0 | 20.4 |
| 1974 | 1.5 | * | * | * | * | * | 0.6 | 1.7 | 3.7 | 7.2 | 12.6 | 20.6 |
| 1973 | 1.6 | * | * | * | * | * | 0.5 | 1.7 | 4.5 | 8.0 | 13.7 | 20.2 |
| 1972 | 2.0 | * | * | * | * | * | 0.6 | 2.4 | 5.4 | 9.3 | 16.2 | 26.5 |
| 1971 | 2.0 | * | * | * | * | * | 1.0 | 2.5 | 5.2 | 9.5 | 16.9 | 25.9 |
| 1970 | 2.5 | * | * | * | * | 0.3 | 1.0 | 2.7 | 6.9 | 11.5 | 22.3 | 30.6 |
| 1969 | 2.7 | * | * | * | * | * | 1.1 | 3.6 | 7.0 | 13.7 | 21.5 | 37.6 |
| 1968 | 3.0 | * | * | * | * | 0.2 | 1.0 | 3.6 | 8.6 | 15.0 | 24.0 | 34.6 |
| WHITE, FEMALE | | | | | | | | | | | | |
| 1978 | 0.5 | * | * | * | * | * | * | 0.4 | 0.7 | 1.7 | 3.4 | 8.7 |
| 1977 | 0.5 | * | * | * | * | * | * | 0.4 | 0.8 | 1.4 | 3.1 | 8.2 |
| 1976 | 0.5 | * | * | * | * | * | 0.2 | 0.3 | 0.9 | 1.8 | 3.5 | 7.2 |
| 1975 | 0.6 | * | * | * | * | * | * | 0.4 | 1.1 | 2.1 | 3.4 | 6.8 |
| 1974 | 0.6 | * | * | * | * | * | 0.2 | 0.5 | 0.9 | 2.2 | 3.5 | 7.9 |
| 1973 | 0.6 | * | * | * | * | * | 0.3 | 0.6 | 1.0 | 2.0 | 4.4 | 6.6 |
| 1972 | 0.7 | * | * | * | * | 0.2 | 0.4 | 0.7 | 1.0 | 2.4 | 4.0 | 8.3 |
| 1971 | 0.7 | * | * | * | * | * | 0.3 | 0.8 | 1.4 | 2.3 | 4.7 | 9.2 |
| 1970 | 0.8 | * | * | * | * | * | 0.5 | 0.9 | 1.4 | 2.7 | 5.4 | 9.1 |
| 1969 | 0.9 | * | * | * | * | 0.2 | 0.5 | 0.9 | 1.4 | 2.8 | 5.2 | 14.9 |
| 1968 | 1.0 | * | * | * | * | * | 0.6 | 1.3 | 1.8 | 2.9 | 7.0 | 12.7 |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|--|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| TUBERCULOSIS OF RESPIRATORY SYSTEM (010-012) | | | | | | | | | | | | |
| ALL OTHER, BOTH SEXES | | | | | | | | | | | | |
| 1978 | 2.3 | * | * | * | * | 0.7 | 2.1 | 4.7 | 8.2 | 10.1 | 13.1 | 26.2 |
| 1977 | 2.5 | * | * | * | * | 1.0 | 2.9 | 5.3 | 7.3 | 12.1 | 14.9 | 26.3 |
| 1976 | 2.7 | * | * | * | * | 0.9 | 3.4 | 5.4 | 7.9 | 11.3 | 21.5 | 24.8 |
| 1975 | 2.8 | * | * | * | * | 1.0 | 3.5 | 6.1 | 9.4 | 13.1 | 15.0 | 21.2 |
| 1974 | 3.2 | * | * | * | * | 1.8 | 3.7 | 6.5 | 11.2 | 13.3 | 19.3 | 28.9 |
| 1973 | 3.6 | * | * | * | * | 1.2 | 5.3 | 9.4 | 11.4 | 15.3 | 17.7 | 28.3 |
| 1972 | 4.2 | * | * | * | * | 2.0 | 5.7 | 10.1 | 11.7 | 20.9 | 20.4 | 37.4 |
| 1971 | 4.4 | * | * | * | * | 2.0 | 6.6 | 10.6 | 12.8 | 19.5 | 24.6 | 43.3 |
| 1970 | 4.9 | * | * | * | * | 2.5 | 6.8 | 11.3 | 14.0 | 21.4 | 29.8 | 30.6 |
| 1969 | 5.4 | * | * | * | 0.5 | 3.2 | 7.1 | 12.5 | 17.0 | 23.2 | 29.7 | 43.1 |
| 1968 | 6.4 | * | * | * | 0.5 | 3.9 | 9.6 | 13.2 | 19.7 | 28.1 | 32.6 | 55.9 |
| ALL OTHER, MALE | | | | | | | | | | | | |
| 1978 | 3.5 | * | * | * | * | 1.0 | 3.1 | 7.3 | 14.5 | 17.0 | 24.1 | 44.8 |
| 1977 | 3.9 | * | * | * | * | 1.7 | 4.8 | 8.8 | 11.8 | 20.9 | 26.8 | 50.0 |
| 1976 | 4.1 | * | * | * | * | 1.1 | 5.1 | 9.4 | 13.5 | 20.2 | 35.8 | 41.8 |
| 1975 | 4.1 | * | * | * | * | 1.8 | 5.5 | 9.7 | 14.0 | 22.4 | 22.7 | 40.7 |
| 1974 | 4.8 | * | * | * | * | 2.5 | 5.6 | 10.8 | 18.2 | 23.4 | 30.1 | 56.0 |
| 1973 | 5.5 | * | * | * | * | 1.8 | 8.3 | 15.7 | 18.8 | 26.1 | 29.5 | 43.8 |
| 1972 | 6.4 | * | * | * | * | 1.9 | 9.7 | 16.3 | 19.6 | 35.8 | 34.1 | 59.6 |
| 1971 | 6.9 | * | * | * | * | 2.4 | 9.4 | 16.9 | 23.3 | 34.1 | 43.5 | 70.2 |
| 1970 | 7.2 | * | * | * | * | 3.1 | 10.0 | 17.6 | 22.0 | 35.3 | 52.3 | 46.9 |
| 1969 | 8.0 | * | * | * | * | 3.4 | 9.4 | 18.7 | 29.0 | 40.2 | 48.9 | 61.4 |
| 1968 | 9.5 | * | * | * | * | 4.9 | 13.7 | 20.2 | 33.0 | 46.3 | 58.6 | 80.5 |
| ALL OTHER, FEMALE | | | | | | | | | | | | |
| 1978 | 1.2 | * | * | * | * | * | 1.2 | 2.5 | 2.9 | 4.8 | 5.7 | * |
| 1977 | 1.2 | * | * | * | * | * | 1.3 | 2.4 | 3.6 | 5.1 | 6.8 | * |
| 1976 | 1.4 | * | * | * | * | * | 1.9 | 2.0 | 3.1 | 4.3 | 11.7 | * |
| 1975 | 1.5 | * | * | * | * | * | 1.8 | 3.1 | 5.5 | 5.7 | 9.6 | * |
| 1974 | 1.7 | * | * | * | * | 1.2 | 2.2 | 2.8 | 5.2 | 5.2 | 11.7 | * |
| 1973 | 1.8 | * | * | * | * | * | 2.7 | 3.9 | 5.0 | 6.7 | 9.4 | * |
| 1972 | 2.2 | * | * | * | * | 2.1 | 2.4 | 4.7 | 4.9 | 8.9 | 10.6 | * |
| 1971 | 2.2 | * | * | * | * | 1.6 | 4.2 | 5.0 | 3.5 | 7.6 | 10.7 | * |
| 1970 | 2.7 | * | * | * | * | 2.0 | 4.2 | 5.8 | 6.9 | 10.0 | 12.6 | * |
| 1969 | 3.1 | * | * | * | * | 3.0 | 5.2 | 7.1 | 6.3 | 9.3 | 15.0 | 30.8 |
| 1968 | 3.5 | * | * | * | * | 3.0 | 6.1 | 7.0 | 7.7 | 13.1 | 12.6 | 39.3 |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|--|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| TUBERCULOSIS OF RESPIRATORY SYSTEM (010-012) | | | | | | | | | | | | |
| BLACK, BOTH SEXES | | | | | | | | | | | | |
| 1978 | 2.4 | * | * | * | * | 0.8 | 2.5 | 5.1 | 8.6 | 10.1 | 11.7 | 26.7 |
| 1977 | 2.7 | * | * | * | * | 1.2 | 3.3 | 5.9 | 8.0 | 12.2 | 14.6 | 24.5 |
| 1976 | 2.8 | * | * | * | * | 1.0 | 3.9 | 5.9 | 8.3 | 11.5 | 20.9 | 24.3 |
| 1975 | 2.9 | * | * | * | * | 1.1 | 3.9 | 6.6 | 9.5 | 13.4 | 14.0 | 18.5 |
| 1974 | 3.4 | * | * | * | * | 2.0 | 4.1 | 6.9 | 11.7 | 13.9 | 18.7 | 26.4 |
| 1973 | 3.8 | * | * | * | * | 1.4 | 5.8 | 10.2 | 12.0 | 14.6 | 16.6 | 25.4 |
| 1972 | 4.3 | * | * | * | * | 2.1 | 6.1 | 10.6 | 12.2 | 21.8 | 18.5 | 36.4 |
| 1971 | 4.7 | * | * | * | * | 2.2 | 7.3 | 11.4 | 13.2 | 20.0 | 22.9 | 41.1 |
| 1970 | 5.1 | * | * | * | * | 2.8 | 7.7 | 11.9 | 14.8 | 21.9 | 29.1 | 28.1 |
| 1969 | 5.6 | * | * | * | 0.5 | 3.4 | 7.7 | 13.0 | 17.3 | 23.5 | 28.2 | 44.3 |
| 1968 | 6.6 | * | * | * | 0.6 | 4.1 | 10.4 | 13.7 | 20.5 | 28.4 | 30.7 | 48.9 |
| BLACK, MALE | | | | | | | | | | | | |
| 1978 | 3.7 | * | * | * | * | 1.1 | 3.7 | 8.0 | 15.7 | 17.6 | 21.8 | 43.1 |
| 1977 | 4.3 | * | * | * | * | 2.0 | 5.5 | 10.0 | 13.3 | 22.2 | 26.8 | 44.9 |
| 1976 | 4.4 | * | * | * | * | 1.2 | 5.9 | 10.4 | 14.6 | 20.7 | 34.2 | * |
| 1975 | 4.3 | * | * | * | * | 1.9 | 6.3 | 10.3 | 14.2 | 23.7 | 21.6 | * |
| 1974 | 5.2 | * | * | * | * | 2.8 | 6.1 | 11.6 | 19.7 | 25.3 | 29.2 | 53.5 |
| 1973 | 5.8 | * | * | * | * | 2.0 | 9.3 | 17.5 | 20.1 | 25.0 | 27.9 | * |
| 1972 | 6.8 | * | * | * | * | 2.1 | 10.4 | 17.8 | 20.9 | 39.2 | 30.5 | 60.0 |
| 1971 | 7.2 | * | * | * | * | 2.7 | 10.5 | 18.4 | 25.1 | 35.1 | 41.9 | 62.5 |
| 1970 | 7.6 | * | * | * | * | 3.6 | 11.4 | 18.5 | 23.4 | 36.9 | 51.0 | 47.8 |
| 1969 | 8.3 | * | * | * | * | 3.7 | 10.4 | 19.9 | 30.2 | 41.3 | 46.7 | 64.9 |
| 1968 | 9.9 | * | * | * | * | 5.4 | 15.2 | 20.9 | 35.2 | 47.9 | 55.8 | 71.4 |
| BLACK, FEMALE | | | | | | | | | | | | |
| 1978 | 1.2 | * | * | * | * | * | 1.4 | 2.7 | 2.8 | 4.6 | * | * |
| 1977 | 1.3 | * | * | * | * | * | 1.4 | 2.4 | 3.6 | 4.6 | 6.5 | * |
| 1976 | 1.5 | * | * | * | * | * | 2.1 | 2.2 | 3.1 | 4.5 | 12.0 | * |
| 1975 | 1.6 | * | * | * | * | * | 1.9 | 3.5 | 5.6 | 5.6 | 8.8 | * |
| 1974 | 1.8 | * | * | * | * | 1.4 | 2.3 | 3.0 | 5.1 | 5.2 | 11.7 | * |
| 1973 | 1.9 | * | * | * | * | * | 2.9 | 4.0 | 5.3 | 6.6 | 8.8 | * |
| 1972 | 2.1 | * | * | * | * | 2.2 | 2.4 | 4.4 | 4.9 | 8.3 | 10.0 | * |
| 1971 | 2.3 | * | * | * | * | 1.9 | 4.6 | 5.3 | 3.2 | 8.1 | 9.3 | * |
| 1970 | 2.9 | * | * | * | * | 2.1 | 4.7 | 6.2 | 7.5 | 10.0 | 13.0 | * |
| 1969 | 3.1 | * | * | * | * | 3.2 | 5.5 | 7.1 | 6.2 | 9.3 | 14.5 | * |
| 1968 | 3.5 | * | * | * | * | 3.0 | 6.3 | 7.4 | 7.7 | 12.6 | 12.2 | 35.7 |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| TUBERCULOSIS, OTHER FORMS (013-019) | | | | | | | | | | | | |
| BOTH SEXES | | | | | | | | | | | | |
| 1978 | 0.3 | * | * | * | * | 0.1 | 0.1 | 0.3 | 0.6 | 1.2 | 2.1 | 2.5 |
| 1977 | 0.3 | * | * | * | * | * | 0.2 | 0.3 | 0.6 | 1.1 | 2.1 | 2.0 |
| 1976 | 0.3 | * | * | * | * | 0.1 | 0.2 | 0.4 | 0.7 | 1.5 | 2.1 | 1.8 |
| 1975 | 0.4 | * | * | * | * | 0.1 | 0.2 | 0.3 | 0.8 | 1.3 | 2.4 | 3.2 |
| 1974 | 0.3 | * | * | * | * | 0.1 | 0.2 | 0.4 | 0.8 | 1.4 | 2.0 | 2.5 |
| 1973 | 0.4 | * | * | * | * | 0.1 | 0.2 | 0.5 | 1.0 | 1.7 | 2.5 | 3.0 |
| 1972 | 0.4 | * | * | * | * | 0.1 | 0.2 | 0.4 | 1.1 | 1.9 | 2.7 | 3.6 |
| 1971 | 0.4 | * | * | * | * | 0.1 | 0.3 | 0.6 | 1.0 | 1.9 | 2.4 | 3.1 |
| 1970 | 0.5 | * | 0.2 | * | 0.1 | 0.2 | 0.3 | 0.7 | 1.2 | 2.2 | 3.0 | 3.0 |
| 1969 | 0.5 | * | * | * | * | 0.1 | 0.4 | 0.8 | 1.4 | 2.0 | 2.8 | 4.3 |
| 1968 | 0.6 | * | 0.2 | * | 0.1 | 0.2 | 0.4 | 0.9 | 1.6 | 2.8 | 3.1 | 5.0 |
| MALE | | | | | | | | | | | | |
| 1978 | 0.4 | * | * | * | * | * | * | 0.4 | 0.7 | 1.8 | 3.1 | 3.8 |
| 1977 | 0.3 | * | * | * | * | * | 0.2 | 0.4 | 0.7 | 1.6 | 3.2 | * |
| 1976 | 0.4 | * | * | * | * | * | 0.2 | 0.5 | 0.9 | 1.9 | 3.5 | 3.3 |
| 1975 | 0.4 | * | * | * | * | * | 0.2 | 0.4 | 1.0 | 2.0 | 4.1 | 4.9 |
| 1974 | 0.4 | * | * | * | * | * | 0.2 | 0.6 | 1.1 | 2.0 | 3.0 | 3.9 |
| 1973 | 0.5 | * | * | * | * | * | 0.2 | 0.6 | 1.3 | 2.4 | 4.4 | 4.6 |
| 1972 | 0.6 | * | * | * | * | * | 0.2 | 0.6 | 1.3 | 3.1 | 5.1 | 6.5 |
| 1971 | 0.6 | * | * | * | * | * | 0.4 | 0.7 | 1.4 | 2.9 | 4.1 | 5.5 |
| 1970 | 0.7 | * | * | * | * | * | 0.3 | 0.9 | 1.7 | 3.0 | 4.7 | 4.6 |
| 1969 | 0.7 | * | * | * | * | 0.2 | 0.4 | 1.0 | 1.8 | 2.9 | 4.9 | 7.6 |
| 1968 | 0.8 | * | * | * | * | * | 0.5 | 1.1 | 2.2 | 4.5 | 5.1 | 7.3 |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| TUBERCULOSIS, OTHER FORMS (013-019) | | | | | | | | | | | | |
| FEMALE | | | | | | | | | | | | |
| 1978 | 0.2 | * | * | * | * | * | * | 0.3 | 0.4 | 0.7 | 1.6 | 1.9 |
| 1977 | 0.2 | * | * | * | * | * | * | 0.2 | 0.5 | 0.7 | 1.5 | 1.6 |
| 1976 | 0.3 | * | * | * | * | * | * | 0.2 | 0.5 | 1.1 | 1.3 | * |
| 1975 | 0.3 | * | * | * | * | 0.1 | * | 0.3 | 0.6 | 0.8 | 1.3 | 2.4 |
| 1974 | 0.3 | * | * | * | * | * | * | 0.3 | 0.5 | 1.0 | 1.4 | 1.8 |
| 1973 | 0.3 | * | * | * | * | * | 0.2 | 0.4 | 0.7 | 1.1 | 1.3 | 2.2 |
| 1972 | 0.3 | * | * | * | * | * | 0.2 | 0.3 | 0.8 | 1.0 | 1.1 | 2.2 |
| 1971 | 0.3 | * | * | * | * | * | * | 0.5 | 0.7 | 1.0 | 1.3 | * |
| 1970 | 0.4 | * | * | * | * | * | 0.3 | 0.5 | 0.8 | 1.5 | 1.8 | 2.1 |
| 1969 | 0.4 | * | * | * | * | * | 0.4 | 0.6 | 0.9 | 1.2 | 1.4 | 2.5 |
| 1968 | 0.5 | * | * | * | * | 0.2 | 0.3 | 0.8 | 1.1 | 1.4 | 1.8 | 3.8 |
| WHITE, BOTH SEXES | | | | | | | | | | | | |
| 1978 | 0.2 | * | * | * | * | * | * | 0.2 | 0.4 | 1.1 | 1.8 | 2.0 |
| 1977 | 0.2 | * | * | * | * | * | * | 0.2 | 0.5 | 0.9 | 1.8 | 1.7 |
| 1976 | 0.3 | * | * | * | * | * | * | 0.2 | 0.6 | 1.2 | 1.8 | 1.7 |
| 1975 | 0.3 | * | * | * | * | * | 0.1 | 0.2 | 0.7 | 1.1 | 2.1 | 2.6 |
| 1974 | 0.3 | * | * | * | * | * | * | 0.3 | 0.7 | 1.2 | 1.8 | 2.2 |
| 1973 | 0.3 | * | * | * | * | * | 0.1 | 0.4 | 0.8 | 1.4 | 2.3 | 3.0 |
| 1972 | 0.4 | * | * | * | * | * | 0.1 | 0.3 | 0.9 | 1.6 | 2.5 | 2.8 |
| 1971 | 0.4 | * | * | * | * | * | 0.1 | 0.4 | 0.9 | 1.6 | 2.2 | 3.1 |
| 1970 | 0.4 | * | * | * | * | * | 0.1 | 0.5 | 1.0 | 2.0 | 2.8 | 2.8 |
| 1969 | 0.4 | * | * | * | * | * | 0.1 | 0.5 | 1.1 | 1.8 | 2.6 | 3.9 |
| 1968 | 0.5 | * | * | * | * | * | 0.2 | 0.7 | 1.3 | 2.3 | 2.9 | 4.4 |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| TUBERCULOSIS, OTHER FORMS (013-019) | | | | | | | | | | | | |
| WHITE, MALE | | | | | | | | | | | | |
| 1978 | 0.3 | * | * | * | * | * | * | 0.3 | 0.5 | 1.7 | 2.7 | * |
| 1977 | 0.3 | * | * | * | * | * | * | 0.2 | 0.6 | 1.2 | 2.6 | * |
| 1976 | 0.3 | * | * | * | * | * | * | 0.3 | 0.7 | 1.6 | 3.0 | * |
| 1975 | 0.4 | * | * | * | * | * | * | 0.2 | 0.8 | 1.6 | 3.8 | 4.3 |
| 1974 | 0.3 | * | * | * | * | * | * | 0.4 | 0.9 | 1.7 | 2.7 | * |
| 1973 | 0.4 | * | * | * | * | * | * | 0.4 | 1.1 | 2.0 | 4.0 | 4.9 |
| 1972 | 0.5 | * | * | * | * | * | * | 0.3 | 1.0 | 2.5 | 4.8 | 5.5 |
| 1971 | 0.5 | * | * | * | * | * | * | 0.4 | 1.2 | 2.5 | 3.8 | 5.4 |
| 1970 | 0.6 | * | * | * | * | * | * | 0.7 | 1.4 | 2.8 | 4.5 | 4.3 |
| 1969 | 0.6 | * | * | * | * | * | * | 0.6 | 1.6 | 2.6 | 4.6 | 7.5 |
| 1968 | 0.7 | * | * | * | * | * | 0.2 | 0.8 | 1.7 | 3.9 | 4.8 | 6.3 |
| WHITE, FEMALE | | | | | | | | | | | | |
| 1978 | 0.2 | * | * | * | * | * | * | * | 0.4 | 0.7 | 1.3 | 1.5 |
| 1977 | 0.2 | * | * | * | * | * | * | * | 0.5 | 0.6 | 1.3 | * |
| 1976 | 0.2 | * | * | * | * | * | * | * | 0.4 | 0.9 | 1.2 | * |
| 1975 | 0.2 | * | * | * | * | * | * | 0.2 | 0.5 | 0.6 | 1.1 | 1.8 |
| 1974 | 0.2 | * | * | * | * | * | * | 0.2 | 0.4 | 0.8 | 1.3 | * |
| 1973 | 0.3 | * | * | * | * | * | * | 0.3 | 0.6 | 1.0 | 1.2 | 2.0 |
| 1972 | 0.2 | * | * | * | * | * | * | 0.2 | 0.7 | 0.8 | 1.0 | * |
| 1971 | 0.3 | * | * | * | * | * | * | 0.4 | 0.6 | 1.0 | 1.1 | * |
| 1970 | 0.3 | * | * | * | * | * | * | 0.4 | 0.6 | 1.4 | 1.7 | * |
| 1969 | 0.3 | * | * | * | * | * | * | 0.5 | 0.8 | 1.2 | 1.2 | * |
| 1968 | 0.4 | * | * | * | * | * | * | 0.5 | 1.0 | 1.1 | 1.6 | 3.4 |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| TUBERCULOSIS, OTHER FORMS (013-019) | | | | | | | | | | | | |
| ALL OTHER, BOTH SEXES | | | | | | | | | | | | |
| 1978 | 0.6 | * | * | * | * | * | * | 1.3 | 1.6 | 2.3 | 5.6 | * |
| 1977 | 0.7 | * | * | * | * | * | 0.9 | 1.3 | 1.7 | 3.3 | 5.3 | * |
| 1976 | 0.8 | * | * | * | * | * | 0.8 | 1.6 | 1.9 | 4.0 | 4.9 | * |
| 1975 | 0.8 | * | * | * | * | * | 0.8 | 1.5 | 2.1 | 3.6 | 5.2 | * |
| 1974 | 0.8 | * | * | * | * | * | 1.2 | 1.5 | 2.4 | 3.9 | 4.5 | * |
| 1973 | 0.9 | * | * | * | * | 0.7 | 1.0 | 1.8 | 2.1 | 4.5 | 5.3 | * |
| 1972 | 1.0 | * | * | * | * | * | 0.8 | 1.8 | 3.0 | 5.7 | 4.6 | * |
| 1971 | 1.0 | * | * | * | * | 0.7 | 1.4 | 2.3 | 2.3 | 4.0 | 5.5 | * |
| 1970 | 1.1 | * | * | * | * | 0.6 | 1.4 | 2.5 | 3.3 | 4.2 | 5.1 | * |
| 1969 | 1.3 | * | * | * | * | 0.8 | 2.3 | 2.9 | 3.5 | 3.9 | 5.8 | * |
| 1968 | 1.5 | * | * | * | * | 0.8 | 1.8 | 3.1 | 4.5 | 7.0 | 6.6 | * |
| ALL OTHER, MALE | | | | | | | | | | | | |
| 1978 | 0.8 | * | * | * | * | * | * | * | 2.2 | 3.5 | * | * |
| 1977 | 0.9 | * | * | * | * | * | * | 2.0 | 2.6 | 5.0 | 8.4 | * |
| 1976 | 1.0 | * | * | * | * | * | * | 2.4 | 2.9 | 5.4 | 8.6 | * |
| 1975 | 1.0 | * | * | * | * | * | * | 2.3 | 2.6 | 5.6 | * | * |
| 1974 | 1.1 | * | * | * | * | * | 1.7 | 2.2 | 3.2 | 5.7 | * | * |
| 1973 | 1.1 | * | * | * | * | * | * | 2.7 | 3.0 | 6.8 | 8.9 | * |
| 1972 | 1.3 | * | * | * | * | * | * | 3.3 | 4.3 | 9.0 | * | * |
| 1971 | 1.4 | * | * | * | * | * | 2.2 | 3.3 | 3.3 | 7.0 | * | * |
| 1970 | 1.4 | * | * | * | * | * | 2.0 | 3.4 | 4.8 | 5.6 | * | * |
| 1969 | 1.6 | * | * | * | * | * | 2.5 | 4.4 | 4.4 | 6.3 | * | * |
| 1968 | 1.9 | * | * | * | * | * | 2.5 | 3.7 | 6.6 | 10.4 | * | * |
| ALL OTHER, FEMALE | | | | | | | | | | | | |
| 1978 | 0.5 | * | * | * | * | * | * | * | * | * | * | * |
| 1977 | 0.5 | * | * | * | * | * | * | * | * | * | * | * |
| 1976 | 0.5 | * | * | * | * | * | * | * | * | 2.9 | * | * |
| 1975 | 0.7 | * | * | * | * | * | * | * | * | * | * | * |
| 1974 | 0.6 | * | * | * | * | * | * | * | * | * | * | * |
| 1973 | 0.7 | * | * | * | * | * | * | * | * | * | * | * |
| 1972 | 0.6 | * | * | * | * | * | * | * | * | 3.1 | * | * |
| 1971 | 0.7 | * | * | * | * | * | * | * | * | * | * | * |
| 1970 | 0.8 | * | * | * | * | * | * | 1.7 | * | 3.2 | * | * |
| 1969 | 1.0 | * | * | * | * | * | 2.1 | * | 2.7 | * | * | * |
| 1968 | 1.2 | * | * | * | * | * | * | 2.6 | 2.6 | 4.1 | * | * |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| TUBERCULOSIS, OTHER FORMS (013-019) | | | | | | | | | | | | |
| BLACK, BOTH SEXES | | | | | | | | | | | | |
| 1978 | 0.6 | * | * | * | * | * | * | 1.3 | 1.6 | 2.3 | 5.5 | * |
| 1977 | 0.7 | * | * | * | * | * | 1.1 | 1.4 | 1.7 | 3.2 | 4.6 | * |
| 1976 | 0.8 | * | * | * | * | * | 0.8 | 1.6 | 1.9 | 3.7 | 4.5 | * |
| 1975 | 0.8 | * | * | * | * | * | 0.9 | 1.7 | 2.2 | 3.3 | 3.6 | * |
| 1974 | 0.8 | * | * | * | * | * | 1.1 | 1.4 | 2.3 | 3.5 | 3.9 | * |
| 1973 | 0.9 | * | * | * | * | 0.8 | 1.1 | 1.8 | 2.1 | 4.2 | 4.8 | * |
| 1972 | 0.9 | * | * | * | * | * | 0.8 | 1.8 | 2.7 | 5.2 | 4.2 | * |
| 1971 | 1.0 | * | * | * | * | * | 1.3 | 2.3 | 2.3 | 4.1 | 4.7 | * |
| 1970 | 1.1 | * | * | * | * | * | 1.5 | 2.7 | 3.2 | 4.2 | * | * |
| 1969 | 1.2 | * | * | * | * | 0.9 | 2.3 | 2.8 | 3.2 | 3.9 | 5.6 | * |
| 1968 | 1.5 | * | * | * | * | 0.8 | 1.8 | 3.1 | 4.3 | 6.9 | 6.7 | * |
| BLACK, MALE | | | | | | | | | | | | |
| 1978 | 0.8 | * | * | * | * | * | * | * | 2.4 | 3.6 | * | * |
| 1977 | 1.0 | * | * | * | * | * | * | 2.2 | 2.8 | 5.0 | * | * |
| 1976 | 1.0 | * | * | * | * | * | * | 2.5 | 3.0 | 4.8 | * | * |
| 1975 | 0.9 | * | * | * | * | * | * | 2.5 | 2.9 | 4.8 | * | * |
| 1974 | 1.1 | * | * | * | * | * | * | 2.3 | 3.3 | 5.3 | * | * |
| 1973 | 1.1 | * | * | * | * | * | * | 2.6 | 3.0 | 6.3 | * | * |
| 1972 | 1.3 | * | * | * | * | * | * | 3.4 | 4.0 | 8.5 | * | * |
| 1971 | 1.4 | * | * | * | * | * | 2.2 | 3.2 | 3.3 | 7.3 | * | * |
| 1970 | 1.4 | * | * | * | * | * | 2.1 | 3.7 | 4.6 | 5.6 | * | * |
| 1969 | 1.6 | * | * | * | * | * | 2.6 | 4.4 | 4.4 | 6.6 | * | * |
| 1968 | 1.9 | * | * | * | * | * | 2.6 | 3.8 | 6.3 | 10.8 | * | * |
| BLACK, FEMALE | | | | | | | | | | | | |
| 1978 | 0.5 | * | * | * | * | * | * | * | * | * | * | * |
| 1977 | 0.5 | * | * | * | * | * | * | * | * | * | * | * |
| 1976 | 0.5 | * | * | * | * | * | * | * | * | 2.9 | * | * |
| 1975 | 0.6 | * | * | * | * | * | * | * | * | * | * | * |
| 1974 | 0.5 | * | * | * | * | * | * | * | * | * | * | * |
| 1973 | 0.7 | * | * | * | * | * | * | * | * | * | * | * |
| 1972 | 0.6 | * | * | * | * | * | * | * | * | * | * | * |
| 1971 | 0.6 | * | * | * | * | * | * | * | * | * | * | * |
| 1970 | 0.8 | * | * | * | * | * | * | 1.8 | * | * | * | * |
| 1969 | 0.9 | * | * | * | * | * | 2.1 | * | * | * | * | * |
| 1968 | 1.1 | * | * | * | * | * | * | 2.5 | 2.5 | 3.7 | * | * |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| WHOOPING COUGH (033) | | | | | | | | | | | | |
| BOTH SEXES | | | | | | | | | | | | |
| 1978 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1977 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1976 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1975 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1974 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1973 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1972 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1971 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1970 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1969 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1968 | 0.0 | 0.8 | * | * | * | * | * | * | * | * | * | * |
| MALE | | | | | | | | | | | | |
| 1978 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1977 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1976 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1975 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1974 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1973 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1972 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1971 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1970 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1969 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1968 | * | * | * | * | * | * | * | * | * | * | * | * |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| WHOOPING COUGH (033) | | | | | | | | | | | | |
| FEMALE | | | | | | | | | | | | |
| 1978 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1977 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1976 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1975 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1974 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1973 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1972 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1971 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1970 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1969 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1968 | * | * | * | * | * | * | * | * | * | * | * | * |
| WHITE, BOTH SEXES | | | | | | | | | | | | |
| 1978 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1977 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1976 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1975 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1974 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1973 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1972 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1971 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1970 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1969 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1968 | * | * | * | * | * | * | * | * | * | * | * | * |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| WHOOPING COUGH (033) | | | | | | | | | | | | |
| WHITE, MALE | | | | | | | | | | | | |
| 1978 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1977 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1976 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1975 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1974 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1973 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1972 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1971 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1970 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1969 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1968 | * | * | * | * | * | * | * | * | * | * | * | * |
| WHITE, FEMALE | | | | | | | | | | | | |
| 1978 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1977 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1976 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1975 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1974 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1973 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1972 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1971 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1970 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1969 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1968 | * | * | * | * | * | * | * | * | * | * | * | * |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| WHOOPING COUGH (033) | | | | | | | | | | | | |
| ALL OTHER, BOTH SEXES | | | | | | | | | | | | |
| 1978 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1977 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1976 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1975 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1974 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1973 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1972 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1971 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1970 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1969 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1968 | * | * | * | * | * | * | * | * | * | * | * | * |
| ALL OTHER, MALE | | | | | | | | | | | | |
| 1978 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1977 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1976 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1975 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1974 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1973 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1972 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1971 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1970 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1969 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1968 | * | * | * | * | * | * | * | * | * | * | * | * |
| ALL OTHER, FEMALE | | | | | | | | | | | | |
| 1978 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1977 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1976 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1975 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1974 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1973 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1972 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1971 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1970 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1969 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1968 | * | * | * | * | * | * | * | * | * | * | * | * |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| WHOOPING COUGH (033) | | | | | | | | | | | | |
| BLACK, BOTH SEXES | | | | | | | | | | | | |
| 1978 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1977 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1976 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1975 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1974 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1973 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1972 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1971 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1970 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1969 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1968 | * | * | * | * | * | * | * | * | * | * | * | * |
| BLACK, MALE | | | | | | | | | | | | |
| 1978 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1977 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1976 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1975 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1974 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1973 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1972 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1971 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1970 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1969 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1968 | * | * | * | * | * | * | * | * | * | * | * | * |
| BLACK, FEMALE | | | | | | | | | | | | |
| 1978 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1977 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1976 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1975 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1974 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1973 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1972 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1971 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1970 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1969 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1968 | * | * | * | * | * | * | * | * | * | * | * | * |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| STREPTOCOCCAL SORE THROAT AND SCARLET FEVER (034) | | | | | | | | | | | | |
| BOTH SEXES | | | | | | | | | | | | |
| 1978 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1977 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1976 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1975 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1974 | 0.0 | * | * | * | * | * | * | * | * | * | * | * |
| 1973 | 0.0 | * | * | * | * | * | * | * | * | * | * | * |
| 1972 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1971 | 0.0 | * | * | * | * | * | * | * | * | * | * | * |
| 1970 | 0.0 | * | * | * | * | * | * | * | * | * | * | * |
| 1969 | 0.0 | * | * | * | * | * | * | * | * | * | * | * |
| 1968 | 0.0 | * | * | * | * | * | * | * | * | * | * | * |
| MALE | | | | | | | | | | | | |
| 1978 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1977 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1976 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1975 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1974 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1973 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1972 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1971 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1970 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1969 | 0.0 | * | * | * | * | * | * | * | * | * | * | * |
| 1968 | * | * | * | * | * | * | * | * | * | * | * | * |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| STREPTOCOCCAL SORE THROAT AND SCARLET FEVER (034) | | | | | | | | | | | | |
| FEMALE | | | | | | | | | | | | |
| 1978 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1977 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1976 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1975 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1974 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1973 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1972 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1971 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1970 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1969 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1968 | * | * | * | * | * | * | * | * | * | * | * | * |
| WHITE, BOTH SEXES | | | | | | | | | | | | |
| 1978 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1977 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1976 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1975 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1974 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1973 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1972 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1971 | 0.0 | * | * | * | * | * | * | * | * | * | * | * |
| 1970 | 0.0 | * | * | * | * | * | * | * | * | * | * | * |
| 1969 | 0.0 | * | * | * | * | * | * | * | * | * | * | * |
| 1968 | 0.0 | * | * | * | * | * | * | * | * | * | * | * |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| STREPTOCOCCAL SORE THROAT AND SCARLET FEVER (034) | | | | | | | | | | | | |
| WHITE, MALE | | | | | | | | | | | | |
| 1978 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1977 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1976 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1975 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1974 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1973 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1972 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1971 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1970 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1969 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1968 | * | * | * | * | * | * | * | * | * | * | * | * |
| WHITE, FEMALE | | | | | | | | | | | | |
| 1978 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1977 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1976 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1975 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1974 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1973 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1972 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1971 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1970 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1969 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1968 | * | * | * | * | * | * | * | * | * | * | * | * |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| STREPTOCOCCAL SORE THROAT AND SCARLET FEVER (034) | | | | | | | | | | | | |
| ALL OTHER, BOTH SEXES | | | | | | | | | | | | |
| 1978 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1977 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1976 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1975 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1974 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1973 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1972 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1971 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1970 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1969 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1968 | * | * | * | * | * | * | * | * | * | * | * | * |
| ALL OTHER, MALE | | | | | | | | | | | | |
| 1978 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1977 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1976 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1975 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1974 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1973 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1972 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1971 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1970 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1969 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1968 | * | * | * | * | * | * | * | * | * | * | * | * |
| ALL OTHER, FEMALE | | | | | | | | | | | | |
| 1978 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1977 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1976 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1975 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1974 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1973 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1972 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1971 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1970 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1969 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1968 | * | * | * | * | * | * | * | * | * | * | * | * |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| STREPTOCOCCAL SORE THROAT AND SCARLET FEVER (034) | | | | | | | | | | | | |
| BLACK, BOTH SEXES | | | | | | | | | | | | |
| 1978 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1977 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1976 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1975 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1974 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1973 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1972 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1971 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1970 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1969 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1968 | * | * | * | * | * | * | * | * | * | * | * | * |
| BLACK, MALE | | | | | | | | | | | | |
| 1978 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1977 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1976 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1975 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1974 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1973 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1972 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1971 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1970 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1969 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1968 | * | * | * | * | * | * | * | * | * | * | * | * |
| BLACK, FEMALE | | | | | | | | | | | | |
| 1978 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1977 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1976 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1975 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1974 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1973 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1972 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1971 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1970 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1969 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1968 | * | * | * | * | * | * | * | * | * | * | * | * |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| MENINGOCOCCAL INFECTIONS (036) | | | | | | | | | | | | |
| BOTH SEXES | | | | | | | | | | | | |
| 1978 | 0.2 | 2.9 | 0.9 | 0.1 | 0.1 | * | * | * | * | * | * | * |
| 1977 | 0.2 | 2.9 | 0.7 | 0.1 | 0.1 | * | * | * | 0.1 | 0.2 | * | * |
| 1976 | 0.2 | 2.7 | 0.7 | 0.1 | 0.1 | * | * | * | * | 0.2 | * | * |
| 1975 | 0.1 | 2.3 | 0.6 | 0.1 | 0.1 | * | * | * | 0.1 | * | * | * |
| 1974 | 0.1 | 1.5 | 0.6 | 0.1 | 0.1 | * | * | * | * | * | * | * |
| 1973 | 0.2 | 2.9 | 0.5 | 0.1 | 0.1 | * | * | * | * | * | * | * |
| 1972 | 0.2 | 2.9 | 0.7 | 0.1 | 0.1 | * | * | * | 0.1 | * | * | * |
| 1971 | 0.2 | 3.5 | 0.7 | 0.2 | 0.3 | * | * | 0.1 | 0.1 | 0.2 | * | * |
| 1970 | 0.3 | 3.6 | 1.0 | 0.2 | 0.3 | * | 0.1 | 0.1 | 0.2 | * | * | * |
| 1969 | 0.4 | 4.2 | 1.4 | 0.3 | 0.4 | 0.1 | 0.1 | 0.1 | 0.2 | 0.2 | * | * |
| 1968 | 0.4 | 4.7 | 1.5 | 0.3 | 0.3 | 0.1 | 0.1 | 0.2 | 0.2 | * | * | * |
| MALE | | | | | | | | | | | | |
| 1978 | 0.2 | 3.3 | 1.0 | 0.1 | 0.1 | * | * | * | * | * | * | * |
| 1977 | 0.2 | 3.6 | 0.7 | * | 0.1 | * | * | * | * | * | * | * |
| 1976 | 0.2 | 3.5 | 0.8 | * | 0.1 | * | * | * | * | * | * | * |
| 1975 | 0.2 | 2.4 | 0.8 | 0.1 | * | * | * | * | * | * | * | * |
| 1974 | 0.2 | 1.8 | 0.7 | 0.2 | 0.1 | * | * | * | * | * | * | * |
| 1973 | 0.2 | 3.4 | 0.5 | 0.1 | 0.1 | * | * | * | * | * | * | * |
| 1972 | 0.2 | 3.0 | 0.8 | * | 0.1 | * | * | * | * | * | * | * |
| 1971 | 0.3 | 3.6 | 0.8 | 0.2 | 0.3 | * | * | * | * | * | * | * |
| 1970 | 0.3 | 4.1 | 1.2 | 0.2 | 0.3 | * | * | * | * | * | * | * |
| 1969 | 0.4 | 5.3 | 1.4 | 0.3 | 0.5 | * | * | * | * | * | * | * |
| 1968 | 0.4 | 5.5 | 1.7 | 0.4 | 0.4 | * | * | * | * | * | * | * |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| MENINGOCOCCAL INFECTIONS (036) | | | | | | | | | | | | |
| FEMALE | | | | | | | | | | | | |
| 1978 | 0.2 | 2.5 | 0.8 | 0.1 | 0.1 | * | * | * | * | * | * | * |
| 1977 | 0.1 | 2.1 | 0.8 | * | * | * | * | * | * | * | * | * |
| 1976 | 0.1 | 2.0 | 0.6 | * | 0.1 | * | * | * | * | * | * | * |
| 1975 | 0.1 | 2.3 | 0.5 | * | * | * | * | * | * | * | * | * |
| 1974 | 0.1 | * | 0.6 | 0.1 | * | * | * | * | * | * | * | * |
| 1973 | 0.1 | 2.3 | 0.5 | 0.1 | * | * | * | * | * | * | * | * |
| 1972 | 0.1 | 2.8 | 0.6 | * | * | * | * | * | * | * | * | * |
| 1971 | 0.2 | 3.4 | 0.7 | 0.2 | 0.2 | * | * | * | * | * | * | * |
| 1970 | 0.2 | 3.0 | 0.9 | 0.2 | 0.2 | * | * | * | * | * | * | * |
| 1969 | 0.3 | 2.9 | 1.4 | 0.3 | 0.3 | * | * | * | * | * | * | * |
| 1968 | 0.3 | 3.8 | 1.3 | 0.2 | 0.2 | * | * | 0.2 | * | * | * | * |
| WHITE, BOTH SEXES | | | | | | | | | | | | |
| 1978 | 0.2 | 2.9 | 0.9 | 0.1 | 0.1 | * | * | * | * | * | * | * |
| 1977 | 0.1 | 2.8 | 0.7 | * | 0.1 | * | * | * | * | 0.2 | * | * |
| 1976 | 0.1 | 2.8 | 0.7 | * | 0.1 | * | * | * | * | * | * | * |
| 1975 | 0.1 | 2.4 | 0.7 | 0.1 | 0.1 | * | * | * | * | * | * | * |
| 1974 | 0.1 | 1.3 | 0.6 | 0.1 | 0.1 | * | * | * | * | * | * | * |
| 1973 | 0.1 | 2.8 | 0.6 | 0.1 | 0.1 | * | * | * | * | * | * | * |
| 1972 | 0.2 | 3.0 | 0.6 | * | 0.1 | * | * | * | 0.1 | * | * | * |
| 1971 | 0.2 | 3.5 | 0.7 | 0.2 | 0.2 | * | * | 0.1 | * | * | * | * |
| 1970 | 0.2 | 3.4 | 1.0 | 0.1 | 0.2 | * | * | 0.1 | * | * | * | * |
| 1969 | 0.3 | 4.1 | 1.2 | 0.2 | 0.4 | * | * | 0.1 | 0.2 | 0.2 | * | * |
| 1968 | 0.3 | 4.5 | 1.2 | 0.2 | 0.2 | * | * | 0.1 | 0.2 | * | * | * |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| MENINGOCOCCAL INFECTIONS (036) | | | | | | | | | | | | |
| WHITE, MALE | | | | | | | | | | | | |
| 1978 | 0.2 | 3.7 | 1.0 | 0.1 | * | * | * | * | * | * | * | * |
| 1977 | 0.1 | 3.5 | 0.6 | * | * | * | * | * | * | * | * | * |
| 1976 | 0.2 | 3.6 | 0.8 | * | * | * | * | * | * | * | * | * |
| 1975 | 0.2 | 2.6 | 0.8 | 0.1 | * | * | * | * | * | * | * | * |
| 1974 | 0.1 | * | 0.6 | 0.1 | * | * | * | * | * | * | * | * |
| 1973 | 0.2 | 3.2 | 0.6 | 0.1 | * | * | * | * | * | * | * | * |
| 1972 | 0.2 | 3.3 | 0.6 | * | 0.1 | * | * | * | * | * | * | * |
| 1971 | 0.2 | 3.5 | 0.7 | 0.2 | 0.3 | * | * | * | * | * | * | * |
| 1970 | 0.3 | 3.9 | 1.2 | 0.1 | 0.3 | * | * | * | * | * | * | * |
| 1969 | 0.4 | 5.5 | 1.1 | 0.2 | 0.5 | * | * | * | * | * | * | * |
| 1968 | 0.4 | 5.1 | 1.5 | 0.3 | 0.3 | * | * | * | * | * | * | * |
| WHITE, FEMALE | | | | | | | | | | | | |
| 1978 | 0.2 | 2.1 | 0.8 | * | * | * | * | * | * | * | * | * |
| 1977 | 0.1 | 2.0 | 0.7 | * | * | * | * | * | * | * | * | * |
| 1976 | 0.1 | 2.0 | 0.6 | * | * | * | * | * | * | * | * | * |
| 1975 | 0.1 | 2.1 | 0.5 | * | * | * | * | * | * | * | * | * |
| 1974 | 0.1 | * | 0.5 | * | * | * | * | * | * | * | * | * |
| 1973 | 0.1 | 2.3 | 0.5 | * | * | * | * | * | * | * | * | * |
| 1972 | 0.1 | 2.8 | 0.7 | * | * | * | * | * | * | * | * | * |
| 1971 | 0.2 | 3.4 | 0.7 | 0.2 | 0.2 | * | * | * | * | * | * | * |
| 1970 | 0.2 | 3.0 | 0.8 | 0.1 | 0.2 | * | * | * | * | * | * | * |
| 1969 | 0.3 | 2.8 | 1.2 | 0.3 | 0.2 | * | * | * | * | * | * | * |
| 1968 | 0.3 | 3.9 | 0.9 | 0.2 | 0.2 | * | * | * | * | * | * | * |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| MENINGOCOCCAL INFECTIONS (036) | | | | | | | | | | | | |
| ALL OTHER, BOTH SEXES | | | | | | | | | | | | |
| 1978 | 0.3 | * | * | * | * | * | * | * | * | * | * | * |
| 1977 | 0.2 | * | 1.0 | * | * | * | * | * | * | * | * | * |
| 1976 | 0.2 | * | * | * | * | * | * | * | * | * | * | * |
| 1975 | 0.2 | * | * | * | * | * | * | * | * | * | * | * |
| 1974 | 0.3 | * | * | * | * | * | * | * | * | * | * | * |
| 1973 | 0.2 | * | * | * | * | * | * | * | * | * | * | * |
| 1972 | 0.2 | * | 1.0 | * | * | * | * | * | * | * | * | * |
| 1971 | 0.4 | 3.9 | * | * | * | * | * | * | * | * | * | * |
| 1970 | 0.5 | 4.2 | 1.1 | 0.4 | 0.4 | * | * | * | * | * | * | * |
| 1969 | 0.8 | 4.3 | 2.6 | 0.5 | 0.8 | * | * | * | * | * | * | * |
| 1968 | 0.8 | 5.5 | 2.8 | 0.7 | 0.6 | * | * | * | * | * | * | * |
| ALL OTHER, MALE | | | | | | | | | | | | |
| 1978 | 0.3 | * | * | * | * | * | * | * | * | * | * | * |
| 1977 | 0.3 | * | * | * | * | * | * | * | * | * | * | * |
| 1976 | 0.3 | * | * | * | * | * | * | * | * | * | * | * |
| 1975 | 0.2 | * | * | * | * | * | * | * | * | * | * | * |
| 1974 | 0.4 | * | * | * | * | * | * | * | * | * | * | * |
| 1973 | 0.3 | * | * | * | * | * | * | * | * | * | * | * |
| 1972 | 0.3 | * | 1.8 | * | * | * | * | * | * | * | * | * |
| 1971 | 0.5 | * | * | * | * | * | * | * | * | * | * | * |
| 1970 | 0.6 | * | * | * | * | * | * | * | * | * | * | * |
| 1969 | 0.9 | * | 3.0 | * | 1.1 | * | * | * | * | * | * | * |
| 1968 | 0.9 | 7.9 | 2.5 | 0.7 | * | * | * | * | * | * | * | * |
| ALL OTHER, FEMALE | | | | | | | | | | | | |
| 1978 | 0.3 | * | * | * | * | * | * | * | * | * | * | * |
| 1977 | 0.2 | * | * | * | * | * | * | * | * | * | * | * |
| 1976 | 0.2 | * | * | * | * | * | * | * | * | * | * | * |
| 1975 | 0.2 | * | * | * | * | * | * | * | * | * | * | * |
| 1974 | 0.2 | * | * | * | * | * | * | * | * | * | * | * |
| 1973 | 0.2 | * | * | * | * | * | * | * | * | * | * | * |
| 1972 | 0.2 | * | * | * | * | * | * | * | * | * | * | * |
| 1971 | 0.3 | * | * | * | * | * | * | * | * | * | * | * |
| 1970 | 0.4 | * | * | * | * | * | * | * | * | * | * | * |
| 1969 | 0.6 | * | 2.3 | * | * | * | * | * | * | * | * | * |
| 1968 | 0.7 | * | 3.2 | * | * | * | * | * | * | * | * | * |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| MENINGOCOCCAL INFECTIONS (036) | | | | | | | | | | | | |
| BLACK, BOTH SEXES | | | | | | | | | | | | |
| 1978 | 0.3 | * | * | * | * | * | * | * | * | * | * | * |
| 1977 | 0.3 | * | * | * | * | * | * | * | * | * | * | * |
| 1976 | 0.2 | * | * | * | * | * | * | * | * | * | * | * |
| 1975 | 0.3 | * | * | * | * | * | * | * | * | * | * | * |
| 1974 | 0.3 | * | * | * | * | * | * | * | * | * | * | * |
| 1973 | 0.2 | * | * | * | * | * | * | * | * | * | * | * |
| 1972 | 0.2 | * | 1.0 | * | * | * | * | * | * | * | * | * |
| 1971 | 0.4 | * | * | * | * | * | * | * | * | * | * | * |
| 1970 | 0.5 | 4.3 | 1.1 | 0.4 | * | * | * | * | * | * | * | * |
| 1969 | 0.8 | 4.4 | 2.8 | 0.5 | 0.8 | * | * | * | * | * | * | * |
| 1968 | 0.8 | 5.7 | 2.8 | 0.7 | 0.6 | * | * | * | * | * | * | * |
| BLACK, MALE | | | | | | | | | | | | |
| 1978 | 0.3 | * | * | * | * | * | * | * | * | * | * | * |
| 1977 | 0.3 | * | * | * | * | * | * | * | * | * | * | * |
| 1976 | 0.3 | * | * | * | * | * | * | * | * | * | * | * |
| 1975 | 0.3 | * | * | * | * | * | * | * | * | * | * | * |
| 1974 | 0.4 | * | * | * | * | * | * | * | * | * | * | * |
| 1973 | 0.3 | * | * | * | * | * | * | * | * | * | * | * |
| 1972 | 0.3 | * | * | * | * | * | * | * | * | * | * | * |
| 1971 | 0.5 | * | * | * | * | * | * | * | * | * | * | * |
| 1970 | 0.6 | * | * | * | * | * | * | * | * | * | * | * |
| 1969 | 1.0 | * | 3.2 | * | 1.3 | * | * | * | * | * | * | * |
| 1968 | 0.9 | * | 2.6 | 0.8 | * | * | * | * | * | * | * | * |
| BLACK, FEMALE | | | | | | | | | | | | |
| 1978 | 0.3 | * | * | * | * | * | * | * | * | * | * | * |
| 1977 | 0.2 | * | * | * | * | * | * | * | * | * | * | * |
| 1976 | 0.2 | * | * | * | * | * | * | * | * | * | * | * |
| 1975 | 0.2 | * | * | * | * | * | * | * | * | * | * | * |
| 1974 | 0.2 | * | * | * | * | * | * | * | * | * | * | * |
| 1973 | 0.2 | * | * | * | * | * | * | * | * | * | * | * |
| 1972 | 0.2 | * | * | * | * | * | * | * | * | * | * | * |
| 1971 | 0.3 | * | * | * | * | * | * | * | * | * | * | * |
| 1970 | 0.4 | * | * | * | * | * | * | * | * | * | * | * |
| 1969 | 0.7 | * | 2.5 | * | * | * | * | * | * | * | * | * |
| 1968 | 0.7 | * | 3.1 | * | * | * | * | * | * | * | * | * |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| SEPTICEMIA 038 | | | | | | | | | | | | |
| BOTH SEXES | | | | | | | | | | | | |
| 1978 | 3.5 | 32.9 | 0.5 | 0.1 | 0.2 | 0.4 | 0.9 | 2.0 | 4.8 | 10.3 | 26.2 | 57.4 |
| 1977 | 3.2 | 33.2 | 0.6 | 0.1 | 0.2 | 0.3 | 0.8 | 1.9 | 4.4 | 9.5 | 23.8 | 51.6 |
| 1976 | 2.9 | 34.6 | 0.6 | 0.1 | 0.2 | 0.4 | 0.8 | 1.8 | 3.9 | 8.5 | 21.6 | 43.1 |
| 1975 | 2.6 | 30.8 | 0.3 | 0.1 | 0.2 | 0.3 | 0.7 | 1.7 | 3.5 | 7.9 | 18.9 | 36.7 |
| 1974 | 2.5 | 29.7 | 0.5 | 0.2 | 0.2 | 0.4 | 0.7 | 1.6 | 3.3 | 7.9 | 17.3 | 33.6 |
| 1973 | 2.2 | 25.7 | 0.6 | 0.1 | 0.2 | 0.3 | 0.6 | 1.5 | 3.0 | 6.9 | 15.2 | 28.1 |
| 1972 | 2.0 | 24.8 | 0.4 | 0.1 | 0.2 | 0.4 | 0.6 | 1.4 | 2.8 | 6.0 | 14.1 | 26.1 |
| 1971 | 1.9 | 23.0 | 0.6 | 0.2 | 0.2 | 0.3 | 0.6 | 1.1 | 2.9 | 5.7 | 11.8 | 23.4 |
| 1970 | 1.7 | 24.8 | 0.5 | 0.1 | 0.2 | 0.3 | 0.6 | 1.2 | 2.6 | 4.9 | 10.5 | 17.0 |
| 1969 | 1.5 | 22.2 | 0.5 | 0.1 | 0.2 | 0.2 | 0.5 | 1.1 | 2.1 | 4.1 | 8.8 | 16.4 |
| 1968 | 1.5 | 23.8 | 0.7 | 0.1 | 0.2 | 0.3 | 0.5 | 0.9 | 2.0 | 4.2 | 8.4 | 17.0 |
| MALE | | | | | | | | | | | | |
| 1978 | 3.8 | 37.9 | 0.6 | 0.1 | 0.2 | 0.4 | 1.0 | 2.5 | 5.7 | 12.9 | 34.8 | 74.4 |
| 1977 | 3.5 | 38.9 | 0.7 | 0.1 | 0.2 | 0.3 | 1.0 | 2.2 | 5.4 | 11.8 | 32.1 | 68.7 |
| 1976 | 3.2 | 37.9 | 0.5 | 0.2 | 0.2 | 0.4 | 0.9 | 1.9 | 4.8 | 10.9 | 28.1 | 56.8 |
| 1975 | 2.9 | 35.2 | 0.3 | 0.1 | 0.2 | 0.4 | 0.7 | 2.0 | 4.3 | 9.8 | 26.1 | 46.5 |
| 1974 | 2.7 | 33.5 | 0.6 | 0.2 | 0.2 | 0.4 | 0.8 | 1.8 | 4.1 | 10.3 | 24.3 | 43.9 |
| 1973 | 2.4 | 29.3 | 0.6 | 0.1 | 0.2 | 0.3 | 0.7 | 1.8 | 3.6 | 9.0 | 20.0 | 33.3 |
| 1972 | 2.3 | 26.5 | 0.5 | 0.1 | 0.2 | 0.4 | 0.7 | 1.6 | 3.5 | 8.0 | 19.4 | 39.8 |
| 1971 | 2.1 | 26.2 | 0.5 | 0.2 | 0.2 | 0.3 | 0.7 | 1.4 | 3.5 | 7.7 | 16.0 | 31.2 |
| 1970 | 2.0 | 26.3 | 0.5 | 0.2 | 0.2 | 0.3 | 0.6 | 1.5 | 3.3 | 6.3 | 14.4 | 21.2 |
| 1969 | 1.7 | 25.1 | 0.5 | 0.1 | 0.2 | 0.2 | 0.6 | 1.3 | 2.7 | 4.9 | 11.7 | 20.3 |
| 1968 | 1.7 | 26.7 | 0.6 | 0.2 | 0.2 | 0.3 | 0.5 | 1.0 | 2.5 | 5.5 | 10.3 | 23.9 |

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| SEPTICEMIA 038 | | | | | | | | | | | | |
| FEMALE | | | | | | | | | | | | |
| 1978 | 3.3 | 27.6 | 0.5 | 0.1 | 0.2 | 0.4 | 0.8 | 1.5 | 4.0 | 8.3 | 21.1 | 49.7 |
| 1977 | 3.0 | 27.1 | 0.4 | 0.1 | 0.2 | 0.4 | 0.7 | 1.6 | 3.5 | 7.7 | 18.9 | 43.7 |
| 1976 | 2.7 | 31.1 | 0.6 | 0.1 | 0.2 | 0.4 | 0.7 | 1.6 | 3.1 | 6.7 | 17.7 | 36.7 |
| 1975 | 2.3 | 26.3 | 0.3 | * | 0.2 | 0.3 | 0.6 | 1.3 | 2.8 | 6.5 | 14.5 | 31.9 |
| 1974 | 2.2 | 25.7 | 0.4 | 0.1 | 0.2 | 0.4 | 0.6 | 1.4 | 2.6 | 6.0 | 13.0 | 28.5 |
| 1973 | 2.0 | 22.0 | 0.5 | 0.1 | 0.2 | 0.3 | 0.6 | 1.3 | 2.5 | 5.4 | 12.1 | 25.5 |
| 1972 | 1.8 | 23.0 | 0.4 | 0.2 | 0.2 | 0.4 | 0.5 | 1.1 | 2.1 | 4.5 | 10.7 | 19.0 |
| 1971 | 1.6 | 19.6 | 0.7 | 0.2 | 0.2 | 0.3 | 0.5 | 0.9 | 2.4 | 4.3 | 9.1 | 19.3 |
| 1970 | 1.5 | 23.3 | 0.5 | * | 0.2 | 0.3 | 0.6 | 0.9 | 1.9 | 3.8 | 7.9 | 14.7 |
| 1969 | 1.3 | 19.2 | 0.5 | 0.1 | 0.2 | 0.2 | 0.4 | 0.9 | 1.6 | 3.5 | 6.9 | 14.4 |
| 1968 | 1.3 | 20.7 | 0.7 | 0.1 | 0.2 | 0.3 | 0.5 | 0.8 | 1.6 | 3.2 | 7.1 | 13.3 |
| WHITE, BOTH SEXES | | | | | | | | | | | | |
| 1978 | 3.2 | 27.6 | 0.5 | 0.1 | 0.2 | 0.3 | 0.6 | 1.5 | 4.0 | 8.9 | 23.9 | 54.5 |
| 1977 | 2.9 | 27.0 | 0.5 | 0.1 | 0.2 | 0.3 | 0.6 | 1.5 | 3.6 | 8.3 | 21.3 | 48.9 |
| 1976 | 2.6 | 28.6 | 0.5 | 0.1 | 0.2 | 0.3 | 0.6 | 1.3 | 3.2 | 7.4 | 19.8 | 39.8 |
| 1975 | 2.3 | 25.2 | 0.3 | 0.1 | 0.2 | 0.3 | 0.5 | 1.3 | 3.0 | 6.7 | 18.0 | 34.6 |
| 1974 | 2.2 | 24.3 | 0.4 | 0.1 | 0.2 | 0.4 | 0.5 | 1.2 | 2.8 | 6.9 | 16.7 | 31.6 |
| 1973 | 1.9 | 21.4 | 0.5 | 0.1 | 0.2 | 0.2 | 0.4 | 1.2 | 2.6 | 6.1 | 14.2 | 26.6 |
| 1972 | 1.8 | 19.6 | 0.4 | 0.1 | 0.2 | 0.3 | 0.4 | 1.1 | 2.2 | 5.1 | 13.2 | 26.1 |
| 1971 | 1.6 | 17.1 | 0.4 | 0.1 | 0.1 | 0.2 | 0.4 | 0.9 | 2.4 | 5.1 | 11.4 | 22.7 |
| 1970 | 1.5 | 20.3 | 0.5 | 0.1 | 0.2 | 0.3 | 0.4 | 1.0 | 2.2 | 4.3 | 10.1 | 16.9 |
| 1969 | 1.3 | 16.5 | 0.5 | 0.1 | 0.1 | 0.1 | 0.4 | 0.9 | 1.8 | 3.7 | 8.2 | 16.4 |
| 1968 | 1.3 | 18.2 | 0.6 | 0.1 | 0.1 | 0.2 | 0.4 | 0.7 | 1.8 | 3.7 | 8.2 | 16.5 |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| SEPTICEMIA 038 | | | | | | | | | | | | |
| WHITE, MALE | | | | | | | | | | | | |
| 1978 | 3.4 | 33.0 | 0.5 | * | 0.2 | 0.3 | 0.7 | 1.8 | 4.8 | 11.2 | 31.8 | 71.4 |
| 1977 | 3.1 | 32.7 | 0.6 | * | 0.2 | 0.3 | 0.7 | 1.7 | 4.5 | 10.5 | 28.9 | 64.3 |
| 1976 | 2.8 | 32.4 | 0.4 | 0.1 | 0.2 | 0.3 | 0.7 | 1.5 | 4.1 | 9.6 | 26.0 | 53.0 |
| 1975 | 2.5 | 29.7 | * | 0.1 | 0.2 | 0.3 | 0.5 | 1.6 | 3.7 | 8.2 | 24.7 | 45.7 |
| 1974 | 2.4 | 27.8 | 0.4 | * | 0.1 | 0.3 | 0.5 | 1.4 | 3.5 | 9.2 | 23.3 | 41.7 |
| 1973 | 2.1 | 24.4 | 0.5 | 0.1 | 0.2 | 0.2 | 0.4 | 1.4 | 3.2 | 8.0 | 18.6 | 31.0 |
| 1972 | 2.0 | 21.6 | 0.6 | * | 0.2 | 0.3 | 0.5 | 1.2 | 2.8 | 7.0 | 18.2 | 40.0 |
| 1971 | 1.9 | 19.2 | 0.4 | 0.2 | 0.2 | 0.2 | 0.5 | 1.2 | 3.0 | 6.7 | 15.5 | 30.9 |
| 1970 | 1.7 | 22.6 | 0.5 | 0.2 | 0.2 | 0.3 | 0.4 | 1.2 | 2.8 | 5.6 | 13.6 | 20.1 |
| 1969 | 1.4 | 18.5 | 0.4 | 0.1 | 0.1 | * | 0.5 | 1.0 | 2.3 | 4.3 | 11.0 | 20.0 |
| 1968 | 1.5 | 20.6 | 0.6 | 0.1 | 0.2 | 0.2 | 0.4 | 0.8 | 2.2 | 4.9 | 9.9 | 23.3 |
| WHITE, FEMALE | | | | | | | | | | | | |
| 1978 | 3.0 | 22.0 | 0.4 | * | 0.2 | 0.3 | 0.5 | 1.1 | 3.4 | 7.1 | 19.3 | 47.0 |
| 1977 | 2.7 | 21.0 | 0.4 | * | 0.2 | 0.3 | 0.5 | 1.3 | 2.8 | 6.6 | 16.9 | 41.9 |
| 1976 | 2.4 | 24.7 | 0.5 | * | 0.2 | 0.3 | 0.6 | 1.2 | 2.3 | 5.8 | 16.1 | 33.7 |
| 1975 | 2.1 | 20.4 | * | * | 0.2 | 0.2 | 0.4 | 1.0 | 2.4 | 5.6 | 14.0 | 29.3 |
| 1974 | 2.0 | 20.6 | * | 0.2 | 0.2 | 0.4 | 0.4 | 1.1 | 2.1 | 5.3 | 12.7 | 26.7 |
| 1973 | 1.8 | 18.3 | 0.4 | * | 0.2 | 0.2 | 0.4 | 0.9 | 2.1 | 4.7 | 11.5 | 24.4 |
| 1972 | 1.5 | 17.5 | * | 0.2 | 0.2 | 0.3 | 0.4 | 0.9 | 1.7 | 3.7 | 10.0 | 19.1 |
| 1971 | 1.4 | 14.8 | 0.5 | 0.1 | 0.1 | 0.3 | 0.4 | 0.7 | 1.9 | 3.9 | 8.8 | 18.5 |
| 1970 | 1.3 | 18.0 | 0.5 | * | * | 0.3 | 0.4 | 0.7 | 1.6 | 3.3 | 7.8 | 15.1 |
| 1969 | 1.2 | 14.4 | 0.6 | * | 0.1 | * | 0.3 | 0.8 | 1.3 | 3.1 | 6.4 | 14.6 |
| 1968 | 1.2 | 15.7 | 0.6 | * | * | 0.2 | 0.4 | 0.6 | 1.4 | 2.8 | 7.0 | 12.9 |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| SEPTICEMIA 038 | | | | | | | | | | | | |
| ALL OTHER, BOTH SEXES | | | | | | | | | | | | |
| 1978 | 5.7 | 58.1 | * | * | 0.3 | 1.0 | 2.5 | 6.1 | 11.5 | 23.2 | 49.7 | 89.9 |
| 1977 | 5.5 | 62.9 | * | * | * | 0.8 | 2.3 | 5.4 | 11.2 | 20.5 | 49.5 | 83.1 |
| 1976 | 5.2 | 63.6 | 1.0 | * | 0.4 | 1.1 | 2.3 | 5.3 | 10.8 | 19.2 | 40.4 | 80.4 |
| 1975 | 4.4 | 59.0 | * | * | * | 0.9 | 2.2 | 4.6 | 8.4 | 19.0 | 28.4 | 61.0 |
| 1974 | 4.2 | 56.5 | 1.0 | * | 0.4 | 1.0 | 1.9 | 4.4 | 8.5 | 16.9 | 24.2 | 56.3 |
| 1973 | 3.7 | 47.1 | 1.0 | * | 0.4 | 1.0 | 2.0 | 4.4 | 7.0 | 14.8 | 25.6 | 46.5 |
| 1972 | 3.8 | 51.4 | * | * | 0.5 | 1.3 | 1.7 | 3.7 | 8.0 | 14.8 | 24.7 | 26.0 |
| 1971 | 3.6 | 54.4 | 1.2 | * | 0.4 | 0.8 | 1.9 | 3.2 | 7.6 | 12.1 | 17.2 | 31.7 |
| 1970 | 3.2 | 48.7 | * | * | 0.5 | 0.8 | 2.1 | 3.1 | 6.2 | 10.6 | 15.2 | 18.6 |
| 1969 | 3.0 | 54.3 | * | * | 0.4 | 0.7 | 1.6 | 3.1 | 5.1 | 8.5 | 16.6 | * |
| 1968 | 2.9 | 53.6 | 1.1 | * | 0.5 | * | 1.3 | 2.8 | 4.1 | 9.0 | 12.2 | 22.5 |
| ALL OTHER, MALE | | | | | | | | | | | | |
| 1978 | 6.3 | 61.9 | * | * | * | 1.0 | 2.9 | 7.7 | 14.1 | 28.5 | 62.4 | 105.2 |
| 1977 | 6.1 | 69.7 | * | * | * | * | 3.0 | 6.8 | 13.5 | 23.7 | 62.5 | 114.3 |
| 1976 | 5.5 | 65.0 | * | * | * | 1.3 | 2.8 | 5.7 | 11.6 | 23.4 | 47.9 | 94.5 |
| 1975 | 5.0 | 62.8 | * | * | * | 1.2 | 2.7 | 5.7 | 10.8 | 24.2 | 40.1 | 53.7 |
| 1974 | 4.9 | 62.5 | * | * | * | 1.1 | 2.3 | 5.4 | 10.5 | 21.1 | 35.4 | 66.0 |
| 1973 | 4.3 | 54.1 | * | * | * | * | 2.7 | 4.8 | 7.7 | 18.5 | 34.8 | 56.3 |
| 1972 | 4.2 | 52.0 | * | * | * | 1.5 | 1.9 | 4.7 | 10.4 | 17.7 | 32.3 | * |
| 1971 | 4.2 | 64.1 | * | * | * | * | 2.3 | 4.1 | 8.3 | 16.8 | 21.7 | * |
| 1970 | 3.6 | 46.6 | * | * | * | * | 2.1 | 3.9 | 7.7 | 13.2 | 24.3 | * |
| 1969 | 3.5 | 63.2 | * | * | * | * | 1.8 | 4.1 | 6.3 | 10.5 | 20.2 | * |
| 1968 | 3.3 | 60.0 | * | * | * | * | * | 2.8 | 5.1 | 11.4 | 16.1 | * |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| SEPTICEMIA 038 | | | | | | | | | | | | |
| ALL OTHER, FEMALE | | | | | | | | | | | | |
| 1978 | 5.2 | 54.3 | * | * | * | 0.9 | 2.2 | 4.7 | 9.4 | 19.0 | 41.1 | 82.6 |
| 1977 | 4.9 | 56.0 | * | * | * | * | 1.8 | 4.3 | 9.2 | 18.0 | 40.5 | 66.3 |
| 1976 | 4.9 | 62.5 | * | * | * | 0.9 | 1.9 | 5.0 | 10.3 | 15.8 | 35.4 | 72.4 |
| 1975 | 3.8 | 55.2 | * | * | * | * | 1.7 | 3.7 | 6.4 | 14.9 | 20.3 | 64.5 |
| 1974 | 3.5 | 50.6 | * | * | * | * | 1.6 | 3.6 | 6.8 | 13.5 | 16.6 | 50.6 |
| 1973 | 3.3 | 40.0 | * | * | * | * | 1.4 | 4.0 | 6.4 | 11.9 | 19.2 | 40.5 |
| 1972 | 3.4 | 50.7 | * | * | 0.7 | 1.1 | 1.5 | 2.9 | 5.9 | 12.5 | 19.3 | * |
| 1971 | 3.0 | 44.5 | * | * | * | * | 1.5 | 2.4 | 7.1 | 8.3 | 13.9 | 30.1 |
| 1970 | 2.8 | 50.8 | * | * | * | * | 2.1 | 2.4 | 5.0 | 8.4 | 8.3 | * |
| 1969 | 2.4 | 45.3 | * | * | * | * | 1.4 | 2.2 | 4.1 | 7.0 | 13.8 | * |
| 1968 | 2.5 | 46.9 | * | * | * | * | * | 2.8 | 3.1 | 7.0 | 9.2 | * |
| BLACK, BOTH SEXES | | | | | | | | | | | | |
| 1978 | 6.3 | 63.3 | * | * | 0.4 | 1.0 | 2.9 | 6.6 | 12.5 | 24.7 | 51.8 | 94.0 |
| 1977 | 5.9 | 68.7 | * | * | * | 0.7 | 2.5 | 6.0 | 12.1 | 21.3 | 49.7 | 85.3 |
| 1976 | 5.6 | 69.0 | 1.1 | * | 0.4 | 1.1 | 2.5 | 6.0 | 11.5 | 20.7 | 41.3 | 84.6 |
| 1975 | 4.7 | 63.2 | * | * | * | 1.0 | 2.4 | 4.9 | 8.7 | 20.1 | 27.8 | 65.4 |
| 1974 | 4.4 | 58.8 | 1.0 | * | 0.5 | 1.1 | 2.2 | 4.7 | 9.0 | 17.9 | 24.5 | 59.5 |
| 1973 | 4.0 | 49.5 | 1.1 | * | 0.4 | 1.0 | 2.2 | 4.8 | 7.6 | 16.1 | 25.3 | 47.4 |
| 1972 | 3.9 | 52.0 | * | * | 0.6 | 1.4 | 1.7 | 3.8 | 8.1 | 15.9 | 25.6 | 27.3 |
| 1971 | 3.7 | 56.2 | 1.1 | * | * | 0.9 | 2.1 | 3.2 | 7.8 | 12.7 | 18.0 | 30.8 |
| 1970 | 3.4 | 51.1 | * | * | 0.5 | 0.9 | 2.3 | 3.4 | 6.7 | 10.9 | 15.6 | 21.3 |
| 1969 | 3.1 | 56.0 | * | * | * | 0.8 | 1.7 | 3.4 | 5.2 | 8.3 | 17.3 | * |
| 1968 | 3.0 | 56.0 | 1.1 | * | 0.5 | * | 1.5 | 3.1 | 4.3 | 9.2 | 11.9 | 22.8 |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| SEPTICEMIA 038 | | | | | | | | | | | | |
| BLACK, MALE | | | | | | | | | | | | |
| 1978 | 6.9 | 68.1 | * | * | * | * | 3.4 | 8.6 | 15.3 | 30.7 | 66.9 | 105.9 |
| 1977 | 6.6 | 76.3 | * | * | * | * | 3.1 | 7.7 | 14.6 | 25.6 | 60.9 | 120.4 |
| 1976 | 5.9 | 70.7 | * | * | * | 1.5 | 2.9 | 6.4 | 12.5 | 25.5 | 50.0 | 97.9 |
| 1975 | 5.4 | 68.7 | * | * | * | 1.4 | 3.0 | 6.3 | 11.3 | 25.5 | 40.5 | 63.0 |
| 1974 | 5.2 | 65.9 | * | * | * | 1.4 | 2.6 | 5.7 | 11.6 | 23.0 | 34.9 | 69.8 |
| 1973 | 4.5 | 55.8 | * | * | * | * | 2.9 | 5.3 | 8.4 | 20.4 | 34.3 | 61.0 |
| 1972 | 4.4 | 53.9 | * | * | * | 1.6 | 2.0 | 4.9 | 10.8 | 19.6 | 33.5 | * |
| 1971 | 4.3 | 65.9 | * | * | * | * | 2.7 | 4.1 | 8.2 | 18.0 | 22.6 | * |
| 1970 | 3.8 | 48.7 | * | * | * | * | 2.4 | 4.3 | 8.1 | 14.1 | 24.9 | * |
| 1969 | 3.6 | 66.4 | * | * | * | * | * | 4.5 | 6.3 | 9.5 | 21.6 | * |
| 1968 | 3.5 | 63.0 | * | * | * | * | * | 3.1 | 5.6 | 11.7 | 15.2 | * |
| BLACK, FEMALE | | | | | | | | | | | | |
| 1978 | 5.7 | 58.4 | * | * | * | 1.0 | 2.5 | 5.0 | 10.3 | 20.3 | 41.6 | 87.9 |
| 1977 | 5.3 | 60.8 | * | * | * | * | 1.9 | 4.5 | 10.2 | 18.1 | 42.4 | 67.0 |
| 1976 | 5.4 | 67.3 | * | * | * | * | 2.1 | 5.6 | 10.7 | 17.1 | 35.5 | 77.5 |
| 1975 | 4.0 | 57.5 | * | * | * | * | 1.9 | 3.8 | 6.6 | 16.0 | 19.2 | 66.7 |
| 1974 | 3.7 | 51.6 | * | * | * | * | 1.8 | 3.9 | 6.8 | 14.1 | 17.5 | 53.8 |
| 1973 | 3.5 | 42.8 | * | * | * | * | 1.6 | 4.4 | 6.9 | 12.7 | 19.3 | 39.7 |
| 1972 | 3.5 | 49.8 | * | * | 0.8 | * | 1.5 | 2.9 | 5.7 | 13.0 | 20.0 | * |
| 1971 | 3.2 | 46.4 | * | * | * | * | 1.7 | 2.5 | 7.4 | 8.6 | 14.7 | 29.9 |
| 1970 | 3.0 | 53.6 | * | * | * | * | 2.2 | 2.6 | 5.4 | 8.4 | 8.7 | * |
| 1969 | 2.5 | 45.6 | * | * | * | * | 1.6 | 2.3 | 4.3 | 7.3 | 14.1 | * |
| 1968 | 2.6 | 48.9 | * | * | * | * | * | 3.0 | 3.3 | 7.1 | 9.5 | * |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| ACUTE POLIOMYELITIS (040-043) | | | | | | | | | | | | |
| BOTH SEXES | | | | | | | | | | | | |
| 1978 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1977 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1976 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1975 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1974 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1973 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1972 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1971 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1970 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1969 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1968 | 0.0 | * | * | * | * | * | * | * | * | * | * | * |
| MALE | | | | | | | | | | | | |
| 1978 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1977 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1976 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1975 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1974 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1973 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1972 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1971 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1970 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1969 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1968 | * | * | * | * | * | * | * | * | * | * | * | * |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| ACUTE POLIOMYELITIS (040-043) | | | | | | | | | | | | |
| FEMALE | | | | | | | | | | | | |
| 1978 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1977 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1976 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1975 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1974 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1973 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1972 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1971 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1970 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1969 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1968 | * | * | * | * | * | * | * | * | * | * | * | * |
| WHITE, BOTH SEXES | | | | | | | | | | | | |
| 1978 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1977 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1976 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1975 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1974 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1973 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1972 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1971 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1970 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1969 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1968 | 0.0 | * | * | * | * | * | * | * | * | * | * | * |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| ACUTE POLIOMYELITIS (040-043) | | | | | | | | | | | | |
| WHITE, MALE | | | | | | | | | | | | |
| 1978 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1977 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1976 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1975 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1974 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1973 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1972 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1971 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1970 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1969 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1968 | * | * | * | * | * | * | * | * | * | * | * | * |
| WHITE, FEMALE | | | | | | | | | | | | |
| 1978 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1977 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1976 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1975 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1974 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1973 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1972 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1971 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1970 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1969 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1968 | * | * | * | * | * | * | * | * | * | * | * | * |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| ACUTE POLIOMYELITIS (040-043) | | | | | | | | | | | | |
| ALL OTHER, BOTH SEXES | | | | | | | | | | | | |
| 1978 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1977 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1976 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1975 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1974 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1973 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1972 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1971 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1970 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1969 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1968 | * | * | * | * | * | * | * | * | * | * | * | * |
| ALL OTHER, MALE | | | | | | | | | | | | |
| 1978 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1977 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1976 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1975 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1974 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1973 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1972 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1971 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1970 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1969 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1968 | * | * | * | * | * | * | * | * | * | * | * | * |
| ALL OTHER, FEMALE | | | | | | | | | | | | |
| 1978 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1977 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1976 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1975 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1974 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1973 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1972 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1971 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1970 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1969 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1968 | * | * | * | * | * | * | * | * | * | * | * | * |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| ACUTE POLIOMYELITIS (040-043) | | | | | | | | | | | | |
| BLACK, BOTH SEXES | | | | | | | | | | | | |
| 1978 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1977 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1976 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1975 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1974 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1973 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1972 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1971 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1970 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1969 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1968 | * | * | * | * | * | * | * | * | * | * | * | * |
| BLACK, MALE | | | | | | | | | | | | |
| 1978 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1977 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1976 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1975 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1974 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1973 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1972 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1971 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1970 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1969 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1968 | * | * | * | * | * | * | * | * | * | * | * | * |
| BLACK, FEMALE | | | | | | | | | | | | |
| 1978 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1977 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1976 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1975 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1974 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1973 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1972 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1971 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1970 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1969 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1968 | * | * | * | * | * | * | * | * | * | * | * | * |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| MEASLES (055) | | | | | | | | | | | | |
| BOTH SEXES | | | | | | | | | | | | |
| 1978 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1977 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1976 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1975 | 0.0 | * | * | * | * | * | * | * | * | * | * | * |
| 1974 | 0.0 | * | * | * | * | * | * | * | * | * | * | * |
| 1973 | 0.0 | * | * | * | * | * | * | * | * | * | * | * |
| 1972 | 0.0 | * | * | * | * | * | * | * | * | * | * | * |
| 1971 | 0.0 | * | 0.3 | 0.1 | * | * | * | * | * | * | * | * |
| 1970 | 0.0 | 0.6 | 0.3 | * | * | * | * | * | * | * | * | * |
| 1969 | 0.0 | * | * | * | * | * | * | * | * | * | * | * |
| 1968 | 0.0 | * | * | * | * | * | * | * | * | * | * | * |
| MALE | | | | | | | | | | | | |
| 1978 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1977 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1976 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1975 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1974 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1973 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1972 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1971 | 0.0 | * | * | * | * | * | * | * | * | * | * | * |
| 1970 | 0.0 | * | 0.3 | * | * | * | * | * | * | * | * | * |
| 1969 | 0.0 | * | * | * | * | * | * | * | * | * | * | * |
| 1968 | * | * | * | * | * | * | * | * | * | * | * | * |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| MEASLES (055) | | | | | | | | | | | | |
| FEMALE | | | | | | | | | | | | |
| 1978 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1977 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1976 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1975 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1974 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1973 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1972 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1971 | 0.1 | * | 0.4 | * | * | * | * | * | * | * | * | * |
| 1970 | 0.0 | * | 0.3 | * | * | * | * | * | * | * | * | * |
| 1969 | 0.0 | * | * | * | * | * | * | * | * | * | * | * |
| 1968 | * | * | * | * | * | * | * | * | * | * | * | * |
| WHITE, BOTH SEXES | | | | | | | | | | | | |
| 1978 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1977 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1976 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1975 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1974 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1973 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1972 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1971 | 0.0 | * | 0.3 | * | * | * | * | * | * | * | * | * |
| 1970 | 0.0 | * | 0.2 | * | * | * | * | * | * | * | * | * |
| 1969 | 0.0 | * | * | * | * | * | * | * | * | * | * | * |
| 1968 | * | * | * | * | * | * | * | * | * | * | * | * |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| MEASLES (055) | | | | | | | | | | | | |
| WHITE, MALE | | | | | | | | | | | | |
| 1978 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1977 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1976 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1975 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1974 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1973 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1972 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1971 | 0.0 | * | * | * | * | * | * | * | * | * | * | * |
| 1970 | 0.0 | * | * | * | * | * | * | * | * | * | * | * |
| 1969 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1968 | * | * | * | * | * | * | * | * | * | * | * | * |
| WHITE, FEMALE | | | | | | | | | | | | |
| 1978 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1977 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1976 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1975 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1974 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1973 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1972 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1971 | 0.0 | * | * | * | * | * | * | * | * | * | * | * |
| 1970 | 0.0 | * | * | * | * | * | * | * | * | * | * | * |
| 1969 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1968 | * | * | * | * | * | * | * | * | * | * | * | * |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| MEASLES (055) | | | | | | | | | | | | |
| ALL OTHER, BOTH SEXES | | | | | | | | | | | | |
| 1978 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1977 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1976 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1975 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1974 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1973 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1972 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1971 | 0.1 | * | * | * | * | * | * | * | * | * | * | * |
| 1970 | 0.1 | * | * | * | * | * | * | * | * | * | * | * |
| 1969 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1968 | * | * | * | * | * | * | * | * | * | * | * | * |
| ALL OTHER, MALE | | | | | | | | | | | | |
| 1978 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1977 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1976 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1975 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1974 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1973 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1972 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1971 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1970 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1969 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1968 | * | * | * | * | * | * | * | * | * | * | * | * |
| ALL OTHER, FEMALE | | | | | | | | | | | | |
| 1978 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1977 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1976 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1975 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1974 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1973 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1972 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1971 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1970 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1969 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1968 | * | * | * | * | * | * | * | * | * | * | * | * |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| MEASLES (055) | | | | | | | | | | | | |
| BLACK, BOTH SEXES | | | | | | | | | | | | |
| 1978 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1977 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1976 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1975 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1974 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1973 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1972 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1971 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1970 | 0.1 | * | * | * | * | * | * | * | * | * | * | * |
| 1969 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1968 | * | * | * | * | * | * | * | * | * | * | * | * |
| BLACK, MALE | | | | | | | | | | | | |
| 1978 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1977 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1976 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1975 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1974 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1973 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1972 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1971 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1970 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1969 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1968 | * | * | * | * | * | * | * | * | * | * | * | * |
| BLACK, FEMALE | | | | | | | | | | | | |
| 1978 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1977 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1976 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1975 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1974 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1973 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1972 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1971 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1970 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1969 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1968 | * | * | * | * | * | * | * | * | * | * | * | * |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| SYPHILIS AND ITS SEQUELAE (090-097) | | | | | | | | | | | | |
| BOTH SEXES | | | | | | | | | | | | |
| 1978 | 0.1 | * | * | * | * | * | * | * | 0.1 | 0.5 | 0.6 | * |
| 1977 | 0.1 | * | * | * | * | * | * | * | 0.2 | 0.5 | 0.8 | * |
| 1976 | 0.1 | * | * | * | * | * | * | * | 0.2 | 0.6 | 0.8 | * |
| 1975 | 0.1 | * | * | * | * | * | * | * | 0.3 | 0.7 | 0.9 | * |
| 1974 | 0.1 | * | * | * | * | * | * | 0.1 | 0.3 | 0.8 | 1.1 | * |
| 1973 | 0.2 | * | * | * | * | * | * | 0.1 | 0.5 | 1.1 | 1.3 | 1.4 |
| 1972 | 0.2 | * | * | * | * | * | * | * | 0.4 | 1.0 | 1.1 | 1.7 |
| 1971 | 0.2 | * | * | * | * | * | * | 0.1 | 0.5 | 1.2 | 1.0 | 1.3 |
| 1970 | 0.2 | * | * | * | * | * | * | 0.1 | 0.6 | 1.3 | 1.7 | * |
| 1969 | 0.3 | 0.6 | * | * | * | * | * | 0.2 | 0.7 | 1.7 | 1.7 | 1.7 |
| 1968 | 0.3 | * | * | * | * | * | * | 0.3 | 1.0 | 1.5 | 1.8 | * |
| MALE | | | | | | | | | | | | |
| 1978 | 0.1 | * | * | * | * | * | * | * | * | 0.6 | 0.8 | * |
| 1977 | 0.1 | * | * | * | * | * | * | * | * | 0.6 | 1.5 | * |
| 1976 | 0.1 | * | * | * | * | * | * | * | 0.2 | 0.8 | 1.4 | * |
| 1975 | 0.2 | * | * | * | * | * | * | * | 0.5 | 1.2 | 1.7 | * |
| 1974 | 0.2 | * | * | * | * | * | * | * | 0.4 | 1.1 | 1.8 | * |
| 1973 | 0.3 | * | * | * | * | * | * | * | 0.7 | 1.9 | 2.2 | * |
| 1972 | 0.2 | * | * | * | * | * | * | * | 0.7 | 1.7 | 2.0 | * |
| 1971 | 0.2 | * | * | * | * | * | * | * | 0.6 | 2.0 | 1.8 | * |
| 1970 | 0.3 | * | * | * | * | * | * | * | 0.9 | 2.1 | 2.5 | * |
| 1969 | 0.4 | * | * | * | * | * | * | 0.3 | 1.0 | 2.7 | 3.5 | * |
| 1968 | 0.4 | * | * | * | * | * | * | 0.4 | 1.5 | 2.6 | 3.4 | * |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| SYPHILIS AND ITS SEQUELAE (090-097) | | | | | | | | | | | | |
| FEMALE | | | | | | | | | | | | |
| 1978 | 0.1 | * | * | * | * | * | * | * | * | 0.4 | 0.5 | * |
| 1977 | 0.1 | * | * | * | * | * | * | * | 0.2 | 0.4 | * | * |
| 1976 | 0.1 | * | * | * | * | * | * | * | 0.2 | 0.4 | * | * |
| 1975 | 0.1 | * | * | * | * | * | * | * | * | 0.3 | * | * |
| 1974 | 0.1 | * | * | * | * | * | * | * | 0.2 | 0.6 | 0.7 | * |
| 1973 | 0.1 | * | * | * | * | * | * | * | 0.3 | 0.5 | 0.8 | * |
| 1972 | 0.1 | * | * | * | * | * | * | * | 0.3 | 0.5 | 0.5 | * |
| 1971 | 0.1 | * | * | * | * | * | * | * | 0.4 | 0.6 | 0.6 | * |
| 1970 | 0.2 | * | * | * | * | * | * | * | 0.4 | 0.7 | 1.2 | * |
| 1969 | 0.2 | * | * | * | * | * | * | * | 0.4 | 0.9 | 0.6 | * |
| 1968 | 0.2 | * | * | * | * | * | * | 0.2 | 0.4 | 0.7 | 0.8 | * |
| WHITE, BOTH SEXES | | | | | | | | | | | | |
| 1978 | 0.1 | * | * | * | * | * | * | * | * | 0.3 | 0.4 | * |
| 1977 | 0.1 | * | * | * | * | * | * | * | * | 0.3 | 0.5 | * |
| 1976 | 0.1 | * | * | * | * | * | * | * | 0.1 | 0.3 | 0.5 | * |
| 1975 | 0.1 | * | * | * | * | * | * | * | 0.1 | 0.5 | 0.7 | * |
| 1974 | 0.1 | * | * | * | * | * | * | * | 0.2 | 0.5 | 0.8 | * |
| 1973 | 0.1 | * | * | * | * | * | * | * | 0.3 | 0.7 | 0.9 | * |
| 1972 | 0.1 | * | * | * | * | * | * | * | 0.2 | 0.6 | 0.9 | * |
| 1971 | 0.1 | * | * | * | * | * | * | * | 0.3 | 0.8 | 0.8 | * |
| 1970 | 0.2 | * | * | * | * | * | * | * | 0.4 | 0.9 | 1.2 | * |
| 1969 | 0.2 | * | * | * | * | * | * | * | 0.4 | 1.0 | 1.3 | * |
| 1968 | 0.2 | * | * | * | * | * | * | 0.1 | 0.6 | 1.0 | 1.4 | * |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| SYPHILIS AND ITS SEQUELAE (090-097) | | | | | | | | | | | | |
| WHITE, MALE | | | | | | | | | | | | |
| 1978 | 0.1 | * | * | * | * | * | * | * | * | 0.4 | * | * |
| 1977 | 0.1 | * | * | * | * | * | * | * | * | 0.4 | 0.8 | * |
| 1976 | 0.1 | * | * | * | * | * | * | * | * | * | 0.9 | * |
| 1975 | 0.1 | * | * | * | * | * | * | * | 0.2 | 0.7 | 1.4 | * |
| 1974 | 0.1 | * | * | * | * | * | * | * | * | 0.8 | 1.3 | * |
| 1973 | 0.2 | * | * | * | * | * | * | * | 0.4 | 1.2 | 1.6 | * |
| 1972 | 0.2 | * | * | * | * | * | * | * | 0.4 | 1.1 | 1.6 | * |
| 1971 | 0.2 | * | * | * | * | * | * | * | 0.4 | 1.3 | 1.3 | * |
| 1970 | 0.2 | * | * | * | * | * | * | * | 0.5 | 1.5 | 1.6 | * |
| 1969 | 0.2 | * | * | * | * | * | * | * | 0.7 | 1.6 | 2.7 | * |
| 1968 | 0.3 | * | * | * | * | * | * | * | 1.0 | 1.6 | 2.5 | * |
| WHITE, FEMALE | | | | | | | | | | | | |
| 1978 | 0.0 | * | * | * | * | * | * | * | * | 0.3 | * | * |
| 1977 | 0.0 | * | * | * | * | * | * | * | * | * | * | * |
| 1976 | 0.1 | * | * | * | * | * | * | * | * | 0.3 | * | * |
| 1975 | 0.1 | * | * | * | * | * | * | * | * | * | * | * |
| 1974 | 0.1 | * | * | * | * | * | * | * | * | 0.3 | 0.5 | * |
| 1973 | 0.1 | * | * | * | * | * | * | * | * | 0.3 | * | * |
| 1972 | 0.1 | * | * | * | * | * | * | * | * | * | * | * |
| 1971 | 0.1 | * | * | * | * | * | * | * | 0.2 | 0.4 | * | * |
| 1970 | 0.1 | * | * | * | * | * | * | * | 0.3 | 0.5 | 0.9 | * |
| 1969 | 0.1 | * | * | * | * | * | * | * | 0.2 | 0.5 | * | * |
| 1968 | 0.1 | * | * | * | * | * | * | * | * | 0.5 | 0.6 | * |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| SYPHILIS AND ITS SEQUELAE (090-097) | | | | | | | | | | | | |
| ALL OTHER, BOTH SEXES | | | | | | | | | | | | |
| 1978 | 0.2 | * | * | * | * | * | * | * | * | 2.0 | * | * |
| 1977 | 0.3 | * | * | * | * | * | * | * | 1.0 | 2.4 | 3.9 | * |
| 1976 | 0.4 | * | * | * | * | * | * | * | 1.1 | 3.2 | 3.6 | * |
| 1975 | 0.4 | * | * | * | * | * | * | * | 1.6 | 3.2 | * | * |
| 1974 | 0.5 | * | * | * | * | * | * | * | 1.8 | 3.5 | 4.5 | * |
| 1973 | 0.7 | * | * | * | * | * | * | * | 2.5 | 5.2 | 6.1 | * |
| 1972 | 0.6 | * | * | * | * | * | * | * | 2.5 | 5.1 | * | * |
| 1971 | 0.6 | * | * | * | * | * | * | * | 2.4 | 5.1 | 4.3 | * |
| 1970 | 0.7 | * | * | * | * | * | * | 0.8 | 2.8 | 4.9 | 8.1 | * |
| 1969 | 1.0 | * | * | * | * | * | * | 1.6 | 3.2 | 8.4 | 6.7 | * |
| 1968 | 1.0 | * | * | * | * | * | * | 1.7 | 4.8 | 7.2 | 8.0 | * |
| ALL OTHER, MALE | | | | | | | | | | | | |
| 1978 | 0.2 | * | * | * | * | * | * | * | * | * | * | * |
| 1977 | 0.4 | * | * | * | * | * | * | * | * | * | 8.0 | * |
| 1976 | 0.5 | * | * | * | * | * | * | * | * | 4.7 | * | * |
| 1975 | 0.6 | * | * | * | * | * | * | * | 2.5 | 5.6 | * | * |
| 1974 | 0.6 | * | * | * | * | * | * | * | 2.5 | 4.1 | * | * |
| 1973 | 0.9 | * | * | * | * | * | * | * | 4.1 | 8.6 | * | * |
| 1972 | 0.8 | * | * | * | * | * | * | * | 3.3 | 7.9 | * | * |
| 1971 | 0.8 | * | * | * | * | * | * | * | 3.0 | 8.8 | * | * |
| 1970 | 1.0 | * | * | * | * | * | * | * | 4.2 | 7.7 | 12.4 | * |
| 1969 | 1.4 | * | * | * | * | * | * | 2.2 | 4.7 | 12.8 | 13.3 | * |
| 1968 | 1.6 | * | * | * | * | * | * | 2.3 | 7.3 | 12.4 | 14.5 | * |
| ALL OTHER, FEMALE | | | | | | | | | | | | |
| 1978 | 0.2 | * | * | * | * | * | * | * | * | * | * | * |
| 1977 | 0.2 | * | * | * | * | * | * | * | * | * | * | * |
| 1976 | 0.2 | * | * | * | * | * | * | * | * | * | * | * |
| 1975 | 0.2 | * | * | * | * | * | * | * | * | * | * | * |
| 1974 | 0.4 | * | * | * | * | * | * | * | * | 3.0 | * | * |
| 1973 | 0.4 | * | * | * | * | * | * | * | * | * | * | * |
| 1972 | 0.4 | * | * | * | * | * | * | * | * | * | * | * |
| 1971 | 0.4 | * | * | * | * | * | * | * | * | * | * | * |
| 1970 | 0.5 | * | * | * | * | * | * | * | * | * | * | * |
| 1969 | 0.6 | * | * | * | * | * | * | * | * | 4.7 | * | * |
| 1968 | 0.6 | * | * | * | * | * | * | * | 2.7 | * | * | * |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| SYPHILIS AND ITS SEQUELAE (090-097) | | | | | | | | | | | | |
| BLACK, BOTH SEXES | | | | | | | | | | | | |
| 1978 | 0.3 | * | * | * | * | * | * | * | * | 2.2 | * | * |
| 1977 | 0.4 | * | * | * | * | * | * | * | 1.2 | 2.7 | 4.1 | * |
| 1976 | 0.4 | * | * | * | * | * | * | * | 1.2 | 3.5 | 3.9 | * |
| 1975 | 0.4 | * | * | * | * | * | * | * | 1.7 | 3.4 | * | * |
| 1974 | 0.5 | * | * | * | * | * | * | * | 2.0 | 3.8 | 4.4 | * |
| 1973 | 0.7 | * | * | * | * | * | * | * | 2.6 | 5.8 | 6.2 | * |
| 1972 | 0.6 | * | * | * | * | * | * | * | 2.8 | 5.6 | * | * |
| 1971 | 0.6 | * | * | * | * | * | * | * | 2.5 | 5.3 | 4.5 | * |
| 1970 | 0.8 | * | * | * | * | * | * | 0.9 | 3.0 | 5.3 | 8.8 | * |
| 1969 | 1.0 | * | * | * | * | * | * | 1.7 | 3.4 | 8.7 | 7.4 | * |
| 1968 | 1.2 | * | * | * | * | * | * | 1.8 | 5.3 | 7.9 | 8.8 | * |
| BLACK, MALE | | | | | | | | | | | | |
| 1978 | 0.3 | * | * | * | * | * | * | * | * | * | * | * |
| 1977 | 0.5 | * | * | * | * | * | * | * | * | * | 8.5 | * |
| 1976 | 0.6 | * | * | * | * | * | * | * | * | 5.2 | * | * |
| 1975 | 0.6 | * | * | * | * | * | * | * | 2.5 | 6.2 | * | * |
| 1974 | 0.6 | * | * | * | * | * | * | * | 2.8 | 4.7 | * | * |
| 1973 | 1.0 | * | * | * | * | * | * | * | 4.3 | 9.8 | * | * |
| 1972 | 0.9 | * | * | * | * | * | * | * | 3.7 | 9.0 | * | * |
| 1971 | 0.9 | * | * | * | * | * | * | * | 3.2 | 9.2 | * | * |
| 1970 | 1.1 | * | * | * | * | * | * | * | 4.5 | 8.5 | 13.6 | * |
| 1969 | 1.5 | * | * | * | * | * | * | 2.4 | 5.1 | 13.2 | 15.0 | * |
| 1968 | 1.7 | * | * | * | * | * | * | 2.5 | 8.1 | 13.7 | 16.4 | * |
| BLACK, FEMALE | | | | | | | | | | | | |
| 1978 | 0.3 | * | * | * | * | * | * | * | * | * | * | * |
| 1977 | 0.3 | * | * | * | * | * | * | * | * | * | * | * |
| 1976 | 0.3 | * | * | * | * | * | * | * | * | * | * | * |
| 1975 | 0.2 | * | * | * | * | * | * | * | * | * | * | * |
| 1974 | 0.4 | * | * | * | * | * | * | * | * | 3.1 | * | * |
| 1973 | 0.5 | * | * | * | * | * | * | * | * | * | * | * |
| 1972 | 0.4 | * | * | * | * | * | * | * | * | * | * | * |
| 1971 | 0.4 | * | * | * | * | * | * | * | * | * | * | * |
| 1970 | 0.6 | * | * | * | * | * | * | * | * | * | * | * |
| 1969 | 0.6 | * | * | * | * | * | * | * | * | 5.1 | * | * |
| 1968 | 0.6 | * | * | * | * | * | * | * | 2.9 | * | * | * |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| OTHER INFECTIVE AND PARASITIC DISEASES (REMAINDER OF 000-136) | | | | | | | | | | | | |
| BOTH SEXES | | | | | | | | | | | | |
| 1978 | 2.2 | 9.6 | 0.9 | 0.4 | 0.6 | 0.8 | 1.3 | 2.2 | 3.4 | 6.3 | 10.4 | 20.7 |
| 1977 | 2.0 | 9.1 | 1.1 | 0.4 | 0.6 | 0.8 | 1.2 | 1.9 | 3.2 | 5.7 | 8.9 | 19.3 |
| 1976 | 2.0 | 10.1 | 1.1 | 0.5 | 0.6 | 0.8 | 1.3 | 2.0 | 3.5 | 5.2 | 9.8 | 19.6 |
| 1975 | 2.0 | 8.9 | 1.0 | 0.4 | 0.6 | 0.8 | 1.3 | 2.0 | 3.6 | 5.4 | 9.6 | 17.4 |
| 1974 | 1.9 | 9.4 | 1.3 | 0.6 | 0.7 | 0.9 | 1.3 | 2.0 | 2.9 | 5.0 | 7.6 | 14.7 |
| 1973 | 1.9 | 9.9 | 1.3 | 0.7 | 0.7 | 0.9 | 1.2 | 1.9 | 3.0 | 4.6 | 8.2 | 16.9 |
| 1972 | 1.9 | 9.4 | 1.2 | 0.7 | 0.8 | 1.0 | 1.2 | 1.8 | 3.0 | 4.5 | 7.4 | 15.7 |
| 1971 | 2.0 | 10.2 | 1.4 | 0.8 | 0.9 | 1.1 | 1.4 | 1.9 | 3.0 | 4.1 | 7.2 | 13.7 |
| 1970 | 2.0 | 10.7 | 1.7 | 0.7 | 1.1 | 1.0 | 1.4 | 2.0 | 3.0 | 4.7 | 7.2 | 13.9 |
| 1969 | 2.0 | 11.4 | 1.9 | 0.8 | 1.0 | 1.0 | 1.3 | 2.1 | 3.1 | 4.2 | 7.6 | 13.5 |
| 1968 | 2.0 | 12.0 | 1.8 | 0.7 | 0.9 | 1.0 | 1.5 | 2.0 | 3.0 | 4.5 | 6.9 | 16.3 |
| MALE | | | | | | | | | | | | |
| 1978 | 2.2 | 10.0 | 0.8 | 0.4 | 0.6 | 0.8 | 1.4 | 2.4 | 3.8 | 7.9 | 13.8 | 24.5 |
| 1977 | 2.1 | 9.9 | 1.1 | 0.5 | 0.6 | 0.9 | 1.2 | 2.0 | 3.9 | 7.1 | 11.8 | 24.6 |
| 1976 | 2.1 | 11.3 | 1.2 | 0.5 | 0.7 | 0.8 | 1.2 | 2.2 | 4.0 | 6.6 | 12.2 | 21.5 |
| 1975 | 2.0 | 9.5 | 1.0 | 0.5 | 0.6 | 0.9 | 1.2 | 2.1 | 4.2 | 6.4 | 12.0 | 18.4 |
| 1974 | 1.9 | 9.8 | 1.3 | 0.6 | 0.6 | 0.8 | 1.3 | 2.2 | 3.2 | 6.6 | 9.7 | 15.9 |
| 1973 | 2.0 | 11.3 | 1.6 | 0.8 | 0.7 | 1.0 | 1.2 | 1.9 | 3.6 | 5.5 | 10.9 | 17.5 |
| 1972 | 2.0 | 10.5 | 1.4 | 0.6 | 0.7 | 0.8 | 1.3 | 1.8 | 3.6 | 5.9 | 10.1 | 19.9 |
| 1971 | 2.0 | 10.7 | 1.5 | 0.8 | 0.9 | 1.2 | 1.5 | 1.9 | 3.3 | 5.1 | 9.3 | 14.7 |
| 1970 | 2.2 | 12.0 | 1.6 | 0.7 | 1.2 | 1.0 | 1.3 | 2.2 | 3.5 | 6.2 | 9.2 | 17.0 |
| 1969 | 2.1 | 13.9 | 2.0 | 0.8 | 1.0 | 0.9 | 1.3 | 2.4 | 3.5 | 4.9 | 9.2 | 14.4 |
| 1968 | 2.1 | 13.2 | 1.8 | 0.8 | 0.8 | 1.0 | 1.6 | 1.8 | 3.3 | 5.6 | 7.8 | 15.5 |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| OTHER INFECTIVE AND PARASITIC DISEASES (REMAINDER OF 000-136) | | | | | | | | | | | | |
| FEMALE | | | | | | | | | | | | |
| 1978 | 2.1 | 9.2 | 1.0 | 0.4 | 0.6 | 0.9 | 1.1 | 1.9 | 3.1 | 5.2 | 8.5 | 19.0 |
| 1977 | 1.9 | 8.3 | 1.0 | 0.4 | 0.5 | 0.7 | 1.1 | 1.8 | 2.7 | 4.6 | 7.2 | 16.8 |
| 1976 | 1.9 | 8.9 | 1.0 | 0.5 | 0.5 | 0.8 | 1.4 | 1.8 | 3.0 | 4.2 | 8.4 | 18.7 |
| 1975 | 2.0 | 8.2 | 1.1 | 0.4 | 0.6 | 0.8 | 1.4 | 1.8 | 3.0 | 4.7 | 8.1 | 17.0 |
| 1974 | 1.8 | 8.9 | 1.2 | 0.5 | 0.7 | 0.9 | 1.3 | 1.9 | 2.5 | 3.8 | 6.3 | 14.0 |
| 1973 | 1.8 | 8.5 | 1.1 | 0.7 | 0.7 | 0.9 | 1.2 | 2.0 | 2.5 | 3.9 | 6.5 | 16.7 |
| 1972 | 1.8 | 8.2 | 1.0 | 0.8 | 0.9 | 1.1 | 1.2 | 1.8 | 2.6 | 3.4 | 5.7 | 13.5 |
| 1971 | 1.9 | 9.8 | 1.4 | 0.8 | 1.0 | 1.1 | 1.3 | 1.9 | 2.7 | 3.3 | 5.9 | 13.1 |
| 1970 | 1.9 | 9.3 | 1.8 | 0.7 | 1.0 | 1.0 | 1.4 | 1.8 | 2.6 | 3.6 | 6.0 | 12.2 |
| 1969 | 1.9 | 8.9 | 1.9 | 0.8 | 1.0 | 1.0 | 1.4 | 1.9 | 2.8 | 3.7 | 6.6 | 13.0 |
| 1968 | 2.0 | 10.8 | 1.8 | 0.6 | 0.9 | 1.0 | 1.5 | 2.3 | 2.7 | 3.7 | 6.4 | 16.8 |
| WHITE, BOTH SEXES | | | | | | | | | | | | |
| 1978 | 2.0 | 8.4 | 0.8 | 0.4 | 0.5 | 0.6 | 0.9 | 1.8 | 3.1 | 6.1 | 10.2 | 20.4 |
| 1977 | 1.8 | 7.4 | 1.0 | 0.5 | 0.5 | 0.6 | 0.9 | 1.6 | 2.8 | 5.2 | 8.6 | 18.9 |
| 1976 | 1.9 | 9.0 | 1.1 | 0.5 | 0.5 | 0.5 | 0.9 | 1.6 | 3.2 | 5.0 | 9.5 | 19.8 |
| 1975 | 1.8 | 7.2 | 1.0 | 0.4 | 0.5 | 0.7 | 1.0 | 1.5 | 3.1 | 5.0 | 9.4 | 17.3 |
| 1974 | 1.7 | 8.8 | 1.2 | 0.6 | 0.6 | 0.7 | 0.9 | 1.7 | 2.5 | 4.7 | 7.3 | 14.7 |
| 1973 | 1.8 | 8.7 | 1.3 | 0.7 | 0.6 | 0.6 | 1.0 | 1.6 | 2.6 | 4.3 | 8.1 | 17.2 |
| 1972 | 1.7 | 7.7 | 1.1 | 0.7 | 0.6 | 0.7 | 1.0 | 1.6 | 2.7 | 4.1 | 7.3 | 15.9 |
| 1971 | 1.8 | 9.0 | 1.4 | 0.8 | 0.8 | 0.9 | 1.1 | 1.5 | 2.7 | 3.8 | 7.3 | 14.1 |
| 1970 | 1.8 | 8.9 | 1.4 | 0.7 | 0.9 | 0.7 | 1.0 | 1.8 | 2.7 | 4.3 | 7.0 | 14.3 |
| 1969 | 1.9 | 9.3 | 1.7 | 0.8 | 0.8 | 0.8 | 1.1 | 1.8 | 2.8 | 4.1 | 7.7 | 14.1 |
| 1968 | 1.8 | 10.6 | 1.7 | 0.7 | 0.7 | 0.8 | 1.1 | 1.7 | 2.7 | 4.4 | 6.7 | 16.7 |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| OTHER INFECTIVE AND PARASITIC DISEASES (REMAINDER OF 000-136) | | | | | | | | | | | | |
| WHITE, MALE | | | | | | | | | | | | |
| 1978 | 2.1 | 8.9 | 0.7 | 0.4 | 0.6 | 0.5 | 1.0 | 2.1 | 3.4 | 7.6 | 13.5 | 22.9 |
| 1977 | 1.9 | 7.9 | 0.9 | 0.5 | 0.6 | 0.7 | 0.9 | 1.8 | 3.4 | 6.4 | 11.6 | 24.0 |
| 1976 | 2.0 | 10.0 | 1.3 | 0.5 | 0.6 | 0.5 | 1.0 | 1.7 | 3.7 | 6.5 | 11.8 | 21.4 |
| 1975 | 1.9 | 7.8 | 1.0 | 0.5 | 0.5 | 0.7 | 1.0 | 1.6 | 3.8 | 5.9 | 11.7 | 18.5 |
| 1974 | 1.8 | 8.9 | 1.3 | 0.6 | 0.6 | 0.8 | 0.9 | 2.0 | 2.8 | 6.2 | 9.6 | 16.1 |
| 1973 | 1.9 | 10.3 | 1.5 | 0.8 | 0.6 | 0.7 | 1.0 | 1.6 | 3.2 | 5.0 | 10.8 | 18.2 |
| 1972 | 1.8 | 8.6 | 1.3 | 0.6 | 0.6 | 0.6 | 1.0 | 1.6 | 3.3 | 5.2 | 9.9 | 20.2 |
| 1971 | 1.9 | 10.0 | 1.4 | 0.7 | 0.7 | 0.9 | 1.2 | 1.6 | 3.0 | 4.8 | 9.4 | 14.7 |
| 1970 | 2.0 | 10.5 | 1.4 | 0.8 | 0.9 | 0.7 | 1.0 | 2.0 | 3.0 | 5.8 | 9.0 | 17.0 |
| 1969 | 2.0 | 11.6 | 1.7 | 0.7 | 0.9 | 0.7 | 1.1 | 2.1 | 3.3 | 4.8 | 9.3 | 14.7 |
| 1968 | 1.9 | 11.9 | 1.7 | 0.8 | 0.7 | 0.8 | 1.2 | 1.6 | 3.0 | 5.4 | 7.4 | 16.3 |
| WHITE, FEMALE | | | | | | | | | | | | |
| 1978 | 1.9 | 7.8 | 0.9 | 0.4 | 0.5 | 0.8 | 0.8 | 1.5 | 2.8 | 4.9 | 8.2 | 19.3 |
| 1977 | 1.7 | 7.0 | 1.1 | 0.4 | 0.5 | 0.5 | 0.8 | 1.5 | 2.3 | 4.2 | 6.9 | 16.6 |
| 1976 | 1.8 | 7.9 | 0.9 | 0.5 | 0.4 | 0.5 | 0.8 | 1.5 | 2.7 | 3.8 | 8.2 | 19.1 |
| 1975 | 1.7 | 6.6 | 0.9 | 0.4 | 0.5 | 0.6 | 0.9 | 1.4 | 2.5 | 4.4 | 8.1 | 16.7 |
| 1974 | 1.6 | 8.6 | 1.1 | 0.6 | 0.6 | 0.7 | 0.9 | 1.5 | 2.3 | 3.5 | 6.0 | 14.0 |
| 1973 | 1.7 | 7.0 | 1.0 | 0.7 | 0.6 | 0.6 | 1.0 | 1.6 | 2.1 | 3.7 | 6.4 | 16.8 |
| 1972 | 1.6 | 6.8 | 1.0 | 0.8 | 0.7 | 0.7 | 0.9 | 1.5 | 2.3 | 3.2 | 5.6 | 13.8 |
| 1971 | 1.7 | 8.0 | 1.3 | 0.8 | 0.8 | 0.8 | 1.0 | 1.5 | 2.4 | 3.1 | 5.9 | 13.8 |
| 1970 | 1.7 | 7.3 | 1.5 | 0.7 | 0.8 | 0.7 | 1.1 | 1.5 | 2.4 | 3.2 | 5.7 | 12.8 |
| 1969 | 1.8 | 6.9 | 1.7 | 0.9 | 0.8 | 0.8 | 1.0 | 1.6 | 2.5 | 3.5 | 6.6 | 13.8 |
| 1968 | 1.8 | 9.3 | 1.6 | 0.6 | 0.7 | 0.8 | 1.1 | 1.8 | 2.4 | 3.6 | 6.3 | 16.9 |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| OTHER INFECTIVE AND PARASITIC DISEASES (REMAINDER OF 000-136) | | | | | | | | | | | | |
| ALL OTHER, BOTH SEXES | | | | | | | | | | | | |
| 1978 | 3.2 | 15.6 | 1.4 | 0.4 | 1.1 | 2.2 | 3.4 | 4.9 | 6.7 | 8.9 | 13.4 | 24.4 |
| 1977 | 3.1 | 17.1 | 1.5 | 0.3 | 0.8 | 2.2 | 3.2 | 4.2 | 6.9 | 10.3 | 11.5 | 23.8 |
| 1976 | 3.0 | 15.7 | 1.1 | 0.6 | 0.9 | 2.4 | 3.9 | 5.1 | 6.2 | 7.5 | 13.0 | 16.3 |
| 1975 | 3.2 | 17.1 | 1.4 | 0.6 | 1.0 | 2.0 | 3.7 | 5.3 | 7.7 | 9.5 | 11.1 | 19.2 |
| 1974 | 2.9 | 12.3 | 1.8 | 0.4 | 1.0 | 2.0 | 4.2 | 4.3 | 6.0 | 8.1 | 10.9 | * |
| 1973 | 3.0 | 16.2 | 1.7 | 0.7 | 1.2 | 2.9 | 2.9 | 4.7 | 7.1 | 7.2 | 8.8 | * |
| 1972 | 3.0 | 17.7 | 1.6 | 0.7 | 1.5 | 2.9 | 3.1 | 3.8 | 5.7 | 8.6 | 9.2 | * |
| 1971 | 3.1 | 16.9 | 1.7 | 0.8 | 2.0 | 3.1 | 3.3 | 4.8 | 5.5 | 6.9 | 6.8 | * |
| 1970 | 3.5 | 20.0 | 3.1 | 0.6 | 2.5 | 2.9 | 3.7 | 4.0 | 5.9 | 8.2 | 10.5 | * |
| 1969 | 3.4 | 23.5 | 3.4 | 0.9 | 1.9 | 2.6 | 3.5 | 4.9 | 6.1 | 5.6 | 6.9 | * |
| 1968 | 3.4 | 19.6 | 2.4 | 0.7 | 1.6 | 2.6 | 4.6 | 5.3 | 5.7 | 6.5 | 9.9 | * |
| ALL OTHER, MALE | | | | | | | | | | | | |
| 1978 | 3.4 | 15.6 | * | * | 1.1 | 2.5 | 4.0 | 4.9 | 7.4 | 10.4 | 16.2 | 41.4 |
| 1977 | 3.2 | 19.7 | 2.0 | * | 0.8 | 2.1 | 3.4 | 3.9 | 7.7 | 12.8 | 13.4 | * |
| 1976 | 3.1 | 17.7 | * | 0.7 | 1.1 | 2.5 | 3.0 | 6.2 | 6.8 | 7.2 | 16.0 | * |
| 1975 | 3.1 | 18.4 | * | 0.7 | 0.8 | 1.8 | 2.7 | 5.8 | 7.3 | 11.1 | 15.3 | * |
| 1974 | 2.8 | 14.3 | 1.8 | * | * | 1.5 | 4.2 | 3.9 | 7.2 | 9.8 | 11.8 | * |
| 1973 | 3.1 | 16.5 | * | 0.8 | 1.0 | 2.9 | 3.0 | 4.1 | 7.5 | 9.9 | 11.6 | * |
| 1972 | 3.2 | 20.5 | 2.0 | 0.6 | 1.2 | 2.6 | 3.0 | 3.7 | 6.4 | 12.1 | 12.0 | * |
| 1971 | 3.3 | 14.6 | 1.9 | 1.0 | 1.9 | 3.7 | 3.4 | 4.5 | 6.0 | 8.2 | * | * |
| 1970 | 3.6 | 20.2 | 2.9 | * | 2.7 | 2.9 | 3.4 | 3.5 | 7.5 | 9.6 | 11.4 | * |
| 1969 | 3.4 | 27.1 | 3.6 | 1.0 | 2.1 | 2.4 | 2.8 | 5.2 | 5.9 | 5.7 | * | * |
| 1968 | 3.2 | 20.4 | 1.9 | 0.7 | 1.6 | 2.5 | 5.2 | 3.6 | 5.9 | 8.0 | 12.4 | * |
| ALL OTHER, FEMALE | | | | | | | | | | | | |
| 1978 | 3.0 | 15.6 | * | * | 1.1 | 1.9 | 3.0 | 4.9 | 6.2 | 7.8 | 11.6 | * |
| 1977 | 2.9 | 14.4 | * | * | 0.8 | 2.2 | 3.0 | 4.4 | 6.2 | 8.3 | 10.2 | 20.2 |
| 1976 | 3.0 | 13.8 | * | * | 0.8 | 2.3 | 4.7 | 4.2 | 5.7 | 7.7 | 10.9 | * |
| 1975 | 3.3 | 15.8 | * | * | 1.2 | 2.2 | 4.5 | 4.9 | 8.0 | 8.2 | 8.2 | * |
| 1974 | 2.9 | 10.3 | 1.8 | * | 1.5 | 2.5 | 4.2 | 4.7 | 5.0 | 6.7 | 10.2 | * |
| 1973 | 3.0 | 15.8 | * | * | 1.4 | 2.9 | 2.8 | 5.1 | 6.7 | 5.0 | 6.9 | * |
| 1972 | 2.9 | 14.9 | * | 0.7 | 1.9 | 3.3 | 3.2 | 3.9 | 5.1 | 5.8 | 7.3 | * |
| 1971 | 3.0 | 19.2 | * | 0.7 | 2.2 | 2.6 | 3.2 | 4.9 | 5.0 | 5.9 | * | * |
| 1970 | 3.3 | 19.7 | 3.3 | * | 2.3 | 2.8 | 3.8 | 4.4 | 4.4 | 7.1 | 9.9 | * |
| 1969 | 3.3 | 19.9 | 3.2 | 0.8 | 1.8 | 2.7 | 4.2 | 4.7 | 6.2 | 5.5 | * | * |
| 1968 | 3.5 | 18.7 | 3.0 | 0.8 | 1.7 | 2.6 | 4.1 | 6.8 | 5.6 | 5.3 | * | * |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| OTHER INFECTIVE AND PARASITIC DISEASES (REMAINDER OF 000-136) | | | | | | | | | | | | |
| BLACK, BOTH SEXES | | | | | | | | | | | | |
| 1978 | 3.3 | 16.5 | 1.3 | 0.4 | 1.1 | 2.5 | 3.7 | 5.3 | 7.0 | 9.0 | 12.9 | 24.0 |
| 1977 | 3.2 | 16.9 | 1.5 | * | 0.9 | 2.4 | 3.6 | 4.7 | 7.1 | 10.2 | 11.7 | 20.3 |
| 1976 | 3.3 | 17.4 | 1.1 | 0.6 | 1.0 | 2.7 | 4.4 | 5.5 | 6.3 | 8.1 | 13.2 | 16.2 |
| 1975 | 3.4 | 15.8 | 1.4 | 0.6 | 1.1 | 2.3 | 4.0 | 5.8 | 7.7 | 9.8 | 10.7 | 18.5 |
| 1974 | 3.0 | 13.2 | 1.7 | 0.4 | 1.2 | 2.2 | 4.6 | 4.4 | 6.2 | 8.5 | 11.0 | * |
| 1973 | 3.2 | 15.6 | 1.8 | 0.7 | 1.3 | 3.2 | 3.1 | 5.0 | 7.3 | 7.1 | 9.2 | * |
| 1972 | 3.1 | 18.9 | 1.7 | 0.6 | 1.7 | 3.2 | 3.3 | 3.8 | 5.8 | 8.4 | 8.0 | * |
| 1971 | 3.2 | 17.4 | 1.6 | 0.8 | 2.3 | 3.5 | 3.5 | 5.0 | 5.5 | 6.9 | 6.3 | * |
| 1970 | 3.6 | 20.3 | 3.1 | 0.7 | 2.6 | 3.1 | 3.9 | 4.1 | 6.2 | 8.2 | 11.0 | * |
| 1969 | 3.5 | 24.9 | 3.2 | 0.9 | 2.0 | 2.8 | 3.8 | 5.0 | 6.4 | 5.6 | 7.1 | * |
| 1968 | 3.4 | 19.9 | 2.4 | 0.8 | 1.7 | 2.8 | 5.0 | 5.3 | 5.9 | 6.5 | 9.8 | * |
| BLACK, MALE | | | | | | | | | | | | |
| 1978 | 3.5 | 16.9 | * | * | 1.2 | 2.9 | 4.2 | 5.4 | 7.7 | 11.1 | 14.2 | 43.1 |
| 1977 | 3.4 | 20.0 | * | * | 0.9 | 2.3 | 3.9 | 4.4 | 8.0 | 13.1 | 13.2 | * |
| 1976 | 3.4 | 19.2 | * | 0.7 | 1.2 | 2.8 | 3.6 | 6.5 | 6.7 | 8.1 | 16.2 | * |
| 1975 | 3.2 | 16.1 | * | * | 0.9 | 2.1 | 2.9 | 6.6 | 7.3 | 11.4 | 14.9 | * |
| 1974 | 2.9 | 15.0 | * | * | * | 1.8 | 4.5 | 4.2 | 7.5 | 10.2 | 11.5 | * |
| 1973 | 3.2 | 15.5 | * | 0.9 | 1.1 | 3.2 | 3.3 | 4.4 | 7.9 | 10.0 | 11.8 | * |
| 1972 | 3.3 | 20.7 | 2.2 | * | 1.3 | 2.7 | 3.5 | 4.0 | 6.3 | 11.5 | 11.2 | * |
| 1971 | 3.3 | 15.7 | * | 0.9 | 2.0 | 4.0 | 3.5 | 4.7 | 6.3 | 7.9 | * | * |
| 1970 | 3.8 | 21.3 | 3.1 | * | 2.8 | 3.1 | 3.7 | 3.7 | 8.1 | 10.0 | 12.5 | * |
| 1969 | 3.5 | 29.3 | 3.3 | 1.0 | 2.2 | 2.8 | 2.9 | 5.3 | 6.2 | 5.3 | * | * |
| 1968 | 3.3 | 20.6 | 2.0 | 0.7 | 1.7 | 2.7 | 5.6 | 3.4 | 6.0 | 8.2 | 12.7 | * |
| BLACK, FEMALE | | | | | | | | | | | | |
| 1978 | 3.1 | 16.0 | * | * | 1.1 | 2.2 | 3.3 | 5.2 | 6.5 | 7.5 | 12.0 | * |
| 1977 | 3.1 | 13.8 | * | * | 0.9 | 2.5 | 3.4 | 5.0 | 6.4 | 8.0 | 10.7 | * |
| 1976 | 3.2 | 15.5 | * | * | 0.8 | 2.6 | 5.1 | 4.6 | 5.9 | 8.2 | 11.2 | * |
| 1975 | 3.5 | 15.5 | * | * | 1.3 | 2.6 | 4.9 | 5.2 | 8.0 | 8.6 | 7.9 | * |
| 1974 | 3.1 | 11.3 | * | * | 1.6 | 2.7 | 4.7 | 4.5 | 5.2 | 7.2 | 10.7 | * |
| 1973 | 3.1 | 15.7 | * | * | 1.6 | 3.2 | 2.9 | 5.6 | 6.8 | 4.8 | 7.4 | * |
| 1972 | 3.0 | 16.9 | * | * | 2.1 | 3.7 | 3.2 | 3.7 | 5.3 | 6.0 | * | * |
| 1971 | 3.2 | 19.2 | * | * | 2.4 | 3.0 | 3.4 | 5.2 | 4.8 | 6.1 | * | * |
| 1970 | 3.4 | 19.4 | 3.1 | * | 2.4 | 3.2 | 4.1 | 4.5 | 4.6 | 6.9 | 10.0 | * |
| 1969 | 3.4 | 20.6 | 3.1 | 0.8 | 1.9 | 2.9 | 4.5 | 4.7 | 6.6 | 5.9 | * | * |
| 1968 | 3.5 | 19.1 | 2.8 | 0.8 | 1.6 | 2.8 | 4.6 | 6.8 | 5.8 | 5.2 | * | * |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|

MALIGNANT NEOPLASMS INCLUDING NEOPLASMS OF LYMPHATIC AND HEMATOPOIETIC TISSUES (140-209)

BOTH SEXES

| | | | | | | | | | | | | |
|------|-------|-----|-----|-----|-----|------|------|-------|-------|-------|---------|---------|
| 1978 | 178.7 | 4.0 | 4.8 | 4.1 | 6.1 | 13.8 | 49.5 | 184.6 | 432.7 | 797.7 | 1,207.8 | 1,527.4 |
| 1977 | 176.0 | 3.7 | 5.1 | 4.7 | 6.4 | 14.2 | 50.9 | 182.6 | 432.4 | 789.6 | 1,192.3 | 1,508.7 |
| 1976 | 173.4 | 3.1 | 5.2 | 4.9 | 6.4 | 14.2 | 51.5 | 182.2 | 431.5 | 784.0 | 1,183.9 | 1,494.7 |
| 1975 | 169.7 | 4.1 | 5.5 | 4.7 | 6.7 | 14.6 | 53.0 | 181.9 | 424.9 | 773.2 | 1,168.0 | 1,452.1 |
| 1974 | 169.0 | 3.8 | 5.8 | 5.2 | 6.8 | 14.8 | 55.1 | 183.9 | 432.5 | 772.4 | 1,169.6 | 1,446.2 |
| 1973 | 166.1 | 4.3 | 6.3 | 5.3 | 7.4 | 15.4 | 56.1 | 181.0 | 426.4 | 766.4 | 1,154.4 | 1,455.9 |
| 1972 | 165.1 | 4.5 | 5.9 | 5.4 | 7.7 | 15.7 | 57.6 | 180.1 | 424.7 | 772.8 | 1,156.2 | 1,441.5 |
| 1971 | 163.1 | 4.2 | 7.1 | 5.8 | 7.8 | 16.9 | 58.2 | 179.7 | 419.9 | 757.8 | 1,154.0 | 1,452.3 |
| 1970 | 162.8 | 4.7 | 7.5 | 6.0 | 8.3 | 16.5 | 59.5 | 182.5 | 423.0 | 754.2 | 1,169.2 | 1,320.7 |
| 1969 | 160.4 | 4.8 | 7.6 | 6.2 | 7.9 | 16.9 | 61.2 | 179.9 | 415.7 | 747.7 | 1,140.4 | 1,466.0 |
| 1968 | 159.8 | 4.9 | 8.0 | 6.4 | 8.2 | 17.3 | 61.0 | 183.0 | 413.5 | 749.2 | 1,133.8 | 1,475.3 |

MALE

| | | | | | | | | | | | | |
|------|-------|-----|-----|-----|------|------|------|-------|-------|---------|---------|---------|
| 1978 | 200.1 | 3.7 | 5.4 | 4.7 | 7.5 | 13.6 | 44.7 | 193.3 | 516.1 | 1,075.7 | 1,741.1 | 2,255.2 |
| 1977 | 196.9 | 4.3 | 5.8 | 5.5 | 7.9 | 14.0 | 46.2 | 189.4 | 515.2 | 1,066.1 | 1,715.6 | 2,196.2 |
| 1976 | 194.0 | 3.3 | 5.5 | 5.7 | 7.8 | 13.7 | 46.3 | 188.8 | 515.7 | 1,058.7 | 1,701.3 | 2,119.5 |
| 1975 | 190.2 | 4.3 | 6.0 | 5.2 | 8.0 | 14.8 | 47.3 | 188.1 | 508.3 | 1,050.6 | 1,662.1 | 2,051.0 |
| 1974 | 189.4 | 4.0 | 6.3 | 5.9 | 8.3 | 14.3 | 48.8 | 191.0 | 518.6 | 1,044.1 | 1,662.0 | 2,026.9 |
| 1973 | 185.8 | 4.3 | 7.3 | 6.0 | 9.0 | 15.3 | 50.7 | 183.9 | 510.9 | 1,032.8 | 1,626.1 | 2,044.2 |
| 1972 | 184.8 | 4.4 | 6.4 | 6.0 | 9.5 | 15.2 | 51.4 | 181.5 | 515.0 | 1,035.5 | 1,623.2 | 1,939.5 |
| 1971 | 182.5 | 4.4 | 7.9 | 6.6 | 9.5 | 17.2 | 51.8 | 180.3 | 504.3 | 1,018.0 | 1,598.6 | 1,968.6 |
| 1970 | 182.1 | 4.4 | 8.3 | 6.7 | 10.4 | 16.3 | 53.0 | 183.5 | 511.8 | 1,006.8 | 1,588.3 | 1,720.8 |
| 1969 | 179.7 | 4.9 | 8.1 | 7.0 | 10.0 | 17.0 | 54.4 | 180.9 | 501.1 | 997.3 | 1,541.1 | 1,952.9 |
| 1968 | 178.8 | 4.7 | 8.8 | 6.9 | 10.0 | 17.3 | 53.7 | 183.0 | 497.9 | 998.2 | 1,520.1 | 1,936.1 |

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|--|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| MALIGNANT NEOPLASMS INCLUDING NEOPLASMS OF LYMPHATIC AND HEMATOPOIETIC TISSUES (140-209) | | | | | | | | | | | | |
| FEMALE | | | | | | | | | | | | |
| 1978 | 158.6 | 4.2 | 4.2 | 3.6 | 4.8 | 13.9 | 54.1 | 176.5 | 359.1 | 585.3 | 890.2 | 1,198.5 |
| 1977 | 156.1 | 3.1 | 4.4 | 3.8 | 4.9 | 14.3 | 55.3 | 176.3 | 359.2 | 578.2 | 879.1 | 1,193.0 |
| 1976 | 153.9 | 3.0 | 5.0 | 4.0 | 5.0 | 14.7 | 56.4 | 176.1 | 357.0 | 573.5 | 871.6 | 1,201.2 |
| 1975 | 150.3 | 3.9 | 4.9 | 4.2 | 5.3 | 14.5 | 58.4 | 176.2 | 351.0 | 560.7 | 866.9 | 1,162.2 |
| 1974 | 149.6 | 3.5 | 5.3 | 4.4 | 5.3 | 15.3 | 61.0 | 177.2 | 356.1 | 564.0 | 866.3 | 1,158.7 |
| 1973 | 147.4 | 4.3 | 5.3 | 4.7 | 5.9 | 15.5 | 61.2 | 178.3 | 351.4 | 561.8 | 859.6 | 1,159.8 |
| 1972 | 146.5 | 4.7 | 5.4 | 4.8 | 5.8 | 16.0 | 63.6 | 178.8 | 344.2 | 570.2 | 858.6 | 1,186.7 |
| 1971 | 144.7 | 4.1 | 6.3 | 4.9 | 6.1 | 16.6 | 64.2 | 179.2 | 344.3 | 556.3 | 865.8 | 1,182.8 |
| 1970 | 144.4 | 5.0 | 6.7 | 5.2 | 6.2 | 16.7 | 65.6 | 181.5 | 343.2 | 557.9 | 891.9 | 1,096.7 |
| 1969 | 142.1 | 4.6 | 7.1 | 5.4 | 6.0 | 16.8 | 67.6 | 178.9 | 338.8 | 552.6 | 871.5 | 1,202.5 |
| 1968 | 141.6 | 5.1 | 7.3 | 5.7 | 6.5 | 17.2 | 67.9 | 183.0 | 337.2 | 553.1 | 869.4 | 1,223.6 |
| WHITE, BOTH SEXES | | | | | | | | | | | | |
| 1978 | 183.4 | 4.0 | 5.0 | 4.2 | 6.1 | 13.6 | 46.5 | 175.8 | 419.9 | 787.0 | 1,211.9 | 1,540.4 |
| 1977 | 180.3 | 3.8 | 5.3 | 4.8 | 6.5 | 14.0 | 47.9 | 173.0 | 419.3 | 778.6 | 1,197.8 | 1,521.8 |
| 1976 | 177.8 | 3.2 | 5.5 | 5.0 | 6.5 | 14.1 | 49.2 | 173.3 | 418.7 | 773.4 | 1,191.5 | 1,510.7 |
| 1975 | 173.9 | 4.3 | 5.6 | 4.8 | 6.8 | 14.5 | 50.3 | 173.4 | 412.1 | 762.2 | 1,176.0 | 1,474.0 |
| 1974 | 172.9 | 3.5 | 6.2 | 5.3 | 6.8 | 14.6 | 52.2 | 175.3 | 419.9 | 759.8 | 1,180.5 | 1,465.9 |
| 1973 | 169.7 | 4.5 | 6.5 | 5.4 | 7.5 | 15.2 | 53.4 | 171.3 | 414.1 | 754.3 | 1,163.7 | 1,477.9 |
| 1972 | 169.2 | 4.4 | 6.2 | 5.7 | 7.9 | 15.2 | 54.4 | 171.4 | 414.6 | 763.3 | 1,167.7 | 1,466.7 |
| 1971 | 166.9 | 4.3 | 7.4 | 5.9 | 7.9 | 16.6 | 55.2 | 171.7 | 408.7 | 749.3 | 1,163.0 | 1,479.2 |
| 1970 | 166.8 | 4.8 | 7.7 | 6.2 | 8.4 | 16.2 | 56.4 | 174.7 | 414.1 | 747.4 | 1,176.1 | 1,354.9 |
| 1969 | 163.9 | 5.1 | 8.0 | 6.4 | 8.1 | 16.4 | 58.0 | 171.4 | 404.1 | 740.6 | 1,149.1 | 1,503.7 |
| 1968 | 163.2 | 4.9 | 8.4 | 6.6 | 8.3 | 16.8 | 58.0 | 175.2 | 402.0 | 742.2 | 1,142.5 | 1,509.7 |

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|--|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| MALIGNANT NEOPLASMS INCLUDING NEOPLASMS OF LYMPHATIC AND HEMATOPOIETIC TISSUES (140-209) | | | | | | | | | | | | |
| WHITE, MALE | | | | | | | | | | | | |
| 1978 | 202.9 | 3.5 | 5.6 | 4.7 | 7.4 | 13.7 | 41.6 | 179.6 | 493.7 | 1,056.2 | 1,745.9 | 2,273.2 |
| 1977 | 199.4 | 4.4 | 6.1 | 5.7 | 8.2 | 14.1 | 43.0 | 175.3 | 492.8 | 1,046.1 | 1,721.3 | 2,208.6 |
| 1976 | 196.6 | 3.0 | 5.8 | 5.9 | 7.9 | 13.7 | 43.7 | 175.7 | 492.7 | 1,039.3 | 1,712.5 | 2,149.2 |
| 1975 | 192.7 | 4.4 | 6.3 | 5.4 | 8.1 | 14.8 | 44.1 | 175.2 | 486.4 | 1,032.0 | 1,672.7 | 2,080.7 |
| 1974 | 192.0 | 3.7 | 6.6 | 6.0 | 8.4 | 14.4 | 46.0 | 178.6 | 498.5 | 1,025.0 | 1,672.2 | 2,053.8 |
| 1973 | 188.3 | 4.4 | 7.5 | 6.1 | 9.2 | 15.4 | 48.5 | 170.8 | 491.5 | 1,015.8 | 1,635.5 | 2,075.1 |
| 1972 | 187.7 | 4.0 | 6.6 | 6.3 | 9.9 | 14.9 | 47.8 | 169.6 | 498.3 | 1,022.6 | 1,636.9 | 1,974.7 |
| 1971 | 185.5 | 4.5 | 8.5 | 6.8 | 9.7 | 17.3 | 49.0 | 169.8 | 487.5 | 1,007.4 | 1,610.5 | 2,010.4 |
| 1970 | 185.1 | 4.3 | 8.5 | 7.0 | 10.6 | 16.2 | 50.1 | 172.0 | 498.1 | 997.0 | 1,592.7 | 1,772.2 |
| 1969 | 182.3 | 5.2 | 8.5 | 7.3 | 10.3 | 16.9 | 51.5 | 169.9 | 484.3 | 986.4 | 1,548.6 | 2,006.5 |
| 1968 | 181.5 | 4.7 | 9.1 | 7.2 | 10.2 | 17.3 | 50.7 | 172.3 | 482.4 | 988.6 | 1,529.7 | 1,985.2 |
| WHITE, FEMALE | | | | | | | | | | | | |
| 1978 | 164.8 | 4.5 | 4.5 | 3.6 | 4.8 | 13.4 | 51.4 | 172.2 | 354.3 | 581.7 | 897.8 | 1,213.8 |
| 1977 | 162.0 | 3.2 | 4.4 | 3.8 | 4.7 | 14.0 | 52.7 | 170.8 | 353.9 | 574.5 | 888.8 | 1,210.9 |
| 1976 | 159.9 | 3.5 | 5.2 | 4.1 | 5.1 | 14.5 | 54.5 | 171.0 | 352.9 | 570.4 | 881.2 | 1,216.9 |
| 1975 | 155.9 | 4.1 | 4.9 | 4.2 | 5.4 | 14.2 | 56.2 | 171.6 | 345.9 | 556.3 | 877.0 | 1,186.4 |
| 1974 | 154.7 | 3.3 | 5.8 | 4.5 | 5.2 | 14.8 | 58.2 | 172.2 | 349.8 | 557.5 | 880.4 | 1,179.2 |
| 1973 | 151.9 | 4.5 | 5.4 | 4.6 | 5.8 | 15.0 | 58.1 | 171.8 | 345.0 | 554.3 | 871.5 | 1,183.5 |
| 1972 | 151.4 | 4.9 | 5.7 | 5.0 | 5.9 | 15.5 | 60.8 | 173.2 | 339.8 | 564.3 | 871.8 | 1,211.0 |
| 1971 | 149.1 | 4.1 | 6.3 | 5.0 | 6.0 | 15.8 | 61.1 | 173.4 | 338.2 | 550.5 | 876.5 | 1,207.1 |
| 1970 | 149.4 | 5.4 | 6.9 | 5.4 | 6.2 | 16.3 | 62.4 | 177.3 | 338.6 | 554.7 | 903.5 | 1,126.6 |
| 1969 | 146.4 | 5.0 | 7.4 | 5.5 | 5.9 | 15.9 | 64.4 | 172.9 | 331.7 | 549.5 | 883.3 | 1,239.1 |
| 1968 | 145.7 | 5.1 | 7.7 | 6.0 | 6.4 | 16.3 | 64.9 | 177.9 | 329.3 | 549.3 | 880.1 | 1,254.9 |

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|--|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| MALIGNANT NEOPLASMS INCLUDING NEOPLASMS OF LYMPHATIC AND HEMATOPOIETIC TISSUES (140-209) | | | | | | | | | | | | |
| ALL OTHER, BOTH SEXES | | | | | | | | | | | | |
| 1978 | 149.2 | 3.9 | 3.9 | 4.1 | 6.1 | 15.2 | 69.7 | 252.9 | 548.0 | 897.6 | 1,165.1 | 1,378.6 |
| 1977 | 148.1 | * | 4.6 | 4.2 | 6.0 | 15.2 | 70.7 | 258.3 | 552.1 | 894.7 | 1,135.6 | 1,358.8 |
| 1976 | 144.8 | * | 4.1 | 4.2 | 6.1 | 14.9 | 67.3 | 253.9 | 549.4 | 884.7 | 1,106.0 | 1,302.6 |
| 1975 | 142.2 | * | 4.9 | 4.3 | 5.9 | 15.5 | 71.8 | 252.1 | 544.0 | 879.7 | 1,082.0 | 1,200.7 |
| 1974 | 142.4 | 4.9 | 4.0 | 4.5 | 6.7 | 16.1 | 75.2 | 255.4 | 550.7 | 895.9 | 1,051.7 | 1,217.8 |
| 1973 | 141.5 | * | 5.4 | 5.1 | 6.9 | 17.1 | 75.1 | 262.8 | 543.5 | 886.6 | 1,047.7 | 1,199.2 |
| 1972 | 137.4 | 5.2 | 4.8 | 4.2 | 6.1 | 18.8 | 80.9 | 254.2 | 520.5 | 866.6 | 1,022.4 | 1,151.2 |
| 1971 | 136.9 | 4.0 | 5.7 | 5.0 | 7.4 | 19.5 | 80.1 | 249.8 | 526.3 | 841.9 | 1,045.0 | 1,146.7 |
| 1970 | 134.4 | 4.0 | 6.3 | 4.5 | 7.5 | 18.4 | 82.3 | 250.3 | 507.1 | 819.9 | 1,081.4 | 969.5 |
| 1969 | 135.4 | * | 5.6 | 4.9 | 6.9 | 20.8 | 85.1 | 255.6 | 527.8 | 817.6 | 1,030.9 | 1,038.5 |
| 1968 | 135.1 | 4.8 | 6.2 | 4.8 | 7.6 | 21.1 | 84.6 | 253.7 | 524.8 | 818.8 | 1,020.7 | 1,091.2 |
| ALL OTHER, MALE | | | | | | | | | | | | |
| 1978 | 181.7 | * | 4.7 | 4.7 | 7.9 | 13.2 | 67.9 | 307.6 | 725.6 | 1,258.2 | 1,689.1 | 2,070.7 |
| 1977 | 180.3 | * | 4.6 | 4.8 | 6.3 | 13.6 | 69.7 | 307.8 | 727.1 | 1,251.6 | 1,662.1 | 2,069.6 |
| 1976 | 176.7 | * | 4.2 | 4.6 | 7.5 | 13.1 | 65.9 | 300.8 | 735.2 | 1,239.7 | 1,601.6 | 1,821.8 |
| 1975 | 173.3 | * | 4.7 | 4.4 | 7.0 | 14.2 | 71.1 | 299.6 | 718.0 | 1,227.0 | 1,557.9 | 1,753.7 |
| 1974 | 171.6 | * | 4.9 | 5.0 | 7.5 | 13.0 | 69.9 | 300.3 | 711.2 | 1,227.9 | 1,557.6 | 1,750.0 |
| 1973 | 168.6 | * | 6.1 | 5.1 | 7.9 | 14.8 | 68.0 | 300.3 | 699.2 | 1,194.9 | 1,521.4 | 1,729.2 |
| 1972 | 164.2 | * | 5.4 | 4.5 | 7.0 | 17.6 | 79.5 | 288.1 | 676.6 | 1,159.3 | 1,475.6 | 1,583.0 |
| 1971 | 161.2 | * | 4.6 | 5.6 | 8.4 | 16.7 | 73.8 | 275.9 | 667.9 | 1,120.0 | 1,473.9 | 1,557.4 |
| 1970 | 161.0 | * | 7.0 | 5.0 | 9.0 | 17.1 | 76.0 | 288.2 | 642.9 | 1,099.2 | 1,536.9 | 1,268.4 |
| 1969 | 160.8 | * | 5.6 | 5.1 | 7.8 | 18.1 | 78.8 | 284.0 | 664.8 | 1,100.2 | 1,451.6 | 1,447.7 |
| 1968 | 159.2 | * | 7.3 | 5.4 | 8.3 | 17.9 | 78.5 | 282.9 | 649.3 | 1,091.6 | 1,404.8 | 1,458.5 |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|--|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| MALIGNANT NEOPLASMS INCLUDING NEOPLASMS OF LYMPHATIC AND HEMATOPOIETIC TISSUES (140-209) | | | | | | | | | | | | |
| ALL OTHER, FEMALE | | | | | | | | | | | | |
| 1978 | 119.7 | * | 3.1 | 3.5 | 4.4 | 16.9 | 71.3 | 207.7 | 400.6 | 618.4 | 809.8 | 1,022.9 |
| 1977 | 118.9 | * | 4.5 | 3.6 | 5.7 | 16.5 | 71.5 | 217.3 | 406.0 | 614.3 | 773.9 | 976.0 |
| 1976 | 115.8 | * | 4.0 | 3.9 | 4.7 | 16.4 | 68.5 | 214.6 | 393.9 | 603.9 | 767.3 | 1,011.2 |
| 1975 | 113.9 | * | 5.0 | 4.3 | 4.7 | 16.5 | 72.4 | 211.9 | 397.1 | 603.5 | 753.7 | 866.7 |
| 1974 | 115.8 | * | 3.2 | 4.0 | 6.0 | 18.8 | 79.7 | 217.4 | 414.0 | 629.9 | 702.7 | 904.7 |
| 1973 | 116.8 | * | 4.7 | 5.0 | 6.0 | 19.0 | 81.2 | 230.7 | 410.8 | 637.6 | 717.3 | 877.2 |
| 1972 | 112.8 | * | 4.2 | 3.9 | 5.2 | 19.8 | 82.0 | 225.2 | 385.9 | 630.3 | 699.0 | 884.2 |
| 1971 | 114.6 | * | 6.8 | 4.3 | 6.6 | 21.9 | 85.4 | 226.9 | 402.5 | 614.5 | 727.9 | 882.2 |
| 1970 | 110.0 | * | 5.5 | 3.9 | 6.2 | 19.6 | 87.5 | 217.1 | 387.1 | 589.6 | 734.7 | 758.9 |
| 1969 | 112.0 | * | 5.5 | 4.6 | 6.1 | 23.1 | 90.3 | 230.8 | 406.3 | 585.1 | 706.5 | 761.5 |
| 1968 | 112.9 | * | 5.0 | 4.3 | 7.0 | 23.8 | 89.8 | 228.1 | 413.0 | 592.5 | 725.9 | 844.3 |
| BLACK, BOTH SEXES | | | | | | | | | | | | |
| 1978 | 162.0 | * | 4.0 | 4.2 | 6.4 | 16.1 | 76.8 | 276.0 | 591.8 | 948.9 | 1,177.9 | 1,408.7 |
| 1977 | 160.7 | * | 4.7 | 4.2 | 6.2 | 16.6 | 77.8 | 283.6 | 592.4 | 942.4 | 1,153.1 | 1,367.1 |
| 1976 | 156.5 | * | 4.3 | 4.3 | 6.3 | 15.9 | 73.6 | 275.9 | 587.4 | 933.6 | 1,118.5 | 1,341.2 |
| 1975 | 152.7 | * | 5.2 | 4.3 | 6.2 | 16.3 | 78.2 | 274.4 | 577.0 | 923.1 | 1,090.5 | 1,200.0 |
| 1974 | 151.8 | 5.4 | 3.8 | 4.5 | 6.9 | 17.2 | 81.6 | 275.5 | 579.5 | 938.9 | 1,058.5 | 1,228.1 |
| 1973 | 150.9 | * | 5.4 | 5.3 | 7.2 | 18.3 | 81.6 | 283.0 | 573.6 | 927.5 | 1,058.5 | 1,221.1 |
| 1972 | 145.3 | 5.0 | 5.1 | 4.4 | 6.3 | 20.0 | 88.3 | 270.5 | 547.0 | 901.5 | 1,029.4 | 1,161.8 |
| 1971 | 144.5 | 4.0 | 5.8 | 4.9 | 7.6 | 20.5 | 85.5 | 266.9 | 551.4 | 875.9 | 1,065.6 | 1,163.6 |
| 1970 | 143.1 | 4.3 | 6.6 | 4.4 | 7.8 | 20.0 | 88.6 | 266.9 | 535.6 | 860.3 | 1,127.8 | 1,024.8 |
| 1969 | 141.7 | * | 5.6 | 4.7 | 7.2 | 22.1 | 90.8 | 268.8 | 550.9 | 845.0 | 1,053.4 | 1,054.6 |
| 1968 | 140.9 | 4.7 | 6.2 | 4.9 | 7.8 | 21.9 | 89.8 | 266.7 | 546.6 | 845.5 | 1,035.9 | 1,079.3 |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|--|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| MALIGNANT NEOPLASMS INCLUDING NEOPLASMS OF LYMPHATIC AND HEMATOPOIETIC TISSUES (140-209) | | | | | | | | | | | | |
| BLACK, MALE | | | | | | | | | | | | |
| 1978 | 198.3 | * | 4.9 | 4.8 | 8.2 | 14.0 | 74.7 | 340.3 | 792.7 | 1,360.1 | 1,720.5 | 2,123.5 |
| 1977 | 196.1 | * | 4.5 | 4.8 | 6.5 | 14.6 | 77.7 | 340.8 | 791.2 | 1,343.3 | 1,712.3 | 2,083.7 |
| 1976 | 191.6 | * | 4.5 | 4.7 | 7.9 | 13.9 | 73.3 | 330.0 | 794.9 | 1,335.3 | 1,624.8 | 1,923.4 |
| 1975 | 187.0 | * | 5.1 | 4.5 | 7.5 | 15.3 | 78.6 | 329.5 | 771.5 | 1,316.5 | 1,582.4 | 1,813.0 |
| 1974 | 183.2 | * | 4.8 | 4.9 | 7.8 | 13.9 | 76.3 | 326.2 | 759.2 | 1,307.5 | 1,587.1 | 1,790.7 |
| 1973 | 179.8 | * | 6.2 | 5.4 | 8.3 | 15.7 | 74.7 | 324.4 | 748.6 | 1,269.2 | 1,554.9 | 1,814.6 |
| 1972 | 173.6 | * | 5.7 | 4.7 | 7.3 | 18.4 | 86.7 | 308.2 | 722.5 | 1,225.6 | 1,495.4 | 1,620.0 |
| 1971 | 170.3 | * | 4.5 | 5.7 | 8.5 | 17.3 | 79.7 | 297.1 | 709.0 | 1,181.2 | 1,526.3 | 1,605.0 |
| 1970 | 171.6 | * | 7.6 | 4.8 | 9.4 | 18.8 | 81.3 | 311.2 | 689.2 | 1,168.9 | 1,624.8 | 1,387.0 |
| 1969 | 168.0 | * | 5.8 | 5.0 | 7.9 | 19.1 | 84.2 | 301.4 | 703.7 | 1,149.2 | 1,493.4 | 1,521.6 |
| 1968 | 165.5 | * | 7.3 | 5.3 | 8.6 | 18.7 | 84.1 | 300.0 | 683.9 | 1,136.0 | 1,440.0 | 1,471.4 |
| BLACK, FEMALE | | | | | | | | | | | | |
| 1978 | 129.3 | * | 3.2 | 3.6 | 4.7 | 18.0 | 78.5 | 223.1 | 427.9 | 642.7 | 815.6 | 1,040.4 |
| 1977 | 128.7 | * | 4.9 | 3.6 | 6.0 | 18.4 | 77.9 | 236.2 | 430.2 | 641.5 | 785.0 | 993.6 |
| 1976 | 124.8 | * | 4.1 | 3.9 | 4.7 | 17.7 | 73.8 | 231.1 | 415.7 | 628.9 | 779.1 | 1,033.7 |
| 1975 | 121.7 | * | 5.3 | 4.2 | 5.1 | 17.3 | 77.8 | 228.3 | 415.6 | 624.8 | 757.6 | 864.3 |
| 1974 | 123.4 | * | 2.7 | 4.1 | 6.0 | 20.1 | 86.0 | 233.0 | 430.5 | 657.5 | 701.0 | 917.9 |
| 1973 | 124.6 | * | 4.6 | 5.2 | 6.2 | 20.5 | 87.3 | 247.7 | 428.1 | 663.4 | 719.9 | 887.7 |
| 1972 | 119.5 | * | 4.5 | 4.0 | 5.4 | 21.5 | 89.8 | 238.3 | 399.8 | 650.5 | 697.9 | 900.0 |
| 1971 | 121.0 | * | 7.1 | 4.2 | 6.7 | 23.2 | 90.4 | 240.7 | 418.0 | 634.1 | 734.7 | 900.0 |
| 1970 | 117.3 | * | 5.7 | 4.0 | 6.4 | 20.9 | 94.6 | 228.6 | 404.8 | 615.8 | 763.3 | 791.5 |
| 1969 | 117.8 | * | 5.5 | 4.5 | 6.6 | 24.6 | 96.3 | 240.6 | 420.7 | 603.0 | 725.1 | 766.7 |
| 1968 | 118.4 | * | 5.1 | 4.5 | 7.1 | 24.6 | 94.6 | 237.6 | 427.8 | 610.3 | 735.6 | 853.6 |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|--|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| MALIGNANT NEOPLASMS OF BUCCAL CAVITY AND PHARYNX (140-149) | | | | | | | | | | | | |
| BOTH SEXES | | | | | | | | | | | | |
| 1978 | 3.8 | * | * | * | 0.1 | 0.2 | 1.0 | 5.5 | 11.5 | 16.3 | 18.4 | 23.3 |
| 1977 | 3.8 | * | * | * | 0.1 | 0.1 | 1.0 | 5.4 | 12.4 | 16.6 | 18.6 | 25.1 |
| 1976 | 3.7 | * | * | * | 0.1 | 0.2 | 1.0 | 5.2 | 12.3 | 16.2 | 17.6 | 24.6 |
| 1975 | 3.8 | * | * | * | 0.1 | 0.2 | 1.0 | 5.5 | 12.3 | 16.8 | 17.9 | 25.5 |
| 1974 | 3.7 | * | * | * | 0.1 | 0.1 | 1.1 | 5.2 | 12.4 | 15.8 | 20.1 | 25.3 |
| 1973 | 3.7 | * | * | * | 0.1 | 0.2 | 1.1 | 5.4 | 12.2 | 16.5 | 18.2 | 27.5 |
| 1972 | 3.6 | * | * | * | 0.1 | 0.1 | 1.0 | 5.1 | 12.4 | 15.2 | 18.3 | 26.6 |
| 1971 | 3.7 | * | * | * | 0.1 | 0.2 | 1.1 | 5.4 | 12.6 | 15.9 | 19.5 | 29.7 |
| 1970 | 3.7 | * | * | * | 0.1 | 0.2 | 1.2 | 5.2 | 12.8 | 15.5 | 21.1 | 29.2 |
| 1969 | 3.8 | * | * | * | 0.1 | 0.2 | 1.3 | 5.4 | 12.8 | 15.2 | 21.0 | 31.8 |
| 1968 | 3.7 | * | * | 0.0 | 0.1 | 0.1 | 1.1 | 5.7 | 12.0 | 14.9 | 20.6 | 33.9 |
| MALE | | | | | | | | | | | | |
| 1978 | 5.4 | * | * | * | 0.1 | 0.2 | 1.4 | 8.4 | 17.9 | 26.7 | 32.1 | 39.1 |
| 1977 | 5.5 | * | * | * | * | 0.2 | 1.5 | 8.2 | 18.9 | 27.6 | 32.5 | 45.6 |
| 1976 | 5.4 | * | * | * | 0.1 | 0.2 | 1.6 | 8.0 | 19.0 | 26.9 | 32.0 | 40.3 |
| 1975 | 5.4 | * | * | * | 0.1 | 0.2 | 1.4 | 8.2 | 18.7 | 28.6 | 30.4 | 42.4 |
| 1974 | 5.5 | * | * | * | 0.1 | 0.2 | 1.7 | 7.8 | 19.4 | 26.8 | 35.7 | 42.8 |
| 1973 | 5.4 | * | * | * | 0.1 | 0.2 | 1.5 | 8.1 | 18.7 | 28.2 | 32.2 | 45.5 |
| 1972 | 5.2 | * | * | * | 0.1 | 0.2 | 1.4 | 7.6 | 19.0 | 26.3 | 32.5 | 46.7 |
| 1971 | 5.5 | * | * | * | 0.1 | 0.3 | 1.6 | 7.9 | 19.6 | 27.6 | 33.6 | 46.1 |
| 1970 | 5.6 | * | * | * | * | 0.2 | 1.7 | 7.9 | 20.1 | 26.8 | 36.6 | 47.0 |
| 1969 | 5.7 | * | * | * | 0.2 | 0.3 | 1.8 | 8.1 | 20.7 | 26.9 | 36.8 | 52.5 |
| 1968 | 5.5 | * | * | * | 0.2 | 0.2 | 1.6 | 8.7 | 19.2 | 26.5 | 35.3 | 56.1 |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|--|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| MALIGNANT NEOPLASMS OF BUCCAL CAVITY AND PHARYNX (140-149) | | | | | | | | | | | | |
| FEMALE | | | | | | | | | | | | |
| 1978 | 2.2 | * | * | * | * | 0.2 | 0.5 | 2.7 | 5.9 | 8.4 | 10.2 | 16.1 |
| 1977 | 2.2 | * | * | * | * | * | 0.5 | 2.8 | 6.7 | 8.1 | 10.3 | 15.7 |
| 1976 | 2.1 | * | * | * | * | 0.2 | 0.6 | 2.6 | 6.4 | 7.9 | 8.9 | 17.3 |
| 1975 | 2.2 | * | * | * | * | 0.2 | 0.6 | 3.0 | 6.6 | 7.8 | 10.2 | 17.3 |
| 1974 | 2.1 | * | * | * | * | * | 0.6 | 2.7 | 6.3 | 7.4 | 10.6 | 16.7 |
| 1973 | 2.1 | * | * | * | * | * | 0.6 | 2.9 | 6.4 | 7.5 | 9.4 | 18.4 |
| 1972 | 2.0 | * | * | * | * | * | 0.7 | 2.8 | 6.5 | 6.7 | 9.3 | 16.3 |
| 1971 | 2.1 | * | * | * | * | * | 0.6 | 3.2 | 6.4 | 6.8 | 10.4 | 21.1 |
| 1970 | 2.0 | * | * | * | * | * | 0.7 | 2.8 | 6.2 | 6.7 | 10.8 | 19.2 |
| 1969 | 1.9 | * | * | * | * | * | 0.8 | 3.0 | 5.6 | 6.0 | 10.4 | 20.6 |
| 1968 | 1.9 | * | * | * | * | * | 0.7 | 3.0 | 5.4 | 5.8 | 10.5 | 21.8 |
| WHITE, BOTH SEXES | | | | | | | | | | | | |
| 1978 | 3.7 | * | * | * | 0.1 | 0.2 | 0.6 | 4.7 | 10.8 | 16.2 | 18.7 | 24.2 |
| 1977 | 3.8 | * | * | * | 0.1 | 0.1 | 0.7 | 4.6 | 11.8 | 16.6 | 18.8 | 26.0 |
| 1976 | 3.7 | * | * | * | 0.1 | 0.2 | 0.8 | 4.3 | 11.8 | 16.0 | 18.1 | 25.5 |
| 1975 | 3.8 | * | * | * | 0.1 | 0.1 | 0.7 | 4.7 | 11.8 | 16.9 | 18.2 | 26.3 |
| 1974 | 3.8 | * | * | * | 0.1 | 0.1 | 0.8 | 4.5 | 12.1 | 15.8 | 20.6 | 26.3 |
| 1973 | 3.7 | * | * | * | 0.1 | 0.1 | 0.8 | 4.6 | 11.8 | 16.4 | 18.6 | 29.1 |
| 1972 | 3.6 | * | * | * | 0.1 | 0.1 | 0.8 | 4.4 | 12.0 | 15.4 | 18.5 | 27.9 |
| 1971 | 3.8 | * | * | * | * | 0.2 | 0.9 | 4.9 | 12.5 | 16.1 | 20.0 | 31.0 |
| 1970 | 3.8 | * | * | * | 0.1 | 0.1 | 1.0 | 4.7 | 12.5 | 15.6 | 21.2 | 30.3 |
| 1969 | 3.8 | * | * | * | 0.1 | 0.2 | 0.9 | 4.8 | 12.6 | 15.4 | 21.6 | 33.2 |
| 1968 | 3.7 | * | * | * | 0.1 | 0.1 | 0.9 | 5.1 | 11.8 | 15.0 | 21.1 | 35.3 |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|--|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| MALIGNANT NEOPLASMS OF BUCCAL CAVITY AND PHARYNX (140-149) | | | | | | | | | | | | |
| WHITE, MALE | | | | | | | | | | | | |
| 1978 | 5.3 | * | * | * | * | 0.2 | 0.9 | 7.2 | 16.7 | 26.2 | 32.9 | 40.1 |
| 1977 | 5.4 | * | * | * | * | 0.2 | 1.0 | 6.8 | 17.7 | 27.4 | 32.8 | 47.3 |
| 1976 | 5.3 | * | * | * | * | 0.2 | 1.1 | 6.4 | 18.0 | 26.5 | 33.0 | 41.9 |
| 1975 | 5.3 | * | * | * | * | 0.2 | 0.9 | 6.9 | 17.6 | 28.4 | 31.1 | 44.4 |
| 1974 | 5.5 | * | * | * | * | 0.2 | 1.2 | 6.9 | 18.7 | 26.9 | 36.4 | 45.0 |
| 1973 | 5.3 | * | * | * | 0.1 | * | 1.2 | 6.9 | 17.9 | 27.8 | 33.2 | 48.8 |
| 1972 | 5.2 | * | * | * | 0.1 | 0.2 | 1.0 | 6.5 | 18.1 | 26.7 | 33.3 | 50.1 |
| 1971 | 5.5 | * | * | * | * | 0.2 | 1.3 | 6.9 | 19.2 | 27.9 | 34.7 | 48.4 |
| 1970 | 5.6 | * | * | * | * | * | 1.4 | 7.1 | 19.7 | 27.1 | 36.7 | 50.3 |
| 1969 | 5.7 | * | * | * | * | 0.2 | 1.3 | 7.0 | 20.3 | 27.4 | 37.8 | 54.7 |
| 1968 | 5.6 | * | * | * | 0.2 | * | 1.3 | 7.7 | 18.8 | 26.8 | 36.4 | 58.6 |
| WHITE, FEMALE | | | | | | | | | | | | |
| 1978 | 2.3 | * | * | * | * | 0.2 | 0.4 | 2.4 | 5.7 | 8.5 | 10.3 | 17.2 |
| 1977 | 2.3 | * | * | * | * | * | 0.4 | 2.4 | 6.6 | 8.3 | 10.6 | 16.3 |
| 1976 | 2.2 | * | * | * | * | 0.2 | 0.5 | 2.2 | 6.3 | 8.0 | 9.2 | 18.0 |
| 1975 | 2.3 | * | * | * | * | * | 0.5 | 2.6 | 6.7 | 8.0 | 10.5 | 17.7 |
| 1974 | 2.1 | * | * | * | * | * | 0.5 | 2.3 | 6.3 | 7.4 | 11.0 | 17.1 |
| 1973 | 2.1 | * | * | * | * | * | 0.5 | 2.6 | 6.4 | 7.6 | 9.5 | 19.3 |
| 1972 | 2.0 | * | * | * | * | * | 0.5 | 2.4 | 6.5 | 6.7 | 9.2 | 16.7 |
| 1971 | 2.2 | * | * | * | * | * | 0.5 | 2.9 | 6.4 | 7.0 | 10.6 | 22.1 |
| 1970 | 2.1 | * | * | * | * | * | 0.6 | 2.6 | 6.1 | 6.8 | 11.0 | 19.3 |
| 1969 | 2.0 | * | * | * | * | * | 0.6 | 2.6 | 5.7 | 6.0 | 10.8 | 21.8 |
| 1968 | 1.9 | * | * | * | * | * | 0.6 | 2.8 | 5.5 | 5.8 | 10.7 | 22.7 |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|--|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| MALIGNANT NEOPLASMS OF BUCCAL CAVITY AND PHARYNX (140-149) | | | | | | | | | | | | |
| ALL OTHER, BOTH SEXES | | | | | | | | | | | | |
| 1978 | 3.9 | * | * | * | * | * | 3.1 | 11.4 | 17.9 | 17.8 | 15.1 | 12.5 |
| 1977 | 4.0 | * | * | * | * | * | 3.0 | 12.1 | 18.2 | 16.2 | 16.5 | 15.0 |
| 1976 | 3.9 | * | * | * | * | * | 2.7 | 12.7 | 16.8 | 17.6 | 12.8 | 14.4 |
| 1975 | 3.8 | * | * | * | * | * | 2.9 | 12.6 | 16.8 | 16.2 | 13.9 | 15.8 |
| 1974 | 3.5 | * | * | * | * | * | 3.3 | 10.5 | 15.4 | 15.8 | 14.6 | * |
| 1973 | 3.7 | * | * | * | * | * | 2.8 | 11.5 | 15.7 | 18.0 | 13.3 | * |
| 1972 | 3.5 | * | * | * | * | * | 2.7 | 10.9 | 16.4 | 13.8 | 16.2 | * |
| 1971 | 3.3 | * | * | * | * | * | 2.7 | 10.4 | 14.5 | 13.9 | 14.2 | * |
| 1970 | 3.4 | * | * | * | * | * | 2.9 | 9.4 | 14.9 | 14.4 | 19.9 | 17.9 |
| 1969 | 3.5 | * | * | * | * | * | 3.7 | 11.7 | 14.3 | 13.3 | 13.1 | * |
| 1968 | 3.3 | * | * | * | * | * | 2.8 | 11.0 | 13.4 | 14.0 | 14.1 | * |
| ALL OTHER, MALE | | | | | | | | | | | | |
| 1978 | 6.2 | * | * | * | * | * | 5.5 | 18.8 | 29.9 | 31.2 | 24.4 | * |
| 1977 | 6.3 | * | * | * | * | * | 5.0 | 20.1 | 30.7 | 29.1 | 30.3 | * |
| 1976 | 6.2 | * | * | * | * | * | 4.6 | 21.1 | 28.1 | 31.3 | 23.0 | * |
| 1975 | 6.0 | * | * | * | * | * | 5.0 | 20.0 | 29.5 | 29.7 | 23.1 | * |
| 1974 | 5.5 | * | * | * | * | * | 5.2 | 16.2 | 26.5 | 26.3 | 27.9 | * |
| 1973 | 5.8 | * | * | * | * | * | 4.4 | 18.7 | 26.2 | 31.6 | 21.9 | * |
| 1972 | 5.3 | * | * | * | * | * | 4.0 | 16.8 | 27.7 | 23.0 | 24.0 | * |
| 1971 | 5.2 | * | * | * | * | * | 4.5 | 16.4 | 23.6 | 24.8 | 21.3 | * |
| 1970 | 5.3 | * | * | * | * | * | 4.4 | 15.1 | 23.8 | 24.4 | 34.7 | * |
| 1969 | 5.6 | * | * | * | * | * | 5.5 | 18.1 | 24.7 | 22.1 | 24.5 | * |
| 1968 | 5.3 | * | * | * | * | * | 4.1 | 17.9 | 23.4 | 24.2 | 22.0 | * |
| ALL OTHER, FEMALE | | | | | | | | | | | | |
| 1978 | 1.9 | * | * | * | * | * | * | 5.4 | 8.0 | 7.3 | 8.7 | * |
| 1977 | 1.8 | * | * | * | * | * | 1.4 | 5.4 | 7.8 | 6.1 | 7.0 | * |
| 1976 | 1.8 | * | * | * | * | * | * | 5.8 | 7.4 | 6.8 | 5.9 | * |
| 1975 | 1.8 | * | * | * | * | * | * | 6.4 | 6.1 | 5.4 | 7.6 | * |
| 1974 | 1.7 | * | * | * | * | * | 1.7 | 5.7 | 5.8 | 7.3 | * | * |
| 1973 | 1.8 | * | * | * | * | * | 1.4 | 5.2 | 6.8 | 7.1 | 7.2 | * |
| 1972 | 1.9 | * | * | * | * | * | 1.6 | 5.7 | 6.7 | 6.4 | 10.6 | * |
| 1971 | 1.6 | * | * | * | * | * | * | 5.3 | 6.6 | 5.0 | 8.9 | * |
| 1970 | 1.7 | * | * | * | * | * | 1.6 | 4.3 | 7.0 | 6.2 | 8.7 | * |
| 1969 | 1.6 | * | * | * | * | * | 2.2 | 6.1 | 5.0 | 6.0 | * | * |
| 1968 | 1.5 | * | * | * | * | * | 1.8 | 5.0 | 4.5 | 5.6 | * | * |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|--|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| MALIGNANT NEOPLASMS OF BUCCAL CAVITY AND PHARYNX (140-149) | | | | | | | | | | | | |
| BLACK, BOTH SEXES | | | | | | | | | | | | |
| 1978 | 4.2 | * | * | * | * | * | 3.5 | 12.7 | 19.5 | 18.6 | 13.9 | * |
| 1977 | 4.3 | * | * | * | * | * | 3.4 | 13.4 | 19.3 | 17.2 | 15.6 | * |
| 1976 | 4.2 | * | * | * | * | * | 2.8 | 13.9 | 18.0 | 18.5 | 12.9 | 15.4 |
| 1975 | 4.0 | * | * | * | * | * | 3.1 | 13.6 | 17.7 | 16.6 | 12.9 | * |
| 1974 | 3.7 | * | * | * | * | * | 3.4 | 11.4 | 15.9 | 15.9 | 14.1 | * |
| 1973 | 3.9 | * | * | * | * | * | 3.0 | 12.5 | 17.0 | 18.0 | 13.8 | * |
| 1972 | 3.7 | * | * | * | * | * | 2.7 | 11.6 | 17.1 | 14.2 | 16.4 | * |
| 1971 | 3.5 | * | * | * | * | * | 2.8 | 11.2 | 14.7 | 13.8 | 14.4 | * |
| 1970 | 3.6 | * | * | * | * | * | 3.2 | 10.2 | 15.7 | 14.1 | 20.6 | 18.7 |
| 1969 | 3.6 | * | * | * | * | * | 3.8 | 12.1 | 14.6 | 13.6 | 13.2 | * |
| 1968 | 3.4 | * | * | * | * | * | 3.2 | 11.3 | 13.8 | 13.9 | 14.0 | * |
| BLACK, MALE | | | | | | | | | | | | |
| 1978 | 6.8 | * | * | * | * | * | 6.1 | 20.9 | 33.2 | 33.6 | 23.4 | * |
| 1977 | 6.8 | * | * | * | * | * | 5.5 | 22.5 | 33.0 | 31.7 | 29.4 | * |
| 1976 | 6.7 | * | * | * | * | * | 4.7 | 23.1 | 30.3 | 34.0 | 23.9 | * |
| 1975 | 6.4 | * | * | * | * | * | 5.3 | 21.6 | 31.6 | 30.9 | 23.0 | * |
| 1974 | 5.7 | * | * | * | * | * | 5.3 | 17.4 | 27.9 | 26.7 | 27.8 | * |
| 1973 | 6.1 | * | * | * | * | * | 4.8 | 20.3 | 28.9 | 31.7 | 23.5 | * |
| 1972 | 5.5 | * | * | * | * | * | 3.8 | 17.8 | 29.4 | 23.9 | 23.4 | * |
| 1971 | 5.4 | * | * | * | * | * | 4.6 | 17.7 | 24.9 | 24.6 | 21.5 | * |
| 1970 | 5.6 | * | * | * | * | * | 4.9 | 16.7 | 25.6 | 23.9 | 36.2 | * |
| 1969 | 5.8 | * | * | * | * | * | 5.6 | 18.8 | 26.1 | 23.3 | 24.6 | * |
| 1968 | 5.4 | * | * | * | * | * | 4.6 | 18.2 | 24.3 | 24.3 | 21.2 | * |
| BLACK, FEMALE | | | | | | | | | | | | |
| 1978 | 2.0 | * | * | * | * | * | * | 5.9 | 8.3 | 7.5 | 7.5 | * |
| 1977 | 1.9 | * | * | * | * | * | 1.7 | 5.8 | 8.1 | 6.3 | 6.5 | * |
| 1976 | 1.9 | * | * | * | * | * | * | 6.4 | 7.8 | 6.7 | * | * |
| 1975 | 1.8 | * | * | * | * | * | * | 6.8 | 6.2 | 5.7 | 6.1 | * |
| 1974 | 1.8 | * | * | * | * | * | 1.7 | 6.3 | 6.0 | 7.7 | * | * |
| 1973 | 1.9 | * | * | * | * | * | 1.5 | 5.9 | 7.2 | 7.4 | 7.1 | * |
| 1972 | 2.0 | * | * | * | * | * | 1.7 | 6.2 | 6.9 | 6.6 | 11.4 | * |
| 1971 | 1.7 | * | * | * | * | * | * | 5.7 | 6.1 | 5.2 | 9.3 | * |
| 1970 | 1.8 | * | * | * | * | * | 1.8 | 4.6 | 7.3 | 6.4 | 9.1 | * |
| 1969 | 1.7 | * | * | * | * | * | 2.4 | 6.4 | 4.8 | 5.9 | * | * |
| 1968 | 1.6 | * | * | * | * | * | 2.0 | 5.3 | 4.7 | 5.5 | * | * |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|--|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| MALIGNANT NEOPLASMS OF DIGESTIVE ORGANS AND PERITONEUM (150-159) | | | | | | | | | | | | |
| BOTH SEXES | | | | | | | | | | | | |
| 1978 | 47.5 | * | 0.2 | 0.1 | 0.3 | 1.6 | 8.3 | 36.8 | 101.6 | 218.9 | 385.0 | 540.5 |
| 1977 | 46.8 | * | 0.2 | 0.1 | 0.4 | 1.8 | 8.9 | 36.1 | 103.6 | 217.2 | 381.4 | 530.6 |
| 1976 | 46.8 | * | 0.2 | 0.1 | 0.3 | 1.7 | 8.8 | 36.9 | 103.4 | 220.3 | 386.5 | 538.1 |
| 1975 | 46.2 | * | * | 0.1 | 0.4 | 1.8 | 8.9 | 37.0 | 103.8 | 220.6 | 389.0 | 519.9 |
| 1974 | 46.4 | * | 0.2 | 0.1 | 0.3 | 1.8 | 9.4 | 37.8 | 107.2 | 223.6 | 390.3 | 530.7 |
| 1973 | 45.7 | * | 0.1 | 0.1 | 0.4 | 1.8 | 9.1 | 37.2 | 106.7 | 222.7 | 392.1 | 528.3 |
| 1972 | 45.8 | * | * | 0.1 | 0.4 | 1.8 | 9.8 | 36.6 | 107.0 | 228.6 | 395.4 | 530.0 |
| 1971 | 45.9 | * | 0.2 | 0.1 | 0.5 | 2.1 | 10.0 | 37.4 | 107.1 | 228.0 | 402.0 | 545.2 |
| 1970 | 46.6 | * | 0.2 | 0.1 | 0.4 | 2.0 | 9.9 | 39.0 | 111.0 | 232.8 | 414.7 | 507.6 |
| 1969 | 46.7 | * | 0.2 | 0.1 | 0.3 | 2.2 | 10.3 | 38.8 | 110.1 | 234.8 | 417.1 | 571.8 |
| 1968 | 46.9 | * | * | 0.1 | 0.4 | 2.1 | 10.3 | 40.3 | 112.0 | 238.8 | 416.8 | 573.5 |
| MALE | | | | | | | | | | | | |
| 1978 | 51.1 | * | * | * | 0.3 | 1.8 | 9.6 | 45.3 | 127.5 | 280.2 | 485.2 | 667.5 |
| 1977 | 50.9 | * | * | * | 0.5 | 2.1 | 10.6 | 44.1 | 130.5 | 279.6 | 479.6 | 672.7 |
| 1976 | 50.8 | * | * | * | 0.4 | 1.9 | 9.9 | 45.0 | 128.9 | 282.2 | 497.1 | 650.8 |
| 1975 | 50.3 | * | * | * | 0.4 | 2.1 | 10.2 | 45.0 | 128.5 | 284.7 | 491.8 | 636.2 |
| 1974 | 50.3 | * | * | 0.1 | 0.4 | 2.0 | 10.4 | 45.3 | 134.2 | 283.4 | 492.0 | 651.0 |
| 1973 | 49.6 | * | * | * | 0.5 | 1.9 | 10.3 | 44.3 | 132.0 | 282.0 | 490.5 | 659.7 |
| 1972 | 50.1 | * | * | 0.1 | 0.5 | 2.1 | 10.8 | 43.9 | 136.4 | 287.4 | 500.1 | 609.2 |
| 1971 | 50.3 | * | * | * | 0.6 | 2.5 | 11.0 | 44.6 | 134.4 | 287.6 | 501.2 | 655.7 |
| 1970 | 51.1 | * | * | * | 0.6 | 2.3 | 11.4 | 45.9 | 139.0 | 293.3 | 507.5 | 583.7 |
| 1969 | 51.5 | * | * | * | 0.4 | 2.5 | 11.3 | 46.3 | 136.6 | 297.9 | 514.3 | 679.5 |
| 1968 | 51.8 | * | * | * | 0.5 | 2.4 | 11.8 | 46.9 | 138.8 | 301.5 | 514.6 | 658.2 |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|--|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| MALIGNANT NEOPLASMS OF DIGESTIVE ORGANS AND PERITONEUM (150-159) | | | | | | | | | | | | |
| FEMALE | | | | | | | | | | | | |
| 1978 | 44.0 | * | * | * | 0.3 | 1.3 | 7.0 | 28.9 | 78.7 | 172.1 | 325.4 | 483.1 |
| 1977 | 43.0 | * | * | * | 0.3 | 1.5 | 7.3 | 28.7 | 79.8 | 169.5 | 322.6 | 465.3 |
| 1976 | 43.0 | * | * | * | 0.3 | 1.4 | 7.7 | 29.3 | 80.8 | 172.9 | 319.7 | 485.1 |
| 1975 | 42.4 | * | * | * | 0.3 | 1.5 | 7.7 | 29.6 | 81.9 | 171.5 | 326.4 | 463.6 |
| 1974 | 42.6 | * | * | * | 0.3 | 1.7 | 8.5 | 30.9 | 83.3 | 177.7 | 327.6 | 471.2 |
| 1973 | 42.0 | * | * | * | 0.3 | 1.7 | 7.9 | 30.6 | 84.3 | 177.2 | 330.7 | 462.1 |
| 1972 | 41.7 | * | * | * | 0.2 | 1.6 | 8.9 | 29.9 | 80.8 | 183.2 | 328.7 | 489.4 |
| 1971 | 41.7 | * | * | * | 0.4 | 1.7 | 9.1 | 30.7 | 82.6 | 181.9 | 337.7 | 487.5 |
| 1970 | 42.3 | * | * | * | 0.3 | 1.6 | 8.6 | 32.5 | 86.0 | 185.8 | 353.3 | 465.0 |
| 1969 | 42.1 | * | * | * | 0.3 | 2.0 | 9.3 | 31.9 | 86.2 | 185.4 | 352.0 | 513.6 |
| 1968 | 42.3 | * | * | * | 0.4 | 1.9 | 9.0 | 34.2 | 87.8 | 189.4 | 349.9 | 527.6 |
| WHITE, BOTH SEXES | | | | | | | | | | | | |
| 1978 | 48.4 | * | * | * | 0.3 | 1.4 | 7.4 | 33.5 | 96.3 | 214.2 | 386.5 | 545.4 |
| 1977 | 47.6 | * | 0.2 | 0.1 | 0.3 | 1.6 | 8.0 | 32.8 | 98.0 | 211.6 | 383.4 | 536.2 |
| 1976 | 47.6 | * | 0.2 | * | 0.3 | 1.5 | 8.1 | 33.6 | 97.9 | 214.7 | 388.7 | 545.0 |
| 1975 | 47.0 | * | * | * | 0.3 | 1.6 | 8.1 | 33.9 | 98.3 | 215.2 | 390.5 | 530.3 |
| 1974 | 47.2 | * | 0.2 | 0.1 | 0.3 | 1.6 | 8.6 | 34.9 | 101.6 | 217.7 | 393.9 | 540.5 |
| 1973 | 46.5 | * | * | * | 0.3 | 1.7 | 8.3 | 34.0 | 101.4 | 217.0 | 395.4 | 540.3 |
| 1972 | 46.7 | * | * | 0.1 | 0.4 | 1.7 | 9.0 | 33.2 | 102.5 | 224.0 | 400.6 | 541.9 |
| 1971 | 46.7 | * | 0.2 | 0.1 | 0.4 | 1.9 | 9.1 | 34.9 | 102.1 | 223.0 | 405.7 | 555.2 |
| 1970 | 47.6 | * | 0.2 | * | 0.4 | 1.8 | 9.0 | 36.1 | 106.5 | 229.3 | 417.4 | 522.9 |
| 1969 | 47.6 | * | 0.2 | 0.1 | 0.3 | 2.0 | 9.4 | 35.9 | 104.9 | 230.5 | 421.0 | 589.9 |
| 1968 | 47.8 | * | * | 0.1 | 0.4 | 1.9 | 9.4 | 37.6 | 106.6 | 234.6 | 421.1 | 590.1 |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|--|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| MALIGNANT NEOPLASMS OF DIGESTIVE ORGANS AND PERITONEUM (150-159) | | | | | | | | | | | | |
| WHITE, MALE | | | | | | | | | | | | |
| 1978 | 51.5 | * | * | * | 0.2 | 1.7 | 8.5 | 40.4 | 119.8 | 274.0 | 486.5 | 674.1 |
| 1977 | 50.9 | * | * | * | 0.4 | 1.9 | 9.2 | 39.5 | 122.2 | 270.7 | 481.0 | 677.9 |
| 1976 | 50.9 | * | * | * | 0.3 | 1.6 | 8.9 | 41.0 | 120.4 | 273.6 | 498.6 | 662.4 |
| 1975 | 50.3 | * | * | * | 0.4 | 1.8 | 9.1 | 40.5 | 120.0 | 276.3 | 493.3 | 649.3 |
| 1974 | 50.5 | * | * | * | 0.3 | 1.8 | 9.4 | 41.1 | 126.3 | 275.0 | 494.1 | 662.1 |
| 1973 | 49.8 | * | * | * | 0.4 | 1.8 | 9.3 | 40.2 | 124.7 | 273.4 | 495.3 | 674.5 |
| 1972 | 50.5 | * | * | 0.1 | 0.5 | 1.7 | 9.6 | 39.3 | 130.2 | 280.8 | 507.7 | 621.1 |
| 1971 | 50.6 | * | * | * | 0.5 | 2.3 | 9.8 | 41.1 | 126.8 | 281.1 | 506.2 | 671.7 |
| 1970 | 51.5 | * | * | * | 0.6 | 2.1 | 10.3 | 41.6 | 132.5 | 288.1 | 508.5 | 601.5 |
| 1969 | 51.7 | * | * | * | 0.4 | 2.2 | 10.3 | 42.1 | 128.9 | 290.6 | 517.4 | 703.1 |
| 1968 | 52.0 | * | * | * | 0.4 | 2.1 | 10.5 | 42.9 | 130.9 | 294.6 | 518.8 | 676.9 |
| WHITE, FEMALE | | | | | | | | | | | | |
| 1978 | 45.5 | * | * | * | 0.3 | 1.2 | 6.3 | 27.0 | 75.4 | 168.6 | 327.7 | 488.1 |
| 1977 | 44.5 | * | * | * | 0.3 | 1.4 | 6.7 | 26.6 | 76.5 | 166.4 | 325.8 | 472.1 |
| 1976 | 44.4 | * | * | * | 0.3 | 1.3 | 7.3 | 26.8 | 77.8 | 169.8 | 323.2 | 491.1 |
| 1975 | 43.9 | * | * | * | 0.3 | 1.4 | 7.1 | 27.8 | 78.9 | 168.6 | 328.6 | 474.2 |
| 1974 | 44.0 | * | * | * | 0.2 | 1.4 | 7.9 | 29.0 | 79.6 | 174.0 | 332.7 | 481.3 |
| 1973 | 43.2 | * | * | * | 0.3 | 1.5 | 7.3 | 28.3 | 80.5 | 173.8 | 333.6 | 474.3 |
| 1972 | 43.1 | * | * | * | 0.2 | 1.7 | 8.5 | 27.5 | 77.7 | 180.3 | 333.0 | 502.1 |
| 1971 | 43.1 | * | * | * | 0.3 | 1.5 | 8.4 | 29.1 | 79.9 | 178.3 | 341.3 | 495.6 |
| 1970 | 43.9 | * | * | * | 0.3 | 1.4 | 7.7 | 30.9 | 83.0 | 183.8 | 357.7 | 479.9 |
| 1969 | 43.6 | * | * | * | 0.3 | 1.8 | 8.6 | 30.0 | 83.3 | 183.7 | 356.9 | 530.7 |
| 1968 | 43.7 | * | * | * | 0.3 | 1.7 | 8.3 | 32.6 | 84.6 | 187.5 | 354.8 | 544.0 |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|--|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| MALIGNANT NEOPLASMS OF DIGESTIVE ORGANS AND PERITONEUM (150-159) | | | | | | | | | | | | |
| ALL OTHER, BOTH SEXES | | | | | | | | | | | | |
| 1978 | 41.4 | * | * | * | 0.5 | 2.5 | 14.2 | 62.1 | 149.1 | 262.7 | 369.8 | 483.9 |
| 1977 | 41.7 | * | * | * | 0.7 | 2.9 | 15.3 | 61.7 | 154.7 | 271.0 | 361.3 | 465.6 |
| 1976 | 41.5 | * | * | * | 0.7 | 3.0 | 13.6 | 63.0 | 154.0 | 273.9 | 364.1 | 456.2 |
| 1975 | 41.0 | * | * | * | 0.6 | 3.0 | 14.9 | 62.4 | 155.3 | 272.8 | 372.9 | 400.0 |
| 1974 | 40.9 | * | * | * | 0.7 | 3.3 | 14.9 | 62.6 | 159.4 | 280.7 | 351.2 | 416.3 |
| 1973 | 40.4 | * | * | * | 0.7 | 2.8 | 14.9 | 63.9 | 157.7 | 279.4 | 354.3 | 388.2 |
| 1972 | 39.3 | * | * | * | 0.5 | 3.1 | 15.6 | 65.9 | 150.2 | 273.9 | 335.3 | 391.9 |
| 1971 | 40.0 | * | * | * | 0.8 | 3.7 | 16.8 | 59.5 | 155.0 | 277.3 | 357.9 | 430.8 |
| 1970 | 39.8 | * | * | * | 0.6 | 3.2 | 16.8 | 64.3 | 154.1 | 267.3 | 381.0 | 351.3 |
| 1969 | 40.4 | * | * | * | 0.6 | 3.7 | 16.8 | 65.5 | 159.7 | 277.7 | 367.7 | 367.9 |
| 1968 | 40.8 | * | * | * | 0.7 | 3.3 | 17.6 | 65.0 | 164.5 | 281.2 | 362.0 | 388.2 |
| ALL OTHER, MALE | | | | | | | | | | | | |
| 1978 | 49.0 | * | * | * | 0.7 | 3.0 | 17.5 | 85.5 | 199.4 | 337.5 | 470.7 | 600.0 |
| 1977 | 50.7 | * | * | * | 0.9 | 3.4 | 20.2 | 82.8 | 208.7 | 362.7 | 467.0 | 619.6 |
| 1976 | 50.0 | * | * | * | 0.9 | 3.9 | 17.2 | 79.6 | 209.4 | 362.5 | 484.4 | 534.5 |
| 1975 | 50.1 | * | * | * | * | 3.9 | 18.9 | 84.3 | 210.0 | 364.0 | 476.9 | 505.6 |
| 1974 | 49.3 | * | * | * | 0.9 | 3.7 | 17.6 | 82.6 | 209.5 | 363.3 | 470.7 | 536.0 |
| 1973 | 48.0 | * | * | * | 0.9 | 3.0 | 18.0 | 80.7 | 203.3 | 364.2 | 437.9 | 508.3 |
| 1972 | 47.3 | * | * | * | 0.8 | 5.1 | 20.4 | 84.8 | 196.5 | 350.3 | 418.4 | 489.4 |
| 1971 | 48.1 | * | * | * | 1.0 | 4.3 | 20.1 | 76.7 | 208.6 | 349.5 | 448.3 | 497.9 |
| 1970 | 48.7 | * | * | * | 0.9 | 4.0 | 19.6 | 85.3 | 200.3 | 342.6 | 496.2 | 427.6 |
| 1969 | 50.2 | * | * | * | * | 4.7 | 20.1 | 85.3 | 212.1 | 368.0 | 476.6 | 456.8 |
| 1968 | 50.5 | * | * | * | * | 4.1 | 22.2 | 83.5 | 216.4 | 368.3 | 464.5 | 475.6 |
| ALL OTHER, FEMALE | | | | | | | | | | | | |
| 1978 | 34.5 | * | * | * | * | 2.2 | 11.5 | 42.9 | 107.4 | 204.8 | 301.8 | 426.6 |
| 1977 | 33.6 | * | * | * | * | 2.5 | 11.1 | 44.2 | 109.7 | 199.0 | 288.3 | 382.7 |
| 1976 | 33.7 | * | * | * | * | 2.2 | 10.6 | 49.1 | 107.6 | 203.9 | 281.9 | 412.2 |
| 1975 | 32.8 | * | * | * | * | 2.2 | 11.6 | 44.0 | 109.0 | 200.3 | 300.8 | 334.4 |
| 1974 | 33.2 | * | * | * | * | 3.1 | 12.6 | 45.7 | 116.7 | 214.6 | 268.7 | 345.9 |
| 1973 | 33.5 | * | * | * | * | 2.7 | 12.3 | 49.6 | 118.8 | 211.0 | 296.5 | 315.2 |
| 1972 | 32.0 | * | * | * | * | 1.4 | 11.6 | 49.7 | 110.3 | 212.2 | 276.4 | 331.6 |
| 1971 | 32.6 | * | * | * | * | 3.1 | 14.0 | 44.6 | 108.1 | 218.2 | 291.1 | 387.7 |
| 1970 | 31.6 | * | * | * | * | 2.5 | 14.4 | 46.0 | 113.2 | 205.3 | 293.3 | 297.5 |
| 1969 | 31.4 | * | * | * | * | 2.9 | 14.0 | 48.2 | 113.3 | 203.4 | 283.4 | 307.7 |
| 1968 | 31.9 | * | * | * | * | 2.7 | 13.8 | 48.7 | 118.0 | 208.9 | 283.7 | 329.5 |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|--|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| MALIGNANT NEOPLASMS OF DIGESTIVE ORGANS AND PERITONEUM (150-159) | | | | | | | | | | | | |
| BLACK, BOTH SEXES | | | | | | | | | | | | |
| 1978 | 44.0 | * | * | * | 0.6 | 2.6 | 14.8 | 66.4 | 157.2 | 273.2 | 365.3 | 486.0 |
| 1977 | 44.4 | * | * | * | 0.7 | 3.2 | 16.3 | 65.9 | 163.1 | 281.0 | 358.5 | 457.3 |
| 1976 | 44.1 | * | * | * | 0.7 | 3.3 | 14.5 | 67.2 | 162.3 | 283.7 | 359.5 | 472.1 |
| 1975 | 43.2 | * | * | * | 0.6 | 2.9 | 15.8 | 66.7 | 161.7 | 281.4 | 371.5 | 383.8 |
| 1974 | 42.8 | * | * | * | 0.7 | 3.6 | 15.6 | 66.3 | 164.6 | 289.8 | 345.9 | 410.7 |
| 1973 | 42.4 | * | * | * | 0.7 | 3.0 | 15.9 | 67.6 | 164.5 | 288.7 | 348.5 | 386.0 |
| 1972 | 40.9 | * | * | * | 0.5 | 3.3 | 16.8 | 68.3 | 156.4 | 281.9 | 329.8 | 374.5 |
| 1971 | 41.6 | * | * | * | 0.9 | 3.7 | 17.4 | 62.9 | 160.4 | 284.4 | 361.3 | 429.0 |
| 1970 | 41.9 | * | * | * | 0.7 | 3.4 | 17.2 | 67.9 | 161.1 | 276.2 | 394.4 | 368.9 |
| 1969 | 41.4 | * | * | * | 0.6 | 4.0 | 17.3 | 67.3 | 165.3 | 280.3 | 366.2 | 359.8 |
| 1968 | 41.9 | * | * | * | 0.7 | 3.4 | 18.4 | 67.1 | 169.9 | 286.5 | 358.7 | 369.6 |
| BLACK, MALE | | | | | | | | | | | | |
| 1978 | 51.9 | * | * | * | 0.8 | 3.1 | 18.1 | 92.0 | 211.2 | 355.4 | 461.5 | 596.1 |
| 1977 | 53.7 | * | * | * | 0.9 | 3.7 | 22.0 | 88.7 | 221.8 | 380.6 | 464.7 | 585.7 |
| 1976 | 53.0 | * | * | * | 1.0 | 4.3 | 18.7 | 86.1 | 222.0 | 379.3 | 475.2 | 555.3 |
| 1975 | 52.6 | * | * | * | * | 4.2 | 19.8 | 90.8 | 220.9 | 380.1 | 474.8 | 484.8 |
| 1974 | 51.3 | * | * | * | 0.9 | 4.0 | 18.2 | 87.7 | 218.3 | 377.4 | 466.0 | 527.9 |
| 1973 | 50.0 | * | * | * | 0.9 | 3.1 | 19.5 | 85.4 | 214.0 | 378.3 | 431.9 | 507.3 |
| 1972 | 48.8 | * | * | * | * | 5.7 | 22.1 | 86.6 | 207.1 | 361.2 | 415.2 | 455.0 |
| 1971 | 49.9 | * | * | * | 1.0 | 4.5 | 21.2 | 81.8 | 217.5 | 362.1 | 458.1 | 485.0 |
| 1970 | 51.1 | * | * | * | 1.0 | 4.4 | 19.9 | 90.7 | 212.4 | 355.8 | 517.1 | 476.1 |
| 1969 | 51.2 | * | * | * | * | 5.1 | 20.5 | 88.2 | 222.1 | 373.6 | 474.9 | 448.6 |
| 1968 | 51.7 | * | * | * | * | 4.5 | 23.6 | 87.3 | 226.0 | 378.6 | 462.4 | 445.7 |
| BLACK, FEMALE | | | | | | | | | | | | |
| 1978 | 36.9 | * | * | * | * | 2.2 | 12.0 | 45.4 | 113.2 | 212.0 | 301.1 | 429.3 |
| 1977 | 36.0 | * | * | * | * | 2.7 | 11.6 | 47.0 | 115.2 | 206.2 | 289.0 | 390.4 |
| 1976 | 36.1 | * | * | * | * | 2.3 | 11.0 | 51.5 | 113.0 | 211.1 | 281.9 | 428.1 |
| 1975 | 34.7 | * | * | * | * | 1.9 | 12.4 | 46.6 | 112.7 | 206.5 | 301.5 | 328.6 |
| 1974 | 35.2 | * | * | * | * | 3.2 | 13.4 | 48.4 | 120.2 | 223.0 | 264.7 | 346.2 |
| 1973 | 35.5 | * | * | * | * | 2.9 | 12.9 | 52.4 | 123.4 | 219.5 | 292.2 | 317.8 |
| 1972 | 33.7 | * | * | * | * | 1.3 | 12.3 | 52.7 | 113.8 | 220.6 | 268.6 | 328.6 |
| 1971 | 34.1 | * | * | * | * | 3.1 | 14.2 | 46.7 | 112.0 | 222.8 | 291.9 | 395.5 |
| 1970 | 33.5 | * | * | * | * | 2.5 | 14.9 | 48.3 | 117.5 | 213.1 | 304.5 | 299.8 |
| 1969 | 32.5 | * | * | * | * | 3.1 | 14.6 | 49.2 | 116.9 | 206.1 | 284.6 | 305.0 |
| 1968 | 32.9 | * | * | * | * | 2.5 | 14.0 | 49.6 | 121.5 | 211.9 | 281.5 | 328.6 |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| MALIGNANT NEOPLASMS OF RESPIRATORY SYSTEM (160-163) | | | | | | | | | | | | |
| BOTH SEXES | | | | | | | | | | | | |
| 1978 | 45.0 | * | * | * | 0.1 | 0.9 | 10.1 | 56.6 | 139.4 | 226.5 | 230.2 | 167.0 |
| 1977 | 43.3 | * | * | * | 0.1 | 0.9 | 10.6 | 55.1 | 134.7 | 218.4 | 223.2 | 163.2 |
| 1976 | 41.9 | * | * | * | 0.1 | 0.8 | 10.7 | 53.5 | 133.4 | 211.8 | 214.5 | 158.1 |
| 1975 | 40.2 | * | * | * | 0.1 | 0.9 | 11.0 | 52.3 | 130.1 | 204.7 | 203.1 | 147.2 |
| 1974 | 39.1 | * | * | * | 0.1 | 0.9 | 11.0 | 51.7 | 129.2 | 198.0 | 199.4 | 144.2 |
| 1973 | 37.5 | * | * | 0.1 | 0.1 | 1.0 | 11.2 | 49.6 | 124.6 | 192.5 | 190.2 | 138.6 |
| 1972 | 36.8 | * | * | * | 0.1 | 0.9 | 11.6 | 49.2 | 122.1 | 190.2 | 188.3 | 129.3 |
| 1971 | 35.2 | * | * | 0.1 | 0.2 | 1.1 | 11.2 | 47.0 | 115.9 | 183.4 | 182.5 | 134.0 |
| 1970 | 34.2 | * | * | * | 0.2 | 1.0 | 11.6 | 46.2 | 116.2 | 174.6 | 175.1 | 113.5 |
| 1969 | 32.8 | * | * | 0.1 | 0.2 | 1.1 | 11.5 | 44.2 | 111.7 | 170.6 | 161.2 | 117.3 |
| 1968 | 31.8 | * | * | 0.1 | 0.2 | 1.1 | 11.0 | 43.3 | 107.8 | 166.3 | 159.5 | 116.8 |
| MALE | | | | | | | | | | | | |
| 1978 | 69.3 | * | * | * | 0.1 | 1.1 | 13.1 | 81.8 | 221.0 | 407.8 | 474.4 | 361.5 |
| 1977 | 67.5 | * | * | * | 0.2 | 1.2 | 13.7 | 80.7 | 215.3 | 400.0 | 467.6 | 345.3 |
| 1976 | 65.8 | * | * | * | 0.2 | 1.0 | 14.5 | 79.1 | 215.4 | 390.5 | 446.7 | 330.2 |
| 1975 | 63.9 | * | * | * | 0.2 | 1.3 | 14.9 | 78.5 | 212.5 | 384.0 | 425.7 | 299.2 |
| 1974 | 62.9 | * | * | * | 0.1 | 1.2 | 15.6 | 78.9 | 214.2 | 374.1 | 416.7 | 293.1 |
| 1973 | 60.9 | * | * | * | 0.2 | 1.2 | 15.8 | 75.9 | 209.8 | 368.4 | 394.8 | 282.9 |
| 1972 | 60.0 | * | * | * | 0.2 | 1.1 | 16.3 | 74.6 | 207.8 | 364.3 | 391.5 | 256.7 |
| 1971 | 57.9 | * | * | * | 0.2 | 1.5 | 16.1 | 71.8 | 199.2 | 353.8 | 374.9 | 262.9 |
| 1970 | 57.0 | * | * | * | 0.2 | 1.5 | 17.0 | 72.1 | 202.3 | 340.7 | 354.2 | 215.3 |
| 1969 | 55.0 | * | * | * | 0.2 | 1.4 | 17.1 | 70.1 | 195.9 | 331.4 | 325.1 | 229.6 |
| 1968 | 53.7 | * | * | * | 0.2 | 1.6 | 16.1 | 69.6 | 192.1 | 324.4 | 313.7 | 217.0 |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| MALIGNANT NEOPLASMS OF RESPIRATORY SYSTEM (160-163) | | | | | | | | | | | | |
| FEMALE | | | | | | | | | | | | |
| 1978 | 22.0 | * | * | * | * | 0.7 | 7.1 | 33.2 | 67.4 | 88.0 | 84.7 | 79.1 |
| 1977 | 20.4 | * | * | * | 0.1 | 0.7 | 7.6 | 31.3 | 63.4 | 79.4 | 77.0 | 79.5 |
| 1976 | 19.3 | * | * | * | * | 0.7 | 7.0 | 29.7 | 60.8 | 75.0 | 74.3 | 77.3 |
| 1975 | 17.7 | * | * | * | * | 0.6 | 7.3 | 28.0 | 57.2 | 67.3 | 67.4 | 73.7 |
| 1974 | 16.6 | * | * | * | * | 0.7 | 6.7 | 26.3 | 53.8 | 62.9 | 65.5 | 70.5 |
| 1973 | 15.3 | * | * | * | 0.1 | 0.8 | 6.8 | 25.1 | 48.8 | 57.5 | 62.2 | 65.9 |
| 1972 | 14.8 | * | * | * | * | 0.6 | 7.2 | 25.6 | 45.8 | 55.9 | 58.9 | 64.1 |
| 1971 | 13.7 | * | * | * | 0.1 | 0.7 | 6.6 | 23.9 | 41.4 | 51.5 | 57.9 | 66.7 |
| 1970 | 12.6 | * | * | * | * | 0.6 | 6.5 | 22.2 | 38.9 | 45.6 | 56.5 | 56.5 |
| 1969 | 11.8 | * | * | * | 0.2 | 0.8 | 6.1 | 20.0 | 35.9 | 44.9 | 51.2 | 56.5 |
| 1968 | 11.1 | * | * | * | * | 0.6 | 6.1 | 18.8 | 31.6 | 41.9 | 53.9 | 61.8 |
| WHITE, BOTH SEXES | | | | | | | | | | | | |
| 1978 | 46.2 | * | * | * | 0.1 | 0.8 | 9.4 | 53.3 | 135.9 | 226.7 | 233.0 | 169.0 |
| 1977 | 44.4 | * | * | * | 0.1 | 0.9 | 9.7 | 51.8 | 131.0 | 217.9 | 227.2 | 163.8 |
| 1976 | 43.1 | * | * | * | 0.2 | 0.8 | 10.0 | 50.3 | 130.5 | 212.1 | 218.5 | 159.3 |
| 1975 | 41.3 | * | * | * | 0.1 | 0.9 | 10.2 | 49.7 | 127.1 | 204.7 | 206.7 | 148.1 |
| 1974 | 40.1 | * | * | * | 0.1 | 0.9 | 10.2 | 48.6 | 126.2 | 197.7 | 202.7 | 145.9 |
| 1973 | 38.4 | * | * | * | 0.1 | 0.9 | 10.5 | 46.3 | 121.9 | 193.3 | 192.8 | 139.6 |
| 1972 | 37.8 | * | * | * | 0.1 | 0.8 | 10.8 | 46.9 | 120.0 | 190.9 | 191.0 | 130.7 |
| 1971 | 36.2 | * | * | * | 0.2 | 1.1 | 10.3 | 44.6 | 114.1 | 184.8 | 185.0 | 137.1 |
| 1970 | 35.1 | * | * | * | 0.1 | 1.0 | 10.7 | 44.1 | 115.0 | 175.9 | 177.0 | 115.6 |
| 1969 | 33.6 | * | * | * | 0.2 | 1.0 | 10.8 | 41.8 | 110.1 | 171.6 | 163.3 | 118.7 |
| 1968 | 32.7 | * | * | 0.1 | 0.2 | 1.0 | 10.2 | 41.2 | 106.6 | 168.3 | 161.3 | 117.4 |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| MALIGNANT NEOPLASMS OF RESPIRATORY SYSTEM (160-163) | | | | | | | | | | | | |
| WHITE, MALE | | | | | | | | | | | | |
| 1978 | 70.5 | * | * | * | 0.1 | 1.0 | 11.7 | 75.1 | 212.2 | 406.1 | 483.7 | 369.0 |
| 1977 | 68.6 | * | * | * | 0.1 | 1.1 | 12.3 | 74.2 | 206.6 | 398.2 | 478.0 | 347.3 |
| 1976 | 67.0 | * | * | * | 0.2 | 1.0 | 13.4 | 72.8 | 207.7 | 390.4 | 458.3 | 335.6 |
| 1975 | 65.1 | * | * | * | 0.2 | 1.2 | 13.3 | 73.0 | 204.8 | 384.1 | 436.3 | 303.1 |
| 1974 | 63.9 | * | * | * | 0.2 | 1.1 | 14.1 | 73.0 | 207.3 | 373.3 | 426.0 | 297.1 |
| 1973 | 61.9 | * | * | * | 0.1 | 1.1 | 14.6 | 69.6 | 203.4 | 370.6 | 402.2 | 286.9 |
| 1972 | 61.2 | * | * | * | 0.2 | 1.0 | 14.9 | 69.9 | 202.2 | 366.2 | 399.8 | 261.5 |
| 1971 | 59.3 | * | * | * | 0.2 | 1.4 | 14.7 | 67.1 | 195.0 | 358.2 | 382.4 | 270.8 |
| 1970 | 58.3 | * | * | * | 0.2 | 1.4 | 15.4 | 67.6 | 199.3 | 344.8 | 360.7 | 221.8 |
| 1969 | 56.1 | * | * | * | 0.2 | 1.2 | 15.8 | 65.2 | 192.6 | 334.5 | 331.7 | 233.0 |
| 1968 | 55.0 | * | * | * | 0.3 | 1.5 | 14.7 | 65.2 | 189.8 | 329.4 | 319.0 | 219.8 |
| WHITE, FEMALE | | | | | | | | | | | | |
| 1978 | 23.2 | * | * | * | * | 0.6 | 7.1 | 32.6 | 68.1 | 89.9 | 85.7 | 79.9 |
| 1977 | 21.3 | * | * | * | * | 0.6 | 7.1 | 30.7 | 63.8 | 80.4 | 79.0 | 80.7 |
| 1976 | 20.2 | * | * | * | * | 0.7 | 6.8 | 29.1 | 61.8 | 75.9 | 75.6 | 77.9 |
| 1975 | 18.6 | * | * | * | * | 0.5 | 7.1 | 27.7 | 57.9 | 67.8 | 68.4 | 74.3 |
| 1974 | 17.3 | * | * | * | * | 0.7 | 6.5 | 25.6 | 54.0 | 63.8 | 66.5 | 72.2 |
| 1973 | 15.9 | * | * | * | * | 0.8 | 6.5 | 24.4 | 49.1 | 57.7 | 63.2 | 66.7 |
| 1972 | 15.4 | * | * | * | * | 0.6 | 7.0 | 25.3 | 46.5 | 56.4 | 59.4 | 64.8 |
| 1971 | 14.2 | * | * | * | 0.1 | 0.7 | 6.1 | 23.4 | 41.6 | 51.2 | 58.7 | 68.6 |
| 1970 | 13.1 | * | * | * | * | 0.6 | 6.0 | 22.1 | 39.3 | 45.4 | 56.8 | 57.4 |
| 1969 | 12.2 | * | * | * | 0.2 | 0.7 | 6.0 | 19.8 | 35.8 | 44.8 | 51.3 | 58.2 |
| 1968 | 11.4 | * | * | * | * | 0.6 | 5.9 | 18.6 | 31.4 | 42.1 | 54.4 | 62.2 |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| MALIGNANT NEOPLASMS OF RESPIRATORY SYSTEM (160-163) | | | | | | | | | | | | |
| ALL OTHER, BOTH SEXES | | | | | | | | | | | | |
| 1978 | 37.0 | * | * | * | * | 1.1 | 14.9 | 82.3 | 170.9 | 224.6 | 200.6 | 143.5 |
| 1977 | 36.4 | * | * | * | * | 1.4 | 16.6 | 81.4 | 168.5 | 222.6 | 182.9 | 155.6 |
| 1976 | 34.3 | * | * | * | * | 1.1 | 15.1 | 79.2 | 160.6 | 209.5 | 173.5 | 143.8 |
| 1975 | 33.2 | * | * | * | * | 1.1 | 16.7 | 74.2 | 158.1 | 205.0 | 164.4 | 137.7 |
| 1974 | 32.9 | * | * | * | * | 1.1 | 16.6 | 77.5 | 156.9 | 200.2 | 162.7 | 124.4 |
| 1973 | 31.6 | * | * | * | * | 1.2 | 16.1 | 77.8 | 150.1 | 185.7 | 159.7 | 126.8 |
| 1972 | 30.1 | * | * | * | * | 1.4 | 17.4 | 68.9 | 142.4 | 183.0 | 156.5 | 113.8 |
| 1971 | 28.6 | * | * | * | * | 1.3 | 17.7 | 67.9 | 133.0 | 169.8 | 152.2 | 99.2 |
| 1970 | 27.7 | * | * | * | * | 1.4 | 18.4 | 65.3 | 127.3 | 162.3 | 150.6 | 92.5 |
| 1969 | 27.2 | * | * | * | * | 1.8 | 16.7 | 65.8 | 126.5 | 160.6 | 135.0 | 100.9 |
| 1968 | 25.7 | * | * | * | * | 1.4 | 17.0 | 62.4 | 119.2 | 146.8 | 135.9 | 109.8 |
| ALL OTHER, MALE | | | | | | | | | | | | |
| 1978 | 61.4 | * | * | * | * | 1.4 | 23.6 | 136.8 | 302.9 | 424.0 | 385.3 | 284.5 |
| 1977 | 60.6 | * | * | * | * | 2.0 | 24.0 | 135.8 | 298.2 | 416.4 | 369.3 | 325.0 |
| 1976 | 57.4 | * | * | * | * | 1.4 | 23.3 | 132.9 | 289.5 | 390.7 | 338.9 | 276.4 |
| 1975 | 56.1 | * | * | * | * | 1.5 | 26.7 | 126.0 | 285.5 | 383.5 | 321.1 | 259.3 |
| 1974 | 55.8 | * | * | * | * | 1.5 | 26.3 | 131.1 | 279.9 | 382.1 | 321.4 | 252.0 |
| 1973 | 53.7 | * | * | * | * | 1.9 | 24.9 | 132.4 | 272.2 | 346.8 | 315.6 | 241.7 |
| 1972 | 51.6 | * | * | * | * | 2.4 | 27.7 | 116.3 | 261.7 | 346.5 | 302.3 | 208.5 |
| 1971 | 48.5 | * | * | * | * | 1.9 | 27.3 | 113.9 | 240.7 | 311.8 | 293.7 | 185.1 |
| 1970 | 47.6 | * | * | * | * | 2.4 | 29.3 | 113.1 | 231.5 | 301.2 | 278.7 | 158.8 |
| 1969 | 47.0 | * | * | * | * | 2.3 | 28.1 | 115.9 | 227.7 | 301.0 | 246.3 | 197.7 |
| 1968 | 44.7 | * | * | * | * | 2.3 | 27.9 | 110.5 | 215.1 | 275.6 | 250.0 | 190.2 |
| ALL OTHER, FEMALE | | | | | | | | | | | | |
| 1978 | 14.8 | * | * | * | * | 0.8 | 7.7 | 37.2 | 61.3 | 70.2 | 74.8 | 69.7 |
| 1977 | 14.4 | * | * | * | * | 0.9 | 10.5 | 36.2 | 60.3 | 70.4 | 55.4 | 64.4 |
| 1976 | 13.2 | * | * | * | * | * | 8.2 | 34.2 | 52.5 | 66.2 | 60.4 | 69.4 |
| 1975 | 12.4 | * | * | * | * | * | 8.4 | 30.6 | 50.5 | 63.1 | 56.8 | 65.6 |
| 1974 | 11.9 | * | * | * | * | * | 8.6 | 32.0 | 52.2 | 54.4 | 53.3 | 49.4 |
| 1973 | 11.4 | * | * | * | * | * | 8.8 | 31.0 | 45.7 | 55.7 | 50.3 | 57.0 |
| 1972 | 10.4 | * | * | * | * | * | 8.7 | 28.1 | 39.6 | 50.5 | 51.8 | 55.3 |
| 1971 | 10.5 | * | * | * | * | * | 9.8 | 28.0 | 39.1 | 53.6 | 47.5 | 43.8 |
| 1970 | 9.5 | * | * | * | * | * | 9.4 | 23.3 | 35.3 | 47.7 | 53.2 | 45.8 |
| 1969 | 9.0 | * | * | * | * | 1.4 | 7.2 | 21.8 | 36.9 | 45.1 | 49.8 | 35.4 |
| 1968 | 8.3 | * | * | * | * | * | 7.8 | 19.8 | 33.1 | 39.9 | 47.7 | 55.7 |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| MALIGNANT NEOPLASMS OF RESPIRATORY SYSTEM (160-163) | | | | | | | | | | | | |
| BLACK, BOTH SEXES | | | | | | | | | | | | |
| 1978 | 40.6 | * | * | * | * | 1.2 | 16.9 | 92.5 | 187.1 | 236.3 | 197.8 | 137.3 |
| 1977 | 40.0 | * | * | * | * | 1.6 | 19.0 | 92.0 | 183.0 | 235.7 | 183.1 | 144.8 |
| 1976 | 37.3 | * | * | * | * | 1.1 | 17.1 | 88.0 | 173.5 | 220.2 | 171.5 | 133.1 |
| 1975 | 36.1 | * | * | * | * | 1.1 | 19.2 | 82.5 | 169.8 | 216.0 | 161.5 | 130.8 |
| 1974 | 35.4 | * | * | * | * | 1.2 | 18.6 | 85.1 | 167.2 | 209.2 | 162.9 | 115.7 |
| 1973 | 33.9 | * | * | * | * | 1.2 | 18.1 | 85.1 | 159.3 | 193.9 | 157.7 | 118.4 |
| 1972 | 32.0 | * | * | * | * | 1.5 | 19.5 | 75.0 | 151.0 | 189.7 | 149.6 | 110.9 |
| 1971 | 30.4 | * | * | * | * | 1.4 | 19.5 | 73.6 | 140.9 | 174.0 | 152.6 | 91.6 |
| 1970 | 29.7 | * | * | * | * | 1.6 | 20.5 | 70.8 | 134.8 | 169.9 | 153.3 | 89.5 |
| 1969 | 28.7 | * | * | * | * | 1.9 | 18.2 | 70.6 | 133.1 | 166.0 | 133.6 | 93.8 |
| 1968 | 26.8 | * | * | * | * | 1.4 | 18.5 | 66.3 | 124.5 | 149.5 | 130.2 | 98.9 |
| BLACK, MALE | | | | | | | | | | | | |
| 1978 | 67.9 | * | * | * | * | 1.6 | 26.8 | 155.0 | 336.1 | 457.9 | 388.3 | 274.5 |
| 1977 | 67.1 | * | * | * | * | 2.2 | 27.6 | 154.0 | 329.1 | 453.0 | 381.7 | 312.2 |
| 1976 | 62.8 | * | * | * | * | 1.4 | 26.7 | 147.9 | 315.9 | 422.2 | 338.0 | 270.2 |
| 1975 | 61.3 | * | * | * | * | 1.5 | 30.7 | 141.1 | 310.5 | 415.9 | 318.9 | 254.3 |
| 1974 | 60.6 | * | * | * | * | 1.8 | 29.8 | 145.1 | 304.1 | 410.2 | 326.8 | 246.5 |
| 1973 | 58.0 | * | * | * | * | 2.1 | 28.0 | 145.7 | 294.1 | 372.1 | 319.1 | 241.5 |
| 1972 | 55.3 | * | * | * | * | 2.5 | 31.1 | 127.4 | 282.3 | 372.7 | 293.4 | 205.0 |
| 1971 | 51.7 | * | * | * | * | 2.0 | 30.5 | 124.5 | 259.2 | 326.8 | 298.9 | 167.5 |
| 1970 | 51.2 | * | * | * | * | 2.9 | 32.6 | 123.5 | 250.3 | 322.2 | 290.6 | 154.4 |
| 1969 | 49.9 | * | * | * | * | 2.6 | 31.1 | 125.8 | 243.8 | 317.6 | 250.3 | 189.2 |
| 1968 | 46.8 | * | * | * | * | 2.4 | 30.3 | 119.1 | 228.8 | 285.0 | 245.5 | 182.9 |
| BLACK, FEMALE | | | | | | | | | | | | |
| 1978 | 16.0 | * | * | * | * | * | 8.7 | 41.1 | 65.5 | 71.6 | 70.7 | 66.7 |
| 1977 | 15.5 | * | * | * | * | 1.1 | 11.9 | 40.6 | 63.5 | 72.5 | 51.7 | 57.4 |
| 1976 | 14.2 | * | * | * | * | * | 9.1 | 38.1 | 56.0 | 67.3 | 59.9 | 60.7 |
| 1975 | 13.3 | * | * | * | * | * | 9.6 | 33.4 | 53.4 | 64.4 | 54.9 | 63.1 |
| 1974 | 12.6 | * | * | * | * | * | 9.4 | 34.7 | 53.6 | 55.3 | 52.1 | 43.6 |
| 1973 | 12.0 | * | * | * | * | * | 9.8 | 33.6 | 47.2 | 56.2 | 47.0 | 49.3 |
| 1972 | 10.8 | * | * | * | * | * | 9.9 | 30.4 | 41.1 | 47.5 | 47.9 | 57.1 |
| 1971 | 11.0 | * | * | * | * | * | 10.4 | 29.9 | 41.0 | 53.5 | 47.5 | 46.3 |
| 1970 | 10.1 | * | * | * | * | * | 10.5 | 25.3 | 36.4 | 49.3 | 52.6 | 47.6 |
| 1969 | 9.4 | * | * | * | * | * | 7.5 | 22.8 | 38.4 | 45.5 | 47.1 | 35.0 |
| 1968 | 8.6 | * | * | * | * | * | 8.6 | 20.4 | 34.3 | 40.1 | 44.6 | 48.2 |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| MALIGNANT NEOPLASMS OF BREAST (174) | | | | | | | | | | | | |
| BOTH SEXES | | | | | | | | | | | | |
| 1978 | 15.6 | * | * | * | 0.1 | 1.6 | 8.7 | 26.4 | 43.1 | 56.3 | 78.3 | 115.3 |
| 1977 | 15.8 | * | * | * | * | 1.8 | 8.9 | 27.0 | 43.3 | 58.1 | 80.1 | 122.1 |
| 1976 | 15.4 | * | * | * | * | 1.6 | 8.9 | 26.5 | 42.9 | 56.5 | 77.7 | 116.8 |
| 1975 | 15.1 | * | * | * | 0.1 | 1.7 | 9.5 | 26.1 | 41.6 | 55.3 | 78.2 | 112.5 |
| 1974 | 15.2 | * | * | * | 0.1 | 1.8 | 9.5 | 26.9 | 42.7 | 55.5 | 78.4 | 117.1 |
| 1973 | 15.2 | * | * | * | 0.1 | 1.7 | 9.9 | 27.1 | 42.7 | 57.6 | 77.3 | 115.4 |
| 1972 | 15.0 | * | * | * | * | 1.9 | 9.9 | 27.5 | 42.6 | 56.0 | 75.9 | 114.5 |
| 1971 | 14.6 | * | * | * | 0.1 | 1.8 | 10.1 | 26.7 | 41.8 | 52.9 | 76.2 | 113.9 |
| 1970 | 14.7 | * | * | * | * | 2.0 | 10.5 | 27.4 | 41.2 | 53.4 | 77.7 | 101.8 |
| 1969 | 14.4 | * | * | * | 0.1 | 1.7 | 10.5 | 27.0 | 41.3 | 52.8 | 72.8 | 111.6 |
| 1968 | 14.6 | * | * | * | * | 2.0 | 10.8 | 27.7 | 40.8 | 53.4 | 74.6 | 117.1 |
| MALE | | | | | | | | | | | | |
| 1978 | 0.3 | * | * | * | * | * | * | 0.3 | 0.6 | 1.3 | 2.2 | 4.6 |
| 1977 | 0.3 | * | * | * | * | * | * | 0.3 | 0.8 | 1.2 | 2.3 | 3.3 |
| 1976 | 0.3 | * | * | * | * | * | * | 0.2 | 0.6 | 1.5 | 2.5 | 4.0 |
| 1975 | 0.3 | * | * | * | * | * | * | 0.2 | 0.8 | 1.3 | 2.2 | 4.4 |
| 1974 | 0.3 | * | * | * | * | * | * | 0.3 | 0.8 | 1.3 | 2.5 | 5.1 |
| 1973 | 0.3 | * | * | * | * | * | * | 0.3 | 0.8 | 1.4 | 2.5 | * |
| 1972 | 0.3 | * | * | * | * | * | * | 0.3 | 0.9 | 1.7 | 2.4 | * |
| 1971 | 0.3 | * | * | * | * | * | * | 0.4 | 0.8 | 1.6 | 2.8 | * |
| 1970 | 0.3 | * | * | * | * | * | * | 0.4 | 0.6 | 1.4 | 2.7 | * |
| 1969 | 0.3 | * | * | * | * | * | * | 0.3 | 0.6 | 1.4 | 2.3 | * |
| 1968 | 0.3 | * | * | * | * | * | * | 0.3 | 0.7 | 1.4 | 2.3 | * |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| MALIGNANT NEOPLASMS OF BREAST (174) | | | | | | | | | | | | |
| FEMALE | | | | | | | | | | | | |
| 1978 | 30.1 | * | * | * | 0.1 | 3.2 | 17.0 | 50.8 | 80.6 | 98.4 | 123.6 | 165.3 |
| 1977 | 30.5 | * | * | * | * | 3.6 | 17.3 | 51.9 | 80.8 | 101.6 | 126.7 | 176.7 |
| 1976 | 29.6 | * | * | * | * | 3.2 | 17.3 | 50.9 | 80.4 | 98.5 | 123.2 | 169.8 |
| 1975 | 29.1 | * | * | * | 0.1 | 3.3 | 18.6 | 50.2 | 77.7 | 96.6 | 124.5 | 164.8 |
| 1974 | 29.4 | * | * | * | 0.1 | 3.6 | 18.6 | 51.7 | 79.9 | 97.0 | 125.2 | 172.6 |
| 1973 | 29.4 | * | * | * | 0.2 | 3.4 | 19.2 | 51.9 | 79.9 | 100.8 | 124.1 | 171.9 |
| 1972 | 29.0 | * | * | * | * | 3.8 | 19.2 | 52.8 | 79.7 | 97.9 | 122.6 | 171.8 |
| 1971 | 28.2 | * | * | * | 0.2 | 3.5 | 19.5 | 51.0 | 78.5 | 92.7 | 123.6 | 171.8 |
| 1970 | 28.4 | * | * | * | * | 3.9 | 20.4 | 52.6 | 77.6 | 93.8 | 127.4 | 157.1 |
| 1969 | 27.9 | * | * | * | 0.1 | 3.4 | 20.3 | 51.9 | 78.0 | 93.0 | 120.1 | 170.2 |
| 1968 | 28.2 | * | * | * | * | 3.8 | 20.9 | 53.3 | 77.0 | 94.3 | 124.1 | 179.4 |
| WHITE, BOTH SEXES | | | | | | | | | | | | |
| 1978 | 16.3 | * | * | * | 0.1 | 1.5 | 8.2 | 26.3 | 43.4 | 57.3 | 79.9 | 117.1 |
| 1977 | 16.5 | * | * | * | * | 1.7 | 8.7 | 26.6 | 43.4 | 58.9 | 82.2 | 124.8 |
| 1976 | 16.1 | * | * | * | * | 1.6 | 8.6 | 26.6 | 43.4 | 57.6 | 79.7 | 118.7 |
| 1975 | 15.8 | * | * | * | * | 1.6 | 9.3 | 26.1 | 41.7 | 56.2 | 80.7 | 116.3 |
| 1974 | 15.8 | * | * | * | * | 1.7 | 9.1 | 26.5 | 43.0 | 56.1 | 81.0 | 119.7 |
| 1973 | 15.9 | * | * | * | 0.1 | 1.6 | 9.5 | 26.8 | 43.1 | 58.2 | 80.0 | 118.3 |
| 1972 | 15.7 | * | * | * | * | 1.8 | 9.4 | 27.4 | 43.2 | 56.7 | 77.9 | 118.7 |
| 1971 | 15.3 | * | * | * | 0.1 | 1.7 | 9.9 | 26.4 | 42.2 | 53.8 | 78.3 | 117.7 |
| 1970 | 15.4 | * | * | * | * | 1.9 | 10.3 | 27.5 | 42.0 | 54.7 | 79.4 | 105.8 |
| 1969 | 15.0 | * | * | * | * | 1.6 | 10.1 | 26.9 | 41.8 | 53.7 | 74.5 | 116.8 |
| 1968 | 15.2 | * | * | * | * | 1.8 | 10.5 | 27.4 | 41.2 | 54.2 | 76.3 | 119.9 |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| MALIGNANT NEOPLASMS OF BREAST (174) | | | | | | | | | | | | |
| WHITE, MALE | | | | | | | | | | | | |
| 1978 | 0.3 | * | * | * | * | * | * | 0.3 | 0.6 | 1.2 | 2.2 | 4.2 |
| 1977 | 0.3 | * | * | * | * | * | * | 0.3 | 0.7 | 1.3 | 2.0 | 3.5 |
| 1976 | 0.3 | * | * | * | * | * | * | 0.2 | 0.5 | 1.5 | 2.3 | 4.0 |
| 1975 | 0.3 | * | * | * | * | * | * | 0.2 | 0.7 | 1.4 | 2.2 | 4.6 |
| 1974 | 0.3 | * | * | * | * | * | * | 0.2 | 0.8 | 1.3 | 2.4 | 4.7 |
| 1973 | 0.3 | * | * | * | * | * | * | 0.3 | 0.7 | 1.4 | 2.6 | * |
| 1972 | 0.3 | * | * | * | * | * | * | 0.2 | 0.7 | 1.6 | 2.5 | * |
| 1971 | 0.3 | * | * | * | * | * | * | 0.4 | 0.9 | 1.5 | 2.9 | * |
| 1970 | 0.3 | * | * | * | * | * | * | 0.4 | 0.6 | 1.4 | 2.7 | * |
| 1969 | 0.3 | * | * | * | * | * | * | 0.3 | 0.6 | 1.4 | 2.3 | * |
| 1968 | 0.3 | * | * | * | * | * | * | 0.3 | 0.6 | 1.3 | 2.2 | * |
| WHITE, FEMALE | | | | | | | | | | | | |
| 1978 | 31.6 | * | * | * | * | 3.1 | 16.2 | 50.8 | 81.5 | 100.1 | 125.6 | 167.4 |
| 1977 | 32.0 | * | * | * | * | 3.4 | 17.0 | 51.5 | 81.4 | 102.8 | 129.5 | 179.8 |
| 1976 | 31.3 | * | * | * | * | 3.1 | 16.9 | 51.4 | 81.6 | 100.5 | 125.9 | 171.8 |
| 1975 | 30.6 | * | * | * | * | 3.3 | 18.2 | 50.5 | 78.2 | 98.1 | 128.1 | 169.6 |
| 1974 | 30.7 | * | * | * | * | 3.4 | 17.9 | 51.2 | 80.7 | 97.9 | 129.0 | 175.8 |
| 1973 | 30.8 | * | * | * | 0.2 | 3.2 | 18.6 | 51.7 | 81.0 | 101.5 | 128.0 | 175.7 |
| 1972 | 30.4 | * | * | * | * | 3.6 | 18.5 | 52.8 | 81.2 | 99.0 | 125.4 | 176.9 |
| 1971 | 29.6 | * | * | * | 0.2 | 3.3 | 19.3 | 50.7 | 79.3 | 94.2 | 126.6 | 176.4 |
| 1970 | 29.9 | * | * | * | * | 3.7 | 20.2 | 53.0 | 79.3 | 95.9 | 129.6 | 161.9 |
| 1969 | 29.1 | * | * | * | * | 3.1 | 19.7 | 51.8 | 78.9 | 94.3 | 122.6 | 177.0 |
| 1968 | 29.4 | * | * | * | * | 3.6 | 20.5 | 53.0 | 77.9 | 95.7 | 126.4 | 183.1 |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| MALIGNANT NEOPLASMS OF BREAST (174) | | | | | | | | | | | | |
| ALL OTHER, BOTH SEXES | | | | | | | | | | | | |
| 1978 | 11.0 | * | * | * | * | 2.1 | 11.9 | 27.9 | 40.2 | 47.4 | 61.7 | 94.6 |
| 1977 | 11.2 | * | * | * | * | 2.4 | 10.2 | 30.1 | 41.8 | 50.8 | 58.8 | 91.3 |
| 1976 | 10.3 | * | * | * | * | 1.8 | 10.9 | 26.0 | 38.3 | 45.4 | 57.2 | 94.1 |
| 1975 | 10.2 | * | * | * | * | 1.7 | 11.5 | 26.5 | 39.9 | 46.1 | 50.8 | 68.5 |
| 1974 | 10.9 | * | * | * | * | 2.4 | 12.5 | 30.4 | 39.9 | 50.0 | 49.4 | 87.4 |
| 1973 | 10.6 | * | * | * | * | 2.4 | 12.7 | 29.0 | 38.2 | 52.0 | 46.4 | 81.9 |
| 1972 | 10.5 | * | * | * | * | 2.5 | 13.3 | 28.5 | 36.7 | 49.6 | 52.0 | 66.7 |
| 1971 | 10.2 | * | * | * | * | 2.8 | 11.4 | 29.0 | 38.2 | 44.0 | 49.9 | 70.0 |
| 1970 | 9.7 | * | * | * | * | 3.1 | 12.2 | 26.5 | 33.2 | 40.8 | 56.4 | 61.2 |
| 1969 | 10.2 | * | * | * | * | 2.7 | 13.7 | 28.4 | 36.7 | 44.4 | 50.0 | 54.1 |
| 1968 | 10.4 | * | * | * | * | 3.0 | 12.8 | 29.5 | 36.5 | 44.9 | 53.8 | 85.3 |
| ALL OTHER, MALE | | | | | | | | | | | | |
| 1978 | 0.3 | * | * | * | * | * | * | * | * | * | * | * |
| 1977 | 0.2 | * | * | * | * | * | * | * | * | * | * | * |
| 1976 | 0.3 | * | * | * | * | * | * | * | * | * | * | * |
| 1975 | 0.2 | * | * | * | * | * | * | * | * | * | * | * |
| 1974 | 0.3 | * | * | * | * | * | * | * | * | * | * | * |
| 1973 | 0.3 | * | * | * | * | * | * | * | * | * | * | * |
| 1972 | 0.4 | * | * | * | * | * | * | * | 2.4 | * | * | * |
| 1971 | 0.2 | * | * | * | * | * | * | * | * | * | * | * |
| 1970 | 0.3 | * | * | * | * | * | * | * | * | * | * | * |
| 1969 | 0.3 | * | * | * | * | * | * | * | * | * | * | * |
| 1968 | 0.3 | * | * | * | * | * | * | * | * | * | * | * |
| ALL OTHER, FEMALE | | | | | | | | | | | | |
| 1978 | 20.8 | * | * | * | * | 4.0 | 21.9 | 50.7 | 72.8 | 82.1 | 102.6 | 141.3 |
| 1977 | 21.2 | * | * | * | * | 4.5 | 18.6 | 54.9 | 75.9 | 90.2 | 95.6 | 139.4 |
| 1976 | 19.3 | * | * | * | * | 3.3 | 20.1 | 47.3 | 69.5 | 79.8 | 93.6 | 144.9 |
| 1975 | 19.2 | * | * | * | * | 3.3 | 21.1 | 48.5 | 72.7 | 81.9 | 84.2 | 106.5 |
| 1974 | 20.6 | * | * | * | * | 4.4 | 22.8 | 55.9 | 72.9 | 88.7 | 81.0 | 132.9 |
| 1973 | 20.1 | * | * | * | * | 4.4 | 23.3 | 53.3 | 69.7 | 92.9 | 78.0 | 126.6 |
| 1972 | 19.7 | * | * | * | * | 4.7 | 24.2 | 52.4 | 66.3 | 87.5 | 88.4 | 107.9 |
| 1971 | 19.3 | * | * | * | * | 5.2 | 20.9 | 53.6 | 70.9 | 78.2 | 85.0 | 113.7 |
| 1970 | 18.3 | * | * | * | * | 5.7 | 22.2 | 49.2 | 61.9 | 73.0 | 97.0 | 103.0 |
| 1969 | 19.3 | * | * | * | * | 5.0 | 25.0 | 53.0 | 68.7 | 79.4 | 85.8 | 87.7 |
| 1968 | 19.7 | * | * | * | * | 5.5 | 23.4 | 55.4 | 68.1 | 80.5 | 92.9 | 134.4 |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| MALIGNANT NEOPLASMS OF BREAST (174) | | | | | | | | | | | | |
| BLACK, BOTH SEXES | | | | | | | | | | | | |
| 1978 | 12.1 | * | * | * | * | 2.4 | 13.4 | 30.0 | 43.4 | 50.7 | 66.2 | 99.3 |
| 1977 | 12.3 | * | * | * | * | 2.7 | 11.5 | 32.6 | 45.1 | 54.6 | 63.2 | 100.7 |
| 1976 | 11.3 | * | * | * | * | 2.1 | 12.2 | 28.4 | 41.0 | 49.5 | 60.9 | 103.7 |
| 1975 | 11.0 | * | * | * | * | 2.0 | 12.8 | 28.5 | 42.5 | 48.4 | 53.5 | 71.5 |
| 1974 | 11.9 | * | * | * | * | 2.5 | 14.0 | 32.8 | 42.2 | 53.9 | 51.4 | 95.9 |
| 1973 | 11.6 | * | * | * | * | 2.7 | 13.9 | 31.5 | 40.8 | 55.9 | 49.5 | 89.5 |
| 1972 | 11.3 | * | * | * | * | 2.9 | 14.7 | 30.1 | 38.9 | 53.5 | 55.9 | 72.7 |
| 1971 | 11.0 | * | * | * | * | 3.0 | 12.5 | 31.3 | 40.6 | 46.8 | 53.7 | 72.0 |
| 1970 | 10.5 | * | * | * | * | 3.2 | 13.4 | 28.3 | 35.4 | 44.0 | 59.7 | 69.0 |
| 1969 | 11.0 | * | * | * | * | 3.0 | 15.0 | 29.8 | 39.2 | 47.6 | 53.7 | 60.8 |
| 1968 | 11.1 | * | * | * | * | 3.1 | 13.5 | 31.2 | 38.5 | 48.3 | 58.4 | 94.6 |
| BLACK, MALE | | | | | | | | | | | | |
| 1978 | 0.3 | * | * | * | * | * | * | * | * | * | * | * |
| 1977 | 0.3 | * | * | * | * | * | * | * | * | * | * | * |
| 1976 | 0.4 | * | * | * | * | * | * | * | * | * | * | * |
| 1975 | 0.2 | * | * | * | * | * | * | * | * | * | * | * |
| 1974 | 0.3 | * | * | * | * | * | * | * | * | * | * | * |
| 1973 | 0.3 | * | * | * | * | * | * | * | * | * | * | * |
| 1972 | 0.4 | * | * | * | * | * | * | * | * | * | * | * |
| 1971 | 0.2 | * | * | * | * | * | * | * | * | * | * | * |
| 1970 | 0.3 | * | * | * | * | * | * | * | * | * | * | * |
| 1969 | 0.3 | * | * | * | * | * | * | * | * | * | * | * |
| 1968 | 0.4 | * | * | * | * | * | * | * | * | * | * | * |
| BLACK, FEMALE | | | | | | | | | | | | |
| 1978 | 22.7 | * | * | * | * | 4.5 | 24.5 | 54.6 | 77.9 | 86.3 | 108.7 | 145.5 |
| 1977 | 23.2 | * | * | * | * | 5.0 | 20.8 | 59.3 | 81.3 | 95.0 | 102.0 | 152.1 |
| 1976 | 21.2 | * | * | * | * | 3.7 | 22.3 | 51.5 | 73.6 | 85.0 | 99.1 | 157.3 |
| 1975 | 20.8 | * | * | * | * | 3.7 | 23.3 | 52.0 | 76.5 | 84.0 | 88.4 | 109.5 |
| 1974 | 22.3 | * | * | * | * | 4.6 | 25.4 | 60.0 | 76.3 | 93.9 | 83.5 | 142.3 |
| 1973 | 21.8 | * | * | * | * | 4.9 | 25.5 | 57.6 | 73.3 | 97.9 | 82.4 | 135.6 |
| 1972 | 21.2 | * | * | * | * | 5.3 | 26.6 | 55.2 | 69.4 | 92.7 | 93.6 | 114.3 |
| 1971 | 20.7 | * | * | * | * | 5.6 | 22.9 | 57.6 | 74.2 | 81.8 | 90.3 | 114.9 |
| 1970 | 19.7 | * | * | * | * | 5.9 | 24.4 | 52.0 | 64.7 | 77.3 | 101.8 | 112.1 |
| 1969 | 20.7 | * | * | * | * | 5.5 | 27.2 | 55.3 | 72.2 | 83.7 | 90.7 | 95.0 |
| 1968 | 21.0 | * | * | * | * | 5.7 | 24.6 | 58.1 | 70.8 | 85.3 | 98.6 | 146.4 |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| MALIGNANT NEOPLASMS OF GENITAL ORGANS (180-187) | | | | | | | | | | | | |
| BOTH SEXES | | | | | | | | | | | | |
| 1978 | 20.3 | * | * | * | 0.6 | 1.7 | 5.0 | 15.2 | 37.6 | 88.4 | 175.2 | 255.3 |
| 1977 | 20.1 | * | * | 0.1 | 0.7 | 1.9 | 5.0 | 15.0 | 39.2 | 88.1 | 174.5 | 247.9 |
| 1976 | 20.4 | * | * | 0.1 | 0.7 | 2.0 | 5.7 | 16.5 | 39.3 | 89.8 | 179.5 | 245.8 |
| 1975 | 20.0 | * | * | 0.1 | 0.7 | 2.0 | 5.9 | 17.0 | 38.4 | 89.3 | 174.4 | 242.1 |
| 1974 | 20.2 | * | * | 0.1 | 0.8 | 2.2 | 6.9 | 17.6 | 40.8 | 90.2 | 175.4 | 242.6 |
| 1973 | 20.2 | * | * | 0.1 | 0.8 | 2.2 | 6.7 | 18.3 | 40.6 | 90.5 | 177.7 | 247.6 |
| 1972 | 20.2 | * | * | 0.1 | 0.9 | 2.4 | 6.9 | 18.8 | 40.8 | 93.2 | 177.8 | 243.8 |
| 1971 | 20.3 | * | * | 0.1 | 0.8 | 2.6 | 7.6 | 19.3 | 41.8 | 91.9 | 178.9 | 248.0 |
| 1970 | 20.3 | * | * | 0.1 | 0.9 | 2.5 | 7.7 | 19.5 | 41.5 | 93.5 | 182.4 | 224.6 |
| 1969 | 20.4 | * | * | 0.1 | 0.9 | 2.7 | 8.4 | 20.4 | 42.4 | 92.6 | 179.2 | 251.8 |
| 1968 | 20.5 | * | * | 0.1 | 0.8 | 2.8 | 8.5 | 20.6 | 43.2 | 94.7 | 179.8 | 259.2 |
| MALE | | | | | | | | | | | | |
| 1978 | 20.8 | * | * | * | 0.7 | 1.1 | 0.9 | 3.3 | 23.1 | 107.0 | 312.2 | 579.0 |
| 1977 | 20.3 | * | * | * | 0.9 | 1.4 | 1.0 | 3.0 | 24.6 | 105.3 | 305.4 | 550.6 |
| 1976 | 20.2 | * | * | * | 0.9 | 1.5 | 1.2 | 3.6 | 23.9 | 106.3 | 306.8 | 534.2 |
| 1975 | 19.5 | * | * | * | 0.9 | 1.5 | 1.2 | 3.3 | 23.1 | 105.3 | 300.5 | 510.8 |
| 1974 | 19.4 | * | * | * | 1.0 | 1.6 | 1.3 | 3.6 | 24.3 | 106.8 | 300.9 | 504.4 |
| 1973 | 19.3 | * | * | * | 1.1 | 1.6 | 1.4 | 3.7 | 24.3 | 105.4 | 301.9 | 521.0 |
| 1972 | 18.8 | * | * | * | 1.1 | 1.6 | 1.0 | 3.2 | 23.9 | 106.1 | 295.8 | 494.6 |
| 1971 | 18.7 | * | * | * | 1.1 | 2.0 | 1.6 | 3.6 | 24.0 | 103.2 | 294.9 | 497.6 |
| 1970 | 18.5 | * | * | * | 1.2 | 1.8 | 1.4 | 3.4 | 22.8 | 103.7 | 299.4 | 434.2 |
| 1969 | 18.2 | * | * | * | 1.0 | 1.8 | 1.5 | 3.5 | 23.1 | 103.0 | 288.5 | 493.2 |
| 1968 | 18.3 | * | * | * | 0.9 | 1.9 | 1.4 | 3.9 | 23.5 | 105.7 | 285.1 | 512.3 |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| MALIGNANT NEOPLASMS OF GENITAL ORGANS (180-187) | | | | | | | | | | | | |
| FEMALE | | | | | | | | | | | | |
| 1978 | 19.8 | * | * | * | 0.5 | 2.3 | 9.0 | 26.3 | 50.3 | 74.2 | 93.6 | 109.0 |
| 1977 | 20.0 | * | * | * | 0.4 | 2.3 | 8.9 | 26.2 | 52.1 | 74.9 | 96.2 | 108.9 |
| 1976 | 20.7 | * | * | * | 0.5 | 2.5 | 10.1 | 28.4 | 52.9 | 77.2 | 102.6 | 110.3 |
| 1975 | 20.4 | * | * | * | 0.5 | 2.4 | 10.3 | 29.8 | 52.0 | 77.1 | 97.7 | 112.0 |
| 1974 | 20.9 | * | * | 0.1 | 0.6 | 2.7 | 12.3 | 30.7 | 55.5 | 77.4 | 98.2 | 112.9 |
| 1973 | 21.0 | * | * | * | 0.5 | 2.8 | 11.8 | 32.0 | 55.0 | 79.1 | 100.1 | 110.0 |
| 1972 | 21.6 | * | * | 0.1 | 0.6 | 3.3 | 12.6 | 33.2 | 55.9 | 83.3 | 102.6 | 115.5 |
| 1971 | 21.8 | * | * | 0.1 | 0.6 | 3.3 | 13.3 | 33.9 | 57.8 | 83.2 | 103.7 | 117.7 |
| 1970 | 22.0 | * | * | 0.2 | 0.6 | 3.3 | 13.6 | 34.4 | 58.2 | 85.6 | 104.9 | 107.3 |
| 1969 | 22.4 | * | * | 0.1 | 0.7 | 3.4 | 14.9 | 36.2 | 59.7 | 84.5 | 105.7 | 121.1 |
| 1968 | 22.6 | * | * | 0.2 | 0.7 | 3.7 | 15.2 | 36.2 | 61.0 | 86.0 | 107.8 | 120.4 |
| WHITE, BOTH SEXES | | | | | | | | | | | | |
| 1978 | 20.1 | * | * | * | 0.6 | 1.6 | 4.6 | 14.5 | 35.2 | 82.4 | 167.4 | 248.7 |
| 1977 | 20.0 | * | * | * | 0.7 | 1.8 | 4.5 | 14.2 | 36.7 | 82.2 | 167.8 | 242.0 |
| 1976 | 20.2 | * | * | 0.1 | 0.7 | 1.9 | 5.3 | 15.6 | 36.5 | 84.0 | 173.2 | 240.4 |
| 1975 | 19.7 | * | * | * | 0.8 | 1.9 | 5.4 | 16.0 | 35.9 | 83.2 | 168.7 | 236.2 |
| 1974 | 19.9 | * | * | 0.1 | 0.8 | 2.1 | 6.3 | 16.8 | 38.0 | 83.6 | 169.9 | 237.4 |
| 1973 | 19.8 | * | * | 0.1 | 0.8 | 2.1 | 6.0 | 16.9 | 37.5 | 84.1 | 171.6 | 242.6 |
| 1972 | 19.9 | * | * | 0.1 | 0.9 | 2.3 | 6.2 | 17.6 | 37.9 | 86.6 | 172.9 | 239.3 |
| 1971 | 19.9 | * | * | 0.1 | 0.9 | 2.5 | 6.7 | 17.9 | 38.6 | 85.5 | 173.2 | 245.1 |
| 1970 | 20.0 | * | * | 0.1 | 0.9 | 2.4 | 6.9 | 18.1 | 38.5 | 87.3 | 177.0 | 224.2 |
| 1969 | 20.0 | * | * | 0.1 | 0.9 | 2.5 | 7.6 | 19.0 | 38.6 | 87.2 | 174.0 | 250.2 |
| 1968 | 20.1 | * | * | 0.1 | 0.8 | 2.6 | 7.7 | 19.1 | 39.3 | 88.4 | 175.0 | 258.2 |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| MALIGNANT NEOPLASMS OF GENITAL ORGANS (180-187) | | | | | | | | | | | | |
| WHITE, MALE | | | | | | | | | | | | |
| 1978 | 20.2 | * | * | * | 0.8 | 1.2 | 1.0 | 3.0 | 19.9 | 96.8 | 295.4 | 565.3 |
| 1977 | 19.8 | * | * | * | 1.1 | 1.5 | 1.0 | 2.6 | 21.4 | 94.6 | 291.9 | 538.0 |
| 1976 | 19.6 | * | * | * | 1.0 | 1.6 | 1.2 | 3.3 | 20.6 | 95.7 | 293.8 | 524.0 |
| 1975 | 18.9 | * | * | * | 1.0 | 1.7 | 1.2 | 2.8 | 20.2 | 94.2 | 287.5 | 497.4 |
| 1974 | 18.9 | * | * | * | 1.1 | 1.8 | 1.4 | 3.3 | 21.1 | 95.7 | 288.4 | 496.1 |
| 1973 | 18.8 | * | * | * | 1.2 | 1.8 | 1.4 | 3.2 | 21.0 | 95.4 | 289.7 | 511.4 |
| 1972 | 18.2 | * | * | * | 1.2 | 1.8 | 1.0 | 2.7 | 20.4 | 95.1 | 284.0 | 485.5 |
| 1971 | 18.2 | * | * | * | 1.2 | 2.2 | 1.7 | 3.1 | 20.6 | 92.9 | 284.0 | 493.3 |
| 1970 | 18.0 | * | * | * | 1.3 | 1.9 | 1.5 | 2.8 | 20.0 | 93.0 | 288.1 | 437.8 |
| 1969 | 17.8 | * | * | * | 1.2 | 2.0 | 1.6 | 3.0 | 19.5 | 93.9 | 277.1 | 495.2 |
| 1968 | 17.8 | * | * | * | 1.0 | 2.0 | 1.5 | 3.2 | 19.9 | 95.7 | 276.2 | 514.8 |
| WHITE, FEMALE | | | | | | | | | | | | |
| 1978 | 20.0 | * | * | * | 0.4 | 2.1 | 8.1 | 25.3 | 48.8 | 71.4 | 92.1 | 107.6 |
| 1977 | 20.1 | * | * | * | 0.4 | 2.1 | 7.9 | 25.1 | 50.3 | 72.7 | 94.5 | 107.9 |
| 1976 | 20.9 | * | * | * | 0.4 | 2.2 | 9.4 | 27.2 | 50.7 | 75.1 | 101.3 | 109.5 |
| 1975 | 20.5 | * | * | * | 0.5 | 2.1 | 9.4 | 28.3 | 49.9 | 74.7 | 97.1 | 112.1 |
| 1974 | 21.0 | * | * | * | 0.5 | 2.4 | 11.1 | 29.5 | 53.1 | 74.4 | 97.6 | 111.3 |
| 1973 | 20.7 | * | * | * | 0.5 | 2.4 | 10.5 | 29.8 | 52.2 | 75.4 | 98.4 | 109.6 |
| 1972 | 21.6 | * | * | 0.1 | 0.5 | 2.8 | 11.3 | 31.6 | 53.6 | 80.0 | 102.8 | 115.5 |
| 1971 | 21.6 | * | * | * | 0.6 | 2.8 | 11.6 | 31.8 | 54.8 | 79.9 | 102.3 | 117.9 |
| 1970 | 21.8 | * | * | 0.2 | 0.5 | 2.9 | 12.1 | 32.5 | 55.1 | 82.9 | 104.3 | 107.3 |
| 1969 | 22.1 | * | * | 0.1 | 0.7 | 2.9 | 13.5 | 34.0 | 55.8 | 81.9 | 105.4 | 120.7 |
| 1968 | 22.2 | * | * | 0.2 | 0.7 | 3.1 | 13.7 | 34.0 | 56.9 | 82.7 | 106.4 | 120.0 |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| MALIGNANT NEOPLASMS OF GENITAL ORGANS (180-187) | | | | | | | | | | | | |
| ALL OTHER, BOTH SEXES | | | | | | | | | | | | |
| 1978 | 21.6 | * | * | * | 0.5 | 2.1 | 8.1 | 20.9 | 58.6 | 144.3 | 255.3 | 331.0 |
| 1977 | 21.4 | * | * | * | 0.4 | 1.9 | 8.5 | 21.7 | 62.1 | 143.9 | 244.0 | 316.3 |
| 1976 | 21.7 | * | * | * | 0.5 | 2.4 | 8.4 | 23.3 | 65.1 | 145.4 | 244.5 | 305.9 |
| 1975 | 21.5 | * | * | * | 0.4 | 2.6 | 9.2 | 25.8 | 62.1 | 149.2 | 236.0 | 308.9 |
| 1974 | 22.0 | * | * | * | 0.6 | 2.8 | 11.2 | 24.8 | 67.0 | 154.9 | 237.4 | 302.2 |
| 1973 | 22.7 | * | * | * | 0.6 | 3.2 | 11.2 | 30.3 | 69.8 | 154.5 | 247.0 | 306.3 |
| 1972 | 22.3 | * | * | * | 0.6 | 3.7 | 12.0 | 28.8 | 68.5 | 158.5 | 234.7 | 295.9 |
| 1971 | 23.0 | * | * | * | 0.7 | 3.7 | 13.9 | 31.0 | 72.2 | 155.1 | 247.4 | 281.7 |
| 1970 | 22.5 | * | * | * | 1.0 | 3.5 | 13.2 | 31.6 | 69.3 | 153.8 | 250.5 | 229.0 |
| 1969 | 22.9 | * | * | * | 0.6 | 3.9 | 14.0 | 33.0 | 78.7 | 146.8 | 246.8 | 267.0 |
| 1968 | 23.7 | * | * | * | 0.7 | 4.6 | 14.3 | 34.1 | 81.0 | 157.7 | 242.0 | 270.6 |
| ALL OTHER, MALE | | | | | | | | | | | | |
| 1978 | 24.8 | * | * | * | * | * | * | 6.0 | 53.0 | 200.8 | 468.4 | 719.0 |
| 1977 | 24.0 | * | * | * | * | * | * | 6.5 | 54.2 | 204.1 | 432.6 | 678.6 |
| 1976 | 23.9 | * | * | * | * | * | * | 6.2 | 56.0 | 205.4 | 431.1 | 636.4 |
| 1975 | 23.5 | * | * | * | * | * | * | 7.2 | 51.5 | 210.4 | 428.5 | 644.4 |
| 1974 | 23.2 | * | * | * | * | * | * | 6.8 | 55.0 | 213.2 | 429.3 | 590.0 |
| 1973 | 22.9 | * | * | * | * | * | * | 7.7 | 56.3 | 201.3 | 428.6 | 618.8 |
| 1972 | 23.1 | * | * | * | * | * | * | 8.2 | 57.7 | 210.9 | 422.1 | 587.2 |
| 1971 | 22.4 | * | * | * | * | * | * | 7.7 | 56.8 | 201.5 | 417.9 | 540.4 |
| 1970 | 21.9 | * | * | * | * | * | * | 9.0 | 49.6 | 204.4 | 431.0 | 402.4 |
| 1969 | 21.7 | * | * | * | * | * | * | 8.1 | 58.4 | 189.7 | 425.5 | 475.0 |
| 1968 | 22.0 | * | * | * | * | * | * | 10.1 | 59.3 | 203.0 | 391.9 | 487.8 |
| ALL OTHER, FEMALE | | | | | | | | | | | | |
| 1978 | 18.7 | * | * | * | 0.6 | 3.6 | 14.3 | 33.2 | 63.2 | 100.6 | 110.3 | 127.5 |
| 1977 | 19.0 | * | * | * | * | 3.4 | 14.9 | 34.4 | 68.8 | 96.7 | 114.9 | 121.2 |
| 1976 | 19.6 | * | * | * | 0.7 | 4.1 | 14.6 | 37.5 | 72.8 | 98.0 | 117.0 | 120.4 |
| 1975 | 19.6 | * | * | * | * | 4.6 | 16.0 | 41.5 | 71.2 | 100.6 | 103.7 | 110.8 |
| 1974 | 20.9 | * | * | * | 1.0 | 5.0 | 20.0 | 40.1 | 77.2 | 108.2 | 105.1 | 132.9 |
| 1973 | 22.6 | * | * | * | 0.9 | 5.5 | 19.9 | 49.6 | 81.4 | 116.7 | 119.8 | 116.5 |
| 1972 | 21.7 | * | * | * | 0.9 | 6.4 | 21.3 | 46.5 | 77.9 | 116.2 | 100.3 | 115.8 |
| 1971 | 23.5 | * | * | * | 1.0 | 6.5 | 24.6 | 51.1 | 85.4 | 117.1 | 121.4 | 115.1 |
| 1970 | 22.9 | * | * | * | 1.2 | 5.9 | 23.7 | 51.3 | 86.7 | 112.0 | 113.1 | 106.8 |
| 1969 | 24.1 | * | * | * | 0.9 | 6.9 | 24.8 | 54.9 | 96.7 | 111.5 | 109.7 | 126.2 |
| 1968 | 25.2 | * | * | * | 1.2 | 7.7 | 25.5 | 55.4 | 100.6 | 120.2 | 126.4 | 124.6 |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| MALIGNANT NEOPLASMS OF GENITAL ORGANS (180-187) | | | | | | | | | | | | |
| BLACK, BOTH SEXES | | | | | | | | | | | | |
| 1978 | 24.0 | * | * | * | 0.5 | 2.2 | 8.9 | 22.4 | 64.0 | 157.2 | 270.4 | 352.0 |
| 1977 | 23.8 | * | * | * | 0.4 | 2.2 | 9.2 | 23.8 | 67.3 | 156.0 | 258.1 | 337.1 |
| 1976 | 24.0 | * | * | * | 0.4 | 2.7 | 9.1 | 25.3 | 70.9 | 157.2 | 257.6 | 330.9 |
| 1975 | 23.7 | * | * | * | * | 2.8 | 9.9 | 28.1 | 67.0 | 162.5 | 246.2 | 333.1 |
| 1974 | 23.9 | * | * | * | 0.7 | 3.1 | 11.9 | 26.5 | 71.9 | 167.0 | 249.2 | 313.2 |
| 1973 | 24.8 | * | * | * | 0.6 | 3.2 | 12.0 | 33.0 | 74.7 | 166.8 | 261.5 | 328.9 |
| 1972 | 24.2 | * | * | * | 0.6 | 4.0 | 13.4 | 30.7 | 73.7 | 170.3 | 247.5 | 307.3 |
| 1971 | 24.7 | * | * | * | 0.6 | 3.9 | 14.8 | 33.0 | 77.1 | 166.6 | 260.0 | 300.9 |
| 1970 | 24.4 | * | * | * | 1.0 | 3.8 | 14.2 | 33.2 | 74.7 | 166.4 | 271.7 | 247.9 |
| 1969 | 24.6 | * | * | * | 0.6 | 4.2 | 15.1 | 35.0 | 83.6 | 156.8 | 261.8 | 291.8 |
| 1968 | 25.4 | * | * | * | 0.7 | 4.8 | 15.2 | 36.2 | 86.7 | 167.5 | 259.4 | 280.4 |
| BLACK, MALE | | | | | | | | | | | | |
| 1978 | 28.0 | * | * | * | * | * | * | 6.8 | 59.5 | 225.6 | 502.5 | 778.4 |
| 1977 | 27.0 | * | * | * | * | * | * | 7.2 | 60.7 | 225.9 | 466.4 | 740.8 |
| 1976 | 26.9 | * | * | * | * | * | * | 6.6 | 62.8 | 228.8 | 459.4 | 721.3 |
| 1975 | 26.4 | * | * | * | * | * | * | 8.2 | 57.2 | 236.9 | 455.4 | 721.7 |
| 1974 | 25.6 | * | * | * | * | * | * | 7.6 | 61.1 | 235.6 | 457.9 | 637.2 |
| 1973 | 25.2 | * | * | * | * | * | * | 8.6 | 61.8 | 223.3 | 459.3 | 700.0 |
| 1972 | 25.2 | * | * | * | * | * | * | 9.1 | 63.5 | 232.0 | 450.8 | 645.0 |
| 1971 | 24.4 | * | * | * | * | * | * | 8.4 | 62.3 | 222.1 | 450.0 | 605.0 |
| 1970 | 24.2 | * | * | * | * | * | * | 9.7 | 54.8 | 228.0 | 476.2 | 456.5 |
| 1969 | 23.5 | * | * | * | * | * | * | 9.0 | 64.1 | 207.3 | 458.1 | 548.6 |
| 1968 | 23.8 | * | * | * | * | * | * | 11.0 | 65.1 | 220.1 | 430.3 | 540.0 |
| BLACK, FEMALE | | | | | | | | | | | | |
| 1978 | 20.4 | * | * | * | * | 3.7 | 15.7 | 35.3 | 67.7 | 106.2 | 115.4 | 132.3 |
| 1977 | 20.9 | * | * | * | * | 3.9 | 16.1 | 37.6 | 72.8 | 103.5 | 120.6 | 126.6 |
| 1976 | 21.4 | * | * | * | 0.7 | 4.6 | 15.7 | 40.8 | 77.5 | 102.9 | 122.3 | 124.7 |
| 1975 | 21.2 | * | * | * | * | 4.9 | 17.3 | 44.8 | 75.0 | 106.2 | 104.6 | 120.2 |
| 1974 | 22.4 | * | * | * | 1.0 | 5.6 | 21.2 | 42.4 | 80.8 | 114.5 | 108.1 | 134.6 |
| 1973 | 24.4 | * | * | * | 1.0 | 5.8 | 21.3 | 53.6 | 85.5 | 123.2 | 126.0 | 120.5 |
| 1972 | 23.3 | * | * | * | 0.9 | 7.1 | 23.6 | 49.1 | 82.2 | 122.6 | 103.6 | 114.3 |
| 1971 | 24.9 | * | * | * | 1.0 | 6.8 | 26.1 | 54.0 | 89.4 | 122.6 | 123.6 | 119.4 |
| 1970 | 24.6 | * | * | * | 1.3 | 6.4 | 25.5 | 53.5 | 91.6 | 117.5 | 121.8 | 113.5 |
| 1969 | 25.6 | * | * | * | 0.9 | 7.4 | 26.7 | 57.5 | 100.4 | 116.6 | 116.3 | 133.3 |
| 1968 | 26.8 | * | * | * | 1.2 | 8.3 | 27.0 | 58.1 | 105.3 | 124.9 | 132.4 | 123.2 |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|

MALIGNANT NEOPLASMS OF URINARY ORGANS (188, 189)

BOTH SEXES

| | | | | | | | | | | | | |
|------|-----|---|-----|-----|-----|-----|-----|-----|------|------|------|------|
| 1978 | 7.9 | * | 0.2 | 0.1 | 0.1 | 0.2 | 1.1 | 5.5 | 15.9 | 36.2 | 66.8 | 99.4 |
| 1977 | 7.9 | * | 0.3 | 0.1 | 0.1 | 0.2 | 1.3 | 5.9 | 15.9 | 37.0 | 67.5 | 95.7 |
| 1976 | 7.8 | * | 0.3 | 0.1 | 0.1 | 0.2 | 1.2 | 5.8 | 16.3 | 36.4 | 66.9 | 96.9 |
| 1975 | 7.7 | * | 0.3 | 0.1 | 0.1 | 0.2 | 1.3 | 5.6 | 16.3 | 37.0 | 66.0 | 94.0 |
| 1974 | 7.7 | * | 0.3 | 0.1 | 0.1 | 0.3 | 1.3 | 5.7 | 16.7 | 38.0 | 67.9 | 89.4 |
| 1973 | 7.5 | * | 0.4 | 0.2 | 0.1 | 0.2 | 1.2 | 5.4 | 16.8 | 36.8 | 66.5 | 94.5 |
| 1972 | 7.5 | * | 0.4 | 0.2 | 0.1 | 0.1 | 1.4 | 5.4 | 16.4 | 38.2 | 66.4 | 96.5 |
| 1971 | 7.4 | * | 0.4 | 0.2 | 0.1 | 0.3 | 1.2 | 5.7 | 16.5 | 37.3 | 66.0 | 91.3 |
| 1970 | 7.6 | * | 0.4 | 0.2 | 0.1 | 0.2 | 1.4 | 5.7 | 17.5 | 37.6 | 71.1 | 88.8 |
| 1969 | 7.4 | * | 0.5 | 0.2 | 0.1 | 0.3 | 1.3 | 5.7 | 17.1 | 37.4 | 67.5 | 91.1 |
| 1968 | 7.4 | * | 0.5 | 0.3 | 0.1 | 0.3 | 1.4 | 5.8 | 16.9 | 38.3 | 66.8 | 92.6 |

MALE

| | | | | | | | | | | | | |
|------|------|---|-----|-----|-----|-----|-----|-----|------|------|-------|-------|
| 1978 | 10.7 | * | * | 0.1 | * | 0.3 | 1.6 | 8.3 | 23.8 | 57.6 | 114.1 | 170.9 |
| 1977 | 10.7 | * | * | * | * | 0.2 | 1.8 | 8.6 | 23.6 | 58.2 | 116.0 | 165.1 |
| 1976 | 10.7 | * | * | 0.1 | * | 0.3 | 1.6 | 8.3 | 25.1 | 58.2 | 113.2 | 166.7 |
| 1975 | 10.5 | * | * | 0.1 | * | 0.2 | 1.7 | 8.1 | 24.6 | 59.5 | 110.5 | 163.8 |
| 1974 | 10.6 | * | 0.4 | 0.1 | 0.1 | 0.3 | 1.8 | 8.6 | 25.0 | 60.4 | 114.0 | 151.3 |
| 1973 | 10.3 | * | 0.4 | 0.2 | * | 0.3 | 1.7 | 7.8 | 25.2 | 59.3 | 110.4 | 155.2 |
| 1972 | 10.4 | * | 0.5 | 0.2 | 0.1 | * | 1.9 | 7.8 | 24.9 | 61.4 | 113.2 | 156.7 |
| 1971 | 10.1 | * | 0.4 | 0.1 | 0.1 | 0.3 | 1.7 | 8.2 | 24.5 | 60.2 | 108.6 | 143.7 |
| 1970 | 10.4 | * | 0.4 | 0.2 | 0.1 | 0.3 | 1.9 | 8.0 | 26.4 | 60.3 | 112.2 | 140.5 |
| 1969 | 10.2 | * | 0.4 | 0.2 | * | 0.3 | 1.8 | 8.4 | 26.0 | 59.4 | 107.1 | 138.1 |
| 1968 | 10.3 | * | 0.5 | 0.3 | 0.1 | 0.3 | 2.0 | 8.5 | 25.4 | 60.7 | 105.7 | 148.2 |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|--|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| MALIGNANT NEOPLASMS OF URINARY ORGANS (188, 189) | | | | | | | | | | | | |
| FEMALE | | | | | | | | | | | | |
| 1978 | 5.3 | * | * | 0.2 | * | 0.2 | 0.7 | 2.9 | 8.9 | 19.9 | 38.7 | 67.1 |
| 1977 | 5.3 | * | * | * | * | 0.2 | 0.9 | 3.4 | 9.1 | 20.7 | 38.5 | 63.9 |
| 1976 | 5.1 | * | 0.3 | 0.2 | * | 0.2 | 0.8 | 3.4 | 8.6 | 19.8 | 38.9 | 64.2 |
| 1975 | 5.0 | * | 0.3 | 0.1 | 0.1 | 0.2 | 0.9 | 3.3 | 8.8 | 19.7 | 38.8 | 60.1 |
| 1974 | 5.0 | * | * | 0.1 | * | 0.2 | 0.8 | 3.0 | 9.4 | 20.9 | 39.6 | 58.7 |
| 1973 | 4.9 | * | 0.3 | 0.2 | * | 0.2 | 0.7 | 3.2 | 9.3 | 19.5 | 39.0 | 63.9 |
| 1972 | 4.8 | * | 0.4 | 0.2 | 0.1 | 0.1 | 0.9 | 3.1 | 8.8 | 20.3 | 36.5 | 65.7 |
| 1971 | 4.8 | * | 0.4 | 0.2 | * | 0.2 | 0.8 | 3.5 | 9.3 | 19.5 | 38.5 | 63.9 |
| 1970 | 5.0 | * | 0.5 | 0.2 | * | 0.2 | 1.0 | 3.5 | 9.4 | 20.1 | 44.0 | 59.9 |
| 1969 | 4.8 | * | 0.5 | 0.2 | * | 0.2 | 0.9 | 3.2 | 9.0 | 20.1 | 40.9 | 65.7 |
| 1968 | 4.7 | * | 0.5 | 0.3 | * | 0.2 | 1.0 | 3.2 | 9.2 | 20.6 | 40.1 | 62.1 |
| WHITE, BOTH SEXES | | | | | | | | | | | | |
| 1978 | 8.4 | * | 0.2 | 0.1 | 0.1 | 0.2 | 1.1 | 5.5 | 16.0 | 36.3 | 68.8 | 103.4 |
| 1977 | 8.4 | * | 0.2 | 0.1 | 0.1 | 0.2 | 1.3 | 5.9 | 16.1 | 37.4 | 69.2 | 98.7 |
| 1976 | 8.2 | * | 0.3 | 0.1 | 0.1 | 0.2 | 1.2 | 5.7 | 16.2 | 36.8 | 68.8 | 99.8 |
| 1975 | 8.1 | * | 0.3 | 0.1 | 0.1 | 0.2 | 1.3 | 5.7 | 16.2 | 37.4 | 67.8 | 96.9 |
| 1974 | 8.2 | * | 0.4 | 0.1 | 0.1 | 0.2 | 1.2 | 5.7 | 16.8 | 38.4 | 70.1 | 92.9 |
| 1973 | 7.9 | * | 0.4 | 0.2 | 0.1 | 0.2 | 1.2 | 5.3 | 16.7 | 37.0 | 68.6 | 98.0 |
| 1972 | 7.9 | * | 0.4 | 0.2 | 0.1 | 0.1 | 1.3 | 5.2 | 16.2 | 38.5 | 68.0 | 99.6 |
| 1971 | 7.8 | * | 0.4 | 0.2 | 0.1 | 0.2 | 1.2 | 5.6 | 16.3 | 37.7 | 67.8 | 95.0 |
| 1970 | 8.0 | * | 0.4 | 0.2 | 0.1 | 0.2 | 1.4 | 5.5 | 17.6 | 38.1 | 73.0 | 93.0 |
| 1969 | 7.8 | * | 0.5 | 0.2 | * | 0.2 | 1.3 | 5.6 | 16.7 | 37.9 | 69.2 | 94.4 |
| 1968 | 7.8 | * | 0.5 | 0.3 | 0.1 | 0.2 | 1.4 | 5.6 | 16.8 | 38.7 | 68.6 | 96.6 |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|--|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| MALIGNANT NEOPLASMS OF URINARY ORGANS (188, 189) | | | | | | | | | | | | |
| WHITE, MALE | | | | | | | | | | | | |
| 1978 | 11.4 | * | * | * | * | 0.3 | 1.5 | 8.3 | 24.1 | 58.2 | 118.6 | 180.1 |
| 1977 | 11.4 | * | * | * | * | 0.2 | 1.8 | 8.6 | 24.0 | 59.5 | 119.9 | 173.6 |
| 1976 | 11.3 | * | * | * | * | 0.3 | 1.6 | 8.3 | 24.9 | 59.4 | 118.6 | 172.8 |
| 1975 | 11.2 | * | * | * | * | 0.2 | 1.8 | 8.3 | 24.8 | 60.9 | 115.1 | 172.0 |
| 1974 | 11.2 | * | 0.4 | * | * | 0.3 | 1.7 | 8.6 | 25.3 | 61.7 | 119.2 | 159.0 |
| 1973 | 10.9 | * | 0.4 | 0.2 | * | 0.3 | 1.8 | 7.7 | 25.3 | 60.6 | 114.7 | 164.5 |
| 1972 | 11.0 | * | 0.5 | 0.2 | * | * | 1.7 | 7.7 | 24.8 | 63.3 | 116.5 | 163.8 |
| 1971 | 10.7 | * | 0.4 | 0.2 | * | 0.3 | 1.7 | 8.1 | 24.6 | 61.5 | 112.2 | 150.3 |
| 1970 | 11.0 | * | 0.4 | 0.2 | * | 0.3 | 1.9 | 7.8 | 26.7 | 61.6 | 116.0 | 148.5 |
| 1969 | 10.7 | * | 0.4 | 0.2 | * | 0.3 | 1.7 | 8.4 | 25.8 | 60.8 | 110.7 | 143.9 |
| 1968 | 10.8 | * | 0.5 | 0.3 | * | 0.3 | 1.8 | 8.3 | 25.4 | 62.1 | 109.4 | 157.6 |
| WHITE, FEMALE | | | | | | | | | | | | |
| 1978 | 5.5 | * | * | 0.1 | * | 0.2 | 0.7 | 2.8 | 8.8 | 19.5 | 39.6 | 69.2 |
| 1977 | 5.5 | * | * | * | * | 0.2 | 0.8 | 3.3 | 9.0 | 20.5 | 39.3 | 64.9 |
| 1976 | 5.3 | * | * | 0.1 | * | 0.2 | 0.8 | 3.3 | 8.5 | 19.5 | 39.1 | 66.1 |
| 1975 | 5.2 | * | * | * | * | 0.2 | 0.8 | 3.3 | 8.6 | 19.5 | 39.3 | 61.2 |
| 1974 | 5.2 | * | * | * | * | 0.2 | 0.8 | 2.9 | 9.1 | 20.7 | 40.1 | 60.7 |
| 1973 | 5.1 | * | * | 0.2 | * | 0.2 | 0.7 | 3.0 | 9.1 | 19.0 | 40.0 | 65.2 |
| 1972 | 4.9 | * | 0.4 | 0.2 | * | * | 0.9 | 2.9 | 8.5 | 19.4 | 37.4 | 67.4 |
| 1971 | 5.0 | * | * | 0.2 | * | 0.2 | 0.7 | 3.2 | 8.8 | 19.3 | 39.4 | 66.7 |
| 1970 | 5.2 | * | 0.4 | 0.1 | * | 0.2 | 0.9 | 3.3 | 9.4 | 19.9 | 44.9 | 62.7 |
| 1969 | 4.9 | * | 0.5 | 0.2 | * | 0.2 | 0.9 | 3.0 | 8.6 | 20.2 | 41.5 | 68.3 |
| 1968 | 4.9 | * | 0.5 | 0.3 | * | 0.2 | 0.9 | 3.1 | 9.0 | 20.4 | 41.0 | 63.7 |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|--|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| MALIGNANT NEOPLASMS OF URINARY ORGANS (188, 189) | | | | | | | | | | | | |
| ALL OTHER, BOTH SEXES | | | | | | | | | | | | |
| 1978 | 4.9 | * | * | * | * | * | 1.5 | 5.7 | 15.2 | 35.7 | 46.2 | 53.6 |
| 1977 | 4.8 | * | * | * | * | * | 1.5 | 6.1 | 14.3 | 32.9 | 49.6 | 61.3 |
| 1976 | 5.0 | * | * | * | * | * | 1.2 | 6.4 | 17.2 | 33.3 | 47.6 | 64.1 |
| 1975 | 4.7 | * | * | * | * | 0.6 | 1.3 | 5.0 | 16.7 | 32.4 | 46.1 | 60.3 |
| 1974 | 4.7 | * | * | * | * | * | 1.7 | 5.9 | 16.7 | 34.3 | 44.6 | 48.1 |
| 1973 | 4.7 | * | * | * | * | * | 0.9 | 6.5 | 17.1 | 34.8 | 42.5 | 52.8 |
| 1972 | 5.1 | * | * | * | * | * | 2.1 | 7.0 | 18.0 | 35.4 | 47.8 | 60.2 |
| 1971 | 4.8 | * | * | * | * | * | 1.4 | 7.2 | 18.2 | 32.9 | 44.6 | 48.3 |
| 1970 | 4.8 | * | * | * | * | * | 1.7 | 7.5 | 16.6 | 33.4 | 47.0 | 45.5 |
| 1969 | 4.8 | * | * | * | * | * | 1.5 | 6.6 | 20.1 | 31.5 | 46.5 | 54.1 |
| 1968 | 4.9 | * | * | * | * | * | 2.2 | 7.7 | 17.9 | 33.7 | 42.3 | 48.0 |
| ALL OTHER, MALE | | | | | | | | | | | | |
| 1978 | 6.4 | * | * | * | * | * | 2.1 | 8.2 | 21.2 | 51.3 | 71.4 | 75.9 |
| 1977 | 6.1 | * | * | * | * | * | 1.4 | 8.3 | 20.0 | 46.1 | 79.3 | 78.6 |
| 1976 | 6.5 | * | * | * | * | * | * | 8.8 | 27.1 | 46.8 | 62.6 | 105.5 |
| 1975 | 5.9 | * | * | * | * | * | 1.6 | 6.7 | 23.3 | 45.6 | 64.5 | 81.5 |
| 1974 | 6.1 | * | * | * | * | * | 2.5 | 8.5 | 22.5 | 47.8 | 60.7 | 72.0 |
| 1973 | 6.0 | * | * | * | * | * | * | 8.5 | 24.0 | 47.5 | 64.3 | 60.4 |
| 1972 | 6.5 | * | * | * | * | * | 3.0 | 8.9 | 26.2 | 43.7 | 77.4 | 85.1 |
| 1971 | 6.2 | * | * | * | * | * | * | 9.1 | 23.4 | 48.0 | 68.1 | 78.7 |
| 1970 | 6.3 | * | * | * | * | * | 2.2 | 10.0 | 23.8 | 47.5 | 67.9 | 70.4 |
| 1969 | 6.3 | * | * | * | * | * | 1.8 | 8.2 | 28.0 | 46.4 | 64.9 | 84.1 |
| 1968 | 6.5 | * | * | * | * | * | 2.9 | 11.2 | 25.5 | 47.1 | 61.3 | 56.1 |
| ALL OTHER, FEMALE | | | | | | | | | | | | |
| 1978 | 3.5 | * | * | * | * | * | * | 3.6 | 10.1 | 23.7 | 29.0 | 42.2 |
| 1977 | 3.6 | * | * | * | * | * | 1.6 | 4.2 | 9.7 | 22.6 | 29.2 | 51.9 |
| 1976 | 3.6 | * | * | * | * | * | * | 4.3 | 8.9 | 22.6 | 37.2 | 40.8 |
| 1975 | 3.6 | * | * | * | * | * | * | 3.6 | 11.0 | 22.0 | 33.3 | 47.3 |
| 1974 | 3.5 | * | * | * | * | * | * | 3.7 | 11.7 | 23.5 | 33.4 | 34.1 |
| 1973 | 3.5 | * | * | * | * | * | * | 4.8 | 11.2 | 24.4 | 27.4 | 48.1 |
| 1972 | 3.8 | * | * | * | * | * | 1.3 | 5.3 | 10.8 | 28.7 | 26.6 | 44.7 |
| 1971 | 3.6 | * | * | * | * | * | * | 5.6 | 13.5 | 20.6 | 27.1 | 28.8 |
| 1970 | 3.4 | * | * | * | * | * | * | 5.4 | 10.2 | 21.7 | 31.1 | 28.0 |
| 1969 | 3.5 | * | * | * | * | * | * | 5.2 | 13.0 | 19.3 | 32.4 | 33.8 |
| 1968 | 3.4 | * | * | * | * | * | 1.6 | 4.7 | 11.0 | 22.5 | 27.6 | 42.6 |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|--|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| MALIGNANT NEOPLASMS OF URINARY ORGANS (188, 189) | | | | | | | | | | | | |
| BLACK, BOTH SEXES | | | | | | | | | | | | |
| 1978 | 5.4 | * | * | * | * | * | 1.8 | 6.3 | 16.4 | 38.4 | 46.7 | 54.7 |
| 1977 | 5.2 | * | * | * | * | * | 1.8 | 6.4 | 15.4 | 35.0 | 51.0 | 61.5 |
| 1976 | 5.3 | * | * | * | * | * | 1.2 | 6.8 | 18.2 | 35.0 | 48.0 | 65.4 |
| 1975 | 5.0 | * | * | * | * | 0.7 | 1.5 | 5.3 | 17.9 | 33.3 | 46.5 | 60.0 |
| 1974 | 5.0 | * | * | * | * | * | 1.8 | 6.5 | 16.9 | 35.7 | 45.2 | 49.6 |
| 1973 | 4.9 | * | * | * | * | * | 1.0 | 6.8 | 18.1 | 35.7 | 41.7 | 50.9 |
| 1972 | 5.3 | * | * | * | * | * | 2.3 | 7.7 | 18.6 | 35.3 | 47.1 | 63.6 |
| 1971 | 5.1 | * | * | * | * | * | 1.5 | 7.8 | 18.7 | 34.2 | 44.0 | 50.5 |
| 1970 | 5.2 | * | * | * | * | * | 1.8 | 8.2 | 17.5 | 35.5 | 50.7 | 44.3 |
| 1969 | 5.1 | * | * | * | * | * | 1.6 | 7.2 | 20.8 | 32.3 | 48.1 | 54.6 |
| 1968 | 5.1 | * | * | * | * | * | 2.3 | 8.4 | 18.4 | 34.7 | 41.6 | 47.8 |
| BLACK, MALE | | | | | | | | | | | | |
| 1978 | 7.0 | * | * | * | * | * | 2.5 | 8.9 | 22.7 | 56.4 | 72.8 | 78.4 |
| 1977 | 6.6 | * | * | * | * | * | 1.7 | 8.7 | 21.6 | 49.8 | 82.1 | 77.6 |
| 1976 | 7.0 | * | * | * | * | * | * | 9.8 | 28.7 | 50.0 | 64.1 | 112.8 |
| 1975 | 6.3 | * | * | * | * | * | 1.8 | 6.9 | 25.4 | 47.6 | 64.9 | 87.0 |
| 1974 | 6.3 | * | * | * | * | * | 2.6 | 9.3 | 22.4 | 50.1 | 62.2 | 74.4 |
| 1973 | 6.2 | * | * | * | * | * | * | 8.7 | 25.7 | 49.2 | 63.2 | 58.5 |
| 1972 | 6.8 | * | * | * | * | * | 3.3 | 9.7 | 28.0 | 44.3 | 78.2 | 95.0 |
| 1971 | 6.5 | * | * | * | * | * | * | 9.9 | 24.3 | 50.3 | 67.7 | 85.0 |
| 1970 | 6.8 | * | * | * | * | * | 2.5 | 11.0 | 25.3 | 51.6 | 73.5 | 69.6 |
| 1969 | 6.6 | * | * | * | * | * | 2.0 | 9.0 | 29.6 | 47.3 | 67.1 | 86.5 |
| 1968 | 6.7 | * | * | * | * | * | 3.1 | 12.1 | 26.7 | 47.9 | 61.2 | 57.1 |
| BLACK, FEMALE | | | | | | | | | | | | |
| 1978 | 3.9 | * | * | * | * | * | * | 4.1 | 11.2 | 24.9 | 29.3 | 42.4 |
| 1977 | 4.0 | * | * | * | * | * | 1.8 | 4.5 | 10.4 | 23.8 | 30.5 | 53.2 |
| 1976 | 3.9 | * | * | * | * | * | * | 4.3 | 9.5 | 23.6 | 37.2 | 40.4 |
| 1975 | 3.8 | * | * | * | * | * | * | 3.9 | 11.7 | 22.5 | 34.1 | 45.2 |
| 1974 | 3.9 | * | * | * | * | * | * | 4.1 | 12.4 | 24.7 | 33.7 | 35.9 |
| 1973 | 3.8 | * | * | * | * | * | * | 5.1 | 11.7 | 25.3 | 27.0 | 46.6 |
| 1972 | 3.9 | * | * | * | * | * | 1.5 | 5.9 | 10.6 | 28.2 | 25.0 | 45.7 |
| 1971 | 3.8 | * | * | * | * | * | * | 6.0 | 13.9 | 21.4 | 27.0 | 29.9 |
| 1970 | 3.6 | * | * | * | * | * | * | 5.7 | 10.9 | 22.7 | 33.9 | 28.0 |
| 1969 | 3.7 | * | * | * | * | * | * | 5.7 | 13.3 | 20.5 | 33.9 | 35.0 |
| 1968 | 3.6 | * | * | * | * | * | 1.7 | 5.1 | 11.2 | 24.0 | 27.0 | 42.9 |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| MALIGNANT NEOPLASMS OF ALL OTHER AND UNSPECIFIED SITES (170-173, 190-199) | | | | | | | | | | | | |
| BOTH SEXES | | | | | | | | | | | | |
| 1978 | 22.0 | 1.8 | 2.3 | 1.5 | 2.2 | 3.8 | 9.1 | 24.7 | 51.7 | 87.7 | 134.6 | 180.9 |
| 1977 | 21.5 | 1.8 | 2.6 | 1.7 | 2.2 | 3.6 | 9.2 | 24.1 | 50.3 | 87.3 | 129.8 | 182.5 |
| 1976 | 21.1 | 1.3 | 2.5 | 1.9 | 2.3 | 4.0 | 9.2 | 23.7 | 51.1 | 85.3 | 126.3 | 179.2 |
| 1975 | 20.7 | 1.6 | 2.3 | 1.8 | 2.3 | 3.9 | 9.0 | 23.9 | 49.5 | 84.8 | 125.5 | 176.6 |
| 1974 | 20.4 | 1.9 | 2.7 | 1.8 | 2.5 | 3.7 | 9.1 | 24.3 | 50.3 | 83.6 | 123.2 | 172.2 |
| 1973 | 20.2 | 2.0 | 2.8 | 1.9 | 2.6 | 3.9 | 9.8 | 23.6 | 50.1 | 82.1 | 121.5 | 174.0 |
| 1972 | 19.9 | 1.7 | 2.6 | 2.0 | 2.6 | 3.8 | 9.5 | 23.0 | 49.8 | 81.9 | 122.2 | 175.9 |
| 1971 | 19.7 | 1.4 | 2.7 | 2.0 | 2.7 | 4.0 | 9.7 | 23.0 | 49.4 | 81.6 | 118.5 | 166.8 |
| 1970 | 19.2 | 1.7 | 2.8 | 2.1 | 2.8 | 3.7 | 9.7 | 23.2 | 48.7 | 79.0 | 116.9 | 145.0 |
| 1969 | 18.8 | 1.9 | 2.9 | 2.1 | 2.6 | 4.1 | 10.0 | 22.8 | 46.7 | 76.9 | 111.8 | 166.2 |
| 1968 | 18.7 | 1.8 | 3.0 | 2.2 | 2.6 | 3.9 | 9.7 | 23.3 | 47.2 | 76.0 | 112.0 | 162.6 |
| MALE | | | | | | | | | | | | |
| 1978 | 23.8 | 1.9 | 2.5 | 1.6 | 2.6 | 4.5 | 10.4 | 29.4 | 63.1 | 109.2 | 164.0 | 226.8 |
| 1977 | 23.2 | 2.1 | 2.9 | 1.9 | 2.8 | 4.4 | 10.4 | 28.1 | 61.0 | 109.0 | 159.2 | 219.9 |
| 1976 | 22.7 | 1.4 | 2.5 | 2.1 | 2.8 | 4.3 | 10.4 | 27.2 | 62.3 | 106.2 | 154.4 | 203.1 |
| 1975 | 22.0 | 1.9 | 2.5 | 1.9 | 2.6 | 4.5 | 10.1 | 27.8 | 58.8 | 104.4 | 150.2 | 200.2 |
| 1974 | 22.1 | 2.0 | 2.8 | 2.0 | 3.0 | 4.2 | 10.2 | 28.4 | 59.6 | 105.0 | 148.6 | 204.1 |
| 1973 | 21.7 | 1.8 | 2.9 | 1.9 | 3.1 | 4.7 | 11.1 | 26.7 | 59.5 | 102.4 | 145.4 | 203.5 |
| 1972 | 21.5 | 1.4 | 2.8 | 2.1 | 3.2 | 4.8 | 10.5 | 26.5 | 59.7 | 101.7 | 143.8 | 195.0 |
| 1971 | 21.1 | 1.5 | 2.9 | 2.3 | 3.3 | 4.6 | 11.1 | 25.5 | 58.6 | 98.9 | 139.3 | 190.8 |
| 1970 | 20.6 | 1.3 | 3.1 | 2.3 | 3.5 | 4.3 | 10.6 | 26.5 | 58.0 | 95.1 | 133.3 | 155.6 |
| 1969 | 20.1 | 2.2 | 2.9 | 2.3 | 3.3 | 4.7 | 10.8 | 25.4 | 56.4 | 92.2 | 125.2 | 192.4 |
| 1968 | 20.3 | 1.6 | 3.3 | 2.3 | 3.2 | 4.5 | 11.0 | 26.0 | 56.3 | 92.7 | 132.5 | 175.9 |

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| MALIGNANT NEOPLASMS OF ALL OTHER AND UNSPECIFIED SITES (170-173, 190-199) | | | | | | | | | | | | |
| FEMALE | | | | | | | | | | | | |
| 1978 | 20.4 | 1.8 | 2.2 | 1.4 | 1.8 | 3.1 | 7.9 | 20.4 | 41.7 | 71.3 | 117.1 | 160.2 |
| 1977 | 19.9 | 1.6 | 2.2 | 1.6 | 1.7 | 2.9 | 8.1 | 20.3 | 40.9 | 70.7 | 112.2 | 165.3 |
| 1976 | 19.6 | * | 2.5 | 1.6 | 1.7 | 3.6 | 8.0 | 20.4 | 41.2 | 69.4 | 109.3 | 168.0 |
| 1975 | 19.4 | 1.4 | 2.1 | 1.8 | 1.9 | 3.2 | 8.0 | 20.3 | 41.2 | 69.7 | 110.5 | 165.1 |
| 1974 | 18.9 | 1.8 | 2.6 | 1.7 | 2.0 | 3.1 | 8.0 | 20.4 | 42.1 | 67.2 | 107.5 | 156.4 |
| 1973 | 18.8 | 2.1 | 2.6 | 1.9 | 2.1 | 3.1 | 8.6 | 20.8 | 41.8 | 66.6 | 106.5 | 159.1 |
| 1972 | 18.5 | 2.0 | 2.3 | 1.9 | 1.9 | 2.9 | 8.5 | 19.7 | 40.9 | 66.7 | 108.4 | 166.1 |
| 1971 | 18.3 | 1.4 | 2.4 | 1.7 | 2.0 | 3.5 | 8.4 | 20.7 | 41.3 | 68.1 | 105.1 | 154.2 |
| 1970 | 17.9 | 2.0 | 2.6 | 1.9 | 2.1 | 3.1 | 8.8 | 20.2 | 40.2 | 66.4 | 106.0 | 139.1 |
| 1969 | 17.5 | 1.6 | 2.8 | 2.0 | 2.0 | 3.4 | 9.2 | 20.4 | 38.1 | 64.9 | 102.9 | 152.0 |
| 1968 | 17.2 | 2.0 | 2.8 | 2.1 | 2.1 | 3.3 | 8.5 | 20.7 | 39.0 | 62.9 | 98.0 | 155.5 |
| WHITE, BOTH SEXES | | | | | | | | | | | | |
| 1978 | 22.7 | 1.9 | 2.4 | 1.5 | 2.2 | 3.9 | 9.1 | 24.1 | 50.5 | 86.0 | 134.9 | 183.0 |
| 1977 | 22.2 | 1.9 | 2.6 | 1.7 | 2.3 | 3.8 | 9.2 | 23.3 | 49.5 | 86.3 | 129.7 | 184.3 |
| 1976 | 21.7 | 1.5 | 2.6 | 1.8 | 2.3 | 4.2 | 9.2 | 23.1 | 50.0 | 84.1 | 126.3 | 182.5 |
| 1975 | 21.3 | 1.7 | 2.4 | 1.8 | 2.3 | 4.0 | 9.0 | 23.3 | 48.3 | 83.4 | 126.4 | 180.7 |
| 1974 | 21.0 | 1.7 | 2.9 | 1.8 | 2.5 | 3.8 | 9.1 | 23.7 | 49.2 | 82.1 | 123.9 | 175.2 |
| 1973 | 20.7 | 2.2 | 2.8 | 1.8 | 2.6 | 4.0 | 9.8 | 23.1 | 49.1 | 80.6 | 122.3 | 175.7 |
| 1972 | 20.5 | 1.7 | 2.7 | 2.1 | 2.7 | 3.9 | 9.4 | 22.4 | 49.0 | 80.6 | 123.3 | 178.4 |
| 1971 | 20.2 | 1.4 | 2.8 | 2.0 | 2.7 | 4.2 | 9.8 | 22.3 | 48.4 | 80.6 | 119.1 | 170.8 |
| 1970 | 19.7 | 1.7 | 2.8 | 2.2 | 2.9 | 3.8 | 9.6 | 22.7 | 47.8 | 77.8 | 117.8 | 148.0 |
| 1969 | 19.2 | 2.1 | 2.9 | 2.2 | 2.7 | 4.2 | 9.9 | 22.3 | 45.7 | 75.8 | 112.5 | 170.3 |
| 1968 | 19.1 | 1.8 | 3.1 | 2.2 | 2.6 | 3.9 | 9.6 | 22.9 | 46.2 | 75.0 | 112.6 | 167.3 |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| MALIGNANT NEOPLASMS OF ALL OTHER AND UNSPECIFIED SITES (170-173, 190-199) | | | | | | | | | | | | |
| WHITE, MALE | | | | | | | | | | | | |
| 1978 | 24.3 | 1.8 | 2.6 | 1.6 | 2.6 | 4.7 | 10.5 | 28.6 | 61.2 | 107.2 | 165.4 | 229.1 |
| 1977 | 23.8 | 2.2 | 3.1 | 1.9 | 2.9 | 4.6 | 10.4 | 27.0 | 59.9 | 108.2 | 159.5 | 222.1 |
| 1976 | 23.2 | 1.5 | 2.6 | 2.1 | 2.8 | 4.6 | 10.3 | 26.5 | 60.6 | 104.5 | 154.5 | 209.4 |
| 1975 | 22.5 | 1.9 | 2.6 | 1.9 | 2.6 | 4.7 | 10.2 | 26.8 | 57.2 | 103.2 | 151.3 | 207.2 |
| 1974 | 22.6 | 1.7 | 2.9 | 2.0 | 3.1 | 4.4 | 10.2 | 27.7 | 58.2 | 103.8 | 150.1 | 207.8 |
| 1973 | 22.1 | 2.1 | 2.9 | 1.9 | 3.2 | 4.8 | 11.1 | 26.0 | 58.3 | 100.5 | 145.6 | 206.3 |
| 1972 | 22.0 | * | 2.9 | 2.2 | 3.3 | 4.9 | 10.3 | 26.0 | 59.1 | 100.4 | 144.5 | 202.5 |
| 1971 | 21.5 | 1.6 | 3.2 | 2.3 | 3.4 | 4.7 | 11.1 | 24.7 | 57.4 | 98.0 | 139.8 | 198.1 |
| 1970 | 21.0 | 1.4 | 3.0 | 2.4 | 3.6 | 4.4 | 10.6 | 25.6 | 56.9 | 94.3 | 134.2 | 162.0 |
| 1969 | 20.6 | 2.4 | 2.9 | 2.4 | 3.4 | 4.8 | 10.6 | 25.0 | 55.2 | 91.8 | 125.9 | 199.5 |
| 1968 | 20.7 | 1.6 | 3.2 | 2.3 | 3.3 | 4.6 | 10.9 | 25.6 | 55.3 | 91.6 | 133.1 | 181.7 |
| WHITE, FEMALE | | | | | | | | | | | | |
| 1978 | 21.1 | 2.1 | 2.3 | 1.4 | 1.8 | 3.2 | 7.7 | 19.9 | 40.9 | 69.8 | 117.0 | 162.4 |
| 1977 | 20.6 | 1.6 | 2.1 | 1.6 | 1.7 | 3.0 | 8.1 | 19.8 | 40.3 | 69.6 | 112.1 | 167.2 |
| 1976 | 20.4 | * | 2.5 | 1.6 | 1.8 | 3.8 | 8.1 | 19.9 | 40.6 | 68.5 | 109.5 | 170.1 |
| 1975 | 20.1 | 1.6 | 2.1 | 1.7 | 2.0 | 3.3 | 7.9 | 20.0 | 40.3 | 68.2 | 111.4 | 168.2 |
| 1974 | 19.5 | 1.7 | 2.9 | 1.7 | 2.0 | 3.3 | 8.1 | 20.0 | 41.1 | 65.6 | 107.9 | 159.3 |
| 1973 | 19.3 | 2.2 | 2.7 | 1.7 | 2.0 | 3.2 | 8.5 | 20.5 | 40.9 | 65.4 | 107.8 | 160.8 |
| 1972 | 19.0 | 2.2 | 2.4 | 2.0 | 2.1 | 2.9 | 8.5 | 19.1 | 39.9 | 65.4 | 109.9 | 166.3 |
| 1971 | 18.9 | * | 2.3 | 1.8 | 2.0 | 3.6 | 8.5 | 20.1 | 40.3 | 67.2 | 105.9 | 156.9 |
| 1970 | 18.5 | 2.0 | 2.7 | 1.9 | 2.2 | 3.2 | 8.7 | 19.9 | 39.6 | 65.0 | 107.0 | 140.4 |
| 1969 | 18.0 | 1.8 | 2.9 | 2.0 | 2.0 | 3.5 | 9.2 | 19.7 | 37.2 | 63.4 | 103.6 | 155.0 |
| 1968 | 17.6 | 2.0 | 2.9 | 2.2 | 2.0 | 3.3 | 8.3 | 20.2 | 38.0 | 62.0 | 98.8 | 159.7 |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| MALIGNANT NEOPLASMS OF ALL OTHER AND UNSPECIFIED SITES (170-173, 190-199) | | | | | | | | | | | | |
| ALL OTHER, BOTH SEXES | | | | | | | | | | | | |
| 1978 | 18.1 | * | 1.8 | 1.6 | 2.1 | 3.1 | 9.2 | 29.2 | 63.0 | 103.8 | 131.1 | 157.1 |
| 1977 | 17.3 | * | 2.3 | 1.8 | 1.8 | 2.6 | 9.0 | 30.0 | 57.6 | 96.6 | 131.3 | 162.5 |
| 1976 | 17.1 | * | 1.9 | 2.0 | 2.2 | 2.6 | 8.9 | 28.0 | 61.3 | 97.2 | 126.5 | 141.2 |
| 1975 | 16.9 | * | 2.3 | 1.9 | 2.3 | 2.9 | 9.0 | 29.3 | 60.4 | 98.0 | 116.6 | 129.5 |
| 1974 | 16.7 | * | 2.0 | 1.9 | 2.2 | 2.6 | 8.6 | 28.9 | 61.3 | 98.2 | 114.8 | 137.8 |
| 1973 | 16.7 | * | 2.4 | 2.2 | 2.2 | 3.3 | 10.1 | 27.9 | 59.9 | 97.6 | 112.0 | 153.5 |
| 1972 | 16.2 | * | 2.2 | 1.8 | 2.0 | 3.4 | 10.2 | 27.9 | 57.1 | 95.1 | 109.4 | 146.3 |
| 1971 | 16.2 | * | 2.2 | 1.9 | 2.4 | 3.2 | 9.2 | 29.0 | 59.8 | 91.0 | 111.7 | 120.8 |
| 1970 | 15.8 | * | 2.8 | 1.9 | 2.3 | 2.9 | 10.3 | 28.0 | 56.8 | 90.5 | 105.4 | 114.1 |
| 1969 | 15.5 | * | 2.7 | 1.8 | 2.1 | 3.3 | 10.9 | 27.5 | 56.7 | 87.4 | 102.5 | 119.3 |
| 1968 | 15.5 | * | 2.8 | 1.9 | 2.4 | 3.6 | 10.6 | 27.1 | 56.9 | 86.1 | 103.8 | 110.8 |
| ALL OTHER, MALE | | | | | | | | | | | | |
| 1978 | 20.4 | * | 2.0 | 1.8 | 2.8 | 3.5 | 9.9 | 36.0 | 80.3 | 128.1 | 150.0 | 203.4 |
| 1977 | 19.2 | * | 1.9 | 2.0 | 1.9 | 3.4 | 10.1 | 37.1 | 72.0 | 115.7 | 156.7 | 198.2 |
| 1976 | 19.4 | * | 1.8 | 2.1 | 2.8 | 2.6 | 11.0 | 32.7 | 79.0 | 121.9 | 153.7 | 140.0 |
| 1975 | 18.7 | * | 2.3 | 1.8 | 2.8 | 3.5 | 9.6 | 36.5 | 73.8 | 115.1 | 139.7 | 129.6 |
| 1974 | 18.5 | * | 2.3 | 2.0 | 2.5 | 2.9 | 9.7 | 35.0 | 73.1 | 117.0 | 133.6 | 166.0 |
| 1973 | 18.7 | * | 2.8 | 1.8 | 2.0 | 3.8 | 11.1 | 33.5 | 71.9 | 120.7 | 142.9 | 175.0 |
| 1972 | 17.8 | * | 2.7 | 1.8 | 2.9 | 4.0 | 12.2 | 31.2 | 65.7 | 114.1 | 136.4 | 119.1 |
| 1971 | 17.8 | * | * | 2.2 | 2.8 | 3.7 | 10.5 | 33.1 | 69.9 | 107.8 | 134.3 | 119.1 |
| 1970 | 17.5 | * | 3.6 | 1.9 | 2.8 | 3.5 | 10.8 | 34.0 | 68.4 | 102.8 | 123.3 | 99.2 |
| 1969 | 16.5 | * | 3.0 | 1.3 | 2.5 | 3.7 | 12.4 | 28.6 | 68.0 | 95.9 | 116.0 | 125.0 |
| 1968 | 17.1 | * | 3.5 | 2.0 | 2.6 | 4.0 | 11.6 | 29.4 | 65.9 | 103.0 | 125.8 | 119.5 |
| ALL OTHER, FEMALE | | | | | | | | | | | | |
| 1978 | 16.0 | * | * | 1.4 | 1.5 | 2.8 | 8.6 | 23.7 | 48.7 | 85.0 | 118.5 | 133.9 |
| 1977 | 15.5 | * | 2.6 | 1.6 | 1.7 | 2.0 | 8.0 | 24.2 | 45.6 | 81.6 | 113.6 | 143.3 |
| 1976 | 15.1 | * | 2.0 | 1.9 | 1.6 | 2.6 | 7.1 | 24.1 | 46.6 | 77.6 | 108.0 | 141.8 |
| 1975 | 15.3 | * | 2.2 | 2.1 | 1.8 | 2.5 | 8.4 | 23.3 | 49.2 | 84.4 | 100.6 | 128.0 |
| 1974 | 15.0 | * | * | 1.7 | 1.9 | 2.3 | 7.7 | 23.6 | 51.2 | 83.1 | 101.8 | 121.2 |
| 1973 | 15.0 | * | 2.1 | 2.6 | 2.4 | 2.8 | 9.3 | 23.2 | 49.7 | 79.0 | 90.6 | 140.5 |
| 1972 | 14.6 | * | * | 1.7 | 1.3 | 2.9 | 8.5 | 25.1 | 49.7 | 79.8 | 90.4 | 163.2 |
| 1971 | 14.7 | * | 2.7 | 1.7 | 2.1 | 2.8 | 8.2 | 25.4 | 51.0 | 77.3 | 95.0 | 121.9 |
| 1970 | 14.2 | * | 1.9 | 1.8 | 1.8 | 2.4 | 9.9 | 22.8 | 46.5 | 80.4 | 91.8 | 124.6 |
| 1969 | 14.6 | * | 2.4 | 2.3 | 1.8 | 3.0 | 9.5 | 26.6 | 46.7 | 80.4 | 91.9 | 115.4 |
| 1968 | 14.0 | * | 2.2 | 1.8 | 2.3 | 3.3 | 9.6 | 25.2 | 48.9 | 72.0 | 87.0 | 104.9 |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| MALIGNANT NEOPLASMS OF ALL OTHER AND UNSPECIFIED SITES (170-173, 190-199) | | | | | | | | | | | | |
| BLACK, BOTH SEXES | | | | | | | | | | | | |
| 1978 | 19.6 | * | 2.1 | 1.6 | 2.3 | 3.4 | 10.3 | 31.5 | 68.3 | 109.3 | 131.5 | 158.0 |
| 1977 | 18.7 | * | 2.4 | 1.8 | 2.0 | 2.8 | 9.6 | 33.1 | 62.1 | 101.4 | 132.4 | 163.6 |
| 1976 | 18.4 | * | 1.9 | 2.0 | 2.4 | 2.5 | 10.0 | 30.2 | 65.1 | 103.3 | 126.9 | 138.2 |
| 1975 | 18.1 | * | 2.3 | 2.0 | 2.4 | 3.2 | 9.2 | 32.2 | 63.7 | 103.2 | 117.6 | 130.0 |
| 1974 | 17.7 | * | 1.7 | 2.0 | 2.3 | 2.7 | 9.5 | 31.3 | 64.8 | 102.6 | 111.4 | 143.8 |
| 1973 | 17.8 | * | 2.6 | 2.2 | 2.3 | 3.7 | 11.1 | 29.7 | 63.0 | 101.8 | 112.8 | 153.5 |
| 1972 | 17.0 | * | 2.3 | 1.8 | 2.0 | 3.5 | 11.0 | 29.9 | 59.4 | 98.5 | 109.2 | 152.7 |
| 1971 | 16.9 | * | 2.2 | 2.0 | 2.5 | 3.3 | 9.6 | 30.8 | 62.4 | 95.6 | 110.3 | 122.4 |
| 1970 | 16.6 | * | 2.9 | 2.0 | 2.2 | 3.1 | 11.0 | 29.7 | 59.7 | 94.5 | 106.1 | 124.4 |
| 1969 | 16.3 | * | 2.9 | 1.8 | 2.2 | 3.4 | 11.8 | 29.2 | 58.4 | 91.1 | 106.1 | 118.6 |
| 1968 | 16.2 | * | 3.0 | 2.0 | 2.5 | 3.7 | 11.4 | 28.8 | 59.0 | 89.3 | 105.7 | 107.6 |
| BLACK, MALE | | | | | | | | | | | | |
| 1978 | 22.1 | * | 2.2 | 1.8 | 2.9 | 3.8 | 11.0 | 39.4 | 87.9 | 138.5 | 151.5 | 198.0 |
| 1977 | 20.9 | * | * | 2.0 | 2.1 | 3.7 | 11.3 | 41.3 | 78.0 | 122.4 | 160.9 | 202.0 |
| 1976 | 21.0 | * | * | 2.1 | 3.0 | 2.5 | 12.7 | 35.8 | 84.6 | 133.4 | 152.6 | 134.0 |
| 1975 | 20.1 | * | 2.5 | 1.9 | 3.0 | 3.9 | 10.5 | 40.3 | 78.3 | 123.3 | 141.4 | 128.3 |
| 1974 | 19.6 | * | 2.1 | 2.0 | 2.6 | 3.0 | 11.0 | 38.2 | 78.0 | 124.0 | 129.2 | 174.4 |
| 1973 | 19.8 | * | 3.0 | 1.8 | 2.3 | 4.4 | 12.3 | 35.3 | 76.6 | 127.1 | 146.1 | 170.7 |
| 1972 | 18.6 | * | 2.8 | 1.9 | 3.0 | 3.9 | 13.0 | 34.0 | 69.8 | 119.0 | 134.0 | 120.0 |
| 1971 | 18.6 | * | * | 2.2 | 2.9 | 3.6 | 11.0 | 35.4 | 73.8 | 114.6 | 134.9 | 125.0 |
| 1970 | 18.4 | * | 3.9 | 2.0 | 2.7 | 3.8 | 11.4 | 36.6 | 72.2 | 107.2 | 126.3 | 110.9 |
| 1969 | 17.3 | * | 3.2 | 1.3 | 2.6 | 3.8 | 13.4 | 30.2 | 71.1 | 100.7 | 119.2 | 137.8 |
| 1968 | 17.8 | * | 3.6 | 2.1 | 2.6 | 4.1 | 12.6 | 31.3 | 68.6 | 107.7 | 127.9 | 108.6 |
| BLACK, FEMALE | | | | | | | | | | | | |
| 1978 | 17.3 | * | * | 1.4 | 1.7 | 3.1 | 9.8 | 25.0 | 52.4 | 87.5 | 118.2 | 137.4 |
| 1977 | 16.7 | * | 2.8 | 1.6 | 1.8 | 2.0 | 8.1 | 26.4 | 49.1 | 85.7 | 113.8 | 143.6 |
| 1976 | 16.1 | * | * | 1.9 | 1.8 | 2.6 | 7.9 | 25.7 | 48.9 | 80.5 | 109.7 | 140.4 |
| 1975 | 16.3 | * | 2.2 | 2.1 | 1.9 | 2.6 | 8.1 | 25.4 | 51.5 | 87.9 | 101.5 | 131.0 |
| 1974 | 16.0 | * | * | 1.9 | 1.9 | 2.5 | 8.2 | 25.6 | 53.9 | 86.3 | 99.4 | 126.9 |
| 1973 | 15.9 | * | 2.1 | 2.6 | 2.4 | 3.2 | 10.2 | 25.0 | 51.6 | 82.3 | 90.2 | 143.8 |
| 1972 | 15.5 | * | * | 1.7 | 1.2 | 3.2 | 9.3 | 26.3 | 50.6 | 82.7 | 91.4 | 171.4 |
| 1971 | 15.4 | * | 2.7 | 1.7 | 2.2 | 3.0 | 8.3 | 26.8 | 52.7 | 80.4 | 92.7 | 120.9 |
| 1970 | 15.1 | * | * | 1.9 | 1.7 | 2.5 | 10.7 | 23.8 | 49.0 | 84.4 | 91.3 | 133.1 |
| 1969 | 15.4 | * | 2.6 | 2.3 | 1.9 | 3.2 | 10.4 | 28.4 | 47.7 | 83.6 | 96.0 | 106.7 |
| 1968 | 14.8 | * | 2.4 | 1.9 | 2.4 | 3.4 | 10.4 | 26.6 | 50.7 | 74.2 | 89.2 | 108.9 |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| LEUKEMIA (204-207) | | | | | | | | | | | | |
| BOTH SEXES | | | | | | | | | | | | |
| 1978 | 6.9 | 1.1 | 1.8 | 1.9 | 1.6 | 1.8 | 2.8 | 5.2 | 10.7 | 24.6 | 50.2 | 73.2 |
| 1977 | 7.0 | 1.0 | 1.8 | 2.2 | 1.7 | 1.7 | 2.7 | 5.2 | 11.3 | 25.1 | 50.2 | 72.7 |
| 1976 | 6.9 | 1.0 | 2.0 | 2.3 | 1.7 | 1.7 | 2.7 | 5.2 | 11.5 | 25.1 | 49.5 | 71.3 |
| 1975 | 6.8 | 1.3 | 2.3 | 2.1 | 1.7 | 1.9 | 2.8 | 5.1 | 11.4 | 24.2 | 50.6 | 72.0 |
| 1974 | 6.8 | 1.3 | 2.1 | 2.4 | 1.7 | 1.8 | 2.7 | 5.1 | 11.4 | 25.0 | 51.5 | 66.8 |
| 1973 | 6.9 | 1.3 | 2.4 | 2.4 | 1.9 | 2.0 | 2.9 | 5.1 | 11.5 | 25.1 | 48.9 | 70.6 |
| 1972 | 6.9 | 1.7 | 2.4 | 2.4 | 1.9 | 1.8 | 2.9 | 4.8 | 11.5 | 26.6 | 51.1 | 66.1 |
| 1971 | 7.0 | 1.6 | 3.4 | 2.7 | 1.8 | 2.0 | 2.8 | 5.2 | 11.4 | 26.2 | 51.0 | 70.1 |
| 1970 | 7.1 | 2.0 | 3.6 | 2.7 | 2.0 | 1.9 | 3.1 | 5.7 | 12.0 | 26.3 | 51.0 | 62.5 |
| 1969 | 7.2 | 1.7 | 3.6 | 2.9 | 1.8 | 1.9 | 3.2 | 5.3 | 11.7 | 27.1 | 52.6 | 73.1 |
| 1968 | 7.2 | 1.6 | 3.8 | 2.9 | 2.1 | 2.0 | 3.3 | 5.6 | 12.2 | 27.4 | 49.2 | 71.6 |
| MALE | | | | | | | | | | | | |
| 1978 | 8.0 | * | 2.0 | 2.2 | 2.1 | 2.2 | 3.1 | 6.2 | 13.6 | 34.1 | 70.6 | 111.3 |
| 1977 | 8.1 | 1.2 | 2.0 | 2.6 | 2.0 | 2.0 | 3.1 | 6.2 | 14.7 | 33.9 | 71.2 | 104.9 |
| 1976 | 8.1 | * | 2.1 | 2.7 | 2.1 | 1.9 | 2.9 | 6.5 | 15.3 | 34.6 | 68.2 | 107.3 |
| 1975 | 8.0 | * | 2.5 | 2.4 | 2.0 | 2.1 | 3.1 | 5.9 | 14.8 | 33.2 | 72.4 | 110.9 |
| 1974 | 7.9 | 1.3 | 2.4 | 2.7 | 2.0 | 2.1 | 2.9 | 6.0 | 14.7 | 33.9 | 72.5 | 98.8 |
| 1973 | 8.0 | 1.3 | 2.9 | 2.8 | 2.4 | 2.2 | 3.5 | 6.0 | 14.5 | 33.7 | 70.3 | 100.6 |
| 1972 | 8.0 | 1.5 | 2.5 | 2.9 | 2.2 | 2.0 | 3.5 | 5.8 | 14.7 | 36.8 | 69.5 | 93.5 |
| 1971 | 8.1 | 1.6 | 3.7 | 3.1 | 2.2 | 2.4 | 3.1 | 6.0 | 14.7 | 34.9 | 70.5 | 101.0 |
| 1970 | 8.2 | 1.6 | 3.9 | 3.0 | 2.4 | 2.2 | 3.4 | 6.6 | 15.4 | 35.3 | 68.3 | 83.3 |
| 1969 | 8.4 | 1.5 | 3.9 | 3.3 | 2.4 | 2.3 | 3.7 | 6.4 | 15.0 | 36.1 | 70.6 | 101.3 |
| 1968 | 8.4 | 1.6 | 4.1 | 3.2 | 2.6 | 2.4 | 3.5 | 6.5 | 15.6 | 37.1 | 65.6 | 100.2 |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| LEUKEMIA (204-207) | | | | | | | | | | | | |
| FEMALE | | | | | | | | | | | | |
| 1978 | 5.9 | 1.4 | 1.6 | 1.6 | 1.2 | 1.4 | 2.5 | 4.3 | 8.1 | 17.4 | 38.1 | 55.9 |
| 1977 | 6.0 | * | 1.5 | 1.7 | 1.4 | 1.5 | 2.4 | 4.2 | 8.4 | 18.3 | 37.6 | 57.9 |
| 1976 | 5.8 | * | 1.8 | 1.8 | 1.4 | 1.5 | 2.5 | 3.9 | 8.1 | 17.9 | 38.2 | 54.4 |
| 1975 | 5.8 | 1.5 | 2.1 | 1.9 | 1.4 | 1.6 | 2.4 | 4.4 | 8.5 | 17.2 | 37.3 | 53.2 |
| 1974 | 5.8 | * | 1.8 | 2.0 | 1.3 | 1.6 | 2.6 | 4.3 | 8.5 | 18.2 | 38.5 | 50.9 |
| 1973 | 5.7 | 1.3 | 1.9 | 2.0 | 1.4 | 1.7 | 2.5 | 4.3 | 8.8 | 18.4 | 35.6 | 55.5 |
| 1972 | 5.8 | 1.9 | 2.3 | 2.0 | 1.5 | 1.6 | 2.4 | 4.0 | 8.7 | 18.7 | 39.4 | 52.2 |
| 1971 | 5.9 | 1.7 | 3.0 | 2.4 | 1.4 | 1.6 | 2.5 | 4.4 | 8.4 | 19.5 | 38.3 | 54.0 |
| 1970 | 6.1 | 2.3 | 3.2 | 2.4 | 1.7 | 1.6 | 2.8 | 4.9 | 9.0 | 19.3 | 39.6 | 50.9 |
| 1969 | 6.0 | 1.9 | 3.2 | 2.5 | 1.3 | 1.5 | 2.7 | 4.3 | 8.7 | 20.0 | 40.6 | 57.8 |
| 1968 | 6.1 | 1.5 | 3.5 | 2.6 | 1.7 | 1.6 | 3.1 | 4.9 | 9.1 | 19.7 | 38.0 | 55.9 |
| WHITE, BOTH SEXES | | | | | | | | | | | | |
| 1978 | 7.3 | 1.0 | 1.9 | 2.0 | 1.7 | 1.8 | 2.7 | 5.2 | 10.7 | 25.3 | 52.0 | 76.1 |
| 1977 | 7.4 | 1.1 | 1.9 | 2.3 | 1.7 | 1.8 | 2.7 | 5.1 | 11.4 | 25.5 | 51.7 | 75.5 |
| 1976 | 7.3 | 0.9 | 2.0 | 2.4 | 1.8 | 1.7 | 2.7 | 5.1 | 11.4 | 25.5 | 51.4 | 73.9 |
| 1975 | 7.2 | 1.4 | 2.4 | 2.2 | 1.8 | 1.8 | 2.8 | 5.1 | 11.5 | 24.6 | 52.4 | 74.9 |
| 1974 | 7.2 | 1.3 | 2.3 | 2.5 | 1.7 | 1.8 | 2.8 | 5.2 | 11.3 | 25.5 | 53.4 | 69.7 |
| 1973 | 7.2 | 1.2 | 2.5 | 2.6 | 1.9 | 2.0 | 3.0 | 5.2 | 11.4 | 25.3 | 50.5 | 73.3 |
| 1972 | 7.3 | 1.9 | 2.5 | 2.6 | 1.9 | 1.8 | 3.0 | 4.8 | 11.8 | 26.9 | 53.1 | 69.5 |
| 1971 | 7.4 | 1.8 | 3.5 | 2.9 | 1.8 | 2.0 | 2.9 | 5.1 | 11.5 | 26.9 | 52.9 | 72.5 |
| 1970 | 7.5 | 2.1 | 3.8 | 2.9 | 2.1 | 1.9 | 3.1 | 5.7 | 12.1 | 26.7 | 52.8 | 65.9 |
| 1969 | 7.6 | 1.8 | 3.9 | 3.1 | 1.9 | 1.9 | 3.3 | 5.3 | 11.7 | 27.8 | 54.6 | 77.1 |
| 1968 | 7.6 | 1.6 | 4.1 | 3.1 | 2.1 | 1.9 | 3.3 | 5.7 | 12.1 | 28.1 | 50.7 | 75.9 |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| LEUKEMIA (204-207) | | | | | | | | | | | | |
| WHITE, MALE | | | | | | | | | | | | |
| 1978 | 8.4 | * | 2.1 | 2.3 | 2.1 | 2.2 | 3.0 | 6.2 | 13.7 | 34.8 | 72.9 | 116.2 |
| 1977 | 8.5 | * | 2.1 | 2.7 | 2.1 | 2.0 | 3.0 | 6.2 | 14.8 | 34.8 | 73.6 | 108.8 |
| 1976 | 8.5 | * | 2.1 | 2.9 | 2.1 | 2.0 | 2.9 | 6.4 | 15.3 | 35.4 | 71.2 | 112.5 |
| 1975 | 8.4 | * | 2.7 | 2.5 | 2.1 | 2.1 | 3.1 | 5.9 | 14.8 | 33.9 | 75.6 | 116.3 |
| 1974 | 8.3 | * | 2.5 | 2.8 | 2.0 | 2.1 | 2.9 | 6.1 | 14.6 | 34.8 | 75.0 | 105.0 |
| 1973 | 8.4 | * | 3.0 | 3.0 | 2.4 | 2.2 | 3.6 | 6.1 | 14.5 | 34.2 | 72.5 | 104.9 |
| 1972 | 8.5 | 1.7 | 2.6 | 3.0 | 2.3 | 1.9 | 3.5 | 5.6 | 15.1 | 37.7 | 72.3 | 99.4 |
| 1971 | 8.6 | 1.7 | 4.0 | 3.2 | 2.2 | 2.4 | 3.1 | 6.0 | 14.8 | 36.0 | 73.6 | 105.4 |
| 1970 | 8.6 | 1.7 | 4.2 | 3.2 | 2.4 | 2.2 | 3.4 | 6.6 | 15.5 | 36.0 | 70.7 | 87.7 |
| 1969 | 8.9 | 1.5 | 4.4 | 3.4 | 2.5 | 2.3 | 3.8 | 6.4 | 15.1 | 37.1 | 73.3 | 108.2 |
| 1968 | 8.8 | 1.7 | 4.3 | 3.4 | 2.6 | 2.4 | 3.5 | 6.5 | 15.6 | 38.3 | 67.7 | 107.8 |
| WHITE, FEMALE | | | | | | | | | | | | |
| 1978 | 6.3 | * | 1.7 | 1.7 | 1.3 | 1.3 | 2.5 | 4.3 | 8.0 | 18.0 | 39.7 | 58.2 |
| 1977 | 6.3 | * | 1.6 | 1.8 | 1.3 | 1.5 | 2.4 | 4.2 | 8.4 | 18.5 | 38.7 | 60.4 |
| 1976 | 6.1 | * | 1.9 | 1.9 | 1.5 | 1.5 | 2.4 | 3.8 | 8.0 | 17.9 | 39.6 | 56.1 |
| 1975 | 6.1 | 1.6 | 2.2 | 1.9 | 1.4 | 1.6 | 2.4 | 4.4 | 8.5 | 17.4 | 38.4 | 55.3 |
| 1974 | 6.1 | * | 2.0 | 2.2 | 1.4 | 1.6 | 2.6 | 4.4 | 8.4 | 18.4 | 40.2 | 52.5 |
| 1973 | 6.0 | * | 1.9 | 2.1 | 1.5 | 1.8 | 2.4 | 4.3 | 8.7 | 18.4 | 36.9 | 57.7 |
| 1972 | 6.1 | 2.1 | 2.5 | 2.1 | 1.5 | 1.6 | 2.4 | 4.1 | 8.8 | 18.7 | 41.0 | 54.4 |
| 1971 | 6.2 | 2.0 | 3.1 | 2.5 | 1.5 | 1.6 | 2.6 | 4.4 | 8.6 | 19.9 | 39.7 | 55.6 |
| 1970 | 6.4 | 2.5 | 3.5 | 2.6 | 1.8 | 1.6 | 2.8 | 4.9 | 9.0 | 19.4 | 41.1 | 53.9 |
| 1969 | 6.3 | 2.1 | 3.4 | 2.7 | 1.3 | 1.5 | 2.8 | 4.2 | 8.7 | 20.5 | 42.1 | 60.8 |
| 1968 | 6.4 | 1.4 | 3.8 | 2.8 | 1.7 | 1.5 | 3.0 | 4.9 | 9.0 | 20.2 | 39.1 | 58.7 |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| LEUKEMIA (204-207) | | | | | | | | | | | | |
| ALL OTHER, BOTH SEXES | | | | | | | | | | | | |
| 1978 | 4.4 | * | 1.3 | 1.7 | 1.3 | 1.9 | 3.3 | 5.2 | 10.9 | 18.7 | 31.7 | 39.9 |
| 1977 | 4.5 | * | 1.4 | 1.6 | 1.7 | 1.6 | 3.0 | 5.3 | 10.7 | 20.9 | 34.7 | 41.3 |
| 1976 | 4.5 | * | 1.6 | 1.6 | 1.4 | 1.6 | 2.8 | 5.6 | 11.8 | 21.4 | 29.9 | 41.2 |
| 1975 | 4.3 | * | 1.8 | 1.6 | 1.2 | 1.9 | 2.8 | 5.1 | 11.1 | 20.4 | 31.8 | 39.0 |
| 1974 | 4.3 | * | 1.3 | 1.7 | 1.4 | 1.8 | 2.7 | 4.6 | 12.4 | 20.2 | 30.1 | 32.6 |
| 1973 | 4.6 | * | 2.0 | 1.7 | 1.6 | 2.1 | 2.5 | 4.8 | 12.0 | 23.2 | 31.5 | 38.6 |
| 1972 | 4.2 | * | 1.6 | 1.6 | 1.5 | 1.9 | 2.8 | 5.0 | 9.3 | 22.9 | 28.5 | 27.6 |
| 1971 | 4.3 | * | 2.4 | 2.0 | 1.6 | 2.1 | 2.4 | 5.5 | 9.9 | 19.8 | 27.5 | 43.3 |
| 1970 | 4.6 | * | 2.0 | 1.6 | 1.9 | 1.9 | 3.1 | 5.9 | 11.7 | 23.0 | 27.5 | 28.3 |
| 1969 | 4.4 | * | 2.0 | 2.0 | 1.6 | 1.9 | 2.9 | 5.5 | 11.5 | 20.2 | 27.9 | 27.5 |
| 1968 | 4.6 | * | 2.1 | 1.9 | 1.9 | 2.2 | 3.4 | 5.4 | 13.4 | 19.6 | 30.5 | 23.5 |
| ALL OTHER, MALE | | | | | | | | | | | | |
| 1978 | 5.4 | * | * | 1.9 | 2.0 | 2.0 | 3.8 | 6.2 | 13.0 | 26.8 | 48.5 | 62.1 |
| 1977 | 5.1 | * | * | 1.8 | 1.6 | 1.5 | 3.1 | 6.1 | 13.6 | 26.0 | 48.3 | 66.1 |
| 1976 | 5.2 | * | 1.9 | 1.8 | 1.8 | 1.5 | 2.4 | 6.8 | 15.8 | 26.8 | 40.1 | 54.5 |
| 1975 | 5.1 | * | * | 1.6 | 1.6 | 2.4 | 3.6 | 5.9 | 14.2 | 27.1 | 41.3 | 57.4 |
| 1974 | 5.1 | * | * | 2.0 | 1.9 | 2.0 | 3.0 | 5.7 | 15.3 | 24.9 | 46.7 | * |
| 1973 | 5.6 | * | 2.3 | 2.2 | 2.4 | 2.7 | 2.5 | 5.7 | 15.0 | 29.0 | 47.3 | 56.3 |
| 1972 | 5.0 | * | 2.0 | 2.1 | 1.8 | 2.3 | 3.6 | 7.0 | 10.6 | 27.9 | 39.6 | * |
| 1971 | 5.2 | * | 2.1 | 2.3 | 2.2 | 2.7 | 3.1 | 6.1 | 12.9 | 25.0 | 36.7 | 57.4 |
| 1970 | 5.5 | * | 2.3 | 1.9 | 2.2 | 2.6 | 3.4 | 6.3 | 15.0 | 28.6 | 39.4 | 45.1 |
| 1969 | 5.2 | * | * | 2.5 | 2.0 | 2.2 | 3.2 | 5.9 | 14.4 | 26.8 | 38.3 | * |
| 1968 | 5.5 | * | 2.6 | 2.5 | 2.2 | 2.7 | 3.4 | 6.5 | 16.3 | 25.1 | 40.9 | * |
| ALL OTHER, FEMALE | | | | | | | | | | | | |
| 1978 | 3.5 | * | * | 1.4 | 0.7 | 1.9 | 2.8 | 4.3 | 9.2 | 12.3 | 20.3 | 28.4 |
| 1977 | 4.0 | * | * | 1.3 | 1.7 | 1.6 | 2.9 | 4.6 | 8.3 | 17.0 | 25.3 | 27.9 |
| 1976 | 3.8 | * | * | 1.3 | 1.0 | 1.7 | 3.1 | 4.7 | 8.4 | 17.2 | 22.9 | 33.7 |
| 1975 | 3.6 | * | 1.8 | 1.5 | 1.0 | 1.4 | 2.1 | 4.4 | 8.5 | 15.1 | 25.1 | 28.0 |
| 1974 | 3.5 | * | * | 1.4 | 0.9 | 1.6 | 2.5 | 3.6 | 9.8 | 16.5 | 18.7 | 31.8 |
| 1973 | 3.7 | * | * | 1.3 | 0.9 | 1.6 | 2.5 | 4.1 | 9.5 | 18.5 | 20.4 | 27.8 |
| 1972 | 3.5 | * | * | 1.1 | 1.3 | 1.6 | 2.1 | 3.3 | 8.2 | 19.0 | 20.6 | * |
| 1971 | 3.6 | * | 2.7 | 1.8 | 1.1 | 1.6 | 1.9 | 4.9 | 7.3 | 15.6 | 20.7 | 34.2 |
| 1970 | 3.7 | * | * | 1.3 | 1.5 | 1.3 | 2.8 | 5.6 | 8.8 | 18.5 | 18.5 | * |
| 1969 | 3.6 | * | 2.3 | 1.4 | 1.3 | 1.7 | 2.7 | 5.2 | 8.9 | 14.8 | 19.8 | * |
| 1968 | 3.8 | * | * | 1.4 | 1.6 | 1.8 | 3.3 | 4.4 | 10.8 | 15.1 | 22.6 | * |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| LEUKEMIA (204-207) | | | | | | | | | | | | |
| BLACK, BOTH SEXES | | | | | | | | | | | | |
| 1978 | 4.7 | * | 1.2 | 1.6 | 1.4 | 2.0 | 3.2 | 5.4 | 11.7 | 20.3 | 32.2 | 42.7 |
| 1977 | 4.7 | * | 1.5 | 1.5 | 1.7 | 1.7 | 3.1 | 5.6 | 11.3 | 21.6 | 34.4 | 41.3 |
| 1976 | 4.7 | * | 1.7 | 1.6 | 1.3 | 1.8 | 2.7 | 5.9 | 12.2 | 22.8 | 31.0 | 41.2 |
| 1975 | 4.6 | * | 1.9 | 1.6 | 1.3 | 2.0 | 3.0 | 5.5 | 11.5 | 21.1 | 32.0 | 36.9 |
| 1974 | 4.5 | * | 1.3 | 1.7 | 1.3 | 1.9 | 2.8 | 4.8 | 13.0 | 20.8 | 31.1 | 33.1 |
| 1973 | 4.8 | * | 1.7 | 1.9 | 1.6 | 2.1 | 2.4 | 5.1 | 12.5 | 24.0 | 31.7 | 42.1 |
| 1972 | 4.3 | * | 1.8 | 1.6 | 1.7 | 1.9 | 2.9 | 5.1 | 9.6 | 22.8 | 29.4 | 27.3 |
| 1971 | 4.5 | * | 2.5 | 2.0 | 1.6 | 2.1 | 2.5 | 5.4 | 10.4 | 20.8 | 28.3 | 44.9 |
| 1970 | 4.8 | * | 2.1 | 1.4 | 1.9 | 2.1 | 3.3 | 6.3 | 12.4 | 24.4 | 27.6 | 30.7 |
| 1969 | 4.4 | * | 1.9 | 1.8 | 1.7 | 2.0 | 2.8 | 5.6 | 11.7 | 20.9 | 28.8 | 28.9 |
| 1968 | 4.7 | * | 2.0 | 1.9 | 1.9 | 2.2 | 3.2 | 5.5 | 13.9 | 20.5 | 31.8 | 22.8 |
| BLACK, MALE | | | | | | | | | | | | |
| 1978 | 5.7 | * | * | 1.9 | 2.1 | 2.1 | 3.8 | 6.5 | 13.9 | 29.4 | 49.4 | 68.6 |
| 1977 | 5.3 | * | * | 1.7 | 1.5 | 1.8 | 3.0 | 6.3 | 14.5 | 26.9 | 47.7 | 69.4 |
| 1976 | 5.5 | * | * | 1.8 | 1.7 | 1.6 | 2.5 | 7.3 | 16.9 | 28.6 | 41.5 | 53.2 |
| 1975 | 5.4 | * | * | 1.6 | 1.6 | 2.5 | 4.0 | 6.2 | 15.0 | 28.5 | 41.4 | 58.7 |
| 1974 | 5.3 | * | * | 1.9 | 1.9 | 2.1 | 3.2 | 5.8 | 16.3 | 26.1 | 48.3 | * |
| 1973 | 5.8 | * | 2.0 | 2.3 | 2.4 | 2.6 | 2.4 | 6.3 | 15.7 | 30.2 | 49.0 | 65.9 |
| 1972 | 5.2 | * | 2.2 | 2.1 | 1.9 | 2.4 | 3.8 | 7.3 | 11.1 | 28.1 | 40.6 | * |
| 1971 | 5.3 | * | 2.1 | 2.4 | 2.1 | 2.8 | 3.2 | 5.9 | 13.6 | 26.6 | 38.2 | 60.0 |
| 1970 | 5.7 | * | 2.4 | 1.5 | 2.3 | 2.9 | 3.8 | 7.2 | 16.1 | 30.4 | 38.5 | 50.0 |
| 1969 | 5.2 | * | * | 2.3 | 2.0 | 2.3 | 3.0 | 6.0 | 14.7 | 27.7 | 38.9 | * |
| 1968 | 5.6 | * | 2.5 | 2.3 | 2.3 | 2.6 | 3.3 | 6.6 | 17.3 | 26.5 | 41.8 | * |
| BLACK, FEMALE | | | | | | | | | | | | |
| 1978 | 3.7 | * | * | 1.4 | 0.7 | 1.9 | 2.7 | 4.5 | 9.8 | 13.5 | 20.7 | 29.3 |
| 1977 | 4.2 | * | * | 1.2 | 1.8 | 1.6 | 3.2 | 5.0 | 8.6 | 17.6 | 25.7 | 26.6 |
| 1976 | 4.0 | * | * | 1.3 | 0.8 | 1.9 | 2.9 | 4.6 | 8.4 | 18.4 | 24.1 | 34.8 |
| 1975 | 3.8 | * | * | 1.5 | 1.0 | 1.5 | 2.2 | 5.0 | 8.6 | 15.6 | 25.6 | 25.0 |
| 1974 | 3.7 | * | * | 1.4 | 0.8 | 1.7 | 2.5 | 4.0 | 10.1 | 16.9 | 19.4 | 33.3 |
| 1973 | 3.8 | * | * | 1.4 | 0.9 | 1.7 | 2.4 | 4.2 | 9.7 | 19.2 | 19.9 | 28.8 |
| 1972 | 3.6 | * | * | 1.2 | 1.5 | 1.5 | 2.1 | 3.2 | 8.4 | 18.6 | 21.4 | * |
| 1971 | 3.7 | * | 2.9 | 1.5 | 1.1 | 1.5 | 1.8 | 5.0 | 7.7 | 16.2 | 21.2 | 35.8 |
| 1970 | 3.9 | * | * | 1.3 | 1.6 | 1.4 | 2.9 | 5.6 | 9.2 | 19.6 | 19.6 | * |
| 1969 | 3.7 | * | 2.3 | 1.3 | 1.4 | 1.7 | 2.6 | 5.2 | 9.2 | 15.6 | 21.1 | * |
| 1968 | 3.9 | * | * | 1.4 | 1.6 | 1.9 | 3.1 | 4.6 | 11.0 | 15.6 | 24.3 | * |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|--|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| OTHER NEOPLASMS OF LYMPHATIC AND HEMATOPOIETIC TISSUES (200-203, 208, 209) | | | | | | | | | | | | |
| BOTH SEXES | | | | | | | | | | | | |
| 1978 | 9.8 | * | 0.2 | 0.4 | 1.1 | 2.0 | 3.4 | 8.6 | 21.2 | 42.7 | 69.1 | 72.6 |
| 1977 | 9.6 | * | 0.2 | 0.5 | 1.1 | 2.1 | 3.3 | 8.8 | 21.6 | 42.0 | 67.0 | 68.8 |
| 1976 | 9.4 | * | 0.2 | 0.4 | 1.1 | 2.0 | 3.4 | 9.1 | 21.2 | 42.5 | 65.4 | 63.9 |
| 1975 | 9.3 | * | 0.3 | 0.5 | 1.3 | 2.2 | 3.6 | 9.2 | 21.5 | 40.6 | 63.4 | 62.4 |
| 1974 | 9.3 | * | 0.3 | 0.6 | 1.2 | 2.2 | 3.9 | 9.5 | 21.6 | 42.7 | 63.4 | 58.0 |
| 1973 | 9.3 | * | 0.5 | 0.6 | 1.4 | 2.4 | 4.2 | 9.3 | 21.4 | 42.5 | 62.0 | 59.5 |
| 1972 | 9.4 | * | 0.3 | 0.5 | 1.6 | 2.7 | 4.6 | 9.7 | 22.1 | 42.9 | 60.9 | 58.8 |
| 1971 | 9.2 | * | 0.4 | 0.6 | 1.6 | 2.8 | 4.4 | 10.0 | 23.3 | 40.5 | 59.4 | 53.4 |
| 1970 | 9.2 | * | 0.4 | 0.7 | 1.7 | 2.9 | 4.4 | 10.5 | 22.1 | 41.4 | 59.2 | 47.6 |
| 1969 | 9.1 | * | 0.4 | 0.7 | 1.8 | 2.8 | 4.8 | 10.2 | 21.9 | 40.4 | 57.2 | 51.3 |
| 1968 | 8.9 | * | 0.4 | 0.7 | 1.9 | 3.1 | 4.9 | 10.6 | 21.5 | 39.5 | 54.4 | 48.0 |
| MALE | | | | | | | | | | | | |
| 1978 | 10.6 | * | 0.4 | 0.6 | 1.4 | 2.5 | 4.5 | 10.4 | 25.5 | 52.0 | 86.4 | 94.5 |
| 1977 | 10.4 | * | * | 0.8 | 1.4 | 2.5 | 4.3 | 10.3 | 25.8 | 51.3 | 81.7 | 88.7 |
| 1976 | 10.2 | * | 0.4 | 0.6 | 1.3 | 2.5 | 4.2 | 10.9 | 25.1 | 52.3 | 80.3 | 83.0 |
| 1975 | 10.3 | * | 0.4 | 0.7 | 1.6 | 2.9 | 4.5 | 11.1 | 26.5 | 49.6 | 78.5 | 83.2 |
| 1974 | 10.4 | * | 0.5 | 0.9 | 1.5 | 2.7 | 4.9 | 12.0 | 26.4 | 52.4 | 79.1 | 76.3 |
| 1973 | 10.3 | * | 0.6 | 0.8 | 1.7 | 3.1 | 5.3 | 11.1 | 26.1 | 52.0 | 78.1 | 72.7 |
| 1972 | 10.5 | * | 0.3 | 0.6 | 2.0 | 3.4 | 5.9 | 11.8 | 27.7 | 49.8 | 74.4 | 84.3 |
| 1971 | 10.5 | * | 0.5 | 0.9 | 2.0 | 3.6 | 5.5 | 12.3 | 28.5 | 50.2 | 72.9 | 67.8 |
| 1970 | 10.5 | * | 0.4 | 0.9 | 2.2 | 3.6 | 5.6 | 12.8 | 27.1 | 50.3 | 74.0 | 58.1 |
| 1969 | 10.4 | * | 0.5 | 1.0 | 2.3 | 3.8 | 6.3 | 12.5 | 26.8 | 48.9 | 71.2 | 62.7 |
| 1968 | 10.2 | * | 0.6 | 0.9 | 2.4 | 4.1 | 6.2 | 12.7 | 26.2 | 48.3 | 65.2 | 64.1 |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|

OTHER NEOPLASMS OF LYMPHATIC AND HEMATOPOIETIC TISSUES (200-203, 208, 209)

FEMALE

| | | | | | | | | | | | | |
|------|-----|---|-----|-----|-----|-----|-----|-----|------|------|------|------|
| 1978 | 9.0 | * | * | 0.3 | 0.7 | 1.5 | 2.4 | 7.0 | 17.4 | 35.6 | 58.8 | 62.7 |
| 1977 | 8.9 | * | * | 0.2 | 0.7 | 1.6 | 2.4 | 7.5 | 17.9 | 34.8 | 58.2 | 59.7 |
| 1976 | 8.6 | * | * | 0.2 | 0.8 | 1.5 | 2.5 | 7.4 | 17.8 | 34.9 | 56.5 | 54.9 |
| 1975 | 8.3 | * | * | 0.3 | 0.9 | 1.6 | 2.8 | 7.4 | 17.1 | 33.7 | 54.1 | 52.4 |
| 1974 | 8.3 | * | * | 0.3 | 0.9 | 1.7 | 2.9 | 7.3 | 17.4 | 35.2 | 53.7 | 48.9 |
| 1973 | 8.3 | * | 0.3 | 0.3 | 1.1 | 1.7 | 3.1 | 7.6 | 17.2 | 35.3 | 52.0 | 52.9 |
| 1972 | 8.4 | * | 0.3 | 0.4 | 1.2 | 2.1 | 3.3 | 7.7 | 17.1 | 37.5 | 52.2 | 45.7 |
| 1971 | 8.1 | * | * | 0.3 | 1.2 | 1.9 | 3.4 | 7.9 | 18.7 | 33.1 | 50.7 | 45.9 |
| 1970 | 8.0 | * | * | 0.4 | 1.2 | 2.3 | 3.2 | 8.3 | 17.7 | 34.6 | 49.4 | 41.7 |
| 1969 | 7.8 | * | * | 0.4 | 1.3 | 1.9 | 3.3 | 8.0 | 17.6 | 33.8 | 47.8 | 45.0 |
| 1968 | 7.7 | * | * | 0.4 | 1.4 | 2.1 | 3.6 | 8.7 | 17.2 | 32.5 | 47.1 | 39.3 |

WHITE, BOTH SEXES

| | | | | | | | | | | | | |
|------|------|---|-----|-----|-----|-----|-----|------|------|------|------|------|
| 1978 | 10.2 | * | 0.2 | 0.4 | 1.0 | 2.1 | 3.4 | 8.7 | 21.1 | 42.7 | 70.6 | 73.5 |
| 1977 | 10.0 | * | 0.2 | 0.5 | 1.1 | 2.1 | 3.3 | 8.7 | 21.4 | 42.2 | 68.0 | 70.5 |
| 1976 | 9.8 | * | 0.3 | 0.4 | 1.1 | 2.0 | 3.3 | 9.0 | 20.9 | 42.6 | 66.9 | 65.8 |
| 1975 | 9.7 | * | 0.2 | 0.5 | 1.3 | 2.4 | 3.6 | 8.9 | 21.3 | 40.7 | 64.7 | 64.3 |
| 1974 | 9.8 | * | 0.3 | 0.6 | 1.3 | 2.3 | 3.9 | 9.5 | 21.6 | 42.8 | 64.9 | 58.2 |
| 1973 | 9.7 | * | 0.5 | 0.6 | 1.4 | 2.5 | 4.2 | 9.0 | 21.2 | 42.6 | 63.9 | 61.0 |
| 1972 | 9.8 | * | 0.3 | 0.6 | 1.7 | 2.8 | 4.5 | 9.5 | 22.1 | 43.7 | 62.5 | 60.6 |
| 1971 | 9.6 | * | 0.4 | 0.6 | 1.7 | 2.9 | 4.4 | 10.0 | 23.1 | 40.8 | 61.0 | 54.7 |
| 1970 | 9.6 | * | 0.3 | 0.7 | 1.8 | 3.1 | 4.5 | 10.3 | 22.0 | 42.1 | 60.5 | 49.3 |
| 1969 | 9.4 | * | 0.4 | 0.7 | 1.9 | 2.9 | 4.8 | 10.0 | 21.8 | 40.9 | 58.5 | 53.0 |
| 1968 | 9.3 | * | 0.5 | 0.7 | 1.9 | 3.2 | 5.0 | 10.6 | 21.4 | 39.9 | 55.8 | 49.1 |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|--|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| OTHER NEOPLASMS OF LYMPHATIC AND HEMATOPOIETIC TISSUES (200-203, 208, 209) | | | | | | | | | | | | |
| WHITE, MALE | | | | | | | | | | | | |
| 1978 | 11.0 | * | 0.4 | 0.5 | 1.4 | 2.5 | 4.4 | 10.5 | 25.5 | 51.6 | 88.3 | 95.1 |
| 1977 | 10.7 | * | * | 0.8 | 1.5 | 2.6 | 4.2 | 10.2 | 25.5 | 51.4 | 82.5 | 90.2 |
| 1976 | 10.6 | * | 0.4 | 0.7 | 1.3 | 2.5 | 4.1 | 10.8 | 24.7 | 52.3 | 82.0 | 86.6 |
| 1975 | 10.6 | * | * | 0.7 | 1.7 | 3.0 | 4.4 | 10.9 | 26.2 | 49.6 | 80.3 | 86.3 |
| 1974 | 10.8 | * | 0.4 | 0.9 | 1.6 | 2.8 | 4.9 | 11.8 | 26.2 | 52.5 | 80.6 | 76.9 |
| 1973 | 10.7 | * | 0.7 | 0.9 | 1.7 | 3.3 | 5.4 | 10.9 | 25.8 | 51.9 | 79.7 | 75.1 |
| 1972 | 10.9 | * | 0.4 | 0.7 | 2.2 | 3.5 | 5.7 | 11.6 | 27.7 | 50.8 | 76.4 | 88.0 |
| 1971 | 10.9 | * | 0.5 | 0.9 | 2.1 | 3.8 | 5.5 | 12.3 | 28.2 | 50.3 | 74.8 | 69.3 |
| 1970 | 10.8 | * | 0.4 | 1.0 | 2.3 | 3.7 | 5.7 | 12.6 | 26.8 | 50.7 | 75.0 | 59.6 |
| 1969 | 10.7 | * | 0.5 | 1.0 | 2.4 | 3.8 | 6.3 | 12.4 | 26.3 | 49.0 | 72.4 | 65.5 |
| 1968 | 10.6 | * | 0.7 | 1.0 | 2.4 | 4.2 | 6.4 | 12.6 | 26.2 | 48.8 | 66.9 | 64.7 |
| WHITE, FEMALE | | | | | | | | | | | | |
| 1978 | 9.5 | * | * | 0.3 | 0.7 | 1.6 | 2.4 | 7.0 | 17.1 | 35.9 | 60.2 | 63.8 |
| 1977 | 9.3 | * | * | 0.2 | 0.8 | 1.6 | 2.4 | 7.3 | 17.7 | 35.2 | 59.4 | 61.5 |
| 1976 | 9.1 | * | * | 0.2 | 0.8 | 1.5 | 2.5 | 7.4 | 17.5 | 35.3 | 58.0 | 56.2 |
| 1975 | 8.7 | * | * | 0.3 | 1.0 | 1.7 | 2.8 | 7.1 | 16.9 | 34.0 | 55.2 | 53.9 |
| 1974 | 8.8 | * | * | 0.3 | 0.9 | 1.8 | 2.9 | 7.3 | 17.5 | 35.4 | 55.3 | 49.1 |
| 1973 | 8.7 | * | * | 0.3 | 1.1 | 1.8 | 3.1 | 7.3 | 17.1 | 35.6 | 54.1 | 54.1 |
| 1972 | 8.8 | * | * | 0.4 | 1.2 | 2.1 | 3.4 | 7.5 | 17.1 | 38.3 | 53.7 | 46.8 |
| 1971 | 8.5 | * | * | 0.3 | 1.2 | 2.0 | 3.4 | 7.8 | 18.5 | 33.5 | 52.2 | 47.2 |
| 1970 | 8.5 | * | * | 0.4 | 1.3 | 2.5 | 3.4 | 8.1 | 17.7 | 35.5 | 51.0 | 43.7 |
| 1969 | 8.2 | * | * | 0.4 | 1.4 | 2.0 | 3.3 | 7.8 | 17.6 | 34.7 | 49.2 | 46.5 |
| 1968 | 8.1 | * | * | 0.4 | 1.4 | 2.1 | 3.6 | 8.6 | 17.1 | 33.0 | 48.3 | 40.7 |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|--|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| OTHER NEOPLASMS OF LYMPHATIC AND HEMATOPOIETIC TISSUES (200-203, 208, 209) | | | | | | | | | | | | |
| ALL OTHER, BOTH SEXES | | | | | | | | | | | | |
| 1978 | 6.8 | * | * | 0.5 | 1.1 | 1.5 | 3.5 | 8.2 | 22.2 | 42.7 | 53.5 | 62.5 |
| 1977 | 6.9 | * | * | 0.5 | 0.8 | 1.8 | 3.6 | 9.8 | 24.0 | 39.6 | 56.6 | 50.0 |
| 1976 | 6.7 | * | * | * | 0.9 | 1.6 | 3.7 | 9.6 | 24.3 | 40.9 | 49.9 | 41.8 |
| 1975 | 6.6 | * | * | 0.5 | 0.9 | 1.3 | 3.5 | 11.0 | 23.6 | 39.4 | 49.6 | 41.1 |
| 1974 | 6.5 | * | * | 0.4 | 1.1 | 1.5 | 3.8 | 10.2 | 21.8 | 41.7 | 46.9 | 54.8 |
| 1973 | 6.5 | * | * | 0.5 | 1.2 | 1.5 | 3.8 | 11.1 | 23.0 | 41.2 | 41.1 | 41.7 |
| 1972 | 6.2 | * | * | 0.4 | 1.1 | 2.0 | 4.8 | 11.3 | 21.9 | 34.4 | 42.0 | 37.4 |
| 1971 | 6.5 | * | * | 0.5 | 1.2 | 1.9 | 4.5 | 10.3 | 25.7 | 38.1 | 39.6 | 38.3 |
| 1970 | 6.2 | * | * | 0.5 | 1.2 | 1.9 | 3.8 | 11.8 | 23.2 | 34.4 | 43.0 | 29.8 |
| 1969 | 6.4 | * | * | 0.6 | 1.3 | 2.4 | 4.8 | 11.5 | 23.6 | 35.6 | 41.2 | 31.2 |
| 1968 | 6.1 | * | * | 0.6 | 1.4 | 2.2 | 3.9 | 11.4 | 21.9 | 34.8 | 36.4 | 36.3 |
| ALL OTHER, MALE | | | | | | | | | | | | |
| 1978 | 7.9 | * | * | 0.7 | 1.4 | 2.1 | 4.8 | 9.8 | 25.0 | 56.0 | 68.0 | 87.9 |
| 1977 | 7.9 | * | * | 0.6 | 1.0 | 2.2 | 4.9 | 10.7 | 28.9 | 51.0 | 73.9 | 73.2 |
| 1976 | 7.8 | * | * | * | 1.3 | 2.2 | 5.1 | 12.0 | 29.2 | 52.4 | 63.8 | 47.3 |
| 1975 | 7.7 | * | * | * | 1.4 | 1.6 | 4.7 | 12.4 | 29.1 | 50.3 | 61.2 | 51.9 |
| 1974 | 7.9 | * | * | * | 1.4 | 1.9 | 4.6 | 14.0 | 28.3 | 51.7 | 63.8 | 70.0 |
| 1973 | 7.8 | * | * | 0.7 | 1.7 | 2.2 | 4.9 | 12.5 | 28.9 | 52.3 | 61.2 | 47.9 |
| 1972 | 7.3 | * | * | * | 1.1 | 2.3 | 7.3 | 14.0 | 28.1 | 39.9 | 53.5 | 46.8 |
| 1971 | 7.7 | * | * | 0.7 | 1.4 | 2.6 | 5.6 | 12.3 | 31.5 | 49.3 | 51.2 | 53.2 |
| 1970 | 8.0 | * | * | 0.7 | 1.8 | 2.8 | 5.5 | 14.7 | 29.6 | 46.0 | 62.7 | 45.1 |
| 1969 | 8.1 | * | * | 0.9 | 1.7 | 3.3 | 6.4 | 13.4 | 30.9 | 48.3 | 56.9 | * |
| 1968 | 7.3 | * | * | 0.7 | 2.0 | 2.9 | 5.0 | 13.5 | 26.3 | 43.3 | 44.6 | 58.5 |
| ALL OTHER, FEMALE | | | | | | | | | | | | |
| 1978 | 5.8 | * | * | * | 0.8 | 1.0 | 2.5 | 6.8 | 19.9 | 32.5 | 43.7 | 49.5 |
| 1977 | 5.9 | * | * | * | 0.7 | 1.4 | 2.5 | 9.1 | 20.0 | 30.7 | 44.6 | 37.5 |
| 1976 | 5.6 | * | * | * | * | 1.1 | 2.6 | 7.6 | 20.2 | 31.8 | 40.4 | 38.8 |
| 1975 | 5.6 | * | * | * | * | 1.1 | 2.5 | 9.7 | 18.9 | 30.8 | 41.5 | 34.4 |
| 1974 | 5.2 | * | * | * | 0.9 | 1.0 | 3.0 | 7.0 | 16.4 | 33.7 | 35.2 | 45.9 |
| 1973 | 5.3 | * | * | * | 0.8 | * | 2.9 | 9.9 | 17.9 | 32.3 | 27.0 | 38.0 |
| 1972 | 5.2 | * | * | * | 1.1 | 1.8 | 2.6 | 9.1 | 16.5 | 30.0 | 33.9 | 31.6 |
| 1971 | 5.3 | * | * | * | 1.0 | 1.3 | 3.5 | 8.5 | 20.6 | 29.0 | 31.1 | 28.8 |
| 1970 | 4.6 | * | * | * | * | * | 2.4 | 9.3 | 17.6 | 24.8 | 28.0 | * |
| 1969 | 4.9 | * | * | * | 0.9 | 1.7 | 3.5 | 9.8 | 17.1 | 25.2 | 29.1 | * |
| 1968 | 5.0 | * | * | * | * | 1.6 | 3.1 | 9.5 | 18.1 | 27.8 | 30.1 | * |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|--|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| OTHER NEOPLASMS OF LYMPHATIC AND HEMATOPOIETIC TISSUES (200-203, 208, 209) | | | | | | | | | | | | |
| BLACK, BOTH SEXES | | | | | | | | | | | | |
| 1978 | 7.4 | * | * | 0.6 | 1.1 | 1.6 | 4.0 | 8.8 | 24.2 | 44.9 | 53.9 | 66.0 |
| 1977 | 7.3 | * | * | 0.5 | 0.9 | 1.9 | 3.9 | 10.7 | 25.8 | 40.0 | 56.8 | 47.6 |
| 1976 | 7.2 | * | * | * | 1.0 | 1.8 | 3.9 | 10.3 | 26.2 | 43.5 | 50.1 | 41.2 |
| 1975 | 7.0 | * | * | 0.5 | 1.0 | 1.4 | 3.8 | 12.0 | 25.1 | 40.6 | 48.9 | 41.5 |
| 1974 | 6.9 | * | * | 0.4 | 1.2 | 1.6 | 4.1 | 10.8 | 23.0 | 44.1 | 47.3 | 52.9 |
| 1973 | 6.8 | * | * | 0.5 | 1.4 | 1.6 | 4.1 | 11.8 | 23.7 | 42.7 | 41.3 | 42.1 |
| 1972 | 6.6 | * | * | 0.4 | 1.1 | 2.3 | 5.1 | 12.1 | 22.3 | 35.3 | 44.5 | 41.8 |
| 1971 | 6.8 | * | * | 0.5 | 1.2 | 2.1 | 5.0 | 10.8 | 26.4 | 39.8 | 40.9 | 38.3 |
| 1970 | 6.5 | * | * | 0.5 | 1.3 | 2.1 | 4.0 | 12.2 | 24.3 | 35.4 | 43.7 | 31.5 |
| 1969 | 6.6 | * | * | 0.6 | 1.3 | 2.6 | 5.1 | 11.9 | 24.2 | 36.2 | 42.0 | 29.9 |
| 1968 | 6.2 | * | * | 0.6 | 1.5 | 2.4 | 4.2 | 11.8 | 21.8 | 35.5 | 36.2 | 39.1 |
| BLACK, MALE | | | | | | | | | | | | |
| 1978 | 8.6 | * | * | 0.9 | 1.5 | 2.2 | 5.6 | 10.7 | 27.0 | 60.6 | 68.6 | 90.2 |
| 1977 | 8.4 | * | * | * | 1.1 | 2.3 | 5.3 | 11.7 | 31.5 | 52.2 | 74.5 | 69.4 |
| 1976 | 8.5 | * | * | * | 1.4 | 2.4 | 5.5 | 12.7 | 32.3 | 56.8 | 66.2 | 46.8 |
| 1975 | 8.2 | * | * | * | 1.6 | 1.6 | 5.3 | 13.7 | 31.4 | 52.0 | 60.8 | 58.7 |
| 1974 | 8.4 | * | * | * | 1.5 | 2.2 | 5.1 | 14.8 | 29.9 | 55.6 | 65.1 | 65.1 |
| 1973 | 8.2 | * | * | * | 1.9 | 2.5 | 5.4 | 13.4 | 29.9 | 55.8 | 60.8 | 48.8 |
| 1972 | 7.7 | * | * | * | 1.1 | 2.7 | 8.0 | 15.4 | 28.8 | 40.9 | 57.9 | 55.0 |
| 1971 | 8.1 | * | * | 0.7 | 1.4 | 2.8 | 6.3 | 12.9 | 32.6 | 51.8 | 54.3 | 52.5 |
| 1970 | 8.4 | * | * | 0.7 | 2.0 | 3.2 | 5.4 | 15.0 | 31.5 | 47.9 | 64.0 | 50.0 |
| 1969 | 8.3 | * | * | 1.0 | 1.6 | 3.4 | 6.9 | 14.0 | 31.7 | 49.7 | 57.5 | * |
| 1968 | 7.4 | * | * | 0.7 | 2.2 | 3.3 | 5.4 | 14.1 | 26.0 | 43.7 | 45.5 | 65.7 |
| BLACK, FEMALE | | | | | | | | | | | | |
| 1978 | 6.4 | * | * | * | 0.8 | 1.1 | 2.7 | 7.2 | 21.9 | 33.2 | 44.1 | 53.5 |
| 1977 | 6.3 | * | * | * | 0.7 | 1.6 | 2.7 | 10.0 | 21.1 | 30.8 | 45.2 | 36.2 |
| 1976 | 6.0 | * | * | * | * | 1.2 | 2.6 | 8.3 | 21.2 | 33.4 | 39.3 | 38.2 |
| 1975 | 6.0 | * | * | * | * | 1.2 | 2.6 | 10.5 | 20.0 | 32.0 | 40.9 | 32.1 |
| 1974 | 5.6 | * | * | * | 0.9 | * | 3.2 | 7.5 | 17.3 | 35.3 | 35.3 | 46.2 |
| 1973 | 5.6 | * | * | * | 0.9 | * | 3.0 | 10.4 | 18.6 | 32.5 | 28.0 | 38.4 |
| 1972 | 5.5 | * | * | * | 1.2 | 1.9 | 2.7 | 9.3 | 16.8 | 30.9 | 35.0 | 34.3 |
| 1971 | 5.6 | * | * | * | 1.0 | 1.5 | 3.9 | 8.9 | 21.1 | 30.2 | 31.3 | 29.9 |
| 1970 | 4.9 | * | * | * | * | * | 2.7 | 9.7 | 18.2 | 25.4 | 28.7 | * |
| 1969 | 5.2 | * | * | * | 1.0 | 1.8 | 3.6 | 10.1 | 17.9 | 25.5 | 30.4 | * |
| 1968 | 5.2 | * | * | * | * | 1.7 | 3.2 | 9.8 | 18.2 | 28.8 | 29.3 | * |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|--|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| BENIGN NEOPLASMS AND NEOPLASMS OF UNSPECIFIED NATURE (210-239) | | | | | | | | | | | | |
| BOTH SEXES | | | | | | | | | | | | |
| 1978 | 2.2 | 1.7 | 0.3 | 0.2 | 0.3 | 0.4 | 0.9 | 2.1 | 4.3 | 8.8 | 14.8 | 22.8 |
| 1977 | 2.2 | 2.4 | 0.2 | 0.3 | 0.3 | 0.5 | 1.1 | 2.1 | 4.4 | 8.6 | 14.6 | 22.3 |
| 1976 | 2.2 | 1.4 | 0.3 | 0.3 | 0.3 | 0.4 | 1.1 | 2.3 | 4.5 | 8.5 | 13.3 | 20.8 |
| 1975 | 2.3 | 1.7 | 0.3 | 0.3 | 0.3 | 0.6 | 1.2 | 2.6 | 5.1 | 8.5 | 13.3 | 22.1 |
| 1974 | 2.4 | 2.0 | 0.4 | 0.4 | 0.4 | 0.6 | 1.3 | 2.7 | 5.1 | 9.1 | 14.3 | 19.6 |
| 1973 | 2.4 | 1.5 | 0.5 | 0.4 | 0.3 | 0.6 | 1.5 | 2.8 | 5.3 | 8.9 | 13.7 | 19.9 |
| 1972 | 2.4 | 1.6 | 0.4 | 0.4 | 0.5 | 0.6 | 1.4 | 2.7 | 5.5 | 9.0 | 13.5 | 17.5 |
| 1971 | 2.4 | 2.1 | 0.4 | 0.5 | 0.5 | 0.7 | 1.4 | 3.2 | 5.3 | 8.6 | 12.4 | 20.6 |
| 1970 | 2.4 | 2.1 | 0.4 | 0.4 | 0.5 | 0.6 | 1.6 | 3.3 | 5.4 | 8.4 | 11.9 | 19.6 |
| 1969 | 2.3 | 2.0 | 0.5 | 0.4 | 0.5 | 0.7 | 1.6 | 3.0 | 5.2 | 8.0 | 12.4 | 20.2 |
| 1968 | 2.5 | 2.7 | 0.6 | 0.5 | 0.4 | 0.9 | 1.7 | 3.4 | 5.5 | 9.0 | 12.0 | 20.6 |
| MALE | | | | | | | | | | | | |
| 1978 | 2.2 | 1.4 | 0.4 | 0.3 | 0.3 | 0.4 | 0.9 | 2.0 | 4.5 | 10.5 | 16.9 | 25.6 |
| 1977 | 2.2 | 2.4 | * | 0.2 | 0.3 | 0.5 | 1.1 | 2.1 | 4.7 | 10.1 | 17.6 | 26.3 |
| 1976 | 2.2 | 1.3 | 0.4 | 0.4 | 0.3 | 0.4 | 1.0 | 2.4 | 5.0 | 10.1 | 16.5 | 22.9 |
| 1975 | 2.3 | 2.0 | 0.4 | 0.4 | 0.4 | 0.6 | 1.1 | 2.5 | 6.0 | 9.7 | 16.3 | 25.8 |
| 1974 | 2.4 | 2.2 | 0.5 | 0.3 | 0.4 | 0.5 | 1.2 | 2.8 | 5.6 | 10.3 | 18.3 | 20.7 |
| 1973 | 2.4 | * | 0.5 | 0.4 | 0.4 | 0.5 | 1.3 | 2.8 | 6.0 | 10.9 | 15.9 | 22.1 |
| 1972 | 2.4 | 1.9 | 0.3 | 0.5 | 0.5 | 0.6 | 1.2 | 2.6 | 5.7 | 11.0 | 17.1 | 17.2 |
| 1971 | 2.4 | 2.1 | 0.5 | 0.5 | 0.5 | 0.7 | 1.3 | 3.1 | 6.1 | 10.5 | 14.5 | 23.1 |
| 1970 | 2.3 | 2.4 | 0.3 | 0.4 | 0.5 | 0.5 | 1.4 | 3.0 | 5.9 | 10.0 | 13.7 | 20.5 |
| 1969 | 2.2 | 2.1 | 0.5 | 0.4 | 0.5 | 0.6 | 1.2 | 2.6 | 5.5 | 8.8 | 15.0 | 19.6 |
| 1968 | 2.4 | 2.9 | 0.7 | 0.5 | 0.5 | 0.7 | 1.5 | 3.1 | 5.9 | 10.3 | 13.2 | 21.6 |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|--|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| BENIGN NEOPLASMS AND NEOPLASMS OF UNSPECIFIED NATURE (210-239) | | | | | | | | | | | | |
| FEMALE | | | | | | | | | | | | |
| 1978 | 2.3 | 2.1 | * | 0.2 | 0.4 | 0.4 | 0.9 | 2.1 | 4.1 | 7.5 | 13.6 | 21.6 |
| 1977 | 2.3 | 2.3 | * | 0.3 | 0.3 | 0.4 | 1.1 | 2.2 | 4.1 | 7.5 | 12.8 | 20.4 |
| 1976 | 2.2 | 1.6 | * | 0.2 | 0.3 | 0.5 | 1.2 | 2.2 | 4.1 | 7.3 | 11.3 | 19.8 |
| 1975 | 2.3 | 1.4 | * | 0.3 | 0.3 | 0.5 | 1.3 | 2.6 | 4.3 | 7.5 | 11.4 | 20.4 |
| 1974 | 2.4 | 1.8 | 0.3 | 0.4 | 0.3 | 0.7 | 1.3 | 2.7 | 4.7 | 8.2 | 11.8 | 19.0 |
| 1973 | 2.4 | 2.0 | 0.4 | 0.3 | 0.3 | 0.7 | 1.6 | 2.9 | 4.7 | 7.3 | 12.4 | 18.7 |
| 1972 | 2.4 | 1.4 | 0.4 | 0.3 | 0.4 | 0.7 | 1.6 | 2.8 | 5.3 | 7.5 | 11.2 | 17.6 |
| 1971 | 2.4 | 2.1 | 0.4 | 0.4 | 0.5 | 0.7 | 1.6 | 3.3 | 4.6 | 7.2 | 11.0 | 19.2 |
| 1970 | 2.4 | 1.8 | 0.5 | 0.4 | 0.5 | 0.7 | 1.9 | 3.6 | 4.9 | 7.1 | 10.6 | 19.1 |
| 1969 | 2.4 | 2.0 | 0.5 | 0.3 | 0.6 | 0.9 | 2.1 | 3.3 | 4.8 | 7.4 | 10.6 | 20.5 |
| 1968 | 2.5 | 2.4 | 0.4 | 0.4 | 0.4 | 1.1 | 1.9 | 3.7 | 5.1 | 7.9 | 11.1 | 20.1 |
| WHITE, BOTH SEXES | | | | | | | | | | | | |
| 1978 | 2.3 | 1.7 | 0.3 | 0.2 | 0.3 | 0.4 | 0.8 | 2.0 | 4.1 | 8.7 | 14.8 | 23.1 |
| 1977 | 2.3 | 2.4 | 0.2 | 0.3 | 0.3 | 0.4 | 1.0 | 2.0 | 4.2 | 8.5 | 14.8 | 22.3 |
| 1976 | 2.2 | 1.4 | 0.4 | 0.3 | 0.3 | 0.4 | 1.0 | 2.1 | 4.3 | 8.3 | 13.4 | 21.2 |
| 1975 | 2.3 | 1.8 | 0.3 | 0.3 | 0.3 | 0.5 | 1.2 | 2.4 | 4.9 | 8.3 | 13.4 | 22.2 |
| 1974 | 2.4 | 2.0 | 0.4 | 0.3 | 0.3 | 0.5 | 1.2 | 2.5 | 4.9 | 8.9 | 14.3 | 20.1 |
| 1973 | 2.4 | 1.6 | 0.4 | 0.4 | 0.3 | 0.6 | 1.3 | 2.6 | 5.1 | 8.4 | 13.7 | 20.0 |
| 1972 | 2.3 | 1.6 | 0.4 | 0.4 | 0.4 | 0.6 | 1.2 | 2.5 | 5.4 | 8.6 | 13.7 | 16.5 |
| 1971 | 2.4 | 2.1 | 0.5 | 0.4 | 0.5 | 0.7 | 1.3 | 3.0 | 5.2 | 8.6 | 12.4 | 21.2 |
| 1970 | 2.4 | 2.1 | 0.4 | 0.4 | 0.5 | 0.6 | 1.5 | 3.0 | 5.2 | 8.2 | 11.9 | 20.2 |
| 1969 | 2.3 | 1.8 | 0.4 | 0.4 | 0.5 | 0.7 | 1.5 | 2.7 | 4.9 | 7.8 | 12.1 | 20.7 |
| 1968 | 2.5 | 2.5 | 0.6 | 0.4 | 0.4 | 0.8 | 1.6 | 3.2 | 5.3 | 8.9 | 11.6 | 21.0 |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|--|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| BENIGN NEOPLASMS AND NEOPLASMS OF UNSPECIFIED NATURE (210-239) | | | | | | | | | | | | |
| WHITE, MALE | | | | | | | | | | | | |
| 1978 | 2.2 | 1.6 | 0.4 | 0.3 | 0.3 | 0.4 | 0.8 | 2.0 | 4.3 | 10.5 | 17.0 | 25.9 |
| 1977 | 2.2 | 2.4 | * | 0.2 | 0.3 | 0.5 | 0.9 | 2.0 | 4.4 | 9.8 | 18.1 | 27.1 |
| 1976 | 2.2 | * | 0.5 | 0.4 | 0.3 | 0.3 | 0.9 | 2.1 | 4.7 | 10.1 | 16.4 | 24.1 |
| 1975 | 2.4 | 2.3 | 0.4 | 0.3 | 0.4 | 0.6 | 1.1 | 2.4 | 5.8 | 9.8 | 16.4 | 26.5 |
| 1974 | 2.4 | 1.9 | 0.5 | 0.3 | 0.4 | 0.5 | 1.1 | 2.5 | 5.4 | 10.5 | 18.4 | 21.9 |
| 1973 | 2.4 | * | 0.5 | 0.4 | 0.4 | 0.5 | 1.2 | 2.7 | 5.8 | 10.4 | 15.9 | 22.9 |
| 1972 | 2.4 | 1.7 | 0.4 | 0.4 | 0.5 | 0.6 | 1.2 | 2.4 | 5.6 | 10.4 | 17.3 | 16.8 |
| 1971 | 2.5 | 2.0 | 0.5 | 0.4 | 0.5 | 0.6 | 1.1 | 3.0 | 6.0 | 10.6 | 14.5 | 24.4 |
| 1970 | 2.3 | 2.5 | * | 0.4 | 0.5 | 0.5 | 1.3 | 2.9 | 5.6 | 9.8 | 13.8 | 21.8 |
| 1969 | 2.2 | 1.8 | 0.5 | 0.4 | 0.5 | 0.6 | 1.2 | 2.4 | 5.3 | 8.7 | 15.1 | 20.2 |
| 1968 | 2.4 | 2.9 | 0.7 | 0.5 | 0.4 | 0.7 | 1.5 | 3.0 | 5.7 | 10.3 | 13.2 | 23.3 |
| WHITE, FEMALE | | | | | | | | | | | | |
| 1978 | 2.3 | 1.9 | * | 0.2 | 0.4 | 0.4 | 0.8 | 2.0 | 3.9 | 7.3 | 13.6 | 21.8 |
| 1977 | 2.3 | 2.3 | * | 0.3 | 0.3 | 0.4 | 1.0 | 2.1 | 4.0 | 7.5 | 12.8 | 20.1 |
| 1976 | 2.2 | 1.7 | * | 0.2 | 0.2 | 0.5 | 1.1 | 2.0 | 3.9 | 6.9 | 11.6 | 19.8 |
| 1975 | 2.3 | * | * | 0.3 | 0.3 | 0.5 | 1.2 | 2.5 | 4.1 | 7.2 | 11.5 | 20.2 |
| 1974 | 2.4 | 2.0 | 0.4 | 0.4 | 0.3 | 0.6 | 1.2 | 2.5 | 4.5 | 7.7 | 11.9 | 19.2 |
| 1973 | 2.3 | 1.9 | 0.4 | 0.4 | 0.3 | 0.7 | 1.4 | 2.5 | 4.5 | 6.9 | 12.4 | 18.6 |
| 1972 | 2.3 | 1.5 | 0.5 | 0.3 | 0.4 | 0.5 | 1.3 | 2.5 | 5.3 | 7.2 | 11.3 | 16.3 |
| 1971 | 2.4 | 2.2 | 0.4 | 0.4 | 0.5 | 0.7 | 1.4 | 3.0 | 4.5 | 7.1 | 11.0 | 19.6 |
| 1970 | 2.4 | 1.7 | 0.4 | 0.4 | 0.4 | 0.6 | 1.6 | 3.1 | 4.8 | 6.9 | 10.7 | 19.3 |
| 1969 | 2.3 | 1.8 | 0.4 | 0.3 | 0.6 | 0.8 | 1.7 | 3.0 | 4.5 | 7.1 | 10.0 | 20.9 |
| 1968 | 2.5 | 2.1 | 0.5 | 0.4 | 0.4 | 0.9 | 1.6 | 3.5 | 4.9 | 7.7 | 10.6 | 19.8 |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|--|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| BENIGN NEOPLASMS AND NEOPLASMS OF UNSPECIFIED NATURE (210-239) | | | | | | | | | | | | |
| ALL OTHER, BOTH SEXES | | | | | | | | | | | | |
| 1978 | 2.0 | * | * | * | 0.3 | 0.6 | 1.4 | 2.7 | 6.3 | 10.0 | 14.5 | 19.6 |
| 1977 | 2.1 | * | * | * | 0.3 | 0.9 | 1.8 | 3.0 | 6.1 | 9.9 | 13.1 | 21.9 |
| 1976 | 2.1 | * | * | 0.4 | 0.4 | 0.5 | 1.6 | 3.8 | 6.9 | 10.9 | 11.8 | 16.3 |
| 1975 | 2.1 | * | * | 0.5 | 0.4 | 0.7 | 1.5 | 3.7 | 6.6 | 9.9 | 12.1 | 21.2 |
| 1974 | 2.4 | * | * | 0.5 | 0.6 | 1.1 | 2.0 | 4.5 | 6.9 | 10.8 | 13.5 | * |
| 1973 | 2.4 | * | * | * | * | 0.8 | 2.5 | 4.5 | 7.3 | 13.1 | 13.4 | 18.1 |
| 1972 | 2.5 | * | * | 0.5 | 0.8 | 1.1 | 2.5 | 4.6 | 5.9 | 13.2 | 11.9 | 29.3 |
| 1971 | 2.3 | * | * | 0.7 | 0.4 | 1.2 | 2.6 | 4.7 | 6.3 | 8.2 | 12.5 | * |
| 1970 | 2.5 | * | * | 0.4 | 0.8 | 0.9 | 2.9 | 5.7 | 7.0 | 10.5 | 11.2 | * |
| 1969 | 2.7 | * | * | 0.5 | 0.7 | 1.1 | 3.1 | 5.3 | 7.9 | 10.2 | 16.1 | * |
| 1968 | 2.7 | * | * | 0.7 | 0.6 | 1.7 | 2.9 | 4.9 | 7.1 | 9.8 | 16.4 | * |
| ALL OTHER, MALE | | | | | | | | | | | | |
| 1978 | 1.9 | * | * | * | * | * | 1.4 | 2.5 | 6.5 | 10.7 | 16.2 | * |
| 1977 | 2.0 | * | * | * | * | * | 1.9 | 2.7 | 7.1 | 12.0 | 12.3 | * |
| 1976 | 2.2 | * | * | * | * | * | 1.6 | 4.4 | 7.9 | 10.8 | 17.5 | * |
| 1975 | 2.0 | * | * | * | * | * | * | 3.4 | 7.7 | 9.6 | 14.9 | * |
| 1974 | 2.2 | * | * | * | * | * | 1.6 | 4.9 | 7.7 | 8.6 | 17.9 | * |
| 1973 | 2.3 | * | * | * | * | * | 1.9 | 3.3 | 8.1 | 16.0 | 15.2 | * |
| 1972 | 2.5 | * | * | 0.6 | 1.0 | * | * | 4.0 | 6.4 | 16.6 | 14.7 | * |
| 1971 | 2.2 | * | * | 0.6 | * | 1.4 | 2.5 | 3.9 | 7.2 | 8.8 | 14.0 | * |
| 1970 | 2.3 | * | * | * | * | * | 1.9 | 3.8 | 8.4 | 11.7 | 12.4 | * |
| 1969 | 2.2 | * | * | 0.7 | * | * | * | 4.4 | 7.4 | 9.7 | 13.8 | * |
| 1968 | 2.2 | * | * | * | * | * | * | 4.1 | 7.0 | 10.2 | 13.4 | * |
| ALL OTHER, FEMALE | | | | | | | | | | | | |
| 1978 | 2.1 | * | * | * | * | * | 1.4 | 2.8 | 6.2 | 9.4 | 13.4 | 18.3 |
| 1977 | 2.1 | * | * | * | * | 1.0 | 1.7 | 3.2 | 5.3 | 8.2 | 13.6 | 24.0 |
| 1976 | 2.1 | * | * | * | * | * | 1.7 | 3.4 | 6.0 | 11.1 | 8.0 | * |
| 1975 | 2.2 | * | * | * | * | * | 1.9 | 3.9 | 5.7 | 10.2 | 10.2 | 22.6 |
| 1974 | 2.5 | * | * | * | * | 1.2 | 2.4 | 4.2 | 6.2 | 12.6 | 10.5 | * |
| 1973 | 2.6 | * | * | * | * | 1.0 | 2.9 | 5.6 | 6.6 | 10.8 | 12.3 | * |
| 1972 | 2.5 | * | * | * | * | 1.4 | 3.4 | 5.1 | 5.5 | 10.4 | 10.0 | 34.2 |
| 1971 | 2.3 | * | * | 0.7 | * | * | 2.7 | 5.3 | 5.5 | 7.8 | 11.4 | * |
| 1970 | 2.8 | * | * | * | 0.8 | * | 3.8 | 7.4 | 5.8 | 9.5 | 10.2 | * |
| 1969 | 3.1 | * | * | * | * | 1.4 | 4.7 | 6.2 | 8.3 | 10.6 | 17.8 | * |
| 1968 | 3.1 | * | * | 0.8 | * | 2.3 | 4.1 | 5.6 | 7.2 | 9.4 | 18.8 | * |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|--|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| BENIGN NEOPLASMS AND NEOPLASMS OF UNSPECIFIED NATURE (210-239) | | | | | | | | | | | | |
| BLACK, BOTH SEXES | | | | | | | | | | | | |
| 1978 | 2.2 | * | * | * | 0.4 | 0.5 | 1.6 | 3.0 | 6.8 | 10.4 | 14.4 | 21.3 |
| 1977 | 2.2 | * | * | * | * | 0.9 | 2.0 | 3.2 | 6.4 | 10.6 | 13.1 | 23.8 |
| 1976 | 2.3 | * | * | 0.4 | 0.4 | 0.6 | 1.8 | 4.3 | 7.3 | 12.0 | 11.1 | 16.9 |
| 1975 | 2.2 | * | * | 0.6 | 0.5 | 0.9 | 1.6 | 3.8 | 7.0 | 10.0 | 11.8 | 22.3 |
| 1974 | 2.5 | * | * | 0.5 | 0.6 | 1.1 | 2.2 | 4.9 | 7.3 | 11.3 | 13.7 | * |
| 1973 | 2.6 | * | * | * | * | 0.9 | 2.6 | 4.9 | 7.9 | 13.5 | 12.8 | 19.3 |
| 1972 | 2.7 | * | * | 0.5 | 0.8 | 1.2 | 2.9 | 4.8 | 6.1 | 14.0 | 13.0 | 30.9 |
| 1971 | 2.4 | * | * | 0.7 | * | 1.3 | 3.0 | 5.0 | 6.6 | 8.6 | 12.6 | * |
| 1970 | 2.7 | * | * | 0.4 | 0.8 | 1.0 | 3.4 | 6.1 | 7.3 | 11.2 | 11.8 | * |
| 1969 | 2.8 | * | * | 0.5 | 0.6 | 1.2 | 3.3 | 5.6 | 8.2 | 10.5 | 16.3 | * |
| 1968 | 2.8 | * | * | 0.7 | 0.6 | 1.9 | 3.1 | 5.2 | 7.4 | 10.1 | 16.0 | * |
| BLACK, MALE | | | | | | | | | | | | |
| 1978 | 2.1 | * | * | * | * | * | * | 2.8 | 7.1 | 11.4 | 15.9 | * |
| 1977 | 2.1 | * | * | * | * | * | 2.1 | 2.8 | 7.4 | 12.7 | 12.3 | * |
| 1976 | 2.3 | * | * | * | * | * | 1.8 | 4.9 | 8.4 | 11.8 | 17.1 | * |
| 1975 | 2.1 | * | * | * | * | * | * | 3.8 | 8.1 | 9.8 | 14.9 | * |
| 1974 | 2.4 | * | * | * | * | * | 1.8 | 5.3 | 8.2 | 9.2 | 18.7 | * |
| 1973 | 2.4 | * | * | * | * | * | 2.1 | 3.5 | 8.9 | 16.3 | 13.7 | * |
| 1972 | 2.7 | * | * | 0.7 | 1.1 | * | * | 4.2 | 6.6 | 18.3 | 16.2 | * |
| 1971 | 2.3 | * | * | 0.7 | * | * | 2.8 | 4.3 | 7.6 | 9.0 | 13.4 | * |
| 1970 | 2.4 | * | * | * | * | * | 2.2 | 4.0 | 8.7 | 12.4 | 12.5 | * |
| 1969 | 2.3 | * | * | * | * | * | * | 4.8 | 7.8 | 9.9 | 12.6 | * |
| 1968 | 2.3 | * | * | * | * | * | * | 4.4 | 7.4 | 10.8 | * | * |
| BLACK, FEMALE | | | | | | | | | | | | |
| 1978 | 2.2 | * | * | * | * | * | 1.6 | 3.2 | 6.6 | 9.7 | 13.4 | 20.2 |
| 1977 | 2.3 | * | * | * | * | 1.1 | 1.9 | 3.4 | 5.6 | 9.0 | 13.6 | 25.5 |
| 1976 | 2.3 | * | * | * | * | * | 1.8 | 3.8 | 6.5 | 12.1 | 7.2 | * |
| 1975 | 2.3 | * | * | * | * | * | 2.1 | 3.8 | 6.0 | 10.1 | 9.8 | * |
| 1974 | 2.7 | * | * | * | * | 1.3 | 2.5 | 4.6 | 6.5 | 13.0 | 10.4 | * |
| 1973 | 2.7 | * | * | * | * | 1.2 | 3.1 | 6.0 | 7.0 | 11.4 | 12.2 | * |
| 1972 | 2.7 | * | * | * | * | 1.5 | 4.0 | 5.4 | 5.7 | 10.6 | 10.7 | 37.1 |
| 1971 | 2.5 | * | * | 0.7 | * | * | 3.1 | 5.6 | 5.7 | 8.3 | 12.0 | * |
| 1970 | 3.0 | * | * | * | * | * | 4.4 | 7.9 | 6.1 | 10.3 | 11.3 | * |
| 1969 | 3.3 | * | * | * | * | 1.4 | 5.2 | 6.4 | 8.6 | 11.0 | 18.9 | * |
| 1968 | 3.3 | * | * | 0.7 | * | 2.5 | 4.7 | 5.9 | 7.5 | 9.6 | 19.4 | * |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| DIABETES MELLITUS (250) | | | | | | | | | | | | |
| BOTH SEXES | | | | | | | | | | | | |
| 1978 | 15.2 | * | * | 0.1 | 0.3 | 1.6 | 3.8 | 9.6 | 26.3 | 64.2 | 135.8 | 223.1 |
| 1977 | 15.0 | * | * | 0.1 | 0.4 | 1.5 | 3.7 | 9.6 | 26.7 | 65.7 | 134.3 | 212.7 |
| 1976 | 15.9 | * | * | 0.1 | 0.4 | 1.8 | 3.9 | 9.8 | 27.9 | 69.8 | 147.7 | 227.3 |
| 1975 | 16.4 | * | * | 0.1 | 0.4 | 1.6 | 4.2 | 10.3 | 28.9 | 75.0 | 153.0 | 231.5 |
| 1974 | 17.5 | * | * | 0.2 | 0.5 | 1.8 | 4.3 | 11.4 | 31.4 | 81.3 | 165.8 | 253.9 |
| 1973 | 18.1 | * | * | 0.2 | 0.5 | 2.0 | 4.6 | 11.8 | 33.8 | 85.2 | 174.6 | 249.4 |
| 1972 | 18.5 | * | 0.2 | 0.2 | 0.6 | 2.4 | 4.9 | 12.3 | 33.6 | 89.5 | 177.8 | 258.1 |
| 1971 | 18.5 | 0.6 | * | 0.2 | 0.6 | 2.3 | 5.1 | 12.4 | 35.0 | 89.4 | 180.2 | 250.0 |
| 1970 | 18.9 | * | 0.2 | 0.3 | 0.7 | 2.2 | 5.3 | 12.8 | 36.7 | 92.1 | 186.8 | 230.2 |
| 1969 | 19.1 | * | 0.1 | 0.2 | 0.7 | 2.5 | 5.4 | 13.1 | 36.7 | 94.0 | 190.6 | 258.4 |
| 1968 | 19.2 | * | 0.2 | 0.2 | 0.7 | 2.6 | 5.3 | 13.1 | 38.7 | 96.3 | 188.2 | 257.9 |
| MALE | | | | | | | | | | | | |
| 1978 | 12.9 | * | * | * | 0.3 | 1.8 | 4.4 | 10.7 | 26.9 | 64.3 | 129.5 | 210.4 |
| 1977 | 12.8 | * | * | * | 0.3 | 1.6 | 4.1 | 10.7 | 27.2 | 66.0 | 126.3 | 206.7 |
| 1976 | 13.2 | * | * | * | 0.4 | 1.9 | 4.6 | 10.0 | 28.0 | 68.8 | 139.5 | 209.7 |
| 1975 | 13.7 | * | * | * | 0.4 | 1.8 | 4.6 | 11.2 | 29.6 | 72.2 | 142.3 | 222.7 |
| 1974 | 14.6 | * | * | 0.1 | 0.5 | 1.9 | 4.7 | 11.7 | 32.0 | 77.0 | 153.7 | 246.4 |
| 1973 | 15.2 | * | * | 0.1 | 0.5 | 2.2 | 5.2 | 12.4 | 33.5 | 81.3 | 163.0 | 237.9 |
| 1972 | 15.5 | * | * | 0.2 | 0.5 | 2.4 | 5.6 | 13.3 | 31.8 | 84.4 | 164.8 | 262.8 |
| 1971 | 15.5 | * | * | 0.2 | 0.6 | 2.6 | 5.7 | 12.8 | 34.2 | 83.1 | 163.8 | 246.1 |
| 1970 | 15.9 | * | * | 0.2 | 0.6 | 2.4 | 6.5 | 13.3 | 35.1 | 84.8 | 169.9 | 220.0 |
| 1969 | 16.0 | * | * | 0.2 | 0.6 | 2.7 | 6.1 | 13.5 | 34.9 | 85.5 | 174.1 | 241.2 |
| 1968 | 16.2 | * | * | 0.2 | 0.7 | 2.9 | 5.9 | 13.9 | 37.5 | 88.3 | 165.9 | 252.5 |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| DIABETES MELLITUS (250) | | | | | | | | | | | | |
| FEMALE | | | | | | | | | | | | |
| 1978 | 17.5 | * | * | 0.1 | 0.4 | 1.4 | 3.2 | 8.6 | 25.8 | 64.1 | 139.6 | 228.8 |
| 1977 | 17.1 | * | * | 0.2 | 0.4 | 1.3 | 3.3 | 8.6 | 26.2 | 65.4 | 139.0 | 215.4 |
| 1976 | 18.3 | * | * | 0.1 | 0.4 | 1.6 | 3.3 | 9.6 | 27.8 | 70.5 | 152.6 | 235.5 |
| 1975 | 18.8 | * | * | 0.2 | 0.4 | 1.4 | 3.7 | 9.4 | 28.3 | 77.1 | 159.5 | 235.7 |
| 1974 | 20.3 | * | * | 0.2 | 0.6 | 1.8 | 3.8 | 11.1 | 30.9 | 84.6 | 173.3 | 257.7 |
| 1973 | 20.8 | * | * | 0.2 | 0.5 | 1.7 | 4.0 | 11.1 | 34.0 | 88.2 | 181.9 | 255.2 |
| 1972 | 21.3 | * | * | 0.2 | 0.7 | 2.3 | 4.2 | 11.3 | 35.2 | 93.5 | 186.1 | 255.7 |
| 1971 | 21.3 | * | * | 0.3 | 0.6 | 2.0 | 4.5 | 12.1 | 35.8 | 94.3 | 190.8 | 252.0 |
| 1970 | 21.7 | * | * | 0.3 | 0.7 | 2.1 | 4.2 | 12.3 | 38.0 | 97.9 | 198.0 | 235.9 |
| 1969 | 22.1 | * | * | 0.3 | 0.8 | 2.3 | 4.8 | 12.8 | 38.4 | 100.7 | 201.8 | 267.7 |
| 1968 | 22.1 | * | * | 0.3 | 0.7 | 2.4 | 4.7 | 12.4 | 39.8 | 102.7 | 203.5 | 261.1 |
| WHITE, BOTH SEXES | | | | | | | | | | | | |
| 1978 | 14.7 | * | * | 0.1 | 0.3 | 1.4 | 3.2 | 8.0 | 22.9 | 58.4 | 130.9 | 222.9 |
| 1977 | 14.4 | * | * | 0.1 | 0.3 | 1.3 | 3.1 | 7.8 | 22.8 | 59.9 | 130.0 | 211.2 |
| 1976 | 15.2 | * | * | 0.1 | 0.3 | 1.6 | 3.2 | 7.9 | 23.6 | 63.4 | 143.8 | 227.4 |
| 1975 | 15.7 | * | * | 0.1 | 0.4 | 1.5 | 3.4 | 8.3 | 24.8 | 68.0 | 148.9 | 232.8 |
| 1974 | 16.9 | * | * | 0.2 | 0.5 | 1.6 | 3.4 | 9.1 | 27.1 | 73.8 | 163.2 | 255.8 |
| 1973 | 17.3 | * | * | 0.2 | 0.5 | 1.7 | 3.7 | 9.3 | 28.6 | 77.0 | 172.4 | 250.0 |
| 1972 | 17.7 | * | * | 0.2 | 0.6 | 2.1 | 3.9 | 9.4 | 28.5 | 81.5 | 176.1 | 258.8 |
| 1971 | 17.8 | * | * | 0.2 | 0.5 | 2.1 | 4.1 | 9.7 | 29.7 | 82.2 | 178.6 | 251.4 |
| 1970 | 18.3 | * | * | 0.3 | 0.6 | 2.0 | 4.3 | 10.4 | 31.4 | 85.5 | 185.4 | 236.5 |
| 1969 | 18.5 | * | * | 0.2 | 0.6 | 2.3 | 4.3 | 10.7 | 30.9 | 87.5 | 189.2 | 264.4 |
| 1968 | 18.6 | * | 0.2 | 0.2 | 0.6 | 2.3 | 4.5 | 10.6 | 33.0 | 89.8 | 187.0 | 260.8 |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| DIABETES MELLITUS (250) | | | | | | | | | | | | |
| WHITE, MALE | | | | | | | | | | | | |
| 1978 | 12.7 | * | * | * | 0.3 | 1.6 | 3.7 | 9.5 | 24.4 | 60.7 | 127.7 | 214.1 |
| 1977 | 12.5 | * | * | * | 0.3 | 1.5 | 3.5 | 9.3 | 24.9 | 62.5 | 124.9 | 207.9 |
| 1976 | 12.9 | * | * | * | 0.3 | 1.7 | 3.8 | 8.4 | 25.2 | 65.1 | 139.1 | 211.3 |
| 1975 | 13.5 | * | * | * | 0.4 | 1.7 | 3.8 | 9.5 | 26.9 | 68.7 | 141.6 | 225.4 |
| 1974 | 14.3 | * | * | * | 0.5 | 1.6 | 4.0 | 10.0 | 29.3 | 73.2 | 153.6 | 251.7 |
| 1973 | 14.8 | * | * | 0.1 | 0.5 | 1.8 | 4.4 | 10.6 | 30.1 | 76.4 | 162.4 | 244.3 |
| 1972 | 15.2 | * | * | 0.2 | 0.5 | 2.2 | 4.5 | 10.8 | 28.8 | 80.1 | 165.6 | 268.6 |
| 1971 | 15.2 | * | * | 0.2 | 0.5 | 2.4 | 4.9 | 10.9 | 31.3 | 79.0 | 164.5 | 252.7 |
| 1970 | 15.7 | * | * | 0.2 | 0.6 | 2.1 | 5.5 | 11.6 | 32.2 | 81.6 | 169.3 | 228.8 |
| 1969 | 15.8 | * | * | 0.2 | 0.5 | 2.5 | 5.0 | 12.0 | 31.8 | 82.6 | 174.8 | 250.4 |
| 1968 | 16.1 | * | * | 0.2 | 0.6 | 2.6 | 5.4 | 12.2 | 34.5 | 85.2 | 166.7 | 259.1 |
| WHITE, FEMALE | | | | | | | | | | | | |
| 1978 | 16.7 | * | * | * | 0.3 | 1.3 | 2.7 | 6.6 | 21.6 | 56.6 | 132.8 | 226.9 |
| 1977 | 16.3 | * | * | 0.2 | 0.3 | 1.2 | 2.6 | 6.4 | 20.9 | 57.9 | 133.1 | 212.7 |
| 1976 | 17.4 | * | * | 0.1 | 0.4 | 1.4 | 2.6 | 7.3 | 22.2 | 62.2 | 146.6 | 235.0 |
| 1975 | 17.9 | * | * | 0.2 | 0.4 | 1.3 | 3.0 | 7.2 | 22.9 | 67.4 | 153.4 | 236.5 |
| 1974 | 19.3 | * | * | 0.2 | 0.5 | 1.6 | 2.8 | 8.2 | 25.2 | 74.2 | 169.0 | 257.9 |
| 1973 | 19.7 | * | * | 0.3 | 0.4 | 1.5 | 3.0 | 8.2 | 27.2 | 77.5 | 178.5 | 253.1 |
| 1972 | 20.1 | * | * | 0.2 | 0.7 | 1.9 | 3.2 | 8.1 | 28.2 | 82.5 | 182.6 | 253.8 |
| 1971 | 20.2 | * | * | 0.3 | 0.5 | 1.8 | 3.3 | 8.6 | 28.2 | 84.7 | 187.7 | 250.8 |
| 1970 | 20.8 | * | * | 0.3 | 0.6 | 1.9 | 3.2 | 9.3 | 30.7 | 88.5 | 196.0 | 240.7 |
| 1969 | 21.1 | * | * | 0.3 | 0.7 | 2.2 | 3.6 | 9.5 | 30.2 | 91.3 | 198.7 | 272.2 |
| 1968 | 21.0 | * | * | 0.3 | 0.7 | 2.0 | 3.6 | 9.0 | 31.6 | 93.5 | 200.8 | 262.0 |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| DIABETES MELLITUS (250) | | | | | | | | | | | | |
| ALL OTHER, BOTH SEXES | | | | | | | | | | | | |
| 1978 | 18.4 | * | * | * | 0.4 | 2.8 | 7.8 | 22.0 | 57.6 | 118.6 | 186.6 | 225.0 |
| 1977 | 18.7 | * | * | * | 0.6 | 2.4 | 7.8 | 23.9 | 62.6 | 120.6 | 178.1 | 229.4 |
| 1976 | 19.9 | * | * | * | 0.7 | 3.1 | 9.0 | 25.4 | 67.5 | 130.6 | 188.2 | 224.8 |
| 1975 | 20.4 | * | * | * | 0.7 | 2.5 | 9.4 | 26.6 | 66.8 | 142.8 | 196.8 | 216.4 |
| 1974 | 21.7 | * | * | * | 0.7 | 3.4 | 10.4 | 30.5 | 71.9 | 155.4 | 195.0 | 231.9 |
| 1973 | 23.2 | * | * | * | 0.7 | 4.1 | 10.9 | 32.2 | 83.1 | 166.0 | 200.0 | 242.5 |
| 1972 | 23.8 | * | * | * | 0.5 | 4.4 | 12.3 | 36.9 | 82.5 | 169.1 | 198.1 | 250.4 |
| 1971 | 23.5 | * | * | * | 1.0 | 3.9 | 12.5 | 36.1 | 86.2 | 160.2 | 199.4 | 233.3 |
| 1970 | 22.9 | * | * | * | 0.9 | 4.0 | 12.6 | 34.0 | 86.7 | 157.3 | 204.4 | 165.6 |
| 1969 | 23.7 | * | * | * | 1.2 | 3.8 | 14.0 | 34.8 | 92.4 | 158.7 | 209.9 | 189.9 |
| 1968 | 23.9 | * | * | * | 1.1 | 4.9 | 11.6 | 35.8 | 94.3 | 161.4 | 203.1 | 225.5 |
| ALL OTHER, MALE | | | | | | | | | | | | |
| 1978 | 14.4 | * | * | * | * | 3.4 | 9.4 | 20.4 | 50.9 | 97.6 | 145.5 | 172.4 |
| 1977 | 14.3 | * | * | * | * | 2.9 | 8.6 | 22.7 | 49.4 | 98.4 | 140.2 | 194.6 |
| 1976 | 15.3 | * | * | * | * | 3.6 | 10.2 | 23.8 | 55.0 | 103.9 | 143.6 | 194.5 |
| 1975 | 15.5 | * | * | * | 0.8 | 2.8 | 10.8 | 26.1 | 54.9 | 105.4 | 149.6 | 196.3 |
| 1974 | 16.1 | * | * | * | * | 4.0 | 10.3 | 26.9 | 57.8 | 112.9 | 154.1 | 192.0 |
| 1973 | 18.0 | * | * | * | * | 5.5 | 11.4 | 28.6 | 67.0 | 128.1 | 168.8 | 172.9 |
| 1972 | 18.0 | * | * | * | * | 4.0 | 13.8 | 35.7 | 61.2 | 125.0 | 155.8 | 204.3 |
| 1971 | 17.4 | * | * | * | 1.0 | 4.5 | 12.1 | 30.4 | 62.3 | 122.1 | 157.0 | 180.9 |
| 1970 | 17.5 | * | * | * | * | 4.5 | 14.5 | 28.7 | 63.5 | 114.7 | 177.7 | 142.5 |
| 1969 | 17.5 | * | * | * | 1.0 | 4.8 | 15.6 | 27.8 | 64.5 | 113.2 | 166.0 | 154.5 |
| 1968 | 17.5 | * | * | * | 1.2 | 5.1 | 10.2 | 29.0 | 67.5 | 118.4 | 155.9 | 187.8 |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| DIABETES MELLITUS (250) | | | | | | | | | | | | |
| ALL OTHER, FEMALE | | | | | | | | | | | | |
| 1978 | 22.0 | * | * | * | * | 2.3 | 6.5 | 23.3 | 63.2 | 134.8 | 215.2 | 255.0 |
| 1977 | 22.6 | * | * | * | 0.8 | 1.9 | 7.2 | 24.8 | 73.6 | 138.0 | 203.4 | 248.1 |
| 1976 | 24.1 | * | * | * | 0.7 | 2.6 | 8.0 | 26.7 | 78.1 | 151.7 | 218.6 | 241.8 |
| 1975 | 24.8 | * | * | * | * | 2.2 | 8.3 | 27.0 | 76.8 | 172.5 | 228.5 | 225.8 |
| 1974 | 26.7 | * | * | * | 0.9 | 2.9 | 10.5 | 33.5 | 83.8 | 189.4 | 223.2 | 255.3 |
| 1973 | 28.0 | * | * | * | 0.8 | 2.8 | 10.5 | 35.2 | 96.9 | 196.6 | 222.6 | 284.8 |
| 1972 | 29.1 | * | * | * | * | 4.8 | 11.0 | 37.9 | 100.9 | 205.2 | 229.2 | 278.9 |
| 1971 | 29.0 | * | * | * | 1.1 | 3.3 | 12.9 | 41.1 | 107.0 | 191.3 | 230.7 | 267.1 |
| 1970 | 27.9 | * | * | * | 1.2 | 3.6 | 11.0 | 38.7 | 107.2 | 192.5 | 224.7 | 181.8 |
| 1969 | 29.4 | * | * | * | 1.3 | 3.0 | 12.7 | 40.9 | 117.2 | 196.1 | 242.5 | 213.8 |
| 1968 | 29.8 | * | * | * | 1.1 | 4.8 | 12.8 | 41.8 | 118.4 | 197.0 | 240.6 | 250.8 |
| BLACK, BOTH SEXES | | | | | | | | | | | | |
| 1978 | 19.9 | * | * | * | 0.5 | 3.2 | 8.8 | 24.4 | 61.1 | 123.7 | 189.3 | 229.3 |
| 1977 | 20.3 | * | * | * | 0.7 | 2.7 | 9.0 | 25.6 | 67.1 | 126.2 | 181.9 | 237.8 |
| 1976 | 21.4 | * | * | * | 0.7 | 3.4 | 10.0 | 28.1 | 71.2 | 136.7 | 186.4 | 234.6 |
| 1975 | 21.8 | * | * | * | 0.7 | 2.8 | 10.5 | 28.6 | 70.5 | 148.8 | 198.9 | 217.7 |
| 1974 | 23.2 | * | * | * | 0.8 | 3.8 | 11.5 | 32.7 | 75.6 | 163.1 | 197.5 | 243.0 |
| 1973 | 24.7 | * | * | * | 0.8 | 4.5 | 11.9 | 34.6 | 86.5 | 174.4 | 201.8 | 249.1 |
| 1972 | 25.2 | * | * | * | 0.6 | 4.7 | 13.4 | 38.8 | 86.5 | 177.4 | 205.5 | 245.5 |
| 1971 | 24.6 | * | * | * | 1.1 | 4.2 | 13.7 | 38.1 | 90.8 | 163.5 | 199.3 | 240.2 |
| 1970 | 24.3 | * | * | * | 1.1 | 4.5 | 13.8 | 36.5 | 90.6 | 164.6 | 209.0 | 173.8 |
| 1969 | 24.9 | * | * | * | 1.3 | 4.2 | 15.3 | 36.6 | 96.1 | 164.3 | 213.0 | 203.1 |
| 1968 | 25.0 | * | * | * | 1.2 | 5.3 | 12.5 | 37.8 | 98.0 | 164.1 | 210.9 | 232.6 |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| DIABETES MELLITUS (250) | | | | | | | | | | | | |
| BLACK, MALE | | | | | | | | | | | | |
| 1978 | 15.4 | * | * | * | * | 3.9 | 10.7 | 22.4 | 53.5 | 102.4 | 143.9 | 172.5 |
| 1977 | 15.3 | * | * | * | * | 3.3 | 10.0 | 24.1 | 52.5 | 102.2 | 142.1 | 198.0 |
| 1976 | 16.2 | * | * | * | * | 3.9 | 11.3 | 26.5 | 58.6 | 109.1 | 136.3 | 204.3 |
| 1975 | 16.2 | * | * | * | 0.8 | 3.3 | 12.0 | 28.1 | 57.8 | 106.0 | 146.4 | 189.1 |
| 1974 | 16.8 | * | * | * | * | 4.4 | 11.2 | 28.5 | 59.6 | 116.7 | 156.5 | 195.3 |
| 1973 | 18.7 | * | * | * | * | 6.2 | 12.3 | 29.9 | 69.9 | 132.5 | 165.2 | 187.8 |
| 1972 | 18.8 | * | * | * | * | 4.6 | 15.4 | 36.6 | 65.1 | 131.3 | 158.4 | 190.0 |
| 1971 | 17.9 | * | * | * | 1.1 | 5.0 | 13.1 | 32.0 | 65.0 | 122.9 | 154.8 | 187.5 |
| 1970 | 18.1 | * | * | * | * | 5.1 | 15.9 | 30.8 | 64.9 | 118.0 | 175.5 | 147.8 |
| 1969 | 17.9 | * | * | * | 1.1 | 5.3 | 17.1 | 29.1 | 66.9 | 114.1 | 161.7 | 170.3 |
| 1968 | 17.8 | * | * | * | 1.2 | 5.3 | 10.9 | 30.4 | 69.8 | 117.2 | 159.4 | 191.4 |
| BLACK, FEMALE | | | | | | | | | | | | |
| 1978 | 24.0 | * | * | * | * | 2.6 | 7.2 | 26.0 | 67.3 | 139.3 | 219.6 | 258.6 |
| 1977 | 24.8 | * | * | * | 0.9 | 2.2 | 8.1 | 26.8 | 79.1 | 144.3 | 208.8 | 258.5 |
| 1976 | 26.1 | * | * | * | 0.8 | 3.0 | 9.0 | 29.5 | 81.5 | 157.4 | 220.1 | 250.6 |
| 1975 | 26.9 | * | * | * | * | 2.4 | 9.3 | 29.0 | 81.0 | 181.3 | 234.5 | 233.3 |
| 1974 | 29.0 | * | * | * | 0.9 | 3.3 | 11.7 | 36.3 | 88.8 | 198.9 | 225.2 | 269.2 |
| 1973 | 30.1 | * | * | * | 0.9 | 3.1 | 11.6 | 38.5 | 100.3 | 206.8 | 227.7 | 283.6 |
| 1972 | 31.0 | * | * | * | * | 4.8 | 11.7 | 40.7 | 104.3 | 213.6 | 237.9 | 277.1 |
| 1971 | 30.7 | * | * | * | 1.2 | 3.6 | 14.3 | 43.2 | 112.4 | 195.1 | 231.3 | 271.6 |
| 1970 | 30.0 | * | * | * | 1.4 | 4.0 | 12.1 | 41.5 | 112.4 | 201.6 | 233.6 | 190.5 |
| 1969 | 31.3 | * | * | * | 1.4 | 3.3 | 13.8 | 43.0 | 121.3 | 204.2 | 249.8 | 223.3 |
| 1968 | 31.6 | * | * | * | 1.2 | 5.3 | 13.8 | 44.3 | 122.3 | 201.6 | 249.1 | 262.5 |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| AVITAMINOSES AND OTHER NUTRITIONAL DEFICIENCIES (260-269) | | | | | | | | | | | | |
| BOTH SEXES | | | | | | | | | | | | |
| 1978 | 1.3 | 2.1 | * | * | 0.0 | 0.1 | 0.1 | 0.4 | 1.0 | 2.7 | 12.4 | 49.6 |
| 1977 | 1.2 | 1.6 | * | * | * | 0.1 | 0.1 | 0.3 | 1.0 | 2.9 | 11.5 | 45.4 |
| 1976 | 1.2 | 2.5 | * | * | 0.1 | 0.1 | 0.1 | 0.4 | 1.0 | 3.2 | 11.8 | 44.5 |
| 1975 | 1.2 | 3.4 | * | * | 0.1 | 0.1 | 0.2 | 0.4 | 0.9 | 3.0 | 11.0 | 49.5 |
| 1974 | 1.2 | 3.2 | * | * | * | 0.1 | 0.2 | 0.4 | 1.2 | 3.3 | 11.3 | 44.8 |
| 1973 | 1.2 | 3.3 | * | * | * | 0.1 | 0.2 | 0.4 | 1.1 | 3.4 | 11.9 | 48.5 |
| 1972 | 1.2 | 3.3 | * | 0.1 | 0.1 | * | 0.2 | 0.5 | 1.1 | 3.3 | 12.1 | 47.3 |
| 1971 | 1.2 | 5.1 | 0.2 | 0.1 | 0.1 | 0.1 | 0.2 | 0.4 | 1.3 | 3.4 | 11.4 | 46.7 |
| 1970 | 1.2 | 4.7 | 0.2 | 0.1 | 0.1 | 0.1 | 0.3 | 0.5 | 1.4 | 3.4 | 11.9 | 41.0 |
| 1969 | 1.3 | 5.8 | 0.3 | 0.1 | 0.1 | 0.1 | 0.2 | 0.6 | 1.3 | 3.5 | 11.1 | 54.2 |
| 1968 | 1.3 | 7.2 | 0.3 | 0.1 | 0.1 | * | 0.2 | 0.5 | 1.3 | 3.3 | 12.3 | 51.9 |
| MALE | | | | | | | | | | | | |
| 1978 | 1.1 | 2.0 | * | * | * | * | * | 0.5 | 1.1 | 3.2 | 13.3 | 51.1 |
| 1977 | 1.0 | 2.0 | * | * | * | * | * | 0.4 | 1.2 | 3.4 | 12.9 | 47.2 |
| 1976 | 1.1 | 2.8 | * | * | * | * | * | 0.4 | 1.0 | 4.1 | 13.4 | 47.0 |
| 1975 | 1.1 | 3.7 | * | * | * | 0.2 | 0.2 | 0.4 | 1.1 | 3.8 | 11.9 | 53.0 |
| 1974 | 1.1 | 4.0 | * | * | * | * | 0.2 | 0.5 | 1.4 | 3.9 | 13.2 | 46.2 |
| 1973 | 1.1 | 3.6 | * | * | * | * | 0.2 | 0.5 | 1.2 | 3.9 | 13.2 | 44.8 |
| 1972 | 1.1 | 3.4 | * | * | * | * | * | 0.6 | 1.3 | 3.8 | 14.7 | 47.5 |
| 1971 | 1.1 | 5.9 | * | * | * | * | 0.2 | 0.5 | 1.6 | 4.2 | 12.3 | 42.7 |
| 1970 | 1.2 | 5.6 | * | * | * | * | 0.3 | 0.5 | 1.5 | 4.0 | 14.0 | 41.1 |
| 1969 | 1.2 | 6.9 | 0.5 | * | * | * | 0.2 | 0.7 | 1.6 | 4.1 | 11.7 | 49.7 |
| 1968 | 1.3 | 8.3 | 0.3 | * | * | * | 0.3 | 0.6 | 1.7 | 4.2 | 14.2 | 51.1 |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| AVITAMINOSES AND OTHER NUTRITIONAL DEFICIENCIES (260-269) | | | | | | | | | | | | |
| FEMALE | | | | | | | | | | | | |
| 1978 | 1.5 | 2.2 | * | * | * | 0.1 | * | 0.4 | 0.9 | 2.4 | 11.8 | 49.0 |
| 1977 | 1.3 | 1.3 | * | * | * | * | * | 0.3 | 0.8 | 2.6 | 10.7 | 44.6 |
| 1976 | 1.3 | 2.2 | * | * | * | * | * | 0.4 | 0.9 | 2.5 | 10.9 | 43.3 |
| 1975 | 1.3 | 3.1 | * | * | * | * | * | 0.5 | 0.8 | 2.4 | 10.4 | 47.8 |
| 1974 | 1.3 | 2.4 | * | * | * | * | 0.2 | 0.4 | 0.9 | 2.8 | 10.1 | 44.2 |
| 1973 | 1.4 | 3.0 | * | * | * | * | 0.3 | 0.4 | 1.1 | 3.0 | 11.1 | 50.4 |
| 1972 | 1.3 | 3.1 | * | * | * | * | 0.3 | 0.5 | 1.0 | 3.0 | 10.5 | 47.3 |
| 1971 | 1.3 | 4.2 | * | * | * | * | 0.2 | 0.4 | 0.9 | 2.8 | 10.9 | 48.8 |
| 1970 | 1.3 | 3.7 | * | * | * | * | 0.2 | 0.5 | 1.2 | 3.0 | 10.5 | 41.0 |
| 1969 | 1.3 | 4.6 | * | * | * | * | 0.2 | 0.5 | 1.0 | 3.1 | 10.6 | 56.7 |
| 1968 | 1.3 | 6.0 | 0.4 | * | * | * | 0.2 | 0.5 | 0.9 | 2.7 | 10.9 | 52.4 |
| WHITE, BOTH SEXES | | | | | | | | | | | | |
| 1978 | 1.3 | 1.3 | * | * | * | 0.1 | 0.1 | 0.3 | 0.9 | 2.4 | 12.0 | 48.1 |
| 1977 | 1.2 | 1.2 | * | * | * | 0.1 | * | 0.3 | 0.9 | 2.7 | 11.0 | 44.7 |
| 1976 | 1.2 | 1.5 | * | * | 0.1 | * | 0.1 | 0.4 | 0.9 | 2.9 | 10.9 | 43.2 |
| 1975 | 1.2 | 1.9 | * | * | * | 0.1 | 0.1 | 0.4 | 0.8 | 2.6 | 10.3 | 48.1 |
| 1974 | 1.1 | 2.1 | * | * | * | 0.1 | 0.2 | 0.4 | 1.0 | 3.0 | 10.7 | 44.6 |
| 1973 | 1.2 | 2.3 | * | * | * | 0.1 | 0.2 | 0.4 | 1.0 | 3.0 | 11.4 | 47.8 |
| 1972 | 1.2 | 1.9 | * | * | 0.1 | * | 0.1 | 0.5 | 1.0 | 2.9 | 11.4 | 47.5 |
| 1971 | 1.2 | 3.0 | * | 0.1 | 0.1 | * | 0.2 | 0.3 | 1.1 | 3.1 | 10.8 | 47.0 |
| 1970 | 1.2 | 2.8 | * | * | * | 0.1 | 0.2 | 0.4 | 1.2 | 3.1 | 11.4 | 41.8 |
| 1969 | 1.2 | 3.1 | 0.2 | * | * | * | 0.2 | 0.4 | 1.1 | 3.0 | 10.6 | 54.9 |
| 1968 | 1.2 | 3.8 | 0.2 | * | * | * | 0.2 | 0.5 | 1.1 | 2.9 | 11.9 | 51.4 |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| AVITAMINOSES AND OTHER NUTRITIONAL DEFICIENCIES (260-269) | | | | | | | | | | | | |
| WHITE, MALE | | | | | | | | | | | | |
| 1978 | 1.0 | * | * | * | * | * | * | 0.3 | 1.0 | 2.8 | 13.1 | 48.5 |
| 1977 | 1.0 | * | * | * | * | * | * | 0.3 | 1.0 | 3.1 | 12.2 | 45.9 |
| 1976 | 1.0 | * | * | * | * | * | * | 0.3 | 0.8 | 3.6 | 12.1 | 45.4 |
| 1975 | 1.0 | 2.2 | * | * | * | 0.2 | * | 0.4 | 1.0 | 3.2 | 11.3 | 51.9 |
| 1974 | 1.0 | 2.4 | * | * | * | * | * | 0.4 | 1.2 | 3.4 | 12.1 | 45.4 |
| 1973 | 1.0 | 2.8 | * | * | * | * | * | 0.3 | 1.0 | 3.4 | 12.7 | 43.9 |
| 1972 | 1.0 | 1.8 | * | * | * | * | * | 0.5 | 1.1 | 3.4 | 13.1 | 48.0 |
| 1971 | 1.0 | 3.1 | * | * | * | * | * | 0.3 | 1.4 | 3.9 | 11.6 | 42.1 |
| 1970 | 1.1 | 3.2 | * | * | * | * | 0.2 | 0.4 | 1.3 | 3.5 | 13.5 | 42.1 |
| 1969 | 1.1 | 4.2 | * | * | * | * | * | 0.5 | 1.4 | 3.6 | 11.0 | 51.1 |
| 1968 | 1.1 | 4.4 | * | * | * | * | 0.2 | 0.5 | 1.5 | 3.6 | 13.6 | 48.1 |
| WHITE, FEMALE | | | | | | | | | | | | |
| 1978 | 1.5 | * | * | * | * | * | * | 0.3 | 0.8 | 2.1 | 11.4 | 47.9 |
| 1977 | 1.4 | * | * | * | * | * | * | 0.3 | 0.8 | 2.3 | 10.3 | 44.1 |
| 1976 | 1.3 | 1.6 | * | * | * | * | * | 0.4 | 0.9 | 2.3 | 10.1 | 42.2 |
| 1975 | 1.3 | 1.6 | * | * | * | * | * | 0.5 | 0.7 | 2.1 | 9.7 | 46.4 |
| 1974 | 1.3 | 1.8 | * | * | * | * | * | 0.3 | 0.8 | 2.6 | 9.9 | 44.2 |
| 1973 | 1.4 | 1.9 | * | * | * | * | 0.2 | 0.4 | 0.9 | 2.8 | 10.6 | 49.8 |
| 1972 | 1.3 | 1.9 | * | * | * | * | 0.3 | 0.4 | 0.9 | 2.6 | 10.4 | 47.2 |
| 1971 | 1.3 | 2.8 | * | * | * | * | * | 0.3 | 0.9 | 2.5 | 10.3 | 49.6 |
| 1970 | 1.2 | 2.4 | * | * | * | * | * | 0.5 | 1.2 | 2.7 | 9.9 | 41.6 |
| 1969 | 1.3 | 2.1 | * | * | * | * | * | 0.4 | 0.9 | 2.6 | 10.3 | 57.0 |
| 1968 | 1.2 | 3.2 | * | * | * | * | 0.2 | 0.5 | 0.7 | 2.3 | 10.8 | 53.2 |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| AVITAMINOSES AND OTHER NUTRITIONAL DEFICIENCIES (260-269) | | | | | | | | | | | | |
| ALL OTHER, BOTH SEXES | | | | | | | | | | | | |
| 1978 | 1.5 | 6.0 | * | * | * | * | * | 1.1 | 2.0 | 5.8 | 15.7 | 67.3 |
| 1977 | 1.3 | 3.7 | * | * | * | * | * | * | 1.5 | 5.6 | 16.5 | 54.4 |
| 1976 | 1.5 | 7.6 | * | * | * | * | * | 0.8 | 1.8 | 6.6 | 21.8 | 59.5 |
| 1975 | 1.6 | 10.7 | * | * | * | * | * | * | 2.0 | 6.7 | 18.0 | 65.1 |
| 1974 | 1.4 | 8.8 | * | * | * | * | * | 1.0 | 2.7 | 6.2 | 17.8 | 47.4 |
| 1973 | 1.5 | 8.2 | * | * | * | * | * | 0.8 | 2.8 | 6.8 | 17.9 | 56.7 |
| 1972 | 1.6 | 10.4 | * | * | * | * | 0.8 | 1.1 | 2.4 | 7.3 | 20.4 | 45.5 |
| 1971 | 1.7 | 16.2 | * | * | * | * | * | 1.3 | 2.4 | 6.7 | 18.3 | 43.3 |
| 1970 | 1.6 | 14.7 | * | * | * | * | 0.9 | 1.2 | 2.6 | 6.8 | 18.4 | 33.6 |
| 1969 | 1.9 | 20.6 | 1.2 | * | * | * | * | 1.5 | 2.7 | 8.7 | 17.5 | 46.8 |
| 1968 | 2.0 | 25.3 | 0.9 | * | * | * | * | 1.2 | 3.1 | 8.1 | 16.7 | 57.8 |
| ALL OTHER, MALE | | | | | | | | | | | | |
| 1978 | 1.5 | * | * | * | * | * | * | 1.7 | 2.3 | 7.3 | 14.7 | 77.6 |
| 1977 | 1.4 | * | * | * | * | * | * | * | 2.6 | 6.5 | 19.5 | 60.7 |
| 1976 | 1.7 | 9.8 | * | * | * | * | * | * | 2.4 | 8.9 | 25.3 | 63.6 |
| 1975 | 1.6 | 11.3 | * | * | * | * | * | * | 2.6 | 9.0 | 17.8 | 64.8 |
| 1974 | 1.7 | 12.0 | * | * | * | * | * | * | 3.5 | 8.1 | 24.5 | 54.0 |
| 1973 | 1.5 | 7.9 | * | * | * | * | * | * | 2.7 | 8.4 | 19.2 | 54.2 |
| 1972 | 1.8 | 11.7 | * | * | * | * | * | * | 3.3 | 7.5 | 32.3 | 42.6 |
| 1971 | 1.9 | 20.6 | * | * | * | * | * | * | 3.7 | 7.6 | 19.3 | 48.9 |
| 1970 | 1.8 | 18.8 | * | * | * | * | * | * | 3.5 | 8.6 | 19.7 | * |
| 1969 | 2.1 | 22.9 | * | * | * | * | * | 1.9 | 3.6 | 9.5 | 20.2 | * |
| 1968 | 2.4 | 29.4 | * | * | * | * | * | * | 3.8 | 10.0 | 21.5 | 80.5 |
| ALL OTHER, FEMALE | | | | | | | | | | | | |
| 1978 | 1.4 | * | * | * | * | * | * | * | 1.8 | 4.8 | 16.5 | 62.4 |
| 1977 | 1.1 | * | * | * | * | * | * | * | * | 4.8 | 14.4 | 51.0 |
| 1976 | 1.4 | * | * | * | * | * | * | * | * | 4.8 | 19.4 | 57.1 |
| 1975 | 1.5 | 10.0 | * | * | * | * | * | * | * | 4.8 | 18.1 | 64.5 |
| 1974 | 1.2 | * | * | * | * | * | * | * | 1.9 | 4.7 | 13.3 | 43.5 |
| 1973 | 1.5 | 8.5 | * | * | * | * | * | * | 2.9 | 5.5 | 17.0 | 58.2 |
| 1972 | 1.3 | 9.0 | * | * | * | * | * | * | * | 7.0 | 12.0 | 47.4 |
| 1971 | 1.5 | 11.7 | * | * | * | * | * | * | * | 5.9 | 17.5 | 39.7 |
| 1970 | 1.4 | 10.6 | * | * | * | * | * | * | * | 5.2 | 17.3 | 34.3 |
| 1969 | 1.7 | 18.4 | * | * | * | * | * | * | * | 8.1 | 15.4 | 53.8 |
| 1968 | 1.7 | 21.0 | * | * | * | * | * | * | 2.6 | 6.5 | 13.0 | 42.6 |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| AVITAMINOSES AND OTHER NUTRITIONAL DEFICIENCIES (260-269) | | | | | | | | | | | | |
| BLACK, BOTH SEXES | | | | | | | | | | | | |
| 1978 | 1.5 | 6.9 | * | * | * | * | * | 1.2 | 2.1 | 6.4 | 15.1 | 66.0 |
| 1977 | 1.3 | * | * | * | * | * | * | * | 1.7 | 5.8 | 15.3 | 53.1 |
| 1976 | 1.7 | 8.4 | * | * | * | * | * | 0.9 | 1.9 | 7.2 | 21.3 | 59.6 |
| 1975 | 1.7 | 11.4 | * | * | * | * | * | * | 2.2 | 7.0 | 17.6 | 65.4 |
| 1974 | 1.5 | 9.2 | * | * | * | * | * | 1.2 | 2.9 | 6.4 | 17.6 | 47.1 |
| 1973 | 1.5 | 8.7 | * | * | * | * | * | 0.9 | 3.0 | 6.7 | 17.4 | 48.2 |
| 1972 | 1.6 | 11.3 | * | * | * | * | 0.9 | 1.2 | 2.4 | 7.3 | 20.6 | 41.8 |
| 1971 | 1.8 | 17.0 | * | * | * | * | * | 1.4 | 2.5 | 7.1 | 18.9 | 43.9 |
| 1970 | 1.6 | 14.8 | * | * | * | * | 1.0 | 1.2 | 2.7 | 7.2 | 18.1 | 32.4 |
| 1969 | 2.0 | 23.0 | 1.1 | * | * | * | * | 1.6 | 2.9 | 9.2 | 17.6 | 45.4 |
| 1968 | 2.0 | 27.1 | * | * | * | * | * | 1.3 | 3.0 | 7.7 | 17.6 | 48.9 |
| BLACK, MALE | | | | | | | | | | | | |
| 1978 | 1.6 | * | * | * | * | * | * | 2.0 | 2.6 | 8.0 | 14.2 | 80.4 |
| 1977 | 1.5 | * | * | * | * | * | * | * | 2.9 | 6.9 | 18.7 | 55.1 |
| 1976 | 1.8 | 10.9 | * | * | * | * | * | * | 2.5 | 9.8 | 23.5 | 66.0 |
| 1975 | 1.7 | 11.7 | * | * | * | * | * | * | 2.8 | 9.4 | 17.6 | 67.4 |
| 1974 | 1.7 | 12.4 | * | * | * | * | * | * | 3.9 | 8.6 | 24.9 | 53.5 |
| 1973 | 1.5 | 8.6 | * | * | * | * | * | * | 2.8 | 8.5 | 19.1 | * |
| 1972 | 1.8 | 12.4 | * | * | * | * | * | * | 3.4 | 7.7 | 32.5 | * |
| 1971 | 2.0 | 21.2 | * | * | * | * | * | * | 3.9 | 7.9 | 19.9 | 50.0 |
| 1970 | 1.9 | 18.8 | * | * | * | * | * | * | 3.7 | 9.5 | 19.6 | * |
| 1969 | 2.2 | 25.3 | * | * | * | * | * | 2.1 | 4.0 | 9.9 | 20.4 | * |
| 1968 | 2.4 | 31.1 | * | * | * | * | * | * | 3.6 | 9.7 | 23.0 | 65.7 |
| BLACK, FEMALE | | | | | | | | | | | | |
| 1978 | 1.5 | * | * | * | * | * | * | * | * | 5.2 | 15.6 | 58.6 |
| 1977 | 1.2 | * | * | * | * | * | * | * | * | 4.9 | 13.0 | 52.1 |
| 1976 | 1.5 | * | * | * | * | * | * | * | * | 5.2 | 19.8 | 56.2 |
| 1975 | 1.6 | 11.1 | * | * | * | * | * | * | * | 5.1 | 17.7 | 64.3 |
| 1974 | 1.3 | * | * | * | * | * | * | * | 2.1 | 4.8 | 12.6 | 43.6 |
| 1973 | 1.5 | 8.7 | * | * | * | * | * | * | 3.2 | 5.3 | 16.2 | 52.1 |
| 1972 | 1.4 | 10.1 | * | * | * | * | * | * | * | 7.0 | 12.1 | 45.7 |
| 1971 | 1.6 | 12.8 | * | * | * | * | * | * | * | 6.4 | 18.1 | 40.3 |
| 1970 | 1.4 | 10.7 | * | * | * | * | * | * | * | 5.3 | 17.0 | 32.2 |
| 1969 | 1.8 | 20.6 | * | * | * | * | * | * | * | 8.6 | 15.4 | 53.3 |
| 1968 | 1.7 | 23.0 | * | * | * | * | * | * | 2.5 | 6.0 | 13.5 | 39.3 |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| ANEMIAS (280-285) | | | | | | | | | | | | |
| BOTH SEXES | | | | | | | | | | | | |
| 1978 | 1.5 | 0.7 | 0.6 | 0.2 | 0.3 | 0.4 | 0.4 | 0.8 | 1.5 | 3.9 | 12.9 | 34.4 |
| 1977 | 1.4 | 1.0 | 0.4 | 0.3 | 0.4 | 0.3 | 0.4 | 0.7 | 1.6 | 4.1 | 12.1 | 31.3 |
| 1976 | 1.5 | 1.4 | 0.4 | 0.3 | 0.3 | 0.4 | 0.5 | 0.7 | 1.5 | 4.2 | 12.9 | 31.6 |
| 1975 | 1.5 | 1.5 | 0.5 | 0.3 | 0.4 | 0.5 | 0.5 | 0.7 | 1.6 | 4.6 | 12.1 | 33.4 |
| 1974 | 1.6 | 1.0 | 0.7 | 0.3 | 0.3 | 0.5 | 0.5 | 0.8 | 1.9 | 4.8 | 13.6 | 32.0 |
| 1973 | 1.6 | 1.8 | 0.6 | 0.3 | 0.4 | 0.5 | 0.5 | 0.8 | 1.8 | 4.6 | 13.6 | 34.1 |
| 1972 | 1.6 | 2.1 | 0.7 | 0.3 | 0.5 | 0.5 | 0.6 | 0.9 | 2.0 | 4.8 | 14.0 | 33.2 |
| 1971 | 1.7 | 1.9 | 0.7 | 0.3 | 0.6 | 0.5 | 0.8 | 1.0 | 2.0 | 5.1 | 13.4 | 34.0 |
| 1970 | 1.7 | 1.7 | 0.8 | 0.3 | 0.5 | 0.5 | 0.6 | 1.0 | 2.0 | 5.4 | 15.1 | 32.8 |
| 1969 | 1.6 | 1.8 | 0.8 | 0.4 | 0.5 | 0.6 | 0.5 | 0.9 | 2.0 | 5.2 | 14.3 | 35.0 |
| 1968 | 1.8 | 1.8 | 0.9 | 0.4 | 0.5 | 0.6 | 0.6 | 0.9 | 2.3 | 5.6 | 15.6 | 35.7 |
| MALE | | | | | | | | | | | | |
| 1978 | 1.4 | * | 0.6 | 0.2 | 0.3 | 0.4 | 0.4 | 0.8 | 1.5 | 4.6 | 14.5 | 41.6 |
| 1977 | 1.4 | * | 0.4 | 0.3 | 0.4 | 0.4 | 0.4 | 0.7 | 1.8 | 4.9 | 14.6 | 37.6 |
| 1976 | 1.4 | 1.4 | 0.5 | 0.3 | 0.4 | 0.4 | 0.3 | 0.7 | 1.6 | 4.9 | 15.9 | 36.5 |
| 1975 | 1.4 | 1.4 | 0.5 | 0.3 | 0.4 | 0.4 | 0.4 | 0.7 | 1.8 | 5.8 | 13.5 | 37.2 |
| 1974 | 1.5 | * | 0.7 | 0.3 | 0.4 | 0.5 | 0.5 | 0.8 | 2.2 | 5.3 | 16.1 | 38.1 |
| 1973 | 1.5 | 2.2 | 0.7 | 0.4 | 0.5 | 0.5 | 0.6 | 0.8 | 2.1 | 5.1 | 15.5 | 40.9 |
| 1972 | 1.6 | 2.4 | 0.7 | 0.4 | 0.5 | 0.6 | 0.5 | 0.8 | 2.2 | 5.7 | 15.7 | 38.3 |
| 1971 | 1.6 | 1.8 | 0.6 | 0.3 | 0.5 | 0.5 | 0.6 | 1.0 | 1.9 | 5.9 | 15.2 | 38.2 |
| 1970 | 1.6 | 1.6 | 0.7 | 0.3 | 0.6 | 0.5 | 0.5 | 1.0 | 2.0 | 6.2 | 17.6 | 39.1 |
| 1969 | 1.6 | 1.7 | 0.8 | 0.4 | 0.6 | 0.6 | 0.5 | 0.8 | 2.1 | 5.8 | 16.6 | 35.5 |
| 1968 | 1.7 | 1.7 | 1.1 | 0.4 | 0.6 | 0.6 | 0.5 | 0.9 | 2.3 | 6.5 | 17.5 | 38.2 |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| ANEMIAS (280-285) | | | | | | | | | | | | |
| FEMALE | | | | | | | | | | | | |
| 1978 | 1.6 | * | 0.6 | 0.3 | 0.3 | 0.4 | 0.3 | 0.8 | 1.6 | 3.4 | 11.9 | 31.2 |
| 1977 | 1.5 | * | 0.4 | 0.2 | 0.4 | 0.3 | 0.4 | 0.6 | 1.4 | 3.5 | 10.6 | 28.4 |
| 1976 | 1.5 | 1.4 | 0.4 | 0.3 | 0.3 | 0.4 | 0.7 | 0.7 | 1.5 | 3.7 | 11.1 | 29.4 |
| 1975 | 1.5 | 1.5 | 0.5 | 0.2 | 0.3 | 0.5 | 0.5 | 0.7 | 1.5 | 3.6 | 11.2 | 31.5 |
| 1974 | 1.6 | * | 0.6 | 0.3 | 0.3 | 0.4 | 0.5 | 0.8 | 1.7 | 4.3 | 12.0 | 29.0 |
| 1973 | 1.6 | 1.4 | 0.6 | 0.3 | 0.4 | 0.5 | 0.5 | 0.9 | 1.6 | 4.3 | 12.4 | 30.7 |
| 1972 | 1.7 | 1.9 | 0.7 | 0.3 | 0.5 | 0.5 | 0.7 | 1.1 | 1.8 | 4.2 | 12.9 | 30.6 |
| 1971 | 1.8 | 1.9 | 0.9 | 0.3 | 0.6 | 0.5 | 0.9 | 1.0 | 2.2 | 4.5 | 12.1 | 31.7 |
| 1970 | 1.7 | 1.8 | 0.8 | 0.3 | 0.5 | 0.6 | 0.8 | 0.9 | 1.9 | 4.8 | 13.5 | 29.3 |
| 1969 | 1.7 | 2.0 | 0.8 | 0.4 | 0.5 | 0.5 | 0.5 | 1.0 | 1.9 | 4.7 | 12.8 | 34.8 |
| 1968 | 1.8 | 1.8 | 0.8 | 0.4 | 0.5 | 0.6 | 0.7 | 1.0 | 2.3 | 4.8 | 14.3 | 34.4 |
| WHITE, BOTH SEXES | | | | | | | | | | | | |
| 1978 | 1.4 | * | * | 0.2 | 0.2 | 0.2 | 0.2 | 0.5 | 1.4 | 3.6 | 12.8 | 35.1 |
| 1977 | 1.3 | * | * | 0.2 | 0.2 | 0.1 | 0.2 | 0.5 | 1.4 | 3.8 | 12.3 | 31.8 |
| 1976 | 1.3 | * | * | 0.2 | 0.2 | 0.2 | 0.2 | 0.5 | 1.4 | 4.0 | 12.9 | 31.8 |
| 1975 | 1.4 | 1.0 | * | 0.2 | 0.2 | 0.3 | 0.2 | 0.5 | 1.5 | 4.3 | 12.1 | 34.3 |
| 1974 | 1.4 | * | 0.3 | 0.2 | 0.2 | 0.2 | 0.3 | 0.6 | 1.7 | 4.4 | 13.5 | 32.0 |
| 1973 | 1.4 | 0.8 | 0.3 | 0.2 | 0.3 | 0.2 | 0.2 | 0.6 | 1.6 | 4.4 | 13.9 | 35.1 |
| 1972 | 1.4 | 0.9 | 0.3 | 0.2 | 0.3 | 0.2 | 0.3 | 0.7 | 1.7 | 4.5 | 14.3 | 33.5 |
| 1971 | 1.5 | 1.1 | 0.3 | 0.2 | 0.3 | 0.2 | 0.4 | 0.8 | 1.8 | 4.9 | 13.6 | 34.5 |
| 1970 | 1.5 | 0.9 | 0.3 | 0.2 | 0.3 | 0.2 | 0.3 | 0.7 | 1.8 | 5.2 | 15.1 | 33.6 |
| 1969 | 1.5 | 1.0 | 0.4 | 0.2 | 0.3 | 0.3 | 0.2 | 0.7 | 1.8 | 5.1 | 14.6 | 36.6 |
| 1968 | 1.6 | 1.0 | 0.3 | 0.3 | 0.3 | 0.2 | 0.3 | 0.7 | 2.0 | 5.3 | 15.8 | 36.4 |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| ANEMIAS (280-285) | | | | | | | | | | | | |
| WHITE, MALE | | | | | | | | | | | | |
| 1978 | 1.3 | * | * | 0.2 | 0.2 | 0.2 | 0.2 | 0.5 | 1.4 | 4.2 | 14.6 | 43.6 |
| 1977 | 1.3 | * | * | 0.2 | 0.2 | 0.2 | 0.2 | 0.5 | 1.5 | 4.6 | 15.0 | 39.4 |
| 1976 | 1.3 | * | * | 0.2 | 0.2 | 0.2 | * | 0.5 | 1.4 | 4.6 | 16.2 | 37.2 |
| 1975 | 1.3 | * | * | 0.2 | 0.2 | 0.2 | * | 0.5 | 1.6 | 5.5 | 13.8 | 38.7 |
| 1974 | 1.3 | * | * | 0.2 | 0.2 | 0.2 | 0.3 | 0.6 | 2.0 | 4.9 | 16.1 | 38.1 |
| 1973 | 1.3 | * | * | 0.2 | 0.3 | 0.2 | 0.2 | 0.6 | 1.8 | 4.9 | 16.0 | 42.0 |
| 1972 | 1.4 | * | * | 0.3 | 0.3 | 0.2 | 0.2 | 0.6 | 2.0 | 5.3 | 16.2 | 36.2 |
| 1971 | 1.4 | * | * | 0.2 | 0.3 | 0.2 | 0.3 | 0.7 | 1.7 | 5.7 | 15.7 | 39.3 |
| 1970 | 1.5 | * | * | 0.2 | 0.3 | 0.2 | 0.2 | 0.8 | 1.8 | 6.0 | 17.6 | 41.3 |
| 1969 | 1.4 | * | 0.5 | 0.2 | 0.3 | 0.3 | 0.2 | 0.7 | 1.9 | 5.7 | 16.9 | 37.8 |
| 1968 | 1.5 | * | 0.4 | 0.2 | 0.3 | 0.2 | 0.3 | 0.7 | 2.1 | 6.2 | 17.8 | 39.1 |
| WHITE, FEMALE | | | | | | | | | | | | |
| 1978 | 1.5 | * | * | 0.2 | 0.2 | 0.1 | * | 0.6 | 1.3 | 3.2 | 11.7 | 31.3 |
| 1977 | 1.3 | * | * | 0.2 | 0.2 | * | * | 0.4 | 1.2 | 3.2 | 10.6 | 28.3 |
| 1976 | 1.4 | * | * | 0.2 | 0.1 | 0.2 | 0.2 | 0.5 | 1.3 | 3.4 | 11.0 | 29.4 |
| 1975 | 1.4 | * | * | 0.2 | 0.2 | 0.3 | 0.3 | 0.5 | 1.3 | 3.3 | 11.0 | 32.2 |
| 1974 | 1.5 | * | * | 0.2 | 0.1 | 0.2 | 0.3 | 0.6 | 1.5 | 4.0 | 11.9 | 29.0 |
| 1973 | 1.5 | * | * | 0.2 | 0.2 | 0.2 | 0.2 | 0.6 | 1.4 | 4.0 | 12.6 | 31.7 |
| 1972 | 1.5 | * | * | 0.1 | 0.2 | 0.2 | 0.4 | 0.7 | 1.4 | 3.9 | 13.2 | 32.2 |
| 1971 | 1.6 | * | 0.4 | 0.2 | 0.3 | 0.2 | 0.5 | 0.8 | 1.9 | 4.3 | 12.3 | 32.1 |
| 1970 | 1.5 | * | 0.4 | 0.1 | 0.2 | 0.2 | 0.4 | 0.6 | 1.7 | 4.5 | 13.5 | 29.3 |
| 1969 | 1.6 | * | 0.3 | 0.3 | 0.3 | 0.2 | 0.2 | 0.8 | 1.7 | 4.6 | 13.0 | 36.0 |
| 1968 | 1.6 | * | * | 0.3 | 0.3 | 0.2 | 0.4 | 0.7 | 1.9 | 4.6 | 14.5 | 35.0 |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| ANEMIAS (280-285) | | | | | | | | | | | | |
| ALL OTHER, BOTH SEXES | | | | | | | | | | | | |
| 1978 | 2.2 | * | 2.7 | 0.6 | 0.9 | 1.8 | 1.8 | 2.5 | 3.3 | 6.1 | 14.0 | 26.8 |
| 1977 | 2.2 | * | 1.9 | 0.6 | 1.4 | 1.6 | 1.5 | 2.1 | 3.4 | 7.1 | 10.6 | 26.3 |
| 1976 | 2.4 | 5.3 | 1.9 | 0.6 | 1.2 | 1.8 | 2.5 | 2.4 | 3.2 | 6.9 | 13.1 | 29.4 |
| 1975 | 2.3 | * | 2.2 | 0.9 | 1.3 | 1.8 | 2.0 | 2.1 | 3.4 | 7.6 | 12.3 | 23.3 |
| 1974 | 2.6 | * | 2.6 | 0.9 | 1.3 | 2.3 | 2.0 | 2.6 | 3.7 | 8.4 | 14.4 | 32.6 |
| 1973 | 2.6 | 6.5 | 2.4 | 0.9 | 1.5 | 2.6 | 2.7 | 2.8 | 3.7 | 7.3 | 10.7 | 22.0 |
| 1972 | 3.0 | 8.5 | 2.9 | 1.0 | 2.1 | 3.2 | 2.3 | 3.2 | 4.6 | 7.9 | 10.0 | 29.3 |
| 1971 | 3.1 | 6.0 | 2.9 | 1.0 | 2.2 | 2.9 | 3.4 | 3.3 | 4.6 | 7.3 | 10.5 | 27.5 |
| 1970 | 3.0 | 5.8 | 3.2 | 1.1 | 2.0 | 2.8 | 2.9 | 3.4 | 3.9 | 8.1 | 15.2 | 25.4 |
| 1969 | 2.8 | 6.2 | 2.7 | 1.5 | 2.3 | 2.9 | 2.4 | 2.5 | 4.0 | 6.3 | 10.8 | * |
| 1968 | 3.2 | 6.1 | 4.0 | 1.1 | 2.0 | 3.2 | 2.5 | 3.2 | 5.0 | 8.5 | 12.7 | 27.5 |
| ALL OTHER, MALE | | | | | | | | | | | | |
| 1978 | 2.2 | * | 2.9 | * | 0.9 | 1.8 | 2.0 | 3.0 | 2.8 | 7.4 | 14.3 | * |
| 1977 | 2.2 | * | 1.9 | 0.7 | 1.6 | 2.0 | 1.5 | 2.2 | 3.8 | 7.8 | 11.1 | * |
| 1976 | 2.3 | * | 2.2 | * | 1.6 | 1.9 | * | 2.7 | 3.3 | 7.9 | 13.6 | * |
| 1975 | 2.4 | * | 2.3 | 1.2 | 1.4 | 2.2 | 2.3 | 1.9 | 4.0 | 8.5 | 10.3 | * |
| 1974 | 2.7 | * | 2.8 | 0.9 | 1.5 | 2.7 | 2.2 | 2.5 | 4.2 | 9.3 | 15.7 | * |
| 1973 | 2.8 | 8.3 | 2.7 | 1.0 | 1.7 | 2.8 | 3.0 | 2.5 | 4.8 | 6.6 | 11.2 | * |
| 1972 | 3.0 | 10.3 | 3.1 | 0.8 | 1.7 | 3.6 | 2.5 | 2.8 | 4.0 | 8.7 | 10.1 | 59.6 |
| 1971 | 3.0 | * | 2.6 | 1.3 | 2.2 | 3.1 | 3.5 | 3.4 | 3.9 | 7.8 | 10.6 | * |
| 1970 | 3.0 | * | 3.2 | 1.2 | 2.0 | 2.6 | 2.5 | 3.2 | 4.2 | 8.4 | 17.6 | * |
| 1969 | 2.9 | * | 2.6 | 1.6 | 2.4 | 3.5 | 2.6 | 1.9 | 4.3 | 6.5 | 12.8 | * |
| 1968 | 3.2 | * | 4.5 | 1.2 | 2.0 | 3.5 | 2.4 | 3.2 | 4.4 | 9.4 | 14.0 | * |
| ALL OTHER, FEMALE | | | | | | | | | | | | |
| 1978 | 2.2 | * | 2.5 | * | 0.9 | 1.8 | 1.6 | 2.1 | 3.6 | 5.1 | 13.9 | 30.3 |
| 1977 | 2.1 | * | * | * | 1.3 | 1.3 | 1.6 | 2.0 | 3.1 | 6.6 | 10.2 | 29.8 |
| 1976 | 2.4 | * | * | * | 0.9 | 1.8 | 3.4 | 2.1 | 3.1 | 6.1 | 12.8 | 29.6 |
| 1975 | 2.2 | * | 2.0 | * | 1.2 | 1.6 | 1.8 | 2.3 | 2.8 | 6.9 | 13.6 | 23.7 |
| 1974 | 2.4 | * | 2.3 | 0.9 | 1.0 | 1.9 | 1.7 | 2.8 | 3.3 | 7.6 | 13.6 | 29.4 |
| 1973 | 2.5 | * | 2.1 | 0.8 | 1.4 | 2.3 | 2.4 | 3.0 | 2.7 | 7.9 | 10.4 | * |
| 1972 | 3.0 | * | 2.7 | 1.1 | 2.6 | 2.9 | 2.1 | 3.6 | 5.1 | 7.3 | 10.0 | * |
| 1971 | 3.1 | * | 3.3 | 0.7 | 2.3 | 2.7 | 3.4 | 3.2 | 5.2 | 6.9 | 10.4 | 27.4 |
| 1970 | 3.1 | * | 3.1 | 0.9 | 1.9 | 2.9 | 3.2 | 3.6 | 3.7 | 7.7 | 13.4 | 29.2 |
| 1969 | 2.8 | * | 2.9 | 1.3 | 2.1 | 2.5 | 2.3 | 3.0 | 3.7 | 6.2 | 9.3 | * |
| 1968 | 3.1 | * | 3.5 | 1.1 | 1.9 | 2.9 | 2.6 | 3.3 | 5.5 | 7.8 | 11.7 | * |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| ANEMIAS (280-285) | | | | | | | | | | | | |
| BLACK, BOTH SEXES | | | | | | | | | | | | |
| 1978 | 2.5 | * | 3.1 | 0.6 | 1.0 | 2.1 | 2.0 | 2.9 | 3.5 | 6.4 | 14.6 | 28.7 |
| 1977 | 2.4 | * | 2.1 | 0.6 | 1.6 | 1.9 | 1.8 | 2.4 | 3.5 | 7.4 | 11.4 | 25.9 |
| 1976 | 2.6 | 6.2 | 2.2 | 0.7 | 1.4 | 2.2 | 2.9 | 2.7 | 3.5 | 7.4 | 13.4 | 30.9 |
| 1975 | 2.6 | * | 2.4 | 0.9 | 1.5 | 2.1 | 2.4 | 2.3 | 3.7 | 7.8 | 12.2 | 26.2 |
| 1974 | 2.8 | * | 2.8 | 1.0 | 1.4 | 2.7 | 2.3 | 2.9 | 4.0 | 8.8 | 15.3 | 33.1 |
| 1973 | 2.9 | 7.2 | 2.7 | 1.0 | 1.7 | 3.0 | 3.1 | 3.1 | 4.1 | 8.0 | 11.2 | 22.8 |
| 1972 | 3.2 | 9.2 | 3.1 | 1.1 | 2.3 | 3.7 | 2.7 | 3.6 | 4.8 | 7.6 | 10.1 | 27.3 |
| 1971 | 3.3 | 6.3 | 3.1 | 1.1 | 2.5 | 3.4 | 3.9 | 3.6 | 4.8 | 7.6 | 11.2 | 29.0 |
| 1970 | 3.3 | 6.2 | 3.4 | 1.2 | 2.2 | 3.2 | 3.3 | 3.6 | 4.3 | 8.5 | 16.3 | 26.4 |
| 1969 | 3.0 | 7.0 | 2.9 | 1.5 | 2.5 | 3.3 | 2.6 | 2.7 | 4.4 | 6.7 | 11.2 | * |
| 1968 | 3.4 | 6.3 | 4.3 | 1.2 | 2.1 | 3.5 | 2.8 | 3.5 | 5.4 | 9.2 | 12.7 | 29.3 |
| BLACK, MALE | | | | | | | | | | | | |
| 1978 | 2.4 | * | 3.3 | * | 1.0 | 2.1 | 2.2 | 3.4 | 3.0 | 7.8 | 14.2 | * |
| 1977 | 2.5 | * | 2.2 | 0.7 | 1.8 | 2.5 | 1.8 | 2.5 | 3.8 | 8.0 | 11.9 | * |
| 1976 | 2.6 | * | 2.5 | * | 1.8 | 2.3 | * | 2.9 | 3.7 | 8.3 | 14.1 | * |
| 1975 | 2.7 | * | 2.7 | 1.3 | 1.6 | 2.6 | 2.7 | 2.1 | 4.5 | 8.4 | 10.4 | * |
| 1974 | 3.0 | * | 3.1 | 0.9 | 1.7 | 3.2 | 2.6 | 2.6 | 4.5 | 10.6 | 17.2 | * |
| 1973 | 3.1 | 9.0 | 3.0 | 1.2 | 1.8 | 3.3 | 3.5 | 2.8 | 5.4 | 7.3 | 11.3 | * |
| 1972 | 3.3 | 11.6 | 3.4 | 0.9 | 1.8 | 4.2 | 2.9 | 3.2 | 4.2 | 9.4 | 10.2 | 60.0 |
| 1971 | 3.3 | * | 2.9 | 1.3 | 2.4 | 3.6 | 4.0 | 3.8 | 4.3 | 8.6 | 11.8 | * |
| 1970 | 3.3 | * | 3.5 | 1.4 | 2.3 | 3.1 | 2.9 | 3.4 | 4.6 | 9.3 | 19.0 | * |
| 1969 | 3.1 | * | 2.8 | 1.6 | 2.7 | 4.0 | 2.7 | 2.1 | 4.8 | 6.8 | 13.8 | * |
| 1968 | 3.5 | * | 4.8 | 1.2 | 2.3 | 3.9 | 2.7 | 3.6 | 4.7 | 10.2 | 15.2 | * |
| BLACK, FEMALE | | | | | | | | | | | | |
| 1978 | 2.5 | * | 2.9 | * | 1.0 | 2.1 | 1.9 | 2.5 | 3.8 | 5.4 | 14.8 | 31.3 |
| 1977 | 2.3 | * | * | * | 1.5 | 1.4 | 1.8 | 2.4 | 3.2 | 7.0 | 11.0 | 29.8 |
| 1976 | 2.7 | * | * | * | 1.0 | 2.1 | 4.0 | 2.4 | 3.3 | 6.7 | 12.9 | 30.3 |
| 1975 | 2.4 | * | 2.2 | * | 1.4 | 1.7 | 2.2 | 2.5 | 3.0 | 7.4 | 13.4 | 26.2 |
| 1974 | 2.6 | * | 2.4 | 1.0 | 1.1 | 2.3 | 2.0 | 3.1 | 3.5 | 7.5 | 13.9 | 29.5 |
| 1973 | 2.7 | * | 2.3 | 0.8 | 1.6 | 2.7 | 2.7 | 3.3 | 2.9 | 8.5 | 11.1 | * |
| 1972 | 3.2 | * | 2.8 | 1.2 | 2.7 | 3.3 | 2.4 | 4.1 | 5.3 | 6.3 | 10.0 | * |
| 1971 | 3.3 | * | 3.4 | 0.8 | 2.5 | 3.1 | 3.8 | 3.4 | 5.3 | 6.7 | 10.8 | * |
| 1970 | 3.3 | * | 3.4 | 1.0 | 2.1 | 3.3 | 3.6 | 3.9 | 4.0 | 7.9 | 14.4 | 32.2 |
| 1969 | 3.0 | * | 3.1 | 1.4 | 2.3 | 2.7 | 2.6 | 3.3 | 4.0 | 6.6 | 9.3 | * |
| 1968 | 3.3 | * | 3.7 | 1.2 | 2.0 | 3.2 | 2.8 | 3.5 | 5.9 | 8.3 | 10.8 | * |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| MENINGITIS (320) | | | | | | | | | | | | |
| BOTH SEXES | | | | | | | | | | | | |
| 1978 | 0.7 | 16.0 | 1.8 | 0.2 | 0.1 | 0.2 | 0.2 | 0.4 | 0.7 | 1.0 | 1.6 | 2.5 |
| 1977 | 0.7 | 15.4 | 1.7 | 0.2 | 0.1 | 0.1 | 0.2 | 0.4 | 0.7 | 1.2 | 1.6 | 2.6 |
| 1976 | 0.7 | 17.7 | 1.8 | 0.2 | 0.1 | 0.1 | 0.3 | 0.4 | 0.8 | 1.0 | 1.6 | 2.5 |
| 1975 | 0.8 | 18.5 | 1.7 | 0.2 | 0.1 | 0.2 | 0.3 | 0.5 | 0.8 | 1.0 | 1.8 | 2.4 |
| 1974 | 0.7 | 18.5 | 1.7 | 0.2 | 0.1 | 0.1 | 0.3 | 0.5 | 0.7 | 1.2 | 1.3 | 2.4 |
| 1973 | 0.7 | 15.7 | 1.6 | 0.2 | 0.1 | 0.1 | 0.3 | 0.6 | 0.7 | 1.2 | 1.4 | 1.7 |
| 1972 | 0.7 | 15.5 | 1.8 | 0.1 | 0.1 | 0.1 | 0.3 | 0.6 | 0.7 | 0.9 | 1.3 | 1.4 |
| 1971 | 0.8 | 14.7 | 1.5 | 0.2 | 0.2 | 0.2 | 0.2 | 0.5 | 0.8 | 1.2 | 1.5 | 2.1 |
| 1970 | 0.8 | 16.8 | 1.9 | 0.3 | 0.2 | 0.2 | 0.3 | 0.6 | 0.9 | 1.2 | 1.3 | 1.7 |
| 1969 | 0.9 | 16.1 | 1.9 | 0.2 | 0.1 | 0.2 | 0.4 | 0.6 | 1.0 | 1.4 | 1.9 | 1.6 |
| 1968 | 0.9 | 15.6 | 1.9 | 0.3 | 0.2 | 0.2 | 0.5 | 0.7 | 0.9 | 1.1 | 1.5 | 1.7 |
| MALE | | | | | | | | | | | | |
| 1978 | 0.8 | 18.3 | 2.3 | 0.2 | 0.2 | 0.2 | 0.3 | 0.4 | 0.7 | 1.2 | 1.9 | * |
| 1977 | 0.8 | 16.3 | 2.1 | 0.3 | 0.2 | * | 0.3 | 0.4 | 0.9 | 1.7 | 1.8 | * |
| 1976 | 0.8 | 19.0 | 2.4 | 0.2 | 0.2 | 0.2 | 0.4 | 0.6 | 0.9 | 1.2 | 1.9 | * |
| 1975 | 0.9 | 20.6 | 1.9 | 0.2 | 0.2 | 0.2 | 0.4 | 0.6 | 0.9 | 1.3 | 1.9 | * |
| 1974 | 0.8 | 20.1 | 1.9 | 0.2 | 0.2 | 0.1 | 0.3 | 0.6 | 0.8 | 1.3 | 1.7 | * |
| 1973 | 0.8 | 17.3 | 1.8 | 0.3 | 0.1 | 0.2 | 0.3 | 0.8 | 0.9 | 1.5 | 1.5 | * |
| 1972 | 0.8 | 17.3 | 1.9 | 0.2 | 0.1 | 0.1 | 0.4 | 0.8 | 1.0 | 1.0 | 1.3 | * |
| 1971 | 0.9 | 17.3 | 1.8 | 0.2 | 0.2 | 0.3 | 0.3 | 0.6 | 1.0 | 1.6 | 2.2 | * |
| 1970 | 1.0 | 19.4 | 2.0 | 0.3 | 0.2 | 0.2 | 0.4 | 0.8 | 1.2 | 1.4 | 1.6 | * |
| 1969 | 1.0 | 17.7 | 2.2 | 0.3 | 0.2 | 0.2 | 0.5 | 0.8 | 1.3 | 1.7 | 2.1 | * |
| 1968 | 1.1 | 19.2 | 2.3 | 0.3 | 0.3 | 0.3 | 0.6 | 0.9 | 1.1 | 1.3 | 2.1 | * |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| MENINGITIS (320) | | | | | | | | | | | | |
| FEMALE | | | | | | | | | | | | |
| 1978 | 0.6 | 13.5 | 1.3 | 0.2 | 0.1 | 0.2 | 0.2 | 0.3 | 0.6 | 0.9 | 1.4 | 2.6 |
| 1977 | 0.6 | 14.4 | 1.3 | 0.2 | 0.1 | 0.1 | * | 0.3 | 0.6 | 0.8 | 1.6 | 2.4 |
| 1976 | 0.6 | 16.3 | 1.1 | 0.2 | 0.1 | * | 0.2 | 0.2 | 0.7 | 0.9 | 1.5 | 2.7 |
| 1975 | 0.7 | 16.4 | 1.5 | 0.2 | * | * | 0.2 | 0.3 | 0.6 | 0.8 | 1.8 | 2.0 |
| 1974 | 0.6 | 16.8 | 1.4 | 0.2 | * | * | 0.2 | 0.3 | 0.6 | 1.1 | 1.1 | 2.8 |
| 1973 | 0.6 | 14.1 | 1.4 | 0.2 | 0.1 | * | 0.3 | 0.4 | 0.6 | 1.0 | 1.4 | * |
| 1972 | 0.6 | 13.7 | 1.7 | 0.1 | * | * | 0.3 | 0.4 | 0.5 | 0.8 | 1.3 | * |
| 1971 | 0.6 | 12.0 | 1.2 | 0.3 | 0.2 | * | * | 0.4 | 0.7 | 0.9 | 1.0 | * |
| 1970 | 0.7 | 14.2 | 1.7 | 0.2 | 0.2 | * | 0.2 | 0.4 | 0.7 | 1.0 | 1.1 | * |
| 1969 | 0.7 | 14.5 | 1.5 | 0.2 | * | * | 0.2 | 0.5 | 0.7 | 1.1 | 1.7 | * |
| 1968 | 0.7 | 11.8 | 1.5 | 0.2 | 0.2 | * | 0.3 | 0.4 | 0.7 | 1.1 | 1.2 | * |
| WHITE, BOTH SEXES | | | | | | | | | | | | |
| 1978 | 0.6 | 13.1 | 1.6 | 0.2 | 0.1 | 0.1 | 0.2 | 0.3 | 0.5 | 0.9 | 1.4 | 2.5 |
| 1977 | 0.6 | 12.3 | 1.6 | 0.2 | 0.1 | 0.1 | 0.1 | 0.3 | 0.6 | 1.1 | 1.3 | 2.5 |
| 1976 | 0.6 | 14.5 | 1.6 | 0.2 | 0.1 | 0.1 | 0.2 | 0.3 | 0.6 | 0.8 | 1.5 | 2.2 |
| 1975 | 0.6 | 14.0 | 1.6 | 0.2 | 0.1 | 0.1 | 0.2 | 0.4 | 0.6 | 0.8 | 1.7 | 2.3 |
| 1974 | 0.6 | 14.3 | 1.4 | 0.1 | 0.1 | 0.1 | 0.2 | 0.4 | 0.6 | 1.0 | 1.1 | 1.8 |
| 1973 | 0.6 | 12.2 | 1.3 | 0.2 | 0.1 | 0.1 | 0.2 | 0.4 | 0.5 | 1.0 | 1.2 | 1.6 |
| 1972 | 0.5 | 11.1 | 1.5 | 0.1 | 0.1 | 0.1 | 0.2 | 0.5 | 0.5 | 0.8 | 1.1 | 1.4 |
| 1971 | 0.6 | 11.6 | 1.4 | 0.2 | 0.2 | 0.1 | 0.1 | 0.3 | 0.7 | 1.0 | 1.3 | 1.8 |
| 1970 | 0.7 | 12.8 | 1.7 | 0.3 | 0.2 | 0.2 | 0.2 | 0.4 | 0.7 | 1.0 | 1.2 | 1.5 |
| 1969 | 0.7 | 12.5 | 1.6 | 0.2 | 0.1 | 0.1 | 0.3 | 0.4 | 0.7 | 1.1 | 1.6 | * |
| 1968 | 0.7 | 11.4 | 1.6 | 0.2 | 0.2 | 0.1 | 0.3 | 0.5 | 0.7 | 0.9 | 1.2 | * |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| MENINGITIS (320) | | | | | | | | | | | | |
| WHITE, MALE | | | | | | | | | | | | |
| 1978 | 0.7 | 15.3 | 2.1 | 0.2 | 0.2 | 0.1 | 0.2 | 0.3 | 0.6 | 1.1 | 1.7 | * |
| 1977 | 0.6 | 13.4 | 2.0 | 0.2 | * | * | * | 0.3 | 0.7 | 1.5 | 1.5 | * |
| 1976 | 0.7 | 16.4 | 2.2 | 0.2 | 0.1 | * | 0.3 | 0.5 | 0.8 | 1.0 | 1.8 | * |
| 1975 | 0.7 | 16.0 | 1.8 | 0.2 | 0.1 | 0.2 | 0.3 | 0.4 | 0.7 | 1.1 | 1.7 | * |
| 1974 | 0.7 | 16.3 | 1.7 | 0.2 | 0.1 | * | * | 0.5 | 0.6 | 1.2 | 1.4 | * |
| 1973 | 0.6 | 13.7 | 1.4 | 0.3 | * | * | * | 0.6 | 0.7 | 1.2 | 1.2 | * |
| 1972 | 0.7 | 13.1 | 1.7 | 0.1 | 0.1 | * | 0.3 | 0.6 | 0.8 | 0.9 | 1.0 | * |
| 1971 | 0.7 | 14.3 | 1.7 | 0.2 | 0.2 | 0.2 | * | 0.4 | 0.8 | 1.3 | 1.8 | * |
| 1970 | 0.8 | 14.9 | 1.8 | 0.3 | 0.2 | 0.2 | 0.3 | 0.5 | 1.0 | 1.2 | 1.6 | * |
| 1969 | 0.8 | 14.4 | 1.9 | 0.3 | 0.2 | * | 0.4 | 0.6 | 1.0 | 1.4 | 2.0 | * |
| 1968 | 0.8 | 14.0 | 1.9 | 0.2 | 0.2 | 0.2 | 0.3 | 0.7 | 0.9 | 1.0 | 1.6 | * |
| WHITE, FEMALE | | | | | | | | | | | | |
| 1978 | 0.5 | 10.7 | 1.2 | 0.2 | * | 0.1 | * | 0.3 | 0.4 | 0.8 | 1.2 | 2.6 |
| 1977 | 0.5 | 11.2 | 1.2 | 0.2 | * | * | * | 0.2 | 0.5 | 0.7 | 1.3 | 2.4 |
| 1976 | 0.5 | 12.5 | 1.0 | 0.2 | * | * | * | * | 0.5 | 0.7 | 1.3 | 2.3 |
| 1975 | 0.5 | 11.8 | 1.4 | 0.2 | * | * | * | 0.3 | 0.5 | 0.7 | 1.7 | 2.0 |
| 1974 | 0.5 | 12.3 | 1.1 | 0.1 | * | * | * | 0.2 | 0.5 | 0.9 | 0.9 | 2.0 |
| 1973 | 0.5 | 10.6 | 1.2 | 0.2 | * | * | 0.2 | 0.2 | 0.4 | 0.8 | 1.2 | * |
| 1972 | 0.4 | 9.0 | 1.2 | * | * | * | 0.2 | 0.3 | 0.3 | 0.8 | 1.2 | * |
| 1971 | 0.5 | 8.7 | 1.2 | 0.2 | 0.1 | * | * | 0.3 | 0.6 | 0.7 | 0.9 | * |
| 1970 | 0.5 | 10.7 | 1.5 | 0.2 | 0.1 | * | * | 0.3 | 0.5 | 0.8 | 0.9 | * |
| 1969 | 0.5 | 10.5 | 1.3 | 0.2 | * | * | * | 0.3 | 0.5 | 0.9 | 1.4 | * |
| 1968 | 0.5 | 8.7 | 1.3 | 0.2 | 0.2 | * | 0.2 | 0.4 | 0.5 | 0.8 | 1.0 | * |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| MENINGITIS (320) | | | | | | | | | | | | |
| ALL OTHER, BOTH SEXES | | | | | | | | | | | | |
| 1978 | 1.4 | 30.1 | 2.5 | 0.4 | * | 0.4 | * | 1.1 | 1.9 | 2.1 | 4.1 | * |
| 1977 | 1.5 | 30.1 | 2.2 | 0.3 | 0.3 | * | 0.8 | 1.2 | 2.1 | 2.7 | 4.7 | * |
| 1976 | 1.6 | 33.3 | 2.3 | 0.4 | * | * | 1.0 | 1.3 | 2.4 | 2.6 | * | * |
| 1975 | 1.7 | 41.3 | 1.8 | * | * | * | 0.9 | 1.3 | 2.4 | 2.8 | * | * |
| 1974 | 1.7 | 39.2 | 3.0 | 0.3 | * | * | 0.8 | 1.3 | 1.7 | 2.9 | 4.5 | * |
| 1973 | 1.8 | 33.1 | 3.0 | * | 0.4 | * | 0.9 | 2.0 | 2.3 | 3.8 | 4.1 | * |
| 1972 | 1.8 | 38.1 | 3.4 | 0.3 | * | * | 0.9 | 1.8 | 2.6 | 1.9 | * | * |
| 1971 | 1.8 | 31.3 | 2.1 | 0.4 | 0.4 | * | 0.9 | 2.0 | 2.3 | 3.9 | * | * |
| 1970 | 2.0 | 38.2 | 2.8 | 0.3 | * | * | 1.0 | 2.1 | 2.9 | 2.9 | * | * |
| 1969 | 2.1 | 36.8 | 3.4 | 0.3 | * | * | 1.2 | 2.4 | 3.1 | 3.8 | 5.1 | * |
| 1968 | 2.3 | 37.8 | 3.4 | 0.4 | 0.6 | * | 1.9 | 1.8 | 3.1 | 3.8 | 5.9 | * |
| ALL OTHER, MALE | | | | | | | | | | | | |
| 1978 | 1.6 | 33.2 | 3.2 | * | * | * | * | * | * | * | * | * |
| 1977 | 1.7 | 30.3 | 2.9 | * | * | * | * | * | 2.4 | 3.7 | * | * |
| 1976 | 1.8 | 32.0 | 3.0 | * | * | * | * | 1.8 | 2.6 | * | * | * |
| 1975 | 2.0 | 43.6 | 2.1 | * | * | * | * | 2.0 | 2.9 | 3.5 | * | * |
| 1974 | 1.9 | 39.8 | 2.8 | * | * | * | * | * | * | * | * | * |
| 1973 | 2.0 | 35.3 | 3.4 | * | * | * | * | 2.6 | 2.9 | 4.0 | * | * |
| 1972 | 2.0 | 38.8 | 3.1 | * | * | * | * | 1.9 | 3.1 | * | * | * |
| 1971 | 2.2 | 33.4 | 2.6 | * | * | * | * | 2.5 | 3.3 | 4.8 | * | * |
| 1970 | 2.2 | 44.1 | 3.2 | * | * | * | * | 2.7 | 3.0 | * | * | * |
| 1969 | 2.4 | 36.8 | 3.8 | * | * | * | 1.8 | 3.2 | 4.1 | 4.7 | * | * |
| 1968 | 3.0 | 47.2 | 4.5 | * | * | * | 2.7 | 2.7 | 3.6 | 4.0 | * | * |
| ALL OTHER, FEMALE | | | | | | | | | | | | |
| 1978 | 1.2 | 27.0 | * | * | * | * | * | * | 1.9 | * | * | * |
| 1977 | 1.3 | 30.0 | * | * | * | * | * | * | 1.9 | * | * | * |
| 1976 | 1.4 | 34.9 | * | * | * | * | * | * | 2.2 | * | * | * |
| 1975 | 1.4 | 39.0 | * | * | * | * | * | * | 1.9 | * | * | * |
| 1974 | 1.6 | 38.7 | 3.2 | * | * | * | * | * | * | 3.0 | * | * |
| 1973 | 1.6 | 30.8 | 2.5 | * | * | * | * | 1.5 | * | 3.6 | * | * |
| 1972 | 1.7 | 37.3 | 3.8 | * | * | * | * | 1.7 | 2.2 | * | * | * |
| 1971 | 1.4 | 29.2 | * | * | * | * | * | 1.6 | * | 3.1 | * | * |
| 1970 | 1.7 | 32.2 | 2.4 | * | * | * | * | 1.6 | 2.8 | * | * | * |
| 1969 | 1.8 | 36.7 | 3.0 | * | * | * | * | 1.6 | 2.3 | * | * | * |
| 1968 | 1.7 | 28.2 | 2.4 | * | * | * | * | * | 2.7 | 3.6 | * | * |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| MENINGITIS (320) | | | | | | | | | | | | |
| BLACK, BOTH SEXES | | | | | | | | | | | | |
| 1978 | 1.6 | 32.6 | 2.7 | 0.5 | * | 0.5 | * | 1.2 | 1.9 | 2.3 | 4.5 | * |
| 1977 | 1.6 | 30.1 | 2.3 | 0.4 | 0.4 | * | 0.9 | 1.2 | 2.2 | 2.9 | 4.7 | * |
| 1976 | 1.7 | 35.4 | 2.6 | * | * | * | 1.1 | 1.5 | 2.4 | 2.6 | * | * |
| 1975 | 1.8 | 42.5 | 1.9 | * | * | * | 1.0 | 1.4 | 2.6 | 3.0 | * | * |
| 1974 | 1.8 | 40.9 | 3.2 | * | * | * | 0.9 | 1.4 | 1.9 | 3.2 | 4.6 | * |
| 1973 | 1.9 | 35.8 | 3.2 | * | 0.4 | * | 1.0 | 2.2 | 2.4 | 4.0 | 4.4 | * |
| 1972 | 1.9 | 40.3 | 3.8 | * | * | * | 1.0 | 1.8 | 2.8 | 2.1 | * | * |
| 1971 | 1.9 | 32.5 | 2.3 | 0.4 | 0.4 | * | 1.0 | 2.1 | 2.5 | 4.0 | * | * |
| 1970 | 2.1 | 38.4 | 3.0 | * | * | * | 1.0 | 2.3 | 3.1 | 3.1 | * | * |
| 1969 | 2.2 | 37.6 | 3.6 | * | * | * | 1.3 | 2.5 | 3.2 | 3.9 | 5.6 | * |
| 1968 | 2.4 | 37.4 | 3.4 | 0.5 | 0.7 | * | 2.0 | 1.9 | 3.4 | 3.9 | 5.9 | * |
| BLACK, MALE | | | | | | | | | | | | |
| 1978 | 1.8 | 36.7 | 3.5 | * | * | * | * | * | * | * | * | * |
| 1977 | 1.8 | 30.2 | 3.0 | * | * | * | * | * | 2.5 | 4.1 | * | * |
| 1976 | 1.9 | 34.1 | 3.3 | * | * | * | * | 2.1 | 2.9 | * | * | * |
| 1975 | 2.2 | 45.2 | 2.3 | * | * | * | * | 2.3 | 3.1 | 4.0 | * | * |
| 1974 | 2.0 | 42.9 | 2.9 | * | * | * | * | * | * | * | * | * |
| 1973 | 2.2 | 37.3 | 3.7 | * | * | * | * | 2.9 | 3.1 | 4.2 | * | * |
| 1972 | 2.1 | 42.3 | 3.4 | * | * | * | * | * | 3.4 | * | * | * |
| 1971 | 2.3 | 33.7 | 2.8 | * | * | * | * | 2.6 | 3.7 | 4.9 | * | * |
| 1970 | 2.3 | 45.0 | 3.4 | * | * | * | * | 3.1 | 3.4 | * | * | * |
| 1969 | 2.5 | 37.6 | 4.0 | * | * | * | 2.0 | 3.4 | 4.3 | 4.6 | * | * |
| 1968 | 3.1 | 46.6 | 4.4 | * | * | * | 2.9 | 3.0 | 4.1 | 4.4 | * | * |
| BLACK, FEMALE | | | | | | | | | | | | |
| 1978 | 1.3 | 28.4 | * | * | * | * | * | * | * | * | * | * |
| 1977 | 1.4 | 30.0 | * | * | * | * | * | * | 2.0 | * | * | * |
| 1976 | 1.5 | 36.7 | * | * | * | * | * | * | 2.1 | * | * | * |
| 1975 | 1.4 | 39.8 | * | * | * | * | * | * | 2.1 | * | * | * |
| 1974 | 1.7 | 38.9 | 3.5 | * | * | * | * | * | * | 3.3 | * | * |
| 1973 | 1.7 | 34.1 | 2.7 | * | * | * | * | * | * | 3.9 | * | * |
| 1972 | 1.8 | 38.0 | 4.3 | * | * | * | * | 1.9 | 2.2 | * | * | * |
| 1971 | 1.5 | 31.2 | * | * | * | * | * | 1.7 | * | * | * | * |
| 1970 | 1.8 | 31.7 | 2.7 | * | * | * | * | * | 2.9 | * | * | * |
| 1969 | 1.9 | 37.7 | 3.1 | * | * | * | * | 1.8 | 2.4 | * | * | * |
| 1968 | 1.7 | 28.1 | 2.4 | * | * | * | * | * | 2.8 | 3.6 | * | * |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| MAJOR CARDIOVASCULAR DISEASES (390-448) | | | | | | | | | | | | |
| BOTH SEXES | | | | | | | | | | | | |
| 1978 | 435.0 | 31.8 | 3.2 | 1.6 | 4.0 | 11.4 | 58.9 | 223.3 | 608.7 | 1,552.4 | 4,098.8 | 10,833.5 |
| 1977 | 437.5 | 28.5 | 2.8 | 1.4 | 4.0 | 12.1 | 62.2 | 231.0 | 624.9 | 1,590.3 | 4,199.1 | 10,924.5 |
| 1976 | 447.9 | 27.7 | 2.6 | 1.5 | 4.2 | 12.5 | 64.8 | 239.6 | 655.7 | 1,651.3 | 4,346.0 | 11,387.8 |
| 1975 | 450.7 | 25.8 | 2.6 | 1.5 | 4.3 | 12.9 | 68.3 | 246.0 | 675.2 | 1,711.2 | 4,464.0 | 11,276.1 |
| 1974 | 473.9 | 27.9 | 2.8 | 1.7 | 4.5 | 13.8 | 72.0 | 259.9 | 713.4 | 1,829.3 | 4,775.5 | 12,425.0 |
| 1973 | 490.9 | 27.9 | 3.2 | 1.7 | 4.6 | 15.1 | 78.0 | 272.7 | 758.1 | 1,915.1 | 5,048.4 | 12,995.5 |
| 1972 | 494.6 | 25.8 | 3.0 | 1.7 | 4.6 | 15.6 | 81.1 | 280.4 | 778.2 | 1,983.7 | 5,134.8 | 12,956.4 |
| 1971 | 491.8 | 20.4 | 3.1 | 1.7 | 5.1 | 16.6 | 84.4 | 283.8 | 782.6 | 1,983.0 | 5,157.4 | 13,005.0 |
| 1970 | 496.0 | 19.2 | 2.8 | 1.6 | 4.9 | 17.4 | 86.8 | 290.9 | 801.8 | 2,054.3 | 5,309.9 | 12,108.5 |
| 1969 | 503.0 | 17.2 | 2.4 | 1.8 | 5.1 | 17.9 | 89.7 | 296.6 | 811.2 | 2,098.1 | 5,385.7 | 13,893.4 |
| 1968 | 513.2 | 18.6 | 2.6 | 1.8 | 5.1 | 18.5 | 92.0 | 307.9 | 844.0 | 2,159.2 | 5,537.3 | 14,403.9 |
| MALE | | | | | | | | | | | | |
| 1978 | 464.7 | 34.5 | 3.3 | 1.7 | 4.7 | 14.7 | 85.8 | 338.1 | 902.8 | 2,170.8 | 5,106.8 | 11,855.7 |
| 1977 | 471.4 | 32.3 | 2.8 | 1.5 | 4.6 | 15.8 | 90.0 | 350.3 | 925.2 | 2,227.3 | 5,245.2 | 11,991.7 |
| 1976 | 483.7 | 30.6 | 3.0 | 1.6 | 5.1 | 16.1 | 94.1 | 363.3 | 971.5 | 2,310.7 | 5,372.8 | 12,397.4 |
| 1975 | 490.2 | 27.7 | 2.9 | 1.6 | 5.1 | 16.2 | 98.2 | 372.9 | 1,000.0 | 2,377.7 | 5,515.3 | 12,285.9 |
| 1974 | 514.4 | 30.3 | 2.8 | 1.7 | 5.2 | 17.1 | 102.5 | 394.0 | 1,055.4 | 2,531.9 | 5,827.2 | 13,388.3 |
| 1973 | 535.5 | 30.8 | 3.5 | 1.7 | 5.3 | 18.2 | 110.5 | 410.6 | 1,120.4 | 2,640.3 | 6,153.9 | 13,929.2 |
| 1972 | 542.8 | 27.2 | 3.3 | 1.9 | 5.6 | 18.9 | 115.0 | 424.5 | 1,149.5 | 2,707.8 | 6,228.2 | 13,869.7 |
| 1971 | 542.7 | 21.7 | 3.3 | 1.7 | 6.1 | 20.2 | 120.2 | 425.5 | 1,149.6 | 2,702.7 | 6,226.8 | 13,998.0 |
| 1970 | 551.0 | 22.2 | 3.2 | 1.8 | 5.8 | 21.2 | 123.9 | 435.0 | 1,175.1 | 2,780.2 | 6,364.4 | 12,558.9 |
| 1969 | 561.3 | 17.5 | 2.7 | 1.8 | 5.6 | 21.8 | 128.2 | 446.9 | 1,191.9 | 2,838.8 | 6,428.4 | 14,720.0 |
| 1968 | 573.7 | 20.1 | 2.6 | 1.9 | 5.6 | 22.3 | 130.6 | 459.2 | 1,230.5 | 2,907.2 | 6,548.9 | 15,292.3 |

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| MAJOR CARDIOVASCULAR DISEASES (390-448) | | | | | | | | | | | | |
| FEMALE | | | | | | | | | | | | |
| 1978 | 406.9 | 28.9 | 3.0 | 1.5 | 3.4 | 8.2 | 33.0 | 116.4 | 349.1 | 1,079.8 | 3,498.7 | 10,371.7 |
| 1977 | 405.4 | 24.5 | 2.8 | 1.3 | 3.3 | 8.5 | 35.6 | 120.0 | 359.3 | 1,103.0 | 3,573.0 | 10,434.4 |
| 1976 | 414.0 | 24.6 | 2.3 | 1.4 | 3.2 | 9.0 | 36.7 | 124.6 | 376.1 | 1,146.3 | 3,726.0 | 10,913.6 |
| 1975 | 413.2 | 23.9 | 2.3 | 1.4 | 3.5 | 9.6 | 39.7 | 128.0 | 387.5 | 1,200.6 | 3,823.6 | 10,787.3 |
| 1974 | 435.3 | 25.3 | 2.8 | 1.6 | 3.7 | 10.5 | 42.8 | 135.3 | 410.1 | 1,290.6 | 4,127.7 | 11,947.9 |
| 1973 | 448.4 | 24.9 | 2.9 | 1.6 | 3.9 | 12.2 | 46.9 | 144.6 | 436.1 | 1,358.2 | 4,358.0 | 12,525.5 |
| 1972 | 448.8 | 24.2 | 2.6 | 1.4 | 3.5 | 12.4 | 48.9 | 146.5 | 447.3 | 1,425.1 | 4,437.5 | 12,489.0 |
| 1971 | 443.4 | 18.9 | 2.9 | 1.7 | 4.2 | 13.1 | 50.3 | 152.0 | 454.4 | 1,425.7 | 4,463.5 | 12,486.6 |
| 1970 | 443.9 | 16.0 | 2.4 | 1.5 | 4.0 | 13.7 | 51.7 | 156.5 | 466.7 | 1,490.3 | 4,612.3 | 11,856.2 |
| 1969 | 447.7 | 16.8 | 2.1 | 1.8 | 4.6 | 14.2 | 53.3 | 156.4 | 468.6 | 1,519.2 | 4,686.5 | 13,446.0 |
| 1968 | 455.8 | 16.9 | 2.6 | 1.7 | 4.5 | 14.8 | 55.5 | 166.1 | 494.7 | 1,570.3 | 4,844.9 | 13,933.3 |
| WHITE, BOTH SEXES | | | | | | | | | | | | |
| 1978 | 451.6 | 26.1 | 2.7 | 1.5 | 3.5 | 9.2 | 50.9 | 205.3 | 578.7 | 1,518.4 | 4,131.8 | 11,086.4 |
| 1977 | 453.6 | 24.7 | 2.5 | 1.3 | 3.4 | 10.0 | 53.7 | 212.7 | 594.7 | 1,553.1 | 4,240.6 | 11,176.3 |
| 1976 | 464.3 | 23.3 | 2.3 | 1.4 | 3.5 | 10.1 | 56.6 | 220.7 | 626.2 | 1,611.5 | 4,401.7 | 11,658.4 |
| 1975 | 466.8 | 22.3 | 2.3 | 1.4 | 3.6 | 10.4 | 59.1 | 228.0 | 645.0 | 1,666.3 | 4,523.4 | 11,558.7 |
| 1974 | 489.8 | 21.7 | 2.4 | 1.5 | 3.9 | 10.9 | 62.4 | 239.0 | 678.8 | 1,775.6 | 4,845.9 | 12,730.2 |
| 1973 | 505.9 | 21.0 | 2.7 | 1.5 | 3.9 | 12.0 | 67.0 | 249.4 | 720.1 | 1,855.0 | 5,114.2 | 13,317.3 |
| 1972 | 509.4 | 18.6 | 2.6 | 1.5 | 3.7 | 12.0 | 69.0 | 256.0 | 738.3 | 1,931.0 | 5,198.4 | 13,310.5 |
| 1971 | 506.2 | 15.4 | 2.7 | 1.5 | 4.3 | 12.9 | 72.2 | 260.5 | 745.2 | 1,928.4 | 5,214.9 | 13,383.6 |
| 1970 | 509.5 | 14.6 | 2.3 | 1.5 | 4.1 | 13.2 | 73.1 | 265.6 | 759.9 | 2,000.3 | 5,352.4 | 12,559.8 |
| 1969 | 515.5 | 14.8 | 2.2 | 1.6 | 4.2 | 13.3 | 75.5 | 270.0 | 767.2 | 2,041.1 | 5,427.2 | 14,324.9 |
| 1968 | 524.4 | 15.1 | 2.1 | 1.6 | 4.2 | 13.8 | 77.0 | 280.5 | 793.6 | 2,097.5 | 5,578.4 | 14,832.3 |

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| MAJOR CARDIOVASCULAR DISEASES (390-448) | | | | | | | | | | | | |
| WHITE, MALE | | | | | | | | | | | | |
| 1978 | 480.9 | 29.2 | 2.9 | 1.6 | 4.1 | 12.0 | 76.1 | 319.1 | 876.3 | 2,157.7 | 5,191.0 | 12,177.1 |
| 1977 | 487.9 | 29.4 | 2.6 | 1.4 | 4.0 | 13.1 | 80.5 | 332.0 | 898.8 | 2,215.1 | 5,348.3 | 12,313.5 |
| 1976 | 500.4 | 26.9 | 2.6 | 1.5 | 4.3 | 13.2 | 84.9 | 344.3 | 947.8 | 2,295.3 | 5,488.2 | 12,750.5 |
| 1975 | 507.0 | 24.2 | 2.5 | 1.5 | 4.3 | 13.0 | 87.8 | 355.5 | 978.7 | 2,359.2 | 5,620.6 | 12,653.5 |
| 1974 | 531.2 | 24.4 | 2.4 | 1.7 | 4.6 | 13.8 | 92.7 | 374.3 | 1,028.5 | 2,505.0 | 5,945.5 | 13,777.7 |
| 1973 | 551.8 | 22.8 | 2.9 | 1.5 | 4.5 | 14.7 | 99.7 | 388.3 | 1,092.0 | 2,611.0 | 6,268.0 | 14,344.9 |
| 1972 | 559.0 | 19.5 | 2.8 | 1.8 | 4.6 | 15.1 | 102.7 | 400.9 | 1,121.2 | 2,687.6 | 6,335.3 | 14,306.1 |
| 1971 | 559.0 | 17.4 | 2.8 | 1.5 | 5.2 | 16.5 | 107.9 | 405.0 | 1,122.3 | 2,679.5 | 6,336.9 | 14,500.4 |
| 1970 | 566.8 | 17.8 | 2.8 | 1.7 | 5.0 | 16.7 | 109.8 | 412.9 | 1,146.2 | 2,756.9 | 6,456.0 | 13,142.4 |
| 1969 | 576.1 | 15.4 | 2.4 | 1.6 | 4.8 | 16.7 | 113.7 | 422.0 | 1,162.5 | 2,811.8 | 6,508.7 | 15,294.5 |
| 1968 | 587.5 | 17.0 | 2.1 | 1.7 | 4.7 | 17.4 | 115.7 | 435.0 | 1,192.2 | 2,876.7 | 6,637.6 | 15,818.8 |
| WHITE, FEMALE | | | | | | | | | | | | |
| 1978 | 423.8 | 22.8 | 2.5 | 1.3 | 2.8 | 6.4 | 26.2 | 97.6 | 314.4 | 1,030.5 | 3,508.4 | 10,600.4 |
| 1977 | 420.9 | 19.7 | 2.4 | 1.1 | 2.7 | 6.8 | 27.6 | 100.0 | 324.0 | 1,047.9 | 3,587.2 | 10,661.4 |
| 1976 | 429.9 | 19.5 | 2.0 | 1.3 | 2.6 | 7.0 | 28.9 | 104.4 | 339.9 | 1,089.2 | 3,754.9 | 11,163.4 |
| 1975 | 428.4 | 20.4 | 2.1 | 1.2 | 2.9 | 7.8 | 31.1 | 108.1 | 347.8 | 1,137.5 | 3,863.5 | 11,047.6 |
| 1974 | 450.3 | 18.8 | 2.5 | 1.4 | 3.1 | 7.9 | 33.0 | 111.9 | 367.2 | 1,219.1 | 4,174.1 | 12,219.3 |
| 1973 | 461.9 | 19.1 | 2.5 | 1.5 | 3.2 | 9.3 | 35.3 | 119.2 | 388.4 | 1,277.1 | 4,399.6 | 12,821.6 |
| 1972 | 462.0 | 17.6 | 2.3 | 1.2 | 2.8 | 9.0 | 36.3 | 120.3 | 395.9 | 1,350.2 | 4,480.9 | 12,809.5 |
| 1971 | 455.9 | 13.3 | 2.6 | 1.5 | 3.4 | 9.4 | 37.6 | 125.0 | 407.2 | 1,349.7 | 4,496.6 | 12,811.6 |
| 1970 | 454.9 | 11.3 | 1.8 | 1.3 | 3.2 | 9.8 | 37.8 | 127.5 | 412.6 | 1,416.1 | 4,630.5 | 12,241.0 |
| 1969 | 457.8 | 14.1 | 1.9 | 1.6 | 3.7 | 9.9 | 38.8 | 127.2 | 410.8 | 1,441.4 | 4,707.6 | 13,829.4 |
| 1968 | 464.3 | 13.1 | 2.2 | 1.6 | 3.6 | 10.3 | 39.9 | 134.8 | 432.9 | 1,487.3 | 4,860.4 | 14,319.8 |

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| MAJOR CARDIOVASCULAR DISEASES (390-448) | | | | | | | | | | | | |
| ALL OTHER, BOTH SEXES | | | | | | | | | | | | |
| 1978 | 329.2 | 59.4 | 5.2 | 2.3 | 7.2 | 25.5 | 112.9 | 362.7 | 878.3 | 1,870.3 | 3,758.8 | 7,932.7 |
| 1977 | 333.9 | 46.9 | 4.4 | 1.9 | 7.5 | 26.2 | 119.5 | 375.8 | 901.2 | 1,943.9 | 3,771.7 | 8,041.9 |
| 1976 | 340.5 | 49.1 | 4.1 | 2.1 | 8.1 | 28.7 | 120.8 | 391.9 | 927.4 | 2,032.9 | 3,773.3 | 8,228.8 |
| 1975 | 344.0 | 43.2 | 4.1 | 2.3 | 8.1 | 29.8 | 131.8 | 394.0 | 954.6 | 2,147.5 | 3,828.6 | 8,033.6 |
| 1974 | 366.4 | 58.9 | 4.8 | 2.3 | 7.9 | 34.0 | 138.8 | 434.3 | 1,038.0 | 2,355.8 | 4,004.6 | 8,873.3 |
| 1973 | 388.7 | 62.0 | 5.8 | 2.7 | 9.1 | 37.5 | 155.3 | 469.8 | 1,117.8 | 2,508.7 | 4,297.2 | 9,244.9 |
| 1972 | 392.3 | 62.5 | 5.2 | 2.7 | 9.7 | 40.8 | 168.5 | 489.5 | 1,158.0 | 2,504.9 | 4,395.0 | 8,871.5 |
| 1971 | 390.7 | 46.8 | 5.3 | 2.8 | 10.5 | 42.7 | 173.9 | 487.0 | 1,140.1 | 2,522.6 | 4,460.2 | 8,691.7 |
| 1970 | 401.8 | 43.4 | 5.3 | 2.4 | 10.3 | 46.4 | 187.6 | 512.5 | 1,197.6 | 2,581.6 | 4,769.8 | 7,474.1 |
| 1969 | 413.9 | 30.7 | 3.8 | 2.9 | 10.6 | 51.0 | 197.5 | 535.7 | 1,235.6 | 2,665.5 | 4,859.2 | 9,019.3 |
| 1968 | 432.6 | 37.3 | 5.0 | 2.8 | 11.0 | 51.9 | 207.4 | 555.7 | 1,333.5 | 2,777.2 | 5,004.7 | 9,619.6 |
| ALL OTHER, MALE | | | | | | | | | | | | |
| 1978 | 359.4 | 60.6 | 4.9 | 2.3 | 7.7 | 33.4 | 157.1 | 496.1 | 1,151.4 | 2,295.7 | 4,295.9 | 8,563.8 |
| 1977 | 362.4 | 46.8 | 4.2 | 1.9 | 8.3 | 34.7 | 159.4 | 505.0 | 1,175.1 | 2,340.9 | 4,274.3 | 8,710.7 |
| 1976 | 371.5 | 49.2 | 4.8 | 2.2 | 9.8 | 37.3 | 162.0 | 527.0 | 1,196.5 | 2,451.5 | 4,300.8 | 8,860.0 |
| 1975 | 376.4 | 45.1 | 4.7 | 2.2 | 9.4 | 39.7 | 175.7 | 524.4 | 1,202.8 | 2,552.7 | 4,474.4 | 8,609.3 |
| 1974 | 399.1 | 60.6 | 5.2 | 2.2 | 8.8 | 41.8 | 176.9 | 567.7 | 1,314.1 | 2,792.7 | 4,610.0 | 9,378.0 |
| 1973 | 421.8 | 70.7 | 6.5 | 3.0 | 10.0 | 44.5 | 193.4 | 608.7 | 1,397.1 | 2,920.2 | 4,932.1 | 9,685.4 |
| 1972 | 428.6 | 67.4 | 6.3 | 2.9 | 11.6 | 48.1 | 210.3 | 636.3 | 1,423.4 | 2,902.4 | 5,077.4 | 9,459.6 |
| 1971 | 426.2 | 44.9 | 6.2 | 2.8 | 12.0 | 48.2 | 217.8 | 611.7 | 1,414.4 | 2,924.6 | 5,031.4 | 9,048.9 |
| 1970 | 438.8 | 46.3 | 5.3 | 2.4 | 11.1 | 55.2 | 236.4 | 636.7 | 1,450.1 | 2,999.9 | 5,300.1 | 7,432.1 |
| 1969 | 454.2 | 29.5 | 4.4 | 2.8 | 11.1 | 60.5 | 247.1 | 679.0 | 1,476.9 | 3,093.5 | 5,465.4 | 9,302.3 |
| 1968 | 472.8 | 37.4 | 5.3 | 3.1 | 11.5 | 60.1 | 254.1 | 686.2 | 1,603.5 | 3,203.2 | 5,484.9 | 10,168.3 |
| ALL OTHER, FEMALE | | | | | | | | | | | | |
| 1978 | 301.9 | 58.2 | 5.5 | 2.3 | 6.6 | 18.7 | 75.9 | 252.5 | 651.7 | 1,540.9 | 3,401.3 | 7,669.7 |
| 1977 | 308.1 | 46.9 | 4.6 | 2.0 | 6.7 | 18.8 | 86.1 | 268.6 | 672.7 | 1,632.0 | 3,419.3 | 7,681.7 |
| 1976 | 312.3 | 49.0 | 3.4 | 1.9 | 6.5 | 21.2 | 86.3 | 278.9 | 702.1 | 1,701.7 | 3,412.8 | 7,874.5 |
| 1975 | 314.5 | 41.3 | 3.5 | 2.3 | 6.8 | 21.3 | 95.0 | 283.8 | 745.1 | 1,825.3 | 3,376.3 | 7,612.9 |
| 1974 | 336.6 | 57.3 | 4.5 | 2.3 | 7.0 | 27.4 | 106.9 | 321.3 | 803.1 | 2,005.9 | 3,587.0 | 8,576.5 |
| 1973 | 358.4 | 53.1 | 5.1 | 2.4 | 8.2 | 31.4 | 123.5 | 350.9 | 879.9 | 2,176.4 | 3,863.5 | 8,977.2 |
| 1972 | 359.1 | 57.5 | 4.0 | 2.5 | 7.9 | 34.5 | 133.5 | 363.3 | 929.1 | 2,185.9 | 3,917.6 | 8,507.9 |
| 1971 | 358.2 | 48.8 | 4.4 | 2.9 | 9.1 | 38.1 | 137.4 | 378.5 | 900.3 | 2,193.9 | 4,037.9 | 8,461.6 |
| 1970 | 367.9 | 40.6 | 5.3 | 2.5 | 9.5 | 38.9 | 147.0 | 403.7 | 974.4 | 2,236.9 | 4,366.2 | 7,503.8 |
| 1969 | 377.1 | 32.0 | 3.2 | 3.0 | 10.1 | 42.9 | 156.1 | 410.1 | 1,021.7 | 2,313.3 | 4,378.1 | 8,827.7 |
| 1968 | 395.6 | 37.0 | 4.6 | 2.6 | 10.7 | 45.0 | 168.2 | 440.8 | 1,091.0 | 2,423.8 | 4,651.9 | 9,250.8 |

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| MAJOR CARDIOVASCULAR DISEASES (390-448) | | | | | | | | | | | | |
| BLACK, BOTH SEXES | | | | | | | | | | | | |
| 1978 | 360.3 | 63.1 | 5.5 | 2.4 | 7.7 | 29.0 | 128.5 | 404.7 | 949.6 | 1,986.0 | 3,859.8 | 8,198.7 |
| 1977 | 365.5 | 51.1 | 4.6 | 2.1 | 7.9 | 29.7 | 135.7 | 418.3 | 971.7 | 2,069.8 | 3,857.1 | 8,362.9 |
| 1976 | 370.9 | 51.9 | 4.6 | 2.1 | 8.7 | 32.6 | 136.2 | 433.5 | 996.8 | 2,157.8 | 3,865.4 | 8,536.0 |
| 1975 | 373.1 | 46.1 | 4.2 | 2.3 | 8.7 | 33.6 | 147.0 | 432.4 | 1,020.3 | 2,278.4 | 3,914.5 | 8,350.8 |
| 1974 | 394.5 | 59.7 | 5.1 | 2.4 | 8.5 | 38.4 | 154.8 | 473.4 | 1,101.2 | 2,492.0 | 4,089.2 | 9,211.6 |
| 1973 | 417.1 | 68.1 | 6.1 | 2.8 | 9.7 | 41.3 | 172.7 | 509.4 | 1,187.7 | 2,656.7 | 4,382.4 | 9,546.5 |
| 1972 | 419.5 | 67.5 | 5.3 | 2.8 | 10.2 | 44.8 | 186.1 | 530.4 | 1,222.0 | 2,648.5 | 4,518.9 | 9,216.4 |
| 1971 | 415.5 | 47.7 | 5.4 | 2.9 | 11.1 | 47.4 | 191.1 | 523.1 | 1,203.3 | 2,656.4 | 4,594.4 | 8,977.6 |
| 1970 | 431.6 | 44.1 | 5.6 | 2.6 | 10.8 | 52.0 | 209.4 | 555.7 | 1,274.0 | 2,730.7 | 5,030.5 | 7,921.9 |
| 1969 | 439.3 | 31.9 | 3.9 | 2.8 | 11.1 | 55.8 | 214.7 | 573.5 | 1,307.5 | 2,794.5 | 5,084.7 | 9,428.9 |
| 1968 | 457.4 | 39.5 | 5.3 | 2.9 | 11.2 | 56.6 | 225.8 | 591.0 | 1,407.2 | 2,908.2 | 5,194.1 | 9,967.4 |
| BLACK, MALE | | | | | | | | | | | | |
| 1978 | 389.1 | 64.1 | 5.5 | 2.3 | 8.2 | 38.2 | 178.5 | 551.1 | 1,245.4 | 2,451.5 | 4,370.7 | 8,680.4 |
| 1977 | 393.1 | 51.8 | 4.3 | 2.0 | 8.8 | 39.5 | 181.1 | 559.9 | 1,270.3 | 2,515.7 | 4,376.2 | 8,955.1 |
| 1976 | 401.1 | 52.0 | 5.4 | 2.2 | 10.5 | 42.5 | 183.7 | 580.0 | 1,288.6 | 2,620.1 | 4,391.9 | 9,225.5 |
| 1975 | 404.2 | 47.8 | 4.8 | 2.3 | 10.2 | 44.7 | 195.2 | 572.9 | 1,286.3 | 2,733.3 | 4,544.6 | 9,041.3 |
| 1974 | 425.9 | 61.5 | 5.3 | 2.3 | 9.4 | 47.0 | 197.1 | 616.4 | 1,400.5 | 2,979.4 | 4,698.1 | 9,830.2 |
| 1973 | 448.9 | 77.7 | 6.8 | 3.1 | 10.5 | 49.7 | 214.4 | 659.3 | 1,494.2 | 3,115.4 | 5,019.6 | 10,161.0 |
| 1972 | 454.9 | 72.2 | 6.5 | 2.7 | 12.3 | 53.7 | 231.4 | 691.4 | 1,515.6 | 3,086.1 | 5,197.0 | 10,040.0 |
| 1971 | 450.0 | 43.9 | 6.6 | 2.9 | 12.7 | 53.9 | 239.7 | 656.6 | 1,503.7 | 3,096.1 | 5,198.9 | 9,472.5 |
| 1970 | 468.0 | 47.0 | 5.3 | 2.4 | 11.9 | 62.7 | 262.8 | 692.5 | 1,555.3 | 3,194.6 | 5,612.8 | 8,024.3 |
| 1969 | 478.0 | 30.6 | 4.4 | 2.7 | 11.5 | 65.6 | 267.0 | 727.6 | 1,576.7 | 3,259.1 | 5,729.3 | 9,970.3 |
| 1968 | 496.2 | 40.8 | 5.7 | 3.2 | 11.2 | 66.2 | 275.8 | 730.5 | 1,705.4 | 3,365.3 | 5,723.0 | 10,845.7 |
| BLACK, FEMALE | | | | | | | | | | | | |
| 1978 | 334.4 | 62.1 | 5.4 | 2.4 | 7.1 | 21.1 | 86.9 | 284.3 | 708.0 | 1,638.1 | 3,518.7 | 7,950.5 |
| 1977 | 340.6 | 50.4 | 4.8 | 2.1 | 7.1 | 21.2 | 97.9 | 301.0 | 728.0 | 1,735.2 | 3,523.4 | 8,054.3 |
| 1976 | 343.6 | 51.8 | 3.8 | 2.0 | 7.0 | 24.1 | 96.8 | 311.8 | 755.3 | 1,805.5 | 3,512.3 | 8,171.9 |
| 1975 | 344.9 | 44.2 | 3.6 | 2.3 | 7.3 | 24.2 | 107.0 | 314.7 | 799.5 | 1,933.4 | 3,488.1 | 7,972.6 |
| 1974 | 366.2 | 57.9 | 4.8 | 2.4 | 7.6 | 30.9 | 119.5 | 353.4 | 852.9 | 2,122.0 | 3,677.3 | 8,870.5 |
| 1973 | 388.3 | 58.1 | 5.4 | 2.4 | 8.8 | 34.2 | 137.9 | 382.0 | 932.8 | 2,302.1 | 3,958.1 | 9,201.4 |
| 1972 | 387.3 | 62.4 | 4.1 | 2.8 | 8.3 | 37.2 | 148.4 | 393.1 | 975.5 | 2,312.0 | 4,025.7 | 8,745.7 |
| 1971 | 384.0 | 51.6 | 4.2 | 2.9 | 9.5 | 41.9 | 150.9 | 408.2 | 948.9 | 2,305.6 | 4,160.2 | 8,682.1 |
| 1970 | 398.5 | 41.2 | 5.9 | 2.7 | 9.8 | 42.9 | 165.1 | 437.5 | 1,034.5 | 2,363.2 | 4,603.4 | 7,855.9 |
| 1969 | 404.1 | 33.3 | 3.3 | 2.9 | 10.8 | 47.6 | 171.3 | 440.1 | 1,078.4 | 2,425.0 | 4,588.1 | 9,095.0 |
| 1968 | 422.1 | 38.3 | 4.9 | 2.6 | 11.1 | 48.6 | 184.0 | 469.5 | 1,149.3 | 2,535.2 | 4,800.9 | 9,596.4 |

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| 390-398, 404, 410-429 Diseases of heart | | | | | | | | | | | | |
| BOTH SEXES | | | | | | | | | | | | |
| 1978 | 328.5 | 25.3 | 2.2 | 1.0 | 2.6 | 8.0 | 46.6 | 188.5 | 511.0 | 1,226.2 | 2,979.3 | 7,459.6 |
| 1977 | 327.1 | 22.3 | 1.9 | 0.8 | 2.5 | 8.3 | 49.3 | 194.3 | 521.0 | 1,245.8 | 3,008.5 | 7,405.7 |
| 1976 | 332.7 | 22.5 | 1.8 | 0.9 | 2.6 | 8.3 | 50.7 | 200.0 | 543.7 | 1,283.0 | 3,094.7 | 7,657.0 |
| 1975 | 332.4 | 19.9 | 1.8 | 0.9 | 2.6 | 8.5 | 53.9 | 205.2 | 557.0 | 1,319.7 | 3,136.0 | 7,505.9 |
| 1974 | 346.0 | 21.6 | 2.0 | 0.9 | 2.7 | 9.3 | 55.7 | 215.6 | 584.6 | 1,396.7 | 3,316.0 | 8,160.9 |
| 1973 | 358.2 | 22.6 | 2.1 | 1.0 | 2.8 | 9.9 | 60.5 | 224.5 | 620.9 | 1,458.3 | 3,507.5 | 8,502.1 |
| 1972 | 361.2 | 20.1 | 2.1 | 1.0 | 2.8 | 10.1 | 62.8 | 229.6 | 633.9 | 1,509.8 | 3,574.2 | 8,479.4 |
| 1971 | 359.4 | 15.2 | 1.8 | 1.0 | 3.0 | 10.7 | 65.1 | 232.1 | 638.3 | 1,509.4 | 3,587.9 | 8,537.3 |
| 1970 | 362.0 | 13.1 | 1.7 | 0.8 | 3.0 | 11.4 | 66.7 | 238.4 | 652.3 | 1,558.2 | 3,683.8 | 7,891.3 |
| 1969 | 367.1 | 11.3 | 1.5 | 1.0 | 3.0 | 11.5 | 69.3 | 243.7 | 661.2 | 1,588.4 | 3,737.9 | 9,035.8 |
| 1968 | 373.5 | 12.3 | 1.6 | 1.0 | 2.8 | 11.9 | 70.8 | 251.7 | 686.9 | 1,632.8 | 3,825.4 | 9,278.3 |
| MALE | | | | | | | | | | | | |
| 1978 | 368.9 | 27.4 | 2.1 | 1.0 | 3.2 | 11.2 | 72.8 | 299.8 | 782.3 | 1,760.0 | 3,826.0 | 8,431.7 |
| 1977 | 371.0 | 25.7 | 1.9 | 0.9 | 3.1 | 11.7 | 76.5 | 309.1 | 797.1 | 1,792.8 | 3,870.7 | 8,412.9 |
| 1976 | 378.4 | 25.4 | 2.1 | 0.9 | 3.3 | 11.7 | 79.7 | 319.1 | 833.1 | 1,845.2 | 3,948.5 | 8,588.9 |
| 1975 | 380.9 | 21.5 | 1.9 | 1.0 | 3.1 | 11.9 | 83.5 | 327.8 | 853.5 | 1,884.6 | 3,996.7 | 8,432.3 |
| 1974 | 396.1 | 23.2 | 1.9 | 0.9 | 3.4 | 12.6 | 85.8 | 344.7 | 895.1 | 1,986.9 | 4,172.0 | 9,086.0 |
| 1973 | 412.2 | 25.4 | 2.3 | 1.1 | 3.4 | 13.3 | 92.8 | 357.4 | 950.1 | 2,066.4 | 4,412.1 | 9,429.7 |
| 1972 | 416.4 | 20.8 | 2.3 | 1.1 | 3.6 | 13.5 | 96.2 | 367.6 | 967.1 | 2,115.2 | 4,453.6 | 9,309.6 |
| 1971 | 416.6 | 16.0 | 1.9 | 0.9 | 3.8 | 14.5 | 100.4 | 367.0 | 968.9 | 2,112.3 | 4,446.0 | 9,481.0 |
| 1970 | 422.5 | 15.1 | 1.9 | 0.9 | 3.7 | 15.2 | 103.2 | 376.4 | 987.2 | 2,170.3 | 4,534.8 | 8,426.2 |
| 1969 | 430.5 | 11.2 | 1.5 | 1.0 | 3.5 | 15.6 | 107.4 | 387.3 | 1,004.4 | 2,208.2 | 4,584.0 | 9,829.2 |
| 1968 | 438.4 | 13.4 | 1.7 | 1.0 | 3.3 | 15.7 | 108.5 | 396.0 | 1,034.7 | 2,257.5 | 4,652.6 | 10,078.0 |

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| 390-398, 404, 410-429 Diseases of heart | | | | | | | | | | | | |
| FEMALE | | | | | | | | | | | | |
| 1978 | 290.2 | 23.2 | 2.4 | 1.0 | 2.0 | 4.9 | 21.5 | 84.9 | 271.6 | 818.3 | 2,475.2 | 7,020.4 |
| 1977 | 285.6 | 18.8 | 1.9 | 0.8 | 1.9 | 5.0 | 23.2 | 87.4 | 276.9 | 827.2 | 2,492.5 | 6,943.0 |
| 1976 | 289.4 | 19.4 | 1.5 | 0.9 | 1.9 | 5.0 | 23.0 | 89.2 | 287.5 | 852.5 | 2,579.1 | 7,219.1 |
| 1975 | 286.4 | 18.2 | 1.6 | 0.8 | 2.1 | 5.2 | 25.5 | 91.1 | 294.4 | 887.0 | 2,611.7 | 7,057.5 |
| 1974 | 298.5 | 19.9 | 2.0 | 0.9 | 2.1 | 6.0 | 26.8 | 95.7 | 309.2 | 944.2 | 2,788.6 | 7,702.8 |
| 1973 | 306.9 | 19.7 | 1.8 | 0.9 | 2.2 | 6.7 | 29.6 | 101.1 | 328.3 | 991.3 | 2,942.4 | 8,035.3 |
| 1972 | 308.7 | 19.4 | 1.8 | 0.9 | 2.1 | 6.7 | 30.9 | 101.3 | 337.1 | 1,042.9 | 3,013.6 | 8,054.5 |
| 1971 | 305.1 | 14.4 | 1.8 | 1.0 | 2.3 | 7.0 | 31.5 | 106.6 | 342.9 | 1,042.4 | 3,031.2 | 8,044.6 |
| 1970 | 304.5 | 10.9 | 1.6 | 0.8 | 2.3 | 7.7 | 32.2 | 109.9 | 351.6 | 1,082.7 | 3,120.8 | 7,591.8 |
| 1969 | 307.0 | 11.3 | 1.4 | 1.0 | 2.5 | 7.5 | 33.3 | 109.7 | 352.3 | 1,104.0 | 3,170.4 | 8,606.4 |
| 1968 | 311.8 | 11.2 | 1.6 | 1.0 | 2.4 | 8.3 | 35.1 | 116.4 | 372.6 | 1,140.9 | 3,259.2 | 8,850.1 |
| WHITE, BOTH SEXES | | | | | | | | | | | | |
| 1978 | 342.6 | 21.3 | 1.8 | 0.9 | 2.1 | 6.5 | 41.3 | 176.7 | 492.9 | 1,212.2 | 3,015.4 | 7,640.4 |
| 1977 | 340.7 | 19.5 | 1.6 | 0.7 | 2.0 | 6.9 | 43.7 | 182.5 | 502.9 | 1,230.1 | 3,050.7 | 7,577.5 |
| 1976 | 346.7 | 18.5 | 1.6 | 0.8 | 2.1 | 6.7 | 45.8 | 188.4 | 526.9 | 1,266.4 | 3,145.9 | 7,847.5 |
| 1975 | 346.1 | 17.2 | 1.5 | 0.8 | 2.1 | 6.9 | 48.0 | 194.5 | 540.6 | 1,300.1 | 3,190.3 | 7,697.7 |
| 1974 | 359.5 | 16.6 | 1.6 | 0.8 | 2.3 | 7.3 | 50.0 | 203.1 | 565.6 | 1,371.7 | 3,375.9 | 8,361.1 |
| 1973 | 371.3 | 16.5 | 1.7 | 0.9 | 2.3 | 7.9 | 54.2 | 210.7 | 600.5 | 1,429.9 | 3,565.1 | 8,713.5 |
| 1972 | 374.4 | 14.6 | 1.8 | 0.8 | 2.2 | 8.0 | 55.6 | 215.2 | 612.4 | 1,488.1 | 3,631.6 | 8,718.0 |
| 1971 | 372.5 | 10.9 | 1.5 | 0.8 | 2.4 | 8.4 | 58.0 | 219.2 | 619.7 | 1,486.2 | 3,638.6 | 8,791.7 |
| 1970 | 374.5 | 9.5 | 1.3 | 0.7 | 2.4 | 8.8 | 58.6 | 224.2 | 630.9 | 1,537.8 | 3,726.5 | 8,185.9 |
| 1969 | 379.1 | 9.6 | 1.3 | 0.8 | 2.4 | 8.7 | 61.0 | 228.7 | 638.3 | 1,566.4 | 3,781.1 | 9,323.9 |
| 1968 | 384.7 | 9.7 | 1.3 | 0.8 | 2.3 | 9.1 | 62.1 | 236.3 | 660.2 | 1,609.4 | 3,868.4 | 9,554.9 |

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| 390-398, 404, 410-429 Diseases of heart | | | | | | | | | | | | |
| WHITE, MALE | | | | | | | | | | | | |
| 1978 | 384.2 | 23.9 | 1.8 | 0.9 | 2.7 | 9.3 | 66.2 | 288.2 | 769.2 | 1,766.3 | 3,904.8 | 8,672.2 |
| 1977 | 386.4 | 23.1 | 1.7 | 0.8 | 2.5 | 9.9 | 70.2 | 297.9 | 784.3 | 1,800.6 | 3,963.4 | 8,650.3 |
| 1976 | 394.2 | 21.8 | 1.8 | 0.8 | 2.7 | 9.8 | 74.0 | 308.1 | 822.9 | 1,851.3 | 4,050.9 | 8,850.6 |
| 1975 | 396.7 | 18.8 | 1.6 | 0.9 | 2.6 | 9.7 | 76.8 | 318.3 | 847.0 | 1,889.1 | 4,090.1 | 8,692.8 |
| 1974 | 411.8 | 18.7 | 1.5 | 0.8 | 2.9 | 10.3 | 80.2 | 334.2 | 885.0 | 1,985.5 | 4,271.9 | 9,359.2 |
| 1973 | 427.9 | 18.2 | 1.9 | 0.9 | 2.9 | 11.0 | 86.6 | 345.3 | 940.0 | 2,064.6 | 4,509.1 | 9,723.3 |
| 1972 | 431.9 | 14.7 | 1.9 | 0.9 | 2.8 | 11.0 | 88.9 | 355.2 | 957.8 | 2,118.5 | 4,546.2 | 9,616.0 |
| 1971 | 432.6 | 12.0 | 1.5 | 0.8 | 3.1 | 12.1 | 93.5 | 357.5 | 961.2 | 2,116.8 | 4,538.5 | 9,828.7 |
| 1970 | 438.3 | 12.0 | 1.5 | 0.8 | 3.0 | 12.3 | 94.6 | 365.7 | 979.3 | 2,177.2 | 4,617.6 | 8,818.0 |
| 1969 | 445.7 | 9.8 | 1.3 | 0.8 | 2.9 | 12.3 | 99.0 | 374.8 | 995.6 | 2,213.8 | 4,658.5 | 10,233.3 |
| 1968 | 453.2 | 11.2 | 1.3 | 0.9 | 2.8 | 12.6 | 100.0 | 384.4 | 1,020.8 | 2,262.9 | 4,736.4 | 10,424.6 |
| WHITE, FEMALE | | | | | | | | | | | | |
| 1978 | 303.1 | 18.5 | 1.8 | 0.9 | 1.6 | 3.8 | 16.8 | 71.2 | 247.6 | 789.3 | 2,492.1 | 7,180.6 |
| 1977 | 297.2 | 15.7 | 1.5 | 0.7 | 1.5 | 3.9 | 17.8 | 73.6 | 252.4 | 794.8 | 2,512.1 | 7,091.7 |
| 1976 | 301.5 | 15.1 | 1.4 | 0.8 | 1.4 | 3.7 | 18.2 | 75.5 | 263.3 | 819.6 | 2,607.1 | 7,390.4 |
| 1975 | 297.9 | 15.6 | 1.4 | 0.7 | 1.6 | 4.1 | 19.9 | 78.0 | 267.7 | 850.6 | 2,648.9 | 7,230.6 |
| 1974 | 309.6 | 14.3 | 1.7 | 0.7 | 1.6 | 4.4 | 20.6 | 80.0 | 281.1 | 903.5 | 2,828.7 | 7,874.3 |
| 1973 | 317.2 | 14.7 | 1.5 | 0.9 | 1.7 | 4.9 | 22.6 | 84.5 | 297.7 | 944.7 | 2,980.5 | 8,222.0 |
| 1972 | 319.4 | 14.5 | 1.7 | 0.7 | 1.6 | 4.9 | 23.2 | 84.0 | 303.7 | 1,004.2 | 3,054.6 | 8,266.1 |
| 1971 | 315.2 | 9.8 | 1.5 | 0.8 | 1.7 | 4.8 | 23.6 | 89.7 | 313.7 | 1,000.4 | 3,062.5 | 8,260.5 |
| 1970 | 313.8 | 7.0 | 1.2 | 0.7 | 1.7 | 5.5 | 23.9 | 91.4 | 317.7 | 1,044.0 | 3,143.5 | 7,839.9 |
| 1969 | 315.7 | 9.5 | 1.2 | 0.8 | 1.8 | 5.3 | 24.5 | 91.4 | 316.1 | 1,062.8 | 3,197.3 | 8,853.9 |
| 1968 | 319.4 | 8.1 | 1.3 | 0.8 | 1.8 | 5.7 | 25.8 | 96.6 | 334.0 | 1,097.6 | 3,280.0 | 9,098.2 |

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| 390-398, 404, 410-429 Diseases of heart | | | | | | | | | | | | |
| ALL OTHER, BOTH SEXES | | | | | | | | | | | | |
| 1978 | 238.3 | 45.0 | 4.3 | 1.6 | 5.2 | 17.7 | 82.9 | 279.7 | 673.8 | 1,357.7 | 2,606.9 | 5,385.7 |
| 1977 | 239.2 | 35.8 | 3.3 | 1.3 | 5.1 | 17.6 | 86.8 | 287.1 | 686.9 | 1,394.9 | 2,574.3 | 5,438.8 |
| 1976 | 241.0 | 41.9 | 2.9 | 1.4 | 5.5 | 18.7 | 84.6 | 293.9 | 698.7 | 1,443.0 | 2,567.6 | 5,435.3 |
| 1975 | 241.3 | 33.0 | 3.0 | 1.4 | 5.3 | 19.5 | 94.7 | 293.3 | 709.4 | 1,510.8 | 2,555.8 | 5,306.2 |
| 1974 | 255.0 | 46.4 | 3.5 | 1.6 | 5.5 | 22.8 | 95.4 | 319.9 | 762.9 | 1,641.1 | 2,657.4 | 5,831.1 |
| 1973 | 268.9 | 52.9 | 4.2 | 1.7 | 6.1 | 24.1 | 105.3 | 341.2 | 813.6 | 1,739.6 | 2,850.5 | 6,038.6 |
| 1972 | 269.9 | 48.4 | 3.4 | 1.7 | 6.7 | 25.1 | 114.7 | 352.9 | 838.9 | 1,724.8 | 2,906.7 | 5,726.8 |
| 1971 | 268.0 | 38.2 | 3.5 | 1.8 | 6.9 | 26.4 | 117.3 | 344.1 | 816.4 | 1,737.7 | 2,972.9 | 5,639.2 |
| 1970 | 274.2 | 31.8 | 3.8 | 1.5 | 6.7 | 29.1 | 126.3 | 363.7 | 854.0 | 1,757.8 | 3,141.6 | 4,866.9 |
| 1969 | 281.2 | 20.4 | 2.5 | 2.0 | 7.0 | 31.0 | 132.3 | 378.8 | 882.3 | 1,806.7 | 3,186.4 | 5,783.5 |
| 1968 | 292.4 | 26.6 | 3.4 | 2.0 | 6.5 | 31.9 | 137.5 | 391.0 | 945.9 | 1,866.7 | 3,268.3 | 6,190.2 |
| ALL OTHER, MALE | | | | | | | | | | | | |
| 1978 | 269.1 | 44.3 | 3.8 | 1.6 | 6.0 | 24.7 | 120.8 | 396.0 | 905.2 | 1,704.1 | 3,070.7 | 5,969.0 |
| 1977 | 269.0 | 38.0 | 3.1 | 1.2 | 6.4 | 24.9 | 122.3 | 403.4 | 918.7 | 1,720.3 | 2,996.9 | 5,992.9 |
| 1976 | 272.6 | 43.2 | 3.5 | 1.5 | 6.7 | 25.5 | 122.0 | 413.5 | 929.7 | 1,784.7 | 2,995.3 | 5,967.3 |
| 1975 | 273.9 | 35.0 | 3.5 | 1.5 | 6.2 | 28.4 | 133.8 | 411.0 | 915.6 | 1,842.4 | 3,073.1 | 5,827.8 |
| 1974 | 287.8 | 45.9 | 3.6 | 1.4 | 6.1 | 30.1 | 128.8 | 437.3 | 992.6 | 2,003.0 | 3,143.7 | 6,272.0 |
| 1973 | 303.3 | 61.3 | 4.7 | 2.0 | 6.9 | 30.9 | 140.4 | 465.1 | 1,048.8 | 2,083.7 | 3,377.2 | 6,433.3 |
| 1972 | 306.7 | 52.7 | 4.2 | 1.8 | 8.7 | 32.6 | 152.7 | 478.6 | 1,057.4 | 2,082.9 | 3,458.1 | 6,212.8 |
| 1971 | 302.9 | 37.6 | 3.9 | 1.9 | 8.1 | 32.3 | 155.8 | 453.5 | 1,043.3 | 2,069.7 | 3,438.2 | 6,055.3 |
| 1970 | 310.2 | 32.2 | 3.7 | 1.5 | 7.7 | 36.9 | 171.0 | 473.2 | 1,062.3 | 2,104.8 | 3,573.1 | 4,983.6 |
| 1969 | 319.6 | 19.4 | 2.9 | 1.8 | 7.3 | 40.5 | 176.4 | 503.3 | 1,089.4 | 2,149.5 | 3,691.5 | 6,018.2 |
| 1968 | 330.0 | 25.7 | 3.7 | 2.2 | 7.0 | 39.4 | 179.0 | 505.3 | 1,169.5 | 2,205.8 | 3,648.4 | 6,704.9 |

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| 390-398, 404, 410-429 Diseases of heart | | | | | | | | | | | | |
| ALL OTHER, FEMALE | | | | | | | | | | | | |
| 1978 | 210.3 | 45.7 | 4.9 | 1.6 | 4.4 | 11.7 | 51.1 | 183.7 | 481.7 | 1,089.6 | 2,296.4 | 5,124.8 |
| 1977 | 212.2 | 33.6 | 3.5 | 1.4 | 4.0 | 11.4 | 57.0 | 190.7 | 493.5 | 1,139.2 | 2,279.6 | 5,140.4 |
| 1976 | 212.3 | 40.6 | 2.2 | 1.2 | 4.3 | 12.9 | 53.3 | 193.9 | 505.3 | 1,172.8 | 2,275.3 | 5,136.7 |
| 1975 | 211.7 | 30.9 | 2.5 | 1.3 | 4.4 | 11.8 | 61.9 | 193.9 | 535.2 | 1,247.0 | 2,194.9 | 4,946.2 |
| 1974 | 225.1 | 47.0 | 3.5 | 1.7 | 4.9 | 16.6 | 67.5 | 220.4 | 567.4 | 1,351.1 | 2,322.0 | 5,571.8 |
| 1973 | 237.5 | 44.2 | 3.6 | 1.4 | 5.4 | 18.2 | 76.0 | 235.2 | 613.2 | 1,461.8 | 2,488.4 | 5,798.7 |
| 1972 | 236.2 | 44.0 | 2.5 | 1.6 | 4.9 | 18.6 | 82.9 | 244.7 | 650.4 | 1,436.7 | 2,518.9 | 5,426.3 |
| 1971 | 235.9 | 38.8 | 3.2 | 1.7 | 5.7 | 21.3 | 85.3 | 248.9 | 618.1 | 1,466.2 | 2,628.9 | 5,371.2 |
| 1970 | 241.0 | 31.4 | 3.9 | 1.6 | 5.8 | 22.4 | 89.1 | 267.6 | 669.8 | 1,471.8 | 2,813.2 | 4,784.7 |
| 1969 | 246.0 | 21.5 | 2.2 | 2.3 | 6.8 | 23.0 | 95.5 | 269.5 | 698.7 | 1,524.5 | 2,789.1 | 5,624.6 |
| 1968 | 257.8 | 27.5 | 3.2 | 1.8 | 6.1 | 25.5 | 102.7 | 290.3 | 745.2 | 1,585.4 | 2,986.2 | 5,844.3 |
| BLACK, BOTH SEXES | | | | | | | | | | | | |
| 1978 | 260.6 | 48.3 | 4.5 | 1.6 | 5.6 | 20.0 | 94.2 | 311.4 | 729.1 | 1,439.9 | 2,672.2 | 5,553.3 |
| 1977 | 261.7 | 39.0 | 3.5 | 1.5 | 5.4 | 20.1 | 98.2 | 319.6 | 740.4 | 1,486.4 | 2,626.3 | 5,645.5 |
| 1976 | 262.4 | 43.7 | 3.2 | 1.3 | 5.8 | 21.4 | 95.2 | 325.2 | 751.0 | 1,529.2 | 2,629.3 | 5,615.4 |
| 1975 | 261.7 | 35.3 | 3.2 | 1.5 | 5.8 | 22.2 | 105.5 | 322.0 | 757.1 | 1,602.8 | 2,608.2 | 5,508.5 |
| 1974 | 274.3 | 46.5 | 3.7 | 1.7 | 5.8 | 25.5 | 105.9 | 347.8 | 808.9 | 1,733.8 | 2,708.9 | 6,059.5 |
| 1973 | 288.5 | 57.7 | 4.4 | 1.8 | 6.5 | 26.4 | 117.2 | 369.9 | 864.3 | 1,841.3 | 2,902.8 | 6,226.3 |
| 1972 | 288.2 | 52.0 | 3.4 | 1.8 | 7.2 | 27.2 | 126.0 | 382.1 | 884.7 | 1,817.7 | 2,989.1 | 5,932.7 |
| 1971 | 284.4 | 38.8 | 3.6 | 1.8 | 7.2 | 29.2 | 128.6 | 368.5 | 859.8 | 1,825.1 | 3,055.3 | 5,827.1 |
| 1970 | 294.0 | 32.4 | 4.1 | 1.6 | 7.1 | 32.4 | 140.3 | 393.7 | 905.9 | 1,855.8 | 3,310.8 | 5,146.4 |
| 1969 | 297.8 | 20.8 | 2.6 | 2.0 | 7.4 | 34.1 | 143.0 | 404.5 | 931.9 | 1,888.2 | 3,328.8 | 6,026.8 |
| 1968 | 308.5 | 28.5 | 3.7 | 2.0 | 6.5 | 34.9 | 149.0 | 414.7 | 995.6 | 1,950.0 | 3,386.3 | 6,397.8 |

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| 390-398, 404, 410-429 Diseases of heart | | | | | | | | | | | | |
| BLACK, MALE | | | | | | | | | | | | |
| 1978 | 291.0 | 46.8 | 4.2 | 1.6 | 6.4 | 27.9 | 137.1 | 438.2 | 979.1 | 1,817.6 | 3,113.8 | 6,023.5 |
| 1977 | 291.6 | 42.0 | 3.2 | 1.4 | 6.7 | 28.1 | 138.5 | 445.7 | 992.1 | 1,851.1 | 3,058.3 | 6,161.2 |
| 1976 | 294.1 | 45.0 | 3.9 | 1.5 | 7.2 | 29.1 | 137.8 | 454.2 | 1,000.9 | 1,905.0 | 3,053.8 | 6,174.5 |
| 1975 | 293.7 | 36.5 | 3.6 | 1.5 | 6.8 | 32.0 | 148.4 | 448.5 | 975.9 | 1,971.7 | 3,112.6 | 6,102.2 |
| 1974 | 306.6 | 46.5 | 3.7 | 1.5 | 6.5 | 33.4 | 142.6 | 472.7 | 1,055.5 | 2,133.8 | 3,201.0 | 6,551.2 |
| 1973 | 322.7 | 67.0 | 4.8 | 2.2 | 7.3 | 34.5 | 155.0 | 502.0 | 1,121.3 | 2,223.5 | 3,436.8 | 6,746.3 |
| 1972 | 325.0 | 55.6 | 4.0 | 1.7 | 9.2 | 36.0 | 166.7 | 519.7 | 1,124.9 | 2,207.2 | 3,541.1 | 6,560.0 |
| 1971 | 319.2 | 36.9 | 4.2 | 1.9 | 8.5 | 36.3 | 170.8 | 485.3 | 1,105.6 | 2,184.8 | 3,548.4 | 6,335.0 |
| 1970 | 330.3 | 33.5 | 3.9 | 1.4 | 8.3 | 41.6 | 189.2 | 512.8 | 1,135.4 | 2,237.8 | 3,783.4 | 5,367.6 |
| 1969 | 335.4 | 19.2 | 3.0 | 1.7 | 7.6 | 44.2 | 189.7 | 537.3 | 1,158.8 | 2,253.6 | 3,858.7 | 6,421.6 |
| 1968 | 345.4 | 28.2 | 4.0 | 2.3 | 6.8 | 43.5 | 192.7 | 535.4 | 1,240.4 | 2,309.5 | 3,797.6 | 7,125.7 |
| BLACK, FEMALE | | | | | | | | | | | | |
| 1978 | 233.2 | 49.8 | 4.8 | 1.7 | 4.9 | 13.1 | 58.6 | 207.2 | 525.0 | 1,157.9 | 2,377.4 | 5,311.1 |
| 1977 | 234.8 | 35.8 | 3.7 | 1.5 | 4.2 | 13.2 | 64.8 | 215.3 | 534.9 | 1,212.6 | 2,346.9 | 5,376.6 |
| 1976 | 233.8 | 42.5 | 2.4 | 1.2 | 4.6 | 14.8 | 59.9 | 218.0 | 544.3 | 1,243.1 | 2,344.7 | 5,320.2 |
| 1975 | 232.7 | 34.1 | 2.7 | 1.4 | 4.8 | 13.7 | 69.9 | 216.0 | 575.5 | 1,323.1 | 2,266.8 | 5,183.3 |
| 1974 | 245.1 | 46.6 | 3.7 | 1.8 | 5.2 | 18.7 | 75.4 | 242.9 | 604.4 | 1,429.7 | 2,376.1 | 5,788.5 |
| 1973 | 257.4 | 48.0 | 3.9 | 1.5 | 5.7 | 19.4 | 85.8 | 257.6 | 650.5 | 1,545.9 | 2,544.6 | 5,934.2 |
| 1972 | 254.8 | 48.1 | 2.8 | 1.8 | 5.4 | 19.8 | 92.2 | 264.8 | 683.1 | 1,517.3 | 2,590.0 | 5,574.3 |
| 1971 | 252.8 | 40.8 | 2.9 | 1.7 | 6.0 | 23.2 | 93.7 | 268.2 | 651.8 | 1,538.8 | 2,701.2 | 5,523.9 |
| 1970 | 261.0 | 31.3 | 4.2 | 1.8 | 6.0 | 24.7 | 99.8 | 290.9 | 710.5 | 1,553.2 | 2,964.1 | 5,003.8 |
| 1969 | 263.7 | 22.4 | 2.3 | 2.3 | 7.1 | 25.8 | 104.3 | 289.4 | 738.7 | 1,597.6 | 2,924.2 | 5,783.3 |
| 1968 | 274.9 | 28.9 | 3.5 | 1.8 | 6.3 | 27.7 | 112.5 | 309.6 | 783.9 | 1,657.4 | 3,080.6 | 6,057.1 |

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|--|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| ACTIVE RHEUMATIC FEVER AND CHRONIC RHEUMATIC HEART DISEASE (390-398) | | | | | | | | | | | | |
| BOTH SEXES | | | | | | | | | | | | |
| 1978 | 6.0 | * | * | 0.1 | 0.3 | 0.7 | 2.2 | 5.6 | 13.0 | 26.9 | 42.5 | 57.9 |
| 1977 | 5.8 | * | * | 0.1 | 0.4 | 0.8 | 2.5 | 5.9 | 13.4 | 26.2 | 38.0 | 47.4 |
| 1976 | 6.0 | * | * | 0.1 | 0.4 | 0.8 | 2.5 | 6.5 | 15.4 | 27.1 | 37.2 | 44.6 |
| 1975 | 6.0 | * | * | 0.1 | 0.4 | 1.0 | 3.1 | 6.7 | 15.5 | 27.4 | 35.2 | 41.4 |
| 1974 | 6.2 | * | * | 0.2 | 0.5 | 1.1 | 3.1 | 7.8 | 16.6 | 28.2 | 33.8 | 42.6 |
| 1973 | 6.5 | * | * | 0.2 | 0.5 | 1.3 | 3.6 | 8.3 | 18.2 | 28.1 | 34.6 | 42.4 |
| 1972 | 6.7 | * | * | 0.1 | 0.4 | 1.4 | 4.3 | 8.9 | 20.1 | 28.2 | 35.3 | 39.4 |
| 1971 | 7.1 | * | 0.2 | 0.2 | 0.6 | 1.5 | 4.4 | 10.0 | 20.5 | 30.1 | 35.1 | 41.6 |
| 1970 | 7.3 | * | * | 0.2 | 0.7 | 1.8 | 4.8 | 11.0 | 21.1 | 31.0 | 34.5 | 37.6 |
| 1969 | 7.7 | * | * | 0.3 | 0.7 | 1.9 | 5.0 | 11.7 | 23.3 | 30.8 | 34.4 | 43.8 |
| 1968 | 8.2 | * | * | 0.3 | 0.7 | 2.3 | 5.7 | 13.3 | 24.9 | 31.6 | 35.9 | 45.3 |
| MALE | | | | | | | | | | | | |
| 1978 | 5.4 | * | * | * | 0.4 | 0.7 | 2.2 | 5.9 | 13.4 | 27.0 | 43.1 | 58.9 |
| 1977 | 5.3 | * | * | * | 0.4 | 0.9 | 2.4 | 6.0 | 14.0 | 26.5 | 38.5 | 48.0 |
| 1976 | 5.5 | * | * | 0.1 | 0.4 | 1.0 | 2.6 | 6.9 | 15.7 | 28.0 | 38.2 | 40.3 |
| 1975 | 5.5 | * | * | 0.2 | 0.4 | 0.9 | 3.1 | 7.3 | 16.1 | 27.0 | 36.0 | 37.5 |
| 1974 | 5.7 | * | * | 0.1 | 0.5 | 1.2 | 2.9 | 8.5 | 17.4 | 28.5 | 32.6 | 39.5 |
| 1973 | 6.0 | * | * | 0.2 | 0.6 | 1.5 | 3.8 | 8.6 | 18.6 | 27.6 | 34.3 | 37.5 |
| 1972 | 6.4 | * | * | 0.1 | 0.5 | 1.4 | 4.5 | 9.7 | 20.6 | 28.5 | 36.7 | 40.2 |
| 1971 | 6.7 | * | * | 0.2 | 0.7 | 1.5 | 4.6 | 10.2 | 21.2 | 30.4 | 36.3 | 39.6 |
| 1970 | 7.0 | * | * | 0.2 | 0.7 | 1.8 | 4.9 | 11.4 | 21.5 | 31.9 | 34.8 | 34.7 |
| 1969 | 7.2 | * | * | 0.3 | 0.8 | 1.9 | 5.1 | 12.0 | 23.7 | 30.5 | 34.2 | 42.0 |
| 1968 | 7.7 | * | * | 0.3 | 0.8 | 2.4 | 5.8 | 13.4 | 25.3 | 31.6 | 33.8 | 46.6 |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|--|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| ACTIVE RHEUMATIC FEVER AND CHRONIC RHEUMATIC HEART DISEASE (390-398) | | | | | | | | | | | | |
| FEMALE | | | | | | | | | | | | |
| 1978 | 6.6 | * | * | * | 0.2 | 0.6 | 2.1 | 5.4 | 12.7 | 26.8 | 42.2 | 57.4 |
| 1977 | 6.3 | * | * | 0.1 | 0.3 | 0.7 | 2.5 | 5.8 | 12.9 | 26.0 | 37.8 | 47.1 |
| 1976 | 6.5 | * | * | * | 0.3 | 0.7 | 2.5 | 6.2 | 15.2 | 26.4 | 36.6 | 46.7 |
| 1975 | 6.5 | * | * | * | 0.4 | 1.0 | 3.2 | 6.1 | 15.0 | 27.7 | 34.7 | 43.2 |
| 1974 | 6.7 | * | * | 0.2 | 0.4 | 1.0 | 3.3 | 7.2 | 15.9 | 28.0 | 34.5 | 44.2 |
| 1973 | 7.0 | * | * | 0.2 | 0.4 | 1.2 | 3.5 | 8.1 | 17.8 | 28.5 | 34.8 | 44.9 |
| 1972 | 7.1 | * | * | 0.1 | 0.3 | 1.4 | 4.1 | 8.1 | 19.5 | 28.0 | 34.4 | 39.0 |
| 1971 | 7.5 | * | * | 0.2 | 0.5 | 1.4 | 4.3 | 9.9 | 19.9 | 29.8 | 34.3 | 42.6 |
| 1970 | 7.7 | * | * | 0.2 | 0.6 | 1.7 | 4.6 | 10.6 | 20.8 | 30.2 | 34.3 | 39.2 |
| 1969 | 8.1 | * | * | 0.3 | 0.7 | 2.0 | 4.8 | 11.3 | 23.0 | 31.1 | 34.6 | 44.7 |
| 1968 | 8.6 | * | * | 0.3 | 0.6 | 2.3 | 5.6 | 13.2 | 24.5 | 31.6 | 37.3 | 44.6 |
| WHITE, BOTH SEXES | | | | | | | | | | | | |
| 1978 | 6.4 | * | * | * | 0.2 | 0.5 | 1.9 | 5.4 | 13.2 | 28.1 | 43.8 | 60.2 |
| 1977 | 6.1 | * | * | 0.1 | 0.3 | 0.7 | 2.2 | 5.7 | 13.7 | 27.2 | 39.6 | 49.3 |
| 1976 | 6.3 | * | * | 0.1 | 0.2 | 0.7 | 2.3 | 6.4 | 15.7 | 28.1 | 38.7 | 45.5 |
| 1975 | 6.3 | * | * | 0.1 | 0.2 | 0.8 | 2.8 | 6.6 | 15.8 | 28.2 | 36.4 | 43.0 |
| 1974 | 6.5 | * | * | 0.1 | 0.3 | 1.0 | 2.9 | 7.7 | 17.0 | 29.3 | 35.0 | 44.3 |
| 1973 | 6.8 | * | * | 0.1 | 0.4 | 1.1 | 3.4 | 8.4 | 18.5 | 29.2 | 36.0 | 43.4 |
| 1972 | 7.0 | * | * | 0.1 | 0.3 | 1.1 | 3.9 | 8.9 | 20.5 | 29.2 | 36.3 | 40.2 |
| 1971 | 7.4 | * | * | 0.1 | 0.4 | 1.3 | 4.1 | 10.0 | 21.0 | 31.2 | 36.4 | 42.5 |
| 1970 | 7.6 | * | * | 0.2 | 0.5 | 1.4 | 4.5 | 10.9 | 21.6 | 32.1 | 35.5 | 39.4 |
| 1969 | 8.0 | * | * | 0.2 | 0.5 | 1.7 | 4.6 | 11.8 | 23.9 | 31.8 | 35.5 | 45.3 |
| 1968 | 8.5 | * | * | 0.2 | 0.6 | 1.9 | 5.4 | 13.3 | 25.6 | 32.9 | 36.9 | 47.1 |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|--|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| ACTIVE RHEUMATIC FEVER AND CHRONIC RHEUMATIC HEART DISEASE (390-398) | | | | | | | | | | | | |
| WHITE, MALE | | | | | | | | | | | | |
| 1978 | 5.6 | * | * | * | 0.3 | 0.6 | 1.9 | 5.8 | 13.6 | 28.0 | 44.6 | 61.8 |
| 1977 | 5.5 | * | * | * | 0.3 | 0.8 | 2.3 | 5.7 | 14.2 | 27.5 | 40.3 | 49.6 |
| 1976 | 5.7 | * | * | * | 0.3 | 0.8 | 2.3 | 6.8 | 15.8 | 29.0 | 39.3 | 42.1 |
| 1975 | 5.7 | * | * | 0.1 | 0.3 | 0.7 | 2.8 | 7.1 | 16.3 | 27.6 | 36.9 | 39.3 |
| 1974 | 5.9 | * | * | * | 0.4 | 1.0 | 2.7 | 8.2 | 17.4 | 29.3 | 33.4 | 41.2 |
| 1973 | 6.2 | * | * | 0.2 | 0.5 | 1.2 | 3.5 | 8.6 | 18.7 | 28.2 | 35.5 | 37.8 |
| 1972 | 6.6 | * | * | * | 0.4 | 1.0 | 4.0 | 9.8 | 21.2 | 29.2 | 37.4 | 40.8 |
| 1971 | 6.9 | * | * | 0.1 | 0.5 | 1.4 | 4.4 | 10.1 | 21.6 | 31.4 | 37.5 | 41.0 |
| 1970 | 7.2 | * | * | 0.2 | 0.5 | 1.4 | 4.6 | 11.3 | 21.8 | 33.1 | 35.9 | 35.5 |
| 1969 | 7.5 | * | * | 0.2 | 0.7 | 1.7 | 4.8 | 12.1 | 24.2 | 31.3 | 34.9 | 44.1 |
| 1968 | 7.9 | * | * | 0.2 | 0.7 | 1.9 | 5.4 | 13.5 | 25.7 | 32.7 | 34.3 | 47.9 |
| WHITE, FEMALE | | | | | | | | | | | | |
| 1978 | 7.1 | * | * | * | 0.2 | 0.5 | 1.9 | 5.1 | 12.8 | 28.1 | 43.4 | 59.5 |
| 1977 | 6.7 | * | * | * | 0.2 | 0.5 | 2.2 | 5.7 | 13.2 | 26.9 | 39.2 | 49.2 |
| 1976 | 6.9 | * | * | * | 0.2 | 0.6 | 2.2 | 6.1 | 15.5 | 27.5 | 38.3 | 47.1 |
| 1975 | 6.9 | * | * | * | 0.2 | 0.8 | 2.8 | 6.1 | 15.4 | 28.7 | 36.1 | 44.8 |
| 1974 | 7.1 | * | * | * | 0.3 | 0.9 | 3.0 | 7.1 | 16.6 | 29.2 | 36.0 | 45.8 |
| 1973 | 7.4 | * | * | 0.1 | 0.3 | 0.9 | 3.3 | 8.1 | 18.3 | 30.0 | 36.4 | 46.3 |
| 1972 | 7.4 | * | * | * | 0.2 | 1.2 | 3.8 | 8.0 | 20.0 | 29.2 | 35.6 | 39.8 |
| 1971 | 7.8 | * | * | 0.1 | 0.4 | 1.1 | 3.8 | 9.8 | 20.5 | 31.0 | 35.6 | 43.3 |
| 1970 | 8.0 | * | * | * | 0.5 | 1.3 | 4.4 | 10.6 | 21.4 | 31.4 | 35.3 | 41.6 |
| 1969 | 8.4 | * | * | 0.1 | 0.4 | 1.6 | 4.4 | 11.4 | 23.7 | 32.3 | 35.9 | 46.0 |
| 1968 | 9.1 | * | * | 0.1 | 0.5 | 1.8 | 5.4 | 13.2 | 25.5 | 33.0 | 38.7 | 46.8 |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|--|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| ACTIVE RHEUMATIC FEVER AND CHRONIC RHEUMATIC HEART DISEASE (390-398) | | | | | | | | | | | | |
| ALL OTHER, BOTH SEXES | | | | | | | | | | | | |
| 1978 | 3.9 | * | * | * | 0.8 | 1.6 | 3.7 | 7.2 | 11.6 | 16.3 | 29.0 | 31.0 |
| 1977 | 3.8 | * | * | * | 0.9 | 2.0 | 4.0 | 7.7 | 10.6 | 17.4 | 21.6 | 25.6 |
| 1976 | 4.0 | * | * | * | 1.0 | 1.9 | 4.4 | 7.2 | 12.9 | 17.2 | 22.0 | 34.6 |
| 1975 | 4.2 | * | * | * | 1.2 | 2.1 | 5.8 | 7.3 | 12.6 | 18.8 | 22.5 | 22.6 |
| 1974 | 4.3 | * | * | 0.5 | 1.3 | 2.3 | 5.1 | 9.2 | 13.4 | 17.7 | 20.5 | 23.0 |
| 1973 | 4.5 | * | * | 0.4 | 1.4 | 3.1 | 5.5 | 8.2 | 15.1 | 17.5 | 18.2 | 30.7 |
| 1972 | 4.9 | * | * | 0.5 | 1.1 | 3.4 | 7.1 | 8.8 | 15.4 | 18.4 | 24.3 | 30.9 |
| 1971 | 5.1 | * | * | 0.5 | 1.6 | 3.0 | 7.0 | 10.6 | 15.6 | 19.3 | 19.7 | 30.8 |
| 1970 | 5.4 | * | * | 0.6 | 1.8 | 4.4 | 6.6 | 11.2 | 16.9 | 19.9 | 20.8 | 18.6 |
| 1969 | 5.6 | * | * | 1.0 | 2.1 | 3.8 | 7.5 | 10.8 | 17.3 | 20.7 | 20.3 | 26.6 |
| 1968 | 6.0 | * | * | 1.0 | 1.7 | 5.8 | 8.3 | 12.9 | 18.0 | 18.8 | 22.3 | 24.5 |
| ALL OTHER, MALE | | | | | | | | | | | | |
| 1978 | 3.7 | * | * | * | 1.0 | 1.7 | 4.3 | 6.9 | 10.8 | 18.3 | 28.6 | * |
| 1977 | 3.7 | * | * | * | 1.1 | 2.2 | 3.6 | 8.3 | 11.6 | 17.7 | 21.1 | * |
| 1976 | 4.2 | * | * | * | 1.1 | 2.5 | 4.5 | 7.9 | 14.3 | 18.7 | 28.0 | * |
| 1975 | 4.4 | * | * | * | 1.2 | 2.5 | 5.9 | 8.6 | 13.7 | 21.0 | 27.7 | * |
| 1974 | 4.7 | * | * | * | 1.2 | 2.7 | 5.1 | 10.3 | 17.5 | 20.2 | 24.5 | * |
| 1973 | 4.9 | * | * | * | 1.4 | 3.2 | 6.4 | 8.9 | 17.8 | 22.0 | 21.4 | * |
| 1972 | 5.2 | * | * | * | 1.2 | 4.3 | 8.2 | 8.9 | 15.6 | 21.8 | 29.5 | * |
| 1971 | 5.0 | * | * | * | 1.7 | 2.4 | 6.4 | 10.5 | 17.7 | 21.3 | 22.7 | * |
| 1970 | 5.5 | * | * | * | 2.0 | 4.6 | 7.0 | 11.8 | 18.1 | 21.1 | 21.2 | * |
| 1969 | 5.4 | * | * | 0.7 | 1.5 | 3.2 | 7.2 | 11.3 | 18.5 | 22.7 | 25.0 | * |
| 1968 | 6.3 | * | * | 1.0 | 1.8 | 5.8 | 9.4 | 12.5 | 21.1 | 21.0 | 27.4 | * |
| ALL OTHER, FEMALE | | | | | | | | | | | | |
| 1978 | 4.1 | * | * | * | * | 1.6 | 3.2 | 7.4 | 12.3 | 14.8 | 29.3 | 32.1 |
| 1977 | 3.9 | * | * | * | 0.7 | 1.8 | 4.4 | 7.3 | 9.8 | 17.1 | 21.9 | 22.1 |
| 1976 | 3.9 | * | * | * | 0.9 | 1.4 | 4.4 | 6.6 | 11.6 | 16.1 | 17.8 | 41.8 |
| 1975 | 4.0 | * | * | * | 1.1 | 1.8 | 5.7 | 6.2 | 11.6 | 17.0 | 18.9 | 23.7 |
| 1974 | 4.0 | * | * | * | 1.3 | 2.0 | 5.2 | 8.3 | 9.8 | 15.6 | 17.8 | 23.5 |
| 1973 | 4.2 | * | * | * | 1.4 | 3.1 | 4.8 | 7.6 | 12.9 | 13.9 | 16.0 | 27.8 |
| 1972 | 4.6 | * | * | * | 1.1 | 2.6 | 6.1 | 8.6 | 15.3 | 15.6 | 20.6 | 28.9 |
| 1971 | 5.1 | * | * | * | 1.5 | 3.5 | 7.6 | 10.7 | 13.8 | 17.6 | 17.5 | 34.2 |
| 1970 | 5.3 | * | * | * | 1.6 | 4.3 | 6.2 | 10.6 | 15.9 | 19.0 | 20.5 | * |
| 1969 | 5.7 | * | * | 1.3 | 2.6 | 4.4 | 7.7 | 10.4 | 16.4 | 19.2 | 16.6 | * |
| 1968 | 5.7 | * | * | 0.9 | 1.7 | 5.8 | 7.3 | 13.2 | 15.3 | 17.1 | 18.4 | * |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|--|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| ACTIVE RHEUMATIC FEVER AND CHRONIC RHEUMATIC HEART DISEASE (390-398) | | | | | | | | | | | | |
| BLACK, BOTH SEXES | | | | | | | | | | | | |
| 1978 | 4.1 | * | * | * | 0.9 | 1.8 | 4.0 | 7.4 | 12.2 | 16.4 | 27.8 | 28.7 |
| 1977 | 4.0 | * | * | * | 0.9 | 2.1 | 4.2 | 8.2 | 10.8 | 17.9 | 21.2 | 25.2 |
| 1976 | 4.2 | * | * | * | 0.9 | 2.1 | 4.7 | 7.7 | 13.4 | 17.7 | 22.0 | 35.3 |
| 1975 | 4.3 | * | * | * | 1.3 | 2.3 | 6.1 | 7.8 | 12.5 | 18.3 | 21.5 | 23.1 |
| 1974 | 4.5 | * | * | 0.5 | 1.3 | 2.4 | 5.6 | 9.5 | 13.7 | 17.7 | 19.3 | 19.8 |
| 1973 | 4.6 | * | * | 0.5 | 1.4 | 3.3 | 6.0 | 8.4 | 15.3 | 17.5 | 18.2 | 28.9 |
| 1972 | 4.9 | * | * | 0.5 | 1.2 | 3.4 | 7.4 | 8.7 | 14.6 | 18.5 | 24.4 | 34.5 |
| 1971 | 5.1 | * | * | 0.5 | 1.6 | 3.1 | 7.3 | 11.0 | 15.3 | 19.5 | 19.6 | 30.8 |
| 1970 | 5.5 | * | * | 0.6 | 1.8 | 4.7 | 6.8 | 11.7 | 16.9 | 20.5 | 20.8 | 19.6 |
| 1969 | 5.7 | * | * | 1.0 | 2.1 | 4.1 | 7.9 | 11.0 | 17.5 | 20.3 | 21.1 | 24.7 |
| 1968 | 6.0 | * | * | 1.0 | 1.6 | 6.1 | 8.4 | 13.4 | 17.9 | 18.6 | 22.0 | 26.1 |
| BLACK, MALE | | | | | | | | | | | | |
| 1978 | 3.9 | * | * | * | 1.1 | 1.8 | 4.7 | 7.4 | 12.0 | 18.0 | 27.6 | * |
| 1977 | 3.9 | * | * | * | 1.1 | 2.3 | 3.9 | 8.9 | 12.1 | 18.5 | 22.6 | * |
| 1976 | 4.3 | * | * | * | 1.1 | 2.8 | 4.6 | 8.3 | 15.2 | 19.1 | 27.8 | * |
| 1975 | 4.5 | * | * | * | 1.3 | 2.8 | 6.4 | 9.4 | 13.5 | 20.9 | 25.2 | * |
| 1974 | 4.7 | * | * | * | 1.1 | 2.8 | 5.6 | 10.4 | 18.1 | 19.8 | 23.4 | * |
| 1973 | 5.0 | * | * | * | 1.6 | 3.6 | 6.8 | 9.1 | 18.0 | 21.9 | 22.5 | * |
| 1972 | 5.3 | * | * | * | 1.2 | 4.5 | 9.1 | 9.3 | 14.6 | 22.2 | 29.4 | * |
| 1971 | 5.2 | * | * | * | 1.7 | 2.7 | 6.9 | 11.2 | 17.9 | 22.5 | 21.5 | * |
| 1970 | 5.6 | * | * | * | 2.1 | 5.0 | 7.3 | 12.6 | 18.3 | 21.7 | 19.6 | * |
| 1969 | 5.4 | * | * | 0.7 | 1.5 | 3.3 | 7.8 | 11.6 | 18.4 | 22.2 | 26.3 | * |
| 1968 | 6.4 | * | * | 1.1 | 1.6 | 6.2 | 9.5 | 13.1 | 20.7 | 21.9 | 27.3 | * |
| BLACK, FEMALE | | | | | | | | | | | | |
| 1978 | 4.2 | * | * | * | * | 1.7 | 3.4 | 7.4 | 12.4 | 15.2 | 27.9 | 29.3 |
| 1977 | 4.0 | * | * | * | * | 1.9 | 4.5 | 7.6 | 9.6 | 17.5 | 20.3 | 24.5 |
| 1976 | 4.1 | * | * | * | 0.8 | 1.5 | 4.7 | 7.2 | 11.8 | 16.6 | 18.1 | 42.7 |
| 1975 | 4.1 | * | * | * | 1.2 | 2.0 | 5.8 | 6.5 | 11.8 | 16.3 | 18.9 | 25.0 |
| 1974 | 4.2 | * | * | * | 1.4 | 2.0 | 5.6 | 8.8 | 9.9 | 16.1 | 16.5 | * |
| 1973 | 4.3 | * | * | * | 1.4 | 3.1 | 5.3 | 7.8 | 13.1 | 14.2 | 15.2 | * |
| 1972 | 4.5 | * | * | * | 1.2 | 2.4 | 5.9 | 8.1 | 14.6 | 15.6 | 20.7 | 31.4 |
| 1971 | 5.1 | * | * | * | 1.5 | 3.5 | 7.6 | 10.8 | 13.1 | 17.0 | 18.1 | 35.8 |
| 1970 | 5.4 | * | * | * | 1.6 | 4.4 | 6.3 | 10.9 | 15.8 | 19.6 | 21.7 | * |
| 1969 | 5.9 | * | * | 1.3 | 2.7 | 4.8 | 8.0 | 10.6 | 16.7 | 18.7 | 17.2 | * |
| 1968 | 5.7 | * | * | 0.9 | 1.6 | 6.1 | 7.4 | 13.6 | 15.6 | 16.0 | 18.0 | * |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| HYPERTENSIVE HEART DISEASE (402) | | | | | | | | | | | | |
| BOTH SEXES | | | | | | | | | | | | |
| 1978 | 3.2 | * | * | * | * | 0.1 | 0.7 | 2.1 | 5.3 | 12.0 | 28.4 | 64.1 |
| 1977 | 3.1 | * | * | * | * | 0.2 | 0.7 | 2.1 | 4.9 | 12.2 | 28.8 | 64.2 |
| 1976 | 3.1 | * | * | * | * | 0.1 | 0.6 | 2.1 | 5.0 | 11.8 | 27.8 | 67.9 |
| 1975 | 3.1 | * | * | * | * | 0.2 | 0.6 | 2.0 | 5.1 | 12.5 | 28.8 | 67.4 |
| 1974 | 3.4 | * | * | * | * | 0.1 | 0.7 | 2.2 | 5.4 | 13.9 | 32.7 | 74.9 |
| 1973 | 3.4 | * | * | * | * | 0.2 | 0.7 | 2.4 | 5.8 | 13.8 | 33.8 | 78.3 |
| 1972 | 3.7 | * | * | * | * | 0.2 | 0.9 | 2.6 | 6.4 | 14.8 | 35.5 | 90.0 |
| 1971 | 3.8 | * | * | * | * | 0.2 | 1.0 | 2.6 | 6.6 | 14.9 | 38.6 | 90.8 |
| 1970 | 4.1 | * | * | * | * | 0.2 | 1.2 | 3.1 | 7.1 | 16.6 | 42.3 | 90.0 |
| 1969 | 4.5 | * | * | * | * | 0.3 | 1.2 | 3.0 | 7.7 | 18.0 | 46.4 | 110.9 |
| 1968 | 4.9 | * | * | * | 0.1 | 0.3 | 1.3 | 3.4 | 8.6 | 20.8 | 50.7 | 123.4 |
| MALE | | | | | | | | | | | | |
| 1978 | 2.7 | * | * | * | * | 0.2 | 0.9 | 2.7 | 6.1 | 13.1 | 25.3 | 47.9 |
| 1977 | 2.6 | * | * | * | * | 0.2 | 0.9 | 2.4 | 5.8 | 12.9 | 25.0 | 49.8 |
| 1976 | 2.6 | * | * | * | * | 0.2 | 0.7 | 2.5 | 6.0 | 12.9 | 24.7 | 51.7 |
| 1975 | 2.6 | * | * | * | * | 0.2 | 0.7 | 2.3 | 6.0 | 13.3 | 24.4 | 48.3 |
| 1974 | 2.8 | * | * | * | * | * | 0.7 | 2.5 | 6.3 | 15.8 | 27.8 | 55.4 |
| 1973 | 2.8 | * | * | * | * | 0.2 | 0.8 | 2.8 | 6.5 | 14.3 | 30.1 | 53.2 |
| 1972 | 3.0 | * | * | * | * | 0.3 | 1.0 | 3.1 | 7.2 | 13.8 | 29.8 | 65.9 |
| 1971 | 3.2 | * | * | * | * | 0.3 | 1.1 | 2.9 | 7.2 | 15.2 | 32.3 | 76.1 |
| 1970 | 3.5 | * | * | * | * | 0.3 | 1.3 | 3.2 | 8.0 | 17.6 | 36.2 | 61.6 |
| 1969 | 3.8 | * | * | * | * | 0.4 | 1.4 | 3.5 | 8.9 | 19.3 | 37.9 | 84.5 |
| 1968 | 4.1 | * | * | * | * | 0.3 | 1.3 | 3.6 | 9.3 | 20.8 | 42.9 | 92.5 |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| HYPERTENSIVE HEART DISEASE (402) | | | | | | | | | | | | |
| FEMALE | | | | | | | | | | | | |
| 1978 | 3.7 | * | * | * | * | * | 0.5 | 1.6 | 4.5 | 11.3 | 30.2 | 71.4 |
| 1977 | 3.6 | * | * | * | * | * | 0.6 | 1.8 | 4.2 | 11.6 | 31.0 | 70.8 |
| 1976 | 3.5 | * | * | * | * | * | 0.5 | 1.7 | 4.2 | 11.0 | 29.7 | 75.5 |
| 1975 | 3.6 | * | * | * | * | * | 0.5 | 1.8 | 4.4 | 11.9 | 31.4 | 76.7 |
| 1974 | 3.9 | * | * | * | * | 0.1 | 0.6 | 1.9 | 4.7 | 12.4 | 35.7 | 84.5 |
| 1973 | 4.0 | * | * | * | * | 0.2 | 0.7 | 2.0 | 5.2 | 13.4 | 36.1 | 91.0 |
| 1972 | 4.4 | * | * | * | * | 0.1 | 0.7 | 2.2 | 5.7 | 15.6 | 39.2 | 102.4 |
| 1971 | 4.4 | * | * | * | * | 0.2 | 0.9 | 2.3 | 6.1 | 14.7 | 42.6 | 98.5 |
| 1970 | 4.8 | * | * | * | * | 0.2 | 1.1 | 3.0 | 6.4 | 15.9 | 46.3 | 105.9 |
| 1969 | 5.1 | * | * | * | * | 0.2 | 1.1 | 2.5 | 6.7 | 17.1 | 52.2 | 125.2 |
| 1968 | 5.7 | * | * | * | * | 0.2 | 1.2 | 3.3 | 8.0 | 20.7 | 56.0 | 140.5 |
| WHITE, BOTH SEXES | | | | | | | | | | | | |
| 1978 | 2.9 | * | * | * | * | 0.1 | 0.4 | 1.5 | 4.0 | 9.9 | 26.2 | 62.8 |
| 1977 | 2.8 | * | * | * | * | 0.1 | 0.4 | 1.3 | 3.7 | 9.9 | 26.5 | 62.3 |
| 1976 | 2.8 | * | * | * | * | * | 0.3 | 1.4 | 3.7 | 9.9 | 26.1 | 66.1 |
| 1975 | 2.8 | * | * | * | * | 0.1 | 0.3 | 1.3 | 3.9 | 10.3 | 27.0 | 65.0 |
| 1974 | 3.0 | * | * | * | * | 0.1 | 0.4 | 1.4 | 4.0 | 11.2 | 30.7 | 72.9 |
| 1973 | 3.1 | * | * | * | * | * | 0.4 | 1.6 | 4.3 | 11.1 | 32.2 | 77.2 |
| 1972 | 3.2 | * | * | * | * | * | 0.4 | 1.6 | 4.5 | 12.0 | 32.8 | 87.7 |
| 1971 | 3.4 | * | * | * | * | 0.1 | 0.4 | 1.6 | 4.9 | 12.4 | 36.8 | 89.0 |
| 1970 | 3.6 | * | * | * | * | * | 0.5 | 1.8 | 5.2 | 13.7 | 39.7 | 89.3 |
| 1969 | 3.9 | * | * | * | * | 0.1 | 0.5 | 1.8 | 5.6 | 14.7 | 43.9 | 107.8 |
| 1968 | 4.3 | * | * | * | * | * | 0.5 | 2.0 | 6.3 | 17.2 | 47.7 | 120.4 |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| HYPERTENSIVE HEART DISEASE (402) | | | | | | | | | | | | |
| WHITE, MALE | | | | | | | | | | | | |
| 1978 | 2.3 | * | * | * | * | 0.1 | 0.5 | 1.9 | 4.8 | 10.9 | 22.6 | 47.0 |
| 1977 | 2.2 | * | * | * | * | 0.2 | 0.5 | 1.6 | 4.5 | 10.8 | 22.4 | 47.3 |
| 1976 | 2.3 | * | * | * | * | * | 0.3 | 1.7 | 4.6 | 11.0 | 23.3 | 49.4 |
| 1975 | 2.2 | * | * | * | * | * | 0.4 | 1.7 | 4.6 | 10.9 | 22.2 | 45.6 |
| 1974 | 2.5 | * | * | * | * | * | 0.4 | 1.8 | 4.9 | 13.1 | 25.6 | 53.4 |
| 1973 | 2.4 | * | * | * | * | * | 0.5 | 2.0 | 5.1 | 11.5 | 28.1 | 50.2 |
| 1972 | 2.5 | * | * | * | * | * | 0.5 | 2.1 | 5.7 | 11.4 | 26.2 | 61.1 |
| 1971 | 2.7 | * | * | * | * | * | 0.5 | 1.8 | 5.7 | 12.9 | 30.2 | 73.9 |
| 1970 | 2.9 | * | * | * | * | * | 0.5 | 2.0 | 5.9 | 14.7 | 33.1 | 59.8 |
| 1969 | 3.2 | * | * | * | * | * | 0.6 | 2.2 | 6.8 | 16.2 | 35.4 | 81.0 |
| 1968 | 3.5 | * | * | * | * | * | 0.6 | 2.3 | 7.1 | 17.7 | 39.2 | 88.0 |
| WHITE, FEMALE | | | | | | | | | | | | |
| 1978 | 3.4 | * | * | * | * | * | 0.2 | 1.0 | 3.3 | 9.2 | 28.3 | 69.9 |
| 1977 | 3.3 | * | * | * | * | * | 0.2 | 1.1 | 3.0 | 9.3 | 29.0 | 69.1 |
| 1976 | 3.2 | * | * | * | * | * | 0.2 | 1.1 | 2.9 | 9.2 | 27.8 | 73.8 |
| 1975 | 3.3 | * | * | * | * | * | 0.3 | 1.0 | 3.2 | 9.8 | 29.9 | 74.3 |
| 1974 | 3.5 | * | * | * | * | * | 0.3 | 1.0 | 3.2 | 9.7 | 33.7 | 82.4 |
| 1973 | 3.7 | * | * | * | * | * | 0.3 | 1.3 | 3.7 | 10.8 | 34.7 | 90.7 |
| 1972 | 3.9 | * | * | * | * | * | 0.3 | 1.2 | 3.5 | 12.5 | 36.9 | 101.1 |
| 1971 | 4.0 | * | * | * | * | * | 0.4 | 1.4 | 4.2 | 12.1 | 41.1 | 96.7 |
| 1970 | 4.3 | * | * | * | * | * | 0.4 | 1.6 | 4.5 | 12.9 | 44.0 | 105.4 |
| 1969 | 4.5 | * | * | * | * | * | 0.4 | 1.4 | 4.5 | 13.6 | 49.6 | 122.1 |
| 1968 | 5.1 | * | * | * | * | * | 0.4 | 1.8 | 5.6 | 16.8 | 53.4 | 138.0 |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| HYPERTENSIVE HEART DISEASE (402) | | | | | | | | | | | | |
| ALL OTHER, BOTH SEXES | | | | | | | | | | | | |
| 1978 | 5.2 | * | * | * | * | 0.6 | 2.7 | 7.2 | 16.6 | 31.6 | 50.6 | 78.0 |
| 1977 | 5.4 | * | * | * | * | * | 3.2 | 7.9 | 16.0 | 33.4 | 51.6 | 86.3 |
| 1976 | 5.1 | * | * | * | * | 0.5 | 2.6 | 7.6 | 17.4 | 29.9 | 45.3 | 88.2 |
| 1975 | 5.3 | * | * | * | * | 0.8 | 2.8 | 7.6 | 16.9 | 33.8 | 47.7 | 95.2 |
| 1974 | 5.9 | * | * | * | * | 0.5 | 2.7 | 8.4 | 19.1 | 40.5 | 54.7 | 97.8 |
| 1973 | 5.9 | * | * | * | * | 1.0 | 2.9 | 8.9 | 19.9 | 40.2 | 51.7 | 91.3 |
| 1972 | 7.1 | * | * | * | * | 1.4 | 4.3 | 11.2 | 24.0 | 43.0 | 67.4 | 117.1 |
| 1971 | 6.7 | * | * | * | * | 1.1 | 5.1 | 11.4 | 22.6 | 39.8 | 59.3 | 111.7 |
| 1970 | 7.9 | * | * | * | * | 1.1 | 6.4 | 14.7 | 25.8 | 45.6 | 75.4 | 97.7 |
| 1969 | 8.5 | * | * | * | * | 1.5 | 6.7 | 13.4 | 28.3 | 51.1 | 79.3 | 145.0 |
| 1968 | 9.5 | * | * | * | * | 1.7 | 7.2 | 16.3 | 31.0 | 56.3 | 89.4 | 156.9 |
| ALL OTHER, MALE | | | | | | | | | | | | |
| 1978 | 5.1 | * | * | * | * | * | 3.4 | 8.9 | 18.2 | 33.0 | 51.5 | 56.9 |
| 1977 | 5.1 | * | * | * | * | * | 3.8 | 9.1 | 17.8 | 33.0 | 49.8 | 75.0 |
| 1976 | 4.8 | * | * | * | * | * | 3.1 | 9.4 | 19.0 | 30.6 | 38.1 | 74.5 |
| 1975 | 5.1 | * | * | * | * | 1.1 | 3.7 | 7.3 | 18.9 | 36.0 | 47.1 | 75.9 |
| 1974 | 5.4 | * | * | * | * | * | 3.1 | 8.3 | 19.7 | 41.7 | 49.8 | 76.0 |
| 1973 | 5.7 | * | * | * | * | * | 3.1 | 9.9 | 20.6 | 41.3 | 50.9 | 83.3 |
| 1972 | 6.6 | * | * | * | * | 2.2 | 5.1 | 12.8 | 21.7 | 37.3 | 68.2 | 114.9 |
| 1971 | 6.3 | * | * | * | * | 1.5 | 5.9 | 12.5 | 21.5 | 37.9 | 56.0 | 97.9 |
| 1970 | 7.5 | * | * | * | * | 1.4 | 7.0 | 14.3 | 27.3 | 45.2 | 72.5 | 77.6 |
| 1969 | 8.0 | * | * | * | * | 1.9 | 7.6 | 15.3 | 28.9 | 49.1 | 67.0 | 118.2 |
| 1968 | 8.7 | * | * | * | * | 2.0 | 7.4 | 16.1 | 30.9 | 51.1 | 86.6 | 136.6 |
| ALL OTHER, FEMALE | | | | | | | | | | | | |
| 1978 | 5.4 | * | * | * | * | * | 2.2 | 5.8 | 15.3 | 30.5 | 50.1 | 89.9 |
| 1977 | 5.7 | * | * | * | * | * | 2.7 | 6.9 | 14.5 | 33.8 | 52.7 | 92.3 |
| 1976 | 5.4 | * | * | * | * | * | 2.2 | 6.2 | 16.1 | 29.3 | 50.3 | 95.9 |
| 1975 | 5.5 | * | * | * | * | * | 2.2 | 7.8 | 15.3 | 32.1 | 48.0 | 105.4 |
| 1974 | 6.4 | * | * | * | * | * | 2.3 | 8.5 | 18.6 | 39.5 | 58.1 | 110.6 |
| 1973 | 6.2 | * | * | * | * | * | 2.7 | 8.1 | 19.4 | 39.4 | 52.5 | 96.2 |
| 1972 | 7.6 | * | * | * | * | * | 3.7 | 9.8 | 25.9 | 47.7 | 67.1 | 118.4 |
| 1971 | 7.2 | * | * | * | * | * | 4.4 | 10.4 | 23.5 | 41.3 | 61.8 | 120.5 |
| 1970 | 8.3 | * | * | * | * | * | 6.0 | 15.1 | 24.5 | 45.8 | 77.7 | 111.9 |
| 1969 | 9.0 | * | * | * | * | * | 5.8 | 11.7 | 27.8 | 52.8 | 88.3 | 163.1 |
| 1968 | 10.3 | * | * | * | * | 1.5 | 7.1 | 16.4 | 31.1 | 60.6 | 92.1 | 170.5 |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| HYPERTENSIVE HEART DISEASE (402) | | | | | | | | | | | | |
| BLACK, BOTH SEXES | | | | | | | | | | | | |
| 1978 | 5.9 | * | * | * | * | 0.6 | 3.2 | 8.3 | 18.4 | 34.4 | 52.8 | 82.0 |
| 1977 | 6.1 | * | * | * | * | * | 3.7 | 9.1 | 17.8 | 36.2 | 53.6 | 89.5 |
| 1976 | 5.7 | * | * | * | * | 0.6 | 2.9 | 8.7 | 19.3 | 32.3 | 47.9 | 95.6 |
| 1975 | 5.9 | * | * | * | * | 0.9 | 3.2 | 8.6 | 18.6 | 36.7 | 50.0 | 100.8 |
| 1974 | 6.5 | * | * | * | * | 0.6 | 2.9 | 9.3 | 20.8 | 43.8 | 57.1 | 105.8 |
| 1973 | 6.5 | * | * | * | * | 1.0 | 3.4 | 9.8 | 21.9 | 43.0 | 54.7 | 95.6 |
| 1972 | 7.8 | * | * | * | * | 1.6 | 4.9 | 12.3 | 26.0 | 46.6 | 70.2 | 129.1 |
| 1971 | 7.3 | * | * | * | * | 1.2 | 5.8 | 12.5 | 24.2 | 42.6 | 63.1 | 115.9 |
| 1970 | 8.7 | * | * | * | * | 1.3 | 7.4 | 16.4 | 27.6 | 49.1 | 82.3 | 105.6 |
| 1969 | 9.3 | * | * | * | * | 1.7 | 7.4 | 14.7 | 30.5 | 54.2 | 86.0 | 154.6 |
| 1968 | 10.3 | * | * | * | * | 2.0 | 8.1 | 17.8 | 33.2 | 59.8 | 94.1 | 166.3 |
| BLACK, MALE | | | | | | | | | | | | |
| 1978 | 5.7 | * | * | * | * | * | 4.0 | 10.3 | 20.3 | 36.7 | 52.7 | 56.9 |
| 1977 | 5.7 | * | * | * | * | * | 4.3 | 10.5 | 20.0 | 36.0 | 51.1 | 73.5 |
| 1976 | 5.4 | * | * | * | * | * | 3.5 | 10.8 | 21.1 | 33.6 | 40.2 | 80.9 |
| 1975 | 5.7 | * | * | * | * | 1.3 | 4.0 | 8.3 | 21.2 | 40.2 | 49.1 | 80.4 |
| 1974 | 5.9 | * | * | * | * | * | 3.3 | 9.2 | 21.4 | 45.2 | 53.1 | 83.7 |
| 1973 | 6.3 | * | * | * | * | * | 3.6 | 11.1 | 23.0 | 45.2 | 54.4 | 90.2 |
| 1972 | 7.2 | * | * | * | * | 2.5 | 5.7 | 13.8 | 23.8 | 40.9 | 70.1 | 135.0 |
| 1971 | 6.8 | * | * | * | * | 1.7 | 6.7 | 13.8 | 23.4 | 40.9 | 60.2 | 92.5 |
| 1970 | 8.3 | * | * | * | * | 1.6 | 8.0 | 15.9 | 29.2 | 50.1 | 80.6 | 87.0 |
| 1969 | 8.7 | * | * | * | * | 2.2 | 8.5 | 16.9 | 31.3 | 51.9 | 73.1 | 127.0 |
| 1968 | 9.4 | * | * | * | * | 2.2 | 8.1 | 17.9 | 33.4 | 54.5 | 92.1 | 145.7 |
| BLACK, FEMALE | | | | | | | | | | | | |
| 1978 | 6.1 | * | * | * | * | * | 2.5 | 6.6 | 16.8 | 32.7 | 52.8 | 94.9 |
| 1977 | 6.4 | * | * | * | * | * | 3.2 | 8.0 | 16.1 | 36.4 | 55.4 | 97.9 |
| 1976 | 6.0 | * | * | * | * | * | 2.4 | 7.1 | 17.8 | 31.2 | 53.0 | 103.4 |
| 1975 | 6.2 | * | * | * | * | * | 2.5 | 8.9 | 16.4 | 34.0 | 50.6 | 111.9 |
| 1974 | 7.1 | * | * | * | * | * | 2.5 | 9.4 | 20.3 | 42.8 | 59.9 | 117.9 |
| 1973 | 6.8 | * | * | * | * | * | 3.2 | 8.8 | 21.1 | 41.2 | 55.1 | 98.6 |
| 1972 | 8.4 | * | * | * | * | * | 4.3 | 11.0 | 27.8 | 51.2 | 70.0 | 125.7 |
| 1971 | 7.8 | * | * | * | * | * | 5.0 | 11.4 | 24.9 | 43.8 | 65.3 | 129.9 |
| 1970 | 9.1 | * | * | * | * | * | 6.9 | 16.8 | 26.3 | 48.3 | 83.5 | 117.7 |
| 1969 | 9.8 | * | * | * | * | * | 6.5 | 12.8 | 29.8 | 56.1 | 95.2 | 171.7 |
| 1968 | 11.1 | * | * | * | * | 1.7 | 8.0 | 17.8 | 33.1 | 63.9 | 95.5 | 182.1 |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| HYPERTENSIVE HEART AND RENAL DISEASE (404) | | | | | | | | | | | | |
| BOTH SEXES | | | | | | | | | | | | |
| 1978 | 1.5 | * | * | * | * | 0.1 | 0.1 | 0.3 | 1.1 | 4.3 | 15.4 | 52.4 |
| 1977 | 1.6 | * | * | * | * | * | 0.1 | 0.4 | 1.3 | 4.5 | 17.3 | 61.3 |
| 1976 | 1.8 | * | * | * | * | * | 0.1 | 0.5 | 1.7 | 5.5 | 20.0 | 68.4 |
| 1975 | 2.0 | * | * | * | * | * | 0.2 | 0.5 | 1.8 | 5.9 | 22.7 | 76.3 |
| 1974 | 2.3 | * | * | * | * | * | 0.2 | 0.7 | 1.9 | 7.0 | 26.2 | 93.7 |
| 1973 | 2.6 | * | * | * | * | * | 0.2 | 0.8 | 2.5 | 8.2 | 30.1 | 104.5 |
| 1972 | 2.8 | * | * | * | * | * | 0.2 | 0.8 | 2.6 | 8.6 | 32.0 | 115.6 |
| 1971 | 3.0 | * | * | * | * | * | 0.3 | 1.0 | 3.1 | 9.7 | 33.8 | 125.8 |
| 1970 | 3.2 | * | * | * | * | 0.1 | 0.4 | 1.2 | 3.2 | 10.7 | 40.4 | 118.6 |
| 1969 | 3.6 | * | * | * | * | 0.1 | 0.5 | 1.4 | 3.8 | 12.1 | 44.8 | 150.3 |
| 1968 | 4.0 | * | * | * | * | 0.1 | 0.4 | 1.3 | 3.9 | 13.4 | 49.2 | 175.6 |
| MALE | | | | | | | | | | | | |
| 1978 | 1.4 | * | * | * | * | * | * | 0.3 | 1.2 | 5.4 | 18.2 | 64.1 |
| 1977 | 1.5 | * | * | * | * | * | * | 0.5 | 1.6 | 5.5 | 22.1 | 68.4 |
| 1976 | 1.7 | * | * | * | * | * | * | 0.6 | 2.0 | 7.0 | 22.2 | 72.6 |
| 1975 | 1.8 | * | * | * | * | * | 0.2 | 0.6 | 2.1 | 7.1 | 25.9 | 80.5 |
| 1974 | 2.1 | * | * | * | * | * | 0.2 | 0.7 | 2.2 | 8.6 | 29.2 | 106.4 |
| 1973 | 2.5 | * | * | * | * | * | * | 1.0 | 3.1 | 9.9 | 36.4 | 115.8 |
| 1972 | 2.6 | * | * | * | * | * | * | 0.9 | 3.7 | 10.4 | 38.4 | 126.8 |
| 1971 | 2.8 | * | * | * | * | * | 0.4 | 1.2 | 3.6 | 11.9 | 38.9 | 136.9 |
| 1970 | 3.0 | * | * | * | * | * | 0.4 | 1.4 | 3.7 | 13.0 | 44.6 | 120.4 |
| 1969 | 3.5 | * | * | * | * | * | 0.5 | 1.6 | 4.5 | 14.4 | 51.6 | 163.0 |
| 1968 | 3.8 | * | * | * | * | * | 0.5 | 1.5 | 4.8 | 16.1 | 55.9 | 188.4 |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| HYPERTENSIVE HEART AND RENAL DISEASE (404) | | | | | | | | | | | | |
| FEMALE | | | | | | | | | | | | |
| 1978 | 1.6 | * | * | * | * | * | * | 0.3 | 1.1 | 3.4 | 13.8 | 47.1 |
| 1977 | 1.7 | * | * | * | * | * | * | 0.4 | 1.0 | 3.7 | 14.5 | 58.1 |
| 1976 | 2.0 | * | * | * | * | * | * | 0.4 | 1.3 | 4.4 | 18.7 | 66.4 |
| 1975 | 2.2 | * | * | * | * | * | * | 0.4 | 1.6 | 5.1 | 20.8 | 74.3 |
| 1974 | 2.5 | * | * | * | * | * | * | 0.6 | 1.7 | 5.8 | 24.4 | 87.5 |
| 1973 | 2.7 | * | * | * | * | * | 0.2 | 0.7 | 1.9 | 6.8 | 26.2 | 98.9 |
| 1972 | 2.9 | * | * | * | * | * | 0.3 | 0.7 | 1.7 | 7.3 | 28.0 | 109.8 |
| 1971 | 3.2 | * | * | * | * | * | 0.3 | 0.9 | 2.6 | 8.0 | 30.6 | 120.0 |
| 1970 | 3.4 | * | * | * | * | * | 0.3 | 0.9 | 2.7 | 8.9 | 37.6 | 117.6 |
| 1969 | 3.8 | * | * | * | * | * | 0.4 | 1.1 | 3.1 | 10.3 | 40.2 | 143.5 |
| 1968 | 4.1 | * | * | * | * | * | 0.3 | 1.2 | 3.2 | 11.2 | 44.6 | 168.8 |
| WHITE, BOTH SEXES | | | | | | | | | | | | |
| 1978 | 1.4 | * | * | * | * | * | * | 0.1 | 0.8 | 3.7 | 14.8 | 53.1 |
| 1977 | 1.6 | * | * | * | * | * | * | 0.3 | 0.9 | 3.6 | 16.9 | 62.9 |
| 1976 | 1.8 | * | * | * | * | * | * | 0.3 | 1.1 | 4.7 | 19.8 | 70.1 |
| 1975 | 2.0 | * | * | * | * | * | * | 0.3 | 1.2 | 5.1 | 22.1 | 78.1 |
| 1974 | 2.3 | * | * | * | * | * | * | 0.4 | 1.4 | 6.0 | 25.8 | 94.8 |
| 1973 | 2.5 | * | * | * | * | * | * | 0.4 | 1.8 | 6.9 | 29.5 | 107.4 |
| 1972 | 2.7 | * | * | * | * | * | * | 0.5 | 2.0 | 7.4 | 31.7 | 118.8 |
| 1971 | 2.9 | * | * | * | * | * | 0.2 | 0.6 | 2.2 | 8.1 | 33.5 | 129.3 |
| 1970 | 3.1 | * | * | * | * | * | 0.2 | 0.7 | 2.4 | 9.2 | 39.4 | 121.9 |
| 1969 | 3.5 | * | * | * | * | * | 0.2 | 0.8 | 2.7 | 10.2 | 44.0 | 154.0 |
| 1968 | 3.8 | * | * | * | * | * | 0.1 | 0.8 | 2.8 | 11.3 | 47.4 | 180.2 |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| HYPERTENSIVE HEART AND RENAL DISEASE (404) | | | | | | | | | | | | |
| WHITE, MALE | | | | | | | | | | | | |
| 1978 | 1.3 | * | * | * | * | * | * | * | 0.9 | 4.8 | 17.4 | 64.8 |
| 1977 | 1.4 | * | * | * | * | * | * | 0.3 | 1.2 | 4.5 | 21.4 | 70.9 |
| 1976 | 1.6 | * | * | * | * | * | * | 0.4 | 1.4 | 6.2 | 22.2 | 74.6 |
| 1975 | 1.7 | * | * | * | * | * | * | 0.4 | 1.6 | 6.2 | 25.3 | 82.6 |
| 1974 | 2.0 | * | * | * | * | * | * | 0.5 | 1.6 | 7.5 | 29.0 | 106.2 |
| 1973 | 2.4 | * | * | * | * | * | * | 0.5 | 2.2 | 8.5 | 36.1 | 119.2 |
| 1972 | 2.5 | * | * | * | * | * | * | 0.6 | 2.9 | 8.9 | 37.3 | 130.9 |
| 1971 | 2.7 | * | * | * | * | * | 0.2 | 0.7 | 2.7 | 9.9 | 38.2 | 140.8 |
| 1970 | 2.9 | * | * | * | * | * | * | 0.9 | 3.0 | 11.3 | 43.2 | 124.0 |
| 1969 | 3.3 | * | * | * | * | * | * | 1.0 | 3.3 | 12.5 | 50.2 | 169.4 |
| 1968 | 3.6 | * | * | * | * | * | * | 1.0 | 3.6 | 14.0 | 53.6 | 195.7 |
| WHITE, FEMALE | | | | | | | | | | | | |
| 1978 | 1.6 | * | * | * | * | * | * | * | 0.8 | 2.9 | 13.2 | 47.9 |
| 1977 | 1.7 | * | * | * | * | * | * | 0.2 | 0.6 | 3.0 | 14.2 | 59.3 |
| 1976 | 2.0 | * | * | * | * | * | * | 0.2 | 0.8 | 3.6 | 18.3 | 68.0 |
| 1975 | 2.2 | * | * | * | * | * | * | 0.2 | 0.9 | 4.2 | 20.3 | 76.0 |
| 1974 | 2.5 | * | * | * | * | * | * | 0.4 | 1.1 | 4.9 | 23.8 | 89.2 |
| 1973 | 2.7 | * | * | * | * | * | * | 0.3 | 1.4 | 5.7 | 25.4 | 101.7 |
| 1972 | 2.9 | * | * | * | * | * | * | 0.4 | 1.2 | 6.2 | 28.1 | 112.7 |
| 1971 | 3.1 | * | * | * | * | * | * | 0.5 | 1.7 | 6.7 | 30.5 | 123.3 |
| 1970 | 3.4 | * | * | * | * | * | * | 0.5 | 1.9 | 7.6 | 36.9 | 120.7 |
| 1969 | 3.6 | * | * | * | * | * | 0.2 | 0.6 | 2.1 | 8.4 | 39.8 | 146.0 |
| 1968 | 3.9 | * | * | * | * | * | * | 0.6 | 2.1 | 9.1 | 43.2 | 172.1 |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| HYPERTENSIVE HEART AND RENAL DISEASE (404) | | | | | | | | | | | | |
| ALL OTHER, BOTH SEXES | | | | | | | | | | | | |
| 1978 | 1.7 | * | * | * | * | * | * | 1.7 | 3.8 | 9.5 | 22.3 | 44.6 |
| 1977 | 1.8 | * | * | * | * | * | * | 1.7 | 4.4 | 12.4 | 22.2 | 43.1 |
| 1976 | 2.1 | * | * | * | * | * | * | 2.4 | 6.7 | 13.2 | 22.1 | 49.0 |
| 1975 | 2.4 | * | * | * | * | * | 0.9 | 2.3 | 7.3 | 14.3 | 29.1 | 56.2 |
| 1974 | 2.6 | * | * | * | * | * | 0.8 | 2.5 | 6.8 | 16.9 | 31.4 | 81.5 |
| 1973 | 3.2 | * | * | * | * | * | 0.8 | 4.1 | 9.6 | 20.3 | 36.8 | 70.9 |
| 1972 | 3.1 | * | * | * | * | * | 1.2 | 3.7 | 8.5 | 21.2 | 36.2 | 78.0 |
| 1971 | 3.8 | * | * | * | * | * | 1.6 | 5.0 | 12.0 | 25.4 | 38.2 | 85.8 |
| 1970 | 4.1 | * | * | * | * | * | 2.0 | 5.5 | 10.6 | 25.3 | 52.8 | 85.0 |
| 1969 | 4.8 | * | * | * | * | * | 2.5 | 6.7 | 14.0 | 31.1 | 56.0 | 109.2 |
| 1968 | 5.3 | * | * | * | * | 0.7 | 2.4 | 6.2 | 14.9 | 34.4 | 72.5 | 123.5 |
| ALL OTHER, MALE | | | | | | | | | | | | |
| 1978 | 1.7 | * | * | * | * | * | * | 1.7 | 4.5 | 10.6 | 25.9 | 56.9 |
| 1977 | 1.9 | * | * | * | * | * | * | 1.8 | 4.8 | 14.1 | 29.1 | 42.9 |
| 1976 | 2.1 | * | * | * | * | * | * | 2.6 | 7.7 | 15.0 | 21.4 | 52.7 |
| 1975 | 2.3 | * | * | * | * | * | * | 2.6 | 6.8 | 15.5 | 32.2 | 59.3 |
| 1974 | 2.6 | * | * | * | * | * | * | 2.7 | 7.3 | 19.5 | 31.0 | 108.0 |
| 1973 | 3.3 | * | * | * | * | * | * | 5.1 | 12.4 | 23.1 | 39.3 | 81.3 |
| 1972 | 3.5 | * | * | * | * | * | * | 4.0 | 11.3 | 24.5 | 49.8 | 85.1 |
| 1971 | 4.0 | * | * | * | * | * | 1.8 | 5.7 | 12.7 | 30.7 | 47.3 | 97.9 |
| 1970 | 4.3 | * | * | * | * | * | 2.5 | 6.2 | 11.1 | 29.1 | 60.6 | 88.4 |
| 1969 | 5.1 | * | * | * | * | * | 3.1 | 8.0 | 16.2 | 32.7 | 68.6 | 102.3 |
| 1968 | 5.4 | * | * | * | * | * | 3.2 | 6.0 | 16.3 | 36.9 | 83.3 | 117.1 |
| ALL OTHER, FEMALE | | | | | | | | | | | | |
| 1978 | 1.6 | * | * | * | * | * | * | 1.8 | 3.2 | 8.7 | 19.8 | 38.5 |
| 1977 | 1.8 | * | * | * | * | * | * | 1.7 | 4.1 | 11.0 | 17.5 | 43.3 |
| 1976 | 2.1 | * | * | * | * | * | * | 2.2 | 5.8 | 11.7 | 22.6 | 46.9 |
| 1975 | 2.4 | * | * | * | * | * | * | 1.9 | 7.7 | 13.4 | 26.8 | 53.8 |
| 1974 | 2.6 | * | * | * | * | * | * | 2.4 | 6.4 | 14.8 | 31.6 | 65.9 |
| 1973 | 3.0 | * | * | * | * | * | * | 3.3 | 7.3 | 18.1 | 35.2 | 64.6 |
| 1972 | 2.8 | * | * | * | * | * | 1.6 | 3.5 | 6.1 | 18.7 | 26.6 | 73.7 |
| 1971 | 3.5 | * | * | * | * | * | 1.4 | 4.5 | 11.4 | 21.2 | 31.4 | 78.1 |
| 1970 | 3.9 | * | * | * | * | * | 1.5 | 4.9 | 10.1 | 22.1 | 46.9 | 82.6 |
| 1969 | 4.6 | * | * | * | * | * | 2.0 | 5.6 | 12.1 | 29.7 | 46.2 | 113.8 |
| 1968 | 5.2 | * | * | * | * | * | 1.8 | 6.3 | 13.7 | 32.3 | 64.4 | 127.9 |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| HYPERTENSIVE HEART AND RENAL DISEASE (404) | | | | | | | | | | | | |
| BLACK, BOTH SEXES | | | | | | | | | | | | |
| 1978 | 1.9 | * | * | * | * | * | * | 2.0 | 4.3 | 10.7 | 24.0 | 46.7 |
| 1977 | 2.0 | * | * | * | * | * | * | 1.9 | 4.9 | 13.5 | 23.2 | 44.8 |
| 1976 | 2.4 | * | * | * | * | * | * | 2.7 | 7.3 | 14.3 | 23.0 | 51.5 |
| 1975 | 2.6 | * | * | * | * | * | 1.0 | 2.5 | 8.0 | 15.7 | 30.4 | 60.0 |
| 1974 | 2.9 | * | * | * | * | * | 0.9 | 2.8 | 7.4 | 18.2 | 33.0 | 87.6 |
| 1973 | 3.5 | * | * | * | * | * | 0.9 | 4.6 | 10.5 | 22.2 | 37.9 | 75.4 |
| 1972 | 3.4 | * | * | * | * | * | 1.2 | 4.2 | 9.2 | 22.6 | 39.1 | 78.2 |
| 1971 | 4.1 | * | * | * | * | * | 1.8 | 5.5 | 12.8 | 27.2 | 39.8 | 88.8 |
| 1970 | 4.4 | * | * | * | * | * | 2.2 | 6.1 | 11.6 | 26.9 | 56.5 | 93.7 |
| 1969 | 5.2 | * | * | * | * | * | 2.8 | 7.4 | 15.2 | 33.2 | 59.0 | 117.5 |
| 1968 | 5.8 | * | * | * | * | 0.8 | 2.7 | 6.8 | 16.1 | 36.6 | 78.0 | 132.6 |
| BLACK, MALE | | | | | | | | | | | | |
| 1978 | 2.0 | * | * | * | * | * | * | 2.0 | 5.2 | 12.2 | 28.5 | 58.8 |
| 1977 | 2.1 | * | * | * | * | * | * | 2.0 | 5.5 | 15.9 | 30.6 | 46.9 |
| 1976 | 2.4 | * | * | * | * | * | * | 2.8 | 8.4 | 16.8 | 22.2 | 55.3 |
| 1975 | 2.6 | * | * | * | * | * | * | 2.9 | 7.4 | 17.3 | 33.8 | 65.2 |
| 1974 | 2.9 | * | * | * | * | * | * | 2.9 | 8.1 | 21.8 | 33.5 | 118.6 |
| 1973 | 3.7 | * | * | * | * | * | * | 5.7 | 13.8 | 25.4 | 40.2 | 90.2 |
| 1972 | 3.7 | * | * | * | * | * | * | 4.5 | 12.4 | 26.4 | 53.8 | 80.0 |
| 1971 | 4.4 | * | * | * | * | * | 2.1 | 6.4 | 13.6 | 33.2 | 49.5 | 110.0 |
| 1970 | 4.6 | * | * | * | * | * | 2.9 | 6.8 | 12.4 | 31.5 | 64.6 | 100.0 |
| 1969 | 5.5 | * | * | * | * | * | 3.4 | 8.9 | 18.1 | 35.4 | 73.1 | 118.9 |
| 1968 | 5.9 | * | * | * | * | * | 3.6 | 6.6 | 17.9 | 39.5 | 91.5 | 134.3 |
| BLACK, FEMALE | | | | | | | | | | | | |
| 1978 | 1.9 | * | * | * | * | * | * | 2.1 | 3.5 | 9.6 | 20.9 | 40.4 |
| 1977 | 2.0 | * | * | * | * | * | * | 1.9 | 4.4 | 11.8 | 18.4 | 43.6 |
| 1976 | 2.3 | * | * | * | * | * | * | 2.5 | 6.4 | 12.4 | 23.5 | 49.4 |
| 1975 | 2.7 | * | * | * | * | * | * | 2.2 | 8.4 | 14.5 | 28.0 | 57.1 |
| 1974 | 2.8 | * | * | * | * | * | * | 2.6 | 6.8 | 15.5 | 32.7 | 70.5 |
| 1973 | 3.3 | * | * | * | * | * | * | 3.7 | 7.9 | 19.6 | 36.5 | 67.1 |
| 1972 | 3.1 | * | * | * | * | * | 1.7 | 3.9 | 6.4 | 19.6 | 28.6 | 77.1 |
| 1971 | 3.8 | * | * | * | * | * | 1.6 | 4.8 | 12.2 | 22.4 | 32.8 | 76.1 |
| 1970 | 4.2 | * | * | * | * | * | 1.7 | 5.5 | 10.8 | 23.2 | 50.5 | 89.7 |
| 1969 | 4.9 | * | * | * | * | * | 2.2 | 6.2 | 12.8 | 31.5 | 48.5 | 116.7 |
| 1968 | 5.7 | * | * | * | * | * | 2.1 | 6.9 | 14.5 | 34.1 | 68.0 | 133.9 |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| ISCHEMIC HEART DISEASE (410-413) | | | | | | | | | | | | |
| BOTH SEXES | | | | | | | | | | | | |
| 1978 | 289.2 | * | * | 0.1 | 0.3 | 4.0 | 35.8 | 159.8 | 445.3 | 1,083.9 | 2,670.5 | 6,694.6 |
| 1977 | 290.5 | * | * | * | 0.4 | 4.1 | 38.4 | 166.4 | 457.9 | 1,112.2 | 2,718.1 | 6,700.5 |
| 1976 | 297.0 | * | * | * | 0.3 | 4.3 | 40.3 | 171.7 | 479.5 | 1,151.9 | 2,810.4 | 6,962.9 |
| 1975 | 298.3 | * | * | * | 0.4 | 4.4 | 42.7 | 177.9 | 494.7 | 1,191.9 | 2,865.4 | 6,844.2 |
| 1974 | 311.6 | * | * | 0.1 | 0.4 | 4.8 | 44.2 | 186.7 | 520.8 | 1,267.3 | 3,042.8 | 7,478.5 |
| 1973 | 323.7 | 0.9 | * | 0.1 | 0.5 | 5.1 | 48.3 | 195.0 | 554.9 | 1,330.4 | 3,229.3 | 7,802.7 |
| 1972 | 327.0 | * | * | 0.1 | 0.6 | 5.1 | 50.1 | 199.6 | 566.3 | 1,380.9 | 3,303.3 | 7,797.0 |
| 1971 | 326.0 | 0.8 | * | 0.1 | 0.7 | 6.1 | 52.1 | 202.3 | 572.4 | 1,384.5 | 3,317.4 | 7,843.2 |
| 1970 | 328.1 | * | * | 0.1 | 0.6 | 6.3 | 52.9 | 206.5 | 585.5 | 1,429.2 | 3,406.3 | 7,249.4 |
| 1969 | 332.6 | 0.7 | * | 0.1 | 0.6 | 6.2 | 55.3 | 211.5 | 592.7 | 1,458.0 | 3,453.6 | 8,284.5 |
| 1968 | 338.4 | 0.9 | * | 0.1 | 0.5 | 6.4 | 56.7 | 218.0 | 616.9 | 1,502.0 | 3,533.2 | 8,483.0 |
| MALE | | | | | | | | | | | | |
| 1978 | 327.6 | * | * | * | 0.5 | 6.4 | 59.2 | 262.0 | 696.4 | 1,575.8 | 3,449.0 | 7,561.5 |
| 1977 | 332.2 | * | * | * | 0.5 | 6.6 | 63.2 | 273.6 | 714.2 | 1,619.7 | 3,512.9 | 7,607.0 |
| 1976 | 340.6 | * | * | * | 0.4 | 6.9 | 67.0 | 282.6 | 749.3 | 1,675.1 | 3,602.9 | 7,832.3 |
| 1975 | 344.9 | * | * | * | 0.6 | 7.0 | 70.2 | 293.3 | 773.0 | 1,722.1 | 3,666.7 | 7,705.2 |
| 1974 | 359.8 | * | * | * | 0.6 | 7.3 | 72.3 | 308.0 | 812.4 | 1,821.9 | 3,848.0 | 8,335.6 |
| 1973 | 375.8 | * | * | 0.1 | 0.7 | 7.7 | 78.4 | 320.0 | 866.2 | 1,904.7 | 4,081.6 | 8,676.0 |
| 1972 | 380.5 | * | * | * | 0.8 | 7.9 | 81.2 | 329.9 | 881.6 | 1,955.6 | 4,135.6 | 8,579.3 |
| 1971 | 381.7 | 1.1 | * | 0.1 | 0.9 | 9.4 | 85.2 | 330.9 | 887.0 | 1,956.9 | 4,134.2 | 8,722.7 |
| 1970 | 387.2 | * | * | * | 0.8 | 9.4 | 87.0 | 338.0 | 904.6 | 2,010.0 | 4,222.7 | 7,781.5 |
| 1969 | 394.7 | * | * | * | 0.8 | 9.7 | 90.7 | 348.1 | 920.6 | 2,050.9 | 4,263.8 | 9,042.0 |
| 1968 | 402.5 | 1.2 | * | * | 0.7 | 9.6 | 92.7 | 356.6 | 950.1 | 2,100.7 | 4,330.3 | 9,259.5 |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| ISCHEMIC HEART DISEASE (410-413) | | | | | | | | | | | | |
| FEMALE | | | | | | | | | | | | |
| 1978 | 252.9 | * | * | * | 0.2 | 1.6 | 13.4 | 64.7 | 223.7 | 708.0 | 2,207.0 | 6,302.9 |
| 1977 | 251.1 | * | * | * | 0.2 | 1.7 | 14.7 | 66.7 | 231.2 | 723.9 | 2,242.3 | 6,284.0 |
| 1976 | 255.6 | * | * | * | 0.2 | 1.8 | 14.7 | 68.7 | 240.6 | 751.2 | 2,331.9 | 6,554.5 |
| 1975 | 254.1 | * | * | * | 0.2 | 1.8 | 16.4 | 70.7 | 248.2 | 785.7 | 2,377.3 | 6,427.3 |
| 1974 | 266.0 | * | * | * | 0.3 | 2.3 | 17.3 | 74.0 | 262.3 | 842.0 | 2,546.8 | 7,054.1 |
| 1973 | 274.1 | * | * | * | 0.3 | 2.5 | 19.5 | 78.8 | 278.2 | 889.3 | 2,696.8 | 7,363.1 |
| 1972 | 276.2 | * | * | * | 0.5 | 2.3 | 20.5 | 78.5 | 285.4 | 937.5 | 2,772.7 | 7,396.7 |
| 1971 | 273.1 | * | * | 0.1 | 0.4 | 2.9 | 20.7 | 82.6 | 291.2 | 941.2 | 2,787.6 | 7,384.0 |
| 1970 | 272.0 | * | * | * | 0.4 | 3.2 | 20.6 | 84.0 | 299.1 | 978.0 | 2,866.3 | 6,951.5 |
| 1969 | 273.8 | * | * | * | 0.4 | 2.9 | 21.8 | 84.1 | 297.5 | 994.6 | 2,910.1 | 7,874.4 |
| 1968 | 277.5 | * | * | * | 0.3 | 3.3 | 22.7 | 88.2 | 315.6 | 1,030.6 | 2,987.6 | 8,066.5 |
| WHITE, BOTH SEXES | | | | | | | | | | | | |
| 1978 | 304.2 | * | * | * | 0.3 | 3.5 | 32.9 | 152.2 | 433.7 | 1,078.6 | 2,715.9 | 6,876.8 |
| 1977 | 305.1 | * | * | * | 0.3 | 3.6 | 35.2 | 158.8 | 445.7 | 1,105.5 | 2,768.7 | 6,875.3 |
| 1976 | 312.1 | * | * | * | 0.3 | 3.8 | 37.7 | 164.1 | 469.1 | 1,144.8 | 2,868.8 | 7,157.8 |
| 1975 | 313.1 | * | * | * | 0.4 | 3.8 | 39.5 | 170.9 | 484.5 | 1,182.4 | 2,926.1 | 7,038.0 |
| 1974 | 326.3 | * | * | * | 0.4 | 4.0 | 40.9 | 178.2 | 508.5 | 1,252.2 | 3,108.2 | 7,683.3 |
| 1973 | 338.1 | 0.8 | * | 0.1 | 0.4 | 4.4 | 44.6 | 185.5 | 541.3 | 1,312.6 | 3,293.9 | 8,014.5 |
| 1972 | 341.8 | * | * | 0.1 | 0.5 | 4.3 | 45.9 | 190.0 | 552.0 | 1,370.1 | 3,367.1 | 8,034.5 |
| 1971 | 340.5 | 0.7 | * | 0.1 | 0.5 | 5.1 | 47.8 | 193.6 | 560.5 | 1,372.1 | 3,374.6 | 8,093.7 |
| 1970 | 342.2 | * | * | 0.1 | 0.5 | 5.2 | 48.0 | 197.1 | 571.3 | 1,419.4 | 3,456.4 | 7,536.0 |
| 1969 | 346.2 | 0.7 | * | 0.1 | 0.5 | 4.9 | 50.2 | 201.2 | 576.9 | 1,447.7 | 3,504.1 | 8,570.4 |
| 1968 | 351.4 | 0.8 | * | 0.1 | 0.4 | 5.2 | 51.2 | 207.3 | 597.9 | 1,490.7 | 3,585.8 | 8,755.2 |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| ISCHEMIC HEART DISEASE (410-413) | | | | | | | | | | | | |
| WHITE, MALE | | | | | | | | | | | | |
| 1978 | 344.1 | * | * | * | 0.4 | 5.6 | 55.7 | 255.4 | 690.3 | 1,591.2 | 3,539.1 | 7,802.4 |
| 1977 | 348.8 | * | * | * | 0.4 | 5.9 | 59.6 | 267.3 | 708.4 | 1,636.5 | 3,614.4 | 7,844.8 |
| 1976 | 357.8 | * | * | * | 0.4 | 6.2 | 64.2 | 276.4 | 746.2 | 1,692.0 | 3,713.4 | 8,098.7 |
| 1975 | 362.3 | * | * | * | 0.5 | 6.2 | 66.8 | 288.1 | 773.5 | 1,738.0 | 3,769.1 | 7,972.6 |
| 1974 | 376.9 | * | * | * | 0.5 | 6.4 | 69.4 | 302.1 | 809.9 | 1,831.7 | 3,953.6 | 8,610.1 |
| 1973 | 393.4 | * | * | * | 0.6 | 6.9 | 75.3 | 313.2 | 863.7 | 1,915.6 | 4,188.7 | 8,975.3 |
| 1972 | 398.1 | * | * | * | 0.6 | 7.1 | 77.4 | 322.9 | 880.1 | 1,972.1 | 4,237.8 | 8,888.8 |
| 1971 | 399.4 | * | * | 0.1 | 0.7 | 8.2 | 81.3 | 326.1 | 886.8 | 1,974.4 | 4,233.9 | 9,067.4 |
| 1970 | 404.9 | * | * | * | 0.7 | 8.2 | 82.3 | 332.5 | 904.8 | 2,028.7 | 4,315.0 | 8,164.2 |
| 1969 | 411.9 | * | * | * | 0.7 | 7.9 | 85.8 | 341.2 | 919.7 | 2,069.9 | 4,348.8 | 9,443.1 |
| 1968 | 419.3 | * | * | * | 0.5 | 8.2 | 87.6 | 349.9 | 945.2 | 2,119.3 | 4,426.3 | 9,598.7 |
| WHITE, FEMALE | | | | | | | | | | | | |
| 1978 | 266.3 | * | * | * | 0.1 | 1.3 | 10.5 | 54.5 | 205.9 | 687.5 | 2,231.5 | 6,464.3 |
| 1977 | 263.4 | * | * | * | 0.2 | 1.3 | 11.2 | 56.3 | 211.8 | 700.3 | 2,269.7 | 6,436.3 |
| 1976 | 268.4 | * | * | * | 0.1 | 1.4 | 11.8 | 58.3 | 222.3 | 726.8 | 2,365.9 | 6,728.9 |
| 1975 | 266.2 | * | * | * | 0.2 | 1.3 | 12.9 | 60.6 | 227.2 | 758.4 | 2,418.9 | 6,599.1 |
| 1974 | 278.0 | * | * | * | 0.2 | 1.7 | 13.2 | 61.9 | 240.1 | 810.2 | 2,591.9 | 7,231.3 |
| 1973 | 285.3 | * | * | * | 0.2 | 1.9 | 14.6 | 65.7 | 253.7 | 851.7 | 2,739.8 | 7,546.5 |
| 1972 | 288.0 | * | * | * | 0.5 | 1.7 | 15.3 | 65.4 | 258.6 | 908.0 | 2,817.8 | 7,604.7 |
| 1971 | 284.3 | * | * | 0.1 | 0.3 | 2.1 | 15.5 | 69.5 | 268.1 | 908.1 | 2,824.6 | 7,595.0 |
| 1970 | 282.5 | * | * | * | 0.3 | 2.2 | 15.1 | 70.0 | 271.6 | 948.9 | 2,894.7 | 7,192.3 |
| 1969 | 283.7 | * | * | * | 0.3 | 1.9 | 16.1 | 69.8 | 267.7 | 963.7 | 2,942.2 | 8,118.8 |
| 1968 | 286.6 | * | * | * | 0.3 | 2.2 | 16.3 | 72.8 | 283.7 | 998.3 | 3,016.1 | 8,311.6 |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| ISCHEMIC HEART DISEASE (410-413) | | | | | | | | | | | | |
| ALL OTHER, BOTH SEXES | | | | | | | | | | | | |
| 1978 | 193.4 | * | * | * | 0.6 | 7.6 | 56.0 | 218.9 | 549.7 | 1,133.6 | 2,203.2 | 4,605.4 |
| 1977 | 196.7 | * | * | * | 0.8 | 7.7 | 60.3 | 226.8 | 569.4 | 1,176.3 | 2,196.9 | 4,698.1 |
| 1976 | 198.3 | * | * | * | 0.6 | 8.2 | 58.1 | 233.3 | 575.4 | 1,220.0 | 2,209.8 | 4,696.1 |
| 1975 | 200.0 | * | * | * | 0.8 | 8.5 | 64.9 | 236.2 | 589.4 | 1,284.4 | 2,216.3 | 4,620.5 |
| 1974 | 212.8 | * | * | * | 0.8 | 10.0 | 66.9 | 257.7 | 636.3 | 1,414.6 | 2,323.9 | 5,094.8 |
| 1973 | 225.0 | * | * | * | 1.1 | 9.9 | 74.5 | 275.3 | 683.3 | 1,506.4 | 2,493.6 | 5,334.6 |
| 1972 | 225.1 | * | * | * | 1.3 | 10.4 | 80.7 | 281.6 | 702.7 | 1,486.8 | 2,561.1 | 5,056.9 |
| 1971 | 225.0 | * | * | * | 1.5 | 13.2 | 83.5 | 277.6 | 686.4 | 1,506.9 | 2,624.0 | 4,989.2 |
| 1970 | 229.4 | * | * | * | 1.4 | 14.0 | 88.6 | 289.3 | 719.1 | 1,524.6 | 2,770.9 | 4,306.8 |
| 1969 | 235.4 | * | * | * | 1.5 | 16.0 | 93.6 | 303.9 | 745.1 | 1,560.4 | 2,805.5 | 5,063.3 |
| 1968 | 244.7 | * | * | * | 1.4 | 15.2 | 98.9 | 315.0 | 800.8 | 1,615.1 | 2,851.4 | 5,443.1 |
| ALL OTHER, MALE | | | | | | | | | | | | |
| 1978 | 219.9 | * | * | * | 0.8 | 12.3 | 84.6 | 316.9 | 754.3 | 1,435.2 | 2,589.5 | 5,094.8 |
| 1977 | 222.1 | * | * | * | 0.9 | 12.0 | 88.9 | 327.0 | 768.6 | 1,463.5 | 2,557.9 | 5,182.1 |
| 1976 | 225.1 | * | * | * | 0.7 | 12.1 | 88.2 | 335.9 | 778.8 | 1,513.0 | 2,571.2 | 5,163.6 |
| 1975 | 227.2 | * | * | * | 1.1 | 12.6 | 95.6 | 338.5 | 769.1 | 1,571.3 | 2,654.5 | 5,031.5 |
| 1974 | 241.4 | * | * | * | 0.8 | 14.2 | 94.1 | 360.1 | 836.5 | 1,732.0 | 2,762.0 | 5,508.0 |
| 1973 | 253.6 | * | * | * | 1.3 | 14.3 | 102.3 | 380.8 | 891.5 | 1,800.6 | 2,943.3 | 5,620.8 |
| 1972 | 256.3 | * | * | * | 2.2 | 14.6 | 110.8 | 392.1 | 895.5 | 1,797.4 | 3,036.9 | 5,451.1 |
| 1971 | 255.8 | * | * | * | 1.9 | 18.7 | 116.3 | 375.0 | 889.6 | 1,789.7 | 3,045.4 | 5,327.7 |
| 1970 | 261.1 | * | * | * | 1.8 | 19.1 | 124.3 | 387.7 | 902.3 | 1,833.2 | 3,149.9 | 4,418.8 |
| 1969 | 269.5 | * | * | * | 1.9 | 23.3 | 131.1 | 413.3 | 929.0 | 1,864.1 | 3,245.7 | 5,259.1 |
| 1968 | 278.8 | * | * | * | 2.0 | 20.7 | 134.9 | 419.4 | 998.1 | 1,920.2 | 3,179.6 | 5,958.5 |
| ALL OTHER, FEMALE | | | | | | | | | | | | |
| 1978 | 169.5 | * | * | * | * | 3.6 | 31.9 | 138.0 | 379.9 | 900.1 | 1,944.7 | 4,387.2 |
| 1977 | 173.6 | * | * | * | 0.7 | 3.9 | 36.4 | 143.7 | 403.3 | 950.6 | 1,945.2 | 4,437.5 |
| 1976 | 174.0 | * | * | * | * | 4.8 | 32.9 | 147.4 | 405.1 | 988.3 | 1,962.8 | 4,433.7 |
| 1975 | 175.3 | * | * | * | * | 5.0 | 39.2 | 149.7 | 437.7 | 1,056.2 | 1,910.5 | 4,332.3 |
| 1974 | 186.7 | * | * | * | 0.8 | 6.4 | 44.2 | 170.9 | 466.0 | 1,160.5 | 2,021.7 | 4,851.8 |
| 1973 | 198.9 | * | * | * | 0.9 | 6.0 | 51.2 | 185.0 | 505.8 | 1,268.9 | 2,184.6 | 5,160.8 |
| 1972 | 196.5 | * | * | * | * | 6.8 | 55.6 | 186.6 | 536.4 | 1,237.0 | 2,226.6 | 4,813.2 |
| 1971 | 196.8 | * | * | * | 1.1 | 8.4 | 56.3 | 193.0 | 508.9 | 1,275.5 | 2,312.5 | 4,771.2 |
| 1970 | 200.4 | * | * | * | 1.0 | 9.6 | 58.9 | 203.0 | 557.1 | 1,270.3 | 2,482.5 | 4,227.9 |
| 1969 | 204.2 | * | * | * | 1.2 | 9.8 | 62.5 | 208.1 | 582.0 | 1,310.4 | 2,459.1 | 4,930.8 |
| 1968 | 213.4 | * | * | * | * | 10.5 | 68.6 | 222.9 | 623.6 | 1,362.1 | 2,607.9 | 5,096.7 |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| ISCHEMIC HEART DISEASE (410-413) | | | | | | | | | | | | |
| BLACK, BOTH SEXES | | | | | | | | | | | | |
| 1978 | 211.1 | * | * | * | 0.7 | 8.7 | 63.5 | 243.6 | 593.9 | 1,198.5 | 2,252.4 | 4,745.3 |
| 1977 | 214.8 | * | * | * | 0.8 | 8.9 | 68.2 | 252.6 | 612.6 | 1,249.9 | 2,234.9 | 4,873.4 |
| 1976 | 215.4 | * | * | * | 0.7 | 9.3 | 65.5 | 257.3 | 617.2 | 1,289.9 | 2,258.7 | 4,841.9 |
| 1975 | 216.5 | * | * | * | 0.9 | 9.7 | 72.5 | 259.0 | 628.1 | 1,360.2 | 2,257.3 | 4,785.4 |
| 1974 | 228.6 | * | * | * | 0.8 | 11.2 | 74.6 | 280.4 | 673.5 | 1,492.2 | 2,364.7 | 5,281.0 |
| 1973 | 241.0 | * | * | * | 1.1 | 11.2 | 83.0 | 297.9 | 724.1 | 1,592.7 | 2,534.3 | 5,501.8 |
| 1972 | 240.0 | * | * | * | 1.4 | 11.6 | 88.8 | 305.3 | 741.1 | 1,562.5 | 2,627.7 | 5,225.5 |
| 1971 | 238.5 | * | * | * | 1.6 | 14.7 | 91.8 | 296.9 | 723.0 | 1,579.6 | 2,688.5 | 5,154.2 |
| 1970 | 245.7 | * | * | * | 1.5 | 15.7 | 98.5 | 312.5 | 762.9 | 1,606.7 | 2,915.3 | 4,545.8 |
| 1969 | 248.9 | * | * | * | 1.6 | 17.6 | 101.4 | 324.4 | 785.1 | 1,627.6 | 2,925.7 | 5,273.2 |
| 1968 | 257.6 | * | * | * | 1.4 | 16.7 | 107.2 | 333.3 | 841.5 | 1,684.0 | 2,944.2 | 5,605.4 |
| BLACK, MALE | | | | | | | | | | | | |
| 1978 | 236.8 | * | * | * | 0.8 | 13.9 | 95.3 | 349.5 | 813.0 | 1,523.6 | 2,616.7 | 5,129.4 |
| 1977 | 239.9 | * | * | * | 1.0 | 13.7 | 100.1 | 360.1 | 826.9 | 1,568.8 | 2,597.4 | 5,324.5 |
| 1976 | 241.9 | * | * | * | 0.8 | 13.6 | 99.4 | 367.7 | 836.5 | 1,607.3 | 2,614.5 | 5,319.1 |
| 1975 | 242.9 | * | * | * | 1.3 | 14.3 | 105.8 | 368.6 | 817.0 | 1,675.3 | 2,681.5 | 5,252.2 |
| 1974 | 256.5 | * | * | * | * | 15.9 | 104.2 | 389.3 | 887.1 | 1,840.1 | 2,804.3 | 5,727.9 |
| 1973 | 269.0 | * | * | * | 1.2 | 16.2 | 112.6 | 409.5 | 950.3 | 1,916.0 | 2,987.3 | 5,870.7 |
| 1972 | 271.0 | * | * | * | 2.3 | 16.3 | 120.7 | 425.9 | 951.6 | 1,897.2 | 3,097.5 | 5,745.0 |
| 1971 | 268.8 | * | * | * | 2.1 | 21.1 | 127.2 | 400.0 | 941.0 | 1,883.1 | 3,132.8 | 5,575.0 |
| 1970 | 277.2 | * | * | * | 1.9 | 21.5 | 137.4 | 418.8 | 963.7 | 1,943.6 | 3,326.8 | 4,743.7 |
| 1969 | 282.0 | * | * | * | 2.0 | 25.5 | 141.0 | 440.4 | 985.2 | 1,948.8 | 3,386.8 | 5,602.7 |
| 1968 | 290.8 | * | * | * | 1.9 | 22.9 | 145.1 | 443.1 | 1,056.7 | 2,004.0 | 3,294.5 | 6,302.9 |
| BLACK, FEMALE | | | | | | | | | | | | |
| 1978 | 188.0 | * | * | * | * | 4.1 | 37.1 | 156.4 | 415.1 | 955.7 | 2,009.2 | 4,547.5 |
| 1977 | 192.2 | * | * | * | 0.7 | 4.7 | 41.8 | 163.6 | 437.8 | 1,010.5 | 2,000.6 | 4,638.3 |
| 1976 | 191.6 | * | * | * | * | 5.6 | 37.4 | 165.6 | 435.8 | 1,048.3 | 2,020.1 | 4,589.9 |
| 1975 | 192.7 | * | * | * | * | 5.8 | 44.9 | 167.2 | 471.4 | 1,121.3 | 1,970.1 | 4,529.8 |
| 1974 | 203.3 | * | * | * | 0.8 | 7.3 | 50.0 | 188.9 | 496.3 | 1,227.7 | 2,067.3 | 5,034.6 |
| 1973 | 215.7 | * | * | * | 0.9 | 6.9 | 58.3 | 203.0 | 535.9 | 1,342.8 | 2,230.7 | 5,294.5 |
| 1972 | 212.0 | * | * | * | * | 7.6 | 62.4 | 202.4 | 564.4 | 1,304.3 | 2,287.9 | 4,928.6 |
| 1971 | 211.0 | * | * | * | 1.2 | 9.3 | 62.5 | 208.3 | 538.5 | 1,337.9 | 2,369.5 | 4,903.0 |
| 1970 | 217.0 | * | * | * | 1.1 | 10.9 | 66.3 | 220.7 | 591.9 | 1,339.8 | 2,613.6 | 4,418.2 |
| 1969 | 218.8 | * | * | * | 1.3 | 10.9 | 68.6 | 224.0 | 614.6 | 1,372.0 | 2,573.6 | 5,070.0 |
| 1968 | 227.5 | * | * | * | * | 11.6 | 75.4 | 237.8 | 655.3 | 1,423.6 | 2,683.8 | 5,269.6 |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| ACUTE MYOCARDIAL INFARCTION (410) | | | | | | | | | | | | |
| BOTH SEXES | | | | | | | | | | | | |
| 1978 | 136.3 | * | * | * | 0.2 | 2.4 | 22.1 | 98.3 | 266.5 | 597.5 | 1,163.6 | 1,974.6 |
| 1977 | 139.4 | * | * | * | 0.2 | 2.5 | 24.4 | 104.1 | 280.1 | 621.2 | 1,189.4 | 1,997.6 |
| 1976 | 146.8 | * | * | * | 0.2 | 2.7 | 27.0 | 111.8 | 304.6 | 658.1 | 1,259.2 | 2,113.3 |
| 1975 | 150.7 | * | * | * | 0.3 | 2.8 | 28.8 | 119.8 | 320.2 | 689.4 | 1,291.9 | 2,118.1 |
| 1974 | 156.6 | * | * | * | 0.2 | 3.1 | 29.9 | 127.4 | 336.3 | 727.9 | 1,352.5 | 2,294.4 |
| 1973 | 166.4 | * | * | * | 0.3 | 3.5 | 33.8 | 134.5 | 365.8 | 773.2 | 1,466.3 | 2,453.0 |
| 1972 | 171.0 | * | * | * | 0.4 | 3.5 | 35.2 | 139.4 | 376.5 | 814.5 | 1,520.2 | 2,490.5 |
| 1971 | 173.0 | 0.7 | * | 0.1 | 0.4 | 4.2 | 36.8 | 142.6 | 384.2 | 822.6 | 1,551.3 | 2,559.7 |
| 1970 | 175.8 | * | * | * | 0.4 | 4.3 | 37.8 | 146.2 | 393.9 | 851.6 | 1,595.8 | 2,412.9 |
| 1969 | 179.5 | * | * | 0.1 | 0.4 | 4.3 | 40.4 | 152.6 | 400.1 | 870.3 | 1,628.8 | 2,768.5 |
| 1968 | 185.4 | * | * | * | 0.4 | 4.6 | 42.3 | 158.5 | 420.8 | 900.5 | 1,687.1 | 2,911.8 |
| MALE | | | | | | | | | | | | |
| 1978 | 170.5 | * | * | * | 0.3 | 3.9 | 37.3 | 164.7 | 424.8 | 893.3 | 1,601.5 | 2,471.5 |
| 1977 | 176.3 | * | * | * | 0.4 | 4.1 | 41.1 | 175.6 | 447.1 | 934.7 | 1,634.1 | 2,525.7 |
| 1976 | 186.5 | * | * | * | 0.3 | 4.5 | 46.0 | 187.7 | 485.9 | 988.5 | 1,724.7 | 2,662.0 |
| 1975 | 192.8 | * | * | * | 0.4 | 4.6 | 48.5 | 201.6 | 512.7 | 1,032.3 | 1,765.8 | 2,653.2 |
| 1974 | 201.1 | * | * | * | 0.3 | 4.9 | 50.6 | 216.2 | 539.7 | 1,090.3 | 1,830.0 | 2,851.2 |
| 1973 | 214.2 | * | * | * | 0.5 | 5.4 | 56.6 | 226.7 | 587.8 | 1,153.2 | 1,981.5 | 3,043.5 |
| 1972 | 220.3 | * | * | * | 0.5 | 5.5 | 58.6 | 236.2 | 604.0 | 1,207.4 | 2,029.3 | 3,066.7 |
| 1971 | 223.9 | * | * | * | 0.5 | 6.8 | 62.2 | 240.1 | 611.9 | 1,214.4 | 2,073.4 | 3,188.0 |
| 1970 | 228.6 | * | * | * | 0.6 | 6.7 | 64.0 | 246.4 | 626.0 | 1,249.4 | 2,123.0 | 2,874.9 |
| 1969 | 235.2 | * | * | * | 0.5 | 6.9 | 68.1 | 258.0 | 640.7 | 1,279.8 | 2,161.1 | 3,392.4 |
| 1968 | 243.0 | * | * | * | 0.5 | 7.2 | 70.9 | 267.1 | 668.3 | 1,315.0 | 2,228.4 | 3,570.7 |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| ACUTE MYOCARDIAL INFARCTION (410) | | | | | | | | | | | | |
| FEMALE | | | | | | | | | | | | |
| 1978 | 103.9 | * | * | * | 0.1 | 1.0 | 7.5 | 36.5 | 126.9 | 371.4 | 902.8 | 1,750.0 |
| 1977 | 104.5 | * | * | * | 0.1 | 1.1 | 8.4 | 37.6 | 132.5 | 381.3 | 923.2 | 1,755.0 |
| 1976 | 109.3 | * | * | * | 0.1 | 1.0 | 8.8 | 41.2 | 144.1 | 405.1 | 978.1 | 1,855.5 |
| 1975 | 110.8 | * | * | * | 0.1 | 1.2 | 10.0 | 43.7 | 149.7 | 426.7 | 1,003.3 | 1,859.1 |
| 1974 | 114.5 | * | * | * | 0.2 | 1.4 | 10.1 | 44.9 | 156.0 | 450.1 | 1,058.3 | 2,018.7 |
| 1973 | 121.0 | * | * | * | 0.2 | 1.6 | 12.0 | 48.9 | 168.6 | 481.3 | 1,144.3 | 2,155.8 |
| 1972 | 124.1 | * | * | * | 0.2 | 1.6 | 12.8 | 49.4 | 173.9 | 511.4 | 1,195.7 | 2,195.7 |
| 1971 | 124.6 | * | * | * | 0.3 | 1.7 | 12.8 | 52.0 | 180.7 | 519.2 | 1,212.9 | 2,231.7 |
| 1970 | 125.8 | * | * | * | 0.2 | 2.1 | 12.9 | 52.7 | 185.6 | 542.6 | 1,247.0 | 2,154.2 |
| 1969 | 126.7 | * | * | * | 0.3 | 1.8 | 14.2 | 54.3 | 183.6 | 550.4 | 1,271.6 | 2,430.8 |
| 1968 | 130.6 | * | * | * | 0.3 | 2.2 | 15.2 | 56.8 | 197.1 | 574.1 | 1,316.5 | 2,553.0 |
| WHITE, BOTH SEXES | | | | | | | | | | | | |
| 1978 | 145.7 | * | * | * | 0.2 | 2.2 | 21.5 | 98.3 | 269.6 | 610.7 | 1,201.0 | 2,034.4 |
| 1977 | 149.1 | * | * | * | 0.2 | 2.3 | 23.8 | 104.4 | 283.2 | 636.3 | 1,229.5 | 2,056.4 |
| 1976 | 156.7 | * | * | * | 0.2 | 2.5 | 26.6 | 111.8 | 307.8 | 671.5 | 1,303.0 | 2,175.4 |
| 1975 | 160.7 | * | * | * | 0.2 | 2.6 | 28.1 | 120.3 | 324.3 | 703.6 | 1,337.1 | 2,178.3 |
| 1974 | 166.8 | * | * | * | 0.2 | 2.8 | 29.4 | 127.3 | 340.6 | 740.6 | 1,399.3 | 2,359.3 |
| 1973 | 176.6 | * | * | * | 0.3 | 3.2 | 33.2 | 133.6 | 369.0 | 786.2 | 1,514.1 | 2,520.1 |
| 1972 | 181.6 | * | * | * | 0.3 | 3.2 | 34.2 | 138.9 | 380.3 | 832.2 | 1,567.0 | 2,562.6 |
| 1971 | 183.6 | * | * | * | 0.3 | 3.7 | 35.8 | 142.6 | 389.5 | 839.1 | 1,595.9 | 2,647.3 |
| 1970 | 186.6 | * | * | * | 0.3 | 3.8 | 36.5 | 145.8 | 398.6 | 871.1 | 1,639.3 | 2,515.0 |
| 1969 | 190.1 | * | * | * | 0.3 | 3.6 | 38.8 | 151.6 | 404.1 | 889.7 | 1,673.3 | 2,869.1 |
| 1968 | 195.9 | * | * | * | 0.3 | 4.1 | 40.3 | 157.6 | 423.9 | 919.8 | 1,732.1 | 3,012.9 |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| ACUTE MYOCARDIAL INFARCTION (410) | | | | | | | | | | | | |
| WHITE, MALE | | | | | | | | | | | | |
| 1978 | 182.8 | * | * | * | 0.3 | 3.5 | 36.8 | 167.3 | 434.9 | 924.1 | 1,668.6 | 2,559.3 |
| 1977 | 189.1 | * | * | * | 0.3 | 3.7 | 40.8 | 178.4 | 458.0 | 969.1 | 1,707.2 | 2,615.2 |
| 1976 | 199.6 | * | * | * | 0.2 | 4.2 | 45.9 | 190.4 | 497.1 | 1,021.2 | 1,805.6 | 2,758.8 |
| 1975 | 206.3 | * | * | * | 0.4 | 4.3 | 48.1 | 205.3 | 527.1 | 1,066.4 | 1,843.8 | 2,752.6 |
| 1974 | 214.7 | * | * | * | 0.3 | 4.5 | 50.7 | 219.6 | 553.8 | 1,125.2 | 1,905.6 | 2,953.8 |
| 1973 | 228.3 | * | * | * | 0.4 | 5.0 | 57.0 | 229.4 | 601.9 | 1,189.5 | 2,061.7 | 3,158.0 |
| 1972 | 234.9 | * | * | * | 0.4 | 5.1 | 58.2 | 239.2 | 619.9 | 1,249.3 | 2,106.0 | 3,183.2 |
| 1971 | 238.6 | * | * | * | 0.4 | 6.2 | 61.8 | 244.5 | 628.6 | 1,255.7 | 2,151.4 | 3,325.5 |
| 1970 | 243.6 | * | * | * | 0.5 | 6.1 | 63.4 | 250.3 | 643.9 | 1,293.5 | 2,198.5 | 3,029.6 |
| 1969 | 250.2 | * | * | * | 0.5 | 6.0 | 67.0 | 261.2 | 658.5 | 1,324.8 | 2,234.0 | 3,559.8 |
| 1968 | 258.0 | * | * | * | 0.4 | 6.5 | 69.6 | 270.4 | 684.3 | 1,360.8 | 2,306.5 | 3,715.3 |
| WHITE, FEMALE | | | | | | | | | | | | |
| 1978 | 110.5 | * | * | * | * | 0.9 | 6.5 | 33.0 | 122.8 | 371.5 | 925.9 | 1,800.5 |
| 1977 | 111.0 | * | * | * | * | 0.9 | 7.1 | 34.5 | 127.6 | 382.3 | 947.5 | 1,803.3 |
| 1976 | 115.7 | * | * | * | * | 0.9 | 7.7 | 37.6 | 139.3 | 404.3 | 1,003.7 | 1,907.6 |
| 1975 | 117.2 | * | * | * | 0.1 | 1.0 | 8.5 | 40.3 | 143.8 | 426.7 | 1,032.2 | 1,906.7 |
| 1974 | 121.0 | * | * | * | * | 1.2 | 8.7 | 40.7 | 150.7 | 447.2 | 1,090.2 | 2,069.3 |
| 1973 | 127.2 | * | * | * | 0.1 | 1.3 | 10.1 | 43.8 | 161.2 | 478.0 | 1,175.0 | 2,206.7 |
| 1972 | 130.7 | * | * | * | 0.2 | 1.2 | 10.8 | 44.8 | 166.1 | 512.1 | 1,227.2 | 2,250.4 |
| 1971 | 131.2 | * | * | * | 0.2 | 1.2 | 10.7 | 47.2 | 175.1 | 518.1 | 1,240.3 | 2,300.0 |
| 1970 | 132.3 | * | * | * | 0.2 | 1.6 | 10.7 | 47.7 | 178.1 | 545.0 | 1,273.6 | 2,233.4 |
| 1969 | 132.9 | * | * | * | 0.2 | 1.3 | 11.6 | 48.6 | 174.8 | 551.3 | 1,300.2 | 2,506.8 |
| 1968 | 136.7 | * | * | * | 0.2 | 1.7 | 12.1 | 51.3 | 188.4 | 574.4 | 1,342.8 | 2,637.8 |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| ACUTE MYOCARDIAL INFARCTION (410) | | | | | | | | | | | | |
| ALL OTHER, BOTH SEXES | | | | | | | | | | | | |
| 1978 | 76.0 | * | * | * | 0.4 | 3.9 | 26.4 | 98.5 | 238.8 | 474.8 | 778.4 | 1,288.7 |
| 1977 | 77.1 | * | * | * | 0.4 | 4.0 | 28.6 | 101.9 | 251.9 | 479.1 | 776.8 | 1,324.4 |
| 1976 | 82.7 | * | * | * | 0.4 | 4.0 | 30.0 | 112.3 | 274.5 | 530.0 | 808.4 | 1,391.5 |
| 1975 | 84.0 | * | * | * | 0.5 | 4.4 | 34.1 | 115.1 | 282.2 | 551.1 | 808.2 | 1,428.1 |
| 1974 | 88.5 | * | * | * | 0.4 | 5.2 | 33.4 | 128.0 | 296.5 | 603.4 | 835.5 | 1,539.3 |
| 1973 | 96.7 | * | * | * | 0.7 | 5.7 | 38.0 | 142.1 | 335.7 | 645.1 | 923.2 | 1,670.9 |
| 1972 | 97.8 | * | * | * | 0.6 | 6.1 | 42.6 | 143.7 | 340.6 | 639.3 | 975.0 | 1,658.5 |
| 1971 | 98.5 | * | * | * | 0.9 | 7.8 | 44.3 | 142.7 | 334.2 | 660.2 | 1,010.9 | 1,561.7 |
| 1970 | 100.2 | * | * | * | 0.9 | 8.1 | 47.0 | 149.4 | 350.1 | 661.1 | 1,042.6 | 1,364.8 |
| 1969 | 104.1 | * | * | * | 1.0 | 9.5 | 52.7 | 162.2 | 362.1 | 677.4 | 1,055.5 | 1,635.8 |
| 1968 | 109.5 | * | * | * | 1.0 | 8.7 | 57.9 | 166.6 | 390.5 | 706.7 | 1,103.1 | 1,782.4 |
| ALL OTHER, MALE | | | | | | | | | | | | |
| 1978 | 90.7 | * | * | * | * | 6.6 | 41.2 | 143.1 | 330.7 | 610.6 | 964.3 | 1,572.4 |
| 1977 | 92.1 | * | * | * | * | 6.5 | 43.1 | 152.0 | 343.2 | 614.8 | 946.4 | 1,612.5 |
| 1976 | 98.9 | * | * | * | * | 6.1 | 46.4 | 164.6 | 378.9 | 679.1 | 966.1 | 1,692.7 |
| 1975 | 100.7 | * | * | * | 0.7 | 6.8 | 51.7 | 168.7 | 376.5 | 707.8 | 994.2 | 1,659.3 |
| 1974 | 106.8 | * | * | * | * | 7.6 | 50.0 | 186.0 | 404.8 | 759.2 | 1,052.0 | 1,794.0 |
| 1973 | 115.6 | * | * | * | 0.9 | 8.5 | 54.2 | 203.0 | 451.6 | 806.8 | 1,132.6 | 1,875.0 |
| 1972 | 117.9 | * | * | * | 1.1 | 8.7 | 61.8 | 209.3 | 449.4 | 805.6 | 1,205.5 | 1,889.4 |
| 1971 | 118.9 | * | * | * | 1.2 | 11.0 | 65.3 | 200.3 | 450.2 | 819.8 | 1,215.0 | 1,834.0 |
| 1970 | 121.6 | * | * | * | 1.3 | 11.3 | 69.2 | 211.1 | 455.7 | 834.0 | 1,245.8 | 1,515.6 |
| 1969 | 126.4 | * | * | * | 1.1 | 14.5 | 76.7 | 228.7 | 468.3 | 843.2 | 1,288.3 | 1,813.6 |
| 1968 | 133.2 | * | * | * | 1.3 | 13.1 | 81.6 | 236.2 | 512.5 | 870.1 | 1,291.4 | 2,163.4 |
| ALL OTHER, FEMALE | | | | | | | | | | | | |
| 1978 | 62.6 | * | * | * | * | 1.5 | 14.0 | 61.7 | 162.5 | 369.7 | 653.2 | 1,149.5 |
| 1977 | 63.6 | * | * | * | * | 2.0 | 16.4 | 60.4 | 175.6 | 372.4 | 659.3 | 1,169.2 |
| 1976 | 67.9 | * | * | * | * | 2.2 | 16.2 | 68.5 | 187.1 | 412.1 | 700.5 | 1,222.4 |
| 1975 | 68.8 | * | * | * | * | 2.3 | 19.3 | 69.8 | 202.6 | 426.4 | 678.8 | 1,278.5 |
| 1974 | 71.9 | * | * | * | * | 3.1 | 19.6 | 78.8 | 204.3 | 478.7 | 686.1 | 1,389.4 |
| 1973 | 79.4 | * | * | * | * | 3.2 | 24.4 | 89.9 | 236.9 | 514.5 | 778.6 | 1,546.8 |
| 1972 | 79.3 | * | * | * | * | 3.8 | 26.6 | 87.3 | 246.7 | 505.2 | 812.0 | 1,515.8 |
| 1971 | 79.8 | * | * | * | * | 5.0 | 26.9 | 92.7 | 232.9 | 529.6 | 860.0 | 1,386.3 |
| 1970 | 80.6 | * | * | * | * | 5.4 | 28.5 | 95.2 | 256.8 | 518.5 | 888.1 | 1,258.5 |
| 1969 | 83.7 | * | * | * | 0.9 | 5.2 | 32.6 | 103.7 | 267.9 | 540.9 | 874.1 | 1,515.4 |
| 1968 | 87.7 | * | * | * | * | 5.0 | 37.9 | 105.3 | 281.0 | 571.2 | 961.1 | 1,526.2 |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| ACUTE MYOCARDIAL INFARCTION (410) | | | | | | | | | | | | |
| BLACK, BOTH SEXES | | | | | | | | | | | | |
| 1978 | 82.1 | * | * | * | 0.4 | 4.4 | 29.5 | 108.4 | 255.1 | 495.8 | 788.1 | 1,316.0 |
| 1977 | 83.4 | * | * | * | 0.5 | 4.7 | 32.3 | 111.9 | 266.6 | 503.6 | 784.2 | 1,353.1 |
| 1976 | 89.2 | * | * | * | 0.5 | 4.6 | 33.6 | 123.0 | 291.1 | 555.9 | 819.4 | 1,432.4 |
| 1975 | 90.1 | * | * | * | 0.5 | 5.0 | 37.8 | 124.4 | 297.6 | 578.5 | 812.4 | 1,468.5 |
| 1974 | 94.4 | * | * | * | 0.4 | 5.9 | 36.8 | 137.8 | 311.1 | 630.7 | 841.9 | 1,581.0 |
| 1973 | 102.8 | * | * | * | 0.7 | 6.5 | 42.0 | 152.7 | 353.1 | 676.4 | 927.1 | 1,701.8 |
| 1972 | 103.1 | * | * | * | 0.7 | 6.7 | 46.2 | 153.9 | 354.3 | 663.4 | 989.5 | 1,689.1 |
| 1971 | 103.6 | * | * | * | 1.0 | 8.7 | 48.4 | 150.6 | 348.0 | 685.8 | 1,027.6 | 1,612.1 |
| 1970 | 106.5 | * | * | * | 0.9 | 9.1 | 51.9 | 159.9 | 369.3 | 688.6 | 1,082.9 | 1,438.0 |
| 1969 | 109.5 | * | * | * | 1.0 | 10.4 | 56.9 | 172.2 | 379.8 | 700.3 | 1,091.1 | 1,696.9 |
| 1968 | 114.6 | * | * | * | 1.0 | 9.6 | 62.8 | 175.2 | 407.1 | 730.8 | 1,125.1 | 1,832.6 |
| BLACK, MALE | | | | | | | | | | | | |
| 1978 | 96.6 | * | * | * | * | 7.5 | 45.4 | 155.8 | 351.6 | 637.9 | 962.8 | 1,582.4 |
| 1977 | 98.2 | * | * | * | * | 7.4 | 48.5 | 164.5 | 362.3 | 650.0 | 950.2 | 1,626.5 |
| 1976 | 105.7 | * | * | * | * | 6.9 | 52.0 | 177.9 | 401.7 | 716.0 | 977.8 | 1,761.7 |
| 1975 | 106.5 | * | * | * | 0.8 | 7.7 | 56.6 | 181.0 | 394.2 | 748.2 | 987.4 | 1,708.7 |
| 1974 | 112.5 | * | * | * | * | 8.7 | 54.6 | 198.1 | 425.6 | 797.8 | 1,057.9 | 1,844.2 |
| 1973 | 121.8 | * | * | * | 0.8 | 9.8 | 59.1 | 216.8 | 476.9 | 851.5 | 1,135.3 | 1,941.5 |
| 1972 | 123.2 | * | * | * | 1.1 | 9.6 | 66.2 | 225.3 | 469.6 | 837.1 | 1,219.3 | 1,945.0 |
| 1971 | 123.7 | * | * | * | 1.3 | 12.5 | 70.7 | 210.3 | 469.7 | 851.0 | 1,238.7 | 1,935.0 |
| 1970 | 128.2 | * | * | * | 1.4 | 12.7 | 76.0 | 226.0 | 484.1 | 875.2 | 1,295.1 | 1,624.0 |
| 1969 | 131.6 | * | * | * | 1.1 | 16.0 | 82.4 | 242.0 | 493.8 | 872.7 | 1,331.1 | 1,932.4 |
| 1968 | 138.1 | * | * | * | 1.3 | 14.5 | 87.8 | 248.4 | 537.3 | 901.3 | 1,321.8 | 2,282.9 |
| BLACK, FEMALE | | | | | | | | | | | | |
| 1978 | 69.1 | * | * | * | * | 1.8 | 16.3 | 69.5 | 176.3 | 389.8 | 671.5 | 1,178.8 |
| 1977 | 70.0 | * | * | * | * | 2.4 | 18.8 | 68.3 | 188.4 | 393.7 | 676.3 | 1,210.6 |
| 1976 | 74.3 | * | * | * | * | 2.6 | 18.4 | 77.3 | 199.6 | 434.1 | 713.2 | 1,258.4 |
| 1975 | 75.3 | * | * | * | * | 2.7 | 22.2 | 77.0 | 217.5 | 449.8 | 693.9 | 1,336.9 |
| 1974 | 77.9 | * | * | * | * | 3.4 | 21.9 | 87.1 | 216.1 | 503.4 | 695.8 | 1,435.9 |
| 1973 | 85.7 | * | * | * | * | 3.7 | 27.8 | 98.2 | 250.1 | 541.1 | 786.8 | 1,567.1 |
| 1972 | 84.8 | * | * | * | * | 4.3 | 29.7 | 93.0 | 257.7 | 529.2 | 824.3 | 1,542.9 |
| 1971 | 85.3 | * | * | * | * | 5.5 | 30.0 | 99.2 | 245.2 | 554.6 | 876.1 | 1,419.4 |
| 1970 | 86.8 | * | * | * | * | 6.0 | 32.0 | 102.9 | 271.6 | 540.8 | 927.3 | 1,318.2 |
| 1969 | 89.4 | * | * | * | 0.9 | 5.8 | 35.7 | 111.7 | 282.5 | 563.1 | 909.7 | 1,551.7 |
| 1968 | 93.1 | * | * | * | * | 5.5 | 41.8 | 111.5 | 294.4 | 592.4 | 978.8 | 1,583.9 |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|--|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| OTHER ACUTE AND SUBACUTE FORMS OF ISCHEMIC HEART DISEASE (411) | | | | | | | | | | | | |
| BOTH SEXES | | | | | | | | | | | | |
| 1978 | 1.8 | * | * | * | * | * | 0.5 | 1.9 | 4.0 | 7.5 | 12.7 | 28.4 |
| 1977 | 1.8 | * | * | * | * | 0.1 | 0.5 | 1.8 | 4.1 | 7.5 | 12.7 | 27.5 |
| 1976 | 1.9 | * | * | * | * | 0.1 | 0.5 | 2.0 | 4.3 | 7.5 | 12.8 | 29.1 |
| 1975 | 1.9 | * | * | * | * | 0.1 | 0.5 | 2.0 | 4.6 | 7.8 | 13.5 | 26.6 |
| 1974 | 2.1 | * | * | * | * | 0.1 | 0.7 | 2.3 | 5.0 | 8.6 | 14.7 | 31.3 |
| 1973 | 2.1 | * | * | * | * | 0.1 | 0.7 | 2.4 | 5.0 | 8.8 | 15.5 | 35.8 |
| 1972 | 2.2 | * | * | * | * | 0.1 | 0.7 | 2.6 | 5.2 | 8.9 | 15.6 | 33.9 |
| 1971 | 1.9 | * | * | * | * | 0.1 | 0.6 | 2.0 | 4.3 | 7.9 | 15.8 | 35.6 |
| 1970 | 2.1 | * | * | * | * | 0.1 | 0.8 | 2.3 | 4.9 | 8.3 | 16.7 | 34.4 |
| 1969 | 2.3 | * | * | * | * | 0.1 | 0.9 | 2.2 | 5.3 | 9.9 | 19.1 | 42.8 |
| 1968 | 2.4 | * | * | * | * | 0.1 | 0.7 | 2.4 | 5.3 | 10.5 | 19.0 | 45.4 |
| MALE | | | | | | | | | | | | |
| 1978 | 2.4 | * | * | * | * | * | 0.8 | 3.1 | 6.5 | 12.4 | 17.8 | 34.2 |
| 1977 | 2.4 | * | * | * | * | * | 0.8 | 3.0 | 6.7 | 11.9 | 19.0 | 34.0 |
| 1976 | 2.5 | * | * | * | * | * | 0.8 | 3.4 | 7.1 | 12.2 | 18.0 | 35.1 |
| 1975 | 2.6 | * | * | * | * | 0.1 | 0.9 | 3.4 | 7.6 | 12.0 | 19.7 | 33.5 |
| 1974 | 2.7 | * | * | * | * | 0.2 | 1.1 | 3.6 | 8.0 | 13.3 | 20.8 | 38.1 |
| 1973 | 2.8 | * | * | * | * | * | 1.1 | 3.9 | 7.8 | 13.7 | 22.2 | 40.9 |
| 1972 | 2.9 | * | * | * | * | 0.2 | 1.1 | 4.3 | 8.5 | 13.9 | 21.1 | 39.1 |
| 1971 | 2.4 | * | * | * | * | * | 1.0 | 3.1 | 6.8 | 11.6 | 20.6 | 38.2 |
| 1970 | 2.7 | * | * | * | * | 0.2 | 1.3 | 3.9 | 7.8 | 11.9 | 21.9 | 41.3 |
| 1969 | 3.0 | * | * | * | * | 0.2 | 1.3 | 3.5 | 8.4 | 14.3 | 25.1 | 48.4 |
| 1968 | 3.0 | * | * | * | * | * | 1.2 | 3.8 | 8.0 | 15.7 | 25.1 | 49.8 |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|--|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| OTHER ACUTE AND SUBACUTE FORMS OF ISCHEMIC HEART DISEASE (411) | | | | | | | | | | | | |
| FEMALE | | | | | | | | | | | | |
| 1978 | 1.3 | * | * | * | * | * | 0.2 | 0.7 | 1.7 | 3.8 | 9.8 | 25.8 |
| 1977 | 1.2 | * | * | * | * | * | 0.2 | 0.7 | 1.8 | 4.1 | 8.9 | 24.5 |
| 1976 | 1.2 | * | * | * | * | * | * | 0.7 | 1.7 | 4.0 | 9.7 | 26.2 |
| 1975 | 1.3 | * | * | * | * | * | 0.2 | 0.8 | 2.0 | 4.5 | 9.7 | 23.2 |
| 1974 | 1.4 | * | * | * | * | * | 0.3 | 1.1 | 2.3 | 4.9 | 10.9 | 28.0 |
| 1973 | 1.5 | * | * | * | * | * | 0.3 | 0.9 | 2.5 | 5.0 | 11.3 | 33.2 |
| 1972 | 1.5 | * | * | * | * | * | 0.3 | 1.0 | 2.4 | 5.1 | 12.1 | 31.2 |
| 1971 | 1.5 | * | * | * | * | * | 0.2 | 1.0 | 2.0 | 5.0 | 12.7 | 34.3 |
| 1970 | 1.5 | * | * | * | * | * | 0.3 | 0.8 | 2.3 | 5.5 | 13.3 | 30.6 |
| 1969 | 1.7 | * | * | * | * | * | 0.4 | 1.0 | 2.6 | 6.5 | 15.0 | 39.9 |
| 1968 | 1.7 | * | * | * | * | * | 0.3 | 1.2 | 2.7 | 6.4 | 14.8 | 43.1 |
| WHITE, BOTH SEXES | | | | | | | | | | | | |
| 1978 | 1.8 | * | * | * | * | * | 0.4 | 1.8 | 3.7 | 7.2 | 12.3 | 28.4 |
| 1977 | 1.8 | * | * | * | * | * | 0.5 | 1.8 | 3.8 | 7.2 | 12.3 | 27.2 |
| 1976 | 1.8 | * | * | * | * | 0.1 | 0.4 | 1.8 | 3.9 | 7.2 | 12.6 | 29.0 |
| 1975 | 1.9 | * | * | * | * | 0.1 | 0.4 | 1.8 | 4.1 | 7.5 | 13.3 | 26.5 |
| 1974 | 2.1 | * | * | * | * | * | 0.6 | 2.1 | 4.6 | 8.1 | 14.6 | 31.3 |
| 1973 | 2.1 | * | * | * | * | 0.1 | 0.6 | 2.2 | 4.6 | 8.2 | 15.5 | 36.1 |
| 1972 | 2.1 | * | * | * | * | 0.1 | 0.6 | 2.3 | 4.9 | 8.7 | 15.2 | 34.2 |
| 1971 | 2.0 | * | * | * | * | 0.1 | 0.6 | 1.8 | 4.0 | 7.5 | 16.0 | 36.6 |
| 1970 | 2.1 | * | * | * | * | 0.1 | 0.6 | 2.0 | 4.5 | 8.0 | 16.4 | 35.6 |
| 1969 | 2.2 | * | * | * | * | 0.1 | 0.7 | 1.8 | 4.7 | 9.0 | 18.6 | 43.6 |
| 1968 | 2.3 | * | * | * | * | * | 0.6 | 2.1 | 4.6 | 9.7 | 18.7 | 44.8 |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|--|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| OTHER ACUTE AND SUBACUTE FORMS OF ISCHEMIC HEART DISEASE (411) | | | | | | | | | | | | |
| WHITE, MALE | | | | | | | | | | | | |
| 1978 | 2.4 | * | * | * | * | * | 0.7 | 2.9 | 6.1 | 12.1 | 17.2 | 33.5 |
| 1977 | 2.4 | * | * | * | * | * | 0.8 | 3.0 | 6.3 | 11.5 | 18.3 | 33.3 |
| 1976 | 2.5 | * | * | * | * | * | 0.7 | 3.1 | 6.7 | 12.0 | 18.1 | 35.8 |
| 1975 | 2.5 | * | * | * | * | * | 0.8 | 3.1 | 6.9 | 11.7 | 19.6 | 33.9 |
| 1974 | 2.7 | * | * | * | * | * | 0.9 | 3.3 | 7.5 | 13.0 | 20.8 | 36.5 |
| 1973 | 2.8 | * | * | * | * | * | 0.9 | 3.7 | 7.3 | 13.0 | 22.1 | 40.2 |
| 1972 | 2.9 | * | * | * | * | 0.2 | 1.0 | 3.9 | 8.2 | 13.7 | 20.3 | 40.0 |
| 1971 | 2.4 | * | * | * | * | * | 1.0 | 2.8 | 6.4 | 11.3 | 21.0 | 39.3 |
| 1970 | 2.7 | * | * | * | * | 0.2 | 1.1 | 3.6 | 7.4 | 11.8 | 21.8 | 42.7 |
| 1969 | 2.9 | * | * | * | * | * | 1.1 | 3.1 | 7.7 | 13.2 | 24.3 | 50.6 |
| 1968 | 2.9 | * | * | * | * | * | 1.0 | 3.4 | 7.2 | 14.9 | 24.8 | 49.1 |
| WHITE, FEMALE | | | | | | | | | | | | |
| 1978 | 1.3 | * | * | * | * | * | * | 0.7 | 1.6 | 3.5 | 9.5 | 26.1 |
| 1977 | 1.2 | * | * | * | * | * | * | 0.6 | 1.6 | 3.9 | 8.7 | 24.5 |
| 1976 | 1.2 | * | * | * | * | * | * | 0.6 | 1.4 | 3.6 | 9.3 | 25.9 |
| 1975 | 1.2 | * | * | * | * | * | * | 0.5 | 1.7 | 4.2 | 9.5 | 23.0 |
| 1974 | 1.4 | * | * | * | * | * | 0.3 | 0.9 | 2.0 | 4.4 | 10.8 | 28.7 |
| 1973 | 1.5 | * | * | * | * | * | 0.2 | 0.7 | 2.3 | 4.6 | 11.4 | 34.2 |
| 1972 | 1.5 | * | * | * | * | * | 0.2 | 0.7 | 2.0 | 4.8 | 12.0 | 31.4 |
| 1971 | 1.5 | * | * | * | * | * | 0.2 | 0.9 | 1.8 | 4.6 | 12.9 | 35.2 |
| 1970 | 1.4 | * | * | * | * | * | 0.2 | 0.6 | 1.9 | 5.0 | 12.9 | 31.7 |
| 1969 | 1.6 | * | * | * | * | * | 0.3 | 0.6 | 2.0 | 5.8 | 14.8 | 40.0 |
| 1968 | 1.6 | * | * | * | * | * | 0.2 | 0.8 | 2.3 | 5.7 | 14.6 | 42.5 |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|--|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| OTHER ACUTE AND SUBACUTE FORMS OF ISCHEMIC HEART DISEASE (411) | | | | | | | | | | | | |
| ALL OTHER, BOTH SEXES | | | | | | | | | | | | |
| 1978 | 1.8 | * | * | * | * | * | 0.9 | 2.7 | 6.5 | 10.5 | 17.2 | 28.6 |
| 1977 | 1.8 | * | * | * | * | * | 0.8 | 2.2 | 6.6 | 10.8 | 16.8 | 30.6 |
| 1976 | 2.0 | * | * | * | * | * | 1.0 | 3.5 | 7.7 | 10.4 | 15.3 | 29.4 |
| 1975 | 2.1 | * | * | * | * | * | 1.0 | 4.0 | 9.4 | 10.8 | 15.8 | 27.4 |
| 1974 | 2.2 | * | * | * | * | * | 1.5 | 3.8 | 8.6 | 12.9 | 15.3 | 31.9 |
| 1973 | 2.2 | * | * | * | * | * | 1.6 | 4.2 | 8.3 | 13.8 | 15.5 | 31.5 |
| 1972 | 2.3 | * | * | * | * | * | 1.3 | 5.4 | 8.2 | 11.6 | 20.0 | 29.3 |
| 1971 | 1.9 | * | * | * | * | * | 1.0 | 3.7 | 7.2 | 11.7 | 12.9 | 25.0 |
| 1970 | 2.3 | * | * | * | * | * | 1.8 | 4.8 | 8.3 | 11.7 | 20.1 | 22.4 |
| 1969 | 3.1 | * | * | * | * | * | 2.3 | 5.9 | 11.4 | 19.0 | 25.3 | 33.9 |
| 1968 | 3.1 | * | * | * | * | * | 2.1 | 5.7 | 11.7 | 18.3 | 22.8 | 52.9 |
| ALL OTHER, MALE | | | | | | | | | | | | |
| 1978 | 2.5 | * | * | * | * | * | * | 4.5 | 10.6 | 15.1 | 23.3 | 41.4 |
| 1977 | 2.5 | * | * | * | * | * | * | 3.2 | 10.5 | 16.2 | 25.7 | 41.1 |
| 1976 | 2.5 | * | * | * | * | * | 1.7 | 5.7 | 11.6 | 14.0 | 17.9 | * |
| 1975 | 2.9 | * | * | * | * | * | 1.8 | 5.7 | 15.2 | 14.3 | 20.7 | * |
| 1974 | 2.9 | * | * | * | * | * | 2.2 | 5.7 | 12.6 | 16.8 | 20.5 | 54.0 |
| 1973 | 3.0 | * | * | * | * | * | 2.2 | 6.0 | 12.2 | 19.8 | 23.2 | 47.9 |
| 1972 | 3.0 | * | * | * | * | * | 1.7 | 8.1 | 11.3 | 15.8 | 29.5 | * |
| 1971 | 2.5 | * | * | * | * | * | 1.8 | 5.9 | 10.8 | 14.9 | 16.9 | * |
| 1970 | 2.8 | * | * | * | * | * | 2.8 | 7.0 | 10.8 | 13.2 | 22.3 | * |
| 1969 | 3.8 | * | * | * | * | * | 3.0 | 7.7 | 15.0 | 24.5 | 35.1 | * |
| 1968 | 3.9 | * | * | * | * | * | 3.3 | 7.3 | 16.6 | 24.0 | 29.0 | 56.1 |
| ALL OTHER, FEMALE | | | | | | | | | | | | |
| 1978 | 1.3 | * | * | * | * | * | * | * | 3.1 | 6.8 | 13.1 | 22.0 |
| 1977 | 1.2 | * | * | * | * | * | * | 1.5 | 3.3 | 6.5 | 10.7 | 25.0 |
| 1976 | 1.5 | * | * | * | * | * | * | 1.7 | 4.5 | 7.6 | 13.6 | 29.6 |
| 1975 | 1.5 | * | * | * | * | * | * | 2.5 | 4.5 | 8.0 | 12.4 | 25.8 |
| 1974 | 1.5 | * | * | * | * | * | * | 2.1 | 5.3 | 9.7 | 11.7 | * |
| 1973 | 1.5 | * | * | * | * | * | * | 2.7 | 5.0 | 8.9 | 10.1 | * |
| 1972 | 1.6 | * | * | * | * | * | * | 3.0 | 5.5 | 8.3 | 13.3 | 28.9 |
| 1971 | 1.3 | * | * | * | * | * | * | 1.9 | 4.2 | 9.0 | 10.0 | * |
| 1970 | 1.8 | * | * | * | * | * | * | 2.8 | 6.1 | 10.4 | 18.5 | * |
| 1969 | 2.4 | * | * | * | * | * | 1.6 | 4.3 | 8.2 | 14.4 | 17.8 | 38.5 |
| 1968 | 2.3 | * | * | * | * | * | * | 4.3 | 7.3 | 13.6 | 18.0 | 50.8 |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|--|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| OTHER ACUTE AND SUBACUTE FORMS OF ISCHEMIC HEART DISEASE (411) | | | | | | | | | | | | |
| BLACK, BOTH SEXES | | | | | | | | | | | | |
| 1978 | 2.0 | * | * | * | * | * | 1.0 | 3.1 | 6.9 | 11.2 | 17.6 | 28.7 |
| 1977 | 2.0 | * | * | * | * | * | 0.9 | 2.5 | 7.1 | 11.6 | 17.3 | 32.2 |
| 1976 | 2.1 | * | * | * | * | * | 1.2 | 4.0 | 8.4 | 11.0 | 16.0 | 29.4 |
| 1975 | 2.4 | * | * | * | * | * | 1.2 | 4.5 | 9.9 | 11.7 | 16.9 | 30.0 |
| 1974 | 2.4 | * | * | * | * | * | 1.6 | 4.2 | 9.2 | 13.5 | 15.8 | 28.9 |
| 1973 | 2.4 | * | * | * | * | * | 1.7 | 4.6 | 8.9 | 14.5 | 16.0 | 31.6 |
| 1972 | 2.5 | * | * | * | * | * | 1.4 | 5.8 | 8.9 | 11.6 | 21.4 | 30.9 |
| 1971 | 2.1 | * | * | * | * | * | 1.1 | 4.1 | 7.8 | 12.3 | 13.7 | 25.2 |
| 1970 | 2.5 | * | * | * | * | * | 2.1 | 5.3 | 9.1 | 12.4 | 20.8 | 23.9 |
| 1969 | 3.4 | * | * | * | * | * | 2.5 | 6.5 | 12.2 | 20.1 | 28.0 | 36.1 |
| 1968 | 3.2 | * | * | * | * | * | 2.2 | 6.0 | 12.2 | 19.4 | 23.5 | 50.0 |
| BLACK, MALE | | | | | | | | | | | | |
| 1978 | 2.7 | * | * | * | * | * | * | 5.0 | 11.2 | 16.3 | 24.3 | 43.1 |
| 1977 | 2.7 | * | * | * | * | * | * | 3.7 | 11.2 | 17.9 | 26.4 | 42.9 |
| 1976 | 2.7 | * | * | * | * | * | 2.0 | 6.5 | 12.6 | 14.9 | 19.2 | * |
| 1975 | 3.2 | * | * | * | * | * | 2.0 | 6.4 | 16.1 | 16.1 | 22.1 | * |
| 1974 | 3.1 | * | * | * | * | * | 2.4 | 6.4 | 13.6 | 17.5 | 21.5 | 48.8 |
| 1973 | 3.2 | * | * | * | * | * | 2.3 | 6.7 | 13.3 | 21.0 | 23.5 | 48.8 |
| 1972 | 3.2 | * | * | * | * | * | 1.8 | 8.7 | 12.7 | 15.8 | 31.5 | * |
| 1971 | 2.8 | * | * | * | * | * | 1.9 | 6.5 | 11.8 | 15.8 | 18.3 | * |
| 1970 | 3.0 | * | * | * | * | * | 3.2 | 7.7 | 12.0 | 14.5 | 23.7 | * |
| 1969 | 4.1 | * | * | * | * | * | 3.3 | 8.5 | 16.3 | 25.9 | 39.5 | * |
| 1968 | 4.1 | * | * | * | * | * | 3.6 | 7.6 | 17.9 | 25.6 | 30.3 | * |
| BLACK, FEMALE | | | | | | | | | | | | |
| 1978 | 1.4 | * | * | * | * | * | * | * | 3.4 | 7.4 | 13.1 | 21.2 |
| 1977 | 1.3 | * | * | * | * | * | * | * | 3.7 | 6.9 | 11.3 | 26.6 |
| 1976 | 1.6 | * | * | * | * | * | * | 1.9 | 4.9 | 8.0 | 13.8 | 30.3 |
| 1975 | 1.6 | * | * | * | * | * | * | 2.9 | 4.8 | 8.3 | 13.4 | 28.6 |
| 1974 | 1.7 | * | * | * | * | * | * | 2.4 | 5.5 | 10.5 | 12.0 | * |
| 1973 | 1.7 | * | * | * | * | * | * | 2.9 | 5.2 | 9.5 | 10.8 | * |
| 1972 | 1.8 | * | * | * | * | * | * | 3.4 | 5.7 | 8.3 | 14.3 | 28.6 |
| 1971 | 1.4 | * | * | * | * | * | * | 2.1 | 4.4 | 9.4 | 10.4 | * |
| 1970 | 2.0 | * | * | * | * | * | * | 3.2 | 6.7 | 10.7 | 18.7 | * |
| 1969 | 2.7 | * | * | * | * | * | 1.8 | 4.7 | 8.7 | 15.4 | 19.4 | 40.0 |
| 1968 | 2.4 | * | * | * | * | * | * | 4.6 | 7.4 | 14.4 | 18.5 | 48.2 |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| CHRONIC ISCHEMIC HEART DISEASE (412) | | | | | | | | | | | | |
| BOTH SEXES | | | | | | | | | | | | |
| 1978 | 151.0 | * | * | * | 0.1 | 1.5 | 13.2 | 59.6 | 174.6 | 478.5 | 1,493.2 | 4,689.8 |
| 1977 | 149.2 | * | * | * | 0.1 | 1.5 | 13.5 | 60.4 | 173.5 | 483.1 | 1,515.3 | 4,673.5 |
| 1976 | 148.2 | * | * | * | 0.1 | 1.5 | 12.8 | 57.8 | 170.5 | 485.9 | 1,537.6 | 4,819.0 |
| 1975 | 145.7 | * | * | * | 0.1 | 1.5 | 13.3 | 56.1 | 169.8 | 494.5 | 1,559.5 | 4,697.7 |
| 1974 | 152.8 | * | * | * | 0.2 | 1.6 | 13.5 | 57.0 | 179.4 | 530.4 | 1,674.9 | 5,151.1 |
| 1973 | 155.1 | * | * | 0.1 | 0.2 | 1.5 | 13.7 | 58.0 | 183.9 | 548.1 | 1,746.8 | 5,311.9 |
| 1972 | 153.8 | * | * | * | 0.2 | 1.4 | 14.2 | 57.5 | 184.4 | 557.1 | 1,766.7 | 5,270.7 |
| 1971 | 151.0 | * | * | 0.1 | 0.2 | 1.8 | 14.6 | 57.6 | 183.8 | 553.5 | 1,749.3 | 5,245.4 |
| 1970 | 150.1 | * | * | * | 0.2 | 1.8 | 14.3 | 58.0 | 186.4 | 568.7 | 1,792.8 | 4,800.4 |
| 1969 | 150.6 | * | * | * | 0.2 | 1.8 | 14.0 | 56.6 | 187.0 | 577.2 | 1,804.8 | 5,470.7 |
| 1968 | 150.6 | * | * | * | 0.1 | 1.6 | 13.6 | 57.0 | 190.6 | 590.4 | 1,826.0 | 5,523.6 |
| MALE | | | | | | | | | | | | |
| 1978 | 154.5 | * | * | * | 0.1 | 2.5 | 21.0 | 94.1 | 264.8 | 669.5 | 1,828.5 | 5,053.7 |
| 1977 | 153.3 | * | * | * | 0.1 | 2.5 | 21.2 | 94.9 | 260.2 | 672.5 | 1,858.7 | 5,045.1 |
| 1976 | 151.5 | * | * | * | 0.2 | 2.4 | 20.2 | 91.3 | 256.2 | 673.9 | 1,859.1 | 5,133.5 |
| 1975 | 149.6 | * | * | * | 0.2 | 2.3 | 20.7 | 88.3 | 252.5 | 677.5 | 1,880.9 | 5,016.7 |
| 1974 | 155.9 | * | * | * | 0.2 | 2.3 | 20.6 | 88.2 | 264.5 | 717.8 | 1,996.3 | 5,443.7 |
| 1973 | 158.7 | * | * | * | 0.2 | 2.1 | 20.7 | 89.3 | 270.4 | 737.2 | 2,076.8 | 5,588.8 |
| 1972 | 157.2 | * | * | * | 0.3 | 2.2 | 21.5 | 89.2 | 269.0 | 733.8 | 2,084.0 | 5,471.3 |
| 1971 | 155.3 | * | * | * | 0.3 | 2.5 | 21.9 | 87.7 | 268.0 | 730.2 | 2,039.1 | 5,493.9 |
| 1970 | 155.7 | * | * | * | 0.2 | 2.5 | 21.6 | 87.6 | 270.5 | 747.9 | 2,076.6 | 4,863.6 |
| 1969 | 156.3 | * | * | * | 0.3 | 2.6 | 21.3 | 86.4 | 271.1 | 756.1 | 2,076.4 | 5,599.1 |
| 1968 | 156.3 | * | * | * | 0.2 | 2.3 | 20.5 | 85.6 | 273.4 | 769.1 | 2,075.5 | 5,636.6 |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| CHRONIC ISCHEMIC HEART DISEASE (412) | | | | | | | | | | | | |
| FEMALE | | | | | | | | | | | | |
| 1978 | 147.6 | * | * | * | * | 0.6 | 5.7 | 27.4 | 95.0 | 332.5 | 1,293.5 | 4,525.4 |
| 1977 | 145.3 | * | * | * | * | 0.6 | 6.1 | 28.3 | 96.9 | 338.2 | 1,309.7 | 4,502.9 |
| 1976 | 145.0 | * | * | * | * | 0.7 | 5.6 | 26.6 | 94.7 | 341.9 | 1,343.4 | 4,671.2 |
| 1975 | 142.0 | * | * | * | * | 0.7 | 6.2 | 26.2 | 96.5 | 354.3 | 1,363.7 | 4,543.3 |
| 1974 | 149.9 | * | * | * | * | 0.9 | 6.8 | 28.0 | 103.9 | 386.8 | 1,477.0 | 5,006.1 |
| 1973 | 151.6 | * | * | * | 0.1 | 0.8 | 7.1 | 29.0 | 107.0 | 402.8 | 1,540.7 | 5,172.6 |
| 1972 | 150.6 | * | * | * | 0.2 | 0.7 | 7.3 | 28.1 | 109.1 | 420.8 | 1,564.3 | 5,168.0 |
| 1971 | 146.9 | * | * | * | 0.1 | 1.0 | 7.7 | 29.6 | 108.4 | 416.8 | 1,561.3 | 5,115.7 |
| 1970 | 144.7 | * | * | * | 0.1 | 1.1 | 7.4 | 30.4 | 111.0 | 429.4 | 1,605.1 | 4,765.0 |
| 1969 | 145.3 | * | * | * | 0.1 | 1.0 | 7.2 | 28.7 | 111.3 | 437.4 | 1,622.9 | 5,401.2 |
| 1968 | 145.1 | * | * | * | * | 1.1 | 7.1 | 30.2 | 115.7 | 449.7 | 1,655.3 | 5,468.4 |
| WHITE, BOTH SEXES | | | | | | | | | | | | |
| 1978 | 156.5 | * | * | * | 0.1 | 1.2 | 10.9 | 52.1 | 160.2 | 460.4 | 1,501.6 | 4,812.2 |
| 1977 | 154.1 | * | * | * | 0.1 | 1.2 | 10.9 | 52.5 | 158.5 | 461.7 | 1,526.2 | 4,789.7 |
| 1976 | 153.5 | * | * | * | 0.1 | 1.2 | 10.6 | 50.4 | 157.3 | 465.7 | 1,552.3 | 4,951.7 |
| 1975 | 150.5 | * | * | * | 0.1 | 1.1 | 11.0 | 48.7 | 156.0 | 471.0 | 1,575.1 | 4,831.3 |
| 1974 | 157.4 | * | * | * | 0.1 | 1.1 | 10.9 | 48.8 | 163.2 | 503.1 | 1,693.4 | 5,291.0 |
| 1973 | 159.3 | * | * | * | 0.1 | 1.1 | 10.8 | 49.6 | 167.5 | 517.8 | 1,763.5 | 5,456.3 |
| 1972 | 158.0 | * | * | * | 0.2 | 1.1 | 11.1 | 48.8 | 166.6 | 528.9 | 1,784.0 | 5,435.5 |
| 1971 | 154.8 | * | * | 0.1 | 0.2 | 1.3 | 11.4 | 49.1 | 166.9 | 525.1 | 1,761.7 | 5,407.2 |
| 1970 | 153.4 | * | * | * | 0.1 | 1.3 | 10.9 | 49.2 | 168.0 | 539.8 | 1,799.6 | 4,983.6 |
| 1969 | 153.8 | * | * | * | 0.2 | 1.2 | 10.8 | 47.8 | 167.8 | 548.5 | 1,811.4 | 5,655.1 |
| 1968 | 153.1 | * | * | * | 0.1 | 1.0 | 10.4 | 47.5 | 169.2 | 560.6 | 1,833.9 | 5,695.3 |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| CHRONIC ISCHEMIC HEART DISEASE (412) | | | | | | | | | | | | |
| WHITE, MALE | | | | | | | | | | | | |
| 1978 | 158.8 | * | * | * | 0.1 | 2.0 | 18.1 | 85.1 | 249.0 | 654.6 | 1,852.0 | 5,207.2 |
| 1977 | 157.2 | * | * | * | * | 2.0 | 18.0 | 85.8 | 243.9 | 655.3 | 1,887.8 | 5,193.9 |
| 1976 | 155.7 | * | * | * | 0.2 | 1.9 | 17.5 | 82.7 | 242.3 | 658.3 | 1,888.7 | 5,302.5 |
| 1975 | 153.4 | * | * | * | 0.1 | 1.8 | 17.9 | 79.5 | 239.4 | 659.4 | 1,905.3 | 5,184.1 |
| 1974 | 159.4 | * | * | * | 0.2 | 1.8 | 17.8 | 79.1 | 248.4 | 693.0 | 2,026.1 | 5,617.3 |
| 1973 | 162.1 | * | * | * | 0.2 | 1.7 | 17.4 | 80.0 | 254.3 | 712.5 | 2,103.5 | 5,774.3 |
| 1972 | 160.2 | * | * | * | 0.2 | 1.8 | 18.1 | 79.6 | 251.9 | 708.6 | 2,110.2 | 5,663.2 |
| 1971 | 158.3 | * | * | * | 0.2 | 1.9 | 18.5 | 78.8 | 251.5 | 706.7 | 2,060.2 | 5,699.8 |
| 1970 | 158.4 | * | * | * | 0.1 | 1.9 | 17.8 | 78.6 | 253.2 | 722.7 | 2,093.6 | 5,090.2 |
| 1969 | 158.7 | * | * | * | 0.2 | 1.8 | 17.6 | 76.7 | 253.2 | 731.2 | 2,089.3 | 5,830.4 |
| 1968 | 158.3 | * | * | * | * | 1.6 | 17.0 | 76.0 | 253.4 | 742.8 | 2,093.7 | 5,831.8 |
| WHITE, FEMALE | | | | | | | | | | | | |
| 1978 | 154.4 | * | * | * | * | 0.4 | 3.8 | 20.8 | 81.4 | 312.1 | 1,295.3 | 4,636.2 |
| 1977 | 151.1 | * | * | * | * | 0.4 | 3.9 | 21.1 | 82.5 | 313.9 | 1,312.9 | 4,606.7 |
| 1976 | 151.4 | * | * | * | * | 0.5 | 4.0 | 20.0 | 81.5 | 318.7 | 1,352.2 | 4,793.7 |
| 1975 | 147.7 | * | * | * | * | 0.4 | 4.2 | 19.7 | 81.7 | 327.3 | 1,376.6 | 4,667.6 |
| 1974 | 155.5 | * | * | * | * | 0.5 | 4.2 | 20.3 | 87.3 | 358.3 | 1,490.1 | 5,131.9 |
| 1973 | 156.6 | * | * | * | * | 0.6 | 4.3 | 21.2 | 90.1 | 369.0 | 1,552.9 | 5,304.2 |
| 1972 | 155.8 | * | * | * | 0.2 | 0.4 | 4.3 | 19.9 | 90.4 | 391.0 | 1,578.0 | 5,321.0 |
| 1971 | 151.5 | * | * | * | * | 0.7 | 4.6 | 21.4 | 91.1 | 385.1 | 1,570.5 | 5,257.3 |
| 1970 | 148.6 | * | * | * | * | 0.6 | 4.2 | 21.7 | 91.4 | 398.5 | 1,607.3 | 4,925.3 |
| 1969 | 149.1 | * | * | * | * | 0.5 | 4.2 | 20.6 | 90.9 | 406.3 | 1,626.5 | 5,569.5 |
| 1968 | 148.2 | * | * | * | * | 0.4 | 4.0 | 20.7 | 93.0 | 417.9 | 1,657.8 | 5,629.4 |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| CHRONIC ISCHEMIC HEART DISEASE (412) | | | | | | | | | | | | |
| ALL OTHER, BOTH SEXES | | | | | | | | | | | | |
| 1978 | 115.5 | * | * | * | * | 3.6 | 28.6 | 117.5 | 304.2 | 647.6 | 1,406.6 | 3,285.7 |
| 1977 | 117.7 | * | * | * | * | 3.5 | 31.0 | 122.5 | 310.7 | 686.0 | 1,402.6 | 3,343.1 |
| 1976 | 113.6 | * | * | * | * | 4.0 | 27.1 | 117.3 | 293.0 | 679.1 | 1,385.8 | 3,274.5 |
| 1975 | 113.9 | * | * | * | * | 4.0 | 29.7 | 117.1 | 297.6 | 722.3 | 1,392.3 | 3,164.4 |
| 1974 | 122.0 | * | * | * | * | 4.6 | 32.0 | 125.8 | 331.2 | 797.8 | 1,472.7 | 3,522.2 |
| 1973 | 126.1 | * | * | * | * | 4.0 | 34.9 | 128.9 | 339.0 | 847.2 | 1,554.9 | 3,629.9 |
| 1972 | 125.0 | * | * | * | 0.6 | 4.0 | 36.8 | 132.5 | 353.8 | 835.8 | 1,566.1 | 3,369.1 |
| 1971 | 124.5 | * | * | * | 0.5 | 5.0 | 38.1 | 131.0 | 344.8 | 834.7 | 1,599.8 | 3,402.5 |
| 1970 | 126.9 | * | * | * | 0.4 | 5.5 | 39.7 | 135.0 | 360.6 | 851.3 | 1,706.3 | 2,919.0 |
| 1969 | 128.1 | * | * | * | 0.5 | 6.2 | 38.7 | 135.8 | 371.4 | 863.4 | 1,724.0 | 3,391.7 |
| 1968 | 132.0 | * | * | * | * | 6.2 | 38.8 | 142.6 | 398.1 | 889.5 | 1,724.6 | 3,605.9 |
| ALL OTHER, MALE | | | | | | | | | | | | |
| 1978 | 126.5 | * | * | * | * | 5.5 | 42.2 | 169.1 | 412.6 | 808.2 | 1,600.8 | 3,481.0 |
| 1977 | 127.5 | * | * | * | * | 5.3 | 44.8 | 171.6 | 414.4 | 831.8 | 1,584.7 | 3,528.6 |
| 1976 | 123.6 | * | * | * | * | 5.9 | 40.1 | 165.5 | 388.2 | 819.4 | 1,586.8 | 3,440.0 |
| 1975 | 123.6 | * | * | * | * | 5.6 | 42.0 | 164.1 | 377.2 | 848.9 | 1,639.7 | 3,342.6 |
| 1974 | 131.6 | * | * | * | * | 6.1 | 41.9 | 168.2 | 419.0 | 955.6 | 1,689.1 | 3,656.0 |
| 1973 | 135.0 | * | * | * | * | 5.7 | 45.9 | 171.6 | 427.5 | 973.9 | 1,787.5 | 3,695.8 |
| 1972 | 135.4 | * | * | * | 1.1 | 5.5 | 47.3 | 174.8 | 434.8 | 975.9 | 1,801.8 | 3,531.9 |
| 1971 | 134.3 | * | * | * | * | 7.1 | 49.2 | 168.7 | 428.5 | 954.5 | 1,813.0 | 3,466.0 |
| 1970 | 136.6 | * | * | * | * | 7.3 | 52.1 | 169.4 | 435.8 | 985.5 | 1,878.8 | 2,872.5 |
| 1969 | 139.2 | * | * | * | * | 8.4 | 51.2 | 176.8 | 445.6 | 995.5 | 1,921.3 | 3,418.2 |
| 1968 | 141.6 | * | * | * | * | 7.2 | 49.8 | 175.8 | 468.6 | 1,025.0 | 1,858.1 | 3,736.6 |
| ALL OTHER, FEMALE | | | | | | | | | | | | |
| 1978 | 105.5 | * | * | * | * | 2.0 | 17.3 | 74.9 | 214.1 | 523.3 | 1,277.4 | 3,211.9 |
| 1977 | 108.8 | * | * | * | * | 1.9 | 19.4 | 81.8 | 224.3 | 571.4 | 1,274.9 | 3,243.3 |
| 1976 | 104.6 | * | * | * | * | 2.4 | 16.2 | 76.9 | 213.2 | 568.2 | 1,248.4 | 3,181.6 |
| 1975 | 105.1 | * | * | * | * | 2.7 | 19.3 | 77.5 | 230.4 | 621.7 | 1,219.2 | 3,026.9 |
| 1974 | 113.2 | * | * | * | * | 3.3 | 23.7 | 90.0 | 256.5 | 671.3 | 1,323.5 | 3,443.5 |
| 1973 | 117.9 | * | * | * | * | 2.6 | 25.8 | 92.4 | 263.7 | 744.9 | 1,395.9 | 3,589.9 |
| 1972 | 115.6 | * | * | * | * | 2.7 | 27.9 | 96.2 | 284.0 | 723.2 | 1,401.3 | 3,268.4 |
| 1971 | 115.6 | * | * | * | * | 3.1 | 28.9 | 98.2 | 271.7 | 736.8 | 1,442.1 | 3,361.6 |
| 1970 | 117.9 | * | * | * | * | 4.0 | 29.3 | 104.9 | 294.2 | 740.7 | 1,575.1 | 2,951.7 |
| 1969 | 118.0 | * | * | * | * | 4.3 | 28.2 | 99.9 | 305.7 | 754.7 | 1,566.8 | 3,373.8 |
| 1968 | 123.3 | * | * | * | * | 5.3 | 29.5 | 113.3 | 334.8 | 777.2 | 1,628.0 | 3,518.0 |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| CHRONIC ISCHEMIC HEART DISEASE (412) | | | | | | | | | | | | |
| BLACK, BOTH SEXES | | | | | | | | | | | | |
| 1978 | 126.9 | * | * | * | * | 4.1 | 33.0 | 132.0 | 331.7 | 690.8 | 1,445.6 | 3,398.0 |
| 1977 | 129.4 | * | * | * | * | 4.0 | 35.1 | 138.1 | 338.7 | 734.3 | 1,432.7 | 3,488.1 |
| 1976 | 124.0 | * | * | * | * | 4.6 | 30.7 | 130.3 | 317.4 | 722.5 | 1,423.0 | 3,380.1 |
| 1975 | 124.0 | * | * | * | * | 4.6 | 33.5 | 130.1 | 320.4 | 769.9 | 1,428.0 | 3,286.2 |
| 1974 | 131.8 | * | * | * | * | 5.1 | 36.1 | 138.3 | 353.1 | 847.4 | 1,506.6 | 3,670.2 |
| 1973 | 135.8 | * | * | * | * | 4.5 | 39.3 | 140.5 | 361.9 | 901.5 | 1,591.2 | 3,767.5 |
| 1972 | 134.5 | * | * | * | 0.7 | 4.6 | 41.1 | 145.5 | 377.7 | 887.3 | 1,616.8 | 3,505.5 |
| 1971 | 132.7 | * | * | * | 0.6 | 5.5 | 42.2 | 142.1 | 367.0 | 881.1 | 1,646.7 | 3,516.8 |
| 1970 | 136.5 | * | * | * | * | 6.2 | 44.4 | 147.2 | 384.4 | 905.2 | 1,809.6 | 3,083.0 |
| 1969 | 135.9 | * | * | * | 0.6 | 6.9 | 42.0 | 145.7 | 392.9 | 906.5 | 1,805.9 | 3,538.1 |
| 1968 | 139.7 | * | * | * | * | 6.9 | 42.0 | 151.9 | 421.7 | 933.1 | 1,794.8 | 3,720.7 |
| BLACK, MALE | | | | | | | | | | | | |
| 1978 | 137.3 | * | * | * | * | 6.2 | 48.5 | 188.6 | 449.8 | 868.1 | 1,628.5 | 3,503.9 |
| 1977 | 138.9 | * | * | * | * | 6.0 | 50.4 | 191.7 | 452.9 | 900.4 | 1,619.6 | 3,655.1 |
| 1976 | 133.4 | * | * | * | * | 6.6 | 45.4 | 183.4 | 422.0 | 875.9 | 1,617.1 | 3,529.8 |
| 1975 | 133.2 | * | * | * | * | 6.4 | 47.1 | 181.2 | 406.4 | 910.8 | 1,672.1 | 3,510.9 |
| 1974 | 140.8 | * | * | * | * | 6.6 | 47.1 | 184.6 | 447.7 | 1,024.4 | 1,724.4 | 3,832.6 |
| 1973 | 144.0 | * | * | * | * | 6.3 | 51.3 | 186.0 | 459.8 | 1,043.5 | 1,828.4 | 3,880.5 |
| 1972 | 144.6 | * | * | * | 1.2 | 6.4 | 52.7 | 191.9 | 469.3 | 1,044.3 | 1,846.7 | 3,765.0 |
| 1971 | 142.2 | * | * | * | * | 8.0 | 54.4 | 183.0 | 459.5 | 1,015.6 | 1,875.3 | 3,612.5 |
| 1970 | 145.8 | * | * | * | * | 8.2 | 58.0 | 185.0 | 467.4 | 1,053.5 | 2,004.4 | 3,084.9 |
| 1969 | 146.2 | * | * | * | * | 9.2 | 55.2 | 189.9 | 474.9 | 1,049.0 | 2,015.0 | 3,640.5 |
| 1968 | 148.4 | * | * | * | * | 8.1 | 53.6 | 186.8 | 501.1 | 1,075.7 | 1,941.2 | 3,962.9 |
| BLACK, FEMALE | | | | | | | | | | | | |
| 1978 | 117.5 | * | * | * | * | 2.3 | 20.1 | 85.3 | 235.2 | 558.4 | 1,323.5 | 3,343.4 |
| 1977 | 120.8 | * | * | * | * | 2.3 | 22.4 | 93.7 | 245.5 | 609.7 | 1,312.7 | 3,401.1 |
| 1976 | 115.5 | * | * | * | * | 2.8 | 18.5 | 86.2 | 231.0 | 605.7 | 1,292.8 | 3,301.1 |
| 1975 | 115.8 | * | * | * | * | 3.1 | 22.2 | 87.3 | 249.0 | 663.1 | 1,262.8 | 3,163.1 |
| 1974 | 123.6 | * | * | * | * | 3.8 | 27.0 | 99.4 | 274.6 | 713.0 | 1,359.2 | 3,580.8 |
| 1973 | 128.3 | * | * | * | * | 3.0 | 29.3 | 101.8 | 280.3 | 791.6 | 1,433.1 | 3,704.1 |
| 1972 | 125.3 | * | * | * | * | 3.0 | 31.5 | 106.0 | 300.8 | 766.4 | 1,449.3 | 3,357.1 |
| 1971 | 124.1 | * | * | * | * | 3.4 | 32.0 | 106.9 | 288.7 | 773.7 | 1,482.6 | 3,459.7 |
| 1970 | 128.1 | * | * | * | * | 4.5 | 33.1 | 114.6 | 313.6 | 787.8 | 1,666.7 | 3,081.8 |
| 1969 | 126.6 | * | * | * | * | 4.9 | 31.0 | 107.4 | 323.1 | 793.2 | 1,644.1 | 3,475.0 |
| 1968 | 131.8 | * | * | * | * | 5.9 | 32.2 | 121.5 | 353.0 | 816.7 | 1,686.0 | 3,635.7 |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| ANGINA PECTORIS (413) | | | | | | | | | | | | |
| BOTH SEXES | | | | | | | | | | | | |
| 1978 | 0.1 | * | * | * | * | * | * | * | 0.2 | 0.4 | 1.0 | 1.9 |
| 1977 | 0.1 | * | * | * | * | * | * | 0.1 | 0.2 | 0.4 | 0.8 | 1.8 |
| 1976 | 0.1 | * | * | * | * | * | * | 0.1 | * | 0.3 | 0.8 | 1.6 |
| 1975 | 0.1 | * | * | * | * | * | * | * | 0.1 | 0.3 | 0.5 | 1.8 |
| 1974 | 0.1 | * | * | * | * | * | * | * | 0.1 | 0.4 | 0.8 | 1.8 |
| 1973 | 0.1 | * | * | * | * | * | * | * | 0.1 | 0.4 | 0.7 | 1.9 |
| 1972 | 0.1 | * | * | * | * | * | * | * | * | 0.3 | 0.9 | 1.9 |
| 1971 | 0.1 | * | * | * | * | * | * | * | 0.2 | 0.4 | 0.9 | 2.4 |
| 1970 | 0.1 | * | * | * | * | * | * | * | 0.2 | 0.5 | 1.0 | 1.7 |
| 1969 | 0.1 | * | * | * | * | * | * | * | 0.2 | 0.5 | 0.9 | 2.4 |
| 1968 | 0.1 | * | * | * | * | * | * | * | 0.2 | 0.6 | 1.1 | 2.2 |
| MALE | | | | | | | | | | | | |
| 1978 | 0.1 | * | * | * | * | * | * | * | 0.3 | 0.6 | 1.3 | * |
| 1977 | 0.1 | * | * | * | * | * | * | * | 0.2 | 0.6 | 1.1 | * |
| 1976 | 0.1 | * | * | * | * | * | * | * | * | 0.5 | 1.0 | * |
| 1975 | 0.1 | * | * | * | * | * | * | * | * | 0.4 | * | * |
| 1974 | 0.1 | * | * | * | * | * | * | * | * | 0.6 | 0.9 | * |
| 1973 | 0.1 | * | * | * | * | * | * | * | * | 0.6 | 1.2 | * |
| 1972 | 0.1 | * | * | * | * | * | * | * | * | 0.5 | 1.2 | * |
| 1971 | 0.1 | * | * | * | * | * | * | * | 0.2 | 0.7 | 1.2 | * |
| 1970 | 0.1 | * | * | * | * | * | * | * | 0.3 | 0.7 | 1.2 | * |
| 1969 | 0.1 | * | * | * | * | * | * | * | 0.3 | 0.7 | 1.1 | * |
| 1968 | 0.1 | * | * | * | * | * | * | * | 0.3 | 0.8 | 1.3 | * |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| ANGINA PECTORIS (413) | | | | | | | | | | | | |
| FEMALE | | | | | | | | | | | | |
| 1978 | 0.1 | * | * | * | * | * | * | * | * | 0.3 | 0.9 | 1.7 |
| 1977 | 0.1 | * | * | * | * | * | * | * | * | 0.3 | 0.6 | 1.6 |
| 1976 | 0.1 | * | * | * | * | * | * | * | * | * | 0.7 | 1.6 |
| 1975 | 0.1 | * | * | * | * | * | * | * | * | * | 0.6 | 1.7 |
| 1974 | 0.1 | * | * | * | * | * | * | * | * | * | 0.7 | * |
| 1973 | 0.1 | * | * | * | * | * | * | * | * | * | * | * |
| 1972 | 0.1 | * | * | * | * | * | * | * | * | * | 0.6 | * |
| 1971 | 0.1 | * | * | * | * | * | * | * | * | * | 0.8 | 2.4 |
| 1970 | 0.1 | * | * | * | * | * | * | * | * | 0.4 | 0.9 | * |
| 1969 | 0.1 | * | * | * | * | * | * | * | * | 0.3 | 0.7 | 2.6 |
| 1968 | 0.1 | * | * | * | * | * | * | * | * | 0.4 | 0.9 | * |
| WHITE, BOTH SEXES | | | | | | | | | | | | |
| 1978 | 0.1 | * | * | * | * | * | * | * | 0.2 | 0.4 | 1.0 | 1.8 |
| 1977 | 0.1 | * | * | * | * | * | * | * | 0.1 | 0.4 | 0.8 | 2.0 |
| 1976 | 0.1 | * | * | * | * | * | * | * | * | 0.3 | 0.9 | 1.7 |
| 1975 | 0.1 | * | * | * | * | * | * | * | 0.1 | 0.3 | 0.6 | 1.9 |
| 1974 | 0.1 | * | * | * | * | * | * | * | 0.1 | 0.3 | 0.8 | 1.8 |
| 1973 | 0.1 | * | * | * | * | * | * | * | 0.1 | 0.4 | 0.8 | 1.9 |
| 1972 | 0.1 | * | * | * | * | * | * | * | * | 0.4 | 0.9 | 2.1 |
| 1971 | 0.1 | * | * | * | * | * | * | * | 0.2 | 0.4 | 1.0 | 2.6 |
| 1970 | 0.1 | * | * | * | * | * | * | * | 0.2 | 0.5 | 1.0 | 1.8 |
| 1969 | 0.1 | * | * | * | * | * | * | * | 0.2 | 0.5 | 0.9 | 2.5 |
| 1968 | 0.1 | * | * | * | * | * | * | * | 0.2 | 0.6 | 1.1 | 2.2 |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| ANGINA PECTORIS (413) | | | | | | | | | | | | |
| WHITE, MALE | | | | | | | | | | | | |
| 1978 | 0.1 | * | * | * | * | * | * | * | 0.3 | 0.5 | 1.3 | * |
| 1977 | 0.1 | * | * | * | * | * | * | * | 0.2 | 0.6 | 1.1 | * |
| 1976 | 0.1 | * | * | * | * | * | * | * | * | 0.5 | 1.1 | * |
| 1975 | 0.1 | * | * | * | * | * | * | * | * | 0.4 | * | * |
| 1974 | 0.1 | * | * | * | * | * | * | * | * | 0.6 | 1.0 | * |
| 1973 | 0.1 | * | * | * | * | * | * | * | * | 0.6 | 1.3 | * |
| 1972 | 0.1 | * | * | * | * | * | * | * | * | 0.6 | 1.3 | * |
| 1971 | 0.1 | * | * | * | * | * | * | * | 0.2 | 0.7 | 1.3 | * |
| 1970 | 0.1 | * | * | * | * | * | * | * | 0.3 | 0.8 | 1.1 | * |
| 1969 | 0.1 | * | * | * | * | * | * | * | 0.4 | 0.7 | 1.2 | * |
| 1968 | 0.1 | * | * | * | * | * | * | * | 0.3 | 0.8 | 1.3 | * |
| WHITE, FEMALE | | | | | | | | | | | | |
| 1978 | 0.1 | * | * | * | * | * | * | * | * | 0.3 | 0.9 | 1.6 |
| 1977 | 0.1 | * | * | * | * | * | * | * | * | 0.3 | 0.6 | 1.7 |
| 1976 | 0.1 | * | * | * | * | * | * | * | * | * | 0.8 | 1.7 |
| 1975 | 0.1 | * | * | * | * | * | * | * | * | * | 0.7 | 1.8 |
| 1974 | 0.1 | * | * | * | * | * | * | * | * | * | 0.7 | * |
| 1973 | 0.1 | * | * | * | * | * | * | * | * | * | * | * |
| 1972 | 0.1 | * | * | * | * | * | * | * | * | * | 0.7 | * |
| 1971 | 0.1 | * | * | * | * | * | * | * | * | * | 0.8 | 2.5 |
| 1970 | 0.1 | * | * | * | * | * | * | * | * | 0.4 | 0.9 | * |
| 1969 | 0.1 | * | * | * | * | * | * | * | * | * | 0.7 | 2.6 |
| 1968 | 0.1 | * | * | * | * | * | * | * | * | 0.4 | 0.9 | * |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| ANGINA PECTORIS (413) | | | | | | | | | | | | |
| ALL OTHER, BOTH SEXES | | | | | | | | | | | | |
| 1978 | 0.1 | * | * | * | * | * | * | * | * | * | * | * |
| 1977 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1976 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1975 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1974 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1973 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1972 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1971 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1970 | 0.1 | * | * | * | * | * | * | * | * | * | * | * |
| 1969 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1968 | 0.1 | * | * | * | * | * | * | * | * | * | * | * |
| ALL OTHER, MALE | | | | | | | | | | | | |
| 1978 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1977 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1976 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1975 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1974 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1973 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1972 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1971 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1970 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1969 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1968 | * | * | * | * | * | * | * | * | * | * | * | * |
| ALL OTHER, FEMALE | | | | | | | | | | | | |
| 1978 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1977 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1976 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1975 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1974 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1973 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1972 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1971 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1970 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1969 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1968 | * | * | * | * | * | * | * | * | * | * | * | * |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| ANGINA PECTORIS (413) | | | | | | | | | | | | |
| BLACK, BOTH SEXES | | | | | | | | | | | | |
| 1978 | 0.1 | * | * | * | * | * | * | * | * | * | * | * |
| 1977 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1976 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1975 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1974 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1973 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1972 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1971 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1970 | 0.1 | * | * | * | * | * | * | * | * | * | * | * |
| 1969 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1968 | 0.1 | * | * | * | * | * | * | * | * | * | * | * |
| BLACK, MALE | | | | | | | | | | | | |
| 1978 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1977 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1976 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1975 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1974 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1973 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1972 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1971 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1970 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1969 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1968 | * | * | * | * | * | * | * | * | * | * | * | * |
| BLACK, FEMALE | | | | | | | | | | | | |
| 1978 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1977 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1976 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1975 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1974 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1973 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1972 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1971 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1970 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1969 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1968 | * | * | * | * | * | * | * | * | * | * | * | * |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| CHRONIC DISEASE OF ENDOCARDIUM AND OTHER MYOCARDIAL INSUFFICIENCY (424,428) | | | | | | | | | | | | |
| BOTH SEXES | | | | | | | | | | | | |
| 1978 | 1.8 | 0.7 | 0.2 | 0.1 | 0.2 | 0.3 | 0.6 | 1.3 | 2.6 | 5.6 | 14.1 | 44.6 |
| 1977 | 1.8 | 1.0 | * | * | 0.2 | 0.3 | 0.6 | 1.1 | 2.6 | 5.5 | 14.8 | 48.6 |
| 1976 | 1.9 | 0.9 | * | 0.1 | 0.2 | 0.3 | 0.6 | 1.4 | 2.7 | 5.7 | 16.3 | 51.8 |
| 1975 | 2.0 | * | * | 0.1 | 0.2 | 0.3 | 0.6 | 1.3 | 2.7 | 6.2 | 17.9 | 59.9 |
| 1974 | 2.3 | 0.9 | 0.2 | 0.1 | 0.2 | 0.3 | 0.7 | 1.4 | 3.0 | 7.3 | 19.9 | 70.8 |
| 1973 | 2.4 | 1.1 | 0.3 | 0.1 | 0.2 | 0.4 | 0.7 | 1.6 | 3.4 | 7.0 | 21.8 | 76.4 |
| 1972 | 2.7 | 1.1 | 0.2 | 0.1 | 0.3 | 0.5 | 0.8 | 1.8 | 4.0 | 8.4 | 23.2 | 82.5 |
| 1971 | 3.0 | 1.1 | 0.2 | 0.1 | 0.2 | 0.4 | 0.9 | 2.0 | 4.0 | 9.4 | 27.6 | 95.8 |
| 1970 | 3.3 | 0.9 | 0.2 | 0.1 | 0.3 | 0.5 | 1.1 | 2.2 | 4.7 | 10.6 | 31.1 | 101.1 |
| 1969 | 3.7 | 1.0 | * | 0.1 | 0.2 | 0.4 | 1.2 | 2.3 | 5.5 | 12.0 | 35.7 | 132.8 |
| 1968 | 3.9 | 1.2 | 0.2 | 0.1 | 0.2 | 0.5 | 1.0 | 2.4 | 5.5 | 12.8 | 39.7 | 150.3 |
| MALE | | | | | | | | | | | | |
| 1978 | 1.9 | * | * | 0.1 | 0.2 | 0.4 | 0.7 | 1.7 | 3.5 | 7.2 | 16.2 | 47.9 |
| 1977 | 1.9 | * | * | * | 0.2 | 0.3 | 0.7 | 1.4 | 3.3 | 7.6 | 18.8 | 56.6 |
| 1976 | 1.9 | * | * | * | 0.3 | 0.4 | 0.7 | 1.7 | 3.4 | 7.2 | 19.3 | 55.0 |
| 1975 | 2.1 | * | * | * | 0.3 | 0.4 | 0.8 | 1.5 | 3.6 | 7.9 | 21.7 | 61.3 |
| 1974 | 2.3 | * | * | * | 0.3 | 0.4 | 0.9 | 1.8 | 4.0 | 9.5 | 22.8 | 74.7 |
| 1973 | 2.4 | * | 0.3 | 0.1 | 0.3 | 0.4 | 0.9 | 1.9 | 4.6 | 9.0 | 24.7 | 75.8 |
| 1972 | 2.7 | 1.2 | * | 0.1 | 0.3 | 0.6 | 1.0 | 2.3 | 5.0 | 10.5 | 27.6 | 80.8 |
| 1971 | 3.0 | * | * | * | 0.3 | 0.5 | 1.0 | 2.5 | 5.3 | 12.1 | 30.9 | 98.0 |
| 1970 | 3.4 | * | * | * | 0.3 | 0.6 | 1.4 | 2.8 | 6.1 | 13.3 | 35.0 | 103.6 |
| 1969 | 3.8 | * | * | * | 0.2 | 0.5 | 1.6 | 3.1 | 7.4 | 14.6 | 39.7 | 133.1 |
| 1968 | 4.0 | 1.2 | * | 0.1 | 0.3 | 0.6 | 1.1 | 2.9 | 7.6 | 16.4 | 44.4 | 146.4 |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|

CHRONIC DISEASE OF ENDOCARDIUM AND OTHER MYOCARDIAL INSUFFICIENCY (424,428)

FEMALE

| | | | | | | | | | | | | |
|------|-----|-----|-----|-----|-----|-----|-----|-----|-----|------|------|-------|
| 1978 | 1.8 | * | * | * | 0.1 | 0.2 | 0.5 | 0.9 | 1.8 | 4.4 | 12.8 | 43.1 |
| 1977 | 1.8 | * | * | * | 0.1 | 0.3 | 0.5 | 0.9 | 2.0 | 3.9 | 12.4 | 45.0 |
| 1976 | 1.9 | * | * | * | 0.1 | 0.2 | 0.4 | 1.0 | 2.0 | 4.5 | 14.6 | 50.3 |
| 1975 | 2.0 | * | * | * | 0.2 | 0.2 | 0.4 | 1.1 | 1.9 | 4.9 | 15.5 | 59.2 |
| 1974 | 2.3 | * | * | * | 0.1 | 0.2 | 0.5 | 1.1 | 2.1 | 5.6 | 18.1 | 68.8 |
| 1973 | 2.5 | * | * | * | 0.2 | 0.3 | 0.6 | 1.4 | 2.4 | 5.5 | 19.9 | 76.6 |
| 1972 | 2.7 | * | 0.3 | 0.1 | 0.2 | 0.5 | 0.7 | 1.4 | 3.0 | 6.7 | 20.3 | 83.3 |
| 1971 | 3.0 | 1.3 | * | 0.1 | 0.1 | 0.3 | 0.9 | 1.6 | 2.8 | 7.4 | 25.5 | 94.7 |
| 1970 | 3.2 | * | * | * | 0.2 | 0.4 | 0.8 | 1.6 | 3.4 | 8.5 | 28.6 | 99.7 |
| 1969 | 3.7 | * | * | * | 0.2 | 0.4 | 0.9 | 1.6 | 3.7 | 10.0 | 33.0 | 132.7 |
| 1968 | 3.9 | * | * | * | 0.2 | 0.4 | 0.8 | 2.0 | 3.6 | 9.9 | 36.4 | 152.6 |

WHITE, BOTH SEXES

| | | | | | | | | | | | | |
|------|-----|-----|-----|-----|-----|-----|-----|-----|-----|------|------|-------|
| 1978 | 1.8 | * | * | 0.1 | 0.2 | 0.2 | 0.4 | 1.2 | 2.4 | 5.4 | 14.4 | 44.7 |
| 1977 | 1.8 | 0.8 | * | * | 0.1 | 0.3 | 0.5 | 1.0 | 2.4 | 5.4 | 14.8 | 49.1 |
| 1976 | 1.9 | * | * | 0.1 | 0.1 | 0.2 | 0.4 | 1.2 | 2.6 | 5.6 | 16.3 | 52.4 |
| 1975 | 2.1 | * | * | * | 0.2 | 0.2 | 0.5 | 1.2 | 2.5 | 6.1 | 18.3 | 61.6 |
| 1974 | 2.3 | * | * | * | 0.2 | 0.3 | 0.6 | 1.4 | 2.8 | 7.0 | 19.8 | 72.2 |
| 1973 | 2.4 | * | 0.2 | 0.1 | 0.2 | 0.3 | 0.6 | 1.4 | 3.2 | 6.6 | 21.4 | 77.7 |
| 1972 | 2.7 | 0.8 | 0.2 | 0.1 | 0.2 | 0.4 | 0.7 | 1.6 | 3.6 | 8.0 | 23.6 | 84.3 |
| 1971 | 3.0 | 0.9 | * | 0.1 | 0.2 | 0.3 | 0.8 | 1.8 | 3.7 | 8.9 | 27.6 | 98.5 |
| 1970 | 3.3 | * | * | 0.1 | 0.2 | 0.3 | 0.9 | 2.0 | 4.4 | 9.9 | 30.8 | 103.3 |
| 1969 | 3.7 | 0.8 | * | * | 0.2 | 0.3 | 1.0 | 2.1 | 5.0 | 11.2 | 35.1 | 135.2 |
| 1968 | 3.8 | 0.8 | * | 0.1 | 0.2 | 0.3 | 0.7 | 2.1 | 4.8 | 11.4 | 38.5 | 149.7 |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| CHRONIC DISEASE OF ENDOCARDIUM AND OTHER MYOCARDIAL INSUFFICIENCY (424,428) | | | | | | | | | | | | |
| WHITE, MALE | | | | | | | | | | | | |
| 1978 | 1.8 | * | * | * | 0.2 | 0.3 | 0.5 | 1.5 | 3.3 | 7.0 | 16.4 | 47.6 |
| 1977 | 1.9 | * | * | * | 0.1 | 0.3 | 0.5 | 1.2 | 3.2 | 7.6 | 19.3 | 58.8 |
| 1976 | 1.9 | * | * | * | 0.2 | 0.3 | 0.5 | 1.5 | 3.3 | 7.0 | 18.9 | 55.0 |
| 1975 | 2.1 | * | * | * | 0.2 | 0.3 | 0.6 | 1.3 | 3.5 | 7.8 | 22.3 | 62.6 |
| 1974 | 2.3 | * | * | * | 0.2 | 0.3 | 0.8 | 1.7 | 3.6 | 9.2 | 23.0 | 76.9 |
| 1973 | 2.3 | * | * | * | 0.2 | 0.3 | 0.7 | 1.7 | 4.4 | 8.4 | 24.3 | 76.1 |
| 1972 | 2.7 | * | * | * | 0.3 | 0.5 | 0.9 | 1.9 | 4.7 | 10.0 | 28.0 | 82.9 |
| 1971 | 2.9 | * | * | * | 0.3 | 0.4 | 0.8 | 2.3 | 5.1 | 11.6 | 31.3 | 98.5 |
| 1970 | 3.3 | * | * | * | 0.3 | 0.4 | 1.2 | 2.5 | 5.8 | 12.5 | 34.6 | 108.0 |
| 1969 | 3.7 | * | * | * | 0.2 | 0.4 | 1.3 | 2.8 | 6.8 | 13.8 | 38.8 | 135.2 |
| 1968 | 3.8 | * | * | * | 0.2 | 0.4 | 0.9 | 2.4 | 6.7 | 14.7 | 43.4 | 146.9 |
| WHITE, FEMALE | | | | | | | | | | | | |
| 1978 | 1.9 | * | * | * | * | 0.2 | 0.4 | 0.8 | 1.6 | 4.3 | 13.1 | 43.4 |
| 1977 | 1.8 | * | * | * | * | 0.3 | 0.4 | 0.7 | 1.7 | 3.7 | 12.1 | 44.6 |
| 1976 | 2.0 | * | * | * | * | 0.2 | 0.4 | 0.9 | 2.0 | 4.5 | 14.7 | 51.3 |
| 1975 | 2.1 | * | * | * | 0.2 | 0.2 | 0.4 | 1.0 | 1.7 | 4.8 | 15.8 | 61.1 |
| 1974 | 2.3 | * | * | * | * | 0.2 | 0.4 | 1.0 | 2.0 | 5.2 | 17.9 | 70.0 |
| 1973 | 2.5 | * | * | * | 0.2 | 0.2 | 0.5 | 1.2 | 2.2 | 5.3 | 19.5 | 78.6 |
| 1972 | 2.7 | * | * | * | 0.1 | 0.4 | 0.5 | 1.3 | 2.5 | 6.4 | 20.8 | 85.0 |
| 1971 | 3.0 | * | * | * | 0.1 | 0.3 | 0.8 | 1.3 | 2.5 | 6.8 | 25.1 | 98.5 |
| 1970 | 3.2 | * | * | * | 0.1 | 0.3 | 0.6 | 1.5 | 3.1 | 7.9 | 28.2 | 100.7 |
| 1969 | 3.7 | * | * | * | 0.2 | 0.2 | 0.7 | 1.4 | 3.3 | 9.2 | 32.7 | 135.4 |
| 1968 | 3.8 | * | * | * | 0.2 | 0.3 | 0.5 | 1.8 | 3.0 | 8.9 | 35.2 | 151.4 |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| CHRONIC DISEASE OF ENDOCARDIUM AND OTHER MYOCARDIAL INSUFFICIENCY (424,428) | | | | | | | | | | | | |
| ALL OTHER, BOTH SEXES | | | | | | | | | | | | |
| 1978 | 1.8 | * | * | * | * | 0.7 | 1.5 | 2.4 | 4.5 | 7.1 | 11.6 | 43.5 |
| 1977 | 1.8 | * | * | * | * | 0.7 | 1.2 | 2.4 | 4.1 | 6.9 | 15.2 | 43.8 |
| 1976 | 1.8 | * | * | * | 0.4 | 0.8 | 1.4 | 2.9 | 3.5 | 7.0 | 16.7 | 44.4 |
| 1975 | 1.7 | * | * | * | 0.4 | 0.8 | 1.4 | 2.0 | 4.0 | 7.3 | 13.6 | 40.4 |
| 1974 | 2.1 | * | * | * | * | 0.8 | 1.2 | 2.2 | 5.3 | 10.5 | 20.5 | 53.3 |
| 1973 | 2.5 | * | * | * | 0.4 | 1.0 | 1.5 | 3.5 | 5.1 | 11.0 | 26.2 | 60.6 |
| 1972 | 2.7 | * | * | * | 0.5 | 1.2 | 1.8 | 4.0 | 7.7 | 12.6 | 17.7 | 61.8 |
| 1971 | 2.9 | * | * | * | 0.6 | 1.1 | 1.9 | 3.8 | 6.8 | 14.8 | 28.3 | 65.8 |
| 1970 | 3.5 | * | * | * | 0.5 | 1.8 | 2.4 | 4.5 | 8.0 | 17.2 | 36.0 | 79.1 |
| 1969 | 4.0 | * | * | * | * | 1.3 | 3.2 | 4.8 | 10.3 | 19.9 | 43.5 | 105.5 |
| 1968 | 4.8 | * | * | * | 0.5 | 1.3 | 2.8 | 5.0 | 12.4 | 26.3 | 54.7 | 156.9 |
| ALL OTHER, MALE | | | | | | | | | | | | |
| 1978 | 2.0 | * | * | * | * | 1.0 | 2.1 | 3.3 | 5.7 | 9.3 | 14.3 | 50.0 |
| 1977 | 1.7 | * | * | * | * | * | 1.7 | 2.9 | 4.2 | 7.8 | 14.6 | * |
| 1976 | 2.2 | * | * | * | * | * | 2.2 | 3.8 | 5.0 | 9.6 | 22.6 | 54.5 |
| 1975 | 2.0 | * | * | * | * | 1.2 | 2.4 | 2.6 | 5.1 | 8.7 | 15.3 | 48.1 |
| 1974 | 2.3 | * | * | * | * | * | 1.5 | 2.6 | 7.3 | 12.0 | 20.5 | 52.0 |
| 1973 | 2.7 | * | * | * | * | * | 1.8 | 4.0 | 6.0 | 14.7 | 28.6 | 72.9 |
| 1972 | 2.9 | * | * | * | * | 1.3 | 1.9 | 6.1 | 7.8 | 15.4 | 23.0 | 59.6 |
| 1971 | 3.1 | * | * | * | 0.9 | 1.5 | 2.1 | 4.3 | 8.0 | 16.8 | 26.1 | 93.6 |
| 1970 | 3.8 | * | * | * | * | 2.4 | 3.0 | 6.0 | 9.6 | 20.5 | 38.8 | 65.0 |
| 1969 | 4.4 | * | * | * | * | 1.5 | 4.0 | 6.0 | 13.3 | 21.9 | 51.1 | 113.6 |
| 1968 | 5.3 | * | * | * | * | 1.6 | 3.2 | 6.9 | 15.9 | 33.1 | 56.5 | 141.5 |
| ALL OTHER, FEMALE | | | | | | | | | | | | |
| 1978 | 1.5 | * | * | * | * | * | * | 1.7 | 3.5 | 5.4 | 9.8 | 40.4 |
| 1977 | 1.8 | * | * | * | * | * | * | 2.0 | 4.0 | 6.1 | 15.7 | 49.0 |
| 1976 | 1.5 | * | * | * | * | * | * | 2.2 | 2.2 | 4.9 | 12.8 | 38.8 |
| 1975 | 1.4 | * | * | * | * | * | * | 1.4 | 3.0 | 6.2 | 12.4 | 35.5 |
| 1974 | 2.0 | * | * | * | * | * | * | 2.0 | 3.6 | 9.3 | 20.5 | 54.1 |
| 1973 | 2.3 | * | * | * | * | 1.0 | * | 3.0 | 4.4 | 8.0 | 24.5 | 53.2 |
| 1972 | 2.4 | * | * | * | * | 1.1 | 1.7 | 2.1 | 7.5 | 10.4 | 14.0 | 63.2 |
| 1971 | 2.8 | * | * | * | * | * | 1.7 | 3.5 | 5.7 | 13.2 | 30.0 | 47.9 |
| 1970 | 3.3 | * | * | * | * | 1.2 | 1.9 | 3.2 | 6.7 | 14.5 | 33.9 | 89.0 |
| 1969 | 3.6 | * | * | * | * | 1.2 | 2.5 | 3.7 | 7.6 | 18.2 | 37.7 | 100.0 |
| 1968 | 4.4 | * | * | * | * | * | 2.5 | 3.3 | 9.2 | 20.7 | 53.6 | 167.2 |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| CHRONIC DISEASE OF ENDOCARDIUM AND OTHER MYOCARDIAL INSUFFICIENCY (424,428) | | | | | | | | | | | | |
| BLACK, BOTH SEXES | | | | | | | | | | | | |
| 1978 | 1.9 | * | * | * | * | 0.7 | 1.7 | 2.6 | 5.0 | 7.6 | 12.1 | 44.7 |
| 1977 | 2.0 | * | * | * | * | 0.8 | 1.2 | 2.6 | 4.5 | 7.4 | 15.9 | 49.0 |
| 1976 | 2.0 | * | * | * | 0.4 | 0.9 | 1.5 | 3.1 | 3.8 | 7.6 | 17.7 | 47.1 |
| 1975 | 1.8 | * | * | * | * | 1.0 | 1.6 | 2.2 | 4.4 | 7.8 | 14.4 | 43.1 |
| 1974 | 2.3 | * | * | * | * | 1.0 | 1.3 | 2.5 | 5.5 | 11.0 | 21.4 | 57.0 |
| 1973 | 2.7 | * | * | * | 0.5 | 1.0 | 1.7 | 3.8 | 5.3 | 11.7 | 26.5 | 62.3 |
| 1972 | 2.9 | * | * | * | 0.6 | 1.2 | 2.0 | 4.4 | 8.2 | 13.2 | 18.5 | 69.1 |
| 1971 | 3.1 | * | * | * | 0.6 | 1.2 | 2.0 | 4.1 | 7.2 | 16.0 | 30.6 | 68.2 |
| 1970 | 3.9 | * | * | * | 0.6 | 1.9 | 2.7 | 5.0 | 8.3 | 18.8 | 39.4 | 88.6 |
| 1969 | 4.3 | * | * | * | * | 1.5 | 3.6 | 5.2 | 11.1 | 21.6 | 46.3 | 113.4 |
| 1968 | 5.2 | * | * | * | * | 1.4 | 3.0 | 5.3 | 13.3 | 28.2 | 58.1 | 167.4 |
| BLACK, MALE | | | | | | | | | | | | |
| 1978 | 2.2 | * | * | * | * | 1.1 | 2.4 | 3.5 | 6.4 | 9.8 | 15.1 | 54.9 |
| 1977 | 1.9 | * | * | * | * | * | 1.7 | 3.2 | 4.8 | 8.4 | 14.5 | * |
| 1976 | 2.4 | * | * | * | * | * | 2.5 | 3.9 | 5.5 | 11.0 | 24.4 | 57.4 |
| 1975 | 2.2 | * | * | * | * | 1.4 | 2.5 | 2.8 | 5.7 | 9.4 | 16.7 | 52.2 |
| 1974 | 2.4 | * | * | * | * | * | * | 2.9 | 7.5 | 12.6 | 21.1 | 53.5 |
| 1973 | 2.9 | * | * | * | * | * | 2.1 | 4.3 | 6.0 | 16.0 | 28.9 | 73.2 |
| 1972 | 3.2 | * | * | * | * | * | 2.2 | 6.7 | 8.7 | 16.6 | 25.4 | 70.0 |
| 1971 | 3.3 | * | * | * | 0.9 | 1.6 | 2.2 | 4.6 | 8.6 | 18.0 | 29.0 | 100.0 |
| 1970 | 4.2 | * | * | * | * | 2.7 | 3.3 | 6.7 | 10.0 | 22.6 | 42.7 | 78.3 |
| 1969 | 4.8 | * | * | * | * | * | 4.4 | 6.6 | 14.7 | 24.2 | 54.5 | 124.3 |
| 1968 | 5.7 | * | * | * | * | * | 3.5 | 7.4 | 17.5 | 36.0 | 60.0 | 160.0 |
| BLACK, FEMALE | | | | | | | | | | | | |
| 1978 | 1.7 | * | * | * | * | * | * | 1.9 | 3.8 | 5.9 | 10.1 | 39.4 |
| 1977 | 2.0 | * | * | * | * | * | * | 2.0 | 4.3 | 6.6 | 16.9 | 54.3 |
| 1976 | 1.6 | * | * | * | * | * | * | 2.4 | 2.5 | 5.1 | 13.2 | 41.6 |
| 1975 | 1.6 | * | * | * | * | * | * | 1.6 | 3.3 | 6.6 | 12.8 | 38.1 |
| 1974 | 2.2 | * | * | * | * | * | * | 2.1 | 3.8 | 9.8 | 21.7 | 59.0 |
| 1973 | 2.5 | * | * | * | * | * | * | 3.3 | 4.7 | 8.4 | 25.0 | 56.2 |
| 1972 | 2.6 | * | * | * | * | * | 1.8 | 2.4 | 7.7 | 10.6 | 13.6 | 68.6 |
| 1971 | 3.0 | * | * | * | * | * | 1.8 | 3.7 | 6.1 | 14.3 | 31.7 | 49.3 |
| 1970 | 3.6 | * | * | * | * | * | 2.1 | 3.5 | 6.9 | 15.8 | 37.0 | 95.3 |
| 1969 | 3.9 | * | * | * | * | 1.4 | 2.8 | 4.0 | 8.1 | 19.6 | 40.1 | 106.7 |
| 1968 | 4.7 | * | * | * | * | * | 2.7 | 3.5 | 9.8 | 21.8 | 56.8 | 175.0 |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|--|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| ALL OTHER FORMS OF HEART DISEASE (420-423,425-427,429) | | | | | | | | | | | | |
| BOTH SEXES | | | | | | | | | | | | |
| 1978 | 26.8 | 24.1 | 1.9 | 0.7 | 1.8 | 2.8 | 7.3 | 19.3 | 43.7 | 93.5 | 208.3 | 546.1 |
| 1977 | 24.2 | 20.7 | 1.7 | 0.6 | 1.6 | 2.8 | 7.0 | 18.3 | 41.0 | 85.2 | 191.5 | 483.6 |
| 1976 | 22.9 | 21.2 | 1.5 | 0.6 | 1.7 | 2.7 | 6.7 | 17.8 | 39.5 | 81.0 | 182.9 | 461.3 |
| 1975 | 20.9 | 18.8 | 1.6 | 0.6 | 1.5 | 2.7 | 6.6 | 16.8 | 37.2 | 75.8 | 166.0 | 416.8 |
| 1974 | 20.2 | 20.0 | 1.6 | 0.6 | 1.6 | 2.9 | 6.9 | 16.8 | 36.8 | 73.0 | 160.6 | 400.5 |
| 1973 | 19.6 | 20.4 | 1.6 | 0.6 | 1.5 | 3.0 | 7.0 | 16.4 | 36.1 | 70.8 | 158.0 | 397.8 |
| 1972 | 18.2 | 18.5 | 1.7 | 0.7 | 1.4 | 2.8 | 6.5 | 15.8 | 34.6 | 68.9 | 144.9 | 354.9 |
| 1971 | 16.6 | 13.2 | 1.3 | 0.5 | 1.5 | 2.4 | 6.3 | 14.2 | 31.7 | 60.7 | 135.3 | 340.1 |
| 1970 | 15.9 | 11.6 | 1.4 | 0.5 | 1.4 | 2.5 | 6.4 | 14.5 | 30.6 | 60.1 | 129.2 | 294.5 |
| 1969 | 15.0 | 9.4 | 1.2 | 0.6 | 1.4 | 2.5 | 6.1 | 13.9 | 28.2 | 57.4 | 123.0 | 313.5 |
| 1968 | 14.1 | 9.9 | 1.2 | 0.6 | 1.3 | 2.3 | 5.7 | 13.2 | 27.2 | 52.3 | 116.8 | 300.8 |
| MALE | | | | | | | | | | | | |
| 1978 | 30.0 | 25.9 | 1.8 | 0.7 | 2.1 | 3.4 | 9.7 | 27.2 | 61.6 | 131.5 | 274.2 | 651.5 |
| 1977 | 27.5 | 23.9 | 1.7 | 0.6 | 1.9 | 3.6 | 9.2 | 25.2 | 58.3 | 120.5 | 253.3 | 583.1 |
| 1976 | 26.1 | 23.7 | 1.8 | 0.6 | 2.2 | 3.2 | 8.6 | 24.8 | 56.6 | 115.0 | 241.3 | 537.1 |
| 1975 | 24.1 | 20.3 | 1.7 | 0.7 | 1.8 | 3.3 | 8.4 | 22.9 | 52.7 | 107.2 | 221.9 | 499.5 |
| 1974 | 23.3 | 21.9 | 1.6 | 0.6 | 2.0 | 3.5 | 8.9 | 23.3 | 52.9 | 102.5 | 211.7 | 474.5 |
| 1973 | 22.7 | 22.9 | 1.8 | 0.6 | 1.8 | 3.5 | 8.8 | 23.1 | 51.1 | 101.0 | 205.0 | 471.4 |
| 1972 | 21.1 | 19.1 | 1.9 | 0.7 | 1.8 | 3.2 | 8.3 | 21.6 | 49.0 | 96.3 | 185.6 | 416.5 |
| 1971 | 19.3 | 13.9 | 1.3 | 0.6 | 1.8 | 2.7 | 8.2 | 19.4 | 44.5 | 85.8 | 173.4 | 407.6 |
| 1970 | 18.5 | 13.7 | 1.5 | 0.5 | 1.7 | 3.0 | 8.3 | 19.6 | 43.3 | 84.5 | 161.6 | 324.5 |
| 1969 | 17.5 | 9.2 | 1.2 | 0.5 | 1.6 | 3.1 | 8.1 | 18.9 | 39.2 | 78.5 | 156.8 | 364.5 |
| 1968 | 16.3 | 10.5 | 1.3 | 0.6 | 1.4 | 2.7 | 7.0 | 18.0 | 37.6 | 71.8 | 145.4 | 344.5 |

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|

ALL OTHER FORMS OF HEART DISEASE (420-423,425-427,429)

FEMALE

| | | | | | | | | | | | | |
|------|------|------|-----|-----|-----|-----|-----|------|------|------|-------|-------|
| 1978 | 23.7 | 22.2 | 2.0 | 0.8 | 1.5 | 2.3 | 5.0 | 12.0 | 27.8 | 64.4 | 169.1 | 498.5 |
| 1977 | 21.1 | 17.3 | 1.6 | 0.7 | 1.2 | 2.1 | 4.9 | 11.9 | 25.7 | 58.1 | 154.5 | 437.9 |
| 1976 | 19.8 | 18.5 | 1.3 | 0.7 | 1.3 | 2.1 | 4.8 | 11.3 | 24.3 | 54.9 | 147.6 | 425.7 |
| 1975 | 18.0 | 17.1 | 1.5 | 0.6 | 1.3 | 2.0 | 4.8 | 11.1 | 23.4 | 51.8 | 132.0 | 376.8 |
| 1974 | 17.2 | 17.9 | 1.7 | 0.6 | 1.3 | 2.3 | 5.0 | 10.8 | 22.5 | 50.4 | 129.1 | 363.8 |
| 1973 | 16.6 | 17.8 | 1.5 | 0.6 | 1.2 | 2.5 | 5.2 | 10.1 | 22.8 | 47.7 | 128.6 | 360.7 |
| 1972 | 15.5 | 17.9 | 1.4 | 0.6 | 1.1 | 2.3 | 4.7 | 10.5 | 21.9 | 47.8 | 118.9 | 323.3 |
| 1971 | 14.0 | 12.5 | 1.4 | 0.5 | 1.2 | 2.1 | 4.5 | 9.3 | 20.3 | 41.2 | 110.6 | 304.9 |
| 1970 | 13.4 | 9.4 | 1.3 | 0.5 | 1.1 | 2.1 | 4.7 | 9.7 | 19.1 | 41.2 | 107.7 | 277.7 |
| 1969 | 12.7 | 9.6 | 1.1 | 0.6 | 1.2 | 2.0 | 4.3 | 9.2 | 18.3 | 40.9 | 100.3 | 285.8 |
| 1968 | 11.9 | 9.3 | 1.2 | 0.6 | 1.1 | 1.9 | 4.5 | 8.6 | 17.7 | 36.8 | 97.2 | 277.1 |

WHITE, BOTH SEXES

| | | | | | | | | | | | | |
|------|------|------|-----|-----|-----|-----|-----|------|------|------|-------|-------|
| 1978 | 25.9 | 20.1 | 1.5 | 0.7 | 1.5 | 2.2 | 5.6 | 16.3 | 38.8 | 86.4 | 200.4 | 542.9 |
| 1977 | 23.4 | 18.0 | 1.4 | 0.6 | 1.3 | 2.2 | 5.4 | 15.5 | 36.5 | 78.5 | 184.2 | 478.5 |
| 1976 | 21.9 | 17.4 | 1.3 | 0.6 | 1.4 | 2.0 | 5.0 | 14.9 | 34.8 | 73.2 | 176.2 | 455.7 |
| 1975 | 19.9 | 16.3 | 1.3 | 0.6 | 1.3 | 2.0 | 4.8 | 14.2 | 32.6 | 68.0 | 160.4 | 412.1 |
| 1974 | 19.1 | 15.2 | 1.3 | 0.6 | 1.4 | 2.0 | 5.2 | 14.1 | 32.0 | 66.1 | 156.5 | 393.6 |
| 1973 | 18.4 | 14.9 | 1.3 | 0.6 | 1.3 | 2.1 | 5.1 | 13.4 | 31.4 | 63.4 | 152.1 | 393.2 |
| 1972 | 17.0 | 13.5 | 1.4 | 0.6 | 1.1 | 2.0 | 4.6 | 12.6 | 29.8 | 61.4 | 140.1 | 352.5 |
| 1971 | 15.4 | 9.3 | 1.2 | 0.5 | 1.2 | 1.6 | 4.7 | 11.7 | 27.4 | 53.5 | 129.7 | 338.8 |
| 1970 | 14.8 | 8.4 | 1.1 | 0.4 | 1.2 | 1.8 | 4.6 | 11.7 | 26.0 | 53.5 | 124.7 | 296.0 |
| 1969 | 13.9 | 8.0 | 1.0 | 0.5 | 1.2 | 1.8 | 4.5 | 11.0 | 24.2 | 50.8 | 118.5 | 311.3 |
| 1968 | 13.0 | 7.8 | 1.0 | 0.5 | 1.1 | 1.6 | 4.2 | 10.7 | 22.8 | 45.9 | 112.0 | 302.2 |

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|--|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| ALL OTHER FORMS OF HEART DISEASE (420-423,425-427,429) | | | | | | | | | | | | |
| WHITE, MALE | | | | | | | | | | | | |
| 1978 | 29.0 | 22.3 | 1.5 | 0.7 | 1.7 | 2.7 | 7.5 | 23.4 | 56.3 | 124.4 | 264.9 | 648.7 |
| 1977 | 26.5 | 21.6 | 1.5 | 0.6 | 1.6 | 2.8 | 7.2 | 21.7 | 52.7 | 113.7 | 245.7 | 578.8 |
| 1976 | 24.9 | 20.3 | 1.5 | 0.5 | 1.8 | 2.4 | 6.6 | 21.4 | 51.6 | 106.1 | 233.8 | 530.9 |
| 1975 | 22.8 | 17.7 | 1.4 | 0.6 | 1.6 | 2.3 | 6.2 | 19.6 | 47.5 | 98.5 | 214.4 | 490.2 |
| 1974 | 22.2 | 17.5 | 1.3 | 0.6 | 1.7 | 2.5 | 6.9 | 19.9 | 47.5 | 94.7 | 207.4 | 471.5 |
| 1973 | 21.2 | 16.5 | 1.4 | 0.5 | 1.5 | 2.5 | 6.5 | 19.3 | 45.9 | 92.5 | 196.5 | 464.7 |
| 1972 | 19.6 | 13.4 | 1.6 | 0.7 | 1.4 | 2.4 | 6.1 | 17.9 | 43.2 | 86.9 | 179.5 | 411.4 |
| 1971 | 18.0 | 10.2 | 1.1 | 0.5 | 1.5 | 2.0 | 6.3 | 16.5 | 39.4 | 76.7 | 167.4 | 407.1 |
| 1970 | 17.2 | 10.7 | 1.3 | 0.5 | 1.5 | 2.2 | 5.9 | 16.5 | 38.0 | 77.0 | 155.8 | 326.5 |
| 1969 | 16.2 | 8.0 | 1.0 | 0.5 | 1.4 | 2.1 | 6.2 | 15.6 | 34.7 | 70.2 | 150.3 | 360.5 |
| 1968 | 15.1 | 8.7 | 1.0 | 0.5 | 1.3 | 1.9 | 5.3 | 15.2 | 32.5 | 64.5 | 139.6 | 347.4 |
| WHITE, FEMALE | | | | | | | | | | | | |
| 1978 | 23.0 | 17.7 | 1.6 | 0.7 | 1.2 | 1.7 | 3.8 | 9.6 | 23.3 | 57.3 | 162.5 | 495.7 |
| 1977 | 20.4 | 14.2 | 1.3 | 0.6 | 1.0 | 1.7 | 3.7 | 9.6 | 22.1 | 51.6 | 147.9 | 433.1 |
| 1976 | 18.9 | 14.4 | 1.1 | 0.6 | 1.0 | 1.5 | 3.5 | 8.9 | 19.8 | 48.0 | 142.0 | 421.3 |
| 1975 | 17.2 | 14.8 | 1.2 | 0.5 | 1.1 | 1.7 | 3.4 | 9.1 | 19.4 | 44.7 | 127.9 | 375.2 |
| 1974 | 16.2 | 12.9 | 1.4 | 0.6 | 1.0 | 1.6 | 3.6 | 8.6 | 18.1 | 44.3 | 125.4 | 355.6 |
| 1973 | 15.6 | 13.2 | 1.2 | 0.6 | 1.0 | 1.8 | 3.7 | 7.9 | 18.4 | 41.2 | 124.6 | 358.2 |
| 1972 | 14.5 | 13.5 | 1.2 | 0.5 | 0.8 | 1.6 | 3.2 | 7.6 | 17.9 | 41.9 | 115.3 | 322.9 |
| 1971 | 13.0 | 8.4 | 1.2 | 0.5 | 0.9 | 1.3 | 3.1 | 7.2 | 16.7 | 35.7 | 105.6 | 303.8 |
| 1970 | 12.4 | 6.1 | 1.0 | 0.4 | 0.8 | 1.5 | 3.3 | 7.2 | 15.3 | 35.3 | 104.4 | 279.3 |
| 1969 | 11.8 | 8.1 | 1.0 | 0.6 | 1.0 | 1.4 | 2.7 | 6.8 | 14.7 | 35.7 | 97.3 | 285.7 |
| 1968 | 10.9 | 6.9 | 1.0 | 0.5 | 0.9 | 1.3 | 3.0 | 6.4 | 14.1 | 31.4 | 93.4 | 278.2 |

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|--|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| ALL OTHER FORMS OF HEART DISEASE (420-423,425-427,429) | | | | | | | | | | | | |
| ALL OTHER, BOTH SEXES | | | | | | | | | | | | |
| 1978 | 32.2 | 43.3 | 3.7 | 1.0 | 3.4 | 7.0 | 18.7 | 42.3 | 87.6 | 159.7 | 290.2 | 583.3 |
| 1977 | 29.7 | 33.9 | 2.8 | 1.0 | 3.1 | 6.8 | 17.7 | 40.5 | 82.3 | 148.6 | 266.7 | 541.9 |
| 1976 | 29.7 | 39.6 | 2.6 | 1.0 | 3.5 | 7.1 | 17.6 | 40.5 | 82.9 | 155.8 | 251.7 | 522.9 |
| 1975 | 27.7 | 31.2 | 2.7 | 0.9 | 2.9 | 7.1 | 18.7 | 38.0 | 79.2 | 152.2 | 226.6 | 471.2 |
| 1974 | 27.3 | 43.5 | 3.0 | 0.8 | 3.0 | 9.0 | 18.6 | 39.9 | 81.9 | 140.9 | 206.4 | 480.7 |
| 1973 | 27.8 | 47.7 | 3.2 | 0.9 | 3.0 | 8.9 | 20.1 | 41.2 | 80.5 | 144.1 | 223.9 | 450.4 |
| 1972 | 27.0 | 44.4 | 3.1 | 1.0 | 3.7 | 8.4 | 19.6 | 43.6 | 80.6 | 142.7 | 200.0 | 382.1 |
| 1971 | 24.5 | 34.2 | 2.3 | 0.9 | 3.1 | 7.9 | 18.1 | 35.6 | 72.9 | 131.5 | 203.3 | 355.8 |
| 1970 | 23.8 | 28.4 | 2.7 | 0.7 | 2.9 | 7.3 | 20.3 | 38.4 | 73.5 | 125.2 | 185.6 | 279.7 |
| 1969 | 22.9 | 17.3 | 1.8 | 0.7 | 2.9 | 7.8 | 18.8 | 39.1 | 67.2 | 123.5 | 181.8 | 333.9 |
| 1968 | 21.9 | 21.5 | 2.4 | 0.7 | 2.6 | 7.2 | 17.9 | 35.7 | 68.9 | 115.7 | 177.9 | 285.3 |
| ALL OTHER, MALE | | | | | | | | | | | | |
| 1978 | 36.7 | 42.9 | 3.4 | 1.0 | 4.0 | 8.6 | 26.2 | 58.4 | 111.7 | 197.8 | 360.9 | 681.0 |
| 1977 | 34.4 | 35.6 | 3.0 | 0.8 | 3.9 | 9.3 | 23.8 | 54.3 | 111.7 | 184.3 | 324.5 | 626.8 |
| 1976 | 34.2 | 40.6 | 3.2 | 1.1 | 4.3 | 9.0 | 23.1 | 53.9 | 105.0 | 197.8 | 314.0 | 600.0 |
| 1975 | 32.8 | 33.8 | 3.0 | 0.8 | 3.3 | 10.8 | 25.0 | 51.4 | 102.0 | 189.9 | 296.3 | 592.6 |
| 1974 | 31.5 | 44.4 | 2.9 | 0.8 | 3.5 | 11.5 | 24.0 | 53.4 | 104.2 | 177.6 | 255.9 | 506.0 |
| 1973 | 33.0 | 54.9 | 3.9 | 1.2 | 3.3 | 11.3 | 26.3 | 56.4 | 100.6 | 182.0 | 293.8 | 539.6 |
| 1972 | 32.2 | 48.4 | 3.8 | 0.9 | 4.6 | 9.7 | 26.0 | 54.6 | 105.4 | 186.4 | 250.7 | 468.1 |
| 1971 | 28.7 | 34.1 | 2.5 | 1.0 | 3.4 | 8.1 | 23.2 | 45.6 | 93.8 | 173.3 | 240.6 | 412.8 |
| 1970 | 28.0 | 30.0 | 2.5 | * | 3.3 | 9.0 | 27.1 | 47.1 | 93.8 | 155.7 | 230.0 | 306.7 |
| 1969 | 27.2 | 16.3 | 2.1 | * | 3.2 | 10.1 | 23.4 | 49.5 | 83.5 | 159.0 | 234.0 | 402.3 |
| 1968 | 25.4 | 20.8 | 2.6 | 0.8 | 2.3 | 8.7 | 21.0 | 44.4 | 87.3 | 143.5 | 215.1 | 317.1 |
| ALL OTHER, FEMALE | | | | | | | | | | | | |
| 1978 | 28.1 | 43.6 | 4.0 | 1.1 | 2.9 | 5.7 | 12.5 | 29.1 | 67.6 | 130.2 | 242.7 | 536.7 |
| 1977 | 25.5 | 32.1 | 2.7 | 1.1 | 2.2 | 4.6 | 12.5 | 29.1 | 57.7 | 120.5 | 226.6 | 496.2 |
| 1976 | 25.5 | 38.7 | 2.0 | 0.9 | 2.6 | 5.5 | 12.9 | 29.3 | 64.4 | 122.5 | 209.0 | 479.6 |
| 1975 | 23.0 | 28.6 | 2.5 | 1.1 | 2.5 | 4.0 | 13.5 | 26.8 | 59.9 | 122.1 | 178.2 | 395.7 |
| 1974 | 23.4 | 42.7 | 3.2 | 0.8 | 2.6 | 6.9 | 14.1 | 28.4 | 62.9 | 111.5 | 172.3 | 465.9 |
| 1973 | 22.9 | 40.4 | 2.5 | * | 2.7 | 6.9 | 14.9 | 28.2 | 63.4 | 113.5 | 175.5 | 396.2 |
| 1972 | 22.2 | 40.3 | 2.4 | 1.1 | 2.8 | 7.2 | 14.3 | 34.1 | 59.1 | 107.3 | 164.1 | 328.9 |
| 1971 | 20.6 | 34.2 | 2.1 | 0.8 | 2.8 | 7.6 | 14.0 | 26.9 | 54.6 | 97.4 | 175.7 | 319.2 |
| 1970 | 20.0 | 26.7 | 2.9 | 0.7 | 2.6 | 5.9 | 14.6 | 30.8 | 55.6 | 100.0 | 151.8 | 260.6 |
| 1969 | 18.9 | 18.4 | * | 0.9 | 2.7 | 5.9 | 15.0 | 30.0 | 52.8 | 94.3 | 141.3 | 287.7 |
| 1968 | 18.8 | 22.1 | 2.3 | 0.7 | 2.9 | 6.0 | 15.3 | 28.1 | 52.4 | 92.7 | 149.8 | 263.9 |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|--|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| ALL OTHER FORMS OF HEART DISEASE (420-423,425-427,429) | | | | | | | | | | | | |
| BLACK, BOTH SEXES | | | | | | | | | | | | |
| 1978 | 35.6 | 46.2 | 3.8 | 1.1 | 3.7 | 8.0 | 21.5 | 47.5 | 95.4 | 172.4 | 303.2 | 606.0 |
| 1977 | 32.9 | 36.7 | 3.0 | 1.1 | 3.2 | 7.7 | 20.3 | 45.2 | 89.7 | 161.4 | 277.5 | 563.6 |
| 1976 | 32.7 | 41.3 | 2.9 | 0.9 | 3.8 | 8.3 | 20.0 | 45.6 | 90.1 | 167.4 | 260.2 | 544.1 |
| 1975 | 30.4 | 33.3 | 2.8 | 1.0 | 3.2 | 8.2 | 21.1 | 41.8 | 85.5 | 164.1 | 234.7 | 496.2 |
| 1974 | 29.6 | 43.4 | 3.2 | 0.8 | 3.3 | 10.1 | 20.6 | 43.3 | 88.1 | 150.9 | 213.3 | 508.3 |
| 1973 | 30.1 | 51.8 | 3.4 | 0.9 | 3.2 | 9.6 | 22.2 | 45.4 | 87.1 | 154.2 | 231.1 | 462.3 |
| 1972 | 29.2 | 47.8 | 3.1 | 1.0 | 3.9 | 9.1 | 21.7 | 47.4 | 85.7 | 154.3 | 209.2 | 396.4 |
| 1971 | 26.3 | 35.4 | 2.4 | 0.9 | 3.3 | 8.7 | 20.0 | 38.4 | 77.2 | 140.3 | 213.7 | 369.2 |
| 1970 | 25.8 | 28.9 | 2.9 | 0.7 | 3.1 | 8.3 | 22.7 | 42.0 | 78.5 | 133.9 | 196.5 | 293.1 |
| 1969 | 24.4 | 18.2 | 1.9 | 0.7 | 3.1 | 8.7 | 20.0 | 41.7 | 72.5 | 131.4 | 190.6 | 343.3 |
| 1968 | 23.6 | 23.3 | 2.6 | 0.8 | 2.8 | 7.9 | 19.7 | 38.1 | 73.6 | 122.9 | 189.9 | 300.0 |
| BLACK, MALE | | | | | | | | | | | | |
| 1978 | 40.5 | 45.2 | 3.6 | 1.0 | 4.2 | 9.8 | 30.4 | 65.5 | 122.2 | 217.4 | 373.2 | 696.1 |
| 1977 | 38.1 | 39.2 | 3.1 | 1.0 | 4.1 | 10.6 | 27.8 | 61.0 | 122.8 | 203.5 | 342.1 | 651.0 |
| 1976 | 37.7 | 41.9 | 3.7 | 1.1 | 4.7 | 10.6 | 26.7 | 60.6 | 114.2 | 217.2 | 324.8 | 640.4 |
| 1975 | 35.9 | 35.2 | 3.0 | 0.8 | 3.6 | 12.2 | 28.2 | 56.5 | 111.2 | 208.6 | 306.3 | 632.6 |
| 1974 | 34.2 | 44.7 | 3.1 | 0.9 | 3.9 | 12.7 | 26.7 | 57.9 | 113.3 | 194.3 | 265.6 | 548.8 |
| 1973 | 35.9 | 59.7 | 4.1 | 1.2 | 3.5 | 12.5 | 29.2 | 62.3 | 110.2 | 199.0 | 303.4 | 587.8 |
| 1972 | 34.6 | 51.5 | 3.6 | 0.7 | 4.9 | 10.7 | 28.3 | 59.3 | 113.8 | 203.8 | 265.0 | 490.0 |
| 1971 | 30.8 | 34.9 | 2.8 | 1.0 | 3.6 | 9.0 | 25.7 | 49.2 | 101.1 | 187.2 | 255.4 | 435.0 |
| 1970 | 30.3 | 31.1 | 2.6 | * | 3.6 | 10.3 | 30.3 | 51.9 | 101.7 | 168.4 | 249.1 | 330.4 |
| 1969 | 28.9 | 16.6 | 2.1 | * | 3.4 | 11.1 | 24.6 | 52.9 | 91.2 | 171.2 | 244.9 | 432.4 |
| 1968 | 27.2 | 22.7 | 2.9 | 0.9 | 2.5 | 9.7 | 23.0 | 47.4 | 94.3 | 153.6 | 232.1 | 345.7 |
| BLACK, FEMALE | | | | | | | | | | | | |
| 1978 | 31.3 | 47.3 | 3.9 | 1.2 | 3.2 | 6.5 | 14.1 | 32.8 | 73.4 | 138.8 | 256.4 | 559.6 |
| 1977 | 28.2 | 34.2 | 2.8 | 1.2 | 2.4 | 5.3 | 14.2 | 32.2 | 62.6 | 129.8 | 235.3 | 518.1 |
| 1976 | 28.1 | 40.7 | 2.2 | 0.8 | 2.9 | 6.3 | 14.5 | 33.2 | 70.1 | 129.6 | 216.9 | 493.3 |
| 1975 | 25.4 | 31.4 | 2.6 | 1.1 | 2.8 | 4.7 | 15.3 | 29.6 | 64.2 | 130.4 | 186.3 | 421.4 |
| 1974 | 25.5 | 42.1 | 3.3 | 0.8 | 2.8 | 7.9 | 15.5 | 31.1 | 67.2 | 117.8 | 178.0 | 485.9 |
| 1973 | 24.8 | 43.7 | 2.7 | * | 3.0 | 7.1 | 16.4 | 31.1 | 67.9 | 119.6 | 182.1 | 391.8 |
| 1972 | 24.2 | 43.9 | 2.6 | 1.2 | 3.0 | 7.7 | 16.1 | 37.1 | 62.1 | 115.9 | 169.3 | 342.9 |
| 1971 | 22.1 | 36.0 | * | 0.8 | 3.0 | 8.4 | 15.2 | 29.2 | 57.1 | 103.2 | 183.8 | 329.9 |
| 1970 | 21.6 | 26.8 | 3.2 | 0.8 | 2.6 | 6.6 | 16.5 | 33.5 | 58.7 | 106.6 | 157.9 | 269.0 |
| 1969 | 20.3 | 19.7 | * | 0.8 | 2.8 | 6.7 | 16.2 | 31.9 | 56.6 | 99.7 | 149.8 | 288.3 |
| 1968 | 20.2 | 23.8 | 2.4 | 0.7 | 3.0 | 6.3 | 16.9 | 29.9 | 55.7 | 97.9 | 158.6 | 276.8 |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| HYPERTENSION (400,401,403) | | | | | | | | | | | | |
| BOTH SEXES | | | | | | | | | | | | |
| 1978 | 2.5 | * | * | * | * | 0.1 | 0.6 | 1.3 | 2.9 | 7.8 | 24.2 | 67.4 |
| 1977 | 2.6 | * | * | * | * | 0.2 | 0.6 | 1.4 | 3.0 | 8.3 | 25.3 | 72.9 |
| 1976 | 2.8 | * | * | * | * | 0.2 | 0.7 | 1.6 | 3.5 | 9.3 | 28.2 | 77.6 |
| 1975 | 2.9 | * | * | * | * | 0.2 | 0.7 | 1.8 | 3.6 | 9.8 | 29.4 | 81.4 |
| 1974 | 3.2 | * | * | * | * | 0.3 | 0.9 | 2.0 | 4.2 | 11.0 | 32.2 | 93.7 |
| 1973 | 3.5 | * | * | * | 0.1 | 0.3 | 1.1 | 2.1 | 4.8 | 12.1 | 36.3 | 98.8 |
| 1972 | 3.7 | * | * | * | 0.1 | 0.5 | 1.4 | 2.5 | 5.2 | 12.4 | 37.7 | 102.7 |
| 1971 | 3.8 | * | * | * | 0.1 | 0.6 | 1.7 | 2.9 | 5.2 | 12.9 | 38.9 | 97.5 |
| 1970 | 4.1 | * | * | * | 0.1 | 0.6 | 2.0 | 3.3 | 5.5 | 13.8 | 42.7 | 99.9 |
| 1969 | 4.2 | * | * | * | 0.1 | 0.7 | 2.0 | 3.4 | 5.7 | 14.7 | 42.5 | 115.1 |
| 1968 | 4.5 | * | * | * | 0.1 | 0.8 | 2.3 | 3.9 | 6.2 | 15.3 | 46.3 | 132.9 |
| MALE | | | | | | | | | | | | |
| 1978 | 2.4 | * | * | * | * | 0.2 | 0.7 | 1.5 | 3.5 | 9.4 | 29.4 | 82.5 |
| 1977 | 2.6 | * | * | * | * | 0.2 | 0.7 | 1.5 | 3.6 | 9.9 | 32.8 | 92.3 |
| 1976 | 2.8 | * | * | * | * | 0.2 | 0.8 | 2.0 | 4.0 | 11.3 | 34.3 | 97.0 |
| 1975 | 2.9 | * | * | * | * | 0.3 | 0.9 | 2.1 | 4.4 | 11.4 | 36.4 | 102.0 |
| 1974 | 3.4 | * | * | * | * | 0.3 | 1.0 | 2.3 | 5.1 | 14.4 | 40.9 | 120.7 |
| 1973 | 3.6 | * | * | * | * | 0.3 | 1.3 | 2.4 | 5.5 | 15.2 | 47.1 | 122.1 |
| 1972 | 3.8 | * | * | * | * | 0.5 | 1.5 | 2.9 | 6.3 | 15.3 | 46.5 | 138.3 |
| 1971 | 4.0 | * | * | * | * | 0.6 | 2.0 | 3.4 | 6.2 | 16.0 | 47.9 | 126.9 |
| 1970 | 4.3 | * | * | * | * | 0.7 | 2.2 | 3.9 | 6.7 | 17.6 | 53.9 | 120.2 |
| 1969 | 4.4 | * | * | * | * | 0.7 | 2.1 | 4.0 | 6.7 | 18.9 | 53.0 | 143.8 |
| 1968 | 4.9 | * | * | * | * | 0.8 | 2.6 | 4.6 | 7.9 | 18.7 | 58.4 | 168.4 |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| HYPERTENSION (400,401,403) | | | | | | | | | | | | |
| FEMALE | | | | | | | | | | | | |
| 1978 | 2.5 | * | * | * | * | * | 0.5 | 1.1 | 2.4 | 6.6 | 21.1 | 60.6 |
| 1977 | 2.6 | * | * | * | * | 0.1 | 0.6 | 1.3 | 2.5 | 7.2 | 20.8 | 64.0 |
| 1976 | 2.8 | * | * | * | * | 0.2 | 0.5 | 1.2 | 3.0 | 7.8 | 24.5 | 68.4 |
| 1975 | 2.9 | * | * | * | * | 0.1 | 0.6 | 1.5 | 2.9 | 8.6 | 25.2 | 71.5 |
| 1974 | 3.1 | * | * | * | * | 0.3 | 0.8 | 1.6 | 3.5 | 8.5 | 26.8 | 80.4 |
| 1973 | 3.4 | * | * | * | * | 0.4 | 1.0 | 1.9 | 4.1 | 9.7 | 29.5 | 87.1 |
| 1972 | 3.6 | * | * | * | * | 0.6 | 1.3 | 2.1 | 4.3 | 10.2 | 32.1 | 84.5 |
| 1971 | 3.6 | * | * | * | * | 0.5 | 1.5 | 2.5 | 4.4 | 10.5 | 33.1 | 82.2 |
| 1970 | 3.8 | * | * | * | * | 0.5 | 1.9 | 2.6 | 4.5 | 10.9 | 35.2 | 88.6 |
| 1969 | 4.0 | * | * | * | 0.1 | 0.8 | 1.9 | 2.9 | 4.8 | 11.4 | 35.5 | 99.6 |
| 1968 | 4.2 | * | * | * | * | 0.9 | 2.0 | 3.2 | 4.7 | 12.7 | 38.0 | 113.5 |
| WHITE, BOTH SEXES | | | | | | | | | | | | |
| 1978 | 2.3 | * | * | * | * | 0.1 | 0.2 | 0.7 | 2.1 | 6.7 | 23.1 | 68.0 |
| 1977 | 2.4 | * | * | * | * | 0.1 | 0.3 | 0.9 | 2.2 | 7.0 | 24.1 | 73.1 |
| 1976 | 2.6 | * | * | * | * | 0.1 | 0.3 | 0.9 | 2.5 | 7.7 | 27.0 | 77.6 |
| 1975 | 2.7 | * | * | * | * | 0.1 | 0.4 | 1.0 | 2.5 | 8.2 | 28.3 | 80.5 |
| 1974 | 2.9 | * | * | * | * | 0.1 | 0.4 | 1.1 | 3.1 | 9.2 | 31.1 | 94.5 |
| 1973 | 3.2 | * | * | * | * | 0.1 | 0.5 | 1.2 | 3.3 | 10.2 | 34.9 | 99.6 |
| 1972 | 3.3 | * | * | * | * | 0.2 | 0.5 | 1.4 | 3.7 | 10.4 | 36.0 | 102.6 |
| 1971 | 3.3 | * | * | * | * | 0.2 | 0.7 | 1.6 | 3.7 | 11.1 | 37.5 | 97.5 |
| 1970 | 3.6 | * | * | * | * | 0.2 | 0.8 | 1.6 | 4.1 | 11.7 | 41.5 | 101.0 |
| 1969 | 3.6 | * | * | * | * | 0.2 | 0.7 | 1.9 | 4.0 | 12.6 | 41.6 | 115.6 |
| 1968 | 3.9 | * | * | * | * | 0.3 | 0.8 | 2.1 | 4.5 | 13.0 | 44.7 | 134.7 |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| HYPERTENSION (400,401,403) | | | | | | | | | | | | |
| WHITE, MALE | | | | | | | | | | | | |
| 1978 | 2.2 | * | * | * | * | * | 0.3 | 0.9 | 2.6 | 8.3 | 28.4 | 83.2 |
| 1977 | 2.3 | * | * | * | * | * | 0.3 | 1.0 | 2.6 | 8.4 | 31.7 | 92.5 |
| 1976 | 2.5 | * | * | * | * | * | 0.3 | 1.2 | 2.8 | 9.7 | 33.7 | 96.2 |
| 1975 | 2.6 | * | * | * | * | * | 0.4 | 1.2 | 3.2 | 9.8 | 35.0 | 102.4 |
| 1974 | 3.0 | * | * | * | * | * | 0.4 | 1.3 | 3.8 | 12.7 | 39.3 | 122.3 |
| 1973 | 3.3 | * | * | * | * | * | 0.6 | 1.6 | 3.9 | 13.6 | 45.4 | 122.7 |
| 1972 | 3.4 | * | * | * | * | 0.2 | 0.5 | 1.8 | 4.8 | 13.4 | 44.4 | 136.8 |
| 1971 | 3.4 | * | * | * | * | 0.2 | 0.8 | 1.8 | 4.4 | 14.2 | 46.1 | 128.3 |
| 1970 | 3.8 | * | * | * | * | 0.3 | 0.9 | 2.0 | 5.1 | 15.6 | 52.3 | 122.4 |
| 1969 | 3.8 | * | * | * | * | 0.2 | 0.8 | 2.2 | 4.7 | 16.5 | 52.0 | 145.3 |
| 1968 | 4.2 | * | * | * | * | 0.3 | 1.0 | 2.5 | 6.2 | 16.4 | 56.3 | 170.7 |
| WHITE, FEMALE | | | | | | | | | | | | |
| 1978 | 2.4 | * | * | * | * | * | 0.2 | 0.6 | 1.7 | 5.4 | 20.0 | 61.3 |
| 1977 | 2.4 | * | * | * | * | * | 0.3 | 0.8 | 1.9 | 5.9 | 19.6 | 64.4 |
| 1976 | 2.6 | * | * | * | * | * | 0.2 | 0.6 | 2.1 | 6.1 | 22.9 | 69.0 |
| 1975 | 2.7 | * | * | * | * | * | 0.3 | 0.8 | 2.0 | 6.9 | 24.3 | 70.2 |
| 1974 | 2.8 | * | * | * | * | * | 0.3 | 0.9 | 2.5 | 6.5 | 26.1 | 81.0 |
| 1973 | 3.1 | * | * | * | * | * | 0.5 | 0.9 | 2.8 | 7.6 | 28.3 | 88.3 |
| 1972 | 3.2 | * | * | * | * | 0.3 | 0.5 | 1.1 | 2.8 | 8.1 | 30.7 | 85.4 |
| 1971 | 3.2 | * | * | * | * | * | 0.6 | 1.3 | 3.0 | 8.6 | 32.0 | 81.7 |
| 1970 | 3.4 | * | * | * | * | 0.2 | 0.7 | 1.3 | 3.2 | 8.8 | 34.5 | 89.3 |
| 1969 | 3.5 | * | * | * | * | 0.3 | 0.7 | 1.6 | 3.3 | 9.6 | 34.7 | 100.0 |
| 1968 | 3.7 | * | * | * | * | 0.3 | 0.7 | 1.7 | 3.0 | 10.5 | 36.8 | 115.4 |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| HYPERTENSION (400,401,403) | | | | | | | | | | | | |
| ALL OTHER, BOTH SEXES | | | | | | | | | | | | |
| 1978 | 3.6 | * | * | * | * | 0.6 | 2.8 | 5.4 | 10.4 | 18.4 | 34.8 | 60.7 |
| 1977 | 3.9 | * | * | * | * | 0.8 | 3.1 | 5.7 | 10.5 | 21.4 | 37.6 | 70.0 |
| 1976 | 4.5 | * | * | * | * | 0.9 | 3.4 | 6.9 | 12.9 | 24.9 | 40.9 | 77.1 |
| 1975 | 4.7 | * | * | * | * | 1.1 | 3.2 | 8.4 | 13.4 | 25.6 | 41.5 | 91.8 |
| 1974 | 5.2 | * | * | * | * | 1.3 | 4.7 | 9.2 | 14.9 | 29.4 | 44.0 | 84.4 |
| 1973 | 5.9 | * | * | * | * | 1.8 | 5.3 | 9.8 | 18.6 | 31.0 | 51.7 | 89.8 |
| 1972 | 6.8 | * | * | * | 0.4 | 2.5 | 7.9 | 11.8 | 19.8 | 32.2 | 57.4 | 104.1 |
| 1971 | 7.2 | * | * | * | * | 3.3 | 9.6 | 14.9 | 20.1 | 31.1 | 56.1 | 97.5 |
| 1970 | 7.6 | * | * | * | * | 3.1 | 11.1 | 17.7 | 19.1 | 34.3 | 57.1 | 88.7 |
| 1969 | 8.1 | * | * | * | * | 4.2 | 11.5 | 17.5 | 22.8 | 35.0 | 54.4 | 109.2 |
| 1968 | 9.0 | * | * | * | * | 4.6 | 13.4 | 19.8 | 23.1 | 38.3 | 67.1 | 112.7 |
| ALL OTHER, MALE | | | | | | | | | | | | |
| 1978 | 3.7 | * | * | * | * | * | 3.3 | 6.3 | 12.1 | 19.6 | 38.0 | 75.9 |
| 1977 | 4.1 | * | * | * | * | 1.0 | 3.5 | 6.5 | 13.3 | 23.4 | 42.9 | 91.1 |
| 1976 | 4.7 | * | * | * | * | 1.3 | 4.6 | 8.6 | 15.4 | 26.3 | 39.7 | 105.5 |
| 1975 | 5.0 | * | * | * | * | 1.6 | 4.5 | 9.8 | 15.4 | 26.4 | 50.8 | 98.1 |
| 1974 | 5.6 | * | * | * | * | 1.4 | 5.5 | 11.4 | 17.2 | 30.2 | 57.6 | 104.0 |
| 1973 | 6.1 | * | * | * | * | 1.9 | 6.3 | 9.8 | 21.6 | 30.8 | 64.7 | 116.7 |
| 1972 | 7.2 | * | * | * | * | 2.7 | 9.3 | 13.5 | 21.5 | 33.5 | 69.1 | 153.2 |
| 1971 | 7.7 | * | * | * | * | 3.4 | 11.3 | 17.5 | 23.1 | 33.0 | 67.6 | 112.8 |
| 1970 | 8.4 | * | * | * | * | 4.2 | 12.3 | 21.5 | 22.2 | 36.4 | 72.0 | 101.0 |
| 1969 | 8.9 | * | * | * | * | 4.3 | 12.6 | 20.7 | 26.0 | 41.6 | 64.9 | 129.5 |
| 1968 | 9.9 | * | * | * | * | 4.7 | 16.5 | 24.1 | 25.1 | 40.9 | 83.9 | 146.3 |
| ALL OTHER, FEMALE | | | | | | | | | | | | |
| 1978 | 3.5 | * | * | * | * | * | 2.3 | 4.6 | 9.0 | 17.5 | 32.6 | 53.2 |
| 1977 | 3.7 | * | * | * | * | * | 2.7 | 5.0 | 8.2 | 19.9 | 33.9 | 58.7 |
| 1976 | 4.3 | * | * | * | * | * | 2.4 | 5.6 | 10.8 | 23.8 | 41.8 | 61.2 |
| 1975 | 4.5 | * | * | * | * | * | 2.2 | 7.2 | 11.7 | 24.9 | 35.0 | 87.1 |
| 1974 | 4.9 | * | * | * | * | 1.1 | 4.0 | 7.3 | 13.0 | 28.7 | 34.6 | 72.9 |
| 1973 | 5.7 | * | * | * | * | 1.8 | 4.5 | 9.7 | 16.0 | 31.1 | 42.8 | 73.4 |
| 1972 | 6.4 | * | * | * | * | 2.4 | 6.6 | 10.4 | 18.3 | 31.2 | 49.2 | 73.7 |
| 1971 | 6.7 | * | * | * | * | 3.2 | 8.1 | 12.5 | 17.5 | 29.6 | 47.5 | 87.7 |
| 1970 | 6.9 | * | * | * | * | 2.3 | 10.1 | 14.2 | 16.4 | 32.6 | 45.7 | 80.1 |
| 1969 | 7.4 | * | * | * | * | 4.1 | 10.6 | 14.7 | 20.0 | 29.5 | 46.2 | 95.4 |
| 1968 | 8.1 | * | * | * | * | 4.5 | 10.9 | 16.1 | 21.3 | 36.1 | 54.4 | 90.2 |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| HYPERTENSION (400,401,403) | | | | | | | | | | | | |
| BLACK, BOTH SEXES | | | | | | | | | | | | |
| 1978 | 4.0 | * | * | * | * | 0.7 | 3.3 | 6.1 | 11.5 | 19.8 | 37.5 | 64.0 |
| 1977 | 4.4 | * | * | * | * | 1.0 | 3.6 | 6.4 | 11.8 | 23.4 | 40.0 | 72.0 |
| 1976 | 5.0 | * | * | * | * | 1.1 | 4.0 | 7.8 | 14.0 | 27.4 | 42.4 | 81.6 |
| 1975 | 5.3 | * | * | * | * | 1.2 | 3.7 | 9.5 | 14.8 | 27.7 | 43.5 | 96.9 |
| 1974 | 5.7 | * | * | * | * | 1.5 | 5.4 | 10.3 | 15.9 | 31.5 | 45.0 | 88.4 |
| 1973 | 6.4 | * | * | * | * | 2.1 | 6.1 | 10.9 | 20.3 | 33.4 | 52.5 | 97.4 |
| 1972 | 7.4 | * | * | * | * | 3.0 | 9.0 | 13.2 | 21.4 | 35.1 | 59.2 | 110.9 |
| 1971 | 7.9 | * | * | * | * | 3.7 | 10.8 | 16.5 | 21.7 | 33.4 | 59.1 | 104.7 |
| 1970 | 8.4 | * | * | * | * | 3.7 | 12.7 | 19.7 | 20.8 | 37.5 | 60.5 | 95.4 |
| 1969 | 8.8 | * | * | * | * | 4.6 | 12.9 | 19.3 | 24.7 | 36.7 | 57.3 | 116.5 |
| 1968 | 9.7 | * | * | * | * | 5.2 | 15.1 | 21.5 | 25.1 | 40.7 | 69.5 | 120.7 |
| BLACK, MALE | | | | | | | | | | | | |
| 1978 | 4.1 | * | * | * | * | * | 4.0 | 6.8 | 13.3 | 21.2 | 41.0 | 76.5 |
| 1977 | 4.7 | * | * | * | * | 1.2 | 4.1 | 7.5 | 15.0 | 26.1 | 46.8 | 95.9 |
| 1976 | 5.3 | * | * | * | * | 1.6 | 5.5 | 9.6 | 16.7 | 29.5 | 40.2 | 117.0 |
| 1975 | 5.6 | * | * | * | * | 1.9 | 5.1 | 11.2 | 17.1 | 29.5 | 53.6 | 104.3 |
| 1974 | 6.1 | * | * | * | * | 1.7 | 6.3 | 12.7 | 19.0 | 32.6 | 58.4 | 116.3 |
| 1973 | 6.7 | * | * | * | * | 2.2 | 7.3 | 11.1 | 24.0 | 33.8 | 65.2 | 131.7 |
| 1972 | 7.9 | * | * | * | * | 3.1 | 10.8 | 15.2 | 23.8 | 37.1 | 71.1 | 175.0 |
| 1971 | 8.5 | * | * | * | * | 3.9 | 12.8 | 19.6 | 25.5 | 36.2 | 71.5 | 120.0 |
| 1970 | 9.4 | * | * | * | * | 4.9 | 14.1 | 24.1 | 24.5 | 40.6 | 76.5 | 117.4 |
| 1969 | 9.6 | * | * | * | * | 4.8 | 14.2 | 22.9 | 28.5 | 43.5 | 67.1 | 148.6 |
| 1968 | 10.8 | * | * | * | * | 5.3 | 18.6 | 26.4 | 27.7 | 44.2 | 88.5 | 162.9 |
| BLACK, FEMALE | | | | | | | | | | | | |
| 1978 | 4.0 | * | * | * | * | * | 2.8 | 5.4 | 10.0 | 18.7 | 35.2 | 57.6 |
| 1977 | 4.2 | * | * | * | * | * | 3.2 | 5.5 | 9.2 | 21.3 | 35.6 | 59.6 |
| 1976 | 4.8 | * | * | * | * | * | 2.7 | 6.4 | 11.7 | 25.8 | 43.8 | 62.9 |
| 1975 | 5.0 | * | * | * | * | * | 2.5 | 8.1 | 12.8 | 26.4 | 36.6 | 92.9 |
| 1974 | 5.3 | * | * | * | * | 1.3 | 4.7 | 8.2 | 13.2 | 30.8 | 35.9 | 73.1 |
| 1973 | 6.3 | * | * | * | * | 2.0 | 5.1 | 10.7 | 17.1 | 33.2 | 43.9 | 78.1 |
| 1972 | 7.0 | * | * | * | * | 2.8 | 7.6 | 11.5 | 19.4 | 33.6 | 50.7 | 74.3 |
| 1971 | 7.3 | * | * | * | * | 3.5 | 9.2 | 13.8 | 18.4 | 31.2 | 50.2 | 95.5 |
| 1970 | 7.6 | * | * | * | * | 2.6 | 11.6 | 15.9 | 17.7 | 35.1 | 48.7 | 81.2 |
| 1969 | 8.1 | * | * | * | * | 4.5 | 11.8 | 16.1 | 21.4 | 31.3 | 49.8 | 96.7 |
| 1968 | 8.8 | * | * | * | * | 5.1 | 12.1 | 17.2 | 22.8 | 37.8 | 55.4 | 96.4 |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| Cerebrovascular disease 430-438 | | | | | | | | | | | | |
| BOTH SEXES | | | | | | | | | | | | |
| 1978 | 79.1 | 5.3 | 0.8 | 0.5 | 1.1 | 2.6 | 9.8 | 27.4 | 72.6 | 242.7 | 849.6 | 2,402.4 |
| 1977 | 82.8 | 5.1 | 0.8 | 0.5 | 1.1 | 3.0 | 10.3 | 28.7 | 78.1 | 258.9 | 913.1 | 2,531.1 |
| 1976 | 86.7 | 4.2 | 0.7 | 0.5 | 1.2 | 3.4 | 11.5 | 31.4 | 84.4 | 279.2 | 961.5 | 2,682.3 |
| 1975 | 90.1 | 5.0 | 0.8 | 0.5 | 1.4 | 3.4 | 11.7 | 32.3 | 90.4 | 302.2 | 1,028.8 | 2,736.4 |
| 1974 | 97.2 | 5.0 | 0.8 | 0.6 | 1.3 | 3.5 | 13.3 | 35.3 | 98.5 | 333.5 | 1,133.9 | 3,063.6 |
| 1973 | 101.4 | 4.3 | 1.0 | 0.6 | 1.4 | 4.2 | 14.0 | 38.8 | 106.1 | 354.3 | 1,198.7 | 3,243.7 |
| 1972 | 101.9 | 4.3 | 0.9 | 0.6 | 1.4 | 4.2 | 14.6 | 40.4 | 111.2 | 366.9 | 1,213.0 | 3,239.7 |
| 1971 | 101.1 | 3.9 | 1.1 | 0.6 | 1.7 | 4.6 | 15.0 | 41.1 | 111.9 | 367.9 | 1,218.7 | 3,217.2 |
| 1970 | 101.9 | 5.0 | 1.0 | 0.7 | 1.6 | 4.5 | 15.6 | 41.6 | 115.8 | 384.1 | 1,254.2 | 3,014.3 |
| 1969 | 102.9 | 4.9 | 0.9 | 0.7 | 1.6 | 4.9 | 15.9 | 41.8 | 115.9 | 395.0 | 1,263.8 | 3,431.1 |
| 1968 | 106.0 | 5.3 | 0.9 | 0.7 | 1.7 | 4.9 | 16.5 | 44.6 | 121.7 | 409.5 | 1,317.6 | 3,605.9 |
| MALE | | | | | | | | | | | | |
| 1978 | 68.2 | 5.6 | 1.0 | 0.6 | 1.2 | 2.6 | 10.1 | 28.8 | 84.3 | 289.8 | 926.8 | 2,368.1 |
| 1977 | 72.4 | 5.1 | 0.8 | 0.6 | 1.2 | 3.1 | 10.4 | 30.9 | 91.5 | 309.9 | 1,005.1 | 2,509.7 |
| 1976 | 76.1 | 4.3 | 0.8 | 0.6 | 1.4 | 3.4 | 11.3 | 33.4 | 98.1 | 334.2 | 1,049.1 | 2,672.1 |
| 1975 | 80.4 | 5.2 | 0.9 | 0.5 | 1.5 | 3.3 | 11.6 | 34.0 | 107.1 | 362.7 | 1,131.5 | 2,735.7 |
| 1974 | 87.0 | 5.8 | 0.9 | 0.7 | 1.4 | 3.4 | 13.2 | 38.0 | 116.5 | 400.0 | 1,237.7 | 3,032.6 |
| 1973 | 91.1 | 4.3 | 1.0 | 0.6 | 1.4 | 3.8 | 13.8 | 41.0 | 125.3 | 424.7 | 1,306.7 | 3,174.5 |
| 1972 | 93.5 | 5.2 | 1.0 | 0.7 | 1.6 | 4.1 | 14.4 | 43.0 | 134.4 | 439.0 | 1,332.8 | 3,225.3 |
| 1971 | 93.5 | 4.3 | 1.2 | 0.6 | 1.9 | 4.4 | 14.6 | 44.7 | 133.6 | 438.8 | 1,345.4 | 3,192.4 |
| 1970 | 94.5 | 5.8 | 1.2 | 0.8 | 1.8 | 4.4 | 15.7 | 44.4 | 138.7 | 449.5 | 1,361.6 | 2,895.2 |
| 1969 | 96.2 | 5.2 | 1.1 | 0.7 | 1.5 | 4.7 | 15.8 | 45.4 | 137.8 | 465.7 | 1,373.9 | 3,396.1 |
| 1968 | 99.6 | 5.6 | 0.8 | 0.7 | 1.8 | 4.8 | 16.4 | 47.9 | 144.2 | 483.1 | 1,418.2 | 3,591.6 |

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| Cerebrovascular disease 430-438 | | | | | | | | | | | | |
| FEMALE | | | | | | | | | | | | |
| 1978 | 89.4 | 4.9 | 0.6 | 0.5 | 1.1 | 2.6 | 9.5 | 26.1 | 62.2 | 206.7 | 803.7 | 2,417.9 |
| 1977 | 92.6 | 5.1 | 0.8 | 0.4 | 1.1 | 2.9 | 10.2 | 26.7 | 66.2 | 219.8 | 858.0 | 2,540.9 |
| 1976 | 96.7 | 4.1 | 0.7 | 0.5 | 1.1 | 3.3 | 11.7 | 29.6 | 72.3 | 237.1 | 908.6 | 2,687.1 |
| 1975 | 99.2 | 4.7 | 0.7 | 0.5 | 1.2 | 3.6 | 11.9 | 30.7 | 75.7 | 255.8 | 966.2 | 2,736.8 |
| 1974 | 106.9 | 4.1 | 0.7 | 0.6 | 1.3 | 3.6 | 13.4 | 32.9 | 82.7 | 282.5 | 1,069.9 | 3,079.0 |
| 1973 | 111.1 | 4.2 | 1.0 | 0.6 | 1.4 | 4.5 | 14.1 | 36.7 | 89.0 | 300.3 | 1,131.5 | 3,278.5 |
| 1972 | 109.9 | 3.3 | 0.8 | 0.5 | 1.2 | 4.3 | 14.7 | 38.0 | 90.5 | 311.3 | 1,136.4 | 3,247.1 |
| 1971 | 108.3 | 3.5 | 1.0 | 0.6 | 1.6 | 4.8 | 15.3 | 37.7 | 92.4 | 313.1 | 1,136.3 | 3,230.2 |
| 1970 | 109.0 | 4.0 | 0.7 | 0.6 | 1.4 | 4.7 | 15.6 | 39.0 | 95.3 | 333.3 | 1,183.1 | 3,081.0 |
| 1969 | 109.3 | 4.6 | 0.7 | 0.7 | 1.7 | 5.1 | 16.0 | 38.6 | 96.1 | 339.8 | 1,190.1 | 3,450.0 |
| 1968 | 112.1 | 4.9 | 0.9 | 0.6 | 1.7 | 4.9 | 16.6 | 41.5 | 101.4 | 351.6 | 1,248.8 | 3,618.3 |
| WHITE, BOTH SEXES | | | | | | | | | | | | |
| 1978 | 80.1 | 3.8 | 0.8 | 0.5 | 1.1 | 2.0 | 7.7 | 22.0 | 62.3 | 223.8 | 843.6 | 2,447.2 |
| 1977 | 83.8 | 4.1 | 0.8 | 0.5 | 1.0 | 2.4 | 7.9 | 23.1 | 67.5 | 239.0 | 908.1 | 2,581.5 |
| 1976 | 87.8 | 3.8 | 0.6 | 0.5 | 1.1 | 2.7 | 8.9 | 25.4 | 73.4 | 257.9 | 960.9 | 2,734.7 |
| 1975 | 91.2 | 4.3 | 0.8 | 0.5 | 1.2 | 2.8 | 9.0 | 26.3 | 78.3 | 279.4 | 1,029.5 | 2,797.3 |
| 1974 | 98.3 | 3.8 | 0.7 | 0.6 | 1.2 | 2.8 | 10.3 | 28.4 | 84.7 | 307.5 | 1,139.2 | 3,129.7 |
| 1973 | 102.0 | 3.6 | 0.9 | 0.5 | 1.2 | 3.2 | 10.2 | 30.8 | 90.6 | 325.8 | 1,201.6 | 3,314.9 |
| 1972 | 102.5 | 3.1 | 0.7 | 0.5 | 1.3 | 3.2 | 11.0 | 32.2 | 95.5 | 339.2 | 1,215.6 | 3,313.3 |
| 1971 | 101.4 | 3.3 | 1.0 | 0.6 | 1.5 | 3.6 | 11.3 | 32.7 | 95.7 | 339.1 | 1,220.7 | 3,299.0 |
| 1970 | 101.9 | 3.9 | 0.9 | 0.7 | 1.4 | 3.3 | 11.6 | 33.0 | 97.9 | 354.1 | 1,249.9 | 3,114.7 |
| 1969 | 102.4 | 4.2 | 0.9 | 0.7 | 1.5 | 3.6 | 11.6 | 32.6 | 97.6 | 363.3 | 1,257.1 | 3,524.4 |
| 1968 | 105.1 | 4.5 | 0.8 | 0.7 | 1.5 | 3.6 | 12.0 | 35.1 | 101.3 | 375.0 | 1,312.5 | 3,698.6 |

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| Cerebrovascular disease 430-438 | | | | | | | | | | | | |
| WHITE, MALE | | | | | | | | | | | | |
| 1978 | 67.8 | 4.0 | 1.0 | 0.6 | 1.1 | 2.0 | 7.5 | 22.5 | 72.5 | 268.9 | 923.6 | 2,415.7 |
| 1977 | 72.1 | 4.6 | 0.8 | 0.5 | 1.1 | 2.5 | 7.7 | 24.7 | 79.5 | 289.7 | 1,004.9 | 2,560.1 |
| 1976 | 75.8 | 4.2 | 0.7 | 0.6 | 1.2 | 2.7 | 8.6 | 26.7 | 86.3 | 312.3 | 1,050.3 | 2,727.2 |
| 1975 | 80.2 | 4.6 | 0.9 | 0.5 | 1.4 | 2.6 | 8.7 | 27.6 | 93.8 | 339.4 | 1,134.9 | 2,807.4 |
| 1974 | 86.8 | 4.4 | 0.8 | 0.7 | 1.3 | 2.7 | 10.2 | 30.4 | 101.4 | 374.4 | 1,246.5 | 3,103.3 |
| 1973 | 90.5 | 3.6 | 0.9 | 0.5 | 1.2 | 2.9 | 10.1 | 32.4 | 108.7 | 397.3 | 1,313.7 | 3,248.4 |
| 1972 | 93.2 | 3.8 | 0.7 | 0.6 | 1.5 | 3.2 | 10.8 | 33.8 | 117.9 | 415.7 | 1,340.5 | 3,303.2 |
| 1971 | 93.0 | 4.0 | 1.1 | 0.6 | 1.6 | 3.5 | 11.0 | 36.2 | 116.5 | 410.7 | 1,353.6 | 3,294.4 |
| 1970 | 93.5 | 4.5 | 1.2 | 0.8 | 1.6 | 3.2 | 11.8 | 35.6 | 119.9 | 420.0 | 1,361.6 | 3,018.1 |
| 1969 | 94.7 | 4.6 | 1.0 | 0.6 | 1.4 | 3.5 | 11.4 | 35.6 | 119.5 | 434.6 | 1,370.2 | 3,504.1 |
| 1968 | 97.8 | 5.0 | 0.7 | 0.7 | 1.5 | 3.6 | 12.0 | 38.1 | 122.8 | 448.8 | 1,416.7 | 3,697.5 |
| WHITE, FEMALE | | | | | | | | | | | | |
| 1978 | 91.9 | 3.7 | 0.6 | 0.4 | 1.0 | 2.1 | 7.8 | 21.6 | 53.2 | 189.4 | 796.5 | 2,461.3 |
| 1977 | 95.0 | 3.6 | 0.8 | 0.4 | 0.9 | 2.4 | 8.1 | 21.6 | 56.9 | 200.3 | 851.1 | 2,591.2 |
| 1976 | 99.2 | 3.3 | 0.6 | 0.5 | 1.0 | 2.7 | 9.1 | 24.1 | 61.9 | 216.4 | 907.8 | 2,740.5 |
| 1975 | 101.6 | 3.9 | 0.7 | 0.5 | 1.1 | 3.0 | 9.3 | 25.0 | 64.5 | 233.6 | 966.2 | 2,794.9 |
| 1974 | 109.3 | 3.1 | 0.7 | 0.6 | 1.2 | 2.9 | 10.5 | 26.5 | 69.8 | 256.5 | 1,073.4 | 3,142.5 |
| 1973 | 113.0 | 3.5 | 1.0 | 0.6 | 1.2 | 3.5 | 10.3 | 29.3 | 74.4 | 271.2 | 1,132.2 | 3,351.3 |
| 1972 | 111.4 | 2.4 | 0.6 | 0.4 | 1.1 | 3.2 | 11.1 | 30.6 | 75.4 | 280.4 | 1,136.6 | 3,318.4 |
| 1971 | 109.5 | 2.6 | 1.0 | 0.6 | 1.4 | 3.7 | 11.6 | 29.5 | 77.1 | 283.9 | 1,135.5 | 3,301.3 |
| 1970 | 109.8 | 3.2 | 0.6 | 0.6 | 1.1 | 3.4 | 11.5 | 30.5 | 78.1 | 303.2 | 1,176.8 | 3,167.6 |
| 1969 | 109.6 | 3.8 | 0.7 | 0.7 | 1.6 | 3.8 | 11.8 | 29.8 | 77.8 | 307.9 | 1,181.9 | 3,539.6 |
| 1968 | 112.1 | 4.1 | 0.9 | 0.6 | 1.4 | 3.6 | 11.9 | 32.2 | 81.9 | 317.2 | 1,241.8 | 3,704.2 |

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| Cerebrovascular disease 430-438 | | | | | | | | | | | | |
| ALL OTHER, BOTH SEXES | | | | | | | | | | | | |
| 1978 | 72.4 | 12.1 | * | 0.7 | 1.5 | 6.2 | 24.0 | 68.8 | 165.2 | 418.7 | 911.4 | 1,888.7 |
| 1977 | 75.9 | 9.8 | * | 0.5 | 1.8 | 6.7 | 26.3 | 72.6 | 174.2 | 447.4 | 964.4 | 1,953.8 |
| 1976 | 79.6 | 6.4 | 1.1 | 0.5 | 2.1 | 7.6 | 29.2 | 80.2 | 186.0 | 483.1 | 967.3 | 2,066.7 |
| 1975 | 82.7 | 8.6 | 1.0 | 0.7 | 2.1 | 8.0 | 30.3 | 81.5 | 202.6 | 524.0 | 1,021.3 | 2,038.4 |
| 1974 | 89.9 | 10.9 | 1.2 | 0.6 | 2.0 | 8.6 | 34.2 | 93.4 | 228.3 | 588.5 | 1,077.0 | 2,294.8 |
| 1973 | 97.2 | 7.6 | 1.5 | 0.8 | 2.4 | 10.6 | 40.7 | 106.2 | 252.5 | 635.7 | 1,164.1 | 2,413.4 |
| 1972 | 98.1 | 10.4 | 1.8 | 0.9 | 2.2 | 11.3 | 40.7 | 111.5 | 260.8 | 641.7 | 1,183.0 | 2,390.2 |
| 1971 | 98.6 | 7.2 | 1.6 | 0.9 | 3.0 | 11.6 | 41.9 | 113.8 | 266.2 | 652.8 | 1,194.5 | 2,285.8 |
| 1970 | 102.6 | 10.7 | 1.4 | 0.7 | 2.9 | 12.9 | 45.3 | 117.4 | 285.1 | 677.4 | 1,308.8 | 1,983.0 |
| 1969 | 106.5 | 8.9 | 1.2 | 0.7 | 2.5 | 14.3 | 48.4 | 124.7 | 292.0 | 710.3 | 1,353.5 | 2,373.4 |
| 1968 | 112.4 | 9.1 | 1.3 | 0.7 | 3.5 | 13.8 | 51.5 | 130.4 | 319.4 | 755.7 | 1,383.8 | 2,570.6 |
| ALL OTHER, MALE | | | | | | | | | | | | |
| 1978 | 71.2 | 13.1 | * | * | 1.3 | 6.7 | 29.0 | 81.4 | 194.7 | 483.0 | 954.1 | 1,881.0 |
| 1977 | 74.0 | 7.4 | * | * | 1.6 | 7.8 | 30.3 | 83.0 | 204.3 | 498.1 | 1,006.5 | 1,996.4 |
| 1976 | 78.2 | * | * | * | 2.5 | 8.8 | 31.0 | 91.1 | 210.7 | 540.2 | 1,041.6 | 2,120.0 |
| 1975 | 81.4 | 8.3 | * | * | 2.3 | 8.4 | 32.9 | 89.4 | 234.2 | 584.0 | 1,097.9 | 2,018.5 |
| 1974 | 88.4 | 12.7 | * | 0.7 | 2.2 | 8.8 | 36.3 | 104.5 | 260.9 | 644.7 | 1,147.2 | 2,304.0 |
| 1973 | 95.3 | 7.9 | * | 0.8 | 2.6 | 10.7 | 42.0 | 116.7 | 286.2 | 686.8 | 1,227.7 | 2,420.8 |
| 1972 | 96.1 | 12.5 | 2.2 | 1.0 | 2.4 | 11.4 | 42.4 | 125.7 | 294.3 | 662.5 | 1,249.8 | 2,438.3 |
| 1971 | 97.7 | * | 2.0 | 0.8 | 3.3 | 10.9 | 43.4 | 122.5 | 300.2 | 707.2 | 1,259.9 | 2,187.2 |
| 1970 | 101.6 | 13.0 | * | 0.7 | 2.9 | 12.8 | 47.0 | 124.2 | 317.7 | 727.4 | 1,361.3 | 1,815.2 |
| 1969 | 106.4 | 8.9 | * | 0.9 | 2.8 | 14.3 | 52.1 | 135.8 | 315.3 | 765.1 | 1,418.1 | 2,377.3 |
| 1968 | 112.5 | 9.4 | * | 0.8 | 3.4 | 14.0 | 52.5 | 139.5 | 352.4 | 816.4 | 1,435.5 | 2,561.0 |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| Cerebrovascular disease 430-438 | | | | | | | | | | | | |
| ALL OTHER, FEMALE | | | | | | | | | | | | |
| 1978 | 73.5 | 11.0 | * | 0.7 | 1.7 | 5.7 | 19.8 | 58.3 | 140.7 | 368.9 | 884.6 | 1,910.1 |
| 1977 | 77.7 | 12.3 | * | * | 2.1 | 5.7 | 23.0 | 64.1 | 149.1 | 407.5 | 933.2 | 1,930.8 |
| 1976 | 81.0 | 8.0 | * | * | 1.8 | 6.6 | 27.8 | 71.2 | 165.5 | 437.9 | 916.5 | 2,036.7 |
| 1975 | 83.8 | 8.9 | * | 0.8 | 1.9 | 7.6 | 28.1 | 74.9 | 176.0 | 476.2 | 966.1 | 2,028.0 |
| 1974 | 91.2 | 9.1 | * | * | 1.8 | 8.4 | 32.5 | 84.1 | 200.6 | 543.4 | 1,028.6 | 2,289.4 |
| 1973 | 99.0 | * | * | 0.9 | 2.3 | 10.6 | 39.5 | 97.1 | 223.9 | 594.5 | 1,123.0 | 2,408.9 |
| 1972 | 100.0 | 8.2 | * | 0.7 | 2.0 | 11.2 | 39.2 | 99.2 | 231.8 | 625.7 | 1,138.9 | 2,360.5 |
| 1971 | 99.5 | 8.2 | * | 1.0 | 2.7 | 12.2 | 40.6 | 106.2 | 236.3 | 608.3 | 1,146.1 | 2,349.3 |
| 1970 | 103.5 | 8.4 | * | 0.7 | 2.9 | 13.1 | 43.8 | 111.4 | 256.3 | 636.1 | 1,268.8 | 2,101.3 |
| 1969 | 106.6 | 9.0 | * | * | 2.3 | 14.3 | 45.4 | 115.0 | 271.3 | 665.3 | 1,298.8 | 2,370.8 |
| 1968 | 112.3 | 8.8 | * | 0.7 | 3.6 | 13.7 | 50.6 | 122.5 | 289.8 | 705.3 | 1,349.4 | 2,577.0 |
| BLACK, BOTH SEXES | | | | | | | | | | | | |
| 1978 | 79.4 | 12.6 | * | 0.7 | 1.6 | 7.1 | 27.2 | 77.3 | 178.1 | 446.4 | 938.9 | 1,951.3 |
| 1977 | 83.2 | 10.7 | * | 0.5 | 1.9 | 7.4 | 30.1 | 81.0 | 188.3 | 476.0 | 992.0 | 2,034.3 |
| 1976 | 86.9 | 7.3 | 1.3 | 0.6 | 2.3 | 8.6 | 33.0 | 88.5 | 200.1 | 514.4 | 994.2 | 2,154.4 |
| 1975 | 89.7 | 9.0 | * | 0.7 | 2.2 | 8.9 | 33.8 | 89.3 | 217.4 | 556.2 | 1,048.5 | 2,122.3 |
| 1974 | 97.1 | 11.6 | 1.2 | 0.6 | 2.1 | 9.9 | 38.4 | 102.5 | 242.9 | 625.3 | 1,107.5 | 2,374.4 |
| 1973 | 104.4 | 8.7 | 1.6 | 0.8 | 2.7 | 11.9 | 44.9 | 115.2 | 268.5 | 674.0 | 1,192.6 | 2,492.1 |
| 1972 | 105.3 | 11.7 | 1.8 | 0.9 | 2.3 | 12.5 | 45.2 | 120.9 | 276.0 | 683.6 | 1,218.1 | 2,485.5 |
| 1971 | 105.3 | 7.5 | 1.7 | 0.9 | 3.1 | 12.8 | 46.1 | 122.8 | 282.4 | 691.3 | 1,234.4 | 2,372.0 |
| 1970 | 110.6 | 10.7 | 1.4 | 0.8 | 3.0 | 14.4 | 50.7 | 127.2 | 305.0 | 720.6 | 1,383.8 | 2,113.6 |
| 1969 | 113.4 | 9.6 | 1.2 | 0.7 | 2.7 | 15.3 | 53.1 | 133.7 | 309.7 | 749.5 | 1,421.1 | 2,485.6 |
| 1968 | 119.4 | 9.3 | 1.3 | 0.8 | 3.6 | 14.7 | 56.4 | 139.4 | 339.5 | 795.9 | 1,442.4 | 2,676.1 |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| Cerebrovascular disease 430-438 | | | | | | | | | | | | |
| BLACK, MALE | | | | | | | | | | | | |
| 1978 | 77.5 | 14.1 | * | * | 1.4 | 7.8 | 32.7 | 92.2 | 211.2 | 518.0 | 977.8 | 1,923.5 |
| 1977 | 80.6 | 8.2 | * | * | 1.7 | 9.1 | 34.8 | 93.3 | 222.0 | 535.1 | 1,040.9 | 2,046.9 |
| 1976 | 84.8 | * | * | * | 2.6 | 10.0 | 35.4 | 100.8 | 227.2 | 579.2 | 1,075.6 | 2,238.3 |
| 1975 | 87.8 | 9.1 | * | * | 2.5 | 9.4 | 36.9 | 98.0 | 253.7 | 626.9 | 1,123.0 | 2,115.2 |
| 1974 | 94.9 | 13.3 | * | * | 2.4 | 10.3 | 40.9 | 115.1 | 280.1 | 692.9 | 1,176.6 | 2,409.3 |
| 1973 | 101.5 | 9.0 | * | 0.8 | 2.7 | 12.0 | 47.0 | 128.0 | 306.7 | 732.5 | 1,251.5 | 2,522.0 |
| 1972 | 102.5 | 14.1 | 2.4 | 1.0 | 2.5 | 12.8 | 47.3 | 137.1 | 315.6 | 710.4 | 1,284.3 | 2,580.0 |
| 1971 | 103.7 | * | 2.2 | 0.9 | 3.6 | 11.9 | 48.3 | 132.3 | 321.5 | 754.0 | 1,306.5 | 2,310.0 |
| 1970 | 108.8 | 12.3 | * | 0.8 | 3.0 | 14.6 | 52.7 | 136.1 | 343.4 | 780.1 | 1,445.7 | 1,963.1 |
| 1969 | 112.7 | 10.0 | * | 0.9 | 2.9 | 15.0 | 56.5 | 146.5 | 339.5 | 815.6 | 1,494.0 | 2,548.6 |
| 1968 | 118.9 | 10.1 | * | 0.9 | 3.3 | 15.1 | 58.0 | 150.1 | 377.9 | 864.9 | 1,507.3 | 2,748.6 |
| BLACK, FEMALE | | | | | | | | | | | | |
| 1978 | 81.1 | 11.1 | * | * | 1.8 | 6.5 | 22.6 | 65.1 | 151.0 | 392.7 | 912.8 | 1,965.7 |
| 1977 | 85.6 | 13.3 | * | * | 2.1 | 6.1 | 26.2 | 70.8 | 160.9 | 431.7 | 962.4 | 2,027.7 |
| 1976 | 88.8 | 8.8 | * | * | 2.0 | 7.3 | 31.0 | 78.3 | 177.5 | 464.7 | 939.5 | 2,110.1 |
| 1975 | 91.4 | 8.8 | * | 0.7 | 1.9 | 8.5 | 31.3 | 81.9 | 187.3 | 502.6 | 998.2 | 2,126.2 |
| 1974 | 99.1 | 10.0 | * | * | 1.9 | 9.5 | 36.2 | 91.9 | 212.2 | 574.4 | 1,060.8 | 2,355.1 |
| 1973 | 107.1 | * | * | 0.8 | 2.6 | 11.8 | 43.2 | 104.2 | 236.8 | 628.8 | 1,156.1 | 2,475.3 |
| 1972 | 107.8 | 9.3 | * | 0.8 | 2.1 | 12.2 | 43.4 | 107.0 | 242.7 | 663.8 | 1,167.1 | 2,431.4 |
| 1971 | 106.8 | 8.8 | * | 1.0 | 2.7 | 13.6 | 44.3 | 114.6 | 249.2 | 640.8 | 1,182.6 | 2,409.0 |
| 1970 | 112.2 | 9.1 | * | 0.8 | 3.0 | 14.3 | 49.1 | 119.4 | 272.4 | 673.5 | 1,338.3 | 2,210.5 |
| 1969 | 114.0 | 9.2 | * | * | 2.5 | 15.5 | 50.2 | 122.7 | 284.5 | 696.9 | 1,361.2 | 2,446.7 |
| 1968 | 119.9 | 8.5 | * | * | 3.9 | 14.4 | 55.1 | 130.1 | 306.3 | 738.9 | 1,394.1 | 2,678.6 |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| CEREBRAL HEMORRHAGE (431) | | | | | | | | | | | | |
| BOTH SEXES | | | | | | | | | | | | |
| 1978 | 10.2 | 3.5 | 0.2 | 0.2 | 0.4 | 0.8 | 3.4 | 9.2 | 18.0 | 37.2 | 84.6 | 160.7 |
| 1977 | 10.8 | 2.9 | * | 0.2 | 0.3 | 0.9 | 3.4 | 9.1 | 18.9 | 40.6 | 94.4 | 179.6 |
| 1976 | 11.4 | 2.0 | * | 0.1 | 0.4 | 1.0 | 3.6 | 9.8 | 19.5 | 44.1 | 99.3 | 198.2 |
| 1975 | 12.5 | 2.1 | 0.2 | 0.1 | 0.4 | 0.9 | 3.6 | 9.9 | 22.0 | 48.9 | 112.0 | 223.9 |
| 1974 | 14.3 | 2.4 | 0.2 | 0.2 | 0.4 | 0.9 | 4.3 | 11.4 | 24.7 | 56.6 | 134.2 | 269.1 |
| 1973 | 16.0 | 1.9 | 0.3 | 0.2 | 0.4 | 1.1 | 4.6 | 12.5 | 28.0 | 63.9 | 152.5 | 321.5 |
| 1972 | 17.6 | 1.4 | 0.3 | 0.2 | 0.4 | 1.1 | 4.7 | 13.6 | 31.5 | 71.8 | 168.8 | 353.6 |
| 1971 | 18.6 | 1.6 | 0.2 | 0.2 | 0.5 | 1.3 | 5.0 | 14.4 | 33.2 | 77.1 | 181.3 | 379.5 |
| 1970 | 20.4 | 2.1 | 0.3 | 0.2 | 0.5 | 1.3 | 5.5 | 14.9 | 35.6 | 85.4 | 206.7 | 399.6 |
| 1969 | 21.5 | 2.2 | 0.2 | 0.2 | 0.5 | 1.4 | 5.4 | 15.3 | 36.4 | 92.3 | 222.9 | 476.1 |
| 1968 | 23.6 | 2.2 | 0.2 | 0.2 | 0.5 | 1.4 | 5.8 | 16.6 | 40.0 | 99.6 | 252.1 | 549.3 |
| MALE | | | | | | | | | | | | |
| 1978 | 9.5 | 3.6 | * | 0.2 | 0.4 | 1.0 | 3.8 | 9.8 | 20.5 | 43.1 | 87.6 | 159.8 |
| 1977 | 10.2 | 2.9 | * | 0.2 | 0.3 | 1.1 | 3.8 | 10.1 | 21.4 | 47.3 | 103.6 | 173.5 |
| 1976 | 10.7 | 2.4 | * | 0.2 | 0.4 | 1.1 | 3.7 | 10.6 | 22.7 | 51.8 | 104.8 | 189.9 |
| 1975 | 11.6 | 2.3 | * | 0.1 | 0.5 | 1.0 | 4.0 | 10.5 | 25.7 | 55.9 | 116.1 | 219.5 |
| 1974 | 13.5 | 3.1 | * | 0.2 | 0.4 | 0.9 | 4.7 | 13.0 | 28.6 | 65.7 | 142.3 | 259.6 |
| 1973 | 15.1 | 1.8 | 0.3 | 0.2 | 0.4 | 1.2 | 4.8 | 13.6 | 32.8 | 74.5 | 164.2 | 301.7 |
| 1972 | 17.0 | 1.4 | 0.3 | 0.2 | 0.5 | 1.4 | 5.3 | 14.9 | 38.5 | 83.6 | 181.6 | 351.0 |
| 1971 | 18.1 | 2.1 | * | 0.2 | 0.5 | 1.3 | 5.5 | 16.4 | 40.4 | 91.3 | 192.1 | 376.3 |
| 1970 | 19.7 | 2.8 | 0.4 | 0.3 | 0.6 | 1.4 | 6.1 | 16.5 | 43.1 | 99.4 | 219.6 | 375.6 |
| 1969 | 20.8 | 2.1 | 0.3 | 0.2 | 0.5 | 1.5 | 5.7 | 16.8 | 43.2 | 106.8 | 239.8 | 467.5 |
| 1968 | 23.2 | 2.0 | * | 0.2 | 0.5 | 1.5 | 6.4 | 18.9 | 48.0 | 116.6 | 271.7 | 548.4 |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| CEREBRAL HEMORRHAGE (431) | | | | | | | | | | | | |
| FEMALE | | | | | | | | | | | | |
| 1978 | 10.9 | 3.4 | * | 0.1 | 0.4 | 0.7 | 2.9 | 8.7 | 15.7 | 32.7 | 82.9 | 161.1 |
| 1977 | 11.5 | 2.9 | * | 0.2 | 0.4 | 0.7 | 3.0 | 8.1 | 16.6 | 35.5 | 88.9 | 182.4 |
| 1976 | 12.1 | 1.5 | * | 0.1 | 0.3 | 1.0 | 3.4 | 8.9 | 16.7 | 38.1 | 96.0 | 202.1 |
| 1975 | 13.3 | 1.9 | * | 0.2 | 0.3 | 0.9 | 3.3 | 9.4 | 18.8 | 43.5 | 109.5 | 226.0 |
| 1974 | 15.1 | 1.7 | * | 0.2 | 0.3 | 0.8 | 3.9 | 9.9 | 21.2 | 49.6 | 129.2 | 273.7 |
| 1973 | 16.9 | 2.0 | * | 0.2 | 0.4 | 1.0 | 4.4 | 11.5 | 23.6 | 55.8 | 145.2 | 331.4 |
| 1972 | 18.1 | 1.4 | * | 0.2 | 0.3 | 0.8 | 4.1 | 12.4 | 25.3 | 62.6 | 160.7 | 354.9 |
| 1971 | 19.1 | 1.2 | * | 0.2 | 0.5 | 1.3 | 4.6 | 12.6 | 26.7 | 66.2 | 174.3 | 381.2 |
| 1970 | 20.9 | 1.4 | * | 0.2 | 0.4 | 1.2 | 4.8 | 13.4 | 28.9 | 74.6 | 198.2 | 413.0 |
| 1969 | 22.1 | 2.3 | * | 0.2 | 0.5 | 1.3 | 5.0 | 13.9 | 30.3 | 81.0 | 211.7 | 480.8 |
| 1968 | 23.9 | 2.4 | * | 0.2 | 0.4 | 1.2 | 5.2 | 14.4 | 32.7 | 86.1 | 238.7 | 550.5 |
| WHITE, BOTH SEXES | | | | | | | | | | | | |
| 1978 | 9.9 | 2.4 | * | 0.2 | 0.4 | 0.6 | 2.3 | 7.2 | 15.7 | 34.9 | 84.3 | 163.6 |
| 1977 | 10.5 | 2.1 | * | 0.2 | 0.3 | 0.6 | 2.4 | 6.9 | 16.4 | 38.2 | 93.9 | 182.8 |
| 1976 | 11.1 | 1.9 | * | 0.1 | 0.3 | 0.8 | 2.4 | 7.5 | 17.0 | 41.3 | 99.7 | 201.4 |
| 1975 | 12.2 | 1.7 | * | 0.1 | 0.4 | 0.7 | 2.5 | 7.8 | 19.4 | 45.5 | 112.7 | 227.5 |
| 1974 | 14.0 | 1.7 | 0.2 | 0.2 | 0.3 | 0.6 | 3.0 | 8.9 | 21.6 | 53.1 | 135.4 | 274.3 |
| 1973 | 15.6 | 1.5 | 0.2 | 0.2 | 0.3 | 0.7 | 2.9 | 9.7 | 24.5 | 59.9 | 154.0 | 328.0 |
| 1972 | 17.2 | 1.0 | 0.2 | 0.1 | 0.3 | 0.7 | 3.2 | 10.7 | 27.4 | 67.5 | 169.6 | 362.1 |
| 1971 | 18.1 | 1.3 | 0.2 | 0.2 | 0.5 | 0.9 | 3.4 | 11.1 | 29.0 | 71.8 | 182.0 | 388.7 |
| 1970 | 19.7 | 1.7 | 0.3 | 0.2 | 0.5 | 0.8 | 3.6 | 11.3 | 30.7 | 79.7 | 206.0 | 412.0 |
| 1969 | 20.8 | 1.7 | 0.2 | 0.2 | 0.5 | 1.0 | 3.3 | 11.6 | 31.3 | 86.2 | 221.9 | 486.6 |
| 1968 | 22.8 | 1.9 | 0.2 | 0.2 | 0.4 | 0.9 | 3.6 | 12.8 | 33.8 | 92.3 | 251.5 | 560.4 |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| CEREBRAL HEMORRHAGE (431) | | | | | | | | | | | | |
| WHITE, MALE | | | | | | | | | | | | |
| 1978 | 8.9 | 2.5 | * | 0.2 | 0.4 | 0.7 | 2.4 | 7.3 | 17.8 | 40.5 | 87.2 | 162.8 |
| 1977 | 9.7 | 2.3 | * | 0.2 | 0.3 | 0.7 | 2.5 | 7.6 | 18.5 | 44.7 | 103.4 | 177.1 |
| 1976 | 10.2 | 2.5 | * | 0.2 | 0.3 | 0.8 | 2.5 | 8.1 | 20.0 | 48.8 | 104.4 | 194.4 |
| 1975 | 11.1 | 1.9 | * | * | 0.5 | 0.7 | 2.6 | 8.0 | 23.0 | 51.8 | 117.6 | 223.7 |
| 1974 | 12.9 | 2.4 | * | 0.2 | 0.3 | 0.6 | 3.3 | 10.2 | 25.1 | 62.0 | 143.9 | 263.5 |
| 1973 | 14.6 | * | * | 0.2 | 0.3 | 0.7 | 3.1 | 10.7 | 29.0 | 70.7 | 166.3 | 308.4 |
| 1972 | 16.4 | * | * | 0.1 | 0.4 | 0.9 | 3.7 | 11.9 | 34.0 | 79.9 | 183.0 | 363.8 |
| 1971 | 17.4 | 1.8 | * | 0.2 | 0.5 | 0.9 | 3.8 | 12.9 | 35.7 | 85.7 | 193.8 | 386.6 |
| 1970 | 18.9 | 2.2 | 0.4 | 0.3 | 0.5 | 0.9 | 4.2 | 12.9 | 37.7 | 93.0 | 220.0 | 392.8 |
| 1969 | 19.9 | 1.8 | * | 0.2 | 0.4 | 1.0 | 3.5 | 13.0 | 38.0 | 100.0 | 239.3 | 477.8 |
| 1968 | 22.2 | 1.7 | * | 0.2 | 0.4 | 1.0 | 3.9 | 14.8 | 41.3 | 109.1 | 272.1 | 561.4 |
| WHITE, FEMALE | | | | | | | | | | | | |
| 1978 | 10.8 | 2.2 | * | * | 0.4 | 0.5 | 2.3 | 7.2 | 13.8 | 30.7 | 82.6 | 163.9 |
| 1977 | 11.3 | 1.8 | * | 0.1 | 0.3 | 0.5 | 2.2 | 6.2 | 14.4 | 33.3 | 88.2 | 185.4 |
| 1976 | 12.0 | * | * | * | 0.3 | 0.8 | 2.3 | 6.9 | 14.4 | 35.5 | 97.0 | 204.8 |
| 1975 | 13.2 | * | * | 0.1 | 0.3 | 0.7 | 2.4 | 7.5 | 16.2 | 40.6 | 109.8 | 229.5 |
| 1974 | 15.0 | * | * | 0.2 | 0.3 | 0.6 | 2.6 | 7.6 | 18.5 | 46.3 | 130.1 | 279.6 |
| 1973 | 16.7 | 1.7 | * | 0.2 | 0.3 | 0.7 | 2.6 | 8.7 | 20.4 | 51.7 | 146.4 | 338.1 |
| 1972 | 17.8 | * | * | 0.1 | 0.2 | 0.5 | 2.8 | 9.6 | 21.6 | 57.9 | 161.1 | 361.2 |
| 1971 | 18.8 | * | * | 0.2 | 0.4 | 0.9 | 3.1 | 9.5 | 22.9 | 61.0 | 174.5 | 389.8 |
| 1970 | 20.5 | * | * | 0.2 | 0.4 | 0.8 | 3.1 | 9.9 | 24.4 | 69.5 | 196.8 | 422.4 |
| 1969 | 21.7 | 1.6 | * | 0.2 | 0.5 | 1.0 | 3.2 | 10.3 | 25.3 | 75.5 | 210.4 | 491.8 |
| 1968 | 23.3 | 2.2 | * | 0.2 | 0.4 | 0.7 | 3.3 | 10.9 | 27.0 | 79.1 | 237.6 | 560.6 |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| CEREBRAL HEMORRHAGE (431) | | | | | | | | | | | | |
| ALL OTHER, BOTH SEXES | | | | | | | | | | | | |
| 1978 | 12.1 | 8.9 | * | * | 0.5 | 2.5 | 10.3 | 24.8 | 38.3 | 58.7 | 87.8 | 128.0 |
| 1977 | 12.9 | 6.8 | * | * | 0.6 | 2.5 | 10.2 | 26.3 | 41.6 | 63.2 | 100.0 | 143.1 |
| 1976 | 13.6 | * | * | * | 0.7 | 2.9 | 11.6 | 28.3 | 42.5 | 70.9 | 95.3 | 160.8 |
| 1975 | 14.5 | 4.2 | * | * | 0.6 | 2.7 | 11.3 | 27.9 | 46.4 | 81.8 | 104.5 | 182.9 |
| 1974 | 16.4 | 5.7 | * | * | 0.7 | 2.8 | 13.4 | 32.4 | 53.9 | 90.8 | 121.6 | 207.4 |
| 1973 | 18.7 | 3.8 | * | * | 0.8 | 3.7 | 16.9 | 36.8 | 61.1 | 103.7 | 135.0 | 244.9 |
| 1972 | 20.4 | * | * | 0.5 | 0.8 | 4.0 | 14.9 | 38.7 | 70.5 | 114.5 | 159.5 | 255.3 |
| 1971 | 22.1 | * | * | 0.4 | 0.9 | 4.0 | 16.8 | 42.9 | 73.2 | 129.6 | 172.5 | 274.2 |
| 1970 | 24.7 | 4.4 | * | * | 1.0 | 4.3 | 18.9 | 46.2 | 82.4 | 141.0 | 216.5 | 272.2 |
| 1969 | 26.3 | 4.7 | * | * | 0.8 | 4.5 | 21.0 | 48.3 | 85.0 | 153.0 | 236.2 | 356.9 |
| 1968 | 29.4 | 3.8 | * | * | 1.1 | 5.0 | 22.5 | 51.2 | 99.6 | 172.8 | 259.4 | 425.5 |
| ALL OTHER, MALE | | | | | | | | | | | | |
| 1978 | 13.1 | 9.0 | * | * | * | 3.0 | 14.3 | 30.8 | 45.8 | 67.8 | 91.4 | 129.3 |
| 1977 | 13.5 | * | * | * | * | 3.5 | 13.1 | 30.7 | 48.2 | 70.9 | 105.4 | 137.5 |
| 1976 | 14.2 | * | * | * | 0.9 | 3.5 | 13.1 | 32.4 | 48.7 | 80.3 | 108.9 | 145.5 |
| 1975 | 14.9 | * | * | * | 0.7 | 3.2 | 14.1 | 32.1 | 51.8 | 94.6 | 101.7 | 177.8 |
| 1974 | 17.1 | * | * | * | 0.8 | 3.2 | 15.0 | 37.5 | 62.7 | 101.3 | 125.8 | 220.0 |
| 1973 | 19.0 | * | * | * | 1.0 | 4.6 | 18.2 | 39.5 | 70.5 | 110.8 | 141.1 | 233.3 |
| 1972 | 20.8 | * | * | * | * | 5.4 | 17.7 | 42.0 | 83.0 | 118.6 | 165.9 | 221.3 |
| 1971 | 23.1 | * | * | * | 1.0 | 4.0 | 19.5 | 47.9 | 85.8 | 144.0 | 173.9 | 274.5 |
| 1970 | 25.6 | * | * | * | 1.2 | 4.7 | 22.0 | 50.0 | 94.2 | 159.3 | 215.0 | 223.7 |
| 1969 | 27.5 | * | * | * | 1.1 | 6.0 | 24.5 | 52.5 | 93.1 | 172.2 | 245.2 | 370.5 |
| 1968 | 30.9 | * | * | * | 1.3 | 5.6 | 26.5 | 57.4 | 113.1 | 189.6 | 267.2 | 422.0 |
| ALL OTHER, FEMALE | | | | | | | | | | | | |
| 1978 | 11.2 | 8.9 | * | * | * | 2.1 | 6.9 | 19.8 | 32.1 | 51.6 | 85.6 | 128.4 |
| 1977 | 12.4 | 7.9 | * | * | 0.8 | 1.7 | 7.7 | 22.6 | 36.1 | 57.1 | 96.1 | 146.2 |
| 1976 | 13.0 | * | * | * | * | 2.4 | 10.3 | 24.8 | 37.2 | 63.4 | 85.9 | 169.4 |
| 1975 | 14.0 | * | * | * | * | 2.2 | 9.0 | 24.3 | 41.9 | 71.6 | 106.2 | 183.9 |
| 1974 | 15.8 | * | * | * | * | 2.5 | 12.1 | 28.1 | 46.3 | 82.4 | 118.7 | 200.0 |
| 1973 | 18.5 | * | * | * | * | 2.9 | 15.8 | 34.4 | 53.2 | 97.9 | 131.1 | 251.9 |
| 1972 | 20.0 | * | * | * | 0.9 | 2.7 | 12.6 | 35.8 | 59.7 | 111.3 | 155.5 | 276.3 |
| 1971 | 21.3 | * | * | * | 0.9 | 3.9 | 14.5 | 38.6 | 62.2 | 117.9 | 171.4 | 274.0 |
| 1970 | 23.9 | * | * | * | 0.9 | 4.0 | 16.4 | 42.9 | 71.9 | 126.0 | 217.6 | 306.4 |
| 1969 | 25.2 | * | * | * | * | 3.2 | 18.1 | 44.7 | 77.8 | 137.2 | 228.3 | 347.7 |
| 1968 | 28.0 | * | * | * | 0.9 | 4.5 | 19.2 | 45.8 | 87.4 | 158.8 | 254.4 | 427.9 |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| CEREBRAL HEMORRHAGE (431) | | | | | | | | | | | | |
| BLACK, BOTH SEXES | | | | | | | | | | | | |
| 1978 | 13.0 | 9.2 | * | * | 0.5 | 2.9 | 11.7 | 27.8 | 40.4 | 59.9 | 88.4 | 126.7 |
| 1977 | 13.9 | 7.4 | * | * | 0.6 | 2.8 | 11.5 | 29.0 | 44.1 | 64.8 | 99.5 | 145.5 |
| 1976 | 14.7 | * | * | * | 0.8 | 3.4 | 13.3 | 31.1 | 45.1 | 72.9 | 94.7 | 169.9 |
| 1975 | 15.6 | 4.6 | * | * | 0.6 | 3.0 | 12.8 | 30.5 | 49.5 | 85.0 | 105.5 | 190.8 |
| 1974 | 17.6 | 6.0 | * | * | 0.8 | 3.2 | 15.1 | 35.3 | 56.6 | 93.6 | 123.4 | 217.4 |
| 1973 | 20.0 | 4.3 | * | * | 0.9 | 4.2 | 18.6 | 39.9 | 63.6 | 108.8 | 135.9 | 255.3 |
| 1972 | 21.8 | * | * | 0.5 | 0.8 | 4.5 | 16.5 | 41.7 | 74.1 | 120.9 | 160.5 | 269.1 |
| 1971 | 23.6 | * | * | 0.4 | 1.0 | 4.5 | 18.7 | 46.1 | 76.6 | 135.8 | 176.4 | 285.0 |
| 1970 | 26.6 | 4.7 | * | * | 1.0 | 4.9 | 21.4 | 49.9 | 87.4 | 150.1 | 225.6 | 293.9 |
| 1969 | 28.0 | 4.8 | * | * | 0.8 | 5.0 | 23.3 | 51.4 | 89.4 | 160.9 | 247.8 | 376.3 |
| 1968 | 31.2 | * | * | * | 1.1 | 5.2 | 24.7 | 55.1 | 105.0 | 180.7 | 269.8 | 440.2 |
| BLACK, MALE | | | | | | | | | | | | |
| 1978 | 14.1 | 9.7 | * | * | * | 3.5 | 16.2 | 35.1 | 48.8 | 69.5 | 89.5 | 131.4 |
| 1977 | 14.5 | * | * | * | * | 4.1 | 14.8 | 34.4 | 51.4 | 72.0 | 106.0 | 134.7 |
| 1976 | 15.2 | * | * | * | 1.0 | 4.2 | 15.1 | 35.7 | 52.1 | 81.3 | 108.5 | 155.3 |
| 1975 | 16.0 | * | * | * | * | 3.6 | 15.8 | 35.5 | 55.4 | 98.4 | 100.5 | 189.1 |
| 1974 | 18.2 | * | * | * | 0.9 | 3.7 | 17.0 | 40.8 | 66.3 | 105.1 | 125.4 | 241.9 |
| 1973 | 20.4 | * | * | * | 1.0 | 5.3 | 20.4 | 43.7 | 73.9 | 117.3 | 144.1 | 251.2 |
| 1972 | 22.0 | * | * | * | * | 6.3 | 19.6 | 45.5 | 87.0 | 127.1 | 164.5 | 230.0 |
| 1971 | 24.3 | * | * | * | 1.1 | 4.5 | 21.8 | 51.0 | 89.8 | 150.3 | 179.0 | 292.5 |
| 1970 | 27.5 | * | * | * | 1.2 | 5.5 | 24.7 | 54.9 | 101.3 | 170.7 | 224.7 | 247.8 |
| 1969 | 29.2 | * | * | * | 1.1 | 6.7 | 27.1 | 56.3 | 99.3 | 182.4 | 258.1 | 400.0 |
| 1968 | 32.5 | * | * | * | 1.3 | 6.0 | 29.1 | 62.3 | 120.0 | 198.9 | 279.4 | 448.6 |
| BLACK, FEMALE | | | | | | | | | | | | |
| 1978 | 12.1 | 8.6 | * | * | * | 2.5 | 7.9 | 21.8 | 33.5 | 52.8 | 87.7 | 124.2 |
| 1977 | 13.4 | 8.3 | * | * | 0.9 | 1.7 | 8.8 | 24.6 | 38.1 | 59.4 | 95.5 | 151.1 |
| 1976 | 14.2 | * | * | * | * | 2.7 | 11.8 | 27.4 | 39.4 | 66.5 | 85.4 | 177.5 |
| 1975 | 15.2 | * | * | * | * | 2.5 | 10.3 | 26.3 | 44.6 | 74.8 | 108.8 | 191.7 |
| 1974 | 17.0 | * | * | * | * | 2.7 | 13.5 | 30.7 | 48.6 | 84.8 | 122.0 | 203.8 |
| 1973 | 19.7 | * | * | * | * | 3.2 | 17.1 | 36.7 | 55.1 | 102.3 | 130.7 | 257.5 |
| 1972 | 21.5 | * | * | * | 0.9 | 2.9 | 14.0 | 38.5 | 63.2 | 116.3 | 157.1 | 291.4 |
| 1971 | 22.9 | * | * | * | 0.9 | 4.5 | 16.2 | 41.9 | 65.4 | 124.1 | 174.5 | 280.6 |
| 1970 | 25.8 | * | * | * | * | 4.4 | 18.6 | 45.6 | 75.6 | 133.7 | 226.2 | 323.6 |
| 1969 | 27.0 | * | * | * | * | 3.6 | 20.2 | 47.2 | 81.0 | 143.7 | 239.2 | 361.7 |
| 1968 | 29.9 | * | * | * | 1.0 | 4.5 | 21.0 | 48.9 | 92.0 | 165.7 | 262.6 | 442.9 |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| CEREBRAL THROMBOSIS (433) | | | | | | | | | | | | |
| BOTH SEXES | | | | | | | | | | | | |
| 1978 | 17.4 | * | * | * | 0.1 | 0.1 | 0.6 | 2.8 | 11.9 | 51.2 | 203.1 | 599.7 |
| 1977 | 19.1 | * | * | * | * | 0.2 | 0.7 | 3.3 | 13.7 | 57.8 | 228.1 | 659.8 |
| 1976 | 20.6 | * | * | * | 0.1 | 0.1 | 0.7 | 3.5 | 14.8 | 64.5 | 248.7 | 725.8 |
| 1975 | 22.5 | * | * | * | 0.1 | 0.2 | 0.7 | 3.6 | 16.7 | 72.5 | 283.7 | 778.1 |
| 1974 | 25.1 | * | * | * | 0.1 | 0.2 | 0.8 | 3.9 | 18.5 | 83.8 | 322.9 | 902.6 |
| 1973 | 26.8 | * | * | * | 0.1 | 0.2 | 0.8 | 4.4 | 20.5 | 92.3 | 350.2 | 978.2 |
| 1972 | 27.5 | * | * | * | 0.1 | 0.2 | 0.9 | 4.0 | 21.9 | 97.7 | 365.4 | 1,000.9 |
| 1971 | 27.7 | * | * | * | 0.1 | 0.2 | 0.9 | 4.7 | 22.0 | 99.5 | 375.3 | 1,011.5 |
| 1970 | 28.5 | * | * | * | 0.1 | 0.2 | 1.0 | 4.6 | 23.4 | 108.1 | 390.9 | 975.8 |
| 1969 | 29.2 | * | * | * | 0.1 | 0.2 | 0.9 | 4.5 | 23.3 | 112.6 | 399.7 | 1,145.9 |
| 1968 | 30.3 | * | * | * | 0.1 | 0.2 | 0.9 | 4.8 | 24.7 | 116.6 | 427.2 | 1,196.9 |
| MALE | | | | | | | | | | | | |
| 1978 | 14.8 | * | * | * | * | * | 0.6 | 3.7 | 15.1 | 63.1 | 219.6 | 574.4 |
| 1977 | 16.5 | * | * | * | * | 0.1 | 0.8 | 4.1 | 17.4 | 70.6 | 252.3 | 644.7 |
| 1976 | 17.9 | * | * | * | * | * | 0.8 | 4.3 | 18.6 | 78.8 | 274.8 | 713.2 |
| 1975 | 19.8 | * | * | * | * | 0.1 | 0.6 | 4.4 | 21.1 | 89.6 | 314.6 | 758.8 |
| 1974 | 22.2 | * | * | * | * | 0.2 | 0.9 | 4.8 | 23.5 | 103.1 | 352.5 | 888.8 |
| 1973 | 23.7 | * | * | * | * | 0.2 | 0.9 | 5.3 | 25.7 | 111.6 | 378.9 | 952.2 |
| 1972 | 24.8 | * | * | * | * | * | 1.1 | 5.0 | 27.9 | 119.6 | 401.1 | 967.8 |
| 1971 | 25.2 | * | * | * | 0.1 | 0.2 | 1.0 | 5.9 | 27.7 | 120.6 | 412.3 | 995.1 |
| 1970 | 26.0 | * | * | * | * | 0.2 | 1.2 | 5.5 | 29.4 | 127.8 | 420.9 | 937.4 |
| 1969 | 26.9 | * | * | * | * | 0.2 | 1.0 | 5.7 | 29.5 | 134.7 | 433.1 | 1,123.7 |
| 1968 | 28.0 | * | * | * | * | * | 1.0 | 5.9 | 30.8 | 139.5 | 460.7 | 1,161.4 |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| CEREBRAL THROMBOSIS (433) | | | | | | | | | | | | |
| FEMALE | | | | | | | | | | | | |
| 1978 | 19.9 | * | * | * | * | 0.1 | 0.6 | 2.1 | 9.0 | 42.1 | 193.3 | 611.2 |
| 1977 | 21.6 | * | * | * | * | 0.2 | 0.5 | 2.5 | 10.3 | 48.0 | 213.5 | 666.7 |
| 1976 | 23.1 | * | * | * | * | * | 0.6 | 2.7 | 11.5 | 53.5 | 233.0 | 731.8 |
| 1975 | 25.0 | * | * | * | 0.1 | 0.2 | 0.7 | 2.7 | 12.7 | 59.4 | 264.9 | 787.5 |
| 1974 | 27.8 | * | * | * | * | 0.2 | 0.7 | 3.1 | 14.1 | 69.0 | 304.7 | 909.4 |
| 1973 | 29.8 | * | * | * | 0.1 | 0.2 | 0.7 | 3.6 | 15.9 | 77.5 | 332.4 | 991.3 |
| 1972 | 30.0 | * | * | * | * | 0.2 | 0.8 | 3.1 | 16.6 | 80.8 | 342.6 | 1,017.8 |
| 1971 | 30.0 | * | * | * | * | 0.2 | 0.8 | 3.6 | 16.9 | 83.1 | 351.2 | 1,020.1 |
| 1970 | 30.8 | * | * | * | * | 0.2 | 0.9 | 3.6 | 17.9 | 92.7 | 371.0 | 997.3 |
| 1969 | 31.3 | * | * | * | * | 0.2 | 0.8 | 3.5 | 17.7 | 95.4 | 377.4 | 1,157.9 |
| 1968 | 32.4 | * | * | * | * | 0.2 | 0.8 | 3.8 | 19.1 | 98.6 | 404.2 | 1,217.9 |
| WHITE, BOTH SEXES | | | | | | | | | | | | |
| 1978 | 18.1 | * | * | * | 0.1 | 0.1 | 0.5 | 2.3 | 10.1 | 48.3 | 203.9 | 614.8 |
| 1977 | 19.8 | * | * | * | * | 0.1 | 0.5 | 2.6 | 11.9 | 54.5 | 228.6 | 678.3 |
| 1976 | 21.4 | * | * | * | * | 0.1 | 0.5 | 2.7 | 13.2 | 60.8 | 251.4 | 744.8 |
| 1975 | 23.4 | * | * | * | 0.1 | 0.2 | 0.5 | 2.8 | 14.5 | 68.3 | 287.0 | 803.0 |
| 1974 | 26.0 | * | * | * | * | 0.2 | 0.6 | 3.1 | 15.9 | 78.8 | 327.6 | 925.8 |
| 1973 | 27.6 | * | * | * | 0.1 | 0.1 | 0.6 | 3.4 | 17.5 | 85.5 | 353.3 | 1,004.7 |
| 1972 | 28.4 | * | * | * | * | 0.1 | 0.7 | 3.1 | 19.2 | 92.7 | 368.6 | 1,027.1 |
| 1971 | 28.5 | * | * | * | 0.1 | 0.1 | 0.6 | 3.7 | 18.7 | 93.1 | 378.5 | 1,043.2 |
| 1970 | 29.2 | * | * | * | 0.1 | 0.1 | 0.8 | 3.6 | 19.7 | 101.3 | 393.3 | 1,011.8 |
| 1969 | 29.9 | * | * | * | * | 0.2 | 0.6 | 3.5 | 19.7 | 105.0 | 400.3 | 1,185.8 |
| 1968 | 30.9 | * | * | * | * | 0.1 | 0.7 | 3.7 | 20.7 | 108.5 | 427.6 | 1,235.4 |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| CEREBRAL THROMBOSIS (433) | | | | | | | | | | | | |
| WHITE, MALE | | | | | | | | | | | | |
| 1978 | 15.0 | * | * | * | * | * | 0.5 | 2.9 | 12.9 | 59.5 | 221.9 | 591.1 |
| 1977 | 16.9 | * | * | * | * | * | 0.5 | 3.2 | 15.2 | 67.8 | 253.6 | 667.1 |
| 1976 | 18.4 | * | * | * | * | * | 0.5 | 3.3 | 16.8 | 75.3 | 278.7 | 733.2 |
| 1975 | 20.3 | * | * | * | * | * | 0.5 | 3.6 | 18.7 | 85.4 | 319.4 | 787.0 |
| 1974 | 22.8 | * | * | * | * | 0.2 | 0.6 | 3.9 | 20.6 | 99.2 | 358.5 | 914.6 |
| 1973 | 24.0 | * | * | * | * | * | 0.6 | 4.0 | 22.3 | 104.8 | 384.6 | 979.0 |
| 1972 | 25.4 | * | * | * | * | * | 0.8 | 3.9 | 25.4 | 116.4 | 407.2 | 989.9 |
| 1971 | 25.8 | * | * | * | * | * | 0.6 | 4.8 | 24.3 | 115.1 | 417.8 | 1,035.4 |
| 1970 | 26.4 | * | * | * | * | * | 0.9 | 4.5 | 25.7 | 122.2 | 423.9 | 978.5 |
| 1969 | 27.2 | * | * | * | * | * | 0.7 | 4.5 | 25.7 | 127.5 | 434.7 | 1,170.1 |
| 1968 | 28.3 | * | * | * | * | * | 0.8 | 4.7 | 26.9 | 132.3 | 461.6 | 1,205.5 |
| WHITE, FEMALE | | | | | | | | | | | | |
| 1978 | 21.0 | * | * | * | * | * | 0.5 | 1.7 | 7.6 | 39.7 | 193.3 | 625.4 |
| 1977 | 22.6 | * | * | * | * | 0.2 | 0.4 | 2.1 | 8.9 | 44.4 | 213.9 | 683.4 |
| 1976 | 24.3 | * | * | * | * | * | 0.4 | 2.2 | 9.9 | 49.7 | 235.2 | 750.8 |
| 1975 | 26.2 | * | * | * | * | 0.2 | 0.6 | 2.0 | 10.7 | 55.2 | 267.6 | 811.4 |
| 1974 | 29.1 | * | * | * | * | 0.2 | 0.6 | 2.4 | 11.6 | 63.2 | 308.6 | 931.3 |
| 1973 | 31.0 | * | * | * | * | 0.2 | 0.6 | 2.8 | 13.3 | 70.8 | 334.0 | 1,018.4 |
| 1972 | 31.2 | * | * | * | * | 0.2 | 0.6 | 2.4 | 13.7 | 74.4 | 344.1 | 1,045.8 |
| 1971 | 31.1 | * | * | * | * | * | 0.7 | 2.7 | 13.7 | 76.2 | 353.3 | 1,047.1 |
| 1970 | 31.9 | * | * | * | * | * | 0.7 | 2.7 | 14.4 | 85.3 | 373.2 | 1,030.1 |
| 1969 | 32.3 | * | * | * | * | * | 0.6 | 2.6 | 14.2 | 87.5 | 377.4 | 1,195.7 |
| 1968 | 33.3 | * | * | * | * | * | 0.7 | 2.8 | 15.2 | 89.8 | 404.5 | 1,253.2 |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| CEREBRAL THROMBOSIS (433) | | | | | | | | | | | | |
| ALL OTHER, BOTH SEXES | | | | | | | | | | | | |
| 1978 | 13.2 | * | * | * | * | * | 1.5 | 7.2 | 27.8 | 78.5 | 195.1 | 426.2 |
| 1977 | 14.6 | * | * | * | * | * | 1.7 | 8.3 | 29.9 | 88.9 | 222.2 | 447.5 |
| 1976 | 15.4 | * | * | * | * | * | 2.5 | 9.7 | 29.7 | 99.7 | 221.5 | 504.6 |
| 1975 | 16.7 | * | * | * | * | * | 1.7 | 9.7 | 37.0 | 113.7 | 248.6 | 492.5 |
| 1974 | 19.0 | * | * | * | * | * | 2.0 | 10.5 | 43.3 | 133.2 | 271.7 | 632.6 |
| 1973 | 21.6 | * | * | * | * | * | 2.3 | 13.1 | 49.2 | 159.3 | 314.7 | 670.1 |
| 1972 | 21.1 | * | * | * | * | * | 2.8 | 11.6 | 48.1 | 147.7 | 329.1 | 699.2 |
| 1971 | 22.0 | * | * | * | * | 0.6 | 2.8 | 13.1 | 53.5 | 162.0 | 336.3 | 650.8 |
| 1970 | 23.1 | * | * | * | * | 0.7 | 2.7 | 13.4 | 57.6 | 173.8 | 360.4 | 605.6 |
| 1969 | 24.3 | * | * | * | * | * | 2.8 | 13.4 | 58.2 | 188.2 | 393.1 | 696.3 |
| 1968 | 25.8 | * | * | * | * | * | 2.5 | 14.6 | 62.7 | 198.4 | 422.1 | 766.7 |
| ALL OTHER, MALE | | | | | | | | | | | | |
| 1978 | 13.0 | * | * | * | * | * | 1.5 | 10.0 | 35.7 | 96.4 | 197.4 | 403.4 |
| 1977 | 14.3 | * | * | * | * | * | 2.7 | 11.7 | 38.0 | 97.1 | 239.8 | 416.1 |
| 1976 | 15.1 | * | * | * | * | * | 3.0 | 12.8 | 35.2 | 111.3 | 239.3 | 512.7 |
| 1975 | 16.5 | * | * | * | * | * | 1.6 | 11.5 | 44.5 | 129.9 | 266.9 | 475.9 |
| 1974 | 18.3 | * | * | * | * | * | 2.6 | 13.0 | 50.8 | 140.8 | 290.0 | 624.0 |
| 1973 | 21.3 | * | * | * | * | * | 3.0 | 16.6 | 59.4 | 176.5 | 317.9 | 679.2 |
| 1972 | 20.1 | * | * | * | * | * | 3.3 | 15.1 | 52.2 | 150.3 | 335.5 | 744.7 |
| 1971 | 21.5 | * | * | * | * | * | 3.7 | 15.9 | 60.4 | 173.3 | 353.6 | 597.9 |
| 1970 | 22.8 | * | * | * | * | * | 3.4 | 15.4 | 64.8 | 181.2 | 385.4 | 575.6 |
| 1969 | 24.4 | * | * | * | * | * | 3.5 | 16.0 | 65.7 | 204.1 | 413.8 | 686.4 |
| 1968 | 25.6 | * | * | * | * | * | 3.1 | 16.8 | 69.5 | 209.2 | 450.0 | 731.7 |
| ALL OTHER, FEMALE | | | | | | | | | | | | |
| 1978 | 13.4 | * | * | * | * | * | 1.5 | 4.9 | 21.2 | 64.7 | 194.1 | 442.2 |
| 1977 | 14.9 | * | * | * | * | * | * | 5.4 | 23.2 | 82.4 | 209.7 | 464.4 |
| 1976 | 15.7 | * | * | * | * | * | 2.0 | 7.1 | 25.1 | 90.5 | 209.3 | 500.0 |
| 1975 | 17.0 | * | * | * | * | * | 1.8 | 8.2 | 30.7 | 100.8 | 235.3 | 496.8 |
| 1974 | 19.6 | * | * | * | * | * | 1.5 | 8.4 | 37.0 | 127.1 | 259.0 | 637.6 |
| 1973 | 21.8 | * | * | * | * | * | 1.7 | 10.1 | 40.4 | 145.3 | 313.5 | 664.6 |
| 1972 | 22.1 | * | * | * | * | * | 2.4 | 8.6 | 44.4 | 145.9 | 325.6 | 671.1 |
| 1971 | 22.5 | * | * | * | * | * | 2.1 | 10.7 | 47.3 | 152.6 | 323.6 | 684.9 |
| 1970 | 23.3 | * | * | * | * | * | 2.1 | 11.6 | 51.3 | 167.7 | 341.3 | 626.7 |
| 1969 | 24.1 | * | * | * | * | * | 2.3 | 11.0 | 51.6 | 175.2 | 375.7 | 703.1 |
| 1968 | 25.9 | * | * | * | * | * | 2.0 | 12.8 | 56.7 | 189.4 | 402.1 | 790.2 |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| CEREBRAL THROMBOSIS (433) | | | | | | | | | | | | |
| BLACK, BOTH SEXES | | | | | | | | | | | | |
| 1978 | 14.5 | * | * | * | * | * | 1.6 | 8.0 | 30.5 | 85.2 | 200.7 | 434.0 |
| 1977 | 16.0 | * | * | * | * | * | 2.0 | 9.5 | 32.4 | 94.8 | 230.7 | 461.5 |
| 1976 | 16.8 | * | * | * | * | * | 2.7 | 10.7 | 32.1 | 106.2 | 228.0 | 533.8 |
| 1975 | 18.1 | * | * | * | * | * | 2.0 | 10.9 | 40.3 | 119.9 | 254.2 | 509.2 |
| 1974 | 20.6 | * | * | * | * | * | 2.3 | 11.7 | 46.6 | 142.8 | 280.5 | 647.9 |
| 1973 | 23.3 | * | * | * | * | * | 2.6 | 14.6 | 53.0 | 170.0 | 323.0 | 692.1 |
| 1972 | 22.8 | * | * | * | * | * | 3.2 | 12.8 | 52.0 | 157.8 | 343.3 | 740.0 |
| 1971 | 23.6 | * | * | * | * | * | 3.1 | 14.3 | 57.5 | 172.0 | 349.2 | 674.8 |
| 1970 | 25.0 | * | * | * | * | 0.8 | 3.0 | 14.8 | 62.2 | 185.8 | 384.1 | 650.9 |
| 1969 | 25.8 | * | * | * | * | * | 3.1 | 14.5 | 62.5 | 198.4 | 412.7 | 728.9 |
| 1968 | 27.4 | * | * | * | * | * | 2.8 | 15.6 | 67.2 | 210.0 | 440.3 | 804.3 |
| BLACK, MALE | | | | | | | | | | | | |
| 1978 | 14.1 | * | * | * | * | * | * | 11.1 | 39.5 | 106.2 | 200.8 | 392.2 |
| 1977 | 15.6 | * | * | * | * | * | 3.3 | 13.3 | 41.5 | 105.2 | 248.9 | 416.3 |
| 1976 | 16.4 | * | * | * | * | * | 3.4 | 14.2 | 38.1 | 119.5 | 244.9 | 563.8 |
| 1975 | 17.8 | * | * | * | * | * | 1.9 | 12.8 | 48.9 | 137.6 | 274.3 | 497.8 |
| 1974 | 19.7 | * | * | * | * | * | 2.9 | 14.8 | 54.7 | 154.0 | 299.5 | 634.9 |
| 1973 | 22.6 | * | * | * | * | * | 3.4 | 18.6 | 64.2 | 189.8 | 322.1 | 692.7 |
| 1972 | 21.5 | * | * | * | * | * | 3.7 | 16.8 | 57.7 | 160.3 | 347.2 | 810.0 |
| 1971 | 23.0 | * | * | * | * | * | 4.1 | 17.7 | 66.0 | 186.1 | 371.5 | 625.0 |
| 1970 | 24.5 | * | * | * | * | * | 3.9 | 17.2 | 70.5 | 195.9 | 413.9 | 619.6 |
| 1969 | 25.8 | * | * | * | * | * | 3.9 | 17.6 | 71.7 | 217.4 | 439.5 | 721.6 |
| 1968 | 27.0 | * | * | * | * | * | 3.5 | 17.9 | 75.4 | 223.6 | 472.1 | 780.0 |
| BLACK, FEMALE | | | | | | | | | | | | |
| 1978 | 14.8 | * | * | * | * | * | 1.6 | 5.6 | 23.1 | 69.4 | 200.6 | 455.6 |
| 1977 | 16.5 | * | * | * | * | * | * | 6.3 | 25.0 | 87.0 | 219.2 | 485.1 |
| 1976 | 17.2 | * | * | * | * | * | 2.1 | 7.7 | 27.1 | 96.1 | 216.6 | 518.0 |
| 1975 | 18.5 | * | * | * | * | * | 2.0 | 9.3 | 33.1 | 106.5 | 240.5 | 515.5 |
| 1974 | 21.3 | * | * | * | * | * | 1.7 | 9.1 | 39.8 | 134.5 | 267.6 | 655.1 |
| 1973 | 23.9 | * | * | * | * | * | 2.0 | 11.2 | 43.7 | 154.8 | 324.7 | 691.8 |
| 1972 | 24.1 | * | * | * | * | * | 2.7 | 9.5 | 47.3 | 156.1 | 339.3 | 700.0 |
| 1971 | 24.1 | * | * | * | * | * | 2.3 | 11.4 | 50.2 | 160.5 | 333.2 | 704.5 |
| 1970 | 25.5 | * | * | * | * | * | 2.3 | 12.8 | 55.1 | 177.9 | 362.3 | 671.0 |
| 1969 | 25.8 | * | * | * | * | * | 2.4 | 11.8 | 54.6 | 183.4 | 391.2 | 733.3 |
| 1968 | 27.8 | * | * | * | * | * | 2.2 | 13.5 | 60.1 | 198.6 | 416.7 | 833.9 |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| CEREBRAL EMBOLISM (434) | | | | | | | | | | | | |
| BOTH SEXES | | | | | | | | | | | | |
| 1978 | 0.3 | * | * | * | * | * | * | 0.1 | 0.4 | 1.2 | 3.6 | 8.4 |
| 1977 | 0.3 | * | * | * | * | * | * | 0.1 | 0.5 | 1.2 | 3.5 | 7.9 |
| 1976 | 0.4 | * | * | * | * | * | * | 0.1 | 0.5 | 1.5 | 4.5 | 7.8 |
| 1975 | 0.4 | * | * | * | * | * | * | 0.1 | 0.6 | 1.3 | 4.2 | 8.9 |
| 1974 | 0.4 | * | * | * | * | * | * | 0.2 | 0.6 | 1.7 | 4.3 | 10.7 |
| 1973 | 0.5 | * | * | * | * | * | * | 0.2 | 0.7 | 1.7 | 5.3 | 11.3 |
| 1972 | 0.5 | * | * | * | * | * | * | 0.2 | 0.7 | 2.1 | 5.0 | 10.4 |
| 1971 | 0.5 | * | * | * | * | * | * | 0.2 | 0.6 | 1.9 | 5.2 | 12.0 |
| 1970 | 0.4 | * | * | * | * | * | * | 0.2 | 0.6 | 1.9 | 5.0 | 10.7 |
| 1969 | 0.5 | * | * | * | * | * | * | 0.2 | 0.7 | 2.2 | 6.1 | 13.2 |
| 1968 | 0.6 | * | * | * | * | * | * | 0.2 | 0.8 | 2.4 | 6.8 | 16.8 |
| MALE | | | | | | | | | | | | |
| 1978 | 0.3 | * | * | * | * | * | * | * | 0.5 | 1.5 | 3.9 | 7.4 |
| 1977 | 0.3 | * | * | * | * | * | * | * | 0.5 | 1.2 | 3.2 | 8.3 |
| 1976 | 0.3 | * | * | * | * | * | * | * | 0.6 | 1.6 | 4.8 | 7.3 |
| 1975 | 0.3 | * | * | * | * | * | * | * | 0.7 | 1.5 | 4.4 | 10.3 |
| 1974 | 0.4 | * | * | * | * | * | * | 0.2 | 0.7 | 2.1 | 4.1 | 12.0 |
| 1973 | 0.4 | * | * | * | * | * | * | 0.2 | 0.7 | 2.0 | 6.1 | 10.6 |
| 1972 | 0.4 | * | * | * | * | * | * | 0.2 | 0.8 | 2.5 | 5.4 | 10.3 |
| 1971 | 0.5 | * | * | * | * | * | * | 0.2 | 0.7 | 2.4 | 6.5 | 12.2 |
| 1970 | 0.4 | * | * | * | * | * | * | 0.3 | 0.7 | 2.0 | 6.1 | 10.1 |
| 1969 | 0.4 | * | * | * | * | * | * | 0.2 | 0.9 | 2.4 | 5.0 | 12.6 |
| 1968 | 0.5 | * | * | * | * | * | * | 0.3 | 1.0 | 2.8 | 7.0 | 17.0 |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| CEREBRAL EMBOLISM (434) | | | | | | | | | | | | |
| FEMALE | | | | | | | | | | | | |
| 1978 | 0.4 | * | * | * | * | * | * | * | 0.3 | 0.9 | 3.4 | 8.9 |
| 1977 | 0.4 | * | * | * | * | * | * | * | 0.4 | 1.1 | 3.7 | 7.8 |
| 1976 | 0.4 | * | * | * | * | * | * | * | 0.3 | 1.5 | 4.3 | 8.1 |
| 1975 | 0.4 | * | * | * | * | * | * | * | 0.6 | 1.2 | 4.0 | 8.2 |
| 1974 | 0.5 | * | * | * | * | * | * | 0.2 | 0.6 | 1.5 | 4.4 | 10.1 |
| 1973 | 0.5 | * | * | * | * | * | * | 0.2 | 0.7 | 1.5 | 4.8 | 11.6 |
| 1972 | 0.5 | * | * | * | * | * | * | 0.2 | 0.5 | 1.9 | 4.8 | 10.4 |
| 1971 | 0.5 | * | * | * | * | * | * | 0.2 | 0.5 | 1.5 | 4.4 | 12.0 |
| 1970 | 0.5 | * | * | * | * | * | * | 0.2 | 0.5 | 1.8 | 4.3 | 10.9 |
| 1969 | 0.6 | * | * | * | * | * | * | 0.3 | 0.5 | 2.0 | 6.8 | 13.4 |
| 1968 | 0.6 | * | * | * | * | * | * | 0.2 | 0.7 | 2.1 | 6.7 | 16.6 |
| WHITE, BOTH SEXES | | | | | | | | | | | | |
| 1978 | 0.4 | * | * | * | * | * | * | 0.1 | 0.4 | 1.2 | 3.8 | 8.8 |
| 1977 | 0.4 | * | * | * | * | * | * | 0.1 | 0.4 | 1.2 | 3.6 | 8.1 |
| 1976 | 0.4 | * | * | * | * | * | * | 0.1 | 0.5 | 1.5 | 4.7 | 8.3 |
| 1975 | 0.4 | * | * | * | * | * | * | 0.1 | 0.6 | 1.4 | 4.4 | 9.3 |
| 1974 | 0.4 | * | * | * | * | * | * | 0.2 | 0.6 | 1.7 | 4.4 | 10.9 |
| 1973 | 0.5 | * | * | * | * | * | * | 0.2 | 0.7 | 1.7 | 5.4 | 11.9 |
| 1972 | 0.5 | * | * | * | * | * | * | 0.2 | 0.6 | 2.1 | 5.2 | 10.6 |
| 1971 | 0.5 | * | * | * | * | * | * | 0.2 | 0.6 | 1.9 | 5.3 | 12.4 |
| 1970 | 0.5 | * | * | * | * | * | * | 0.2 | 0.6 | 2.0 | 5.1 | 11.2 |
| 1969 | 0.5 | * | * | * | * | * | * | 0.2 | 0.7 | 2.2 | 6.1 | 13.6 |
| 1968 | 0.6 | * | * | * | * | * | * | 0.2 | 0.8 | 2.3 | 7.0 | 17.3 |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| CEREBRAL EMBOLISM (434) | | | | | | | | | | | | |
| WHITE, MALE | | | | | | | | | | | | |
| 1978 | 0.3 | * | * | * | * | * | * | * | 0.5 | 1.6 | 4.2 | 7.4 |
| 1977 | 0.3 | * | * | * | * | * | * | * | 0.5 | 1.2 | 3.4 | 8.4 |
| 1976 | 0.3 | * | * | * | * | * | * | * | 0.6 | 1.5 | 5.1 | 7.6 |
| 1975 | 0.4 | * | * | * | * | * | * | * | 0.7 | 1.6 | 4.5 | 10.6 |
| 1974 | 0.4 | * | * | * | * | * | * | * | 0.7 | 2.0 | 4.2 | 12.6 |
| 1973 | 0.4 | * | * | * | * | * | * | * | 0.7 | 2.0 | 6.1 | 11.0 |
| 1972 | 0.5 | * | * | * | * | * | * | 0.2 | 0.8 | 2.6 | 5.8 | 10.5 |
| 1971 | 0.5 | * | * | * | * | * | * | 0.2 | 0.7 | 2.5 | 6.6 | 12.3 |
| 1970 | 0.4 | * | * | * | * | * | * | 0.2 | 0.7 | 2.1 | 6.4 | 10.7 |
| 1969 | 0.4 | * | * | * | * | * | * | * | 0.9 | 2.5 | 4.9 | 12.8 |
| 1968 | 0.6 | * | * | * | * | * | * | 0.3 | 1.0 | 2.8 | 7.3 | 17.3 |
| WHITE, FEMALE | | | | | | | | | | | | |
| 1978 | 0.4 | * | * | * | * | * | * | * | 0.3 | 1.0 | 3.6 | 9.4 |
| 1977 | 0.4 | * | * | * | * | * | * | * | 0.4 | 1.2 | 3.8 | 7.9 |
| 1976 | 0.5 | * | * | * | * | * | * | * | 0.3 | 1.5 | 4.4 | 8.6 |
| 1975 | 0.4 | * | * | * | * | * | * | * | 0.5 | 1.2 | 4.3 | 8.7 |
| 1974 | 0.5 | * | * | * | * | * | * | 0.2 | 0.5 | 1.5 | 4.6 | 10.1 |
| 1973 | 0.5 | * | * | * | * | * | * | 0.2 | 0.6 | 1.4 | 5.0 | 12.3 |
| 1972 | 0.5 | * | * | * | * | * | * | 0.2 | 0.5 | 1.8 | 4.9 | 10.6 |
| 1971 | 0.5 | * | * | * | * | * | * | 0.2 | 0.4 | 1.5 | 4.5 | 12.5 |
| 1970 | 0.5 | * | * | * | * | * | * | * | 0.5 | 1.9 | 4.3 | 11.5 |
| 1969 | 0.6 | * | * | * | * | * | * | 0.2 | 0.5 | 1.9 | 7.0 | 14.0 |
| 1968 | 0.6 | * | * | * | * | * | * | * | 0.6 | 1.9 | 6.7 | 17.3 |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| CEREBRAL EMBOLISM (434) | | | | | | | | | | | | |
| ALL OTHER, BOTH SEXES | | | | | | | | | | | | |
| 1978 | 0.2 | * | * | * | * | * | * | * | * | * | * | * |
| 1977 | 0.2 | * | * | * | * | * | * | * | * | * | * | * |
| 1976 | 0.2 | * | * | * | * | * | * | * | * | 1.8 | * | * |
| 1975 | 0.2 | * | * | * | * | * | * | * | * | * | * | * |
| 1974 | 0.3 | * | * | * | * | * | * | * | * | 2.0 | * | * |
| 1973 | 0.3 | * | * | * | * | * | * | * | * | 2.2 | 3.7 | * |
| 1972 | 0.3 | * | * | * | * | * | * | * | * | 2.4 | * | * |
| 1971 | 0.3 | * | * | * | * | * | * | * | 1.2 | 1.9 | * | * |
| 1970 | 0.2 | * | * | * | * | * | * | * | * | * | * | * |
| 1969 | 0.3 | * | * | * | * | * | * | * | * | 2.0 | 5.3 | * |
| 1968 | 0.4 | * | * | * | * | * | * | * | * | 3.4 | 5.2 | * |
| ALL OTHER, MALE | | | | | | | | | | | | |
| 1978 | 0.2 | * | * | * | * | * | * | * | * | * | * | * |
| 1977 | 0.2 | * | * | * | * | * | * | * | * | * | * | * |
| 1976 | 0.3 | * | * | * | * | * | * | * | * | * | * | * |
| 1975 | 0.2 | * | * | * | * | * | * | * | * | * | * | * |
| 1974 | 0.3 | * | * | * | * | * | * | * | * | * | * | * |
| 1973 | 0.3 | * | * | * | * | * | * | * | * | * | * | * |
| 1972 | 0.2 | * | * | * | * | * | * | * | * | * | * | * |
| 1971 | 0.3 | * | * | * | * | * | * | * | * | * | * | * |
| 1970 | 0.2 | * | * | * | * | * | * | * | * | * | * | * |
| 1969 | 0.3 | * | * | * | * | * | * | * | * | * | * | * |
| 1968 | 0.4 | * | * | * | * | * | * | * | * | * | * | * |
| ALL OTHER, FEMALE | | | | | | | | | | | | |
| 1978 | 0.2 | * | * | * | * | * | * | * | * | * | * | * |
| 1977 | 0.2 | * | * | * | * | * | * | * | * | * | * | * |
| 1976 | 0.2 | * | * | * | * | * | * | * | * | * | * | * |
| 1975 | 0.2 | * | * | * | * | * | * | * | * | * | * | * |
| 1974 | 0.3 | * | * | * | * | * | * | * | * | * | * | * |
| 1973 | 0.3 | * | * | * | * | * | * | * | * | * | * | * |
| 1972 | 0.4 | * | * | * | * | * | * | * | * | 3.1 | * | * |
| 1971 | 0.3 | * | * | * | * | * | * | * | * | * | * | * |
| 1970 | 0.2 | * | * | * | * | * | * | * | * | * | * | * |
| 1969 | 0.3 | * | * | * | * | * | * | * | * | * | * | * |
| 1968 | 0.5 | * | * | * | * | * | * | * | * | 3.6 | * | * |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| CEREBRAL EMBOLISM (434) | | | | | | | | | | | | |
| BLACK, BOTH SEXES | | | | | | | | | | | | |
| 1978 | 0.2 | * | * | * | * | * | * | * | * | * | * | * |
| 1977 | 0.2 | * | * | * | * | * | * | * | * | * | * | * |
| 1976 | 0.2 | * | * | * | * | * | * | * | * | 1.8 | * | * |
| 1975 | 0.2 | * | * | * | * | * | * | * | * | * | * | * |
| 1974 | 0.3 | * | * | * | * | * | * | * | * | 2.2 | * | * |
| 1973 | 0.3 | * | * | * | * | * | * | * | * | 2.4 | * | * |
| 1972 | 0.3 | * | * | * | * | * | * | * | * | 2.4 | * | * |
| 1971 | 0.4 | * | * | * | * | * | * | * | 1.3 | 2.0 | * | * |
| 1970 | 0.2 | * | * | * | * | * | * | * | * | * | * | * |
| 1969 | 0.3 | * | * | * | * | * | * | * | * | 2.1 | 5.3 | * |
| 1968 | 0.5 | * | * | * | * | * | * | * | * | 3.6 | 5.7 | * |
| BLACK, MALE | | | | | | | | | | | | |
| 1978 | 0.2 | * | * | * | * | * | * | * | * | * | * | * |
| 1977 | 0.2 | * | * | * | * | * | * | * | * | * | * | * |
| 1976 | 0.3 | * | * | * | * | * | * | * | * | * | * | * |
| 1975 | 0.2 | * | * | * | * | * | * | * | * | * | * | * |
| 1974 | 0.4 | * | * | * | * | * | * | * | * | * | * | * |
| 1973 | 0.4 | * | * | * | * | * | * | * | * | * | * | * |
| 1972 | 0.2 | * | * | * | * | * | * | * | * | * | * | * |
| 1971 | 0.4 | * | * | * | * | * | * | * | * | * | * | * |
| 1970 | 0.2 | * | * | * | * | * | * | * | * | * | * | * |
| 1969 | 0.3 | * | * | * | * | * | * | * | * | * | * | * |
| 1968 | 0.4 | * | * | * | * | * | * | * | * | * | * | * |
| BLACK, FEMALE | | | | | | | | | | | | |
| 1978 | 0.2 | * | * | * | * | * | * | * | * | * | * | * |
| 1977 | 0.2 | * | * | * | * | * | * | * | * | * | * | * |
| 1976 | 0.2 | * | * | * | * | * | * | * | * | * | * | * |
| 1975 | 0.2 | * | * | * | * | * | * | * | * | * | * | * |
| 1974 | 0.3 | * | * | * | * | * | * | * | * | * | * | * |
| 1973 | 0.3 | * | * | * | * | * | * | * | * | * | * | * |
| 1972 | 0.5 | * | * | * | * | * | * | * | * | * | * | * |
| 1971 | 0.4 | * | * | * | * | * | * | * | * | * | * | * |
| 1970 | 0.3 | * | * | * | * | * | * | * | * | * | * | * |
| 1969 | 0.4 | * | * | * | * | * | * | * | * | * | * | * |
| 1968 | 0.5 | * | * | * | * | * | * | * | * | 3.9 | * | * |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|--|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| ALL OTHER CEREBROVASCULAR DISEASES (430,432,435-438) | | | | | | | | | | | | |
| BOTH SEXES | | | | | | | | | | | | |
| 1978 | 51.1 | 1.6 | 0.6 | 0.4 | 0.7 | 1.6 | 5.7 | 15.2 | 42.3 | 153.0 | 558.2 | 1,633.6 |
| 1977 | 52.5 | 2.0 | 0.6 | 0.3 | 0.8 | 1.9 | 6.2 | 16.2 | 45.1 | 159.3 | 587.1 | 1,683.7 |
| 1976 | 54.3 | 2.1 | 0.6 | 0.4 | 0.8 | 2.2 | 7.1 | 18.1 | 49.7 | 169.2 | 608.9 | 1,750.4 |
| 1975 | 54.7 | 2.6 | 0.6 | 0.4 | 0.9 | 2.4 | 7.3 | 18.6 | 51.1 | 179.5 | 628.9 | 1,725.5 |
| 1974 | 57.4 | 2.4 | 0.5 | 0.4 | 0.9 | 2.4 | 8.2 | 19.9 | 54.7 | 191.4 | 672.5 | 1,881.2 |
| 1973 | 58.1 | 2.2 | 0.7 | 0.4 | 0.9 | 2.9 | 8.5 | 21.6 | 56.9 | 196.4 | 690.7 | 1,932.7 |
| 1972 | 56.4 | 2.8 | 0.6 | 0.4 | 0.9 | 2.9 | 8.9 | 22.6 | 57.1 | 195.3 | 673.7 | 1,874.8 |
| 1971 | 54.3 | 2.1 | 0.9 | 0.4 | 1.1 | 3.1 | 9.0 | 21.7 | 56.1 | 189.4 | 656.9 | 1,814.2 |
| 1970 | 52.7 | 2.7 | 0.6 | 0.4 | 0.9 | 3.1 | 9.1 | 21.9 | 56.2 | 188.7 | 651.6 | 1,628.3 |
| 1969 | 51.7 | 2.6 | 0.6 | 0.4 | 1.0 | 3.3 | 9.6 | 21.8 | 55.5 | 187.9 | 635.0 | 1,795.9 |
| 1968 | 51.6 | 2.7 | 0.6 | 0.5 | 1.2 | 3.3 | 9.7 | 22.9 | 56.3 | 191.0 | 631.5 | 1,842.9 |
| MALE | | | | | | | | | | | | |
| 1978 | 43.7 | 1.6 | 0.8 | 0.4 | 0.7 | 1.5 | 5.6 | 15.2 | 48.2 | 182.0 | 615.7 | 1,626.5 |
| 1977 | 45.4 | 2.0 | 0.6 | 0.3 | 0.8 | 1.9 | 5.8 | 16.6 | 52.1 | 190.8 | 646.0 | 1,683.3 |
| 1976 | 47.2 | 1.7 | 0.7 | 0.4 | 0.9 | 2.2 | 6.7 | 18.4 | 56.2 | 202.0 | 664.6 | 1,761.7 |
| 1975 | 48.6 | 2.6 | 0.7 | 0.4 | 1.0 | 2.2 | 6.9 | 18.8 | 59.5 | 215.8 | 696.4 | 1,747.1 |
| 1974 | 50.9 | 2.6 | 0.5 | 0.5 | 1.0 | 2.2 | 7.6 | 20.0 | 63.6 | 229.2 | 738.8 | 1,872.0 |
| 1973 | 51.9 | 2.4 | 0.6 | 0.4 | 0.9 | 2.5 | 8.0 | 21.9 | 66.1 | 236.5 | 757.5 | 1,910.0 |
| 1972 | 51.4 | 3.7 | 0.7 | 0.5 | 1.1 | 2.6 | 8.0 | 22.9 | 67.2 | 233.4 | 744.7 | 1,896.2 |
| 1971 | 49.7 | 2.1 | 0.9 | 0.4 | 1.2 | 2.9 | 8.0 | 22.2 | 64.8 | 224.4 | 734.4 | 1,808.8 |
| 1970 | 48.4 | 2.8 | 0.7 | 0.4 | 1.0 | 2.8 | 8.4 | 22.1 | 65.5 | 220.3 | 715.0 | 1,572.1 |
| 1969 | 48.0 | 3.0 | 0.7 | 0.4 | 0.9 | 3.0 | 9.0 | 22.7 | 64.3 | 221.8 | 695.9 | 1,792.2 |
| 1968 | 47.8 | 3.1 | 0.6 | 0.5 | 1.1 | 3.1 | 9.0 | 22.9 | 64.4 | 224.2 | 678.8 | 1,864.8 |

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|--|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| ALL OTHER CEREBROVASCULAR DISEASES (430,432,435-438) | | | | | | | | | | | | |
| FEMALE | | | | | | | | | | | | |
| 1978 | 58.2 | 1.5 | 0.4 | 0.3 | 0.6 | 1.8 | 5.9 | 15.2 | 37.2 | 130.9 | 524.0 | 1,636.7 |
| 1977 | 59.2 | 2.1 | 0.6 | 0.2 | 0.7 | 2.0 | 6.6 | 15.9 | 38.9 | 135.2 | 551.9 | 1,684.0 |
| 1976 | 61.0 | 2.4 | 0.5 | 0.4 | 0.7 | 2.2 | 7.6 | 17.8 | 43.8 | 144.0 | 575.3 | 1,745.1 |
| 1975 | 60.5 | 2.6 | 0.5 | 0.4 | 0.8 | 2.5 | 7.7 | 18.4 | 43.6 | 151.8 | 587.8 | 1,715.0 |
| 1974 | 63.5 | 2.2 | 0.5 | 0.4 | 0.8 | 2.6 | 8.7 | 19.7 | 46.7 | 162.5 | 631.6 | 1,885.8 |
| 1973 | 63.9 | 2.0 | 0.8 | 0.4 | 0.9 | 3.2 | 9.0 | 21.4 | 48.8 | 165.5 | 649.1 | 1,944.2 |
| 1972 | 61.3 | 2.0 | 0.4 | 0.3 | 0.8 | 3.3 | 9.8 | 22.3 | 48.0 | 166.0 | 628.4 | 1,863.9 |
| 1971 | 58.7 | 2.1 | 0.8 | 0.4 | 1.0 | 3.2 | 9.9 | 21.3 | 48.3 | 162.3 | 606.5 | 1,817.0 |
| 1970 | 56.8 | 2.6 | 0.4 | 0.4 | 0.8 | 3.3 | 9.8 | 21.8 | 47.9 | 164.2 | 609.6 | 1,659.7 |
| 1969 | 55.2 | 2.2 | 0.5 | 0.5 | 1.1 | 3.6 | 10.1 | 20.9 | 47.6 | 161.5 | 594.3 | 1,797.9 |
| 1968 | 55.2 | 2.2 | 0.6 | 0.4 | 1.2 | 3.5 | 10.5 | 23.0 | 48.9 | 164.8 | 599.2 | 1,833.3 |
| WHITE, BOTH SEXES | | | | | | | | | | | | |
| 1978 | 51.8 | 1.3 | 0.6 | 0.3 | 0.6 | 1.3 | 4.8 | 12.4 | 36.1 | 139.4 | 551.5 | 1,660.0 |
| 1977 | 53.2 | 1.9 | 0.6 | 0.3 | 0.7 | 1.6 | 5.0 | 13.5 | 38.9 | 145.0 | 582.0 | 1,712.3 |
| 1976 | 54.9 | 1.7 | 0.5 | 0.4 | 0.7 | 1.8 | 6.0 | 15.1 | 42.8 | 154.4 | 605.1 | 1,780.3 |
| 1975 | 55.2 | 2.3 | 0.6 | 0.4 | 0.8 | 2.0 | 5.9 | 15.5 | 43.8 | 164.3 | 625.4 | 1,757.4 |
| 1974 | 57.9 | 1.9 | 0.5 | 0.4 | 0.9 | 2.0 | 6.7 | 16.2 | 46.6 | 174.0 | 671.8 | 1,918.6 |
| 1973 | 58.3 | 1.9 | 0.6 | 0.3 | 0.8 | 2.3 | 6.7 | 17.6 | 48.0 | 178.7 | 688.8 | 1,970.3 |
| 1972 | 56.5 | 2.0 | 0.4 | 0.4 | 0.9 | 2.4 | 7.0 | 18.1 | 48.2 | 177.0 | 672.2 | 1,913.6 |
| 1971 | 54.3 | 1.8 | 0.8 | 0.4 | 1.0 | 2.5 | 7.2 | 17.7 | 47.4 | 172.2 | 654.8 | 1,854.6 |
| 1970 | 52.4 | 2.0 | 0.5 | 0.4 | 0.8 | 2.4 | 7.2 | 17.9 | 46.9 | 171.0 | 645.5 | 1,679.8 |
| 1969 | 51.2 | 2.3 | 0.6 | 0.4 | 0.9 | 2.5 | 7.6 | 17.2 | 45.9 | 169.9 | 628.7 | 1,838.4 |
| 1968 | 50.9 | 2.3 | 0.5 | 0.4 | 1.0 | 2.6 | 7.6 | 18.4 | 46.0 | 172.0 | 626.4 | 1,885.5 |

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|--|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| ALL OTHER CEREBROVASCULAR DISEASES (430,432,435-438) | | | | | | | | | | | | |
| WHITE, MALE | | | | | | | | | | | | |
| 1978 | 43.5 | * | 0.7 | 0.4 | 0.7 | 1.2 | 4.6 | 12.1 | 41.3 | 167.4 | 610.2 | 1,654.4 |
| 1977 | 45.3 | 2.1 | 0.6 | 0.4 | 0.8 | 1.6 | 4.7 | 13.8 | 45.2 | 176.0 | 644.5 | 1,707.5 |
| 1976 | 47.0 | 1.6 | 0.6 | 0.4 | 0.8 | 1.8 | 5.6 | 15.2 | 48.9 | 186.6 | 662.1 | 1,792.0 |
| 1975 | 48.4 | 2.4 | 0.7 | 0.4 | 0.9 | 1.8 | 5.6 | 15.8 | 51.4 | 200.6 | 693.5 | 1,786.1 |
| 1974 | 50.7 | 2.0 | 0.5 | 0.5 | 1.0 | 1.8 | 6.2 | 16.2 | 55.1 | 211.3 | 739.9 | 1,912.6 |
| 1973 | 51.5 | 2.2 | 0.5 | 0.3 | 0.9 | 2.1 | 6.4 | 17.6 | 56.8 | 219.8 | 756.7 | 1,950.0 |
| 1972 | 50.8 | 2.5 | 0.5 | 0.5 | 1.0 | 2.2 | 6.3 | 17.8 | 57.8 | 216.8 | 744.4 | 1,938.9 |
| 1971 | 49.3 | 2.0 | 0.8 | 0.4 | 1.1 | 2.4 | 6.5 | 18.2 | 55.7 | 207.3 | 735.4 | 1,860.0 |
| 1970 | 47.7 | 2.1 | 0.7 | 0.4 | 1.0 | 2.2 | 6.7 | 18.1 | 55.8 | 202.8 | 711.4 | 1,636.1 |
| 1969 | 47.1 | 2.6 | 0.7 | 0.4 | 0.8 | 2.4 | 7.2 | 17.9 | 54.9 | 204.6 | 691.3 | 1,843.4 |
| 1968 | 46.8 | 2.8 | 0.5 | 0.5 | 1.0 | 2.5 | 7.3 | 18.4 | 53.6 | 204.6 | 675.8 | 1,913.3 |
| WHITE, FEMALE | | | | | | | | | | | | |
| 1978 | 59.7 | * | * | 0.3 | 0.5 | 1.5 | 5.0 | 12.7 | 31.5 | 118.0 | 516.9 | 1,662.6 |
| 1977 | 60.7 | 1.7 | 0.6 | 0.2 | 0.6 | 1.7 | 5.4 | 13.2 | 33.2 | 121.4 | 545.2 | 1,714.4 |
| 1976 | 62.5 | 1.7 | 0.4 | 0.4 | 0.6 | 1.9 | 6.4 | 15.0 | 37.3 | 129.7 | 571.3 | 1,776.3 |
| 1975 | 61.7 | 2.2 | 0.5 | 0.3 | 0.7 | 2.1 | 6.3 | 15.4 | 37.1 | 136.5 | 584.6 | 1,745.3 |
| 1974 | 64.8 | 1.9 | 0.4 | 0.4 | 0.8 | 2.1 | 7.2 | 16.3 | 39.1 | 145.5 | 630.2 | 1,921.5 |
| 1973 | 64.8 | 1.7 | 0.7 | 0.3 | 0.8 | 2.6 | 7.0 | 17.5 | 40.1 | 147.3 | 646.8 | 1,982.4 |
| 1972 | 61.8 | 1.5 | 0.4 | 0.3 | 0.7 | 2.5 | 7.6 | 18.4 | 39.7 | 146.3 | 626.5 | 1,900.8 |
| 1971 | 59.1 | 1.7 | 0.7 | 0.4 | 0.9 | 2.6 | 7.8 | 17.1 | 40.1 | 145.1 | 603.2 | 1,851.9 |
| 1970 | 56.9 | 2.0 | 0.4 | 0.3 | 0.7 | 2.6 | 7.6 | 17.7 | 38.9 | 146.6 | 602.5 | 1,703.6 |
| 1969 | 55.0 | 2.1 | 0.5 | 0.5 | 1.0 | 2.5 | 8.0 | 16.6 | 37.8 | 143.0 | 587.1 | 1,838.1 |
| 1968 | 54.8 | 1.7 | 0.6 | 0.4 | 1.0 | 2.7 | 7.9 | 18.4 | 39.0 | 146.5 | 593.0 | 1,873.1 |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|--|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| ALL OTHER CEREBROVASCULAR DISEASES (430,432,435-438) | | | | | | | | | | | | |
| ALL OTHER, BOTH SEXES | | | | | | | | | | | | |
| 1978 | 46.9 | * | * | 0.4 | 0.9 | 3.5 | 12.2 | 36.6 | 98.2 | 280.6 | 627.1 | 1,329.8 |
| 1977 | 48.2 | * | * | * | 1.1 | 3.8 | 14.4 | 37.9 | 102.0 | 294.1 | 640.0 | 1,356.9 |
| 1976 | 50.4 | 4.0 | 1.0 | 0.4 | 1.3 | 4.5 | 15.0 | 42.0 | 113.4 | 310.8 | 648.0 | 1,398.7 |
| 1975 | 51.3 | 4.0 | * | 0.5 | 1.3 | 5.1 | 17.1 | 43.8 | 118.1 | 328.0 | 666.1 | 1,358.9 |
| 1974 | 54.2 | 4.9 | * | 0.3 | 1.2 | 5.3 | 18.7 | 50.3 | 130.1 | 362.5 | 680.9 | 1,446.7 |
| 1973 | 56.6 | * | 1.0 | 0.5 | 1.5 | 6.5 | 21.4 | 55.9 | 141.5 | 370.6 | 710.7 | 1,494.5 |
| 1972 | 56.3 | 7.0 | 1.1 | 0.3 | 1.3 | 6.9 | 22.8 | 61.0 | 141.2 | 377.1 | 691.7 | 1,427.6 |
| 1971 | 54.2 | 3.5 | 1.2 | 0.5 | 1.8 | 6.9 | 22.2 | 57.5 | 138.4 | 359.3 | 681.7 | 1,353.3 |
| 1970 | 54.6 | 6.2 | * | 0.4 | 1.8 | 7.8 | 23.6 | 57.4 | 144.4 | 361.4 | 728.4 | 1,100.0 |
| 1969 | 55.6 | 4.3 | 0.9 | 0.4 | 1.6 | 9.1 | 24.5 | 62.6 | 147.9 | 367.1 | 718.9 | 1,311.9 |
| 1968 | 56.8 | 4.9 | 1.0 | 0.5 | 2.2 | 8.4 | 26.3 | 64.2 | 155.9 | 381.1 | 697.2 | 1,367.6 |
| ALL OTHER, MALE | | | | | | | | | | | | |
| 1978 | 44.9 | * | * | * | 0.8 | 3.5 | 13.1 | 40.3 | 112.1 | 317.5 | 664.7 | 1,341.4 |
| 1977 | 46.1 | * | * | * | 1.1 | 4.0 | 14.4 | 40.3 | 117.3 | 328.6 | 660.5 | 1,435.7 |
| 1976 | 48.7 | * | * | * | 1.5 | 5.1 | 14.7 | 45.7 | 126.1 | 346.1 | 691.1 | 1,458.2 |
| 1975 | 49.8 | * | * | * | 1.5 | 4.9 | 16.8 | 45.8 | 136.8 | 359.5 | 725.6 | 1,357.4 |
| 1974 | 52.7 | * | * | * | 1.3 | 5.0 | 18.6 | 53.8 | 146.1 | 400.0 | 727.9 | 1,454.0 |
| 1973 | 54.6 | * | * | * | 1.4 | 5.6 | 20.6 | 60.1 | 155.7 | 397.1 | 762.9 | 1,502.1 |
| 1972 | 55.1 | 9.5 | * | * | 1.6 | 5.6 | 21.2 | 68.5 | 158.4 | 392.1 | 747.5 | 1,463.8 |
| 1971 | 52.7 | * | * | * | 2.1 | 6.4 | 20.0 | 58.6 | 153.0 | 387.8 | 727.1 | 1,304.3 |
| 1970 | 53.0 | * | * | * | 1.6 | 7.1 | 21.5 | 58.4 | 157.8 | 385.8 | 757.8 | 1,010.4 |
| 1969 | 54.2 | * | * | * | 1.5 | 7.4 | 24.0 | 66.8 | 155.4 | 387.6 | 752.1 | 1,309.1 |
| 1968 | 55.7 | * | * | * | 1.8 | 7.7 | 22.8 | 65.0 | 168.8 | 414.4 | 714.5 | 1,392.7 |
| ALL OTHER, FEMALE | | | | | | | | | | | | |
| 1978 | 48.7 | * | * | * | 1.0 | 3.4 | 11.5 | 33.5 | 86.6 | 252.0 | 603.1 | 1,335.8 |
| 1977 | 50.2 | * | * | * | 1.2 | 3.7 | 14.3 | 35.9 | 89.3 | 267.1 | 624.3 | 1,314.4 |
| 1976 | 52.0 | * | * | * | 1.2 | 4.0 | 15.3 | 39.0 | 102.9 | 282.8 | 618.6 | 1,365.3 |
| 1975 | 52.6 | * | * | * | 1.2 | 5.3 | 17.3 | 42.2 | 102.4 | 302.9 | 623.4 | 1,345.2 |
| 1974 | 55.5 | * | * | * | 1.1 | 5.5 | 18.8 | 47.4 | 116.5 | 332.5 | 648.5 | 1,442.4 |
| 1973 | 58.4 | * | * | * | 1.5 | 7.3 | 22.0 | 52.4 | 129.5 | 349.2 | 676.1 | 1,489.9 |
| 1972 | 57.5 | * | * | * | 1.0 | 8.0 | 24.1 | 54.5 | 126.4 | 365.4 | 653.8 | 1,405.3 |
| 1971 | 55.5 | * | * | * | 1.6 | 7.4 | 24.0 | 56.4 | 125.5 | 336.0 | 648.2 | 1,384.9 |
| 1970 | 56.1 | * | * | * | 1.9 | 8.5 | 25.3 | 56.5 | 132.6 | 341.4 | 706.0 | 1,163.1 |
| 1969 | 56.9 | * | * | * | 1.6 | 10.6 | 25.0 | 58.9 | 141.3 | 350.3 | 690.7 | 1,313.8 |
| 1968 | 57.8 | * | * | * | 2.6 | 8.9 | 29.3 | 63.6 | 144.4 | 353.5 | 686.6 | 1,350.8 |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|--|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| ALL OTHER CEREBROVASCULAR DISEASES (430,432,435-438) | | | | | | | | | | | | |
| BLACK, BOTH SEXES | | | | | | | | | | | | |
| 1978 | 51.6 | * | * | 0.5 | 1.0 | 3.9 | 13.9 | 41.3 | 106.3 | 300.5 | 648.2 | 1,385.3 |
| 1977 | 53.1 | * | * | * | 1.1 | 4.3 | 16.5 | 42.3 | 111.1 | 315.2 | 659.8 | 1,421.0 |
| 1976 | 55.1 | 4.4 | 1.1 | 0.4 | 1.4 | 4.9 | 16.9 | 46.4 | 122.3 | 333.4 | 668.8 | 1,449.3 |
| 1975 | 55.8 | * | * | 0.5 | 1.4 | 5.6 | 18.9 | 47.7 | 126.7 | 350.8 | 686.5 | 1,418.5 |
| 1974 | 58.6 | 5.1 | * | * | 1.2 | 6.1 | 20.9 | 55.2 | 138.7 | 386.7 | 700.8 | 1,500.8 |
| 1973 | 60.8 | * | 1.1 | 0.6 | 1.6 | 7.2 | 23.6 | 60.3 | 151.1 | 392.8 | 730.3 | 1,540.4 |
| 1972 | 60.3 | 8.0 | 1.0 | * | 1.4 | 7.6 | 25.3 | 66.0 | 148.8 | 402.4 | 711.3 | 1,469.1 |
| 1971 | 57.8 | * | 1.3 | 0.5 | 1.9 | 7.6 | 24.1 | 62.1 | 147.0 | 381.6 | 704.5 | 1,403.7 |
| 1970 | 58.7 | 5.7 | * | 0.4 | 1.8 | 8.7 | 26.2 | 62.1 | 154.8 | 383.6 | 770.0 | 1,162.9 |
| 1969 | 59.2 | 4.8 | 1.0 | 0.4 | 1.6 | 9.6 | 26.6 | 67.4 | 156.9 | 388.0 | 755.2 | 1,372.2 |
| 1968 | 60.4 | 4.9 | 1.0 | 0.5 | 2.3 | 9.0 | 28.8 | 68.3 | 166.2 | 401.6 | 726.6 | 1,419.6 |
| BLACK, MALE | | | | | | | | | | | | |
| 1978 | 49.1 | * | * | * | 0.7 | 4.1 | 14.9 | 45.8 | 121.6 | 341.0 | 686.6 | 1,392.2 |
| 1977 | 50.3 | * | * | * | 1.1 | 4.6 | 16.7 | 45.3 | 128.2 | 356.3 | 685.1 | 1,487.8 |
| 1976 | 53.0 | * | * | * | 1.5 | 5.6 | 16.8 | 50.6 | 136.3 | 375.7 | 719.7 | 1,517.0 |
| 1975 | 53.9 | * | * | * | 1.7 | 5.4 | 18.8 | 49.6 | 148.4 | 390.8 | 744.1 | 1,421.7 |
| 1974 | 56.6 | * | * | * | 1.4 | 5.8 | 20.9 | 59.2 | 157.7 | 430.8 | 748.3 | 1,525.6 |
| 1973 | 58.1 | * | * | * | 1.5 | 6.1 | 23.0 | 65.1 | 168.0 | 422.7 | 779.9 | 1,570.7 |
| 1972 | 58.7 | 10.8 | * | * | 1.7 | 6.1 | 23.9 | 74.6 | 170.1 | 421.3 | 771.6 | 1,535.0 |
| 1971 | 56.0 | * | * | * | 2.2 | 6.9 | 22.2 | 63.5 | 164.4 | 415.4 | 750.0 | 1,380.0 |
| 1970 | 56.5 | * | * | * | 1.7 | 8.0 | 24.0 | 63.7 | 170.6 | 412.4 | 803.5 | 1,089.2 |
| 1969 | 57.3 | * | * | * | 1.5 | 7.5 | 25.5 | 72.1 | 167.4 | 414.5 | 788.6 | 1,416.2 |
| 1968 | 59.0 | * | * | * | 1.8 | 8.3 | 25.3 | 69.6 | 181.6 | 439.1 | 751.5 | 1,502.9 |
| BLACK, FEMALE | | | | | | | | | | | | |
| 1978 | 53.9 | * | * | * | 1.2 | 3.8 | 13.0 | 37.5 | 93.7 | 270.0 | 622.6 | 1,381.8 |
| 1977 | 55.5 | * | * | * | 1.2 | 4.0 | 16.4 | 39.8 | 97.2 | 284.3 | 644.9 | 1,386.2 |
| 1976 | 57.1 | * | * | * | 1.2 | 4.3 | 17.0 | 42.9 | 110.7 | 301.0 | 634.7 | 1,413.5 |
| 1975 | 57.6 | * | * | * | 1.2 | 5.8 | 19.0 | 46.1 | 108.6 | 320.4 | 647.6 | 1,416.7 |
| 1974 | 60.4 | * | * | * | 1.1 | 6.3 | 20.9 | 51.9 | 123.1 | 353.4 | 668.6 | 1,487.2 |
| 1973 | 63.2 | * | * | * | 1.7 | 8.2 | 24.1 | 56.1 | 137.1 | 369.7 | 698.6 | 1,523.3 |
| 1972 | 61.7 | * | * | * | 1.0 | 8.9 | 26.5 | 58.7 | 130.8 | 388.4 | 666.4 | 1,431.4 |
| 1971 | 59.5 | * | * | * | 1.5 | 8.2 | 25.8 | 60.9 | 132.2 | 354.3 | 671.8 | 1,417.9 |
| 1970 | 60.7 | * | * | * | 2.0 | 9.3 | 28.1 | 60.8 | 141.3 | 360.7 | 745.5 | 1,210.3 |
| 1969 | 60.9 | * | * | * | 1.8 | 11.4 | 27.5 | 63.2 | 148.2 | 367.0 | 727.3 | 1,345.0 |
| 1968 | 61.6 | * | * | * | 2.8 | 9.6 | 31.7 | 67.2 | 152.9 | 370.7 | 708.1 | 1,392.9 |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| ATHEROSCLEROSIS 440 | | | | | | | | | | | | |
| BOTH SEXES | | | | | | | | | | | | |
| 1978 | 13.0 | * | * | * | * | * | 0.2 | 0.9 | 4.7 | 24.2 | 134.3 | 672.2 |
| 1977 | 13.1 | * | * | * | * | * | 0.2 | 1.1 | 5.2 | 25.3 | 137.3 | 688.0 |
| 1976 | 13.5 | * | * | * | * | * | 0.2 | 0.9 | 5.1 | 25.7 | 144.6 | 740.7 |
| 1975 | 13.4 | * | * | * | * | * | 0.2 | 1.1 | 5.1 | 26.5 | 152.4 | 728.5 |
| 1974 | 15.1 | * | * | * | * | * | 0.2 | 1.1 | 5.8 | 31.5 | 172.1 | 869.6 |
| 1973 | 15.4 | * | * | * | * | * | 0.2 | 1.1 | 6.0 | 32.3 | 185.2 | 903.0 |
| 1972 | 15.5 | * | * | * | * | * | 0.2 | 1.3 | 6.6 | 34.9 | 190.1 | 900.6 |
| 1971 | 15.2 | * | * | * | * | * | 0.2 | 1.2 | 6.4 | 34.4 | 188.1 | 911.9 |
| 1970 | 15.6 | * | * | * | * | * | 0.3 | 1.3 | 6.8 | 37.3 | 197.8 | 880.2 |
| 1969 | 16.4 | * | * | * | * | * | 0.2 | 1.1 | 6.6 | 39.0 | 212.0 | 1,060.8 |
| 1968 | 16.8 | * | * | * | * | * | 0.2 | 1.2 | 6.8 | 40.7 | 220.7 | 1,121.7 |
| MALE | | | | | | | | | | | | |
| 1978 | 10.7 | * | * | * | * | * | 0.3 | 1.3 | 6.4 | 30.9 | 152.4 | 689.4 |
| 1977 | 10.9 | * | * | * | * | * | 0.2 | 1.6 | 6.9 | 32.1 | 160.0 | 697.8 |
| 1976 | 11.2 | * | * | * | * | * | 0.2 | 1.2 | 7.1 | 32.4 | 163.0 | 755.1 |
| 1975 | 11.3 | * | * | * | * | * | 0.3 | 1.4 | 6.7 | 33.2 | 172.9 | 745.8 |
| 1974 | 12.7 | * | * | * | * | * | 0.2 | 1.4 | 7.8 | 39.7 | 195.8 | 868.3 |
| 1973 | 13.0 | * | * | * | * | * | * | 1.5 | 8.0 | 40.0 | 207.4 | 901.7 |
| 1972 | 13.5 | * | * | * | * | * | 0.2 | 1.9 | 8.9 | 42.3 | 213.8 | 930.7 |
| 1971 | 13.2 | * | * | * | * | * | 0.3 | 1.6 | 8.7 | 42.2 | 211.1 | 919.4 |
| 1970 | 13.9 | * | * | * | * | * | 0.3 | 1.8 | 9.3 | 45.5 | 226.2 | 869.5 |
| 1969 | 14.6 | * | * | * | * | * | 0.3 | 1.4 | 8.8 | 48.9 | 239.8 | 1,067.1 |
| 1968 | 15.1 | * | * | * | * | * | 0.2 | 1.5 | 8.9 | 50.8 | 244.3 | 1,150.0 |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| ATHEROSCLEROSIS 440 | | | | | | | | | | | | |
| FEMALE | | | | | | | | | | | | |
| 1978 | 15.2 | * | * | * | * | * | * | 0.6 | 3.2 | 19.1 | 123.5 | 664.4 |
| 1977 | 15.2 | * | * | * | * | * | * | 0.6 | 3.6 | 20.0 | 123.7 | 683.4 |
| 1976 | 15.7 | * | * | * | * | * | 0.2 | 0.7 | 3.4 | 20.5 | 133.5 | 734.0 |
| 1975 | 15.4 | * | * | * | * | * | * | 0.7 | 3.6 | 21.4 | 139.8 | 720.1 |
| 1974 | 17.4 | * | * | * | * | * | * | 0.7 | 4.1 | 25.2 | 157.4 | 870.2 |
| 1973 | 17.7 | * | * | * | * | * | 0.2 | 0.8 | 4.2 | 26.4 | 171.4 | 903.6 |
| 1972 | 17.5 | * | * | * | * | * | * | 0.7 | 4.7 | 29.1 | 174.9 | 885.3 |
| 1971 | 17.1 | * | * | * | * | * | * | 0.8 | 4.4 | 28.4 | 173.2 | 908.0 |
| 1970 | 17.2 | * | * | * | * | * | 0.2 | 0.8 | 4.5 | 30.9 | 179.0 | 886.2 |
| 1969 | 18.1 | * | * | * | * | * | * | 0.8 | 4.6 | 31.2 | 193.4 | 1,057.4 |
| 1968 | 18.5 | * | * | * | * | * | * | 0.9 | 5.0 | 32.8 | 204.5 | 1,107.5 |
| WHITE, BOTH SEXES | | | | | | | | | | | | |
| 1978 | 14.0 | * | * | * | * | * | 0.2 | 1.0 | 4.3 | 23.4 | 136.2 | 694.4 |
| 1977 | 14.0 | * | * | * | * | * | 0.1 | 1.0 | 4.8 | 24.4 | 139.6 | 712.6 |
| 1976 | 14.5 | * | * | * | * | * | 0.2 | 0.9 | 4.8 | 24.8 | 147.9 | 763.9 |
| 1975 | 14.3 | * | * | * | * | * | 0.2 | 1.0 | 4.7 | 25.5 | 155.0 | 753.9 |
| 1974 | 16.1 | * | * | * | * | * | 0.2 | 0.9 | 5.4 | 29.9 | 175.0 | 900.7 |
| 1973 | 16.4 | * | * | * | * | * | 0.2 | 1.0 | 5.5 | 30.4 | 188.3 | 934.9 |
| 1972 | 16.4 | * | * | * | * | * | 0.2 | 1.2 | 6.0 | 32.7 | 191.9 | 935.2 |
| 1971 | 16.2 | * | * | * | * | * | 0.2 | 1.0 | 5.9 | 32.8 | 191.2 | 947.5 |
| 1970 | 16.6 | * | * | * | * | * | 0.2 | 1.1 | 6.2 | 35.2 | 200.0 | 927.9 |
| 1969 | 17.4 | * | * | * | * | * | 0.1 | 0.9 | 6.0 | 36.9 | 214.6 | 1,102.8 |
| 1968 | 17.8 | * | * | * | * | * | 0.2 | 1.0 | 5.9 | 38.3 | 222.8 | 1,172.0 |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| ATHEROSCLEROSIS 440 | | | | | | | | | | | | |
| WHITE, MALE | | | | | | | | | | | | |
| 1978 | 11.3 | * | * | * | * | * | 0.3 | 1.3 | 5.9 | 30.3 | 155.0 | 712.3 |
| 1977 | 11.5 | * | * | * | * | * | 0.2 | 1.5 | 6.5 | 31.3 | 163.3 | 722.4 |
| 1976 | 11.8 | * | * | * | * | * | 0.2 | 1.2 | 6.9 | 31.6 | 167.9 | 780.8 |
| 1975 | 11.9 | * | * | * | * | * | 0.2 | 1.3 | 6.3 | 32.5 | 175.8 | 770.7 |
| 1974 | 13.4 | * | * | * | * | * | * | 1.3 | 7.3 | 38.5 | 199.2 | 899.0 |
| 1973 | 13.7 | * | * | * | * | * | * | 1.3 | 7.6 | 38.2 | 211.2 | 936.3 |
| 1972 | 14.1 | * | * | * | * | * | 0.2 | 1.7 | 8.2 | 40.0 | 215.7 | 972.6 |
| 1971 | 14.0 | * | * | * | * | * | 0.3 | 1.5 | 8.1 | 41.0 | 215.9 | 961.1 |
| 1970 | 14.6 | * | * | * | * | * | 0.3 | 1.6 | 8.8 | 43.7 | 229.4 | 923.9 |
| 1969 | 15.3 | * | * | * | * | * | 0.2 | 1.2 | 8.2 | 46.6 | 243.4 | 1,115.7 |
| 1968 | 15.7 | * | * | * | * | * | 0.2 | 1.3 | 8.0 | 48.0 | 247.5 | 1,207.5 |
| WHITE, FEMALE | | | | | | | | | | | | |
| 1978 | 16.5 | * | * | * | * | * | * | 0.6 | 2.9 | 18.1 | 125.1 | 686.4 |
| 1977 | 16.4 | * | * | * | * | * | * | 0.5 | 3.2 | 19.1 | 125.6 | 708.1 |
| 1976 | 17.0 | * | * | * | * | * | * | 0.6 | 3.0 | 19.6 | 136.1 | 756.7 |
| 1975 | 16.7 | * | * | * | * | * | * | 0.7 | 3.3 | 20.2 | 142.5 | 746.5 |
| 1974 | 18.7 | * | * | * | * | * | * | 0.6 | 3.7 | 23.3 | 160.2 | 901.5 |
| 1973 | 19.0 | * | * | * | * | * | * | 0.7 | 3.6 | 24.4 | 174.1 | 935.1 |
| 1972 | 18.7 | * | * | * | * | * | * | 0.7 | 4.0 | 27.2 | 176.9 | 916.3 |
| 1971 | 18.3 | * | * | * | * | * | * | 0.6 | 3.8 | 26.5 | 175.4 | 940.5 |
| 1970 | 18.4 | * | * | * | * | * | * | 0.7 | 3.8 | 28.6 | 180.8 | 930.0 |
| 1969 | 19.3 | * | * | * | * | * | * | 0.7 | 4.0 | 29.3 | 195.4 | 1,097.4 |
| 1968 | 19.7 | * | * | * | * | * | * | 0.7 | 4.0 | 30.6 | 206.0 | 1,154.4 |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| ATHEROSCLEROSIS 440 | | | | | | | | | | | | |
| ALL OTHER, BOTH SEXES | | | | | | | | | | | | |
| 1978 | 7.1 | * | * | * | * | * | * | 0.9 | 8.2 | 32.0 | 115.1 | 417.9 |
| 1977 | 7.1 | * | * | * | * | * | * | 1.7 | 8.5 | 33.4 | 113.2 | 406.3 |
| 1976 | 7.2 | * | * | * | * | * | * | 1.3 | 7.7 | 34.2 | 110.4 | 471.9 |
| 1975 | 7.4 | * | * | * | * | * | * | 1.9 | 8.5 | 36.3 | 124.4 | 437.7 |
| 1974 | 8.4 | * | * | * | * | * | * | 2.1 | 10.0 | 47.3 | 140.1 | 507.4 |
| 1973 | 8.8 | * | * | * | * | * | * | 1.9 | 10.7 | 51.1 | 150.8 | 531.5 |
| 1972 | 9.3 | * | * | * | * | * | * | 2.3 | 12.9 | 56.0 | 168.4 | 502.4 |
| 1971 | 8.6 | * | * | * | * | * | * | 2.4 | 12.0 | 50.4 | 150.5 | 506.7 |
| 1970 | 8.9 | * | * | * | * | * | * | 2.6 | 12.9 | 57.6 | 169.4 | 390.8 |
| 1969 | 9.7 | * | * | * | * | * | * | 2.7 | 12.5 | 59.9 | 179.7 | 589.0 |
| 1968 | 10.2 | * | * | * | * | * | * | 2.8 | 16.1 | 65.6 | 193.2 | 559.8 |
| ALL OTHER, MALE | | | | | | | | | | | | |
| 1978 | 6.7 | * | * | * | * | * | * | * | 10.8 | 36.1 | 127.4 | 455.2 |
| 1977 | 6.8 | * | * | * | * | * | * | 2.0 | 10.5 | 39.1 | 128.4 | 446.4 |
| 1976 | 6.7 | * | * | * | * | * | * | 1.7 | 8.9 | 39.9 | 117.5 | 498.2 |
| 1975 | 7.3 | * | * | * | * | * | * | 2.5 | 10.5 | 40.2 | 144.2 | 496.3 |
| 1974 | 8.2 | * | * | * | * | * | * | 2.2 | 12.6 | 51.5 | 161.1 | 552.0 |
| 1973 | 8.4 | * | * | * | * | * | * | 3.0 | 11.8 | 57.2 | 167.0 | 547.9 |
| 1972 | 9.3 | * | * | * | * | * | * | 3.7 | 15.4 | 64.8 | 193.5 | 506.4 |
| 1971 | 8.1 | * | * | * | * | * | * | 2.8 | 14.2 | 53.3 | 158.9 | 508.5 |
| 1970 | 8.9 | * | * | * | * | * | * | 3.5 | 14.9 | 62.5 | 190.1 | 391.5 |
| 1969 | 9.8 | * | * | * | * | * | * | 3.6 | 14.5 | 70.4 | 196.8 | 609.1 |
| 1968 | 10.3 | * | * | * | * | * | * | 3.3 | 17.8 | 77.8 | 205.4 | 590.2 |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| ATHEROSCLEROSIS 440 | | | | | | | | | | | | |
| ALL OTHER, FEMALE | | | | | | | | | | | | |
| 1978 | 7.4 | * | * | * | * | * | * | * | 6.1 | 28.8 | 106.9 | 401.8 |
| 1977 | 7.3 | * | * | * | * | * | * | 1.5 | 6.8 | 28.8 | 102.6 | 384.6 |
| 1976 | 7.7 | * | * | * | * | * | * | * | 6.7 | 29.7 | 105.6 | 457.1 |
| 1975 | 7.4 | * | * | * | * | * | * | 1.4 | 6.8 | 33.2 | 110.5 | 398.9 |
| 1974 | 8.6 | * | * | * | * | * | * | 2.0 | 7.8 | 43.8 | 125.6 | 481.2 |
| 1973 | 9.1 | * | * | * | * | * | * | * | 9.7 | 46.2 | 139.9 | 521.5 |
| 1972 | 9.2 | * | * | * | * | * | * | * | 10.8 | 48.9 | 150.8 | 500.0 |
| 1971 | 9.0 | * | * | * | * | * | * | 2.2 | 10.1 | 48.0 | 144.3 | 505.5 |
| 1970 | 8.9 | * | * | * | * | * | * | 1.8 | 11.2 | 53.6 | 153.7 | 390.3 |
| 1969 | 9.5 | * | * | * | * | * | * | 1.9 | 10.8 | 51.3 | 166.0 | 575.4 |
| 1968 | 10.0 | * | * | * | * | * | * | 2.4 | 14.7 | 55.5 | 184.5 | 539.3 |
| BLACK, BOTH SEXES | | | | | | | | | | | | |
| 1978 | 7.8 | * | * | * | * | * | * | 1.1 | 8.9 | 34.1 | 120.1 | 440.7 |
| 1977 | 7.8 | * | * | * | * | * | * | 1.9 | 9.0 | 36.2 | 118.0 | 432.2 |
| 1976 | 7.9 | * | * | * | * | * | * | 1.5 | 8.5 | 37.2 | 113.7 | 494.9 |
| 1975 | 8.0 | * | * | * | * | * | * | 2.2 | 9.3 | 38.7 | 128.9 | 461.5 |
| 1974 | 9.0 | * | * | * | * | * | * | 2.4 | 10.4 | 50.3 | 141.7 | 524.8 |
| 1973 | 9.5 | * | * | * | * | * | * | 2.2 | 11.4 | 55.2 | 156.1 | 553.5 |
| 1972 | 9.9 | * | * | * | * | * | * | 2.4 | 13.5 | 60.1 | 172.7 | 536.4 |
| 1971 | 9.1 | * | * | * | * | * | * | 2.6 | 12.8 | 54.3 | 158.0 | 514.0 |
| 1970 | 9.5 | * | * | * | * | * | * | 2.9 | 13.9 | 61.5 | 180.2 | 415.7 |
| 1969 | 10.4 | * | * | * | * | * | * | 2.9 | 13.7 | 64.4 | 190.8 | 628.9 |
| 1968 | 10.8 | * | * | * | * | * | * | 3.1 | 17.2 | 69.0 | 203.4 | 583.7 |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| ATHEROSCLEROSIS 440 | | | | | | | | | | | | |
| BLACK, MALE | | | | | | | | | | | | |
| 1978 | 7.2 | * | * | * | * | * | * | * | 11.4 | 38.7 | 133.9 | 464.7 |
| 1977 | 7.4 | * | * | * | * | * | * | 2.2 | 11.3 | 42.5 | 134.0 | 469.4 |
| 1976 | 7.2 | * | * | * | * | * | * | * | 10.0 | 43.6 | 119.7 | 519.1 |
| 1975 | 7.9 | * | * | * | * | * | * | 2.8 | 11.5 | 44.0 | 148.2 | 547.8 |
| 1974 | 8.7 | * | * | * | * | * | * | 2.5 | 13.4 | 54.8 | 161.7 | 593.0 |
| 1973 | 9.0 | * | * | * | * | * | * | 3.3 | 12.9 | 62.5 | 172.5 | 582.9 |
| 1972 | 9.9 | * | * | * | * | * | * | 4.0 | 16.1 | 70.4 | 195.9 | 555.0 |
| 1971 | 8.5 | * | * | * | * | * | * | 3.0 | 15.4 | 57.8 | 165.1 | 530.0 |
| 1970 | 9.5 | * | * | * | * | * | * | 3.9 | 16.5 | 67.7 | 202.8 | 430.5 |
| 1969 | 10.5 | * | * | * | * | * | * | 3.8 | 16.0 | 76.3 | 212.6 | 670.3 |
| 1968 | 10.8 | * | * | * | * | * | * | 3.7 | 19.3 | 82.8 | 217.0 | 634.3 |
| BLACK, FEMALE | | | | | | | | | | | | |
| 1978 | 8.3 | * | * | * | * | * | * | * | 6.9 | 30.7 | 110.9 | 428.3 |
| 1977 | 8.2 | * | * | * | * | * | * | 1.6 | 7.1 | 31.5 | 107.6 | 412.8 |
| 1976 | 8.5 | * | * | * | * | * | * | * | 7.3 | 32.2 | 109.7 | 482.0 |
| 1975 | 8.1 | * | * | * | * | * | * | 1.6 | 7.5 | 34.7 | 115.9 | 414.3 |
| 1974 | 9.3 | * | * | * | * | * | * | 2.3 | 8.0 | 46.9 | 128.2 | 487.2 |
| 1973 | 9.9 | * | * | * | * | * | * | * | 10.2 | 49.6 | 145.3 | 537.0 |
| 1972 | 10.0 | * | * | * | * | * | * | * | 11.3 | 52.2 | 155.7 | 525.7 |
| 1971 | 9.6 | * | * | * | * | * | * | 2.3 | 10.7 | 51.4 | 152.9 | 504.5 |
| 1970 | 9.6 | * | * | * | * | * | * | 2.0 | 11.7 | 56.5 | 163.5 | 406.2 |
| 1969 | 10.3 | * | * | * | * | * | * | 2.1 | 11.6 | 54.9 | 174.0 | 603.3 |
| 1968 | 10.7 | * | * | * | * | * | * | 2.6 | 15.5 | 57.7 | 193.2 | 562.5 |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| OTHER DISEASES OF ARTERIES, ARTERIOLES, AND CAPILLARIES (441-448) | | | | | | | | | | | | |
| BOTH SEXES | | | | | | | | | | | | |
| 1978 | 12.0 | 1.1 | * | 0.1 | 0.3 | 0.6 | 1.7 | 5.2 | 17.5 | 51.5 | 111.4 | 231.8 |
| 1977 | 12.0 | 1.1 | * | 0.1 | 0.3 | 0.6 | 1.9 | 5.6 | 17.6 | 52.1 | 115.0 | 227.0 |
| 1976 | 12.1 | 0.9 | * | 0.1 | 0.3 | 0.6 | 1.7 | 5.7 | 19.0 | 54.1 | 117.1 | 230.3 |
| 1975 | 11.9 | 1.0 | * | 0.1 | 0.3 | 0.7 | 1.8 | 5.7 | 19.0 | 53.0 | 117.4 | 223.8 |
| 1974 | 12.3 | 1.3 | * | 0.1 | 0.3 | 0.7 | 1.9 | 5.9 | 20.2 | 56.5 | 121.5 | 237.2 |
| 1973 | 12.3 | 1.0 | * | 0.1 | 0.3 | 0.7 | 2.2 | 6.1 | 20.4 | 58.1 | 120.7 | 247.9 |
| 1972 | 12.3 | 1.3 | * | 0.1 | 0.3 | 0.7 | 2.2 | 6.6 | 21.2 | 59.6 | 119.8 | 234.0 |
| 1971 | 12.2 | 1.1 | 0.1 | 0.1 | 0.3 | 0.8 | 2.4 | 6.5 | 20.7 | 58.4 | 123.8 | 241.1 |
| 1970 | 12.5 | 1.1 | * | 0.1 | 0.3 | 0.8 | 2.2 | 6.3 | 21.3 | 60.9 | 131.5 | 222.7 |
| 1969 | 12.5 | 0.9 | * | 0.1 | 0.4 | 0.7 | 2.3 | 6.6 | 21.9 | 61.0 | 129.4 | 250.6 |
| 1968 | 12.4 | 0.9 | * | 0.1 | 0.4 | 0.8 | 2.2 | 6.6 | 22.3 | 60.8 | 127.3 | 265.1 |
| MALE | | | | | | | | | | | | |
| 1978 | 14.5 | 1.5 | * | * | 0.3 | 0.7 | 2.0 | 6.7 | 26.4 | 80.8 | 172.2 | 283.9 |
| 1977 | 14.6 | 1.6 | * | * | 0.3 | 0.6 | 2.1 | 7.3 | 26.1 | 82.6 | 176.7 | 278.9 |
| 1976 | 15.1 | * | * | * | 0.4 | 0.7 | 2.1 | 7.6 | 29.2 | 87.5 | 177.9 | 284.2 |
| 1975 | 14.7 | * | * | * | 0.4 | 0.7 | 2.0 | 7.6 | 28.2 | 85.7 | 177.7 | 270.0 |
| 1974 | 15.2 | 1.3 | * | 0.1 | 0.3 | 0.8 | 2.2 | 7.6 | 31.0 | 90.9 | 180.7 | 280.7 |
| 1973 | 15.5 | * | * | * | 0.4 | 0.7 | 2.4 | 8.3 | 31.4 | 94.0 | 180.6 | 301.1 |
| 1972 | 15.6 | 1.2 | * | 0.1 | 0.3 | 0.7 | 2.6 | 9.0 | 32.8 | 96.1 | 181.5 | 265.9 |
| 1971 | 15.3 | 1.4 | * | 0.1 | 0.4 | 0.8 | 2.9 | 8.8 | 32.2 | 93.4 | 176.4 | 278.4 |
| 1970 | 15.8 | 1.2 | * | 0.1 | 0.3 | 0.9 | 2.5 | 8.5 | 33.2 | 97.4 | 187.8 | 247.8 |
| 1969 | 15.7 | * | * | 0.1 | 0.4 | 0.7 | 2.7 | 8.9 | 34.2 | 97.1 | 177.7 | 283.9 |
| 1968 | 15.8 | * | * | * | 0.4 | 1.0 | 2.8 | 9.2 | 34.8 | 97.2 | 175.4 | 304.3 |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| OTHER DISEASES OF ARTERIES, ARTERIOLES, AND CAPILLARIES (441-448) | | | | | | | | | | | | |
| FEMALE | | | | | | | | | | | | |
| 1978 | 9.5 | * | * | * | 0.2 | 0.6 | 1.4 | 3.7 | 9.6 | 29.1 | 75.2 | 208.2 |
| 1977 | 9.5 | * | * | * | 0.3 | 0.5 | 1.6 | 4.0 | 10.2 | 28.7 | 78.0 | 203.1 |
| 1976 | 9.3 | * | * | * | 0.2 | 0.5 | 1.4 | 3.9 | 9.9 | 28.5 | 80.4 | 205.0 |
| 1975 | 9.2 | * | * | * | 0.2 | 0.7 | 1.7 | 3.9 | 10.8 | 27.9 | 80.7 | 201.5 |
| 1974 | 9.5 | 1.3 | * | * | 0.3 | 0.6 | 1.6 | 4.3 | 10.7 | 30.2 | 85.0 | 215.6 |
| 1973 | 9.3 | * | * | * | 0.3 | 0.7 | 1.9 | 4.2 | 10.6 | 30.5 | 83.3 | 221.0 |
| 1972 | 9.1 | 1.4 | * | * | 0.2 | 0.7 | 1.8 | 4.4 | 10.8 | 31.5 | 80.5 | 217.6 |
| 1971 | 9.3 | * | * | * | 0.3 | 0.8 | 1.8 | 4.4 | 10.3 | 31.3 | 89.7 | 221.6 |
| 1970 | 9.3 | * | * | * | 0.3 | 0.8 | 1.9 | 4.2 | 10.7 | 32.5 | 94.2 | 208.7 |
| 1969 | 9.4 | * | * | * | 0.3 | 0.7 | 1.9 | 4.4 | 10.8 | 32.8 | 97.0 | 232.5 |
| 1968 | 9.2 | * | * | * | 0.3 | 0.7 | 1.7 | 4.1 | 11.0 | 32.2 | 94.4 | 243.9 |
| WHITE, BOTH SEXES | | | | | | | | | | | | |
| 1978 | 12.6 | 0.9 | * | 0.1 | 0.3 | 0.6 | 1.5 | 4.8 | 17.1 | 52.4 | 113.4 | 236.3 |
| 1977 | 12.6 | 1.1 | * | 0.1 | 0.3 | 0.5 | 1.7 | 5.2 | 17.3 | 52.6 | 118.1 | 231.7 |
| 1976 | 12.8 | 0.9 | * | 0.1 | 0.3 | 0.5 | 1.5 | 5.2 | 18.6 | 54.8 | 120.0 | 234.7 |
| 1975 | 12.5 | 0.8 | * | 0.1 | 0.3 | 0.6 | 1.6 | 5.3 | 18.8 | 53.2 | 120.4 | 229.4 |
| 1974 | 12.9 | 1.3 | * | 0.1 | 0.3 | 0.6 | 1.6 | 5.5 | 20.0 | 57.2 | 124.7 | 244.2 |
| 1973 | 13.0 | 0.9 | * | 0.1 | 0.3 | 0.7 | 2.0 | 5.6 | 20.2 | 58.7 | 124.3 | 254.4 |
| 1972 | 12.9 | 0.9 | * | 0.1 | 0.2 | 0.6 | 1.8 | 6.1 | 20.7 | 60.6 | 123.3 | 241.4 |
| 1971 | 12.8 | 1.1 | * | 0.1 | 0.3 | 0.7 | 2.0 | 5.9 | 20.2 | 59.2 | 126.9 | 248.0 |
| 1970 | 13.0 | 1.1 | * | 0.1 | 0.3 | 0.8 | 1.9 | 5.7 | 20.8 | 61.5 | 134.5 | 230.3 |
| 1969 | 13.0 | 0.9 | * | 0.1 | 0.3 | 0.6 | 2.0 | 5.9 | 21.4 | 61.8 | 132.8 | 258.2 |
| 1968 | 12.9 | 0.8 | * | 0.1 | 0.3 | 0.8 | 1.9 | 6.0 | 21.6 | 61.8 | 130.0 | 272.2 |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| OTHER DISEASES OF ARTERIES, ARTERIOLES, AND CAPILLARIES (441-448) | | | | | | | | | | | | |
| WHITE, MALE | | | | | | | | | | | | |
| 1978 | 15.5 | * | * | * | 0.3 | 0.6 | 1.8 | 6.3 | 26.1 | 83.8 | 179.2 | 293.8 |
| 1977 | 15.6 | 1.6 | * | * | 0.4 | 0.6 | 2.0 | 6.9 | 25.9 | 85.1 | 184.9 | 288.3 |
| 1976 | 16.0 | * | * | * | 0.3 | 0.6 | 1.8 | 7.1 | 28.9 | 90.4 | 185.5 | 295.6 |
| 1975 | 15.6 | * | * | * | 0.3 | 0.6 | 1.7 | 7.1 | 28.4 | 88.4 | 184.7 | 280.2 |
| 1974 | 16.1 | * | * | * | 0.3 | 0.7 | 1.7 | 7.1 | 31.0 | 93.8 | 188.5 | 293.8 |
| 1973 | 16.5 | * | * | * | 0.4 | 0.7 | 2.2 | 7.6 | 31.7 | 97.3 | 188.6 | 314.3 |
| 1972 | 16.5 | * | * | 0.2 | 0.3 | 0.7 | 2.3 | 8.4 | 32.6 | 100.0 | 188.4 | 277.5 |
| 1971 | 16.1 | 1.4 | * | * | 0.4 | 0.7 | 2.4 | 8.1 | 32.1 | 96.8 | 182.8 | 287.9 |
| 1970 | 16.6 | * | * | 0.1 | 0.3 | 0.8 | 2.2 | 7.9 | 33.2 | 100.4 | 195.1 | 260.0 |
| 1969 | 16.5 | * | * | * | 0.4 | 0.7 | 2.4 | 8.2 | 34.5 | 100.3 | 184.7 | 296.1 |
| 1968 | 16.6 | * | * | * | 0.4 | 0.9 | 2.5 | 8.7 | 34.4 | 100.8 | 180.7 | 318.5 |
| WHITE, FEMALE | | | | | | | | | | | | |
| 1978 | 9.9 | * | * | * | 0.2 | 0.5 | 1.3 | 3.5 | 9.1 | 28.3 | 74.7 | 210.7 |
| 1977 | 9.8 | * | * | * | 0.2 | 0.4 | 1.3 | 3.6 | 9.6 | 27.9 | 78.8 | 206.0 |
| 1976 | 9.6 | * | * | * | 0.2 | 0.4 | 1.2 | 3.4 | 9.5 | 27.5 | 81.0 | 206.8 |
| 1975 | 9.5 | * | * | * | 0.2 | 0.6 | 1.5 | 3.6 | 10.3 | 26.2 | 81.7 | 205.5 |
| 1974 | 9.9 | * | * | * | 0.3 | 0.5 | 1.4 | 3.9 | 10.3 | 29.3 | 85.7 | 220.0 |
| 1973 | 9.6 | * | * | * | 0.2 | 0.7 | 1.8 | 3.7 | 9.9 | 29.3 | 84.5 | 225.0 |
| 1972 | 9.4 | * | * | * | 0.1 | 0.5 | 1.4 | 3.9 | 10.0 | 30.3 | 82.2 | 223.3 |
| 1971 | 9.6 | * | * | * | 0.3 | 0.7 | 1.7 | 3.9 | 9.5 | 30.3 | 91.2 | 227.5 |
| 1970 | 9.6 | * | * | * | 0.2 | 0.7 | 1.7 | 3.7 | 9.7 | 31.5 | 94.9 | 214.1 |
| 1969 | 9.7 | * | * | * | 0.2 | 0.6 | 1.6 | 3.8 | 9.7 | 31.8 | 98.3 | 238.4 |
| 1968 | 9.4 | * | * | * | 0.3 | 0.7 | 1.4 | 3.5 | 10.0 | 31.3 | 95.7 | 247.5 |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| OTHER DISEASES OF ARTERIES, ARTERIOLES, AND CAPILLARIES (441-448) | | | | | | | | | | | | |
| ALL OTHER, BOTH SEXES | | | | | | | | | | | | |
| 1978 | 7.8 | * | * | * | 0.3 | 1.0 | 3.1 | 7.9 | 20.8 | 43.5 | 90.7 | 179.8 |
| 1977 | 7.8 | * | * | * | 0.4 | 1.0 | 3.1 | 8.7 | 21.1 | 46.9 | 82.1 | 173.1 |
| 1976 | 8.1 | * | * | * | 0.4 | 1.3 | 3.3 | 9.5 | 22.0 | 47.6 | 87.0 | 177.8 |
| 1975 | 7.9 | * | * | * | 0.5 | 1.2 | 3.3 | 8.8 | 20.7 | 50.9 | 85.5 | 159.6 |
| 1974 | 7.9 | * | * | * | 0.4 | 1.3 | 4.0 | 9.6 | 21.9 | 49.6 | 86.1 | 155.6 |
| 1973 | 7.9 | * | * | * | 0.4 | 0.9 | 3.7 | 10.7 | 22.4 | 51.2 | 80.1 | 171.7 |
| 1972 | 8.3 | * | * | * | 0.4 | 1.9 | 5.0 | 11.0 | 25.6 | 50.3 | 79.4 | 148.0 |
| 1971 | 8.4 | * | * | * | 0.4 | 1.5 | 4.7 | 11.8 | 25.4 | 50.6 | 86.2 | 162.5 |
| 1970 | 8.6 | * | * | * | 0.5 | 1.3 | 4.4 | 11.2 | 26.5 | 54.5 | 92.9 | 144.7 |
| 1969 | 8.4 | * | * | * | 0.6 | 1.5 | 4.7 | 12.1 | 26.0 | 53.6 | 85.3 | 164.2 |
| 1968 | 8.7 | * | * | * | 0.6 | 1.5 | 4.7 | 11.6 | 28.9 | 51.0 | 92.3 | 186.3 |
| ALL OTHER, MALE | | | | | | | | | | | | |
| 1978 | 8.6 | * | * | * | * | 1.2 | 3.8 | 10.9 | 28.6 | 53.0 | 105.6 | 182.8 |
| 1977 | 8.5 | * | * | * | * | 1.0 | 3.0 | 10.2 | 28.3 | 60.1 | 99.6 | 183.9 |
| 1976 | 9.2 | * | * | * | * | 1.6 | 4.1 | 12.1 | 31.8 | 60.4 | 106.6 | 169.1 |
| 1975 | 8.8 | * | * | * | 0.7 | 1.2 | 4.0 | 11.6 | 27.1 | 59.7 | 108.3 | 168.5 |
| 1974 | 9.0 | * | * | * | * | 1.4 | 5.6 | 12.2 | 30.9 | 63.1 | 100.4 | 146.0 |
| 1973 | 8.8 | * | * | * | * | * | 4.5 | 14.1 | 28.7 | 61.7 | 95.5 | 166.7 |
| 1972 | 9.3 | * | * | * | * | 1.4 | 5.5 | 14.7 | 34.8 | 58.8 | 106.9 | 148.9 |
| 1971 | 9.7 | * | * | * | * | 1.6 | 6.6 | 15.3 | 33.5 | 61.3 | 106.8 | 185.1 |
| 1970 | 9.6 | * | * | * | * | 1.4 | 5.6 | 14.2 | 33.1 | 68.6 | 103.6 | 140.7 |
| 1969 | 9.4 | * | * | * | * | 1.5 | 5.3 | 15.6 | 31.7 | 66.9 | 94.1 | 168.2 |
| 1968 | 10.0 | * | * | * | * | 2.0 | 5.7 | 14.0 | 38.8 | 62.3 | 111.8 | 165.9 |
| ALL OTHER, FEMALE | | | | | | | | | | | | |
| 1978 | 7.2 | * | * | * | * | 1.0 | 2.4 | 5.4 | 14.2 | 36.1 | 80.7 | 179.8 |
| 1977 | 7.2 | * | * | * | * | 1.0 | 3.1 | 7.4 | 15.1 | 36.6 | 70.0 | 167.3 |
| 1976 | 7.1 | * | * | * | * | 1.1 | 2.6 | 7.3 | 13.8 | 37.4 | 73.7 | 182.7 |
| 1975 | 7.1 | * | * | * | * | 1.2 | 2.8 | 6.4 | 15.4 | 44.0 | 69.8 | 152.7 |
| 1974 | 6.9 | * | * | * | * | 1.2 | 2.6 | 7.4 | 14.3 | 38.8 | 76.2 | 161.2 |
| 1973 | 7.1 | * | * | * | * | * | 3.1 | 7.8 | 17.0 | 42.8 | 69.5 | 174.7 |
| 1972 | 7.3 | * | * | * | * | 2.3 | 4.6 | 7.9 | 17.7 | 43.4 | 59.8 | 147.4 |
| 1971 | 7.1 | * | * | * | * | 1.4 | 3.1 | 8.7 | 18.3 | 41.9 | 71.1 | 147.9 |
| 1970 | 7.6 | * | * | * | * | 1.2 | 3.4 | 8.6 | 20.7 | 42.8 | 84.7 | 147.5 |
| 1969 | 7.5 | * | * | * | * | 1.5 | 4.1 | 9.0 | 20.9 | 42.7 | 78.1 | 161.5 |
| 1968 | 7.4 | * | * | * | * | * | 3.7 | 9.5 | 20.0 | 41.6 | 77.4 | 200.0 |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| OTHER DISEASES OF ARTERIES, ARTERIOLES, AND CAPILLARIES (441-448) | | | | | | | | | | | | |
| BLACK, BOTH SEXES | | | | | | | | | | | | |
| 1978 | 8.5 | * | * | * | 0.4 | 1.3 | 3.6 | 8.8 | 22.0 | 45.8 | 91.1 | 189.3 |
| 1977 | 8.3 | * | * | * | 0.5 | 1.1 | 3.5 | 9.4 | 22.2 | 47.8 | 80.8 | 179.0 |
| 1976 | 8.7 | * | * | * | 0.5 | 1.5 | 3.7 | 10.5 | 23.3 | 49.6 | 85.8 | 189.7 |
| 1975 | 8.4 | * | * | * | 0.5 | 1.3 | 3.6 | 9.5 | 21.7 | 52.8 | 85.5 | 161.5 |
| 1974 | 8.4 | * | * | * | 0.4 | 1.4 | 4.5 | 10.5 | 23.0 | 51.1 | 86.1 | 164.5 |
| 1973 | 8.3 | * | * | * | * | 0.9 | 4.1 | 11.3 | 23.2 | 52.7 | 78.4 | 177.2 |
| 1972 | 8.7 | * | * | * | * | 2.1 | 5.6 | 11.8 | 26.4 | 52.1 | 79.8 | 150.9 |
| 1971 | 8.7 | * | * | * | 0.4 | 1.7 | 5.1 | 12.6 | 26.5 | 52.2 | 87.6 | 159.8 |
| 1970 | 9.0 | * | * | * | 0.5 | 1.5 | 4.9 | 12.2 | 28.3 | 55.2 | 95.3 | 150.8 |
| 1969 | 8.9 | * | * | * | 0.7 | 1.6 | 5.1 | 13.1 | 27.5 | 55.8 | 86.8 | 171.1 |
| 1968 | 9.0 | * | * | * | 0.6 | 1.7 | 4.9 | 12.3 | 29.8 | 52.6 | 92.5 | 189.1 |
| BLACK, MALE | | | | | | | | | | | | |
| 1978 | 9.2 | * | * | * | * | 1.4 | 4.4 | 12.1 | 30.4 | 56.1 | 104.2 | 192.2 |
| 1977 | 8.9 | * | * | * | * | 1.2 | 3.5 | 11.3 | 29.9 | 60.8 | 96.2 | 181.6 |
| 1976 | 9.8 | * | * | * | * | 1.7 | 4.7 | 13.6 | 33.8 | 62.7 | 102.6 | 176.6 |
| 1975 | 9.1 | * | * | * | * | 1.3 | 4.1 | 12.3 | 28.1 | 61.2 | 107.2 | 171.7 |
| 1974 | 9.6 | * | * | * | * | 1.6 | 6.5 | 13.4 | 32.6 | 65.4 | 100.5 | 160.5 |
| 1973 | 9.0 | * | * | * | * | * | 5.1 | 14.9 | 29.4 | 63.1 | 93.6 | 178.0 |
| 1972 | 9.7 | * | * | * | * | 1.6 | 6.2 | 15.4 | 35.2 | 61.0 | 104.6 | 170.0 |
| 1971 | 10.1 | * | * | * | * | 1.8 | 7.1 | 16.3 | 35.7 | 63.4 | 107.5 | 177.5 |
| 1970 | 10.0 | * | * | * | * | 1.6 | 6.2 | 15.6 | 35.6 | 68.3 | 104.4 | 145.7 |
| 1969 | 10.0 | * | * | * | * | * | 5.9 | 17.0 | 33.7 | 70.1 | 97.0 | 181.1 |
| 1968 | 10.3 | * | * | * | * | 2.1 | 6.0 | 14.9 | 40.2 | 64.0 | 112.7 | 174.3 |
| BLACK, FEMALE | | | | | | | | | | | | |
| 1978 | 7.9 | * | * | * | * | 1.2 | 2.9 | 6.1 | 15.1 | 38.1 | 82.4 | 187.9 |
| 1977 | 7.8 | * | * | * | * | 1.1 | 3.5 | 7.8 | 15.9 | 38.1 | 70.9 | 177.7 |
| 1976 | 7.8 | * | * | * | * | 1.3 | 2.8 | 8.0 | 14.5 | 39.7 | 74.5 | 196.6 |
| 1975 | 7.7 | * | * | * | * | 1.3 | 3.1 | 7.0 | 16.3 | 46.5 | 70.7 | 156.0 |
| 1974 | 7.3 | * | * | * | * | 1.3 | 2.9 | 8.0 | 15.2 | 40.3 | 76.4 | 166.7 |
| 1973 | 7.5 | * | * | * | * | * | 3.3 | 8.2 | 18.1 | 44.6 | 68.2 | 176.7 |
| 1972 | 7.8 | * | * | * | * | 2.4 | 5.0 | 8.8 | 19.0 | 45.2 | 62.1 | 140.0 |
| 1971 | 7.5 | * | * | * | * | 1.5 | 3.4 | 9.3 | 18.8 | 43.3 | 73.4 | 149.3 |
| 1970 | 8.2 | * | * | * | * | 1.4 | 3.8 | 9.3 | 22.1 | 44.9 | 88.7 | 154.1 |
| 1969 | 8.0 | * | * | * | * | 1.7 | 4.5 | 9.7 | 22.2 | 44.4 | 78.9 | 165.0 |
| 1968 | 7.8 | * | * | * | * | * | 4.0 | 10.0 | 20.8 | 43.3 | 77.5 | 201.8 |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| ACUTE BRONCHITIS AND BRONCHIOLITIS (466) | | | | | | | | | | | | |
| BOTH SEXES | | | | | | | | | | | | |
| 1978 | 0.3 | 2.8 | 0.3 | * | * | 0.1 | 0.1 | 0.2 | 0.3 | 1.0 | 2.2 | 7.2 |
| 1977 | 0.3 | 2.5 | 0.4 | * | * | * | * | 0.2 | 0.4 | 0.8 | 2.0 | 5.9 |
| 1976 | 0.4 | 3.0 | 0.4 | * | * | 0.1 | 0.1 | 0.2 | 0.5 | 1.1 | 2.5 | 8.0 |
| 1975 | 0.3 | 3.3 | 0.4 | * | 0.0 | * | 0.1 | 0.2 | 0.4 | 0.9 | 2.2 | 4.8 |
| 1974 | 0.4 | 3.8 | 0.4 | 0.1 | * | * | * | 0.2 | 0.4 | 1.1 | 2.1 | 5.5 |
| 1973 | 0.4 | 5.4 | 0.5 | * | * | 0.1 | 0.1 | 0.3 | 0.5 | 1.1 | 2.5 | 6.0 |
| 1972 | 0.5 | 8.4 | 0.9 | 0.1 | 0.1 | 0.1 | 0.1 | 0.3 | 0.5 | 1.5 | 2.6 | 7.4 |
| 1971 | 0.6 | 11.8 | 0.7 | 0.1 | 0.1 | 0.1 | 0.2 | 0.3 | 0.5 | 0.9 | 2.4 | 5.7 |
| 1970 | 0.6 | 13.5 | 1.0 | 0.1 | 0.1 | 0.1 | 0.2 | 0.3 | 0.6 | 1.2 | 2.2 | 5.6 |
| 1969 | 0.6 | 13.2 | 1.0 | 0.1 | * | * | 0.2 | 0.3 | 0.7 | 1.2 | 2.3 | 7.1 |
| 1968 | 0.7 | 11.5 | 1.0 | 0.1 | 0.1 | 0.1 | 0.2 | 0.4 | 0.9 | 1.5 | 3.1 | 10.4 |
| MALE | | | | | | | | | | | | |
| 1978 | 0.4 | 3.4 | * | * | * | * | * | 0.2 | 0.4 | 1.4 | 3.3 | 9.0 |
| 1977 | 0.4 | 2.9 | 0.4 | * | * | * | * | 0.2 | 0.5 | 1.2 | 3.3 | 6.5 |
| 1976 | 0.5 | 4.0 | 0.5 | * | * | * | * | 0.2 | 0.6 | 1.7 | 3.4 | 12.0 |
| 1975 | 0.4 | 3.7 | 0.5 | * | * | * | * | 0.3 | 0.5 | 1.2 | 3.2 | 5.6 |
| 1974 | 0.4 | 4.5 | 0.5 | * | * | * | * | 0.2 | 0.5 | 1.4 | 3.3 | 6.5 |
| 1973 | 0.5 | 6.1 | 0.6 | * | * | * | * | 0.3 | 0.6 | 1.6 | 3.7 | 6.1 |
| 1972 | 0.7 | 9.6 | 0.9 | * | * | * | * | 0.4 | 0.7 | 2.1 | 4.1 | 11.9 |
| 1971 | 0.6 | 12.6 | 0.9 | * | * | * | * | 0.3 | 0.7 | 1.4 | 3.3 | 6.5 |
| 1970 | 0.7 | 14.9 | 1.2 | 0.1 | * | * | 0.2 | 0.3 | 0.9 | 1.8 | 2.6 | 7.0 |
| 1969 | 0.7 | 14.1 | 1.1 | 0.1 | * | * | * | 0.4 | 0.9 | 1.9 | 2.8 | 8.5 |
| 1968 | 0.9 | 13.4 | 1.1 | 0.1 | 0.1 | * | 0.3 | 0.5 | 1.2 | 2.3 | 4.3 | 10.5 |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| ACUTE BRONCHITIS AND BRONCHIOLITIS (466) | | | | | | | | | | | | |
| FEMALE | | | | | | | | | | | | |
| 1978 | 0.3 | 2.2 | * | * | * | * | * | * | 0.2 | 0.6 | 1.5 | 6.4 |
| 1977 | 0.3 | 2.1 | 0.4 | * | * | * | * | 0.2 | 0.2 | 0.5 | 1.1 | 5.6 |
| 1976 | 0.3 | 2.0 | * | * | * | * | * | 0.2 | 0.4 | 0.7 | 2.0 | 6.1 |
| 1975 | 0.3 | 2.9 | 0.3 | * | * | * | * | * | 0.3 | 0.7 | 1.5 | 4.5 |
| 1974 | 0.3 | 3.0 | * | * | * | * | * | 0.2 | 0.3 | 0.8 | 1.4 | 5.0 |
| 1973 | 0.4 | 4.8 | 0.5 | * | * | * | * | 0.2 | 0.5 | 0.7 | 1.7 | 5.9 |
| 1972 | 0.4 | 7.2 | 0.8 | * | * | * | * | 0.2 | 0.4 | 1.0 | 1.6 | 5.1 |
| 1971 | 0.5 | 11.0 | 0.6 | * | * | * | 0.2 | 0.3 | 0.3 | 0.5 | 1.9 | 5.3 |
| 1970 | 0.5 | 12.1 | 0.8 | * | * | * | 0.2 | 0.3 | 0.4 | 0.8 | 1.9 | 4.9 |
| 1969 | 0.6 | 12.2 | 0.8 | * | * | * | 0.2 | 0.3 | 0.5 | 0.7 | 2.0 | 6.4 |
| 1968 | 0.6 | 9.5 | 0.8 | * | * | * | 0.2 | 0.4 | 0.6 | 0.8 | 2.3 | 10.4 |
| WHITE, BOTH SEXES | | | | | | | | | | | | |
| 1978 | 0.3 | 2.2 | 0.3 | * | * | * | * | 0.2 | 0.3 | 1.0 | 2.2 | 7.5 |
| 1977 | 0.3 | 2.1 | 0.3 | * | * | * | * | 0.2 | 0.4 | 0.8 | 2.1 | 6.2 |
| 1976 | 0.4 | 2.5 | 0.3 | * | * | * | 0.1 | 0.2 | 0.4 | 1.2 | 2.7 | 8.2 |
| 1975 | 0.3 | 2.9 | 0.3 | * | * | * | * | 0.2 | 0.4 | 1.0 | 2.2 | 5.1 |
| 1974 | 0.3 | 3.2 | 0.3 | * | * | * | * | 0.2 | 0.4 | 1.1 | 2.2 | 5.9 |
| 1973 | 0.4 | 4.7 | 0.5 | * | * | * | 0.1 | 0.3 | 0.5 | 1.1 | 2.6 | 6.1 |
| 1972 | 0.5 | 5.5 | 0.8 | * | * | 0.1 | 0.1 | 0.3 | 0.5 | 1.5 | 2.7 | 7.8 |
| 1971 | 0.5 | 6.8 | 0.5 | 0.1 | * | * | 0.1 | 0.3 | 0.5 | 0.9 | 2.5 | 5.9 |
| 1970 | 0.5 | 7.6 | 0.8 | 0.1 | * | 0.1 | 0.2 | 0.3 | 0.6 | 1.2 | 2.3 | 6.0 |
| 1969 | 0.6 | 9.2 | 0.8 | 0.1 | * | * | 0.2 | 0.3 | 0.7 | 1.2 | 2.4 | 7.3 |
| 1968 | 0.7 | 8.7 | 0.8 | 0.1 | 0.1 | * | 0.2 | 0.3 | 0.8 | 1.6 | 3.2 | 10.7 |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| ACUTE BRONCHITIS AND BRONCHIOLITIS (466) | | | | | | | | | | | | |
| WHITE, MALE | | | | | | | | | | | | |
| 1978 | 0.4 | 2.3 | * | * | * | * | * | 0.2 | 0.4 | 1.4 | 3.4 | 9.6 |
| 1977 | 0.4 | 2.4 | 0.4 | * | * | * | * | 0.2 | 0.5 | 1.3 | 3.6 | 7.0 |
| 1976 | 0.4 | 3.2 | 0.4 | * | * | * | * | * | 0.6 | 1.7 | 3.6 | 12.2 |
| 1975 | 0.4 | 3.3 | 0.5 | * | * | * | * | 0.2 | 0.6 | 1.3 | 3.3 | 5.9 |
| 1974 | 0.4 | 3.8 | 0.4 | * | * | * | * | * | 0.5 | 1.4 | 3.4 | 7.0 |
| 1973 | 0.5 | 5.3 | 0.5 | * | * | * | * | 0.3 | 0.6 | 1.6 | 4.0 | 6.5 |
| 1972 | 0.6 | 6.6 | 0.8 | * | * | * | * | 0.4 | 0.7 | 2.2 | 4.5 | 12.2 |
| 1971 | 0.5 | 7.3 | 0.7 | * | * | * | * | 0.2 | 0.8 | 1.4 | 3.4 | 6.9 |
| 1970 | 0.6 | 9.1 | 1.0 | * | * | * | 0.2 | 0.3 | 0.8 | 1.9 | 2.7 | 7.6 |
| 1969 | 0.6 | 10.9 | 1.0 | 0.1 | * | * | * | 0.3 | 0.8 | 1.8 | 2.9 | 8.7 |
| 1968 | 0.8 | 10.7 | 1.0 | * | 0.2 | * | 0.2 | 0.4 | 1.1 | 2.4 | 4.4 | 10.5 |
| WHITE, FEMALE | | | | | | | | | | | | |
| 1978 | 0.3 | 2.2 | * | * | * | * | * | * | 0.2 | 0.7 | 1.6 | 6.6 |
| 1977 | 0.3 | 1.8 | * | * | * | * | * | 0.2 | 0.3 | 0.5 | 1.2 | 5.9 |
| 1976 | 0.3 | 1.7 | * | * | * | * | * | * | 0.3 | 0.8 | 2.1 | 6.4 |
| 1975 | 0.3 | 2.4 | * | * | * | * | * | * | 0.3 | 0.7 | 1.6 | 4.8 |
| 1974 | 0.3 | 2.5 | * | * | * | * | * | 0.2 | 0.3 | 0.9 | 1.5 | 5.3 |
| 1973 | 0.4 | 4.2 | 0.4 | * | * | * | * | 0.2 | 0.5 | 0.7 | 1.8 | 6.0 |
| 1972 | 0.4 | 4.3 | 0.7 | * | * | * | * | 0.2 | 0.3 | 1.0 | 1.6 | 5.5 |
| 1971 | 0.4 | 6.3 | 0.4 | * | * | * | * | 0.3 | 0.3 | 0.5 | 1.9 | 5.3 |
| 1970 | 0.4 | 6.1 | 0.6 | * | * | * | * | 0.3 | 0.4 | 0.8 | 2.0 | 5.2 |
| 1969 | 0.5 | 7.4 | 0.7 | * | * | * | 0.2 | 0.3 | 0.5 | 0.7 | 2.0 | 6.5 |
| 1968 | 0.5 | 6.6 | 0.6 | * | * | * | * | 0.3 | 0.6 | 0.9 | 2.4 | 10.8 |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| ACUTE BRONCHITIS AND BRONCHIOLITIS (466) | | | | | | | | | | | | |
| ALL OTHER, BOTH SEXES | | | | | | | | | | | | |
| 1978 | 0.3 | 5.8 | * | * | * | * | * | * | * | * | * | * |
| 1977 | 0.3 | 4.3 | * | * | * | * | * | * | * | * | * | * |
| 1976 | 0.4 | 5.7 | * | * | * | * | * | * | * | * | * | * |
| 1975 | 0.3 | 5.3 | * | * | * | * | * | * | * | * | * | * |
| 1974 | 0.4 | 6.8 | 1.0 | * | * | * | * | * | * | * | * | * |
| 1973 | 0.5 | 8.9 | * | * | * | * | * | * | * | * | * | * |
| 1972 | 0.9 | 23.3 | 1.5 | * | * | * | * | * | * | * | * | * |
| 1971 | 1.3 | 38.4 | 1.8 | * | * | * | * | * | * | * | * | * |
| 1970 | 1.5 | 44.9 | 2.2 | * | * | * | * | * | * | * | * | * |
| 1969 | 1.2 | 35.8 | 1.8 | * | * | * | * | * | * | * | * | * |
| 1968 | 1.2 | 26.6 | 1.7 | * | * | * | * | 1.0 | 1.2 | * | * | * |
| ALL OTHER, MALE | | | | | | | | | | | | |
| 1978 | 0.5 | 9.0 | * | * | * | * | * | * | * | * | * | * |
| 1977 | 0.3 | * | * | * | * | * | * | * | * | * | * | * |
| 1976 | 0.5 | 7.9 | * | * | * | * | * | * | * | * | * | * |
| 1975 | 0.4 | * | * | * | * | * | * | * | * | * | * | * |
| 1974 | 0.4 | 8.1 | * | * | * | * | * | * | * | * | * | * |
| 1973 | 0.5 | 10.2 | * | * | * | * | * | * | * | * | * | * |
| 1972 | 1.1 | 24.9 | * | * | * | * | * | * | * | * | * | * |
| 1971 | 1.5 | 41.1 | 2.1 | * | * | * | * | * | * | * | * | * |
| 1970 | 1.7 | 46.3 | 2.2 | * | * | * | * | * | * | * | * | * |
| 1969 | 1.3 | 32.6 | 1.8 | * | * | * | * | * | * | * | * | * |
| 1968 | 1.3 | 28.3 | 1.7 | * | * | * | * | * | * | * | * | * |
| ALL OTHER, FEMALE | | | | | | | | | | | | |
| 1978 | 0.2 | * | * | * | * | * | * | * | * | * | * | * |
| 1977 | 0.3 | * | * | * | * | * | * | * | * | * | * | * |
| 1976 | 0.3 | * | * | * | * | * | * | * | * | * | * | * |
| 1975 | 0.3 | * | * | * | * | * | * | * | * | * | * | * |
| 1974 | 0.3 | * | * | * | * | * | * | * | * | * | * | * |
| 1973 | 0.4 | 7.7 | * | * | * | * | * | * | * | * | * | * |
| 1972 | 0.8 | 21.6 | * | * | * | * | * | * | * | * | * | * |
| 1971 | 1.1 | 35.6 | * | * | * | * | * | * | * | * | * | * |
| 1970 | 1.3 | 43.5 | 2.2 | * | * | * | * | * | * | * | * | * |
| 1969 | 1.1 | 39.1 | * | * | * | * | * | * | * | * | * | * |
| 1968 | 1.1 | 24.8 | * | * | * | * | * | * | * | * | * | * |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| ACUTE BRONCHITIS AND BRONCHIOLITIS (466) | | | | | | | | | | | | |
| BLACK, BOTH SEXES | | | | | | | | | | | | |
| 1978 | 0.3 | 5.7 | * | * | * | * | * | * | * | * | * | * |
| 1977 | 0.3 | 4.3 | * | * | * | * | * | * | * | * | * | * |
| 1976 | 0.4 | 5.7 | * | * | * | * | * | * | * | * | * | * |
| 1975 | 0.4 | 5.9 | * | * | * | * | * | * | * | * | * | * |
| 1974 | 0.4 | 7.4 | * | * | * | * | * | * | * | * | * | * |
| 1973 | 0.5 | 9.3 | * | * | * | * | * | * | * | * | * | * |
| 1972 | 0.9 | 25.2 | 1.4 | * | * | * | * | * | * | * | * | * |
| 1971 | 1.4 | 42.6 | 1.9 | * | * | * | * | * | * | * | * | * |
| 1970 | 1.6 | 49.1 | 2.4 | * | * | * | * | * | * | * | * | * |
| 1969 | 1.3 | 38.7 | 1.8 | * | * | * | * | * | * | * | * | * |
| 1968 | 1.2 | 26.6 | 1.7 | * | * | * | * | 1.1 | 1.3 | * | * | * |
| BLACK, MALE | | | | | | | | | | | | |
| 1978 | 0.5 | 8.9 | * | * | * | * | * | * | * | * | * | * |
| 1977 | 0.3 | * | * | * | * | * | * | * | * | * | * | * |
| 1976 | 0.5 | * | * | * | * | * | * | * | * | * | * | * |
| 1975 | 0.5 | * | * | * | * | * | * | * | * | * | * | * |
| 1974 | 0.4 | * | * | * | * | * | * | * | * | * | * | * |
| 1973 | 0.6 | 10.3 | * | * | * | * | * | * | * | * | * | * |
| 1972 | 1.1 | 26.6 | * | * | * | * | * | * | * | * | * | * |
| 1971 | 1.6 | 45.5 | 2.4 | * | * | * | * | * | * | * | * | * |
| 1970 | 1.8 | 50.3 | 2.3 | * | * | * | * | * | * | * | * | * |
| 1969 | 1.4 | 35.8 | * | * | * | * | * | * | * | * | * | * |
| 1968 | 1.3 | 26.5 | * | * | * | * | * | * | * | * | * | * |
| BLACK, FEMALE | | | | | | | | | | | | |
| 1978 | 0.2 | * | * | * | * | * | * | * | * | * | * | * |
| 1977 | 0.3 | * | * | * | * | * | * | * | * | * | * | * |
| 1976 | 0.3 | * | * | * | * | * | * | * | * | * | * | * |
| 1975 | 0.3 | * | * | * | * | * | * | * | * | * | * | * |
| 1974 | 0.3 | * | * | * | * | * | * | * | * | * | * | * |
| 1973 | 0.4 | * | * | * | * | * | * | * | * | * | * | * |
| 1972 | 0.7 | 23.6 | * | * | * | * | * | * | * | * | * | * |
| 1971 | 1.2 | 39.6 | * | * | * | * | * | * | * | * | * | * |
| 1970 | 1.4 | 47.8 | 2.5 | * | * | * | * | * | * | * | * | * |
| 1969 | 1.2 | 41.7 | * | * | * | * | * | * | * | * | * | * |
| 1968 | 1.2 | 26.8 | * | * | * | * | * | * | * | * | * | * |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| PNEUMONIA AND INFLUENZA 470-474, 480-486 | | | | | | | | | | | | |
| BOTH SEXES | | | | | | | | | | | | |
| 1978 | 26.3 | 46.1 | 2.9 | 0.8 | 1.3 | 2.1 | 4.4 | 10.0 | 23.5 | 65.5 | 244.6 | 884.3 |
| 1977 | 23.3 | 51.3 | 3.0 | 0.9 | 1.2 | 2.0 | 4.4 | 9.6 | 21.5 | 58.3 | 222.2 | 763.1 |
| 1976 | 28.4 | 63.0 | 3.8 | 1.0 | 1.5 | 2.4 | 5.4 | 11.6 | 25.9 | 69.9 | 274.3 | 994.6 |
| 1975 | 25.8 | 69.8 | 4.1 | 0.9 | 1.7 | 2.4 | 5.3 | 11.1 | 26.7 | 70.0 | 252.7 | 801.0 |
| 1974 | 25.7 | 85.3 | 4.6 | 1.3 | 1.7 | 2.5 | 5.7 | 11.8 | 26.4 | 68.6 | 249.9 | 824.9 |
| 1973 | 29.6 | 113.9 | 5.8 | 1.4 | 2.0 | 3.1 | 6.9 | 14.5 | 31.4 | 81.9 | 287.3 | 923.4 |
| 1972 | 29.9 | 131.0 | 6.5 | 1.4 | 2.1 | 3.3 | 7.8 | 16.0 | 32.8 | 86.6 | 283.4 | 870.4 |
| 1971 | 27.7 | 150.0 | 7.1 | 1.4 | 2.2 | 3.3 | 6.9 | 13.4 | 29.0 | 74.7 | 254.7 | 846.7 |
| 1970 | 30.9 | 180.8 | 7.6 | 1.6 | 2.4 | 3.8 | 8.5 | 17.3 | 36.2 | 90.2 | 272.8 | 814.5 |
| 1969 | 33.9 | 209.2 | 8.5 | 1.8 | 2.5 | 4.1 | 9.0 | 18.1 | 39.0 | 99.0 | 302.9 | 1,026.2 |
| 1968 | 36.9 | 234.9 | 9.9 | 1.8 | 2.8 | 4.4 | 9.3 | 19.9 | 41.9 | 106.0 | 330.7 | 1,167.8 |
| MALE | | | | | | | | | | | | |
| 1978 | 28.5 | 50.0 | 2.8 | 0.9 | 1.5 | 2.6 | 5.4 | 13.1 | 32.7 | 96.0 | 349.4 | 1,159.4 |
| 1977 | 25.8 | 57.6 | 3.4 | 0.9 | 1.5 | 2.4 | 5.5 | 12.9 | 30.5 | 85.9 | 319.7 | 1,010.4 |
| 1976 | 30.7 | 68.2 | 3.9 | 0.9 | 1.6 | 2.7 | 6.7 | 14.9 | 36.2 | 101.1 | 385.4 | 1,270.0 |
| 1975 | 29.0 | 76.3 | 4.1 | 0.9 | 2.0 | 2.7 | 6.5 | 15.2 | 37.2 | 102.0 | 366.0 | 1,036.0 |
| 1974 | 28.7 | 93.2 | 4.8 | 1.3 | 1.9 | 2.9 | 7.1 | 14.9 | 37.3 | 99.8 | 355.3 | 1,042.7 |
| 1973 | 33.2 | 126.3 | 6.1 | 1.4 | 2.4 | 3.7 | 8.2 | 18.9 | 43.7 | 119.5 | 405.8 | 1,138.7 |
| 1972 | 34.0 | 149.9 | 6.8 | 1.4 | 2.4 | 3.7 | 9.4 | 21.3 | 46.3 | 125.8 | 397.1 | 1,075.1 |
| 1971 | 31.4 | 169.6 | 7.5 | 1.3 | 2.6 | 3.8 | 8.3 | 17.5 | 41.0 | 107.5 | 352.0 | 1,046.1 |
| 1970 | 35.5 | 203.4 | 8.6 | 1.6 | 2.8 | 4.5 | 10.4 | 22.6 | 50.2 | 129.9 | 376.3 | 959.8 |
| 1969 | 38.6 | 235.3 | 9.3 | 1.8 | 2.9 | 4.4 | 10.7 | 23.7 | 54.0 | 142.3 | 406.0 | 1,214.4 |
| 1968 | 41.8 | 260.9 | 10.3 | 1.8 | 3.3 | 5.0 | 11.5 | 25.9 | 58.3 | 152.2 | 431.5 | 1,375.5 |

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| PNEUMONIA AND INFLUENZA 470-474, 480-486 | | | | | | | | | | | | |
| FEMALE | | | | | | | | | | | | |
| 1978 | 24.1 | 42.0 | 2.9 | 0.8 | 1.1 | 1.6 | 3.3 | 7.0 | 15.4 | 42.2 | 182.3 | 760.1 |
| 1977 | 20.9 | 44.7 | 2.7 | 1.0 | 1.0 | 1.5 | 3.3 | 6.5 | 13.5 | 37.1 | 163.9 | 649.5 |
| 1976 | 26.3 | 57.5 | 3.8 | 1.0 | 1.3 | 2.1 | 4.2 | 8.5 | 16.7 | 45.9 | 207.2 | 865.3 |
| 1975 | 22.8 | 63.1 | 4.0 | 1.0 | 1.4 | 2.2 | 4.2 | 7.3 | 17.3 | 45.5 | 183.7 | 687.3 |
| 1974 | 22.8 | 76.9 | 4.4 | 1.2 | 1.4 | 2.1 | 4.4 | 8.9 | 16.6 | 44.6 | 184.9 | 717.1 |
| 1973 | 26.1 | 100.8 | 5.5 | 1.4 | 1.6 | 2.5 | 5.7 | 10.5 | 20.4 | 53.0 | 213.2 | 815.1 |
| 1972 | 26.0 | 111.3 | 6.3 | 1.4 | 1.9 | 2.9 | 6.3 | 11.1 | 20.8 | 56.3 | 210.9 | 765.7 |
| 1971 | 24.1 | 129.5 | 6.7 | 1.4 | 1.9 | 2.8 | 5.6 | 9.5 | 18.3 | 49.4 | 191.5 | 742.7 |
| 1970 | 26.5 | 157.4 | 6.6 | 1.6 | 2.1 | 3.2 | 6.6 | 12.3 | 23.6 | 59.3 | 204.4 | 733.2 |
| 1969 | 29.5 | 182.2 | 7.7 | 1.9 | 2.1 | 3.8 | 7.4 | 12.9 | 25.4 | 65.2 | 233.7 | 924.4 |
| 1968 | 32.2 | 207.8 | 9.5 | 1.8 | 2.4 | 3.9 | 7.3 | 14.3 | 27.1 | 69.6 | 261.7 | 1,055.1 |
| WHITE, BOTH SEXES | | | | | | | | | | | | |
| 1978 | 26.8 | 35.0 | 2.4 | 0.8 | 1.1 | 1.6 | 3.2 | 8.0 | 21.1 | 62.8 | 247.1 | 911.3 |
| 1977 | 23.5 | 38.3 | 2.5 | 0.9 | 1.1 | 1.4 | 3.2 | 7.6 | 19.2 | 55.6 | 224.2 | 783.3 |
| 1976 | 29.0 | 47.5 | 3.1 | 0.9 | 1.3 | 1.8 | 4.1 | 9.5 | 23.3 | 67.3 | 278.0 | 1,029.0 |
| 1975 | 26.1 | 53.7 | 3.5 | 0.9 | 1.6 | 1.9 | 4.0 | 9.2 | 24.3 | 67.3 | 256.2 | 821.7 |
| 1974 | 25.7 | 64.4 | 3.7 | 1.2 | 1.6 | 1.9 | 4.2 | 9.7 | 23.9 | 64.9 | 253.4 | 849.4 |
| 1973 | 29.4 | 86.4 | 4.9 | 1.3 | 1.7 | 2.3 | 5.2 | 11.8 | 28.3 | 77.3 | 290.2 | 951.7 |
| 1972 | 29.5 | 98.1 | 5.5 | 1.4 | 1.9 | 2.4 | 5.8 | 13.3 | 29.8 | 82.5 | 286.5 | 899.1 |
| 1971 | 26.9 | 114.1 | 5.9 | 1.3 | 1.9 | 2.4 | 5.1 | 11.0 | 25.8 | 71.1 | 255.7 | 872.6 |
| 1970 | 29.7 | 135.3 | 6.0 | 1.4 | 2.1 | 2.9 | 6.2 | 14.0 | 32.3 | 84.8 | 272.9 | 847.0 |
| 1969 | 32.6 | 153.2 | 6.6 | 1.7 | 2.2 | 2.9 | 6.5 | 15.1 | 35.0 | 93.6 | 302.7 | 1,059.5 |
| 1968 | 35.1 | 169.2 | 7.4 | 1.6 | 2.6 | 3.2 | 6.5 | 16.5 | 37.1 | 100.4 | 330.1 | 1,197.7 |

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL 1 YEAR | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-----------------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| PNEUMONIA AND INFLUENZA 470-474, 480-486 | | | | | | | | | | | | |
| WHITE, MALE | | | | | | | | | | | | |
| 1978 | 28.4 | 37.8 | 2.3 | 0.8 | 1.4 | 1.9 | 3.8 | 10.2 | 28.9 | 91.4 | 352.1 | 1,194.6 |
| 1977 | 25.5 | 43.9 | 2.8 | 0.9 | 1.3 | 1.7 | 3.7 | 10.0 | 27.0 | 81.9 | 322.8 | 1,030.3 |
| 1976 | 30.6 | 51.1 | 3.1 | 0.9 | 1.4 | 2.0 | 4.9 | 11.8 | 32.0 | 96.9 | 390.1 | 1,317.8 |
| 1975 | 28.7 | 58.3 | 3.3 | 0.8 | 1.9 | 2.0 | 4.6 | 12.2 | 33.4 | 97.9 | 370.2 | 1,062.6 |
| 1974 | 28.1 | 69.6 | 4.0 | 1.3 | 1.8 | 2.1 | 4.9 | 11.8 | 33.3 | 94.8 | 359.8 | 1,075.1 |
| 1973 | 32.4 | 98.0 | 4.9 | 1.3 | 2.2 | 2.7 | 6.0 | 15.0 | 39.0 | 113.2 | 408.7 | 1,174.9 |
| 1972 | 33.0 | 115.9 | 5.9 | 1.4 | 2.2 | 2.5 | 6.7 | 17.2 | 41.7 | 119.3 | 400.0 | 1,118.3 |
| 1971 | 30.0 | 130.6 | 6.1 | 1.2 | 2.1 | 2.6 | 5.7 | 14.1 | 36.2 | 102.3 | 353.3 | 1,085.3 |
| 1970 | 33.7 | 154.8 | 6.6 | 1.4 | 2.5 | 3.3 | 7.5 | 17.9 | 44.7 | 123.1 | 376.5 | 1,003.0 |
| 1969 | 36.5 | 177.9 | 6.8 | 1.7 | 2.6 | 3.0 | 7.3 | 19.2 | 48.6 | 134.9 | 404.4 | 1,258.6 |
| 1968 | 39.3 | 193.3 | 7.7 | 1.6 | 2.9 | 3.4 | 7.9 | 21.1 | 51.8 | 145.0 | 429.1 | 1,415.8 |
| WHITE, FEMALE | | | | | | | | | | | | |
| 1978 | 25.3 | 32.0 | 2.4 | 0.7 | 0.9 | 1.2 | 2.7 | 6.0 | 14.2 | 41.0 | 185.4 | 785.0 |
| 1977 | 21.7 | 32.5 | 2.2 | 0.9 | 0.9 | 1.2 | 2.6 | 5.4 | 12.3 | 35.6 | 166.1 | 671.5 |
| 1976 | 27.4 | 43.7 | 3.0 | 0.9 | 1.2 | 1.6 | 3.3 | 7.4 | 15.5 | 44.6 | 211.2 | 896.4 |
| 1975 | 23.6 | 48.8 | 3.6 | 0.9 | 1.2 | 1.8 | 3.4 | 6.3 | 16.1 | 43.9 | 187.5 | 707.7 |
| 1974 | 23.4 | 59.0 | 3.4 | 1.2 | 1.3 | 1.7 | 3.5 | 7.8 | 15.6 | 42.1 | 188.4 | 739.3 |
| 1973 | 26.5 | 74.3 | 4.8 | 1.3 | 1.3 | 2.0 | 4.4 | 8.8 | 18.7 | 49.8 | 216.8 | 842.1 |
| 1972 | 26.2 | 79.5 | 5.1 | 1.4 | 1.6 | 2.4 | 4.9 | 9.7 | 19.2 | 54.3 | 214.9 | 788.8 |
| 1971 | 24.0 | 96.7 | 5.6 | 1.3 | 1.7 | 2.3 | 4.4 | 8.0 | 16.4 | 47.0 | 193.2 | 763.6 |
| 1970 | 25.9 | 114.9 | 5.3 | 1.4 | 1.7 | 2.6 | 5.0 | 10.4 | 21.2 | 55.2 | 205.1 | 761.7 |
| 1969 | 28.9 | 127.3 | 6.3 | 1.7 | 1.9 | 2.8 | 5.8 | 11.2 | 22.7 | 61.4 | 235.1 | 955.4 |
| 1968 | 31.2 | 143.9 | 7.1 | 1.6 | 2.2 | 2.9 | 5.3 | 12.2 | 23.8 | 65.4 | 262.9 | 1,081.6 |

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| PNEUMONIA AND INFLUENZA 470-474, 480-486 | | | | | | | | | | | | |
| ALL OTHER, BOTH SEXES | | | | | | | | | | | | |
| 1978 | 22.9 | 99.8 | 5.2 | 1.2 | 2.3 | 5.7 | 12.1 | 25.0 | 45.2 | 90.6 | 218.9 | 575.6 |
| 1977 | 21.7 | 114.1 | 5.5 | 1.2 | 2.0 | 5.4 | 12.5 | 24.8 | 41.9 | 83.8 | 201.4 | 531.9 |
| 1976 | 25.1 | 138.6 | 7.6 | 1.2 | 2.5 | 6.2 | 14.7 | 28.4 | 49.4 | 94.9 | 236.8 | 596.1 |
| 1975 | 24.3 | 150.7 | 7.0 | 1.4 | 2.4 | 6.3 | 14.7 | 27.3 | 48.9 | 95.8 | 216.3 | 564.4 |
| 1974 | 25.4 | 188.7 | 9.1 | 1.6 | 2.2 | 6.5 | 16.6 | 28.8 | 49.2 | 104.5 | 211.1 | 540.0 |
| 1973 | 31.3 | 249.6 | 10.7 | 1.7 | 3.4 | 8.3 | 19.6 | 37.8 | 60.6 | 127.7 | 253.2 | 593.7 |
| 1972 | 32.7 | 299.4 | 12.1 | 1.7 | 3.5 | 9.3 | 22.5 | 39.2 | 61.6 | 126.3 | 247.0 | 539.8 |
| 1971 | 32.8 | 341.7 | 13.7 | 1.7 | 4.3 | 9.2 | 20.7 | 34.5 | 60.3 | 111.1 | 241.9 | 552.5 |
| 1970 | 39.1 | 423.9 | 16.3 | 2.5 | 4.7 | 10.1 | 24.7 | 45.8 | 72.8 | 142.2 | 272.2 | 481.0 |
| 1969 | 43.3 | 525.5 | 18.5 | 2.6 | 4.3 | 12.0 | 27.7 | 45.3 | 77.3 | 153.6 | 306.0 | 650.5 |
| 1968 | 49.4 | 589.9 | 23.5 | 3.2 | 4.8 | 13.3 | 30.7 | 50.9 | 88.5 | 161.9 | 338.5 | 834.3 |
| ALL OTHER, MALE | | | | | | | | | | | | |
| 1978 | 29.4 | 109.7 | 5.3 | 1.4 | 2.5 | 7.4 | 17.6 | 37.8 | 68.4 | 138.5 | 322.2 | 798.3 |
| 1977 | 28.2 | 125.0 | 6.1 | 1.1 | 2.4 | 7.4 | 18.4 | 37.7 | 63.2 | 123.9 | 290.8 | 807.1 |
| 1976 | 31.7 | 153.4 | 7.7 | 1.2 | 2.6 | 7.7 | 20.0 | 41.8 | 75.8 | 140.2 | 342.4 | 790.9 |
| 1975 | 31.5 | 166.9 | 7.9 | 1.2 | 2.9 | 8.0 | 20.9 | 41.7 | 73.8 | 140.0 | 324.8 | 770.4 |
| 1974 | 32.5 | 212.4 | 9.0 | 1.7 | 2.4 | 8.9 | 24.1 | 42.0 | 76.0 | 147.8 | 308.7 | 708.0 |
| 1973 | 39.4 | 268.0 | 12.5 | 1.7 | 3.5 | 10.8 | 25.7 | 53.8 | 89.6 | 179.8 | 373.2 | 768.8 |
| 1972 | 41.6 | 326.7 | 11.6 | 1.5 | 3.6 | 12.5 | 30.2 | 58.3 | 91.0 | 187.9 | 365.9 | 638.3 |
| 1971 | 41.3 | 380.8 | 15.0 | 1.5 | 5.6 | 13.1 | 28.7 | 48.8 | 87.4 | 156.4 | 339.6 | 659.6 |
| 1970 | 48.9 | 467.0 | 19.5 | 2.3 | 5.3 | 13.4 | 34.1 | 65.7 | 102.6 | 193.3 | 373.5 | 581.0 |
| 1969 | 53.8 | 565.5 | 22.2 | 2.5 | 5.0 | 14.5 | 38.6 | 65.0 | 106.8 | 213.4 | 425.5 | 797.7 |
| 1968 | 60.2 | 631.3 | 24.7 | 3.0 | 5.9 | 17.0 | 41.2 | 70.9 | 121.1 | 221.8 | 459.7 | 982.9 |
| ALL OTHER, FEMALE | | | | | | | | | | | | |
| 1978 | 16.9 | 89.7 | 5.0 | 1.1 | 2.1 | 4.2 | 7.5 | 14.4 | 25.9 | 53.5 | 148.8 | 462.4 |
| 1977 | 15.9 | 102.9 | 4.8 | 1.2 | 1.5 | 3.7 | 7.5 | 14.1 | 24.1 | 52.3 | 139.9 | 383.7 |
| 1976 | 19.0 | 124.1 | 7.4 | 1.3 | 2.4 | 4.9 | 10.2 | 17.2 | 27.4 | 59.0 | 164.6 | 486.7 |
| 1975 | 17.8 | 134.0 | 6.1 | 1.6 | 2.0 | 4.8 | 9.5 | 15.1 | 27.8 | 60.7 | 141.5 | 438.7 |
| 1974 | 18.9 | 165.2 | 9.3 | 1.5 | 2.1 | 4.4 | 10.4 | 17.5 | 26.5 | 69.9 | 143.7 | 441.2 |
| 1973 | 23.9 | 230.8 | 9.0 | 1.6 | 3.3 | 6.1 | 14.5 | 24.2 | 35.8 | 85.6 | 169.5 | 487.3 |
| 1972 | 24.5 | 271.6 | 12.6 | 1.8 | 3.4 | 6.5 | 16.0 | 22.8 | 36.3 | 76.5 | 162.1 | 478.9 |
| 1971 | 25.0 | 301.8 | 12.5 | 2.0 | 3.1 | 5.8 | 14.0 | 22.1 | 36.6 | 74.0 | 169.6 | 483.6 |
| 1970 | 30.1 | 380.2 | 13.1 | 2.6 | 4.2 | 7.3 | 17.0 | 28.3 | 46.4 | 100.2 | 195.1 | 410.6 |
| 1969 | 33.7 | 485.2 | 14.8 | 2.7 | 3.6 | 10.0 | 18.6 | 28.0 | 51.3 | 104.4 | 213.8 | 550.8 |
| 1968 | 39.5 | 545.8 | 22.2 | 3.5 | 3.8 | 10.2 | 21.9 | 33.2 | 59.3 | 112.3 | 245.6 | 734.4 |

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| PNEUMONIA AND INFLUENZA 470-474, 480-486 | | | | | | | | | | | | |
| BLACK, BOTH SEXES | | | | | | | | | | | | |
| 1978 | 24.3 | 108.1 | 5.4 | 1.3 | 2.4 | 6.4 | 13.9 | 27.8 | 48.6 | 94.7 | 214.1 | 548.0 |
| 1977 | 23.0 | 123.1 | 5.9 | 1.2 | 2.1 | 6.1 | 13.9 | 27.2 | 45.2 | 86.5 | 193.9 | 514.0 |
| 1976 | 26.5 | 148.8 | 7.8 | 1.3 | 2.6 | 6.7 | 16.2 | 31.1 | 52.5 | 99.3 | 232.4 | 569.9 |
| 1975 | 25.5 | 158.6 | 7.0 | 1.4 | 2.6 | 6.8 | 16.0 | 29.1 | 52.2 | 99.7 | 212.5 | 534.6 |
| 1974 | 26.8 | 203.8 | 9.7 | 1.6 | 2.4 | 7.3 | 18.3 | 31.5 | 51.9 | 109.0 | 208.9 | 504.1 |
| 1973 | 32.5 | 267.2 | 11.3 | 1.7 | 3.6 | 9.3 | 21.5 | 40.6 | 62.5 | 130.9 | 243.3 | 567.5 |
| 1972 | 34.2 | 317.0 | 12.4 | 1.7 | 3.7 | 9.9 | 25.3 | 42.7 | 65.1 | 131.0 | 244.5 | 500.0 |
| 1971 | 34.1 | 363.6 | 14.5 | 1.8 | 4.6 | 10.0 | 22.7 | 36.8 | 62.7 | 113.1 | 234.6 | 526.2 |
| 1970 | 41.4 | 454.6 | 17.6 | 2.6 | 5.1 | 11.2 | 27.6 | 49.5 | 76.7 | 147.7 | 275.2 | 475.4 |
| 1969 | 45.3 | 559.5 | 19.3 | 2.7 | 4.5 | 13.1 | 29.9 | 48.1 | 80.8 | 159.2 | 307.9 | 647.4 |
| 1968 | 51.5 | 621.4 | 24.3 | 3.3 | 4.9 | 14.3 | 33.3 | 54.1 | 93.5 | 167.6 | 339.8 | 807.6 |
| BLACK, MALE | | | | | | | | | | | | |
| 1978 | 31.4 | 119.0 | 5.7 | 1.5 | 2.6 | 8.3 | 20.2 | 42.9 | 73.8 | 147.7 | 313.8 | 754.9 |
| 1977 | 29.7 | 133.5 | 6.7 | 1.2 | 2.6 | 8.4 | 20.7 | 41.7 | 68.7 | 129.1 | 281.7 | 765.3 |
| 1976 | 33.5 | 163.8 | 8.0 | 1.4 | 2.7 | 8.3 | 22.4 | 46.4 | 81.5 | 149.0 | 337.2 | 742.6 |
| 1975 | 33.2 | 180.0 | 8.2 | 1.2 | 3.2 | 8.7 | 23.0 | 45.0 | 80.0 | 148.0 | 319.8 | 741.3 |
| 1974 | 34.2 | 227.4 | 9.6 | 1.8 | 2.5 | 9.9 | 26.8 | 46.5 | 81.1 | 155.2 | 306.2 | 658.1 |
| 1973 | 40.8 | 287.1 | 13.3 | 1.8 | 3.7 | 12.2 | 28.3 | 58.3 | 94.0 | 186.0 | 357.4 | 726.8 |
| 1972 | 43.7 | 343.6 | 12.1 | 1.6 | 3.8 | 13.6 | 34.7 | 64.1 | 98.7 | 198.7 | 361.4 | 570.0 |
| 1971 | 42.9 | 403.1 | 16.0 | 1.6 | 6.1 | 14.4 | 31.7 | 52.2 | 92.4 | 161.0 | 329.6 | 622.5 |
| 1970 | 52.1 | 503.1 | 21.3 | 2.5 | 5.7 | 15.1 | 38.4 | 72.1 | 109.3 | 204.1 | 389.0 | 571.8 |
| 1969 | 56.3 | 605.7 | 23.1 | 2.6 | 5.4 | 15.9 | 41.8 | 69.5 | 113.2 | 224.4 | 428.1 | 816.2 |
| 1968 | 62.5 | 660.9 | 25.4 | 3.0 | 6.2 | 18.4 | 44.7 | 76.3 | 130.2 | 230.5 | 463.6 | 945.7 |
| BLACK, FEMALE | | | | | | | | | | | | |
| 1978 | 18.0 | 97.1 | 5.1 | 1.1 | 2.2 | 4.7 | 8.6 | 15.5 | 28.0 | 55.3 | 147.5 | 441.4 |
| 1977 | 16.9 | 112.5 | 5.0 | 1.3 | 1.6 | 4.1 | 8.3 | 15.2 | 25.9 | 54.5 | 136.2 | 383.0 |
| 1976 | 20.1 | 133.6 | 7.6 | 1.3 | 2.5 | 5.4 | 11.1 | 18.3 | 28.6 | 61.7 | 162.2 | 478.7 |
| 1975 | 18.5 | 136.7 | 5.8 | 1.5 | 2.1 | 5.3 | 10.3 | 15.8 | 29.1 | 63.1 | 139.9 | 421.4 |
| 1974 | 20.1 | 179.6 | 9.8 | 1.5 | 2.3 | 5.1 | 11.3 | 18.9 | 27.7 | 73.8 | 143.0 | 419.2 |
| 1973 | 25.0 | 245.9 | 9.3 | 1.7 | 3.5 | 6.9 | 15.9 | 25.5 | 36.2 | 88.2 | 165.5 | 478.1 |
| 1972 | 25.6 | 288.6 | 12.8 | 1.9 | 3.6 | 6.7 | 17.5 | 24.5 | 36.9 | 78.4 | 161.4 | 460.0 |
| 1971 | 26.1 | 323.2 | 13.0 | 2.0 | 3.2 | 6.3 | 15.2 | 23.7 | 37.6 | 75.2 | 166.4 | 468.7 |
| 1970 | 31.7 | 405.9 | 13.9 | 2.6 | 4.6 | 8.0 | 18.6 | 29.9 | 48.8 | 102.9 | 191.8 | 413.2 |
| 1969 | 35.3 | 513.2 | 15.6 | 2.7 | 3.7 | 10.9 | 20.0 | 29.6 | 53.2 | 107.3 | 218.1 | 543.3 |
| 1968 | 41.4 | 581.3 | 23.1 | 3.7 | 3.8 | 10.9 | 23.8 | 34.7 | 61.8 | 116.7 | 247.7 | 735.7 |

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| INFLUENZA (470-474) | | | | | | | | | | | | |
| BOTH SEXES | | | | | | | | | | | | |
| 1978 | 1.8 | 1.0 | * | * | 0.1 | 0.1 | 0.2 | 0.5 | 1.2 | 4.0 | 18.0 | 74.7 |
| 1977 | 0.6 | * | * | * | * | * | * | 0.2 | 0.5 | 1.4 | 5.2 | 24.3 |
| 1976 | 3.6 | 1.1 | 0.2 | 0.1 | 0.1 | 0.2 | 0.4 | 0.7 | 2.2 | 7.7 | 36.1 | 174.7 |
| 1975 | 2.0 | 1.0 | * | 0.1 | 0.1 | 0.1 | 0.3 | 0.7 | 1.9 | 5.9 | 21.9 | 62.4 |
| 1974 | 1.0 | 1.2 | 0.2 | 0.1 | 0.1 | 0.1 | 0.2 | 0.4 | 0.9 | 2.6 | 10.5 | 39.4 |
| 1973 | 2.4 | 1.7 | 0.2 | 0.1 | 0.1 | 0.2 | 0.5 | 1.0 | 2.5 | 7.4 | 25.7 | 85.1 |
| 1972 | 2.4 | 1.8 | 0.3 | 0.1 | 0.1 | 0.2 | 0.5 | 1.5 | 3.0 | 8.6 | 23.2 | 69.3 |
| 1971 | 0.7 | 1.2 | 0.1 | * | 0.1 | 0.1 | 0.1 | 0.3 | 0.7 | 2.0 | 6.9 | 31.2 |
| 1970 | 1.8 | 2.4 | 0.3 | 0.1 | 0.1 | 0.3 | 0.5 | 1.2 | 2.9 | 6.8 | 15.4 | 48.0 |
| 1969 | 3.0 | 3.7 | 0.4 | 0.1 | 0.2 | 0.3 | 0.8 | 2.2 | 4.6 | 11.1 | 25.0 | 89.2 |
| 1968 | 3.5 | 4.0 | 0.5 | 0.1 | 0.3 | 0.4 | 0.8 | 2.1 | 4.6 | 11.3 | 33.3 | 140.7 |
| MALE | | | | | | | | | | | | |
| 1978 | 1.5 | * | * | * | * | * | 0.2 | 0.6 | 1.3 | 5.1 | 20.9 | 71.8 |
| 1977 | 0.5 | * | * | * | * | * | * | 0.2 | 0.5 | 1.6 | 5.1 | 23.0 |
| 1976 | 3.1 | 1.3 | * | * | * | 0.2 | 0.4 | 0.8 | 2.3 | 9.9 | 41.6 | 188.9 |
| 1975 | 1.8 | 1.4 | * | * | 0.1 | * | 0.3 | 0.7 | 2.0 | 7.4 | 26.0 | 66.8 |
| 1974 | 1.0 | * | * | * | * | * | 0.2 | 0.4 | 1.0 | 3.4 | 12.1 | 41.1 |
| 1973 | 2.3 | 1.8 | * | 0.1 | 0.2 | 0.2 | 0.5 | 1.0 | 2.9 | 9.7 | 30.7 | 92.6 |
| 1972 | 2.3 | 1.9 | 0.4 | * | 0.1 | * | 0.5 | 1.6 | 3.6 | 11.9 | 26.8 | 67.8 |
| 1971 | 0.7 | 1.3 | * | * | * | * | * | 0.3 | 0.8 | 2.5 | 7.8 | 30.2 |
| 1970 | 1.9 | 2.6 | 0.4 | * | 0.1 | 0.2 | 0.5 | 1.3 | 3.6 | 9.1 | 19.3 | 49.0 |
| 1969 | 3.1 | 4.2 | 0.4 | * | 0.3 | 0.3 | 0.7 | 2.3 | 6.1 | 14.9 | 30.2 | 88.0 |
| 1968 | 3.4 | 4.1 | 0.5 | 0.1 | 0.3 | 0.3 | 0.8 | 2.2 | 5.6 | 13.9 | 36.8 | 140.9 |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| INFLUENZA (470-474) | | | | | | | | | | | | |
| FEMALE | | | | | | | | | | | | |
| 1978 | 2.1 | * | * | * | * | * | 0.2 | 0.4 | 1.1 | 3.2 | 16.3 | 76.0 |
| 1977 | 0.7 | * | * | * | * | * | * | * | 0.5 | 1.2 | 5.3 | 25.0 |
| 1976 | 4.1 | * | * | * | 0.1 | 0.2 | 0.4 | 0.6 | 2.0 | 6.1 | 32.7 | 168.0 |
| 1975 | 2.1 | * | * | * | 0.1 | 0.1 | 0.3 | 0.7 | 1.8 | 4.8 | 19.4 | 60.2 |
| 1974 | 1.1 | * | * | 0.1 | * | * | * | 0.5 | 0.7 | 1.9 | 9.4 | 38.6 |
| 1973 | 2.5 | 1.5 | * | * | 0.1 | 0.2 | 0.6 | 1.0 | 2.1 | 5.5 | 22.6 | 81.4 |
| 1972 | 2.4 | 1.7 | * | * | * | 0.3 | 0.6 | 1.4 | 2.4 | 6.1 | 20.9 | 70.0 |
| 1971 | 0.8 | * | * | * | * | * | * | 0.2 | 0.5 | 1.6 | 6.4 | 31.7 |
| 1970 | 1.7 | 2.1 | * | * | 0.1 | 0.3 | 0.5 | 1.1 | 2.2 | 5.0 | 12.8 | 47.4 |
| 1969 | 2.9 | 3.1 | * | 0.2 | 0.2 | 0.4 | 0.8 | 2.1 | 3.2 | 8.2 | 21.5 | 89.9 |
| 1968 | 3.7 | 3.8 | 0.6 | 0.1 | 0.3 | 0.5 | 0.8 | 2.1 | 3.7 | 9.2 | 30.8 | 140.8 |
| WHITE, BOTH SEXES | | | | | | | | | | | | |
| 1978 | 2.0 | * | * | * | 0.1 | 0.1 | 0.1 | 0.5 | 1.2 | 4.1 | 19.0 | 78.4 |
| 1977 | 0.6 | * | * | * | * | * | * | 0.1 | 0.4 | 1.4 | 5.3 | 25.1 |
| 1976 | 4.0 | 1.0 | * | 0.1 | 0.1 | 0.2 | 0.4 | 0.7 | 2.2 | 7.9 | 38.1 | 185.4 |
| 1975 | 2.1 | * | * | 0.1 | 0.1 | 0.1 | 0.3 | 0.7 | 1.9 | 6.0 | 23.0 | 66.0 |
| 1974 | 1.1 | 1.1 | 0.2 | 0.1 | 0.1 | 0.1 | 0.1 | 0.4 | 0.9 | 2.5 | 10.9 | 41.6 |
| 1973 | 2.6 | 1.2 | 0.2 | 0.1 | 0.1 | 0.2 | 0.5 | 0.9 | 2.4 | 7.2 | 26.7 | 88.3 |
| 1972 | 2.5 | 1.2 | 0.3 | 0.1 | 0.1 | 0.2 | 0.5 | 1.5 | 2.9 | 8.8 | 24.0 | 71.3 |
| 1971 | 0.8 | 0.8 | * | * | * | 0.1 | 0.1 | 0.3 | 0.6 | 2.0 | 7.2 | 32.9 |
| 1970 | 1.8 | 1.6 | 0.2 | 0.1 | 0.1 | 0.3 | 0.4 | 1.1 | 2.7 | 6.4 | 15.3 | 48.7 |
| 1969 | 3.0 | 2.3 | 0.3 | 0.1 | 0.2 | 0.3 | 0.7 | 2.0 | 4.2 | 10.6 | 25.1 | 93.0 |
| 1968 | 3.6 | 2.1 | 0.5 | 0.1 | 0.3 | 0.4 | 0.6 | 1.9 | 4.3 | 11.0 | 33.5 | 144.6 |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| INFLUENZA (470-474) | | | | | | | | | | | | |
| WHITE, MALE | | | | | | | | | | | | |
| 1978 | 1.7 | * | * | * | * | * | * | 0.6 | 1.4 | 5.2 | 21.9 | 76.4 |
| 1977 | 0.5 | * | * | * | * | * | * | * | 0.5 | 1.7 | 5.0 | 23.8 |
| 1976 | 3.4 | * | * | * | * | 0.2 | 0.4 | 0.7 | 2.3 | 10.1 | 44.0 | 204.2 |
| 1975 | 2.0 | * | * | * | * | * | 0.2 | 0.7 | 2.0 | 7.5 | 27.3 | 70.7 |
| 1974 | 1.0 | * | * | * | * | * | * | 0.4 | 1.0 | 3.3 | 12.8 | 42.9 |
| 1973 | 2.4 | 1.5 | * | 0.1 | 0.2 | 0.2 | 0.4 | 0.9 | 2.9 | 9.6 | 32.1 | 96.7 |
| 1972 | 2.4 | * | 0.3 | * | * | * | 0.5 | 1.6 | 3.6 | 12.2 | 27.5 | 70.3 |
| 1971 | 0.7 | * | * | * | * | * | * | 0.3 | 0.8 | 2.6 | 8.2 | 32.2 |
| 1970 | 1.9 | 1.9 | * | * | 0.2 | 0.2 | 0.4 | 1.1 | 3.4 | 8.7 | 19.0 | 50.3 |
| 1969 | 3.1 | 2.7 | 0.4 | * | 0.3 | 0.2 | 0.6 | 2.1 | 5.7 | 14.4 | 30.3 | 93.0 |
| 1968 | 3.4 | 2.1 | 0.4 | 0.1 | 0.3 | 0.3 | 0.7 | 1.9 | 5.2 | 13.8 | 37.0 | 146.1 |
| WHITE, FEMALE | | | | | | | | | | | | |
| 1978 | 2.3 | * | * | * | * | * | * | 0.4 | 1.0 | 3.2 | 17.2 | 79.3 |
| 1977 | 0.7 | * | * | * | * | * | * | * | 0.4 | 1.1 | 5.5 | 25.7 |
| 1976 | 4.5 | * | * | * | 0.1 | 0.2 | 0.4 | 0.7 | 2.0 | 6.3 | 34.6 | 176.9 |
| 1975 | 2.3 | * | * | * | 0.1 | * | 0.3 | 0.7 | 1.7 | 4.8 | 20.5 | 63.8 |
| 1974 | 1.2 | * | * | 0.1 | * | * | * | 0.5 | 0.7 | 1.9 | 9.8 | 40.9 |
| 1973 | 2.7 | * | * | * | * | 0.2 | 0.5 | 0.9 | 1.9 | 5.4 | 23.4 | 84.2 |
| 1972 | 2.6 | * | * | * | * | 0.4 | 0.5 | 1.4 | 2.2 | 6.1 | 21.7 | 71.8 |
| 1971 | 0.8 | * | * | * | * | * | * | 0.2 | 0.5 | 1.6 | 6.6 | 33.3 |
| 1970 | 1.7 | * | * | * | * | 0.3 | 0.4 | 1.0 | 2.1 | 4.5 | 12.9 | 47.9 |
| 1969 | 2.9 | 1.9 | * | 0.2 | 0.2 | 0.4 | 0.7 | 1.9 | 2.9 | 7.7 | 21.7 | 93.1 |
| 1968 | 3.8 | 2.2 | 0.6 | * | 0.3 | 0.5 | 0.6 | 2.0 | 3.4 | 8.8 | 31.2 | 144.0 |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| INFLUENZA (470-474) | | | | | | | | | | | | |
| ALL OTHER, BOTH SEXES | | | | | | | | | | | | |
| 1978 | 0.8 | * | * | * | * | * | * | * | 1.6 | 3.4 | 7.9 | 31.5 |
| 1977 | 0.4 | * | * | * | * | * | * | * | 1.2 | * | 4.2 | 15.6 |
| 1976 | 1.3 | * | * | * | * | * | * | 1.0 | 2.1 | 5.8 | 15.2 | 51.0 |
| 1975 | 1.0 | * | * | * | * | * | * | 1.0 | 2.3 | 5.3 | 10.4 | 21.2 |
| 1974 | 0.6 | * | * | * | * | * | * | * | 1.1 | 3.0 | 5.2 | * |
| 1973 | 1.5 | 4.2 | * | * | * | * | 1.0 | 1.6 | 3.1 | 8.5 | 14.0 | 48.0 |
| 1972 | 1.6 | 4.8 | * | * | * | * | 0.9 | 1.9 | 4.3 | 7.1 | 14.3 | 45.5 |
| 1971 | 0.4 | * | * | * | * | * | * | * | * | * | 4.1 | * |
| 1970 | 1.9 | 6.4 | * | * | * | * | 1.0 | 2.6 | 4.3 | 10.9 | 16.3 | 40.3 |
| 1969 | 2.9 | 11.5 | * | * | * | * | 1.4 | 3.9 | 8.2 | 16.1 | 23.5 | 46.8 |
| 1968 | 3.3 | 13.9 | * | * | * | 0.8 | 2.2 | 3.7 | 7.9 | 13.8 | 30.0 | 97.1 |
| ALL OTHER, MALE | | | | | | | | | | | | |
| 1978 | 0.8 | * | * | * | * | * | * | * | * | 4.3 | 11.7 | * |
| 1977 | 0.4 | * | * | * | * | * | * | * | * | * | * | * |
| 1976 | 1.3 | * | * | * | * | * | * | * | 2.8 | 7.4 | 19.5 | 36.4 |
| 1975 | 1.0 | * | * | * | * | * | * | * | 2.5 | 5.7 | 13.6 | * |
| 1974 | 0.6 | * | * | * | * | * | * | * | * | 4.5 | * | * |
| 1973 | 1.6 | * | * | * | * | * | * | * | 3.0 | 11.0 | 16.1 | 50.0 |
| 1972 | 1.7 | * | * | * | * | * | * | 2.3 | 4.0 | 8.7 | 19.4 | 42.6 |
| 1971 | 0.4 | * | * | * | * | * | * | * | * | * | * | * |
| 1970 | 2.1 | * | * | * | * | * | * | 2.9 | 5.0 | 12.8 | 23.3 | 37.9 |
| 1969 | 3.2 | 13.2 | * | * | * | * | * | 4.1 | 10.2 | 19.9 | 28.7 | * |
| 1968 | 3.4 | 15.1 | * | * | * | * | 1.8 | 4.5 | 9.3 | 15.0 | 34.4 | 90.2 |
| ALL OTHER, FEMALE | | | | | | | | | | | | |
| 1978 | 0.8 | * | * | * | * | * | * | * | 1.9 | 2.8 | 5.4 | 35.8 |
| 1977 | 0.5 | * | * | * | * | * | * | * | * | * | * | * |
| 1976 | 1.2 | * | * | * | * | * | * | * | * | 4.5 | 12.2 | 59.2 |
| 1975 | 0.9 | * | * | * | * | * | * | * | 2.2 | 5.0 | 8.2 | * |
| 1974 | 0.5 | * | * | * | * | * | * | * | * | * | * | * |
| 1973 | 1.5 | * | * | * | * | * | * | 1.5 | 3.2 | 6.5 | 12.6 | 46.8 |
| 1972 | 1.5 | * | * | * | * | * | * | 1.7 | 4.5 | 5.8 | 10.6 | 47.4 |
| 1971 | 0.4 | * | * | * | * | * | * | * | * | * | * | * |
| 1970 | 1.8 | * | * | * | * | * | * | 2.4 | 3.7 | 9.3 | 11.0 | 41.9 |
| 1969 | 2.6 | 9.8 | * | * | * | * | 1.5 | 3.7 | 6.3 | 13.0 | 19.4 | 50.8 |
| 1968 | 3.2 | 12.6 | * | * | * | * | 2.5 | 2.9 | 6.7 | 12.9 | 26.8 | 101.6 |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| INFLUENZA (470-474) | | | | | | | | | | | | |
| BLACK, BOTH SEXES | | | | | | | | | | | | |
| 1978 | 0.9 | * | * | * | * | * | * | * | 1.7 | 3.6 | 8.4 | 28.0 |
| 1977 | 0.4 | * | * | * | * | * | * | * | 1.3 | * | 3.9 | 15.4 |
| 1976 | 1.2 | * | * | * | * | * | * | 1.1 | 2.1 | 5.8 | 13.9 | 46.3 |
| 1975 | 1.0 | * | * | * | * | * | * | 1.0 | 2.5 | 5.8 | 10.4 | 18.5 |
| 1974 | 0.6 | * | * | * | * | * | * | * | * | 2.9 | 5.6 | * |
| 1973 | 1.5 | * | * | * | * | * | 1.1 | 1.5 | 2.8 | 8.7 | 12.8 | 40.4 |
| 1972 | 1.6 | 5.0 | * | * | * | * | 0.8 | 2.0 | 4.7 | 7.3 | 13.9 | 40.0 |
| 1971 | 0.4 | * | * | * | * | * | * | * | * | * | * | * |
| 1970 | 2.1 | 7.2 | * | * | * | * | 1.1 | 2.9 | 4.7 | 11.5 | 16.1 | 42.6 |
| 1969 | 3.0 | 12.7 | * | * | * | * | 1.4 | 4.1 | 8.3 | 17.3 | 22.9 | 52.6 |
| 1968 | 3.4 | 14.0 | * | * | * | * | 2.3 | 3.9 | 8.2 | 13.9 | 31.5 | 100.0 |
| BLACK, MALE | | | | | | | | | | | | |
| 1978 | 0.9 | * | * | * | * | * | * | * | * | 4.4 | 12.1 | * |
| 1977 | 0.4 | * | * | * | * | * | * | * | * | * | * | * |
| 1976 | 1.2 | * | * | * | * | * | * | * | 2.9 | 7.3 | 16.7 | * |
| 1975 | 1.1 | * | * | * | * | * | * | * | 2.8 | 6.6 | 13.5 | * |
| 1974 | 0.6 | * | * | * | * | * | * | * | * | 4.1 | * | * |
| 1973 | 1.5 | * | * | * | * | * | * | * | 2.6 | 11.0 | 14.7 | * |
| 1972 | 1.7 | * | * | * | * | * | * | 2.4 | 4.5 | 9.0 | 19.3 | * |
| 1971 | 0.4 | * | * | * | * | * | * | * | * | * | * | * |
| 1970 | 2.3 | * | * | * | * | * | * | 3.3 | 5.4 | 14.1 | 24.3 | * |
| 1969 | 3.3 | 14.4 | * | * | * | * | * | 4.2 | 10.0 | 22.0 | 28.1 | * |
| 1968 | 3.4 | 15.5 | * | * | * | * | * | 4.8 | 9.8 | 14.6 | 35.8 | 91.4 |
| BLACK, FEMALE | | | | | | | | | | | | |
| 1978 | 0.9 | * | * | * | * | * | * | * | 2.2 | 3.0 | 5.9 | 30.3 |
| 1977 | 0.5 | * | * | * | * | * | * | * | * | * | * | * |
| 1976 | 1.3 | * | * | * | * | * | * | * | * | 4.7 | 12.0 | 56.2 |
| 1975 | 0.9 | * | * | * | * | * | * | * | 2.3 | 5.1 | 8.2 | * |
| 1974 | 0.5 | * | * | * | * | * | * | * | * | * | * | * |
| 1973 | 1.5 | * | * | * | * | * | * | * | 2.9 | 6.9 | 11.5 | 42.5 |
| 1972 | 1.5 | * | * | * | * | * | * | 1.7 | 4.9 | 6.0 | 10.0 | 42.9 |
| 1971 | 0.4 | * | * | * | * | * | * | * | * | * | * | * |
| 1970 | 1.9 | * | * | * | * | * | * | 2.6 | 4.0 | 9.5 | 10.0 | 44.8 |
| 1969 | 2.8 | 11.0 | * | * | * | * | 1.5 | 3.9 | 6.8 | 13.6 | 18.9 | 55.0 |
| 1968 | 3.4 | 12.3 | * | * | * | * | 2.8 | 3.1 | 6.9 | 13.3 | 28.4 | 107.1 |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| PNEUMONIA (480-486) | | | | | | | | | | | | |
| BOTH SEXES | | | | | | | | | | | | |
| 1978 | 24.4 | 45.1 | 2.7 | 0.8 | 1.2 | 2.0 | 4.2 | 9.5 | 22.3 | 61.5 | 226.7 | 809.7 |
| 1977 | 22.7 | 50.8 | 2.9 | 0.9 | 1.2 | 1.9 | 4.3 | 9.4 | 20.9 | 56.9 | 217.0 | 738.8 |
| 1976 | 24.8 | 61.9 | 3.7 | 0.9 | 1.4 | 2.2 | 5.0 | 10.9 | 23.7 | 62.1 | 238.3 | 819.9 |
| 1975 | 23.8 | 68.9 | 3.9 | 0.9 | 1.6 | 2.3 | 5.0 | 10.4 | 24.7 | 64.1 | 230.8 | 738.7 |
| 1974 | 24.6 | 84.1 | 4.4 | 1.2 | 1.6 | 2.4 | 5.6 | 11.3 | 25.5 | 66.0 | 239.4 | 785.5 |
| 1973 | 27.2 | 112.2 | 5.6 | 1.3 | 1.8 | 2.9 | 6.4 | 13.5 | 28.9 | 74.5 | 261.6 | 838.3 |
| 1972 | 27.5 | 129.2 | 6.2 | 1.4 | 2.0 | 3.1 | 7.3 | 14.5 | 29.8 | 77.9 | 260.2 | 801.2 |
| 1971 | 26.9 | 148.8 | 7.0 | 1.3 | 2.2 | 3.2 | 6.8 | 13.1 | 28.4 | 72.8 | 247.7 | 815.5 |
| 1970 | 29.0 | 178.5 | 7.3 | 1.5 | 2.3 | 3.6 | 8.0 | 16.1 | 33.3 | 83.4 | 257.4 | 766.6 |
| 1969 | 31.0 | 205.5 | 8.1 | 1.7 | 2.3 | 3.7 | 8.2 | 15.9 | 34.4 | 87.9 | 277.9 | 937.0 |
| 1968 | 33.3 | 231.0 | 9.4 | 1.7 | 2.6 | 4.0 | 8.5 | 17.8 | 37.3 | 94.7 | 297.4 | 1,027.2 |
| MALE | | | | | | | | | | | | |
| 1978 | 27.0 | 48.9 | 2.7 | 0.8 | 1.5 | 2.5 | 5.3 | 12.5 | 31.4 | 90.9 | 328.5 | 1,087.6 |
| 1977 | 25.4 | 57.1 | 3.3 | 0.8 | 1.5 | 2.4 | 5.4 | 12.7 | 29.9 | 84.3 | 314.6 | 987.4 |
| 1976 | 27.6 | 66.9 | 3.7 | 0.8 | 1.5 | 2.6 | 6.3 | 14.1 | 33.9 | 91.2 | 343.8 | 1,081.0 |
| 1975 | 27.2 | 74.9 | 3.9 | 0.8 | 1.9 | 2.6 | 6.2 | 14.5 | 35.2 | 94.6 | 340.0 | 969.2 |
| 1974 | 27.7 | 92.0 | 4.6 | 1.2 | 1.8 | 2.8 | 6.9 | 14.5 | 36.3 | 96.4 | 343.2 | 1,001.6 |
| 1973 | 30.9 | 124.5 | 5.9 | 1.3 | 2.2 | 3.5 | 7.7 | 17.9 | 40.8 | 109.8 | 375.1 | 1,046.1 |
| 1972 | 31.7 | 148.0 | 6.4 | 1.3 | 2.2 | 3.5 | 8.8 | 19.6 | 42.7 | 114.0 | 370.3 | 1,007.3 |
| 1971 | 30.7 | 168.3 | 7.3 | 1.2 | 2.5 | 3.7 | 8.2 | 17.2 | 40.2 | 105.0 | 344.2 | 1,015.9 |
| 1970 | 33.6 | 200.7 | 8.2 | 1.5 | 2.7 | 4.2 | 10.0 | 21.3 | 46.6 | 120.7 | 356.9 | 910.8 |
| 1969 | 35.6 | 231.1 | 8.8 | 1.7 | 2.6 | 4.1 | 10.0 | 21.4 | 47.9 | 127.4 | 375.9 | 1,126.4 |
| 1968 | 38.4 | 256.9 | 9.8 | 1.7 | 3.0 | 4.6 | 10.7 | 23.7 | 52.7 | 138.3 | 394.7 | 1,234.5 |

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| PNEUMONIA (480-486) | | | | | | | | | | | | |
| FEMALE | | | | | | | | | | | | |
| 1978 | 22.0 | 41.1 | 2.7 | 0.7 | 1.0 | 1.6 | 3.2 | 6.6 | 14.2 | 39.0 | 166.0 | 684.1 |
| 1977 | 20.2 | 44.2 | 2.6 | 0.9 | 1.0 | 1.5 | 3.2 | 6.3 | 13.0 | 36.0 | 158.6 | 624.5 |
| 1976 | 22.2 | 56.6 | 3.6 | 0.9 | 1.2 | 1.8 | 3.8 | 7.9 | 14.7 | 39.8 | 174.5 | 697.3 |
| 1975 | 20.7 | 62.6 | 3.9 | 0.9 | 1.2 | 2.0 | 3.9 | 6.6 | 15.5 | 40.7 | 164.3 | 627.1 |
| 1974 | 21.7 | 75.8 | 4.2 | 1.1 | 1.4 | 2.0 | 4.3 | 8.4 | 15.9 | 42.7 | 175.5 | 678.5 |
| 1973 | 23.6 | 99.3 | 5.3 | 1.3 | 1.5 | 2.3 | 5.1 | 9.5 | 18.3 | 47.5 | 190.6 | 733.7 |
| 1972 | 23.5 | 109.6 | 6.1 | 1.4 | 1.8 | 2.6 | 5.8 | 9.7 | 18.4 | 50.2 | 190.0 | 695.7 |
| 1971 | 23.4 | 128.4 | 6.6 | 1.4 | 1.8 | 2.7 | 5.5 | 9.3 | 17.8 | 47.8 | 185.2 | 711.0 |
| 1970 | 24.7 | 155.3 | 6.4 | 1.5 | 1.9 | 2.9 | 6.1 | 11.2 | 21.3 | 54.3 | 191.6 | 685.8 |
| 1969 | 26.6 | 179.0 | 7.4 | 1.7 | 1.9 | 3.4 | 6.5 | 10.8 | 22.2 | 57.0 | 212.1 | 834.6 |
| 1968 | 28.5 | 203.9 | 8.9 | 1.7 | 2.1 | 3.4 | 6.4 | 12.3 | 23.4 | 60.4 | 230.8 | 914.4 |
| WHITE, BOTH SEXES | | | | | | | | | | | | |
| 1978 | 24.8 | 34.4 | 2.2 | 0.7 | 1.1 | 1.5 | 3.1 | 7.5 | 19.9 | 58.8 | 228.2 | 832.8 |
| 1977 | 22.9 | 37.9 | 2.4 | 0.8 | 1.1 | 1.4 | 3.1 | 7.5 | 18.8 | 54.2 | 218.9 | 758.2 |
| 1976 | 25.0 | 46.5 | 2.9 | 0.8 | 1.2 | 1.6 | 3.7 | 8.9 | 21.1 | 59.3 | 239.9 | 843.6 |
| 1975 | 23.9 | 53.1 | 3.3 | 0.8 | 1.5 | 1.8 | 3.7 | 8.5 | 22.4 | 61.3 | 233.1 | 755.7 |
| 1974 | 24.6 | 63.3 | 3.5 | 1.1 | 1.5 | 1.8 | 4.0 | 9.3 | 23.1 | 62.4 | 242.5 | 807.8 |
| 1973 | 26.8 | 85.2 | 4.7 | 1.2 | 1.6 | 2.1 | 4.7 | 10.9 | 25.9 | 70.0 | 263.5 | 863.4 |
| 1972 | 27.0 | 96.9 | 5.2 | 1.3 | 1.8 | 2.2 | 5.3 | 11.8 | 26.9 | 73.8 | 262.5 | 827.8 |
| 1971 | 26.1 | 113.3 | 5.7 | 1.2 | 1.8 | 2.3 | 4.9 | 10.7 | 25.2 | 69.0 | 248.5 | 839.6 |
| 1970 | 27.9 | 133.7 | 5.8 | 1.4 | 2.0 | 2.7 | 5.8 | 13.0 | 29.6 | 78.5 | 257.6 | 798.3 |
| 1969 | 29.7 | 150.8 | 6.3 | 1.6 | 2.0 | 2.6 | 5.9 | 13.1 | 30.8 | 82.9 | 277.6 | 966.6 |
| 1968 | 31.5 | 167.0 | 6.9 | 1.5 | 2.3 | 2.8 | 5.9 | 14.5 | 32.8 | 89.4 | 296.6 | 1,053.1 |

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| PNEUMONIA (480-486) | | | | | | | | | | | | |
| WHITE, MALE | | | | | | | | | | | | |
| 1978 | 26.7 | 37.2 | 2.2 | 0.7 | 1.3 | 1.9 | 3.7 | 9.6 | 27.5 | 86.3 | 330.3 | 1,118.2 |
| 1977 | 25.0 | 43.5 | 2.7 | 0.8 | 1.3 | 1.7 | 3.7 | 9.8 | 26.5 | 80.1 | 317.7 | 1,006.5 |
| 1976 | 27.1 | 49.8 | 2.9 | 0.8 | 1.4 | 1.9 | 4.5 | 11.1 | 29.7 | 86.8 | 346.1 | 1,113.6 |
| 1975 | 26.7 | 57.7 | 3.1 | 0.7 | 1.8 | 1.9 | 4.3 | 11.4 | 31.4 | 90.4 | 342.9 | 991.9 |
| 1974 | 27.1 | 68.4 | 3.8 | 1.2 | 1.8 | 2.0 | 4.7 | 11.4 | 32.3 | 91.5 | 347.1 | 1,032.2 |
| 1973 | 29.9 | 96.5 | 4.7 | 1.2 | 2.0 | 2.5 | 5.5 | 14.1 | 36.1 | 103.6 | 376.7 | 1,078.2 |
| 1972 | 30.5 | 114.8 | 5.5 | 1.3 | 2.1 | 2.4 | 6.2 | 15.6 | 38.1 | 107.1 | 372.5 | 1,048.0 |
| 1971 | 29.3 | 129.8 | 6.0 | 1.2 | 2.1 | 2.4 | 5.6 | 13.8 | 35.5 | 99.8 | 345.1 | 1,053.1 |
| 1970 | 31.8 | 152.9 | 6.4 | 1.4 | 2.3 | 3.1 | 7.1 | 16.7 | 41.3 | 114.4 | 357.5 | 952.7 |
| 1969 | 33.5 | 175.2 | 6.5 | 1.6 | 2.3 | 2.8 | 6.7 | 17.1 | 42.9 | 120.6 | 374.1 | 1,165.5 |
| 1968 | 35.9 | 191.2 | 7.3 | 1.5 | 2.6 | 3.1 | 7.2 | 19.1 | 46.6 | 131.2 | 392.2 | 1,269.7 |
| WHITE, FEMALE | | | | | | | | | | | | |
| 1978 | 23.0 | 31.4 | 2.2 | 0.7 | 0.8 | 1.1 | 2.5 | 5.6 | 13.1 | 37.8 | 168.1 | 705.7 |
| 1977 | 20.9 | 32.1 | 2.1 | 0.9 | 0.9 | 1.1 | 2.6 | 5.3 | 11.9 | 34.5 | 160.6 | 645.8 |
| 1976 | 22.9 | 43.0 | 2.8 | 0.9 | 1.0 | 1.4 | 2.9 | 6.8 | 13.5 | 38.3 | 176.6 | 719.5 |
| 1975 | 21.3 | 48.3 | 3.5 | 0.8 | 1.1 | 1.6 | 3.1 | 5.7 | 14.4 | 39.1 | 167.1 | 643.9 |
| 1974 | 22.2 | 58.0 | 3.2 | 1.0 | 1.2 | 1.7 | 3.4 | 7.3 | 14.8 | 40.2 | 178.6 | 698.4 |
| 1973 | 23.8 | 73.5 | 4.6 | 1.2 | 1.2 | 1.8 | 3.8 | 7.8 | 16.7 | 44.3 | 193.4 | 757.8 |
| 1972 | 23.6 | 78.1 | 4.8 | 1.3 | 1.5 | 2.0 | 4.4 | 8.3 | 17.0 | 48.2 | 193.2 | 716.9 |
| 1971 | 23.2 | 95.9 | 5.5 | 1.3 | 1.6 | 2.3 | 4.3 | 7.8 | 15.9 | 45.3 | 186.7 | 730.3 |
| 1970 | 24.2 | 113.5 | 5.1 | 1.4 | 1.6 | 2.3 | 4.6 | 9.4 | 19.1 | 50.7 | 192.2 | 713.8 |
| 1969 | 26.0 | 125.4 | 6.0 | 1.6 | 1.7 | 2.5 | 5.1 | 9.3 | 19.8 | 53.7 | 213.4 | 862.3 |
| 1968 | 27.4 | 141.7 | 6.5 | 1.4 | 1.9 | 2.5 | 4.6 | 10.2 | 20.4 | 56.6 | 231.8 | 937.6 |

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| PNEUMONIA (480-486) | | | | | | | | | | | | |
| ALL OTHER, BOTH SEXES | | | | | | | | | | | | |
| 1978 | 22.0 | 96.7 | 5.0 | 1.1 | 2.2 | 5.5 | 11.7 | 24.4 | 43.6 | 87.1 | 211.0 | 544.0 |
| 1977 | 21.3 | 113.0 | 5.2 | 1.2 | 1.9 | 5.3 | 12.3 | 24.6 | 40.7 | 82.6 | 197.2 | 516.3 |
| 1976 | 23.8 | 137.3 | 7.3 | 1.2 | 2.4 | 6.0 | 14.2 | 27.4 | 47.4 | 89.1 | 221.6 | 545.1 |
| 1975 | 23.3 | 147.6 | 6.9 | 1.3 | 2.3 | 6.2 | 14.2 | 26.3 | 46.6 | 90.5 | 205.9 | 543.2 |
| 1974 | 24.9 | 187.1 | 8.9 | 1.5 | 2.2 | 6.4 | 16.3 | 28.3 | 48.2 | 101.5 | 205.9 | 525.9 |
| 1973 | 29.7 | 245.4 | 10.6 | 1.5 | 3.3 | 8.0 | 18.6 | 36.3 | 57.5 | 119.2 | 239.2 | 545.7 |
| 1972 | 31.1 | 294.6 | 11.6 | 1.7 | 3.3 | 9.1 | 21.6 | 37.3 | 57.4 | 119.2 | 232.8 | 494.3 |
| 1971 | 32.4 | 338.4 | 13.6 | 1.7 | 4.2 | 9.0 | 20.6 | 34.3 | 59.3 | 109.6 | 237.8 | 540.8 |
| 1970 | 37.1 | 417.5 | 15.6 | 2.4 | 4.5 | 9.8 | 23.7 | 43.2 | 68.4 | 131.3 | 255.8 | 440.8 |
| 1969 | 40.4 | 514.0 | 18.0 | 2.4 | 4.0 | 11.4 | 26.2 | 41.4 | 69.2 | 137.5 | 282.5 | 603.7 |
| 1968 | 46.1 | 576.0 | 22.7 | 3.0 | 4.4 | 12.5 | 28.6 | 47.2 | 80.6 | 148.1 | 308.5 | 737.3 |
| ALL OTHER, MALE | | | | | | | | | | | | |
| 1978 | 28.6 | 105.9 | 5.1 | 1.2 | 2.4 | 7.2 | 17.2 | 37.2 | 67.3 | 134.2 | 310.5 | 774.1 |
| 1977 | 27.8 | 123.9 | 5.8 | 1.1 | 2.3 | 7.3 | 18.3 | 37.4 | 62.2 | 123.4 | 284.7 | 792.9 |
| 1976 | 30.4 | 152.3 | 7.3 | 1.1 | 2.5 | 7.5 | 19.6 | 40.3 | 73.0 | 132.8 | 323.0 | 754.5 |
| 1975 | 30.4 | 161.7 | 7.8 | 1.2 | 2.8 | 8.0 | 20.3 | 40.9 | 71.3 | 134.3 | 311.2 | 742.6 |
| 1974 | 31.9 | 211.2 | 8.8 | 1.6 | 2.3 | 8.9 | 23.4 | 41.6 | 74.4 | 143.3 | 303.5 | 686.0 |
| 1973 | 37.8 | 264.7 | 12.2 | 1.6 | 3.3 | 10.6 | 24.6 | 52.1 | 86.5 | 168.8 | 357.1 | 718.8 |
| 1972 | 39.9 | 320.9 | 10.8 | 1.5 | 3.3 | 12.4 | 29.5 | 56.0 | 87.0 | 179.3 | 346.5 | 595.7 |
| 1971 | 40.8 | 376.7 | 14.8 | 1.5 | 5.5 | 13.0 | 28.6 | 48.6 | 86.0 | 154.9 | 335.3 | 648.9 |
| 1970 | 46.7 | 460.5 | 18.4 | 2.3 | 5.1 | 13.2 | 33.0 | 62.8 | 97.5 | 180.4 | 350.2 | 543.1 |
| 1969 | 50.6 | 552.3 | 21.5 | 2.3 | 4.7 | 13.8 | 37.3 | 60.9 | 96.6 | 193.5 | 396.8 | 756.8 |
| 1968 | 56.8 | 616.2 | 23.6 | 2.8 | 5.6 | 16.4 | 39.5 | 66.4 | 111.9 | 206.8 | 425.3 | 892.7 |
| ALL OTHER, FEMALE | | | | | | | | | | | | |
| 1978 | 16.1 | 87.2 | 4.9 | 1.1 | 2.0 | 4.1 | 7.2 | 13.8 | 24.0 | 50.7 | 143.4 | 426.6 |
| 1977 | 15.4 | 101.8 | 4.5 | 1.2 | 1.5 | 3.6 | 7.3 | 13.9 | 22.6 | 50.6 | 137.1 | 367.3 |
| 1976 | 17.8 | 122.6 | 7.3 | 1.3 | 2.3 | 4.7 | 9.6 | 16.6 | 25.9 | 54.5 | 152.4 | 427.6 |
| 1975 | 16.9 | 133.2 | 5.9 | 1.5 | 1.9 | 4.7 | 9.1 | 14.0 | 25.6 | 55.7 | 133.3 | 421.5 |
| 1974 | 18.5 | 163.2 | 9.0 | 1.4 | 2.0 | 4.3 | 10.3 | 17.0 | 25.9 | 68.1 | 138.6 | 431.8 |
| 1973 | 22.3 | 225.8 | 8.9 | 1.5 | 3.2 | 5.8 | 13.6 | 22.7 | 32.6 | 79.1 | 156.9 | 440.5 |
| 1972 | 23.0 | 267.9 | 12.4 | 1.8 | 3.3 | 6.4 | 15.1 | 21.1 | 31.8 | 70.6 | 151.5 | 431.6 |
| 1971 | 24.6 | 299.3 | 12.5 | 1.9 | 3.0 | 5.6 | 13.9 | 21.9 | 36.0 | 72.6 | 165.7 | 471.2 |
| 1970 | 28.3 | 374.0 | 12.8 | 2.5 | 4.0 | 6.9 | 16.1 | 25.9 | 42.7 | 90.9 | 184.1 | 368.6 |
| 1969 | 31.1 | 475.4 | 14.4 | 2.5 | 3.4 | 9.3 | 17.1 | 24.3 | 44.9 | 91.4 | 194.3 | 500.0 |
| 1968 | 36.3 | 533.2 | 21.9 | 3.2 | 3.4 | 9.2 | 19.4 | 30.3 | 52.5 | 99.3 | 218.8 | 632.8 |

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| PNEUMONIA (480-486) | | | | | | | | | | | | |
| BLACK, BOTH SEXES | | | | | | | | | | | | |
| 1978 | 23.4 | 104.5 | 5.3 | 1.2 | 2.3 | 6.2 | 13.5 | 27.3 | 46.8 | 91.2 | 205.7 | 520.0 |
| 1977 | 22.6 | 121.9 | 5.6 | 1.2 | 2.1 | 5.9 | 13.8 | 27.0 | 43.8 | 85.3 | 190.0 | 498.6 |
| 1976 | 25.2 | 147.5 | 7.6 | 1.3 | 2.5 | 6.6 | 15.7 | 30.0 | 50.5 | 93.5 | 218.5 | 523.5 |
| 1975 | 24.4 | 155.0 | 7.0 | 1.3 | 2.5 | 6.8 | 15.5 | 28.1 | 49.7 | 94.0 | 202.2 | 516.2 |
| 1974 | 26.2 | 202.2 | 9.4 | 1.6 | 2.3 | 7.2 | 17.9 | 31.1 | 50.9 | 106.1 | 203.3 | 490.1 |
| 1973 | 31.0 | 263.1 | 11.1 | 1.6 | 3.5 | 9.0 | 20.5 | 39.0 | 59.7 | 122.2 | 230.5 | 527.2 |
| 1972 | 32.6 | 311.9 | 12.0 | 1.7 | 3.6 | 9.8 | 24.5 | 40.7 | 60.4 | 123.7 | 230.7 | 460.0 |
| 1971 | 33.7 | 359.8 | 14.4 | 1.8 | 4.6 | 9.8 | 22.5 | 36.6 | 61.7 | 111.9 | 230.8 | 515.9 |
| 1970 | 39.3 | 447.5 | 16.9 | 2.5 | 4.9 | 10.9 | 26.4 | 46.6 | 72.0 | 136.2 | 259.2 | 432.8 |
| 1969 | 42.3 | 546.8 | 18.7 | 2.5 | 4.3 | 12.5 | 28.4 | 44.1 | 72.5 | 141.9 | 285.0 | 594.8 |
| 1968 | 48.1 | 607.4 | 23.4 | 3.1 | 4.5 | 13.6 | 31.0 | 50.2 | 85.3 | 153.7 | 308.3 | 707.6 |
| BLACK, MALE | | | | | | | | | | | | |
| 1978 | 30.5 | 114.5 | 5.6 | 1.3 | 2.5 | 8.0 | 19.8 | 42.3 | 72.6 | 143.4 | 301.7 | 731.4 |
| 1977 | 29.3 | 132.2 | 6.4 | 1.2 | 2.5 | 8.2 | 20.6 | 41.5 | 67.6 | 128.5 | 276.6 | 749.0 |
| 1976 | 32.3 | 162.9 | 7.7 | 1.3 | 2.6 | 8.1 | 22.0 | 44.8 | 78.7 | 141.7 | 320.5 | 714.9 |
| 1975 | 32.0 | 173.9 | 8.1 | 1.2 | 3.0 | 8.6 | 22.5 | 44.2 | 77.2 | 141.4 | 306.3 | 717.4 |
| 1974 | 33.6 | 226.1 | 9.4 | 1.7 | 2.5 | 9.8 | 26.1 | 46.1 | 79.4 | 151.1 | 300.5 | 637.2 |
| 1973 | 39.3 | 284.1 | 13.0 | 1.6 | 3.5 | 11.9 | 27.1 | 56.6 | 91.3 | 175.0 | 342.6 | 690.2 |
| 1972 | 42.0 | 336.9 | 11.3 | 1.6 | 3.6 | 13.4 | 33.8 | 61.7 | 94.2 | 189.8 | 342.1 | 535.0 |
| 1971 | 42.4 | 398.4 | 15.8 | 1.6 | 6.1 | 14.2 | 31.6 | 52.0 | 90.9 | 160.0 | 326.3 | 610.0 |
| 1970 | 49.8 | 495.7 | 20.1 | 2.4 | 5.5 | 14.8 | 37.2 | 68.9 | 103.9 | 190.0 | 364.7 | 532.6 |
| 1969 | 53.0 | 591.3 | 22.3 | 2.4 | 5.1 | 15.1 | 40.5 | 65.3 | 103.2 | 202.4 | 400.0 | 767.6 |
| 1968 | 59.1 | 645.4 | 24.1 | 2.8 | 5.9 | 17.9 | 42.9 | 71.5 | 120.4 | 215.9 | 427.9 | 854.3 |
| BLACK, FEMALE | | | | | | | | | | | | |
| 1978 | 17.1 | 94.2 | 5.0 | 1.1 | 2.1 | 4.6 | 8.2 | 15.0 | 25.8 | 52.4 | 141.6 | 411.1 |
| 1977 | 16.4 | 111.3 | 4.8 | 1.2 | 1.6 | 3.9 | 8.1 | 15.0 | 24.4 | 52.8 | 133.1 | 368.1 |
| 1976 | 18.9 | 131.9 | 7.4 | 1.3 | 2.4 | 5.2 | 10.5 | 17.8 | 27.2 | 57.0 | 150.1 | 422.5 |
| 1975 | 17.6 | 135.8 | 5.8 | 1.4 | 2.1 | 5.2 | 9.8 | 14.7 | 26.8 | 58.0 | 131.7 | 406.0 |
| 1974 | 19.6 | 177.8 | 9.5 | 1.4 | 2.2 | 5.0 | 11.1 | 18.5 | 27.2 | 71.7 | 137.5 | 409.0 |
| 1973 | 23.5 | 240.6 | 9.2 | 1.6 | 3.4 | 6.5 | 14.9 | 24.1 | 33.3 | 81.3 | 154.1 | 435.6 |
| 1972 | 24.1 | 285.2 | 12.8 | 1.9 | 3.5 | 6.7 | 16.7 | 22.8 | 32.0 | 72.4 | 151.4 | 417.1 |
| 1971 | 25.7 | 320.4 | 13.0 | 1.9 | 3.2 | 6.1 | 15.1 | 23.4 | 36.9 | 73.9 | 162.2 | 459.7 |
| 1970 | 29.8 | 398.9 | 13.6 | 2.5 | 4.4 | 7.5 | 17.6 | 27.3 | 44.8 | 93.5 | 181.8 | 368.4 |
| 1969 | 32.5 | 502.2 | 15.2 | 2.6 | 3.5 | 10.2 | 18.4 | 25.7 | 46.4 | 93.7 | 199.1 | 488.3 |
| 1968 | 38.1 | 568.9 | 22.7 | 3.4 | 3.3 | 10.0 | 21.0 | 31.6 | 55.0 | 103.4 | 219.4 | 628.6 |

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|

Chronic obstructive pulmonary diseases and allied conditions 490-493, *519.3

BOTH SEXES

| | | | | | | | | | | | | |
|------|------|-----|-----|-----|-----|-----|-----|------|------|-------|-------|-------|
| 1978 | 22.7 | 2.1 | 0.4 | 0.2 | 0.3 | 0.5 | 1.7 | 10.0 | 42.5 | 121.8 | 204.5 | 232.8 |
| 1977 | 21.1 | 2.2 | 0.4 | 0.2 | 0.3 | 0.4 | 1.8 | 9.6 | 40.2 | 114.5 | 192.2 | 208.1 |
| 1976 | 21.1 | 2.4 | 0.5 | 0.2 | 0.2 | 0.5 | 1.9 | 10.4 | 42.5 | 114.9 | 188.6 | 212.7 |
| 1975 | 20.0 | 3.3 | 0.6 | 0.2 | 0.3 | 0.6 | 2.1 | 10.1 | 42.3 | 110.9 | 178.1 | 180.3 |
| 1974 | 19.3 | 2.9 | 0.5 | 0.2 | 0.4 | 0.5 | 2.1 | 10.3 | 41.8 | 109.5 | 172.5 | 171.5 |
| 1973 | 19.4 | 3.2 | 0.5 | 0.2 | 0.4 | 0.6 | 2.4 | 10.6 | 43.6 | 110.0 | 171.1 | 183.6 |
| 1972 | 18.6 | 3.9 | 0.6 | 0.3 | 0.4 | 0.8 | 2.6 | 10.8 | 42.5 | 107.1 | 162.0 | 163.4 |
| 1971 | 17.7 | 3.7 | 0.7 | 0.3 | 0.4 | 0.7 | 2.7 | 11.2 | 41.2 | 100.8 | 152.3 | 157.8 |
| 1970 | 17.4 | 4.3 | 0.9 | 0.3 | 0.5 | 0.8 | 2.9 | 11.1 | 42.0 | 99.9 | 147.0 | 136.5 |
| 1969 | 16.8 | 4.7 | 0.8 | 0.3 | 0.5 | 0.9 | 2.9 | 11.1 | 40.7 | 95.7 | 141.1 | 153.9 |
| 1968 | 16.6 | 5.2 | 0.9 | 0.3 | 0.5 | 1.0 | 2.9 | 10.9 | 40.8 | 94.4 | 139.1 | 154.5 |

MALE

| | | | | | | | | | | | | |
|------|------|-----|-----|-----|-----|-----|-----|------|------|-------|-------|-------|
| 1978 | 33.4 | 2.8 | 0.5 | 0.2 | 0.3 | 0.5 | 1.7 | 12.4 | 60.8 | 207.5 | 411.7 | 500.0 |
| 1977 | 31.8 | 2.8 | 0.5 | 0.2 | 0.4 | 0.4 | 2.0 | 12.6 | 60.2 | 199.3 | 393.8 | 456.5 |
| 1976 | 32.2 | 2.9 | 0.6 | 0.2 | 0.3 | 0.5 | 2.0 | 13.5 | 64.3 | 204.1 | 386.5 | 475.2 |
| 1975 | 30.9 | 4.2 | 0.7 | 0.2 | 0.4 | 0.7 | 2.2 | 13.1 | 64.7 | 200.2 | 371.0 | 393.6 |
| 1974 | 30.1 | 4.0 | 0.6 | 0.2 | 0.4 | 0.5 | 2.2 | 13.8 | 65.1 | 199.2 | 357.6 | 372.2 |
| 1973 | 30.6 | 4.5 | 0.7 | 0.3 | 0.4 | 0.6 | 2.4 | 14.3 | 69.9 | 203.2 | 356.9 | 384.2 |
| 1972 | 29.6 | 4.6 | 0.5 | 0.3 | 0.5 | 0.8 | 2.6 | 14.8 | 68.7 | 198.8 | 337.4 | 344.4 |
| 1971 | 28.3 | 4.5 | 0.8 | 0.3 | 0.4 | 0.6 | 3.0 | 15.2 | 67.7 | 189.8 | 313.9 | 327.6 |
| 1970 | 28.1 | 5.1 | 1.0 | 0.4 | 0.5 | 0.7 | 3.3 | 15.0 | 69.4 | 190.1 | 304.4 | 272.1 |
| 1969 | 27.4 | 5.1 | 0.9 | 0.3 | 0.6 | 0.8 | 3.4 | 15.6 | 68.5 | 184.7 | 288.6 | 295.9 |
| 1968 | 27.0 | 5.7 | 1.0 | 0.3 | 0.5 | 0.9 | 3.2 | 15.5 | 68.5 | 182.2 | 281.3 | 298.4 |

*519.3 Code introduced by NCHS in 1969

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|

Chronic obstructive pulmonary diseases and allied conditions 490-493, *519.3

FEMALE

| | | | | | | | | | | | | |
|------|------|-----|-----|-----|-----|-----|-----|-----|------|------|------|-------|
| 1978 | 12.7 | 1.5 | 0.3 | 0.1 | 0.3 | 0.4 | 1.6 | 7.7 | 26.4 | 56.3 | 81.2 | 112.1 |
| 1977 | 10.9 | 1.6 | * | 0.2 | 0.2 | 0.4 | 1.6 | 6.7 | 22.6 | 49.5 | 71.5 | 94.0 |
| 1976 | 10.6 | 2.0 | 0.4 | 0.2 | 0.2 | 0.5 | 1.8 | 7.5 | 23.2 | 46.5 | 69.1 | 89.3 |
| 1975 | 9.7 | 2.3 | 0.5 | 0.2 | 0.3 | 0.6 | 2.0 | 7.4 | 22.6 | 42.5 | 60.6 | 77.1 |
| 1974 | 9.1 | 1.9 | 0.4 | 0.2 | 0.4 | 0.4 | 1.9 | 7.1 | 21.2 | 40.7 | 58.5 | 72.1 |
| 1973 | 8.8 | 1.9 | 0.4 | 0.2 | 0.4 | 0.6 | 2.4 | 7.1 | 20.3 | 38.3 | 55.0 | 82.7 |
| 1972 | 8.2 | 3.1 | 0.7 | 0.2 | 0.3 | 0.8 | 2.7 | 7.1 | 19.2 | 36.4 | 50.2 | 70.8 |
| 1971 | 7.6 | 2.8 | 0.6 | 0.3 | 0.4 | 0.8 | 2.4 | 7.5 | 17.6 | 32.0 | 47.6 | 69.2 |
| 1970 | 7.2 | 3.5 | 0.7 | 0.3 | 0.6 | 1.0 | 2.5 | 7.5 | 17.4 | 29.8 | 42.9 | 60.6 |
| 1969 | 6.8 | 4.3 | 0.8 | 0.2 | 0.5 | 1.0 | 2.4 | 6.9 | 15.7 | 26.2 | 42.1 | 77.1 |
| 1968 | 6.7 | 4.7 | 0.8 | 0.3 | 0.6 | 1.2 | 2.5 | 6.5 | 15.8 | 25.3 | 41.8 | 75.5 |

WHITE, BOTH SEXES

| | | | | | | | | | | | | |
|------|------|-----|-----|-----|-----|-----|-----|------|------|-------|-------|-------|
| 1978 | 24.6 | 1.9 | 0.3 | 0.2 | 0.3 | 0.3 | 1.4 | 9.6 | 43.2 | 127.5 | 214.4 | 241.0 |
| 1977 | 22.8 | 1.8 | 0.3 | 0.2 | 0.2 | 0.3 | 1.6 | 9.2 | 41.1 | 119.9 | 201.4 | 215.0 |
| 1976 | 22.8 | 2.2 | 0.4 | 0.2 | 0.2 | 0.4 | 1.7 | 10.0 | 43.3 | 120.2 | 197.9 | 220.6 |
| 1975 | 21.6 | 2.7 | 0.5 | 0.1 | 0.3 | 0.4 | 1.7 | 9.7 | 43.5 | 116.4 | 186.4 | 187.9 |
| 1974 | 20.8 | 2.3 | 0.4 | 0.1 | 0.3 | 0.4 | 1.7 | 10.2 | 42.8 | 114.0 | 181.3 | 178.2 |
| 1973 | 20.8 | 2.7 | 0.4 | 0.2 | 0.3 | 0.4 | 2.1 | 10.1 | 44.5 | 115.1 | 178.7 | 190.5 |
| 1972 | 19.9 | 3.0 | 0.5 | 0.2 | 0.3 | 0.7 | 2.2 | 10.3 | 43.3 | 112.1 | 168.0 | 169.3 |
| 1971 | 18.8 | 3.1 | 0.6 | 0.2 | 0.3 | 0.5 | 2.2 | 10.7 | 42.0 | 105.3 | 158.5 | 164.3 |
| 1970 | 18.5 | 3.4 | 0.7 | 0.3 | 0.4 | 0.6 | 2.4 | 10.8 | 42.6 | 104.6 | 152.0 | 142.8 |
| 1969 | 17.8 | 3.6 | 0.6 | 0.2 | 0.3 | 0.7 | 2.5 | 10.5 | 41.5 | 100.0 | 146.2 | 160.2 |
| 1968 | 17.5 | 4.5 | 0.7 | 0.3 | 0.4 | 0.7 | 2.3 | 10.1 | 41.6 | 99.0 | 144.0 | 161.6 |

*519.3 Code introduced by NCHS in 1969

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|

Chronic obstructive pulmonary diseases and allied conditions 490-493, *519.3

WHITE, MALE

| | | | | | | | | | | | | |
|------|------|-----|-----|-----|-----|-----|-----|------|------|-------|-------|-------|
| 1978 | 35.9 | 2.5 | * | 0.2 | 0.3 | 0.4 | 1.3 | 11.6 | 61.1 | 215.9 | 434.8 | 521.9 |
| 1977 | 34.3 | 2.4 | 0.5 | 0.2 | 0.3 | 0.4 | 1.8 | 11.9 | 60.8 | 208.3 | 416.0 | 475.0 |
| 1976 | 34.6 | 2.6 | 0.5 | 0.2 | 0.2 | 0.4 | 1.6 | 12.8 | 65.1 | 213.3 | 407.7 | 497.3 |
| 1975 | 33.2 | 3.3 | 0.6 | 0.2 | 0.3 | 0.5 | 1.8 | 12.5 | 65.9 | 210.4 | 390.7 | 413.5 |
| 1974 | 32.3 | 3.2 | 0.5 | 0.1 | 0.3 | 0.4 | 1.7 | 13.5 | 66.1 | 207.5 | 377.7 | 388.9 |
| 1973 | 32.8 | 3.5 | 0.5 | 0.2 | 0.3 | 0.4 | 2.1 | 13.6 | 71.1 | 212.8 | 374.8 | 401.0 |
| 1972 | 31.5 | 3.5 | 0.5 | 0.3 | 0.4 | 0.7 | 2.2 | 13.9 | 69.7 | 208.2 | 351.8 | 361.7 |
| 1971 | 30.2 | 4.0 | 0.6 | 0.2 | 0.3 | 0.4 | 2.5 | 14.5 | 68.9 | 198.9 | 329.2 | 343.6 |
| 1970 | 29.9 | 3.9 | 0.9 | 0.3 | 0.3 | 0.5 | 2.7 | 14.6 | 70.2 | 199.9 | 316.9 | 287.9 |
| 1969 | 29.1 | 4.0 | 0.7 | 0.2 | 0.4 | 0.6 | 3.0 | 14.7 | 69.8 | 193.5 | 301.0 | 311.6 |
| 1968 | 28.7 | 5.0 | 0.8 | 0.3 | 0.3 | 0.6 | 2.6 | 14.3 | 70.0 | 191.7 | 292.7 | 316.5 |

WHITE, FEMALE

| | | | | | | | | | | | | |
|------|------|-----|-----|-----|-----|-----|-----|-----|------|------|------|-------|
| 1978 | 13.9 | * | * | * | 0.3 | 0.3 | 1.5 | 7.8 | 27.3 | 60.1 | 84.9 | 115.8 |
| 1977 | 11.9 | * | * | 0.2 | 0.2 | 0.3 | 1.3 | 6.7 | 23.5 | 52.5 | 74.7 | 97.3 |
| 1976 | 11.5 | 1.7 | * | 0.2 | 0.2 | 0.4 | 1.7 | 7.4 | 23.9 | 49.1 | 72.9 | 93.0 |
| 1975 | 10.4 | 2.0 | 0.4 | * | 0.2 | 0.4 | 1.7 | 7.1 | 23.5 | 44.7 | 63.4 | 80.6 |
| 1974 | 9.8 | * | 0.4 | 0.1 | 0.3 | 0.3 | 1.6 | 7.1 | 22.1 | 42.7 | 61.5 | 75.4 |
| 1973 | 9.4 | 1.7 | * | 0.2 | 0.3 | 0.4 | 2.0 | 6.9 | 20.8 | 40.4 | 57.2 | 86.5 |
| 1972 | 8.7 | 2.4 | 0.6 | * | 0.2 | 0.6 | 2.3 | 7.0 | 19.6 | 38.3 | 52.2 | 72.5 |
| 1971 | 8.0 | 2.0 | 0.6 | 0.3 | 0.3 | 0.6 | 1.9 | 7.1 | 17.9 | 33.2 | 49.3 | 72.5 |
| 1970 | 7.6 | 2.9 | 0.5 | 0.2 | 0.4 | 0.7 | 2.1 | 7.2 | 17.8 | 31.1 | 44.2 | 63.4 |
| 1969 | 7.0 | 3.2 | 0.6 | 0.2 | 0.3 | 0.7 | 2.0 | 6.6 | 15.9 | 27.3 | 43.3 | 80.2 |
| 1968 | 6.9 | 4.0 | 0.7 | 0.3 | 0.4 | 0.8 | 2.0 | 6.1 | 16.0 | 26.4 | 43.3 | 78.2 |

*519.3 Code introduced by NCHS in 1969

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|

Chronic obstructive pulmonary diseases and allied conditions 490-493, *519.3

ALL OTHER, BOTH SEXES

| | | | | | | | | | | | | |
|------|------|------|-----|-----|-----|-----|-----|------|------|------|-------|-------|
| 1978 | 10.9 | 3.5 | 0.9 | * | 0.7 | 1.3 | 3.4 | 12.7 | 36.3 | 68.5 | 102.4 | 138.7 |
| 1977 | 10.0 | 4.3 | * | * | 0.7 | 0.8 | 3.5 | 12.3 | 32.6 | 63.1 | 96.7 | 128.8 |
| 1976 | 10.1 | 3.8 | 0.9 | * | 0.6 | 1.2 | 3.5 | 13.3 | 34.6 | 63.6 | 92.7 | 120.3 |
| 1975 | 9.6 | 6.3 | 0.9 | 0.5 | 0.5 | 1.9 | 4.5 | 13.2 | 32.0 | 57.5 | 89.2 | 93.8 |
| 1974 | 9.3 | 6.0 | * | 0.4 | 1.0 | 1.1 | 4.8 | 11.3 | 32.2 | 65.2 | 75.8 | 94.1 |
| 1973 | 9.9 | 6.1 | 1.0 | 0.5 | 0.9 | 1.7 | 4.8 | 14.4 | 35.1 | 59.7 | 86.0 | 103.1 |
| 1972 | 10.0 | 8.5 | 1.3 | 0.7 | 0.9 | 1.5 | 5.5 | 14.8 | 35.2 | 57.5 | 91.3 | 95.9 |
| 1971 | 9.7 | 6.9 | 1.5 | 0.5 | 0.9 | 2.2 | 6.2 | 15.8 | 33.7 | 56.9 | 76.4 | 84.2 |
| 1970 | 9.9 | 9.1 | 1.8 | 0.7 | 1.6 | 2.4 | 6.4 | 13.9 | 36.4 | 53.4 | 83.3 | 72.3 |
| 1969 | 9.7 | 10.7 | 1.9 | 0.7 | 1.6 | 2.6 | 5.8 | 16.0 | 33.4 | 53.3 | 75.1 | 83.5 |
| 1968 | 9.8 | 9.1 | 1.7 | 0.7 | 1.5 | 3.2 | 7.5 | 17.7 | 33.2 | 48.6 | 75.4 | 74.5 |

ALL OTHER, MALE

| | | | | | | | | | | | | |
|------|------|------|-----|-----|-----|-----|-----|------|------|-------|-------|-------|
| 1978 | 17.0 | * | * | * | 0.7 | 1.3 | 4.3 | 19.1 | 57.5 | 130.3 | 193.2 | 275.9 |
| 1977 | 15.7 | * | * | * | 0.9 | 1.0 | 3.8 | 18.5 | 54.3 | 115.9 | 184.3 | 267.9 |
| 1976 | 16.0 | * | * | * | 0.7 | 1.2 | 4.5 | 19.5 | 56.3 | 117.7 | 186.8 | 254.5 |
| 1975 | 14.8 | 8.6 | * | * | 0.7 | 2.3 | 5.3 | 17.6 | 52.9 | 103.7 | 176.4 | 194.4 |
| 1974 | 14.9 | 7.7 | * | * | 1.0 | * | 6.0 | 16.6 | 54.7 | 120.8 | 150.7 | 200.0 |
| 1973 | 15.5 | 9.4 | * | * | 1.1 | 1.7 | 4.5 | 21.0 | 58.7 | 111.7 | 167.9 | 212.5 |
| 1972 | 15.7 | 10.3 | * | * | 1.0 | 1.3 | 5.9 | 22.6 | 58.9 | 108.5 | 182.5 | 170.2 |
| 1971 | 14.8 | 7.0 | 1.9 | * | 0.8 | 2.5 | 7.1 | 21.8 | 55.4 | 102.3 | 145.4 | 170.2 |
| 1970 | 15.2 | 11.6 | 2.1 | 0.7 | 1.6 | 1.9 | 7.8 | 18.6 | 61.6 | 97.8 | 160.1 | 133.5 |
| 1969 | 14.9 | 10.9 | 2.0 | 0.8 | 1.7 | 1.9 | 6.8 | 23.6 | 56.0 | 99.4 | 140.4 | 147.7 |
| 1968 | 14.8 | 9.8 | 1.7 | 0.7 | 1.4 | 2.9 | 8.6 | 26.7 | 54.1 | 89.6 | 144.1 | 122.0 |

ALL OTHER, FEMALE

| | | | | | | | | | | | | |
|------|-----|------|-----|-----|-----|-----|-----|------|------|------|------|------|
| 1978 | 5.3 | * | * | * | 0.6 | 1.2 | 2.8 | 7.4 | 18.6 | 20.6 | 40.6 | 67.0 |
| 1977 | 4.8 | * | * | * | * | * | 3.3 | 7.1 | 14.5 | 21.7 | 36.8 | 53.8 |
| 1976 | 4.7 | * | * | * | * | 1.2 | 2.7 | 8.1 | 16.5 | 20.9 | 28.5 | 44.9 |
| 1975 | 4.8 | * | * | * | * | 1.6 | 3.8 | 9.5 | 14.3 | 20.7 | 29.4 | 34.4 |
| 1974 | 4.3 | * | * | * | 1.0 | 1.0 | 3.7 | 6.8 | 13.0 | 20.6 | 24.1 | 31.8 |
| 1973 | 4.7 | * | * | * | 0.7 | 1.6 | 5.0 | 8.9 | 14.9 | 17.6 | 28.6 | 36.7 |
| 1972 | 4.8 | * | * | 0.8 | 0.9 | 1.6 | 5.2 | 8.0 | 14.9 | 16.2 | 25.9 | 50.0 |
| 1971 | 5.1 | * | * | * | 0.9 | 2.0 | 5.5 | 10.6 | 14.8 | 19.8 | 25.4 | 28.8 |
| 1970 | 5.0 | * | * | * | 1.6 | 2.8 | 5.2 | 9.8 | 14.2 | 16.8 | 24.8 | 29.2 |
| 1969 | 5.0 | 10.5 | 1.9 | * | 1.5 | 3.2 | 4.9 | 9.4 | 13.4 | 15.3 | 25.1 | 40.0 |
| 1968 | 5.2 | 8.4 | 1.7 | 0.7 | 1.6 | 3.4 | 6.6 | 9.8 | 14.5 | 14.6 | 22.2 | 42.6 |

*519.3 Code introduced by NCHS in 1969

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|

Chronic obstructive pulmonary diseases and allied conditions 490-493, *519.3

BLACK, BOTH SEXES

| | | | | | | | | | | | | |
|------|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| 1978 | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- |
| 1977 | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- |
| 1976 | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- |
| 1975 | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- |
| 1974 | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- |
| 1973 | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- |
| 1972 | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- |
| 1971 | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- |
| 1970 | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- |
| 1969 | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- |
| 1968 | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- |

BLACK, MALE

| | | | | | | | | | | | | |
|------|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| 1978 | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- |
| 1977 | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- |
| 1976 | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- |
| 1975 | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- |
| 1974 | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- |
| 1973 | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- |
| 1972 | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- |
| 1971 | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- |
| 1970 | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- |
| 1969 | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- |
| 1968 | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- |

BLACK, FEMALE

| | | | | | | | | | | | | |
|------|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| 1978 | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- |
| 1977 | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- |
| 1976 | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- |
| 1975 | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- |
| 1974 | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- |
| 1973 | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- |
| 1972 | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- |
| 1971 | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- |
| 1970 | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- |
| 1969 | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- |
| 1968 | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- |

--- Data not available

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|--|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| BRONCHITIS, EMPHYSEMA, AND ASTHMAS (490-493) | | | | | | | | | | | | |
| BOTH SEXES | | | | | | | | | | | | |
| 1978 | 9.8 | 2.0 | 0.4 | 0.2 | 0.3 | 0.4 | 1.1 | 5.0 | 19.3 | 51.0 | 84.8 | 94.2 |
| 1977 | 10.2 | 2.1 | 0.4 | 0.2 | 0.3 | 0.4 | 1.3 | 5.2 | 20.0 | 54.2 | 88.7 | 95.7 |
| 1976 | 11.2 | 2.3 | 0.5 | 0.2 | 0.2 | 0.5 | 1.3 | 6.0 | 22.6 | 60.5 | 96.2 | 112.5 |
| 1975 | 11.8 | 3.1 | 0.5 | 0.2 | 0.3 | 0.6 | 1.6 | 6.4 | 25.0 | 65.2 | 101.5 | 104.4 |
| 1974 | 12.6 | 2.7 | 0.4 | 0.2 | 0.4 | 0.4 | 1.7 | 7.0 | 27.0 | 70.8 | 110.7 | 110.8 |
| 1973 | 14.1 | 3.1 | 0.5 | 0.2 | 0.4 | 0.5 | 2.0 | 7.7 | 31.1 | 79.2 | 122.6 | 135.1 |
| 1972 | 14.7 | 3.8 | 0.6 | 0.2 | 0.4 | 0.7 | 2.3 | 8.7 | 33.3 | 84.2 | 125.6 | 130.5 |
| 1971 | 14.6 | 3.4 | 0.7 | 0.3 | 0.4 | 0.7 | 2.3 | 9.3 | 33.7 | 83.1 | 125.7 | 131.4 |
| 1970 | 15.2 | 4.2 | 0.9 | 0.3 | 0.5 | 0.8 | 2.7 | 9.7 | 36.2 | 86.6 | 129.1 | 121.2 |
| 1969 | 15.5 | 4.6 | 0.8 | 0.3 | 0.5 | 0.9 | 2.7 | 10.1 | 37.2 | 87.7 | 130.0 | 144.5 |
| 1968 | 16.6 | 5.2 | 0.9 | 0.3 | 0.5 | 1.0 | 2.9 | 10.9 | 40.8 | 94.4 | 139.1 | 154.5 |
| MALE | | | | | | | | | | | | |
| 1978 | 14.1 | 2.8 | 0.5 | 0.2 | 0.3 | 0.4 | 1.0 | 5.9 | 27.0 | 86.5 | 168.7 | 191.4 |
| 1977 | 15.1 | 2.7 | 0.5 | 0.2 | 0.3 | 0.4 | 1.4 | 6.6 | 29.3 | 94.0 | 179.6 | 199.5 |
| 1976 | 16.8 | 2.8 | 0.5 | 0.2 | 0.3 | 0.4 | 1.3 | 7.5 | 33.8 | 107.6 | 194.3 | 237.8 |
| 1975 | 18.1 | 4.0 | 0.6 | 0.2 | 0.3 | 0.6 | 1.6 | 8.1 | 37.7 | 118.1 | 209.9 | 224.6 |
| 1974 | 19.4 | 3.6 | 0.5 | 0.2 | 0.4 | 0.5 | 1.8 | 9.2 | 41.5 | 128.5 | 228.6 | 233.6 |
| 1973 | 22.1 | 4.3 | 0.6 | 0.2 | 0.4 | 0.5 | 1.9 | 10.3 | 49.4 | 146.7 | 255.2 | 281.4 |
| 1972 | 23.2 | 4.5 | 0.5 | 0.3 | 0.5 | 0.7 | 2.2 | 11.9 | 53.6 | 155.8 | 261.6 | 268.2 |
| 1971 | 23.3 | 4.3 | 0.8 | 0.2 | 0.4 | 0.6 | 2.5 | 12.4 | 55.0 | 156.1 | 259.4 | 268.0 |
| 1970 | 24.4 | 4.8 | 1.0 | 0.4 | 0.5 | 0.6 | 2.9 | 13.1 | 59.5 | 164.8 | 266.8 | 238.9 |
| 1969 | 25.2 | 4.9 | 0.9 | 0.3 | 0.5 | 0.8 | 3.2 | 14.2 | 62.4 | 169.6 | 265.5 | 275.2 |
| 1968 | 27.0 | 5.7 | 1.0 | 0.3 | 0.5 | 0.9 | 3.2 | 15.5 | 68.5 | 182.2 | 281.3 | 298.4 |

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|--|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| BRONCHITIS, EMPHYSEMA, AND ASTHMAS (490-493) | | | | | | | | | | | | |
| FEMALE | | | | | | | | | | | | |
| 1978 | 5.8 | 1.3 | * | * | 0.3 | 0.4 | 1.1 | 4.2 | 12.6 | 23.9 | 34.9 | 50.3 |
| 1977 | 5.5 | 1.6 | * | 0.2 | 0.2 | 0.3 | 1.2 | 3.8 | 11.7 | 23.8 | 34.3 | 48.0 |
| 1976 | 5.9 | 1.8 | 0.4 | 0.2 | 0.2 | 0.5 | 1.4 | 4.6 | 12.8 | 24.5 | 36.9 | 53.6 |
| 1975 | 5.9 | 2.1 | 0.5 | 0.2 | 0.2 | 0.5 | 1.5 | 4.9 | 13.8 | 24.6 | 35.4 | 46.2 |
| 1974 | 6.1 | 1.8 | 0.4 | 0.2 | 0.4 | 0.4 | 1.6 | 5.0 | 14.2 | 26.5 | 38.1 | 50.0 |
| 1973 | 6.4 | 1.8 | 0.4 | 0.2 | 0.4 | 0.5 | 2.0 | 5.2 | 14.8 | 27.3 | 39.7 | 61.5 |
| 1972 | 6.6 | 3.1 | 0.7 | 0.2 | 0.3 | 0.8 | 2.3 | 5.8 | 15.3 | 28.9 | 39.0 | 60.0 |
| 1971 | 6.4 | 2.6 | 0.6 | 0.3 | 0.4 | 0.8 | 2.1 | 6.3 | 14.7 | 26.5 | 39.2 | 60.1 |
| 1970 | 6.4 | 3.5 | 0.7 | 0.3 | 0.6 | 1.0 | 2.4 | 6.6 | 15.2 | 25.9 | 38.0 | 55.2 |
| 1969 | 6.3 | 4.3 | 0.8 | 0.2 | 0.5 | 1.0 | 2.3 | 6.3 | 14.4 | 23.7 | 39.0 | 73.8 |
| 1968 | 6.7 | 4.7 | 0.8 | 0.3 | 0.6 | 1.2 | 2.5 | 6.5 | 15.8 | 25.3 | 41.8 | 75.5 |
| WHITE, BOTH SEXES | | | | | | | | | | | | |
| 1978 | 10.6 | 1.8 | 0.3 | 0.2 | 0.3 | 0.3 | 0.9 | 4.8 | 19.7 | 53.7 | 89.3 | 97.9 |
| 1977 | 11.0 | 1.7 | 0.3 | 0.2 | 0.2 | 0.3 | 1.1 | 4.9 | 20.4 | 57.1 | 93.3 | 99.9 |
| 1976 | 12.1 | 2.0 | 0.4 | 0.1 | 0.2 | 0.4 | 1.2 | 5.8 | 23.1 | 63.5 | 101.6 | 117.4 |
| 1975 | 12.7 | 2.6 | 0.5 | 0.1 | 0.3 | 0.4 | 1.3 | 6.1 | 25.7 | 68.6 | 106.6 | 109.3 |
| 1974 | 13.6 | 2.1 | 0.4 | 0.1 | 0.3 | 0.3 | 1.4 | 6.9 | 27.9 | 74.1 | 116.6 | 115.7 |
| 1973 | 15.1 | 2.5 | 0.4 | 0.2 | 0.3 | 0.4 | 1.7 | 7.3 | 31.8 | 83.4 | 128.5 | 140.8 |
| 1972 | 15.8 | 2.9 | 0.5 | 0.2 | 0.3 | 0.6 | 1.9 | 8.4 | 34.2 | 88.6 | 131.0 | 135.2 |
| 1971 | 15.6 | 2.9 | 0.6 | 0.2 | 0.3 | 0.5 | 1.9 | 8.8 | 34.6 | 87.1 | 131.4 | 137.0 |
| 1970 | 16.2 | 3.3 | 0.7 | 0.3 | 0.4 | 0.6 | 2.2 | 9.5 | 36.9 | 91.1 | 133.9 | 127.1 |
| 1969 | 16.4 | 3.6 | 0.6 | 0.2 | 0.3 | 0.6 | 2.3 | 9.6 | 37.9 | 91.8 | 135.0 | 151.1 |
| 1968 | 17.5 | 4.5 | 0.7 | 0.3 | 0.4 | 0.7 | 2.3 | 10.1 | 41.6 | 99.0 | 144.0 | 161.6 |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|--|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| BRONCHITIS, EMPHYSEMA, AND ASTHMAS (490-493) | | | | | | | | | | | | |
| WHITE, MALE | | | | | | | | | | | | |
| 1978 | 15.2 | 2.5 | * | 0.2 | 0.3 | 0.3 | 0.8 | 5.5 | 27.2 | 90.5 | 179.1 | 201.9 |
| 1977 | 16.2 | 2.3 | 0.4 | 0.1 | 0.2 | 0.3 | 1.2 | 6.2 | 29.7 | 98.9 | 190.7 | 209.8 |
| 1976 | 18.1 | 2.4 | 0.5 | 0.1 | 0.2 | 0.4 | 1.1 | 7.1 | 34.3 | 112.8 | 206.5 | 250.3 |
| 1975 | 19.5 | 3.3 | 0.6 | 0.1 | 0.3 | 0.4 | 1.3 | 7.7 | 38.5 | 124.6 | 221.9 | 237.2 |
| 1974 | 20.9 | 2.9 | 0.4 | 0.1 | 0.3 | 0.4 | 1.4 | 9.0 | 42.5 | 134.7 | 242.3 | 245.6 |
| 1973 | 23.7 | 3.2 | 0.5 | 0.2 | 0.2 | 0.4 | 1.7 | 9.8 | 50.4 | 154.7 | 268.9 | 295.3 |
| 1972 | 24.9 | 3.4 | 0.4 | 0.2 | 0.4 | 0.6 | 1.9 | 11.4 | 55.0 | 164.3 | 274.2 | 281.7 |
| 1971 | 25.0 | 3.9 | 0.6 | 0.2 | 0.3 | 0.4 | 2.1 | 11.9 | 56.3 | 164.4 | 273.1 | 281.6 |
| 1970 | 26.1 | 3.7 | 0.9 | 0.3 | 0.3 | 0.5 | 2.4 | 12.8 | 60.7 | 174.1 | 278.5 | 253.8 |
| 1969 | 26.8 | 3.9 | 0.7 | 0.2 | 0.4 | 0.6 | 2.8 | 13.4 | 63.8 | 178.1 | 277.5 | 291.1 |
| 1968 | 28.7 | 5.0 | 0.8 | 0.3 | 0.3 | 0.6 | 2.6 | 14.3 | 70.0 | 191.7 | 292.7 | 316.5 |
| WHITE, FEMALE | | | | | | | | | | | | |
| 1978 | 6.3 | * | * | * | 0.2 | 0.2 | 1.0 | 4.1 | 12.9 | 25.6 | 36.6 | 51.6 |
| 1977 | 6.0 | * | * | 0.2 | 0.1 | 0.3 | 1.0 | 3.8 | 12.2 | 25.1 | 35.8 | 50.1 |
| 1976 | 6.4 | 1.6 | * | 0.1 | 0.2 | 0.4 | 1.3 | 4.6 | 13.1 | 25.8 | 39.0 | 56.1 |
| 1975 | 6.3 | 2.0 | 0.4 | * | 0.2 | 0.4 | 1.3 | 4.6 | 14.3 | 25.8 | 37.1 | 48.4 |
| 1974 | 6.6 | * | * | 0.1 | 0.3 | 0.3 | 1.3 | 5.0 | 14.9 | 27.9 | 40.0 | 52.4 |
| 1973 | 6.9 | 1.7 | * | 0.1 | 0.3 | 0.4 | 1.7 | 5.0 | 15.2 | 28.8 | 41.5 | 64.4 |
| 1972 | 7.0 | 2.4 | 0.6 | * | 0.2 | 0.6 | 2.0 | 5.7 | 15.7 | 30.4 | 40.8 | 61.4 |
| 1971 | 6.7 | 1.9 | 0.5 | 0.3 | 0.3 | 0.6 | 1.6 | 6.0 | 15.1 | 27.6 | 40.7 | 62.9 |
| 1970 | 6.7 | 2.9 | 0.5 | 0.2 | 0.4 | 0.7 | 2.0 | 6.4 | 15.5 | 27.0 | 39.3 | 57.8 |
| 1969 | 6.5 | 3.2 | 0.6 | 0.2 | 0.3 | 0.6 | 1.9 | 6.1 | 14.6 | 24.7 | 40.2 | 77.1 |
| 1968 | 6.9 | 4.0 | 0.7 | 0.3 | 0.4 | 0.8 | 2.0 | 6.1 | 16.0 | 26.4 | 43.3 | 78.2 |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|--|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| BRONCHITIS, EMPHYSEMA, AND ASTHMAS (490-493) | | | | | | | | | | | | |
| ALL OTHER, BOTH SEXES | | | | | | | | | | | | |
| 1978 | 4.9 | * | 0.9 | * | 0.6 | 1.2 | 2.4 | 6.9 | 16.4 | 26.2 | 38.1 | 51.8 |
| 1977 | 4.9 | 4.3 | * | * | 0.7 | 0.7 | 2.6 | 6.8 | 15.8 | 27.6 | 41.4 | 47.5 |
| 1976 | 5.4 | * | 0.9 | * | 0.5 | 1.1 | 2.4 | 7.9 | 18.5 | 32.3 | 40.9 | 55.6 |
| 1975 | 5.8 | 5.3 | * | 0.5 | 0.5 | 1.8 | 3.5 | 8.9 | 18.6 | 31.8 | 47.2 | 48.6 |
| 1974 | 6.0 | 5.8 | * | 0.4 | 1.0 | 1.0 | 4.2 | 7.5 | 19.2 | 37.8 | 45.3 | 53.3 |
| 1973 | 6.9 | 5.9 | 1.0 | 0.5 | 0.9 | 1.6 | 4.1 | 10.4 | 24.4 | 38.4 | 56.5 | 68.5 |
| 1972 | 7.5 | 8.5 | 1.3 | 0.7 | 0.9 | 1.5 | 4.7 | 11.2 | 24.7 | 40.5 | 63.2 | 76.4 |
| 1971 | 7.7 | 6.2 | 1.5 | 0.5 | 0.9 | 2.2 | 5.5 | 13.1 | 25.7 | 43.1 | 57.5 | 67.5 |
| 1970 | 8.4 | 8.9 | 1.8 | 0.7 | 1.6 | 2.4 | 6.1 | 12.0 | 29.3 | 42.7 | 68.7 | 60.4 |
| 1969 | 8.8 | 10.7 | 1.9 | 0.7 | 1.6 | 2.5 | 5.5 | 14.4 | 30.1 | 47.0 | 65.0 | 70.6 |
| 1968 | 9.8 | 9.1 | 1.7 | 0.7 | 1.5 | 3.2 | 7.5 | 17.7 | 33.2 | 48.6 | 75.4 | 74.5 |
| ALL OTHER, MALE | | | | | | | | | | | | |
| 1978 | 7.2 | * | * | * | 0.6 | 1.2 | 2.9 | 9.8 | 24.9 | 49.4 | 69.5 | 84.5 |
| 1977 | 7.3 | * | * | * | 0.9 | * | 2.5 | 10.3 | 25.5 | 48.9 | 75.1 | 94.6 |
| 1976 | 8.1 | * | * | * | 0.7 | 1.1 | 2.8 | 11.1 | 28.7 | 58.6 | 80.2 | 112.7 |
| 1975 | 8.5 | 7.5 | * | * | * | 2.2 | 3.8 | 11.1 | 30.4 | 55.8 | 91.7 | 98.1 |
| 1974 | 9.1 | * | * | * | 1.0 | * | 4.9 | 11.0 | 32.2 | 69.2 | 88.2 | 110.0 |
| 1973 | 10.6 | 9.4 | * | * | 1.1 | 1.6 | 3.4 | 15.0 | 40.5 | 70.5 | 111.2 | 139.6 |
| 1972 | 11.2 | 10.3 | * | * | 1.0 | 1.3 | 4.9 | 16.1 | 40.2 | 74.2 | 126.3 | 131.9 |
| 1971 | 11.5 | * | 1.9 | * | 0.8 | 2.4 | 5.8 | 17.7 | 42.2 | 76.8 | 107.7 | 134.0 |
| 1970 | 12.5 | 11.2 | 2.1 | 0.7 | 1.6 | 1.9 | 7.3 | 15.5 | 48.5 | 76.9 | 131.1 | 108.3 |
| 1969 | 13.3 | 10.9 | 2.0 | 0.8 | 1.7 | 1.8 | 6.4 | 21.0 | 49.6 | 87.4 | 121.8 | 125.0 |
| 1968 | 14.8 | 9.8 | 1.7 | 0.7 | 1.4 | 2.9 | 8.6 | 26.7 | 54.1 | 89.6 | 144.1 | 122.0 |
| ALL OTHER, FEMALE | | | | | | | | | | | | |
| 1978 | 2.8 | * | * | * | * | 1.2 | 1.9 | 4.5 | 9.4 | 8.3 | 16.7 | 34.9 |
| 1977 | 2.7 | * | * | * | * | * | 2.7 | 3.9 | 7.6 | 11.0 | 18.3 | 22.1 |
| 1976 | 2.9 | * | * | * | * | 1.2 | 2.1 | 5.2 | 10.0 | 11.5 | 14.1 | 23.5 |
| 1975 | 3.3 | * | * | * | * | 1.5 | 3.3 | 7.1 | 8.7 | 12.7 | 16.7 | * |
| 1974 | 3.0 | * | * | * | 0.9 | 1.0 | 3.7 | 4.6 | 8.2 | 12.6 | 15.7 | * |
| 1973 | 3.6 | * | * | * | 0.7 | 1.6 | 4.6 | 6.5 | 10.7 | 12.4 | 18.2 | 25.3 |
| 1972 | 4.0 | * | * | 0.8 | 0.9 | 1.6 | 4.5 | 6.9 | 11.4 | 13.1 | 17.9 | 42.1 |
| 1971 | 4.3 | * | * | * | 0.9 | 2.0 | 5.3 | 9.0 | 11.2 | 15.6 | 20.4 | * |
| 1970 | 4.6 | * | * | * | 1.6 | 2.7 | 5.0 | 8.9 | 12.4 | 14.5 | 21.3 | 26.7 |
| 1969 | 4.7 | 10.5 | 1.9 | * | 1.5 | 3.2 | 4.7 | 8.6 | 12.9 | 13.8 | 21.5 | 33.8 |
| 1968 | 5.2 | 8.4 | * | 0.7 | 1.6 | 3.4 | 6.6 | 9.8 | 14.5 | 14.6 | 22.2 | 42.6 |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|--|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| BRONCHITIS, EMPHYSEMA, AND ASTHMAS (490-493) | | | | | | | | | | | | |
| BLACK, BOTH SEXES | | | | | | | | | | | | |
| 1978 | 5.2 | * | 1.1 | * | 0.6 | 1.4 | 2.8 | 7.7 | 17.8 | 26.1 | 34.8 | 50.7 |
| 1977 | 5.2 | 4.3 | * | * | 0.7 | 0.8 | 3.1 | 7.6 | 16.9 | 28.7 | 40.0 | 41.3 |
| 1976 | 5.8 | * | * | * | 0.6 | 1.3 | 2.7 | 8.8 | 19.7 | 33.4 | 39.1 | 55.1 |
| 1975 | 6.1 | 5.0 | * | 0.4 | 0.6 | 2.0 | 4.0 | 9.7 | 19.9 | 32.8 | 45.3 | 40.8 |
| 1974 | 6.2 | 5.6 | * | 0.5 | 1.0 | 1.2 | 4.8 | 8.2 | 20.3 | 38.2 | 41.5 | 43.8 |
| 1973 | 7.2 | 5.9 | 1.1 | 0.5 | 0.9 | 1.8 | 4.5 | 11.4 | 25.4 | 39.1 | 52.9 | 54.4 |
| 1972 | 7.8 | 8.8 | 1.3 | 0.7 | 1.1 | 1.6 | 5.2 | 12.3 | 25.7 | 42.0 | 62.2 | 65.5 |
| 1971 | 8.0 | 6.3 | 1.5 | 0.5 | 0.9 | 2.4 | 6.1 | 13.6 | 26.7 | 44.1 | 55.7 | 63.6 |
| 1970 | 8.8 | 9.0 | 1.9 | 0.7 | 1.7 | 2.6 | 6.6 | 13.1 | 31.3 | 43.8 | 66.0 | 53.7 |
| 1969 | 9.1 | 11.6 | 1.8 | 0.7 | 1.7 | 2.8 | 5.8 | 15.2 | 31.6 | 47.1 | 62.8 | 63.9 |
| 1968 | 10.1 | 9.1 | 1.9 | 0.7 | 1.6 | 3.3 | 7.9 | 18.4 | 34.8 | 48.9 | 72.9 | 68.5 |
| BLACK, MALE | | | | | | | | | | | | |
| 1978 | 7.6 | * | * | * | * | 1.3 | 3.6 | 11.0 | 27.2 | 50.1 | 65.3 | 82.4 |
| 1977 | 7.8 | * | * | * | 1.0 | * | 2.9 | 11.6 | 28.1 | 51.9 | 75.7 | 75.5 |
| 1976 | 8.7 | * | * | * | * | 1.2 | 3.1 | 12.4 | 30.8 | 62.5 | 77.4 | 110.6 |
| 1975 | 9.1 | * | * | * | * | 2.4 | 4.4 | 12.0 | 33.0 | 59.4 | 91.0 | 80.4 |
| 1974 | 9.5 | * | * | * | 1.1 | * | 5.6 | 12.3 | 34.5 | 72.1 | 82.3 | 90.7 |
| 1973 | 11.0 | 9.0 | * | * | 1.2 | 1.8 | 3.8 | 16.6 | 42.3 | 73.8 | 103.4 | 119.5 |
| 1972 | 11.7 | 10.8 | * | * | 1.1 | 1.5 | 5.3 | 17.8 | 42.6 | 78.9 | 124.9 | 110.0 |
| 1971 | 12.0 | * | * | * | * | 2.7 | 6.4 | 18.9 | 44.7 | 80.3 | 107.0 | 130.0 |
| 1970 | 13.2 | 11.9 | 2.3 | 0.8 | 1.7 | 2.1 | 8.1 | 17.0 | 53.0 | 80.5 | 129.3 | 95.7 |
| 1969 | 13.7 | 11.8 | 2.0 | 0.9 | 1.7 | 2.1 | 6.5 | 22.7 | 52.9 | 88.6 | 121.0 | 116.2 |
| 1968 | 15.2 | 10.1 | 1.9 | 0.7 | 1.5 | 3.0 | 9.0 | 28.3 | 57.5 | 92.3 | 140.0 | 114.3 |
| BLACK, FEMALE | | | | | | | | | | | | |
| 1978 | 3.0 | * | * | * | * | 1.4 | 2.1 | 5.1 | 10.2 | 8.3 | 14.5 | 34.3 |
| 1977 | 2.9 | * | * | * | * | * | 3.2 | 4.4 | 7.7 | 11.3 | 16.4 | 23.4 |
| 1976 | 3.1 | * | * | * | * | 1.4 | 2.4 | 5.9 | 10.5 | 11.4 | 13.5 | 25.8 |
| 1975 | 3.4 | * | * | * | * | 1.7 | 3.7 | 7.7 | 9.1 | 12.7 | 14.3 | * |
| 1974 | 3.2 | * | * | * | 1.0 | 1.2 | 4.1 | 4.8 | 8.5 | 12.2 | 13.9 | * |
| 1973 | 3.8 | * | * | * | * | 1.8 | 5.0 | 7.1 | 11.4 | 12.4 | 18.2 | * |
| 1972 | 4.3 | * | * | 0.9 | 1.0 | 1.8 | 5.2 | 7.6 | 11.5 | 13.3 | 17.9 | 40.0 |
| 1971 | 4.4 | * | * | * | 0.9 | 2.1 | 5.7 | 9.1 | 11.6 | 15.5 | 18.9 | * |
| 1970 | 4.8 | * | * | * | 1.6 | 3.1 | 5.3 | 9.7 | 12.8 | 14.8 | 19.6 | * |
| 1969 | 4.8 | 11.4 | * | * | 1.6 | 3.4 | 5.1 | 8.8 | 13.4 | 14.2 | 19.8 | * |
| 1968 | 5.3 | * | * | * | 1.7 | 3.6 | 6.9 | 9.7 | 15.1 | 13.9 | 23.0 | 41.1 |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| CHRONIC AND UNQUALIFIED BRONCHITIS (490, 491) | | | | | | | | | | | | |
| BOTH SEXES | | | | | | | | | | | | |
| 1978 | 2.0 | 1.7 | 0.3 | 0.1 | 0.1 | 0.1 | 0.2 | 0.8 | 3.5 | 9.3 | 17.1 | 27.2 |
| 1977 | 2.0 | 1.6 | 0.3 | 0.1 | 0.1 | 0.1 | 0.2 | 0.9 | 3.5 | 9.6 | 16.7 | 27.0 |
| 1976 | 2.1 | 1.8 | 0.3 | * | * | 0.1 | 0.2 | 1.1 | 4.0 | 10.2 | 18.4 | 31.4 |
| 1975 | 2.2 | 2.4 | 0.4 | 0.1 | 0.0 | 0.1 | 0.3 | 1.0 | 4.4 | 10.8 | 19.0 | 27.5 |
| 1974 | 2.4 | 2.4 | 0.2 | 0.1 | 0.1 | 0.1 | 0.3 | 1.3 | 4.7 | 12.0 | 21.9 | 29.6 |
| 1973 | 2.6 | 2.1 | 0.3 | 0.1 | 0.1 | 0.1 | 0.4 | 1.3 | 5.2 | 13.1 | 24.0 | 38.1 |
| 1972 | 2.8 | 3.0 | 0.3 | 0.1 | 0.1 | 0.2 | 0.5 | 1.7 | 5.6 | 15.1 | 23.6 | 34.1 |
| 1971 | 2.7 | 2.6 | 0.4 | 0.1 | 0.1 | 0.1 | 0.4 | 1.5 | 5.8 | 13.9 | 24.0 | 34.0 |
| 1970 | 2.9 | 3.3 | 0.5 | 0.1 | 0.1 | 0.1 | 0.4 | 1.8 | 6.3 | 14.4 | 25.7 | 31.8 |
| 1969 | 2.9 | 3.5 | 0.5 | 0.1 | 0.1 | 0.1 | 0.4 | 1.9 | 6.1 | 14.9 | 25.4 | 40.2 |
| 1968 | 3.1 | 4.3 | 0.5 | 0.1 | 0.1 | 0.1 | 0.4 | 1.8 | 7.0 | 16.0 | 27.4 | 43.2 |
| MALE | | | | | | | | | | | | |
| 1978 | 2.6 | 2.2 | 0.3 | * | * | 0.1 | * | 0.8 | 4.6 | 14.5 | 30.3 | 45.9 |
| 1977 | 2.6 | 1.8 | 0.3 | * | 0.1 | * | 0.2 | 1.2 | 4.8 | 15.5 | 30.2 | 42.9 |
| 1976 | 2.9 | 2.1 | 0.3 | * | * | * | 0.2 | 1.4 | 5.7 | 17.1 | 33.8 | 55.3 |
| 1975 | 3.1 | 2.9 | 0.4 | * | * | * | 0.3 | 1.2 | 6.4 | 18.2 | 35.9 | 54.5 |
| 1974 | 3.4 | 3.2 | * | * | * | * | 0.3 | 1.7 | 6.6 | 21.0 | 41.1 | 55.0 |
| 1973 | 3.8 | 3.1 | 0.4 | 0.1 | * | 0.1 | 0.4 | 1.5 | 7.8 | 22.5 | 46.0 | 68.4 |
| 1972 | 4.1 | 3.4 | * | 0.1 | * | 0.2 | 0.6 | 2.3 | 8.9 | 26.0 | 45.7 | 60.5 |
| 1971 | 4.0 | 3.0 | 0.5 | * | 0.1 | * | 0.5 | 1.9 | 9.1 | 24.8 | 45.9 | 60.8 |
| 1970 | 4.3 | 3.7 | 0.7 | 0.1 | 0.1 | * | 0.5 | 2.3 | 10.0 | 25.9 | 49.9 | 53.5 |
| 1969 | 4.4 | 3.8 | 0.6 | * | 0.1 | * | 0.6 | 2.6 | 9.9 | 27.4 | 48.1 | 66.4 |
| 1968 | 4.7 | 4.7 | 0.6 | * | * | * | 0.5 | 2.6 | 11.2 | 28.8 | 51.7 | 74.5 |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| CHRONIC AND UNQUALIFIED BRONCHITIS (490, 491) | | | | | | | | | | | | |
| FEMALE | | | | | | | | | | | | |
| 1978 | 1.4 | * | * | * | * | * | 0.2 | 0.8 | 2.5 | 5.3 | 9.3 | 18.8 |
| 1977 | 1.4 | 1.4 | * | * | * | * | 0.2 | 0.7 | 2.4 | 5.0 | 8.6 | 19.7 |
| 1976 | 1.4 | 1.5 | * | * | * | * | 0.2 | 0.9 | 2.5 | 4.9 | 9.1 | 20.2 |
| 1975 | 1.3 | 1.8 | 0.3 | * | * | * | 0.2 | 0.9 | 2.7 | 5.0 | 8.8 | 14.4 |
| 1974 | 1.4 | 1.5 | * | * | 0.1 | * | 0.3 | 0.9 | 2.9 | 5.1 | 10.0 | 17.0 |
| 1973 | 1.5 | * | * | * | * | * | 0.3 | 1.0 | 2.9 | 5.9 | 10.3 | 22.9 |
| 1972 | 1.5 | 2.6 | 0.5 | * | * | * | 0.4 | 1.1 | 2.7 | 6.7 | 9.4 | 20.6 |
| 1971 | 1.4 | 2.2 | 0.4 | * | * | * | 0.3 | 1.1 | 2.8 | 5.5 | 9.8 | 20.1 |
| 1970 | 1.5 | 2.9 | 0.4 | * | 0.1 | * | 0.4 | 1.4 | 3.0 | 5.5 | 9.7 | 19.6 |
| 1969 | 1.4 | 3.3 | 0.5 | * | * | * | 0.3 | 1.2 | 2.6 | 5.1 | 10.1 | 25.9 |
| 1968 | 1.6 | 3.9 | 0.5 | * | * | * | 0.4 | 1.1 | 3.2 | 5.9 | 10.8 | 26.0 |
| WHITE, BOTH SEXES | | | | | | | | | | | | |
| 1978 | 2.1 | 1.5 | 0.2 | * | 0.1 | 0.1 | 0.2 | 0.8 | 3.6 | 9.9 | 18.1 | 28.6 |
| 1977 | 2.1 | 1.2 | 0.3 | 0.1 | 0.1 | 0.1 | 0.2 | 1.0 | 3.6 | 10.1 | 17.6 | 28.3 |
| 1976 | 2.3 | 1.7 | 0.3 | * | * | 0.1 | 0.2 | 1.1 | 4.1 | 10.7 | 19.5 | 33.4 |
| 1975 | 2.4 | 2.1 | 0.4 | 0.1 | * | * | 0.2 | 1.1 | 4.6 | 11.3 | 20.2 | 28.8 |
| 1974 | 2.6 | 1.8 | 0.2 | * | 0.1 | 0.1 | 0.3 | 1.3 | 4.8 | 12.4 | 23.1 | 31.1 |
| 1973 | 2.8 | 1.8 | 0.3 | 0.1 | * | 0.1 | 0.3 | 1.2 | 5.3 | 13.9 | 25.1 | 39.7 |
| 1972 | 3.0 | 2.3 | 0.3 | 0.1 | 0.1 | 0.2 | 0.4 | 1.6 | 5.7 | 15.7 | 24.8 | 34.8 |
| 1971 | 2.9 | 2.2 | 0.4 | 0.1 | 0.1 | 0.1 | 0.3 | 1.4 | 6.0 | 14.7 | 25.0 | 35.3 |
| 1970 | 3.1 | 2.6 | 0.5 | 0.1 | 0.1 | 0.1 | 0.4 | 1.8 | 6.4 | 15.2 | 26.8 | 33.4 |
| 1969 | 3.1 | 2.7 | 0.4 | 0.1 | 0.1 | 0.1 | 0.5 | 1.8 | 6.2 | 15.6 | 26.3 | 42.4 |
| 1968 | 3.3 | 3.7 | 0.4 | 0.1 | * | 0.1 | 0.3 | 1.7 | 7.1 | 16.7 | 28.5 | 45.1 |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| CHRONIC AND UNQUALIFIED BRONCHITIS (490, 491) | | | | | | | | | | | | |
| WHITE, MALE | | | | | | | | | | | | |
| 1978 | 2.8 | 2.1 | * | * | * | * | * | 0.8 | 4.8 | 15.3 | 32.3 | 49.2 |
| 1977 | 2.8 | 1.4 | * | * | * | * | * | 1.2 | 4.9 | 16.4 | 31.9 | 44.7 |
| 1976 | 3.2 | 1.8 | * | * | * | * | * | 1.2 | 5.8 | 17.9 | 36.0 | 59.2 |
| 1975 | 3.4 | 2.4 | 0.5 | * | * | * | 0.2 | 1.3 | 6.5 | 19.2 | 38.3 | 58.0 |
| 1974 | 3.7 | 2.5 | * | * | * | * | 0.2 | 1.7 | 6.7 | 21.8 | 43.8 | 57.7 |
| 1973 | 4.1 | 2.4 | * | * | * | * | 0.3 | 1.4 | 7.9 | 23.9 | 48.3 | 71.4 |
| 1972 | 4.4 | 2.8 | * | * | * | 0.2 | 0.5 | 2.1 | 9.0 | 27.2 | 48.3 | 62.7 |
| 1971 | 4.3 | 2.7 | 0.4 | * | * | * | 0.4 | 1.8 | 9.3 | 26.3 | 48.2 | 64.1 |
| 1970 | 4.6 | 2.7 | 0.6 | 0.1 | * | * | 0.5 | 2.2 | 10.1 | 27.4 | 52.4 | 56.5 |
| 1969 | 4.7 | 3.0 | 0.5 | * | * | * | 0.6 | 2.5 | 10.1 | 28.9 | 50.0 | 70.6 |
| 1968 | 5.0 | 4.1 | 0.4 | * | * | * | 0.4 | 2.4 | 11.4 | 30.2 | 54.0 | 78.2 |
| WHITE, FEMALE | | | | | | | | | | | | |
| 1978 | 1.5 | * | * | * | * | * | 0.2 | 0.8 | 2.6 | 5.7 | 9.8 | 19.4 |
| 1977 | 1.5 | * | * | * | * | * | 0.2 | 0.8 | 2.5 | 5.3 | 9.1 | 20.9 |
| 1976 | 1.5 | * | * | * | * | * | 0.2 | 0.9 | 2.5 | 5.2 | 9.7 | 21.5 |
| 1975 | 1.4 | 1.7 | * | * | * | * | 0.2 | 0.9 | 2.9 | 5.3 | 9.3 | 15.0 |
| 1974 | 1.5 | * | * | * | * | * | 0.3 | 1.0 | 3.1 | 5.3 | 10.5 | 18.2 |
| 1973 | 1.6 | * | * | * | * | * | 0.3 | 1.0 | 3.0 | 6.3 | 10.8 | 24.1 |
| 1972 | 1.6 | 1.8 | 0.4 | * | * | * | 0.4 | 1.1 | 2.8 | 6.9 | 10.0 | 20.8 |
| 1971 | 1.5 | 1.6 | 0.4 | * | * | * | 0.3 | 1.1 | 2.9 | 5.7 | 10.2 | 20.6 |
| 1970 | 1.6 | 2.4 | 0.4 | * | * | * | 0.3 | 1.5 | 3.1 | 5.8 | 10.1 | 20.8 |
| 1969 | 1.5 | 2.4 | 0.3 | * | * | * | 0.3 | 1.2 | 2.7 | 5.3 | 10.5 | 27.5 |
| 1968 | 1.7 | 3.4 | 0.5 | * | * | * | 0.3 | 1.1 | 3.3 | 6.2 | 11.2 | 27.3 |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| CHRONIC AND UNQUALIFIED BRONCHITIS (490, 491) | | | | | | | | | | | | |
| ALL OTHER, BOTH SEXES | | | | | | | | | | | | |
| 1978 | 0.8 | * | * | * | * | * | * | 1.1 | 2.5 | 4.1 | 6.4 | * |
| 1977 | 0.9 | 3.6 | * | * | * | * | * | 0.8 | 2.3 | 4.7 | 7.5 | 12.5 |
| 1976 | 1.0 | * | * | * | * | * | * | 1.5 | 3.3 | 5.4 | 7.1 | * |
| 1975 | 0.9 | 4.0 | * | * | * | * | * | * | 2.8 | 5.3 | 6.9 | * |
| 1974 | 1.2 | 5.3 | * | * | * | * | * | 1.0 | 3.7 | 7.8 | 8.4 | * |
| 1973 | 1.3 | 4.0 | * | * | * | * | 0.7 | 1.7 | 4.2 | 5.0 | 11.8 | 19.7 |
| 1972 | 1.6 | 6.7 | * | * | * | * | 0.8 | 2.2 | 4.5 | 8.8 | 9.2 | 26.0 |
| 1971 | 1.4 | 4.8 | * | * | * | * | 1.0 | 1.7 | 4.2 | 6.8 | 11.5 | 19.2 |
| 1970 | 1.5 | 7.3 | * | * | * | * | * | 1.9 | 5.7 | 6.4 | 11.9 | 14.9 |
| 1969 | 1.6 | 8.4 | 1.0 | * | * | * | * | 2.4 | 5.0 | 7.6 | 13.6 | * |
| 1968 | 1.8 | 7.4 | 1.0 | * | * | * | 1.1 | 2.6 | 6.0 | 8.4 | 13.8 | 21.6 |
| ALL OTHER, MALE | | | | | | | | | | | | |
| 1978 | 1.1 | * | * | * | * | * | * | * | 3.6 | 6.9 | 11.3 | * |
| 1977 | 1.2 | * | * | * | * | * | * | * | 3.8 | 7.1 | 13.8 | * |
| 1976 | 1.5 | * | * | * | * | * | * | 2.6 | 4.8 | 9.4 | 13.2 | * |
| 1975 | 1.4 | * | * | * | * | * | * | * | 4.9 | 9.4 | 12.8 | * |
| 1974 | 1.8 | * | * | * | * | * | * | 1.7 | 5.9 | 13.2 | 14.0 | * |
| 1973 | 2.0 | * | * | * | * | * | * | 2.2 | 6.7 | 8.8 | 22.8 | * |
| 1972 | 2.2 | * | * | * | * | * | * | 3.5 | 7.8 | 14.3 | 18.4 | * |
| 1971 | 2.0 | * | * | * | * | * | * | 2.3 | 6.8 | 10.7 | 20.8 | * |
| 1970 | 2.3 | 9.0 | * | * | * | * | * | 3.1 | 8.9 | 11.7 | 21.2 | * |
| 1969 | 2.5 | 8.5 | * | * | * | * | * | 3.7 | 8.6 | 13.2 | 26.1 | * |
| 1968 | 2.8 | 8.3 | * | * | * | * | * | 4.0 | 10.0 | 15.4 | 24.2 | * |
| ALL OTHER, FEMALE | | | | | | | | | | | | |
| 1978 | 0.6 | * | * | * | * | * | * | * | * | * | * | * |
| 1977 | 0.5 | * | * | * | * | * | * | * | * | 2.8 | * | * |
| 1976 | 0.5 | * | * | * | * | * | * | * | 1.9 | * | * | * |
| 1975 | 0.5 | * | * | * | * | * | * | * | * | * | * | * |
| 1974 | 0.6 | * | * | * | * | * | * | * | * | 3.4 | * | * |
| 1973 | 0.6 | * | * | * | * | * | * | * | 2.0 | * | * | * |
| 1972 | 1.0 | * | * | * | * | * | * | * | * | 4.3 | * | * |
| 1971 | 0.9 | * | * | * | * | * | * | * | * | 3.6 | * | * |
| 1970 | 0.8 | * | * | * | * | * | * | * | 2.9 | * | * | * |
| 1969 | 0.8 | 8.2 | * | * | * | * | * | * | * | * | * | * |
| 1968 | 1.0 | * | * | * | * | * | * | * | 2.4 | * | * | * |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| CHRONIC AND UNQUALIFIED BRONCHITIS (490, 491) | | | | | | | | | | | | |
| BLACK, BOTH SEXES | | | | | | | | | | | | |
| 1978 | 0.9 | * | * | * | * | * | * | 1.1 | 2.8 | 3.9 | 5.0 | * |
| 1977 | 0.9 | * | * | * | * | * | * | * | 2.3 | 5.2 | 7.8 | * |
| 1976 | 1.0 | * | * | * | * | * | * | 1.6 | 3.5 | 5.5 | 6.3 | * |
| 1975 | 1.0 | * | * | * | * | * | * | * | 3.1 | 5.4 | 6.4 | * |
| 1974 | 1.2 | 5.1 | * | * | * | * | * | 1.1 | 3.9 | 7.2 | 6.9 | * |
| 1973 | 1.2 | * | * | * | * | * | 0.8 | 1.9 | 4.3 | 4.9 | 9.8 | * |
| 1972 | 1.6 | 6.7 | * | * | * | * | 0.9 | 2.4 | 4.2 | 9.3 | 10.1 | 20.0 |
| 1971 | 1.5 | 5.0 | * | * | * | * | 1.1 | 1.8 | 4.2 | 7.0 | 11.9 | * |
| 1970 | 1.6 | 7.2 | * | * | * | * | * | 2.1 | 6.0 | 6.6 | 10.3 | * |
| 1969 | 1.6 | 9.0 | * | * | * | * | * | 2.4 | 5.3 | 7.5 | 12.7 | * |
| 1968 | 1.9 | 7.2 | 1.1 | * | * | * | 1.1 | 2.7 | 6.5 | 8.1 | 13.4 | * |
| BLACK, MALE | | | | | | | | | | | | |
| 1978 | 1.1 | * | * | * | * | * | * | * | 4.0 | 6.5 | 9.2 | * |
| 1977 | 1.3 | * | * | * | * | * | * | * | 3.9 | 8.0 | 14.5 | * |
| 1976 | 1.5 | * | * | * | * | * | * | 2.7 | 5.1 | 10.0 | 12.0 | * |
| 1975 | 1.5 | * | * | * | * | * | * | * | 5.5 | 9.6 | 12.6 | * |
| 1974 | 1.8 | * | * | * | * | * | * | 2.0 | 6.3 | 13.0 | 12.0 | * |
| 1973 | 1.9 | * | * | * | * | * | * | 2.5 | 7.1 | 9.0 | 17.6 | * |
| 1972 | 2.3 | * | * | * | * | * | * | 3.8 | 7.4 | 16.2 | 20.3 | * |
| 1971 | 2.1 | * | * | * | * | * | * | 2.5 | 7.0 | 11.1 | 22.0 | * |
| 1970 | 2.4 | 9.4 | * | * | * | * | * | 3.3 | 9.7 | 12.1 | 18.4 | * |
| 1969 | 2.5 | 9.2 | * | * | * | * | * | 3.9 | 9.3 | 13.2 | 25.7 | * |
| 1968 | 2.9 | 8.4 | * | * | * | * | * | 4.4 | 11.0 | 15.0 | 23.6 | * |
| BLACK, FEMALE | | | | | | | | | | | | |
| 1978 | 0.6 | * | * | * | * | * | * | * | * | * | * | * |
| 1977 | 0.6 | * | * | * | * | * | * | * | * | 3.1 | * | * |
| 1976 | 0.5 | * | * | * | * | * | * | * | 2.2 | * | * | * |
| 1975 | 0.5 | * | * | * | * | * | * | * | * | * | * | * |
| 1974 | 0.6 | * | * | * | * | * | * | * | * | * | * | * |
| 1973 | 0.7 | * | * | * | * | * | * | * | * | * | * | * |
| 1972 | 1.0 | * | * | * | * | * | * | * | * | 4.0 | * | * |
| 1971 | 0.9 | * | * | * | * | * | * | * | * | 3.7 | * | * |
| 1970 | 0.8 | * | * | * | * | * | * | * | 2.8 | * | * | * |
| 1969 | 0.8 | 8.8 | * | * | * | * | * | * | * | * | * | * |
| 1968 | 1.0 | * | * | * | * | * | * | * | 2.5 | * | * | * |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| EMPHYSEMA (492) | | | | | | | | | | | | |
| BOTH SEXES | | | | | | | | | | | | |
| 1978 | 7.0 | * | * | * | * | * | 0.5 | 3.3 | 14.1 | 38.7 | 63.0 | 60.8 |
| 1977 | 7.5 | * | * | * | * | 0.1 | 0.6 | 3.4 | 15.0 | 41.6 | 67.6 | 64.2 |
| 1976 | 8.2 | * | * | * | * | 0.1 | 0.6 | 3.8 | 16.5 | 47.0 | 73.2 | 75.6 |
| 1975 | 8.7 | * | * | * | * | 0.1 | 0.7 | 4.2 | 18.5 | 51.2 | 77.7 | 72.0 |
| 1974 | 9.3 | * | * | * | * | 0.1 | 0.8 | 4.6 | 20.3 | 55.6 | 84.3 | 76.6 |
| 1973 | 10.5 | * | * | * | * | 0.1 | 0.8 | 5.2 | 23.9 | 63.0 | 94.5 | 91.8 |
| 1972 | 10.8 | * | * | * | * | 0.1 | 0.9 | 5.6 | 25.0 | 65.2 | 96.8 | 90.7 |
| 1971 | 10.9 | * | * | * | * | 0.1 | 1.0 | 6.3 | 25.6 | 65.5 | 97.5 | 91.5 |
| 1970 | 11.2 | * | * | * | * | 0.2 | 1.2 | 6.2 | 27.2 | 68.7 | 98.6 | 84.4 |
| 1969 | 11.4 | * | * | * | 0.1 | 0.2 | 1.3 | 6.6 | 28.2 | 69.1 | 99.5 | 98.5 |
| 1968 | 12.1 | * | * | * | * | 0.2 | 1.4 | 7.1 | 30.7 | 74.0 | 105.8 | 103.5 |
| MALE | | | | | | | | | | | | |
| 1978 | 10.9 | * | * | * | * | * | 0.6 | 4.4 | 21.1 | 69.0 | 133.5 | 139.4 |
| 1977 | 11.8 | * | * | * | * | * | 0.9 | 4.7 | 23.2 | 75.4 | 144.9 | 150.9 |
| 1976 | 13.1 | * | * | * | * | * | 0.7 | 5.4 | 26.2 | 86.8 | 155.5 | 176.4 |
| 1975 | 14.2 | * | * | * | * | * | 0.9 | 6.0 | 29.4 | 96.2 | 169.2 | 165.5 |
| 1974 | 15.2 | * | * | * | * | * | 1.1 | 6.6 | 32.9 | 104.1 | 182.1 | 173.1 |
| 1973 | 17.5 | * | * | * | * | * | 1.1 | 7.7 | 39.8 | 120.8 | 204.6 | 208.0 |
| 1972 | 18.1 | * | * | * | * | 0.2 | 1.0 | 8.4 | 42.2 | 125.6 | 209.9 | 202.3 |
| 1971 | 18.4 | * | * | * | * | 0.2 | 1.4 | 9.3 | 43.8 | 127.4 | 208.7 | 199.6 |
| 1970 | 19.1 | * | * | * | * | 0.2 | 1.6 | 9.5 | 47.0 | 135.2 | 211.3 | 178.8 |
| 1969 | 19.6 | * | * | * | * | 0.2 | 1.8 | 10.3 | 49.7 | 137.8 | 210.6 | 202.6 |
| 1968 | 21.0 | * | * | * | * | 0.3 | 1.9 | 11.2 | 54.1 | 148.1 | 222.8 | 215.2 |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| EMPHYSEMA (492) | | | | | | | | | | | | |
| FEMALE | | | | | | | | | | | | |
| 1978 | 3.4 | * | * | * | * | * | 0.3 | 2.2 | 7.9 | 15.6 | 21.1 | 25.3 |
| 1977 | 3.4 | * | * | * | * | * | 0.3 | 2.1 | 7.8 | 15.8 | 21.3 | 24.3 |
| 1976 | 3.5 | * | * | * | * | * | 0.4 | 2.3 | 7.9 | 16.4 | 23.4 | 28.2 |
| 1975 | 3.6 | * | * | * | * | * | 0.5 | 2.6 | 8.9 | 16.7 | 22.0 | 26.7 |
| 1974 | 3.8 | * | * | * | * | * | 0.5 | 2.7 | 9.1 | 18.4 | 24.0 | 28.7 |
| 1973 | 3.9 | * | * | * | * | * | 0.6 | 2.8 | 9.7 | 18.6 | 25.6 | 33.3 |
| 1972 | 3.9 | * | * | * | * | * | 0.7 | 3.0 | 9.6 | 18.6 | 24.8 | 33.5 |
| 1971 | 3.8 | * | * | * | * | * | 0.6 | 3.5 | 9.2 | 17.5 | 25.4 | 35.0 |
| 1970 | 3.7 | * | * | * | * | * | 0.8 | 3.2 | 9.4 | 17.0 | 24.0 | 31.5 |
| 1969 | 3.6 | * | * | * | * | * | 0.8 | 3.2 | 8.9 | 15.4 | 24.8 | 42.1 |
| 1968 | 3.7 | * | * | * | * | * | 0.9 | 3.3 | 9.6 | 15.7 | 25.7 | 42.1 |
| WHITE, BOTH SEXES | | | | | | | | | | | | |
| 1978 | 7.7 | * | * | * | * | * | 0.4 | 3.3 | 14.5 | 40.9 | 66.7 | 63.1 |
| 1977 | 8.2 | * | * | * | * | * | 0.6 | 3.4 | 15.6 | 44.1 | 71.5 | 67.0 |
| 1976 | 9.0 | * | * | * | * | 0.1 | 0.6 | 3.8 | 17.2 | 49.6 | 77.5 | 78.8 |
| 1975 | 9.5 | * | * | * | * | 0.1 | 0.6 | 4.2 | 19.3 | 54.2 | 81.7 | 75.6 |
| 1974 | 10.2 | * | * | * | * | 0.1 | 0.7 | 4.7 | 21.3 | 58.7 | 88.9 | 79.9 |
| 1973 | 11.5 | * | * | * | * | * | 0.8 | 5.2 | 24.7 | 66.6 | 99.2 | 95.9 |
| 1972 | 11.8 | * | * | * | * | 0.1 | 0.8 | 5.7 | 26.1 | 69.2 | 101.0 | 94.3 |
| 1971 | 11.8 | * | * | * | * | 0.1 | 0.9 | 6.2 | 26.5 | 69.0 | 102.2 | 95.6 |
| 1970 | 12.1 | * | * | * | * | 0.1 | 1.1 | 6.3 | 28.2 | 72.6 | 102.4 | 88.4 |
| 1969 | 12.3 | * | * | * | * | 0.2 | 1.2 | 6.6 | 29.3 | 72.7 | 103.7 | 102.8 |
| 1968 | 13.1 | * | * | * | * | 0.2 | 1.3 | 6.9 | 31.8 | 78.1 | 109.9 | 108.6 |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| EMPHYSEMA (492) | | | | | | | | | | | | |
| WHITE, MALE | | | | | | | | | | | | |
| 1978 | 11.8 | * | * | * | * | * | 0.5 | 4.2 | 21.4 | 72.5 | 142.2 | 146.5 |
| 1977 | 12.8 | * | * | * | * | * | 0.8 | 4.5 | 23.6 | 79.5 | 154.4 | 159.4 |
| 1976 | 14.3 | * | * | * | * | * | 0.7 | 5.3 | 27.0 | 91.5 | 165.5 | 185.5 |
| 1975 | 15.4 | * | * | * | * | * | 0.8 | 5.8 | 30.3 | 102.1 | 178.8 | 175.2 |
| 1974 | 16.5 | * | * | * | * | * | 0.9 | 6.6 | 34.2 | 109.8 | 192.9 | 182.5 |
| 1973 | 19.0 | * | * | * | * | * | 1.0 | 7.6 | 40.9 | 127.8 | 216.1 | 218.8 |
| 1972 | 19.6 | * | * | * | * | * | 0.9 | 8.4 | 43.8 | 133.4 | 220.2 | 213.5 |
| 1971 | 19.9 | * | * | * | * | * | 1.3 | 9.1 | 45.2 | 134.5 | 220.3 | 209.9 |
| 1970 | 20.7 | * | * | * | * | * | 1.4 | 9.6 | 48.5 | 143.3 | 220.8 | 190.2 |
| 1969 | 21.2 | * | * | * | * | 0.2 | 1.6 | 10.1 | 51.4 | 145.2 | 220.8 | 214.0 |
| 1968 | 22.6 | * | * | * | * | 0.2 | 1.7 | 10.7 | 56.0 | 156.5 | 232.5 | 229.6 |
| WHITE, FEMALE | | | | | | | | | | | | |
| 1978 | 3.8 | * | * | * | * | * | 0.3 | 2.4 | 8.3 | 16.8 | 22.3 | 26.0 |
| 1977 | 3.7 | * | * | * | * | * | 0.3 | 2.3 | 8.4 | 17.0 | 22.5 | 25.2 |
| 1976 | 3.9 | * | * | * | * | * | 0.5 | 2.5 | 8.4 | 17.5 | 25.1 | 29.5 |
| 1975 | 3.9 | * | * | * | * | * | 0.5 | 2.6 | 9.6 | 17.7 | 23.1 | 28.2 |
| 1974 | 4.2 | * | * | * | * | * | 0.5 | 2.9 | 9.8 | 19.7 | 25.4 | 29.8 |
| 1973 | 4.3 | * | * | * | * | * | 0.6 | 2.9 | 10.2 | 19.8 | 26.9 | 35.2 |
| 1972 | 4.3 | * | * | * | * | * | 0.7 | 3.1 | 10.3 | 19.9 | 25.9 | 34.3 |
| 1971 | 4.2 | * | * | * | * | * | 0.6 | 3.5 | 9.8 | 18.5 | 26.6 | 37.1 |
| 1970 | 4.0 | * | * | * | * | * | 0.8 | 3.3 | 10.0 | 18.0 | 24.9 | 32.7 |
| 1969 | 3.9 | * | * | * | * | * | 0.7 | 3.3 | 9.3 | 16.4 | 25.8 | 43.9 |
| 1968 | 4.0 | * | * | * | * | * | 0.8 | 3.4 | 10.0 | 16.6 | 26.8 | 43.4 |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| EMPHYSEMA (492) | | | | | | | | | | | | |
| ALL OTHER, BOTH SEXES | | | | | | | | | | | | |
| 1978 | 2.8 | * | * | * | * | * | 0.8 | 3.4 | 10.6 | 18.4 | 25.2 | 34.5 |
| 1977 | 2.8 | * | * | * | * | * | 0.7 | 3.7 | 10.2 | 18.3 | 27.5 | 31.3 |
| 1976 | 3.0 | * | * | * | * | * | * | 3.5 | 10.3 | 21.7 | 28.4 | 38.6 |
| 1975 | 3.3 | * | * | * | * | * | 1.3 | 4.7 | 10.8 | 21.8 | 35.8 | 30.8 |
| 1974 | 3.3 | * | * | * | * | * | 1.4 | 3.6 | 11.2 | 25.4 | 33.3 | 37.8 |
| 1973 | 4.0 | * | * | * | * | * | 1.0 | 4.9 | 15.9 | 28.0 | 40.3 | 43.3 |
| 1972 | 4.0 | * | * | * | * | * | 1.3 | 5.0 | 14.4 | 26.3 | 47.8 | 48.8 |
| 1971 | 4.4 | * | * | * | * | * | 1.6 | 6.9 | 16.1 | 30.7 | 39.8 | 44.2 |
| 1970 | 4.6 | * | * | * | * | * | 2.0 | 5.4 | 17.4 | 30.6 | 50.8 | 43.3 |
| 1969 | 4.7 | * | * | * | * | * | 2.1 | 7.1 | 18.0 | 32.7 | 44.9 | 50.5 |
| 1968 | 5.2 | * | * | * | * | * | 2.3 | 8.9 | 19.6 | 33.3 | 52.3 | 46.1 |
| ALL OTHER, MALE | | | | | | | | | | | | |
| 1978 | 4.8 | * | * | * | * | * | 1.4 | 6.2 | 18.3 | 37.1 | 51.1 | 67.2 |
| 1977 | 4.9 | * | * | * | * | * | * | 6.6 | 18.7 | 36.5 | 55.9 | 64.3 |
| 1976 | 5.2 | * | * | * | * | * | * | 6.2 | 18.6 | 42.3 | 61.5 | 85.5 |
| 1975 | 5.6 | * | * | * | * | * | 1.8 | 7.8 | 20.6 | 40.2 | 74.0 | 68.5 |
| 1974 | 5.9 | * | * | * | * | * | 2.5 | 6.7 | 21.3 | 49.9 | 70.3 | 76.0 |
| 1973 | 7.0 | * | * | * | * | * | * | 8.7 | 28.8 | 54.3 | 83.5 | 97.9 |
| 1972 | 7.1 | * | * | * | * | * | 1.6 | 9.1 | 27.0 | 51.2 | 98.6 | 89.4 |
| 1971 | 7.7 | * | * | * | * | * | 2.4 | 11.0 | 30.4 | 59.0 | 80.2 | 97.9 |
| 1970 | 8.1 | * | * | * | * | * | 3.4 | 9.0 | 32.6 | 58.9 | 102.0 | 79.4 |
| 1969 | 8.5 | * | * | * | * | * | 3.2 | 12.5 | 33.2 | 66.1 | 88.8 | 95.5 |
| 1968 | 9.2 | * | * | * | * | * | 3.6 | 16.1 | 35.1 | 66.1 | 106.5 | 75.6 |
| ALL OTHER, FEMALE | | | | | | | | | | | | |
| 1978 | 0.9 | * | * | * | * | * | * | * | 4.2 | 3.9 | 7.5 | * |
| 1977 | 0.9 | * | * | * | * | * | * | * | 3.2 | 4.0 | 8.1 | * |
| 1976 | 0.9 | * | * | * | * | * | * | * | 3.4 | 5.5 | 5.9 | * |
| 1975 | 1.2 | * | * | * | * | * | * | 2.1 | 2.5 | 7.2 | 9.6 | * |
| 1974 | 0.9 | * | * | * | * | * | * | * | 2.5 | 5.7 | 7.8 | * |
| 1973 | 1.2 | * | * | * | * | * | * | 1.7 | 4.9 | 6.7 | 10.1 | * |
| 1972 | 1.2 | * | * | * | * | * | * | 1.5 | 3.7 | 6.1 | 11.3 | * |
| 1971 | 1.3 | * | * | * | * | * | * | 3.3 | 3.6 | 7.5 | 10.0 | * |
| 1970 | 1.4 | * | * | * | * | * | * | 2.1 | 3.9 | 7.3 | 11.8 | * |
| 1969 | 1.3 | * | * | * | * | * | * | 2.3 | 4.5 | 5.2 | 11.3 | * |
| 1968 | 1.4 | * | * | * | * | * | * | 2.5 | 5.6 | 6.1 | 10.5 | * |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| EMPHYSEMA (492) | | | | | | | | | | | | |
| BLACK, BOTH SEXES | | | | | | | | | | | | |
| 1978 | 3.0 | * | * | * | * | * | 0.9 | 3.8 | 11.5 | 18.6 | 23.5 | 36.7 |
| 1977 | 3.0 | * | * | * | * | * | 0.8 | 4.1 | 11.1 | 19.0 | 26.4 | 28.0 |
| 1976 | 3.2 | * | * | * | * | * | * | 3.9 | 11.0 | 22.6 | 27.6 | 40.4 |
| 1975 | 3.5 | * | * | * | * | * | 1.5 | 5.1 | 11.6 | 22.5 | 34.9 | 26.9 |
| 1974 | 3.5 | * | * | * | * | * | 1.6 | 3.9 | 12.0 | 26.4 | 30.7 | 32.2 |
| 1973 | 4.1 | * | * | * | * | * | 1.0 | 5.4 | 16.4 | 28.7 | 38.7 | 37.7 |
| 1972 | 4.2 | * | * | * | * | * | 1.5 | 5.6 | 15.3 | 27.4 | 45.8 | 43.6 |
| 1971 | 4.5 | * | * | * | * | * | 1.7 | 7.2 | 16.9 | 31.7 | 37.5 | 43.0 |
| 1970 | 4.8 | * | * | * | * | * | 2.2 | 5.8 | 18.7 | 31.4 | 49.4 | 40.9 |
| 1969 | 4.9 | * | * | * | * | * | 2.2 | 7.5 | 18.8 | 32.7 | 43.5 | 45.4 |
| 1968 | 5.3 | * | * | * | * | * | 2.4 | 9.2 | 20.0 | 34.0 | 50.1 | 44.6 |
| BLACK, MALE | | | | | | | | | | | | |
| 1978 | 5.1 | * | * | * | * | * | 1.7 | 6.9 | 19.8 | 38.1 | 49.0 | 72.5 |
| 1977 | 5.3 | * | * | * | * | * | * | 7.5 | 20.9 | 39.0 | 55.7 | 55.1 |
| 1976 | 5.7 | * | * | * | * | * | * | 7.0 | 20.1 | 45.2 | 60.3 | 91.5 |
| 1975 | 6.0 | * | * | * | * | * | 2.1 | 8.4 | 22.3 | 42.8 | 73.9 | 60.9 |
| 1974 | 6.2 | * | * | * | * | * | 2.9 | 7.3 | 23.2 | 53.2 | 66.0 | 67.4 |
| 1973 | 7.3 | * | * | * | * | * | * | 9.5 | 29.9 | 57.1 | 81.4 | 90.2 |
| 1972 | 7.4 | * | * | * | * | * | 1.8 | 10.1 | 29.4 | 54.2 | 95.4 | 80.0 |
| 1971 | 8.0 | * | * | * | * | * | 2.8 | 11.9 | 32.5 | 62.5 | 77.4 | 97.5 |
| 1970 | 8.5 | * | * | * | * | * | 3.7 | 9.9 | 35.7 | 61.8 | 103.2 | 73.9 |
| 1969 | 8.7 | * | * | * | * | * | 3.3 | 13.4 | 35.3 | 67.5 | 88.0 | 89.2 |
| 1968 | 9.5 | * | * | * | * | * | 3.7 | 17.2 | 36.7 | 69.1 | 102.4 | 74.3 |
| BLACK, FEMALE | | | | | | | | | | | | |
| 1978 | 1.0 | * | * | * | * | * | * | * | 4.6 | 4.0 | 6.4 | * |
| 1977 | 0.9 | * | * | * | * | * | * | * | 3.1 | 3.9 | 7.1 | * |
| 1976 | 1.0 | * | * | * | * | * | * | * | 3.5 | 5.5 | 5.7 | * |
| 1975 | 1.2 | * | * | * | * | * | * | 2.4 | 2.7 | 7.1 | 8.5 | * |
| 1974 | 1.0 | * | * | * | * | * | * | * | 2.7 | 5.9 | 6.8 | * |
| 1973 | 1.3 | * | * | * | * | * | * | 1.8 | 5.2 | 6.8 | 9.5 | * |
| 1972 | 1.3 | * | * | * | * | * | * | 1.7 | 3.5 | 6.6 | 10.7 | * |
| 1971 | 1.3 | * | * | * | * | * | * | 3.2 | 3.8 | 7.4 | 8.9 | * |
| 1970 | 1.4 | * | * | * | * | * | * | 2.2 | 4.1 | 7.2 | 10.0 | * |
| 1969 | 1.4 | * | * | * | * | * | * | 2.3 | 4.7 | 5.1 | 10.6 | * |
| 1968 | 1.4 | * | * | * | * | * | * | 2.3 | 5.6 | 5.7 | 11.3 | * |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| ASTHMA (493) | | | | | | | | | | | | |
| BOTH SEXES | | | | | | | | | | | | |
| 1978 | 0.8 | * | * | 0.1 | 0.2 | 0.3 | 0.5 | 0.9 | 1.7 | 3.0 | 4.7 | 6.2 |
| 1977 | 0.8 | * | * | 0.1 | 0.2 | 0.2 | 0.5 | 0.8 | 1.5 | 3.0 | 4.4 | 4.5 |
| 1976 | 0.9 | * | 0.2 | 0.1 | 0.2 | 0.3 | 0.6 | 1.1 | 2.2 | 3.4 | 4.6 | 5.5 |
| 1975 | 0.9 | * | * | 0.1 | 0.2 | 0.4 | 0.6 | 1.2 | 2.1 | 3.2 | 4.7 | 4.9 |
| 1974 | 0.9 | * | 0.2 | 0.1 | 0.2 | 0.3 | 0.6 | 1.1 | 2.1 | 3.2 | 4.6 | 4.6 |
| 1973 | 0.9 | * | 0.1 | 0.1 | 0.3 | 0.3 | 0.8 | 1.2 | 2.0 | 3.1 | 4.1 | 5.2 |
| 1972 | 1.1 | * | 0.2 | 0.1 | 0.3 | 0.4 | 0.9 | 1.5 | 2.7 | 3.8 | 5.3 | 5.7 |
| 1971 | 1.0 | * | 0.2 | 0.2 | 0.3 | 0.5 | 0.9 | 1.5 | 2.4 | 3.7 | 4.3 | 5.9 |
| 1970 | 1.1 | * | 0.3 | 0.2 | 0.4 | 0.5 | 1.0 | 1.7 | 2.7 | 3.5 | 4.8 | 5.0 |
| 1969 | 1.2 | * | 0.3 | 0.2 | 0.3 | 0.6 | 1.0 | 1.6 | 2.9 | 3.8 | 5.2 | 5.9 |
| 1968 | 1.3 | * | 0.3 | 0.2 | 0.4 | 0.7 | 1.1 | 1.9 | 3.2 | 4.4 | 5.9 | 7.8 |
| MALE | | | | | | | | | | | | |
| 1978 | 0.7 | * | * | 0.1 | 0.2 | 0.2 | 0.3 | 0.7 | 1.3 | 3.0 | 4.9 | 6.1 |
| 1977 | 0.7 | * | * | 0.1 | 0.2 | 0.2 | 0.3 | 0.7 | 1.4 | 3.1 | 4.5 | 5.7 |
| 1976 | 0.8 | * | * | 0.1 | 0.2 | 0.2 | 0.4 | 0.7 | 1.9 | 3.7 | 5.0 | 6.1 |
| 1975 | 0.8 | * | * | * | 0.2 | 0.4 | 0.5 | 0.8 | 2.0 | 3.6 | 4.9 | 4.5 |
| 1974 | 0.8 | * | * | * | 0.2 | 0.2 | 0.4 | 0.9 | 2.0 | 3.4 | 5.5 | 5.5 |
| 1973 | 0.8 | * | * | 0.1 | 0.3 | 0.3 | 0.4 | 1.1 | 1.8 | 3.4 | 4.6 | 5.0 |
| 1972 | 1.0 | * | * | 0.2 | 0.3 | 0.3 | 0.6 | 1.2 | 2.5 | 4.2 | 6.0 | 5.4 |
| 1971 | 0.9 | * | * | 0.1 | 0.2 | 0.3 | 0.6 | 1.3 | 2.1 | 3.9 | 4.7 | 7.6 |
| 1970 | 1.0 | * | 0.4 | 0.2 | 0.3 | 0.3 | 0.8 | 1.3 | 2.5 | 3.7 | 5.6 | 6.6 |
| 1969 | 1.1 | * | * | 0.2 | 0.3 | 0.4 | 0.7 | 1.2 | 2.9 | 4.4 | 6.8 | 6.1 |
| 1968 | 1.3 | * | 0.4 | 0.2 | 0.3 | 0.5 | 0.8 | 1.8 | 3.2 | 5.3 | 6.8 | 8.6 |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| ASTHMA (493) | | | | | | | | | | | | |
| FEMALE | | | | | | | | | | | | |
| 1978 | 1.0 | * | * | * | 0.3 | 0.3 | 0.6 | 1.1 | 2.2 | 3.0 | 4.5 | 6.2 |
| 1977 | 0.8 | * | * | * | 0.2 | 0.2 | 0.7 | 0.9 | 1.5 | 3.0 | 4.4 | 4.0 |
| 1976 | 1.0 | * | * | * | 0.2 | 0.3 | 0.8 | 1.4 | 2.4 | 3.1 | 4.4 | 5.3 |
| 1975 | 1.0 | * | * | * | 0.2 | 0.4 | 0.8 | 1.5 | 2.2 | 2.9 | 4.6 | 5.1 |
| 1974 | 1.0 | * | * | 0.1 | 0.2 | 0.3 | 0.8 | 1.3 | 2.2 | 3.0 | 4.1 | 4.2 |
| 1973 | 1.0 | * | * | 0.1 | 0.3 | 0.4 | 1.1 | 1.3 | 2.2 | 2.8 | 3.9 | 5.2 |
| 1972 | 1.2 | * | * | 0.1 | 0.2 | 0.6 | 1.2 | 1.7 | 2.9 | 3.6 | 4.8 | 5.9 |
| 1971 | 1.2 | * | * | 0.2 | 0.3 | 0.6 | 1.2 | 1.8 | 2.6 | 3.4 | 4.0 | 5.0 |
| 1970 | 1.3 | * | * | 0.2 | 0.4 | 0.7 | 1.3 | 2.0 | 2.8 | 3.3 | 4.3 | 4.1 |
| 1969 | 1.2 | * | * | 0.1 | 0.3 | 0.7 | 1.2 | 2.0 | 2.9 | 3.3 | 4.1 | 5.8 |
| 1968 | 1.4 | * | 0.3 | 0.2 | 0.5 | 0.9 | 1.3 | 2.1 | 3.1 | 3.7 | 5.3 | 7.4 |
| WHITE, BOTH SEXES | | | | | | | | | | | | |
| 1978 | 0.8 | * | * | 0.1 | 0.2 | 0.2 | 0.3 | 0.7 | 1.6 | 2.9 | 4.5 | 6.2 |
| 1977 | 0.7 | * | * | 0.1 | 0.1 | 0.2 | 0.4 | 0.6 | 1.3 | 2.9 | 4.2 | 4.6 |
| 1976 | 0.8 | * | * | 0.1 | 0.1 | 0.2 | 0.4 | 0.9 | 1.9 | 3.2 | 4.5 | 5.3 |
| 1975 | 0.8 | * | * | * | 0.2 | 0.2 | 0.4 | 0.9 | 1.8 | 3.0 | 4.7 | 4.8 |
| 1974 | 0.8 | * | * | 0.1 | 0.2 | 0.2 | 0.4 | 0.9 | 1.9 | 3.0 | 4.7 | 4.7 |
| 1973 | 0.8 | * | * | 0.1 | 0.2 | 0.2 | 0.5 | 0.9 | 1.8 | 2.9 | 4.1 | 5.1 |
| 1972 | 1.0 | * | * | 0.1 | 0.2 | 0.3 | 0.7 | 1.2 | 2.4 | 3.7 | 5.2 | 6.1 |
| 1971 | 0.9 | * | * | 0.1 | 0.2 | 0.3 | 0.6 | 1.2 | 2.1 | 3.4 | 4.1 | 6.1 |
| 1970 | 1.0 | * | 0.2 | 0.1 | 0.2 | 0.3 | 0.7 | 1.3 | 2.3 | 3.3 | 4.7 | 5.3 |
| 1969 | 1.0 | * | * | 0.1 | 0.2 | 0.4 | 0.7 | 1.3 | 2.4 | 3.5 | 5.1 | 6.0 |
| 1968 | 1.1 | * | 0.3 | 0.1 | 0.3 | 0.5 | 0.7 | 1.4 | 2.7 | 4.2 | 5.6 | 7.9 |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| ASTHMA (493) | | | | | | | | | | | | |
| WHITE, MALE | | | | | | | | | | | | |
| 1978 | 0.6 | * | * | 0.1 | 0.2 | 0.2 | * | 0.5 | 1.1 | 2.7 | 4.7 | 6.2 |
| 1977 | 0.6 | * | * | * | 0.1 | 0.2 | 0.3 | 0.5 | 1.2 | 2.9 | 4.4 | 5.8 |
| 1976 | 0.7 | * | * | * | * | 0.2 | 0.3 | 0.6 | 1.5 | 3.4 | 5.0 | 5.6 |
| 1975 | 0.7 | * | * | * | 0.2 | 0.2 | 0.3 | 0.7 | 1.7 | 3.3 | 4.8 | 4.1 |
| 1974 | 0.7 | * | * | * | 0.1 | 0.2 | 0.3 | 0.7 | 1.7 | 3.1 | 5.6 | 5.4 |
| 1973 | 0.7 | * | * | * | 0.2 | 0.2 | 0.4 | 0.8 | 1.5 | 3.0 | 4.6 | 5.1 |
| 1972 | 0.9 | * | * | 0.1 | 0.2 | 0.3 | 0.4 | 0.9 | 2.2 | 3.7 | 5.7 | 5.5 |
| 1971 | 0.8 | * | * | * | 0.2 | 0.2 | 0.4 | 1.0 | 1.8 | 3.6 | 4.6 | 7.6 |
| 1970 | 0.8 | * | * | 0.1 | 0.2 | 0.2 | 0.5 | 1.1 | 2.1 | 3.4 | 5.4 | 7.2 |
| 1969 | 0.9 | * | * | 0.1 | 0.2 | 0.3 | 0.5 | 0.8 | 2.4 | 4.0 | 6.7 | 6.5 |
| 1968 | 1.1 | * | 0.3 | 0.2 | 0.2 | 0.3 | 0.5 | 1.2 | 2.6 | 5.0 | 6.2 | 8.8 |
| WHITE, FEMALE | | | | | | | | | | | | |
| 1978 | 0.9 | * | * | * | 0.2 | 0.2 | 0.5 | 0.9 | 2.0 | 3.1 | 4.4 | 6.2 |
| 1977 | 0.8 | * | * | * | * | 0.2 | 0.5 | 0.7 | 1.3 | 2.8 | 4.2 | 4.0 |
| 1976 | 1.0 | * | * | * | 0.1 | 0.2 | 0.6 | 1.1 | 2.2 | 3.0 | 4.3 | 5.1 |
| 1975 | 0.9 | * | * | * | 0.2 | 0.3 | 0.6 | 1.1 | 1.9 | 2.8 | 4.7 | 5.2 |
| 1974 | 0.9 | * | * | * | 0.2 | 0.2 | 0.5 | 1.1 | 2.0 | 3.0 | 4.1 | 4.4 |
| 1973 | 0.9 | * | * | * | 0.2 | 0.2 | 0.7 | 1.1 | 2.0 | 2.7 | 3.8 | 5.2 |
| 1972 | 1.1 | * | * | * | 0.2 | 0.4 | 0.9 | 1.4 | 2.6 | 3.6 | 4.9 | 6.4 |
| 1971 | 1.0 | * | * | 0.1 | 0.2 | 0.4 | 0.8 | 1.4 | 2.3 | 3.3 | 3.9 | 5.3 |
| 1970 | 1.1 | * | * | 0.2 | 0.3 | 0.4 | 0.9 | 1.6 | 2.5 | 3.1 | 4.3 | 4.3 |
| 1969 | 1.1 | * | * | * | 0.2 | 0.5 | 0.9 | 1.6 | 2.5 | 3.0 | 3.9 | 5.7 |
| 1968 | 1.2 | * | * | 0.1 | 0.4 | 0.6 | 0.9 | 1.6 | 2.7 | 3.5 | 5.2 | 7.4 |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| ASTHMA (493) | | | | | | | | | | | | |
| ALL OTHER, BOTH SEXES | | | | | | | | | | | | |
| 1978 | 1.3 | * | * | * | 0.5 | 0.9 | 1.4 | 2.4 | 3.4 | 3.8 | 6.6 | * |
| 1977 | 1.2 | * | * | * | 0.5 | 0.5 | 1.5 | 2.4 | 3.2 | 4.6 | 6.4 | * |
| 1976 | 1.5 | * | * | * | 0.5 | 0.9 | 1.5 | 2.9 | 4.9 | 5.1 | 5.4 | * |
| 1975 | 1.6 | * | * | 0.3 | 0.4 | 1.3 | 1.8 | 3.5 | 5.0 | 4.7 | 4.5 | * |
| 1974 | 1.5 | * | * | * | 0.7 | 0.8 | 2.1 | 3.0 | 4.4 | 4.6 | 3.6 | * |
| 1973 | 1.7 | * | * | 0.4 | 0.8 | 1.1 | 2.3 | 3.8 | 4.4 | 5.4 | 4.4 | * |
| 1972 | 1.9 | * | * | 0.5 | 0.8 | 1.0 | 2.6 | 4.0 | 5.8 | 5.4 | 6.2 | * |
| 1971 | 2.0 | * | * | 0.3 | 0.6 | 1.5 | 2.9 | 4.6 | 5.4 | 5.7 | 6.2 | * |
| 1970 | 2.3 | * | * | 0.6 | 1.2 | 1.9 | 3.3 | 4.7 | 6.3 | 5.7 | 6.0 | * |
| 1969 | 2.5 | * | * | 0.6 | 1.3 | 1.9 | 3.0 | 4.9 | 7.2 | 6.8 | 6.5 | * |
| 1968 | 2.8 | * | * | 0.5 | 1.2 | 2.5 | 4.1 | 6.3 | 7.7 | 6.9 | 9.2 | * |
| ALL OTHER, MALE | | | | | | | | | | | | |
| 1978 | 1.3 | * | * | * | * | * | * | 2.4 | 3.0 | 5.4 | * | * |
| 1977 | 1.2 | * | * | * | * | * | * | 2.6 | 3.1 | 5.2 | * | * |
| 1976 | 1.4 | * | * | * | * | * | * | 2.3 | 5.3 | 6.9 | * | * |
| 1975 | 1.5 | * | * | * | * | 1.7 | 1.5 | 2.2 | 4.8 | 6.3 | * | * |
| 1974 | 1.4 | * | * | * | 0.8 | * | * | 2.7 | 4.9 | 6.1 | * | * |
| 1973 | 1.6 | * | * | * | 0.9 | * | * | 4.1 | 5.0 | 7.3 | * | * |
| 1972 | 1.9 | * | * | * | 1.0 | * | 2.2 | 3.5 | 5.4 | 8.7 | 9.2 | * |
| 1971 | 1.8 | * | * | * | * | * | 2.0 | 4.5 | 5.0 | 7.0 | * | * |
| 1970 | 2.1 | * | * | * | 1.1 | * | 3.2 | 3.4 | 7.1 | 6.3 | * | * |
| 1969 | 2.4 | * | * | 0.7 | 1.3 | * | 2.5 | 4.8 | 7.7 | 8.1 | * | * |
| 1968 | 2.8 | * | * | * | 1.0 | 1.8 | 3.8 | 6.6 | 9.0 | 8.2 | 13.4 | * |
| ALL OTHER, FEMALE | | | | | | | | | | | | |
| 1978 | 1.3 | * | * | * | * | 1.1 | 1.5 | 2.5 | 3.6 | 2.6 | 6.2 | * |
| 1977 | 1.3 | * | * | * | * | * | 2.1 | 2.2 | 3.3 | 4.2 | 7.0 | * |
| 1976 | 1.5 | * | * | * | * | 1.0 | 1.7 | 3.5 | 4.6 | 3.7 | 5.3 | * |
| 1975 | 1.6 | * | * | * | * | 1.0 | 2.1 | 4.5 | 5.1 | 3.5 | * | * |
| 1974 | 1.5 | * | * | * | * | * | 2.7 | 3.3 | 3.9 | 3.4 | * | * |
| 1973 | 1.7 | * | * | * | * | 1.4 | 3.5 | 3.5 | 3.8 | 3.9 | * | * |
| 1972 | 1.9 | * | * | * | * | 1.4 | 2.9 | 4.4 | 6.1 | * | * | * |
| 1971 | 2.1 | * | * | * | 0.8 | 1.8 | 3.7 | 4.6 | 5.7 | 4.5 | * | * |
| 1970 | 2.4 | * | * | * | 1.3 | 2.5 | 3.4 | 5.9 | 5.6 | 5.2 | * | * |
| 1969 | 2.5 | * | * | * | 1.3 | 2.4 | 3.5 | 5.0 | 6.7 | 5.7 | * | * |
| 1968 | 2.8 | * | * | * | 1.4 | 3.1 | 4.4 | 6.0 | 6.5 | 5.8 | * | * |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| ASTHMA (493) | | | | | | | | | | | | |
| BLACK, BOTH SEXES | | | | | | | | | | | | |
| 1978 | 1.4 | * | * | * | 0.5 | 1.0 | 1.5 | 2.8 | 3.6 | 3.7 | 6.4 | * |
| 1977 | 1.3 | * | * | * | 0.6 | 0.6 | 1.8 | 2.7 | 3.5 | 4.6 | 5.8 | * |
| 1976 | 1.6 | * | * | * | 0.5 | 1.0 | 1.7 | 3.3 | 5.2 | 5.3 | 5.1 | * |
| 1975 | 1.7 | * | * | * | 0.5 | 1.5 | 2.1 | 3.8 | 5.2 | 4.9 | 4.0 | * |
| 1974 | 1.6 | * | * | * | 0.8 | 1.0 | 2.5 | 3.2 | 4.4 | 4.5 | 3.9 | * |
| 1973 | 1.8 | * | * | 0.4 | 0.8 | 1.2 | 2.6 | 4.1 | 4.6 | 5.5 | 4.4 | * |
| 1972 | 2.0 | * | * | 0.5 | 0.8 | 1.1 | 2.8 | 4.4 | 6.1 | 5.2 | 6.3 | * |
| 1971 | 2.0 | * | * | * | 0.7 | 1.7 | 3.2 | 4.6 | 5.6 | 5.4 | 6.3 | * |
| 1970 | 2.5 | * | * | 0.6 | 1.2 | 2.1 | 3.7 | 5.2 | 6.7 | 5.8 | 6.3 | * |
| 1969 | 2.6 | * | * | 0.6 | 1.4 | 2.1 | 3.2 | 5.3 | 7.5 | 6.9 | 6.6 | * |
| 1968 | 2.9 | * | * | 0.5 | 1.3 | 2.6 | 4.3 | 6.4 | 8.3 | 6.8 | 9.3 | * |
| BLACK, MALE | | | | | | | | | | | | |
| 1978 | 1.4 | * | * | * | * | * | * | 2.6 | 3.4 | 5.4 | * | * |
| 1977 | 1.3 | * | * | * | * | * | * | 2.9 | 3.3 | 4.9 | * | * |
| 1976 | 1.5 | * | * | * | * | * | * | 2.6 | 5.6 | 7.3 | * | * |
| 1975 | 1.6 | * | * | * | * | 2.0 | 1.8 | 2.3 | 5.2 | 7.0 | * | * |
| 1974 | 1.5 | * | * | * | 0.8 | * | * | 3.0 | 5.0 | 5.9 | * | * |
| 1973 | 1.8 | * | * | * | 1.0 | * | * | 4.5 | 5.2 | 7.7 | * | * |
| 1972 | 2.0 | * | * | * | 1.1 | * | 2.2 | 4.0 | 5.8 | 8.5 | * | * |
| 1971 | 1.9 | * | * | * | * | * | 2.2 | 4.5 | 5.2 | 6.6 | * | * |
| 1970 | 2.3 | * | * | * | 1.1 | * | 3.7 | 3.8 | 7.6 | 6.5 | * | * |
| 1969 | 2.5 | * | * | 0.8 | 1.3 | * | 2.5 | 5.4 | 8.4 | 7.9 | * | * |
| 1968 | 2.9 | * | * | * | 1.1 | 1.9 | 4.0 | 6.7 | 9.8 | 8.2 | 13.9 | * |
| BLACK, FEMALE | | | | | | | | | | | | |
| 1978 | 1.4 | * | * | * | * | 1.3 | 1.5 | 2.9 | 3.8 | * | 5.9 | * |
| 1977 | 1.4 | * | * | * | * | * | 2.5 | 2.5 | 3.6 | 4.3 | 5.9 | * |
| 1976 | 1.7 | * | * | * | * | 1.2 | 2.1 | 3.9 | 4.8 | 3.8 | * | * |
| 1975 | 1.7 | * | * | * | * | 1.1 | 2.3 | 5.0 | 5.2 | 3.3 | * | * |
| 1974 | 1.6 | * | * | * | * | * | 3.1 | 3.4 | 3.9 | 3.4 | * | * |
| 1973 | 1.8 | * | * | * | * | 1.5 | 3.9 | 3.8 | 4.1 | 3.9 | * | * |
| 1972 | 2.0 | * | * | * | * | 1.5 | 3.3 | 4.7 | 6.4 | * | * | * |
| 1971 | 2.2 | * | * | * | * | 1.9 | 4.1 | 4.7 | 5.9 | 4.4 | * | * |
| 1970 | 2.6 | * | * | * | 1.3 | 2.8 | 3.7 | 6.4 | 5.9 | 5.3 | * | * |
| 1969 | 2.6 | * | * | * | 1.5 | 2.5 | 3.7 | 5.3 | 6.8 | 6.1 | * | * |
| 1968 | 2.9 | * | * | * | 1.4 | 3.2 | 4.6 | 6.2 | 7.0 | 5.7 | * | * |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| PEPTIC ULCER (531-533) | | | | | | | | | | | | |
| BOTH SEXES | | | | | | | | | | | | |
| 1978 | 2.5 | 0.7 | * | * | 0.1 | 0.2 | 0.6 | 1.7 | 4.3 | 9.6 | 21.6 | 43.9 |
| 1977 | 2.7 | * | * | * | 0.1 | 0.2 | 0.7 | 1.9 | 4.9 | 10.4 | 22.8 | 48.3 |
| 1976 | 3.0 | 0.8 | * | * | * | 0.3 | 0.9 | 2.5 | 5.5 | 12.1 | 24.2 | 48.3 |
| 1975 | 3.1 | 0.9 | * | * | 0.1 | 0.3 | 0.9 | 2.5 | 6.1 | 12.8 | 26.3 | 50.9 |
| 1974 | 3.3 | 0.9 | * | * | 0.1 | 0.3 | 1.1 | 2.7 | 7.0 | 13.6 | 28.1 | 51.2 |
| 1973 | 3.6 | 1.2 | * | * | 0.1 | 0.4 | 1.2 | 3.3 | 7.6 | 15.7 | 29.9 | 55.0 |
| 1972 | 3.8 | 1.8 | * | * | 0.1 | 0.4 | 1.3 | 3.6 | 8.0 | 16.4 | 30.6 | 55.8 |
| 1971 | 3.9 | 1.5 | * | * | 0.1 | 0.4 | 1.4 | 3.9 | 8.4 | 16.7 | 32.7 | 58.0 |
| 1970 | 4.2 | 1.7 | * | * | 0.1 | 0.5 | 1.4 | 4.0 | 9.6 | 18.0 | 35.5 | 59.6 |
| 1969 | 4.6 | 1.1 | * | * | 0.1 | 0.5 | 1.8 | 4.2 | 10.2 | 20.1 | 41.2 | 64.8 |
| 1968 | 4.7 | 1.7 | * | * | 0.1 | 0.5 | 1.7 | 4.5 | 10.8 | 21.5 | 39.5 | 69.0 |
| MALE | | | | | | | | | | | | |
| 1978 | 3.0 | * | * | * | * | 0.3 | 0.8 | 2.5 | 6.3 | 13.9 | 30.9 | 58.6 |
| 1977 | 3.3 | * | * | * | * | 0.4 | 0.9 | 2.8 | 7.3 | 16.0 | 33.0 | 61.9 |
| 1976 | 3.8 | * | * | * | * | 0.4 | 1.2 | 3.6 | 8.1 | 18.8 | 37.2 | 67.8 |
| 1975 | 4.0 | * | * | * | 0.1 | 0.3 | 1.3 | 3.6 | 8.8 | 20.7 | 39.6 | 72.7 |
| 1974 | 4.3 | 1.4 | * | * | 0.1 | 0.5 | 1.5 | 4.1 | 10.3 | 21.0 | 42.7 | 73.1 |
| 1973 | 4.8 | 1.6 | * | * | 0.1 | 0.4 | 1.7 | 4.7 | 11.7 | 25.4 | 46.4 | 75.3 |
| 1972 | 5.1 | 2.2 | * | * | 0.2 | 0.5 | 2.0 | 5.5 | 12.0 | 26.0 | 47.4 | 83.9 |
| 1971 | 5.3 | 1.9 | * | * | 0.2 | 0.6 | 2.1 | 5.7 | 13.0 | 26.4 | 49.6 | 84.5 |
| 1970 | 5.8 | 1.6 | * | * | 0.2 | 0.7 | 2.0 | 6.0 | 14.8 | 28.7 | 54.9 | 81.7 |
| 1969 | 6.4 | 1.7 | * | * | 0.2 | 0.7 | 2.5 | 6.6 | 16.0 | 31.9 | 63.0 | 87.8 |
| 1968 | 6.6 | 1.9 | * | * | 0.2 | 0.8 | 2.4 | 6.8 | 16.9 | 35.0 | 61.3 | 96.4 |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| PEPTIC ULCER (531-533) | | | | | | | | | | | | |
| FEMALE | | | | | | | | | | | | |
| 1978 | 2.0 | * | * | * | * | * | 0.4 | 1.1 | 2.4 | 6.3 | 16.1 | 37.3 |
| 1977 | 2.1 | * | * | * | * | 0.1 | 0.5 | 1.0 | 2.7 | 6.2 | 16.8 | 42.1 |
| 1976 | 2.2 | * | * | * | * | 0.2 | 0.5 | 1.5 | 3.2 | 7.0 | 16.3 | 39.1 |
| 1975 | 2.3 | * | * | * | * | 0.2 | 0.5 | 1.6 | 3.7 | 6.8 | 18.2 | 40.3 |
| 1974 | 2.4 | * | * | * | * | 0.2 | 0.6 | 1.4 | 4.1 | 8.0 | 19.0 | 40.4 |
| 1973 | 2.5 | * | * | * | * | 0.3 | 0.8 | 1.9 | 3.9 | 8.3 | 19.6 | 44.8 |
| 1972 | 2.5 | 1.2 | * | * | * | 0.3 | 0.6 | 1.9 | 4.4 | 8.9 | 19.9 | 41.4 |
| 1971 | 2.6 | * | * | * | * | 0.3 | 0.8 | 2.2 | 4.3 | 9.1 | 21.8 | 44.1 |
| 1970 | 2.8 | 1.9 | * | * | 0.1 | 0.3 | 0.9 | 2.2 | 5.0 | 9.7 | 22.6 | 47.3 |
| 1969 | 3.0 | * | * | * | * | 0.3 | 1.1 | 2.0 | 5.0 | 10.9 | 26.5 | 52.4 |
| 1968 | 3.0 | 1.4 | * | * | * | 0.3 | 1.0 | 2.4 | 5.2 | 10.8 | 24.6 | 54.0 |
| WHITE, BOTH SEXES | | | | | | | | | | | | |
| 1978 | 2.6 | * | * | * | 0.1 | 0.2 | 0.6 | 1.7 | 4.1 | 9.6 | 22.2 | 45.5 |
| 1977 | 2.8 | * | * | * | * | 0.2 | 0.6 | 1.8 | 4.7 | 10.5 | 23.4 | 50.0 |
| 1976 | 3.1 | * | * | * | * | 0.2 | 0.7 | 2.3 | 5.5 | 12.0 | 24.8 | 49.7 |
| 1975 | 3.2 | 0.8 | * | * | 0.1 | 0.2 | 0.7 | 2.4 | 6.0 | 12.8 | 26.9 | 52.5 |
| 1974 | 3.4 | * | * | * | 0.1 | 0.3 | 0.9 | 2.5 | 6.9 | 13.5 | 29.2 | 52.8 |
| 1973 | 3.8 | * | * | * | 0.1 | 0.3 | 1.1 | 3.1 | 7.5 | 15.8 | 30.6 | 56.8 |
| 1972 | 3.9 | 1.4 | * | * | 0.1 | 0.3 | 1.0 | 3.3 | 7.8 | 16.4 | 31.3 | 58.9 |
| 1971 | 4.1 | 0.9 | * | * | 0.1 | 0.3 | 1.2 | 3.6 | 8.3 | 16.9 | 33.6 | 61.4 |
| 1970 | 4.4 | 1.2 | * | * | 0.1 | 0.4 | 1.2 | 3.8 | 9.6 | 18.2 | 36.5 | 63.1 |
| 1969 | 4.8 | * | * | * | 0.1 | 0.5 | 1.6 | 4.1 | 10.1 | 20.2 | 41.8 | 68.1 |
| 1968 | 4.9 | 1.1 | * | * | 0.1 | 0.4 | 1.4 | 4.3 | 10.6 | 21.7 | 40.7 | 71.4 |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| PEPTIC ULCER (531-533) | | | | | | | | | | | | |
| WHITE, MALE | | | | | | | | | | | | |
| 1978 | 3.1 | * | * | * | * | 0.2 | 0.8 | 2.3 | 6.1 | 13.8 | 31.6 | 61.6 |
| 1977 | 3.4 | * | * | * | * | 0.3 | 0.8 | 2.7 | 7.0 | 16.2 | 34.1 | 64.6 |
| 1976 | 3.8 | * | * | * | * | 0.2 | 1.0 | 3.3 | 8.1 | 18.7 | 38.2 | 69.3 |
| 1975 | 4.1 | * | * | * | * | 0.3 | 1.1 | 3.3 | 8.6 | 20.8 | 40.6 | 75.7 |
| 1974 | 4.4 | * | * | * | * | 0.4 | 1.3 | 3.8 | 10.1 | 20.8 | 44.5 | 75.5 |
| 1973 | 4.9 | * | * | * | * | 0.3 | 1.4 | 4.4 | 11.5 | 25.5 | 47.3 | 78.8 |
| 1972 | 5.2 | 1.7 | * | * | 0.2 | 0.4 | 1.7 | 5.0 | 11.5 | 25.8 | 48.6 | 89.3 |
| 1971 | 5.4 | * | * | * | 0.2 | 0.4 | 1.7 | 5.2 | 12.9 | 27.0 | 51.2 | 89.8 |
| 1970 | 5.9 | * | * | * | 0.1 | 0.5 | 1.7 | 5.6 | 14.9 | 28.9 | 56.2 | 88.3 |
| 1969 | 6.5 | * | * | * | 0.2 | 0.7 | 2.2 | 6.3 | 15.8 | 32.4 | 63.8 | 94.7 |
| 1968 | 6.8 | * | * | * | * | 0.6 | 2.0 | 6.4 | 16.7 | 35.3 | 63.7 | 102.3 |
| WHITE, FEMALE | | | | | | | | | | | | |
| 1978 | 2.2 | * | * | * | * | * | 0.4 | 1.0 | 2.3 | 6.4 | 16.7 | 38.3 |
| 1977 | 2.2 | * | * | * | * | * | 0.5 | 0.9 | 2.6 | 6.1 | 17.1 | 43.4 |
| 1976 | 2.3 | * | * | * | * | 0.1 | 0.4 | 1.4 | 3.2 | 6.9 | 16.8 | 40.6 |
| 1975 | 2.4 | * | * | * | * | 0.2 | 0.4 | 1.5 | 3.7 | 6.6 | 18.7 | 41.5 |
| 1974 | 2.5 | * | * | * | * | 0.2 | 0.5 | 1.4 | 4.1 | 7.9 | 19.9 | 41.8 |
| 1973 | 2.7 | * | * | * | * | 0.3 | 0.8 | 1.8 | 4.0 | 8.4 | 20.3 | 46.0 |
| 1972 | 2.7 | * | * | * | * | 0.2 | 0.4 | 1.7 | 4.5 | 9.2 | 20.5 | 43.6 |
| 1971 | 2.7 | * | * | * | * | 0.2 | 0.7 | 2.0 | 4.1 | 9.1 | 22.3 | 46.8 |
| 1970 | 2.9 | 1.4 | * | * | * | 0.2 | 0.8 | 2.1 | 4.8 | 10.0 | 23.6 | 49.3 |
| 1969 | 3.1 | * | * | * | * | 0.3 | 1.0 | 1.9 | 4.9 | 10.8 | 27.1 | 54.2 |
| 1968 | 3.1 | * | * | * | * | 0.2 | 0.8 | 2.4 | 5.1 | 11.0 | 25.2 | 54.8 |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| PEPTIC ULCER (531-533) | | | | | | | | | | | | |
| ALL OTHER, BOTH SEXES | | | | | | | | | | | | |
| 1978 | 1.8 | * | * | * | * | * | 0.9 | 2.4 | 5.9 | 9.6 | 15.9 | 25.6 |
| 1977 | 1.9 | * | * | * | * | 0.5 | 1.4 | 2.5 | 6.5 | 10.1 | 16.5 | 29.4 |
| 1976 | 2.3 | * | * | * | * | 0.9 | 1.8 | 4.2 | 5.7 | 12.6 | 18.0 | 32.0 |
| 1975 | 2.4 | * | * | * | * | * | 1.9 | 4.0 | 7.4 | 13.0 | 19.3 | 32.2 |
| 1974 | 2.4 | * | * | * | * | 0.6 | 2.0 | 4.2 | 7.6 | 14.8 | 15.2 | 32.6 |
| 1973 | 2.7 | * | * | * | * | 0.9 | 2.5 | 4.6 | 8.0 | 15.1 | 21.9 | 33.9 |
| 1972 | 3.1 | 3.7 | * | * | * | 0.9 | 3.2 | 6.4 | 9.9 | 16.0 | 21.6 | 19.5 |
| 1971 | 3.0 | 4.9 | * | * | * | 1.2 | 2.9 | 6.4 | 9.2 | 14.3 | 22.2 | 19.2 |
| 1970 | 3.1 | 4.5 | * | * | * | 1.1 | 2.9 | 6.0 | 10.3 | 16.0 | 23.1 | 23.9 |
| 1969 | 3.5 | 4.3 | * | * | * | 1.0 | 3.3 | 5.6 | 11.2 | 18.5 | 34.1 | 27.5 |
| 1968 | 3.8 | 4.6 | * | * | * | 1.4 | 4.2 | 6.6 | 12.7 | 19.7 | 23.5 | 42.2 |
| ALL OTHER, MALE | | | | | | | | | | | | |
| 1978 | 2.4 | * | * | * | * | * | 1.5 | 4.0 | 8.8 | 14.5 | 24.4 | * |
| 1977 | 2.6 | * | * | * | * | * | 2.1 | 4.0 | 9.9 | 14.8 | 21.8 | * |
| 1976 | 3.3 | * | * | * | * | 1.5 | 2.8 | 6.5 | 8.8 | 19.2 | 28.0 | 52.7 |
| 1975 | 3.3 | * | * | * | * | * | 3.1 | 5.9 | 11.3 | 19.3 | 29.3 | 42.6 |
| 1974 | 3.6 | * | * | * | * | * | 3.1 | 6.8 | 12.4 | 22.4 | 24.9 | 48.0 |
| 1973 | 4.1 | * | * | * | * | 1.3 | 4.3 | 7.1 | 13.1 | 25.1 | 36.6 | * |
| 1972 | 4.7 | * | * | * | * | * | 4.8 | 9.8 | 16.8 | 27.5 | 34.1 | * |
| 1971 | 4.3 | * | * | * | * | 2.0 | 4.7 | 10.0 | 13.5 | 21.3 | 32.9 | * |
| 1970 | 4.6 | * | * | * | * | 1.7 | 4.7 | 9.4 | 14.1 | 26.8 | 40.4 | * |
| 1969 | 5.0 | * | * | * | * | * | 4.5 | 8.9 | 17.6 | 27.0 | 53.7 | * |
| 1968 | 5.5 | * | * | * | * | 2.4 | 6.1 | 10.7 | 19.3 | 32.3 | 33.3 | * |
| ALL OTHER, FEMALE | | | | | | | | | | | | |
| 1978 | 1.2 | * | * | * | * | * | * | * | 3.5 | 5.9 | 10.0 | 24.8 |
| 1977 | 1.4 | * | * | * | * | * | * | * | 3.8 | 6.5 | 12.8 | 26.9 |
| 1976 | 1.4 | * | * | * | * | * | * | 2.3 | 3.1 | 7.5 | 11.2 | 20.4 |
| 1975 | 1.6 | * | * | * | * | * | * | 2.4 | 4.1 | 8.0 | 12.4 | 25.8 |
| 1974 | 1.4 | * | * | * | * | * | * | 2.0 | 3.6 | 8.7 | 8.4 | 23.5 |
| 1973 | 1.5 | * | * | * | * | * | * | 2.5 | 3.7 | 7.0 | 11.6 | 30.4 |
| 1972 | 1.6 | * | * | * | * | * | 1.9 | 3.5 | 3.9 | 6.7 | 12.6 | * |
| 1971 | 1.8 | * | * | * | * | * | 1.4 | 3.3 | 5.5 | 8.6 | 14.3 | * |
| 1970 | 1.8 | * | * | * | * | * | 1.5 | 3.0 | 6.9 | 7.1 | 9.9 | * |
| 1969 | 2.1 | * | * | * | * | * | 2.4 | 2.8 | 5.6 | 11.5 | 19.0 | 30.8 |
| 1968 | 2.1 | * | * | * | * | * | 2.6 | 2.9 | 6.7 | 9.3 | 15.9 | 44.3 |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| PEPTIC ULCER (531-533) | | | | | | | | | | | | |
| BLACK, BOTH SEXES | | | | | | | | | | | | |
| 1978 | 1.9 | * | * | * | * | * | 1.0 | 2.5 | 5.9 | 9.7 | 15.4 | 26.7 |
| 1977 | 2.1 | * | * | * | * | 0.6 | 1.6 | 2.8 | 7.0 | 10.3 | 14.6 | 28.7 |
| 1976 | 2.4 | * | * | * | * | 0.9 | 1.9 | 4.4 | 5.9 | 12.5 | 16.6 | 31.6 |
| 1975 | 2.5 | * | * | * | * | * | 2.2 | 4.4 | 7.8 | 13.0 | 17.3 | 32.3 |
| 1974 | 2.5 | * | * | * | * | 0.6 | 2.2 | 4.6 | 7.9 | 15.0 | 13.9 | 28.9 |
| 1973 | 2.8 | * | * | * | * | 1.0 | 2.8 | 4.8 | 8.2 | 15.2 | 19.4 | 33.3 |
| 1972 | 3.1 | * | * | * | * | 0.9 | 3.7 | 6.8 | 9.9 | 15.9 | 18.9 | 18.2 |
| 1971 | 3.1 | 5.3 | * | * | * | 1.4 | 3.2 | 6.7 | 9.9 | 13.6 | 20.4 | * |
| 1970 | 3.3 | 5.1 | * | * | * | 1.2 | 3.3 | 6.2 | 11.0 | 15.7 | 22.8 | 23.0 |
| 1969 | 3.5 | 4.4 | * | * | * | 1.1 | 3.6 | 5.9 | 11.5 | 17.9 | 32.3 | * |
| 1968 | 3.8 | 4.4 | * | * | * | 1.5 | 4.5 | 6.8 | 13.0 | 19.7 | 21.7 | 33.7 |
| BLACK, MALE | | | | | | | | | | | | |
| 1978 | 2.5 | * | * | * | * | * | * | 4.1 | 9.0 | 14.9 | 23.8 | * |
| 1977 | 2.7 | * | * | * | * | * | 2.5 | 4.5 | 10.6 | 15.5 | 18.3 | * |
| 1976 | 3.4 | * | * | * | * | 1.6 | 3.0 | 6.8 | 9.2 | 18.9 | 26.5 | 48.9 |
| 1975 | 3.5 | * | * | * | * | * | 3.5 | 6.5 | 12.1 | 19.5 | 25.7 | 45.7 |
| 1974 | 3.7 | * | * | * | * | * | 3.2 | 7.3 | 12.9 | 23.4 | 23.9 | * |
| 1973 | 4.2 | * | * | * | * | 1.4 | 4.8 | 7.4 | 13.4 | 25.6 | 30.9 | * |
| 1972 | 4.7 | * | * | * | * | * | 5.5 | 10.7 | 16.9 | 27.3 | 29.4 | * |
| 1971 | 4.4 | * | * | * | * | 2.2 | 5.2 | 10.6 | 14.7 | 19.9 | 30.6 | * |
| 1970 | 4.7 | * | * | * | * | 2.0 | 5.3 | 10.0 | 15.3 | 26.0 | 40.9 | * |
| 1969 | 4.9 | * | * | * | * | * | 4.9 | 9.4 | 18.1 | 25.7 | 49.1 | * |
| 1968 | 5.6 | * | * | * | * | 2.7 | 6.7 | 11.4 | 20.3 | 32.5 | 29.7 | * |
| BLACK, FEMALE | | | | | | | | | | | | |
| 1978 | 1.3 | * | * | * | * | * | * | * | 3.4 | 5.8 | 9.8 | 26.3 |
| 1977 | 1.5 | * | * | * | * | * | * | * | 4.0 | 6.4 | 12.1 | 28.7 |
| 1976 | 1.5 | * | * | * | * | * | * | 2.4 | 3.2 | 7.6 | 10.0 | 22.5 |
| 1975 | 1.6 | * | * | * | * | * | * | 2.7 | 4.2 | 8.0 | 11.6 | 25.0 |
| 1974 | 1.5 | * | * | * | * | * | * | 2.2 | 3.8 | 8.6 | 7.1 | * |
| 1973 | 1.6 | * | * | * | * | * | * | 2.6 | 3.9 | 7.1 | 11.5 | 31.5 |
| 1972 | 1.7 | * | * | * | * | * | 2.1 | 3.5 | 4.0 | 7.0 | 11.4 | * |
| 1971 | 1.9 | * | * | * | * | * | 1.5 | 3.3 | 5.9 | 8.6 | 13.1 | * |
| 1970 | 1.9 | * | * | * | * | * | 1.6 | 2.8 | 7.3 | 7.6 | 9.6 | * |
| 1969 | 2.2 | * | * | * | * | * | 2.5 | 3.0 | 5.9 | 11.7 | 19.8 | * |
| 1968 | 2.1 | * | * | * | * | * | 2.7 | 2.8 | 6.6 | 9.4 | 15.8 | 35.7 |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| APPENDICITIS (540-543) | | | | | | | | | | | | |
| BOTH SEXES | | | | | | | | | | | | |
| 1978 | 0.3 | * | * | 0.1 | 0.1 | 0.1 | 0.1 | 0.3 | 0.5 | 1.2 | 2.3 | 4.2 |
| 1977 | 0.3 | * | * | 0.1 | * | * | 0.1 | 0.3 | 0.6 | 1.1 | 2.5 | 4.6 |
| 1976 | 0.3 | * | * | 0.1 | 0.1 | 0.1 | 0.2 | 0.3 | 0.6 | 1.2 | 2.2 | 3.9 |
| 1975 | 0.4 | * | * | 0.1 | 0.1 | 0.1 | 0.2 | 0.3 | 0.7 | 1.3 | 2.6 | 5.2 |
| 1974 | 0.4 | * | * | 0.1 | 0.1 | 0.1 | 0.2 | 0.4 | 0.7 | 1.4 | 2.9 | 5.0 |
| 1973 | 0.5 | * | * | 0.1 | 0.1 | 0.1 | 0.2 | 0.4 | 1.0 | 2.0 | 3.4 | 6.1 |
| 1972 | 0.6 | * | 0.1 | 0.2 | 0.2 | 0.2 | 0.2 | 0.5 | 1.1 | 1.9 | 3.7 | 7.9 |
| 1971 | 0.6 | * | 0.2 | 0.2 | 0.2 | 0.2 | 0.4 | 0.5 | 1.1 | 2.2 | 4.2 | 7.5 |
| 1970 | 0.7 | * | 0.2 | 0.2 | 0.2 | 0.2 | 0.3 | 0.5 | 1.3 | 2.6 | 5.0 | 8.0 |
| 1969 | 0.7 | * | 0.1 | 0.2 | 0.2 | 0.2 | 0.4 | 0.6 | 1.3 | 2.3 | 5.1 | 7.8 |
| 1968 | 0.7 | * | 0.2 | 0.2 | 0.2 | 0.2 | 0.3 | 0.6 | 1.4 | 2.8 | 5.6 | 9.1 |
| MALE | | | | | | | | | | | | |
| 1978 | 0.4 | * | * | * | 0.1 | * | * | 0.4 | 0.6 | 1.7 | 3.4 | 5.4 |
| 1977 | 0.4 | * | * | 0.1 | * | * | 0.2 | 0.3 | 0.9 | 1.6 | 3.6 | 4.8 |
| 1976 | 0.4 | * | * | 0.1 | 0.1 | * | 0.3 | 0.4 | 0.9 | 1.8 | 2.9 | 5.8 |
| 1975 | 0.5 | * | * | 0.1 | * | * | 0.2 | 0.4 | 0.9 | 1.9 | 3.9 | 6.7 |
| 1974 | 0.5 | * | * | 0.1 | 0.1 | * | 0.2 | 0.5 | 1.0 | 2.0 | 4.0 | 8.3 |
| 1973 | 0.6 | * | * | 0.1 | 0.2 | * | 0.3 | 0.6 | 1.4 | 3.0 | 5.1 | 7.4 |
| 1972 | 0.7 | * | * | 0.2 | 0.2 | * | 0.3 | 0.8 | 1.5 | 2.8 | 4.6 | 11.1 |
| 1971 | 0.7 | * | * | 0.2 | 0.2 | 0.2 | 0.5 | 0.7 | 1.6 | 3.2 | 5.7 | 8.2 |
| 1970 | 0.9 | * | * | 0.2 | 0.2 | 0.2 | 0.4 | 0.8 | 1.9 | 4.0 | 6.8 | 11.1 |
| 1969 | 0.9 | * | * | 0.3 | 0.2 | 0.2 | 0.6 | 0.9 | 1.9 | 3.5 | 7.2 | 11.1 |
| 1968 | 0.9 | * | * | 0.2 | 0.3 | 0.2 | 0.4 | 0.8 | 2.0 | 3.9 | 7.9 | 10.9 |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| APPENDICITIS (540-543) | | | | | | | | | | | | |
| FEMALE | | | | | | | | | | | | |
| 1978 | 0.3 | * | * | * | * | * | * | 0.2 | 0.4 | 0.8 | 1.7 | 3.7 |
| 1977 | 0.3 | * | * | * | * | * | * | 0.3 | 0.4 | 0.8 | 1.8 | 4.5 |
| 1976 | 0.3 | * | * | * | * | * | * | 0.2 | 0.3 | 0.8 | 1.8 | 2.9 |
| 1975 | 0.3 | * | * | * | * | * | * | 0.2 | 0.6 | 0.8 | 1.8 | 4.5 |
| 1974 | 0.3 | * | * | * | * | * | * | 0.3 | 0.5 | 0.9 | 2.2 | 3.4 |
| 1973 | 0.4 | * | * | 0.1 | * | 0.2 | 0.2 | 0.2 | 0.6 | 1.1 | 2.3 | 5.4 |
| 1972 | 0.5 | * | * | 0.2 | 0.1 | 0.2 | 0.2 | 0.3 | 0.7 | 1.2 | 3.0 | 6.3 |
| 1971 | 0.5 | * | * | 0.2 | 0.1 | * | 0.2 | 0.4 | 0.7 | 1.5 | 3.2 | 7.1 |
| 1970 | 0.5 | * | * | 0.2 | * | * | * | 0.3 | 0.8 | 1.6 | 3.8 | 6.3 |
| 1969 | 0.5 | * | * | 0.1 | 0.1 | * | 0.3 | 0.3 | 0.8 | 1.4 | 3.6 | 6.0 |
| 1968 | 0.6 | * | * | 0.2 | 0.1 | 0.2 | 0.2 | 0.5 | 0.8 | 2.0 | 4.0 | 8.1 |
| WHITE, BOTH SEXES | | | | | | | | | | | | |
| 1978 | 0.3 | * | * | 0.1 | 0.1 | * | * | 0.2 | 0.4 | 1.2 | 2.2 | 4.3 |
| 1977 | 0.3 | * | * | * | * | * | 0.1 | 0.3 | 0.5 | 1.0 | 2.3 | 4.3 |
| 1976 | 0.3 | * | * | * | 0.1 | 0.1 | 0.2 | 0.2 | 0.5 | 1.1 | 2.1 | 3.8 |
| 1975 | 0.4 | * | * | 0.1 | 0.1 | * | 0.1 | 0.2 | 0.6 | 1.2 | 2.5 | 5.3 |
| 1974 | 0.4 | * | * | 0.1 | 0.1 | * | 0.1 | 0.3 | 0.6 | 1.2 | 2.9 | 5.1 |
| 1973 | 0.5 | * | * | 0.1 | 0.1 | * | 0.2 | 0.3 | 0.8 | 1.7 | 3.3 | 6.1 |
| 1972 | 0.5 | * | * | 0.2 | 0.1 | 0.1 | 0.2 | 0.4 | 1.0 | 1.8 | 3.4 | 8.0 |
| 1971 | 0.6 | * | * | 0.2 | 0.1 | 0.1 | 0.3 | 0.5 | 1.0 | 2.1 | 4.1 | 7.8 |
| 1970 | 0.7 | * | * | 0.2 | 0.1 | 0.1 | 0.2 | 0.5 | 1.2 | 2.5 | 4.8 | 7.8 |
| 1969 | 0.6 | * | * | 0.2 | 0.1 | 0.1 | 0.3 | 0.5 | 1.2 | 2.1 | 5.0 | 7.8 |
| 1968 | 0.7 | * | 0.2 | 0.2 | 0.2 | 0.2 | 0.2 | 0.5 | 1.2 | 2.7 | 5.7 | 9.1 |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| APPENDICITIS (540-543) | | | | | | | | | | | | |
| WHITE, MALE | | | | | | | | | | | | |
| 1978 | 0.4 | * | * | * | * | * | * | 0.3 | 0.6 | 1.6 | 3.3 | 5.2 |
| 1977 | 0.4 | * | * | * | * | * | * | 0.2 | 0.8 | 1.4 | 3.1 | 4.7 |
| 1976 | 0.4 | * | * | * | * | * | 0.3 | 0.3 | 0.7 | 1.5 | 2.9 | 5.3 |
| 1975 | 0.4 | * | * | * | * | * | * | 0.3 | 0.8 | 1.8 | 3.6 | 6.9 |
| 1974 | 0.5 | * | * | * | * | * | * | 0.4 | 0.9 | 1.7 | 4.1 | 8.2 |
| 1973 | 0.6 | * | * | * | 0.1 | * | * | 0.5 | 1.1 | 2.7 | 4.9 | 7.6 |
| 1972 | 0.7 | * | * | 0.2 | 0.2 | * | 0.2 | 0.7 | 1.4 | 2.6 | 4.4 | 11.4 |
| 1971 | 0.7 | * | * | 0.2 | 0.2 | * | 0.4 | 0.7 | 1.4 | 3.1 | 5.6 | 8.6 |
| 1970 | 0.8 | * | * | 0.2 | 0.2 | 0.2 | 0.3 | 0.7 | 1.8 | 3.9 | 6.6 | 10.5 |
| 1969 | 0.8 | * | * | 0.2 | 0.1 | 0.2 | 0.5 | 0.7 | 1.6 | 3.2 | 7.2 | 10.8 |
| 1968 | 0.9 | * | * | 0.2 | 0.3 | 0.2 | 0.3 | 0.7 | 1.8 | 3.7 | 8.2 | 10.8 |
| WHITE, FEMALE | | | | | | | | | | | | |
| 1978 | 0.3 | * | * | * | * | * | * | * | 0.3 | 0.8 | 1.6 | 3.8 |
| 1977 | 0.3 | * | * | * | * | * | * | 0.3 | 0.3 | 0.6 | 1.8 | 4.1 |
| 1976 | 0.3 | * | * | * | * | * | * | * | 0.3 | 0.8 | 1.6 | 3.1 |
| 1975 | 0.3 | * | * | * | * | * | * | * | 0.4 | 0.7 | 1.8 | 4.5 |
| 1974 | 0.3 | * | * | * | * | * | * | 0.2 | 0.4 | 0.8 | 2.2 | 3.6 |
| 1973 | 0.3 | * | * | * | * | * | * | * | 0.5 | 1.0 | 2.3 | 5.5 |
| 1972 | 0.4 | * | * | 0.2 | * | 0.2 | * | 0.2 | 0.7 | 1.1 | 2.8 | 6.4 |
| 1971 | 0.5 | * | * | 0.1 | * | * | 0.2 | 0.3 | 0.6 | 1.4 | 3.1 | 7.3 |
| 1970 | 0.5 | * | * | 0.1 | * | * | * | 0.3 | 0.7 | 1.4 | 3.7 | 6.4 |
| 1969 | 0.5 | * | * | 0.1 | * | * | 0.2 | 0.3 | 0.8 | 1.3 | 3.5 | 6.3 |
| 1968 | 0.5 | * | * | 0.2 | * | * | * | 0.3 | 0.6 | 2.0 | 4.0 | 8.3 |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| APPENDICITIS (540-543) | | | | | | | | | | | | |
| ALL OTHER, BOTH SEXES | | | | | | | | | | | | |
| 1978 | 0.5 | * | * | * | * | * | * | 0.8 | 1.2 | 1.6 | 3.4 | * |
| 1977 | 0.6 | * | * | * | * | * | * | * | 1.5 | 2.9 | 4.8 | * |
| 1976 | 0.6 | * | * | * | * | * | * | 0.9 | 1.4 | 2.7 | * | * |
| 1975 | 0.6 | * | * | * | * | * | * | 0.9 | 1.7 | 2.1 | 4.4 | * |
| 1974 | 0.5 | * | * | * | * | * | * | 0.9 | 1.6 | 3.5 | * | * |
| 1973 | 0.8 | * | * | * | * | * | 0.7 | 1.2 | 2.5 | 4.0 | 4.6 | * |
| 1972 | 0.8 | * | * | * | * | * | 0.7 | 1.5 | 1.8 | 2.9 | 6.6 | * |
| 1971 | 0.9 | * | * | 0.4 | * | * | 0.8 | 1.0 | 2.4 | 3.6 | 5.1 | * |
| 1970 | 0.9 | * | * | * | * | * | * | 1.2 | 2.2 | 4.0 | 6.9 | * |
| 1969 | 1.1 | * | * | 0.4 | * | * | 1.1 | 1.7 | 2.9 | 4.4 | 5.5 | * |
| 1968 | 1.0 | * | * | 0.4 | * | * | 1.0 | 1.9 | 3.1 | 3.3 | * | * |
| ALL OTHER, MALE | | | | | | | | | | | | |
| 1978 | 0.5 | * | * | * | * | * | * | * | * | * | * | * |
| 1977 | 0.6 | * | * | * | * | * | * | * | * | * | 8.8 | * |
| 1976 | 0.7 | * | * | * | * | * | * | * | 2.2 | 4.5 | * | * |
| 1975 | 0.7 | * | * | * | * | * | * | * | * | * | * | * |
| 1974 | 0.7 | * | * | * | * | * | * | * | 2.4 | 4.8 | * | * |
| 1973 | 1.1 | * | * | * | * | * | * | * | 3.6 | 5.9 | * | * |
| 1972 | 0.9 | * | * | * | * | * | * | 1.9 | 2.8 | 4.1 | * | * |
| 1971 | 0.9 | * | * | * | * | * | * | * | 2.9 | 4.4 | * | * |
| 1970 | 1.1 | * | * | * | * | * | * | 1.9 | 2.9 | 4.4 | * | * |
| 1969 | 1.5 | * | * | * | * | * | 1.6 | 2.4 | 4.3 | 6.7 | * | * |
| 1968 | 1.1 | * | * | * | * | * | * | 1.9 | 4.1 | 5.0 | * | * |
| ALL OTHER, FEMALE | | | | | | | | | | | | |
| 1978 | 0.4 | * | * | * | * | * | * | * | * | * | * | * |
| 1977 | 0.5 | * | * | * | * | * | * | * | * | 2.8 | * | * |
| 1976 | 0.4 | * | * | * | * | * | * | * | * | * | * | * |
| 1975 | 0.5 | * | * | * | * | * | * | * | * | * | * | * |
| 1974 | 0.4 | * | * | * | * | * | * | * | * | * | * | * |
| 1973 | 0.6 | * | * | * | * | * | * | * | * | * | * | * |
| 1972 | 0.7 | * | * | * | * | * | * | * | * | * | * | * |
| 1971 | 0.8 | * | * | * | * | * | * | * | * | * | * | * |
| 1970 | 0.7 | * | * | * | * | * | * | * | * | 3.6 | * | * |
| 1969 | 0.7 | * | * | * | * | * | * | * | * | * | * | * |
| 1968 | 0.8 | * | * | * | * | * | * | 2.0 | * | * | * | * |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| APPENDICITIS (540-543) | | | | | | | | | | | | |
| BLACK, BOTH SEXES | | | | | | | | | | | | |
| 1978 | 0.5 | * | * | * | * | * | * | 0.9 | 1.3 | 1.8 | 3.4 | * |
| 1977 | 0.6 | * | * | * | * | * | * | * | 1.7 | 3.0 | 4.9 | * |
| 1976 | 0.6 | * | * | * | * | * | * | 1.0 | 1.4 | 2.9 | * | * |
| 1975 | 0.6 | * | * | * | * | * | * | 1.0 | 1.8 | 2.3 | 4.2 | * |
| 1974 | 0.5 | * | * | * | * | * | * | 0.9 | 1.6 | 3.7 | * | * |
| 1973 | 0.9 | * | * | * | * | * | * | 1.3 | 2.7 | 4.1 | 4.6 | * |
| 1972 | 0.8 | * | * | * | * | * | 0.8 | 1.6 | 1.9 | 3.0 | 6.3 | * |
| 1971 | 0.9 | * | * | * | * | * | 0.9 | 1.1 | 2.5 | 3.8 | 5.2 | * |
| 1970 | 0.9 | * | * | * | * | * | * | 1.3 | 2.2 | 4.1 | 7.5 | * |
| 1969 | 1.2 | * | * | 0.5 | * | * | 1.2 | 1.8 | 3.0 | 4.7 | 5.9 | * |
| 1968 | 1.0 | * | * | 0.4 | * | * | 1.0 | 2.1 | 3.3 | 3.6 | * | * |
| BLACK, MALE | | | | | | | | | | | | |
| 1978 | 0.5 | * | * | * | * | * | * | * | * | * | * | * |
| 1977 | 0.7 | * | * | * | * | * | * | * | * | * | 8.9 | * |
| 1976 | 0.8 | * | * | * | * | * | * | * | * | 5.2 | * | * |
| 1975 | 0.7 | * | * | * | * | * | * | * | * | * | * | * |
| 1974 | 0.8 | * | * | * | * | * | * | * | * | 5.3 | * | * |
| 1973 | 1.1 | * | * | * | * | * | * | * | 4.1 | 6.0 | * | * |
| 1972 | 0.9 | * | * | * | * | * | * | 2.2 | 3.2 | 4.3 | * | * |
| 1971 | 1.0 | * | * | * | * | * | * | * | 3.1 | 4.5 | * | * |
| 1970 | 1.2 | * | * | * | * | * | * | 2.0 | 3.0 | 4.8 | * | * |
| 1969 | 1.6 | * | * | * | * | * | * | 2.6 | 4.5 | 7.3 | * | * |
| 1968 | 1.2 | * | * | * | * | * | * | * | 4.5 | 5.5 | * | * |
| BLACK, FEMALE | | | | | | | | | | | | |
| 1978 | 0.4 | * | * | * | * | * | * | * | * | * | * | * |
| 1977 | 0.5 | * | * | * | * | * | * | * | * | 3.1 | * | * |
| 1976 | 0.5 | * | * | * | * | * | * | * | * | * | * | * |
| 1975 | 0.5 | * | * | * | * | * | * | * | * | * | * | * |
| 1974 | 0.4 | * | * | * | * | * | * | * | * | * | * | * |
| 1973 | 0.6 | * | * | * | * | * | * | * | * | * | * | * |
| 1972 | 0.7 | * | * | * | * | * | * | * | * | * | * | * |
| 1971 | 0.8 | * | * | * | * | * | * | * | * | * | * | * |
| 1970 | 0.7 | * | * | * | * | * | * | * | * | 3.6 | * | * |
| 1969 | 0.8 | * | * | * | * | * | * | * | * | * | * | * |
| 1968 | 0.9 | * | * | * | * | * | * | 2.2 | * | * | * | * |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|--|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| HERNIA AND INTESTINAL OBSTRUCTION (550-553, 560) | | | | | | | | | | | | |
| BOTH SEXES | | | | | | | | | | | | |
| 1978 | 2.5 | 14.9 | 0.2 | * | 0.1 | 0.1 | 0.4 | 1.0 | 2.7 | 7.0 | 22.9 | 60.9 |
| 1977 | 2.6 | 14.4 | 0.2 | 0.1 | 0.1 | 0.2 | 0.4 | 1.1 | 2.6 | 7.6 | 23.8 | 67.1 |
| 1976 | 2.7 | 15.3 | 0.3 | * | 0.1 | 0.2 | 0.5 | 1.1 | 2.8 | 9.0 | 24.8 | 67.8 |
| 1975 | 2.9 | 14.1 | 0.2 | 0.1 | 0.2 | 0.2 | 0.4 | 1.4 | 3.3 | 9.2 | 27.3 | 68.6 |
| 1974 | 3.0 | 16.2 | 0.3 | 0.1 | 0.1 | 0.2 | 0.6 | 1.5 | 3.6 | 10.0 | 29.9 | 71.9 |
| 1973 | 3.2 | 15.2 | 0.3 | 0.1 | 0.2 | 0.3 | 0.7 | 1.7 | 4.2 | 10.7 | 30.0 | 76.5 |
| 1972 | 3.2 | 14.8 | 0.3 | 0.1 | 0.1 | 0.3 | 0.6 | 1.8 | 4.5 | 11.0 | 31.5 | 76.3 |
| 1971 | 3.4 | 14.4 | 0.3 | 0.1 | 0.2 | 0.3 | 0.7 | 1.8 | 4.8 | 11.6 | 32.3 | 81.3 |
| 1970 | 3.6 | 18.0 | 0.3 | 0.1 | 0.2 | 0.4 | 0.8 | 1.8 | 5.2 | 12.7 | 34.3 | 72.8 |
| 1969 | 3.7 | 18.7 | 0.3 | 0.1 | 0.2 | 0.3 | 0.8 | 1.9 | 5.0 | 14.0 | 36.3 | 86.0 |
| 1968 | 3.9 | 19.0 | 0.4 | 0.1 | 0.2 | 0.4 | 0.9 | 2.3 | 5.5 | 14.3 | 39.0 | 86.9 |
| MALE | | | | | | | | | | | | |
| 1978 | 2.2 | 17.4 | * | * | 0.1 | 0.1 | 0.4 | 1.1 | 2.8 | 7.9 | 22.7 | 58.4 |
| 1977 | 2.2 | 16.1 | * | * | 0.1 | 0.2 | 0.4 | 1.0 | 2.7 | 8.0 | 24.5 | 66.5 |
| 1976 | 2.4 | 17.9 | 0.4 | * | 0.1 | 0.2 | 0.6 | 1.0 | 3.0 | 10.0 | 24.9 | 63.2 |
| 1975 | 2.5 | 15.7 | * | 0.1 | 0.1 | 0.2 | 0.4 | 1.3 | 3.6 | 10.1 | 27.6 | 64.3 |
| 1974 | 2.6 | 16.8 | 0.4 | * | 0.2 | 0.2 | 0.5 | 1.4 | 3.5 | 10.9 | 30.0 | 71.0 |
| 1973 | 2.9 | 17.9 | 0.4 | 0.1 | 0.2 | 0.4 | 0.6 | 1.7 | 4.5 | 11.5 | 31.1 | 82.7 |
| 1972 | 2.9 | 17.9 | 0.4 | 0.1 | 0.1 | 0.3 | 0.6 | 1.4 | 4.7 | 11.6 | 33.0 | 76.2 |
| 1971 | 3.1 | 15.4 | 0.4 | * | 0.2 | 0.3 | 0.7 | 1.7 | 5.1 | 12.5 | 34.9 | 76.7 |
| 1970 | 3.4 | 20.0 | 0.4 | 0.1 | 0.2 | 0.5 | 0.8 | 1.8 | 5.8 | 14.1 | 35.3 | 75.6 |
| 1969 | 3.5 | 21.5 | 0.5 | 0.1 | 0.2 | 0.3 | 0.7 | 1.7 | 5.6 | 15.9 | 38.4 | 84.5 |
| 1968 | 3.7 | 22.4 | 0.6 | 0.1 | 0.3 | 0.4 | 0.8 | 2.1 | 5.8 | 15.8 | 41.1 | 81.1 |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|--|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| HERNIA AND INTESTINAL OBSTRUCTION (550-553, 560) | | | | | | | | | | | | |
| FEMALE | | | | | | | | | | | | |
| 1978 | 2.8 | 12.3 | * | * | * | 0.1 | 0.5 | 1.0 | 2.5 | 6.3 | 23.0 | 62.0 |
| 1977 | 2.9 | 12.7 | * | * | 0.1 | * | 0.4 | 1.1 | 2.4 | 7.4 | 23.4 | 67.4 |
| 1976 | 3.0 | 12.7 | * | * | * | 0.2 | 0.4 | 1.3 | 2.7 | 8.2 | 24.8 | 70.0 |
| 1975 | 3.2 | 12.5 | * | * | 0.2 | 0.2 | 0.4 | 1.5 | 3.0 | 8.5 | 27.1 | 70.7 |
| 1974 | 3.4 | 15.6 | * | * | * | 0.2 | 0.6 | 1.6 | 3.7 | 9.4 | 29.8 | 72.4 |
| 1973 | 3.5 | 12.5 | * | * | 0.1 | 0.3 | 0.8 | 1.7 | 4.0 | 10.1 | 29.3 | 73.3 |
| 1972 | 3.6 | 11.5 | * | * | 0.1 | 0.4 | 0.6 | 2.1 | 4.3 | 10.6 | 30.5 | 76.3 |
| 1971 | 3.7 | 13.4 | * | 0.1 | 0.2 | 0.3 | 0.7 | 1.8 | 4.6 | 10.9 | 30.7 | 83.7 |
| 1970 | 3.7 | 15.9 | * | * | 0.2 | 0.3 | 0.8 | 1.9 | 4.7 | 11.6 | 33.6 | 71.2 |
| 1969 | 3.9 | 15.9 | * | * | 0.2 | 0.3 | 1.0 | 2.1 | 4.5 | 12.6 | 34.9 | 86.8 |
| 1968 | 4.1 | 15.5 | * | 0.1 | 0.1 | 0.4 | 1.0 | 2.4 | 5.2 | 13.1 | 37.5 | 90.3 |
| WHITE, BOTH SEXES | | | | | | | | | | | | |
| 1978 | 2.5 | 14.4 | 0.2 | * | 0.1 | 0.1 | 0.3 | 0.8 | 2.4 | 6.6 | 22.7 | 62.7 |
| 1977 | 2.6 | 14.5 | * | 0.1 | 0.1 | 0.1 | 0.3 | 1.0 | 2.3 | 7.1 | 23.8 | 68.7 |
| 1976 | 2.7 | 14.8 | 0.3 | * | 0.1 | 0.2 | 0.3 | 0.9 | 2.5 | 8.4 | 25.2 | 69.0 |
| 1975 | 2.9 | 13.5 | * | 0.1 | 0.1 | 0.2 | 0.4 | 1.2 | 3.0 | 8.5 | 27.6 | 70.0 |
| 1974 | 3.1 | 16.5 | 0.3 | 0.1 | 0.1 | 0.2 | 0.5 | 1.3 | 3.3 | 9.5 | 30.2 | 74.0 |
| 1973 | 3.2 | 14.6 | 0.2 | 0.1 | 0.1 | 0.3 | 0.6 | 1.5 | 3.7 | 10.1 | 30.4 | 78.0 |
| 1972 | 3.3 | 14.3 | 0.3 | 0.1 | 0.1 | 0.2 | 0.4 | 1.5 | 4.3 | 10.3 | 32.3 | 77.9 |
| 1971 | 3.3 | 13.4 | 0.3 | 0.1 | 0.1 | 0.3 | 0.6 | 1.5 | 4.4 | 10.7 | 32.6 | 83.3 |
| 1970 | 3.6 | 17.5 | 0.3 | 0.1 | 0.2 | 0.3 | 0.6 | 1.6 | 4.8 | 12.2 | 34.3 | 76.0 |
| 1969 | 3.7 | 17.9 | 0.3 | 0.1 | 0.2 | 0.2 | 0.7 | 1.7 | 4.7 | 13.6 | 36.2 | 88.2 |
| 1968 | 3.9 | 18.0 | 0.4 | 0.1 | 0.2 | 0.3 | 0.7 | 2.0 | 5.0 | 13.8 | 39.3 | 89.9 |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|--|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| HERNIA AND INTESTINAL OBSTRUCTION (550-553, 560) | | | | | | | | | | | | |
| WHITE, MALE | | | | | | | | | | | | |
| 1978 | 2.1 | 16.1 | * | * | 0.1 | * | 0.3 | 0.9 | 2.6 | 7.3 | 22.2 | 59.3 |
| 1977 | 2.2 | 15.8 | * | * | * | 0.2 | 0.3 | 0.8 | 2.4 | 7.3 | 23.7 | 67.4 |
| 1976 | 2.3 | 17.0 | * | * | * | 0.2 | 0.4 | 0.9 | 2.7 | 9.3 | 24.6 | 63.2 |
| 1975 | 2.4 | 15.1 | * | * | 0.1 | 0.2 | 0.4 | 1.1 | 3.4 | 9.1 | 27.5 | 65.0 |
| 1974 | 2.6 | 16.7 | 0.3 | * | 0.2 | 0.2 | 0.4 | 1.2 | 3.1 | 10.3 | 29.7 | 71.7 |
| 1973 | 2.8 | 17.0 | * | 0.1 | 0.2 | 0.3 | 0.5 | 1.4 | 3.9 | 10.8 | 31.0 | 84.3 |
| 1972 | 2.8 | 17.0 | 0.4 | * | 0.1 | 0.2 | 0.4 | 1.3 | 4.3 | 10.6 | 33.4 | 78.3 |
| 1971 | 2.9 | 14.1 | * | * | 0.1 | 0.3 | 0.5 | 1.4 | 4.5 | 11.4 | 34.4 | 78.6 |
| 1970 | 3.3 | 19.7 | 0.4 | * | 0.2 | 0.4 | 0.6 | 1.5 | 5.4 | 13.3 | 35.0 | 77.6 |
| 1969 | 3.4 | 19.9 | 0.4 | 0.1 | 0.2 | 0.3 | 0.6 | 1.5 | 5.2 | 15.2 | 37.4 | 85.1 |
| 1968 | 3.6 | 21.0 | 0.6 | 0.1 | 0.2 | 0.3 | 0.6 | 1.9 | 5.1 | 15.2 | 40.9 | 81.2 |
| WHITE, FEMALE | | | | | | | | | | | | |
| 1978 | 2.9 | 12.7 | * | * | * | * | 0.4 | 0.8 | 2.3 | 6.1 | 23.0 | 64.3 |
| 1977 | 3.0 | 13.1 | * | * | 0.1 | * | 0.3 | 1.1 | 2.2 | 7.0 | 23.8 | 69.3 |
| 1976 | 3.1 | 12.5 | * | * | * | * | 0.3 | 1.0 | 2.4 | 7.6 | 25.5 | 71.7 |
| 1975 | 3.3 | 11.7 | * | * | 0.2 | 0.2 | 0.4 | 1.3 | 2.8 | 8.0 | 27.7 | 72.5 |
| 1974 | 3.6 | 16.2 | * | * | * | 0.2 | 0.5 | 1.4 | 3.5 | 8.9 | 30.5 | 75.1 |
| 1973 | 3.5 | 12.2 | * | * | * | 0.2 | 0.7 | 1.5 | 3.5 | 9.6 | 30.0 | 74.9 |
| 1972 | 3.7 | 11.4 | * | * | * | 0.3 | 0.5 | 1.7 | 4.3 | 10.1 | 31.5 | 77.8 |
| 1971 | 3.7 | 12.7 | * | * | 0.1 | 0.3 | 0.6 | 1.6 | 4.2 | 10.2 | 31.5 | 85.7 |
| 1970 | 3.8 | 15.3 | * | * | 0.2 | 0.3 | 0.7 | 1.7 | 4.3 | 11.4 | 33.9 | 75.1 |
| 1969 | 4.0 | 15.8 | * | * | 0.2 | 0.2 | 0.7 | 1.8 | 4.3 | 12.3 | 35.4 | 90.0 |
| 1968 | 4.2 | 14.8 | * | 0.1 | * | 0.3 | 0.8 | 2.2 | 4.8 | 12.7 | 38.2 | 94.7 |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|--|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| HERNIA AND INTESTINAL OBSTRUCTION (550-553, 560) | | | | | | | | | | | | |
| ALL OTHER, BOTH SEXES | | | | | | | | | | | | |
| 1978 | 2.4 | 17.0 | * | * | * | * | 1.1 | 2.6 | 4.7 | 10.6 | 24.8 | 39.9 |
| 1977 | 2.4 | 14.3 | * | * | * | * | 1.0 | 2.2 | 4.8 | 12.5 | 24.6 | 48.8 |
| 1976 | 2.7 | 17.8 | * | * | * | * | 1.4 | 2.7 | 5.4 | 15.2 | 21.2 | 54.2 |
| 1975 | 2.8 | 17.5 | * | * | * | 0.6 | 0.9 | 3.2 | 5.4 | 15.8 | 24.4 | 52.7 |
| 1974 | 2.8 | 15.0 | * | * | * | 0.6 | 1.4 | 3.3 | 6.0 | 15.4 | 26.4 | 48.1 |
| 1973 | 3.3 | 18.3 | * | * | * | 0.7 | 1.6 | 3.7 | 9.3 | 16.8 | 25.2 | 59.1 |
| 1972 | 3.1 | 17.4 | * | * | * | 0.9 | 1.6 | 3.9 | 6.8 | 17.4 | 22.4 | 56.9 |
| 1971 | 3.6 | 19.9 | * | * | 0.4 | 0.8 | 1.7 | 3.9 | 9.4 | 20.0 | 28.3 | 58.3 |
| 1970 | 3.5 | 20.5 | * | * | 0.4 | 1.0 | 1.8 | 4.0 | 9.0 | 17.3 | 34.0 | 40.3 |
| 1969 | 3.7 | 23.3 | * | * | 0.6 | 0.8 | 2.3 | 3.9 | 8.0 | 18.5 | 37.6 | 60.6 |
| 1968 | 3.9 | 24.3 | * | * | * | 1.0 | 2.3 | 4.5 | 10.3 | 18.9 | 35.0 | 53.9 |
| ALL OTHER, MALE | | | | | | | | | | | | |
| 1978 | 2.6 | 23.5 | * | * | * | * | * | 3.3 | 5.2 | 12.8 | 27.1 | 50.0 |
| 1977 | 2.7 | 17.6 | * | * | * | * | * | 2.6 | 5.6 | 14.4 | 32.2 | 57.1 |
| 1976 | 2.8 | 22.2 | * | * | * | * | 1.9 | 2.2 | 6.3 | 16.7 | 28.0 | 63.6 |
| 1975 | 2.9 | 18.8 | * | * | * | * | * | 3.1 | 6.0 | 19.1 | 29.3 | 57.4 |
| 1974 | 3.1 | 17.4 | * | * | * | * | 1.6 | 3.6 | 7.1 | 16.3 | 33.6 | 64.0 |
| 1973 | 3.5 | 22.6 | * | * | * | * | 1.7 | 4.4 | 10.1 | 18.2 | 32.1 | 66.7 |
| 1972 | 3.3 | 22.7 | * | * | * | * | 1.9 | 2.6 | 9.0 | 20.3 | 28.6 | 55.3 |
| 1971 | 4.0 | 22.6 | * | * | * | * | 1.9 | 4.1 | 10.9 | 22.7 | 40.6 | 57.4 |
| 1970 | 3.8 | 21.7 | * | * | * | * | 2.5 | 4.4 | 9.7 | 21.1 | 39.4 | 57.7 |
| 1969 | 4.4 | 30.6 | * | * | * | * | 2.1 | 3.9 | 10.1 | 22.7 | 50.0 | 79.5 |
| 1968 | 4.5 | 29.4 | * | * | * | * | 2.2 | 4.6 | 12.3 | 21.6 | 44.1 | 80.5 |
| ALL OTHER, FEMALE | | | | | | | | | | | | |
| 1978 | 2.2 | 10.3 | * | * | * | * | * | 1.9 | 4.3 | 8.9 | 23.4 | 34.9 |
| 1977 | 2.2 | 10.8 | * | * | * | * | * | 1.9 | 4.2 | 11.0 | 19.3 | 44.2 |
| 1976 | 2.5 | 13.4 | * | * | * | * | * | 3.1 | 4.7 | 14.0 | 16.5 | 49.0 |
| 1975 | 2.7 | 16.2 | * | * | * | * | * | 3.3 | 4.8 | 13.1 | 20.9 | 49.5 |
| 1974 | 2.6 | 12.6 | * | * | * | * | * | 3.0 | 5.2 | 14.6 | 21.4 | 38.8 |
| 1973 | 3.1 | 13.8 | * | * | * | * | 1.4 | 3.1 | 8.6 | 15.7 | 20.4 | 54.4 |
| 1972 | 2.8 | 11.9 | * | * | * | * | 1.3 | 5.0 | 4.9 | 15.0 | 17.9 | 57.9 |
| 1971 | 3.2 | 17.1 | * | * | * | * | 1.5 | 3.8 | 8.1 | 17.8 | 19.3 | 58.9 |
| 1970 | 3.2 | 19.4 | * | * | * | * | 1.3 | 3.7 | 8.4 | 14.2 | 30.0 | 28.0 |
| 1969 | 3.2 | 16.0 | * | * | * | * | 2.4 | 4.0 | 6.2 | 15.1 | 27.9 | 47.7 |
| 1968 | 3.4 | 19.1 | * | * | * | * | 2.4 | 4.3 | 8.6 | 16.7 | 28.0 | 36.1 |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|--|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| HERNIA AND INTESTINAL OBSTRUCTION (550-553, 560) | | | | | | | | | | | | |
| BLACK, BOTH SEXES | | | | | | | | | | | | |
| 1978 | 2.6 | 17.9 | * | * | * | * | 1.2 | 2.8 | 5.2 | 11.3 | 26.3 | 40.0 |
| 1977 | 2.7 | 15.1 | * | * | * | * | 1.1 | 2.4 | 5.2 | 13.6 | 24.6 | 52.4 |
| 1976 | 2.9 | 17.8 | * | * | * | * | 1.5 | 3.1 | 5.6 | 16.4 | 21.8 | 55.9 |
| 1975 | 3.0 | 18.6 | * | * | * | 0.7 | 1.0 | 3.5 | 5.9 | 16.8 | 24.7 | 54.6 |
| 1974 | 3.0 | 15.0 | * | * | * | 0.7 | 1.5 | 3.7 | 6.4 | 16.3 | 27.8 | 49.6 |
| 1973 | 3.5 | 20.4 | * | * | * | 0.8 | 1.8 | 4.0 | 10.0 | 17.6 | 26.3 | 57.9 |
| 1972 | 3.3 | 17.2 | * | * | * | 1.0 | 1.8 | 4.1 | 7.5 | 18.3 | 23.9 | 63.6 |
| 1971 | 3.8 | 21.0 | * | * | * | 0.9 | 2.0 | 4.3 | 10.0 | 20.8 | 29.0 | 64.5 |
| 1970 | 3.7 | 22.0 | * | * | * | 1.1 | 2.0 | 4.5 | 9.3 | 17.9 | 35.1 | 40.0 |
| 1969 | 3.9 | 25.4 | * | * | 0.6 | 1.0 | 2.5 | 4.1 | 8.5 | 19.3 | 38.9 | 57.7 |
| 1968 | 4.1 | 25.4 | * | * | * | 1.1 | 2.5 | 4.8 | 10.9 | 19.7 | 36.4 | 55.4 |
| BLACK, MALE | | | | | | | | | | | | |
| 1978 | 2.9 | 24.6 | * | * | * | * | * | 3.7 | 6.0 | 13.6 | 28.0 | 49.0 |
| 1977 | 2.9 | 18.4 | * | * | * | * | * | 3.0 | 5.9 | 16.2 | 31.9 | 61.2 |
| 1976 | 3.0 | 22.3 | * | * | * | * | 2.0 | 2.5 | 6.6 | 18.3 | 29.5 | 66.0 |
| 1975 | 3.1 | 19.6 | * | * | * | * | * | 3.3 | 6.8 | 20.7 | 30.2 | 60.9 |
| 1974 | 3.3 | 17.3 | * | * | * | * | 1.9 | 4.1 | 7.9 | 17.7 | 34.9 | 67.4 |
| 1973 | 3.8 | 24.9 | * | * | * | * | 2.0 | 4.7 | 11.2 | 19.0 | 33.8 | 65.9 |
| 1972 | 3.6 | 22.4 | * | * | * | * | 2.2 | 3.0 | 10.1 | 21.7 | 31.5 | 65.0 |
| 1971 | 4.2 | 23.9 | * | * | * | * | 2.2 | 4.4 | 11.9 | 23.3 | 41.9 | 65.0 |
| 1970 | 4.1 | 23.7 | * | * | * | * | 2.7 | 4.8 | 10.3 | 22.1 | 42.7 | 60.9 |
| 1969 | 4.6 | 33.2 | * | * | * | * | 2.4 | 4.3 | 10.8 | 23.7 | 52.7 | 81.1 |
| 1968 | 4.7 | 30.3 | * | * | * | * | 2.5 | 4.9 | 13.3 | 23.0 | 47.9 | 85.7 |
| BLACK, FEMALE | | | | | | | | | | | | |
| 1978 | 2.4 | 11.1 | * | * | * | * | * | 2.1 | 4.6 | 9.6 | 25.1 | 35.4 |
| 1977 | 2.4 | 11.7 | * | * | * | * | * | 1.9 | 4.6 | 11.6 | 19.8 | 47.9 |
| 1976 | 2.7 | 13.3 | * | * | * | * | * | 3.5 | 4.8 | 14.9 | 16.6 | 50.6 |
| 1975 | 2.9 | 17.7 | * | * | * | * | * | 3.7 | 5.2 | 13.7 | 21.0 | 51.2 |
| 1974 | 2.7 | 12.7 | * | * | * | * | * | 3.4 | 5.2 | 15.2 | 23.0 | 39.7 |
| 1973 | 3.3 | 15.7 | * | * | * | * | 1.7 | 3.3 | 9.1 | 16.6 | 21.3 | 53.4 |
| 1972 | 3.0 | 11.8 | * | * | * | * | 1.5 | 5.1 | 5.3 | 15.6 | 18.6 | 62.9 |
| 1971 | 3.5 | 18.0 | * | * | * | * | 1.8 | 4.1 | 8.4 | 18.7 | 19.7 | 64.2 |
| 1970 | 3.3 | 20.2 | * | * | * | * | 1.5 | 4.1 | 8.5 | 14.6 | 29.6 | * |
| 1969 | 3.3 | 17.5 | * | * | * | * | 2.6 | 3.8 | 6.5 | 15.7 | 28.6 | 43.3 |
| 1968 | 3.6 | 20.4 | * | * | * | * | 2.6 | 4.6 | 8.8 | 17.1 | 27.9 | 37.5 |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| Chronic liver disease and cirrhosis 571 | | | | | | | | | | | | |
| BOTH SEXES | | | | | | | | | | | | |
| 1978 | 13.5 | 0.8 | * | * | 0.3 | 3.4 | 14.4 | 32.2 | 42.6 | 41.4 | 28.8 | 18.9 |
| 1977 | 14.0 | 1.0 | * | * | 0.3 | 3.7 | 15.3 | 33.8 | 44.6 | 42.4 | 28.4 | 17.7 |
| 1976 | 14.5 | 1.1 | * | * | 0.3 | 3.6 | 16.9 | 35.0 | 46.9 | 42.4 | 27.8 | 18.6 |
| 1975 | 14.7 | 1.1 | * | * | 0.4 | 3.7 | 16.7 | 36.0 | 48.3 | 42.7 | 27.7 | 17.5 |
| 1974 | 15.6 | 1.1 | * | 0.1 | 0.4 | 4.2 | 18.6 | 38.7 | 50.4 | 45.7 | 28.4 | 18.9 |
| 1973 | 15.8 | 1.2 | * | 0.1 | 0.4 | 4.2 | 19.4 | 39.1 | 52.1 | 44.5 | 28.0 | 20.2 |
| 1972 | 15.6 | 0.8 | 0.2 | 0.1 | 0.3 | 4.3 | 20.3 | 38.9 | 50.0 | 43.0 | 28.9 | 20.5 |
| 1971 | 15.4 | 0.8 | 0.2 | 0.1 | 0.5 | 4.2 | 19.3 | 38.5 | 49.6 | 42.2 | 29.7 | 21.8 |
| 1970 | 15.5 | 0.7 | 0.2 | 0.2 | 0.4 | 4.4 | 19.6 | 38.3 | 49.3 | 42.2 | 30.9 | 20.3 |
| 1969 | 14.8 | 1.3 | 0.3 | 0.1 | 0.5 | 4.3 | 19.0 | 36.7 | 47.2 | 39.8 | 29.5 | 21.5 |
| 1968 | 14.6 | 1.5 | 0.2 | 0.1 | 0.4 | 4.2 | 18.1 | 36.8 | 46.1 | 40.3 | 29.6 | 22.7 |
| MALE | | | | | | | | | | | | |
| 1978 | 18.2 | * | * | * | 0.3 | 4.6 | 19.6 | 43.4 | 60.8 | 63.6 | 44.8 | 31.3 |
| 1977 | 18.9 | * | * | * | 0.4 | 5.1 | 20.6 | 46.5 | 63.9 | 62.6 | 44.6 | 27.9 |
| 1976 | 19.5 | 1.3 | * | * | 0.3 | 4.9 | 22.8 | 48.1 | 67.1 | 64.4 | 42.4 | 29.7 |
| 1975 | 19.9 | * | * | * | 0.4 | 5.1 | 22.7 | 49.2 | 68.4 | 66.3 | 44.2 | 28.3 |
| 1974 | 21.0 | 1.3 | * | * | 0.4 | 5.4 | 24.4 | 52.6 | 71.0 | 71.5 | 45.3 | 29.0 |
| 1973 | 21.2 | * | * | * | 0.4 | 5.5 | 25.4 | 52.2 | 74.3 | 69.0 | 43.4 | 33.6 |
| 1972 | 21.0 | * | * | * | 0.3 | 5.3 | 26.8 | 53.1 | 72.2 | 66.4 | 44.2 | 35.6 |
| 1971 | 20.5 | * | * | * | 0.4 | 5.4 | 24.8 | 51.4 | 71.3 | 64.6 | 45.4 | 34.5 |
| 1970 | 20.6 | * | * | 0.1 | 0.4 | 5.6 | 24.8 | 50.8 | 70.7 | 65.9 | 46.2 | 30.6 |
| 1969 | 19.8 | 1.2 | * | * | 0.5 | 5.2 | 24.2 | 48.2 | 68.7 | 61.2 | 43.5 | 35.3 |
| 1968 | 19.4 | 1.5 | * | 0.1 | 0.3 | 5.1 | 22.6 | 48.6 | 66.4 | 61.1 | 44.3 | 30.9 |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| Chronic liver disease and cirrhosis 571 | | | | | | | | | | | | |
| FEMALE | | | | | | | | | | | | |
| 1978 | 9.1 | * | * | * | 0.3 | 2.2 | 9.4 | 21.7 | 26.6 | 24.4 | 19.2 | 13.4 |
| 1977 | 9.5 | * | * | * | 0.3 | 2.4 | 10.2 | 22.0 | 27.4 | 27.0 | 18.7 | 13.0 |
| 1976 | 9.7 | * | * | * | 0.3 | 2.3 | 11.2 | 22.8 | 29.0 | 25.6 | 18.9 | 13.4 |
| 1975 | 9.8 | 1.4 | * | * | 0.4 | 2.4 | 11.0 | 23.7 | 30.6 | 24.6 | 17.7 | 12.2 |
| 1974 | 10.5 | * | * | * | 0.3 | 3.0 | 13.1 | 25.8 | 32.0 | 25.9 | 18.1 | 13.8 |
| 1973 | 10.7 | 1.4 | * | * | 0.3 | 2.9 | 13.6 | 26.9 | 32.3 | 25.7 | 18.3 | 13.5 |
| 1972 | 10.4 | * | * | * | 0.3 | 3.3 | 14.0 | 25.7 | 30.3 | 24.9 | 19.2 | 12.7 |
| 1971 | 10.5 | * | * | 0.1 | 0.5 | 3.1 | 14.1 | 26.4 | 30.3 | 24.8 | 19.6 | 15.1 |
| 1970 | 10.6 | * | * | 0.2 | 0.5 | 3.2 | 14.6 | 26.7 | 30.1 | 23.8 | 20.8 | 14.5 |
| 1969 | 10.1 | 1.3 | * | 0.2 | 0.5 | 3.4 | 14.0 | 26.0 | 27.9 | 23.0 | 20.1 | 14.0 |
| 1968 | 10.1 | 1.5 | 0.3 | 0.1 | 0.5 | 3.3 | 13.7 | 25.9 | 27.8 | 23.9 | 19.5 | 18.3 |
| WHITE, BOTH SEXES | | | | | | | | | | | | |
| 1978 | 13.0 | * | * | * | 0.2 | 2.2 | 10.9 | 29.0 | 41.1 | 41.7 | 29.7 | 19.5 |
| 1977 | 13.4 | 0.8 | * | * | 0.2 | 2.3 | 11.8 | 30.0 | 42.9 | 43.2 | 29.0 | 18.3 |
| 1976 | 13.8 | 0.9 | * | * | 0.2 | 2.2 | 12.9 | 31.5 | 45.4 | 42.9 | 28.9 | 19.6 |
| 1975 | 14.1 | 1.2 | * | * | 0.3 | 2.2 | 13.0 | 32.8 | 47.0 | 43.1 | 28.6 | 17.6 |
| 1974 | 14.9 | 1.0 | * | * | 0.3 | 2.6 | 14.4 | 34.9 | 49.1 | 46.6 | 29.2 | 19.5 |
| 1973 | 15.1 | 1.0 | * | 0.1 | 0.2 | 2.6 | 15.0 | 35.6 | 51.1 | 45.2 | 28.7 | 20.6 |
| 1972 | 14.8 | 0.9 | 0.2 | 0.1 | 0.2 | 2.4 | 15.7 | 35.4 | 49.5 | 43.2 | 29.9 | 20.9 |
| 1971 | 14.9 | 0.8 | * | 0.1 | 0.3 | 2.6 | 15.2 | 35.6 | 49.4 | 43.0 | 30.6 | 22.3 |
| 1970 | 14.9 | * | * | 0.1 | 0.3 | 2.4 | 15.2 | 35.7 | 49.1 | 43.1 | 31.8 | 21.6 |
| 1969 | 14.2 | 1.3 | 0.3 | 0.1 | 0.4 | 2.5 | 14.8 | 34.2 | 46.8 | 40.3 | 30.4 | 21.9 |
| 1968 | 14.1 | 1.4 | 0.2 | 0.1 | 0.3 | 2.4 | 14.5 | 34.1 | 45.9 | 41.2 | 30.6 | 23.6 |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| Chronic liver disease and cirrhosis 571 | | | | | | | | | | | | |
| WHITE, MALE | | | | | | | | | | | | |
| 1978 | 17.5 | * | * | * | 0.2 | 3.1 | 15.0 | 39.4 | 58.7 | 64.4 | 46.5 | 31.5 |
| 1977 | 18.0 | * | * | * | 0.3 | 3.4 | 16.0 | 41.3 | 61.5 | 63.8 | 46.0 | 29.2 |
| 1976 | 18.7 | * | * | * | 0.2 | 3.1 | 17.8 | 43.3 | 64.8 | 65.4 | 44.7 | 31.6 |
| 1975 | 19.1 | * | * | * | 0.2 | 3.2 | 18.2 | 44.7 | 66.3 | 67.2 | 45.6 | 29.1 |
| 1974 | 20.2 | * | * | * | 0.4 | 3.4 | 19.1 | 47.7 | 69.4 | 73.3 | 46.8 | 30.5 |
| 1973 | 20.4 | * | * | * | 0.2 | 3.6 | 20.0 | 48.0 | 73.1 | 70.6 | 44.8 | 33.1 |
| 1972 | 20.1 | * | * | * | 0.1 | 3.3 | 21.3 | 48.1 | 71.2 | 66.7 | 45.5 | 36.2 |
| 1971 | 20.0 | * | * | * | 0.3 | 3.5 | 20.0 | 47.9 | 71.2 | 66.0 | 46.9 | 35.9 |
| 1970 | 20.1 | * | * | 0.1 | 0.3 | 3.4 | 19.7 | 47.3 | 70.6 | 67.8 | 48.1 | 33.1 |
| 1969 | 19.2 | * | * | * | 0.3 | 3.3 | 19.2 | 45.3 | 68.2 | 62.1 | 45.2 | 36.4 |
| 1968 | 18.9 | 1.4 | * | 0.1 | 0.2 | 3.0 | 18.8 | 45.1 | 66.2 | 62.5 | 45.9 | 32.6 |
| WHITE, FEMALE | | | | | | | | | | | | |
| 1978 | 8.6 | * | * | * | 0.2 | 1.2 | 6.9 | 19.2 | 25.4 | 24.4 | 19.8 | 14.1 |
| 1977 | 9.0 | * | * | * | 0.2 | 1.3 | 7.6 | 19.4 | 26.4 | 27.5 | 19.0 | 13.3 |
| 1976 | 9.1 | * | * | * | 0.1 | 1.3 | 8.2 | 20.4 | 28.1 | 25.7 | 19.5 | 14.0 |
| 1975 | 9.3 | * | * | * | 0.3 | 1.2 | 8.0 | 21.5 | 29.7 | 24.7 | 18.4 | 12.1 |
| 1974 | 9.9 | * | * | * | 0.2 | 1.7 | 9.9 | 22.9 | 31.0 | 26.3 | 18.6 | 14.2 |
| 1973 | 10.0 | * | * | * | 0.2 | 1.5 | 10.1 | 24.0 | 31.4 | 25.7 | 18.7 | 14.5 |
| 1972 | 9.8 | * | * | * | 0.2 | 1.6 | 10.3 | 23.5 | 30.0 | 25.1 | 20.1 | 13.1 |
| 1971 | 10.0 | * | * | 0.1 | 0.4 | 1.7 | 10.6 | 24.1 | 29.9 | 25.3 | 20.2 | 15.4 |
| 1970 | 10.0 | * | * | 0.2 | 0.4 | 1.6 | 10.8 | 24.7 | 29.6 | 24.1 | 21.2 | 15.4 |
| 1969 | 9.5 | * | * | 0.2 | 0.4 | 1.8 | 10.6 | 23.7 | 27.5 | 23.3 | 20.6 | 14.2 |
| 1968 | 9.6 | * | * | * | 0.4 | 1.8 | 10.4 | 23.8 | 27.6 | 24.5 | 20.2 | 18.8 |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| Chronic liver disease and cirrhosis 571 | | | | | | | | | | | | |
| ALL OTHER, BOTH SEXES | | | | | | | | | | | | |
| 1978 | 17.2 | * | * | * | 0.7 | 11.4 | 37.8 | 56.7 | 56.4 | 38.7 | 19.1 | 13.1 |
| 1977 | 18.3 | * | * | * | 1.0 | 13.0 | 39.2 | 64.0 | 59.5 | 35.1 | 21.9 | * |
| 1976 | 18.7 | * | * | * | 1.0 | 12.9 | 43.9 | 63.1 | 60.4 | 38.2 | 16.1 | * |
| 1975 | 18.7 | * | * | * | 1.0 | 14.1 | 42.3 | 62.4 | 61.0 | 38.9 | 18.5 | 16.4 |
| 1974 | 20.2 | * | * | * | 1.0 | 15.5 | 47.8 | 70.0 | 62.2 | 36.9 | 19.6 | * |
| 1973 | 20.5 | * | * | * | 1.3 | 15.6 | 50.7 | 68.8 | 61.1 | 38.3 | 19.9 | 15.7 |
| 1972 | 20.5 | * | * | * | 1.2 | 17.3 | 52.9 | 68.7 | 55.3 | 41.0 | 17.3 | 16.3 |
| 1971 | 19.0 | * | * | * | 1.3 | 15.8 | 48.9 | 63.5 | 52.2 | 34.3 | 18.9 | * |
| 1970 | 19.4 | * | * | * | 0.9 | 18.1 | 51.9 | 61.7 | 51.8 | 32.8 | 19.5 | * |
| 1969 | 19.1 | * | * | * | 1.6 | 16.9 | 50.5 | 59.6 | 51.8 | 34.3 | 18.2 | * |
| 1968 | 18.2 | * | * | * | 1.3 | 16.9 | 45.4 | 61.6 | 47.8 | 31.2 | 16.9 | * |
| ALL OTHER, MALE | | | | | | | | | | | | |
| 1978 | 22.9 | * | * | * | 0.8 | 15.4 | 53.3 | 77.1 | 80.4 | 56.6 | 28.9 | * |
| 1977 | 24.5 | * | * | * | 0.9 | 17.4 | 53.7 | 90.6 | 86.9 | 51.3 | 31.4 | * |
| 1976 | 25.1 | * | * | * | 1.0 | 17.8 | 59.8 | 89.0 | 88.9 | 54.9 | 21.4 | * |
| 1975 | 25.0 | * | * | * | 1.1 | 19.3 | 56.2 | 88.2 | 87.7 | 57.0 | 30.6 | * |
| 1974 | 26.5 | * | * | * | 0.9 | 20.3 | 64.1 | 95.4 | 87.1 | 55.3 | 29.7 | * |
| 1973 | 26.4 | * | * | * | 1.5 | 20.3 | 67.5 | 90.0 | 85.4 | 54.5 | 29.5 | * |
| 1972 | 27.3 | * | * | * | 1.3 | 20.7 | 69.5 | 97.9 | 81.1 | 63.3 | 29.5 | * |
| 1971 | 24.3 | * | * | * | 1.5 | 20.0 | 62.8 | 83.3 | 72.6 | 51.8 | 28.5 | * |
| 1970 | 24.6 | * | * | * | 1.0 | 22.7 | 65.6 | 82.6 | 71.3 | 47.5 | 24.9 | * |
| 1969 | 24.2 | * | * | * | 1.9 | 20.0 | 65.3 | 75.4 | 73.8 | 52.1 | 23.9 | * |
| 1968 | 23.0 | * | * | * | 1.2 | 21.0 | 54.4 | 81.1 | 67.9 | 46.9 | 25.8 | * |
| ALL OTHER, FEMALE | | | | | | | | | | | | |
| 1978 | 12.1 | * | * | * | 0.7 | 7.9 | 24.8 | 39.8 | 36.4 | 24.9 | 12.3 | * |
| 1977 | 12.7 | * | * | * | 1.1 | 9.3 | 26.9 | 41.8 | 36.6 | 22.4 | 15.4 | * |
| 1976 | 12.9 | * | * | * | 0.9 | 8.6 | 30.6 | 41.4 | 36.6 | 25.0 | 12.5 | * |
| 1975 | 13.0 | * | * | * | 0.9 | 9.6 | 30.6 | 40.6 | 38.5 | 24.5 | 10.2 | * |
| 1974 | 14.5 | * | * | * | 1.0 | 11.3 | 34.2 | 48.5 | 41.1 | 22.2 | 12.7 | * |
| 1973 | 15.0 | * | * | * | 1.0 | 11.5 | 36.7 | 50.7 | 40.4 | 25.2 | 13.2 | * |
| 1972 | 14.3 | * | * | * | 1.1 | 14.5 | 39.0 | 43.7 | 33.0 | 22.9 | 8.6 | * |
| 1971 | 14.1 | * | * | * | 1.1 | 12.2 | 37.4 | 46.2 | 34.3 | 19.9 | 11.8 | * |
| 1970 | 14.6 | * | * | * | 0.9 | 14.2 | 40.5 | 43.4 | 34.6 | 20.7 | 15.4 | * |
| 1969 | 14.3 | * | * | * | 1.2 | 14.2 | 38.1 | 45.8 | 32.3 | 19.6 | 13.8 | * |
| 1968 | 13.9 | * | * | * | 1.3 | 13.5 | 37.8 | 44.3 | 29.7 | 18.2 | 10.0 | * |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| Chronic liver disease and cirrhosis 571 | | | | | | | | | | | | |
| BLACK, BOTH SEXES | | | | | | | | | | | | |
| 1978 | 17.9 | * | * | * | 0.7 | 12.3 | 40.1 | 60.3 | 58.3 | 39.2 | 16.6 | 13.3 |
| 1977 | 19.2 | * | * | * | 1.0 | 13.4 | 42.1 | 68.6 | 62.1 | 36.7 | 20.8 | * |
| 1976 | 19.4 | * | * | * | 0.9 | 13.5 | 46.1 | 67.0 | 62.8 | 38.2 | 14.8 | * |
| 1975 | 19.5 | * | * | * | 1.0 | 14.9 | 44.8 | 66.1 | 62.1 | 39.8 | 16.9 | 16.2 |
| 1974 | 20.9 | * | * | * | 0.9 | 16.3 | 50.6 | 73.6 | 63.7 | 38.2 | 16.2 | * |
| 1973 | 21.1 | * | * | * | 1.4 | 16.1 | 53.5 | 71.7 | 62.6 | 38.5 | 18.6 | * |
| 1972 | 21.3 | * | * | * | 1.3 | 18.6 | 56.0 | 72.6 | 55.3 | 40.1 | 17.2 | * |
| 1971 | 19.3 | * | * | * | 1.3 | 16.4 | 51.5 | 65.2 | 51.3 | 33.5 | 18.7 | * |
| 1970 | 20.1 | * | * | * | 1.0 | 19.4 | 54.6 | 64.3 | 53.5 | 32.3 | 19.1 | * |
| 1969 | 19.3 | * | * | * | 1.5 | 17.2 | 52.5 | 61.6 | 51.4 | 34.3 | 16.3 | * |
| 1968 | 18.5 | * | * | * | 1.3 | 17.7 | 47.0 | 63.3 | 47.5 | 30.3 | 15.5 | * |
| BLACK, MALE | | | | | | | | | | | | |
| 1978 | 24.0 | * | * | * | 0.7 | 16.8 | 57.0 | 82.5 | 84.4 | 58.3 | 24.7 | * |
| 1977 | 26.0 | * | * | * | 0.8 | 18.4 | 58.4 | 98.0 | 91.7 | 54.7 | 30.2 | * |
| 1976 | 26.4 | * | * | * | 1.0 | 19.0 | 63.8 | 95.5 | 94.3 | 56.0 | 19.7 | * |
| 1975 | 26.3 | * | * | * | 1.1 | 21.3 | 60.7 | 93.6 | 90.4 | 59.6 | 28.4 | * |
| 1974 | 27.7 | * | * | * | 0.8 | 21.5 | 68.7 | 102.2 | 90.3 | 58.2 | 25.8 | * |
| 1973 | 27.4 | * | * | * | 1.7 | 21.1 | 72.1 | 94.5 | 88.1 | 55.4 | 27.0 | * |
| 1972 | 28.3 | * | * | * | 1.4 | 22.5 | 75.0 | 104.5 | 80.7 | 61.0 | 29.4 | * |
| 1971 | 25.0 | * | * | * | 1.6 | 21.6 | 67.1 | 86.7 | 72.5 | 51.0 | 29.6 | * |
| 1970 | 25.8 | * | * | * | 1.1 | 24.8 | 69.7 | 87.4 | 75.4 | 47.5 | 25.5 | * |
| 1969 | 24.8 | * | * | * | 1.9 | 21.2 | 68.9 | 78.9 | 74.3 | 52.7 | 20.4 | * |
| 1968 | 23.7 | * | * | * | 1.2 | 22.6 | 57.7 | 85.0 | 68.6 | 45.9 | 23.6 | * |
| BLACK, FEMALE | | | | | | | | | | | | |
| 1978 | 12.5 | * | * | * | 0.7 | 8.4 | 26.1 | 42.0 | 37.1 | 25.0 | 11.2 | * |
| 1977 | 13.2 | * | * | * | 1.1 | 9.1 | 28.5 | 44.2 | 37.9 | 23.2 | 14.7 | * |
| 1976 | 13.2 | * | * | * | 0.8 | 8.8 | 31.5 | 43.3 | 36.8 | 24.8 | 11.5 | * |
| 1975 | 13.3 | * | * | * | 0.9 | 9.3 | 31.7 | 43.0 | 38.7 | 24.8 | 9.1 | * |
| 1974 | 14.8 | * | * | * | 1.1 | 11.8 | 35.4 | 49.7 | 41.6 | 22.8 | 9.7 | * |
| 1973 | 15.4 | * | * | * | 1.0 | 11.9 | 38.1 | 52.2 | 41.4 | 25.4 | 12.8 | * |
| 1972 | 14.8 | * | * | * | 1.2 | 15.2 | 40.3 | 45.4 | 34.0 | 23.9 | 8.6 | * |
| 1971 | 14.1 | * | * | * | 1.1 | 12.0 | 38.5 | 46.7 | 33.3 | 19.7 | 10.8 | * |
| 1970 | 14.9 | * | * | * | 1.0 | 14.8 | 42.0 | 44.5 | 34.9 | 20.3 | 14.4 | * |
| 1969 | 14.4 | * | * | * | 1.1 | 13.9 | 38.9 | 46.5 | 31.8 | 19.6 | 13.2 | * |
| 1968 | 13.8 | * | * | * | 1.3 | 13.7 | 38.0 | 44.5 | 29.2 | 17.8 | 9.5 | * |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| CHOLELITHIASIS, CHOLECYSTITIS AND CHOLANGITIS (574,575) | | | | | | | | | | | | |
| BOTH SEXES | | | | | | | | | | | | |
| 1978 | 1.3 | * | * | * | * | * | 0.2 | 0.6 | 1.3 | 4.8 | 13.5 | 35.6 |
| 1977 | 1.3 | * | * | * | * | 0.1 | 0.2 | 0.6 | 1.4 | 4.9 | 13.1 | 39.3 |
| 1976 | 1.4 | * | * | * | * | 0.1 | 0.2 | 0.6 | 1.5 | 4.8 | 14.7 | 36.7 |
| 1975 | 1.4 | * | * | * | * | 0.1 | 0.2 | 0.5 | 1.7 | 5.2 | 15.2 | 36.7 |
| 1974 | 1.5 | * | * | * | * | 0.1 | 0.3 | 0.7 | 1.8 | 5.7 | 16.3 | 39.3 |
| 1973 | 1.6 | * | * | * | * | 0.1 | 0.3 | 0.7 | 2.0 | 6.2 | 18.2 | 44.7 |
| 1972 | 1.8 | * | * | * | * | 0.1 | 0.3 | 0.7 | 2.2 | 7.1 | 20.2 | 47.3 |
| 1971 | 1.8 | * | * | * | * | 0.1 | 0.3 | 0.8 | 2.4 | 7.9 | 20.2 | 46.1 |
| 1970 | 2.0 | * | * | * | * | 0.2 | 0.3 | 0.9 | 2.5 | 8.2 | 23.8 | 46.3 |
| 1969 | 2.1 | * | * | * | * | 0.2 | 0.3 | 1.0 | 2.9 | 9.1 | 25.2 | 54.9 |
| 1968 | 2.2 | * | * | * | 0.1 | 0.1 | 0.4 | 1.0 | 3.1 | 9.7 | 26.0 | 56.9 |
| MALE | | | | | | | | | | | | |
| 1978 | 1.2 | * | * | * | * | * | 0.2 | 0.7 | 1.5 | 6.1 | 14.7 | 37.7 |
| 1977 | 1.2 | * | * | * | * | * | * | 0.6 | 1.5 | 5.9 | 15.7 | 39.1 |
| 1976 | 1.2 | * | * | * | * | * | 0.2 | 0.6 | 1.6 | 6.2 | 17.0 | 38.0 |
| 1975 | 1.2 | * | * | * | * | * | * | 0.5 | 1.8 | 6.2 | 17.4 | 36.5 |
| 1974 | 1.3 | * | * | * | * | * | 0.2 | 0.7 | 2.0 | 6.6 | 18.9 | 37.0 |
| 1973 | 1.5 | * | * | * | * | * | 0.2 | 0.7 | 2.2 | 7.1 | 21.3 | 46.3 |
| 1972 | 1.6 | * | * | * | * | * | 0.2 | 0.8 | 2.4 | 8.4 | 22.7 | 42.9 |
| 1971 | 1.6 | * | * | * | * | * | 0.3 | 0.8 | 2.4 | 8.9 | 22.2 | 47.1 |
| 1970 | 1.7 | * | * | * | * | * | 0.2 | 0.9 | 2.6 | 8.9 | 25.5 | 44.4 |
| 1969 | 1.9 | * | * | * | * | * | 0.3 | 0.9 | 2.8 | 9.7 | 28.2 | 51.6 |
| 1968 | 2.0 | * | * | * | * | * | 0.3 | 1.2 | 3.3 | 9.8 | 27.7 | 55.2 |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| CHOLELITHIASIS, CHOLECYSTITIS AND CHOLANGITIS (574,575) | | | | | | | | | | | | |
| FEMALE | | | | | | | | | | | | |
| 1978 | 1.4 | * | * | * | * | * | * | 0.5 | 1.1 | 3.8 | 12.7 | 34.6 |
| 1977 | 1.5 | * | * | * | * | * | 0.3 | 0.5 | 1.2 | 4.1 | 11.5 | 39.4 |
| 1976 | 1.5 | * | * | * | * | * | 0.2 | 0.5 | 1.5 | 3.6 | 13.4 | 36.1 |
| 1975 | 1.5 | * | * | * | * | * | 0.3 | 0.6 | 1.6 | 4.4 | 13.9 | 36.8 |
| 1974 | 1.6 | * | * | * | * | 0.1 | 0.3 | 0.7 | 1.7 | 5.0 | 14.7 | 40.5 |
| 1973 | 1.7 | * | * | * | * | * | 0.3 | 0.7 | 1.8 | 5.6 | 16.3 | 44.0 |
| 1972 | 1.9 | * | * | * | * | 0.1 | 0.4 | 0.7 | 1.9 | 6.1 | 18.6 | 49.6 |
| 1971 | 2.0 | * | * | * | * | * | 0.3 | 0.8 | 2.4 | 7.1 | 18.9 | 45.6 |
| 1970 | 2.2 | * | * | * | * | 0.3 | 0.4 | 0.8 | 2.3 | 7.7 | 22.6 | 47.3 |
| 1969 | 2.3 | * | * | * | * | 0.2 | 0.4 | 1.1 | 3.0 | 8.6 | 23.1 | 56.6 |
| 1968 | 2.4 | * | * | * | * | 0.2 | 0.5 | 0.9 | 3.0 | 9.6 | 24.9 | 57.9 |
| WHITE, BOTH SEXES | | | | | | | | | | | | |
| 1978 | 1.4 | * | * | * | * | * | 0.2 | 0.6 | 1.2 | 4.7 | 14.0 | 36.8 |
| 1977 | 1.4 | * | * | * | * | * | 0.2 | 0.5 | 1.4 | 4.9 | 13.3 | 40.6 |
| 1976 | 1.4 | * | * | * | * | * | 0.2 | 0.5 | 1.5 | 4.8 | 15.3 | 37.8 |
| 1975 | 1.5 | * | * | * | * | * | 0.2 | 0.5 | 1.7 | 5.2 | 15.6 | 38.1 |
| 1974 | 1.6 | * | * | * | * | 0.1 | 0.2 | 0.7 | 1.8 | 5.7 | 16.8 | 40.7 |
| 1973 | 1.7 | * | * | * | * | 0.1 | 0.2 | 0.7 | 1.9 | 6.3 | 18.6 | 46.6 |
| 1972 | 1.9 | * | * | * | * | 0.1 | 0.2 | 0.8 | 2.1 | 7.2 | 20.8 | 49.9 |
| 1971 | 1.9 | * | * | * | * | 0.1 | 0.3 | 0.7 | 2.3 | 8.0 | 21.0 | 47.9 |
| 1970 | 2.1 | * | * | * | * | 0.1 | 0.3 | 0.9 | 2.4 | 8.6 | 24.5 | 49.1 |
| 1969 | 2.3 | * | * | * | * | 0.1 | 0.3 | 0.9 | 2.9 | 9.3 | 26.2 | 57.5 |
| 1968 | 2.3 | * | * | * | * | 0.1 | 0.4 | 1.0 | 3.2 | 9.8 | 26.8 | 59.6 |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| CHOLELITHIASIS, CHOLECYSTITIS AND CHOLANGITIS (574,575) | | | | | | | | | | | | |
| WHITE, MALE | | | | | | | | | | | | |
| 1978 | 1.3 | * | * | * | * | * | 0.2 | 0.7 | 1.5 | 6.2 | 15.4 | 39.4 |
| 1977 | 1.3 | * | * | * | * | * | * | 0.6 | 1.5 | 6.1 | 16.4 | 39.6 |
| 1976 | 1.3 | * | * | * | * | * | 0.2 | 0.6 | 1.6 | 6.4 | 17.7 | 39.4 |
| 1975 | 1.3 | * | * | * | * | * | * | 0.5 | 1.8 | 6.5 | 18.0 | 37.4 |
| 1974 | 1.4 | * | * | * | * | * | * | 0.7 | 1.9 | 6.8 | 19.6 | 38.3 |
| 1973 | 1.6 | * | * | * | * | * | * | 0.7 | 2.3 | 7.2 | 22.0 | 48.0 |
| 1972 | 1.7 | * | * | * | * | * | * | 0.8 | 2.3 | 8.8 | 23.7 | 45.5 |
| 1971 | 1.7 | * | * | * | * | * | 0.3 | 0.8 | 2.4 | 9.2 | 23.3 | 49.9 |
| 1970 | 1.9 | * | * | * | * | * | 0.3 | 0.9 | 2.7 | 9.4 | 26.3 | 48.3 |
| 1969 | 2.0 | * | * | * | * | * | 0.3 | 0.9 | 3.0 | 10.1 | 29.6 | 54.5 |
| 1968 | 2.1 | * | * | * | * | * | 0.3 | 1.1 | 3.4 | 10.0 | 28.9 | 58.4 |
| WHITE, FEMALE | | | | | | | | | | | | |
| 1978 | 1.5 | * | * | * | * | * | * | 0.5 | 1.0 | 3.6 | 13.3 | 35.6 |
| 1977 | 1.5 | * | * | * | * | * | 0.2 | 0.4 | 1.2 | 4.0 | 11.5 | 41.0 |
| 1976 | 1.5 | * | * | * | * | * | 0.3 | 0.4 | 1.3 | 3.6 | 13.8 | 37.1 |
| 1975 | 1.6 | * | * | * | * | * | 0.3 | 0.5 | 1.6 | 4.3 | 14.2 | 38.4 |
| 1974 | 1.7 | * | * | * | * | * | 0.3 | 0.7 | 1.7 | 4.9 | 15.1 | 42.0 |
| 1973 | 1.8 | * | * | * | * | * | 0.2 | 0.7 | 1.6 | 5.6 | 16.4 | 46.0 |
| 1972 | 2.0 | * | * | * | * | * | 0.3 | 0.7 | 1.9 | 6.0 | 19.0 | 52.1 |
| 1971 | 2.1 | * | * | * | * | * | 0.3 | 0.7 | 2.2 | 7.1 | 19.4 | 46.9 |
| 1970 | 2.3 | * | * | * | * | 0.2 | 0.4 | 0.8 | 2.3 | 7.9 | 23.3 | 49.6 |
| 1969 | 2.5 | * | * | * | * | 0.2 | 0.3 | 1.0 | 2.9 | 8.8 | 23.9 | 59.1 |
| 1968 | 2.6 | * | * | * | * | * | 0.5 | 0.8 | 2.9 | 9.6 | 25.4 | 60.4 |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| CHOLELITHIASIS, CHOLECYSTITIS AND CHOLANGITIS (574,575) | | | | | | | | | | | | |
| ALL OTHER, BOTH SEXES | | | | | | | | | | | | |
| 1978 | 0.8 | * | * | * | * | * | * | 1.0 | 1.7 | 5.5 | 7.8 | 21.4 |
| 1977 | 0.9 | * | * | * | * | * | * | 1.1 | 1.5 | 4.6 | 10.6 | 25.0 |
| 1976 | 0.8 | * | * | * | * | * | * | 0.8 | 2.1 | 4.4 | 9.0 | 24.2 |
| 1975 | 0.8 | * | * | * | * | * | * | 0.9 | 1.7 | 4.7 | 10.6 | 21.2 |
| 1974 | 0.9 | * | * | * | * | * | * | 1.1 | 1.9 | 5.6 | 11.2 | 23.0 |
| 1973 | 1.0 | * | * | * | * | * | * | 1.0 | 2.5 | 6.0 | 14.0 | 22.8 |
| 1972 | 0.9 | * | * | * | * | * | * | * | 2.6 | 5.9 | 12.7 | 17.9 |
| 1971 | 1.0 | * | * | * | * | * | * | 1.1 | 3.1 | 7.2 | 11.1 | 25.8 |
| 1970 | 0.9 | * | * | * | * | * | * | 0.9 | 2.6 | 4.9 | 14.5 | 17.2 |
| 1969 | 1.1 | * | * | * | * | * | * | 1.7 | 2.5 | 6.5 | 12.2 | 25.7 |
| 1968 | 1.2 | * | * | * | * | * | * | 1.6 | 2.9 | 8.4 | 15.7 | 26.5 |
| ALL OTHER, MALE | | | | | | | | | | | | |
| 1978 | 0.7 | * | * | * | * | * | * | * | * | 5.7 | 8.6 | * |
| 1977 | 0.7 | * | * | * | * | * | * | * | * | 3.9 | 8.8 | * |
| 1976 | 0.6 | * | * | * | * | * | * | * | * | 4.2 | 9.7 | * |
| 1975 | 0.6 | * | * | * | * | * | * | * | * | * | 10.7 | * |
| 1974 | 0.8 | * | * | * | * | * | * | * | 2.5 | 4.8 | 11.8 | * |
| 1973 | 0.8 | * | * | * | * | * | * | * | * | 6.4 | 13.8 | * |
| 1972 | 0.8 | * | * | * | * | * | * | * | 3.1 | 4.1 | 11.1 | * |
| 1971 | 0.8 | * | * | * | * | * | * | * | * | 6.9 | 9.7 | * |
| 1970 | 0.7 | * | * | * | * | * | * | * | * | 4.2 | 16.1 | * |
| 1969 | 0.8 | * | * | * | * | * | * | * | * | 5.7 | 11.2 | * |
| 1968 | 1.0 | * | * | * | * | * | * | * | * | 7.2 | 12.9 | * |
| ALL OTHER, FEMALE | | | | | | | | | | | | |
| 1978 | 0.9 | * | * | * | * | * | * | * | 2.0 | 5.4 | 7.2 | 22.0 |
| 1977 | 1.0 | * | * | * | * | * | * | * | * | 5.2 | 11.7 | 20.2 |
| 1976 | 1.0 | * | * | * | * | * | * | * | 2.9 | 4.5 | 8.5 | 24.5 |
| 1975 | 1.0 | * | * | * | * | * | * | * | * | 5.8 | 10.5 | * |
| 1974 | 1.0 | * | * | * | * | * | * | * | * | 6.2 | 10.8 | * |
| 1973 | 1.2 | * | * | * | * | * | * | * | 3.4 | 5.6 | 14.2 | * |
| 1972 | 1.1 | * | * | * | * | * | * | * | 2.2 | 7.3 | 14.0 | * |
| 1971 | 1.3 | * | * | * | * | * | * | * | 3.9 | 7.5 | 12.1 | 30.1 |
| 1970 | 1.1 | * | * | * | * | * | * | * | 2.9 | 5.5 | 13.4 | * |
| 1969 | 1.3 | * | * | * | * | * | * | 2.4 | 3.7 | 7.1 | 13.0 | * |
| 1968 | 1.5 | * | * | * | * | * | * | 1.7 | 3.6 | 9.4 | 18.0 | * |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| CHOLELITHIASIS, CHOLECYSTITIS AND CHOLANGITIS (574,575) | | | | | | | | | | | | |
| BLACK, BOTH SEXES | | | | | | | | | | | | |
| 1978 | 0.8 | * | * | * | * | * | * | 1.0 | 1.7 | 5.6 | 6.4 | 18.7 |
| 1977 | 0.8 | * | * | * | * | * | * | 1.1 | 1.4 | 4.6 | 9.3 | 22.4 |
| 1976 | 0.7 | * | * | * | * | * | * | * | 2.2 | 3.6 | 7.5 | 22.8 |
| 1975 | 0.7 | * | * | * | * | * | * | 0.9 | 1.6 | 4.6 | 7.5 | 17.7 |
| 1974 | 0.8 | * | * | * | * | * | * | 1.0 | 1.9 | 5.3 | 8.9 | 17.4 |
| 1973 | 0.9 | * | * | * | * | * | * | 1.0 | 2.4 | 5.2 | 11.6 | 20.2 |
| 1972 | 0.9 | * | * | * | * | * | * | * | 2.3 | 5.4 | 11.3 | * |
| 1971 | 0.9 | * | * | * | * | * | * | 1.2 | 2.9 | 6.3 | 7.0 | 20.6 |
| 1970 | 0.8 | * | * | * | * | * | * | * | 2.4 | 4.2 | 11.0 | * |
| 1969 | 0.9 | * | * | * | * | * | * | 1.7 | 2.0 | 5.8 | 10.4 | 22.7 |
| 1968 | 1.1 | * | * | * | * | * | * | 1.6 | 2.8 | 7.5 | 13.4 | * |
| BLACK, MALE | | | | | | | | | | | | |
| 1978 | 0.7 | * | * | * | * | * | * | * | * | 5.8 | * | * |
| 1977 | 0.7 | * | * | * | * | * | * | * | * | 3.7 | * | * |
| 1976 | 0.6 | * | * | * | * | * | * | * | * | * | * | * |
| 1975 | 0.5 | * | * | * | * | * | * | * | * | * | * | * |
| 1974 | 0.7 | * | * | * | * | * | * | * | * | 4.5 | * | * |
| 1973 | 0.7 | * | * | * | * | * | * | * | * | 5.2 | 9.8 | * |
| 1972 | 0.7 | * | * | * | * | * | * | * | 2.6 | * | * | * |
| 1971 | 0.6 | * | * | * | * | * | * | * | * | 5.6 | * | * |
| 1970 | 0.5 | * | * | * | * | * | * | * | * | * | * | * |
| 1969 | 0.6 | * | * | * | * | * | * | * | * | 5.1 | * | * |
| 1968 | 0.8 | * | * | * | * | * | * | * | * | 5.5 | * | * |
| BLACK, FEMALE | | | | | | | | | | | | |
| 1978 | 0.9 | * | * | * | * | * | * | * | 2.0 | 5.5 | 6.4 | * |
| 1977 | 1.0 | * | * | * | * | * | * | * | * | 5.3 | 10.5 | * |
| 1976 | 0.9 | * | * | * | * | * | * | * | 3.0 | 3.5 | 7.4 | 23.6 |
| 1975 | 0.9 | * | * | * | * | * | * | * | * | 5.7 | 8.2 | * |
| 1974 | 0.9 | * | * | * | * | * | * | * | * | 5.9 | 8.7 | * |
| 1973 | 1.1 | * | * | * | * | * | * | * | 3.4 | 5.2 | 12.8 | * |
| 1972 | 1.0 | * | * | * | * | * | * | * | * | 7.0 | 12.9 | * |
| 1971 | 1.2 | * | * | * | * | * | * | * | 3.7 | 6.9 | 9.3 | * |
| 1970 | 1.0 | * | * | * | * | * | * | * | 2.6 | 5.2 | 11.7 | * |
| 1969 | 1.3 | * | * | * | * | * | * | 2.5 | 2.9 | 6.5 | 12.3 | * |
| 1968 | 1.4 | * | * | * | * | * | * | * | 3.4 | 9.1 | 16.2 | * |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|--|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| Nephritis, nephrotic syndrome, & nephrosis 580-584 | | | | | | | | | | | | |
| BOTH SEXES | | | | | | | | | | | | |
| 1978 | 4.0 | 1.9 | 0.2 | 0.1 | 0.2 | 0.6 | 1.1 | 2.8 | 6.5 | 15.1 | 34.0 | 66.0 |
| 1977 | 3.9 | 1.9 | 0.2 | 0.1 | 0.3 | 0.6 | 1.2 | 2.8 | 6.4 | 14.8 | 32.8 | 64.4 |
| 1976 | 3.9 | 1.1 | 0.2 | 0.1 | 0.3 | 0.7 | 1.4 | 3.1 | 6.5 | 15.1 | 32.3 | 67.0 |
| 1975 | 3.7 | 0.6 | * | 0.2 | 0.3 | 0.6 | 1.6 | 3.0 | 6.9 | 15.0 | 29.4 | 58.6 |
| 1974 | 3.8 | 0.7 | * | 0.1 | 0.4 | 0.7 | 1.7 | 3.4 | 7.2 | 14.5 | 30.3 | 57.0 |
| 1973 | 3.9 | 0.6 | 0.2 | 0.2 | 0.5 | 0.9 | 1.9 | 3.6 | 7.5 | 15.2 | 30.1 | 63.0 |
| 1972 | 4.1 | 0.7 | 0.1 | 0.2 | 0.5 | 0.9 | 2.0 | 4.4 | 8.0 | 15.6 | 29.9 | 63.9 |
| 1971 | 4.1 | 1.1 | 0.2 | 0.3 | 0.6 | 1.2 | 2.3 | 4.2 | 8.3 | 14.8 | 28.5 | 63.7 |
| 1970 | 4.4 | 1.0 | 0.3 | 0.3 | 0.8 | 1.4 | 2.7 | 4.6 | 8.5 | 15.9 | 30.9 | 58.4 |
| 1969 | 4.7 | 1.1 | 0.3 | 0.4 | 1.0 | 1.6 | 3.0 | 4.8 | 9.5 | 16.0 | 33.3 | 68.7 |
| 1968 | 4.7 | 0.8 | 0.2 | 0.4 | 1.0 | 1.5 | 3.4 | 5.2 | 9.2 | 15.8 | 32.1 | 74.5 |
| MALE | | | | | | | | | | | | |
| 1978 | 4.4 | 2.2 | * | * | 0.2 | 0.7 | 1.3 | 3.2 | 7.3 | 19.5 | 50.0 | 100.0 |
| 1977 | 4.3 | 2.3 | * | * | 0.3 | 0.7 | 1.4 | 3.0 | 7.3 | 18.8 | 46.8 | 103.0 |
| 1976 | 4.3 | 1.4 | * | 0.1 | 0.4 | 0.8 | 1.4 | 3.7 | 7.7 | 18.7 | 45.2 | 98.8 |
| 1975 | 4.1 | * | * | 0.1 | 0.4 | 0.8 | 1.8 | 3.2 | 8.5 | 19.3 | 40.0 | 82.3 |
| 1974 | 4.2 | * | * | 0.1 | 0.5 | 0.8 | 1.9 | 3.7 | 8.1 | 18.5 | 42.9 | 83.0 |
| 1973 | 4.3 | * | * | 0.2 | 0.6 | 1.1 | 2.0 | 4.1 | 8.3 | 19.7 | 41.3 | 89.8 |
| 1972 | 4.6 | * | * | 0.2 | 0.6 | 1.0 | 2.3 | 4.9 | 9.5 | 20.1 | 41.6 | 91.6 |
| 1971 | 4.5 | 1.2 | * | 0.3 | 0.7 | 1.3 | 2.5 | 4.6 | 10.2 | 18.1 | 37.2 | 88.6 |
| 1970 | 4.9 | 1.2 | 0.3 | 0.3 | 0.9 | 1.7 | 3.3 | 5.4 | 10.2 | 20.1 | 41.0 | 77.8 |
| 1969 | 5.2 | 1.3 | 0.3 | 0.3 | 1.1 | 2.0 | 3.5 | 5.8 | 11.5 | 19.2 | 43.6 | 86.3 |
| 1968 | 5.3 | * | * | 0.4 | 1.2 | 1.7 | 3.8 | 6.4 | 10.9 | 20.6 | 41.8 | 95.2 |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|--|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| Nephritis, nephrotic syndrome, & nephrosis 580-584 | | | | | | | | | | | | |
| FEMALE | | | | | | | | | | | | |
| 1978 | 3.6 | 1.5 | * | * | 0.2 | 0.5 | 0.9 | 2.5 | 5.8 | 11.7 | 24.4 | 50.6 |
| 1977 | 3.5 | 1.6 | * | * | 0.2 | 0.4 | 1.1 | 2.5 | 5.6 | 11.8 | 24.4 | 46.7 |
| 1976 | 3.6 | * | * | 0.1 | 0.3 | 0.5 | 1.3 | 2.6 | 5.4 | 12.4 | 24.5 | 52.1 |
| 1975 | 3.4 | * | * | 0.2 | 0.2 | 0.4 | 1.3 | 2.8 | 5.5 | 11.7 | 23.0 | 47.2 |
| 1974 | 3.4 | * | * | 0.2 | 0.3 | 0.5 | 1.5 | 3.1 | 6.5 | 11.4 | 22.5 | 44.2 |
| 1973 | 3.6 | * | * | 0.2 | 0.3 | 0.8 | 1.7 | 3.2 | 6.8 | 11.9 | 23.1 | 49.5 |
| 1972 | 3.6 | * | * | 0.2 | 0.4 | 0.8 | 1.7 | 3.9 | 6.7 | 12.1 | 22.5 | 49.8 |
| 1971 | 3.7 | * | * | 0.3 | 0.6 | 1.0 | 2.0 | 3.8 | 6.6 | 12.2 | 22.8 | 50.7 |
| 1970 | 3.9 | * | * | 0.3 | 0.7 | 1.2 | 2.2 | 3.9 | 7.0 | 12.6 | 24.2 | 47.5 |
| 1969 | 4.2 | * | 0.4 | 0.4 | 0.9 | 1.2 | 2.6 | 3.9 | 7.6 | 13.5 | 26.4 | 59.2 |
| 1968 | 4.1 | * | 0.3 | 0.3 | 0.8 | 1.3 | 2.9 | 4.1 | 7.7 | 12.0 | 25.4 | 63.1 |
| WHITE, BOTH SEXES | | | | | | | | | | | | |
| 1978 | 3.4 | 1.6 | * | 0.1 | 0.2 | 0.4 | 0.6 | 1.9 | 4.6 | 12.3 | 30.0 | 61.7 |
| 1977 | 3.3 | 1.4 | * | 0.1 | 0.2 | 0.4 | 0.8 | 1.8 | 4.4 | 12.0 | 29.2 | 61.7 |
| 1976 | 3.3 | 1.0 | * | 0.1 | 0.3 | 0.4 | 0.9 | 2.0 | 4.8 | 11.8 | 28.6 | 63.2 |
| 1975 | 3.2 | * | * | 0.1 | 0.2 | 0.4 | 1.0 | 2.0 | 5.0 | 12.1 | 27.0 | 56.7 |
| 1974 | 3.2 | * | * | 0.1 | 0.3 | 0.4 | 1.1 | 2.2 | 5.3 | 11.7 | 27.8 | 54.0 |
| 1973 | 3.3 | * | * | 0.1 | 0.4 | 0.7 | 1.2 | 2.4 | 5.5 | 11.9 | 27.5 | 58.3 |
| 1972 | 3.4 | 0.7 | * | 0.2 | 0.4 | 0.6 | 1.1 | 3.1 | 6.0 | 12.3 | 27.8 | 60.6 |
| 1971 | 3.5 | 1.0 | 0.2 | 0.2 | 0.5 | 0.7 | 1.5 | 3.1 | 6.2 | 12.4 | 26.3 | 59.5 |
| 1970 | 3.7 | 0.9 | 0.3 | 0.3 | 0.6 | 1.0 | 1.7 | 3.3 | 6.6 | 13.3 | 28.5 | 55.5 |
| 1969 | 4.0 | 1.0 | 0.3 | 0.3 | 0.8 | 1.2 | 2.0 | 3.4 | 7.3 | 13.6 | 31.2 | 66.1 |
| 1968 | 4.0 | 0.7 | 0.2 | 0.3 | 0.7 | 1.1 | 2.2 | 3.8 | 7.4 | 13.4 | 29.3 | 70.0 |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|--|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| Nephritis, nephrotic syndrome, & nephrosis 580-584 | | | | | | | | | | | | |
| WHITE, MALE | | | | | | | | | | | | |
| 1978 | 3.9 | 2.2 | * | * | 0.2 | 0.4 | 0.8 | 2.3 | 5.4 | 16.4 | 45.5 | 95.1 |
| 1977 | 3.7 | 1.8 | * | * | 0.3 | 0.5 | 0.9 | 2.1 | 5.3 | 15.7 | 42.6 | 98.1 |
| 1976 | 3.7 | * | * | * | 0.3 | 0.6 | 1.0 | 2.6 | 6.0 | 15.4 | 41.4 | 94.9 |
| 1975 | 3.5 | * | * | * | 0.3 | 0.6 | 1.1 | 2.2 | 6.4 | 16.1 | 37.3 | 78.1 |
| 1974 | 3.5 | * | * | * | 0.3 | 0.6 | 1.2 | 2.6 | 6.2 | 15.4 | 39.3 | 78.1 |
| 1973 | 3.7 | * | * | 0.1 | 0.5 | 0.8 | 1.3 | 2.9 | 6.5 | 16.1 | 38.7 | 82.0 |
| 1972 | 3.9 | * | * | 0.2 | 0.5 | 0.6 | 1.2 | 3.8 | 7.8 | 16.0 | 37.9 | 84.2 |
| 1971 | 3.8 | * | * | 0.2 | 0.5 | 0.8 | 1.7 | 3.5 | 8.0 | 15.4 | 35.2 | 81.0 |
| 1970 | 4.2 | * | * | 0.3 | 0.7 | 1.1 | 2.2 | 3.9 | 8.4 | 17.2 | 38.1 | 73.1 |
| 1969 | 4.5 | * | * | 0.3 | 0.9 | 1.6 | 2.4 | 4.2 | 9.5 | 16.5 | 40.6 | 83.1 |
| 1968 | 4.6 | * | * | 0.3 | 0.9 | 1.4 | 2.7 | 4.9 | 9.1 | 17.9 | 38.0 | 91.0 |
| WHITE, FEMALE | | | | | | | | | | | | |
| 1978 | 3.0 | * | * | * | 0.2 | 0.4 | 0.5 | 1.5 | 3.9 | 9.2 | 20.9 | 46.8 |
| 1977 | 2.9 | * | * | * | 0.1 | 0.3 | 0.6 | 1.5 | 3.6 | 9.1 | 21.2 | 45.2 |
| 1976 | 2.9 | * | * | * | 0.2 | 0.3 | 0.8 | 1.5 | 3.6 | 9.2 | 21.0 | 48.7 |
| 1975 | 2.9 | * | * | 0.2 | 0.2 | 0.3 | 0.9 | 1.8 | 3.7 | 9.1 | 20.9 | 46.5 |
| 1974 | 2.8 | * | * | * | 0.2 | 0.3 | 0.9 | 1.8 | 4.5 | 8.8 | 20.8 | 42.2 |
| 1973 | 2.9 | * | * | 0.1 | 0.3 | 0.5 | 1.0 | 1.9 | 4.7 | 8.7 | 20.5 | 46.6 |
| 1972 | 3.0 | * | * | 0.1 | 0.4 | 0.6 | 1.0 | 2.5 | 4.4 | 9.4 | 21.4 | 48.7 |
| 1971 | 3.1 | * | * | 0.3 | 0.5 | 0.6 | 1.2 | 2.7 | 4.6 | 10.0 | 20.6 | 48.6 |
| 1970 | 3.2 | * | * | 0.2 | 0.5 | 0.8 | 1.2 | 2.7 | 5.0 | 10.2 | 22.2 | 45.9 |
| 1969 | 3.6 | * | * | 0.4 | 0.7 | 0.9 | 1.7 | 2.7 | 5.4 | 11.3 | 24.9 | 57.2 |
| 1968 | 3.4 | * | * | 0.3 | 0.6 | 0.9 | 1.8 | 2.8 | 5.9 | 9.8 | 23.5 | 58.7 |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|--|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| Nephritis, nephrotic syndrome, & nephrosis 580-584 | | | | | | | | | | | | |
| ALL OTHER, BOTH SEXES | | | | | | | | | | | | |
| 1978 | 7.7 | * | * | * | 0.5 | 1.9 | 4.0 | 10.2 | 23.2 | 40.7 | 74.5 | 114.9 |
| 1977 | 7.7 | 4.6 | * | * | 0.7 | 1.7 | 4.3 | 10.7 | 24.9 | 42.0 | 69.8 | 95.6 |
| 1976 | 7.9 | * | * | * | 0.6 | 2.2 | 4.7 | 11.8 | 22.2 | 46.5 | 70.5 | 109.8 |
| 1975 | 7.3 | * | * | * | 0.6 | 1.9 | 5.5 | 11.6 | 24.7 | 42.6 | 54.6 | 81.5 |
| 1974 | 7.8 | * | * | 0.3 | 1.3 | 2.3 | 5.9 | 13.4 | 25.3 | 42.1 | 58.1 | 92.6 |
| 1973 | 8.4 | * | * | 0.4 | 0.9 | 2.8 | 6.8 | 14.2 | 26.4 | 47.8 | 59.7 | 117.3 |
| 1972 | 8.6 | * | * | 0.3 | 1.0 | 3.2 | 8.0 | 14.8 | 27.0 | 48.9 | 55.1 | 102.4 |
| 1971 | 8.5 | * | * | 0.6 | 1.5 | 4.4 | 8.2 | 14.1 | 28.2 | 38.7 | 54.2 | 110.8 |
| 1970 | 9.1 | * | * | 0.6 | 2.2 | 4.9 | 10.3 | 16.4 | 26.6 | 41.4 | 61.8 | 88.0 |
| 1969 | 9.4 | * | * | 0.6 | 2.3 | 4.3 | 10.8 | 17.5 | 30.1 | 40.3 | 61.1 | 96.3 |
| 1968 | 9.7 | * | * | 0.6 | 2.4 | 4.6 | 11.9 | 18.5 | 26.6 | 40.5 | 67.6 | 124.5 |
| ALL OTHER, MALE | | | | | | | | | | | | |
| 1978 | 8.1 | * | * | * | * | 2.4 | 5.2 | 10.9 | 24.7 | 47.5 | 92.5 | 150.0 |
| 1977 | 8.1 | * | * | * | 0.7 | 2.3 | 4.7 | 11.2 | 26.4 | 47.2 | 85.8 | 153.6 |
| 1976 | 8.1 | * | * | * | 0.8 | 2.9 | 4.8 | 13.5 | 23.4 | 49.8 | 81.7 | 138.2 |
| 1975 | 8.1 | * | * | * | 0.7 | 2.6 | 6.8 | 12.5 | 28.7 | 49.6 | 66.1 | 124.1 |
| 1974 | 8.4 | * | * | * | 1.6 | 2.9 | 6.6 | 13.9 | 26.4 | 47.9 | 79.5 | 134.0 |
| 1973 | 8.7 | * | * | * | 1.3 | 3.3 | 7.7 | 15.0 | 26.2 | 53.4 | 68.3 | 168.8 |
| 1972 | 9.6 | * | * | * | 1.3 | 4.5 | 10.5 | 14.7 | 26.2 | 59.9 | 82.0 | 166.0 |
| 1971 | 9.0 | * | * | 0.7 | 1.7 | 4.9 | 9.1 | 14.6 | 32.1 | 43.8 | 59.4 | 163.8 |
| 1970 | 10.1 | * | * | 0.6 | 2.7 | 5.8 | 12.1 | 19.1 | 27.3 | 47.4 | 75.6 | 119.1 |
| 1969 | 10.4 | * | * | * | 2.8 | 5.4 | 12.7 | 20.9 | 31.2 | 45.8 | 79.3 | 115.9 |
| 1968 | 10.4 | * | * | * | 3.0 | 4.6 | 12.7 | 21.2 | 27.6 | 47.7 | 87.1 | 136.6 |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|--|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| Nephritis, nephrotic syndrome, & nephrosis 580-584 | | | | | | | | | | | | |
| ALL OTHER, FEMALE | | | | | | | | | | | | |
| 1978 | 7.3 | * | * | * | * | 1.5 | 2.9 | 9.7 | 21.9 | 35.4 | 62.5 | 97.2 |
| 1977 | 7.3 | * | * | * | * | 1.2 | 4.1 | 10.2 | 23.6 | 37.9 | 58.7 | 64.4 |
| 1976 | 7.7 | * | * | * | * | 1.5 | 4.5 | 10.4 | 21.2 | 43.8 | 62.8 | 93.9 |
| 1975 | 6.7 | * | * | * | * | 1.3 | 4.4 | 10.8 | 21.3 | 37.1 | 46.6 | 55.9 |
| 1974 | 7.3 | * | * | * | 1.0 | 1.7 | 5.3 | 13.0 | 24.3 | 37.4 | 43.4 | 68.2 |
| 1973 | 8.1 | * | * | * | * | 2.3 | 6.1 | 13.5 | 26.5 | 43.3 | 53.8 | 86.1 |
| 1972 | 7.6 | * | * | * | 0.7 | 2.1 | 6.0 | 15.0 | 27.7 | 40.1 | 35.9 | 63.2 |
| 1971 | 7.9 | * | * | * | 1.4 | 4.0 | 7.6 | 13.6 | 24.9 | 34.6 | 50.4 | 76.7 |
| 1970 | 8.3 | * | * | * | 1.7 | 4.0 | 8.8 | 14.0 | 26.0 | 36.5 | 51.2 | 66.1 |
| 1969 | 8.6 | * | * | 0.7 | 1.8 | 3.4 | 9.1 | 14.5 | 29.0 | 35.9 | 47.0 | 83.1 |
| 1968 | 9.0 | * | * | 0.7 | 1.8 | 4.5 | 11.3 | 16.1 | 25.7 | 34.6 | 52.7 | 116.4 |
| BLACK, BOTH SEXES | | | | | | | | | | | | |
| 1978 | 8.4 | * | * | * | 0.6 | 2.2 | 4.5 | 11.4 | 25.3 | 43.5 | 76.0 | 116.0 |
| 1977 | 8.5 | 5.2 | * | * | 0.7 | 2.0 | 5.0 | 12.1 | 26.9 | 44.6 | 72.0 | 100.7 |
| 1976 | 8.7 | * | * | * | 0.7 | 2.4 | 5.2 | 13.1 | 24.3 | 49.5 | 72.9 | 114.7 |
| 1975 | 8.0 | * | * | * | 0.7 | 2.1 | 6.1 | 13.0 | 26.8 | 45.4 | 54.5 | 77.7 |
| 1974 | 8.6 | * | * | 0.4 | 1.4 | 2.4 | 6.7 | 15.0 | 27.3 | 45.1 | 58.5 | 96.7 |
| 1973 | 9.1 | * | * | 0.4 | 1.0 | 3.2 | 7.5 | 15.6 | 28.2 | 50.4 | 61.3 | 121.1 |
| 1972 | 9.3 | * | * | 0.4 | 1.1 | 3.6 | 9.1 | 16.3 | 28.8 | 51.3 | 56.3 | 109.1 |
| 1971 | 9.1 | * | * | 0.6 | 1.6 | 4.9 | 8.9 | 14.8 | 29.8 | 41.3 | 56.6 | 121.5 |
| 1970 | 9.9 | * | * | 0.6 | 2.4 | 5.2 | 11.5 | 17.9 | 28.2 | 43.9 | 65.5 | 93.7 |
| 1969 | 10.0 | * | * | 0.7 | 2.3 | 4.6 | 11.7 | 18.6 | 31.7 | 42.1 | 64.1 | 104.1 |
| 1968 | 10.2 | * | * | 0.7 | 2.5 | 4.8 | 13.1 | 19.6 | 27.5 | 42.6 | 70.8 | 128.3 |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|--|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| Nephritis, nephrotic syndrome, & nephrosis 580-584 | | | | | | | | | | | | |
| BLACK, MALE | | | | | | | | | | | | |
| 1978 | 8.9 | * | * | * | * | 2.6 | 6.2 | 12.4 | 27.2 | 51.9 | 93.7 | 149.0 |
| 1977 | 9.0 | * | * | * | 0.9 | 2.8 | 5.5 | 12.7 | 28.8 | 51.1 | 91.1 | 165.3 |
| 1976 | 8.9 | * | * | * | 0.9 | 3.2 | 5.7 | 15.1 | 25.3 | 54.6 | 83.3 | 148.9 |
| 1975 | 8.7 | * | * | * | 0.8 | 3.0 | 7.5 | 14.1 | 31.2 | 53.0 | 64.0 | 117.4 |
| 1974 | 9.2 | * | * | * | 1.7 | 3.2 | 7.5 | 15.6 | 29.1 | 52.3 | 78.5 | 144.2 |
| 1973 | 9.4 | * | * | * | 1.4 | 3.8 | 8.3 | 16.7 | 28.3 | 56.7 | 71.1 | 178.0 |
| 1972 | 10.4 | * | * | * | 1.5 | 4.9 | 11.9 | 16.4 | 28.6 | 62.3 | 86.3 | 195.0 |
| 1971 | 9.6 | * | * | * | 1.8 | 5.5 | 9.9 | 15.5 | 34.4 | 46.9 | 61.8 | 187.5 |
| 1970 | 10.9 | * | * | * | 3.0 | 6.1 | 13.6 | 20.8 | 29.8 | 51.0 | 80.6 | 132.6 |
| 1969 | 11.0 | * | * | * | 2.9 | 5.6 | 13.9 | 22.4 | 33.3 | 48.1 | 84.4 | 132.4 |
| 1968 | 10.8 | * | * | * | 3.1 | 4.9 | 13.7 | 22.5 | 28.9 | 50.3 | 89.7 | 142.9 |
| BLACK, FEMALE | | | | | | | | | | | | |
| 1978 | 8.0 | * | * | * | * | 1.8 | 3.2 | 10.5 | 23.8 | 37.3 | 64.2 | 99.0 |
| 1977 | 8.0 | * | * | * | * | 1.4 | 4.7 | 11.5 | 25.3 | 39.6 | 59.6 | 67.0 |
| 1976 | 8.5 | * | * | * | * | 1.8 | 4.8 | 11.6 | 23.4 | 45.6 | 65.9 | 96.6 |
| 1975 | 7.4 | * | * | * | * | 1.3 | 5.0 | 12.1 | 23.1 | 39.6 | 48.2 | 56.0 |
| 1974 | 8.0 | * | * | * | 1.1 | 1.8 | 6.1 | 14.5 | 25.8 | 39.7 | 45.0 | 70.5 |
| 1973 | 8.8 | * | * | * | * | 2.7 | 6.9 | 14.7 | 28.1 | 45.6 | 54.7 | 89.0 |
| 1972 | 8.2 | * | * | * | 0.8 | 2.4 | 6.8 | 16.2 | 29.0 | 42.9 | 35.0 | 60.0 |
| 1971 | 8.5 | * | * | * | 1.5 | 4.3 | 8.0 | 14.2 | 25.9 | 36.8 | 52.9 | 82.1 |
| 1970 | 8.9 | * | * | * | 1.9 | 4.5 | 9.8 | 15.4 | 27.0 | 38.3 | 54.4 | 68.6 |
| 1969 | 9.1 | * | * | 0.8 | 1.8 | 3.9 | 9.8 | 15.4 | 30.3 | 37.2 | 48.9 | 86.7 |
| 1968 | 9.6 | * | * | * | 1.9 | 4.7 | 12.5 | 17.2 | 26.3 | 36.2 | 56.8 | 121.4 |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|--|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| ACUTE NEPHRITIS AND NEPHROTIC SYNDROME (580,581) | | | | | | | | | | | | |
| BOTH SEXES | | | | | | | | | | | | |
| 1978 | 0.9 | 1.7 | * | * | * | 0.1 | 0.2 | 0.4 | 1.3 | 3.5 | 8.8 | 16.8 |
| 1977 | 0.8 | 1.6 | * | * | * | 0.1 | 0.2 | 0.4 | 1.1 | 3.3 | 7.6 | 15.9 |
| 1976 | 0.8 | 1.0 | * | * | * | 0.1 | 0.2 | 0.4 | 1.2 | 3.1 | 7.4 | 15.6 |
| 1975 | 0.7 | * | * | * | 0.1 | 0.1 | 0.2 | 0.5 | 1.2 | 3.2 | 5.9 | 12.1 |
| 1974 | 0.7 | * | * | * | 0.1 | 0.1 | 0.3 | 0.5 | 1.3 | 3.0 | 6.3 | 9.8 |
| 1973 | 0.7 | 0.6 | * | 0.1 | 0.1 | 0.1 | 0.3 | 0.5 | 1.4 | 2.9 | 6.1 | 10.1 |
| 1972 | 0.7 | 0.6 | * | * | 0.1 | 0.2 | 0.3 | 0.7 | 1.4 | 2.6 | 5.2 | 9.7 |
| 1971 | 0.7 | 0.9 | 0.2 | 0.0 | 0.1 | 0.2 | 0.3 | 0.6 | 1.4 | 2.5 | 4.7 | 8.7 |
| 1970 | 0.7 | * | 0.2 | 0.1 | 0.2 | 0.2 | 0.3 | 0.6 | 1.4 | 2.5 | 4.7 | 6.4 |
| 1969 | 0.7 | 0.9 | 0.2 | 0.1 | 0.2 | 0.2 | 0.4 | 0.6 | 1.4 | 2.3 | 4.9 | 7.4 |
| 1968 | 0.6 | 0.6 | 0.2 | 0.1 | 0.1 | 0.2 | 0.4 | 0.7 | 1.3 | 2.2 | 4.3 | 7.3 |
| MALE | | | | | | | | | | | | |
| 1978 | 1.0 | 2.0 | * | * | * | * | 0.2 | 0.5 | 1.6 | 4.8 | 13.0 | 24.2 |
| 1977 | 0.9 | 1.8 | * | * | * | * | 0.2 | 0.4 | 1.4 | 4.3 | 10.9 | 24.9 |
| 1976 | 0.9 | 1.3 | * | * | * | * | 0.2 | 0.6 | 1.5 | 3.8 | 10.6 | 21.3 |
| 1975 | 0.8 | * | * | * | * | 0.2 | 0.2 | 0.5 | 1.4 | 4.4 | 8.4 | 15.3 |
| 1974 | 0.8 | * | * | * | * | 0.1 | 0.3 | 0.5 | 1.3 | 4.2 | 9.3 | 13.3 |
| 1973 | 0.8 | * | * | * | * | 0.1 | 0.3 | 0.6 | 1.5 | 4.0 | 9.1 | 16.0 |
| 1972 | 0.8 | * | * | * | 0.1 | 0.2 | 0.4 | 0.9 | 1.7 | 3.7 | 7.7 | 12.6 |
| 1971 | 0.7 | * | * | * | * | * | 0.3 | 0.7 | 1.7 | 3.1 | 5.9 | 13.1 |
| 1970 | 0.8 | * | * | * | 0.1 | 0.2 | 0.4 | 0.7 | 1.8 | 3.3 | 6.2 | 10.1 |
| 1969 | 0.7 | * | * | 0.1 | 0.2 | 0.2 | 0.4 | 0.8 | 1.6 | 2.8 | 6.1 | 9.2 |
| 1968 | 0.7 | * | * | 0.1 | 0.2 | 0.2 | 0.5 | 0.9 | 1.4 | 2.9 | 6.5 | 9.3 |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|--|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| ACUTE NEPHRITIS AND NEPHROTIC SYNDROME (580,581) | | | | | | | | | | | | |
| FEMALE | | | | | | | | | | | | |
| 1978 | 0.8 | 1.4 | * | * | * | * | 0.2 | 0.3 | 1.1 | 2.6 | 6.3 | 13.5 |
| 1977 | 0.7 | 1.3 | * | * | * | * | * | 0.4 | 0.9 | 2.6 | 5.6 | 11.7 |
| 1976 | 0.7 | * | * | * | * | * | * | 0.3 | 0.9 | 2.7 | 5.6 | 12.9 |
| 1975 | 0.7 | * | * | * | * | * | 0.2 | 0.5 | 1.0 | 2.4 | 4.5 | 10.6 |
| 1974 | 0.6 | * | * | * | * | * | 0.2 | 0.5 | 1.3 | 2.1 | 4.5 | 8.1 |
| 1973 | 0.6 | * | * | * | * | * | 0.3 | 0.4 | 1.2 | 2.1 | 4.2 | 7.2 |
| 1972 | 0.6 | * | * | * | * | 0.2 | 0.2 | 0.5 | 1.1 | 1.8 | 3.6 | 8.2 |
| 1971 | 0.6 | * | * | * | 0.1 | 0.2 | 0.2 | 0.5 | 1.1 | 2.0 | 3.9 | 6.4 |
| 1970 | 0.6 | * | * | 0.1 | 0.2 | * | 0.2 | 0.6 | 1.0 | 1.8 | 3.6 | 4.2 |
| 1969 | 0.6 | * | * | 0.1 | 0.2 | 0.2 | 0.3 | 0.5 | 1.2 | 1.9 | 4.1 | 6.5 |
| 1968 | 0.6 | * | * | * | 0.1 | 0.2 | 0.4 | 0.6 | 1.1 | 1.7 | 2.8 | 6.3 |
| WHITE, BOTH SEXES | | | | | | | | | | | | |
| 1978 | 0.9 | 1.4 | * | * | * | * | 0.2 | 0.3 | 1.1 | 3.3 | 8.3 | 16.3 |
| 1977 | 0.8 | 1.0 | * | * | * | * | 0.1 | 0.3 | 1.0 | 2.9 | 7.3 | 16.0 |
| 1976 | 0.8 | 0.9 | * | * | * | * | 0.2 | 0.3 | 1.0 | 2.8 | 6.9 | 15.7 |
| 1975 | 0.7 | * | * | * | * | 0.1 | 0.1 | 0.4 | 1.0 | 2.8 | 5.7 | 12.2 |
| 1974 | 0.7 | * | * | * | * | * | 0.2 | 0.4 | 1.1 | 2.8 | 6.2 | 9.2 |
| 1973 | 0.7 | * | * | * | * | 0.1 | 0.2 | 0.4 | 1.2 | 2.5 | 5.8 | 9.8 |
| 1972 | 0.6 | * | * | * | 0.1 | 0.1 | 0.2 | 0.6 | 1.1 | 2.4 | 5.1 | 9.9 |
| 1971 | 0.6 | 0.8 | * | * | 0.1 | * | 0.2 | 0.5 | 1.2 | 2.3 | 4.6 | 8.3 |
| 1970 | 0.6 | * | 0.2 | 0.1 | 0.1 | 0.1 | 0.3 | 0.5 | 1.3 | 2.2 | 4.4 | 6.0 |
| 1969 | 0.6 | 0.7 | * | 0.1 | 0.1 | 0.1 | 0.3 | 0.5 | 1.2 | 2.0 | 4.7 | 7.3 |
| 1968 | 0.6 | * | * | 0.1 | 0.1 | 0.1 | 0.3 | 0.6 | 1.1 | 2.1 | 4.1 | 6.5 |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|--|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| ACUTE NEPHRITIS AND NEPHROTIC SYNDROME (580,581) | | | | | | | | | | | | |
| WHITE, MALE | | | | | | | | | | | | |
| 1978 | 1.0 | 1.9 | * | * | * | * | * | 0.4 | 1.4 | 4.4 | 12.4 | 22.9 |
| 1977 | 0.9 | * | * | * | * | * | * | 0.3 | 1.2 | 3.7 | 10.6 | 25.6 |
| 1976 | 0.8 | * | * | * | * | * | * | 0.5 | 1.4 | 3.5 | 10.0 | 21.4 |
| 1975 | 0.8 | * | * | * | * | * | * | 0.4 | 1.2 | 4.0 | 8.0 | 15.2 |
| 1974 | 0.8 | * | * | * | * | * | 0.2 | 0.4 | 1.2 | 3.8 | 9.1 | 12.6 |
| 1973 | 0.8 | * | * | * | * | * | * | 0.5 | 1.3 | 3.6 | 8.9 | 15.1 |
| 1972 | 0.8 | * | * | * | * | * | 0.2 | 0.7 | 1.5 | 3.4 | 7.3 | 12.6 |
| 1971 | 0.7 | * | * | * | * | * | * | 0.6 | 1.5 | 2.8 | 5.9 | 12.1 |
| 1970 | 0.7 | * | * | * | * | * | 0.4 | 0.5 | 1.7 | 3.1 | 6.0 | 9.2 |
| 1969 | 0.7 | * | * | * | 0.1 | * | 0.3 | 0.6 | 1.4 | 2.6 | 5.8 | 8.9 |
| 1968 | 0.7 | * | * | * | * | * | 0.3 | 0.7 | 1.3 | 2.8 | 6.1 | 7.8 |
| WHITE, FEMALE | | | | | | | | | | | | |
| 1978 | 0.8 | * | * | * | * | * | * | 0.3 | 0.9 | 2.4 | 5.9 | 13.4 |
| 1977 | 0.7 | * | * | * | * | * | * | 0.3 | 0.8 | 2.3 | 5.4 | 11.7 |
| 1976 | 0.7 | * | * | * | * | * | * | 0.2 | 0.7 | 2.3 | 5.1 | 13.1 |
| 1975 | 0.6 | * | * | * | * | * | * | 0.3 | 0.8 | 2.0 | 4.3 | 10.8 |
| 1974 | 0.6 | * | * | * | * | * | * | 0.4 | 1.1 | 1.9 | 4.4 | 7.6 |
| 1973 | 0.6 | * | * | * | * | * | * | 0.3 | 1.1 | 1.7 | 3.9 | 7.2 |
| 1972 | 0.5 | * | * | * | * | * | * | 0.4 | 0.8 | 1.7 | 3.7 | 8.5 |
| 1971 | 0.5 | * | * | * | * | * | * | 0.4 | 0.9 | 1.9 | 3.7 | 6.3 |
| 1970 | 0.5 | * | * | * | * | * | * | 0.4 | 0.9 | 1.5 | 3.4 | 4.2 |
| 1969 | 0.6 | * | * | 0.1 | 0.1 | * | 0.3 | 0.4 | 1.0 | 1.6 | 4.0 | 6.4 |
| 1968 | 0.5 | * | * | * | * | * | 0.3 | 0.4 | 0.9 | 1.6 | 2.7 | 5.8 |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|--|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| ACUTE NEPHRITIS AND NEPHROTIC SYNDROME (580,581) | | | | | | | | | | | | |
| ALL OTHER, BOTH SEXES | | | | | | | | | | | | |
| 1978 | 1.2 | * | * | * | * | * | * | 1.1 | 3.3 | 6.1 | 14.0 | 23.2 |
| 1977 | 1.1 | 4.5 | * | * | * | * | * | 1.1 | 2.6 | 7.6 | 10.3 | 13.8 |
| 1976 | 1.1 | * | * | * | * | * | * | 1.1 | 2.7 | 6.4 | 13.0 | 14.4 |
| 1975 | 1.1 | * | * | * | * | * | 0.7 | 1.6 | 3.2 | 7.2 | 8.6 | * |
| 1974 | 1.0 | * | * | * | * | * | 0.9 | 1.3 | 2.8 | 5.6 | 8.0 | 16.3 |
| 1973 | 1.2 | * | * | * | * | * | 1.1 | 1.6 | 3.3 | 7.0 | 9.4 | * |
| 1972 | 1.1 | * | * | * | * | 0.6 | 1.2 | 1.8 | 4.1 | 4.4 | 6.9 | * |
| 1971 | 1.2 | * | * | * | * | 0.9 | 1.1 | 1.7 | 3.3 | 4.8 | 6.8 | * |
| 1970 | 1.1 | * | * | * | 0.4 | * | 0.9 | 2.0 | 2.9 | 4.9 | 7.4 | * |
| 1969 | 1.3 | * | * | * | * | * | 1.4 | 2.0 | 3.2 | 5.3 | 7.6 | * |
| 1968 | 1.2 | * | * | * | * | 0.8 | 1.3 | 2.1 | 2.8 | 3.8 | 7.5 | * |
| ALL OTHER, MALE | | | | | | | | | | | | |
| 1978 | 1.5 | * | * | * | * | * | * | * | 3.7 | 8.2 | 19.2 | 37.9 |
| 1977 | 1.3 | * | * | * | * | * | * | * | 2.9 | 10.2 | 13.8 | * |
| 1976 | 1.2 | * | * | * | * | * | * | * | 2.9 | 6.6 | 16.0 | * |
| 1975 | 1.2 | * | * | * | * | * | * | * | 3.3 | 8.3 | 12.0 | * |
| 1974 | 1.2 | * | * | * | * | * | * | * | 2.7 | 7.5 | 11.4 | * |
| 1973 | 1.3 | * | * | * | * | * | * | * | 3.7 | 8.1 | 11.2 | * |
| 1972 | 1.3 | * | * | * | * | * | 1.6 | 2.3 | 3.5 | 6.8 | 12.0 | * |
| 1971 | 1.2 | * | * | * | * | * | * | 1.8 | 3.7 | 6.3 | * | * |
| 1970 | 1.2 | * | * | * | * | * | * | 2.2 | 3.2 | 5.2 | * | * |
| 1969 | 1.3 | * | * | * | * | * | 1.9 | 2.5 | 3.7 | 4.7 | * | * |
| 1968 | 1.3 | * | * | * | * | * | 1.9 | 2.4 | * | 4.4 | 10.8 | * |
| ALL OTHER, FEMALE | | | | | | | | | | | | |
| 1978 | 1.0 | * | * | * | * | * | * | * | 2.9 | 4.5 | 10.5 | * |
| 1977 | 1.0 | * | * | * | * | * | * | * | 2.3 | 5.6 | 7.8 | * |
| 1976 | 1.0 | * | * | * | * | * | * | * | 2.5 | 6.3 | 10.9 | * |
| 1975 | 1.1 | * | * | * | * | * | * | 1.9 | 3.0 | 6.2 | 6.2 | * |
| 1974 | 0.9 | * | * | * | * | * | * | * | 2.9 | 4.0 | * | * |
| 1973 | 1.1 | * | * | * | * | * | * | 1.5 | 2.9 | 6.2 | 8.2 | * |
| 1972 | 0.9 | * | * | * | * | * | * | * | 4.5 | * | * | * |
| 1971 | 1.1 | * | * | * | * | * | * | 1.5 | 3.0 | 3.6 | * | * |
| 1970 | 1.0 | * | * | * | * | * | * | 1.9 | 2.5 | 4.6 | * | * |
| 1969 | 1.2 | * | * | * | * | * | * | * | 2.8 | 5.7 | * | * |
| 1968 | 1.1 | * | * | * | * | * | * | 1.8 | 3.1 | 3.3 | * | * |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|--|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| ACUTE NEPHRITIS AND NEPHROTIC SYNDROME (580,581) | | | | | | | | | | | | |
| BLACK, BOTH SEXES | | | | | | | | | | | | |
| 1978 | 1.3 | * | * | * | * | * | * | 1.2 | 3.6 | 6.7 | 13.4 | 20.0 |
| 1977 | 1.3 | 4.9 | * | * | * | * | * | 1.3 | 2.8 | 8.3 | 10.3 | * |
| 1976 | 1.2 | * | * | * | * | * | * | 1.2 | 3.0 | 6.8 | 13.4 | 16.2 |
| 1975 | 1.2 | * | * | * | * | * | * | 1.8 | 3.4 | 7.4 | 8.2 | * |
| 1974 | 1.1 | * | * | * | * | * | 1.0 | 1.4 | 3.2 | 5.7 | 8.1 | 17.4 |
| 1973 | 1.3 | * | * | * | * | * | 1.3 | 1.8 | 3.5 | 7.3 | 9.8 | * |
| 1972 | 1.2 | * | * | * | * | 0.7 | 1.3 | 1.8 | 4.1 | 4.9 | 7.1 | * |
| 1971 | 1.2 | * | * | * | * | 0.9 | 1.0 | 1.7 | 3.6 | 5.2 | 6.7 | * |
| 1970 | 1.2 | * | * | * | 0.5 | * | 1.1 | 2.3 | 3.0 | 5.2 | 7.3 | * |
| 1969 | 1.3 | * | * | * | * | * | 1.5 | 2.1 | 3.4 | 5.6 | 7.6 | * |
| 1968 | 1.3 | * | * | * | * | 0.9 | 1.5 | 2.2 | 3.0 | 3.8 | 7.8 | * |
| BLACK, MALE | | | | | | | | | | | | |
| 1978 | 1.6 | * | * | * | * | * | * | * | 4.1 | 9.4 | 18.8 | * |
| 1977 | 1.5 | * | * | * | * | * | * | * | 3.2 | 11.4 | 15.3 | * |
| 1976 | 1.3 | * | * | * | * | * | * | * | 3.2 | 7.1 | 15.8 | * |
| 1975 | 1.2 | * | * | * | * | * | * | * | 3.7 | 8.2 | 11.3 | * |
| 1974 | 1.3 | * | * | * | * | * | * | * | 3.1 | 7.7 | 11.0 | * |
| 1973 | 1.4 | * | * | * | * | * | * | * | 4.1 | 8.3 | 12.3 | * |
| 1972 | 1.5 | * | * | * | * | * | * | 2.6 | 3.4 | 7.7 | 13.2 | * |
| 1971 | 1.3 | * | * | * | * | * | * | * | 4.0 | 6.9 | * | * |
| 1970 | 1.3 | * | * | * | * | * | * | 2.5 | 3.5 | 5.6 | * | * |
| 1969 | 1.4 | * | * | * | * | * | 2.0 | 2.6 | 4.0 | 5.3 | * | * |
| 1968 | 1.3 | * | * | * | * | * | 2.1 | 2.5 | * | * | * | * |
| BLACK, FEMALE | | | | | | | | | | | | |
| 1978 | 1.1 | * | * | * | * | * | * | * | 3.2 | 4.7 | 9.8 | * |
| 1977 | 1.1 | * | * | * | * | * | * | * | 2.5 | 6.0 | 7.1 | * |
| 1976 | 1.1 | * | * | * | * | * | * | * | 2.8 | 6.6 | 11.7 | * |
| 1975 | 1.2 | * | * | * | * | * | * | 2.2 | 3.2 | 6.8 | 6.1 | * |
| 1974 | 1.0 | * | * | * | * | * | * | * | 3.2 | 4.2 | * | * |
| 1973 | 1.1 | * | * | * | * | * | * | 1.7 | 2.9 | 6.4 | 8.1 | * |
| 1972 | 0.9 | * | * | * | * | * | * | * | 4.6 | * | * | * |
| 1971 | 1.1 | * | * | * | * | * | * | * | 3.3 | 3.9 | * | * |
| 1970 | 1.1 | * | * | * | * | * | * | 2.1 | 2.6 | 4.8 | * | * |
| 1969 | 1.3 | * | * | * | * | * | * | * | 2.8 | 5.9 | * | * |
| 1968 | 1.2 | * | * | * | * | * | * | 2.0 | 3.3 | 3.6 | * | * |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| CHRONIC AND UNQUALIFIED NEPHRITIS AND RENAL SCLEROSIS (582-584) | | | | | | | | | | | | |
| BOTH SEXES | | | | | | | | | | | | |
| 1978 | 3.1 | * | * | 0.1 | 0.2 | 0.5 | 0.9 | 2.4 | 5.2 | 11.5 | 25.1 | 49.1 |
| 1977 | 3.0 | * | * | 0.1 | 0.2 | 0.5 | 1.1 | 2.4 | 5.2 | 11.5 | 25.2 | 48.5 |
| 1976 | 3.1 | * | * | 0.1 | 0.3 | 0.6 | 1.2 | 2.7 | 5.3 | 12.0 | 24.9 | 51.4 |
| 1975 | 3.0 | * | * | 0.1 | 0.2 | 0.5 | 1.3 | 2.5 | 5.7 | 11.7 | 23.5 | 46.5 |
| 1974 | 3.1 | * | * | 0.1 | 0.3 | 0.6 | 1.4 | 2.9 | 5.9 | 11.5 | 24.0 | 47.2 |
| 1973 | 3.2 | * | * | 0.1 | 0.4 | 0.8 | 1.6 | 3.1 | 6.1 | 12.3 | 24.0 | 52.8 |
| 1972 | 3.4 | * | * | 0.2 | 0.4 | 0.7 | 1.7 | 3.7 | 6.6 | 13.0 | 24.7 | 54.2 |
| 1971 | 3.4 | * | * | 0.2 | 0.5 | 1.0 | 2.0 | 3.6 | 6.9 | 12.3 | 23.7 | 54.9 |
| 1970 | 3.7 | * | * | 0.2 | 0.7 | 1.3 | 2.4 | 4.0 | 7.1 | 13.4 | 26.3 | 52.0 |
| 1969 | 4.0 | * | 0.2 | 0.2 | 0.8 | 1.4 | 2.6 | 4.2 | 8.1 | 13.7 | 28.4 | 61.3 |
| 1968 | 4.0 | * | * | 0.3 | 0.8 | 1.3 | 2.9 | 4.5 | 8.0 | 13.6 | 27.8 | 67.1 |
| MALE | | | | | | | | | | | | |
| 1978 | 3.4 | * | * | * | 0.2 | 0.6 | 1.1 | 2.7 | 5.7 | 14.7 | 37.0 | 75.8 |
| 1977 | 3.4 | * | * | * | 0.3 | 0.6 | 1.2 | 2.6 | 5.9 | 14.5 | 35.9 | 78.1 |
| 1976 | 3.4 | * | * | * | 0.3 | 0.8 | 1.2 | 3.1 | 6.2 | 14.9 | 34.7 | 77.6 |
| 1975 | 3.3 | * | * | 0.1 | 0.3 | 0.6 | 1.5 | 2.7 | 7.1 | 14.9 | 31.6 | 67.0 |
| 1974 | 3.4 | * | * | 0.1 | 0.4 | 0.7 | 1.5 | 3.2 | 6.7 | 14.3 | 33.6 | 69.7 |
| 1973 | 3.5 | * | * | 0.1 | 0.5 | 0.9 | 1.7 | 3.5 | 6.8 | 15.6 | 32.3 | 73.8 |
| 1972 | 3.7 | * | * | 0.2 | 0.5 | 0.8 | 1.9 | 4.0 | 7.8 | 16.4 | 34.0 | 78.9 |
| 1971 | 3.7 | * | * | 0.2 | 0.6 | 1.1 | 2.2 | 3.9 | 8.5 | 14.9 | 31.3 | 75.5 |
| 1970 | 4.1 | * | * | 0.2 | 0.8 | 1.4 | 2.9 | 4.7 | 8.4 | 16.8 | 34.8 | 67.7 |
| 1969 | 4.4 | * | * | 0.2 | 0.9 | 1.8 | 3.0 | 5.1 | 9.9 | 16.5 | 37.5 | 77.1 |
| 1968 | 4.5 | * | * | 0.3 | 1.0 | 1.6 | 3.3 | 5.6 | 9.5 | 17.7 | 35.3 | 85.9 |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|

CHRONIC AND UNQUALIFIED NEPHRITIS AND RENAL SCLEROSIS (582-584)

FEMALE

| | | | | | | | | | | | | |
|------|-----|---|---|-----|-----|-----|-----|-----|-----|------|------|------|
| 1978 | 2.8 | * | * | * | 0.2 | 0.4 | 0.7 | 2.1 | 4.7 | 9.1 | 18.1 | 37.1 |
| 1977 | 2.8 | * | * | * | 0.2 | 0.4 | 1.0 | 2.1 | 4.7 | 9.2 | 18.8 | 34.9 |
| 1976 | 2.8 | * | * | * | 0.2 | 0.4 | 1.1 | 2.3 | 4.5 | 9.7 | 19.0 | 39.1 |
| 1975 | 2.7 | * | * | 0.1 | 0.2 | 0.4 | 1.2 | 2.4 | 4.5 | 9.3 | 18.5 | 36.6 |
| 1974 | 2.8 | * | * | * | 0.3 | 0.4 | 1.2 | 2.6 | 5.2 | 9.3 | 18.0 | 36.1 |
| 1973 | 3.0 | * | * | 0.1 | 0.3 | 0.7 | 1.4 | 2.7 | 5.5 | 9.8 | 18.8 | 42.3 |
| 1972 | 3.0 | * | * | 0.1 | 0.3 | 0.6 | 1.5 | 3.4 | 5.6 | 10.4 | 18.8 | 41.6 |
| 1971 | 3.1 | * | * | 0.2 | 0.5 | 0.8 | 1.8 | 3.3 | 5.5 | 10.2 | 18.8 | 44.2 |
| 1970 | 3.3 | * | * | 0.2 | 0.5 | 1.1 | 2.0 | 3.4 | 6.0 | 10.8 | 20.6 | 43.3 |
| 1969 | 3.6 | * | * | 0.3 | 0.7 | 1.1 | 2.2 | 3.4 | 6.5 | 11.6 | 22.3 | 52.7 |
| 1968 | 3.5 | * | * | 0.3 | 0.6 | 1.1 | 2.6 | 3.5 | 6.6 | 10.3 | 22.6 | 56.9 |

WHITE, BOTH SEXES

| | | | | | | | | | | | | |
|------|-----|---|---|-----|-----|-----|-----|-----|-----|------|------|------|
| 1978 | 2.5 | * | * | 0.1 | 0.2 | 0.3 | 0.5 | 1.5 | 3.5 | 9.1 | 21.7 | 45.4 |
| 1977 | 2.5 | * | * | * | 0.2 | 0.3 | 0.6 | 1.5 | 3.4 | 9.1 | 21.8 | 45.6 |
| 1976 | 2.6 | * | * | 0.1 | 0.2 | 0.4 | 0.7 | 1.7 | 3.7 | 9.0 | 21.7 | 47.5 |
| 1975 | 2.5 | * | * | 0.1 | 0.2 | 0.4 | 0.8 | 1.6 | 4.0 | 9.3 | 21.3 | 44.5 |
| 1974 | 2.5 | * | * | 0.1 | 0.2 | 0.4 | 0.9 | 1.8 | 4.2 | 8.9 | 21.6 | 44.7 |
| 1973 | 2.6 | * | * | 0.1 | 0.3 | 0.6 | 1.0 | 2.0 | 4.3 | 9.4 | 21.7 | 48.5 |
| 1972 | 2.8 | * | * | 0.1 | 0.3 | 0.5 | 1.0 | 2.6 | 4.9 | 9.8 | 22.7 | 50.7 |
| 1971 | 2.9 | * | * | 0.2 | 0.4 | 0.6 | 1.3 | 2.6 | 5.0 | 10.1 | 21.8 | 51.3 |
| 1970 | 3.1 | * | * | 0.2 | 0.5 | 0.8 | 1.5 | 2.8 | 5.3 | 11.1 | 24.0 | 49.5 |
| 1969 | 3.4 | * | * | 0.2 | 0.6 | 1.1 | 1.7 | 2.9 | 6.1 | 11.6 | 26.5 | 58.9 |
| 1968 | 3.4 | * | * | 0.2 | 0.6 | 1.0 | 1.9 | 3.2 | 6.3 | 11.3 | 25.3 | 63.5 |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| CHRONIC AND UNQUALIFIED NEPHRITIS AND RENAL SCLEROSIS (582-584) | | | | | | | | | | | | |
| WHITE, MALE | | | | | | | | | | | | |
| 1978 | 2.9 | * | * | * | 0.2 | 0.4 | 0.6 | 1.9 | 4.0 | 12.0 | 33.1 | 72.2 |
| 1977 | 2.8 | * | * | * | 0.2 | 0.4 | 0.8 | 1.7 | 4.0 | 12.1 | 32.0 | 72.5 |
| 1976 | 2.9 | * | * | * | 0.3 | 0.5 | 0.8 | 2.1 | 4.7 | 11.9 | 31.4 | 73.5 |
| 1975 | 2.8 | * | * | * | 0.3 | 0.4 | 0.9 | 1.7 | 5.2 | 12.1 | 29.3 | 63.0 |
| 1974 | 2.8 | * | * | * | 0.3 | 0.5 | 1.0 | 2.2 | 5.0 | 11.6 | 30.3 | 65.4 |
| 1973 | 2.9 | * | * | * | 0.4 | 0.7 | 1.1 | 2.4 | 5.2 | 12.5 | 29.9 | 66.9 |
| 1972 | 3.1 | * | * | 0.2 | 0.4 | 0.5 | 1.0 | 3.1 | 6.2 | 12.6 | 30.6 | 71.6 |
| 1971 | 3.1 | * | * | 0.2 | 0.5 | 0.7 | 1.5 | 2.9 | 6.4 | 12.6 | 29.4 | 68.9 |
| 1970 | 3.5 | * | * | 0.2 | 0.6 | 1.0 | 1.8 | 3.3 | 6.7 | 14.1 | 32.1 | 63.9 |
| 1969 | 3.8 | * | * | 0.2 | 0.7 | 1.4 | 2.1 | 3.6 | 8.1 | 13.9 | 34.8 | 74.2 |
| 1968 | 3.9 | * | * | 0.2 | 0.8 | 1.3 | 2.4 | 4.2 | 7.9 | 15.1 | 31.9 | 83.2 |
| WHITE, FEMALE | | | | | | | | | | | | |
| 1978 | 2.2 | * | * | * | 0.2 | 0.3 | 0.4 | 1.2 | 3.1 | 6.8 | 15.0 | 33.5 |
| 1977 | 2.2 | * | * | * | 0.1 | 0.2 | 0.6 | 1.2 | 2.8 | 6.8 | 15.8 | 33.5 |
| 1976 | 2.2 | * | * | * | 0.2 | 0.3 | 0.6 | 1.3 | 2.9 | 6.8 | 15.9 | 35.6 |
| 1975 | 2.3 | * | * | 0.1 | 0.2 | 0.3 | 0.8 | 1.5 | 3.0 | 7.1 | 16.5 | 35.7 |
| 1974 | 2.2 | * | * | * | 0.2 | 0.3 | 0.8 | 1.5 | 3.4 | 6.9 | 16.3 | 34.7 |
| 1973 | 2.3 | * | * | * | 0.2 | 0.4 | 0.8 | 1.6 | 3.6 | 7.1 | 16.6 | 39.4 |
| 1972 | 2.5 | * | * | * | 0.3 | 0.4 | 0.9 | 2.1 | 3.7 | 7.7 | 17.7 | 40.3 |
| 1971 | 2.6 | * | * | 0.2 | 0.4 | 0.5 | 1.1 | 2.3 | 3.7 | 8.2 | 16.9 | 42.3 |
| 1970 | 2.7 | * | * | 0.1 | 0.4 | 0.7 | 1.1 | 2.3 | 4.1 | 8.7 | 18.8 | 41.7 |
| 1969 | 3.0 | * | * | 0.2 | 0.6 | 0.8 | 1.4 | 2.2 | 4.4 | 9.7 | 20.9 | 50.8 |
| 1968 | 2.9 | * | * | 0.2 | 0.5 | 0.7 | 1.5 | 2.3 | 4.9 | 8.3 | 20.8 | 52.9 |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| CHRONIC AND UNQUALIFIED NEPHRITIS AND RENAL SCLEROSIS (582-584) | | | | | | | | | | | | |
| ALL OTHER, BOTH SEXES | | | | | | | | | | | | |
| 1978 | 6.4 | * | * | * | 0.4 | 1.6 | 3.6 | 9.2 | 19.9 | 34.5 | 60.5 | 91.7 |
| 1977 | 6.6 | * | * | * | 0.6 | 1.5 | 3.9 | 9.6 | 22.3 | 34.4 | 59.6 | 81.9 |
| 1976 | 6.8 | * | * | * | 0.6 | 1.9 | 4.2 | 10.7 | 19.5 | 40.1 | 57.5 | 95.4 |
| 1975 | 6.2 | * | * | * | 0.5 | 1.5 | 4.8 | 10.0 | 21.5 | 35.4 | 46.1 | 69.9 |
| 1974 | 6.8 | * | * | * | 1.1 | 1.9 | 5.0 | 12.1 | 22.5 | 36.5 | 50.1 | 76.3 |
| 1973 | 7.2 | * | * | * | 0.8 | 2.4 | 5.7 | 12.7 | 23.1 | 40.7 | 50.3 | 103.1 |
| 1972 | 7.5 | * | * | * | 0.8 | 2.6 | 6.8 | 13.1 | 23.0 | 44.5 | 48.2 | 94.3 |
| 1971 | 7.3 | * | * | 0.5 | 1.2 | 3.6 | 7.2 | 12.4 | 24.9 | 33.9 | 47.4 | 96.7 |
| 1970 | 8.0 | * | * | 0.4 | 1.8 | 4.3 | 9.3 | 14.4 | 23.7 | 36.6 | 54.4 | 77.6 |
| 1969 | 8.2 | * | * | 0.3 | 1.9 | 3.7 | 9.4 | 15.5 | 26.8 | 35.1 | 53.5 | 87.2 |
| 1968 | 8.5 | * | * | 0.4 | 2.0 | 3.8 | 10.6 | 16.3 | 23.8 | 36.7 | 60.1 | 107.8 |
| ALL OTHER, MALE | | | | | | | | | | | | |
| 1978 | 6.6 | * | * | * | * | 1.9 | 4.7 | 9.5 | 21.0 | 39.3 | 73.3 | 112.1 |
| 1977 | 6.8 | * | * | * | * | 2.0 | 4.1 | 10.1 | 23.5 | 37.0 | 72.0 | 135.7 |
| 1976 | 6.9 | * | * | * | * | 2.7 | 4.2 | 11.9 | 20.6 | 43.3 | 65.8 | 118.2 |
| 1975 | 6.9 | * | * | * | * | 2.2 | 6.1 | 11.4 | 25.3 | 41.2 | 54.1 | 107.4 |
| 1974 | 7.3 | * | * | * | 1.3 | 2.5 | 5.6 | 12.5 | 23.7 | 40.4 | 68.1 | 114.0 |
| 1973 | 7.4 | * | * | * | 1.1 | 2.8 | 6.3 | 13.4 | 22.5 | 45.3 | 57.1 | 143.8 |
| 1972 | 8.2 | * | * | * | 1.2 | 3.4 | 8.9 | 12.4 | 22.7 | 53.1 | 70.0 | 153.2 |
| 1971 | 7.8 | * | * | * | 1.4 | 4.3 | 7.8 | 12.8 | 28.3 | 37.5 | 52.7 | 140.4 |
| 1970 | 8.8 | * | * | * | 2.2 | 4.9 | 10.9 | 17.0 | 24.1 | 42.2 | 67.3 | 101.0 |
| 1969 | 9.1 | * | * | * | 2.4 | 4.9 | 10.8 | 18.4 | 27.6 | 41.0 | 69.7 | 104.5 |
| 1968 | 9.1 | * | * | * | 2.6 | 3.9 | 10.8 | 18.7 | 25.3 | 43.3 | 76.3 | 112.2 |
| ALL OTHER, FEMALE | | | | | | | | | | | | |
| 1978 | 6.3 | * | * | * | * | 1.3 | 2.7 | 8.9 | 19.0 | 30.9 | 51.9 | 81.7 |
| 1977 | 6.3 | * | * | * | * | 1.1 | 3.7 | 9.1 | 21.3 | 32.3 | 50.9 | 52.9 |
| 1976 | 6.7 | * | * | * | * | 1.2 | 4.3 | 9.8 | 18.7 | 37.5 | 51.9 | 82.7 |
| 1975 | 5.6 | * | * | * | * | * | 3.7 | 8.9 | 18.3 | 30.8 | 40.4 | 47.3 |
| 1974 | 6.4 | * | * | * | 0.9 | 1.4 | 4.4 | 11.8 | 21.4 | 33.4 | 37.7 | 54.1 |
| 1973 | 7.1 | * | * | * | * | 2.1 | 5.2 | 12.0 | 23.6 | 37.0 | 45.6 | 78.5 |
| 1972 | 6.7 | * | * | * | * | 1.9 | 5.0 | 13.6 | 23.2 | 37.6 | 32.6 | 57.9 |
| 1971 | 6.9 | * | * | * | 1.0 | 3.0 | 6.7 | 12.0 | 21.9 | 31.0 | 43.6 | 68.5 |
| 1970 | 7.3 | * | * | * | 1.3 | 3.7 | 8.0 | 12.1 | 23.4 | 31.9 | 44.5 | 61.0 |
| 1969 | 7.3 | * | * | * | 1.3 | 2.8 | 8.2 | 13.0 | 26.2 | 30.2 | 40.9 | 75.4 |
| 1968 | 7.9 | * | * | * | 1.4 | 3.6 | 10.5 | 14.3 | 22.6 | 31.3 | 47.7 | 104.9 |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| CHRONIC AND UNQUALIFIED NEPHRITIS AND RENAL SCLEROSIS (582-584) | | | | | | | | | | | | |
| BLACK, BOTH SEXES | | | | | | | | | | | | |
| 1978 | 7.1 | * | * | * | 0.5 | 1.8 | 4.1 | 10.2 | 21.7 | 36.8 | 62.6 | 96.0 |
| 1977 | 7.2 | * | * | * | 0.6 | 1.8 | 4.5 | 10.8 | 24.1 | 36.2 | 61.7 | 88.1 |
| 1976 | 7.5 | * | * | * | 0.6 | 2.2 | 4.7 | 11.9 | 21.3 | 42.7 | 59.5 | 98.5 |
| 1975 | 6.8 | * | * | * | 0.5 | 1.6 | 5.4 | 11.2 | 23.4 | 38.0 | 46.4 | 67.7 |
| 1974 | 7.4 | * | * | * | 1.2 | 2.1 | 5.7 | 13.5 | 24.2 | 39.4 | 50.4 | 79.3 |
| 1973 | 7.8 | * | * | * | 0.9 | 2.8 | 6.3 | 13.8 | 24.7 | 43.1 | 51.5 | 107.0 |
| 1972 | 8.1 | * | * | * | 0.9 | 2.9 | 7.8 | 14.5 | 24.7 | 46.5 | 49.2 | 100.0 |
| 1971 | 7.8 | * | * | 0.5 | 1.3 | 3.9 | 7.8 | 13.1 | 26.2 | 36.1 | 49.9 | 105.6 |
| 1970 | 8.7 | * | * | 0.4 | 1.9 | 4.6 | 10.4 | 15.7 | 25.2 | 38.7 | 58.2 | 82.6 |
| 1969 | 8.7 | * | * | 0.4 | 1.9 | 4.0 | 10.2 | 16.5 | 28.3 | 36.4 | 56.5 | 93.8 |
| 1968 | 8.9 | * | * | 0.4 | 2.1 | 3.9 | 11.6 | 17.4 | 24.5 | 38.7 | 63.0 | 109.8 |
| BLACK, MALE | | | | | | | | | | | | |
| 1978 | 7.3 | * | * | * | * | 2.0 | 5.5 | 10.8 | 23.1 | 42.5 | 74.9 | 119.6 |
| 1977 | 7.5 | * | * | * | * | 2.4 | 4.8 | 11.4 | 25.6 | 39.7 | 75.7 | 149.0 |
| 1976 | 7.6 | * | * | * | * | 3.0 | 4.9 | 13.2 | 22.1 | 47.5 | 67.5 | 125.5 |
| 1975 | 7.4 | * | * | * | * | 2.5 | 6.8 | 12.8 | 27.6 | 44.8 | 52.7 | 104.3 |
| 1974 | 7.9 | * | * | * | 1.4 | 2.7 | 6.4 | 14.0 | 26.0 | 44.6 | 67.5 | 123.3 |
| 1973 | 8.0 | * | * | * | 1.2 | 3.3 | 6.8 | 14.8 | 24.3 | 48.3 | 58.8 | 153.7 |
| 1972 | 8.9 | * | * | * | 1.3 | 3.7 | 10.2 | 13.8 | 25.1 | 54.6 | 73.1 | 180.0 |
| 1971 | 8.3 | * | * | * | 1.4 | 4.8 | 8.6 | 13.7 | 30.3 | 40.0 | 55.4 | 160.0 |
| 1970 | 9.5 | * | * | * | 2.4 | 5.1 | 12.2 | 18.4 | 26.2 | 45.3 | 72.3 | 113.0 |
| 1969 | 9.6 | * | * | * | 2.5 | 5.0 | 11.9 | 19.8 | 29.4 | 42.9 | 74.3 | 118.9 |
| 1968 | 9.5 | * | * | * | 2.6 | 4.1 | 11.6 | 20.0 | 26.3 | 46.1 | 78.8 | 114.3 |
| BLACK, FEMALE | | | | | | | | | | | | |
| 1978 | 6.9 | * | * | * | * | 1.5 | 2.9 | 9.7 | 20.6 | 32.6 | 54.5 | 83.8 |
| 1977 | 7.0 | * | * | * | * | 1.3 | 4.2 | 10.3 | 22.8 | 33.6 | 52.5 | 56.4 |
| 1976 | 7.4 | * | * | * | * | 1.4 | 4.5 | 10.9 | 20.6 | 39.1 | 54.2 | 84.3 |
| 1975 | 6.1 | * | * | * | * | * | 4.2 | 9.9 | 19.9 | 32.8 | 42.1 | 47.6 |
| 1974 | 7.0 | * | * | * | 1.0 | 1.5 | 5.1 | 13.2 | 22.6 | 35.5 | 38.8 | 55.1 |
| 1973 | 7.7 | * | * | * | * | 2.4 | 5.8 | 13.0 | 25.1 | 39.1 | 46.6 | 80.8 |
| 1972 | 7.3 | * | * | * | * | 2.2 | 5.8 | 15.0 | 24.3 | 40.2 | 32.1 | 54.3 |
| 1971 | 7.4 | * | * | * | 1.2 | 3.2 | 7.2 | 12.6 | 22.6 | 32.9 | 45.9 | 73.1 |
| 1970 | 7.9 | * | * | * | 1.4 | 4.2 | 8.9 | 13.3 | 24.3 | 33.5 | 47.8 | 63.0 |
| 1969 | 7.8 | * | * | * | 1.4 | 3.2 | 8.7 | 13.7 | 27.5 | 31.3 | 43.2 | 78.3 |
| 1968 | 8.4 | * | * | * | 1.5 | 3.7 | 11.6 | 15.2 | 23.1 | 32.7 | 51.4 | 108.9 |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| INFECTIONS OF KIDNEY (590) | | | | | | | | | | | | |
| BOTH SEXES | | | | | | | | | | | | |
| 1978 | 1.4 | * | * | * | * | 0.1 | 0.2 | 0.5 | 1.3 | 4.2 | 14.8 | 42.6 |
| 1977 | 1.7 | * | * | * | * | 0.1 | 0.2 | 0.6 | 1.6 | 5.2 | 18.0 | 53.5 |
| 1976 | 1.8 | * | * | * | 0.0 | 0.1 | 0.3 | 0.6 | 2.1 | 6.1 | 19.8 | 54.3 |
| 1975 | 2.1 | * | * | * | * | 0.1 | 0.3 | 0.8 | 2.3 | 7.3 | 23.2 | 61.0 |
| 1974 | 2.6 | * | * | * | 0.1 | 0.2 | 0.4 | 1.0 | 2.8 | 9.7 | 28.7 | 70.6 |
| 1973 | 2.9 | * | * | * | 0.1 | 0.2 | 0.6 | 1.1 | 3.7 | 11.1 | 31.3 | 82.8 |
| 1972 | 3.3 | * | * | * | 0.1 | 0.2 | 0.6 | 1.8 | 4.0 | 12.8 | 38.0 | 88.3 |
| 1971 | 3.7 | * | * | * | 0.2 | 0.3 | 0.7 | 2.0 | 5.1 | 13.8 | 40.4 | 96.6 |
| 1970 | 4.0 | 0.6 | * | 0.1 | 0.2 | 0.4 | 1.0 | 2.4 | 6.0 | 15.6 | 43.2 | 99.5 |
| 1969 | 4.3 | * | * | 0.1 | 0.2 | 0.4 | 1.1 | 2.5 | 6.2 | 17.3 | 46.8 | 122.3 |
| 1968 | 4.7 | 0.9 | * | 0.1 | 0.2 | 0.5 | 1.3 | 3.3 | 7.6 | 18.6 | 48.4 | 126.9 |
| MALE | | | | | | | | | | | | |
| 1978 | 1.2 | * | * | * | * | * | * | 0.4 | 1.0 | 4.3 | 16.2 | 51.8 |
| 1977 | 1.4 | * | * | * | * | * | * | 0.5 | 1.5 | 5.2 | 20.2 | 58.5 |
| 1976 | 1.6 | * | * | * | * | * | * | 0.4 | 1.8 | 6.5 | 23.5 | 64.4 |
| 1975 | 1.7 | * | * | * | * | * | 0.3 | 0.6 | 2.1 | 6.9 | 25.3 | 71.5 |
| 1974 | 2.2 | * | * | * | * | * | 0.3 | 0.8 | 2.7 | 10.4 | 31.3 | 83.7 |
| 1973 | 2.5 | * | * | * | * | 0.1 | 0.4 | 0.9 | 3.3 | 11.5 | 35.8 | 94.2 |
| 1972 | 3.0 | * | * | * | * | 0.2 | 0.6 | 1.4 | 3.4 | 14.4 | 44.0 | 106.1 |
| 1971 | 3.4 | * | * | * | 0.2 | 0.3 | 0.4 | 1.6 | 4.5 | 15.5 | 47.1 | 122.2 |
| 1970 | 3.7 | * | * | * | 0.1 | 0.3 | 0.6 | 1.9 | 5.6 | 16.5 | 49.8 | 121.5 |
| 1969 | 4.1 | * | * | * | 0.2 | 0.2 | 0.8 | 1.9 | 6.1 | 18.7 | 55.9 | 151.6 |
| 1968 | 4.5 | 1.2 | * | * | 0.2 | 0.3 | 1.0 | 2.6 | 7.0 | 21.1 | 58.4 | 159.3 |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| INFECTIONS OF KIDNEY (590) | | | | | | | | | | | | |
| FEMALE | | | | | | | | | | | | |
| 1978 | 1.6 | * | * | * | * | * | 0.2 | 0.6 | 1.5 | 4.0 | 14.0 | 38.5 |
| 1977 | 2.0 | * | * | * | * | * | 0.3 | 0.7 | 1.7 | 5.3 | 16.7 | 51.1 |
| 1976 | 2.1 | * | * | * | * | 0.1 | 0.4 | 0.8 | 2.3 | 5.9 | 17.7 | 49.5 |
| 1975 | 2.4 | * | * | * | * | * | 0.4 | 0.9 | 2.4 | 7.5 | 21.9 | 55.9 |
| 1974 | 2.9 | * | * | * | 0.1 | 0.2 | 0.6 | 1.1 | 2.8 | 9.2 | 27.1 | 64.1 |
| 1973 | 3.3 | * | * | * | 0.1 | 0.2 | 0.8 | 1.4 | 4.1 | 10.8 | 28.5 | 77.0 |
| 1972 | 3.6 | * | * | * | 0.1 | 0.2 | 0.6 | 2.2 | 4.5 | 11.5 | 34.2 | 79.2 |
| 1971 | 3.9 | * | * | * | 0.2 | 0.4 | 1.0 | 2.3 | 5.6 | 12.6 | 36.1 | 83.3 |
| 1970 | 4.4 | * | * | 0.1 | 0.2 | 0.4 | 1.3 | 2.8 | 6.4 | 14.9 | 38.9 | 87.2 |
| 1969 | 4.6 | * | * | * | 0.2 | 0.6 | 1.4 | 3.1 | 6.3 | 16.2 | 40.6 | 106.5 |
| 1968 | 4.9 | * | * | 0.1 | 0.2 | 0.6 | 1.6 | 3.9 | 8.1 | 16.7 | 41.5 | 109.3 |
| WHITE, BOTH SEXES | | | | | | | | | | | | |
| 1978 | 1.4 | * | * | * | * | * | 0.2 | 0.4 | 1.1 | 3.7 | 14.3 | 42.5 |
| 1977 | 1.6 | * | * | * | * | * | 0.1 | 0.5 | 1.4 | 4.5 | 17.4 | 52.7 |
| 1976 | 1.8 | * | * | * | * | 0.1 | 0.2 | 0.5 | 1.8 | 5.3 | 19.2 | 53.8 |
| 1975 | 2.0 | * | * | * | * | * | 0.3 | 0.6 | 1.9 | 6.2 | 23.0 | 61.3 |
| 1974 | 2.5 | * | * | * | 0.1 | 0.1 | 0.3 | 0.8 | 2.3 | 8.6 | 28.4 | 71.2 |
| 1973 | 2.8 | * | * | * | 0.1 | 0.1 | 0.5 | 0.8 | 3.0 | 10.0 | 30.7 | 82.7 |
| 1972 | 3.2 | * | * | * | 0.1 | 0.2 | 0.4 | 1.2 | 3.3 | 10.9 | 37.1 | 88.9 |
| 1971 | 3.5 | * | * | * | 0.1 | 0.3 | 0.5 | 1.4 | 4.1 | 12.1 | 39.3 | 98.0 |
| 1970 | 3.8 | * | * | 0.1 | 0.2 | 0.3 | 0.6 | 1.8 | 4.7 | 13.4 | 42.0 | 101.0 |
| 1969 | 4.1 | * | * | 0.1 | 0.1 | 0.3 | 0.7 | 1.8 | 4.9 | 15.3 | 45.4 | 124.0 |
| 1968 | 4.4 | 0.8 | * | 0.1 | 0.2 | 0.4 | 0.9 | 2.4 | 6.0 | 16.1 | 46.8 | 128.8 |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| INFECTIONS OF KIDNEY (590) | | | | | | | | | | | | |
| WHITE, MALE | | | | | | | | | | | | |
| 1978 | 1.1 | * | * | * | * | * | * | 0.3 | 0.9 | 3.8 | 15.3 | 51.0 |
| 1977 | 1.3 | * | * | * | * | * | * | 0.3 | 1.3 | 4.3 | 19.3 | 57.8 |
| 1976 | 1.5 | * | * | * | * | * | * | 0.4 | 1.6 | 5.5 | 23.0 | 63.0 |
| 1975 | 1.7 | * | * | * | * | * | 0.2 | 0.4 | 1.7 | 5.8 | 24.8 | 71.7 |
| 1974 | 2.2 | * | * | * | * | * | 0.2 | 0.6 | 2.3 | 9.2 | 30.6 | 83.9 |
| 1973 | 2.4 | * | * | * | * | * | 0.3 | 0.7 | 2.7 | 10.2 | 34.2 | 94.5 |
| 1972 | 2.9 | * | * | * | * | 0.2 | 0.5 | 0.9 | 2.8 | 12.6 | 43.4 | 106.1 |
| 1971 | 3.2 | * | * | * | 0.1 | 0.3 | 0.3 | 1.2 | 3.6 | 13.6 | 45.6 | 124.0 |
| 1970 | 3.4 | * | * | * | 0.1 | 0.3 | 0.4 | 1.4 | 4.4 | 14.1 | 47.7 | 121.8 |
| 1969 | 3.8 | * | * | * | 0.1 | * | 0.6 | 1.5 | 4.9 | 16.4 | 53.9 | 153.3 |
| 1968 | 4.2 | * | * | * | 0.2 | 0.3 | 0.7 | 2.0 | 5.5 | 18.2 | 55.7 | 161.7 |
| WHITE, FEMALE | | | | | | | | | | | | |
| 1978 | 1.6 | * | * | * | * | * | * | 0.6 | 1.3 | 3.7 | 13.8 | 38.7 |
| 1977 | 2.0 | * | * | * | * | * | * | 0.6 | 1.4 | 4.7 | 16.3 | 50.4 |
| 1976 | 2.1 | * | * | * | * | * | 0.3 | 0.7 | 2.0 | 5.2 | 17.0 | 49.6 |
| 1975 | 2.4 | * | * | * | * | * | 0.3 | 0.7 | 2.1 | 6.5 | 21.9 | 56.4 |
| 1974 | 2.8 | * | * | * | * | * | 0.4 | 0.9 | 2.2 | 8.1 | 27.1 | 65.0 |
| 1973 | 3.2 | * | * | * | * | 0.2 | 0.6 | 1.0 | 3.3 | 9.8 | 28.5 | 76.9 |
| 1972 | 3.4 | * | * | * | 0.1 | * | 0.3 | 1.5 | 3.8 | 9.7 | 33.2 | 80.3 |
| 1971 | 3.7 | * | * | * | 0.2 | 0.3 | 0.6 | 1.6 | 4.5 | 10.9 | 35.2 | 84.7 |
| 1970 | 4.2 | * | * | 0.1 | 0.2 | 0.3 | 0.8 | 2.2 | 5.0 | 12.9 | 38.3 | 89.7 |
| 1969 | 4.4 | * | * | * | 0.1 | 0.5 | 0.8 | 2.1 | 5.0 | 14.5 | 39.9 | 108.7 |
| 1968 | 4.6 | * | * | * | 0.1 | 0.5 | 1.0 | 2.8 | 6.4 | 14.4 | 40.8 | 111.2 |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| INFECTIONS OF KIDNEY (590) | | | | | | | | | | | | |
| ALL OTHER, BOTH SEXES | | | | | | | | | | | | |
| 1978 | 1.4 | * | * | * | * | * | * | 1.1 | 2.6 | 8.4 | 19.7 | 44.0 |
| 1977 | 2.0 | * | * | * | * | * | * | 1.8 | 3.7 | 11.9 | 24.0 | 61.9 |
| 1976 | 2.1 | * | * | * | * | * | 0.7 | 1.3 | 4.7 | 14.1 | 26.2 | 59.5 |
| 1975 | 2.4 | * | * | * | * | * | 1.0 | 2.2 | 5.8 | 17.2 | 25.2 | 57.5 |
| 1974 | 2.9 | * | * | * | * | 0.6 | 1.2 | 2.7 | 7.6 | 20.8 | 31.7 | 63.7 |
| 1973 | 3.5 | * | * | * | * | * | 1.6 | 3.8 | 10.1 | 22.5 | 37.9 | 83.5 |
| 1972 | 4.4 | * | * | * | * | * | 2.2 | 6.8 | 10.3 | 31.2 | 48.9 | 81.3 |
| 1971 | 5.0 | * | * | * | * | 0.9 | 2.4 | 7.0 | 14.7 | 31.3 | 54.6 | 80.8 |
| 1970 | 5.7 | * | * | * | * | 1.0 | 3.5 | 7.4 | 18.3 | 37.0 | 58.2 | 84.3 |
| 1969 | 6.1 | * | * | * | * | 1.3 | 4.0 | 8.8 | 18.3 | 37.1 | 63.8 | 102.8 |
| 1968 | 7.1 | * | * | * | * | 1.2 | 4.5 | 11.2 | 23.1 | 44.2 | 68.8 | 105.9 |
| ALL OTHER, MALE | | | | | | | | | | | | |
| 1978 | 1.4 | * | * | * | * | * | * | * | 2.2 | 9.5 | 24.4 | 60.3 |
| 1977 | 1.8 | * | * | * | * | * | * | 1.9 | 3.2 | 12.8 | 28.7 | 66.1 |
| 1976 | 2.0 | * | * | * | * | * | * | * | 3.9 | 15.8 | 28.0 | 78.2 |
| 1975 | 2.2 | * | * | * | * | * | * | 2.0 | 5.9 | 17.0 | 30.2 | 70.4 |
| 1974 | 2.7 | * | * | * | * | * | * | 2.1 | 7.0 | 21.5 | 37.6 | 82.0 |
| 1973 | 3.3 | * | * | * | * | * | * | 2.8 | 8.8 | 24.0 | 52.2 | 91.7 |
| 1972 | 4.0 | * | * | * | * | * | 1.7 | 6.0 | 9.0 | 32.4 | 50.7 | 106.4 |
| 1971 | 4.5 | * | * | * | * | * | * | 5.2 | 13.4 | 33.1 | 64.7 | 104.3 |
| 1970 | 5.6 | * | * | * | * | * | 2.2 | 6.6 | 17.3 | 39.1 | 74.1 | 119.1 |
| 1969 | 5.8 | * | * | * | * | * | 2.5 | 6.0 | 17.8 | 40.6 | 80.3 | 136.4 |
| 1968 | 6.9 | * | * | * | * | * | 3.1 | 8.7 | 21.5 | 49.3 | 90.9 | 136.6 |
| ALL OTHER, FEMALE | | | | | | | | | | | | |
| 1978 | 1.5 | * | * | * | * | * | * | * | 2.9 | 7.6 | 16.5 | 35.8 |
| 1977 | 2.1 | * | * | * | * | * | * | 1.6 | 4.0 | 11.2 | 20.6 | 59.6 |
| 1976 | 2.2 | * | * | * | * | * | * | 1.5 | 5.5 | 12.8 | 25.0 | 49.0 |
| 1975 | 2.5 | * | * | * | * | * | * | 2.4 | 5.7 | 17.3 | 21.8 | 49.5 |
| 1974 | 3.1 | * | * | * | * | * | 1.5 | 3.2 | 8.2 | 20.3 | 27.7 | 52.9 |
| 1973 | 3.6 | * | * | * | * | * | 1.9 | 4.6 | 11.3 | 21.3 | 28.0 | 78.5 |
| 1972 | 4.7 | * | * | * | * | * | 2.6 | 7.6 | 11.4 | 30.3 | 47.8 | 65.8 |
| 1971 | 5.3 | * | * | * | * | * | 3.7 | 8.6 | 15.8 | 29.8 | 47.1 | 65.8 |
| 1970 | 5.8 | * | * | * | * | 1.4 | 4.6 | 8.1 | 19.3 | 35.2 | 46.1 | 59.7 |
| 1969 | 6.3 | * | * | * | * | 1.7 | 5.1 | 11.2 | 18.8 | 34.3 | 51.0 | 80.0 |
| 1968 | 7.3 | * | * | * | * | 1.6 | 5.8 | 13.4 | 24.6 | 39.9 | 51.9 | 85.2 |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| INFECTIONS OF KIDNEY (590) | | | | | | | | | | | | |
| BLACK, BOTH SEXES | | | | | | | | | | | | |
| 1978 | 1.6 | * | * | * | * | * | * | 1.2 | 2.9 | 9.0 | 19.3 | 44.0 |
| 1977 | 2.1 | * | * | * | * | * | * | 2.0 | 4.1 | 12.6 | 24.2 | 64.3 |
| 1976 | 2.3 | * | * | * | * | * | * | 1.5 | 4.9 | 15.2 | 27.1 | 58.8 |
| 1975 | 2.5 | * | * | * | * | * | 1.0 | 2.4 | 6.2 | 17.9 | 25.6 | 60.8 |
| 1974 | 3.1 | * | * | * | * | 0.7 | 1.2 | 2.9 | 8.3 | 22.2 | 32.4 | 63.6 |
| 1973 | 3.7 | * | * | * | * | * | 1.8 | 4.1 | 10.8 | 23.9 | 38.9 | 85.1 |
| 1972 | 4.7 | * | * | * | * | * | 2.4 | 7.4 | 11.1 | 32.1 | 50.0 | 85.5 |
| 1971 | 5.3 | * | * | * | * | 0.9 | 2.7 | 7.6 | 15.9 | 32.7 | 56.6 | 81.3 |
| 1970 | 6.1 | * | * | * | * | 1.1 | 3.9 | 8.0 | 19.4 | 39.7 | 61.5 | 87.7 |
| 1969 | 6.4 | * | * | * | * | 1.4 | 4.3 | 9.3 | 19.4 | 39.3 | 66.4 | 104.1 |
| 1968 | 7.5 | * | * | * | * | 1.2 | 4.9 | 11.7 | 24.7 | 46.3 | 70.3 | 104.3 |
| BLACK, MALE | | | | | | | | | | | | |
| 1978 | 1.5 | * | * | * | * | * | * | * | 2.5 | 10.5 | 23.8 | 62.7 |
| 1977 | 2.0 | * | * | * | * | * | * | 2.2 | 3.7 | 13.8 | 28.5 | 73.5 |
| 1976 | 2.2 | * | * | * | * | * | * | * | 4.2 | 17.6 | 29.5 | 87.2 |
| 1975 | 2.4 | * | * | * | * | * | * | 2.3 | 6.5 | 18.5 | 30.2 | 80.4 |
| 1974 | 2.9 | * | * | * | * | * | * | 2.3 | 7.9 | 22.6 | 39.7 | 81.4 |
| 1973 | 3.6 | * | * | * | * | * | * | 3.2 | 9.8 | 25.6 | 53.9 | 104.9 |
| 1972 | 4.3 | * | * | * | * | * | 2.0 | 6.3 | 10.1 | 34.5 | 50.8 | 120.0 |
| 1971 | 4.8 | * | * | * | * | * | * | 5.6 | 14.8 | 35.1 | 67.2 | 105.0 |
| 1970 | 6.0 | * | * | * | * | * | 2.4 | 7.2 | 18.5 | 42.7 | 81.8 | 123.9 |
| 1969 | 6.1 | * | * | * | * | * | 2.8 | 6.4 | 19.8 | 43.5 | 85.0 | 137.8 |
| 1968 | 7.3 | * | * | * | * | * | 3.4 | 9.4 | 23.6 | 51.9 | 94.5 | 148.6 |
| BLACK, FEMALE | | | | | | | | | | | | |
| 1978 | 1.6 | * | * | * | * | * | * | * | 3.1 | 7.9 | 16.2 | 34.3 |
| 1977 | 2.3 | * | * | * | * | * | * | 1.8 | 4.4 | 11.6 | 21.5 | 59.6 |
| 1976 | 2.4 | * | * | * | * | * | * | 1.6 | 5.5 | 13.4 | 25.5 | 43.8 |
| 1975 | 2.6 | * | * | * | * | * | * | 2.5 | 5.9 | 17.4 | 22.6 | 50.0 |
| 1974 | 3.4 | * | * | * | * | * | 1.5 | 3.4 | 8.6 | 21.9 | 27.5 | 53.8 |
| 1973 | 3.9 | * | * | * | * | * | 2.1 | 4.9 | 11.7 | 22.5 | 28.7 | 74.0 |
| 1972 | 5.0 | * | * | * | * | * | 2.7 | 8.3 | 11.9 | 30.2 | 49.3 | 65.7 |
| 1971 | 5.7 | * | * | * | * | * | 4.1 | 9.3 | 16.8 | 30.7 | 49.0 | 67.2 |
| 1970 | 6.2 | * | * | * | * | 1.4 | 5.1 | 8.6 | 20.2 | 37.3 | 46.5 | 64.4 |
| 1969 | 6.7 | * | * | * | * | 1.8 | 5.6 | 11.7 | 19.2 | 36.0 | 52.4 | 83.3 |
| 1968 | 7.6 | * | * | * | * | 1.6 | 6.3 | 13.7 | 25.7 | 41.7 | 52.3 | 78.6 |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| HYPERPLASIA OF PROSTATE (600) | | | | | | | | | | | | |
| BOTH SEXES | | | | | | | | | | | | |
| 1978 | 0.4 | * | * | * | * | * | * | * | 0.1 | 1.1 | 5.0 | 13.8 |
| 1977 | 0.5 | * | * | * | * | * | * | * | 0.2 | 1.3 | 5.8 | 16.7 |
| 1976 | 0.5 | * | * | * | * | * | * | * | 0.2 | 1.6 | 6.1 | 19.5 |
| 1975 | 0.6 | * | * | * | * | * | * | * | 0.3 | 1.9 | 7.4 | 20.6 |
| 1974 | 0.6 | * | * | * | * | * | * | * | 0.3 | 2.2 | 8.4 | 25.2 |
| 1973 | 0.8 | * | * | * | * | * | * | * | 0.4 | 2.5 | 10.4 | 30.2 |
| 1972 | 0.9 | * | * | * | * | * | * | * | 0.4 | 3.0 | 13.2 | 33.5 |
| 1971 | 0.9 | * | * | * | * | * | * | * | 0.5 | 3.2 | 12.9 | 38.7 |
| 1970 | 1.1 | * | * | * | * | * | * | * | 0.6 | 4.0 | 15.5 | 40.0 |
| 1969 | 1.2 | * | * | * | * | * | * | 0.1 | 0.7 | 4.9 | 17.3 | 53.1 |
| 1968 | 1.3 | * | * | * | * | * | * | * | 0.9 | 5.5 | 19.1 | 54.1 |
| MALE | | | | | | | | | | | | |
| 1978 | 0.8 | * | * | * | * | * | * | * | 0.3 | 2.5 | 13.5 | 44.3 |
| 1977 | 0.9 | * | * | * | * | * | * | * | 0.5 | 3.1 | 15.5 | 53.0 |
| 1976 | 1.0 | * | * | * | * | * | * | * | 0.4 | 3.7 | 16.2 | 61.1 |
| 1975 | 1.2 | * | * | * | * | * | * | * | 0.6 | 4.3 | 19.6 | 63.3 |
| 1974 | 1.3 | * | * | * | * | * | * | * | 0.7 | 5.2 | 22.0 | 76.1 |
| 1973 | 1.5 | * | * | * | * | * | * | * | 0.8 | 5.8 | 26.9 | 90.3 |
| 1972 | 1.8 | * | * | * | * | * | * | * | 0.9 | 6.9 | 33.8 | 98.9 |
| 1971 | 1.9 | * | * | * | * | * | * | * | 1.0 | 7.4 | 32.8 | 112.7 |
| 1970 | 2.2 | * | * | * | * | * | * | * | 1.2 | 9.2 | 38.8 | 111.5 |
| 1969 | 2.6 | * | * | * | * | * | * | 0.2 | 1.5 | 11.1 | 43.0 | 151.2 |
| 1968 | 2.7 | * | * | * | * | * | * | * | 1.8 | 12.4 | 47.0 | 152.7 |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| HYPERPLASIA OF PROSTATE (600) | | | | | | | | | | | | |
| FEMALE | | | | | | | | | | | | |
| 1978 | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 1977 | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 1976 | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 1975 | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 1974 | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 1973 | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 1972 | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 1971 | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 1970 | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 1969 | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 1968 | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| WHITE, BOTH SEXES | | | | | | | | | | | | |
| 1978 | 0.4 | * | * | * | * | * | * | * | * | 0.9 | 4.8 | 13.6 |
| 1977 | 0.4 | * | * | * | * | * | * | * | 0.2 | 1.1 | 5.5 | 16.3 |
| 1976 | 0.5 | * | * | * | * | * | * | * | 0.2 | 1.4 | 5.9 | 19.7 |
| 1975 | 0.6 | * | * | * | * | * | * | * | 0.2 | 1.6 | 7.4 | 20.8 |
| 1974 | 0.7 | * | * | * | * | * | * | * | 0.3 | 2.0 | 8.1 | 25.7 |
| 1973 | 0.8 | * | * | * | * | * | * | * | 0.3 | 2.3 | 10.1 | 30.5 |
| 1972 | 0.9 | * | * | * | * | * | * | * | 0.3 | 2.7 | 12.7 | 33.3 |
| 1971 | 0.9 | * | * | * | * | * | * | * | 0.4 | 2.8 | 12.5 | 39.1 |
| 1970 | 1.1 | * | * | * | * | * | * | * | 0.5 | 3.6 | 14.8 | 41.7 |
| 1969 | 1.2 | * | * | * | * | * | * | * | 0.6 | 4.1 | 16.6 | 53.8 |
| 1968 | 1.3 | * | * | * | * | * | * | * | 0.7 | 4.8 | 18.3 | 54.8 |

... Category not applicable

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| HYPERPLASIA OF PROSTATE (600) | | | | | | | | | | | | |
| WHITE, MALE | | | | | | | | | | | | |
| 1978 | 0.8 | * | * | * | * | * | * | * | * | 2.1 | 13.0 | 44.3 |
| 1977 | 0.9 | * | * | * | * | * | * | * | 0.5 | 2.6 | 14.7 | 52.4 |
| 1976 | 1.0 | * | * | * | * | * | * | * | 0.4 | 3.3 | 15.8 | 62.3 |
| 1975 | 1.2 | * | * | * | * | * | * | * | 0.5 | 3.8 | 19.6 | 64.4 |
| 1974 | 1.3 | * | * | * | * | * | * | * | 0.6 | 4.6 | 21.3 | 78.4 |
| 1973 | 1.6 | * | * | * | * | * | * | * | 0.7 | 5.3 | 26.5 | 92.0 |
| 1972 | 1.8 | * | * | * | * | * | * | * | 0.7 | 6.2 | 32.9 | 99.4 |
| 1971 | 1.9 | * | * | * | * | * | * | * | 0.9 | 6.5 | 32.1 | 115.3 |
| 1970 | 2.2 | * | * | * | * | * | * | * | 1.0 | 8.2 | 37.3 | 117.9 |
| 1969 | 2.5 | * | * | * | * | * | * | * | 1.2 | 9.4 | 41.6 | 155.4 |
| 1968 | 2.7 | * | * | * | * | * | * | * | 1.4 | 11.0 | 45.3 | 156.4 |
| WHITE, FEMALE | | | | | | | | | | | | |
| 1978 | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 1977 | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 1976 | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 1975 | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 1974 | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 1973 | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 1972 | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 1971 | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 1970 | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 1969 | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 1968 | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |

* Figure does not meet standards of reliability or precision.

... Category not applicable

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| HYPERPLASIA OF PROSTATE (600) | | | | | | | | | | | | |
| ALL OTHER, BOTH SEXES | | | | | | | | | | | | |
| 1978 | 0.4 | * | * | * | * | * | * | * | * | 2.7 | 7.5 | 15.5 |
| 1977 | 0.5 | * | * | * | * | * | * | * | * | 3.3 | 9.2 | 20.6 |
| 1976 | 0.5 | * | * | * | * | * | * | * | * | 3.4 | 8.2 | 17.6 |
| 1975 | 0.5 | * | * | * | * | * | * | * | * | 4.2 | 8.2 | 19.2 |
| 1974 | 0.6 | * | * | * | * | * | * | * | * | 4.7 | 11.8 | 19.3 |
| 1973 | 0.7 | * | * | * | * | * | * | * | * | 4.9 | 12.7 | 27.6 |
| 1972 | 0.9 | * | * | * | * | * | * | * | 1.2 | 5.7 | 18.1 | 35.8 |
| 1971 | 0.9 | * | * | * | * | * | * | * | * | 7.5 | 17.5 | 34.2 |
| 1970 | 1.1 | * | * | * | * | * | * | * | 1.6 | 8.4 | 24.4 | 23.1 |
| 1969 | 1.4 | * | * | * | * | * | * | * | 1.9 | 12.5 | 25.8 | 45.0 |
| 1968 | 1.5 | * | * | * | * | * | * | * | 2.9 | 11.7 | 29.3 | 47.1 |
| ALL OTHER, MALE | | | | | | | | | | | | |
| 1978 | 0.9 | * | * | * | * | * | * | * | * | 6.2 | 18.4 | 44.8 |
| 1977 | 1.0 | * | * | * | * | * | * | * | * | 7.5 | 22.6 | 58.9 |
| 1976 | 1.0 | * | * | * | * | * | * | * | * | 7.7 | 20.2 | 49.1 |
| 1975 | 1.1 | * | * | * | * | * | * | * | * | 9.4 | 20.2 | 51.9 |
| 1974 | 1.3 | * | * | * | * | * | * | * | * | 10.6 | 28.8 | 52.0 |
| 1973 | 1.4 | * | * | * | * | * | * | * | * | 11.0 | 30.8 | 72.9 |
| 1972 | 1.9 | * | * | * | * | * | * | * | 2.6 | 12.8 | 43.3 | 93.6 |
| 1971 | 1.9 | * | * | * | * | * | * | * | * | 16.8 | 41.1 | 87.2 |
| 1970 | 2.2 | * | * | * | * | * | * | * | 3.5 | 18.6 | 56.5 | 55.9 |
| 1969 | 2.9 | * | * | * | * | * | * | * | 4.1 | 27.6 | 59.6 | 111.4 |
| 1968 | 3.1 | * | * | * | * | * | * | * | 6.1 | 25.7 | 67.2 | 117.1 |
| ALL OTHER, FEMALE | | | | | | | | | | | | |
| 1978 | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 1977 | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 1976 | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 1975 | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 1974 | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 1973 | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 1972 | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 1971 | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 1970 | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 1969 | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 1968 | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |

* Figure does not meet standards of reliability or precision.

... Category not applicable

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| HYPERPLASIA OF PROSTATE (600) | | | | | | | | | | | | |
| BLACK, BOTH SEXES | | | | | | | | | | | | |
| 1978 | 0.5 | * | * | * | * | * | * | * | * | 2.9 | 8.2 | 14.0 |
| 1977 | 0.5 | * | * | * | * | * | * | * | * | 3.4 | 9.3 | 19.6 |
| 1976 | 0.5 | * | * | * | * | * | * | * | * | 3.8 | 8.4 | 17.6 |
| 1975 | 0.6 | * | * | * | * | * | * | * | * | 4.4 | 8.5 | 19.2 |
| 1974 | 0.7 | * | * | * | * | * | * | * | * | 5.1 | 12.2 | 20.7 |
| 1973 | 0.7 | * | * | * | * | * | * | * | * | 5.3 | 12.8 | 28.9 |
| 1972 | 1.0 | * | * | * | * | * | * | * | 1.3 | 6.2 | 18.9 | 40.0 |
| 1971 | 1.0 | * | * | * | * | * | * | * | * | 8.0 | 18.4 | 35.5 |
| 1970 | 1.1 | * | * | * | * | * | * | * | 1.8 | 9.2 | 24.8 | 24.7 |
| 1969 | 1.5 | * | * | * | * | * | * | * | 2.0 | 13.0 | 27.2 | 48.5 |
| 1968 | 1.6 | * | * | * | * | * | * | * | 3.1 | 12.1 | 31.5 | 47.8 |
| BLACK, MALE | | | | | | | | | | | | |
| 1978 | 1.0 | * | * | * | * | * | * | * | * | 6.9 | 20.5 | 41.2 |
| 1977 | 1.1 | * | * | * | * | * | * | * | * | 7.8 | 23.4 | 57.1 |
| 1976 | 1.1 | * | * | * | * | * | * | * | * | 8.9 | 20.9 | 51.1 |
| 1975 | 1.2 | * | * | * | * | * | * | * | * | 10.2 | 21.2 | 54.3 |
| 1974 | 1.4 | * | * | * | * | * | * | * | * | 11.8 | 30.1 | 58.1 |
| 1973 | 1.5 | * | * | * | * | * | * | * | * | 12.1 | 31.4 | 80.5 |
| 1972 | 2.1 | * | * | * | * | * | * | * | 2.9 | 14.1 | 45.7 | 110.0 |
| 1971 | 2.0 | * | * | * | * | * | * | * | * | 18.2 | 44.1 | 95.0 |
| 1970 | 2.4 | * | * | * | * | * | * | * | 3.9 | 20.8 | 58.7 | 63.0 |
| 1969 | 3.1 | * | * | * | * | * | * | * | 4.4 | 29.5 | 64.1 | 127.0 |
| 1968 | 3.3 | * | * | * | * | * | * | * | 6.7 | 27.2 | 73.9 | 125.7 |
| BLACK, FEMALE | | | | | | | | | | | | |
| 1978 | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 1977 | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 1976 | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 1975 | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 1974 | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 1973 | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 1972 | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 1971 | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 1970 | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 1969 | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 1968 | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |

* Figure does not meet standards of reliability or precision.

... Category not applicable

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|--|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| COMPLICATIONS OF PREGNANCY, CHILDBIRTH, AND THE PUERPERIUM (630-678) | | | | | | | | | | | | |
| BOTH SEXES | | | | | | | | | | | | |
| 1978 | 0.1 | * | * | * | 0.3 | 0.4 | 0.2 | * | * | * | * | * |
| 1977 | 0.2 | * | * | * | 0.3 | 0.5 | 0.3 | * | * | * | * | * |
| 1976 | 0.2 | * | * | * | 0.3 | 0.5 | 0.3 | * | * | * | * | * |
| 1975 | 0.2 | * | * | * | 0.4 | 0.6 | 0.3 | * | * | * | * | * |
| 1974 | 0.2 | * | * | * | 0.4 | 0.7 | 0.3 | * | * | * | * | * |
| 1973 | 0.2 | * | * | * | 0.4 | 0.7 | 0.4 | * | * | * | * | * |
| 1972 | 0.3 | * | * | * | 0.7 | 0.9 | 0.5 | * | * | * | * | * |
| 1971 | 0.3 | * | * | * | 0.7 | 1.0 | 0.7 | * | * | * | * | * |
| 1970 | 0.4 | * | * | * | 0.9 | 1.2 | 0.8 | * | * | * | * | * |
| 1969 | 0.4 | * | * | * | 0.8 | 1.3 | 0.8 | * | * | * | * | * |
| 1968 | 0.4 | * | * | * | 0.8 | 1.5 | 0.9 | * | * | * | * | * |
| MALE | | | | | | | | | | | | |
| 1978 | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 1977 | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 1976 | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 1975 | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 1974 | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 1973 | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 1972 | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 1971 | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 1970 | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 1969 | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 1968 | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |

* Figure does not meet standards of reliability or precision.

... Category not applicable

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|--|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| COMPLICATIONS OF PREGNANCY, CHILDBIRTH, AND THE PUERPERIUM (630-678) | | | | | | | | | | | | |
| FEMALE | | | | | | | | | | | | |
| 1978 | 0.3 | * | * | * | 0.6 | 0.8 | 0.4 | * | * | * | * | * |
| 1977 | 0.3 | * | * | * | 0.6 | 1.0 | 0.5 | * | * | * | * | * |
| 1976 | 0.3 | * | * | * | 0.7 | 1.1 | 0.6 | * | * | * | * | * |
| 1975 | 0.4 | * | * | * | 0.7 | 1.1 | 0.6 | * | * | * | * | * |
| 1974 | 0.4 | * | * | * | 0.9 | 1.4 | 0.6 | * | * | * | * | * |
| 1973 | 0.4 | * | * | * | 0.9 | 1.5 | 0.7 | * | * | * | * | * |
| 1972 | 0.6 | * | * | * | 1.3 | 1.8 | 0.9 | * | * | * | * | * |
| 1971 | 0.6 | * | * | * | 1.4 | 2.0 | 1.3 | * | * | * | * | * |
| 1970 | 0.8 | * | * | * | 1.7 | 2.4 | 1.6 | * | * | * | * | * |
| 1969 | 0.8 | * | * | * | 1.6 | 2.5 | 1.6 | * | * | * | * | * |
| 1968 | 0.8 | * | * | * | 1.6 | 2.8 | 1.8 | * | * | * | * | * |
| WHITE, BOTH SEXES | | | | | | | | | | | | |
| 1978 | 0.1 | * | * | * | 0.2 | 0.3 | 0.1 | * | * | * | * | * |
| 1977 | 0.1 | * | * | * | 0.2 | 0.4 | 0.2 | * | * | * | * | * |
| 1976 | 0.1 | * | * | * | 0.2 | 0.4 | 0.2 | * | * | * | * | * |
| 1975 | 0.1 | * | * | * | 0.2 | 0.4 | 0.2 | * | * | * | * | * |
| 1974 | 0.1 | * | * | * | 0.3 | 0.5 | 0.2 | * | * | * | * | * |
| 1973 | 0.1 | * | * | * | 0.3 | 0.5 | 0.2 | * | * | * | * | * |
| 1972 | 0.2 | * | * | * | 0.5 | 0.7 | 0.3 | * | * | * | * | * |
| 1971 | 0.2 | * | * | * | 0.4 | 0.8 | 0.4 | * | * | * | * | * |
| 1970 | 0.3 | * | * | * | 0.5 | 0.8 | 0.6 | * | * | * | * | * |
| 1969 | 0.3 | * | * | * | 0.5 | 0.9 | 0.5 | * | * | * | * | * |
| 1968 | 0.3 | * | * | * | 0.5 | 0.9 | 0.7 | * | * | * | * | * |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|--|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| COMPLICATIONS OF PREGNANCY, CHILDBIRTH, AND THE PUERPERIUM (630-678) | | | | | | | | | | | | |
| WHITE, MALE | | | | | | | | | | | | |
| 1978 | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 1977 | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 1976 | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 1975 | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 1974 | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 1973 | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 1972 | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 1971 | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 1970 | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 1969 | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 1968 | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| WHITE, FEMALE | | | | | | | | | | | | |
| 1978 | 0.2 | * | * | * | 0.3 | 0.5 | 0.3 | * | * | * | * | * |
| 1977 | 0.2 | * | * | * | 0.4 | 0.7 | 0.3 | * | * | * | * | * |
| 1976 | 0.2 | * | * | * | 0.5 | 0.7 | 0.4 | * | * | * | * | * |
| 1975 | 0.2 | * | * | * | 0.5 | 0.7 | 0.5 | * | * | * | * | * |
| 1974 | 0.3 | * | * | * | 0.5 | 0.9 | 0.5 | * | * | * | * | * |
| 1973 | 0.3 | * | * | * | 0.6 | 1.1 | 0.4 | * | * | * | * | * |
| 1972 | 0.4 | * | * | * | 1.0 | 1.3 | 0.6 | * | * | * | * | * |
| 1971 | 0.4 | * | * | * | 0.8 | 1.5 | 0.8 | * | * | * | * | * |
| 1970 | 0.5 | * | * | * | 1.1 | 1.5 | 1.1 | * | * | * | * | * |
| 1969 | 0.5 | * | * | * | 1.0 | 1.8 | 1.1 | * | * | * | * | * |
| 1968 | 0.5 | * | * | * | 1.0 | 1.8 | 1.3 | * | * | * | * | * |

... Category not applicable

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|--|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| COMPLICATIONS OF PREGNANCY, CHILDBIRTH, AND THE PUERPERIUM (630-678) | | | | | | | | | | | | |
| ALL OTHER, BOTH SEXES | | | | | | | | | | | | |
| 1978 | 0.5 | * | * | * | 0.9 | 1.4 | 0.7 | * | * | * | * | * |
| 1977 | 0.6 | * | * | * | 1.0 | 1.6 | 0.9 | * | * | * | * | * |
| 1976 | 0.6 | * | * | * | 0.9 | 1.7 | 1.0 | * | * | * | * | * |
| 1975 | 0.6 | * | * | * | 1.1 | 1.9 | 0.9 | * | * | * | * | * |
| 1974 | 0.7 | * | * | * | 1.6 | 2.3 | 0.9 | * | * | * | * | * |
| 1973 | 0.8 | * | * | * | 1.5 | 2.1 | 1.4 | * | * | * | * | * |
| 1972 | 0.9 | * | * | * | 1.8 | 2.6 | 1.7 | * | * | * | * | * |
| 1971 | 1.1 | * | * | * | 2.5 | 2.6 | 2.7 | * | * | * | * | * |
| 1970 | 1.4 | * | * | * | 2.9 | 4.4 | 2.8 | * | * | * | * | * |
| 1969 | 1.4 | * | * | * | 2.7 | 3.9 | 3.2 | * | * | * | * | * |
| 1968 | 1.5 | * | * | * | 3.0 | 5.4 | 3.1 | * | * | * | * | * |
| ALL OTHER, MALE | | | | | | | | | | | | |
| 1978 | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 1977 | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 1976 | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 1975 | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 1974 | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 1973 | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 1972 | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 1971 | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 1970 | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 1969 | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 1968 | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| ALL OTHER, FEMALE | | | | | | | | | | | | |
| 1978 | 0.9 | * | * | * | 1.9 | 2.6 | 1.4 | * | * | * | * | * |
| 1977 | 1.1 | * | * | * | 1.9 | 3.1 | 1.7 | * | * | * | * | * |
| 1976 | 1.1 | * | * | * | 1.8 | 3.2 | 1.8 | * | * | * | * | * |
| 1975 | 1.2 | * | * | * | 2.2 | 3.5 | 1.7 | * | * | * | * | * |
| 1974 | 1.4 | * | * | * | 3.1 | 4.2 | 1.6 | * | * | * | * | * |
| 1973 | 1.4 | * | * | * | 2.9 | 4.0 | 2.5 | * | * | * | * | * |
| 1972 | 1.7 | * | * | * | 3.5 | 4.8 | 3.2 | * | * | * | * | * |
| 1971 | 2.1 | * | * | * | 4.9 | 4.8 | 4.9 | * | * | * | * | * |
| 1970 | 2.7 | * | * | * | 5.6 | 8.2 | 5.1 | * | * | * | * | * |
| 1969 | 2.6 | * | * | * | 5.3 | 7.3 | 5.9 | * | * | * | * | * |
| 1968 | 3.0 | * | * | * | 5.8 | 9.9 | 5.7 | * | * | * | * | * |

* Figure does not meet standards of reliability or precision.

... Category not applicable

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|--|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| COMPLICATIONS OF PREGNANCY, CHILDBIRTH, AND THE PUERPERIUM (630-678) | | | | | | | | | | | | |
| BLACK, BOTH SEXES | | | | | | | | | | | | |
| 1978 | 0.5 | * | * | * | 1.0 | 1.5 | 0.8 | * | * | * | * | * |
| 1977 | 0.6 | * | * | * | 1.1 | 1.9 | 1.1 | * | * | * | * | * |
| 1976 | 0.6 | * | * | * | 1.0 | 2.0 | 1.1 | * | * | * | * | * |
| 1975 | 0.6 | * | * | * | 1.3 | 1.9 | 1.1 | * | * | * | * | * |
| 1974 | 0.8 | * | * | * | 1.8 | 2.4 | 0.9 | * | * | * | * | * |
| 1973 | 0.8 | * | * | * | 1.6 | 2.4 | 1.6 | * | * | * | * | * |
| 1972 | 0.9 | * | * | * | 1.9 | 2.8 | 1.8 | * | * | * | * | * |
| 1971 | 1.2 | * | * | * | 2.7 | 2.7 | 3.0 | * | * | * | * | * |
| 1970 | 1.5 | * | * | * | 3.2 | 4.8 | 3.1 | * | * | * | * | * |
| 1969 | 1.4 | * | * | * | 3.0 | 4.2 | 3.4 | * | * | * | * | * |
| 1968 | 1.6 | * | * | * | 3.2 | 5.6 | 3.2 | * | * | * | * | * |
| BLACK, MALE | | | | | | | | | | | | |
| 1978 | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 1977 | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 1976 | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 1975 | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 1974 | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 1973 | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 1972 | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 1971 | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 1970 | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 1969 | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 1968 | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| BLACK, FEMALE | | | | | | | | | | | | |
| 1978 | 1.0 | * | * | * | 2.0 | 2.8 | 1.5 | * | * | * | * | * |
| 1977 | 1.2 | * | * | * | 2.1 | 3.5 | 2.0 | * | * | * | * | * |
| 1976 | 1.2 | * | * | * | 1.9 | 3.6 | 2.1 | * | * | * | * | * |
| 1975 | 1.2 | * | * | * | 2.5 | 3.5 | 2.0 | * | * | * | * | * |
| 1974 | 1.5 | * | * | * | 3.5 | 4.5 | 1.7 | * | * | * | * | * |
| 1973 | 1.6 | * | * | * | 3.1 | 4.4 | 2.9 | * | * | * | * | * |
| 1972 | 1.7 | * | * | * | 3.6 | 5.2 | 3.3 | * | * | * | * | * |
| 1971 | 2.3 | * | * | * | 5.3 | 5.0 | 5.5 | * | * | * | * | * |
| 1970 | 2.9 | * | * | * | 6.1 | 8.9 | 5.7 | * | * | * | * | * |
| 1969 | 2.8 | * | * | * | 5.8 | 7.7 | 6.2 | * | * | * | * | * |
| 1968 | 3.0 | * | * | * | 6.2 | 10.2 | 5.9 | * | * | * | * | * |

* Figure does not meet standards of reliability or precision.

... Category not applicable

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| ABORTIONS (640-645) | | | | | | | | | | | | |
| BOTH SEXES | | | | | | | | | | | | |
| 1978 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1977 | 0.0 | * | * | * | * | * | * | * | * | * | * | * |
| 1976 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1975 | 0.0 | * | * | * | * | * | * | * | * | * | * | * |
| 1974 | 0.0 | * | * | * | * | * | * | * | * | * | * | * |
| 1973 | 0.0 | * | * | * | * | * | * | * | * | * | * | * |
| 1972 | 0.0 | * | * | * | 0.1 | 0.1 | * | * | * | * | * | * |
| 1971 | 0.0 | * | * | * | 0.1 | 0.1 | 0.1 | * | * | * | * | * |
| 1970 | 0.1 | * | * | * | 0.2 | 0.2 | 0.1 | * | * | * | * | * |
| 1969 | 0.1 | * | * | * | 0.1 | 0.2 | 0.1 | * | * | * | * | * |
| 1968 | 0.1 | * | * | * | 0.1 | 0.3 | 0.1 | * | * | * | * | * |
| MALE | | | | | | | | | | | | |
| 1978 | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 1977 | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 1976 | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 1975 | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 1974 | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 1973 | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 1972 | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 1971 | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 1970 | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 1969 | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 1968 | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |

* Figure does not meet standards of reliability or precision.

... Category not applicable

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| ABORTIONS (640-645) | | | | | | | | | | | | |
| FEMALE | | | | | | | | | | | | |
| 1978 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1977 | 0.0 | * | * | * | * | * | * | * | * | * | * | * |
| 1976 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1975 | 0.0 | * | * | * | * | * | * | * | * | * | * | * |
| 1974 | 0.0 | * | * | * | * | * | * | * | * | * | * | * |
| 1973 | 0.0 | * | * | * | * | * | * | * | * | * | * | * |
| 1972 | 0.1 | * | * | * | 0.2 | 0.2 | * | * | * | * | * | * |
| 1971 | 0.1 | * | * | * | 0.3 | 0.2 | 0.2 | * | * | * | * | * |
| 1970 | 0.1 | * | * | * | 0.3 | 0.4 | 0.2 | * | * | * | * | * |
| 1969 | 0.1 | * | * | * | 0.3 | 0.4 | 0.2 | * | * | * | * | * |
| 1968 | 0.1 | * | * | * | 0.3 | 0.5 | 0.2 | * | * | * | * | * |
| WHITE, BOTH SEXES | | | | | | | | | | | | |
| 1978 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1977 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1976 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1975 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1974 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1973 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1972 | 0.0 | * | * | * | 0.1 | * | * | * | * | * | * | * |
| 1971 | 0.0 | * | * | * | 0.1 | * | * | * | * | * | * | * |
| 1970 | 0.0 | * | * | * | 0.1 | * | * | * | * | * | * | * |
| 1969 | 0.0 | * | * | * | 0.1 | 0.2 | * | * | * | * | * | * |
| 1968 | 0.0 | * | * | * | * | 0.2 | * | * | * | * | * | * |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| ABORTIONS (640-645) | | | | | | | | | | | | |
| WHITE, MALE | | | | | | | | | | | | |
| 1978 | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 1977 | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 1976 | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 1975 | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 1974 | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 1973 | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 1972 | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 1971 | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 1970 | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 1969 | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 1968 | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| WHITE, FEMALE | | | | | | | | | | | | |
| 1978 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1977 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1976 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1975 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1974 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1973 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1972 | 0.0 | * | * | * | 0.1 | * | * | * | * | * | * | * |
| 1971 | 0.0 | * | * | * | 0.1 | * | * | * | * | * | * | * |
| 1970 | 0.1 | * | * | * | 0.2 | * | * | * | * | * | * | * |
| 1969 | 0.1 | * | * | * | 0.1 | 0.3 | * | * | * | * | * | * |
| 1968 | 0.1 | * | * | * | * | 0.3 | * | * | * | * | * | * |

... Category not applicable

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| ABORTIONS (640-645) | | | | | | | | | | | | |
| ALL OTHER, BOTH SEXES | | | | | | | | | | | | |
| 1978 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1977 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1976 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1975 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1974 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1973 | 0.1 | * | * | * | * | * | * | * | * | * | * | * |
| 1972 | 0.1 | * | * | * | * | * | * | * | * | * | * | * |
| 1971 | 0.2 | * | * | * | 0.5 | * | * | * | * | * | * | * |
| 1970 | 0.3 | * | * | * | 0.7 | 0.9 | * | * | * | * | * | * |
| 1969 | 0.3 | * | * | * | 0.6 | 0.7 | * | * | * | * | * | * |
| 1968 | 0.3 | * | * | * | 0.7 | 1.1 | * | * | * | * | * | * |
| ALL OTHER, MALE | | | | | | | | | | | | |
| 1978 | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 1977 | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 1976 | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 1975 | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 1974 | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 1973 | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 1972 | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 1971 | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 1970 | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 1969 | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 1968 | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| ALL OTHER, FEMALE | | | | | | | | | | | | |
| 1978 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1977 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1976 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1975 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1974 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1973 | 0.1 | * | * | * | * | * | * | * | * | * | * | * |
| 1972 | 0.2 | * | * | * | * | * | * | * | * | * | * | * |
| 1971 | 0.4 | * | * | * | 1.0 | * | * | * | * | * | * | * |
| 1970 | 0.5 | * | * | * | 1.3 | 1.7 | * | * | * | * | * | * |
| 1969 | 0.5 | * | * | * | 1.2 | 1.4 | * | * | * | * | * | * |
| 1968 | 0.6 | * | * | * | 1.3 | 2.0 | * | * | * | * | * | * |

* Figure does not meet standards of reliability or precision.

... Category not applicable

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| ABORTIONS (640-645) | | | | | | | | | | | | |
| BLACK, BOTH SEXES | | | | | | | | | | | | |
| 1978 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1977 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1976 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1975 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1974 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1973 | 0.1 | * | * | * | * | * | * | * | * | * | * | * |
| 1972 | 0.1 | * | * | * | * | * | * | * | * | * | * | * |
| 1971 | 0.2 | * | * | * | 0.6 | * | * | * | * | * | * | * |
| 1970 | 0.3 | * | * | * | 0.8 | 1.0 | * | * | * | * | * | * |
| 1969 | 0.3 | * | * | * | 0.7 | 0.8 | * | * | * | * | * | * |
| 1968 | 0.3 | * | * | * | 0.8 | 1.2 | * | * | * | * | * | * |
| BLACK, MALE | | | | | | | | | | | | |
| 1978 | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 1977 | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 1976 | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 1975 | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 1974 | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 1973 | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 1972 | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 1971 | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 1970 | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 1969 | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 1968 | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| BLACK, FEMALE | | | | | | | | | | | | |
| 1978 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1977 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1976 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1975 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1974 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1973 | 0.2 | * | * | * | * | * | * | * | * | * | * | * |
| 1972 | 0.2 | * | * | * | * | * | * | * | * | * | * | * |
| 1971 | 0.5 | * | * | * | 1.1 | * | * | * | * | * | * | * |
| 1970 | 0.6 | * | * | * | 1.5 | 1.9 | * | * | * | * | * | * |
| 1969 | 0.6 | * | * | * | 1.3 | 1.4 | * | * | * | * | * | * |
| 1968 | 0.6 | * | * | * | 1.5 | 2.2 | * | * | * | * | * | * |

* Figure does not meet standards of reliability or precision.

... Category not applicable

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR TOTAL UNDER 1 YEAR 1-4 YEARS 5-14 YEARS 15-24 YEARS 25-34 YEARS 35-44 YEARS 45-54 YEARS 55-64 YEARS 65-74 YEARS 75-84 YEARS 85+ YEARS

OTHER COMPLICATIONS OF PREGNANCY, CHILDBIRTH, AND THE PUERPERIUM (630-639, 650-678)

BOTH SEXES

| | | | | | | | | | | | | |
|------|-----|---|---|---|-----|-----|-----|---|---|---|---|---|
| 1978 | 0.1 | * | * | * | 0.3 | 0.4 | 0.2 | * | * | * | * | * |
| 1977 | 0.2 | * | * | * | 0.3 | 0.5 | 0.3 | * | * | * | * | * |
| 1976 | 0.2 | * | * | * | 0.3 | 0.5 | 0.3 | * | * | * | * | * |
| 1975 | 0.2 | * | * | * | 0.3 | 0.5 | 0.3 | * | * | * | * | * |
| 1974 | 0.2 | * | * | * | 0.4 | 0.6 | 0.3 | * | * | * | * | * |
| 1973 | 0.2 | * | * | * | 0.4 | 0.7 | 0.3 | * | * | * | * | * |
| 1972 | 0.3 | * | * | * | 0.6 | 0.8 | 0.4 | * | * | * | * | * |
| 1971 | 0.3 | * | * | * | 0.6 | 0.9 | 0.6 | * | * | * | * | * |
| 1970 | 0.3 | * | * | * | 0.7 | 1.0 | 0.7 | * | * | * | * | * |
| 1969 | 0.3 | * | * | * | 0.7 | 1.0 | 0.7 | * | * | * | * | * |
| 1968 | 0.4 | * | * | * | 0.7 | 1.2 | 0.8 | * | * | * | * | * |

MALE

| | | | | | | | | | | | | |
|------|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| 1978 | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 1977 | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 1976 | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 1975 | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 1974 | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 1973 | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 1972 | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 1971 | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 1970 | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 1969 | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 1968 | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |

* Figure does not meet standards of reliability or precision.

... Category not applicable

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| OTHER COMPLICATIONS OF PREGNANCY, CHILDBIRTH, AND THE PUERPERIUM (630-639, 650-678) | | | | | | | | | | | | |
| FEMALE | | | | | | | | | | | | |
| 1978 | 0.3 | * | * | * | 0.5 | 0.8 | 0.4 | * | * | * | * | * |
| 1977 | 0.3 | * | * | * | 0.6 | 1.0 | 0.5 | * | * | * | * | * |
| 1976 | 0.3 | * | * | * | 0.6 | 1.0 | 0.6 | * | * | * | * | * |
| 1975 | 0.3 | * | * | * | 0.7 | 1.0 | 0.6 | * | * | * | * | * |
| 1974 | 0.4 | * | * | * | 0.8 | 1.3 | 0.6 | * | * | * | * | * |
| 1973 | 0.4 | * | * | * | 0.8 | 1.4 | 0.6 | * | * | * | * | * |
| 1972 | 0.5 | * | * | * | 1.1 | 1.6 | 0.8 | * | * | * | * | * |
| 1971 | 0.5 | * | * | * | 1.1 | 1.7 | 1.1 | * | * | * | * | * |
| 1970 | 0.6 | * | * | * | 1.4 | 2.0 | 1.4 | * | * | * | * | * |
| 1969 | 0.6 | * | * | * | 1.3 | 2.1 | 1.4 | * | * | * | * | * |
| 1968 | 0.7 | * | * | * | 1.4 | 2.3 | 1.6 | * | * | * | * | * |
| WHITE, BOTH SEXES | | | | | | | | | | | | |
| 1978 | 0.1 | * | * | * | 0.2 | 0.3 | 0.1 | * | * | * | * | * |
| 1977 | 0.1 | * | * | * | 0.2 | 0.3 | 0.2 | * | * | * | * | * |
| 1976 | 0.1 | * | * | * | 0.2 | 0.4 | 0.2 | * | * | * | * | * |
| 1975 | 0.1 | * | * | * | 0.2 | 0.4 | 0.2 | * | * | * | * | * |
| 1974 | 0.1 | * | * | * | 0.2 | 0.4 | 0.2 | * | * | * | * | * |
| 1973 | 0.1 | * | * | * | 0.3 | 0.5 | 0.2 | * | * | * | * | * |
| 1972 | 0.2 | * | * | * | 0.4 | 0.6 | 0.3 | * | * | * | * | * |
| 1971 | 0.2 | * | * | * | 0.3 | 0.7 | 0.4 | * | * | * | * | * |
| 1970 | 0.2 | * | * | * | 0.5 | 0.7 | 0.5 | * | * | * | * | * |
| 1969 | 0.2 | * | * | * | 0.5 | 0.7 | 0.5 | * | * | * | * | * |
| 1968 | 0.2 | * | * | * | 0.5 | 0.8 | 0.6 | * | * | * | * | * |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| OTHER COMPLICATIONS OF PREGNANCY, CHILDBIRTH, AND THE PUERPERIUM (630-639, 650-678) | | | | | | | | | | | | |
| WHITE, MALE | | | | | | | | | | | | |
| 1978 | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 1977 | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 1976 | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 1975 | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 1974 | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 1973 | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 1972 | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 1971 | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 1970 | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 1969 | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 1968 | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| WHITE, FEMALE | | | | | | | | | | | | |
| 1978 | 0.2 | * | * | * | 0.3 | 0.5 | 0.3 | * | * | * | * | * |
| 1977 | 0.2 | * | * | * | 0.4 | 0.7 | 0.3 | * | * | * | * | * |
| 1976 | 0.2 | * | * | * | 0.5 | 0.7 | 0.4 | * | * | * | * | * |
| 1975 | 0.2 | * | * | * | 0.5 | 0.7 | 0.4 | * | * | * | * | * |
| 1974 | 0.3 | * | * | * | 0.5 | 0.9 | 0.4 | * | * | * | * | * |
| 1973 | 0.3 | * | * | * | 0.5 | 1.0 | 0.4 | * | * | * | * | * |
| 1972 | 0.4 | * | * | * | 0.8 | 1.2 | 0.5 | * | * | * | * | * |
| 1971 | 0.4 | * | * | * | 0.6 | 1.4 | 0.7 | * | * | * | * | * |
| 1970 | 0.4 | * | * | * | 0.9 | 1.3 | 1.0 | * | * | * | * | * |
| 1969 | 0.4 | * | * | * | 0.9 | 1.5 | 1.0 | * | * | * | * | * |
| 1968 | 0.5 | * | * | * | 0.9 | 1.5 | 1.2 | * | * | * | * | * |

... Category not applicable

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| OTHER COMPLICATIONS OF PREGNANCY, CHILDBIRTH, AND THE PUERPERIUM (630-639, 650-678) | | | | | | | | | | | | |
| ALL OTHER, BOTH SEXES | | | | | | | | | | | | |
| 1978 | 0.5 | * | * | * | 0.9 | 1.3 | 0.6 | * | * | * | * | * |
| 1977 | 0.5 | * | * | * | 0.9 | 1.6 | 0.8 | * | * | * | * | * |
| 1976 | 0.5 | * | * | * | 0.8 | 1.7 | 0.9 | * | * | * | * | * |
| 1975 | 0.6 | * | * | * | 1.0 | 1.7 | 0.8 | * | * | * | * | * |
| 1974 | 0.7 | * | * | * | 1.5 | 2.1 | 0.8 | * | * | * | * | * |
| 1973 | 0.7 | * | * | * | 1.3 | 1.9 | 1.2 | * | * | * | * | * |
| 1972 | 0.8 | * | * | * | 1.4 | 2.3 | 1.6 | * | * | * | * | * |
| 1971 | 0.9 | * | * | * | 2.0 | 2.2 | 2.2 | * | * | * | * | * |
| 1970 | 1.1 | * | * | * | 2.2 | 3.5 | 2.5 | * | * | * | * | * |
| 1969 | 1.1 | * | * | * | 2.1 | 3.2 | 2.7 | * | * | * | * | * |
| 1968 | 1.2 | * | * | * | 2.3 | 4.3 | 2.6 | * | * | * | * | * |
| ALL OTHER, MALE | | | | | | | | | | | | |
| 1978 | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 1977 | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 1976 | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 1975 | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 1974 | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 1973 | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 1972 | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 1971 | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 1970 | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 1969 | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 1968 | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| ALL OTHER, FEMALE | | | | | | | | | | | | |
| 1978 | 0.9 | * | * | * | 1.7 | 2.4 | 1.2 | * | * | * | * | * |
| 1977 | 1.0 | * | * | * | 1.7 | 2.9 | 1.5 | * | * | * | * | * |
| 1976 | 1.0 | * | * | * | 1.6 | 3.1 | 1.6 | * | * | * | * | * |
| 1975 | 1.1 | * | * | * | 1.9 | 3.3 | 1.5 | * | * | * | * | * |
| 1974 | 1.3 | * | * | * | 2.9 | 3.8 | 1.5 | * | * | * | * | * |
| 1973 | 1.3 | * | * | * | 2.5 | 3.6 | 2.3 | * | * | * | * | * |
| 1972 | 1.4 | * | * | * | 2.8 | 4.3 | 2.9 | * | * | * | * | * |
| 1971 | 1.7 | * | * | * | 3.9 | 4.0 | 3.9 | * | * | * | * | * |
| 1970 | 2.2 | * | * | * | 4.3 | 6.5 | 4.5 | * | * | * | * | * |
| 1969 | 2.1 | * | * | * | 4.1 | 5.9 | 4.9 | * | * | * | * | * |
| 1968 | 2.4 | * | * | * | 4.5 | 7.9 | 4.8 | * | * | * | * | * |

* Figure does not meet standards of reliability or precision.

... Category not applicable

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR TOTAL UNDER 1 YEAR 1-4 YEARS 5-14 YEARS 15-24 YEARS 25-34 YEARS 35-44 YEARS 45-54 YEARS 55-64 YEARS 65-74 YEARS 75-84 YEARS 85+ YEARS

OTHER COMPLICATIONS OF PREGNANCY, CHILDBIRTH, AND THE PUERPERIUM (630-639, 650-678)

BLACK, BOTH SEXES

| | | | | | | | | | | | | |
|------|-----|---|---|---|-----|-----|-----|---|---|---|---|---|
| 1978 | 0.5 | * | * | * | 0.9 | 1.4 | 0.8 | * | * | * | * | * |
| 1977 | 0.6 | * | * | * | 1.0 | 1.8 | 1.0 | * | * | * | * | * |
| 1976 | 0.6 | * | * | * | 0.9 | 1.9 | 1.0 | * | * | * | * | * |
| 1975 | 0.6 | * | * | * | 1.1 | 1.8 | 1.0 | * | * | * | * | * |
| 1974 | 0.7 | * | * | * | 1.7 | 2.2 | 0.9 | * | * | * | * | * |
| 1973 | 0.7 | * | * | * | 1.4 | 2.2 | 1.4 | * | * | * | * | * |
| 1972 | 0.8 | * | * | * | 1.5 | 2.5 | 1.7 | * | * | * | * | * |
| 1971 | 0.9 | * | * | * | 2.2 | 2.2 | 2.4 | * | * | * | * | * |
| 1970 | 1.2 | * | * | * | 2.4 | 3.8 | 2.7 | * | * | * | * | * |
| 1969 | 1.2 | * | * | * | 2.3 | 3.4 | 2.8 | * | * | * | * | * |
| 1968 | 1.3 | * | * | * | 2.5 | 4.4 | 2.7 | * | * | * | * | * |

BLACK, MALE

| | | | | | | | | | | | | |
|------|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| 1978 | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 1977 | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 1976 | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 1975 | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 1974 | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 1973 | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 1972 | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 1971 | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 1970 | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 1969 | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 1968 | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |

BLACK, FEMALE

| | | | | | | | | | | | | |
|------|-----|---|---|---|-----|-----|-----|---|---|---|---|---|
| 1978 | 0.9 | * | * | * | 1.8 | 2.6 | 1.4 | * | * | * | * | * |
| 1977 | 1.1 | * | * | * | 2.0 | 3.3 | 1.8 | * | * | * | * | * |
| 1976 | 1.1 | * | * | * | 1.7 | 3.5 | 1.8 | * | * | * | * | * |
| 1975 | 1.1 | * | * | * | 2.2 | 3.3 | 1.8 | * | * | * | * | * |
| 1974 | 1.4 | * | * | * | 3.2 | 4.1 | 1.7 | * | * | * | * | * |
| 1973 | 1.4 | * | * | * | 2.7 | 4.1 | 2.6 | * | * | * | * | * |
| 1972 | 1.5 | * | * | * | 3.0 | 4.7 | 3.0 | * | * | * | * | * |
| 1971 | 1.8 | * | * | * | 4.2 | 4.1 | 4.4 | * | * | * | * | * |
| 1970 | 2.3 | * | * | * | 4.7 | 7.1 | 5.0 | * | * | * | * | * |
| 1969 | 2.2 | * | * | * | 4.4 | 6.3 | 5.1 | * | * | * | * | * |
| 1968 | 2.4 | * | * | * | 4.7 | 8.0 | 5.0 | * | * | * | * | * |

* Figure does not meet standards of reliability or precision.

... Category not applicable

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER | 1-4 | 5-14 | 15-24 | 25-34 | 35-44 | 45-54 | 55-64 | 65-74 | 75-84 | 85+ |
|---|--------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|
| | 1 YEAR | YEARS | YEARS | YEARS | YEARS | YEARS | YEARS | YEARS | YEARS | YEARS | YEARS | YEARS |
| Congenital anomalies 740-759 | | | | | | | | | | | | |
| BOTH SEXES | | | | | | | | | | | | |
| 1978 | 5.8 | 252.7 | 8.3 | 1.8 | 1.5 | 1.2 | 1.2 | 1.6 | 2.1 | 2.4 | 3.1 | 3.7 |
| 1977 | 5.9 | 256.8 | 8.7 | 1.8 | 1.6 | 1.3 | 1.3 | 1.6 | 2.0 | 2.4 | 2.5 | 3.8 |
| 1976 | 6.0 | 267.9 | 8.9 | 2.0 | 1.4 | 1.3 | 1.3 | 1.8 | 2.3 | 2.4 | 2.7 | 3.6 |
| 1975 | 6.1 | 272.3 | 8.8 | 1.9 | 1.6 | 1.4 | 1.4 | 1.7 | 2.0 | 2.2 | 2.8 | 2.9 |
| 1974 | 6.3 | 280.8 | 8.9 | 2.1 | 1.7 | 1.4 | 1.4 | 1.8 | 2.3 | 2.7 | 2.8 | 3.2 |
| 1973 | 6.7 | 286.2 | 9.5 | 2.1 | 1.7 | 1.4 | 1.6 | 1.9 | 2.3 | 2.7 | 2.8 | 3.7 |
| 1972 | 7.0 | 286.1 | 10.3 | 2.3 | 1.6 | 1.5 | 1.3 | 2.0 | 2.6 | 2.6 | 2.8 | 4.4 |
| 1971 | 7.7 | 296.1 | 9.5 | 2.3 | 1.9 | 1.5 | 1.7 | 2.2 | 2.4 | 2.6 | 3.1 | 3.4 |
| 1970 | 8.3 | 323.0 | 9.7 | 2.2 | 2.1 | 1.7 | 1.8 | 2.5 | 2.8 | 2.9 | 3.3 | 4.8 |
| 1969 | 8.4 | 331.4 | 10.2 | 2.5 | 2.0 | 1.8 | 1.8 | 2.5 | 2.7 | 2.9 | 3.8 | 5.4 |
| 1968 | 8.4 | 328.3 | 9.8 | 2.5 | 2.1 | 1.8 | 1.8 | 2.6 | 3.0 | 2.8 | 3.5 | 5.3 |
| MALE | | | | | | | | | | | | |
| 1978 | 6.4 | 264.4 | 8.8 | 1.8 | 1.7 | 1.3 | 1.3 | 1.8 | 2.3 | 2.5 | 3.8 | 5.4 |
| 1977 | 6.5 | 271.9 | 8.7 | 1.9 | 1.8 | 1.5 | 1.5 | 1.7 | 2.2 | 2.6 | 2.9 | 5.6 |
| 1976 | 6.5 | 281.0 | 8.6 | 1.9 | 1.6 | 1.5 | 1.4 | 1.9 | 2.5 | 2.6 | 3.0 | 4.6 |
| 1975 | 6.8 | 291.4 | 8.4 | 2.0 | 1.9 | 1.6 | 1.5 | 1.9 | 2.1 | 2.5 | 3.0 | 4.4 |
| 1974 | 7.1 | 302.6 | 9.0 | 2.1 | 1.8 | 1.6 | 1.6 | 2.0 | 2.4 | 3.0 | 4.0 | 4.2 |
| 1973 | 7.3 | 304.8 | 9.7 | 2.2 | 1.9 | 1.6 | 1.9 | 2.0 | 2.6 | 3.1 | 3.4 | 4.3 |
| 1972 | 7.7 | 304.8 | 10.1 | 2.3 | 1.8 | 2.0 | 1.2 | 2.5 | 2.7 | 2.7 | 3.0 | 6.9 |
| 1971 | 8.5 | 310.4 | 9.7 | 2.4 | 2.2 | 1.7 | 2.0 | 2.4 | 2.5 | 3.0 | 3.2 | 4.3 |
| 1970 | 9.0 | 338.5 | 10.0 | 2.3 | 2.3 | 1.8 | 1.9 | 2.8 | 2.9 | 3.0 | 3.9 | 6.3 |
| 1969 | 9.3 | 354.5 | 9.7 | 2.7 | 2.3 | 1.9 | 2.0 | 2.8 | 2.7 | 3.2 | 4.8 | 8.3 |
| 1968 | 9.3 | 349.7 | 10.1 | 2.6 | 2.4 | 2.1 | 2.0 | 2.8 | 3.3 | 2.9 | 4.4 | 8.9 |

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| Congenital anomalies 740-759 | | | | | | | | | | | | |
| FEMALE | | | | | | | | | | | | |
| 1978 | 5.3 | 240.4 | 7.7 | 1.7 | 1.3 | 1.0 | 1.1 | 1.5 | 2.0 | 2.4 | 2.7 | 2.9 |
| 1977 | 5.3 | 240.9 | 8.7 | 1.8 | 1.3 | 1.0 | 1.2 | 1.5 | 1.8 | 2.3 | 2.2 | 3.0 |
| 1976 | 5.5 | 254.0 | 9.2 | 2.0 | 1.2 | 1.1 | 1.3 | 1.7 | 2.0 | 2.3 | 2.6 | 3.1 |
| 1975 | 5.5 | 252.2 | 9.2 | 1.9 | 1.3 | 1.2 | 1.4 | 1.5 | 2.0 | 2.0 | 2.6 | 2.1 |
| 1974 | 5.7 | 258.0 | 8.7 | 2.1 | 1.5 | 1.3 | 1.2 | 1.6 | 2.2 | 2.5 | 2.1 | 2.6 |
| 1973 | 6.0 | 266.9 | 9.3 | 2.1 | 1.4 | 1.3 | 1.3 | 1.9 | 2.1 | 2.3 | 2.5 | 3.4 |
| 1972 | 6.4 | 266.5 | 10.5 | 2.4 | 1.4 | 1.1 | 1.4 | 1.6 | 2.5 | 2.5 | 2.6 | 3.1 |
| 1971 | 7.0 | 281.1 | 9.3 | 2.1 | 1.5 | 1.2 | 1.4 | 2.1 | 2.4 | 2.4 | 3.1 | 3.0 |
| 1970 | 7.6 | 306.9 | 9.5 | 2.1 | 1.9 | 1.5 | 1.7 | 2.3 | 2.7 | 2.8 | 2.9 | 3.9 |
| 1969 | 7.6 | 307.4 | 10.8 | 2.3 | 1.7 | 1.6 | 1.6 | 2.2 | 2.7 | 2.6 | 3.1 | 3.9 |
| 1968 | 7.5 | 306.1 | 9.5 | 2.3 | 1.7 | 1.5 | 1.6 | 2.4 | 2.8 | 2.7 | 2.9 | 3.4 |
| WHITE, BOTH SEXES | | | | | | | | | | | | |
| 1978 | 5.6 | 247.9 | 8.2 | 1.7 | 1.5 | 1.1 | 1.2 | 1.6 | 2.1 | 2.4 | 3.1 | 3.8 |
| 1977 | 5.6 | 252.8 | 8.4 | 1.8 | 1.6 | 1.2 | 1.3 | 1.5 | 2.1 | 2.4 | 2.5 | 3.6 |
| 1976 | 5.7 | 262.7 | 8.7 | 2.0 | 1.4 | 1.3 | 1.3 | 1.8 | 2.3 | 2.4 | 2.8 | 3.6 |
| 1975 | 5.9 | 269.7 | 8.6 | 2.0 | 1.5 | 1.4 | 1.4 | 1.7 | 2.1 | 2.2 | 2.9 | 3.0 |
| 1974 | 6.1 | 278.8 | 8.6 | 2.1 | 1.6 | 1.4 | 1.4 | 1.8 | 2.3 | 2.7 | 2.9 | 3.2 |
| 1973 | 6.4 | 286.5 | 9.4 | 2.1 | 1.7 | 1.4 | 1.6 | 1.9 | 2.3 | 2.6 | 2.9 | 3.8 |
| 1972 | 6.8 | 285.5 | 10.1 | 2.3 | 1.6 | 1.5 | 1.2 | 2.1 | 2.5 | 2.6 | 2.8 | 4.5 |
| 1971 | 7.5 | 298.3 | 9.3 | 2.3 | 1.8 | 1.5 | 1.6 | 2.2 | 2.4 | 2.6 | 3.3 | 3.5 |
| 1970 | 8.1 | 325.4 | 9.3 | 2.2 | 2.1 | 1.6 | 1.8 | 2.5 | 2.8 | 2.8 | 3.4 | 4.9 |
| 1969 | 8.1 | 328.5 | 9.8 | 2.5 | 2.0 | 1.7 | 1.7 | 2.4 | 2.6 | 2.9 | 3.8 | 5.6 |
| 1968 | 8.2 | 331.2 | 9.6 | 2.5 | 2.1 | 1.8 | 1.7 | 2.5 | 3.0 | 2.8 | 3.5 | 5.3 |

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| Congenital anomalies 740-759 | | | | | | | | | | | | |
| WHITE, MALE | | | | | | | | | | | | |
| 1978 | 6.1 | 259.8 | 9.1 | 1.8 | 1.7 | 1.3 | 1.3 | 1.7 | 2.2 | 2.4 | 3.6 | 5.4 |
| 1977 | 6.3 | 269.3 | 8.4 | 1.9 | 1.9 | 1.4 | 1.4 | 1.6 | 2.3 | 2.5 | 3.0 | 5.3 |
| 1976 | 6.2 | 276.7 | 8.4 | 2.0 | 1.5 | 1.5 | 1.3 | 1.8 | 2.6 | 2.7 | 3.0 | 5.1 |
| 1975 | 6.5 | 288.6 | 8.3 | 2.0 | 1.7 | 1.6 | 1.4 | 1.8 | 2.1 | 2.5 | 3.1 | 4.4 |
| 1974 | 6.7 | 298.6 | 8.8 | 2.2 | 1.8 | 1.5 | 1.6 | 1.9 | 2.4 | 2.9 | 4.2 | 4.5 |
| 1973 | 7.1 | 306.8 | 9.6 | 2.1 | 1.9 | 1.6 | 1.9 | 1.9 | 2.5 | 3.0 | 3.6 | 4.5 |
| 1972 | 7.5 | 307.1 | 9.7 | 2.3 | 1.7 | 1.9 | 1.3 | 2.4 | 2.6 | 2.8 | 3.1 | 6.7 |
| 1971 | 8.2 | 310.7 | 9.6 | 2.5 | 2.2 | 1.8 | 1.8 | 2.4 | 2.4 | 2.8 | 3.3 | 4.5 |
| 1970 | 8.7 | 341.2 | 9.4 | 2.3 | 2.3 | 1.7 | 1.8 | 2.7 | 2.9 | 2.8 | 4.0 | 6.4 |
| 1969 | 9.0 | 351.3 | 9.3 | 2.7 | 2.3 | 1.7 | 1.9 | 2.7 | 2.7 | 3.4 | 4.8 | 8.9 |
| 1968 | 9.1 | 352.3 | 9.9 | 2.6 | 2.4 | 2.0 | 1.9 | 2.8 | 3.2 | 2.8 | 4.3 | 8.8 |
| WHITE, FEMALE | | | | | | | | | | | | |
| 1978 | 5.1 | 235.2 | 7.3 | 1.7 | 1.3 | 1.0 | 1.1 | 1.6 | 2.0 | 2.4 | 2.8 | 3.1 |
| 1977 | 5.1 | 235.4 | 8.3 | 1.7 | 1.3 | 1.0 | 1.2 | 1.5 | 1.9 | 2.3 | 2.2 | 2.9 |
| 1976 | 5.2 | 248.0 | 8.9 | 2.0 | 1.2 | 1.1 | 1.3 | 1.7 | 2.0 | 2.2 | 2.7 | 2.9 |
| 1975 | 5.4 | 249.6 | 9.0 | 1.9 | 1.3 | 1.2 | 1.3 | 1.5 | 2.0 | 2.0 | 2.7 | 2.3 |
| 1974 | 5.5 | 258.0 | 8.5 | 2.1 | 1.5 | 1.3 | 1.2 | 1.6 | 2.2 | 2.5 | 2.1 | 2.6 |
| 1973 | 5.8 | 265.2 | 9.2 | 2.1 | 1.4 | 1.3 | 1.3 | 1.9 | 2.2 | 2.3 | 2.4 | 3.4 |
| 1972 | 6.1 | 262.9 | 10.6 | 2.4 | 1.4 | 1.1 | 1.2 | 1.7 | 2.5 | 2.4 | 2.6 | 3.4 |
| 1971 | 6.9 | 285.2 | 9.1 | 2.1 | 1.5 | 1.3 | 1.3 | 2.0 | 2.4 | 2.4 | 3.2 | 3.0 |
| 1970 | 7.4 | 308.9 | 9.2 | 2.1 | 1.9 | 1.5 | 1.7 | 2.3 | 2.7 | 2.8 | 3.0 | 4.0 |
| 1969 | 7.3 | 304.7 | 10.3 | 2.2 | 1.6 | 1.7 | 1.5 | 2.2 | 2.6 | 2.5 | 3.1 | 3.8 |
| 1968 | 7.4 | 309.0 | 9.3 | 2.3 | 1.8 | 1.6 | 1.6 | 2.3 | 2.8 | 2.7 | 2.9 | 3.4 |

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER | 1-4 | 5-14 | 15-24 | 25-34 | 35-44 | 45-54 | 55-64 | 65-74 | 75-84 | 85+ |
|---|--------|--------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|
| | 1 YEAR | 1 YEAR | YEARS | YEARS | YEARS | YEARS | YEARS | YEARS | YEARS | YEARS | YEARS | YEARS |
| Congenital anomalies 740-759 | | | | | | | | | | | | |
| ALL OTHER, BOTH SEXES | | | | | | | | | | | | |
| 1978 | 7.5 | 275.7 | 8.5 | 2.1 | 1.6 | 1.5 | 1.3 | 1.6 | 2.1 | 2.6 | 3.0 | * |
| 1977 | 7.6 | 275.9 | 10.1 | 2.0 | 1.5 | 1.5 | 1.5 | 1.9 | 1.8 | 2.6 | * | * |
| 1976 | 7.7 | 292.8 | 10.1 | 2.0 | 1.5 | 1.5 | 1.4 | 1.9 | 2.2 | 2.3 | * | * |
| 1975 | 7.7 | 285.0 | 9.6 | 1.8 | 1.9 | 1.5 | 1.9 | 1.9 | 1.6 | 2.2 | * | * |
| 1974 | 8.0 | 290.1 | 10.0 | 2.0 | 1.8 | 1.8 | 1.8 | 2.0 | 2.5 | 2.7 | * | * |
| 1973 | 8.2 | 284.2 | 10.0 | 2.3 | 1.6 | 1.6 | 1.6 | 2.4 | 2.3 | 3.4 | * | * |
| 1972 | 8.7 | 289.5 | 11.2 | 2.3 | 1.8 | 1.9 | 1.7 | 1.5 | 3.1 | 2.5 | * | * |
| 1971 | 9.1 | 284.3 | 10.6 | 2.4 | 2.1 | 1.3 | 2.3 | 2.9 | 2.6 | 3.3 | * | * |
| 1970 | 9.8 | 310.3 | 11.9 | 2.1 | 2.1 | 2.0 | 2.1 | 3.2 | 2.9 | 3.2 | * | * |
| 1969 | 10.6 | 347.7 | 12.5 | 2.6 | 2.1 | 2.3 | 2.3 | 2.8 | 2.9 | 2.6 | * | * |
| 1968 | 10.1 | 313.1 | 11.0 | 2.7 | 2.1 | 1.8 | 2.1 | 3.3 | 3.5 | 3.3 | * | * |
| ALL OTHER, MALE | | | | | | | | | | | | |
| 1978 | 8.3 | 285.8 | 7.3 | 2.3 | 1.7 | 1.8 | 1.4 | 2.0 | 3.0 | 3.9 | * | * |
| 1977 | 8.4 | 284.5 | 9.8 | 1.9 | 1.8 | 1.9 | 1.8 | 2.5 | * | 3.6 | * | * |
| 1976 | 8.3 | 302.6 | 9.7 | 1.6 | 1.8 | 1.7 | 1.9 | 2.7 | 2.5 | * | * | * |
| 1975 | 8.7 | 304.1 | 9.2 | 1.8 | 2.7 | 1.5 | 2.3 | 2.7 | * | * | * | * |
| 1974 | 9.3 | 322.8 | 10.3 | 2.0 | 2.2 | 2.5 | 2.2 | 2.6 | 2.5 | * | * | * |
| 1973 | 8.9 | 293.2 | 9.9 | 2.3 | 1.7 | 2.2 | 1.7 | 2.7 | 2.8 | 4.8 | * | * |
| 1972 | 9.5 | 293.8 | 12.3 | 2.3 | 2.2 | 3.1 | * | 2.6 | 3.5 | * | * | * |
| 1971 | 10.4 | 308.7 | 10.6 | 2.3 | 2.6 | 1.6 | 3.1 | 2.9 | 3.0 | 5.1 | * | * |
| 1970 | 10.9 | 323.9 | 12.7 | 2.2 | 2.7 | 2.6 | 2.3 | 3.6 | 2.8 | 4.0 | * | * |
| 1969 | 11.7 | 373.3 | 11.7 | 2.9 | 2.2 | 3.4 | 2.4 | 3.7 | * | * | * | * |
| 1968 | 11.5 | 334.0 | 11.6 | 2.9 | 2.7 | 2.9 | 2.4 | 3.3 | 4.4 | * | * | * |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| Congenital anomalies 740-759 | | | | | | | | | | | | |
| ALL OTHER, FEMALE | | | | | | | | | | | | |
| 1978 | 6.8 | 265.2 | 9.8 | 2.0 | 1.4 | 1.3 | 1.2 | * | * | * | * | * |
| 1977 | 6.9 | 267.1 | 10.5 | 2.1 | 1.2 | 1.2 | 1.3 | 1.4 | * | * | * | * |
| 1976 | 7.2 | 283.9 | 10.5 | 2.4 | 1.2 | 1.4 | * | * | 1.8 | 2.7 | * | * |
| 1975 | 6.7 | 265.3 | 9.9 | 1.9 | 1.1 | 1.4 | 1.5 | * | * | * | * | * |
| 1974 | 6.8 | 257.7 | 9.7 | 1.9 | 1.4 | 1.3 | 1.4 | 1.5 | 2.4 | * | * | * |
| 1973 | 7.5 | 275.0 | 10.1 | 2.3 | 1.6 | 1.0 | 1.4 | 2.1 | * | * | * | * |
| 1972 | 8.0 | 285.1 | 10.0 | 2.4 | 1.4 | * | 2.2 | * | 2.7 | 3.4 | * | * |
| 1971 | 8.0 | 259.4 | 10.7 | 2.4 | 1.8 | * | 1.5 | 2.9 | 2.2 | * | * | * |
| 1970 | 8.8 | 296.5 | 11.1 | 1.9 | 1.5 | 1.5 | 2.0 | 2.8 | 3.1 | * | * | * |
| 1969 | 9.6 | 321.9 | 13.2 | 2.4 | 2.1 | 1.5 | 2.2 | 2.0 | 3.6 | * | * | * |
| 1968 | 8.9 | 290.8 | 10.4 | 2.5 | 1.6 | * | 1.8 | 3.3 | 2.8 | * | * | * |
| BLACK, BOTH SEXES | | | | | | | | | | | | |
| 1978 | 7.7 | 280.2 | 9.0 | 2.3 | 1.7 | 1.6 | 1.3 | 1.7 | 2.2 | 2.6 | * | * |
| 1977 | 8.0 | 286.2 | 10.9 | 2.1 | 1.6 | 1.6 | 1.6 | 2.1 | 2.0 | 2.6 | * | * |
| 1976 | 8.0 | 302.2 | 10.6 | 2.2 | 1.5 | 1.7 | 1.4 | 2.1 | 2.4 | 2.5 | * | * |
| 1975 | 8.0 | 297.1 | 9.9 | 1.9 | 2.0 | 1.5 | 2.0 | 2.0 | 1.8 | 2.1 | * | * |
| 1974 | 8.3 | 302.5 | 10.6 | 2.1 | 1.9 | 2.0 | 2.0 | 2.0 | 2.6 | 2.7 | * | * |
| 1973 | 8.3 | 290.5 | 10.6 | 2.3 | 1.7 | 1.7 | 1.6 | 2.5 | 2.3 | 3.6 | * | * |
| 1972 | 8.9 | 297.3 | 11.5 | 2.3 | 1.9 | 2.1 | 1.8 | 1.7 | 2.9 | 2.6 | * | * |
| 1971 | 9.3 | 285.5 | 11.0 | 2.5 | 2.3 | 1.4 | 2.5 | 3.0 | 2.3 | 3.2 | * | * |
| 1970 | 10.2 | 321.8 | 12.2 | 2.1 | 2.2 | 2.2 | 2.1 | 3.3 | 3.2 | 3.0 | * | * |
| 1969 | 10.9 | 359.3 | 13.0 | 2.6 | 2.3 | 2.5 | 2.5 | 3.1 | 3.0 | 2.8 | * | * |
| 1968 | 10.3 | 319.7 | 11.2 | 2.8 | 2.1 | 2.0 | 2.3 | 3.5 | 3.4 | 3.5 | * | * |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| Congenital anomalies 740-759 | | | | | | | | | | | | |
| BLACK, MALE | | | | | | | | | | | | |
| 1978 | 8.6 | 296.8 | 7.3 | 2.3 | 1.9 | 1.9 | * | 2.2 | 3.1 | 4.0 | * | * |
| 1977 | 8.9 | 297.6 | 10.6 | 2.0 | 1.9 | 2.0 | 2.0 | 2.8 | * | 4.1 | * | * |
| 1976 | 8.6 | 313.5 | 9.9 | 1.7 | 1.9 | 1.9 | 1.9 | 3.0 | 2.9 | * | * | * |
| 1975 | 9.1 | 317.4 | 9.6 | 1.8 | 3.0 | 1.6 | 2.4 | 2.9 | * | * | * | * |
| 1974 | 9.7 | 336.7 | 11.0 | 2.1 | 2.3 | 2.7 | 2.5 | 2.6 | * | * | * | * |
| 1973 | 9.3 | 302.1 | 10.2 | 2.5 | 1.8 | 2.6 | 1.8 | 2.7 | 3.0 | 5.2 | * | * |
| 1972 | 9.7 | 296.3 | 12.9 | 2.1 | 2.3 | 3.3 | * | 3.0 | 3.7 | * | * | * |
| 1971 | 10.5 | 305.9 | 11.0 | 2.4 | 2.8 | 1.8 | 3.5 | 3.2 | * | 5.1 | * | * |
| 1970 | 11.3 | 335.0 | 12.7 | 2.3 | 2.9 | 2.9 | 2.4 | 3.9 | 3.1 | * | * | * |
| 1969 | 12.0 | 386.5 | 11.8 | 2.7 | 2.4 | 3.6 | 2.4 | 4.0 | * | * | * | * |
| 1968 | 11.7 | 340.3 | 11.7 | 3.0 | 2.7 | 3.3 | 2.7 | 3.2 | 4.5 | * | * | * |
| BLACK, FEMALE | | | | | | | | | | | | |
| 1978 | 6.9 | 263.4 | 10.7 | 2.2 | 1.6 | 1.4 | * | * | * | * | * | * |
| 1977 | 7.2 | 274.6 | 11.3 | 2.2 | 1.3 | 1.3 | * | * | * | * | * | * |
| 1976 | 7.4 | 290.7 | 11.4 | 2.7 | 1.2 | 1.5 | * | * | * | * | * | * |
| 1975 | 7.0 | 276.5 | 10.3 | 2.0 | 1.1 | 1.4 | 1.6 | * | * | * | * | * |
| 1974 | 7.1 | 267.4 | 10.3 | 2.1 | 1.5 | 1.5 | 1.5 | * | 2.7 | * | * | * |
| 1973 | 7.5 | 277.3 | 11.0 | 2.2 | 1.5 | * | 1.5 | 2.2 | * | * | * | * |
| 1972 | 8.2 | 297.0 | 10.2 | 2.5 | 1.5 | * | 2.4 | * | 2.2 | 3.7 | * | * |
| 1971 | 8.1 | 264.8 | 11.0 | 2.6 | 1.8 | * | 1.6 | 2.8 | * | * | * | * |
| 1970 | 9.1 | 308.6 | 11.7 | 2.0 | 1.5 | 1.6 | 1.9 | 2.8 | 3.2 | * | * | * |
| 1969 | 9.9 | 332.0 | 14.1 | 2.5 | 2.2 | 1.5 | 2.5 | 2.2 | 3.8 | * | * | * |
| 1968 | 9.1 | 298.7 | 10.7 | 2.6 | 1.5 | * | 1.9 | 3.6 | 2.5 | * | * | * |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|

Certain causes of mortality in early infancy 760-769.2, 769.4-772, 774-778

BOTH SEXES

| | | | | | | | | | | | | |
|------|------|---------|-----|-----|---|---|---|---|---|---|---|---|
| 1978 | 9.9 | 661.1 | 0.2 | * | * | * | * | * | * | * | * | * |
| 1977 | 10.6 | 712.3 | 0.3 | * | * | * | * | * | * | * | * | * |
| 1976 | 11.4 | 795.1 | * | * | * | * | * | * | * | * | * | * |
| 1975 | 12.4 | 843.4 | * | * | * | * | * | * | * | * | * | * |
| 1974 | 13.5 | 936.8 | * | * | * | * | * | * | * | * | * | * |
| 1973 | 14.4 | 971.6 | * | * | * | * | * | * | * | * | * | * |
| 1972 | 16.1 | 1,020.4 | * | * | * | * | * | * | * | * | * | * |
| 1971 | 18.6 | 1,066.3 | 0.2 | * | * | * | * | * | * | * | * | * |
| 1970 | 21.3 | 1,238.8 | * | * | * | * | * | * | * | * | * | * |
| 1969 | 21.4 | 1,261.6 | 0.2 | * | * | * | * | * | * | * | * | * |
| 1968 | 22.0 | 1,298.5 | 0.3 | 0.1 | * | * | * | * | * | * | * | * |

MALE

| | | | | | | | | | | | | |
|------|------|---------|-----|---|---|---|---|---|---|---|---|---|
| 1978 | 11.8 | 744.1 | * | * | * | * | * | * | * | * | * | * |
| 1977 | 12.6 | 799.0 | * | * | * | * | * | * | * | * | * | * |
| 1976 | 13.4 | 890.0 | * | * | * | * | * | * | * | * | * | * |
| 1975 | 14.6 | 949.3 | * | * | * | * | * | * | * | * | * | * |
| 1974 | 16.1 | 1,062.5 | * | * | * | * | * | * | * | * | * | * |
| 1973 | 17.3 | 1,109.9 | * | * | * | * | * | * | * | * | * | * |
| 1972 | 19.4 | 1,167.3 | * | * | * | * | * | * | * | * | * | * |
| 1971 | 22.4 | 1,222.1 | * | * | * | * | * | * | * | * | * | * |
| 1970 | 25.5 | 1,418.4 | * | * | * | * | * | * | * | * | * | * |
| 1969 | 25.9 | 1,453.0 | * | * | * | * | * | * | * | * | * | * |
| 1968 | 26.6 | 1,502.2 | 0.3 | * | * | * | * | * | * | * | * | * |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|

Certain causes of mortality in early infancy 760-769.2, 769.4-772, 774-778

FEMALE

| | | | | | | | | | | | | |
|------|------|---------|---|---|---|---|---|---|---|---|---|---|
| 1978 | 8.2 | 574.1 | * | * | * | * | * | * | * | * | * | * |
| 1977 | 8.8 | 621.2 | * | * | * | * | * | * | * | * | * | * |
| 1976 | 9.5 | 695.3 | * | * | * | * | * | * | * | * | * | * |
| 1975 | 10.2 | 732.4 | * | * | * | * | * | * | * | * | * | * |
| 1974 | 11.0 | 804.9 | * | * | * | * | * | * | * | * | * | * |
| 1973 | 11.7 | 827.1 | * | * | * | * | * | * | * | * | * | * |
| 1972 | 13.1 | 866.9 | * | * | * | * | * | * | * | * | * | * |
| 1971 | 15.0 | 902.9 | * | * | * | * | * | * | * | * | * | * |
| 1970 | 17.2 | 1,051.9 | * | * | * | * | * | * | * | * | * | * |
| 1969 | 17.2 | 1,062.8 | * | * | * | * | * | * | * | * | * | * |
| 1968 | 17.6 | 1,086.1 | * | * | * | * | * | * | * | * | * | * |

WHITE, BOTH SEXES

| | | | | | | | | | | | | |
|------|------|---------|-----|---|---|---|---|---|---|---|---|---|
| 1978 | 7.8 | 544.4 | * | * | * | * | * | * | * | * | * | * |
| 1977 | 8.5 | 592.8 | 0.2 | * | * | * | * | * | * | * | * | * |
| 1976 | 9.1 | 664.8 | * | * | * | * | * | * | * | * | * | * |
| 1975 | 10.1 | 717.0 | * | * | * | * | * | * | * | * | * | * |
| 1974 | 11.2 | 815.6 | * | * | * | * | * | * | * | * | * | * |
| 1973 | 12.0 | 849.2 | * | * | * | * | * | * | * | * | * | * |
| 1972 | 13.4 | 886.5 | * | * | * | * | * | * | * | * | * | * |
| 1971 | 15.8 | 939.4 | * | * | * | * | * | * | * | * | * | * |
| 1970 | 18.1 | 1,096.8 | * | * | * | * | * | * | * | * | * | * |
| 1969 | 18.3 | 1,112.1 | * | * | * | * | * | * | * | * | * | * |
| 1968 | 18.9 | 1,159.2 | 0.3 | * | * | * | * | * | * | * | * | * |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR TOTAL UNDER 1 YEAR 1-4 YEARS 5-14 YEARS 15-24 YEARS 25-34 YEARS 35-44 YEARS 45-54 YEARS 55-64 YEARS 65-74 YEARS 75-84 YEARS 85+ YEARS

Certain causes of mortality in early infancy 760-769.2, 769.4-772, 774-778

WHITE, MALE

| | | | | | | | | | | | | |
|------|------|---------|---|---|---|---|---|---|---|---|---|---|
| 1978 | 9.3 | 616.1 | * | * | * | * | * | * | * | * | * | * |
| 1977 | 10.1 | 672.2 | * | * | * | * | * | * | * | * | * | * |
| 1976 | 10.8 | 750.5 | * | * | * | * | * | * | * | * | * | * |
| 1975 | 12.1 | 819.8 | * | * | * | * | * | * | * | * | * | * |
| 1974 | 13.6 | 937.0 | * | * | * | * | * | * | * | * | * | * |
| 1973 | 14.5 | 978.3 | * | * | * | * | * | * | * | * | * | * |
| 1972 | 16.3 | 1,026.0 | * | * | * | * | * | * | * | * | * | * |
| 1971 | 19.2 | 1,087.8 | * | * | * | * | * | * | * | * | * | * |
| 1970 | 21.9 | 1,262.7 | * | * | * | * | * | * | * | * | * | * |
| 1969 | 22.3 | 1,290.3 | * | * | * | * | * | * | * | * | * | * |
| 1968 | 23.2 | 1,359.1 | * | * | * | * | * | * | * | * | * | * |

WHITE, FEMALE

| | | | | | | | | | | | | |
|------|------|-------|---|---|---|---|---|---|---|---|---|---|
| 1978 | 6.4 | 468.3 | * | * | * | * | * | * | * | * | * | * |
| 1977 | 6.9 | 509.0 | * | * | * | * | * | * | * | * | * | * |
| 1976 | 7.5 | 574.7 | * | * | * | * | * | * | * | * | * | * |
| 1975 | 8.1 | 608.1 | * | * | * | * | * | * | * | * | * | * |
| 1974 | 9.0 | 687.5 | * | * | * | * | * | * | * | * | * | * |
| 1973 | 9.6 | 713.6 | * | * | * | * | * | * | * | * | * | * |
| 1972 | 10.7 | 740.1 | * | * | * | * | * | * | * | * | * | * |
| 1971 | 12.5 | 783.0 | * | * | * | * | * | * | * | * | * | * |
| 1970 | 14.5 | 923.1 | * | * | * | * | * | * | * | * | * | * |
| 1969 | 14.5 | 925.8 | * | * | * | * | * | * | * | * | * | * |
| 1968 | 14.7 | 949.5 | * | * | * | * | * | * | * | * | * | * |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR TOTAL UNDER 1 YEAR 1-4 YEARS 5-14 YEARS 15-24 YEARS 25-34 YEARS 35-44 YEARS 45-54 YEARS 55-64 YEARS 65-74 YEARS 75-84 YEARS 85+ YEARS

Certain causes of mortality in early infancy 760-769.2, 769.4-772, 774-778

ALL OTHER, BOTH SEXES

| | | | | | | | | | | | | |
|------|------|---------|---|---|---|---|---|---|---|---|---|---|
| 1978 | 23.2 | 1,224.2 | * | * | * | * | * | * | * | * | * | * |
| 1977 | 24.6 | 1,291.1 | * | * | * | * | * | * | * | * | * | * |
| 1976 | 26.2 | 1,432.0 | * | * | * | * | * | * | * | * | * | * |
| 1975 | 27.5 | 1,476.2 | * | * | * | * | * | * | * | * | * | * |
| 1974 | 28.7 | 1,538.2 | * | * | * | * | * | * | * | * | * | * |
| 1973 | 30.8 | 1,575.5 | * | * | * | * | * | * | * | * | * | * |
| 1972 | 35.0 | 1,706.5 | * | * | * | * | * | * | * | * | * | * |
| 1971 | 38.4 | 1,744.0 | * | * | * | * | * | * | * | * | * | * |
| 1970 | 43.2 | 1,996.4 | * | * | * | * | * | * | * | * | * | * |
| 1969 | 43.8 | 2,105.3 | * | * | * | * | * | * | * | * | * | * |
| 1968 | 44.5 | 2,050.6 | * | * | * | * | * | * | * | * | * | * |

ALL OTHER, MALE

| | | | | | | | | | | | | |
|------|------|---------|---|---|---|---|---|---|---|---|---|---|
| 1978 | 27.6 | 1,367.5 | * | * | * | * | * | * | * | * | * | * |
| 1977 | 28.8 | 1,422.2 | * | * | * | * | * | * | * | * | * | * |
| 1976 | 30.7 | 1,586.1 | * | * | * | * | * | * | * | * | * | * |
| 1975 | 31.7 | 1,602.3 | * | * | * | * | * | * | * | * | * | * |
| 1974 | 33.5 | 1,697.3 | * | * | * | * | * | * | * | * | * | * |
| 1973 | 36.5 | 1,765.0 | * | * | * | * | * | * | * | * | * | * |
| 1972 | 41.2 | 1,904.0 | * | * | * | * | * | * | * | * | * | * |
| 1971 | 45.4 | 1,950.5 | * | * | * | * | * | * | * | * | * | * |
| 1970 | 51.4 | 2,263.0 | * | * | * | * | * | * | * | * | * | * |
| 1969 | 52.1 | 2,388.8 | * | * | * | * | * | * | * | * | * | * |
| 1968 | 52.1 | 2,281.9 | * | * | * | * | * | * | * | * | * | * |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|

Certain causes of mortality in early infancy 760-769.2, 769.4-772, 774-778

ALL OTHER, FEMALE

| | | | | | | | | | | | | |
|------|------|---------|---|---|---|---|---|---|---|---|---|---|
| 1978 | 19.3 | 1,077.3 | * | * | * | * | * | * | * | * | * | * |
| 1977 | 20.8 | 1,156.7 | * | * | * | * | * | * | * | * | * | * |
| 1976 | 22.2 | 1,280.5 | * | * | * | * | * | * | * | * | * | * |
| 1975 | 23.6 | 1,346.7 | * | * | * | * | * | * | * | * | * | * |
| 1974 | 24.3 | 1,381.4 | * | * | * | * | * | * | * | * | * | * |
| 1973 | 25.5 | 1,381.5 | * | * | * | * | * | * | * | * | * | * |
| 1972 | 29.3 | 1,505.2 | * | * | * | * | * | * | * | * | * | * |
| 1971 | 32.0 | 1,533.1 | * | * | * | * | * | * | * | * | * | * |
| 1970 | 35.6 | 1,726.7 | * | * | * | * | * | * | * | * | * | * |
| 1969 | 36.2 | 1,819.5 | * | * | * | * | * | * | * | * | * | * |
| 1968 | 37.5 | 1,808.8 | * | * | * | * | * | * | * | * | * | * |

BLACK, BOTH SEXES

| | | | | | | | | | | | | |
|------|------|---------|---|---|---|---|---|---|---|---|---|---|
| 1978 | 25.5 | 1,340.5 | * | * | * | * | * | * | * | * | * | * |
| 1977 | 26.9 | 1,412.2 | * | * | * | * | * | * | * | * | * | * |
| 1976 | 28.6 | 1,574.5 | * | * | * | * | * | * | * | * | * | * |
| 1975 | 29.8 | 1,611.4 | * | * | * | * | * | * | * | * | * | * |
| 1974 | 30.8 | 1,672.3 | * | * | * | * | * | * | * | * | * | * |
| 1973 | 33.0 | 1,709.5 | * | * | * | * | * | * | * | * | * | * |
| 1972 | 37.4 | 1,844.4 | * | * | * | * | * | * | * | * | * | * |
| 1971 | 41.1 | 1,878.4 | * | * | * | * | * | * | * | * | * | * |
| 1970 | 46.4 | 2,147.2 | * | * | * | * | * | * | * | * | * | * |
| 1969 | 46.4 | 2,259.5 | * | * | * | * | * | * | * | * | * | * |
| 1968 | 47.0 | 2,175.9 | * | * | * | * | * | * | * | * | * | * |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

CAUSE OF DEATH CODE, TOTAL UNDER 1-4 5-14 15-24 25-34 35-44 45-54 55-64 65-74 75-84 85+
 RACE, SEX, AND YEAR 1 YEAR YEARS YEARS YEARS YEARS YEARS YEARS YEARS YEARS YEARS YEARS YEARS YEARS YEARS YEARS

Certain causes of mortality in early infancy 760-769.2, 769.4-772, 774-778

BLACK, MALE

| | | | | | | | | | | | | | | |
|------|------|---------|---|---|---|---|---|---|---|---|---|---|---|---|
| 1978 | 30.3 | 1,495.2 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1977 | 31.6 | 1,556.7 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1976 | 33.7 | 1,747.6 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1975 | 34.4 | 1,748.7 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1974 | 36.0 | 1,837.2 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1973 | 39.3 | 1,912.0 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1972 | 44.1 | 2,047.3 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1971 | 48.6 | 2,091.8 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1970 | 55.5 | 2,436.8 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1969 | 55.3 | 2,563.3 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1968 | 55.2 | 2,423.9 | * | * | * | * | * | * | * | * | * | * | * | * |

BLACK, FEMALE

| | | | | | | | | | | | | | | |
|------|------|---------|---|---|---|---|---|---|---|---|---|---|---|---|
| 1978 | 21.1 | 1,182.7 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1977 | 22.7 | 1,264.6 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1976 | 24.1 | 1,399.1 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1975 | 25.7 | 1,471.7 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1974 | 26.1 | 1,503.6 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1973 | 27.3 | 1,496.1 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1972 | 31.4 | 1,630.4 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1971 | 34.3 | 1,660.8 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1970 | 38.1 | 1,855.4 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1969 | 38.2 | 1,954.4 | * | * | * | * | * | * | * | * | * | * | * | * |
| 1968 | 39.5 | 1,924.7 | * | * | * | * | * | * | * | * | * | * | * | * |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|--|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| BIRTH INJURY, DIFFICULT LABOR, AND OTHER ANOXIC AND HYPOXIC CONDITIONS (764-768, 772, 776) | | | | | | | | | | | | |
| BOTH SEXES | | | | | | | | | | | | |
| 1978 | 5.3 | 351.0 | 0.2 | * | * | * | * | * | * | * | * | * |
| 1977 | 5.7 | 384.3 | 0.2 | * | * | * | * | * | * | * | * | * |
| 1976 | 6.2 | 430.4 | * | * | * | * | * | * | * | * | * | * |
| 1975 | 6.8 | 462.3 | * | * | * | * | * | * | * | * | * | * |
| 1974 | 7.5 | 519.7 | * | * | * | * | * | * | * | * | * | * |
| 1973 | 8.0 | 539.7 | * | * | * | * | * | * | * | * | * | * |
| 1972 | 8.9 | 564.9 | * | * | * | * | * | * | * | * | * | * |
| 1971 | 10.1 | 581.7 | * | * | * | * | * | * | * | * | * | * |
| 1970 | 11.2 | 653.8 | * | * | * | * | * | * | * | * | * | * |
| 1969 | 10.9 | 640.8 | * | * | * | * | * | * | * | * | * | * |
| 1968 | 11.1 | 655.1 | 0.2 | * | * | * | * | * | * | * | * | * |
| MALE | | | | | | | | | | | | |
| 1978 | 6.5 | 408.8 | * | * | * | * | * | * | * | * | * | * |
| 1977 | 6.9 | 438.9 | * | * | * | * | * | * | * | * | * | * |
| 1976 | 7.4 | 489.9 | * | * | * | * | * | * | * | * | * | * |
| 1975 | 8.2 | 534.0 | * | * | * | * | * | * | * | * | * | * |
| 1974 | 9.1 | 603.3 | * | * | * | * | * | * | * | * | * | * |
| 1973 | 10.0 | 639.7 | * | * | * | * | * | * | * | * | * | * |
| 1972 | 11.1 | 666.9 | * | * | * | * | * | * | * | * | * | * |
| 1971 | 12.6 | 686.3 | * | * | * | * | * | * | * | * | * | * |
| 1970 | 13.8 | 765.6 | * | * | * | * | * | * | * | * | * | * |
| 1969 | 13.6 | 760.4 | * | * | * | * | * | * | * | * | * | * |
| 1968 | 14.0 | 787.0 | * | * | * | * | * | * | * | * | * | * |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|--|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| BIRTH INJURY, DIFFICULT LABOR, AND OTHER ANOXIC AND HYPOXIC CONDITIONS (764-768, 772, 776) | | | | | | | | | | | | |
| FEMALE | | | | | | | | | | | | |
| 1978 | 4.1 | 290.5 | * | * | * | * | * | * | * | * | * | * |
| 1977 | 4.6 | 326.9 | * | * | * | * | * | * | * | * | * | * |
| 1976 | 5.0 | 367.8 | * | * | * | * | * | * | * | * | * | * |
| 1975 | 5.4 | 387.2 | * | * | * | * | * | * | * | * | * | * |
| 1974 | 5.9 | 432.0 | * | * | * | * | * | * | * | * | * | * |
| 1973 | 6.2 | 435.2 | * | * | * | * | * | * | * | * | * | * |
| 1972 | 6.9 | 458.3 | * | * | * | * | * | * | * | * | * | * |
| 1971 | 7.8 | 472.0 | * | * | * | * | * | * | * | * | * | * |
| 1970 | 8.8 | 537.3 | * | * | * | * | * | * | * | * | * | * |
| 1969 | 8.4 | 516.5 | * | * | * | * | * | * | * | * | * | * |
| 1968 | 8.4 | 517.6 | * | * | * | * | * | * | * | * | * | * |
| WHITE, BOTH SEXES | | | | | | | | | | | | |
| 1978 | 4.3 | 300.0 | * | * | * | * | * | * | * | * | * | * |
| 1977 | 4.7 | 328.4 | 0.2 | * | * | * | * | * | * | * | * | * |
| 1976 | 5.1 | 369.6 | * | * | * | * | * | * | * | * | * | * |
| 1975 | 5.7 | 405.4 | * | * | * | * | * | * | * | * | * | * |
| 1974 | 6.4 | 463.6 | * | * | * | * | * | * | * | * | * | * |
| 1973 | 6.8 | 481.3 | * | * | * | * | * | * | * | * | * | * |
| 1972 | 7.5 | 498.0 | * | * | * | * | * | * | * | * | * | * |
| 1971 | 8.8 | 522.9 | * | * | * | * | * | * | * | * | * | * |
| 1970 | 9.8 | 591.7 | * | * | * | * | * | * | * | * | * | * |
| 1969 | 9.5 | 576.8 | * | * | * | * | * | * | * | * | * | * |
| 1968 | 9.7 | 598.6 | 0.2 | * | * | * | * | * | * | * | * | * |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|--|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| BIRTH INJURY, DIFFICULT LABOR, AND OTHER ANOXIC AND HYPOXIC CONDITIONS (764-768, 772, 776) | | | | | | | | | | | | |
| WHITE, MALE | | | | | | | | | | | | |
| 1978 | 5.3 | 349.2 | * | * | * | * | * | * | * | * | * | * |
| 1977 | 5.7 | 378.9 | * | * | * | * | * | * | * | * | * | * |
| 1976 | 6.1 | 424.8 | * | * | * | * | * | * | * | * | * | * |
| 1975 | 7.0 | 475.7 | * | * | * | * | * | * | * | * | * | * |
| 1974 | 7.9 | 546.0 | * | * | * | * | * | * | * | * | * | * |
| 1973 | 8.5 | 575.5 | * | * | * | * | * | * | * | * | * | * |
| 1972 | 9.5 | 597.9 | * | * | * | * | * | * | * | * | * | * |
| 1971 | 11.0 | 624.6 | * | * | * | * | * | * | * | * | * | * |
| 1970 | 12.1 | 699.0 | * | * | * | * | * | * | * | * | * | * |
| 1969 | 11.9 | 690.8 | * | * | * | * | * | * | * | * | * | * |
| 1968 | 12.4 | 729.6 | * | * | * | * | * | * | * | * | * | * |
| WHITE, FEMALE | | | | | | | | | | | | |
| 1978 | 3.4 | 247.8 | * | * | * | * | * | * | * | * | * | * |
| 1977 | 3.7 | 275.1 | * | * | * | * | * | * | * | * | * | * |
| 1976 | 4.1 | 311.6 | * | * | * | * | * | * | * | * | * | * |
| 1975 | 4.4 | 331.1 | * | * | * | * | * | * | * | * | * | * |
| 1974 | 4.9 | 376.7 | * | * | * | * | * | * | * | * | * | * |
| 1973 | 5.2 | 382.4 | * | * | * | * | * | * | * | * | * | * |
| 1972 | 5.7 | 393.3 | * | * | * | * | * | * | * | * | * | * |
| 1971 | 6.7 | 415.9 | * | * | * | * | * | * | * | * | * | * |
| 1970 | 7.6 | 479.4 | * | * | * | * | * | * | * | * | * | * |
| 1969 | 7.2 | 457.5 | * | * | * | * | * | * | * | * | * | * |
| 1968 | 7.2 | 461.2 | * | * | * | * | * | * | * | * | * | * |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|--|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| BIRTH INJURY, DIFFICULT LABOR, AND OTHER ANOXIC AND HYPOXIC CONDITIONS (764-768, 772, 776) | | | | | | | | | | | | |
| ALL OTHER, BOTH SEXES | | | | | | | | | | | | |
| 1978 | 11.3 | 597.4 | * | * | * | * | * | * | * | * | * | * |
| 1977 | 12.5 | 654.7 | * | * | * | * | * | * | * | * | * | * |
| 1976 | 13.3 | 727.3 | * | * | * | * | * | * | * | * | * | * |
| 1975 | 13.9 | 746.9 | * | * | * | * | * | * | * | * | * | * |
| 1974 | 14.9 | 797.9 | * | * | * | * | * | * | * | * | * | * |
| 1973 | 16.2 | 827.8 | * | * | * | * | * | * | * | * | * | * |
| 1972 | 18.6 | 907.2 | * | * | * | * | * | * | * | * | * | * |
| 1971 | 19.7 | 895.2 | * | * | * | * | * | * | * | * | * | * |
| 1970 | 21.3 | 984.9 | * | * | * | * | * | * | * | * | * | * |
| 1969 | 20.9 | 1,002.3 | * | * | * | * | * | * | * | * | * | * |
| 1968 | 20.9 | 960.3 | * | * | * | * | * | * | * | * | * | * |
| ALL OTHER, MALE | | | | | | | | | | | | |
| 1978 | 14.1 | 699.0 | * | * | * | * | * | * | * | * | * | * |
| 1977 | 14.9 | 733.5 | * | * | * | * | * | * | * | * | * | * |
| 1976 | 15.8 | 814.7 | * | * | * | * | * | * | * | * | * | * |
| 1975 | 16.4 | 827.1 | * | * | * | * | * | * | * | * | * | * |
| 1974 | 17.7 | 893.1 | * | * | * | * | * | * | * | * | * | * |
| 1973 | 19.8 | 959.0 | * | * | * | * | * | * | * | * | * | * |
| 1972 | 22.2 | 1,027.1 | * | * | * | * | * | * | * | * | * | * |
| 1971 | 23.8 | 1,021.3 | * | * | * | * | * | * | * | * | * | * |
| 1970 | 25.6 | 1,127.4 | * | * | * | * | * | * | * | * | * | * |
| 1969 | 25.3 | 1,160.9 | * | * | * | * | * | * | * | * | * | * |
| 1968 | 25.1 | 1,099.2 | * | * | * | * | * | * | * | * | * | * |
| ALL OTHER, FEMALE | | | | | | | | | | | | |
| 1978 | 8.8 | 493.3 | * | * | * | * | * | * | * | * | * | * |
| 1977 | 10.3 | 574.0 | * | * | * | * | * | * | * | * | * | * |
| 1976 | 11.1 | 641.0 | * | * | * | * | * | * | * | * | * | * |
| 1975 | 11.7 | 664.5 | * | * | * | * | * | * | * | * | * | * |
| 1974 | 12.4 | 703.6 | * | * | * | * | * | * | * | * | * | * |
| 1973 | 12.8 | 693.5 | * | * | * | * | * | * | * | * | * | * |
| 1972 | 15.3 | 785.1 | * | * | * | * | * | * | * | * | * | * |
| 1971 | 16.0 | 766.5 | * | * | * | * | * | * | * | * | * | * |
| 1970 | 17.3 | 840.9 | * | * | * | * | * | * | * | * | * | * |
| 1969 | 16.8 | 842.6 | * | * | * | * | * | * | * | * | * | * |
| 1968 | 17.0 | 816.0 | * | * | * | * | * | * | * | * | * | * |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|--|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| BIRTH INJURY, DIFFICULT LABOR, AND OTHER ANOXIC AND HYPOXIC CONDITIONS (764-768, 772, 776) | | | | | | | | | | | | |
| BLACK, BOTH SEXES | | | | | | | | | | | | |
| 1978 | 12.3 | 647.7 | * | * | * | * | * | * | * | * | * | * |
| 1977 | 13.6 | 714.8 | * | * | * | * | * | * | * | * | * | * |
| 1976 | 14.5 | 797.8 | * | * | * | * | * | * | * | * | * | * |
| 1975 | 15.1 | 814.0 | * | * | * | * | * | * | * | * | * | * |
| 1974 | 15.9 | 864.4 | * | * | * | * | * | * | * | * | * | * |
| 1973 | 17.3 | 898.0 | * | * | * | * | * | * | * | * | * | * |
| 1972 | 19.8 | 976.1 | * | * | * | * | * | * | * | * | * | * |
| 1971 | 21.0 | 961.2 | * | * | * | * | * | * | * | * | * | * |
| 1970 | 22.8 | 1,056.4 | * | * | * | * | * | * | * | * | * | * |
| 1969 | 22.0 | 1,073.7 | * | * | * | * | * | * | * | * | * | * |
| 1968 | 21.8 | 1,009.7 | * | * | * | * | * | * | * | * | * | * |
| BLACK, MALE | | | | | | | | | | | | |
| 1978 | 15.3 | 755.6 | * | * | * | * | * | * | * | * | * | * |
| 1977 | 16.3 | 800.8 | * | * | * | * | * | * | * | * | * | * |
| 1976 | 17.3 | 899.1 | * | * | * | * | * | * | * | * | * | * |
| 1975 | 17.7 | 901.7 | * | * | * | * | * | * | * | * | * | * |
| 1974 | 18.9 | 962.8 | * | * | * | * | * | * | * | * | * | * |
| 1973 | 21.3 | 1,039.1 | * | * | * | * | * | * | * | * | * | * |
| 1972 | 23.7 | 1,097.9 | * | * | * | * | * | * | * | * | * | * |
| 1971 | 25.3 | 1,090.2 | * | * | * | * | * | * | * | * | * | * |
| 1970 | 27.5 | 1,209.0 | * | * | * | * | * | * | * | * | * | * |
| 1969 | 26.8 | 1,242.8 | * | * | * | * | * | * | * | * | * | * |
| 1968 | 26.4 | 1,158.8 | * | * | * | * | * | * | * | * | * | * |
| BLACK, FEMALE | | | | | | | | | | | | |
| 1978 | 9.6 | 537.4 | * | * | * | * | * | * | * | * | * | * |
| 1977 | 11.3 | 627.1 | * | * | * | * | * | * | * | * | * | * |
| 1976 | 12.0 | 695.1 | * | * | * | * | * | * | * | * | * | * |
| 1975 | 12.6 | 724.8 | * | * | * | * | * | * | * | * | * | * |
| 1974 | 13.3 | 763.8 | * | * | * | * | * | * | * | * | * | * |
| 1973 | 13.7 | 750.7 | * | * | * | * | * | * | * | * | * | * |
| 1972 | 16.3 | 848.1 | * | * | * | * | * | * | * | * | * | * |
| 1971 | 17.1 | 829.6 | * | * | * | * | * | * | * | * | * | * |
| 1970 | 18.5 | 902.8 | * | * | * | * | * | * | * | * | * | * |
| 1969 | 17.7 | 903.9 | * | * | * | * | * | * | * | * | * | * |
| 1968 | 17.6 | 858.7 | * | * | * | * | * | * | * | * | * | * |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| OTHER CAUSES OF MORTALITY IN EARLY INFANCY (REMAINDER OF 760-778) | | | | | | | | | | | | |
| BOTH SEXES | | | | | | | | | | | | |
| 1978 | 4.6 | 310.0 | * | * | * | * | * | * | * | * | * | * |
| 1977 | 4.9 | 328.0 | * | * | * | * | * | * | * | * | * | * |
| 1976 | 5.2 | 364.7 | * | * | * | * | * | * | * | * | * | * |
| 1975 | 5.6 | 381.1 | * | * | * | * | * | * | * | * | * | * |
| 1974 | 6.0 | 417.1 | * | * | * | * | * | * | * | * | * | * |
| 1973 | 6.4 | 431.9 | * | * | * | * | * | * | * | * | * | * |
| 1972 | 7.2 | 455.5 | * | * | * | * | * | * | * | * | * | * |
| 1971 | 8.5 | 484.6 | * | * | * | * | * | * | * | * | * | * |
| 1970 | 10.0 | 585.1 | * | * | * | * | * | * | * | * | * | * |
| 1969 | 10.5 | 620.8 | * | * | * | * | * | * | * | * | * | * |
| 1968 | 10.9 | 643.4 | * | * | * | * | * | * | * | * | * | * |
| MALE | | | | | | | | | | | | |
| 1978 | 5.3 | 335.3 | * | * | * | * | * | * | * | * | * | * |
| 1977 | 5.7 | 360.1 | * | * | * | * | * | * | * | * | * | * |
| 1976 | 6.0 | 400.1 | * | * | * | * | * | * | * | * | * | * |
| 1975 | 6.4 | 415.4 | * | * | * | * | * | * | * | * | * | * |
| 1974 | 6.9 | 459.2 | * | * | * | * | * | * | * | * | * | * |
| 1973 | 7.3 | 470.2 | * | * | * | * | * | * | * | * | * | * |
| 1972 | 8.3 | 500.4 | * | * | * | * | * | * | * | * | * | * |
| 1971 | 9.8 | 535.8 | * | * | * | * | * | * | * | * | * | * |
| 1970 | 11.7 | 652.8 | * | * | * | * | * | * | * | * | * | * |
| 1969 | 12.3 | 692.6 | * | * | * | * | * | * | * | * | * | * |
| 1968 | 12.7 | 715.2 | * | * | * | * | * | * | * | * | * | * |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| OTHER CAUSES OF MORTALITY IN EARLY INFANCY (REMAINDER OF 760-778) | | | | | | | | | | | | |
| FEMALE | | | | | | | | | | | | |
| 1978 | 4.0 | 283.6 | * | * | * | * | * | * | * | * | * | * |
| 1977 | 4.2 | 294.3 | * | * | * | * | * | * | * | * | * | * |
| 1976 | 4.5 | 327.4 | * | * | * | * | * | * | * | * | * | * |
| 1975 | 4.8 | 345.2 | * | * | * | * | * | * | * | * | * | * |
| 1974 | 5.1 | 372.9 | * | * | * | * | * | * | * | * | * | * |
| 1973 | 5.5 | 391.9 | * | * | * | * | * | * | * | * | * | * |
| 1972 | 6.2 | 408.7 | * | * | * | * | * | * | * | * | * | * |
| 1971 | 7.2 | 430.9 | * | * | * | * | * | * | * | * | * | * |
| 1970 | 8.4 | 514.5 | * | * | * | * | * | * | * | * | * | * |
| 1969 | 8.8 | 546.3 | * | * | * | * | * | * | * | * | * | * |
| 1968 | 9.2 | 568.5 | * | * | * | * | * | * | * | * | * | * |
| WHITE, BOTH SEXES | | | | | | | | | | | | |
| 1978 | 3.5 | 244.4 | * | * | * | * | * | * | * | * | * | * |
| 1977 | 3.8 | 264.4 | * | * | * | * | * | * | * | * | * | * |
| 1976 | 4.1 | 295.2 | * | * | * | * | * | * | * | * | * | * |
| 1975 | 4.4 | 311.5 | * | * | * | * | * | * | * | * | * | * |
| 1974 | 4.8 | 351.9 | * | * | * | * | * | * | * | * | * | * |
| 1973 | 5.2 | 367.9 | * | * | * | * | * | * | * | * | * | * |
| 1972 | 5.9 | 388.4 | * | * | * | * | * | * | * | * | * | * |
| 1971 | 7.0 | 416.4 | * | * | * | * | * | * | * | * | * | * |
| 1970 | 8.3 | 505.1 | * | * | * | * | * | * | * | * | * | * |
| 1969 | 8.8 | 535.3 | * | * | * | * | * | * | * | * | * | * |
| 1968 | 9.1 | 560.6 | * | * | * | * | * | * | * | * | * | * |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| OTHER CAUSES OF MORTALITY IN EARLY INFANCY (REMAINDER OF 760-778) | | | | | | | | | | | | |
| WHITE, MALE | | | | | | | | | | | | |
| 1978 | 4.0 | 266.9 | * | * | * | * | * | * | * | * | * | * |
| 1977 | 4.4 | 293.3 | * | * | * | * | * | * | * | * | * | * |
| 1976 | 4.7 | 325.7 | * | * | * | * | * | * | * | * | * | * |
| 1975 | 5.1 | 344.1 | * | * | * | * | * | * | * | * | * | * |
| 1974 | 5.7 | 391.0 | * | * | * | * | * | * | * | * | * | * |
| 1973 | 6.0 | 402.8 | * | * | * | * | * | * | * | * | * | * |
| 1972 | 6.8 | 428.1 | * | * | * | * | * | * | * | * | * | * |
| 1971 | 8.2 | 463.2 | * | * | * | * | * | * | * | * | * | * |
| 1970 | 9.8 | 563.8 | * | * | * | * | * | * | * | * | * | * |
| 1969 | 10.3 | 599.5 | * | * | * | * | * | * | * | * | * | * |
| 1968 | 10.7 | 629.5 | * | * | * | * | * | * | * | * | * | * |
| WHITE, FEMALE | | | | | | | | | | | | |
| 1978 | 3.0 | 220.5 | * | * | * | * | * | * | * | * | * | * |
| 1977 | 3.2 | 233.9 | * | * | * | * | * | * | * | * | * | * |
| 1976 | 3.4 | 263.1 | * | * | * | * | * | * | * | * | * | * |
| 1975 | 3.7 | 277.0 | * | * | * | * | * | * | * | * | * | * |
| 1974 | 4.1 | 310.8 | * | * | * | * | * | * | * | * | * | * |
| 1973 | 4.5 | 331.3 | * | * | * | * | * | * | * | * | * | * |
| 1972 | 5.0 | 346.8 | * | * | * | * | * | * | * | * | * | * |
| 1971 | 5.9 | 367.1 | * | * | * | * | * | * | * | * | * | * |
| 1970 | 7.0 | 443.7 | * | * | * | * | * | * | * | * | * | * |
| 1969 | 7.3 | 468.3 | * | * | * | * | * | * | * | * | * | * |
| 1968 | 7.6 | 488.3 | * | * | * | * | * | * | * | * | * | * |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| OTHER CAUSES OF MORTALITY IN EARLY INFANCY (REMAINDER OF 760-778) | | | | | | | | | | | | |
| ALL OTHER, BOTH SEXES | | | | | | | | | | | | |
| 1978 | 11.9 | 626.8 | * | * | * | * | * | * | * | * | * | * |
| 1977 | 12.1 | 636.4 | * | * | * | * | * | * | * | * | * | * |
| 1976 | 12.9 | 704.7 | * | * | * | * | * | * | * | * | * | * |
| 1975 | 13.6 | 729.3 | * | * | * | * | * | * | * | * | * | * |
| 1974 | 13.8 | 740.4 | * | * | * | * | * | * | * | * | * | * |
| 1973 | 14.6 | 747.7 | * | * | * | * | * | * | * | * | * | * |
| 1972 | 16.4 | 799.3 | * | * | * | * | * | * | * | * | * | * |
| 1971 | 18.7 | 848.8 | * | * | * | * | * | * | * | * | * | * |
| 1970 | 21.9 | 1,011.5 | * | * | * | * | * | * | * | * | * | * |
| 1969 | 22.9 | 1,102.9 | * | * | * | * | * | * | * | * | * | * |
| 1968 | 23.7 | 1,090.3 | * | * | * | * | * | * | * | * | * | * |
| ALL OTHER, MALE | | | | | | | | | | | | |
| 1978 | 13.5 | 668.5 | * | * | * | * | * | * | * | * | * | * |
| 1977 | 13.9 | 688.7 | * | * | * | * | * | * | * | * | * | * |
| 1976 | 14.9 | 771.4 | * | * | * | * | * | * | * | * | * | * |
| 1975 | 15.3 | 775.2 | * | * | * | * | * | * | * | * | * | * |
| 1974 | 15.8 | 804.2 | * | * | * | * | * | * | * | * | * | * |
| 1973 | 16.7 | 806.0 | * | * | * | * | * | * | * | * | * | * |
| 1972 | 19.0 | 876.9 | * | * | * | * | * | * | * | * | * | * |
| 1971 | 21.6 | 929.3 | * | * | * | * | * | * | * | * | * | * |
| 1970 | 25.8 | 1,135.7 | * | * | * | * | * | * | * | * | * | * |
| 1969 | 26.8 | 1,227.9 | * | * | * | * | * | * | * | * | * | * |
| 1968 | 27.0 | 1,182.6 | * | * | * | * | * | * | * | * | * | * |
| ALL OTHER, FEMALE | | | | | | | | | | | | |
| 1978 | 10.4 | 584.0 | * | * | * | * | * | * | * | * | * | * |
| 1977 | 10.5 | 582.7 | * | * | * | * | * | * | * | * | * | * |
| 1976 | 11.1 | 639.5 | * | * | * | * | * | * | * | * | * | * |
| 1975 | 12.0 | 682.2 | * | * | * | * | * | * | * | * | * | * |
| 1974 | 11.9 | 677.9 | * | * | * | * | * | * | * | * | * | * |
| 1973 | 12.7 | 688.1 | * | * | * | * | * | * | * | * | * | * |
| 1972 | 14.0 | 720.1 | * | * | * | * | * | * | * | * | * | * |
| 1971 | 16.0 | 766.5 | * | * | * | * | * | * | * | * | * | * |
| 1970 | 18.3 | 885.8 | * | * | * | * | * | * | * | * | * | * |
| 1969 | 19.4 | 977.0 | * | * | * | * | * | * | * | * | * | * |
| 1968 | 20.6 | 992.7 | * | * | * | * | * | * | * | * | * | * |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| OTHER CAUSES OF MORTALITY IN EARLY INFANCY (REMAINDER OF 760-778) | | | | | | | | | | | | |
| BLACK, BOTH SEXES | | | | | | | | | | | | |
| 1978 | 13.2 | 692.9 | * | * | * | * | * | * | * | * | * | * |
| 1977 | 13.3 | 697.3 | * | * | * | * | * | * | * | * | * | * |
| 1976 | 14.1 | 776.7 | * | * | * | * | * | * | * | * | * | * |
| 1975 | 14.7 | 797.4 | * | * | * | * | * | * | * | * | * | * |
| 1974 | 14.9 | 807.8 | * | * | * | * | * | * | * | * | * | * |
| 1973 | 15.7 | 811.5 | * | * | * | * | * | * | * | * | * | * |
| 1972 | 17.6 | 868.3 | * | * | * | * | * | * | * | * | * | * |
| 1971 | 20.1 | 917.2 | * | * | * | * | * | * | * | * | * | * |
| 1970 | 23.5 | 1,090.7 | * | * | * | * | * | * | * | * | * | * |
| 1969 | 24.3 | 1,185.8 | * | * | * | * | * | * | * | * | * | * |
| 1968 | 25.1 | 1,166.2 | * | * | * | * | * | * | * | * | * | * |
| BLACK, MALE | | | | | | | | | | | | |
| 1978 | 15.0 | 739.5 | * | * | * | * | * | * | * | * | * | * |
| 1977 | 15.3 | 755.9 | * | * | * | * | * | * | * | * | * | * |
| 1976 | 16.3 | 848.5 | * | * | * | * | * | * | * | * | * | * |
| 1975 | 16.6 | 847.0 | * | * | * | * | * | * | * | * | * | * |
| 1974 | 17.1 | 874.3 | * | * | * | * | * | * | * | * | * | * |
| 1973 | 18.0 | 873.0 | * | * | * | * | * | * | * | * | * | * |
| 1972 | 20.4 | 949.4 | * | * | * | * | * | * | * | * | * | * |
| 1971 | 23.3 | 1,001.6 | * | * | * | * | * | * | * | * | * | * |
| 1970 | 27.9 | 1,227.8 | * | * | * | * | * | * | * | * | * | * |
| 1969 | 28.5 | 1,320.5 | * | * | * | * | * | * | * | * | * | * |
| 1968 | 28.8 | 1,265.1 | * | * | * | * | * | * | * | * | * | * |
| BLACK, FEMALE | | | | | | | | | | | | |
| 1978 | 11.5 | 645.3 | * | * | * | * | * | * | * | * | * | * |
| 1977 | 11.4 | 637.5 | * | * | * | * | * | * | * | * | * | * |
| 1976 | 12.1 | 704.0 | * | * | * | * | * | * | * | * | * | * |
| 1975 | 13.0 | 746.9 | * | * | * | * | * | * | * | * | * | * |
| 1974 | 12.8 | 739.8 | * | * | * | * | * | * | * | * | * | * |
| 1973 | 13.6 | 745.4 | * | * | * | * | * | * | * | * | * | * |
| 1972 | 15.1 | 782.3 | * | * | * | * | * | * | * | * | * | * |
| 1971 | 17.2 | 831.2 | * | * | * | * | * | * | * | * | * | * |
| 1970 | 19.6 | 952.6 | * | * | * | * | * | * | * | * | * | * |
| 1969 | 20.5 | 1,050.4 | * | * | * | * | * | * | * | * | * | * |
| 1968 | 21.8 | 1,066.0 | * | * | * | * | * | * | * | * | * | * |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| SYMPTOMS AND ILL-DEFINED CONDITIONS (780-796) | | | | | | | | | | | | |
| BOTH SEXES | | | | | | | | | | | | |
| 1978 | 14.1 | 171.8 | 2.4 | 0.4 | 1.9 | 3.5 | 5.5 | 10.4 | 19.5 | 37.1 | 71.4 | 207.3 |
| 1977 | 14.6 | 170.2 | 2.5 | 0.4 | 2.4 | 3.7 | 6.2 | 11.7 | 20.8 | 39.8 | 74.7 | 199.4 |
| 1976 | 14.2 | 167.2 | 2.4 | 0.4 | 2.5 | 3.9 | 5.9 | 11.5 | 20.5 | 38.8 | 71.6 | 204.0 |
| 1975 | 14.7 | 162.5 | 2.7 | 0.7 | 3.7 | 4.8 | 7.1 | 12.6 | 22.3 | 38.9 | 70.1 | 199.0 |
| 1974 | 14.6 | 153.3 | 2.8 | 0.7 | 4.4 | 5.2 | 8.0 | 12.7 | 22.3 | 38.3 | 67.5 | 199.1 |
| 1973 | 14.4 | 134.5 | 2.7 | 0.6 | 3.9 | 4.9 | 8.3 | 13.3 | 23.3 | 38.6 | 69.8 | 202.7 |
| 1972 | 13.1 | 114.7 | 2.7 | 0.5 | 2.4 | 3.5 | 6.1 | 12.2 | 22.7 | 39.1 | 69.0 | 191.7 |
| 1971 | 12.8 | 99.5 | 2.3 | 0.4 | 2.5 | 3.5 | 6.6 | 11.8 | 22.4 | 37.9 | 67.6 | 191.6 |
| 1970 | 12.7 | 100.9 | 2.1 | 0.5 | 2.1 | 3.5 | 6.5 | 11.9 | 23.1 | 37.8 | 67.1 | 176.7 |
| 1969 | 13.0 | 94.0 | 2.3 | 0.4 | 2.3 | 3.7 | 7.3 | 12.9 | 23.8 | 39.7 | 70.3 | 193.9 |
| 1968 | 11.9 | 87.9 | 2.1 | 0.4 | 1.6 | 2.9 | 6.0 | 11.1 | 21.6 | 36.3 | 68.3 | 212.4 |
| MALE | | | | | | | | | | | | |
| 1978 | 16.7 | 198.9 | 2.7 | 0.5 | 2.5 | 4.5 | 7.0 | 14.3 | 28.4 | 53.4 | 98.2 | 240.3 |
| 1977 | 17.4 | 198.4 | 2.7 | 0.5 | 3.2 | 5.0 | 7.5 | 15.9 | 30.1 | 56.8 | 103.1 | 235.2 |
| 1976 | 17.0 | 191.7 | 3.0 | 0.5 | 3.5 | 5.0 | 7.8 | 15.9 | 29.2 | 55.7 | 98.0 | 242.1 |
| 1975 | 17.9 | 188.3 | 2.9 | 0.8 | 5.2 | 6.4 | 9.1 | 17.0 | 32.2 | 56.0 | 96.0 | 233.2 |
| 1974 | 17.8 | 178.1 | 3.2 | 0.9 | 6.3 | 7.0 | 9.7 | 17.2 | 31.6 | 55.1 | 91.4 | 233.1 |
| 1973 | 17.7 | 156.4 | 3.0 | 0.7 | 5.3 | 6.3 | 10.5 | 18.0 | 34.2 | 56.1 | 95.4 | 247.8 |
| 1972 | 16.1 | 132.7 | 3.1 | 0.6 | 3.3 | 4.4 | 7.5 | 17.3 | 33.0 | 56.5 | 95.8 | 225.3 |
| 1971 | 15.7 | 114.7 | 2.3 | 0.5 | 3.4 | 4.3 | 8.2 | 16.2 | 32.6 | 55.3 | 92.4 | 222.9 |
| 1970 | 15.5 | 117.6 | 2.0 | 0.4 | 2.5 | 4.0 | 8.0 | 16.2 | 33.3 | 55.4 | 91.1 | 197.5 |
| 1969 | 16.1 | 106.8 | 2.6 | 0.4 | 3.0 | 4.7 | 9.3 | 17.8 | 34.9 | 58.1 | 94.3 | 220.9 |
| 1968 | 14.6 | 99.0 | 2.4 | 0.5 | 2.0 | 3.3 | 7.5 | 15.2 | 31.5 | 53.3 | 91.7 | 255.5 |

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| SYMPTOMS AND ILL-DEFINED CONDITIONS (780-796) | | | | | | | | | | | | |
| FEMALE | | | | | | | | | | | | |
| 1978 | 11.7 | 143.3 | 2.0 | 0.4 | 1.4 | 2.4 | 4.0 | 6.9 | 11.7 | 24.7 | 55.4 | 192.4 |
| 1977 | 12.0 | 140.5 | 2.3 | 0.3 | 1.7 | 2.5 | 5.0 | 7.9 | 12.6 | 26.8 | 57.6 | 183.0 |
| 1976 | 11.5 | 141.6 | 1.8 | 0.4 | 1.5 | 2.8 | 4.2 | 7.5 | 12.7 | 25.8 | 55.7 | 186.1 |
| 1975 | 11.8 | 135.3 | 2.4 | 0.5 | 2.3 | 3.2 | 5.1 | 8.6 | 13.4 | 25.9 | 54.3 | 182.5 |
| 1974 | 11.6 | 127.3 | 2.5 | 0.6 | 2.5 | 3.4 | 6.3 | 8.6 | 14.0 | 25.3 | 52.7 | 182.2 |
| 1973 | 11.3 | 111.6 | 2.4 | 0.6 | 2.4 | 3.5 | 6.1 | 8.9 | 13.6 | 25.2 | 53.8 | 180.1 |
| 1972 | 10.3 | 95.9 | 2.2 | 0.4 | 1.6 | 2.6 | 4.8 | 7.5 | 13.5 | 25.7 | 51.9 | 174.5 |
| 1971 | 10.1 | 83.6 | 2.2 | 0.3 | 1.6 | 2.8 | 5.1 | 7.7 | 13.3 | 24.5 | 51.5 | 175.2 |
| 1970 | 10.1 | 83.5 | 2.3 | 0.5 | 1.7 | 3.0 | 5.2 | 8.0 | 13.9 | 24.1 | 51.3 | 165.1 |
| 1969 | 10.1 | 80.7 | 2.0 | 0.4 | 1.6 | 2.8 | 5.3 | 8.2 | 13.8 | 25.3 | 54.2 | 179.2 |
| 1968 | 9.3 | 76.3 | 1.7 | 0.3 | 1.2 | 2.4 | 4.7 | 7.2 | 12.5 | 22.9 | 52.4 | 189.0 |
| WHITE, BOTH SEXES | | | | | | | | | | | | |
| 1978 | 11.9 | 135.2 | 1.9 | 0.3 | 1.7 | 2.7 | 4.1 | 8.1 | 16.1 | 30.7 | 60.8 | 182.5 |
| 1977 | 12.3 | 134.7 | 1.9 | 0.3 | 2.2 | 2.9 | 4.5 | 9.0 | 16.8 | 32.8 | 64.3 | 176.5 |
| 1976 | 11.8 | 132.4 | 2.0 | 0.3 | 2.1 | 3.0 | 4.2 | 9.0 | 16.7 | 31.8 | 60.7 | 177.6 |
| 1975 | 12.1 | 125.6 | 2.0 | 0.6 | 3.1 | 3.5 | 5.1 | 9.8 | 18.0 | 31.6 | 58.4 | 168.6 |
| 1974 | 11.9 | 118.4 | 2.1 | 0.6 | 3.6 | 3.7 | 5.5 | 9.9 | 18.0 | 30.1 | 57.1 | 170.2 |
| 1973 | 11.5 | 101.6 | 1.8 | 0.5 | 3.0 | 3.5 | 5.6 | 10.1 | 18.6 | 30.1 | 58.0 | 172.1 |
| 1972 | 10.5 | 87.0 | 2.1 | 0.4 | 1.8 | 2.5 | 4.1 | 9.3 | 17.7 | 29.1 | 58.0 | 163.4 |
| 1971 | 10.1 | 71.3 | 1.6 | 0.3 | 1.9 | 2.4 | 4.5 | 8.8 | 17.5 | 29.5 | 55.1 | 163.1 |
| 1970 | 9.9 | 71.2 | 1.5 | 0.4 | 1.6 | 2.4 | 4.4 | 8.8 | 17.4 | 29.1 | 54.4 | 146.6 |
| 1969 | 9.9 | 63.5 | 1.4 | 0.3 | 1.6 | 2.4 | 4.7 | 9.3 | 18.0 | 29.1 | 56.4 | 160.6 |
| 1968 | 8.8 | 55.8 | 1.4 | 0.3 | 1.2 | 1.7 | 3.8 | 7.9 | 15.9 | 26.1 | 53.4 | 169.4 |

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| SYMPTOMS AND ILL-DEFINED CONDITIONS (780-796) | | | | | | | | | | | | |
| WHITE, MALE | | | | | | | | | | | | |
| 1978 | 14.1 | 159.7 | 2.1 | 0.3 | 2.2 | 3.5 | 5.1 | 11.2 | 24.1 | 45.7 | 83.0 | 209.3 |
| 1977 | 14.7 | 161.3 | 2.1 | 0.4 | 2.9 | 3.9 | 5.3 | 12.2 | 24.8 | 48.7 | 88.8 | 206.1 |
| 1976 | 14.2 | 156.0 | 2.4 | 0.3 | 3.0 | 3.8 | 5.4 | 12.5 | 24.2 | 47.3 | 83.8 | 211.6 |
| 1975 | 14.7 | 149.8 | 2.2 | 0.6 | 4.4 | 4.7 | 6.3 | 13.1 | 26.7 | 46.7 | 80.2 | 198.9 |
| 1974 | 14.4 | 143.4 | 2.2 | 0.7 | 5.1 | 5.0 | 6.6 | 13.4 | 26.2 | 44.1 | 77.3 | 197.1 |
| 1973 | 14.2 | 120.8 | 2.1 | 0.6 | 4.1 | 4.4 | 7.0 | 13.8 | 28.0 | 45.4 | 79.1 | 207.1 |
| 1972 | 13.0 | 104.7 | 2.5 | 0.5 | 2.5 | 3.1 | 5.0 | 13.6 | 26.8 | 43.9 | 80.7 | 192.0 |
| 1971 | 12.5 | 85.9 | 1.7 | 0.4 | 2.7 | 3.0 | 5.5 | 12.1 | 26.3 | 45.0 | 75.3 | 186.2 |
| 1970 | 12.3 | 85.7 | 1.4 | 0.4 | 2.0 | 2.8 | 5.5 | 12.3 | 26.2 | 45.2 | 75.0 | 161.6 |
| 1969 | 12.5 | 76.2 | 1.6 | 0.3 | 2.2 | 3.0 | 6.0 | 13.3 | 27.4 | 44.7 | 77.8 | 182.9 |
| 1968 | 11.0 | 64.5 | 1.6 | 0.3 | 1.5 | 2.0 | 4.9 | 11.6 | 24.1 | 40.2 | 72.3 | 200.8 |
| WHITE, FEMALE | | | | | | | | | | | | |
| 1978 | 9.9 | 109.3 | 1.6 | 0.3 | 1.2 | 2.0 | 3.0 | 5.3 | 9.0 | 19.3 | 47.7 | 170.6 |
| 1977 | 10.0 | 106.7 | 1.8 | 0.3 | 1.5 | 1.9 | 3.8 | 6.0 | 9.7 | 20.6 | 49.8 | 163.0 |
| 1976 | 9.6 | 107.5 | 1.6 | 0.3 | 1.3 | 2.1 | 3.1 | 5.6 | 9.9 | 20.0 | 47.0 | 162.1 |
| 1975 | 9.6 | 100.0 | 1.9 | 0.5 | 1.9 | 2.4 | 3.8 | 6.7 | 10.3 | 20.1 | 45.3 | 154.3 |
| 1974 | 9.4 | 92.2 | 2.0 | 0.5 | 2.0 | 2.4 | 4.5 | 6.6 | 10.6 | 19.4 | 44.8 | 157.1 |
| 1973 | 8.9 | 81.4 | 1.5 | 0.4 | 1.9 | 2.5 | 4.3 | 6.7 | 10.2 | 18.4 | 44.9 | 154.9 |
| 1972 | 8.0 | 68.3 | 1.6 | 0.4 | 1.2 | 1.9 | 3.3 | 5.2 | 9.6 | 17.8 | 43.7 | 148.9 |
| 1971 | 7.8 | 56.0 | 1.5 | 0.3 | 1.2 | 1.8 | 3.5 | 5.7 | 9.5 | 17.5 | 42.1 | 151.3 |
| 1970 | 7.5 | 56.0 | 1.6 | 0.4 | 1.1 | 2.1 | 3.4 | 5.5 | 9.4 | 16.8 | 40.9 | 138.4 |
| 1969 | 7.4 | 50.3 | 1.2 | 0.3 | 1.1 | 1.8 | 3.4 | 5.7 | 9.5 | 17.0 | 42.2 | 149.0 |
| 1968 | 6.6 | 46.8 | 1.2 | 0.2 | 0.9 | 1.4 | 2.8 | 4.5 | 8.4 | 15.0 | 40.6 | 152.6 |

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| SYMPTOMS AND ILL-DEFINED CONDITIONS (780-796) | | | | | | | | | | | | |
| ALL OTHER, BOTH SEXES | | | | | | | | | | | | |
| 1978 | 28.0 | 348.2 | 4.6 | 0.9 | 3.3 | 8.1 | 15.1 | 28.2 | 50.0 | 96.8 | 180.3 | 491.7 |
| 1977 | 29.6 | 341.9 | 5.3 | 0.8 | 3.7 | 8.9 | 17.7 | 33.0 | 57.8 | 106.1 | 181.6 | 462.5 |
| 1976 | 29.3 | 337.7 | 4.3 | 0.8 | 4.8 | 9.9 | 17.4 | 32.4 | 55.6 | 105.4 | 183.4 | 503.3 |
| 1975 | 32.2 | 347.0 | 5.8 | 1.2 | 7.3 | 13.3 | 20.9 | 35.6 | 61.6 | 109.9 | 194.6 | 547.9 |
| 1974 | 33.0 | 326.7 | 6.7 | 1.4 | 9.2 | 15.6 | 25.1 | 36.3 | 62.5 | 118.6 | 182.0 | 534.8 |
| 1973 | 34.1 | 297.3 | 7.2 | 1.4 | 9.3 | 14.9 | 27.0 | 40.5 | 67.8 | 122.5 | 203.9 | 559.8 |
| 1972 | 31.5 | 256.6 | 5.8 | 0.8 | 6.1 | 10.3 | 20.1 | 37.8 | 69.8 | 137.8 | 196.5 | 518.7 |
| 1971 | 31.7 | 250.0 | 5.6 | 0.8 | 6.4 | 11.5 | 22.2 | 37.9 | 69.9 | 121.6 | 219.3 | 515.8 |
| 1970 | 32.4 | 259.2 | 5.6 | 0.9 | 5.5 | 11.0 | 22.0 | 39.6 | 76.8 | 122.5 | 228.8 | 485.5 |
| 1969 | 35.3 | 265.6 | 6.7 | 1.0 | 6.6 | 13.1 | 27.0 | 44.5 | 79.7 | 144.5 | 251.2 | 557.8 |
| 1968 | 34.0 | 260.8 | 5.6 | 1.1 | 4.4 | 11.3 | 23.0 | 39.6 | 76.7 | 138.6 | 262.0 | 693.1 |
| ALL OTHER, MALE | | | | | | | | | | | | |
| 1978 | 33.6 | 390.3 | 5.3 | 1.1 | 4.3 | 11.6 | 21.2 | 40.2 | 68.5 | 124.0 | 241.0 | 558.6 |
| 1977 | 35.3 | 380.6 | 5.8 | 0.9 | 4.9 | 12.1 | 24.0 | 47.0 | 80.7 | 132.3 | 237.9 | 532.1 |
| 1976 | 35.2 | 369.5 | 6.0 | 1.1 | 6.6 | 13.6 | 25.4 | 45.6 | 77.1 | 134.8 | 233.1 | 547.3 |
| 1975 | 39.3 | 383.1 | 6.5 | 1.7 | 10.0 | 19.0 | 29.9 | 50.7 | 85.0 | 143.8 | 251.7 | 575.9 |
| 1974 | 40.5 | 354.1 | 8.1 | 1.7 | 13.5 | 22.2 | 33.5 | 50.3 | 83.4 | 160.6 | 236.7 | 604.0 |
| 1973 | 42.1 | 334.6 | 7.3 | 1.6 | 13.0 | 20.9 | 37.5 | 55.8 | 94.3 | 158.9 | 266.1 | 662.5 |
| 1972 | 37.9 | 278.4 | 6.3 | 1.0 | 8.4 | 14.0 | 26.9 | 50.3 | 93.1 | 177.8 | 258.1 | 561.7 |
| 1971 | 38.3 | 270.7 | 5.7 | 1.0 | 8.7 | 14.6 | 29.6 | 53.7 | 93.8 | 153.7 | 282.6 | 585.1 |
| 1970 | 37.9 | 290.2 | 5.6 | 0.7 | 6.1 | 13.7 | 27.7 | 51.9 | 100.6 | 151.7 | 278.7 | 512.4 |
| 1969 | 42.3 | 282.6 | 7.7 | 1.1 | 8.5 | 17.2 | 36.1 | 60.2 | 108.0 | 187.2 | 292.0 | 579.5 |
| 1968 | 40.7 | 287.5 | 6.8 | 1.4 | 5.9 | 13.7 | 28.7 | 49.7 | 103.6 | 180.8 | 323.7 | 787.8 |
| ALL OTHER, FEMALE | | | | | | | | | | | | |
| 1978 | 23.0 | 305.0 | 3.8 | 0.7 | 2.3 | 5.1 | 9.9 | 18.3 | 34.7 | 75.8 | 139.3 | 460.6 |
| 1977 | 24.3 | 302.2 | 4.7 | 0.7 | 2.6 | 6.2 | 12.5 | 21.5 | 38.7 | 85.6 | 142.8 | 425.0 |
| 1976 | 24.0 | 306.5 | 2.5 | * | 3.0 | 6.6 | 10.8 | 21.3 | 37.6 | 82.0 | 149.5 | 478.6 |
| 1975 | 25.8 | 310.0 | 5.1 | 0.7 | 4.7 | 8.4 | 13.3 | 22.8 | 41.8 | 82.8 | 155.1 | 525.8 |
| 1974 | 26.1 | 300.0 | 5.2 | 1.1 | 5.1 | 9.9 | 18.1 | 24.4 | 44.7 | 85.0 | 144.3 | 494.1 |
| 1973 | 26.8 | 259.2 | 7.1 | 1.2 | 5.8 | 9.7 | 18.3 | 27.3 | 45.2 | 93.0 | 160.7 | 497.5 |
| 1972 | 25.5 | 234.3 | 5.3 | * | 3.8 | 7.0 | 14.4 | 27.0 | 49.7 | 105.5 | 152.8 | 492.1 |
| 1971 | 25.6 | 228.8 | 5.4 | * | 4.2 | 8.9 | 16.0 | 24.2 | 49.1 | 95.3 | 172.5 | 471.2 |
| 1970 | 27.3 | 227.8 | 5.6 | 1.0 | 5.0 | 8.7 | 17.2 | 28.8 | 55.7 | 98.5 | 190.8 | 466.5 |
| 1969 | 28.9 | 248.4 | 5.8 | 0.9 | 4.9 | 9.6 | 19.4 | 30.6 | 54.6 | 109.4 | 219.0 | 543.1 |
| 1968 | 27.9 | 232.8 | 4.5 | 0.8 | 3.0 | 9.2 | 18.2 | 30.7 | 52.5 | 103.6 | 215.1 | 629.5 |

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| SYMPTOMS AND ILL-DEFINED CONDITIONS (780-796) | | | | | | | | | | | | |
| BLACK, BOTH SEXES | | | | | | | | | | | | |
| 1978 | 30.9 | 378.0 | 4.9 | 0.9 | 3.3 | 9.2 | 16.8 | 31.7 | 55.8 | 104.3 | 187.9 | 526.7 |
| 1977 | 32.7 | 370.1 | 5.7 | 0.8 | 3.9 | 10.0 | 20.1 | 37.3 | 63.3 | 115.0 | 191.9 | 492.3 |
| 1976 | 32.1 | 363.3 | 4.5 | 0.8 | 5.1 | 11.1 | 19.8 | 36.1 | 60.0 | 115.1 | 192.8 | 530.9 |
| 1975 | 35.1 | 372.4 | 6.4 | 1.1 | 7.8 | 14.8 | 23.4 | 38.9 | 66.4 | 118.6 | 201.8 | 585.4 |
| 1974 | 35.8 | 349.2 | 7.2 | 1.5 | 9.9 | 17.4 | 27.8 | 40.0 | 67.3 | 128.0 | 189.0 | 567.8 |
| 1973 | 37.0 | 320.8 | 7.6 | 1.4 | 10.1 | 17.0 | 30.4 | 43.9 | 72.6 | 131.7 | 211.2 | 588.6 |
| 1972 | 34.1 | 274.6 | 5.9 | 0.9 | 6.6 | 11.0 | 22.6 | 41.7 | 75.3 | 147.6 | 206.7 | 552.7 |
| 1971 | 34.0 | 265.1 | 5.7 | 0.8 | 6.8 | 12.7 | 24.3 | 41.2 | 74.6 | 130.2 | 232.4 | 544.9 |
| 1970 | 35.2 | 279.4 | 6.0 | 0.9 | 6.1 | 12.1 | 23.9 | 43.0 | 82.8 | 131.6 | 248.9 | 527.3 |
| 1969 | 37.6 | 283.6 | 7.0 | 1.0 | 7.1 | 13.9 | 29.3 | 48.1 | 85.1 | 153.0 | 264.6 | 586.6 |
| 1968 | 36.0 | 274.6 | 5.5 | 1.0 | 4.6 | 12.4 | 24.5 | 42.2 | 81.5 | 146.0 | 275.5 | 725.0 |
| BLACK, MALE | | | | | | | | | | | | |
| 1978 | 36.9 | 420.2 | 5.7 | 1.1 | 4.2 | 13.0 | 23.6 | 45.4 | 77.0 | 135.6 | 251.5 | 600.0 |
| 1977 | 39.0 | 410.2 | 6.5 | 0.9 | 5.2 | 13.8 | 27.4 | 52.6 | 88.6 | 146.3 | 254.0 | 569.4 |
| 1976 | 38.6 | 400.9 | 6.2 | 1.1 | 7.1 | 15.3 | 29.1 | 51.2 | 83.5 | 151.0 | 243.2 | 574.5 |
| 1975 | 42.8 | 411.7 | 7.4 | 1.6 | 10.8 | 21.0 | 34.2 | 56.1 | 92.2 | 157.8 | 260.8 | 628.3 |
| 1974 | 44.1 | 381.0 | 8.6 | 1.8 | 14.7 | 25.1 | 37.1 | 55.3 | 91.1 | 177.4 | 246.9 | 662.8 |
| 1973 | 45.7 | 360.1 | 7.6 | 1.7 | 14.1 | 23.9 | 42.5 | 60.7 | 102.2 | 173.3 | 277.0 | 722.0 |
| 1972 | 41.1 | 297.9 | 6.5 | 1.1 | 9.4 | 14.9 | 30.5 | 55.8 | 101.9 | 192.3 | 271.1 | 635.0 |
| 1971 | 41.2 | 288.2 | 6.0 | 1.0 | 9.2 | 16.4 | 32.4 | 59.0 | 101.1 | 166.8 | 301.1 | 637.5 |
| 1970 | 41.4 | 312.9 | 5.9 | 0.8 | 6.8 | 15.4 | 30.5 | 56.2 | 110.9 | 165.5 | 307.2 | 589.2 |
| 1969 | 45.1 | 302.2 | 8.0 | 1.0 | 9.2 | 18.8 | 39.5 | 65.4 | 117.1 | 199.6 | 310.8 | 627.0 |
| 1968 | 43.0 | 304.6 | 6.5 | 1.1 | 6.1 | 15.1 | 30.5 | 53.0 | 111.6 | 192.3 | 343.0 | 862.9 |
| BLACK, FEMALE | | | | | | | | | | | | |
| 1978 | 25.5 | 335.0 | 4.1 | 0.8 | 2.4 | 5.8 | 11.1 | 20.5 | 38.5 | 80.9 | 145.5 | 488.9 |
| 1977 | 27.0 | 329.2 | 4.9 | * | 2.7 | 6.8 | 14.2 | 24.6 | 42.6 | 91.6 | 151.1 | 452.1 |
| 1976 | 26.3 | 325.2 | 2.8 | * | 3.2 | 7.5 | 12.1 | 23.6 | 40.6 | 87.9 | 159.0 | 507.9 |
| 1975 | 28.1 | 332.3 | 5.4 | * | 4.9 | 9.5 | 14.4 | 24.6 | 45.0 | 89.0 | 161.9 | 561.9 |
| 1974 | 28.2 | 316.7 | 5.8 | 1.2 | 5.4 | 10.8 | 20.1 | 27.1 | 47.6 | 90.3 | 149.8 | 515.4 |
| 1973 | 29.0 | 279.5 | 7.7 | 1.1 | 6.3 | 11.0 | 20.3 | 29.7 | 47.9 | 99.5 | 166.6 | 513.7 |
| 1972 | 27.6 | 249.8 | 5.3 | * | 4.0 | 7.7 | 16.0 | 29.7 | 53.0 | 113.0 | 160.7 | 505.7 |
| 1971 | 27.5 | 241.6 | 5.4 | * | 4.6 | 9.6 | 17.6 | 26.0 | 52.1 | 101.2 | 183.0 | 489.6 |
| 1970 | 29.6 | 245.6 | 6.1 | 1.0 | 5.4 | 9.3 | 18.5 | 31.6 | 58.9 | 104.7 | 206.2 | 487.5 |
| 1969 | 30.8 | 264.9 | 6.1 | 0.9 | 5.2 | 9.8 | 20.9 | 33.1 | 57.7 | 115.9 | 229.5 | 561.7 |
| 1968 | 29.6 | 244.3 | 4.4 | 0.9 | 3.3 | 10.1 | 19.4 | 32.9 | 55.4 | 108.5 | 225.2 | 651.8 |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| ALL OTHER DISEASES (RESIDUAL) | | | | | | | | | | | | |
| BOTH SEXES | | | | | | | | | | | | |
| 1978 | 61.9 | 56.2 | 7.3 | 3.5 | 6.2 | 10.0 | 21.5 | 47.7 | 104.5 | 233.9 | 477.8 | 876.0 |
| 1977 | 59.4 | 56.4 | 7.4 | 3.4 | 6.4 | 10.7 | 22.2 | 47.6 | 102.8 | 226.9 | 452.7 | 817.0 |
| 1976 | 57.6 | 57.8 | 7.1 | 3.5 | 7.1 | 11.6 | 23.2 | 47.6 | 103.8 | 220.1 | 432.8 | 771.7 |
| 1975 | 55.8 | 59.2 | 7.6 | 3.4 | 7.9 | 12.2 | 23.7 | 48.0 | 101.8 | 214.5 | 415.5 | 718.0 |
| 1974 | 55.5 | 62.3 | 7.8 | 4.1 | 8.2 | 12.7 | 25.1 | 49.8 | 102.4 | 214.1 | 408.8 | 716.5 |
| 1973 | 54.3 | 64.3 | 7.9 | 3.8 | 8.1 | 12.9 | 26.7 | 50.5 | 102.4 | 207.9 | 396.0 | 704.9 |
| 1972 | 52.9 | 62.5 | 8.1 | 3.9 | 9.4 | 13.7 | 27.6 | 50.2 | 100.6 | 201.5 | 379.6 | 660.1 |
| 1971 | 50.6 | 62.8 | 8.5 | 3.9 | 9.7 | 14.0 | 27.2 | 49.9 | 96.4 | 188.5 | 356.1 | 639.9 |
| 1970 | 49.8 | 69.8 | 8.3 | 3.9 | 8.4 | 13.3 | 27.0 | 49.7 | 97.1 | 187.3 | 358.0 | 568.5 |
| 1969 | 48.7 | 68.1 | 8.9 | 4.4 | 8.3 | 13.1 | 27.3 | 48.7 | 92.6 | 182.7 | 347.2 | 645.1 |
| 1968 | 48.8 | 67.7 | 8.8 | 4.0 | 8.2 | 12.9 | 26.8 | 48.9 | 94.3 | 188.9 | 348.2 | 654.1 |
| MALE | | | | | | | | | | | | |
| 1978 | 71.7 | 61.8 | 8.4 | 3.7 | 7.4 | 11.4 | 25.0 | 57.8 | 135.9 | 330.0 | 702.8 | 1,196.8 |
| 1977 | 69.0 | 63.6 | 8.2 | 3.7 | 7.7 | 12.2 | 26.4 | 58.1 | 132.6 | 318.8 | 663.1 | 1,122.8 |
| 1976 | 67.6 | 65.1 | 8.0 | 3.7 | 8.6 | 13.5 | 27.3 | 58.6 | 135.2 | 309.6 | 635.3 | 1,058.1 |
| 1975 | 65.2 | 67.5 | 8.0 | 3.4 | 9.6 | 14.7 | 28.3 | 58.4 | 131.7 | 300.7 | 597.2 | 969.4 |
| 1974 | 64.1 | 71.4 | 8.3 | 4.4 | 10.2 | 14.6 | 28.6 | 59.2 | 132.3 | 295.3 | 577.5 | 931.9 |
| 1973 | 62.3 | 72.7 | 8.1 | 3.9 | 10.0 | 14.8 | 30.8 | 60.3 | 132.1 | 283.2 | 546.7 | 904.1 |
| 1972 | 60.8 | 70.6 | 8.2 | 3.9 | 11.7 | 15.6 | 30.8 | 60.0 | 132.2 | 270.9 | 517.6 | 834.5 |
| 1971 | 58.1 | 69.0 | 9.3 | 4.2 | 12.0 | 15.3 | 30.1 | 59.9 | 124.3 | 253.7 | 481.1 | 811.2 |
| 1970 | 56.7 | 80.2 | 8.7 | 4.2 | 10.0 | 14.7 | 31.0 | 58.3 | 125.6 | 250.7 | 463.1 | 679.0 |
| 1969 | 55.1 | 74.5 | 9.4 | 4.7 | 9.5 | 14.0 | 30.9 | 58.0 | 118.5 | 242.2 | 444.0 | 772.3 |
| 1968 | 55.5 | 76.6 | 9.5 | 4.2 | 9.3 | 13.7 | 30.0 | 58.6 | 121.6 | 248.1 | 443.4 | 796.8 |

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| ALL OTHER DISEASES (RESIDUAL) | | | | | | | | | | | | |
| FEMALE | | | | | | | | | | | | |
| 1978 | 52.5 | 50.3 | 6.2 | 3.4 | 5.0 | 8.5 | 18.2 | 38.4 | 76.7 | 160.4 | 343.8 | 731.0 |
| 1977 | 50.2 | 48.8 | 6.6 | 3.2 | 5.1 | 9.2 | 18.2 | 37.8 | 76.5 | 156.6 | 326.8 | 676.6 |
| 1976 | 48.1 | 50.1 | 6.1 | 3.2 | 5.5 | 9.6 | 19.3 | 37.3 | 76.1 | 151.6 | 310.5 | 637.2 |
| 1975 | 46.9 | 50.6 | 7.3 | 3.3 | 6.3 | 9.8 | 19.4 | 38.4 | 75.2 | 148.4 | 304.8 | 596.3 |
| 1974 | 47.3 | 52.8 | 7.3 | 3.7 | 6.2 | 10.8 | 21.7 | 41.0 | 75.9 | 151.9 | 304.9 | 609.8 |
| 1973 | 46.7 | 55.4 | 7.7 | 3.7 | 6.3 | 11.1 | 22.8 | 41.3 | 76.1 | 150.1 | 301.9 | 604.6 |
| 1972 | 45.4 | 54.1 | 8.0 | 3.9 | 7.0 | 11.8 | 24.5 | 41.0 | 72.5 | 147.9 | 291.7 | 570.8 |
| 1971 | 43.5 | 56.4 | 7.7 | 3.7 | 7.3 | 12.8 | 24.3 | 40.6 | 71.4 | 138.1 | 275.0 | 550.6 |
| 1970 | 43.1 | 58.9 | 8.0 | 3.7 | 6.8 | 12.0 | 23.2 | 41.6 | 71.6 | 138.1 | 288.5 | 506.5 |
| 1969 | 42.6 | 61.4 | 8.4 | 4.0 | 7.0 | 12.3 | 23.8 | 40.1 | 69.4 | 136.1 | 282.2 | 576.2 |
| 1968 | 42.5 | 58.4 | 8.0 | 3.7 | 7.1 | 12.2 | 23.9 | 39.9 | 69.6 | 142.3 | 283.0 | 576.4 |
| WHITE, BOTH SEXES | | | | | | | | | | | | |
| 1978 | 61.9 | 48.4 | 7.0 | 3.5 | 5.6 | 7.6 | 16.4 | 41.7 | 98.1 | 230.6 | 480.8 | 883.3 |
| 1977 | 59.1 | 48.9 | 6.9 | 3.5 | 5.7 | 8.2 | 16.9 | 41.3 | 96.8 | 223.2 | 455.1 | 823.0 |
| 1976 | 57.1 | 49.3 | 6.8 | 3.4 | 6.2 | 8.8 | 17.5 | 41.1 | 97.9 | 216.3 | 436.2 | 781.4 |
| 1975 | 55.2 | 51.4 | 7.4 | 3.3 | 6.9 | 9.2 | 18.3 | 42.1 | 96.2 | 210.3 | 418.7 | 723.6 |
| 1974 | 54.7 | 54.2 | 7.5 | 4.1 | 7.2 | 9.7 | 19.4 | 43.3 | 96.6 | 209.3 | 412.1 | 727.9 |
| 1973 | 53.3 | 55.7 | 7.5 | 3.7 | 6.9 | 9.6 | 20.9 | 43.8 | 96.6 | 202.7 | 398.8 | 714.3 |
| 1972 | 51.6 | 54.9 | 7.9 | 4.0 | 7.9 | 9.9 | 21.7 | 43.6 | 94.4 | 195.6 | 382.6 | 671.9 |
| 1971 | 49.1 | 54.7 | 8.2 | 3.9 | 8.0 | 10.0 | 20.7 | 43.5 | 90.5 | 183.6 | 357.2 | 650.5 |
| 1970 | 48.6 | 61.9 | 7.8 | 3.9 | 7.2 | 10.0 | 21.0 | 43.5 | 91.7 | 182.8 | 359.9 | 583.4 |
| 1969 | 47.5 | 59.6 | 8.3 | 4.3 | 7.2 | 9.8 | 21.6 | 42.9 | 86.7 | 179.5 | 349.1 | 659.7 |
| 1968 | 47.5 | 58.9 | 8.3 | 3.9 | 7.1 | 9.5 | 20.9 | 43.1 | 88.6 | 184.4 | 350.5 | 670.2 |

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| ALL OTHER DISEASES (RESIDUAL) | | | | | | | | | | | | |
| WHITE, MALE | | | | | | | | | | | | |
| 1978 | 71.4 | 52.4 | 7.8 | 3.6 | 7.0 | 8.7 | 18.6 | 50.3 | 126.8 | 327.1 | 714.3 | 1,211.1 |
| 1977 | 68.4 | 55.0 | 7.7 | 3.7 | 7.1 | 9.3 | 19.6 | 49.7 | 124.6 | 316.3 | 672.1 | 1,132.9 |
| 1976 | 66.6 | 55.7 | 7.4 | 3.7 | 7.9 | 10.2 | 20.1 | 49.9 | 127.6 | 305.3 | 643.0 | 1,076.8 |
| 1975 | 64.2 | 59.2 | 7.8 | 3.4 | 8.6 | 10.7 | 21.3 | 50.7 | 124.9 | 296.9 | 604.0 | 979.4 |
| 1974 | 62.9 | 62.4 | 7.9 | 4.4 | 9.1 | 10.8 | 21.8 | 51.1 | 124.7 | 291.7 | 585.0 | 945.0 |
| 1973 | 60.8 | 63.9 | 7.5 | 3.7 | 8.7 | 10.7 | 23.6 | 51.9 | 124.8 | 278.8 | 551.8 | 919.4 |
| 1972 | 58.8 | 61.1 | 8.3 | 4.1 | 10.0 | 11.2 | 23.9 | 51.8 | 123.9 | 263.4 | 522.7 | 847.6 |
| 1971 | 56.0 | 59.6 | 8.6 | 4.1 | 10.0 | 10.7 | 22.8 | 52.0 | 117.3 | 248.4 | 482.7 | 831.1 |
| 1970 | 55.0 | 71.5 | 8.3 | 4.2 | 8.8 | 10.7 | 23.8 | 51.1 | 118.7 | 245.5 | 465.7 | 700.3 |
| 1969 | 53.2 | 64.4 | 8.4 | 4.7 | 8.3 | 10.3 | 24.0 | 51.0 | 110.5 | 238.7 | 445.7 | 791.3 |
| 1968 | 53.6 | 66.5 | 9.0 | 4.2 | 8.4 | 10.0 | 22.7 | 51.4 | 114.5 | 241.9 | 444.8 | 816.3 |
| WHITE, FEMALE | | | | | | | | | | | | |
| 1978 | 52.9 | 44.2 | 6.1 | 3.3 | 4.2 | 6.4 | 14.3 | 33.7 | 72.6 | 156.9 | 343.4 | 737.2 |
| 1977 | 50.3 | 42.4 | 6.1 | 3.2 | 4.4 | 7.2 | 14.2 | 33.3 | 72.2 | 152.2 | 327.0 | 682.7 |
| 1976 | 48.1 | 42.7 | 6.1 | 3.1 | 4.5 | 7.4 | 14.9 | 32.7 | 71.5 | 148.3 | 312.9 | 645.5 |
| 1975 | 46.7 | 43.1 | 7.0 | 3.3 | 5.3 | 7.7 | 15.4 | 34.1 | 70.6 | 144.3 | 307.2 | 602.4 |
| 1974 | 47.0 | 45.5 | 7.1 | 3.7 | 5.2 | 8.5 | 17.2 | 36.0 | 71.6 | 146.5 | 306.7 | 622.0 |
| 1973 | 46.1 | 47.0 | 7.5 | 3.7 | 5.1 | 8.6 | 18.3 | 36.2 | 71.4 | 144.6 | 304.1 | 613.4 |
| 1972 | 44.7 | 48.5 | 7.4 | 3.9 | 5.7 | 8.7 | 19.6 | 35.9 | 68.0 | 143.5 | 294.3 | 583.5 |
| 1971 | 42.5 | 49.6 | 7.7 | 3.6 | 5.9 | 9.4 | 18.7 | 35.6 | 66.4 | 133.7 | 276.9 | 558.1 |
| 1970 | 42.4 | 52.0 | 7.4 | 3.7 | 5.7 | 9.4 | 18.3 | 36.3 | 67.3 | 134.4 | 290.7 | 519.4 |
| 1969 | 42.0 | 54.6 | 8.2 | 3.9 | 6.1 | 9.4 | 19.4 | 35.3 | 65.3 | 133.5 | 284.8 | 590.8 |
| 1968 | 41.8 | 50.9 | 7.6 | 3.6 | 5.9 | 9.1 | 19.2 | 35.3 | 65.2 | 139.4 | 286.6 | 592.3 |

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| ALL OTHER DISEASES (RESIDUAL) | | | | | | | | | | | | |
| ALL OTHER, BOTH SEXES | | | | | | | | | | | | |
| 1978 | 61.7 | 93.7 | 9.0 | 3.8 | 9.6 | 25.4 | 56.3 | 93.9 | 162.0 | 265.0 | 447.1 | 792.3 |
| 1977 | 60.8 | 92.9 | 9.9 | 3.1 | 10.0 | 27.0 | 58.1 | 97.4 | 157.4 | 261.6 | 427.8 | 748.1 |
| 1976 | 60.7 | 99.1 | 8.3 | 3.8 | 12.1 | 30.0 | 62.0 | 100.3 | 158.3 | 256.9 | 397.6 | 656.9 |
| 1975 | 59.6 | 98.5 | 8.8 | 3.4 | 13.8 | 32.8 | 61.1 | 96.7 | 153.6 | 255.1 | 380.8 | 654.1 |
| 1974 | 60.6 | 102.9 | 9.3 | 4.1 | 14.5 | 33.8 | 64.7 | 103.5 | 156.5 | 261.2 | 372.9 | 583.7 |
| 1973 | 61.3 | 106.7 | 9.9 | 4.1 | 15.3 | 36.0 | 67.3 | 107.3 | 157.6 | 258.9 | 363.7 | 594.5 |
| 1972 | 61.8 | 101.3 | 9.4 | 3.6 | 18.7 | 40.5 | 70.2 | 106.6 | 160.0 | 259.9 | 344.5 | 523.6 |
| 1971 | 61.3 | 106.2 | 10.6 | 4.1 | 20.6 | 41.8 | 74.5 | 105.5 | 152.6 | 236.8 | 342.1 | 519.2 |
| 1970 | 58.1 | 111.4 | 10.8 | 4.0 | 15.8 | 36.3 | 70.9 | 104.4 | 148.9 | 231.1 | 333.5 | 415.4 |
| 1969 | 56.9 | 115.8 | 11.9 | 4.6 | 15.4 | 36.5 | 70.0 | 101.0 | 149.9 | 214.0 | 323.5 | 478.0 |
| 1968 | 57.9 | 115.4 | 11.2 | 4.5 | 15.2 | 37.0 | 72.2 | 101.6 | 149.4 | 233.5 | 317.8 | 473.5 |
| ALL OTHER, MALE | | | | | | | | | | | | |
| 1978 | 73.9 | 107.6 | 11.0 | 4.1 | 10.1 | 30.5 | 72.1 | 120.1 | 221.3 | 357.7 | 592.1 | 1,050.0 |
| 1977 | 73.1 | 106.0 | 10.5 | 3.3 | 11.0 | 32.9 | 75.9 | 128.7 | 208.8 | 341.7 | 578.2 | 1,019.6 |
| 1976 | 74.3 | 112.0 | 10.5 | 3.9 | 13.2 | 37.5 | 80.1 | 133.4 | 207.6 | 348.8 | 564.6 | 870.9 |
| 1975 | 72.4 | 109.4 | 8.8 | 3.3 | 15.9 | 44.0 | 80.3 | 125.3 | 196.9 | 337.0 | 530.2 | 868.5 |
| 1974 | 72.6 | 117.4 | 10.1 | 4.6 | 17.2 | 42.9 | 80.8 | 130.8 | 204.9 | 330.8 | 500.4 | 796.0 |
| 1973 | 73.1 | 116.5 | 11.0 | 4.6 | 17.7 | 46.1 | 85.7 | 135.6 | 202.3 | 325.0 | 490.6 | 747.9 |
| 1972 | 74.8 | 120.1 | 7.8 | 3.2 | 23.1 | 49.9 | 85.2 | 134.0 | 212.8 | 343.1 | 462.7 | 702.1 |
| 1971 | 73.0 | 119.9 | 13.1 | 4.4 | 25.4 | 50.1 | 88.7 | 132.1 | 192.0 | 304.4 | 464.7 | 614.9 |
| 1970 | 68.9 | 127.2 | 10.6 | 4.1 | 17.8 | 44.0 | 88.5 | 124.1 | 191.2 | 299.5 | 432.5 | 492.6 |
| 1969 | 68.3 | 132.2 | 14.4 | 5.0 | 17.9 | 41.7 | 87.6 | 122.8 | 196.3 | 276.1 | 423.9 | 593.2 |
| 1968 | 69.4 | 131.7 | 12.4 | 4.5 | 15.6 | 42.2 | 89.9 | 125.9 | 190.6 | 307.8 | 426.9 | 607.3 |
| ALL OTHER, FEMALE | | | | | | | | | | | | |
| 1978 | 50.6 | 79.4 | 7.0 | 3.5 | 9.2 | 21.1 | 43.1 | 72.3 | 112.8 | 193.2 | 349.1 | 662.4 |
| 1977 | 49.7 | 79.4 | 9.2 | 3.0 | 9.2 | 21.9 | 43.2 | 71.5 | 114.6 | 198.7 | 324.3 | 601.9 |
| 1976 | 48.4 | 86.2 | 6.1 | 3.8 | 11.1 | 23.5 | 46.9 | 72.6 | 117.1 | 184.2 | 283.5 | 536.7 |
| 1975 | 48.0 | 87.3 | 8.7 | 3.5 | 11.8 | 23.2 | 45.1 | 72.4 | 117.1 | 189.9 | 277.7 | 522.6 |
| 1974 | 49.6 | 88.5 | 8.6 | 3.7 | 11.9 | 25.9 | 51.2 | 80.3 | 115.3 | 205.4 | 284.9 | 458.8 |
| 1973 | 50.5 | 96.5 | 8.7 | 3.7 | 13.1 | 27.3 | 52.0 | 83.1 | 119.4 | 205.6 | 275.5 | 501.3 |
| 1972 | 49.9 | 82.1 | 11.1 | 3.9 | 14.6 | 32.3 | 57.7 | 83.1 | 114.6 | 192.7 | 260.5 | 413.2 |
| 1971 | 50.5 | 92.2 | 8.1 | 3.8 | 15.9 | 34.7 | 62.6 | 82.3 | 118.1 | 181.6 | 251.4 | 457.5 |
| 1970 | 48.2 | 95.4 | 11.0 | 3.9 | 13.9 | 29.6 | 56.3 | 87.1 | 111.4 | 174.8 | 258.2 | 361.0 |
| 1969 | 46.5 | 99.2 | 9.4 | 4.2 | 13.0 | 32.0 | 55.3 | 81.9 | 108.8 | 162.8 | 245.7 | 400.0 |
| 1968 | 47.3 | 98.5 | 10.1 | 4.6 | 14.7 | 32.6 | 57.3 | 80.3 | 112.5 | 171.9 | 234.3 | 383.6 |

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| ALL OTHER DISEASES (RESIDUAL) | | | | | | | | | | | | |
| BLACK, BOTH SEXES | | | | | | | | | | | | |
| 1978 | 66.9 | 98.8 | 9.5 | 4.2 | 10.4 | 28.7 | 62.4 | 104.3 | 175.3 | 277.1 | 453.9 | 798.0 |
| 1977 | 65.4 | 98.6 | 10.1 | 3.3 | 10.3 | 29.8 | 64.5 | 106.7 | 168.9 | 276.6 | 428.3 | 739.9 |
| 1976 | 65.4 | 105.9 | 8.8 | 4.1 | 13.1 | 33.2 | 68.9 | 109.9 | 168.5 | 270.7 | 398.8 | 665.4 |
| 1975 | 63.8 | 105.7 | 9.2 | 3.5 | 14.7 | 36.3 | 66.7 | 104.9 | 163.1 | 266.7 | 380.4 | 662.3 |
| 1974 | 64.5 | 110.1 | 9.9 | 4.2 | 15.3 | 37.5 | 70.2 | 111.4 | 166.0 | 274.6 | 373.2 | 586.8 |
| 1973 | 64.7 | 109.5 | 10.1 | 4.2 | 16.4 | 39.5 | 73.5 | 113.8 | 164.4 | 271.3 | 364.3 | 586.8 |
| 1972 | 65.0 | 107.3 | 9.3 | 3.4 | 20.1 | 44.4 | 76.2 | 112.8 | 166.5 | 272.2 | 344.1 | 510.9 |
| 1971 | 64.5 | 111.7 | 10.7 | 4.2 | 22.2 | 45.2 | 79.9 | 112.3 | 159.3 | 244.7 | 345.2 | 522.4 |
| 1970 | 61.7 | 117.2 | 11.3 | 4.1 | 16.8 | 40.4 | 77.7 | 111.3 | 156.2 | 241.4 | 343.7 | 422.5 |
| 1969 | 59.5 | 120.1 | 12.0 | 4.6 | 16.2 | 39.1 | 75.5 | 105.9 | 156.5 | 221.1 | 331.6 | 482.5 |
| 1968 | 60.2 | 118.2 | 11.4 | 4.6 | 15.9 | 39.4 | 77.1 | 105.8 | 154.9 | 240.8 | 322.7 | 475.0 |
| BLACK, MALE | | | | | | | | | | | | |
| 1978 | 80.1 | 112.9 | 11.8 | 4.5 | 10.7 | 34.3 | 80.5 | 134.3 | 241.2 | 381.7 | 597.9 | 1,043.1 |
| 1977 | 78.6 | 115.5 | 10.8 | 3.5 | 11.4 | 36.1 | 84.8 | 141.3 | 226.1 | 368.8 | 575.3 | 1,010.2 |
| 1976 | 80.0 | 119.7 | 10.9 | 4.0 | 14.1 | 41.8 | 90.2 | 146.8 | 222.3 | 372.8 | 567.5 | 891.5 |
| 1975 | 77.3 | 117.8 | 9.6 | 3.4 | 16.8 | 49.1 | 87.4 | 135.2 | 211.3 | 358.4 | 529.3 | 876.1 |
| 1974 | 77.2 | 124.3 | 10.8 | 4.7 | 18.1 | 47.6 | 87.7 | 140.9 | 220.8 | 351.3 | 503.8 | 811.6 |
| 1973 | 77.4 | 120.6 | 11.1 | 4.8 | 19.2 | 50.8 | 94.4 | 143.8 | 214.6 | 346.0 | 494.6 | 763.4 |
| 1972 | 78.8 | 128.6 | 8.3 | 2.9 | 25.0 | 55.4 | 93.6 | 141.2 | 224.9 | 367.2 | 451.8 | 705.0 |
| 1971 | 76.8 | 129.8 | 13.4 | 4.4 | 27.7 | 55.0 | 96.5 | 141.9 | 201.9 | 315.4 | 466.1 | 630.0 |
| 1970 | 73.3 | 135.4 | 11.0 | 4.1 | 19.2 | 49.8 | 97.7 | 133.9 | 203.8 | 314.8 | 451.9 | 502.2 |
| 1969 | 71.5 | 138.4 | 14.7 | 5.1 | 19.1 | 44.9 | 95.0 | 129.6 | 207.0 | 288.4 | 436.5 | 618.9 |
| 1968 | 72.4 | 136.6 | 12.9 | 4.6 | 16.1 | 45.7 | 96.7 | 133.2 | 199.6 | 320.8 | 444.8 | 620.0 |
| BLACK, FEMALE | | | | | | | | | | | | |
| 1978 | 55.0 | 84.4 | 7.2 | 3.8 | 10.1 | 23.8 | 47.5 | 79.5 | 121.5 | 199.2 | 357.8 | 671.7 |
| 1977 | 53.6 | 81.3 | 9.4 | 3.0 | 9.3 | 24.4 | 47.7 | 78.2 | 122.2 | 207.4 | 331.9 | 598.9 |
| 1976 | 52.2 | 92.0 | 6.6 | 4.1 | 12.1 | 25.7 | 51.3 | 79.2 | 123.9 | 193.3 | 285.7 | 546.1 |
| 1975 | 51.6 | 93.4 | 8.9 | 3.6 | 12.6 | 25.3 | 49.6 | 79.5 | 123.1 | 197.1 | 279.6 | 545.2 |
| 1974 | 53.0 | 95.5 | 9.0 | 3.6 | 12.6 | 28.8 | 55.5 | 86.7 | 120.6 | 216.3 | 284.8 | 462.8 |
| 1973 | 53.2 | 97.8 | 9.0 | 3.6 | 13.9 | 29.8 | 56.2 | 88.2 | 122.7 | 213.5 | 275.7 | 487.7 |
| 1972 | 52.5 | 85.2 | 10.4 | 3.8 | 15.4 | 35.0 | 61.8 | 88.4 | 117.6 | 198.7 | 267.1 | 400.0 |
| 1971 | 53.2 | 93.2 | 8.0 | 4.1 | 17.0 | 36.8 | 66.2 | 86.9 | 123.3 | 188.5 | 258.3 | 458.2 |
| 1970 | 51.1 | 98.9 | 11.5 | 4.1 | 14.5 | 32.5 | 61.3 | 91.7 | 115.8 | 183.2 | 264.4 | 371.2 |
| 1969 | 48.7 | 101.8 | 9.3 | 4.2 | 13.5 | 34.3 | 59.3 | 85.3 | 113.4 | 167.7 | 252.9 | 398.3 |
| 1968 | 49.1 | 99.6 | 9.9 | 4.6 | 15.6 | 34.1 | 60.7 | 81.9 | 116.3 | 176.0 | 232.0 | 392.9 |

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| E800-E949- ACCIDENTS | | | | | | | | | | | | |
| BOTH SEXES | | | | | | | | | | | | |
| 1978 | 47.5 | 37.9 | 28.2 | 16.9 | 63.1 | 44.6 | 38.9 | 39.6 | 45.5 | 60.5 | 120.8 | 291.5 |
| 1977 | 47.0 | 35.8 | 26.8 | 17.0 | 61.3 | 43.1 | 37.7 | 40.4 | 46.9 | 61.5 | 126.4 | 296.6 |
| 1976 | 46.3 | 40.2 | 27.5 | 16.7 | 58.9 | 42.5 | 37.0 | 39.9 | 46.9 | 62.0 | 127.5 | 318.0 |
| 1975 | 47.8 | 42.4 | 27.8 | 17.8 | 59.5 | 44.1 | 39.8 | 42.1 | 48.1 | 66.2 | 134.8 | 326.6 |
| 1974 | 49.0 | 47.4 | 28.9 | 18.2 | 60.9 | 44.7 | 40.0 | 43.5 | 50.6 | 68.7 | 139.8 | 368.3 |
| 1973 | 54.8 | 54.1 | 31.7 | 20.6 | 68.2 | 50.1 | 45.4 | 48.8 | 58.9 | 77.3 | 156.0 | 410.0 |
| 1972 | 55.2 | 54.0 | 31.6 | 20.6 | 67.6 | 49.4 | 45.1 | 50.8 | 60.5 | 80.8 | 165.5 | 416.0 |
| 1971 | 54.8 | 62.5 | 31.1 | 20.1 | 66.1 | 49.9 | 46.6 | 49.9 | 60.7 | 82.7 | 164.3 | 416.5 |
| 1970 | 56.4 | 65.8 | 31.5 | 20.1 | 68.7 | 51.6 | 48.2 | 53.5 | 63.2 | 85.6 | 174.7 | 392.6 |
| 1969 | 57.8 | 71.1 | 32.6 | 20.0 | 71.7 | 53.1 | 49.2 | 53.5 | 64.6 | 86.5 | 179.9 | 482.2 |
| 1968 | 57.6 | 74.9 | 32.6 | 20.5 | 69.2 | 52.5 | 47.6 | 53.6 | 64.7 | 90.0 | 188.8 | 513.7 |
| MALE | | | | | | | | | | | | |
| 1978 | 68.4 | 40.5 | 32.9 | 22.7 | 98.2 | 72.1 | 61.2 | 60.7 | 67.9 | 86.8 | 165.8 | 375.8 |
| 1977 | 67.3 | 40.0 | 31.3 | 22.8 | 95.5 | 70.3 | 58.8 | 61.2 | 69.9 | 88.4 | 170.0 | 370.5 |
| 1976 | 66.4 | 44.6 | 32.7 | 22.5 | 92.4 | 69.3 | 58.1 | 60.6 | 70.0 | 89.2 | 171.2 | 398.2 |
| 1975 | 69.0 | 44.6 | 32.3 | 24.6 | 95.2 | 72.0 | 62.6 | 64.6 | 72.7 | 96.0 | 181.8 | 389.1 |
| 1974 | 70.5 | 52.5 | 34.4 | 24.8 | 97.8 | 73.1 | 63.2 | 66.9 | 76.1 | 99.1 | 180.6 | 429.4 |
| 1973 | 78.2 | 61.0 | 37.3 | 27.9 | 108.4 | 81.5 | 71.0 | 74.2 | 87.9 | 109.1 | 202.0 | 480.9 |
| 1972 | 78.2 | 62.2 | 38.2 | 27.3 | 106.8 | 80.3 | 69.6 | 76.5 | 90.2 | 114.0 | 217.0 | 489.7 |
| 1971 | 77.6 | 71.4 | 36.3 | 26.9 | 105.2 | 80.3 | 72.5 | 75.5 | 90.5 | 117.6 | 207.8 | 480.0 |
| 1970 | 80.6 | 70.4 | 36.8 | 27.4 | 110.5 | 84.4 | 76.1 | 81.2 | 95.9 | 123.1 | 220.6 | 426.6 |
| 1969 | 82.4 | 77.8 | 37.5 | 27.2 | 116.2 | 87.9 | 77.2 | 81.5 | 97.2 | 122.6 | 223.7 | 528.1 |
| 1968 | 81.8 | 80.2 | 37.3 | 28.0 | 112.8 | 87.0 | 74.7 | 82.6 | 97.6 | 127.9 | 232.0 | 533.6 |

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| E800-E949- ACCIDENTS | | | | | | | | | | | | |
| FEMALE | | | | | | | | | | | | |
| 1978 | 27.8 | 35.2 | 23.3 | 10.9 | 27.4 | 17.6 | 17.5 | 19.9 | 25.7 | 40.4 | 94.1 | 253.4 |
| 1977 | 27.7 | 31.3 | 22.2 | 11.0 | 26.6 | 16.5 | 17.5 | 20.9 | 26.5 | 41.0 | 100.3 | 262.6 |
| 1976 | 27.3 | 35.7 | 22.1 | 10.6 | 24.9 | 16.3 | 16.8 | 20.7 | 26.5 | 41.1 | 101.1 | 280.3 |
| 1975 | 27.7 | 40.2 | 23.2 | 10.8 | 23.4 | 16.9 | 18.0 | 21.1 | 26.4 | 43.5 | 106.2 | 296.4 |
| 1974 | 28.7 | 42.1 | 23.2 | 11.3 | 23.6 | 16.9 | 17.9 | 21.8 | 28.1 | 45.4 | 114.7 | 338.1 |
| 1973 | 32.6 | 46.9 | 25.8 | 13.0 | 27.8 | 19.4 | 21.0 | 25.3 | 33.1 | 53.0 | 127.3 | 374.4 |
| 1972 | 33.3 | 45.4 | 24.7 | 13.7 | 28.3 | 19.4 | 21.8 | 27.0 | 34.0 | 55.3 | 132.6 | 378.2 |
| 1971 | 33.2 | 53.1 | 25.8 | 13.0 | 27.2 | 20.5 | 22.0 | 26.1 | 34.0 | 55.7 | 136.2 | 383.4 |
| 1970 | 33.4 | 61.0 | 25.9 | 12.5 | 27.6 | 19.9 | 21.8 | 27.6 | 33.8 | 56.4 | 144.4 | 373.6 |
| 1969 | 34.5 | 64.0 | 27.5 | 12.6 | 28.4 | 19.8 | 22.8 | 27.3 | 35.4 | 58.3 | 150.5 | 457.4 |
| 1968 | 34.7 | 69.3 | 27.8 | 12.8 | 26.9 | 19.5 | 22.1 | 26.5 | 35.0 | 60.1 | 159.3 | 503.4 |
| WHITE, BOTH SEXES | | | | | | | | | | | | |
| 1978 | 47.1 | 31.9 | 25.5 | 16.2 | 66.3 | 43.6 | 36.2 | 36.7 | 42.6 | 58.0 | 120.9 | 298.3 |
| 1977 | 46.4 | 30.9 | 24.7 | 16.1 | 63.9 | 41.4 | 35.2 | 37.4 | 43.9 | 59.4 | 127.1 | 303.3 |
| 1976 | 45.7 | 34.7 | 25.5 | 16.0 | 61.2 | 40.4 | 34.2 | 36.9 | 44.4 | 59.7 | 128.6 | 327.3 |
| 1975 | 46.9 | 35.3 | 25.3 | 16.8 | 61.1 | 41.9 | 36.6 | 38.9 | 45.2 | 63.9 | 135.9 | 336.4 |
| 1974 | 48.1 | 40.5 | 26.8 | 17.2 | 62.3 | 42.4 | 36.8 | 40.2 | 47.6 | 65.8 | 142.4 | 378.2 |
| 1973 | 53.4 | 44.2 | 28.4 | 19.5 | 69.0 | 47.3 | 41.6 | 45.1 | 56.0 | 74.5 | 157.9 | 422.8 |
| 1972 | 53.7 | 46.5 | 28.8 | 19.7 | 68.0 | 46.4 | 40.9 | 47.2 | 57.4 | 77.8 | 167.3 | 425.2 |
| 1971 | 53.1 | 51.9 | 28.1 | 18.8 | 65.6 | 46.5 | 42.9 | 46.3 | 57.8 | 79.4 | 165.6 | 429.0 |
| 1970 | 54.6 | 54.0 | 28.5 | 18.8 | 68.2 | 47.9 | 44.1 | 49.5 | 60.0 | 83.0 | 175.9 | 409.9 |
| 1969 | 56.0 | 58.9 | 29.0 | 18.5 | 71.1 | 49.8 | 45.4 | 49.8 | 61.3 | 84.0 | 181.0 | 500.2 |
| 1968 | 55.7 | 61.3 | 28.4 | 19.0 | 68.9 | 48.7 | 43.4 | 49.8 | 61.1 | 87.0 | 190.5 | 533.6 |

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| E800-E949- ACCIDENTS | | | | | | | | | | | | |
| WHITE, MALE | | | | | | | | | | | | |
| 1978 | 67.1 | 34.5 | 30.3 | 21.6 | 102.7 | 69.9 | 56.1 | 55.5 | 62.9 | 82.2 | 164.0 | 383.5 |
| 1977 | 65.7 | 35.0 | 28.8 | 21.3 | 99.1 | 66.8 | 53.9 | 55.8 | 64.8 | 84.2 | 170.7 | 380.2 |
| 1976 | 64.7 | 38.9 | 30.4 | 21.3 | 95.5 | 65.5 | 53.0 | 55.0 | 65.3 | 84.9 | 171.5 | 411.4 |
| 1975 | 66.9 | 37.2 | 29.7 | 22.9 | 97.5 | 67.7 | 56.9 | 59.1 | 67.5 | 91.6 | 182.9 | 400.9 |
| 1974 | 68.5 | 44.9 | 32.4 | 23.4 | 99.6 | 69.3 | 57.7 | 61.1 | 70.9 | 93.8 | 183.0 | 441.9 |
| 1973 | 75.4 | 51.5 | 33.9 | 26.1 | 109.3 | 76.5 | 64.3 | 67.6 | 81.9 | 104.1 | 203.1 | 495.9 |
| 1972 | 75.2 | 54.6 | 35.3 | 26.1 | 106.5 | 74.8 | 62.0 | 69.9 | 84.2 | 108.7 | 218.4 | 503.6 |
| 1971 | 74.4 | 60.5 | 32.9 | 25.0 | 103.9 | 74.3 | 66.2 | 69.2 | 85.3 | 112.2 | 208.3 | 495.0 |
| 1970 | 77.2 | 57.2 | 33.8 | 25.8 | 108.9 | 77.9 | 68.8 | 74.0 | 90.0 | 119.1 | 222.0 | 448.9 |
| 1969 | 79.0 | 66.8 | 33.8 | 25.0 | 114.8 | 81.8 | 70.2 | 75.1 | 90.9 | 118.4 | 224.6 | 552.5 |
| 1968 | 78.0 | 65.6 | 32.7 | 25.9 | 111.4 | 80.0 | 67.4 | 75.6 | 91.4 | 122.4 | 233.8 | 556.4 |
| WHITE, FEMALE | | | | | | | | | | | | |
| 1978 | 28.0 | 29.2 | 20.5 | 10.5 | 29.0 | 17.3 | 16.6 | 18.9 | 24.5 | 39.5 | 95.6 | 260.4 |
| 1977 | 28.0 | 26.6 | 20.3 | 10.7 | 27.8 | 16.0 | 16.9 | 20.0 | 25.3 | 40.4 | 101.4 | 268.4 |
| 1976 | 27.5 | 30.4 | 20.3 | 10.3 | 26.0 | 15.4 | 15.9 | 19.8 | 25.8 | 40.5 | 103.0 | 288.7 |
| 1975 | 27.7 | 33.3 | 20.8 | 10.3 | 23.9 | 16.3 | 16.8 | 20.0 | 25.4 | 42.7 | 107.7 | 306.0 |
| 1974 | 28.7 | 35.9 | 20.8 | 10.7 | 24.3 | 15.8 | 16.5 | 20.6 | 27.0 | 44.4 | 117.5 | 347.2 |
| 1973 | 32.5 | 36.5 | 22.7 | 12.5 | 28.0 | 18.2 | 19.5 | 23.9 | 32.8 | 51.9 | 129.9 | 387.0 |
| 1972 | 33.2 | 38.1 | 22.1 | 12.9 | 28.9 | 18.4 | 20.3 | 25.9 | 33.4 | 54.2 | 135.0 | 385.8 |
| 1971 | 32.9 | 42.9 | 23.1 | 12.3 | 27.2 | 19.2 | 20.4 | 24.8 | 33.1 | 54.1 | 138.2 | 395.2 |
| 1970 | 33.1 | 50.8 | 23.0 | 11.5 | 28.0 | 18.6 | 20.2 | 26.6 | 33.1 | 55.1 | 145.7 | 388.6 |
| 1969 | 34.1 | 50.6 | 23.9 | 11.7 | 28.2 | 18.6 | 21.6 | 26.1 | 34.5 | 57.3 | 152.0 | 473.1 |
| 1968 | 34.3 | 56.7 | 23.9 | 11.8 | 27.3 | 18.2 | 20.4 | 25.5 | 33.6 | 59.2 | 161.2 | 522.1 |

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| E800-E949- ACCIDENTS | | | | | | | | | | | | |
| ALL OTHER, BOTH SEXES | | | | | | | | | | | | |
| 1978 | 50.3 | 66.9 | 41.1 | 20.3 | 45.3 | 51.1 | 57.8 | 61.9 | 71.7 | 84.1 | 119.7 | 213.1 |
| 1977 | 50.7 | 59.5 | 37.2 | 21.5 | 46.8 | 54.8 | 54.6 | 63.6 | 74.2 | 82.1 | 118.5 | 220.0 |
| 1976 | 50.5 | 67.0 | 37.3 | 20.6 | 46.0 | 56.7 | 56.2 | 64.8 | 70.4 | 83.6 | 116.3 | 209.8 |
| 1975 | 54.1 | 78.1 | 40.0 | 23.3 | 50.0 | 59.2 | 61.8 | 67.9 | 75.1 | 89.4 | 122.9 | 214.4 |
| 1974 | 55.4 | 81.7 | 39.7 | 23.3 | 52.3 | 60.2 | 62.3 | 71.2 | 78.6 | 97.1 | 112.3 | 253.3 |
| 1973 | 64.0 | 102.7 | 48.3 | 26.8 | 63.6 | 70.0 | 72.6 | 80.7 | 86.5 | 105.0 | 135.0 | 261.4 |
| 1972 | 65.0 | 92.1 | 46.1 | 25.9 | 65.7 | 70.7 | 75.6 | 82.4 | 89.9 | 110.5 | 144.5 | 308.9 |
| 1971 | 66.8 | 119.0 | 47.3 | 27.5 | 69.3 | 73.8 | 73.7 | 81.4 | 88.7 | 115.9 | 149.3 | 274.2 |
| 1970 | 68.8 | 128.7 | 47.1 | 27.4 | 71.7 | 76.9 | 78.9 | 88.1 | 93.2 | 111.1 | 160.0 | 214.8 |
| 1969 | 70.5 | 139.7 | 51.5 | 29.0 | 76.1 | 76.8 | 78.2 | 86.0 | 97.2 | 111.6 | 165.7 | 280.7 |
| 1968 | 71.6 | 148.3 | 55.1 | 29.2 | 71.3 | 79.9 | 80.5 | 88.4 | 100.0 | 120.5 | 166.7 | 291.2 |
| ALL OTHER, MALE | | | | | | | | | | | | |
| 1978 | 76.9 | 69.9 | 45.4 | 27.9 | 72.6 | 87.8 | 98.9 | 103.9 | 114.7 | 129.8 | 182.3 | 296.6 |
| 1977 | 77.9 | 64.8 | 43.3 | 30.6 | 74.5 | 95.6 | 94.6 | 106.8 | 118.5 | 127.8 | 162.8 | 271.4 |
| 1976 | 77.6 | 72.9 | 43.6 | 28.8 | 74.2 | 97.0 | 96.4 | 108.4 | 114.9 | 129.6 | 168.5 | 265.5 |
| 1975 | 83.1 | 82.0 | 45.2 | 33.4 | 81.3 | 103.3 | 104.8 | 112.9 | 122.1 | 138.3 | 170.7 | 270.4 |
| 1974 | 84.4 | 90.7 | 44.4 | 32.4 | 86.2 | 101.9 | 104.8 | 117.8 | 126.0 | 149.6 | 155.9 | 300.0 |
| 1973 | 97.7 | 107.9 | 55.0 | 37.8 | 102.6 | 119.6 | 122.9 | 132.8 | 145.4 | 156.3 | 189.7 | 327.1 |
| 1972 | 99.3 | 101.8 | 54.0 | 33.9 | 108.4 | 122.7 | 128.4 | 135.7 | 148.2 | 165.0 | 201.8 | 348.9 |
| 1971 | 100.8 | 130.7 | 54.8 | 37.8 | 113.5 | 126.3 | 122.5 | 132.9 | 141.3 | 169.1 | 202.9 | 331.9 |
| 1970 | 105.0 | 142.4 | 53.1 | 36.6 | 121.0 | 133.4 | 134.5 | 146.9 | 152.0 | 160.9 | 204.1 | 231.0 |
| 1969 | 106.9 | 141.5 | 57.1 | 40.1 | 126.0 | 135.0 | 134.6 | 140.9 | 157.8 | 162.9 | 213.3 | 297.7 |
| 1968 | 109.2 | 160.0 | 62.4 | 40.4 | 122.3 | 141.3 | 135.2 | 148.8 | 157.5 | 181.6 | 210.8 | 312.2 |

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| E800-E949- ACCIDENTS | | | | | | | | | | | | |
| ALL OTHER, FEMALE | | | | | | | | | | | | |
| 1978 | 26.3 | 63.8 | 36.7 | 12.5 | 18.9 | 19.4 | 23.3 | 27.2 | 35.9 | 48.7 | 77.1 | 170.6 |
| 1977 | 26.1 | 54.2 | 30.9 | 12.2 | 20.0 | 19.6 | 21.0 | 27.7 | 37.3 | 46.2 | 88.0 | 192.3 |
| 1976 | 25.9 | 61.3 | 30.8 | 12.3 | 18.8 | 22.0 | 22.7 | 28.2 | 33.1 | 47.1 | 80.6 | 178.6 |
| 1975 | 27.6 | 74.1 | 34.6 | 13.0 | 20.1 | 21.2 | 25.8 | 29.8 | 35.3 | 50.5 | 89.8 | 179.6 |
| 1974 | 29.0 | 72.7 | 34.9 | 14.1 | 20.0 | 24.4 | 26.8 | 31.6 | 38.4 | 55.2 | 82.2 | 225.9 |
| 1973 | 33.3 | 97.3 | 41.5 | 15.7 | 26.5 | 27.3 | 30.5 | 36.1 | 36.1 | 63.6 | 96.9 | 221.5 |
| 1972 | 33.6 | 82.1 | 38.0 | 17.9 | 25.1 | 26.1 | 31.3 | 36.4 | 39.6 | 66.4 | 103.7 | 284.2 |
| 1971 | 35.6 | 107.1 | 39.8 | 17.2 | 27.6 | 29.0 | 33.2 | 36.7 | 42.9 | 72.3 | 109.6 | 237.0 |
| 1970 | 35.5 | 114.8 | 41.1 | 18.1 | 25.5 | 28.6 | 32.8 | 36.6 | 41.2 | 70.0 | 126.5 | 203.4 |
| 1969 | 37.2 | 137.9 | 45.9 | 17.9 | 29.8 | 27.4 | 31.3 | 37.8 | 43.5 | 69.3 | 128.7 | 269.2 |
| 1968 | 37.0 | 135.9 | 47.8 | 18.0 | 23.8 | 28.1 | 34.6 | 35.0 | 48.4 | 69.9 | 133.1 | 277.0 |
| BLACK, BOTH SEXES | | | | | | | | | | | | |
| 1978 | 50.6 | 71.1 | 41.8 | 20.9 | 42.3 | 51.2 | 60.4 | 64.6 | 73.7 | 84.1 | 115.1 | 204.0 |
| 1977 | 51.5 | 64.3 | 38.3 | 22.0 | 44.1 | 56.5 | 57.0 | 66.4 | 76.8 | 84.9 | 115.1 | 218.2 |
| 1976 | 50.8 | 69.9 | 38.9 | 21.0 | 42.9 | 57.8 | 58.0 | 67.1 | 72.0 | 85.6 | 113.4 | 213.2 |
| 1975 | 54.6 | 83.3 | 41.7 | 23.4 | 47.5 | 61.0 | 65.2 | 70.8 | 75.9 | 90.0 | 118.2 | 206.9 |
| 1974 | 56.1 | 86.8 | 40.8 | 23.9 | 50.7 | 62.2 | 64.5 | 72.1 | 80.2 | 98.9 | 110.6 | 253.7 |
| 1973 | 64.0 | 108.5 | 49.6 | 27.7 | 60.8 | 69.2 | 73.9 | 83.0 | 87.3 | 104.8 | 129.5 | 260.5 |
| 1972 | 65.3 | 99.0 | 46.2 | 26.3 | 63.5 | 70.5 | 78.9 | 84.5 | 90.2 | 111.8 | 141.6 | 309.1 |
| 1971 | 67.4 | 124.8 | 48.6 | 28.2 | 68.0 | 75.7 | 76.6 | 82.5 | 89.2 | 116.1 | 147.9 | 265.4 |
| 1970 | 70.0 | 134.6 | 48.1 | 28.2 | 70.9 | 80.1 | 82.0 | 90.8 | 94.9 | 110.9 | 162.8 | 219.8 |
| 1969 | 70.4 | 147.9 | 52.2 | 29.5 | 74.6 | 76.1 | 79.6 | 86.4 | 96.9 | 109.4 | 165.4 | 279.4 |
| 1968 | 72.0 | 156.7 | 54.7 | 30.0 | 70.7 | 79.7 | 83.2 | 89.5 | 99.5 | 120.4 | 167.7 | 285.9 |

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| E800-E949- ACCIDENTS | | | | | | | | | | | | |
| BLACK, MALE | | | | | | | | | | | | |
| 1978 | 77.4 | 73.0 | 45.6 | 29.0 | 68.6 | 89.0 | 105.3 | 109.3 | 118.9 | 130.5 | 171.5 | 282.4 |
| 1977 | 79.5 | 69.4 | 45.0 | 31.5 | 71.3 | 99.8 | 100.7 | 112.4 | 123.9 | 132.8 | 158.7 | 259.2 |
| 1976 | 78.5 | 76.4 | 45.6 | 29.6 | 70.4 | 99.8 | 101.6 | 113.2 | 118.0 | 133.8 | 162.0 | 266.0 |
| 1975 | 84.2 | 88.7 | 47.0 | 33.7 | 78.1 | 107.8 | 111.4 | 119.1 | 124.2 | 141.8 | 162.2 | 250.0 |
| 1974 | 85.9 | 98.2 | 46.2 | 33.3 | 84.1 | 106.8 | 110.2 | 120.8 | 129.3 | 154.6 | 152.2 | 297.7 |
| 1973 | 97.9 | 115.9 | 56.8 | 39.4 | 98.9 | 119.4 | 126.6 | 137.5 | 147.8 | 157.9 | 177.9 | 336.6 |
| 1972 | 100.1 | 111.2 | 53.9 | 34.4 | 106.1 | 122.8 | 135.8 | 140.9 | 149.7 | 169.3 | 198.0 | 350.0 |
| 1971 | 102.1 | 136.9 | 55.6 | 38.6 | 112.2 | 132.0 | 127.7 | 134.5 | 144.0 | 170.7 | 201.1 | 325.0 |
| 1970 | 107.6 | 150.5 | 53.7 | 37.9 | 121.4 | 141.8 | 141.4 | 152.9 | 156.1 | 161.0 | 204.0 | 245.7 |
| 1969 | 106.8 | 148.9 | 57.1 | 40.9 | 124.6 | 135.3 | 136.9 | 142.7 | 159.4 | 158.9 | 207.8 | 297.3 |
| 1968 | 110.4 | 167.6 | 62.0 | 41.6 | 122.2 | 142.5 | 139.9 | 152.7 | 158.9 | 183.0 | 214.5 | 314.3 |
| BLACK, FEMALE | | | | | | | | | | | | |
| 1978 | 26.4 | 69.1 | 38.0 | 12.7 | 17.2 | 18.5 | 23.1 | 27.8 | 36.8 | 49.7 | 77.4 | 163.6 |
| 1977 | 26.2 | 59.2 | 31.4 | 12.3 | 18.1 | 19.2 | 20.6 | 28.4 | 38.3 | 48.9 | 86.4 | 196.8 |
| 1976 | 25.8 | 63.3 | 32.2 | 12.3 | 16.6 | 21.7 | 21.8 | 28.7 | 34.0 | 49.1 | 80.8 | 185.4 |
| 1975 | 27.8 | 77.9 | 36.3 | 13.0 | 18.5 | 20.7 | 26.8 | 30.3 | 35.9 | 50.8 | 88.4 | 183.3 |
| 1974 | 29.1 | 75.1 | 35.3 | 14.4 | 19.0 | 24.0 | 26.4 | 31.1 | 39.4 | 56.4 | 82.5 | 229.5 |
| 1973 | 33.2 | 100.4 | 42.4 | 16.0 | 24.8 | 26.3 | 30.1 | 36.8 | 37.0 | 63.8 | 96.6 | 217.8 |
| 1972 | 33.6 | 86.1 | 38.4 | 18.1 | 23.1 | 26.0 | 31.5 | 36.5 | 40.4 | 67.1 | 101.4 | 285.7 |
| 1971 | 36.0 | 112.4 | 41.5 | 17.7 | 26.4 | 28.0 | 34.3 | 37.9 | 43.0 | 73.0 | 109.7 | 229.9 |
| 1970 | 35.9 | 118.7 | 42.4 | 18.5 | 24.0 | 28.1 | 32.8 | 37.1 | 42.7 | 71.3 | 132.7 | 203.1 |
| 1969 | 37.4 | 146.9 | 47.3 | 18.2 | 28.3 | 26.5 | 32.0 | 37.7 | 43.5 | 70.1 | 133.5 | 268.3 |
| 1968 | 37.2 | 145.5 | 47.3 | 18.4 | 23.1 | 27.2 | 35.7 | 34.6 | 48.2 | 69.8 | 132.9 | 273.2 |

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| MOTOR VEHICLE ACCIDENTS (E810-E823) | | | | | | | | | | | | |
| BOTH SEXES | | | | | | | | | | | | |
| 1978 | 23.6 | 7.9 | 10.4 | 8.6 | 45.4 | 27.7 | 20.2 | 18.0 | 18.4 | 21.5 | 29.1 | 25.3 |
| 1977 | 22.5 | 7.7 | 9.9 | 8.5 | 43.3 | 25.4 | 18.9 | 17.8 | 18.5 | 20.9 | 29.7 | 26.6 |
| 1976 | 21.6 | 7.7 | 10.3 | 8.4 | 40.3 | 24.2 | 18.3 | 17.4 | 17.9 | 21.6 | 30.6 | 27.0 |
| 1975 | 21.3 | 8.1 | 10.2 | 8.6 | 38.7 | 24.5 | 18.8 | 17.2 | 17.8 | 21.9 | 31.1 | 24.9 |
| 1974 | 21.8 | 7.3 | 9.9 | 8.6 | 40.0 | 25.0 | 19.0 | 18.3 | 19.3 | 22.6 | 30.4 | 27.4 |
| 1973 | 26.3 | 10.4 | 12.2 | 10.5 | 46.3 | 29.9 | 23.7 | 22.9 | 24.6 | 29.4 | 39.6 | 35.2 |
| 1972 | 26.9 | 9.1 | 11.6 | 10.7 | 47.1 | 30.0 | 24.2 | 24.7 | 26.0 | 32.0 | 43.9 | 35.7 |
| 1971 | 26.3 | 9.9 | 11.2 | 10.5 | 45.7 | 30.5 | 23.9 | 23.6 | 26.0 | 32.4 | 41.9 | 36.6 |
| 1970 | 26.9 | 9.8 | 11.5 | 10.2 | 47.2 | 30.9 | 24.9 | 25.5 | 27.9 | 32.8 | 43.5 | 34.2 |
| 1969 | 27.7 | 10.8 | 12.2 | 9.9 | 50.7 | 32.3 | 25.7 | 25.4 | 28.1 | 34.2 | 44.0 | 34.0 |
| 1968 | 27.5 | 9.5 | 11.5 | 10.1 | 49.8 | 32.6 | 24.9 | 25.8 | 28.5 | 35.0 | 47.4 | 39.7 |
| MALE | | | | | | | | | | | | |
| 1978 | 35.3 | 7.2 | 11.4 | 11.0 | 68.9 | 44.0 | 31.0 | 26.9 | 26.7 | 30.7 | 47.0 | 53.4 |
| 1977 | 33.5 | 8.0 | 10.8 | 10.6 | 65.5 | 40.4 | 28.5 | 26.6 | 27.2 | 29.9 | 47.3 | 50.4 |
| 1976 | 32.2 | 7.0 | 11.8 | 10.5 | 61.6 | 38.4 | 28.0 | 26.1 | 26.1 | 30.6 | 48.7 | 53.6 |
| 1975 | 32.0 | 8.1 | 11.7 | 11.1 | 60.0 | 39.1 | 29.3 | 25.8 | 26.1 | 32.1 | 50.0 | 49.7 |
| 1974 | 32.9 | 6.5 | 10.7 | 10.9 | 62.6 | 40.1 | 29.6 | 28.0 | 28.5 | 33.4 | 48.0 | 56.5 |
| 1973 | 38.8 | 10.7 | 14.1 | 13.3 | 71.2 | 47.2 | 35.8 | 33.8 | 35.4 | 41.7 | 62.8 | 69.3 |
| 1972 | 39.4 | 9.4 | 13.0 | 13.1 | 72.2 | 47.9 | 36.0 | 35.7 | 37.4 | 44.1 | 72.9 | 71.6 |
| 1971 | 38.6 | 10.8 | 12.0 | 13.3 | 70.4 | 48.0 | 35.8 | 34.6 | 37.6 | 45.8 | 65.7 | 71.2 |
| 1970 | 39.7 | 9.3 | 12.9 | 13.1 | 73.2 | 49.4 | 37.7 | 37.2 | 40.9 | 47.3 | 68.2 | 63.1 |
| 1969 | 41.0 | 10.8 | 13.6 | 12.5 | 79.8 | 52.2 | 38.7 | 37.4 | 40.4 | 49.0 | 68.9 | 64.1 |
| 1968 | 41.0 | 9.6 | 12.7 | 13.0 | 78.8 | 52.7 | 37.7 | 38.6 | 41.5 | 50.0 | 75.3 | 74.1 |

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| MOTOR VEHICLE ACCIDENTS (E810-E823) | | | | | | | | | | | | |
| FEMALE | | | | | | | | | | | | |
| 1978 | 12.5 | 8.7 | 9.3 | 6.2 | 21.6 | 11.8 | 9.8 | 9.7 | 11.1 | 14.4 | 18.5 | 12.6 |
| 1977 | 12.1 | 7.4 | 9.0 | 6.3 | 20.7 | 10.8 | 9.6 | 9.5 | 10.9 | 14.1 | 19.3 | 15.7 |
| 1976 | 11.6 | 8.5 | 8.8 | 6.2 | 18.8 | 10.3 | 9.1 | 9.4 | 10.6 | 14.8 | 19.7 | 14.4 |
| 1975 | 11.1 | 8.1 | 8.6 | 6.0 | 17.0 | 10.3 | 8.9 | 9.2 | 10.5 | 14.1 | 19.5 | 13.0 |
| 1974 | 11.2 | 8.2 | 9.0 | 6.2 | 17.2 | 10.2 | 9.0 | 9.3 | 11.1 | 14.4 | 19.6 | 13.1 |
| 1973 | 14.4 | 10.1 | 10.2 | 7.6 | 21.3 | 13.1 | 12.1 | 12.8 | 14.9 | 19.9 | 25.1 | 18.1 |
| 1972 | 15.0 | 8.8 | 10.1 | 8.1 | 21.9 | 12.7 | 13.0 | 14.4 | 15.9 | 22.7 | 25.5 | 17.3 |
| 1971 | 14.6 | 9.0 | 10.3 | 7.7 | 21.1 | 13.5 | 12.5 | 13.3 | 15.5 | 22.0 | 26.5 | 18.5 |
| 1970 | 14.7 | 10.4 | 10.0 | 7.2 | 21.6 | 13.0 | 12.9 | 14.5 | 16.2 | 21.6 | 27.2 | 18.0 |
| 1969 | 15.1 | 10.9 | 10.8 | 7.2 | 22.4 | 13.3 | 13.4 | 14.2 | 17.0 | 22.7 | 27.3 | 17.8 |
| 1968 | 14.7 | 9.3 | 10.2 | 7.0 | 21.5 | 13.4 | 12.9 | 13.9 | 16.8 | 23.2 | 28.2 | 20.9 |
| WHITE, BOTH SEXES | | | | | | | | | | | | |
| 1978 | 23.9 | 7.7 | 9.6 | 8.7 | 48.9 | 27.6 | 19.5 | 17.2 | 17.6 | 21.1 | 29.0 | 25.5 |
| 1977 | 22.8 | 7.9 | 9.4 | 8.4 | 46.3 | 25.1 | 18.3 | 17.0 | 17.6 | 20.7 | 30.0 | 26.8 |
| 1976 | 21.8 | 7.4 | 9.7 | 8.3 | 43.1 | 23.7 | 17.5 | 16.5 | 17.2 | 21.2 | 31.1 | 27.4 |
| 1975 | 21.3 | 8.0 | 9.4 | 8.4 | 41.2 | 23.9 | 18.0 | 16.2 | 16.9 | 21.2 | 31.5 | 25.0 |
| 1974 | 21.7 | 7.4 | 9.3 | 8.3 | 42.3 | 24.4 | 18.1 | 17.4 | 18.4 | 21.8 | 30.8 | 27.1 |
| 1973 | 26.0 | 10.3 | 11.1 | 10.3 | 48.3 | 28.9 | 22.2 | 21.4 | 23.6 | 28.6 | 40.5 | 36.2 |
| 1972 | 26.7 | 8.5 | 10.8 | 10.5 | 49.0 | 28.9 | 22.6 | 23.1 | 25.0 | 31.8 | 44.6 | 35.1 |
| 1971 | 26.0 | 9.5 | 10.4 | 10.2 | 47.2 | 29.2 | 22.6 | 22.4 | 24.8 | 31.7 | 42.5 | 37.2 |
| 1970 | 26.7 | 9.6 | 10.9 | 9.8 | 48.8 | 29.7 | 23.6 | 24.1 | 26.9 | 32.6 | 44.4 | 34.9 |
| 1969 | 27.3 | 10.9 | 11.5 | 9.5 | 52.0 | 31.1 | 24.2 | 24.0 | 26.8 | 34.1 | 44.3 | 35.8 |
| 1968 | 27.2 | 9.5 | 10.7 | 9.8 | 51.2 | 31.1 | 23.5 | 24.5 | 27.5 | 34.6 | 47.8 | 40.6 |

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| MOTOR VEHICLE ACCIDENTS (E810-E823) | | | | | | | | | | | | |
| WHITE, MALE | | | | | | | | | | | | |
| 1978 | 35.6 | 7.2 | 10.7 | 11.0 | 73.7 | 43.4 | 29.6 | 25.3 | 25.1 | 29.3 | 45.9 | 54.0 |
| 1977 | 33.5 | 8.3 | 10.0 | 10.2 | 69.6 | 39.3 | 27.2 | 25.0 | 25.3 | 28.7 | 47.3 | 50.8 |
| 1976 | 32.1 | 6.6 | 11.1 | 10.3 | 65.4 | 37.2 | 26.2 | 24.3 | 24.6 | 29.4 | 48.8 | 54.6 |
| 1975 | 31.8 | 7.8 | 10.8 | 10.7 | 63.7 | 37.6 | 27.6 | 24.0 | 24.2 | 30.2 | 50.5 | 49.6 |
| 1974 | 32.7 | 6.6 | 10.1 | 10.5 | 65.8 | 39.0 | 27.8 | 26.2 | 26.8 | 31.3 | 48.2 | 56.1 |
| 1973 | 38.2 | 11.3 | 12.6 | 13.1 | 74.0 | 45.3 | 32.9 | 31.1 | 33.2 | 39.7 | 63.7 | 71.8 |
| 1972 | 38.7 | 9.2 | 11.7 | 12.9 | 74.6 | 45.3 | 33.1 | 32.9 | 34.9 | 43.0 | 73.7 | 71.2 |
| 1971 | 37.8 | 10.2 | 11.1 | 12.7 | 72.3 | 45.7 | 33.7 | 32.5 | 35.1 | 43.9 | 66.0 | 74.3 |
| 1970 | 39.1 | 9.1 | 12.2 | 12.6 | 75.2 | 47.0 | 35.2 | 34.6 | 39.0 | 46.2 | 69.2 | 65.5 |
| 1969 | 40.1 | 10.6 | 12.7 | 11.9 | 81.6 | 49.7 | 35.6 | 34.7 | 37.6 | 48.0 | 68.8 | 68.0 |
| 1968 | 40.2 | 10.0 | 11.5 | 12.6 | 80.5 | 49.8 | 35.2 | 36.0 | 39.3 | 48.6 | 75.7 | 76.9 |
| WHITE, FEMALE | | | | | | | | | | | | |
| 1978 | 12.8 | 8.2 | 8.4 | 6.3 | 23.4 | 11.9 | 9.7 | 9.5 | 10.9 | 14.8 | 19.1 | 12.8 |
| 1977 | 12.5 | 7.6 | 8.7 | 6.5 | 22.3 | 10.9 | 9.6 | 9.5 | 10.8 | 14.6 | 19.8 | 15.9 |
| 1976 | 11.9 | 8.2 | 8.3 | 6.3 | 20.3 | 10.2 | 8.9 | 9.2 | 10.6 | 14.9 | 20.5 | 14.8 |
| 1975 | 11.3 | 8.2 | 8.0 | 6.0 | 18.2 | 10.3 | 8.7 | 8.8 | 10.3 | 14.3 | 20.0 | 13.2 |
| 1974 | 11.3 | 8.2 | 8.4 | 6.1 | 18.4 | 9.9 | 8.7 | 9.2 | 10.8 | 14.5 | 20.1 | 13.0 |
| 1973 | 14.4 | 9.4 | 9.5 | 7.5 | 22.2 | 12.6 | 11.7 | 12.2 | 15.0 | 20.2 | 26.1 | 18.6 |
| 1972 | 15.2 | 7.7 | 9.8 | 7.9 | 23.1 | 12.6 | 12.5 | 14.0 | 16.1 | 23.2 | 26.2 | 16.9 |
| 1971 | 14.7 | 8.9 | 9.7 | 7.6 | 21.9 | 13.0 | 12.0 | 12.9 | 15.5 | 22.2 | 27.4 | 18.3 |
| 1970 | 14.8 | 10.2 | 9.6 | 6.9 | 22.7 | 12.7 | 12.3 | 14.3 | 16.1 | 22.1 | 28.1 | 18.1 |
| 1969 | 15.2 | 11.2 | 10.3 | 6.9 | 22.9 | 13.1 | 13.2 | 13.9 | 17.0 | 23.2 | 28.0 | 18.8 |
| 1968 | 14.9 | 8.9 | 9.9 | 6.8 | 22.5 | 13.0 | 12.2 | 13.7 | 16.8 | 23.7 | 28.8 | 21.0 |

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| MOTOR VEHICLE ACCIDENTS (E810-E823) | | | | | | | | | | | | |
| ALL OTHER, BOTH SEXES | | | | | | | | | | | | |
| 1978 | 21.4 | 9.1 | 14.0 | 8.3 | 26.3 | 28.2 | 24.8 | 23.9 | 25.7 | 24.9 | 29.9 | 22.6 |
| 1977 | 20.9 | 6.6 | 12.4 | 9.2 | 26.4 | 27.7 | 22.5 | 23.6 | 26.7 | 23.1 | 26.9 | 24.4 |
| 1976 | 20.6 | 9.5 | 13.2 | 8.8 | 24.6 | 27.6 | 24.3 | 24.7 | 24.8 | 25.9 | 25.9 | 21.6 |
| 1975 | 21.1 | 8.6 | 13.9 | 9.7 | 23.8 | 28.5 | 24.6 | 25.8 | 26.8 | 28.9 | 26.7 | 24.7 |
| 1974 | 21.8 | 6.8 | 12.6 | 10.0 | 26.2 | 28.7 | 25.6 | 25.7 | 27.9 | 30.9 | 26.7 | 31.1 |
| 1973 | 27.8 | 10.8 | 17.8 | 11.3 | 34.4 | 37.4 | 34.7 | 35.6 | 33.9 | 37.0 | 29.7 | 23.6 |
| 1972 | 28.4 | 12.2 | 15.7 | 11.7 | 35.2 | 38.2 | 35.4 | 37.9 | 36.0 | 34.1 | 35.8 | 42.3 |
| 1971 | 28.3 | 12.1 | 15.5 | 12.1 | 36.5 | 39.3 | 32.8 | 34.5 | 37.3 | 39.3 | 34.9 | 29.2 |
| 1970 | 28.5 | 11.1 | 14.7 | 12.4 | 36.9 | 39.0 | 35.2 | 37.2 | 37.0 | 35.4 | 32.7 | 26.8 |
| 1969 | 30.3 | 10.5 | 15.9 | 12.3 | 42.4 | 41.0 | 37.3 | 38.2 | 40.3 | 35.8 | 40.6 | * |
| 1968 | 29.6 | 9.3 | 15.5 | 11.6 | 40.1 | 43.3 | 36.0 | 37.9 | 38.4 | 38.6 | 42.3 | 30.4 |
| ALL OTHER, MALE | | | | | | | | | | | | |
| 1978 | 33.5 | 7.3 | 14.7 | 10.8 | 41.1 | 48.3 | 41.6 | 40.0 | 41.3 | 44.0 | 57.1 | 46.6 |
| 1977 | 33.2 | * | 14.6 | 12.6 | 41.5 | 48.1 | 38.1 | 40.1 | 44.4 | 40.9 | 46.7 | 46.4 |
| 1976 | 32.5 | 8.6 | 14.8 | 11.7 | 39.2 | 47.2 | 41.0 | 41.5 | 41.0 | 42.1 | 47.9 | 43.6 |
| 1975 | 33.4 | 9.4 | 16.3 | 13.2 | 38.0 | 50.0 | 41.8 | 41.9 | 44.1 | 49.9 | 45.5 | 50.0 |
| 1974 | 34.2 | * | 13.7 | 13.1 | 42.7 | 48.0 | 43.3 | 43.4 | 44.7 | 53.1 | 46.3 | 60.0 |
| 1973 | 43.1 | 7.9 | 22.2 | 14.6 | 53.8 | 62.2 | 58.3 | 57.2 | 57.2 | 61.1 | 53.6 | 43.8 |
| 1972 | 44.4 | 10.3 | 19.9 | 13.9 | 56.7 | 67.6 | 58.4 | 61.1 | 61.5 | 54.6 | 63.6 | 76.6 |
| 1971 | 43.7 | 14.3 | 17.2 | 16.2 | 58.0 | 65.7 | 52.7 | 54.3 | 62.1 | 63.2 | 62.8 | * |
| 1970 | 44.3 | 10.8 | 17.0 | 15.7 | 60.3 | 67.4 | 57.7 | 60.6 | 59.9 | 57.5 | 56.5 | 41.5 |
| 1969 | 47.5 | 11.6 | 18.4 | 16.1 | 67.7 | 71.6 | 63.8 | 62.7 | 66.8 | 57.8 | 70.7 | * |
| 1968 | 46.8 | * | 18.8 | 15.3 | 67.1 | 75.3 | 58.2 | 62.9 | 62.9 | 62.9 | 70.4 | * |
| ALL OTHER, FEMALE | | | | | | | | | | | | |
| 1978 | 10.4 | 11.0 | 13.2 | 5.8 | 12.1 | 10.9 | 10.6 | 10.7 | 12.7 | 10.1 | 11.3 | * |
| 1977 | 9.6 | * | 10.2 | 5.6 | 12.0 | 10.1 | 9.4 | 9.9 | 11.9 | 9.1 | 13.3 | * |
| 1976 | 9.9 | 10.3 | 11.5 | 5.8 | 10.5 | 10.8 | 10.3 | 10.6 | 11.2 | 13.2 | 10.9 | * |
| 1975 | 9.9 | 7.7 | 11.4 | 6.1 | 10.3 | 10.0 | 10.2 | 12.1 | 12.3 | 12.2 | 13.8 | * |
| 1974 | 10.5 | 7.9 | 11.5 | 6.8 | 10.6 | 12.1 | 10.8 | 10.6 | 13.5 | 13.0 | 13.3 | * |
| 1973 | 13.9 | 13.8 | 13.5 | 7.9 | 16.0 | 16.1 | 15.0 | 17.1 | 13.9 | 17.5 | 12.9 | * |
| 1972 | 13.6 | 14.2 | 11.5 | 9.5 | 14.6 | 13.0 | 16.1 | 17.8 | 14.1 | 17.4 | 15.9 | * |
| 1971 | 14.3 | 10.0 | 13.8 | 7.9 | 16.2 | 16.7 | 16.3 | 17.4 | 15.7 | 19.8 | 14.3 | * |
| 1970 | 13.9 | 11.3 | 12.3 | 9.1 | 14.9 | 14.7 | 16.6 | 16.6 | 16.7 | 17.2 | 14.6 | * |
| 1969 | 14.4 | 9.4 | 13.5 | 8.6 | 19.0 | 15.1 | 15.2 | 16.7 | 16.9 | 17.7 | 17.4 | * |
| 1968 | 13.9 | 11.5 | 12.1 | 7.9 | 15.0 | 16.2 | 17.4 | 15.8 | 16.4 | 18.4 | 20.5 | * |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| MOTOR VEHICLE ACCIDENTS (E810-E823) | | | | | | | | | | | | |
| BLACK, BOTH SEXES | | | | | | | | | | | | |
| 1978 | 20.3 | 8.6 | 13.6 | 8.3 | 23.7 | 26.7 | 24.5 | 24.2 | 25.5 | 23.0 | 27.0 | 19.3 |
| 1977 | 20.0 | 7.0 | 12.2 | 9.0 | 23.7 | 27.0 | 22.4 | 23.6 | 26.8 | 22.8 | 24.7 | 23.1 |
| 1976 | 19.7 | 7.9 | 13.5 | 9.0 | 21.3 | 26.7 | 23.6 | 24.6 | 25.2 | 26.4 | 23.8 | 20.6 |
| 1975 | 20.2 | 8.3 | 13.8 | 9.4 | 20.7 | 27.6 | 24.8 | 26.2 | 26.6 | 28.3 | 25.3 | 21.5 |
| 1974 | 21.1 | 6.0 | 12.9 | 10.1 | 24.2 | 28.1 | 25.5 | 25.2 | 27.9 | 30.2 | 25.5 | 29.8 |
| 1973 | 26.9 | 9.8 | 18.0 | 11.5 | 31.3 | 36.0 | 35.0 | 36.1 | 33.3 | 36.1 | 25.3 | 23.7 |
| 1972 | 27.5 | 11.7 | 15.3 | 11.5 | 32.4 | 37.2 | 35.9 | 38.3 | 35.3 | 34.0 | 32.4 | 41.8 |
| 1971 | 27.9 | 12.3 | 16.0 | 12.3 | 34.4 | 38.7 | 33.3 | 34.8 | 37.2 | 38.3 | 34.6 | 29.0 |
| 1970 | 28.1 | 11.3 | 14.8 | 12.7 | 34.9 | 39.4 | 35.8 | 37.3 | 37.8 | 33.4 | 30.1 | 27.3 |
| 1969 | 29.6 | 9.6 | 16.0 | 12.5 | 40.6 | 39.6 | 37.7 | 38.4 | 39.5 | 34.5 | 39.4 | * |
| 1968 | 29.3 | 9.5 | 14.8 | 12.1 | 39.3 | 42.2 | 36.9 | 37.8 | 37.7 | 37.7 | 43.2 | 27.2 |
| BLACK, MALE | | | | | | | | | | | | |
| 1978 | 32.3 | * | 14.0 | 10.9 | 37.4 | 47.1 | 42.6 | 40.7 | 41.6 | 42.1 | 52.3 | 43.1 |
| 1977 | 32.6 | * | 14.5 | 12.7 | 38.1 | 48.3 | 39.3 | 41.1 | 45.6 | 41.0 | 45.1 | 42.9 |
| 1976 | 31.4 | * | 15.1 | 12.0 | 34.8 | 46.5 | 41.6 | 41.4 | 41.7 | 43.6 | 45.3 | 42.6 |
| 1975 | 32.4 | 8.7 | 16.4 | 12.9 | 33.4 | 49.7 | 43.2 | 43.2 | 44.7 | 50.4 | 42.8 | 43.5 |
| 1974 | 33.6 | * | 14.1 | 13.4 | 39.6 | 48.2 | 44.3 | 43.6 | 45.7 | 52.7 | 44.0 | 55.8 |
| 1973 | 42.2 | * | 22.6 | 15.1 | 49.3 | 61.0 | 60.2 | 59.2 | 56.3 | 62.1 | 47.1 | * |
| 1972 | 43.5 | 10.0 | 19.2 | 13.9 | 53.2 | 66.0 | 60.0 | 62.9 | 61.1 | 55.4 | 59.9 | 80.0 |
| 1971 | 43.4 | 14.9 | 17.5 | 16.6 | 55.2 | 66.6 | 54.0 | 55.0 | 63.4 | 63.2 | 64.5 | * |
| 1970 | 44.3 | 10.6 | 16.9 | 16.1 | 58.1 | 70.4 | 59.5 | 61.4 | 62.1 | 54.9 | 51.6 | 45.7 |
| 1969 | 46.8 | 10.5 | 18.4 | 16.3 | 65.1 | 70.3 | 64.8 | 63.6 | 66.3 | 56.5 | 68.9 | * |
| 1968 | 46.9 | * | 18.5 | 16.1 | 66.4 | 74.9 | 60.3 | 63.8 | 62.8 | 62.7 | 74.5 | * |
| BLACK, FEMALE | | | | | | | | | | | | |
| 1978 | 9.5 | 10.7 | 13.2 | 5.7 | 10.6 | 9.1 | 9.5 | 10.6 | 12.4 | 8.9 | 10.1 | * |
| 1977 | 8.6 | * | 9.9 | 5.3 | 9.9 | 8.7 | 8.3 | 9.1 | 11.5 | 9.1 | 11.3 | * |
| 1976 | 9.1 | 8.8 | 11.8 | 6.0 | 8.5 | 9.7 | 8.8 | 10.7 | 11.5 | 13.4 | 9.5 | * |
| 1975 | 9.1 | * | 11.2 | 5.9 | 8.6 | 8.6 | 9.6 | 12.0 | 11.6 | 11.5 | 13.4 | * |
| 1974 | 9.9 | * | 11.6 | 6.9 | 9.6 | 10.8 | 9.9 | 9.6 | 13.2 | 13.0 | 12.9 | * |
| 1973 | 13.0 | 12.7 | 13.4 | 7.8 | 14.2 | 14.7 | 14.0 | 16.4 | 14.1 | 16.1 | 10.5 | * |
| 1972 | 12.9 | 13.5 | 11.4 | 9.1 | 12.6 | 12.7 | 15.8 | 17.2 | 13.7 | 17.3 | 12.9 | * |
| 1971 | 13.8 | 9.6 | 14.6 | 8.0 | 14.7 | 15.2 | 16.2 | 17.4 | 15.2 | 18.7 | 13.1 | * |
| 1970 | 13.4 | 11.9 | 12.6 | 9.3 | 13.4 | 13.3 | 16.1 | 16.4 | 17.0 | 16.3 | 14.4 | * |
| 1969 | 14.0 | 8.8 | 13.5 | 8.7 | 17.9 | 13.8 | 15.2 | 16.6 | 16.6 | 17.0 | 17.6 | * |
| 1968 | 13.4 | 11.9 | 11.0 | 8.1 | 14.3 | 14.9 | 17.3 | 15.2 | 15.9 | 17.6 | 19.8 | * |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| ALL OTHER ACCIDENTS (E800-E807, E825-E949) | | | | | | | | | | | | |
| BOTH SEXES | | | | | | | | | | | | |
| 1978 | 23.9 | 30.0 | 17.9 | 8.2 | 17.7 | 16.9 | 18.7 | 21.6 | 27.1 | 39.0 | 91.7 | 266.2 |
| 1977 | 24.4 | 28.1 | 16.9 | 8.5 | 18.0 | 17.7 | 18.9 | 22.6 | 28.3 | 40.6 | 96.6 | 270.0 |
| 1976 | 24.7 | 32.5 | 17.2 | 8.3 | 18.6 | 18.3 | 18.7 | 22.5 | 29.0 | 40.3 | 96.9 | 291.0 |
| 1975 | 26.5 | 34.3 | 17.7 | 9.2 | 20.8 | 19.6 | 20.9 | 24.9 | 30.3 | 44.4 | 103.8 | 301.7 |
| 1974 | 27.3 | 40.1 | 19.1 | 9.6 | 20.9 | 19.7 | 21.0 | 25.2 | 31.3 | 46.1 | 109.4 | 340.9 |
| 1973 | 28.5 | 43.6 | 19.5 | 10.1 | 21.9 | 20.1 | 21.7 | 25.9 | 34.3 | 47.9 | 116.4 | 374.8 |
| 1972 | 28.3 | 44.9 | 20.0 | 10.0 | 20.5 | 19.4 | 20.9 | 26.2 | 34.5 | 48.8 | 121.6 | 380.3 |
| 1971 | 28.6 | 52.5 | 19.9 | 9.6 | 20.4 | 19.5 | 22.8 | 26.3 | 34.7 | 50.3 | 122.4 | 380.0 |
| 1970 | 29.5 | 56.0 | 20.0 | 9.9 | 21.5 | 20.7 | 23.3 | 28.0 | 35.3 | 52.8 | 131.2 | 358.5 |
| 1969 | 30.1 | 60.2 | 20.3 | 10.1 | 21.0 | 20.8 | 23.5 | 28.0 | 36.6 | 52.3 | 135.9 | 448.2 |
| 1968 | 30.1 | 65.4 | 21.1 | 10.5 | 19.5 | 19.9 | 22.7 | 27.8 | 36.2 | 55.0 | 141.5 | 474.0 |
| MALE | | | | | | | | | | | | |
| 1978 | 33.1 | 33.3 | 21.5 | 11.7 | 29.4 | 28.1 | 30.2 | 33.8 | 41.2 | 56.1 | 118.8 | 322.4 |
| 1977 | 33.8 | 32.0 | 20.4 | 12.2 | 30.0 | 29.9 | 30.3 | 34.6 | 42.7 | 58.6 | 122.7 | 320.1 |
| 1976 | 34.2 | 37.6 | 20.9 | 12.0 | 30.8 | 30.9 | 30.2 | 34.5 | 43.8 | 58.6 | 122.5 | 344.6 |
| 1975 | 37.0 | 36.5 | 20.6 | 13.5 | 35.2 | 32.9 | 33.3 | 38.8 | 46.6 | 63.9 | 131.8 | 339.4 |
| 1974 | 37.6 | 46.0 | 23.7 | 13.9 | 35.2 | 33.0 | 33.7 | 38.9 | 47.5 | 65.7 | 132.6 | 372.9 |
| 1973 | 39.4 | 50.3 | 23.2 | 14.6 | 37.1 | 34.3 | 35.2 | 40.4 | 52.4 | 67.4 | 139.2 | 411.5 |
| 1972 | 38.8 | 52.8 | 25.2 | 14.2 | 34.6 | 32.4 | 33.6 | 40.7 | 52.8 | 69.9 | 144.2 | 418.0 |
| 1971 | 39.1 | 60.6 | 24.3 | 13.6 | 34.8 | 32.3 | 36.7 | 40.9 | 52.9 | 71.8 | 142.0 | 408.8 |
| 1970 | 40.9 | 61.1 | 23.9 | 14.4 | 37.3 | 35.0 | 38.4 | 44.0 | 55.0 | 75.8 | 152.4 | 363.6 |
| 1969 | 41.3 | 67.0 | 23.9 | 14.7 | 36.4 | 35.7 | 38.5 | 44.1 | 56.8 | 73.6 | 154.8 | 464.1 |
| 1968 | 40.8 | 70.6 | 24.6 | 15.0 | 34.0 | 34.3 | 37.0 | 44.1 | 56.1 | 78.0 | 156.7 | 459.5 |

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| ALL OTHER ACCIDENTS (E800-E807, E825-E949) | | | | | | | | | | | | |
| FEMALE | | | | | | | | | | | | |
| 1978 | 15.3 | 26.5 | 14.1 | 4.7 | 5.8 | 5.9 | 7.7 | 10.2 | 14.6 | 26.0 | 75.6 | 240.8 |
| 1977 | 15.6 | 24.0 | 13.2 | 4.7 | 5.9 | 5.7 | 7.9 | 11.4 | 15.6 | 26.9 | 81.0 | 247.0 |
| 1976 | 15.7 | 27.1 | 13.3 | 4.4 | 6.1 | 6.0 | 7.7 | 11.4 | 15.9 | 26.3 | 81.4 | 265.9 |
| 1975 | 16.6 | 32.0 | 14.6 | 4.8 | 6.3 | 6.7 | 9.1 | 11.9 | 15.9 | 29.4 | 86.7 | 283.5 |
| 1974 | 17.5 | 34.0 | 14.2 | 5.1 | 6.4 | 6.8 | 8.9 | 12.5 | 17.0 | 31.0 | 95.1 | 325.1 |
| 1973 | 18.2 | 36.7 | 15.6 | 5.4 | 6.5 | 6.3 | 8.8 | 12.5 | 18.2 | 33.0 | 102.2 | 356.3 |
| 1972 | 18.3 | 36.6 | 14.6 | 5.5 | 6.4 | 6.7 | 8.8 | 12.6 | 18.1 | 32.6 | 107.1 | 361.0 |
| 1971 | 18.6 | 44.1 | 15.4 | 5.4 | 6.1 | 7.0 | 9.5 | 12.8 | 18.5 | 33.7 | 109.7 | 364.9 |
| 1970 | 18.7 | 50.7 | 15.9 | 5.3 | 6.0 | 6.9 | 9.0 | 13.1 | 17.6 | 34.8 | 117.2 | 355.6 |
| 1969 | 19.4 | 53.1 | 16.7 | 5.4 | 6.0 | 6.5 | 9.4 | 13.1 | 18.4 | 35.6 | 123.2 | 439.6 |
| 1968 | 19.9 | 60.0 | 17.5 | 5.8 | 5.3 | 6.1 | 9.2 | 12.5 | 18.2 | 36.9 | 131.0 | 482.5 |
| WHITE, BOTH SEXES | | | | | | | | | | | | |
| 1978 | 23.1 | 24.2 | 15.9 | 7.5 | 17.5 | 16.0 | 16.6 | 19.5 | 25.0 | 36.9 | 91.9 | 272.8 |
| 1977 | 23.6 | 22.9 | 15.3 | 7.8 | 17.6 | 16.3 | 16.9 | 20.4 | 26.2 | 38.7 | 97.1 | 276.5 |
| 1976 | 23.9 | 27.4 | 15.7 | 7.6 | 18.1 | 16.7 | 16.8 | 20.3 | 27.2 | 38.5 | 97.5 | 299.9 |
| 1975 | 25.6 | 27.3 | 15.9 | 8.4 | 19.9 | 18.0 | 18.6 | 22.8 | 28.4 | 42.7 | 104.5 | 311.5 |
| 1974 | 26.3 | 33.1 | 17.5 | 8.9 | 20.0 | 18.0 | 18.8 | 22.8 | 29.3 | 44.0 | 111.6 | 351.1 |
| 1973 | 27.4 | 33.9 | 17.4 | 9.1 | 20.7 | 18.4 | 19.4 | 23.7 | 32.4 | 45.9 | 117.4 | 386.6 |
| 1972 | 27.1 | 38.1 | 18.1 | 9.2 | 18.9 | 17.6 | 18.3 | 24.0 | 32.4 | 46.0 | 122.7 | 390.1 |
| 1971 | 27.1 | 42.4 | 17.7 | 8.6 | 18.4 | 17.3 | 20.3 | 24.0 | 33.0 | 47.7 | 123.1 | 391.8 |
| 1970 | 28.0 | 44.4 | 17.6 | 9.0 | 19.4 | 18.2 | 20.5 | 25.4 | 33.1 | 50.4 | 131.5 | 375.1 |
| 1969 | 28.7 | 48.0 | 17.5 | 9.0 | 19.1 | 18.7 | 21.3 | 25.8 | 34.5 | 49.9 | 136.7 | 464.4 |
| 1968 | 28.4 | 51.8 | 17.7 | 9.2 | 17.7 | 17.6 | 19.9 | 25.3 | 33.6 | 52.3 | 142.8 | 493.1 |

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| ALL OTHER ACCIDENTS (E800-E807, E825-E949) | | | | | | | | | | | | |
| WHITE, MALE | | | | | | | | | | | | |
| 1978 | 31.5 | 27.3 | 19.6 | 10.6 | 29.0 | 26.5 | 26.5 | 30.2 | 37.7 | 52.9 | 118.1 | 329.5 |
| 1977 | 32.2 | 26.6 | 18.7 | 11.1 | 29.4 | 27.5 | 26.8 | 30.8 | 39.4 | 55.5 | 123.4 | 329.4 |
| 1976 | 32.6 | 32.3 | 19.3 | 11.1 | 30.1 | 28.3 | 26.8 | 30.7 | 40.7 | 55.5 | 122.8 | 356.8 |
| 1975 | 35.1 | 29.4 | 18.9 | 12.3 | 33.8 | 30.1 | 29.3 | 35.1 | 43.3 | 61.4 | 132.5 | 351.3 |
| 1974 | 35.8 | 38.2 | 22.3 | 12.9 | 33.8 | 30.2 | 30.0 | 34.8 | 44.0 | 62.5 | 134.8 | 385.8 |
| 1973 | 37.2 | 40.3 | 21.3 | 13.1 | 35.3 | 31.2 | 31.4 | 36.4 | 48.8 | 64.5 | 139.5 | 424.1 |
| 1972 | 36.5 | 45.4 | 23.5 | 13.2 | 31.9 | 29.5 | 29.0 | 37.0 | 49.3 | 65.7 | 144.7 | 432.4 |
| 1971 | 36.5 | 50.3 | 21.8 | 12.3 | 31.6 | 28.6 | 32.6 | 36.7 | 50.2 | 68.3 | 142.3 | 420.7 |
| 1970 | 38.2 | 48.1 | 21.6 | 13.2 | 33.7 | 30.9 | 33.6 | 39.3 | 51.1 | 72.9 | 152.8 | 383.4 |
| 1969 | 38.8 | 56.1 | 21.1 | 13.1 | 33.2 | 32.1 | 34.6 | 40.4 | 53.3 | 70.3 | 155.8 | 484.6 |
| 1968 | 37.9 | 55.6 | 21.1 | 13.3 | 30.9 | 30.2 | 32.2 | 39.6 | 52.1 | 73.8 | 158.1 | 479.4 |
| WHITE, FEMALE | | | | | | | | | | | | |
| 1978 | 15.1 | 21.0 | 12.0 | 4.3 | 5.6 | 5.4 | 7.0 | 9.4 | 13.7 | 24.7 | 76.5 | 247.6 |
| 1977 | 15.4 | 19.0 | 11.6 | 4.3 | 5.5 | 5.1 | 7.3 | 10.5 | 14.5 | 25.8 | 81.6 | 252.5 |
| 1976 | 15.6 | 22.2 | 12.0 | 4.0 | 5.8 | 5.2 | 7.0 | 10.5 | 15.2 | 25.6 | 82.5 | 273.8 |
| 1975 | 16.5 | 25.1 | 12.8 | 4.4 | 5.7 | 5.9 | 8.1 | 11.2 | 15.1 | 28.4 | 87.6 | 292.8 |
| 1974 | 17.3 | 27.7 | 12.4 | 4.6 | 5.9 | 5.9 | 7.8 | 11.4 | 16.1 | 29.9 | 97.4 | 334.2 |
| 1973 | 18.1 | 27.2 | 13.2 | 5.0 | 5.8 | 5.6 | 7.8 | 11.7 | 17.8 | 31.7 | 103.8 | 368.4 |
| 1972 | 18.0 | 30.4 | 12.3 | 5.0 | 5.8 | 5.8 | 7.9 | 11.9 | 17.4 | 31.0 | 108.7 | 368.9 |
| 1971 | 18.2 | 34.0 | 13.4 | 4.7 | 5.2 | 6.2 | 8.5 | 12.0 | 17.5 | 31.8 | 110.8 | 377.0 |
| 1970 | 18.3 | 40.6 | 13.4 | 4.7 | 5.3 | 5.8 | 7.9 | 12.3 | 16.9 | 33.0 | 117.6 | 370.5 |
| 1969 | 19.0 | 39.5 | 13.7 | 4.7 | 5.2 | 5.6 | 8.4 | 12.2 | 17.5 | 34.1 | 124.0 | 454.3 |
| 1968 | 19.5 | 47.8 | 14.1 | 5.0 | 4.8 | 5.2 | 8.1 | 11.8 | 16.8 | 35.5 | 132.4 | 501.1 |

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| ALL OTHER ACCIDENTS (E800-E807, E825-E949) | | | | | | | | | | | | |
| ALL OTHER, BOTH SEXES | | | | | | | | | | | | |
| 1978 | 29.0 | 57.8 | 27.1 | 11.9 | 18.9 | 22.9 | 33.0 | 37.9 | 46.0 | 59.2 | 89.8 | 190.5 |
| 1977 | 29.9 | 52.9 | 24.8 | 12.3 | 20.3 | 27.1 | 32.0 | 40.0 | 47.5 | 59.0 | 91.6 | 195.6 |
| 1976 | 29.9 | 57.6 | 24.1 | 11.8 | 21.4 | 29.1 | 31.9 | 40.0 | 45.6 | 57.6 | 90.4 | 188.2 |
| 1975 | 32.9 | 69.5 | 26.1 | 13.6 | 26.2 | 30.7 | 37.2 | 42.1 | 48.2 | 60.5 | 96.1 | 189.7 |
| 1974 | 33.6 | 74.9 | 27.1 | 13.3 | 26.1 | 31.5 | 36.7 | 45.5 | 50.8 | 66.3 | 85.6 | 222.2 |
| 1973 | 36.2 | 91.8 | 30.5 | 15.5 | 29.2 | 32.6 | 37.9 | 45.1 | 52.6 | 68.0 | 105.3 | 237.8 |
| 1972 | 36.7 | 79.9 | 30.4 | 14.3 | 30.5 | 32.5 | 40.2 | 44.5 | 53.9 | 76.4 | 108.7 | 266.7 |
| 1971 | 38.5 | 106.9 | 31.8 | 15.4 | 32.8 | 34.5 | 40.9 | 46.8 | 51.4 | 76.5 | 114.4 | 245.0 |
| 1970 | 40.3 | 117.6 | 32.5 | 14.9 | 34.8 | 38.0 | 43.7 | 51.0 | 56.2 | 75.6 | 127.4 | 187.9 |
| 1969 | 40.2 | 129.2 | 35.6 | 16.7 | 33.7 | 35.8 | 40.9 | 47.8 | 56.9 | 75.8 | 125.1 | 266.1 |
| 1968 | 42.0 | 139.0 | 39.7 | 17.6 | 31.2 | 36.7 | 44.5 | 50.5 | 61.6 | 82.0 | 124.4 | 260.8 |
| ALL OTHER, MALE | | | | | | | | | | | | |
| 1978 | 43.4 | 62.6 | 30.7 | 17.1 | 31.5 | 39.5 | 57.2 | 63.9 | 73.5 | 85.8 | 125.2 | 250.0 |
| 1977 | 44.7 | 58.1 | 28.7 | 17.9 | 33.0 | 47.5 | 56.5 | 66.7 | 74.0 | 86.9 | 116.1 | 225.0 |
| 1976 | 45.1 | 64.3 | 28.8 | 17.1 | 35.0 | 49.8 | 55.3 | 66.9 | 73.8 | 87.5 | 120.6 | 221.8 |
| 1975 | 49.6 | 72.6 | 28.9 | 20.2 | 43.4 | 53.4 | 63.0 | 71.0 | 78.0 | 88.3 | 125.2 | 220.4 |
| 1974 | 50.2 | 84.9 | 30.7 | 19.3 | 43.5 | 53.9 | 61.5 | 74.4 | 81.2 | 96.4 | 109.6 | 240.0 |
| 1973 | 54.5 | 100.0 | 32.9 | 23.2 | 48.8 | 57.4 | 64.6 | 75.6 | 88.2 | 95.2 | 136.2 | 283.3 |
| 1972 | 54.8 | 91.6 | 34.1 | 20.0 | 51.6 | 55.2 | 70.0 | 74.6 | 86.8 | 110.4 | 138.2 | 272.3 |
| 1971 | 57.2 | 116.4 | 37.6 | 21.6 | 55.5 | 60.5 | 69.8 | 78.5 | 79.2 | 105.9 | 140.1 | 291.5 |
| 1970 | 60.7 | 131.6 | 36.1 | 20.9 | 60.6 | 66.1 | 76.9 | 86.3 | 92.1 | 103.3 | 147.6 | 189.5 |
| 1969 | 59.4 | 129.8 | 38.8 | 24.0 | 58.3 | 63.4 | 70.8 | 78.2 | 91.0 | 105.1 | 142.6 | 270.5 |
| 1968 | 62.5 | 152.8 | 43.6 | 25.1 | 55.3 | 66.1 | 77.0 | 85.9 | 94.6 | 118.8 | 140.3 | 265.9 |
| ALL OTHER, FEMALE | | | | | | | | | | | | |
| 1978 | 15.9 | 52.8 | 23.5 | 6.7 | 6.8 | 8.5 | 12.6 | 16.5 | 23.2 | 38.6 | 65.8 | 160.6 |
| 1977 | 16.4 | 47.7 | 20.7 | 6.6 | 8.1 | 9.5 | 11.6 | 17.8 | 25.4 | 37.1 | 74.7 | 179.8 |
| 1976 | 16.0 | 51.0 | 19.4 | 6.5 | 8.2 | 11.2 | 12.3 | 17.6 | 21.9 | 34.0 | 69.7 | 169.4 |
| 1975 | 17.7 | 66.4 | 23.3 | 6.9 | 9.8 | 11.2 | 15.6 | 17.8 | 23.1 | 38.3 | 76.0 | 169.9 |
| 1974 | 18.5 | 64.8 | 23.4 | 7.3 | 9.4 | 12.3 | 16.0 | 21.0 | 24.8 | 42.1 | 69.0 | 211.8 |
| 1973 | 19.4 | 83.5 | 28.0 | 7.8 | 10.5 | 11.3 | 15.5 | 19.0 | 22.2 | 46.1 | 84.0 | 210.1 |
| 1972 | 20.0 | 67.9 | 26.6 | 8.4 | 10.4 | 13.0 | 15.2 | 18.6 | 25.5 | 48.9 | 87.7 | 263.2 |
| 1971 | 21.3 | 97.2 | 26.0 | 9.3 | 11.4 | 12.3 | 16.8 | 19.3 | 27.2 | 52.5 | 95.4 | 215.1 |
| 1970 | 21.6 | 103.5 | 28.8 | 8.9 | 10.6 | 13.9 | 16.2 | 20.0 | 24.5 | 52.8 | 111.9 | 186.9 |
| 1969 | 22.7 | 128.5 | 32.3 | 9.3 | 10.9 | 12.3 | 16.0 | 21.1 | 26.6 | 51.6 | 111.3 | 263.1 |
| 1968 | 23.1 | 124.4 | 35.7 | 10.1 | 8.8 | 11.8 | 17.2 | 19.3 | 32.0 | 51.5 | 112.6 | 257.4 |

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| ALL OTHER ACCIDENTS (E800-E807, E825-E949) | | | | | | | | | | | | |
| BLACK, BOTH SEXES | | | | | | | | | | | | |
| 1978 | 30.3 | 62.5 | 28.2 | 12.6 | 18.6 | 24.5 | 35.9 | 40.4 | 48.2 | 61.1 | 88.1 | 184.7 |
| 1977 | 31.5 | 57.3 | 26.1 | 12.9 | 20.4 | 29.5 | 34.6 | 42.9 | 50.0 | 62.1 | 90.3 | 195.1 |
| 1976 | 31.1 | 62.0 | 25.5 | 12.0 | 21.5 | 31.1 | 34.3 | 42.5 | 46.8 | 59.2 | 89.5 | 192.6 |
| 1975 | 34.4 | 75.0 | 27.8 | 14.0 | 26.8 | 33.4 | 40.3 | 44.6 | 49.3 | 61.8 | 92.9 | 185.4 |
| 1974 | 34.9 | 80.8 | 27.9 | 13.7 | 26.6 | 34.1 | 39.0 | 46.9 | 52.2 | 68.7 | 85.1 | 224.0 |
| 1973 | 37.1 | 98.7 | 31.6 | 16.2 | 29.5 | 33.2 | 39.0 | 47.0 | 54.1 | 68.7 | 104.2 | 236.8 |
| 1972 | 37.8 | 87.2 | 30.9 | 14.7 | 31.1 | 33.3 | 43.0 | 46.3 | 54.9 | 77.8 | 109.2 | 267.3 |
| 1971 | 39.6 | 112.5 | 32.5 | 15.9 | 33.6 | 36.9 | 43.3 | 47.8 | 52.0 | 77.8 | 113.3 | 236.4 |
| 1970 | 41.9 | 123.4 | 33.3 | 15.5 | 36.0 | 40.7 | 46.2 | 53.5 | 57.1 | 77.6 | 132.7 | 192.5 |
| 1969 | 40.8 | 138.3 | 36.3 | 17.0 | 34.0 | 36.5 | 41.9 | 48.0 | 57.4 | 75.0 | 126.0 | 266.0 |
| 1968 | 42.7 | 147.1 | 39.9 | 17.9 | 31.4 | 37.5 | 46.3 | 51.7 | 61.9 | 82.7 | 124.5 | 258.7 |
| BLACK, MALE | | | | | | | | | | | | |
| 1978 | 45.1 | 66.5 | 31.6 | 18.1 | 31.2 | 41.9 | 62.7 | 68.6 | 77.3 | 88.4 | 119.2 | 239.2 |
| 1977 | 46.9 | 62.0 | 30.5 | 18.9 | 33.2 | 51.5 | 61.4 | 71.3 | 78.3 | 91.8 | 113.6 | 216.3 |
| 1976 | 47.1 | 69.4 | 30.5 | 17.6 | 35.6 | 53.3 | 60.0 | 71.7 | 76.3 | 90.2 | 116.7 | 223.4 |
| 1975 | 51.8 | 80.0 | 30.5 | 20.8 | 44.6 | 58.0 | 68.2 | 75.9 | 79.5 | 91.4 | 119.4 | 206.5 |
| 1974 | 52.3 | 92.9 | 32.1 | 19.9 | 44.5 | 58.5 | 65.9 | 77.2 | 83.7 | 101.8 | 108.1 | 241.9 |
| 1973 | 55.8 | 109.0 | 34.1 | 24.3 | 49.6 | 58.4 | 66.4 | 78.4 | 91.5 | 95.8 | 130.9 | 290.2 |
| 1972 | 56.6 | 101.2 | 34.7 | 20.5 | 52.9 | 56.9 | 75.9 | 77.9 | 88.6 | 113.9 | 138.1 | 270.0 |
| 1971 | 58.7 | 122.0 | 38.1 | 22.0 | 57.0 | 65.4 | 73.8 | 79.6 | 80.6 | 107.5 | 136.6 | 282.5 |
| 1970 | 63.3 | 139.9 | 36.8 | 21.8 | 63.3 | 71.4 | 81.9 | 91.5 | 94.0 | 106.1 | 152.4 | 200.0 |
| 1969 | 60.0 | 138.4 | 38.7 | 24.5 | 59.5 | 64.9 | 72.1 | 79.0 | 93.1 | 102.4 | 138.9 | 270.3 |
| 1968 | 63.5 | 160.5 | 43.5 | 25.5 | 55.8 | 67.6 | 79.6 | 88.8 | 96.1 | 120.3 | 140.0 | 271.4 |
| BLACK, FEMALE | | | | | | | | | | | | |
| 1978 | 16.9 | 58.4 | 24.8 | 7.0 | 6.6 | 9.4 | 13.6 | 17.1 | 24.4 | 40.8 | 67.3 | 156.6 |
| 1977 | 17.5 | 52.5 | 21.6 | 7.0 | 8.2 | 10.5 | 12.3 | 19.3 | 26.8 | 39.8 | 75.1 | 184.0 |
| 1976 | 16.7 | 54.4 | 20.3 | 6.3 | 8.1 | 12.0 | 13.0 | 18.1 | 22.4 | 35.7 | 71.3 | 176.4 |
| 1975 | 18.6 | 69.9 | 25.1 | 7.2 | 9.8 | 12.2 | 17.2 | 18.3 | 24.3 | 39.3 | 75.0 | 173.8 |
| 1974 | 19.2 | 68.3 | 23.6 | 7.5 | 9.5 | 13.1 | 16.5 | 21.5 | 26.1 | 43.4 | 69.6 | 214.1 |
| 1973 | 20.3 | 87.8 | 29.0 | 8.2 | 10.6 | 11.6 | 16.1 | 20.3 | 22.9 | 47.7 | 86.1 | 206.8 |
| 1972 | 20.7 | 72.6 | 27.0 | 9.0 | 10.5 | 13.3 | 15.7 | 19.2 | 26.7 | 49.8 | 88.6 | 265.7 |
| 1971 | 22.2 | 102.8 | 26.9 | 9.7 | 11.6 | 12.8 | 18.0 | 20.5 | 27.8 | 54.3 | 96.5 | 209.0 |
| 1970 | 22.5 | 106.7 | 29.8 | 9.1 | 10.6 | 14.8 | 16.7 | 20.7 | 25.7 | 55.0 | 118.3 | 187.7 |
| 1969 | 23.4 | 138.2 | 33.8 | 9.5 | 10.5 | 12.6 | 16.8 | 21.1 | 26.9 | 53.1 | 115.9 | 263.3 |
| 1968 | 23.8 | 133.6 | 36.3 | 10.3 | 8.8 | 12.4 | 18.4 | 19.5 | 32.2 | 52.2 | 113.1 | 255.4 |

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| Suicide E950-E959 | | | | | | | | | | | | |
| BOTH SEXES | | | | | | | | | | | | |
| 1978 | 12.3 | ... | ... | 0.4 | 12.1 | 16.2 | 15.8 | 17.2 | 17.8 | 18.8 | 21.1 | 19.6 |
| 1977 | 13.1 | ... | ... | 0.5 | 13.3 | 17.2 | 16.8 | 18.9 | 19.0 | 20.0 | 20.3 | 18.1 |
| 1976 | 12.3 | ... | ... | 0.4 | 11.5 | 15.5 | 16.3 | 19.2 | 19.6 | 19.5 | 19.7 | 19.6 |
| 1975 | 12.6 | ... | ... | 0.4 | 11.7 | 16.1 | 17.4 | 20.1 | 19.8 | 19.6 | 19.7 | 18.6 |
| 1974 | 12.0 | ... | ... | 0.5 | 10.8 | 15.5 | 16.8 | 19.6 | 19.5 | 18.8 | 20.2 | 17.9 |
| 1973 | 11.9 | ... | ... | 0.4 | 10.5 | 14.7 | 16.4 | 19.5 | 20.2 | 19.8 | 20.8 | 20.2 |
| 1972 | 11.9 | ... | ... | 0.3 | 10.1 | 14.6 | 16.8 | 19.9 | 21.3 | 20.4 | 21.7 | 20.8 |
| 1971 | 11.6 | ... | ... | 0.3 | 9.3 | 13.7 | 17.1 | 19.8 | 21.4 | 20.9 | 21.3 | 19.1 |
| 1970 | 11.6 | ... | ... | 0.3 | 8.8 | 14.1 | 16.9 | 20.0 | 21.4 | 20.8 | 21.2 | 19.0 |
| 1969 | 11.1 | ... | ... | 0.3 | 7.9 | 12.9 | 16.6 | 19.5 | 21.1 | 20.1 | 22.7 | 20.4 |
| 1968 | 10.7 | ... | ... | 0.3 | 7.1 | 12.0 | 16.1 | 19.7 | 21.5 | 19.0 | 21.3 | 22.1 |
| MALE | | | | | | | | | | | | |
| 1978 | 18.7 | ... | ... | 0.6 | 19.5 | 24.8 | 21.7 | 23.6 | 27.2 | 33.1 | 44.8 | 50.9 |
| 1977 | 19.8 | ... | ... | 0.8 | 21.3 | 25.9 | 23.6 | 25.8 | 28.9 | 34.9 | 42.9 | 47.8 |
| 1976 | 18.4 | ... | ... | 0.7 | 18.2 | 23.0 | 22.7 | 26.3 | 29.5 | 33.9 | 40.7 | 49.3 |
| 1975 | 18.7 | ... | ... | 0.7 | 18.6 | 23.9 | 23.4 | 28.0 | 29.9 | 33.6 | 40.4 | 47.6 |
| 1974 | 17.9 | ... | ... | 0.7 | 16.9 | 22.9 | 22.8 | 26.6 | 30.1 | 32.9 | 42.2 | 46.2 |
| 1973 | 17.6 | ... | ... | 0.6 | 16.8 | 21.6 | 21.8 | 26.9 | 30.4 | 34.5 | 42.5 | 50.9 |
| 1972 | 17.4 | ... | ... | 0.4 | 15.6 | 20.7 | 22.1 | 28.0 | 31.4 | 35.9 | 44.2 | 50.6 |
| 1971 | 16.7 | ... | ... | 0.5 | 14.0 | 19.1 | 22.2 | 26.8 | 32.4 | 35.6 | 43.4 | 48.8 |
| 1970 | 16.8 | ... | ... | 0.5 | 13.5 | 19.8 | 22.1 | 27.9 | 32.7 | 36.0 | 42.8 | 42.4 |
| 1969 | 16.2 | ... | ... | 0.5 | 12.2 | 18.3 | 21.7 | 27.2 | 32.4 | 34.5 | 45.7 | 50.5 |
| 1968 | 15.8 | ... | ... | 0.5 | 10.9 | 17.2 | 22.0 | 27.3 | 33.6 | 33.6 | 43.0 | 54.1 |

... Category not applicable

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| Suicide E950-E959 | | | | | | | | | | | | |
| FEMALE | | | | | | | | | | | | |
| 1978 | 6.2 | ... | ... | 0.2 | 4.6 | 7.8 | 10.1 | 11.2 | 9.4 | 7.8 | 6.9 | 5.4 |
| 1977 | 6.7 | ... | ... | 0.2 | 5.2 | 8.7 | 10.4 | 12.5 | 10.3 | 8.6 | 6.7 | 4.4 |
| 1976 | 6.6 | ... | ... | 0.2 | 4.7 | 8.2 | 10.2 | 12.7 | 10.9 | 8.4 | 7.0 | 5.6 |
| 1975 | 6.7 | ... | ... | 0.2 | 4.7 | 8.4 | 11.6 | 12.7 | 10.8 | 8.9 | 7.0 | 4.6 |
| 1974 | 6.5 | ... | ... | 0.2 | 4.6 | 8.2 | 11.1 | 13.0 | 10.1 | 7.9 | 6.6 | 3.9 |
| 1973 | 6.5 | ... | ... | 0.2 | 4.2 | 8.0 | 11.3 | 12.5 | 11.1 | 8.4 | 7.2 | 4.7 |
| 1972 | 6.7 | ... | ... | 0.2 | 4.7 | 8.7 | 11.6 | 12.4 | 12.3 | 8.5 | 7.4 | 5.5 |
| 1971 | 6.8 | ... | ... | 0.2 | 4.6 | 8.5 | 12.1 | 13.4 | 11.6 | 9.5 | 7.0 | 3.6 |
| 1970 | 6.6 | ... | ... | 0.2 | 4.2 | 8.6 | 11.9 | 12.6 | 11.4 | 9.0 | 7.0 | 5.9 |
| 1969 | 6.3 | ... | ... | 0.2 | 3.8 | 7.8 | 11.7 | 12.3 | 10.9 | 8.9 | 7.3 | 4.1 |
| 1968 | 5.9 | ... | ... | 0.1 | 3.4 | 7.1 | 10.6 | 12.5 | 10.6 | 7.5 | 6.5 | 4.5 |
| WHITE, BOTH SEXES | | | | | | | | | | | | |
| 1978 | 13.2 | ... | ... | 0.4 | 12.8 | 16.7 | 16.7 | 18.3 | 19.0 | 20.0 | 22.4 | 20.6 |
| 1977 | 14.0 | ... | ... | 0.6 | 14.0 | 17.6 | 17.9 | 20.2 | 20.2 | 21.5 | 21.6 | 19.0 |
| 1976 | 13.2 | ... | ... | 0.4 | 11.9 | 15.8 | 17.2 | 20.5 | 21.0 | 20.6 | 21.0 | 20.1 |
| 1975 | 13.4 | ... | ... | 0.5 | 12.2 | 16.4 | 18.4 | 21.5 | 21.1 | 20.9 | 20.9 | 19.7 |
| 1974 | 12.9 | ... | ... | 0.5 | 11.2 | 15.8 | 17.9 | 21.0 | 20.8 | 19.9 | 21.3 | 18.8 |
| 1973 | 12.7 | ... | ... | 0.4 | 10.8 | 15.0 | 17.4 | 20.8 | 21.5 | 21.1 | 22.0 | 21.1 |
| 1972 | 12.7 | ... | ... | 0.3 | 10.0 | 14.9 | 17.7 | 21.3 | 22.7 | 21.9 | 22.9 | 22.3 |
| 1971 | 12.5 | ... | ... | 0.4 | 9.5 | 14.0 | 18.0 | 21.3 | 22.9 | 22.2 | 22.6 | 20.0 |
| 1970 | 12.4 | ... | ... | 0.3 | 9.0 | 14.4 | 18.1 | 21.2 | 23.1 | 22.2 | 22.4 | 20.0 |
| 1969 | 11.9 | ... | ... | 0.4 | 8.1 | 13.2 | 17.5 | 20.8 | 22.6 | 21.5 | 23.7 | 21.4 |
| 1968 | 11.5 | ... | ... | 0.3 | 7.3 | 12.4 | 17.2 | 21.0 | 23.0 | 20.2 | 22.3 | 23.6 |

... Category not applicable

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| Suicide E950-E959 | | | | | | | | | | | | |
| WHITE, MALE | | | | | | | | | | | | |
| 1978 | 19.9 | ... | ... | 0.7 | 20.4 | 25.1 | 22.4 | 24.8 | 29.0 | 35.3 | 48.3 | 54.5 |
| 1977 | 21.0 | ... | ... | 0.9 | 22.4 | 26.1 | 24.7 | 27.4 | 30.6 | 37.4 | 46.2 | 51.0 |
| 1976 | 19.5 | ... | ... | 0.7 | 18.9 | 23.2 | 23.5 | 27.7 | 31.3 | 36.1 | 43.8 | 50.8 |
| 1975 | 19.9 | ... | ... | 0.8 | 19.3 | 24.0 | 24.4 | 29.7 | 31.9 | 36.0 | 43.2 | 51.1 |
| 1974 | 19.0 | ... | ... | 0.8 | 17.6 | 23.0 | 23.8 | 28.3 | 31.9 | 34.8 | 44.7 | 48.9 |
| 1973 | 18.7 | ... | ... | 0.7 | 17.3 | 21.5 | 22.8 | 28.4 | 32.3 | 36.9 | 45.3 | 54.9 |
| 1972 | 18.4 | ... | ... | 0.5 | 15.4 | 20.7 | 22.9 | 29.7 | 33.4 | 38.4 | 47.2 | 54.7 |
| 1971 | 17.9 | ... | ... | 0.5 | 14.4 | 19.3 | 23.2 | 28.6 | 34.5 | 37.8 | 46.6 | 52.1 |
| 1970 | 18.0 | ... | ... | 0.5 | 13.9 | 19.9 | 23.3 | 29.5 | 35.0 | 38.7 | 45.5 | 45.8 |
| 1969 | 17.3 | ... | ... | 0.5 | 12.5 | 18.4 | 22.5 | 28.8 | 34.7 | 36.8 | 48.0 | 54.0 |
| 1968 | 17.0 | ... | ... | 0.5 | 11.3 | 17.4 | 23.2 | 29.0 | 35.8 | 35.8 | 45.5 | 58.9 |
| WHITE, FEMALE | | | | | | | | | | | | |
| 1978 | 6.7 | ... | ... | 0.2 | 4.9 | 8.2 | 11.0 | 12.1 | 10.1 | 8.4 | 7.3 | 5.4 |
| 1977 | 7.2 | ... | ... | 0.2 | 5.4 | 9.1 | 11.2 | 13.5 | 10.9 | 9.4 | 7.1 | 4.5 |
| 1976 | 7.1 | ... | ... | 0.2 | 4.8 | 8.5 | 11.1 | 13.8 | 11.8 | 8.9 | 7.4 | 6.0 |
| 1975 | 7.3 | ... | ... | 0.2 | 4.9 | 8.8 | 12.6 | 13.8 | 11.5 | 9.4 | 7.5 | 4.8 |
| 1974 | 7.0 | ... | ... | 0.2 | 4.7 | 8.6 | 12.1 | 14.1 | 10.9 | 8.5 | 7.0 | 4.1 |
| 1973 | 7.0 | ... | ... | 0.2 | 4.3 | 8.4 | 12.2 | 13.7 | 11.8 | 9.0 | 7.5 | 4.4 |
| 1972 | 7.3 | ... | ... | 0.2 | 4.5 | 9.2 | 12.6 | 13.4 | 13.2 | 9.2 | 7.6 | 5.9 |
| 1971 | 7.3 | ... | ... | 0.2 | 4.5 | 8.8 | 13.0 | 14.5 | 12.5 | 10.2 | 7.2 | 3.7 |
| 1970 | 7.1 | ... | ... | 0.1 | 4.2 | 9.0 | 13.0 | 13.5 | 12.3 | 9.6 | 7.2 | 5.8 |
| 1969 | 6.8 | ... | ... | 0.2 | 3.7 | 8.1 | 12.7 | 13.3 | 11.8 | 9.6 | 7.6 | 4.1 |
| 1968 | 6.4 | ... | ... | * | 3.4 | 7.5 | 11.5 | 13.5 | 11.5 | 7.9 | 6.6 | 4.6 |

... Category not applicable

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| Suicide E950-E959 | | | | | | | | | | | | |
| ALL OTHER, BOTH SEXES | | | | | | | | | | | | |
| 1978 | 6.8 | ... | ... | * | 8.6 | 13.3 | 10.1 | 8.7 | 6.6 | 6.7 | 6.9 | * |
| 1977 | 7.1 | ... | ... | * | 9.4 | 14.9 | 9.8 | 8.3 | 8.2 | 6.4 | 6.5 | * |
| 1976 | 6.9 | ... | ... | 0.4 | 9.1 | 13.5 | 10.0 | 8.7 | 7.0 | 8.3 | 6.0 | 13.1 |
| 1975 | 6.7 | ... | ... | * | 8.9 | 14.4 | 9.8 | 8.5 | 7.3 | 6.9 | 6.4 | * |
| 1974 | 6.4 | ... | ... | * | 8.1 | 13.6 | 9.4 | 7.7 | 7.5 | 7.9 | 7.8 | * |
| 1973 | 6.3 | ... | ... | * | 8.8 | 13.0 | 9.4 | 8.0 | 7.9 | 6.6 | 7.2 | * |
| 1972 | 6.6 | ... | ... | * | 10.9 | 12.5 | 9.9 | 8.2 | 7.4 | 6.1 | 7.7 | * |
| 1971 | 5.9 | ... | ... | 0.3 | 8.2 | 11.6 | 10.0 | 7.0 | 7.1 | 8.1 | 5.7 | * |
| 1970 | 5.6 | ... | ... | * | 7.6 | 12.2 | 8.1 | 9.0 | 6.1 | 6.5 | 6.9 | * |
| 1969 | 5.4 | ... | ... | * | 6.9 | 11.2 | 9.6 | 7.6 | 6.0 | 6.8 | 9.2 | * |
| 1968 | 4.8 | ... | ... | * | 5.6 | 9.6 | 8.0 | 7.7 | 7.1 | 7.0 | 8.0 | * |
| ALL OTHER, MALE | | | | | | | | | | | | |
| 1978 | 10.9 | ... | ... | * | 14.4 | 22.5 | 16.5 | 13.7 | 10.2 | 12.0 | 12.4 | * |
| 1977 | 11.2 | ... | ... | * | 15.1 | 24.4 | 15.6 | 12.1 | 13.0 | 12.0 | 11.9 | * |
| 1976 | 10.9 | ... | ... | * | 14.4 | 21.8 | 16.4 | 13.9 | 12.1 | 13.6 | 12.1 | * |
| 1975 | 10.5 | ... | ... | * | 14.1 | 23.7 | 15.7 | 13.2 | 11.3 | 11.5 | 12.4 | * |
| 1974 | 10.1 | ... | ... | * | 12.6 | 22.2 | 15.3 | 12.2 | 12.4 | 14.7 | 16.2 | * |
| 1973 | 9.9 | ... | ... | * | 13.8 | 22.1 | 14.1 | 13.5 | 12.0 | 11.7 | 12.5 | * |
| 1972 | 10.2 | ... | ... | * | 16.5 | 20.6 | 15.7 | 13.3 | 11.8 | 11.7 | 12.0 | * |
| 1971 | 8.5 | ... | ... | * | 11.5 | 17.4 | 14.9 | 10.3 | 11.8 | 15.0 | * | * |
| 1970 | 8.5 | ... | ... | * | 11.3 | 19.8 | 12.6 | 14.1 | 10.5 | 10.2 | 11.9 | * |
| 1969 | 8.2 | ... | ... | * | 10.0 | 17.6 | 15.5 | 12.0 | 10.2 | 12.2 | 17.6 | * |
| 1968 | 7.4 | ... | ... | * | 8.4 | 15.8 | 12.3 | 12.3 | 12.0 | 12.4 | 12.9 | * |

... Category not applicable

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| Suicide E950-E959 | | | | | | | | | | | | |
| ALL OTHER, FEMALE | | | | | | | | | | | | |
| 1978 | 3.0 | ... | ... | * | 3.1 | 5.4 | 4.7 | 4.6 | 3.6 | 2.7 | * | * |
| 1977 | 3.4 | ... | ... | * | 3.9 | 6.6 | 5.0 | 5.2 | 4.2 | * | * | * |
| 1976 | 3.2 | ... | ... | * | 3.9 | 6.3 | 4.7 | 4.3 | 2.7 | 4.0 | * | * |
| 1975 | 3.2 | ... | ... | * | 3.9 | 6.3 | 4.9 | 4.5 | 4.0 | 3.3 | * | * |
| 1974 | 3.0 | ... | ... | * | 3.8 | 6.2 | 4.5 | 4.0 | 3.3 | * | * | * |
| 1973 | 3.0 | ... | ... | * | 4.1 | 5.3 | 5.4 | 3.2 | 4.3 | * | * | * |
| 1972 | 3.2 | ... | ... | * | 5.6 | 5.6 | 5.2 | 3.8 | 3.7 | * | * | * |
| 1971 | 3.4 | ... | ... | * | 5.2 | 6.7 | 5.9 | 4.1 | 3.0 | * | * | * |
| 1970 | 2.9 | ... | ... | * | 4.1 | 5.8 | 4.3 | 4.5 | 2.2 | 3.5 | * | * |
| 1969 | 2.7 | ... | ... | * | 4.1 | 5.8 | 4.7 | 3.7 | 2.2 | * | * | * |
| 1968 | 2.4 | ... | ... | * | 3.1 | 4.4 | 4.3 | 3.7 | 2.7 | * | * | * |
| BLACK, BOTH SEXES | | | | | | | | | | | | |
| 1978 | 6.5 | ... | ... | * | 7.7 | 13.6 | 10.2 | 8.8 | 6.2 | 6.3 | 6.2 | * |
| 1977 | 6.6 | ... | ... | * | 8.3 | 14.8 | 9.3 | 7.6 | 7.5 | 5.5 | 5.3 | * |
| 1976 | 6.4 | ... | ... | * | 8.2 | 13.4 | 9.7 | 8.5 | 6.5 | 7.8 | 4.3 | * |
| 1975 | 6.1 | ... | ... | * | 7.8 | 13.7 | 9.4 | 7.9 | 6.7 | 6.7 | 5.5 | * |
| 1974 | 5.9 | ... | ... | * | 7.1 | 13.5 | 9.3 | 7.6 | 7.0 | 6.9 | 5.6 | * |
| 1973 | 5.8 | ... | ... | * | 7.9 | 12.8 | 8.7 | 7.9 | 6.9 | 5.8 | 5.4 | * |
| 1972 | 6.0 | ... | ... | * | 9.6 | 12.3 | 9.4 | 8.1 | 6.9 | 5.4 | 4.2 | * |
| 1971 | 5.3 | ... | ... | * | 7.2 | 10.8 | 9.4 | 6.6 | 6.5 | 7.6 | * | * |
| 1970 | 5.2 | ... | ... | * | 7.0 | 11.9 | 7.8 | 8.4 | 5.9 | 5.5 | * | * |
| 1969 | 4.9 | ... | ... | * | 6.5 | 10.2 | 9.5 | 7.0 | 5.2 | 5.9 | 6.4 | * |
| 1968 | 4.3 | ... | ... | * | 5.0 | 9.1 | 7.4 | 7.3 | 6.6 | 6.3 | 5.9 | * |

... Category not applicable

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| Suicide E950-E959 | | | | | | | | | | | | |
| BLACK, MALE | | | | | | | | | | | | |
| 1978 | 10.7 | ... | ... | * | 13.0 | 23.1 | 16.9 | 14.5 | 10.1 | 11.6 | 11.7 | * |
| 1977 | 10.6 | ... | ... | * | 13.0 | 25.1 | 14.8 | 12.1 | 12.5 | 10.8 | 10.6 | * |
| 1976 | 10.4 | ... | ... | * | 13.0 | 22.1 | 16.4 | 14.0 | 11.1 | 13.5 | 9.4 | * |
| 1975 | 9.9 | ... | ... | * | 12.7 | 23.4 | 16.0 | 12.4 | 10.7 | 11.6 | 11.7 | * |
| 1974 | 9.7 | ... | ... | * | 11.1 | 22.8 | 16.1 | 12.7 | 11.7 | 13.2 | 12.9 | * |
| 1973 | 9.4 | ... | ... | * | 12.7 | 21.9 | 13.6 | 13.4 | 11.4 | 11.3 | 12.3 | * |
| 1972 | 9.4 | ... | ... | * | 14.7 | 20.3 | 15.0 | 12.9 | 11.1 | 11.1 | * | * |
| 1971 | 7.8 | ... | ... | * | 9.7 | 16.6 | 14.6 | 10.0 | 10.8 | 14.3 | * | * |
| 1970 | 8.0 | ... | ... | * | 10.5 | 19.2 | 12.6 | 13.8 | 10.6 | 8.7 | * | * |
| 1969 | 7.6 | ... | ... | * | 9.2 | 16.6 | 15.8 | 11.1 | 9.5 | 11.2 | 12.6 | * |
| 1968 | 6.9 | ... | ... | * | 7.2 | 15.4 | 11.8 | 12.1 | 11.5 | 11.7 | 12.1 | * |
| BLACK, FEMALE | | | | | | | | | | | | |
| 1978 | 2.7 | ... | ... | * | 2.7 | 5.4 | 4.6 | 4.2 | 3.0 | * | * | * |
| 1977 | 3.0 | ... | ... | * | 3.7 | 6.0 | 4.8 | 4.0 | 3.4 | * | * | * |
| 1976 | 2.9 | ... | ... | * | 3.7 | 5.9 | 4.1 | 4.0 | 2.7 | 3.5 | * | * |
| 1975 | 2.7 | ... | ... | * | 3.2 | 5.4 | 4.0 | 4.0 | 3.4 | 3.0 | * | * |
| 1974 | 2.5 | ... | ... | * | 3.4 | 5.4 | 3.8 | 3.4 | 3.1 | * | * | * |
| 1973 | 2.5 | ... | ... | * | 3.3 | 5.0 | 4.5 | 3.2 | 3.2 | * | * | * |
| 1972 | 2.9 | ... | ... | * | 4.7 | 5.5 | 4.7 | 4.1 | 3.3 | * | * | * |
| 1971 | 3.0 | ... | ... | * | 4.8 | 6.0 | 5.1 | 3.6 | 2.8 | * | * | * |
| 1970 | 2.6 | ... | ... | * | 3.8 | 5.7 | 3.7 | 3.7 | * | * | * | * |
| 1969 | 2.4 | ... | ... | * | 4.1 | 4.8 | 4.2 | 3.5 | * | * | * | * |
| 1968 | 2.0 | ... | ... | * | 2.9 | 4.0 | 3.7 | 3.2 | 2.4 | * | * | * |

... Category not applicable

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| HOMICIDE E960-E978 | | | | | | | | | | | | |
| BOTH SEXES | | | | | | | | | | | | |
| 1978 | 9.2 | 4.8 | 2.5 | 1.3 | 12.9 | 16.4 | 14.0 | 10.0 | 6.6 | 5.0 | 4.4 | 4.7 |
| 1977 | 9.1 | 5.4 | 2.7 | 1.2 | 12.4 | 16.0 | 14.4 | 9.8 | 7.2 | 5.2 | 4.6 | 4.9 |
| 1976 | 9.0 | 5.5 | 2.4 | 1.0 | 12.2 | 16.2 | 14.3 | 10.0 | 7.2 | 5.3 | 5.1 | 5.1 |
| 1975 | 9.9 | 5.6 | 2.4 | 1.0 | 13.5 | 18.1 | 15.8 | 11.6 | 7.9 | 6.0 | 5.4 | 4.8 |
| 1974 | 10.1 | 5.4 | 2.2 | 1.0 | 14.0 | 19.0 | 16.4 | 11.7 | 7.9 | 5.8 | 5.1 | 4.3 |
| 1973 | 9.7 | 5.1 | 2.5 | 1.1 | 13.3 | 18.8 | 15.9 | 11.1 | 7.8 | 5.8 | 4.9 | 3.4 |
| 1972 | 9.4 | 5.2 | 1.8 | 0.9 | 13.4 | 18.3 | 16.2 | 11.0 | 7.8 | 4.6 | 3.9 | 4.8 |
| 1971 | 9.1 | 5.2 | 2.2 | 1.0 | 12.5 | 18.5 | 15.6 | 10.4 | 7.4 | 5.1 | 4.2 | 4.6 |
| 1970 | 8.3 | 4.3 | 1.9 | 0.9 | 11.7 | 16.6 | 13.7 | 10.1 | 7.1 | 5.0 | 4.0 | 4.2 |
| 1969 | 7.7 | 4.4 | 1.7 | 0.7 | 11.0 | 15.9 | 13.4 | 8.8 | 6.4 | 4.6 | 2.9 | 2.9 |
| 1968 | 7.4 | 4.8 | 1.5 | 0.7 | 10.1 | 15.3 | 12.9 | 9.1 | 6.3 | 4.4 | 3.4 | 3.7 |
| MALE | | | | | | | | | | | | |
| 1978 | 14.7 | 4.8 | 2.7 | 1.4 | 19.9 | 26.9 | 22.9 | 16.6 | 11.1 | 7.9 | 5.9 | 6.9 |
| 1977 | 14.4 | 5.7 | 2.9 | 1.2 | 19.0 | 26.1 | 23.5 | 16.2 | 11.9 | 8.4 | 6.8 | 8.3 |
| 1976 | 14.3 | 5.5 | 2.4 | 1.2 | 18.8 | 26.4 | 23.7 | 16.5 | 11.9 | 8.9 | 6.7 | 6.8 |
| 1975 | 15.8 | 6.3 | 2.8 | 1.1 | 20.9 | 29.7 | 25.6 | 19.4 | 13.4 | 10.0 | 7.9 | 7.2 |
| 1974 | 16.1 | 5.0 | 2.4 | 1.3 | 21.8 | 30.9 | 26.9 | 19.8 | 13.4 | 9.5 | 8.0 | 6.4 |
| 1973 | 15.4 | 5.1 | 2.7 | 1.3 | 20.5 | 30.8 | 25.3 | 18.3 | 13.4 | 9.6 | 7.2 | 5.6 |
| 1972 | 15.3 | 5.4 | 2.0 | 1.0 | 21.3 | 30.5 | 27.3 | 18.7 | 13.3 | 7.6 | 5.5 | 8.4 |
| 1971 | 14.7 | 5.0 | 2.3 | 1.1 | 20.5 | 30.7 | 25.5 | 17.3 | 12.2 | 8.7 | 6.8 | 6.1 |
| 1970 | 13.4 | 4.5 | 1.9 | 1.0 | 19.0 | 27.6 | 22.2 | 17.0 | 12.2 | 8.6 | 6.0 | 7.4 |
| 1969 | 12.4 | 5.0 | 1.8 | 0.8 | 18.0 | 26.1 | 22.1 | 14.5 | 10.7 | 7.9 | 4.6 | * |
| 1968 | 11.9 | 4.7 | 1.5 | 0.9 | 16.4 | 25.3 | 20.8 | 15.1 | 10.5 | 7.6 | 5.4 | 5.9 |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| HOMICIDE E960-E978 | | | | | | | | | | | | |
| FEMALE | | | | | | | | | | | | |
| 1978 | 4.0 | 4.9 | 2.3 | 1.1 | 5.8 | 6.2 | 5.4 | 3.7 | 2.6 | 2.7 | 3.4 | 3.7 |
| 1977 | 4.1 | 5.1 | 2.4 | 1.1 | 5.8 | 6.2 | 5.7 | 3.8 | 3.1 | 2.7 | 3.3 | 3.3 |
| 1976 | 3.9 | 5.4 | 2.5 | 0.8 | 5.5 | 6.1 | 5.3 | 4.0 | 3.0 | 2.6 | 4.1 | 4.3 |
| 1975 | 4.3 | 4.9 | 2.0 | 0.9 | 6.1 | 6.8 | 6.4 | 4.5 | 3.1 | 2.8 | 3.9 | 3.6 |
| 1974 | 4.3 | 5.9 | 2.0 | 0.8 | 6.2 | 7.4 | 6.5 | 4.2 | 3.0 | 3.1 | 3.3 | 3.3 |
| 1973 | 4.3 | 5.2 | 2.3 | 0.9 | 6.1 | 7.0 | 6.9 | 4.3 | 2.9 | 2.8 | 3.5 | 2.3 |
| 1972 | 3.7 | 4.9 | 1.6 | 0.8 | 5.4 | 6.4 | 5.6 | 3.9 | 2.9 | 2.2 | 2.9 | 2.9 |
| 1971 | 3.7 | 5.4 | 2.0 | 0.9 | 4.6 | 6.6 | 6.2 | 4.0 | 3.1 | 2.3 | 2.5 | 3.8 |
| 1970 | 3.4 | 4.1 | 1.9 | 0.7 | 4.6 | 6.0 | 5.7 | 3.7 | 2.5 | 2.2 | 2.7 | 2.5 |
| 1969 | 3.2 | 3.8 | 1.6 | 0.6 | 4.2 | 6.2 | 5.2 | 3.5 | 2.5 | 2.0 | 1.8 | * |
| 1968 | 3.1 | 5.0 | 1.5 | 0.6 | 4.0 | 5.7 | 5.4 | 3.3 | 2.4 | 1.8 | 2.1 | 2.5 |
| WHITE, BOTH SEXES | | | | | | | | | | | | |
| 1978 | 5.8 | 3.4 | 1.5 | 0.9 | 8.1 | 9.4 | 8.7 | 6.6 | 4.6 | 3.6 | 3.7 | 4.2 |
| 1977 | 5.6 | 4.0 | 1.7 | 0.9 | 7.6 | 9.0 | 8.8 | 6.1 | 4.9 | 3.8 | 3.8 | 4.1 |
| 1976 | 5.4 | 3.8 | 1.6 | 0.8 | 7.0 | 8.7 | 8.3 | 6.1 | 4.6 | 3.7 | 4.2 | 4.6 |
| 1975 | 5.9 | 4.2 | 1.5 | 0.8 | 7.5 | 9.6 | 9.2 | 7.0 | 5.4 | 4.3 | 4.7 | 4.0 |
| 1974 | 5.7 | 3.6 | 1.5 | 0.8 | 7.5 | 9.9 | 9.0 | 6.7 | 5.0 | 4.1 | 4.3 | 3.8 |
| 1973 | 5.4 | 3.8 | 1.7 | 0.8 | 7.1 | 9.5 | 8.3 | 6.2 | 5.1 | 4.0 | 4.2 | 3.1 |
| 1972 | 4.9 | 3.7 | 1.3 | 0.6 | 6.4 | 8.5 | 8.0 | 6.0 | 4.9 | 3.2 | 3.3 | 4.7 |
| 1971 | 4.7 | 4.0 | 1.4 | 0.6 | 5.5 | 8.7 | 7.7 | 5.7 | 4.8 | 3.8 | 3.5 | 4.0 |
| 1970 | 4.4 | 2.9 | 1.3 | 0.5 | 5.3 | 8.1 | 7.1 | 5.5 | 4.7 | 3.4 | 3.5 | 3.5 |
| 1969 | 4.0 | 3.5 | 1.4 | 0.5 | 5.0 | 7.4 | 6.3 | 4.6 | 4.2 | 3.2 | 2.7 | 2.6 |
| 1968 | 3.9 | 4.0 | 1.1 | 0.5 | 4.8 | 6.9 | 6.3 | 5.0 | 4.0 | 3.3 | 2.9 | 3.3 |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| HOMICIDE E960-E978 | | | | | | | | | | | | |
| WHITE, MALE | | | | | | | | | | | | |
| 1978 | 9.0 | 3.4 | 1.7 | 1.0 | 12.2 | 14.9 | 13.9 | 10.4 | 7.5 | 5.5 | 4.4 | 6.6 |
| 1977 | 8.6 | 4.0 | 1.8 | 0.9 | 11.3 | 14.1 | 13.8 | 9.7 | 7.7 | 5.6 | 5.2 | 6.1 |
| 1976 | 8.2 | 3.8 | 1.7 | 0.9 | 10.4 | 13.9 | 13.6 | 9.6 | 7.2 | 5.8 | 5.3 | 5.6 |
| 1975 | 9.0 | 4.9 | 1.9 | 0.8 | 11.0 | 15.3 | 14.4 | 11.3 | 8.8 | 7.0 | 6.4 | 5.7 |
| 1974 | 8.8 | 3.6 | 1.7 | 1.0 | 11.3 | 15.6 | 14.1 | 10.9 | 8.0 | 6.2 | 6.5 | 5.4 |
| 1973 | 8.2 | 4.0 | 1.8 | 0.8 | 10.4 | 15.2 | 12.7 | 9.8 | 8.5 | 6.3 | 5.6 | 5.1 |
| 1972 | 7.6 | 4.1 | 1.5 | 0.6 | 9.5 | 13.8 | 13.0 | 9.6 | 8.1 | 5.2 | 4.5 | 8.8 |
| 1971 | 7.3 | 3.7 | 1.5 | 0.6 | 8.6 | 13.9 | 12.5 | 9.1 | 7.6 | 6.2 | 5.2 | 4.5 |
| 1970 | 6.8 | 2.9 | 1.4 | 0.5 | 7.9 | 13.0 | 11.0 | 9.0 | 7.7 | 5.6 | 5.1 | 6.4 |
| 1969 | 6.1 | 3.4 | 1.4 | 0.5 | 7.6 | 11.6 | 10.1 | 7.1 | 6.9 | 5.2 | 4.2 | * |
| 1968 | 6.0 | 3.9 | 1.0 | 0.5 | 7.5 | 11.1 | 9.8 | 8.0 | 6.4 | 5.5 | 4.4 | 5.0 |
| WHITE, FEMALE | | | | | | | | | | | | |
| 1978 | 2.8 | 3.4 | 1.4 | 0.8 | 4.0 | 3.9 | 3.6 | 3.0 | 1.9 | 2.1 | 3.2 | 3.1 |
| 1977 | 2.9 | 3.9 | 1.6 | 0.9 | 3.9 | 3.8 | 4.0 | 2.7 | 2.4 | 2.3 | 3.0 | 3.3 |
| 1976 | 2.6 | 3.9 | 1.5 | 0.7 | 3.5 | 3.5 | 3.2 | 2.8 | 2.2 | 2.2 | 3.6 | 4.2 |
| 1975 | 2.9 | 3.4 | 1.2 | 0.8 | 4.0 | 4.0 | 4.0 | 3.0 | 2.3 | 2.3 | 3.6 | 3.2 |
| 1974 | 2.8 | 3.6 | 1.3 | 0.6 | 3.7 | 4.3 | 3.9 | 2.8 | 2.3 | 2.6 | 3.0 | 2.9 |
| 1973 | 2.7 | 3.6 | 1.6 | 0.7 | 3.8 | 3.8 | 4.1 | 2.8 | 2.1 | 2.3 | 3.3 | 2.1 |
| 1972 | 2.3 | 3.3 | 1.2 | 0.6 | 3.2 | 3.3 | 3.2 | 2.6 | 2.0 | 1.7 | 2.6 | 2.5 |
| 1971 | 2.3 | 4.3 | 1.3 | 0.6 | 2.5 | 3.6 | 3.2 | 2.5 | 2.3 | 1.9 | 2.5 | 3.7 |
| 1970 | 2.1 | 2.9 | 1.2 | 0.5 | 2.7 | 3.4 | 3.2 | 2.2 | 2.0 | 1.7 | 2.5 | * |
| 1969 | 2.0 | 3.5 | 1.3 | 0.6 | 2.4 | 3.3 | 2.6 | 2.3 | 1.8 | 1.7 | 1.7 | * |
| 1968 | 1.9 | 4.2 | 1.2 | 0.5 | 2.2 | 2.8 | 2.9 | 2.2 | 1.8 | 1.5 | 1.9 | * |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| HOMICIDE E960-E978 | | | | | | | | | | | | |
| ALL OTHER, BOTH SEXES | | | | | | | | | | | | |
| 1978 | 30.6 | 11.7 | 7.2 | 2.9 | 39.4 | 61.8 | 49.7 | 35.9 | 24.7 | 18.0 | 11.7 | * |
| 1977 | 31.3 | 12.3 | 7.3 | 2.6 | 39.6 | 62.9 | 52.3 | 38.3 | 28.4 | 18.5 | 12.9 | 13.1 |
| 1976 | 32.7 | 13.4 | 6.4 | 2.2 | 42.1 | 66.1 | 54.6 | 42.1 | 31.4 | 20.6 | 13.6 | * |
| 1975 | 36.6 | 13.0 | 6.8 | 2.2 | 48.7 | 75.8 | 61.5 | 49.9 | 31.3 | 21.8 | 13.8 | 13.7 |
| 1974 | 39.2 | 14.4 | 5.7 | 2.4 | 52.7 | 81.9 | 68.8 | 53.5 | 34.7 | 22.7 | 13.7 | * |
| 1973 | 38.8 | 11.8 | 6.7 | 2.8 | 51.0 | 84.4 | 69.6 | 52.2 | 33.6 | 22.9 | 13.6 | * |
| 1972 | 40.3 | 12.9 | 4.2 | 2.8 | 56.7 | 87.9 | 74.7 | 54.4 | 35.2 | 18.0 | 10.4 | * |
| 1971 | 39.5 | 11.6 | 6.2 | 2.8 | 57.2 | 87.4 | 73.3 | 52.1 | 32.5 | 18.1 | 12.1 | * |
| 1970 | 35.5 | 11.8 | 5.2 | 2.9 | 53.0 | 75.9 | 62.6 | 50.8 | 30.0 | 20.6 | 10.3 | * |
| 1969 | 34.2 | 9.7 | 3.5 | 1.9 | 50.8 | 77.1 | 67.2 | 46.4 | 27.6 | 17.9 | 6.5 | * |
| 1968 | 32.4 | 9.1 | 3.5 | 2.1 | 45.9 | 75.6 | 63.6 | 45.6 | 28.4 | 15.6 | 10.1 | * |
| ALL OTHER, MALE | | | | | | | | | | | | |
| 1978 | 51.7 | 11.8 | 7.8 | 3.2 | 64.0 | 110.1 | 89.2 | 68.1 | 44.7 | 30.3 | 19.9 | * |
| 1977 | 52.8 | 14.1 | 8.2 | 2.9 | 63.7 | 111.5 | 94.8 | 70.9 | 51.3 | 33.6 | 22.2 | * |
| 1976 | 55.1 | 14.3 | 5.9 | 3.0 | 68.5 | 117.2 | 98.1 | 76.4 | 56.3 | 37.9 | 20.2 | * |
| 1975 | 61.9 | 13.5 | 7.3 | 2.6 | 80.3 | 135.4 | 109.3 | 89.8 | 56.8 | 38.4 | 23.1 | * |
| 1974 | 66.5 | 12.0 | 6.0 | 2.8 | 86.7 | 145.4 | 123.5 | 98.6 | 64.4 | 40.8 | 23.1 | * |
| 1973 | 65.3 | 10.5 | 7.3 | 3.9 | 83.9 | 150.0 | 122.4 | 93.6 | 61.1 | 40.7 | 24.1 | * |
| 1972 | 69.7 | 12.5 | 4.5 | 3.7 | 96.9 | 158.4 | 138.6 | 100.0 | 63.6 | 31.3 | 16.6 | * |
| 1971 | 67.5 | 11.8 | 6.7 | 3.5 | 98.9 | 158.1 | 129.6 | 92.3 | 56.9 | 33.0 | 25.1 | * |
| 1970 | 60.8 | 13.4 | 4.6 | 3.9 | 92.0 | 137.3 | 110.8 | 90.3 | 55.1 | 37.0 | 17.1 | * |
| 1969 | 58.7 | 14.0 | 4.1 | 2.8 | 88.7 | 137.3 | 120.2 | 83.2 | 48.2 | 33.7 | * | * |
| 1968 | 55.1 | 9.1 | 3.8 | 2.9 | 78.2 | 135.6 | 112.0 | 81.9 | 50.9 | 28.5 | 17.2 | * |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| HOMICIDE E960-E978 | | | | | | | | | | | | |
| ALL OTHER, FEMALE | | | | | | | | | | | | |
| 1978 | 11.5 | 11.7 | 6.6 | 2.6 | 15.6 | 20.2 | 16.7 | 9.3 | 8.2 | 8.5 | 6.2 | * |
| 1977 | 11.8 | 10.5 | 6.3 | 2.3 | 16.5 | 21.1 | 16.7 | 11.3 | 9.2 | 6.6 | 6.5 | * |
| 1976 | 12.3 | 12.6 | 7.0 | 1.5 | 16.6 | 22.1 | 18.2 | 13.5 | 10.5 | 6.9 | 9.0 | * |
| 1975 | 13.6 | 12.4 | 6.3 | 1.8 | 18.5 | 24.4 | 21.5 | 16.2 | 9.7 | 8.6 | 7.3 | * |
| 1974 | 14.3 | 17.0 | 5.4 | 2.0 | 20.2 | 27.4 | 23.0 | 15.1 | 9.3 | 8.2 | 7.2 | * |
| 1973 | 14.5 | 13.1 | 6.1 | 1.7 | 19.6 | 27.9 | 25.5 | 16.8 | 10.2 | 8.4 | 6.3 | * |
| 1972 | 13.3 | 13.4 | 3.8 | 2.0 | 18.4 | 27.4 | 21.3 | 15.1 | 10.8 | 7.3 | * | * |
| 1971 | 13.8 | 11.4 | 5.8 | 2.2 | 17.8 | 27.0 | 26.5 | 17.1 | 11.1 | 5.9 | * | * |
| 1970 | 12.3 | 10.2 | 5.8 | 1.9 | 16.3 | 23.3 | 22.6 | 16.1 | 7.8 | 7.1 | * | * |
| 1969 | 11.7 | * | 3.0 | 1.0 | 15.6 | 26.0 | 23.2 | 14.0 | 9.4 | 4.9 | * | * |
| 1968 | 11.6 | 9.2 | 3.2 | 1.3 | 15.9 | 24.8 | 23.0 | 13.4 | 8.2 | 4.8 | * | * |
| BLACK, BOTH SEXES | | | | | | | | | | | | |
| 1978 | 34.2 | 13.0 | 8.1 | 3.1 | 43.4 | 71.7 | 57.8 | 40.6 | 27.2 | 19.3 | 12.2 | * |
| 1977 | 34.8 | 13.6 | 7.8 | 2.9 | 43.8 | 72.9 | 60.3 | 42.5 | 31.0 | 19.9 | 13.1 | 14.0 |
| 1976 | 36.3 | 14.5 | 6.9 | 2.3 | 46.5 | 76.3 | 62.3 | 47.1 | 34.2 | 22.4 | 14.1 | * |
| 1975 | 40.6 | 14.3 | 7.2 | 2.4 | 53.8 | 87.4 | 70.4 | 56.2 | 33.9 | 22.9 | 13.5 | * |
| 1974 | 43.0 | 15.4 | 6.2 | 2.5 | 57.9 | 92.9 | 77.8 | 59.1 | 37.3 | 23.9 | 14.1 | * |
| 1973 | 42.5 | 12.6 | 7.1 | 3.1 | 55.7 | 95.9 | 79.0 | 57.5 | 36.0 | 24.5 | 14.2 | * |
| 1972 | 44.0 | 13.0 | 4.5 | 3.1 | 62.1 | 99.3 | 83.6 | 59.7 | 37.8 | 19.2 | 9.2 | * |
| 1971 | 43.1 | 12.5 | 6.5 | 3.1 | 62.5 | 98.6 | 81.8 | 57.2 | 34.2 | 18.7 | 12.6 | * |
| 1970 | 39.1 | 12.5 | 5.7 | 3.1 | 58.6 | 86.4 | 70.9 | 56.0 | 31.9 | 22.2 | 11.3 | * |
| 1969 | 36.9 | 10.3 | 3.7 | 2.1 | 55.0 | 85.1 | 74.6 | 50.2 | 29.1 | 18.5 | 6.4 | * |
| 1968 | 35.0 | 9.5 | 3.8 | 2.2 | 49.8 | 83.7 | 70.2 | 49.1 | 30.2 | 16.0 | 10.1 | * |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| HOMICIDE E960-E978 | | | | | | | | | | | | |
| BLACK, MALE | | | | | | | | | | | | |
| 1978 | 58.1 | 12.9 | 8.7 | 3.4 | 70.7 | 127.8 | 104.5 | 77.2 | 49.5 | 33.6 | 20.9 | * |
| 1977 | 59.1 | 16.3 | 8.8 | 3.2 | 70.5 | 129.3 | 110.6 | 79.3 | 57.1 | 37.1 | 22.1 | * |
| 1976 | 61.6 | 15.3 | 6.3 | 3.1 | 76.1 | 135.6 | 113.6 | 85.8 | 62.3 | 42.1 | 20.9 | * |
| 1975 | 69.0 | 14.8 | 7.9 | 2.7 | 89.0 | 156.9 | 125.5 | 102.0 | 62.3 | 41.2 | 23.0 | * |
| 1974 | 73.3 | 13.3 | 6.6 | 2.9 | 95.4 | 165.5 | 140.1 | 109.7 | 70.7 | 44.2 | 23.4 | * |
| 1973 | 72.1 | 10.7 | 7.7 | 4.3 | 92.2 | 172.0 | 139.7 | 103.4 | 66.4 | 44.8 | 25.0 | * |
| 1972 | 76.6 | 12.4 | 4.6 | 3.9 | 106.7 | 180.3 | 155.6 | 110.4 | 69.0 | 34.1 | 17.3 | * |
| 1971 | 74.0 | 12.2 | 7.0 | 3.8 | 108.2 | 180.1 | 145.3 | 102.1 | 61.1 | 34.5 | 26.9 | * |
| 1970 | 67.6 | 14.3 | 5.1 | 4.2 | 102.5 | 158.5 | 126.2 | 100.5 | 59.8 | 40.6 | 19.0 | * |
| 1969 | 63.8 | 14.8 | 4.3 | 3.1 | 96.3 | 152.7 | 133.8 | 90.7 | 51.6 | 35.4 | * | * |
| 1968 | 60.0 | 9.2 | 4.2 | 3.1 | 85.4 | 151.7 | 123.5 | 89.1 | 55.2 | 29.6 | 17.6 | * |
| BLACK, FEMALE | | | | | | | | | | | | |
| 1978 | 12.8 | 13.2 | 7.6 | 2.7 | 17.2 | 23.1 | 19.0 | 10.5 | 8.9 | 8.7 | 6.4 | * |
| 1977 | 12.9 | 10.8 | 6.7 | 2.6 | 18.2 | 24.3 | 18.5 | 12.0 | 9.7 | 7.0 | 7.1 | * |
| 1976 | 13.4 | 13.7 | 7.6 | 1.5 | 18.3 | 25.1 | 19.7 | 14.8 | 11.0 | 7.4 | 9.5 | * |
| 1975 | 14.9 | 13.7 | 6.6 | 2.0 | 20.3 | 27.6 | 24.6 | 17.8 | 10.4 | 9.1 | 7.0 | * |
| 1974 | 15.5 | 17.6 | 5.9 | 2.1 | 22.2 | 30.6 | 25.8 | 16.6 | 9.6 | 8.4 | 7.8 | * |
| 1973 | 15.6 | 14.4 | 6.5 | 1.9 | 21.1 | 30.9 | 28.3 | 18.4 | 10.7 | 8.9 | 6.8 | * |
| 1972 | 14.3 | 13.5 | 4.3 | 2.2 | 19.8 | 30.3 | 23.7 | 16.4 | 11.7 | 7.6 | * | * |
| 1971 | 14.9 | 12.8 | 6.0 | 2.3 | 19.5 | 29.6 | 29.1 | 18.6 | 11.5 | 6.2 | * | * |
| 1970 | 13.3 | 10.7 | 6.3 | 2.0 | 17.7 | 25.6 | 25.1 | 17.5 | 8.1 | 7.7 | * | * |
| 1969 | 12.5 | * | 3.0 | 1.1 | 16.7 | 28.4 | 25.5 | 15.1 | 9.9 | 5.1 | * | * |
| 1968 | 12.3 | 9.8 | 3.5 | 1.4 | 16.9 | 26.8 | 25.5 | 14.4 | 8.7 | 5.0 | * | * |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| OTHER EXTERNAL CAUSES (E980-E999) | | | | | | | | | | | | |
| BOTH SEXES | | | | | | | | | | | | |
| 1978 | 1.9 | 2.5 | 0.9 | 0.3 | 2.0 | 2.8 | 2.3 | 2.3 | 2.0 | 1.9 | 2.1 | 2.9 |
| 1977 | 2.0 | 3.2 | 1.1 | 0.4 | 2.1 | 2.7 | 2.7 | 2.7 | 2.5 | 2.0 | 2.3 | 3.2 |
| 1976 | 2.2 | 3.3 | 1.2 | 0.4 | 2.3 | 2.8 | 2.9 | 2.9 | 2.7 | 2.3 | 2.9 | 2.9 |
| 1975 | 2.3 | 3.0 | 1.1 | 0.3 | 2.5 | 3.1 | 3.0 | 3.0 | 2.6 | 2.5 | 2.5 | 2.7 |
| 1974 | 2.4 | 3.5 | 1.3 | 0.4 | 2.7 | 3.1 | 3.4 | 3.2 | 2.8 | 2.8 | 2.6 | 3.8 |
| 1973 | 2.4 | 2.9 | 1.1 | 0.4 | 2.5 | 2.9 | 3.3 | 3.3 | 3.2 | 2.9 | 3.1 | 3.7 |
| 1972 | 2.4 | 2.8 | 1.3 | 0.4 | 2.5 | 3.2 | 3.3 | 3.3 | 3.2 | 2.5 | 3.4 | 5.3 |
| 1971 | 2.5 | 3.1 | 1.3 | 0.3 | 2.5 | 3.2 | 3.3 | 3.4 | 3.4 | 2.8 | 3.2 | 4.6 |
| 1970 | 2.7 | 3.6 | 1.3 | 0.4 | 3.1 | 3.4 | 3.6 | 3.6 | 3.4 | 3.5 | 3.4 | 4.0 |
| 1969 | 2.6 | 3.7 | 1.3 | 0.4 | 3.0 | 3.2 | 3.4 | 3.5 | 3.3 | 3.1 | 3.2 | 3.4 |
| 1968 | 2.2 | 3.0 | 1.1 | 0.3 | 2.1 | 2.8 | 3.1 | 3.3 | 3.1 | 2.6 | 2.9 | 3.9 |
| MALE | | | | | | | | | | | | |
| 1978 | 2.7 | 2.7 | 1.2 | 0.4 | 3.0 | 4.0 | 3.3 | 3.1 | 2.7 | 2.9 | 3.7 | 5.5 |
| 1977 | 2.9 | 4.2 | 1.2 | 0.5 | 3.0 | 4.0 | 3.5 | 3.8 | 3.5 | 3.1 | 3.7 | 6.5 |
| 1976 | 3.0 | 3.3 | 1.1 | 0.6 | 3.3 | 4.1 | 3.9 | 3.9 | 3.9 | 3.3 | 5.0 | 4.3 |
| 1975 | 3.2 | 2.9 | 1.3 | 0.5 | 3.7 | 4.5 | 4.1 | 3.9 | 3.7 | 3.8 | 4.1 | 5.6 |
| 1974 | 3.4 | 3.7 | 1.3 | 0.6 | 4.2 | 4.5 | 4.7 | 4.2 | 4.1 | 4.3 | 4.0 | 7.1 |
| 1973 | 3.3 | 3.4 | 1.3 | 0.6 | 3.6 | 4.3 | 4.5 | 4.3 | 4.6 | 4.4 | 5.3 | 6.9 |
| 1972 | 3.4 | 3.2 | 1.2 | 0.6 | 3.8 | 4.5 | 4.6 | 4.4 | 4.6 | 3.9 | 6.1 | 7.7 |
| 1971 | 3.5 | 3.9 | 1.4 | 0.5 | 3.9 | 4.7 | 4.4 | 4.6 | 5.0 | 4.3 | 5.3 | 7.5 |
| 1970 | 3.9 | 3.7 | 1.5 | 0.6 | 4.9 | 5.0 | 5.0 | 4.8 | 5.1 | 5.7 | 5.8 | 5.3 |
| 1969 | 3.7 | 3.7 | 1.3 | 0.5 | 4.8 | 4.9 | 4.9 | 4.8 | 5.2 | 4.9 | 4.9 | 5.4 |
| 1968 | 3.1 | 3.5 | 1.1 | 0.4 | 3.2 | 4.1 | 4.2 | 4.4 | 4.8 | 4.0 | 5.0 | 6.1 |

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| OTHER EXTERNAL CAUSES (E980-E999) | | | | | | | | | | | | |
| FEMALE | | | | | | | | | | | | |
| 1978 | 1.1 | 2.3 | 0.6 | 0.1 | 1.0 | 1.6 | 1.4 | 1.6 | 1.4 | 1.2 | 1.2 | 1.7 |
| 1977 | 1.2 | 2.2 | 0.9 | 0.2 | 1.1 | 1.4 | 1.8 | 1.7 | 1.7 | 1.2 | 1.4 | 1.6 |
| 1976 | 1.4 | 3.3 | 1.2 | 0.2 | 1.2 | 1.5 | 2.0 | 2.0 | 1.6 | 1.6 | 1.6 | 2.2 |
| 1975 | 1.4 | 3.1 | 0.9 | 0.2 | 1.3 | 1.6 | 2.0 | 2.2 | 1.6 | 1.5 | 1.6 | * |
| 1974 | 1.5 | 3.3 | 1.2 | 0.2 | 1.2 | 1.8 | 2.1 | 2.2 | 1.7 | 1.6 | 1.8 | 2.1 |
| 1973 | 1.5 | 2.4 | 0.9 | 0.2 | 1.3 | 1.6 | 2.2 | 2.4 | 1.9 | 1.8 | 1.7 | 2.2 |
| 1972 | 1.5 | 2.5 | 1.3 | 0.2 | 1.3 | 1.9 | 2.1 | 2.3 | 2.0 | 1.5 | 1.7 | 4.1 |
| 1971 | 1.5 | 2.3 | 1.2 | 0.2 | 1.2 | 1.7 | 2.3 | 2.4 | 1.9 | 1.7 | 1.9 | 3.2 |
| 1970 | 1.5 | 3.6 | 1.1 | 0.2 | 1.3 | 1.9 | 2.2 | 2.5 | 1.9 | 1.8 | 1.9 | 3.2 |
| 1969 | 1.5 | 3.8 | 1.3 | 0.2 | 1.3 | 1.6 | 2.0 | 2.3 | 1.7 | 1.7 | 2.0 | 2.4 |
| 1968 | 1.3 | 2.4 | 1.0 | 0.2 | 1.0 | 1.5 | 2.0 | 2.1 | 1.6 | 1.4 | 1.4 | 2.8 |
| WHITE, BOTH SEXES | | | | | | | | | | | | |
| 1978 | 1.7 | 1.7 | 0.6 | 0.2 | 1.8 | 2.4 | 2.0 | 2.0 | 2.0 | 1.8 | 1.9 | 2.8 |
| 1977 | 1.8 | 2.1 | 0.8 | 0.3 | 1.9 | 2.2 | 2.3 | 2.4 | 2.4 | 1.8 | 2.2 | 3.1 |
| 1976 | 1.9 | 2.2 | 0.7 | 0.3 | 2.0 | 2.2 | 2.4 | 2.4 | 2.5 | 2.2 | 2.7 | 2.8 |
| 1975 | 2.0 | 1.9 | 0.7 | 0.3 | 2.2 | 2.5 | 2.5 | 2.7 | 2.4 | 2.3 | 2.4 | 2.6 |
| 1974 | 2.1 | 2.7 | 0.8 | 0.3 | 2.3 | 2.5 | 2.7 | 2.8 | 2.5 | 2.5 | 2.5 | 3.6 |
| 1973 | 2.0 | 2.3 | 0.8 | 0.3 | 2.1 | 2.3 | 2.6 | 2.9 | 2.9 | 2.8 | 2.9 | 3.4 |
| 1972 | 2.0 | 1.7 | 0.8 | 0.3 | 2.0 | 2.5 | 2.6 | 2.9 | 3.0 | 2.3 | 3.3 | 5.4 |
| 1971 | 2.0 | 2.1 | 0.8 | 0.3 | 1.9 | 2.3 | 2.6 | 3.1 | 3.1 | 2.7 | 3.1 | 4.7 |
| 1970 | 2.2 | 2.4 | 0.8 | 0.3 | 2.3 | 2.4 | 2.8 | 3.1 | 3.3 | 3.3 | 3.4 | 4.1 |
| 1969 | 2.1 | 2.7 | 0.8 | 0.3 | 2.4 | 2.4 | 2.5 | 3.0 | 3.1 | 2.9 | 3.1 | 3.3 |
| 1968 | 1.8 | 2.0 | 0.7 | 0.3 | 1.7 | 2.1 | 2.4 | 2.9 | 3.0 | 2.4 | 2.9 | 3.5 |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| OTHER EXTERNAL CAUSES (E980-E999) | | | | | | | | | | | | |
| WHITE, MALE | | | | | | | | | | | | |
| 1978 | 2.3 | 1.8 | 0.8 | 0.3 | 2.7 | 3.5 | 2.7 | 2.5 | 2.6 | 2.7 | 3.5 | 5.7 |
| 1977 | 2.5 | 2.4 | 0.8 | 0.4 | 2.8 | 3.2 | 2.9 | 3.3 | 3.2 | 2.7 | 3.5 | 6.5 |
| 1976 | 2.6 | 2.3 | 0.7 | 0.5 | 3.0 | 3.3 | 3.1 | 3.1 | 3.6 | 3.1 | 4.7 | 4.0 |
| 1975 | 2.7 | 1.9 | 0.9 | 0.4 | 3.4 | 3.6 | 3.2 | 3.2 | 3.3 | 3.5 | 3.7 | 5.4 |
| 1974 | 2.9 | 2.9 | 0.9 | 0.5 | 3.6 | 3.5 | 3.5 | 3.6 | 3.5 | 3.7 | 3.8 | 6.6 |
| 1973 | 2.8 | 2.6 | 0.9 | 0.5 | 3.1 | 3.3 | 3.5 | 3.5 | 4.0 | 4.0 | 5.1 | 5.7 |
| 1972 | 2.8 | 1.7 | 0.7 | 0.4 | 3.1 | 3.3 | 3.4 | 3.7 | 4.2 | 3.6 | 6.0 | 7.6 |
| 1971 | 2.8 | 2.6 | 0.9 | 0.4 | 2.9 | 3.4 | 3.2 | 3.9 | 4.4 | 4.0 | 5.1 | 7.3 |
| 1970 | 3.1 | 2.3 | 1.0 | 0.4 | 3.7 | 3.4 | 3.8 | 3.9 | 4.7 | 5.3 | 5.7 | 6.0 |
| 1969 | 3.0 | 2.6 | 0.7 | 0.4 | 3.9 | 3.4 | 3.4 | 3.8 | 4.7 | 4.6 | 4.6 | 5.3 |
| 1968 | 2.6 | 2.5 | 0.7 | 0.4 | 2.7 | 3.0 | 3.1 | 3.8 | 4.5 | 3.7 | 5.1 | 6.0 |
| WHITE, FEMALE | | | | | | | | | | | | |
| 1978 | 1.0 | 1.6 | * | * | 0.9 | 1.4 | 1.3 | 1.6 | 1.4 | 1.1 | 1.0 | * |
| 1977 | 1.1 | 1.8 | 0.7 | * | 0.9 | 1.2 | 1.7 | 1.5 | 1.7 | 1.1 | 1.4 | * |
| 1976 | 1.2 | 2.1 | 0.7 | 0.2 | 1.0 | 1.2 | 1.8 | 1.8 | 1.5 | 1.5 | 1.5 | 2.2 |
| 1975 | 1.3 | 1.9 | 0.5 | 0.2 | 1.1 | 1.4 | 1.8 | 2.1 | 1.5 | 1.4 | 1.5 | * |
| 1974 | 1.3 | 2.6 | 0.7 | * | 1.0 | 1.5 | 1.8 | 2.1 | 1.6 | 1.5 | 1.7 | 2.2 |
| 1973 | 1.3 | 2.0 | 0.7 | 0.1 | 1.0 | 1.4 | 1.8 | 2.3 | 1.8 | 1.8 | 1.5 | 2.3 |
| 1972 | 1.3 | 1.6 | 0.8 | 0.2 | 0.9 | 1.7 | 1.9 | 2.1 | 2.0 | 1.3 | 1.6 | 4.2 |
| 1971 | 1.3 | 1.7 | 0.7 | 0.1 | 1.0 | 1.3 | 2.0 | 2.3 | 1.8 | 1.6 | 1.8 | 3.3 |
| 1970 | 1.4 | 2.6 | 0.6 | 0.1 | 0.9 | 1.5 | 1.9 | 2.4 | 2.0 | 1.7 | 1.9 | 3.1 |
| 1969 | 1.3 | 2.8 | 0.8 | 0.1 | 0.9 | 1.4 | 1.6 | 2.3 | 1.6 | 1.6 | 2.0 | * |
| 1968 | 1.1 | 1.6 | 0.7 | 0.1 | 0.8 | 1.2 | 1.7 | 2.0 | 1.6 | 1.4 | 1.4 | * |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL 1 YEAR | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-----------------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| OTHER EXTERNAL CAUSES (E980-E999) | | | | | | | | | | | | |
| ALL OTHER, BOTH SEXES | | | | | | | | | | | | |
| 1978 | 3.2 | 6.3 | 2.5 | 0.6 | 2.9 | 4.8 | 4.7 | 4.7 | 2.6 | 3.0 | 4.1 | * |
| 1977 | 3.6 | 8.4 | 2.5 | 0.8 | 3.2 | 5.9 | 5.0 | 4.9 | 3.9 | 4.3 | * | * |
| 1976 | 4.2 | 8.7 | 3.5 | 0.8 | 3.9 | 6.3 | 6.4 | 6.8 | 4.3 | 3.7 | 5.1 | * |
| 1975 | 4.2 | 8.4 | 3.2 | 0.6 | 4.0 | 6.8 | 6.3 | 6.0 | 4.4 | 4.5 | 4.4 | * |
| 1974 | 4.9 | 7.4 | 3.4 | 0.8 | 5.1 | 7.3 | 8.5 | 5.8 | 5.6 | 6.0 | 4.5 | * |
| 1973 | 4.8 | 5.9 | 2.6 | 1.1 | 4.9 | 7.1 | 8.2 | 7.0 | 6.1 | 4.5 | 5.5 | * |
| 1972 | 5.1 | 8.9 | 3.9 | 0.9 | 5.9 | 8.1 | 8.3 | 7.1 | 5.0 | 4.4 | 4.6 | * |
| 1971 | 5.4 | 8.3 | 4.0 | 0.8 | 6.4 | 8.9 | 8.7 | 6.7 | 6.3 | 4.4 | 4.3 | * |
| 1970 | 6.0 | 10.0 | 3.9 | 1.0 | 8.1 | 10.1 | 9.1 | 8.1 | 5.0 | 5.5 | 4.5 | * |
| 1969 | 5.9 | 9.7 | 4.0 | 1.0 | 7.0 | 9.2 | 10.5 | 8.0 | 6.0 | 4.8 | 4.8 | * |
| 1968 | 4.5 | 8.0 | 3.1 | 0.6 | 4.5 | 7.7 | 8.6 | 6.8 | 4.7 | 4.1 | * | * |
| ALL OTHER, MALE | | | | | | | | | | | | |
| 1978 | 4.7 | 6.9 | 3.0 | 0.9 | 4.3 | 7.2 | 7.8 | 7.9 | 4.2 | 4.6 | * | * |
| 1977 | 5.4 | 12.7 | 3.1 | 1.0 | 4.3 | 9.3 | 8.3 | 7.8 | 6.7 | 6.5 | * | * |
| 1976 | 6.0 | 8.6 | 3.2 | 1.1 | 5.4 | 9.7 | 9.7 | 10.9 | 6.5 | 5.4 | 7.8 | * |
| 1975 | 6.3 | 7.9 | 3.4 | 0.8 | 6.0 | 11.2 | 10.2 | 9.9 | 7.0 | 6.8 | * | * |
| 1974 | 7.3 | 7.7 | 3.1 | 1.1 | 7.6 | 11.8 | 13.9 | 9.3 | 9.5 | 10.0 | * | * |
| 1973 | 7.3 | 7.5 | 3.3 | 1.6 | 6.8 | 11.6 | 12.5 | 11.5 | 10.3 | 8.3 | * | * |
| 1972 | 7.5 | 11.0 | 3.6 | 1.4 | 8.2 | 13.5 | 13.3 | 10.3 | 9.0 | 6.0 | * | * |
| 1971 | 8.3 | 10.8 | 4.3 | 1.1 | 10.3 | 14.6 | 13.9 | 11.0 | 10.6 | 7.0 | * | * |
| 1970 | 9.4 | 11.2 | 4.4 | 1.4 | 13.2 | 16.5 | 14.9 | 13.3 | 8.7 | 9.6 | * | * |
| 1969 | 9.2 | 10.1 | 4.5 | 1.2 | 10.7 | 16.4 | 17.6 | 14.0 | 10.3 | 7.3 | * | * |
| 1968 | 6.9 | 9.1 | 3.2 | 0.9 | 7.3 | 12.7 | 13.7 | 10.7 | 8.3 | 7.0 | * | * |
| ALL OTHER, FEMALE | | | | | | | | | | | | |
| 1978 | 1.8 | * | 2.1 | * | 1.6 | 2.7 | 2.2 | 2.0 | * | * | * | * |
| 1977 | 2.0 | * | 1.9 | 0.7 | 2.2 | 3.0 | 2.3 | 2.6 | * | 2.6 | * | * |
| 1976 | 2.7 | 8.8 | 3.7 | * | 2.5 | 3.3 | 3.6 | 3.5 | 2.4 | * | * | * |
| 1975 | 2.3 | 8.9 | 3.0 | * | 2.2 | 3.1 | 3.1 | 2.6 | 2.3 | 2.8 | * | * |
| 1974 | 2.6 | * | 3.6 | * | 2.7 | 3.4 | 4.1 | 2.8 | 2.3 | * | * | * |
| 1973 | 2.6 | * | 2.0 | * | 3.0 | 3.2 | 4.6 | 3.2 | 2.5 | * | * | * |
| 1972 | 2.9 | * | 4.2 | * | 3.8 | 3.4 | 4.1 | 4.4 | * | 3.1 | * | * |
| 1971 | 2.7 | * | 3.8 | * | 2.7 | 4.0 | 4.3 | 3.0 | 2.6 | * | * | * |
| 1970 | 2.9 | 8.8 | 3.3 | * | 3.3 | 4.6 | 4.2 | 3.5 | * | * | * | * |
| 1969 | 2.8 | 9.4 | 3.6 | 0.8 | 3.6 | 3.0 | 4.5 | 2.8 | * | * | * | * |
| 1968 | 2.3 | * | 3.0 | * | 2.0 | 3.4 | 4.3 | 3.3 | * | * | * | * |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.

82/10/14 TABLE 290. DEATH RATES FOR 69 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1968-78

PAGE

[Rates per 100,000 population for specified group. Deaths based on a 50-percent sample in 1972]

| CAUSE OF DEATH CODE, RACE, SEX, AND YEAR | TOTAL 1 YEAR | UNDER 1 YEAR | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85+ YEARS |
|---|-----------------|-----------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|
| OTHER EXTERNAL CAUSES (E980-E999) | | | | | | | | | | | | |
| BLACK, BOTH SEXES | | | | | | | | | | | | |
| 1978 | 3.3 | 7.3 | 2.8 | 0.7 | 3.1 | 5.1 | 5.3 | 4.9 | 2.8 | 2.9 | 3.9 | * |
| 1977 | 3.8 | 9.7 | 2.8 | 0.9 | 3.0 | 6.7 | 5.7 | 5.4 | 3.9 | 4.2 | * | * |
| 1976 | 4.5 | 9.7 | 3.7 | 0.9 | 4.0 | 6.8 | 7.0 | 7.5 | 4.3 | 3.3 | 4.1 | * |
| 1975 | 4.5 | 9.6 | 3.4 | 0.6 | 4.2 | 7.7 | 6.8 | 6.6 | 4.5 | 4.5 | 4.4 | * |
| 1974 | 5.0 | 7.8 | 3.5 | 0.8 | 5.3 | 7.7 | 9.3 | 6.0 | 5.7 | 5.7 | 4.2 | * |
| 1973 | 5.0 | 5.4 | 2.8 | 1.2 | 5.0 | 7.5 | 8.9 | 7.7 | 6.1 | 4.5 | 4.4 | * |
| 1972 | 5.5 | 10.1 | 4.1 | 1.0 | 6.4 | 8.6 | 9.1 | 7.7 | 5.1 | 4.1 | 4.6 | * |
| 1971 | 5.7 | 9.1 | 4.1 | 0.8 | 6.7 | 9.5 | 9.4 | 7.3 | 6.7 | 4.3 | * | * |
| 1970 | 6.4 | 11.1 | 4.1 | 1.0 | 8.6 | 11.4 | 9.7 | 8.7 | 5.2 | 5.4 | * | * |
| 1969 | 6.2 | 10.5 | 4.5 | 1.1 | 7.4 | 10.0 | 11.2 | 8.2 | 5.9 | 4.9 | * | * |
| 1968 | 4.7 | 8.7 | 3.2 | 0.7 | 4.6 | 8.1 | 9.3 | 6.9 | 4.6 | 4.0 | * | * |
| BLACK, MALE | | | | | | | | | | | | |
| 1978 | 5.0 | 8.1 | 3.3 | 1.0 | 4.5 | 7.7 | 8.7 | 8.4 | 4.6 | 4.5 | * | * |
| 1977 | 5.6 | 14.7 | 3.5 | 1.1 | 4.0 | 10.4 | 9.4 | 8.4 | 6.6 | 6.0 | * | * |
| 1976 | 6.3 | 9.2 | 3.4 | 1.2 | 5.5 | 10.8 | 10.7 | 12.0 | 6.9 | 4.8 | * | * |
| 1975 | 6.7 | 9.1 | 3.7 | 0.8 | 6.3 | 12.5 | 11.2 | 11.2 | 7.3 | 6.6 | * | * |
| 1974 | 7.6 | * | 3.3 | 1.2 | 8.1 | 12.5 | 15.3 | 9.6 | 9.8 | 9.6 | * | * |
| 1973 | 7.6 | * | 3.7 | 1.7 | 7.1 | 12.7 | 13.4 | 12.7 | 10.4 | 8.3 | * | * |
| 1972 | 8.0 | 12.4 | 4.0 | 1.6 | 8.9 | 14.3 | 14.4 | 10.9 | 9.3 | 5.1 | * | * |
| 1971 | 8.8 | 11.8 | 4.6 | 1.2 | 10.7 | 15.9 | 15.0 | 12.3 | 11.2 | 6.9 | * | * |
| 1970 | 10.1 | 12.3 | 4.8 | 1.4 | 14.3 | 18.9 | 16.0 | 14.4 | 9.3 | 9.5 | * | * |
| 1969 | 9.7 | 10.9 | 4.9 | 1.3 | 11.4 | 18.1 | 19.2 | 14.4 | 10.3 | 7.3 | * | * |
| 1968 | 7.2 | 10.1 | 3.3 | 0.9 | 7.4 | 13.7 | 14.7 | 11.1 | 8.1 | 7.1 | * | * |
| BLACK, FEMALE | | | | | | | | | | | | |
| 1978 | 1.9 | * | 2.2 | * | 1.8 | 2.8 | 2.5 | 2.1 | * | * | * | * |
| 1977 | 2.2 | * | 2.2 | * | 2.1 | 3.5 | 2.5 | 2.8 | * | 2.8 | * | * |
| 1976 | 2.8 | 10.2 | 4.1 | * | 2.6 | 3.4 | 3.8 | 3.8 | 2.3 | * | * | * |
| 1975 | 2.4 | 10.2 | 3.1 | * | 2.2 | 3.5 | 3.1 | 2.7 | 2.2 | * | * | * |
| 1974 | 2.7 | * | 3.7 | * | 2.6 | 3.7 | 4.4 | 3.0 | 2.2 | * | * | * |
| 1973 | 2.6 | * | * | * | 2.9 | 3.1 | 5.1 | 3.4 | 2.5 | * | * | * |
| 1972 | 3.2 | * | 4.3 | * | 4.0 | 3.7 | 4.7 | 4.9 | * | 3.3 | * | * |
| 1971 | 2.8 | * | 3.6 | * | 2.9 | 4.1 | 4.7 | 3.0 | 2.8 | * | * | * |
| 1970 | 3.0 | 9.9 | 3.4 | * | 3.4 | 5.1 | 4.5 | 3.7 | * | * | * | * |
| 1969 | 3.0 | 10.1 | 4.0 | 0.9 | 3.8 | 3.2 | 4.6 | 3.0 | * | * | * | * |
| 1968 | 2.4 | * | 3.1 | * | 2.0 | 3.4 | 4.7 | 3.3 | * | * | * | * |

* Figure does not meet standards of reliability or precision.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished trend table 290 for 1968-78.