

**Table 19. Reported AIDS cases for male adults and adolescents, by transmission category and race/ethnicity, 2006 and cumulative—United States and dependent areas**

Transmission category	2006		Cumulative <sup>a</sup>	
	No.	%	No.	%
<b>White, not Hispanic</b>				
Male-to-male sexual contact	6,251	65	253,927	73
Injection drug use	733	8	31,737	9
Male-to-male sexual contact and injection drug use	732	8	31,434	9
Hemophilia/coagulation disorder	26	0	3,989	1
High-risk heterosexual contact <sup>b</sup>	416	4	7,850	2
Sex with injection drug user	74	1	2,310	1
Sex with person with hemophilia	0	0	36	0
Sex with HIV-infected transfusion recipient	2	0	176	0
Sex with HIV-infected person, risk factor not specified	340	4	5,328	2
Receipt of blood transfusion, blood components, or tissue	20	0	3,194	1
Other/risk factor not reported or identified	1,400	15	16,469	5
<b>Total</b>	<b>9,578</b>	<b>100</b>	<b>348,600</b>	<b>100</b>
<b>Black, not Hispanic</b>				
Male-to-male sexual contact	4,309	36	103,297	37
Injection drug use	1,635	14	83,369	30
Male-to-male sexual contact and injection drug use	526	4	22,134	8
Hemophilia/coagulation disorder	10	0	604	0
High-risk heterosexual contact <sup>b</sup>	1,875	16	29,952	11
Sex with injection drug user	167	1	6,823	2
Sex with person with hemophilia	1	0	36	0
Sex with HIV-infected transfusion recipient	15	0	237	0
Sex with HIV-infected person, risk factor not specified	1,692	14	22,856	8
Receipt of blood transfusion, blood components, or tissue	24	0	1,218	0
Other/risk factor not reported or identified	3,579	30	40,268	14
<b>Total</b>	<b>11,958</b>	<b>100</b>	<b>280,842</b>	<b>100</b>
<b>Hispanic</b>				
Male-to-male sexual contact	2,816	47	64,893	44
Injection drug use	965	16	46,337	31
Male-to-male sexual contact and injection drug use	287	5	10,884	7
Hemophilia/coagulation disorder	6	0	466	0
High-risk heterosexual contact <sup>b</sup>	638	11	10,950	7
Sex with injection drug user	84	1	2,409	2
Sex with person with hemophilia	0	0	11	0
Sex with HIV-infected transfusion recipient	5	0	125	0
Sex with HIV-infected person, risk factor not specified	549	9	8,405	6
Receipt of blood transfusion, blood components, or tissue	17	0	657	0
Other/risk factor not reported or identified	1,326	22	14,832	10
<b>Total</b>	<b>6,055</b>	<b>100</b>	<b>149,019</b>	<b>100</b>

**Table 19. Reported AIDS cases for male adults and adolescents, by transmission category and race/ethnicity, 2006 and cumulative—United States and dependent areas (cont)**

Transmission category	2006		Cumulative <sup>a</sup>	
	No.	%	No.	%
<b>Asian/Pacific Islander</b>				
Male-to-male sexual contact	225	54	4,475	68
Injection drug use	20	5	317	5
Male-to-male sexual contact and injection drug use	17	4	284	4
Hemophilia/coagulation disorder	1	0	70	1
High-risk heterosexual contact <sup>b</sup>	34	8	376	6
Sex with injection drug user	1	0	54	1
Sex with person with hemophilia	0	0	1	0
Sex with HIV-infected transfusion recipient	0	0	8	0
Sex with HIV-infected person, risk factor not specified	33	8	313	5
Receipt of blood transfusion, blood components, or tissue	2	0	112	2
Other/risk factor not reported or identified	114	28	949	14
<b>Total</b>	<b>413</b>	<b>100</b>	<b>6,583</b>	<b>100</b>
<b>American Indian/Alaska Native</b>				
Male-to-male sexual contact	61	49	1,403	55
Injection drug use	15	12	395	15
Male-to-male sexual contact and injection drug use	23	19	457	18
Hemophilia/coagulation disorder	0	0	27	1
High-risk heterosexual contact <sup>b</sup>	9	7	118	5
Sex with injection drug user	1	1	33	1
Sex with person with hemophilia	0	0	0	0
Sex with HIV-infected transfusion recipient	0	0	3	0
Sex with HIV-infected person, risk factor not specified	8	6	82	3
Receipt of blood transfusion, blood components, or tissue	0	0	8	0
Other/risk factor not reported or identified	16	13	155	6
<b>Total</b>	<b>124</b>	<b>100</b>	<b>2,563</b>	<b>100</b>
<b>Total cases</b>				
Male-to-male sexual contact	13,775	49	429,897	54
Injection drug use	3,413	12	163,070	21
Male-to-male sexual contact and injection drug use	1,603	6	65,524	8
Hemophilia/coagulation disorder	43	0	5,176	1
High-risk heterosexual contact <sup>b</sup>	2,997	11	49,610	6
Sex with injection drug user	330	1	11,683	1
Sex with person with hemophilia	1	0	84	0
Sex with HIV-infected transfusion recipient	22	0	553	0
Sex with HIV-infected person, risk factor not specified	2,644	9	37,290	5
Receipt of blood transfusion, blood components, or tissue	63	0	5,213	1
Other/risk factor not reported or identified	6,484	23	73,136	9
<b>Total</b>	<b>28,378<sup>c</sup></b>	<b>100</b>	<b>791,626<sup>d</sup></b>	<b>100</b>

<sup>a</sup> From the beginning of the epidemic through 2006.

<sup>b</sup> Heterosexual contact with a person known to have, or to be at high risk for, HIV infection.

<sup>c</sup> Includes 250 males of unknown race or multiple races.

<sup>d</sup> Includes 4,019 males of unknown race or multiple races.