

**Table 13. Proportion of persons surviving for more than 12, 24, and 36 months after an AIDS diagnosis in 2002, by selected characteristics—United States and dependent areas**

	No. of persons	Survival in months		
		>12	>24	>36
<b>Data for 50 states and the District of Columbia</b>				
<b>Age at diagnosis (yrs)</b>				
<13	101	0.96	0.94	0.94
13–14	60	0.98	0.98	0.95
15–19	305	0.98	0.96	0.95
20–24	1,351	0.96	0.93	0.92
25–29	3,034	0.95	0.92	0.91
30–34	5,499	0.94	0.91	0.89
35–39	7,888	0.93	0.90	0.87
40–44	7,085	0.92	0.88	0.85
45–49	5,090	0.89	0.85	0.82
50–54	3,067	0.86	0.81	0.78
55–59	1,505	0.83	0.78	0.74
60–64	806	0.80	0.73	0.69
≥65	614	0.73	0.68	0.65
<b>Race/ethnicity</b>				
White, not Hispanic	10,706	0.92	0.89	0.87
Black, not Hispanic	18,406	0.90	0.86	0.82
Hispanic	6,398	0.92	0.90	0.88
Asian/Pacific Islander	404	0.94	0.92	0.90
American Indian/Alaska Native	167	0.84	0.81	0.75
<b>Transmission category</b>				
<b>Male adult or adolescent</b>				
Male-to-male sexual contact	12,830	0.93	0.90	0.88
Injection drug use	4,048	0.89	0.84	0.81
Male-to-male sexual contact and injection drug use	1,701	0.93	0.89	0.86
High-risk heterosexual contact <sup>a</sup>	3,170	0.91	0.88	0.85
Other <sup>b</sup>	5,019	0.86	0.83	0.80
Subtotal	26,768	0.91	0.88	0.85
<b>Female adult or adolescent</b>				
Injection drug use	2,101	0.89	0.83	0.78
High-risk heterosexual contact <sup>a</sup>	4,686	0.93	0.89	0.86
Other <sup>b</sup>	2,749	0.90	0.85	0.83
Subtotal	9,536	0.91	0.87	0.83
<b>Child (&lt;13 yrs at diagnosis)</b>				
Perinatal	98	0.96	0.95	0.95
Other <sup>c</sup>	3	1.00	0.67	0.67
Subtotal	101	0.96	0.94	0.94
<b>Subtotal for 50 states and the District of Columbia</b>	<b>36,405</b>	<b>0.91</b>	<b>0.87</b>	<b>0.85</b>
<b>Data for U.S. dependent areas</b>	<b>891</b>	<b>0.83</b>	<b>0.78</b>	<b>0.75</b>
<b>Total</b>	<b>37,352<sup>d</sup></b>	<b>0.91</b>	<b>0.87</b>	<b>0.84</b>

Note. Excludes persons whose date of death was before, or in the same month as, date of diagnosis. Excludes persons whose month of diagnosis or month of death is unknown.

<sup>a</sup> Heterosexual contact with a person known to have, or to be at high risk for, HIV infection.

<sup>b</sup> Includes hemophilia, blood transfusion, perinatal exposure, and risk factor not reported or not identified.

<sup>c</sup> Includes hemophilia, blood transfusion, and risk factor not reported or not identified.

<sup>d</sup> Includes 325 persons of unknown race or multiple races.