

Table 11. Estimated numbers of persons living with AIDS at the end of 2006, by race/ethnicity, sex, and transmission category—50 states and the District of Columbia

Transmission category	White, not Hispanic		Black, not Hispanic		Hispanic		Asian/Pacific Islander		American Indian/Alaska Native		Total ^a	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent												
Male-to-male sexual contact	101,299	75	57,520	45	36,087	57	2,715	72	704	57	199,693	60
Injection drug use	13,168	10	33,911	27	14,795	23	325	9	201	16	62,998	19
Male-to-male sexual contact and injection drug use	12,393	9	10,121	8	4,352	7	198	5	220	18	27,523	8
High-risk heterosexual contact ^b	6,081	5	24,205	19	7,816	12	444	12	105	8	39,004	12
Other ^c	1,743	1	1,602	1	574	1	69	2	16	1	4,026	1
Subtotal	134,683	100	127,359	100	63,623	100	3,752	100	1,246	100	333,244	100
Female adult or adolescent												
Injection drug use	7,407	38	18,531	30	5,025	30	115	15	153	39	31,575	32
High-risk heterosexual contact ^b	11,350	59	41,741	68	11,092	67	588	78	227	58	65,631	66
Other ^c	508	3	1,456	2	413	2	51	7	13	3	2,465	2
Subtotal	19,264	100	61,728	100	16,531	100	753	100	393	100	99,671	100
Child (<13 yrs at diagnosis)												
Perinatal	500	92	2,468	99	629	95	20	94	12	100	3,660	97
Other ^d	46	8	34	1	33	5	1	6	0	0	115	3
Subtotal	546	100	2,503	100	662	100	21	100	12	100	3,775	100
Total^e	154,495		191,590		80,815		4,526		1,651		436,693	

Note. These numbers do not represent reported case counts. Rather, these numbers are point estimates, which result from adjustments of reported case counts. The reported case counts have been adjusted for reporting delays and for redistribution of cases in persons initially reported without an identified risk factor, but not for incomplete reporting.

^a Includes 3,616 persons of unknown race or multiple races.

^b Heterosexual contact with a person known to have, or to be at high risk for, HIV infection.

^c Includes hemophilia, blood transfusion, perinatal exposure, and risk factor not reported or not identified.

^d Includes hemophilia, blood transfusion, and risk factor not reported or not identified.

^e Includes 3 person of unknown sex. Because column totals were calculated independently of the values for the subpopulations, the values in each column may not sum to the column total.