

Table 1. Estimated numbers of cases of HIV/AIDS, by year of diagnosis and selected characteristics, 2003–2006—33 states and 5 U.S. dependent areas with confidential name-based HIV infection reporting

	Year of diagnosis			
	2003	2004	2005	2006
Data for 33 states				
Age at diagnosis (yrs)				
<13	211	183	169	135
13–14	53	36	40	41
15–19	993	993	1,126	1,332
20–24	3,163	3,368	3,592	3,886
25–29	4,023	4,057	4,236	4,603
30–34	5,189	4,820	4,676	4,466
35–39	6,369	5,807	5,535	5,442
40–44	5,786	5,429	5,529	5,718
45–49	4,028	3,877	4,028	4,204
50–54	2,451	2,401	2,547	2,718
55–59	1,279	1,363	1,455	1,438
60–64	655	702	692	714
≥65	570	624	613	618
Race/ethnicity				
White, not Hispanic	10,033	10,181	10,528	10,758
Black, not Hispanic	17,668	16,718	16,629	17,356
Hispanic	6,355	6,010	6,217	6,481
Asian/Pacific Islander	338	339	373	397
American Indian/Alaska Native	179	171	182	166
Transmission category				
Male adult or adolescent				
Male-to-male sexual contact	15,409	15,880	16,833	17,465
Injection drug use	3,514	3,083	2,978	3,016
Male-to-male sexual contact and injection drug use	1,349	1,299	1,247	1,180
High-risk heterosexual contact ^a	4,269	3,959	3,871	4,152
Other ^b	125	110	107	114
Subtotal	24,666	24,331	25,036	25,928
Female adult or adolescent				
Injection drug use	2,027	1,856	1,720	1,712
High-risk heterosexual contact ^a	7,731	7,182	7,216	7,432
Other ^b	134	107	97	109
Subtotal	9,892	9,145	9,033	9,252
Child (<13 yrs at diagnosis)				
Perinatal	190	157	147	115
Other ^c	23	27	23	20
Subtotal	213	184	170	135
Subtotal for 33 states	34,770	33,659	34,239	35,314
Data for U.S. dependent areas	1,331	1,234	1,395	1,503
Total^d	36,102	34,894	35,634	36,817

Note. These numbers do not represent reported case counts. Rather, these numbers are point estimates, which result from adjustments of reported case counts. The reported case counts have been adjusted for reporting delays and for redistribution of cases in persons initially reported without an identified risk factor, but not for incomplete reporting.

Data include persons with a diagnosis of HIV infection (not AIDS), a diagnosis of HIV infection and a later diagnosis of AIDS, or concurrent diagnoses of HIV infection and AIDS.

See Technical Notes for the list of areas that have had laws or regulations requiring confidential name-based HIV infection reporting since at least 2003.

^a Heterosexual contact with a person known to have, or to be at high risk for, HIV infection.

^b Includes hemophilia, blood transfusion, perinatal exposure, and risk factor not reported or not identified.

^c Includes hemophilia, blood transfusion, and risk factor not reported or not identified.

^d Includes persons of unknown race or multiple races and persons of unknown sex. Because column totals were calculated independently of the values for the subpopulations, the values in each column may not sum to the column total.