

Submitted to:

National Toxicology Program
National Institute of Environmental Health Sciences
P.O. Box 12233
Research Triangle Park, North Carolina 27709

FINAL REPORT

**TWO-GENERATION REPRODUCTION TOXICITY STUDY OF
PROPYLTHIOURACIL WHEN ADMINISTERED TO SPRAGUE-DAWLEY RATS IN
THE DRINKING WATER**

Authors:

Gary W. Wolfe, Ph.D., D.A.B.T.
Stefanie Pepperl, B.S.
Jennifer Greenlaw, M.H.S.

Study Completion Date:

To be added at report finalization

Submitted by:

TherImmune Research Corporation
15 Firstfield Road
Gaithersburg, Maryland 20878

Project Identification:

TRC Study No. 7244-601

Volume I of

TABLE OF CONTENTS

QUALITY ASSURANCE STATEMENT 10

COMPLIANCE STATEMENT..... 12

SIGNATURE PAGE 13

ACKNOWLEDGMENTS 14

ABSTRACT

Summary of Results..... 15

INTRODUCTION 25

METHODS..... 25

 General Study Design..... 25

 Two-Generation Reproduction Toxicity Study Flow Diagram for Propylthiouracil 28

 Study Animals 29

 Animal Husbandry and Environmental Conditions 30

 Test Article..... 30

 Dosage Formulation and Administration..... 31

 F₀ Cohabitation..... 32

 F₁ PND 21 Necropsy and Terminal Procedures 33

 F₀ Adult Necropsy and Terminal Procedures 34

 Allocation of F₁ Weanlings 35

 F₁ Cohabitation..... 36

 F₂ PND 21 Necropsy and Terminal Procedures 37

 F₁ Adult Necropsy and Terminal Procedures 38

 Statistical Analyses..... 39

 Specimen, Raw Data and Final Report Storage..... 40

RESULTS..... 42

 F₀ Cohabitation..... 42

 F₁ PND 21 Necropsy 44

 F₀ Thyroid Hormone Analysis..... 45

 F₀ Adult Necropsy 45

 F₁ Growing Phase..... 46

 F₁ Cohabitation..... 47

 F₂ PND 21 Necropsy 48

 F₁ Thyroid Hormone Analysis..... 48

 F₁ Adult Necropsy 49

DISCUSSION..... 50

TABLE OF CONTENTS (continued)

REFERENCES	54
------------------	----

FIGURES

1 - Two-Generation Reproduction Toxicity Study Flow Diagram for Propylthiouracil	28
--	----

TABLES

0-1A - F ₀ : SUMMARY OF MALE BODY WEIGHTS (GRAMS).....	56
0-1B - F ₀ : SUMMARY OF FEMALE BODY WEIGHTS (GRAMS).....	57
0-2A - F ₀ : SUMMARY OF DAILY FEED CONSUMPTION (G/KG BODY WEIGHT/DAY).....	58
0-2B - F ₀ : SUMMARY OF DAILY FEED CONSUMPTION (G/ANIMAL/DAY).....	59
0-3A - F ₀ : SUMMARY OF DAILY WATER CONSUMPTION (G/KG BODY WEIGHT/DAY).....	60
0-3B - F ₀ : SUMMARY OF DAILY WATER CONSUMPTION (G/ANIMAL/DAY)	61
0-4 - F ₀ : SUMMARY OF ESTIMATED DOSAGE (MG/KG BODY WEIGHT/DAY).....	62
0-5 - F ₀ : SUMMARY OF REPRODUCTIVE PERFORMANCE OF COHABITED PAIRS	63
0-6 - F ₀ : SUMMARY OF DAM WEIGHTS (GRAMS) DURING LACTATION OF THE F ₁ LITTER	64
0-7A - F ₀ : SUMMARY OF DAM DAILY FEED CONSUMPTION (G/KG BODY WEIGHT/DAY) DURING LACTATION OF THE F ₁ LITTER	65
0-7B - F ₀ : SUMMARY OF DAM DAILY FEED CONSUMPTION (G/ANIMAL/DAY) DURING LACTATION OF THE F ₁ LITTER	66
0-8A - F ₀ : SUMMARY OF DAM DAILY WATER CONSUMPTION (G/KG BODY WEIGHT/DAY) DURING LACTATION OF THE F ₁ LITTER.....	67
0-8B - F ₀ : SUMMARY OF DAM DAILY WATER CONSUMPTION (G/ANIMAL/DAY) DURING LACTATION OF THE F ₁ LITTER	68
0-9 - F ₀ : SUMMARY OF ESTIMATED DOSAGE (G/KG BODY WEIGHT/DAY) DURING LACTATION OF THE F ₁ LITTER.....	69
0-10 - F ₀ : SUMMARY OF ANOGENITAL DISTANCE AND PUP WEIGHT FOR THE F ₁ LITTER	70

TABLE OF CONTENTS (continued)

0-11	-	F ₀ : SUMMARY OF AVERAGE PUP WEIGHTS FOR THE F ₁ LITTER	71
0-12	-	F ₀ : SUMMARY OF PUP SURVIVAL FOR THE F ₁ LITTER.....	72
0-13	-	F ₀ : SUMMARY OF NIPPLE RETENTION FOR THE F ₁ LITTER.....	73
0-14	-	F ₀ : SUMMARY OF PINNA DETACHMENT AND EYE OPENING FOR THE F ₁ LITTER	74
0-15	-	F ₀ : SUMMARY OF F ₁ SEXUAL DEVELOPMENT DATA	75
0-16	-	F ₀ : SUMMARY OF F ₁ MALE PND 21 NECROPSY GROSS OBSERVATIONS	76
0-17	-	F ₀ : SUMMARY OF F ₁ MALE PND 21 ABSOLUTE ORGAN WEIGHTS AND ORGAN-TO-BODY WEIGHT RATIOS.....	78
0-18	-	F ₀ : SUMMARY OF F ₁ FEMALE PND 21 NECROPSY GROSS OBSERVATIONS.....	79
0-19	-	F ₀ : SUMMARY OF F ₁ FEMALE PND 21 ABSOLUTE ORGAN WEIGHTS AND ORGAN-TO-BODY WEIGHT RATIOS	81
0-20	-	F ₀ : SUMMARY OF STAGES AND LENGTH OF ESTROUS CYCLE.....	82
0-21	-	F ₀ : SUMMARY OF THYROID HORMONES	83
0-22	-	F ₀ : SUMMARY OF MALE GROSS OBSERVATIONS	84
0-23	-	F ₀ : SUMMARY OF MALE ABSOLUTE ORGAN WEIGHTS AND ORGAN-TO-BODY WEIGHT RATIOS	86
0-24A	-	F ₀ : SUMMARY OF SPERM ANALYSIS DATA.....	87
0-24B	-	F ₀ : SUMMARY OF HAMILTON-THORNE SPERM ANALYSIS DATA.....	88
0-25	-	F ₀ : SUMMARY OF FEMALE GROSS OBSERVATIONS.....	89
0-26	-	F ₀ : SUMMARY OF FEMALE ABSOLUTE ORGAN WEIGHTS AND ORGAN-TO-BODY WEIGHT RATIOS.....	91
0-27	-	F ₀ : SUMMARY OF MORTALITY DATA	92
1A-1	-	F ₁ : SUMMARY OF ADOLESCENT BODY WEIGHTS (GRAMS).....	93
1A-2	-	F ₁ : SUMMARY OF ADOLESCENT MORTALITY DATA	94

TABLE OF CONTENTS (continued)

1-1A	-	F ₁ : SUMMARY OF MALE BODY WEIGHTS (GRAMS).....	95
1-1B	-	F ₁ : SUMMARY OF FEMALE BODY WEIGHTS (GRAMS).....	96
1-2A	-	F ₁ : SUMMARY OF DAILY FEED CONSUMPTION (G/KG BODY WEIGHT/DAY).....	97
1-2B	-	F ₁ : SUMMARY OF DAILY FEED CONSUMPTION (G/ANIMAL/DAY).....	98
1-3A	-	F ₁ : SUMMARY OF DAILY WATER CONSUMPTION (G/KG BODY WEIGHT/DAY).....	99
1-3B	-	F ₁ : SUMMARY OF DAILY WATER CONSUMPTION (G/ANIMAL/DAY)	100
1-4	-	F ₁ : SUMMARY OF ESTIMATED DOSAGE (MG/KG BODY WEIGHT/DAY).....	101
1-5	-	F ₁ : SUMMARY OF REPRODUCTIVE PERFORMANCE OF COHABITED PAIRS	102
1-6	-	F ₁ : SUMMARY OF DAM WEIGHTS (GRAMS) DURING LACTATION OF THE F ₂ LITTER.....	103
1-7A	-	F ₁ : SUMMARY OF DAM DAILY FEED CONSUMPTION (G/KG BODY WEIGHT/DAY) DURING LACTATION OF THE F ₂ LITTER	104
1-7B	-	F ₁ : SUMMARY OF DAM DAILY FEED CONSUMPTION (G/ANIMAL/DAY) DURING LACTATION OF THE F ₂ LITTER	105
1-8A	-	F ₁ : SUMMARY OF DAM DAILY WATER CONSUMPTION (G/KG BODY WEIGHT/DAY) DURING LACTATION OF THE F ₂ LITTER.....	106
1-8B	-	F ₁ : SUMMARY OF DAM DAILY WATER CONSUMPTION (G/ANIMAL/DAY) DURING LACTATION OF THE F ₂ LITTER	107
1-9	-	F ₁ : SUMMARY OF ESTIMATED DOSAGE (G/KG BODY WEIGHT/DAY) DURING LACTATION OF THE F ₂ LITTER.....	108
1-10	-	F ₁ : SUMMARY OF ANOGENITAL DISTANCE AND PUP WEIGHT FOR THE F ₂ LITTER	109
1-11	-	F ₁ : SUMMARY OF AVERAGE PUP WEIGHTS FOR THE F ₂ LITTER	110
1-12	-	F ₁ : SUMMARY OF PUP SURVIVAL FOR THE F ₂ LITTER.....	111
1-13	-	F ₁ : SUMMARY OF NIPPLE RETENTION FOR THE F ₂ LITTER.....	112
1-14	-	F ₁ : SUMMARY OF PINNA DETACHMENT AND EYE OPENING FOR THE F ₂ LITTER	113
1-15	-	F ₁ : SUMMARY OF F ₂ MALE PND 21 NECROPSY GROSS OBSERVATIONS	114

TABLE OF CONTENTS (continued)

1-16	-	F ₁ : SUMMARY OF F ₂ MALE PND 21 ABSOLUTE ORGAN WEIGHTS AND ORGAN-TO-BODY WEIGHT RATIOS.....	116
1-17	-	F ₁ : SUMMARY OF F ₂ FEMALE PND 21 NECROPSY GROSS OBSERVATIONS.....	117
1-18	-	F ₁ : SUMMARY OF F ₂ FEMALE PND 21 ABSOLUTE ORGAN WEIGHTS AND ORGAN-TO-BODY WEIGHT RATIOS	119
1-19	-	F ₁ : SUMMARY OF STAGES AND LENGTH OF ESTROUS CYCLE.....	120
1-20	-	F ₁ : SUMMARY OF THYROID HORMONES	121
1-21	-	F ₁ : SUMMARY OF MALE GROSS OBSERVATIONS	122
1-22	-	F ₁ : SUMMARY OF MALE ABSOLUTE ORGAN WEIGHTS AND ORGAN-TO-BODY WEIGHT RATIOS	124
1-23A	-	F ₁ : SUMMARY OF SPERM ANALYSIS DATA.....	125
1-23B	-	F ₁ : SUMMARY OF HAMILTON-THORNE SPERM ANALYSIS DATA.....	126
1-24	-	F ₁ : SUMMARY OF FEMALE GROSS OBSERVATIONS.....	127
1-25	-	F ₁ : SUMMARY OF FEMALE ABSOLUTE ORGAN WEIGHTS AND ORGAN-TO-BODY WEIGHT RATIOS.....	128
1-26	-	F ₁ : SUMMARY OF MORTALITY DATA	129

APPENDIX TABLES

A0-1	-	F ₀ : INDIVIDUAL BODY WEIGHTS (GRAMS).....	130
A0-2	-	F ₀ : INDIVIDUAL FEED CONSUMPTION (GRAMS)	146
A0-3	-	F ₀ : INDIVIDUAL WATER CONSUMPTION (GRAMS).....	154
A0-4A	-	F ₀ : SUMMARY OF MALE CLINICAL OBSERVATIONS.....	162
A0-4B	-	F ₀ : SUMMARY OF FEMALE CLINICAL OBSERVATIONS	164
A0-5	-	F ₀ : INDIVIDUAL REPRODUCTIVE PERFORMANCE DURING THE F ₁ LITTER	165

TABLE OF CONTENTS (continued)

A0-6	- F ₀ : INDIVIDUAL DAM FEED CONSUMPTION (GRAMS/DAY) DURING LACTATION.....	177
A0-7	- F ₀ : INDIVIDUAL DAM WATER CONSUMPTION (GRAMS) DURING LACTATION.....	181
A0-8	- F ₀ : INDIVIDUAL ANOGENITAL DISTANCE AND PUP WEIGHTS FOR THE F ₁ LITTER.....	185
A0-9	- F ₀ : INDIVIDUAL NIPPLE RETENTION FOR THE F ₁ LITTER.....	202
A0-10	- F ₀ : INDIVIDUAL PINNA DETACHMENT AND EYE OPENING FOR THE F ₁ LITTER.....	210
A0-11	- F ₀ : INDIVIDUAL SEXUAL DEVELOPMENT DATA FOR THE F ₁ LITTER.....	214
A0-12	- F ₀ : INDIVIDUAL F ₁ MALE PND 21 GROSS OBSERVATIONS.....	217
A0-13	- F ₀ INDIVIDUAL F ₁ MALE PND 21 ORGAN WEIGHTS AND ORGAN-TO-BODY WEIGHT RATIOS.....	225
A0-14	- F ₀ : INDIVIDUAL F ₁ FEMALE PND 21 GROSS OBSERVATIONS.....	233
A0-15	- F ₀ INDIVIDUAL F ₁ FEMALE PND 21 ORGAN WEIGHTS AND ORGAN-TO-BODY WEIGHT RATIOS.....	240
A0-16	- F ₀ : INDIVIDUAL STAGES AND LENGTH OF ESTROUS CYCLE.....	247
A0-17	- F ₀ : INDIVIDUAL THYROID HORMONE DATA.....	251
A0-18	- F ₀ : INDIVIDUAL MALE GROSS OBSERVATIONS.....	259
A0-19A	- F ₀ : INDIVIDUAL MALE ORGAN WEIGHTS (GRAMS).....	264
A0-19B	- F ₀ : INDIVIDUAL MALE ORGAN-TO-BODY-WEIGHT RATIOS.....	268
A0-20A	- F ₀ : INDIVIDUAL SPERM ANALYSIS DATA.....	272
A0-20B	- F ₀ : INDIVIDUAL HAMILTON-THORNE SPERM ANALYSIS DATA.....	276
A0-21	- F ₀ : INDIVIDUAL FEMALE GROSS OBSERVATIONS.....	280
A0-22A	- F ₀ : INDIVIDUAL FEMALE ORGAN WEIGHTS (GRAMS).....	284
A0-22B	- F ₀ : INDIVIDUAL FEMALE ORGAN-TO-BODY-WEIGHT RATIOS.....	288
A0-23	- F ₀ : INDIVIDUAL MORTALITY DATA.....	292
A1-1	- F ₁ : INDIVIDUAL ADOLESCENT BODY WEIGHTS (GRAMS).....	296

TABLE OF CONTENTS (continued)

A1-2A	-	F ₁ : SUMMARY OF ADOLESCENT MALE CLINICAL OBSERVATIONS.....	302
A1-2B	-	F ₁ : SUMMARY OF ADOLESCENT FEMALE CLINICAL OBSERVATIONS	303
A1-3	-	F ₁ : INDIVIDUAL ADOLESCENT MORTALITY DATA	304
A1-4	-	F ₁ : INDIVIDUAL BODY WEIGHTS (GRAMS).....	308
A1-5	-	F ₁ : INDIVIDUAL FEED CONSUMPTION (GRAMS)	314
A1-6	-	F ₁ : INDIVIDUAL WATER CONSUMPTION (GRAMS).....	320
A1-7A	-	F ₁ : SUMMARY OF MALE CLINICAL OBSERVATIONS.....	326
A1-7B	-	F ₁ : SUMMARY OF FEMALE CLINICAL OBSERVATIONS	327
A1-8	-	F ₁ : INDIVIDUAL REPRODUCTIVE PERFORMANCE DURING THE F ₂ LITTER.....	328
A1-9	-	F ₁ : INDIVIDUAL DAM FEED CONSUMPTION (GRAMS) DURING LACTATION.....	337
A1-10	-	F ₁ : INDIVIDUAL DAM WATER CONSUMPTION (GRAMS) DURING LACTATION.....	340
A1-11	-	F ₁ : INDIVIDUAL ANOGENITAL DISTANCE AND PUP WEIGHTS FOR THE F ₂ LITTER.....	343
A1-12	-	F ₁ : INDIVIDUAL NIPPLE RETENTION FOR THE F ₂ LITTER.....	355
A1-13	-	F ₁ : INDIVIDUAL PINNA DETACHMENT AND EYE OPENING FOR THE F ₂ LITTER	361
A1-14	-	F ₁ : INDIVIDUAL F ₂ MALE PND 21 GROSS OBSERVATIONS	364
A1-15	-	F ₁ : INDIVIDUAL F ₂ MALE PND 21 ORGAN WEIGHTS AND ORGAN-TO-BODY WEIGHT RATIOS	370
A1-16	-	F ₁ : INDIVIDUAL F ₂ FEMALE PND 21 GROSS OBSERVATIONS.....	376
A1-17	-	F ₁ : INDIVIDUAL F ₂ FEMALE PND 21 ORGAN WEIGHTS AND ORGAN-TO-BODY WEIGHT RATIOS	381
A1-18	-	F ₁ : INDIVIDUAL STAGES AND LENGTH OF ESTROUS CYCLE.....	387

TABLE OF CONTENTS (continued)

A1-19	-	F ₁ : INDIVIDUAL THYROID HORMONE DATA.....	390
A1-20	-	F ₁ : INDIVIDUAL MALE GROSS OBSERVATIONS	396
A1-21A	-	F ₁ : INDIVIDUAL MALE ORGAN WEIGHTS (GRAMS).....	399
A1-21B	-	F ₁ : INDIVIDUAL MALE ORGAN-TO-BODY WEIGHT RATIOS	402
A1-22A	-	F ₁ : INDIVIDUAL SPERM ANALYSIS DATA	405
A1-22B	-	F ₁ : INDIVIDUAL HAMILTON-THORNE SPERM ANALYSIS DATA.....	408
A1-23	-	F ₁ : INDIVIDUAL FEMALE GROSS OBSERVATIONS.....	411
A1-24A	-	F ₁ : INDIVIDUAL FEMALE ORGAN WEIGHTS (GRAMS).....	414
A1-24B	-	F ₁ : INDIVIDUAL FEMALE ORGAN-TO-BODY WEIGHT RATIOS.....	417
A1-25	-	F ₁ : INDIVIDUAL MORTALITY DATA	420
A2-1	-	DEVIATIONS IN THE ANIMAL ROOM TEMPERATURE AND/OR HUMIDITY	423
A2-2	-	VERIFICATION OF TEST ARTICLE CONCENTRATION	425
A2-3	-	PURITY OF BULK TEST ARTICLE	426
A2-4	-	SEROLOGY RESULTS	427
Appendix 2	-	Study Protocol and Amendments.....	428
Appendix 3	-	Milestone Schedule.....	494
Appendix 4	-	Analytical Chemistry Reports	496
Appendix 5	-	Dose Formulation Procedure	549
Appendix 6	-	Pathology Report (PAI)	550
Appendix 7	-	Thyroid Hormone Analysis Methods.....	637

QUALITY ASSURANCE STATEMENT

Page 1 of 2

(To be completed at the time of report finalization)

QUALITY ASSURANCE STATEMENT

Page 2 of 2

COMPLIANCE STATEMENT

This study was conducted in compliance with the Good Laboratory Practice Regulations as set forth in Title 21 of the U.S. Code of Federal Regulations Part 58, issued December 22, 1978 (effective June 20, 1979). Standard Operating Procedure and protocol deviations occurred during the study and are documented in the study data; however, none of these deviations affected the quality or integrity of the study. To the best of my knowledge, this final report accurately describes the study methods and procedures used, and the reported results accurately reflect the raw data.

Study Director:

Gary W. Wolfe, PhD., DABT/Date

SIGNATURE PAGE

Study Director:

Report Manager:

Gary W. Wolfe, Ph.D., D.A.B.T./Date

Stefanie Pepperl, B.S. /Date

Project Officer:

Reproductive Toxicologist:

Jack Bishop, Ph.D./Date

Yefan Wang, M.S. /Date

ACKNOWLEDGMENTS

The present study was conducted at TherImmune Research Corporation in Gaithersburg, Maryland, under contract to the National Toxicology Program and the National Institute for Environmental Health Sciences (NTP/NIEHS Contract No. NO1-ES-75409). The following personnel contributed to the completion of this investigation:

Study Director:	Gary W. Wolfe, PhD, DABT
Pathologists at Necropsy/Histopathologist:	Jerry L. Quance, DVM, DACVP
Project Coordinator:	Julie Delaney, BS, LATG (PAI)
Reproductive Toxicologist:	Yefan Wang, MS
Report Manager:	Stefanie Pepperl, BS
Technical Supervisors:	Roland Naawu, BS, LATG Meredith James, BS
Health and Safety Officer/ Facility Manager:	Robert K. Blackford, AA
Statistical Support:	Julie Scott, MS (ASI)
Veterinarian:	Edward Greenstein, DVM, ACLAM
Quality Assurance Officers:	Cecilia Matos-Rosa, BS David Wilson, BS
Directors of Quality Assurance:	James Carignan, BS Carol Hoffman (PAI)
Dose Preparation Supervisor:	Gary Holley, BS
Head Technician:	Larissa Nehrebeckyj, BS

ABSTRACT

Propylthiouracil was evaluated for potential reproductive toxicity using a Two-Generation study model. Beginning on Study Day (SD) 1, propylthiouracil was administered in the drinking water at dose levels of 0, 0.0001, 0.0004, and 0.0015 % (weight/volume) to adult male and female rats (N=20). The F₀ cohabitation period began on SD 71. Mating pairs were allowed to produce one litter. Dosing of the F₁ generation was initiated on post-natal day (PND) 21 (i.e., at weaning). On PND 99 ± 10, F₁ animals were assigned to mating pairs and allowed to produce one litter. Due to high mortality in the adolescent F₁ 0.0015 % animals, this group was not available for the F₁ cohabitation. Endpoints evaluated included body weight, food and water consumption, clinical signs of toxicity, number and weight of pups, anogenital distance (AGD), sexual development endpoints, thyroid hormone levels, sperm parameters, vaginal cytology, organ weights, and gross and microscopic pathology.

In the F₀ Cohabitation there were significant decreases at the 0.0015% dose level in mean body weights (6-12%), mean food consumption (3-35%) expressed as g/kg body weight/day and g/animal/day, and water consumption (13-38%) expressed as g/animal/day. The pregnancy index was comparable in all groups, but the number of female and total pups was decreased by 43 and 30%, respectively, at 0.0015%. Pup weights were comparable among groups but by PND 14 there was a significant decrease (15-37%) in the 0.0015% male and female pup weights. Survival was also comparable until weaning on PND 21, after which an increase in mortality was observed at 0.0015%. By PND 25, the 0.0015% pups were smaller with domed heads and misshapen snouts. During necropsy on PND 21 a delay in eruption of teeth was noted. Microscopically the jaws had damage (depletion and vacuolation) to the odontoblasts and ameloblasts resulting in a delay in cellular maturation and subsequent tooth eruption.

In the F₁ offspring, eye opening was delayed by 1.9 days at 0.0015%. Prepuce separation was delayed by 2.0 days and vaginal opening by 1.4 days at 0.0004%. Because of the mortality observed in the F₁ animals at 0.0015%, only the 0, 0.0001, and 0.0004% groups continued to the F₁ cohabitation. In the F₁ cohabitation, no changes were noted in body weights, food consumption, water consumption, or reproductive endpoints, except there was a decrease in anogenital distance noted in the males at 0.0001 and 0.0004%.

In the necropsy of F₀ parents there were significant decreases at 0.0015% in absolute adrenal, brain, kidney, liver, and spleen weights and relative right testis and seminal vesicle weights. The absolute thyroid/parathyroid weights were increased 67-373% in the 0.0004 and 0.0015% males and females; relative thyroid/parathyroid weights were increased 34-443% in the 0.0004 and 0.0015% males and females and 18% in the 0.0001% females. In the necropsy of the F₁ parents there were significant increases in the absolute and relative thyroid weights in the 0.0004% males and females and also in relative thyroid weight at 0.0001% (males only). The findings in the F₀ parents correlated with enlarged thyroids/parathyroids observed at necropsy in the 0.0004% males and females (7/20 and 1/20, respectively) and 0.0015% males and females (20/20 and 18/20, respectively). There were no gross pathology findings in the F₁ parents. Upon microscopic examination of the thyroid, follicular cell hyperplasia was observed in 10/10 0.0015% F₀ males and females, 7/10 0.0004% F₀ males, and 1/10 0.0004% F₁ males. There was also degeneration of the germinal epithelium of the testes in 2/10, 3/10, and 3/10 0.0001%, 0.0004%, and 0.0015% F₀ males respectively, and 1/10 0.0001% and 0.0004% F₁ males. TSH levels were increased in the 0.0004% and 0.0015% F₀ males and females and 0.0004% F₁ males and females. T4 levels were decreased in the 0.0004% and 0.0015% F₀ males and females and the 0.0004% F₁ males and females.

Based on the findings of this two generation study with one litter per generation, Propylthiouracil would be considered to be a reproductive/developmental toxicant in females at dose levels greater than or equal to 0.0004% based on decreased total pups per litter, delayed vaginal opening, delayed eye opening, and changes in estrous cyclicity. PTU would be considered a male reproductive toxicant at dose levels greater than or equal to 0.0001% based on delayed eye opening, delayed preputial separation, degeneration of the germinal epithelium of the testes, and decreased anogenital distance. Propylthiouracil would also be considered a general toxicant at 0.0015 and 0.0004% based upon decreased body weight and food consumption (mostly at 0.0015%) and increased thyroid weights, changes in thyroid hormone levels, and /or thyroid follicular cell hyperplasia at 0.0004 and 0.0015%.

SUMMARY OF RESULTS
TWO-GENERATION REPRODUCTION TOXICITY STUDY OF PROPYLTHIOURACIL WHEN ADMINISTERED TO SPRAGUE-DAWLEY RATS IN THE DRINKING WATER

Parameters	Treatment Group (%)		
	Low (0.0001)	Mid (0.0004)	High (0.0015)
<u>General Toxicity - F₀ Cohabitation</u>			
Body Weight			
Male (Weeks 3 - 20)	-	-	↓
Female (Weeks 6, 7 and 9 - 20)	-	-	↓
Female (Week 16)	↓	-	↓
Sire at Delivery	-	-	↓
Dam at Delivery	-	↓	↓
Lactating Females (PND 4, 7 and 14)	-	-	↓
Feed Consumption (g/kg body weight/day)			
Male (Weeks 2 - 5, 8 and 10)	-	-	↓
Male (Week 13)	↓	↓	↓
Male (Weeks 6 - 7)	↓	-	↓
Male (Week 17)	↓	-	-
Female (Weeks 2 - 10)	-	-	↓
Lactating Females (PND 4 - 7)	↓	-	↓
Lactating Females (PND 11 - 14)	-	↑	-
Lactating Females (PND 18 - 21)	-	-	↓

KEY: M = Male ↑ = Significant increase in the parameter.
 F = Female ↓ = Significant decrease in the parameter.
 - = No observed effect.

SUMMARY OF RESULTS (CONTINUED)
TWO-GENERATION REPRODUCTION TOXICITY STUDY OF PROPYLTHIOURACIL WHEN ADMINISTERED TO SPRAGUE-DAWLEY RATS IN THE DRINKING WATER

Parameters	Treatment Group (%)		
	Low (0.0001)	Mid (0.0004)	High (0.0015)
Feed Consumption (g/animal/day)			
Male (Weeks 2 - 10 and 13 - 19)	-	-	↓
Male (Week 13)	-	↓	↓
Female (Weeks 3 - 10 and 13 - 19)	-	-	↓
Female (Week 6)	-	↓	↓
Lactating Females (PND 1 - 4)	↓	-	↓
Lactating Females (PND 4 - 7)	↓	↓	↓
Lactating Females (PND 7 - 11 and 18 - 21)	-	-	↓
Lactating Females (PND 11 - 14)	-	↑	-
Water Consumption (g/kg body weight/day)			
Female (Week 18)	-	↓	↓
Lactating Females (PND 4 - 7, 11 - 14 and 18 - 21)	-	-	↓
Water Consumption (g/animal/day)			
Male (Weeks 3 - 10, 13 - 15 and 18 - 19)	-	-	↓
Male (Week 17)	↓	↓	↓
Female (Weeks 4, 8, 10, 13 and 19)	-	-	↓
Female (Week 18)	-	↓	↓
Lactating Females (PND 1 - 4, 4 - 7, 7 - 11 and 11 - 14)	-	-	↓
Lactating Females (PND 14 - 18 and 18 - 21)	↓	-	↓
Estrous Cyclicity			
Number of Females with Regular Cycle	-	-	↓
Relative Amount of Time Spent in Estrous Stages	-	-	*

KEY: M = Male 8 = Significant increase in the parameter.
 F = Female ↓ = Significant decrease in the parameter.
 * = Significant change in the parameter
 - = No observed effect.

SUMMARY OF RESULTS (continued)
TWO-GENERATION REPRODUCTION TOXICITY STUDY OF PROPYLTHIOURACIL WHEN ADMINISTERED TO SPRAGUE-DAWLEY RATS IN
THE DRINKING WATER

Parameters	Treatment Group (%)		
	Low (0.0001)	Mid (0.0004)	High (0.0015)
Male Necropsy			
Terminal Body Weight	-	-	↓
Absolute Thyroid/Parathyroid Weight	-	↑	↑
Relative Brain Weight	-	-	↑
Relative Pituitary Weight	-	↑	↑
Relative Right Testis Weight	-	-	↑
Relative Seminal Vesicle Weight	-	-	↑
Relative Thyroid/Parathyroid Weight	-	↑	↑
Female Necropsy			
Terminal Body Weight	-	-	↓
Absolute Adrenal Weight	-	-	↓
Absolute Brain Weight	-	-	↓
Absolute Kidney Weight	-	-	↓
Absolute Liver Weight	-	-	↓

KEY: M = Male ↑ = Significant increase in the parameter.
 F = Female ↓ = Significant decrease in the parameter.
 - = No observed effect.

SUMMARY OF RESULTS (CONTINUED)
 TWO-GENERATION REPRODUCTION TOXICITY STUDY OF PROPYLTHIOURACIL WHEN ADMINISTERED TO SPRAGUE-DAWLEY RATS IN THE DRINKING WATER

Parameters	Treatment Group (%)		
	Low (0.0001)	Mid (0.0004)	High (0.0015)
Female Necropsy			
Absolute Spleen Weight	-	-	↓
Absolute Thyroid/Parathyroid Weight	-	-	↑
Relative Thyroid/Parathyroid Weight	↑	↑	↑
Microscopic Findings			
Male Thyroid - Follicular Cell Hyperplasia	-	↑	↑
Male Testes - Germinal Epithelium Degeneration	↑	↑	↑
Female Thyroid - Follicular Cell Hyperplasia	-	↑	↑
Thyroid Hormones			
Male TSH	-	↑	↑
Male T4	-	↓	↓
Female TSH	-	↑	↑
Female T4	-	↓*	↓*
Sperm Data			
Morphology - % of Abnormal Sperm	↑	-	-
Mortality	(Control) 0M,0F 1M,0F	0M,0F	0M,2F

KEY: M = Male ↑ = Significant increase in the parameter.
 F = Female ↓ = Significant decrease in the parameter.
 * = These decreases were not statistically significant at p<0.05.
 - = No observed effect.

SUMMARY OF RESULTS (CONTINUED)
TWO-GENERATION REPRODUCTION TOXICITY STUDY OF PROPYLTHIOURACIL WHEN ADMINISTERED TO SPRAGUE-DAWLEY RATS IN THE DRINKING WATER

Parameters	Treatment Group (%)		
	Low (0.0001)	Mid (0.0004)	High (0.0015)
<u>Reproductive and Developmental Parameters - F₀ Cohabitation</u>			
Live Pups per Litter			
Female	-	-	↓
Male and Female	-	-	↓
Average Pup Weights			
Male (PND 14 and 21)	-	-	↓
Female (PND 14 and 21)	-	-	↓
Day of Pup Eye Opening	-	↑	a
Day of Pup Vaginal Opening	-	↑	a
Day of Pup Preputial Separation	-	↑	a
F ₁ Male PND 21 Necropsy			
Terminal Body Weight	-	-	↓
Absolute Brain Weight	-	-	↓
Relative Brain Weight	-	-	↑
Relative Spleen Weight	-	-	↓
Relative Thymus Weight	-	-	↓

KEY: M = Male ↑ = Significant increase in the parameter.
 F = Female ↓ = Significant decrease in the parameter.
 - = No observed effect.

a These parameters were not evaluated in the 0.0015% F₁ animals due to increased mortality.

SUMMARY OF RESULTS (CONTINUED)
 TWO-GENERATION REPRODUCTION TOXICITY STUDY OF PROPYLTHIOURACIL WHEN ADMINISTERED TO SPRAGUE-DAWLEY RATS IN THE DRINKING WATER

Parameters	Treatment Group (%)		
	Low (0.0001)	Mid (0.0004)	High (0.0015) ^a
F ₁ Female PND 21 Necropsy			
Terminal Body Weight	-	-	↓
Absolute Brain Weight	-	-	↓
Relative Brain Weight	-	-	↑
Relative Spleen Weight	-	-	↓
Relative Thymus Weight	-	-	↓
<u>General Toxicity - F₁ Cohabitation</u>			
Feed Consumption (g/animal/day)			
Male (Week 3)	↓	-	
Lactating Female (PND 4 - 7 and 7 - 11)	↓	-	
Lactating Female (PND 18 - 21)	-	↓	
Water Consumption (g/animal/day)			
Lactating Female (PND 4 - 7)	↓	↓	
Lactating Female (PND 1 - 4, 7 - 11 and 11 - 14)	↓	-	
Male Necropsy			
Absolute Thyroid/Parathyroid Weight	-	↑	
Relative Thyroid/Parathyroid Weight	↑	↑	

KEY: M = Male ↑ = Significant increase in the parameter.
 F = Female ↓ = Significant decrease in the parameter.
 - = No observed effect.

^a This dose level was discontinued for the F₁ Cohabitation due to high mortality observed among F₁ adolescents.

SUMMARY OF RESULTS (CONTINUED)
TWO-GENERATION REPRODUCTION TOXICITY STUDY OF PROPYLTHIOURACIL WHEN ADMINISTERED TO SPRAGUE-DAWLEY RATS IN THE DRINKING WATER

Parameters	Treatment Group (%)		
	Low (0.0001)	Mid (0.0004)	High (0.0015)
Female Necropsy			
Absolute Thyroid/Parathyroid Weight	-	↑	
Relative Thyroid/Parathyroid Weight	-	↑	
Microscopic Findings			
Male Thyroid – Follicular Cell Hyperplasia	-	↑	
Male Testes – Germinal Epithelium Degeneration	↑	↑	
Thyroid Hormones			
Male TSH	-	↑	
Male T4	-	↓	
Female TSH	-	↑	
Female T4	-	↓	
Mortality			
Adolescent	(Control) 0M,0F	0M,0F	0M,0F
Adult	(Control) 1M,0F	0M,0F	23M,20F
<u>Reproductive and Developmental Parameters – F₁ Cohabitation</u>			
Anogenital Distance (AGD)			
Male AGD	↓	↓	
AGD/Male Pup Weight Ratio	-	↓	

KEY: M = Male ↑ = Significant increase in the parameter.
 F = Female ↓ = Significant decrease in the parameter.
 - = No observed effect.

SUMMARY OF RESULTS (CONTINUED)
TWO-GENERATION REPRODUCTION TOXICITY STUDY OF PROPYLTHIOURACIL WHEN ADMINISTERED TO SPRAGUE-DAWLEY RATS IN THE DRINKING WATER

Parameters	Treatment Group (%)	
	Low (0.0001)	Mid (0.0004)
F ₁ Male PND 21 Necropsy		
Relative Spleen Weight	↑	-
F ₁ Female PND 21 Necropsy		
Absolute Thymus Weight	-	↓
Relative Thymus Weight	-	↓

The F₁ cohabitation started on PND 99 ±10.

KEY: M = Male ↑ = Significant increase in the parameter.
 F = Female ↓ = Significant decrease in the parameter.
 - = No observed effect.

REPRODUCTIVE TOXICANT:
 Male - Yes
 Female - Yes

CLASSIFICATION:

		<u>0.0001 %</u>	<u>0.0004 %</u>	<u>0.0015 %</u>
Males	Reproductive Toxicity	+	+	+
	General Toxicity	-	+	+
Females	Reproductive/Developmental Toxicity	-	+	+
	General Toxicity	-	+	+

INTRODUCTION

Propylthiouracil (CAS No. 51-52-5) is a thyroid hormone-synthesis inhibitor and antithyroid agent for the treatment of hyperthyroidism. Propylthiouracil (PTU) has been shown to decrease T3 and T4 while increasing TSH. This study was designed to validate a Two-Generation Study Model proposed to identify potent and weak thyroid toxicants. The dose levels selected for the study were 0.0001, 0.0004, and 0.0015 % PTU (weight/volume). The high dose of 0.0015% was expected to result in increased TSH, decreased T3 and T4, thyroid pathology, and decreased growth. The remaining dose levels were expected to result in less toxicity with no effect at 0.0001% PTU.

METHODS

General Study Design

A schematic diagram of this study is presented in Figure 1. The F₀ Cohabitation consisted of a control group and three treated groups (20 pairs/group). F₀ males and females were administered propylthiouracil in the drinking water starting on SD 1 and continuing until necropsy. F₀ animals had body weights collected at randomization, weekly, and at littering and feed and water consumption measured weekly when housed individually. During the lactation period for the F₁ litter, F₀ females also had body weights measured on PND 1, 4, 7, 14, and 21 and feed and water consumption measured for PND 1-4, 4-7, 7-11, 11-14, 14-18 and 18-21. Physical examinations were performed at randomization, at initiation of dosing, and weekly thereafter. Prior to cohabitation, vaginal cytology was conducted for 14 days on the F₀ dams. Following ten weeks of pre-mating exposure to propylthiouracil, the F₀ animals were housed as breeding pairs (1:1 ratio). Vaginal smears were examined daily for confirmation of mating. When sperm or plug positive or after 14 days of cohabitation, the females were separated from the males. The F₁ pups were reared by the dam until weaning on PND 21. On PND 1, 4, 7, 14, and 21 the pups were counted and weighed. All pups had their anogenital distance (AGD) and individual weight recorded on PND 1 and were examined for pinna detachment and eye opening beginning on PND 2. All male pups were examined for retained nipples on PND 12 and 13.

On PND 16, 1-2 males and 1-2 females from each litter were randomly selected for rearing for the F₁ mating trial. These animals were assigned a unique identification number which was tattooed on the tail. Observations of testicular descent (starting on PND 16), vaginal opening (starting on PND 25), and preputial separation (starting on PND 35) were conducted. On PND 21, animals selected for the F₁ cohabitation were separated from the dam and housed. Pups selected for the F₁ cohabitation period were administered propylthiouracil in the drinking water starting on PND 21 (dose levels of 0.0001, 0.0004, and 0.0015 % PTU). Body weights were collected weekly starting on PND 21.

Three additional males and three additional females from all F₁ litters (when available) were randomly selected on PND 16 for the PND 21 necropsy. On PND 21 these animals were necropsied, terminal body weights and organ weights were obtained, and tissues were saved for histopathologic evaluation. The remaining F₁ pups were euthanized and discarded without necropsy on PND 21.

After the lactation phase was completed, terminal body weights were obtained for all F₀ animals, then the animals were euthanized and necropsied, organ weights were obtained, sperm analyses were performed, and tissues were saved for possible histopathologic evaluation.

F₁ weanlings selected for the F₁ cohabitation were reared in same sex groups until PND 99 ±10 days of age, when twenty animals of each sex in each dose group were randomly assigned to breeding pairs (avoiding sibling matings) and cohabited. Vaginal smears were examined daily for confirmation of mating. When sperm or plug positive or after 14 days of cohabitation, the females were separated from the males. Once cohabited, F₁ animals had body weights collected weekly and at littering. Feed and water consumption were measured weekly when the animals were housed individually (i.e., not during cohabitation). During the lactation period for the F₂ litter, F₁ females also had body weights measured on PND 1, 4, 7, 14, and 21 and feed and water consumption measured for PND 1-4, 4-7, 7-11, 11-14, 14-18 and 18-21. Physical examinations were performed weekly.

The F₂ pups were reared by the dam until PND 21. On PND 1, 4, 7, 14, and 21, the pups were counted and weighed. All pups had their AGD and individual weight recorded on PND 1 and were examined for pinna detachment and eye opening beginning on PND 2. All male pups were


examined for retained nipples on PND 12 and 13. Vaginal cytology was conducted on the F₁ dams for 14 days beginning at least 4 days after the last F₂ PND 21.

Three males and three females from all litters (where available) were randomly selected on PND 16 for the PND 21 necropsy. On PND 21 these animals were necropsied, terminal body weights and organ weights were obtained, and tissues were saved for histopathologic evaluation. The remaining F₂ pups were euthanized and discarded without necropsy on PND 21.

The F₁ animals were retained until completion of vaginal cytology. Terminal body weights were obtained for all F₁ animals, the animals were euthanized and necropsied, organ weights were obtained, sperm analyses were performed, and tissues were saved for possible histopathologic evaluation.

Figure 1

Two-Generation Reproduction Toxicity Study Flow Diagram for Propylthiouracil



Study Animals

The Sprague-Dawley rat was selected as the test animal due to its established quality as a breeder and the availability of historical toxicologic data for reference. For the F₀ cohabitation, 170 (85 male and 85 female) Sprague-Dawley CrI:CD® BR rats were received from Charles River Laboratories (Raleigh, NC) on May 9, 2000, and assigned temporary animal numbers. Seven days after receipt, all of the animals were weighed and randomly assigned to one of four groups by a computer-generated randomization procedure that ensured equal weight distribution between the groups. All study animals were assigned unique animal numbers and individually identified with the animal number on their tail by tattooing. Dosing was initiated after a 14 day acclimation period. During this period, two males and two females were forwarded to Anmed Biosafe, Inc., Rockville, Maryland, for determination of viral antibody titers. All sera were negative for antibodies at that time (Appendix table A2-4). Extra animals were subsequently removed from the study room. The study animals were approximately 7-8 weeks of age at the initiation of dosing and the body weight range of the animals was 232.3 – 293.9 g for males and 160.1 – 215.3 g for females.

The F₁ and F₂ animals were born at TherImmune Research Corporation. During the F₁ and F₂ litters, pup selection occurred on PND 16. On PND 16 for the F₁ litter, at least two males and two females from each litter, when available, were randomly selected for the F₁ cohabitation and identified with tail tattoos. On PND 21, these animals were housed separately from the dam, and dosing of the selected F₁ animals was initiated. On PND 16 for the F₁ and F₂ animals, three males and three females from each litter, when available, were randomly selected for the PND 21 necropsy. F₁ and F₂ pups not selected for PND 21 necropsy were euthanized by carbon dioxide asphyxiation and discarded without necropsy. For the F₁ animals, approximately one week before PND 99 ± 10 (initiation of F₁ cohabitation), one male was assigned to one female, avoiding sibling matings, to form twenty mating pairs per group for the F₁ cohabitation. Animals not selected for the F₁ cohabitation were euthanized via carbon dioxide asphyxiation and discarded without necropsy one week after the initiation of the F₁ cohabitation.

Animal Husbandry and Environmental Conditions

Animals were housed two per cage by sex from receipt to randomization, two per cage (one male and one female) during the F₀ and F₁ cohabitations, two per cage (same sex) during the F₁ growth phase, and individually at all other times. All animals were housed in polycarbonate cages (19"L x 10.5"W x 8"H) suspended on stainless-steel racks. Racks were equipped with filter paper liners. Polycarbonate caging contained Sani Chip certified heat treated hardwood laboratory bedding. Pelleted Harlan Teklad™ NIH-07 Small Animal Feed was available *ad libitum* in stainless-steel hanging feeders, and was used within five months of the milling date. The feed was analyzed for nutrients, aflatoxins, nitrosamines, heavy metals, chlorinated hydrocarbons, organophosphates, PCBs, nitrites, nitrates, BHA, BHT, total bacterial plates, coliforms, *E. coli*, and *Salmonella* by the vendor. Deionized water was provided in water bottles, which were changed at least weekly. The test article was given to the animals in the deionized water. The water is routinely analyzed for total dissolved solids, heavy metals, chlorinated hydrocarbons, organophosphates, nitrates, nitrites, microbiological content and total trihalomethanes at least semi-annually to conform with the Safe Drinking Water Act. None of the feed, water, or bedding contaminants was at levels believed sufficient to interfere with the study.

A 12-hour light/12-hour dark cycle was maintained throughout the study. During the study, the temperature range in the animal room was 68-74°F and the relative percent humidity was 30-70%, except as noted in Appendix Table A2-1.

Test Article

6-Propyl-2-thiouracil, lot no. 47H2500, was received from Battelle, Columbus, OH, on April 20, 2000 and June 16, 2000. It was described as a white powder with a purity of 99.8%. Prior to use, two 0.5 g samples of bulk test article were collected into glass bottles with Teflon® coated lids, sealed, and stored at approximately -20°C protected from light for possible future reanalysis. At the start of the F₁ cohabitation, and again after the end of the in-life portion of the study, five grams of bulk sample were sent to Battelle for analysis.

Each time a new mix or batch was prepared, two 50 mL archival samples of each dose level formulation were collected and stored at TherImmune in amber glass bottles with Teflon® coated

lids protected from light in the refrigerator. One sample of each dose level formulation from mixes 3, 4, 5, 11, 27, 51 was sent to Battelle for analysis. Archival samples which were not selected for analysis were discarded as hazardous waste at least 90 days after preparation.

Dosage Formulation and Administration

Formulations of 0, 0.0001, 0.0004, and 0.0015 % (w/v) were prepared at least weekly throughout the study. For Group 1 (control), a precalibrated carboy was filled with the required volume of deionized water. For all other groups, precalibrated carboys were initially filled to approximately 75% of the required volume with deionized water. The required quantity of 6-propyl-2-thiouracil was weighed into a weigh boat and poured into a volumetric flask (approximately half-filled with deionized water). The weigh boat was rinsed at least three times with deionized water and the rinse was added to the flask. The flasks were mixed until the propylthiouracil was completely dissolved. The solution was then poured into a carboy, the flask was rinsed at least three times with deionized water and the rinse was added to the carboy, deionized water was added to the carboy to achieve the required volume, and the solution was mixed with a variable speed stirrer to ensure complete dissolution.

The formulations were stored refrigerated and protected from light. Under these conditions, formulations are reported to be stable for 35 days. The formulations were dispensed into amber glass bottles with neoprene stoppers and stainless steel sipper tubes. Dispensed formulations are stable for seven days. The stability and storage of formulations conditions were based upon the Dose Formulation Developmental Study Report provided by Battelle.

F₀ Cohabitation

The F₀ cohabitation consisted of a control group and three treated groups (20 pairs/group). F₀ animals were administered propylthiouracil in their drinking water at doses of 0 (control), 0.0001, 0.0004, or 0.0015 % from SD 1 until necropsy. Prior to cohabitation, vaginal cytology was conducted for 14 days on the F₀ dams. Following ten weeks of pre-mating exposure to propylthiouracil (i.e., on SD 71), the animals were housed as breeding pairs (1:1 ratio). Vaginal smears were examined daily for confirmation of mating. When sperm or plug positive or after 14

days of cohabitation, the females were separated from the males. The F₁ litter was reared by the dam until weaning on PND 21. The total number of pups, number of live and dead pups, the number of live male and female pups, and total body weight of live male and female pups were recorded on PND 1, 4, 7, 14, and 21. On PND 1, all pups had individual pup body weights and AGD measured, and the dam and sire weights were recorded. The dam was also weighed on PND 4, 7, 14 and 21, and feed and water consumption were measured for PND 1-4, 4-7, 7-11, 11-14, 14-18 and 18-21. All pups were examined for pinna detachment and eye opening beginning on PND 2. All male pups were examined for retained nipples on PND 12 and 13.

On PND 16, 1-2 males and 1-2 females from each litter were randomly selected for rearing for the F₁ mating trial. These animals were assigned a unique identification number and tail tattooed. Observations of testicular descent (starting on PND 16), vaginal opening (starting on PND 25), and preputial separation (starting on PND 35) were conducted. On PND 21, animals selected for the F₁ cohabitation were housed separately from the dam in same-sex pairs and drinking water dosing was initiated. Body weights were collected weekly starting on PND 21.

Three additional males and females from all litters (when available) were randomly selected on PND 16 for the PND 21 necropsy. On PND 21 these animals were necropsied, terminal body weights and organ weights were obtained, and tissues were saved for histopathologic evaluation. Pups not selected were euthanized by carbon dioxide asphyxiation and discarded without necropsy on PND 21.

Approximately one week before PND 99 ± 10 (i.e., the initiation of F₁ cohabitation), one male was assigned to one female, avoiding sibling matings, to form twenty mating pairs per group for the F₁ cohabitation. Animals not selected for the F₁ cohabitation were euthanized by carbon dioxide asphyxiation and discarded without necropsy.

All animals were observed twice daily for mortality and signs of toxicity. In males and non-lactating females, body weights were collected weekly and at littering (males only), and feed and water consumption was measured weekly when animals were housed individually. Physical examinations were performed weekly.

F₁ PND 21 Necropsy and Terminal Procedures

On PND 21, up to three surviving F₁ males and females per litter were weighed, sacrificed by carbon dioxide asphyxiation, and exsanguinated. A gross necropsy was performed on all animals. Necropsies were performed by trained personnel from Pathology Associates International (PAI, Frederick, MD) under the direct supervision of a Board-certified pathologist. Necropsies included examination of the external surface of the body, all orifices, and the cranial, thoracic, and abdominal cavities and their contents.

The following organs were weighed:

brain
spleen
thymus

The adrenals, kidneys, liver, pituitary, ventral prostate, dorsolateral prostate, seminal vesicles with coagulating glands, spleen, thyroid/parathyroids, vagina/cervix/uterus and gross lesions from all necropsied animals were preserved in 10% neutral-buffered formalin (NBF) for histopathological examination. The left testis and epididymis were fixed in a 2% para-formaldehyde/3% glutaraldehyde solution for 3-5 days and then transferred to phosphate buffered saline. The right testis was frozen at approximately -80°C to be used for spermatid head counts. (Spermatid head counts for PND 21 animals were discontinued because no sperm were found to be present in the testes of these animals.) The ovaries were preserved in Bouin's for 24-48 hours and then transferred to 70% ethanol. Histopathology was not required for these tissues.

F₀ Adult Necropsy and Terminal Procedures

After the end of lactation for all F₁ pups, terminal body weights were obtained from all surviving F₀ animals. Blood was collected from the orbital sinus under 70% O₂/30% CO₂ anesthesia. The blood was centrifuged and separated to obtain plasma. The plasma was frozen at approximately -80° and was forwarded to Anilytics, Inc. (Gaithersburg, MD) for determination of TSH, T3, and T4 levels. The methods used for the thyroid hormone analyses are presented in Appendix 7. Following blood collection, the animals were sacrificed by carbon dioxide asphyxiation and exsanguinated. A gross necropsy was performed on all animals. Necropsies were performed by trained personnel from PAI under the direct supervision of a Board-certified pathologist. Necropsies included examination of the external surface of the body, all orifices, and the cranial, thoracic, and abdominal cavities and their contents.

The following organs were weighed:

adrenals (paired)	brain
cauda epididymis (right)	epididymis (right)
kidneys (paired)	liver
ovaries (paired)	pituitary
prostate (ventral)	prostate (dorsolateral)
seminal vesicles with coagulating glands	spleen
testis (right)	thyroids/parathyroids
uterus/vagina/cervix	

The adrenals, brain, kidneys, liver, pituitary, ventral prostate, dorsolateral prostate, seminal vesicles with coagulating glands, spleen, thyroid/parathyroids, vagina/cervix/uterus, and gross lesions from all necropsied animals were preserved in 10% NBF for histopathological examination. The left testis and epididymis were fixed in a 2% para-formaldehyde/3% glutaraldehyde solution for 3-5 days and then transferred to phosphate buffered saline. The right testis was frozen at approximately -80°C and used for spermatid head counts. The ovaries were preserved in Bouin's for 24-48 hours and then transferred to 70% ethanol. The thyroid/parathyroids, ovaries, uterus, cervix, vagina, and gross lesions from the first ten surviving males and females per group were embedded in paraffin, sectioned, stained with hematoxylin and eosin, and examined microscopically by a pathologist. The left testis and epididymis from the first ten surviving males per group were

processed through glycol methacrylate (GMA), sectioned, stained with periodic acid-Schiff (PAS) and hematoxylin, and microscopically evaluated by a pathologist.

Computer-assisted sperm motion analysis using Hamilton-Thorne Research Version 10 IVOS Sperm Analysis System (Hamilton-Thorne Research, Beverly, Massachusetts), epididymal sperm density, sperm morphology, and testicular spermatid head count data were measured on all males surviving to termination. The right vas deferens was used for sperm motility analysis; the right cauda epididymis was used for sperm density and morphology; and the right testis was used for evaluation of spermatid head counts. After conducting the epididymal sperm density counts, approximately 5mL of each sperm suspension was saved, frozen, and sent to an NTP subcontractor.

Animals found dead or euthanized *in extremis* during the study (all generations) were subject to a gross necropsy. The following tissues were retained in NBF for possible histopathological examination:

adrenals	brain
kidneys	liver
left epididymis	left testis
ovaries (transferred to 70% ethanol within 24-48 hours)	pituitary
seminal vesicles/coagulating glands	prostate (ventral and dorso-lateral lobes)
stomach	spleen
vagina/uterus/cervix	thyroid/parathyroids
	gross lesions

The lower and upper jaws of ten F₁ 0.0015% (Group 4) animals (one animal/sex/litter) were embedded in paraffin, step sectioned, and examined microscopically by the study pathologist.

Allocation of F₁ Weanlings

On PND 16, 1 - 2 males and 1 - 2 females per litter, when available, were randomly selected for the F₁ cohabitation. These animals were assigned a unique identification number and tail tattooed. Observations of testicular descent (starting on PND 16), vaginal opening (starting on PND 25), and preputial separation (starting on PND 35) were conducted until the endpoint was seen. On PND 21, animals selected for the F₁ cohabitation were housed separately from the dam, and administration of propylthiouracil in drinking water was initiated. On PND 21, pups not selected for

the F₁ cohabitation or PND 21 necropsy were euthanized and discarded without necropsy. All animals were observed twice daily for mortality and signs of toxicity. Following weaning of the final F₁ litter, all weekly body weights and physical examinations were adjusted to occur on the same day and began on PND 29 ± 10.

Due to a high level of mortality in the 0.0015 % F₁ animals following weaning, all surviving animals in this group were removed from the study and euthanized.

F₁ Cohabitation

The F₁ cohabitation consisted of a control group and two treated groups (20 pairs/group). F₁ weanlings selected for the F₁ cohabitation were reared in same sex groups until PND 99 ± 10 days of age when twenty animals of each sex in each dose group were randomly assigned to breeding pairs (avoiding sibling matings) and cohabited. Vaginal smears were examined daily for confirmation of mating. When sperm or plug positive or after 14 days of cohabitation, the females were separated from the males. The F₂ litter was reared by the dam until PND 21. The total number of pups, number of live and dead pups, the number of live male and female pups, and total body weight of live male and female pups were recorded on PND 1, 4, 7, 14, and 21. On PND 1, all pups had individual pup body weights and AGD measured, and the dam and sire weights were recorded. The dam was also weighed on PND 4, 7, 14 and 21, and feed and water consumption were measured for PND 1-4, 4-7, 7-11, 11-14, 14-18 and 18-21. All pups were examined for pinna detachment and eye opening beginning on PND 2. All male pups were examined for retained nipples on PND 12 and 13. Weekly physical examinations and body weights were continued. In males and non-lactating females, body weights were collected weekly and at littering (males only), and feed and water consumption was measured weekly, when animals were housed individually. Physical examinations were performed weekly. All F₁ adults were observed twice daily for mortality and signs of toxicity. All animals not selected for the F₁ cohabitation were euthanized and discarded without necropsy on PND 106 ± 10.

Three males and three females from all litters (where available) were randomly selected on PND 16 for the PND 21 necropsy. On PND 21 for these animals, terminal body weights were

recorded, the animals euthanized and necropsied, organ weights were obtained, and tissues were saved for histopathologic evaluation.

Pups not selected were euthanized and discarded without necropsy on PND 21.

At least 4 days after the delivery of the last F₂ litters, vaginal cytology was conducted for 14 days on the F₁ dams.

F₂ PND 21 Necropsy and Terminal Procedures

Up to three surviving F₁ males and females per litter were weighed, sacrificed by carbon dioxide asphyxiation, and exsanguinated. A gross necropsy was performed on all animals. Necropsies were performed by trained personnel from PAI under the direct supervision of a Board-certified pathologist. Necropsies included examination of the external surface of the body, all orifices, and the cranial, thoracic, and abdominal cavities and their contents.

The following organs were weighed:

- brain
- spleen
- thymus

The adrenals, kidneys, liver, pituitary, ventral prostate, dorsolateral prostate, seminal vesicles with coagulating glands, spleen, thyroid/parathyroids, vagina/cervix/uterus and gross lesions from all necropsied animals were preserved in 10% NBF for histopathological examination. The left testis and epididymis were fixed in a 2% para-formaldehyde/3% glutaraldehyde solution for 3-5 days and then transferred to phosphate buffered saline. The right testis was frozen at approximately -80°C to be used for spermatid head counts. (Spermatid head counts for PND 21 animals were discontinued because no sperm were found in the testes of these animals.) The ovaries were preserved in Bouin's for 24-48 hours and then transferred to 70% ethanol. Histopathology was not performed on these tissues.

F₁ Adult Necropsy and Terminal Procedures

After the conclusion of vaginal cytology for all F₁ females, terminal body weights were obtained from all surviving F₁ animals. Blood was collected from all animals from the orbital sinus under 70% O₂/30% CO₂ anesthesia. The blood was centrifuged and separated to obtain plasma. The plasma from each animal was then split into aliquots and frozen at approximately -80°. One aliquot was sent to Anilytics, Inc. (Gaithersburg, MD) for determination of TSH, T3, and T4 levels. The other aliquot was retained at TherImmune for possible future reanalysis. The methods used for the thyroid hormone analyses are presented in Appendix 7. Following blood collection, the animals were euthanized by carbon dioxide asphyxiation and exsanguinated. A gross necropsy was performed on all animals. Necropsies were performed by trained personnel from PAI under the direct supervision of a Board-certified pathologist. Necropsies included examination of the external surface of the body, all orifices, and the cranial, thoracic, and abdominal cavities and their contents.

The following organs were weighed:

adrenals (paired)	brain
cauda epididymis (right)	epididymis (right)
kidneys (paired)	liver
ovaries (paired)	pituitary
prostate (ventral)	prostate (dorsolateral)
seminal vesicles with coagulating glands	spleen
testis (right)	thyroids/parathyroids
uterus/vagina/cervix	

The adrenals, brain, kidneys, liver, pituitary, ventral prostate, dorsolateral prostate, seminal vesicles with coagulating glands, spleen, thyroid/parathyroids, vagina/cervix/uterus, and gross lesions from all necropsied animals were preserved in 10% neutral-buffered formalin for histopathological examination. The left testis and epididymis were fixed in a 2% paraformaldehyde/3% glutaraldehyde solution for 3-5 days and then transferred to phosphate buffered saline. The right testis was frozen at approximately -80°C and used for spermatid head counts. The ovaries were preserved in Bouin's for 24-48 hours and then transferred to 70% ethanol. The thyroid/parathyroids, ovaries, uterus, cervix, vagina, and gross lesions from the first ten surviving males and females per group were embedded in paraffin, sectioned, stained with hematoxylin and eosin, and examined microscopically by a pathologist. The left testis and epididymis from the first

ten surviving males per group were processed through glycol methacrylate (GMA), sectioned, stained with periodic acid-Schiff (PAS) and hematoxylin, and microscopically evaluated.

Computer-assisted sperm motion analysis using Hamilton-Thorne Research Version 10 IVOS Sperm Analysis System (Hamilton-Thorne Research, Beverly, Massachusetts), epididymal sperm density, sperm morphology, and testicular spermatid head count data were measured on all males surviving to termination. The right vas deferens was used for sperm motility analysis; the right cauda epididymis was used for sperm density and morphology; and the right testis was used for evaluation of spermatid head counts. After conducting the epididymal sperm density counts, approximately 5 mL of each sperm suspension was saved, frozen, and sent to an NTP subcontractor.

Statistical Analyses

Data were statistically analyzed by Analytical Sciences Inc. (Durham, NC). Most hypotheses (including but not limited to pup weights, body weights, feed and water consumption, organ weights, sperm parameters, and estrous cycle length) were tested using the nonparametric multiple comparisons procedure of Dunn (1964) or Shirley (1977), as modified by Williams (1986). Shirley's test was designed to detect treatment-related differences when the response to treatment consistently increased (or decreased) with increasing dose. Although the test employs a smoothing algorithm to adjust for dose-response inversions, Dunn's test was more appropriate if the departure from monotonicity was severe. Jonckheere's test (1954) was used to ascertain whether there was sufficient evidence of a dose-related response to apply Shirley's test. If the p-value from Jonckheere's test was less than 0.01, Shirley's test was used; otherwise, Dunn's test was applied. Reproductive and developmental data were analyzed nested by dam.

For data expressed as a proportion, such as number fertile/number cohabitated, the Cochran-Armitage test (Armitage, 1971) was used to test for a dose-related trend, and pairwise comparisons were performed using a chi-squared test (Conover, 1971).

The ratio of the number of pups born alive to the total number of pups carried to full term was computed for all fertile pairs. The sex ratio, expressed as the proportion of males, was computed for all fertile pairs with at least one live pup. Shirley's or Dunn's test was used to compare dosed groups to controls based on Jonckheere's test, as described above.

Since the number of pups in a litter may influence the average pup weight, a parametric analysis of covariance (Neter and Wasserman, 1974) was used to test overall equality in average pup weight, after adjustment for average litter size. The covariant used was average litter size, including live and dead pups. Least squares estimates of dose group means adjusted for litter size were computed and tested for overall equality using an F-test and pairwise equality using Dunnett's test (1955). Unadjusted weights were analyzed with Shirley's or Dunn's test.

Absolute organ weights were analyzed using body weight as a covariate. Adjusted mean dose effects were compared to the control with Dunnett's test.

Litter sizes and number of litters in dose groups were compared to controls using Dunn's or Shirley's test. To examine potential differences in treatment effects on males and females, number of male pups, number of female pups, and total number of pups in litters in treated groups were compared to controls.

Using either Shirley's or Dunn's test, feed and water consumption data were analyzed as g/animal/day and g/kg body weight/day. Sperm parameters were analyzed by Shirley's or Dunn's test.

The relative time spent in estrous cycle stages was analyzed using a multivariate analysis of variance (Wilks criterion) to test for the simultaneous equality of measurements across dose levels (Morrison, 1976). Before applying the test, an arcsine transformation was performed to bring the data into closer conformance with normality assumptions.

Thyroid hormone data were analyzed using Dunnett's one-tailed test followed by Jonckheere's trend test.

All findings described in this report as "increased" or "decreased" were statistically significant as compared to the control group at the 95% confidence level ($p < 0.05$).

Specimen, Raw Data and Final Report Storage

Upon submission of the final report, all original study records, including all original data sheets; the original final report; all biological samples; tissues; sperm morphology slides; sperm data optical disks; computer printouts generated in the statistical analysis of the sperm data; and copies of the final report, will be forwarded to the contracting agency, the NIEHS, Research Triangle Park, NC. Copies of the final study report will also be filed with TherImmune Research Corporation.

RESULTS

F₀ Cohabitation

During Weeks 3-20 of the F₀ Cohabitation, mean body weights of the 0.0015% males were decreased by 6-21% (Table 0-1A). Mean body weights of the 0.0015% females were decreased by 8-17% during Weeks 6-7 and 9-20. Mean body weights for the 0.0001% females were decreased by 6% during Week 16 (Table 0-1B).

Feed consumption was decreased by 4-11% during Weeks 6-7, 13, and 17 in the 0.0001% males, by 11% during Week 13 in the 0.0004% males, by 6-18% during Weeks 2-8, 10, and 13 in the 0.0015% males, and by 3-17% during Weeks 2-10 in the 0.0015% females, when examined as g/kg body weight/day (Table 0-2A). Feed consumption examined as g/animal/day was decreased by 12% during Week 13 in the 0.0004% males, by 12-35% during Weeks 2-10 and 13-19 in the 0.0015% males, by 9% during Week 6 in the 0.0004% females, and by 12-25% during Weeks 3-10, 13, and 18-19 in the 0.0015% females (Table 0-2B).

Water consumption was decreased in the 0.0004 and 0.0015% females by 15% and 31%, respectively, during Week 18 when examined as g/kg body weight/day (Table 0-3A). When examined as g/animal/day, water consumption was decreased by 16% during Week 17 in the 0.0001 and 0.0004% males, by 14-23% during Weeks 2-10, 13-15, and 17-19 in the 0.0015% males, by 18% during Week 18 in the 0.0004% females, and by 13-38% during Weeks 4, 8, 10, 13, and 18-19 in the 0.0015% females (Table 0-3B).

Clinical signs noted in the F₀ animals included abrasions, alopecia, anorexia, discharge from the eye, hunched posture, lacrimation, languid behavior, loss of appetite, rough haircoat, swelling, vaginal discharge, and thinness. The incidence of these observations was low (0-10%), with the exception of alopecia (10-45%), and was not considered to be treatment-related (Appendix Tables A0-4A and A0-4B).

The estimated dosage for the parental generation was 0.1 mg/kg/day for Group 2, 0.2 to 0.5 mg/kg/day for Group 3, and 0.9 to 1.8 mg/kg/day for Group 4 (Table 0-4). The estimated dosage levels for the F₁ offspring were 0.1 to 0.3 mg/kg/day for Group 2, 0.5 to 1.3 mg/kg/day for Group 3, and 1.9 to 3.4 mg/kg/day for Group 4 (Table 0-9).

The following changes were observed in the reproductive data for the F₀ cohabitation:

- # Live pups per litter were decreased by 43 and 30% for female and total pups, respectively, in the 0.0015% group (Table 0-5).
- # Average pup weights at PND 14 and 21 were decreased by 15-33% for males and 15-37% for females in the 0.0015% group (Table 0-11).
- # Sire weights at delivery for the 0.0015% males were decreased by 20% (Table 0-5).
- # Body weights for lactating 0.0015% dams were decreased by 14-18% on PND 1, 4, 7, and 14. Body weights for lactating 0.0004% dams were decreased by 7% on PND 1 (Table 0-5 and Table 0-6).
- # Feed consumption in lactating dams was decreased by 26% on PND 4-7 for 0.0001% dams and by 32-42% on PND 4-7 and 18-21 for 0.0015% dams and was increased by 25% on PND 11-14 for 0.0004% dams when examined as g/kg body weight/day (Table 0-7A).
- # Feed consumption in lactating dams was decreased by 32-36% on PND 1-4 and 4-7 for 0.0001% dams, by 16% on PND 4-7 for 0.0004% dams, and by 31-49% on PND 1-4, 4-7, 7-11, and 18-21 for 0.0015% dams and was increased by 22% on PND 11-14 for 0.0004% dams when examined as g/animal/day (Table 0-7B).
- # Water consumption in lactating dams was decreased by 28-40% on PND 4-7, 11-14, and 18-21 for 0.0015% dams when examined as g/kg body weight/day (Table 0-8A).
- # Water consumption in lactating dams was decreased by 11-16% on PND 14-18 and 18-21 for 0.0001% dams and by 25-39% throughout lactation for 0.0015% dams when examined as g/animal/day (Table 0-8B).
- # The day of eye opening, day of vaginal opening and day of preputial separation were all delayed in the 0.0004% group (1.9 days, 1.4 days and 2.0 days, respectively) (Tables 0-14 and 0-15).
- # There was no change in anogenital distance in the F₁ males or females.

Vaginal cytology was performed on F₀ dams for 14 days prior to cohabitation. The percentage of females with regular cycles was decreased from 95% in the controls to 70% in the 0.0015 % females, and these animals also differed from the controls in the relative amount of time spent in estrous stages. No changes were seen in cycle length, number of cycles, or number of cycling females across the dose groups as compared to the control females (Table 0-20).

Mortality was observed in one F₀ males and two F₀ females. One 0.0001 % male was sacrificed as moribund on SD 113 after observations of anorexia, thinness, rough haircoat, languid behavior, and hunched posture. Two 0.0015 % females were found dead on SD 98 and 110. These females appeared normal prior to death (Table 0-27).

F₁ PND 21 Necropsy

In the 0.0015% males, terminal body weights decreased by 23%. Absolute brain weights were decreased by 6%, and relative spleen and thymus weights were decreased by 15 and 41%, respectively. Relative brain weights were increased by 16% (Table 0-16). All absolute and relative organ weights in the 0.0001 and 0.0004% males were comparable to controls. Incidental gross findings at necropsy consisted of dilated kidneys, a discoloration on a mammary gland, and masses and a nodule on the thymus (Table 0-17).

In the 0.0015% females, terminal body weights decreased by 26%. Absolute brain weights were decreased by 7%, and relative spleen and thymus weights were decreased by 17 and 42%, respectively. Relative brain weights were increased by 17% (Table 0-18). All weights in the 0.0001 and 0.0004% females were comparable to controls. Incidental gross findings at necropsy consisted of dilated kidneys and a small spleen (Table 0-19).

Histopathologic examination was not performed on these tissues.

F₀ Thyroid Hormone Analysis

TSH levels in the 0.0004% and 0.0015% males increased by 126% and 510%, respectively. T4 levels decreased by 46% in the 0.0004% males and by 78% in the 0.0015% males. TSH levels in the 0.0004% and 0.0015 % females increased by 91% and 476%, respectively. T4 levels decreased

by 52% in the 0.0004% females and by 49% in the 0.0015% females (non-significant). T3 levels for all treated groups were comparable to controls (Table 0-21).

F₀ Adult Necropsy

In the F₀ 0.0004 and 0.0015% males, the absolute thyroid/parathyroid weight was increased by 67% and 301%, respectively. In the 0.0015% males, terminal body weight was decreased by 19%, while relative brain (15%), right testis (23%) and seminal vesicle (16%) weights were all increased. In the F₀ females, the absolute thyroid/parathyroid weight was increased in the 0.0015% group by 373%, while relative thyroid/parathyroid weights were increased in the 0.0001, 0.0004, and 0.0015% animals (18, 34, and 443% respectively). In the 0.0015% females, terminal body weights (13%) and absolute adrenal (22%), brain (6%), kidney (12%), liver (10%), and spleen (15%) weights were all decreased. All other weights (absolute and relative) for both males and females were comparable to the controls (Tables 0-23 and 0-26).

The percentage of abnormal sperm increased from 0.3% in the controls to 1.0% in the 0.0001% males. The sperm per mg of cauda, total sperm per cauda, spermatids per mg of testis, and total spermatids per testis for all treated groups were comparable to controls (Table 0-24A). Computer-assisted sperm motion analysis using Hamilton Thorne Integrated Visual Optic System revealed no changes in mean path velocity, progressive velocity, track speed, lateral amplitude, beat frequency, straightness, linearity, or motile percentage of the treated groups as compared to the controls (Table 0-24B).

Enlarged thyroids/parathyroids were seen in 0.0004% males (7 of 20), 0.0015% males (20 of 20), 0.0004% females (1 of 20), and 0.0015% females (18 of 20). Incidental gross findings in the males at necropsy included a discoloration on adipose tissue, a kidney cyst, and a discoloration on a mammary gland. Incidental findings in the females at necropsy included an enlarged kidney, a discoloration on a mammary gland, enlarged mammary glands, an enlarged pituitary, an enlarged urinary bladder, and distended, enlarged or fluid-filled vagina/cervix/uterus. All findings except the enlarged thyroids/parathyroids were spread throughout the dose groups and the incidence and severity were not related to dose (Tables 0-22 and 0-25).

Microscopic findings are described in the Pathology Report (Appendix 6). Thyroid follicular cell hyperplasia was seen in 0.0015% males (10 of 10) and females (10 of 10) and in 0.0004% males (7 of 10) and females (1 of 10). In addition, a follicular cell adenoma was seen in one of the 0.0015% females with follicular cell hyperplasia. Degeneration of the germinal epithelium of the testes was seen in 0.0015% males (2 of 10), 0.0004% males (3 of 10), and 0.0001% males (3 of 10). Multinucleated giant cells were only seen in 1 of the 0.0015% males exhibiting degeneration of the germinal epithelium.

F₁ Growing Phase

Increased mortality among the F₁ offspring was observed during the growing phase (i.e., after weaning): 23/29 males and 20/26 females in the 0.0015% group were either found dead or sacrificed as moribund.

By PND 25, the 0.0015% pups appeared to be smaller than pups in other groups, with domed heads and misshapen snouts. Six male and eight female pups had been found dead and 17 males and 12 females had been sacrificed as moribund (Table 1A-2). During necropsy of these unscheduled deaths, a delay in eruption of teeth was noted. The upper and lower jaws were evaluated microscopically. Changes were seen which seemed to represent damage (depletion and vacuolation) to the odontoblasts and ameloblasts, resulting in a delay in cellular maturation and subsequent tooth eruption. Tissues as examined, although not normal, seemed to be progressing toward normal, in spite of the delayed tooth eruption.

Because of the high level of mortality, the remainder of the 0.0015% pups were removed from the study and euthanized, and the growing phase and F₁ cohabitation were conducted with only three groups: control, 0.0001%, and 0.0004%.

There were no changes in male or female body weights during the F₁ growing phase (Table 1A-1).

Clinical signs noted during the growing phase included abrasions and alopecia. The incidence of these observations was low (0-17%), and was not considered to be treatment-related (Appendix Tables A1-2A and A1-2B). With the exception of the high-dose animals discussed above, there was no mortality seen during the growing phase (Table 1A-2).

F₁ Cohabitation

Mean body weights for all treated males and females were comparable to controls (Tables 1-1A and 1-1B).

All feed consumption values were comparable to controls when examined as g/kg body weight/day (Table 1-2A). When examined as g/animal/day, the feed consumption for 0.0001 % males decreased 8% during Week 4 (Table 1-2B). With the exception of lactating females, all water consumption values were comparable to controls when examined as either g/kg body weight/day or g/animal/day (Tables 1-3A or 1-3B).

Clinical signs noted during this portion of the study included abrasions, alopecia, swelling, a small stationary tissue mass, and an ulceration. The incidence of these observations was low to moderate (0-30%), and was not considered to be treatment-related (Appendix Tables A1-7A and A1-7B).

The following changes were observed in the reproductive data for the F₁ Cohabitation:

- # Male anogenital distance (AGD) in the 0.0001 and 0.0004% pups was decreased by 8% and 7% respectively. Male AGD/Pup Weight Ratio in the 0.0004% pups was decreased by 8% (Table 1-10).
- # Feed consumption in lactating dams was decreased by 11-12% on PND 4-11 for 0.0001% dams and by 9% on PND 18-21 for 0.0004% dams when examined as g/animal/day (Table 1-7B).
- # Water consumption in lactating dams was decreased by 10-16% on PND 1-14 for 0.0001% dams and by 12% on PND 4-7 for 0.0004% dams when examined as g/animal/day (Table 1-8B).

Vaginal cytology was performed on F₁ dams for 14 days after the last F₂ litters. No changes were revealed in the number of females with regular cycles, cycle length, number of cycles, or number of cycling females across the dose groups as compared to the control females. Treated females did not differ from the control females in the relative amount of time spent in estrous stages (Table 1-19).

Mortality was observed in one F₁ animal. One control male was found dead on SD 67 of the F₁ cohabitation. The animal appeared normal prior to death (Table 1-26).

F₂ PND 21 Necropsy

In the 0.0001% males, the absolute spleen weight was increased by 14%, and the relative spleen weight was increased by 12%. All weights in the 0.0004% males were comparable to controls (Table 1-16). The only gross finding at necropsy was an enlarged liver in one control male (Table 1-15).

In the 0.0004% females, absolute and relative thymus weights were decreased by 9%. All weights in the 0.0001% females were comparable to controls (Table 1-18). There were no gross findings at necropsy (Table 1-17).

Histopathologic examination was not performed on these tissues.

F₁ Thyroid Hormone Analysis

TSH levels in the 0.0004% males increased by 172%. T4 levels in the 0.0004% males decreased by 61%. TSH levels in the 0.0004% females increased by 144%. T4 levels in the 0.0004% females decreased by 59%. T3 levels for all treated groups were comparable to controls (Table 1-20).

F₁ Adult Necropsy

Absolute thyroid/parathyroid weights increased by 80% in the 0.0004% males. Relative thyroid/parathyroid weights in the 0.0001 and 0.0004% males increased by 26 and 84%, respectively. Absolute and relative thyroid/parathyroid weights increased by 72 and 74%, respectively, in the 0.0004% females. All other weights (absolute and relative) for both males and females were comparable to the controls (Tables 1-22 and 1-25).

Computer-assisted sperm motion analysis using Hamilton Thorne Integrated Visual Optic System revealed no changes in mean path velocity, progressive velocity, track speed, lateral amplitude, beat frequency, straightness, linearity, or motile percentage of the treated groups as compared to the control (Table 1-23B). The total sperm per cauda, sperm per mg of cauda, spermatids per mg of testis, total spermatids per testis, and percent of abnormal sperm were also all comparable to controls (Table 1-23A).

The only gross finding seen in a male at necropsy was a distended, red urinary bladder in one 0.0001% animal (Table 1-21). There were no gross findings in the females at necropsy (Table 1-24).

Microscopic findings are described in the Pathology Report (Appendix 6). Thyroid follicular cell hyperplasia was seen in one 0.0004 % male. Degeneration of the germinal epithelium of the testes was seen in one 0.0001 % male and one 0.0004 % male. There were no treatment-related microscopic changes in any of the F₁ females evaluated (Appendix 6).

DISCUSSION

In this two-generation study, propylthiouracil (PTU) was administered to male and female Sprague-Dawley rats in the drinking water at dose levels of 0.0001, 0.0004, or 0.0015 % (w/v). The corresponding dose levels were 0.1, 0.2 to 0.5, and 0.9 to 1.8 mg/kg/day, respectively. The F₀ generation was exposed to PTU only through their drinking water; the F₁ generation was exposed during gestation and lactation and subsequently through the drinking water; and the F₂ pups were exposed during gestation and lactation. Each generation was evaluated for general and reproductive/developmental toxicity endpoints.

In the F₀ males and females, body weights were decreased at 0.0015% but were comparable to controls at 0.0001 and 0.0004%. This decrease in body weight was previously observed by O'Connor *et al.* (1999) in males following 15 doses of PTU at 10 mg/kg. A similar effect was not seen in females following five doses. The body weight loss noted in the 0.0015% males and females was accompanied by decreased food and water consumption. Comparable decreases in body weight and food and water consumption were seen in lactating F₀ dams at 0.0015% and, to a much lesser extent, at 0.0001%. In general, less remarkable changes in body weight, food consumption, and water consumption were noted in the 0.0004% and, sporadically, the 0.0001% groups at some time points, indicating mild to moderate general toxicity at these doses.

Changes in estrous cyclicity were observed in the F₀ females at 0.0015% but not at the lower dose levels. These changes consisted of a decrease in the number of females with a regular cycle and changes in the amount of time spent in the various estrous stages. These changes did not affect female fertility, as no decrease in pregnancy index was noted among these animals.

Even though fertility was unaffected in the F₀ parents, there was a decrease in the number of female and total F₁ pups at 0.0015%. F₁ pup survival at this dose level was comparable through PND 21, although a marked decrease in male and female pup body weights was observed on PND 14 and became severe by PND 21. A high level of mortality was observed in these animals after weaning. By PND 25 the 0.0015% pups were noticeably smaller with domed heads and misshapen snouts. During the PND 21 necropsy a delay in eruption of the teeth was noted and the jaws, when evaluated microscopically, revealed depletion and vacuolization of the odontoblasts and ameloblasts, resulting in a delay in cellular maturation and subsequent tooth eruption. Tissues examined at PND

21, although not normal, seemed to be progressing toward normal, in spite of the delayed tooth eruption. Also noted at PND 21 were decreases in relative spleen, thymus, and brain weights and absolute brain weights in the 0.0015% males and females. These data reflect the overall diminished growth that was observed in these animals.

Due to the increased mortality observed among the F₁ juveniles at 0.0015%, surviving animals in this group were removed from study, and the remainder of the study was conducted using only 0, 0.0001, and 0.0004% dose levels.

In the F₁ offspring there were delays in eye opening (0.0015 %), preputial separation (0.0004 %), and vaginal opening (0.0004 %). No animals could be evaluated at 0.0015% for preputial separation and vaginal opening due to mortality. Preputial separation was delayed 2.0 days from 42.7 days in controls to 44.7 days at 0.0004%, and vaginal opening was delayed 1.4 days from 32.4 days in controls to 33.8 days at 0.0004 %. Marty *et al.* (2001) confirmed a delay in preputial separation in males dosed at 240 mg/kg/day from 44.4 days in controls to greater than 50 days in treated animals. Marty *et al.* (1999) confirmed that vaginal opening was delayed in females dosed at 240/mg/kg/day to 34.2 days from 32.3 or 33.5 days in controls. Wilen *et al.* (1981) also noted a delay in vaginal opening at 0.1% in immature rats fed PTU from weaning through the day of vaginal opening.

Delayed vaginal opening indicates delayed pubertal onset in juvenile female rats (Marty, 1999) and may occur with compounds that cause endocrine disruption or other reproductive toxicity. However, decreases in body weight may also delay vaginal opening, thereby making it more difficult to attribute delayed vaginal opening to reproductive/developmental toxicity or endocrine disruption (Marty, 1999). For compounds that have not been as thoroughly evaluated as PTU, for example, it is important to evaluate the significance of delayed vaginal opening in context with other observations of toxicity.

The observed delays in preputial separation and vaginal opening did not affect the subsequent fertility of the F₁ animals; no decreases in pregnancy index were noted among these animals during the F₁ Cohabitation. In contrast to the F₀ Cohabitation, no decreases in number of pups or pup weights were observed in the F₁ Cohabitation. However, decreased anogenital distance was noted in the 0.0004% and 0.0001% F₂ male pups, which was not observed in the F₁ male pups.

The AGD/pup weight ratio was also decreased in the 0.0004% pups. These observations are a definitive indicator of male reproductive toxicity at 0.0004% and possibly 0.0001%. Moreover, the decrease in AGD that was only seen in the F₂ pups supports a two-generation study design.

As expected, TSH levels were increased and T4 levels were decreased in the PTU-treated F₀ and F₁ animals. The TSH and T4 levels were correlated with the thyroid weights, discussed below, by being most affected at 0.0015%, less in the females, and more in the F₁ males and females. T3 levels were not affected by PTU administration. Analysis of T3 levels may be useful when evaluating other compounds, although it was not essential here.

In the PTU-treated F₀ and F₁ animals there were dose-related increases in absolute and relative thyroid weights. At 0.0015% the thyroid weight increases were 3 to 4 times the control weight, while at 0.0004% the increases were much less remarkable. The observed changes in thyroid weight were less severe in females than males and more severe in both sexes in the F₁ than the F₀ adults. At 0.0001% there was an increase in relative thyroid weight in the F₁ males and F₀ females. O'Connor *et al.* (1999) concluded that relative thyroid weight was a more sensitive indicator of thyroid toxicity than absolute thyroid weight: Relative thyroid weight removes the potentially confounding effect of body weight and allows detection of a true compound effect, particularly when decreases in body weight are observed. Also of interest is the observation of increased relative pituitary weights in the 0.0015% and 0.0004% F₀ males. The significance of this particular observation is unclear at this time, but may reflect increased demand on the pituitary for TRH.

At the F₀ gross necropsy, enlarged thyroids/parathyroids were seen in most 0.0015% animals and in 7/20 and 1/20 0.0004% males and females, respectively. No changes were noted in the F₁ gross necropsy findings.

Histopathology findings in the thyroid and testes were generally consistent between the F₀ and F₁ animals. Thyroid follicular cell hyperplasia was seen in all 0.0015% F₀ animals and in 7/10 0.0004% F₀ males, but in no F₁ females. In the 0.0004% animals, only 1/10 F₁ males was noted with thyroid follicular cell hyperplasia. The paucity of microscopic findings in the F₁ animals was surprising in light of the changes seen in thyroid weight and thyroid hormones and the microscopic findings in the F₀ animals. The pathology, thyroid hormone, and thyroid weight findings are in

general agreement with those of O'Connor *et al.* (1999), Marty *et al.* (1999), and Marty *et al.* (2001). O'Connor *et al.* (1999) observed these effects in females at 10 mg/kg/day when treated for five days and in males at 0.25 mg/kg/day when treated for 15 days. Marty *et al.* (1999) and Marty *et al.* (2001) treated females for 20 days at 240 mg/kg/day and males for 30 days at 240 mg/kg/day and observed similar findings. However, the belief that histopathology of the thyroid is the most sensitive indicator of thyroid toxicity (O'Connor, 1999) is not supported by the data presented here.

Degeneration of the germinal epithelium of the testes was also noted in all available treated male groups in the F₀ and F₁ generations. Degeneration was noted in 2/10, 3/10, and 3/10 F₀ males at 0.0015, 0.0004 and 0.0001%, respectively. In the F₁ males, degeneration was noted in 1/10 males at 0.0004 and 0.0001%.

Stoker *et al.* (2000) concluded that the induction of hypothyroidism by administration of PTU in polychlorinated biphenyls from birth to weaning increased testis size, number of Sertoli cells and Leydig cells, and daily sperm production in the adult. Other reports cited by Stoker *et al.* (2000) show that the critical period for this effect is the first two weeks after birth with no effects observed if PTU treatment was started during late lactation. Whether these results are attributable to the PTU or the PCBs is unknown. Regardless, the results in our study are in general agreement with these authors: There was no difference noted in the sperm parameters in either the F₀ males of the F₁ males.

Based on the findings of this two generation study with one litter per generation, propylthiouracil is considered to be a reproductive/developmental toxicant in females at dose levels greater than or equal to 0.0004% based on decreased total pups per litter, delayed vaginal opening, delayed eye opening, and changes in estrous cyclicity. PTU is considered a male reproductive toxicant at dose levels greater than or equal to 0.0001% based on delayed eye opening, delayed preputial separation, degeneration of the germinal epithelium of the testes, and decreased anogenital distance. Propylthiouracil is also considered a general toxicant at 0.0015 and 0.0004% based upon decreased body weight and food consumption (mostly at 0.0015%) and increased thyroid weights, changes in thyroid hormone levels, and/or thyroid follicular cell hyperplasia at 0.0004 and 0.0015%.

REFERENCES

- Armitage, P. (1971). *Statistical Methods in Medical Research*. John Wiley & Sons, New York.
- Conover, W.J. (1971). *Practical Nonparametric Statistics*. John Wiley & Sons, New York.
- Dunn, O.J. (1964). Multiple Comparisons Using Rank Sums. *Technometrics* 6 241-252.
- Dunnett, W. (1955). A Multiple Comparison Procedure for Comparing Several Treatments with a Control. *JASA*, 50:1096-1211.
- Health Effects Test Guidelines, OPPTS 870.3800, Reproduction and Fertility Effects, EPA 712-C-98-208.
- Jonckheere, A.R. (1954). A distribution-free K-sample Test Against Ordered Alternatives. *Biometrika*, 41:133-145.
- Kruskal, W.H. and Wallis, W.A. (1952). Use of Ranks in one-criterion Variance Analysis. *JASA*, 47:583-621.
- Marty, M.S., Crissman J.W. and Carney, E.W. (2001). Evaluation of the Male Pubertal Assay's Ability to Detect Thyroid Inhibitors and Dopaminergic Agents. *Toxicol Sci*, 60(1):63-76.
- Marty, M.S., Crissman J.W. and Carney, E.W. (1999). Evaluation of the EDSTAC female pubertal assay in CD rats using 17 β -estradiol, steroid biosynthesis inhibitors, and a thyroid inhibitor. *Toxicol Sci*, 52(2):269-77.
- Morrison, D.F. (1976). *Multivariate Statistical Methods*. McGraw Hill, New York.
- Neter, J. and Wasserman, W. (1974). *Applied Linear Statistical Models*. Richard Irwin, Inc., Homewood, IL.
- O'Connor, J.C., Frame, S.R., Davis, L.G. and Cook, J.C. (1999). Detection of thyroid toxicants in a tier I screening battery and alterations in thyroid endpoints over 28 days of exposure. *Toxicol Sci*, 51(1):54-70.
- Shirley, E. (1977). A non-parametric equivalent of William's test for contrasting increasing dose levels of a treatment. *Biometrics*, 33:386-389.
- Stoker, T.E, Parks, L.G., Gray, L.E. and Cooper, R.L. (2000). Endocrine-Disrupting Chemicals: Prepubertal Exposures and Effects on Sexual Maturation and Thyroid Function in the Male Rat. A Focus on the EDSTAC Recommendations. *Critical Reviews in Toxicology*, 30(2):197-252.

Wilens, R., Bastomsky, C.H. and Naftolin, F. (1981). Control of Puberty in Female Rats: the Effects of PTU-Induced Hypothyroidism and Systemic Undernutrition. *Pediatr Res*, 15:169-171.

Williams, D.A. (1986). A Note on Shirley's Nonparametric Test for Comparing Several Dose Levels with a Zero-Dose Control. *Biometrics*, 42:183-186.

TABLE 0-1A
 F₀: SUMMARY OF MALE BODY WEIGHTS (GRAMS)
 TWO GENERATION REPRODUCTION TOXICITY STUDY OF PROPYLTHIOURACIL WHEN ADMINISTERED TO SPRAGUE-DAWLEY RATS IN THE DRINKING WATER

WEEK OF STUDY	DOSE GROUPS ^a				TREND (b)
	0.0000 %	0.0001 %	0.0004 %	0.0015 %	
1	260.5 ± 2.55 (20)	261.3 ± 2.70 (20)	262.3 ± 3.21 (20)	259.1 ± 3.45 (20)	P=0.833
2	312.7 ± 3.57 (20)	315.1 ± 3.87 (20)	314.8 ± 4.44 (20)	312.2 ± 4.47 (20)	P=0.767
3	355.5 ± 5.02 (20)	357.9 ± 5.34 (20)	356.0 ± 5.71 (20)	*335.1 ± 5.17 (20)	P=0.003
4	391.9 ± 6.23 (20)	391.5 ± 6.01 (20)	388.3 ± 6.57 (20)	*342.2 ± 5.93 (20)	P<0.001
5	419.9 ± 7.11 (20)	424.2 ± 6.90 (20)	415.8 ± 7.58 (20)	*351.2 ± 6.25 (20)	P<0.001
6	439.8 ± 8.03 (20)	448.1 ± 7.90 (20)	437.0 ± 8.74 (20)	*352.7 ± 6.76 (20)	P<0.001
7	469.2 ± 9.23 (20)	470.8 ± 8.88 (20)	459.1 ± 8.81 (20)	*368.9 ± 7.43 (20)	P<0.001
8	489.2 ± 10.00 (20)	488.9 ± 9.79 (20)	478.8 ± 9.86 (20)	*381.2 ± 8.00 (20)	P<0.001
9	504.3 ± 10.59 (20)	506.1 ± 10.05 (20)	491.2 ± 10.12 (20)	*390.3 ± 8.71 (20)	P<0.001
10	518.6 ± 11.14 (20)	522.0 ± 10.18 (20)	508.0 ± 10.80 (20)	*405.5 ± 9.44 (20)	P<0.001
11	532.7 ± 11.62 (20)	532.1 ± 13.26 (20)	523.4 ± 11.54 (20)	*414.3 ± 10.00 (20)	P<0.001
12	539.2 ± 11.77 (20)	538.4 ± 11.59 (20)	528.3 ± 11.82 (20)	*417.2 ± 9.84 (20)	P<0.001
13	548.0 ± 12.58 (20)	552.5 ± 10.86 (20)	541.2 ± 12.18 (20)	*429.0 ± 10.32 (20)	P<0.001
14	559.6 ± 13.20 (20)	559.2 ± 15.32 (20)	553.3 ± 12.33 (20)	*439.8 ± 10.40 (20)	P<0.001
15	574.1 ± 13.85 (20)	569.6 ± 18.11 (20)	566.3 ± 13.32 (20)	*450.4 ± 10.78 (20)	P<0.001
16	584.5 ± 13.82 (20)	580.0 ± 18.27 (20)	573.3 ± 13.88 (20)	*460.8 ± 11.27 (20)	P<0.001
17	598.6 ± 14.32 (20)	594.5 ± 16.74 (20)	578.3 ± 14.53 (20)	*472.4 ± 11.93 (20)	P<0.001
18	611.0 ± 15.02 (20)	615.5 ± 11.98 (19)	588.9 ± 13.87 (20)	*483.2 ± 12.34 (20)	P<0.001
19	619.1 ± 15.50 (20)	626.3 ± 12.64 (19)	596.4 ± 13.96 (20)	*494.3 ± 12.80 (20)	P<0.001
20	623.8 ± 17.62 (20)	637.6 ± 13.39 (19)	603.2 ± 14.42 (20)	*506.8 ± 12.92 (20)	P<0.001

a Mean ± standard error (number of animals).

b Each dose group is compared to the control with Shirley's test when a trend is present (P<0.01 from Jonckheere's trend test), otherwise Dunn's test is applied (*= P<0.05).

TABLE 0-1B
 F₀: SUMMARY OF FEMALE BODY WEIGHTS (GRAMS)
 TWO GENERATION REPRODUCTION TOXICITY STUDY OF PROPYLTHIOURACIL WHEN ADMINISTERED TO SPRAGUE-DAWLEY RATS IN THE DRINKING WATER

WEEK OF STUDY	DOSE GROUPS ^a				TREND (b)
	0.0000 %	0.0001 %	0.0004 %	0.0015 %	
1	192.3 ± 3.42 (20)	188.9 ± 2.91 (20)	191.8 ± 3.28 (20)	190.3 ± 2.78 (20)	P=0.693
2	215.8 ± 4.10 (20)	213.1 ± 3.63 (20)	212.9 ± 3.88 (20)	212.4 ± 3.73 (20)	P=0.556
3	232.5 ± 4.72 (20)	231.6 ± 4.14 (20)	230.4 ± 4.38 (20)	227.0 ± 3.91 (20)	P=0.422
4	248.1 ± 5.37 (20)	246.3 ± 4.53 (20)	243.4 ± 4.96 (20)	234.5 ± 4.13 (20)	P=0.033
5	261.5 ± 6.15 (20)	258.6 ± 4.83 (20)	256.4 ± 5.24 (20)	244.2 ± 4.74 (20)	P=0.036
6	269.9 ± 5.97 (20)	275.8 ± 5.34 (20)	256.0 ± 5.66 (20)	*244.1 ± 4.58 (20)	P<0.001
7	279.4 ± 6.73 (20)	279.1 ± 5.25 (20)	275.3 ± 5.62 (20)	*255.8 ± 5.01 (20)	P=0.006
8	283.0 ± 7.55 (20)	288.2 ± 5.78 (20)	284.5 ± 6.61 (20)	261.5 ± 5.26 (20)	P=0.017
9	295.3 ± 7.50 (20)	293.3 ± 5.62 (20)	288.1 ± 6.50 (20)	*266.3 ± 5.26 (20)	P=0.002
10	300.6 ± 8.02 (20)	298.3 ± 5.65 (20)	293.6 ± 6.40 (20)	*273.2 ± 5.37 (20)	P=0.006
11	304.7 ± 8.68 (20)	302.7 ± 5.90 (20)	296.8 ± 6.38 (20)	*275.0 ± 5.71 (20)	P=0.004
12	324.7 ± 8.11 (20)	319.1 ± 5.81 (20)	318.3 ± 6.72 (20)	*288.1 ± 6.26 (20)	P<0.001
13	348.4 ± 7.99 (20)	340.1 ± 6.58 (20)	335.1 ± 9.35 (20)	*296.9 ± 7.73 (20)	P<0.001
14	398.1 ± 9.86 (20)	386.4 ± 9.59 (20)	396.5 ± 11.10 (20)	*336.5 ± 7.30 (20)	P<0.001
15	350.2 ± 8.19 (20)	333.3 ± 7.11 (20)	333.1 ± 7.06 (20)	*290.4 ± 5.94 (19)	P<0.001
16	367.2 ± 8.48 (20)	*346.7 ± 7.82 (20)	353.4 ± 7.08 (20)	*306.4 ± 4.94 (19)	P<0.001
17	341.1 ± 8.35 (20)	340.7 ± 7.78 (20)	347.4 ± 5.54 (20)	*314.6 ± 5.22 (18)	P=0.043
18	326.4 ± 8.21 (20)	316.8 ± 6.30 (20)	313.0 ± 5.31 (20)	*292.4 ± 6.21 (18)	P=0.002
19	335.2 ± 9.12 (20)	325.8 ± 5.80 (20)	321.8 ± 6.35 (20)	*293.6 ± 6.36 (18)	P<0.001
20	346.0 ± 9.83 (20)	333.4 ± 6.14 (20)	329.6 ± 7.20 (20)	*300.9 ± 7.42 (18)	P<0.001

a Mean ± standard error (number of animals).

b Each dose group is compared to the control with Shirley's test when a trend is present (P<0.01 from Jonckheere's trend test), otherwise Dunn's test is applied (*= P<0.05).

TABLE 0-2A
 F₀: SUMMARY OF DAILY FEED CONSUMPTION (G/KG BODY WEIGHT/DAY)
 TWO GENERATION REPRODUCTION TOXICITY STUDY OF PROPYLTHIOURACIL WHEN ADMINISTERED TO SPRAGUE-DAWLEY RATS IN THE DRINKING WATER

WEEK OF STUDY	DOSE GROUPS ^a				TREND (b)
	0.0000 %	0.0001 %	0.0004 %	0.0015 %	
<u>MALE</u>					
1	101.0 ± 1.57 (20)	99.1 ± 1.21 (20)	100.3 ± 1.25 (20)	98.6 ± 1.11 (20)	P=0.336
2	83.9 ± 1.09 (20)	84.2 ± 0.99 (20)	84.6 ± 0.92 (20)	*74.3 ± 1.14 (20)	P<0.001
3	77.0 ± 1.09 (20)	76.3 ± 0.95 (20)	75.7 ± 0.89 (20)	*63.8 ± 0.92 (20)	P<0.001
4	75.5 ± 0.77 (20)	75.8 ± 1.00 (20)	75.3 ± 0.91 (20)	*64.4 ± 1.02 (20)	P<0.001
5	74.2 ± 0.99 (20)	72.5 ± 0.75 (20)	74.0 ± 1.14 (20)	*64.2 ± 1.21 (20)	P<0.001
6	72.0 ± 0.76 (20)	*69.2 ± 0.92 (20)	71.0 ± 1.02 (20)	*58.8 ± 1.02 (20)	P<0.001
7	66.2 ± 0.73 (20)	*63.2 ± 0.95 (20)	66.4 ± 0.88 (20)	*57.8 ± 0.95 (20)	P<0.001
8	60.6 ± 0.67 (20)	61.6 ± 0.92 (20)	62.0 ± 0.73 (20)	*56.7 ± 1.08 (20)	P=0.045
9	56.2 ± 0.52 (20)	56.0 ± 0.53 (20)	57.9 ± 0.69 (20)	53.3 ± 0.88 (20)	P=0.239
10	55.3 ± 0.50 (20)	53.3 ± 1.08 (20)	53.6 ± 0.63 (20)	*50.9 ± 0.84 (20)	P<0.001
13	57.0 ± 1.01 (20)	*50.5 ± 2.43 (20)	*50.9 ± 2.00 (20)	*52.2 ± 1.51 (20)	P=0.017
14	54.3 ± 0.93 (20)	52.1 ± 1.04 (20)	52.6 ± 0.97 (19)	53.7 ± 1.07 (20)	P=0.827
15	51.5 ± 0.70 (20)	52.9 ± 2.69 (19)	50.4 ± 0.70 (20)	53.0 ± 0.80 (20)	P=0.372
16	53.6 ± 0.99 (20)	53.9 ± 3.45 (19)	48.3 ± 3.24 (20)	52.7 ± 0.76 (20)	P=0.753
17	50.6 ± 1.60 (20)	*47.0 ± 1.39 (19)	48.6 ± 1.82 (20)	50.7 ± 1.56 (20)	P=0.478
18	50.5 ± 0.89 (20)	49.8 ± 1.20 (19)	50.2 ± 0.88 (20)	54.0 ± 1.20 (20)	P=0.034
19	47.2 ± 2.02 (20)	45.6 ± 1.85 (19)	47.7 ± 0.61 (20)	50.6 ± 0.87 (20)	P=0.062
<u>FEMALE</u>					
1	96.7 ± 1.59 (20)	98.7 ± 1.93 (20)	95.9 ± 1.78 (20)	93.0 ± 2.39 (20)	P=0.279
2	87.4 ± 2.94 (20)	91.2 ± 1.31 (20)	89.9 ± 1.51 (20)	*84.4 ± 1.95 (20)	P=0.006
3	87.7 ± 1.99 (19)	86.8 ± 1.18 (20)	85.2 ± 1.25 (20)	*75.9 ± 1.76 (20)	P<0.001
4	85.3 ± 2.68 (20)	85.6 ± 1.61 (20)	85.7 ± 1.39 (20)	*73.7 ± 1.15 (20)	P<0.001
5	86.4 ± 2.36 (20)	83.1 ± 1.32 (20)	90.6 ± 1.24 (20)	*73.8 ± 1.62 (20)	P=0.012
6	85.0 ± 2.26 (20)	85.3 ± 1.40 (20)	81.2 ± 1.32 (20)	*78.1 ± 6.00 (20)	P<0.001
7	76.0 ± 3.73 (19)	77.8 ± 0.87 (20)	75.2 ± 1.01 (20)	*67.0 ± 2.27 (20)	P<0.001
8	77.4 ± 4.17 (20)	72.1 ± 1.36 (20)	69.5 ± 1.31 (20)	*64.0 ± 1.42 (19)	P<0.001
9	64.0 ± 1.70 (19)	62.6 ± 1.23 (20)	63.8 ± 1.23 (20)	*58.4 ± 1.47 (19)	P=0.037
10	63.7 ± 2.90 (20)	62.7 ± 1.36 (20)	61.0 ± 1.62 (20)	*56.7 ± 2.46 (20)	P<0.001
13	65.1 ± 3.96 (20)	64.9 ± 3.17 (19)	71.4 ± 2.38 (20)	59.5 ± 2.97 (20)	P=0.130
18	71.6 ± 4.26 (20)	81.3 ± 4.59 (18)	82.3 ± 4.13 (20)	60.3 ± 6.03 (18)	P=0.254
19	72.5 ± 2.38 (20)	73.1 ± 2.66 (20)	70.7 ± 2.02 (20)	72.8 ± 6.57 (18)	P=0.035

a Mean ± standard error (number of animals).

b Each dose group is compared to the control with Shirley's test when a trend is present (P<0.01 from Jonckheere's trend test), otherwise Dunn's test is applied (*= P<0.05).

TABLE 0-2B
 F₀: SUMMARY OF DAILY FEED CONSUMPTION (G/ANIMAL/DAY)
 TWO GENERATION REPRODUCTION TOXICITY STUDY OF PROPYLTHIOURACIL WHEN ADMINISTERED TO SPRAGUE-DAWLEY RATS IN THE DRINKING WATER

WEEK OF STUDY	DOSE GROUPS ^a				TREND (b)
	0.0000 %	0.0001 %	0.0004 %	0.0015 %	
<u>MALE</u>					
1	26.3 ± 0.43 (20)	25.9 ± 0.43 (20)	26.3 ± 0.50 (20)	25.5 ± 0.45 (20)	P=0.219
2	26.3 ± 0.54 (20)	26.5 ± 0.51 (20)	26.7 ± 0.56 (20)	*23.2 ± 0.51 (20)	P<0.001
3	27.4 ± 0.63 (20)	27.3 ± 0.59 (20)	27.0 ± 0.61 (20)	*21.4 ± 0.53 (20)	P<0.001
4	29.6 ± 0.62 (20)	29.7 ± 0.64 (20)	29.3 ± 0.67 (20)	*22.0 ± 0.53 (20)	P<0.001
5	31.2 ± 0.69 (20)	30.8 ± 0.68 (20)	30.7 ± 0.68 (20)	*22.6 ± 0.64 (20)	P<0.001
6	31.7 ± 0.73 (20)	31.0 ± 0.69 (20)	31.0 ± 0.69 (20)	*20.7 ± 0.52 (20)	P<0.001
7	31.1 ± 0.76 (20)	29.8 ± 0.77 (20)	30.5 ± 0.73 (20)	*21.3 ± 0.54 (20)	P<0.001
8	29.7 ± 0.73 (20)	30.0 ± 0.59 (20)	29.7 ± 0.65 (20)	*21.6 ± 0.58 (20)	P<0.001
9	28.3 ± 0.66 (20)	28.3 ± 0.54 (20)	28.4 ± 0.67 (20)	*20.8 ± 0.64 (20)	P<0.001
10	28.7 ± 0.67 (20)	27.8 ± 0.80 (20)	27.2 ± 0.66 (20)	*20.7 ± 0.64 (20)	P<0.001
13	31.2 ± 0.80 (20)	28.1 ± 1.48 (20)	*27.4 ± 1.18 (20)	*22.3 ± 0.78 (20)	P<0.001
14	30.3 ± 0.79 (20)	29.1 ± 0.95 (20)	29.2 ± 0.94 (19)	*23.6 ± 0.75 (20)	P<0.001
15	29.5 ± 0.65 (20)	29.4 ± 0.58 (19)	28.6 ± 0.83 (20)	*23.9 ± 0.67 (20)	P<0.001
16	31.3 ± 0.83 (20)	30.3 ± 1.03 (19)	27.8 ± 2.06 (20)	*24.3 ± 0.68 (20)	P<0.001
17	30.2 ± 1.15 (20)	28.5 ± 0.92 (19)	27.8 ± 0.88 (20)	*24.0 ± 1.03 (20)	P<0.001
18	30.8 ± 0.77 (20)	30.6 ± 0.82 (19)	29.5 ± 0.68 (20)	*26.1 ± 0.83 (20)	P<0.001
19	29.3 ± 1.38 (20)	28.6 ± 1.32 (19)	28.4 ± 0.70 (20)	*25.0 ± 0.74 (20)	P<0.001
<u>FEMALE</u>					
1	18.6 ± 0.47 (20)	18.6 ± 0.39 (20)	18.4 ± 0.48 (20)	17.7 ± 0.54 (20)	P=0.310
2	18.8 ± 0.69 (20)	19.4 ± 0.46 (20)	19.1 ± 0.46 (20)	17.9 ± 0.41 (20)	P=0.038
3	20.4 ± 0.67 (19)	20.1 ± 0.45 (20)	19.6 ± 0.50 (20)	*17.2 ± 0.44 (20)	P<0.001
4	21.1 ± 0.80 (20)	21.1 ± 0.57 (20)	20.9 ± 0.53 (20)	*17.3 ± 0.36 (20)	P<0.001
5	22.5 ± 0.68 (20)	21.5 ± 0.56 (20)	23.2 ± 0.47 (20)	*18.0 ± 0.38 (20)	P<0.001
6	22.9 ± 0.70 (20)	23.5 ± 0.51 (20)	*20.8 ± 0.51 (20)	*19.0 ± 1.42 (20)	P<0.001
7	21.1 ± 1.08 (19)	21.7 ± 0.45 (20)	20.7 ± 0.52 (20)	*17.1 ± 0.55 (20)	P<0.001
8	21.8 ± 1.23 (20)	20.7 ± 0.45 (20)	19.7 ± 0.51 (20)	*16.7 ± 0.51 (19)	P<0.001
9	19.0 ± 0.69 (19)	18.3 ± 0.45 (20)	18.3 ± 0.36 (20)	*15.5 ± 0.44 (19)	P<0.001
10	19.2 ± 1.01 (20)	18.7 ± 0.46 (20)	17.8 ± 0.44 (20)	*15.5 ± 0.74 (20)	P<0.001
13	22.7 ± 1.44 (20)	22.2 ± 1.26 (19)	23.7 ± 0.73 (20)	*17.6 ± 0.90 (20)	P=0.003
18	23.4 ± 1.49 (20)	25.7 ± 1.41 (18)	25.7 ± 1.34 (20)	*17.6 ± 1.76 (18)	P=0.022
19	24.3 ± 0.95 (20)	23.8 ± 0.90 (20)	22.8 ± 0.86 (20)	*21.3 ± 1.85 (18)	P<0.001

a Mean ± standard error (number of animals).

b Each dose group is compared to the control with Shirley's test when a trend is present (P<0.01 from Jonckheere's trend test), otherwise Dunn's test is applied (*= P<0.05).

TABLE 0-3A
 F₀: SUMMARY OF DAILY WATER CONSUMPTION (G/KG BODY WEIGHT/DAY)
 TWO GENERATION REPRODUCTION TOXICITY STUDY OF PROPYLTHIOURACIL WHEN ADMINISTERED TO SPRAGUE-DAWLEY RATS IN THE DRINKING WATER

WEEK OF STUDY	DOSE GROUPS ^a				TREND (b)
	0.0000 %	0.0001 %	0.0004 %	0.0015 %	
<u>MALE</u>					
1	121.9 ± 3.03 (20)	125.6 ± 4.36 (20)	127.8 ± 3.57 (20)	120.1 ± 2.67 (20)	P=0.830
2	106.2 ± 3.17 (20)	109.4 ± 6.00 (20)	109.0 ± 4.47 (20)	96.7 ± 2.89 (20)	P=0.200
3	101.0 ± 4.20 (20)	98.9 ± 5.01 (20)	99.7 ± 4.96 (20)	90.6 ± 2.94 (19)	P=0.080
4	87.5 ± 3.38 (20)	82.5 ± 2.42 (20)	89.1 ± 3.70 (20)	84.8 ± 3.15 (20)	P=0.830
5	86.8 ± 3.49 (20)	79.0 ± 2.78 (20)	83.7 ± 2.41 (20)	81.3 ± 3.47 (20)	P=0.643
6	76.9 ± 3.54 (19)	71.0 ± 2.53 (20)	80.3 ± 3.34 (20)	79.8 ± 3.65 (20)	P=0.611
7	75.3 ± 3.20 (20)	68.1 ± 2.21 (19)	76.8 ± 3.05 (20)	75.8 ± 3.44 (20)	P=0.420
8	73.2 ± 3.09 (20)	74.7 ± 7.27 (20)	73.9 ± 3.76 (20)	74.7 ± 4.06 (20)	P=0.803
9	70.5 ± 3.60 (20)	67.0 ± 3.68 (20)	71.9 ± 3.51 (20)	73.1 ± 3.75 (20)	P=0.476
10	65.9 ± 2.75 (20)	63.8 ± 4.01 (20)	64.4 ± 3.29 (20)	66.7 ± 2.87 (20)	P=0.757
13	64.9 ± 3.05 (19)	63.6 ± 8.30 (20)	62.7 ± 4.05 (19)	64.5 ± 3.52 (20)	P=0.911
14	60.6 ± 3.29 (20)	60.0 ± 5.49 (20)	59.6 ± 2.97 (20)	63.7 ± 4.03 (20)	P=0.486
15	58.0 ± 3.43 (20)	59.1 ± 7.89 (20)	54.7 ± 2.32 (20)	56.8 ± 2.75 (20)	P=0.790
16	58.9 ± 2.76 (20)	52.4 ± 2.33 (19)	57.2 ± 3.23 (19)	66.1 ± 4.16 (18)	P=0.244
17	64.4 ± 4.24 (20)	52.7 ± 2.78 (19)	55.5 ± 3.48 (20)	63.8 ± 3.00 (20)	P=0.473
18	58.1 ± 3.23 (20)	51.4 ± 2.42 (19)	56.6 ± 2.96 (20)	59.2 ± 2.89 (20)	P=0.376
19	53.7 ± 3.76 (20)	51.0 ± 2.40 (19)	56.1 ± 2.57 (20)	58.8 ± 2.62 (20)	P=0.115
<u>FEMALE</u>					
1	122.7 ± 4.55 (20)	129.4 ± 4.45 (20)	118.7 ± 2.31 (20)	113.6 ± 5.32 (20)	P=0.072
2	107.0 ± 5.13 (20)	113.0 ± 3.10 (20)	109.3 ± 2.59 (20)	105.5 ± 4.54 (20)	P=0.371
3	108.8 ± 4.75 (20)	108.2 ± 4.01 (20)	103.1 ± 2.84 (20)	97.8 ± 4.89 (20)	P=0.033
4	100.1 ± 3.87 (20)	100.4 ± 3.64 (20)	99.4 ± 2.53 (20)	92.0 ± 3.82 (20)	P=0.108
5	93.8 ± 3.23 (20)	102.0 ± 4.16 (20)	104.5 ± 2.75 (20)	93.0 ± 4.40 (20)	P=0.973
6	94.1 ± 3.05 (20)	99.0 ± 3.23 (20)	98.2 ± 3.04 (20)	90.5 ± 5.21 (20)	P=0.331
7	86.7 ± 5.50 (20)	97.0 ± 3.77 (20)	88.9 ± 2.48 (20)	95.8 ± 10.04 (19)	P=0.436
8	94.2 ± 4.69 (20)	95.7 ± 5.45 (20)	83.5 ± 2.48 (20)	85.4 ± 4.25 (20)	P=0.084
9	84.4 ± 3.87 (20)	85.1 ± 2.97 (20)	81.1 ± 2.41 (20)	81.2 ± 3.89 (20)	P=0.315
10	80.8 ± 3.08 (20)	82.4 ± 3.37 (20)	73.8 ± 2.48 (20)	74.0 ± 3.87 (20)	P=0.028
13	114.9 ± 5.95 (18)	122.7 ± 4.97 (20)	117.5 ± 7.94 (20)	110.0 ± 8.81 (20)	P=0.280
18	112.9 ± 7.55 (20)	101.3 ± 4.98 (20)	*95.5 ± 3.27 (20)	*77.8 ± 5.31 (18)	P<0.001
19	100.6 ± 6.62 (20)	95.0 ± 4.72 (19)	88.0 ± 2.77 (20)	87.9 ± 6.54 (18)	P=0.040

a Mean ± standard error (number of animals).

b Each dose group is compared to the control with Shirley's test when a trend is present (P<0.01 from Jonckheere's trend test), otherwise Dunn's test is applied (*= P<0.05).

TABLE 0-3B
 F₀: SUMMARY OF DAILY WATER CONSUMPTION (G/ANIMAL/DAY)
 TWO GENERATION REPRODUCTION TOXICITY STUDY OF PROPYLTHIOURACIL WHEN ADMINISTERED TO SPRAGUE-DAWLEY RATS IN THE DRINKING WATER

WEEK OF STUDY	DOSE GROUPS ^a				TREND (b)
	0.0000 %	0.0001 %	0.0004 %	0.0015 %	
<u>MALE</u>					
1	31.7 ± 0.86 (20)	32.8 ± 1.12 (20)	33.5 ± 1.08 (20)	31.1 ± 0.71 (20)	P=0.810
2	33.2 ± 1.10 (20)	34.3 ± 1.59 (20)	34.3 ± 1.54 (20)	30.1 ± 0.84 (20)	P=0.088
3	35.9 ± 1.55 (20)	35.2 ± 1.51 (20)	35.4 ± 1.82 (20)	*30.2 ± 0.90 (19)	P=0.006
4	34.2 ± 1.31 (20)	32.3 ± 1.07 (20)	34.5 ± 1.48 (20)	*28.8 ± 0.87 (20)	P=0.004
5	36.3 ± 1.42 (20)	33.4 ± 1.07 (20)	34.6 ± 0.89 (20)	*28.3 ± 0.96 (20)	P<0.001
6	33.7 ± 1.47 (19)	31.7 ± 1.15 (20)	34.8 ± 1.22 (20)	*27.9 ± 1.03 (20)	P=0.006
7	35.2 ± 1.44 (20)	32.2 ± 1.03 (19)	35.2 ± 1.48 (20)	*27.7 ± 1.00 (20)	P=0.001
8	35.6 ± 1.41 (20)	35.8 ± 2.78 (20)	35.2 ± 1.80 (20)	*28.1 ± 1.26 (20)	P<0.001
9	35.4 ± 1.78 (20)	33.6 ± 1.52 (20)	35.1 ± 1.70 (20)	*28.2 ± 1.15 (20)	P=0.004
10	34.0 ± 1.36 (20)	32.9 ± 1.62 (20)	32.6 ± 1.69 (20)	*26.7 ± 0.89 (20)	P<0.001
13	35.3 ± 1.54 (19)	34.3 ± 3.74 (20)	33.4 ± 1.82 (19)	*27.5 ± 1.49 (20)	P=0.002
14	33.6 ± 1.70 (20)	32.2 ± 1.64 (20)	32.8 ± 1.64 (20)	*27.6 ± 1.52 (20)	P=0.011
15	32.9 ± 1.73 (20)	31.5 ± 2.03 (20)	30.7 ± 1.17 (20)	*25.3 ± 0.97 (20)	P<0.001
16	34.1 ± 1.42 (20)	31.0 ± 1.14 (19)	32.6 ± 1.64 (19)	29.8 ± 1.42 (18)	P=0.031
17	37.7 ± 1.88 (20)	*31.8 ± 1.54 (19)	*31.5 ± 1.59 (20)	*29.9 ± 1.48 (20)	P=0.002
18	35.1 ± 1.78 (20)	31.4 ± 1.30 (19)	33.0 ± 1.53 (20)	*28.2 ± 1.16 (20)	P=0.005
19	33.1 ± 2.21 (20)	31.8 ± 1.43 (19)	33.2 ± 1.37 (20)	*28.6 ± 0.92 (20)	P=0.026
<u>FEMALE</u>					
1	23.6 ± 0.93 (20)	24.4 ± 0.79 (20)	22.7 ± 0.56 (20)	21.6 ± 1.00 (20)	P=0.126
2	23.0 ± 1.13 (20)	24.1 ± 0.77 (20)	23.2 ± 0.61 (20)	22.3 ± 0.86 (20)	P=0.609
3	25.3 ± 1.20 (20)	25.0 ± 0.97 (20)	23.8 ± 0.83 (20)	22.1 ± 1.07 (20)	P=0.011
4	24.8 ± 1.06 (20)	24.7 ± 0.93 (20)	24.2 ± 0.75 (20)	*21.5 ± 0.95 (20)	P=0.010
5	24.4 ± 0.93 (20)	26.4 ± 1.16 (20)	26.7 ± 0.69 (20)	22.6 ± 0.96 (20)	P=0.246
6	25.4 ± 1.04 (20)	27.2 ± 0.93 (20)	25.0 ± 0.75 (20)	21.9 ± 1.14 (20)	P=0.017
7	24.1 ± 1.57 (20)	27.0 ± 1.06 (20)	24.5 ± 0.85 (20)	24.1 ± 2.23 (19)	P=0.065
8	26.4 ± 1.12 (20)	27.7 ± 1.96 (20)	23.7 ± 0.76 (20)	*22.2 ± 1.02 (20)	P=0.002
9	24.9 ± 1.30 (20)	24.9 ± 0.89 (20)	23.2 ± 0.60 (20)	21.4 ± 0.86 (20)	P=0.024
10	24.2 ± 1.05 (20)	24.5 ± 0.97 (20)	21.5 ± 0.68 (20)	*20.0 ± 0.92 (20)	P<0.001
13	39.1 ± 1.99 (18)	41.6 ± 1.81 (20)	39.1 ± 2.67 (20)	*31.8 ± 2.19 (20)	P=0.006
18	36.4 ± 2.29 (20)	31.9 ± 1.43 (20)	*29.9 ± 1.13 (20)	*22.7 ± 1.53 (18)	P<0.001
19	33.4 ± 2.04 (20)	31.0 ± 1.58 (19)	28.3 ± 1.05 (20)	*25.5 ± 1.66 (18)	P=0.001

a Mean ± standard error (number of animals).

b Each dose group is compared to the control with Shirley's test when a trend is present (P<0.01 from Jonckheere's trend test), otherwise Dunn's test is applied (*= P<0.05).

TABLE 0-4
F₀: SUMMARY OF ESTIMATED DOSAGE (MG/KG BODY WEIGHT/DAY)
TWO GENERATION REPRODUCTION TOXICITY STUDY OF PROPYLTHIOURACIL WHEN ADMINISTERED TO SPRAGUE-DAWLEY RATS IN THE DRINKING WATER

WEEK OF STUDY	DOSE GROUPS ^a		
	0.0001 %	0.0004 %	0.0015 %
	<u>MALE</u>		
1	0.1 ± 0.00 ((20))	0.5 ± 0.01 (20)	1.8 ± 0.04 (20)
2	0.1 ± 0.01 ((20))	0.4 ± 0.02 (20)	1.5 ± 0.04 (20)
3	0.1 ± 0.01 ((20))	0.4 ± 0.02 (20)	1.4 ± 0.04 (19)
4	0.1 ± 0.00 ((20))	0.4 ± 0.01 (20)	1.3 ± 0.05 (20)
5	0.1 ± 0.00 ((20))	0.3 ± 0.01 (20)	1.2 ± 0.05 (20)
6	0.1 ± 0.00 ((20))	0.3 ± 0.01 (20)	1.2 ± 0.05 (20)
7	0.1 ± 0.00 ((19))	0.3 ± 0.01 (20)	1.1 ± 0.05 (20)
8	0.1 ± 0.01 ((20))	0.3 ± 0.02 (20)	1.1 ± 0.06 (20)
9	0.1 ± 0.00 ((20))	0.3 ± 0.01 (20)	1.1 ± 0.06 (20)
10	0.1 ± 0.00 ((20))	0.3 ± 0.01 (20)	1.0 ± 0.04 (20)
13	0.1 ± 0.01 ((20))	0.3 ± 0.02 (19)	1.0 ± 0.05 (20)
14	0.1 ± 0.01 ((20))	0.2 ± 0.01 (20)	1.0 ± 0.06 (20)
15	0.1 ± 0.01 ((20))	0.2 ± 0.01 (20)	0.9 ± 0.04 (20)
16	0.1 ± 0.00 ((19))	0.2 ± 0.01 (19)	1.0 ± 0.06 (18)
17	0.1 ± 0.00 ((19))	0.2 ± 0.01 (20)	1.0 ± 0.05 (20)
18	0.1 ± 0.00 ((19))	0.2 ± 0.01 (20)	0.9 ± 0.04 (20)
19	0.1 ± 0.00 ((19))	0.2 ± 0.01 (20)	0.9 ± 0.04 (20)
	<u>FEMALE</u>		
1	0.1 ± 0.00 ((20))	0.5 ± 0.01 (20)	1.7 ± 0.08 (20)
2	0.1 ± 0.00 ((20))	0.4 ± 0.01 (20)	1.6 ± 0.07 (20)
3	0.1 ± 0.00 ((20))	0.4 ± 0.01 (20)	1.5 ± 0.07 (20)
4	0.1 ± 0.00 ((20))	0.4 ± 0.01 (20)	1.4 ± 0.06 (20)
5	0.1 ± 0.00 ((20))	0.4 ± 0.01 (20)	1.4 ± 0.07 (20)
6	0.1 ± 0.00 ((20))	0.4 ± 0.01 (20)	1.4 ± 0.08 (20)
7	0.1 ± 0.00 ((20))	0.4 ± 0.01 (20)	1.4 ± 0.15 (19)
8	0.1 ± 0.01 ((20))	0.3 ± 0.01 (20)	1.3 ± 0.06 (20)
9	0.1 ± 0.00 ((20))	0.3 ± 0.01 (20)	1.2 ± 0.06 (20)
10	0.1 ± 0.00 ((20))	0.3 ± 0.01 (20)	1.1 ± 0.06 (20)
13	0.1 ± 0.00 ((20))	0.5 ± 0.03 (20)	1.6 ± 0.13 (20)
18	0.1 ± 0.00 ((20))	0.4 ± 0.01 (20)	1.2 ± 0.08 (18)
19	0.1 ± 0.00 ((19))	0.4 ± 0.01 (20)	1.3 ± 0.10 (18)

a Mean ± standard error (number of animals).

TABLE 0-5
F₀: SUMMARY OF REPRODUCTIVE PERFORMANCE OF COHABITED PAIRS
TWO GENERATION REPRODUCTION TOXICITY STUDY OF PROPYLTHIOURACIL WHEN ADMINISTERED TO SPRAGUE-DAWLEY RATS IN THE DRINKING WATER

REPRODUCTIVE PARAMETER	DOSE GROUP ^a								TREND
	0.0000 %		0.0001 %		0.0004 %		0.0015 %		
Pregnancy Index ^{b,e,g}	16/20	(80)	18/20	(90)	18/20	(90)	15/20	(75)	P=0.058
Live Pups Per Litter ^{c,h,i}									
Male	6.2 ± 0.46	(16)	5.6 ± 0.48	(18)	5.4 ± 0.62	(18)	5.5 ± 0.41	(15)	P=0.343
Female	8.6 ± 0.53	(16)	7.0 ± 0.68	(18)	7.6 ± 0.68	(18)	*4.9 ± 0.59	(15)	P=0.001
Combined	14.8 ± 0.51	(16)	12.6 ± 0.95	(18)	13.8 ± 0.58	(17)	*10.3 ± 0.73	(15)	P<0.001
Proportion Pups Born Alive ^{c,h,i}	0.99 ± 0.008	(16)	0.99 ± 0.007	(18)	0.98 ± 0.015	(17)	0.95 ± 0.019	(15)	P=0.068
Sex of Pups Born Alive (Males/Total) ^{c,j,i}	0.42 ± 0.030	(16)	0.43 ± 0.037	(18)	0.41 ± 0.036	(17)	0.55 ± 0.040	(15)	P=0.045
Live Pup Weight (grams) ^{c,j,i}									
Male	6.77 ± 0.135	(16)	6.84 ± 0.174	(17)	6.82 ± 0.166	(17)	6.87 ± 0.220	(15)	P=0.967
Female	6.44 ± 0.116	(16)	6.56 ± 0.183	(18)	6.50 ± 0.134	(17)	6.54 ± 0.228	(15)	P=0.963
Combined	6.58 ± 0.115	(16)	6.71 ± 0.180	(18)	6.66 ± 0.133	(17)	6.74 ± 0.227	(15)	P=0.810
Adjusted Live Pup Weight (grams) ^{c,k,l}									OVERALL
Male	6.90 ± 0.175	(16)	6.83 ± 0.163	(17)	6.89 ± 0.165	(17)	6.68 ± 0.189	(15)	P=0.849
Female	6.60 ± 0.164	(16)	6.51 ± 0.149	(18)	6.59 ± 0.155	(17)	6.33 ± 0.173	(15)	P=0.686
Combined	6.75 ± 0.161	(16)	6.66 ± 0.146	(18)	6.75 ± 0.152	(17)	6.52 ± 0.169	(15)	P=0.760
Average Dam Weight (grams) ^{c,m,i}	345.0 ± 9.94	(16)	338.7 ± 7.55	(18)	*322.0 ± 9.89	(17)	*281.7 ± 6.33	(15)	P<0.001
Average Sire Weight (grams) ^{n,i}	559.7 ± 16.47	(16)	579.2 ± 10.93	(18)	562.9 ± 14.69	(17)	*449.3 ± 13.46	(15)	P<0.001
Average Days to Litter ^{c,m,i}	25.3 ± 0.31	(16)	25.4 ± 0.33	(18)	25.3 ± 0.77	(18)	25.7 ± 0.33	(15)	P=0.676

a Statistical significance for comparisons of dosed groups to controls: * = P<0.05.

b Only those pairs surviving cohabitation were included for statistical analysis of data.

c Only those females surviving to the end of the parturition period were included for statistical analysis of the data.

d Number of females with vaginal sperm / number of cohabiting pairs (percent with vaginal sperm).

e Number of fertile pairs / number of cohabiting pairs (percent pregnant).

f Number of fertile pairs / number of females with vaginal sperm (percent fertile).

g P-value from Cochran-Armitage trend test. Dosed groups compared to control group by Chi-squared test. If fertility index > 1, then it is set equal to 1 for statistical analysis.

h Mean ± standard error (number of pregnant pairs).

i Each dose group is compared to the control group by Shirley's test if P<0.01 from Jonckheere's trend test. Otherwise, Dunn's test is applied.

j Mean ± standard error (number of pregnant pairs producing live pups).

k Least squares estimate of mean pup weight adjusted for average litter size ± standard error (number of pregnant pairs producing live pups).

l Overall differences tested with an F-test, pairwise comparisons from Dunnett's test.

m Mean ± standard error (number of dams).

n Mean ± standard error (number of sires).

TABLE 0-6
F₀: SUMMARY OF DAM WEIGHTS (GRAMS) DURING LACTATION OF THE F₁ LITTER
TWO GENERATION REPRODUCTION TOXICITY STUDY OF PROPYLTHIOURACIL WHEN ADMINISTERED TO SPRAGUE-DAWLEY RATS IN THE DRINKING WATER

AGE (DAYS)	DOSE GROUPS ^a				TREND ^b
	0.0000 %	0.0001 %	0.0004 %	0.0015 %	
1	345.0 ± 9.94 (16)	338.7 ± 7.55 (18)	*322.0 ± 9.89 (17)	*281.7 ± 6.33 (15)	P<0.001
4	348.2 ± 9.96 (14)	338.0 ± 7.93 (16)	332.6 ± 8.94 (11)	*288.0 ± 7.15 (14)	P<0.001
7	367.8 ± 13.01 (16)	355.3 ± 13.40 (18)	353.9 ± 8.28 (17)	*311.8 ± 11.51 (15)	P=0.012
14	368.0 ± 8.05 (16)	358.3 ± 6.69 (18)	357.3 ± 7.61 (17)	*317.8 ± 6.43 (15)	P<0.001
21	321.6 ± 10.43 (14)	334.8 ± 8.24 (15)	337.2 ± 5.76 (16)	320.9 ± 5.60 (14)	P=0.807

a Mean ± standard error (number of dams).

b Each dose group is compared to the control with Shirley's test when a trend is present (P<0.01 from Jonckheere's trend test), otherwise Dunn's test is applied (* = P<0.05).

TABLE 0-7A
 F₀: SUMMARY OF DAM DAILY FEED CONSUMPTION (G/KG BODY WEIGHT/DAY) DURING LACTATION OF THE F₁ LITTER
 TWO GENERATION REPRODUCTION TOXICITY STUDY OF PROPYLTHIOURACIL WHEN ADMINISTERED TO SPRAGUE-DAWLEY RATS IN THE DRINKING WATER

POSTNATAL DAYS	DOSE GROUPS ^a				TREND ^b
	0.0000 %	0.0001 %	0.0004 %	0.0015 %	
	<u>FEMALE</u>				
1-4	123.2 ± 20.64 (14)	90.5 ± 6.69 (15)	129.0 ± 19.19 (11)	83.3 ± 3.13 (14)	P=0.153
4-7	134.2 ± 7.00 (16)	*99.6 ± 10.87 (18)	115.0 ± 10.14 (17)	*77.7 ± 16.39 (14)	P=0.002
11-14	190.3 ± 7.85 (16)	187.8 ± 10.65 (18)	*237.6 ± 12.53 (17)	170.8 ± 16.23 (15)	P=0.814
18-21	266.0 ± 9.42 (14)	246.6 ± 6.12 (15)	270.3 ± 6.52 (16)	*181.4 ± 13.07 (14)	P=0.002

a Mean ± standard error (number of dams).

b Each dose group is compared to the control with Shirley's test when a trend is present (P<0.01 from Jonckheere's trend test), otherwise Dunn's test is applied (* = P<0.05).

TABLE 0-7B
 F₀: SUMMARY OF DAM DAILY FEED CONSUMPTION (G/ANIMAL/DAY) DURING LACTATION OF THE F₁ LITTER
 TWO GENERATION REPRODUCTION TOXICITY STUDY OF PROPYLTHIOURACIL WHEN ADMINISTERED TO SPRAGUE-DAWLEY RATS IN THE DRINKING WATER

POSTNATAL DAYS	DOSE GROUPS ^a				TREND ^b
	0.0000 %	0.0001 %	0.0004 %	0.0015 %	
	<u>FEMALE</u>				
1-4	45.5 ± 7.47 (16)	*29.1 ± 2.01 (17)	38.9 ± 4.94 (15)	*28.7 ± 4.91 (15)	P=0.004
4-7	48.8 ± 2.77 (16)	*33.4 ± 3.51 (18)	*40.8 ± 3.57 (17)	*25.0 ± 6.00 (14)	P<0.001
7-11	68.3 ± 3.54 (16)	61.3 ± 4.61 (18)	58.5 ± 5.27 (17)	*46.9 ± 5.89 (15)	P=0.009
11-14	69.6 ± 2.64 (16)	66.5 ± 3.13 (18)	*84.6 ± 4.37 (17)	54.4 ± 5.37 (15)	P=0.740
14-18	69.2 ± 2.16 (16)	68.3 ± 2.12 (18)	76.3 ± 1.51 (17)	62.5 ± 3.81 (14)	P=0.953
18-21	85.0 ± 2.17 (16)	78.6 ± 3.62 (18)	90.7 ± 2.40 (17)	*58.7 ± 4.71 (14)	P=0.014

a Mean ± standard error (number of dams).

b Each dose group is compared to the control with Shirley's test when a trend is present (P<0.01 from Jonckheere's trend test), otherwise Dunn's test is applied (* = P<0.05).

TABLE 0-8A
 F₀: SUMMARY OF DAM DAILY WATER CONSUMPTION (G/KG BODY WEIGHT/DAY) DURING LACTATION OF THE F₁ LITTER
 TWO GENERATION REPRODUCTION TOXICITY STUDY OF PROPYLTHIOURACIL WHEN ADMINISTERED TO SPRAGUE-DAWLEY RATS IN THE DRINKING WATER

POSTNATAL DAYS	DOSE GROUPS ^a				TREND ^b
	0.0000 %	0.0001 %	0.0004 %	0.0015 %	
	<u>FEMALE</u>				
1-4	137.0 ± 10.78 (14)	123.2 ± 11.03 (15)	131.2 ± 9.80 (11)	125.1 ± 8.84 (14)	P=0.799
4-7	181.9 ± 19.53 (16)	152.4 ± 13.19 (18)	156.9 ± 9.16 (17)	*131.3 ± 9.48 (15)	P=0.027
11-14	254.9 ± 14.61 (16)	235.5 ± 15.70 (18)	270.5 ± 11.95 (17)	*183.6 ± 12.74 (15)	P=0.041
18-21	375.3 ± 24.22 (13)	316.8 ± 15.39 (15)	334.6 ± 11.16 (15)	*224.9 ± 14.34 (14)	P<0.001

a Mean ± standard error (number of dams).

b Each dose group is compared to the control with Shirley's test when a trend is present (P<0.01 from Jonckheere's trend test), otherwise Dunn's test is applied (* = P<0.05).

TABLE 0-8B
 F₀: SUMMARY OF DAM DAILY WATER CONSUMPTION (G/ANIMAL/DAY) DURING LACTATION OF THE F₁ LITTER
 TWO GENERATION REPRODUCTION TOXICITY STUDY OF PROPYLTHIOURACIL WHEN ADMINISTERED TO SPRAGUE-DAWLEY RATS IN THE DRINKING WATER

POSTNATAL DAYS	DOSE GROUPS ^a				TREND ^b
	0.0000 %	0.0001 %	0.0004 %	0.0015 %	
	<u>FEMALE</u>				
1-4	46.3 ± 2.70 (16)	39.7 ± 3.26 (17)	43.9 ± 2.15 (17)	*34.9 ± 2.38 (15)	P=0.023
4-7	65.1 ± 5.59 (16)	52.6 ± 4.08 (18)	55.0 ± 3.04 (17)	*40.2 ± 2.75 (15)	P<0.001
7-11	76.7 ± 3.96 (16)	68.8 ± 4.46 (18)	66.4 ± 5.54 (17)	*51.6 ± 5.80 (15)	P=0.003
11-14	92.7 ± 3.88 (16)	83.4 ± 4.88 (18)	96.3 ± 4.07 (17)	*58.5 ± 4.41 (15)	P=0.003
14-18	93.7 ± 2.19 (16)	*83.3 ± 3.85 (18)	89.8 ± 2.38 (17)	*70.3 ± 5.05 (13)	P=0.001
18-21	118.6 ± 4.02 (15)	*99.3 ± 5.71 (18)	112.4 ± 3.89 (16)	*72.6 ± 5.35 (14)	P<0.001

a Mean ± standard error (number of dams).

b Each dose group is compared to the control with Shirley's test when a trend is present (P<0.01 from Jonckheere's trend test), otherwise Dunn's test is applied (* = P<0.05).

TABLE 0-9
F₀: SUMMARY OF ESTIMATED DOSAGE (MG/KG BODY WEIGHT/DAY) DURING LACTATION OF THE F₁ LITTER
TWO GENERATION REPRODUCTION TOXICITY STUDY OF PROPYLTHIOURACIL WHEN ADMINISTERED TO SPRAGUE-DAWLEY RATS IN THE DRINKING WATER

POSTNATAL DAYS	DOSE GROUPS ^a		
	0.0001 %	0.0004 %	0.0015 %
	<u>FEMALE</u>		
1-4	0.1 ± 0.01 (15)	0.5 ± 0.04 (11)	1.9 ± 0.13 (14)
4-7	0.2 ± 0.01 (18)	0.6 ± 0.04 (17)	2.0 ± 0.14 (15)
11-18	0.2 ± 0.02 (18)	1.1 ± 0.05 (17)	2.8 ± 0.19 (15)
18-21	0.3 ± 0.02 (15)	1.3 ± 0.04 (15)	3.4 ± 0.22 (14)

a Mean ± standard error (number of dams).

b Each dose group is compared to the control with Shirley's test when a trend is present (P<0.01 from Jonckheere's trend test), otherwise Dunn's test is applied (* = P<0.05).

TABLE 0-10
F₀: SUMMARY OF ANOGENITAL DISTANCE AND PUP WEIGHT FOR THE F₁ LITTER
TWO GENERATION REPRODUCTION TOXICITY STUDY OF PROPYLTHIOURACIL WHEN ADMINISTERED TO SPRAGUE-DAWLEY RATS IN THE DRINKING WATER

Endpoint	DOSE GROUPS ^a				TREND ^b
	0.0000 %	0.0001 %	0.0004 %	0.0015 %	
<u>MALE</u>					
Anogenital Distance (PND1)	2.58 ± 0.041 (16)	2.58 ± 0.030 (17)	2.59 ± 0.038 (17)	2.61 ± 0.043 (15)	P=0.586
Average Pup Weight (PND1)	6.75 ± 0.125 (16)	6.83 ± 0.175 (17)	6.84 ± 0.176 (17)	6.63 ± 0.155 (15)	P=0.554
Ratio AGD/Pup Weight (PND1)	0.39 ± 0.010 (16)	0.38 ± 0.011 (17)	0.38 ± 0.008 (17)	0.40 ± 0.010 (15)	P=0.274
<u>FEMALE</u>					
Anogenital Distance (PND1)	1.14 ± 0.024 (16)	1.14 ± 0.020 (18)	1.15 ± 0.019 (17)	1.12 ± 0.013 (15)	P=0.688
Average Pup Weight (PND1)	6.42 ± 0.115 (16)	6.53 ± 0.177 (18)	6.51 ± 0.136 (17)	6.24 ± 0.145 (15)	P=0.381
Ratio AGD/Pup Weight (PND1)	0.18 ± 0.006 (16)	0.18 ± 0.003 (18)	0.18 ± 0.004 (17)	0.18 ± 0.005 (15)	P=0.509

a Mean ± standard error (number of dams).

b Each dose group is compared to the control with Shirley's test when a trend is present (P<0.01 from Jonckheere's trend test), otherwise Dunn's test is applied (* = P<0.05).

TABLE 0-11
 F₀: SUMMARY OF AVERAGE PUP WEIGHTS FOR THE F₁ LITTER
 TWO GENERATION REPRODUCTION TOXICITY STUDY OF PROPYLTHIOURACIL WHEN ADMINISTERED TO SPRAGUE-DAWLEY RATS IN THE DRINKING WATER

AGE (DAYS)	DOSE GROUPS ^a				TREND ^b
	0.0000 %	0.0001 %	0.0004 %	0.0015 %	
<u>MALE</u>					
1	6.77 ± 0.135 (16)	6.84 ± 0.174 (17)	6.82 ± 0.166 (17)	6.87 ± 0.220 (15)	P=0.967
4	9.55 ± 0.245 (14)	9.80 ± 0.392 (15)	10.29 ± 0.575 (12)	10.48 ± 1.070 (14)	P=0.946
7	15.83 ± 1.564 (16)	17.38 ± 2.221 (17)	17.85 ± 2.424 (17)	17.75 ± 2.716 (13)	P=0.572
14	27.53 ± 0.701 (16)	27.69 ± 0.796 (17)	28.20 ± 1.383 (17)	*23.39 ± 0.690 (15)	P<0.001
21	45.63 ± 5.280 (16)	56.45 ± 9.869 (17)	39.96 ± 1.217 (16)	*30.77 ± 1.039 (14)	P<0.001
<u>FEMALE</u>					
1	6.44 ± 0.116 (16)	6.56 ± 0.183 (18)	6.50 ± 0.134 (17)	6.54 ± 0.228 (15)	P=0.963
4	9.25 ± 0.223 (14)	9.50 ± 0.328 (16)	10.06 ± 0.460 (12)	10.35 ± 1.145 (14)	P=0.650
7	13.69 ± 0.779 (16)	16.43 ± 1.967 (18)	15.89 ± 1.438 (17)	17.27 ± 3.549 (13)	P=0.971
14	26.67 ± 0.768 (16)	26.78 ± 0.795 (18)	27.20 ± 1.293 (17)	*22.75 ± 0.752 (15)	P=0.003
21	47.16 ± 5.859 (16)	68.76 ± 16.683 (18)	38.76 ± 1.334 (16)	*29.71 ± 0.830 (14)	P<0.001

a Mean ± standard error (number of litters).

b Each dose group is compared to the control with Shirley's test when a trend is present (P<0.01 from Jonckheere's trend test), otherwise Dunn's test is applied (* = P<0.05).

TABLE 0-12
 F₀: SUMMARY OF PUP SURVIVAL FOR THE F₁ LITTER
 TWO GENERATION REPRODUCTION TOXICITY STUDY OF PROPYLTHIOURACIL WHEN ADMINISTERED TO SPRAGUE-DAWLEY RATS IN THE DRINKING WATER

AGE (DAYS)	DOSE GROUPS ^a				TREND ^b
	0.0000 %	0.0001 %	0.0004 %	0.0015 %	
<u>MALE</u>					
4	1.00 ± 0.000 (16)	0.97 ± 0.014 (17)	0.99 ± 0.007 (17)	0.96 ± 0.028 (15)	P=0.381
7	1.00 ± 0.000 (16)	0.97 ± 0.014 (17)	0.99 ± 0.010 (17)	0.96 ± 0.028 (15)	P=0.310
14	0.97 ± 0.015 (16)	0.97 ± 0.014 (17)	0.99 ± 0.010 (17)	0.96 ± 0.028 (15)	P=0.633
21	0.97 ± 0.015 (16)	0.97 ± 0.014 (17)	0.99 ± 0.010 (17)	0.99 ± 0.012 (14)	P=0.315
<u>FEMALE</u>					
4	0.99 ± 0.006 (16)	0.98 ± 0.018 (18)	1.00 ± 0.000 (17)	0.94 ± 0.030 (15)	P=0.266
7	0.98 ± 0.014 (16)	0.98 ± 0.019 (18)	1.00 ± 0.000 (17)	0.94 ± 0.030 (15)	P=0.764
14	0.97 ± 0.015 (16)	0.97 ± 0.019 (18)	1.00 ± 0.000 (17)	0.94 ± 0.030 (15)	P=0.766
21	0.97 ± 0.016 (16)	0.97 ± 0.019 (18)	1.00 ± 0.000 (17)	0.96 ± 0.024 (14)	P=0.211
<u>COMBINED</u>					
4	1.00 ± 0.004 (16)	0.98 ± 0.015 (18)	1.00 ± 0.005 (17)	0.95 ± 0.026 (15)	P=0.194
7	0.99 ± 0.009 (16)	0.97 ± 0.015 (18)	0.99 ± 0.006 (17)	0.95 ± 0.026 (15)	P=0.443
14	0.97 ± 0.011 (16)	0.97 ± 0.015 (18)	0.99 ± 0.006 (17)	0.95 ± 0.026 (15)	P=0.636
21	0.97 ± 0.012 (16)	0.97 ± 0.015 (18)	0.99 ± 0.006 (17)	0.98 ± 0.014 (14)	P=0.320

a Mean of (number of live pups/number of pups born alive) ± standard error (number of litters). Ratios greater than 1 or increases in survival over time indicate missexing of pups.

b Each dose group is compared to the control with Shirley's test when a trend is present (P<0.01 from Jonckheere's trend test), otherwise Dunn's test is applied (* = P<0.05).

TABLE 0-13
F₀: SUMMARY OF NIPPLE RETENTION FOR THE F₁ LITTER
TWO GENERATION REPRODUCTION TOXICITY STUDY OF PROPYLTHIOURACIL WHEN ADMINISTERED TO SPRAGUE-DAWLEY RATS IN THE DRINKING WATER

AGE (DAYS)	DOSE GROUPS ^a			
	0.0000 %	0.0001 %	0.0004 %	0.0015 %
12	0.00 ± 0.00 (16)	0.00 ± 0.00 (18)	0.00 ± 0.00 (17)	0.00 ± 0.00 (15)
13	0.00 ± 0.00 (16)	0.00 ± 0.00 (18)	0.00 ± 0.00 (17)	0.00 ± 0.00 (15)

a Mean ± standard error (number of animals).

TABLE 0-14
 F₀: SUMMARY OF PINNA DETACHMENT AND EYE OPENING FOR THE F₁ LITTER
 TWO GENERATION REPRODUCTION TOXICITY STUDY OF PROPYLTHIOURACIL WHEN ADMINISTERED TO SPRAGUE-DAWLEY RATS IN THE DRINKING WATER

ENDPOINT ^b	DOSE GROUPS ^a				TREND ^c
	0.0000 %	0.0001 %	0.0004 %	0.0015 %	
DAY OF EYE OPENING	15.7 ± 0.20 (16)	15.6 ± 0.22 (18)	16.2 ± 0.18 (17)	*17.6 ± 0.36 (14)	P<0.001
DAY OF PINNA DETACHMENT	4.2 ± 0.21 (16)	4.1 ± 0.16 (18)	4.0 ± 0.19 (17)	4.5 ± 0.19 (15)	P=0.538

a Mean ± standard error (number of animals).

b Mean = age (days)

c Each dose group is compared to the control with Shirley's test when a trend is present (P<0.01 from Jonckheere's trend test), otherwise Dunn's test is applied (* = P<0.05).

TABLE 0-15
 F₀: SUMMARY OF F₁ SEXUAL DEVELOPMENT DATA
 TWO GENERATION REPRODUCTION TOXICITY STUDY OF PROPYLTHIOURACIL WHEN ADMINISTERED TO SPRAGUE-DAWLEY RATS IN THE DRINKING WATER

ENDPOINT ^B	DOSE GROUPS ^a			TREND ^c
	0.0000 %	0.0001 %	0.0004 %	
	<u>MALE</u>			
DAY OF PREPUCE SEPARATION	42.7 ± 0.31 (28)	43.3 ± 0.41 (30)	*44.7 ± 0.74 (31)	P=0.009
DAY OF TESTICULAR DESCENT	28.7 ± 0.36 (28)	29.4 ± 0.46 (30)	30.2 ± 0.82 (31)	P=0.175
	<u>FEMALE</u>			
DAY OF VAGINAL OPENING	32.4 ± 0.35 (30)	32.9 ± 0.39 (33)	*33.8 ± 0.43 (33)	P=0.020

a Mean ± standard error (number of animals).

b Mean = age (days)

c Each dose group is compared to the control with Shirley's test when a trend is present (P<0.01 from Jonckheere's trend test), otherwise Dunn's test is applied (* = P<0.05).

TABLE 0-16
 F₀: SUMMARY OF F₁ MALE PND 21 NECROPSY GROSS OBSERVATIONS
 TWO GENERATION REPRODUCTION TOXICITY STUDY OF PROPYLTHIOURACIL WHEN ADMINISTERED TO SPRAGUE-DAWLEY RATS IN THE DRINKING WATER

	GROUP :	1	2	3	4
	DOSE (%) :	0.0000	0.0001	0.0004	0.0015
	NUMBER OF ANIMALS ON STUDY:	47	50	46	41
	NUMBER OF ANIMALS COMPLETED:	(46)	(50)	(46)	(41)
RIGHT TESTIS;					
SUBMITTED.....	(47)	(50)	(46)	(40)	
NO VISIBLE LESIONS.....	47	50	46	41	
LIVER;					
SUBMITTED.....	(47)	(50)	(46)	(41)	
NO VISIBLE LESIONS.....	47	50	46	41	
LEFT TESTES W/EPIDIDYMIDES;					
SUBMITTED.....	(47)	(49)	(46)	(40)	
NO VISIBLE LESIONS.....	47	50	46	41	
THYROID/PARATHYROID;					
SUBMITTED.....	(47)	(50)	(46)	(41)	
NO VISIBLE LESIONS.....	47	50	46	41	
SEMINAL VESICLES W/ COAGULATING GLANDS;					
SUBMITTED.....	(47)	(50)	(46)	(40)	
NO VISIBLE LESIONS.....	47	50	46	41	
ADRENAL GLAND(S);					
SUBMITTED.....	(47)	(50)	(46)	(41)	
NO VISIBLE LESIONS.....	47	50	46	41	
BRAIN;					
SUBMITTED.....	(0)	(0)	(0)	(0)	
NO VISIBLE LESIONS.....	47	50	46	41	
KIDNEY(S);					
SUBMITTED.....	(47)	(50)	(46)	(41)	
NO VISIBLE LESIONS.....	47	49	45	40	
DILATION; RIGHT	0	1	1	2	

TABLE 0-16 (CONTINUED)
 F₀: SUMMARY OF F₁ MALE PND 21 NECROPSY GROSS OBSERVATIONS
 TWO GENERATION REPRODUCTION TOXICITY STUDY OF PROPYLTHIOURACIL WHEN ADMINISTERED TO SPRAGUE-DAWLEY RATS IN THE DRINKING WATER

	GROUP :	1	2	3	4
	DOSE (%) :	0.0000	0.0001	0.0004	0.0015
MAMMARY GLAND(S);					
SUBMITTED.....		(0)	(1)	(0)	(0)
NO VISIBLE LESIONS.....		0	0	0	0
DISCOLORATION; BLUE; LEFT; FOCAL		0	1	0	0
PITUITARY GLAND;					
SUBMITTED.....		(47)	(50)	(46)	(41)
NO VISIBLE LESIONS.....		47	50	46	41
SPLEEN;					
SUBMITTED.....		(0)	(0)	(1)	(0)
NO VISIBLE LESIONS.....		47	50	46	41
THYMUS;					
SUBMITTED.....		(44)	(47)	(42)	(36)
NO VISIBLE LESIONS.....		47	50	45	41
MASS (2)		0	0	1	0
MASS (4)		0	0	1	0
NODULE(S)		0	0	1	0
VENTRAL PROSTATE;					
SUBMITTED.....		(47)	(50)	(46)	(40)
NO VISIBLE LESIONS.....		47	50	46	41
DORSOLATERAL PROSTATE;					
SUBMITTED.....		(47)	(50)	(46)	(40)
NO VISIBLE LESIONS.....		47	50	46	41

TABLE 0-17
 F₀: SUMMARY OF F₁ MALE PND 21 ABSOLUTE ORGAN WEIGHTS AND ORGAN-TO-BODY WEIGHT RATIOS
 TWO GENERATION REPRODUCTION TOXICITY STUDY OF PROPYLTHIOURACIL WHEN ADMINISTERED TO SPRAGUE-DAWLEY RATS IN THE DRINKING WATER

ORGAN	DOSE GROUPS				TREND ^a
	0.0000 %	0.0001 %	0.0004 %	0.0015 %	
	<u>ORGAN WEIGHTS (g)^b</u>				
BODY	40.3 ± 0.87 (47)	42.9 ± 1.01 (50)	41.1 ± 0.73 (46)	*30.9 ± 0.86 (41)	P<0.001
BRAIN	1.389 ± 0.0109 (47)	1.394 ± 0.0110 (50)	1.425 ± 0.0111 (46)	*1.306 ± 0.0138 (41)	P=0.026
SPLEEN	0.160 ± 0.0048 (47)	0.170 ± 0.0049 (50)	0.154 ± 0.0049 (46)	0.154 ± 0.0061 (41)	P=0.245
THYMUS	0.155 ± 0.0186 (47)	0.214 ± 0.0189 (50)	0.154 ± 0.0190 (46)	0.103 ± 0.0237 (41)	P=0.105
	<u>ORGAN-TO-BODY WEIGHT RATIOS^c</u>				
BRAIN	35.342 ± 0.7379 (47)	33.874 ± 0.6416 (50)	35.474 ± 0.5677 (46)	*41.089 ± 0.9854 (41)	P<0.001
SPLEEN	4.090 ± 0.1275 (47)	4.409 ± 0.1237 (50)	3.986 ± 0.0971 (46)	*3.465 ± 0.2631 (41)	P<0.001
THYMUS	3.883 ± 0.1434 (47)	5.406 ± 0.8582 (50)	3.925 ± 0.1042 (46)	*2.296 ± 0.0854 (41)	P<0.001

a Each dose group is compared with the control group by Shirley's test if P<0.01 from Jonckheere's trend test, otherwise Dunn's test is applied (* = P<0.05).

b Mean organ weight ± standard error (number of animals).

c Mean ratio (mg/g body weight) ± standard error (number of animals).

Note: Absolute organ weights were covaried by body weight.

TABLE 0-18
 F₀: SUMMARY OF F₁ FEMALE PND 21 NECROPSY GROSS OBSERVATIONS
 TWO GENERATION REPRODUCTION TOXICITY STUDY OF PROPYLTHIOURACIL WHEN ADMINISTERED TO SPRAGUE-DAWLEY RATS IN THE DRINKING WATER

	GROUP :	1	2	3	4
	DOSE (%) :	0.0000	0.0001	0.0004	0.0015
	NUMBER OF ANIMALS ON STUDY:	48	49	51	32
	NUMBER OF ANIMALS COMPLETED:	(48)	(49)	(51)	(32)
VAGINA/UTERUS/CERVIX;					
SUBMITTED.....	(48)	(49)	(51)	(32)	
NO VISIBLE LESIONS.....	48	49	51	32	
THYROID/PARATHYROID;					
SUBMITTED.....	(48)	(49)	(51)	(32)	
NO VISIBLE LESIONS.....	48	49	51	32	
LIVER;					
SUBMITTED.....	(48)	(49)	(51)	(32)	
NO VISIBLE LESIONS.....	48	49	51	32	
ADRENAL GLAND(S);					
SUBMITTED.....	(48)	(49)	(51)	(32)	
NO VISIBLE LESIONS.....	48	49	51	32	
BRAIN;					
SUBMITTED.....	(0)	(0)	(0)	(0)	
NO VISIBLE LESIONS.....	48	49	51	32	
KIDNEY(S);					
SUBMITTED.....	(48)	(49)	(51)	(32)	
NO VISIBLE LESIONS.....	47	48	51	32	
DILATION; RIGHT	1	1	0	0	
OVARY(IES);					
SUBMITTED.....	(48)	(49)	(51)	(32)	
NO VISIBLE LESIONS.....	48	49	51	32	
PITUITARY GLAND;					
SUBMITTED.....	(48)	(49)	(51)	(32)	
NO VISIBLE LESIONS.....	48	49	51	32	

TABLE 0-18 (CONTINUED)
 F₀: SUMMARY OF F₁ FEMALE PND 21 NECROPSY GROSS OBSERVATIONS
 TWO GENERATION REPRODUCTION TOXICITY STUDY OF PROPYLTHIOURACIL WHEN ADMINISTERED TO SPRAGUE-DAWLEY RATS IN THE DRINKING WATER

	GROUP:	1	2	3	4
	DOSE (%):	0.0000	0.0001	0.0004	0.0015
SPLEEN;					
SUBMITTED.....		(0)	(1)	(0)	(0)
NO VISIBLE LESIONS.....		47	49	51	32
REDUCTION IN SIZE		1	0	0	0
THYMUS;					
SUBMITTED.....		(45)	(44)	(51)	(32)
NO VISIBLE LESIONS.....		48	49	51	32

TABLE 0-19
 F₀: SUMMARY OF F₁ FEMALE PND 21 ABSOLUTE ORGAN WEIGHTS AND ORGAN-TO-BODY WEIGHT RATIOS
 TWO GENERATION REPRODUCTION TOXICITY STUDY OF PROPYLTHIOURACIL WHEN ADMINISTERED TO SPRAGUE-DAWLEY RATS IN THE DRINKING WATER

ORGAN	DOSE GROUPS				TREND ^a
	0.0000 %	0.0001 %	0.0004 %	0.0015 %	
	<u>ORGAN WEIGHTS (g)^b</u>				
BODY	39.6 ± 0.81 (48)	40.9 ± 1.02 (49)	39.4 ± 0.74 (51)	*29.4 ± 0.80 (32)	P<0.001
BRAIN	1.350 ± 0.0116 (48)	1.338 ± 0.0118 (49)	1.387 ± 0.0113 (51)	*1.260 ± 0.0167 (32)	P=0.026
SPLEEN	0.160 ± 0.0049 (48)	0.164 ± 0.0049 (49)	0.154 ± 0.0047 (51)	0.156 ± 0.0070 (32)	P=0.245
THYMUS	0.159 ± 0.0137 (48)	0.159 ± 0.0138 (49)	0.181 ± 0.0132 (51)	0.127 ± 0.0196 (32)	P=0.105
	<u>ORGAN-TO-BODY WEIGHT RATIOS^c</u>				
BRAIN	34.935 ± 0.6078 (48)	34.128 ± 0.7531 (49)	35.981 ± 0.5943 (51)	*40.948 ± 0.9223 (32)	P<0.001
SPLEEN	4.212 ± 0.1151 (48)	4.395 ± 0.1502 (49)	4.054 ± 0.1348 (51)	*3.510 ± 0.2928 (32)	P<0.001
THYMUS	4.200 ± 0.1369 (48)	4.295 ± 0.1167 (49)	4.716 ± 0.5545 (51)	*2.450 ± 0.0929 (32)	P<0.001

a Each dose group is compared with the control group by Shirley's test if P<0.01 from Jonckheere's trend test, otherwise Dunn's test is applied (* = P<0.05).

b Mean organ weight ± standard error (number of animals).

c Mean ratio (mg/g body weight) ± standard error (number of animals).\

Note: Absolute organ weights were covaried by body weight.

TABLE 0-20
F₀: SUMMARY OF STAGES AND LENGTH OF ESTROUS CYCLE
TWO GENERATION REPRODUCTION TOXICITY STUDY OF PROPYLTHIOURACIL WHEN ADMINISTERED TO SPRAGUE-DAWLEY RATS IN THE DRINKING WATER

DOSE %	ESTROUS STAGE (%)					CYCLE LENGTH ^{a,d} (DAYS)	NUMBER OF CYCLES ^{a,d}	NUMBER OF CYCLING FEMALES ^{b,c}	NUMBER OF FEMALES W/REGULAR CYCLE ^{c,e}
	P	E	M	D	N				
0.0000	15.0	29.3	20.0	35.4	0.4	4.1 ± 0.05 (19) ^f	2.6 ± 0.14 (20)	20/20 (100)	19/20 (95)
0.0001	8.6	30.7	21.4	39.3	0.0	4.4 ± 0.16 (20)	2.3 ± 0.12 (20)	20/20 (100)	19/20 (95)
0.0004	7.9	26.8	21.4	42.9	1.1	4.4 ± 0.17 (20)	2.4 ± 0.13 (20)	20/20 (100)	18/20 (90)
0.0015	3.6	33.2	19.3	41.8	2.1	4.5 ± 0.14 (19) ^g	2.4 ± 0.14 (19)	20/20 (100)	*14/20 (70)

P=Proestrus, E=Estrus, M=Metestrus, D=Diestrus, N=Not Clear, Poor Quality, or Insufficient Number of Cells

a Mean ± standard error (number of animals)

b Number of cycling females / number of females sampled (percent cycling).

c Number of females with a regular cycle / number of females sampled (percent regular).

d Each dose group is compared to the control group by Shirley's test if P<0.01 from Jonckheere's trend test. Otherwise, Dunn's test is applied.

e Dose group compared to the control group using the Chi-squared test (* = P<0.05).

f Estrous cycle ten days or longer, not cycling, not available, or unclear in 1 of 20 animals.

g Estrous cycle ten days or longer, not cycling, not available, or unclear in 1 of 20 animals.

NOTE: There is significant evidence (Wilks' criterion P=0.0210) that females in the high dose group differed from the control group in the relative amount of time spent in estrous stages.

TABLE 0-21
 F₀: SUMMARY OF THYROID HORMONES
 TWO GENERATION REPRODUCTION TOXICITY STUDY OF PROPYLTHIOURACIL WHEN ADMINISTERED TO SPRAGUE-DAWLEY RATS IN THE DRINKING WATER

ENDPOINT	DOSE GROUPS ^a				TREND ^b
	0.0000 %	0.0001 %	0.0004 %	0.0015 %	
<u>MALE</u>					
T3 (NG/DL)	96.08 ± 3.634 (20)	96.19 ± 3.913 (19)	95.08 ± 3.744 (20)	88.33 ± 3.358 (20)	P=0.156
T4 (UG/DL)	2.09 ± 0.312 (20)	2.96 ± 0.410 (19)	*1.13 ± 0.217 (20)	*0.47 ± 0.177 (20)	P<0.001
TSH (NG/ML)	3.63 ± 0.644 (20)	3.79 ± 0.563 (19)	*8.20 ± 1.309 (20)	*22.15 ± 2.560 (20)	P<0.001
<u>FEMALE</u>					
T3 (NG/DL)	105.29 ± 3.828 (20)	113.78 ± 4.364 (20)	111.51 ± 4.827 (20)	90.93 ± 7.690 (18)	P=0.294
T4 (UG/DL)	1.19 ± 0.260 (20)	1.41 ± 0.229 (20)	0.57 ± 0.186 (20)	0.61 ± 0.239 (18)	P=0.004
TSH (NG/ML)	3.76 ± 0.837 (20)	4.61 ± 0.747 (20)	*7.19 ± 0.834 (20)	*21.64 ± 1.651 (18)	P<0.001

a Mean ± standard error (number of animals).

b Each dose group is compared to the control with Dunnett's one-tailed low test for T3 and T4 or Dunnett's one-tailed high test for TSH.
 (* = P<0.05 from Jonckheere's trend test.)

TABLE 0-22
 F₀: SUMMARY OF MALE GROSS OBSERVATIONS
 TWO GENERATION REPRODUCTION TOXICITY STUDY OF PROPYLTHIOURACIL WHEN ADMINISTERED TO SPRAGUE-DAWLEY RATS IN THE DRINKING WATER

	GROUP: DOSE (%):	1 0.0000	2 0.0001	3 0.0004	4 0.0015
	NUMBER OF ANIMALS ON STUDY:	20	20	20	20
	NUMBER OF ANIMALS COMPLETED:	(20)	(20)	(20)	(20)
THYROID/PARATHYROID;					
SUBMITTED.....	(20)	(19)	(20)	(20)	
NO VISIBLE LESIONS.....	20	19	13	0	
ENLARGEMENT; BILATERAL	0	0	7	20	
VENTRAL PROSTATE;					
SUBMITTED.....	(20)	(19)	(20)	(20)	
NO VISIBLE LESIONS.....	20	19	20	20	
RIGHT TESTIS;					
SUBMITTED.....	(20)	(19)	(20)	(20)	
NO VISIBLE LESIONS.....	20	19	20	20	
ADIPOSE TISSUE;					
SUBMITTED.....	(0)	(0)	(1)	(0)	
NO VISIBLE LESIONS.....	0	0	0	0	
DISCOLORATION; BLACK; FOCAL	0	0	1	0	
ADRENAL GLAND(S);					
SUBMITTED.....	(20)	(19)	(20)	(20)	
NO VISIBLE LESIONS.....	20	19	20	20	
BRAIN;					
SUBMITTED.....	(20)	(19)	(20)	(20)	
NO VISIBLE LESIONS.....	20	19	20	20	
KIDNEY(S);					
SUBMITTED.....	(20)	(19)	(20)	(20)	
NO VISIBLE LESIONS.....	19	19	20	20	
CYST(S)	1	0	0	0	
LIVER;					
SUBMITTED.....	(20)	(19)	(20)	(20)	
NO VISIBLE LESIONS.....	20	19	20	20	

TABLE 0-22 (CONTINUED)
 F₀: SUMMARY OF MALE GROSS OBSERVATIONS
 TWO GENERATION REPRODUCTION TOXICITY STUDY OF PROPYLTHIOURACIL WHEN ADMINISTERED TO SPRAGUE-DAWLEY RATS IN THE DRINKING WATER

	GROUP:	1	2	3	4
	DOSE (%):	0.0000	0.0001	0.0004	0.0015
MAMMARY GLAND(S);					
SUBMITTED.....		(0)	(0)	(0)	(1)
NO VISIBLE LESIONS.....		0	0	0	0
DISCOLORATION; GREEN; LEFT; FOCAL		0	0	0	1
PITUITARY GLAND;					
SUBMITTED.....		(20)	(19)	(20)	(20)
NO VISIBLE LESIONS.....		20	19	20	20
SPLEEN;					
SUBMITTED.....		(20)	(19)	(20)	(20)
NO VISIBLE LESIONS.....		20	19	20	20
DORSOLATERAL PROSTATE;					
SUBMITTED.....		(20)	(19)	(20)	(20)
NO VISIBLE LESIONS.....		20	19	20	20
LEFT TESTIS W/ EPIDIDYMIS;					
SUBMITTED.....		(20)	(19)	(20)	(20)
NO VISIBLE LESIONS.....		20	18	20	20
SEMINAL VESICLES W/COAGULATING GLANDS;					
SUBMITTED.....		(20)	(19)	(20)	(20)
NO VISIBLE LESIONS.....		20	19	20	20

TABLE 0-23
 F₀: SUMMARY OF MALE ABSOLUTE ORGAN WEIGHTS (G) AND ORGAN-TO-BODY WEIGHT RATIOS
 TWO GENERATION REPRODUCTION TOXICITY STUDY OF PROPYLTHIOURACIL WHEN ADMINISTERED TO SPRAGUE-DAWLEY RATS IN THE DRINKING WATER

ORGAN	DOSE GROUPS				TREND ^a
	0.0000 %	0.0001 %	0.0004 %	0.0015 %	
ORGAN WEIGHTS (g) ^b					
BODY	628.9 ± 17.82 (20)	643.2 ± 13.19 (19)	608.6 ± 14.59 (20)	*511.7 ± 13.16 (20)	P<0.001
ADRENALS	0.0590 ± 0.00228 (20)	0.0531 ± 0.00240 (19)	0.0576 ± 0.00222 (20)	0.0509 ± 0.00266 (20)	P=0.111
BRAIN	2.123 ± 0.0224 (20)	2.068 ± 0.0237 (19)	2.117 ± 0.0219 (20)	2.070 ± 0.0263 (20)	P=0.326
KIDNEYS	3.9069 ± 0.06175 (20)	4.0242 ± 0.06516 (19)	4.0177 ± 0.06020 (20)	3.7460 ± 0.07223 (20)	P=0.413
LIVER	19.9736 ± 0.35780 (20)	22.6335 ± 0.37756 (19)	20.3357 ± 0.34881 (20)	21.2523 ± 0.41850 (20)	P=0.069
PITUITARY	0.0123 ± 0.00165 (20)	0.0126 ± 0.00174 (19)	0.0161 ± 0.00161 (20)	0.0123 ± 0.00193 (20)	P=0.495
DORSOLATERAL PROSTATE	0.7294 ± 0.03209 (20)	0.6560 ± 0.03386 (19)	0.6424 ± 0.03128 (20)	0.6329 ± 0.03753 (20)	P=0.041
VENTRAL PROSTATE	0.6876 ± 0.05257 (20)	0.6777 ± 0.05548 (19)	0.6781 ± 0.05125 (20)	0.6947 ± 0.06149 (20)	P=0.979
RIGHT CAUDA EPIDIDYMIS	0.3060 ± 0.01021 (20)	0.2979 ± 0.01078 (19)	0.3256 ± 0.09951 (20)	0.2931 ± 0.01194 (20)	P=0.949
RIGHT EPIDIDYMIS	0.7562 ± 0.02063 (20)	0.7202 ± 0.02177 (19)	0.7896 ± 0.02011 (20)	0.7197 ± 0.02413 (20)	P=0.865
RIGHT TESTIS	1.7087 ± 0.03673 (20)	1.7520 ± 0.03876 (19)	1.7110 ± 0.03581 (20)	1.7843 ± 0.04296 (20)	P=0.375
SEMINAL VESICLES	2.3010 ± 0.09383 (20)	2.2072 ± 0.09901 (19)	2.5045 ± 0.09147 (20)	2.1866 ± 0.10975 (20)	P=0.859
SPLEEN	0.818 ± 0.0254 (20)	0.888 ± 0.0268 (19)	0.845 ± 0.0248 (20)	0.726 ± 0.0298 (20)	P=0.124
THYROID	0.0323 ± 0.00474 (20)	0.0294 ± 0.00500 (19)	*0.0539 ± 0.00462 (20)	*0.1295 ± 0.00554 (20)	P<0.001
ORGAN-TO-BODY WEIGHT RATIOS ^c					
ADRENALS	0.0971 ± 0.00400 (20)	0.0853 ± 0.00343 (19)	0.0957 ± 0.00364 (20)	0.0946 ± 0.00566 (20)	P=0.779
BRAIN	3.452 ± 0.0905 (20)	3.280 ± 0.0678 (19)	3.521 ± 0.0765 (20)	*3.984 ± 0.1043 (20)	P<0.001
KIDNEYS	6.4801 ± 0.06584 (20)	6.6240 ± 0.11151 (19)	6.7313 ± 0.13322 (20)	6.4730 ± 0.10702 (20)	P=0.909
LIVER	33.7161 ± 0.76665 (20)	35.0990 ± 0.53426 (19)	34.1839 ± 0.59443 (20)	34.0647 ± 0.66415 (20)	P=0.726
PITUITARY	0.0196 ± 0.00064 (20)	0.0195 ± 0.00091 (19)	*0.0270 ± 0.00542 (20)	*0.0247 ± 0.00094 (20)	P<0.001
DORSOLATERAL PROSTATE	1.1656 ± 0.04758 (20)	1.0228 ± 0.05987 (19)	1.0681 ± 0.06376 (20)	1.2794 ± 0.06814 (20)	P=0.285
VENTRAL PROSTATE	1.1137 ± 0.08972 (20)	1.0625 ± 0.08258 (19)	1.1270 ± 0.06223 (20)	1.3617 ± 0.12883 (20)	P=0.049
RIGHT CAUDA EPIDIDYMIS	0.4998 ± 0.01831 (20)	0.4754 ± 0.01530 (19)	0.5421 ± 0.02030 (20)	0.5546 ± 0.02593 (20)	P=0.010
RIGHT EPIDIDYMIS	1.2280 ± 0.03113 (20)	1.1457 ± 0.03450 (19)	1.3137 ± 0.04020 (20)	1.3775 ± 0.06483 (20)	P=0.029
RIGHT TESTIS	2.7829 ± 0.09072 (20)	2.7819 ± 0.07505 (19)	2.8413 ± 0.07289 (20)	*3.4314 ± 0.10726 (20)	P<0.001
SEMINAL VESICLES	3.7062 ± 0.13912 (20)	3.4884 ± 0.18279 (19)	4.1495 ± 0.21360 (20)	*4.3077 ± 0.18792 (20)	P=0.009
SPLEEN	1.361 ± 0.0498 (20)	1.459 ± 0.0347 (19)	1.416 ± 0.0373 (20)	1.229 ± 0.0418 (20)	P=0.041
THYROID	0.0518 ± 0.00281 (20)	0.0455 ± 0.00330 (19)	*0.0884 ± 0.00561 (20)	*0.2601 ± 0.01930 (20)	P<0.001

a Each dose group is compared with the control group by Shirley's test if P<0.01 from Jonckheere's trend test, otherwise Dunn's test is applied (*= P<0.05).

b Mean organ weight ± standard error (number of animals).

c Mean ratio (mg/g body weight) ± standard error (number of animals).

Note: Absolute organ weights were covaried by body weight.

TABLE 0-24A
F₀: SUMMARY OF SPERM ANALYSIS DATA
TWO GENERATION REPRODUCTION TOXICITY STUDY OF PROPYLTHIOURACIL WHEN ADMINISTERED TO SPRAGUE-DAWLEY RATS IN THE DRINKING WATER

ENDPOINT	DOSE GROUPS ^a				TREND ^b
	0.0000 %	0.0001 %	0.0004 %	0.0015 %	
EPIDIDYMAL SPERM DENSITY ^c	423.64 ± 16.385 (20)	390.00 ± 21.443 (19)	398.55 ± 12.948 (20)	375.07 ± 28.224 (20)	P=0.349
SPERM MORPHOLOGY (% ABNORMAL) ^d	0.3 ± 0.11 (20)	*1.0 ± 0.18 (19)	0.4 ± 0.10 (20)	0.7 ± 0.17 (18)	P=0.354
SPERMATIDS/MG TESTIS ^e	63.36 ± 2.670 (20)	57.87 ± 2.184 (19)	64.43 ± 3.065 (20)	59.10 ± 2.828 (20)	P=0.563
TOTAL SPERMATIDS/TESTIS ^f	101.31 ± 3.445 (20)	95.33 ± 3.513 (19)	101.47 ± 4.143 (20)	94.25 ± 4.553 (20)	P=0.324
TOTAL SPERMAT/CAUDA ^f	130.03 ± 4.474 (20)	117.63 ± 6.469 (19)	129.25 ± 4.419 (20)	111.10 ± 9.620 (20)	P=0.170

a Endpoint mean ± standard error (number of animals).

b Each dose group is compared with the control group by Shirley's test if P<0.01 from Jonckheere's trend test, otherwise Dunn's test is applied (* = P<0.05).

c Sperm density expressed as 1000 sperm per mg right caudal tissue.

d Dose group means and standard errors are computed only from samples with at least 200 sperm.

e Spermatid heads per mg testis (x 1,000) = (total number of spermatid heads per right testis) / (right testis weight in mg)

f Total number of spermatid heads or sperm (x 1,000,000) = (average number of spermatid heads or sperm x 2.5 x 100) / 0.0001

TABLE 0-24B
F₀: SUMMARY OF HAMILTON-THORNE SPERM ANALYSIS DATA
TWO GENERATION REPRODUCTION TOXICITY STUDY OF PROPYLTHIOURACIL WHEN ADMINISTERED TO SPRAGUE-DAWLEY RATS IN THE DRINKING WATER

ENDPOINT	DOSE GROUPS ^a				TREND ^b
	0.0000 %	0.0001 %	0.0004 %	0.0015 %	
AMPLITUDE	18.6 ± 0.38 (20)	18.1 ± 0.35 (19)	18.1 ± 0.37 (20)	18.6 ± 0.47 (18)	P=0.920
BEAT/CROSS FREQUENCY (hz/sec)	16.1 ± 0.22 (20)	16.2 ± 0.36 (19)	16.1 ± 0.32 (20)	16.4 ± 0.43 (18)	P=0.792
LINEARITY	37.4 ± 0.48 (20)	37.7 ± 0.55 (19)	37.9 ± 0.42 (20)	37.8 ± 0.61 (18)	P=0.649
MOTILE PERCENTAGE	93.0 ± 1.62 (20)	93.6 ± 0.62 (19)	94.8 ± 0.49 (20)	88.3 ± 4.46 (18)	P=0.246
PROGRESSIVE VELOCITY	103.6 ± 2.03 (20)	102.2 ± 1.68 (19)	105.2 ± 2.79 (20)	104.7 ± 2.82 (18)	P=0.392
PATH VELOCITY	163.1 ± 3.12 (20)	159.7 ± 2.30 (19)	163.4 ± 3.68 (20)	163.5 ± 4.68 (18)	P=0.630
STRAIGHTNESS	63.1 ± 0.80 (20)	63.8 ± 0.83 (19)	63.5 ± 0.63 (20)	63.6 ± 0.75 (18)	P=0.844
TRACK SPEED	284.5 ± 5.27 (20)	279.7 ± 4.36 (19)	283.2 ± 6.11 (20)	286.6 ± 8.14 (18)	P=0.551

a Endpoint mean ± standard error (number of animals).

b Each dose group is compared with the control group by Shirley's test if P<0.01 from Jonckheere's trend test, otherwise Dunn's test is applied (* = P<0.05).

TABLE 0-25
 F₀: SUMMARY OF FEMALE GROSS OBSERVATIONS
 TWO GENERATION REPRODUCTION TOXICITY STUDY OF PROPYLTHIOURACIL WHEN ADMINISTERED TO SPRAGUE-DAWLEY RATS IN THE DRINKING WATER

	GROUP: DOSE (%):	1 0.0000	2 0.0001	3 0.0004	4 0.0015
	NUMBER OF ANIMALS ON STUDY :	20	20	20	20
	NUMBER OF ANIMALS COMPLETED:	(20)	(20)	(20)	(20)
THYROID/PARATHYROID;					
SUBMITTED.....		(20)	(20)	(20)	(18)
NO VISIBLE LESIONS.....		19	20	19	0
DISCOLORATION; TAN; BILATERAL; FOCAL		1	0	0	0
ENLARGEMENT; BILATERAL		0	0	1	18
ADRENAL GLAND(S);					
SUBMITTED.....		(20)	(20)	(20)	(18)
NO VISIBLE LESIONS.....		20	20	20	18
BRAIN;					
SUBMITTED.....		(20)	(20)	(20)	(18)
NO VISIBLE LESIONS.....		20	20	20	18
KIDNEY(S);					
SUBMITTED.....		(20)	(20)	(20)	(18)
NO VISIBLE LESIONS.....		19	20	20	18
ENLARGEMENT; RIGHT		1	0	0	0
LIVER;					
SUBMITTED.....		(20)	(20)	(20)	(18)
NO VISIBLE LESIONS.....		20	20	20	18
MAMMARY GLAND(S);					
SUBMITTED.....		(0)	(0)	(1)	(1)
NO VISIBLE LESIONS.....		0	0	0	0
DISCOLORATION; YELLOW; RIGHT; FOCAL		0	0	0	1
ENLARGEMENT; RIGHT		0	0	1	0
ENLARGEMENT; RIGHT; FOCAL		0	0	0	1
OVARY(IES);					
SUBMITTED.....		(20)	(20)	(20)	(18)
NO VISIBLE LESIONS.....		20	20	20	18

TABLE 0-25 (CONTINUED)
 F₀: SUMMARY OF FEMALE GROSS OBSERVATIONS
 TWO GENERATION REPRODUCTION TOXICITY STUDY OF PROPYLTHIOURACIL WHEN ADMINISTERED TO SPRAGUE-DAWLEY RATS IN THE DRINKING WATER

	GROUP:	1	2	3	4
	DOSE (%):	0.0000	0.0001	0.0004	0.0015
PITUITARY GLAND;					
SUBMITTED.....		(20)	(20)	(20)	(18)
NO VISIBLE LESIONS.....		20	19	20	18
ENLARGEMENT		0	1	0	0
SPLEEN;					
SUBMITTED.....		(20)	(20)	(20)	(18)
NO VISIBLE LESIONS.....		20	20	20	18
URINARY BLADDER;					
SUBMITTED.....		(1)	(0)	(0)	(0)
NO VISIBLE LESIONS.....		0	0	0	0
ENLARGEMENT		1	0	0	0
VAGINA/CERVIX/UTERUS;					
SUBMITTED.....		(20)	(20)	(20)	(18)
NO VISIBLE LESIONS.....		19	19	20	18
DISTENSION; RIGHT		0	1	0	0
ENLARGEMENT; RIGHT		1	0	0	0
FLUID-FILLED		0	1	0	0

TABLE 0-26
 F₀: SUMMARY OF FEMALE ABSOLUTE ORGAN WEIGHTS (G) AND ORGAN-TO-BODY WEIGHT RATIOS
 TWO GENERATION REPRODUCTION TOXICITY STUDY OF PROPYLTHIOURACIL WHEN ADMINISTERED TO SPRAGUE-DAWLEY RATS IN THE DRINKING WATER

ORGAN	DOSE GROUPS				TREND ^a
	0.0000 %	0.0001 %	0.0004 %	0.0015 %	
<u>ORGAN WEIGHTS (g)^b</u>					
BODY	349.3 ± 10.09 (20)	338.9 ± 6.56 (20)	333.6 ± 7.38 (20)	*303.5 ± 6.93 (18)	P<0.001
ADRENALS	0.0754 ± 0.00399 (20)	0.0828 ± 0.00388 (20)	0.0716 ± 0.00386 (20)	*0.0586 ± 0.00439 (18)	P=0.005
BRAIN	1.963 ± 0.0236 (20)	1.934 ± 0.0230 (20)	1.901 ± 0.0229 (20)	*1.842 ± 0.0260 (18)	P=0.001
KIDNEYS	2.4098 ± 0.06308 (20)	2.3157 ± 0.06143 (20)	2.3415 ± 0.06112 (20)	*2.1244 ± 0.06945 (18)	P=0.013
LIVER	12.7812 ± 0.30173 (20)	12.9263 ± 0.29381 (20)	12.4662 ± 0.29236 (20)	*11.5513 ± 0.33217 (18)	P=0.010
OVARIES	0.1311 ± 0.00716 (20)	0.1354 ± 0.00697 (20)	0.1348 ± 0.00694 (20)	*0.1173 ± 0.00788 (18)	P=0.300
PITUITARY	0.0170 ± 0.00079 (20)	0.0174 ± 0.00077 (20)	0.0156 ± 0.00077 (20)	0.0165 ± 0.00087 (18)	P=0.315
SPLEEN	0.609 ± 0.0188 (20)	0.567 ± 0.0183 (20)	0.599 ± 0.0182 (20)	*0.520 ± 0.0207 (18)	P=0.024
THYROID	0.0245 ± 0.00566 (20)	0.0281 ± 0.00551 (20)	*0.0319 ± 0.00548 (20)	*0.1159 ± 0.00623 (18)	P<0.001
UTERUS	1.045 ± 0.0796 (20)	0.994 ± 0.0775 (20)	1.097 ± 0.0772 (20)	0.937 ± 0.0877 (18)	P=0.661
<u>ORGAN-TO-BODY WEIGHT RATIOS^c</u>					
ADRENALS	0.2199 ± 0.00982 (20)	0.2462 ± 0.01801 (20)	0.2163 ± 0.00867 (20)	0.1936 ± 0.01161 (18)	P=0.028
BRAIN	5.717 ± 0.2072 (20)	5.731 ± 0.0873 (20)	5.754 ± 0.1565 (20)	6.136 ± 0.1549 (18)	P=0.023
KIDNEYS	7.1608 ± 0.34801 (20)	6.9189 ± 0.13672 (20)	7.0396 ± 0.14600 (20)	6.7208 ± 0.17960 (18)	P=0.288
LIVER	38.1986 ± 0.95614 (20)	38.8161 ± 0.80916 (20)	37.5723 ± 0.65854 (20)	35.1795 ± 1.04268 (18)	P=0.016
OVARIES	0.3866 ± 0.01899 (20)	0.4044 ± 0.02178 (20)	0.4067 ± 0.02374 (20)	0.3696 ± 0.01943 (18)	P=0.668
PITUITARY	0.0502 ± 0.00219 (20)	0.0522 ± 0.00260 (20)	0.0470 ± 0.00248 (20)	0.0523 ± 0.00230 (18)	P=0.755
SPLEEN	1.805 ± 0.0715 (20)	1.688 ± 0.0550 (20)	1.805 ± 0.0438 (20)	1.655 ± 0.0730 (18)	P=0.401
THYROID	0.0710 ± 0.00288 (20)	*0.0837 ± 0.00397 (20)	*0.0958 ± 0.00474 (20)	*0.3858 ± 0.04129 (18)	P<0.001
UTERUS	3.048 ± 0.2530 (20)	2.932 ± 0.1577 (20)	3.313 ± 0.3827 (20)	3.164 ± 0.1565 (18)	P=0.172

a Each dose group is compared with the control group by Shirley's test if P<0.01 from Jonckheere's trend test, otherwise Dunn's test is applied (*= P<0.05).

b Mean organ weight ± standard error (number of animals).

c Mean ratio (mg/g body weight) ± standard error (number of animals).

Note: Absolute organ weights were covaried by body weight.

TABLE 0-27
 F₀: SUMMARY OF MORTALITY DATA
 TWO GENERATION REPRODUCTION TOXICITY STUDY OF PROPYLTHIURACIL WHEN ADMINISTERED TO SPRAGUE-DAWLEY RATS IN THE DRINKING WATER

	DOSE GROUPS ^a			
	0.0000 %	0.0001 %	0.0004 %	0.0015 %
<u>MALE</u>				
FOUND DEAD	0/20	0/20	0/20	0/20
MORIBUND KILLS	0/20	1/20	0/20	0/20
REMOVED FROM STUDY	0/20	0/20	0/20	0/20
<u>FEMALE</u>				
FOUND DEAD	0/20	0/20	0/20	2/20
MORIBUND KILLS	0/20	0/20	0/20	0/20
REMOVED FROM STUDY	0/20	0/20	0/20	0/20

a Number of animals found dead, moribund, or removed / total number of animals in group.

TABLE 1A-1
 F₁: SUMMARY OF ADOLESCENT BODY WEIGHTS (GRAMS)
 TWO GENERATION REPRODUCTION TOXICITY STUDY OF PROPYLTHIOURACIL WHEN ADMINISTERED TO SPRAGUE-DAWLEY RATS IN THE DRINKING WATER

POSTNATAL DAY	DOSE GROUP ^a			TREND ^b
	0.0000 %	0.0001 %	0.0004 %	
<u>MALE</u>				
32	120.4 ± 3.42 (28)	113.3 ± 3.37 (30)	112.3 ± 5.29 (31)	P=0.210
39	182.4 ± 4.31 (28)	172.5 ± 4.32 (30)	169.7 ± 7.13 (31)	P=0.196
46	247.6 ± 5.52 (28)	234.8 ± 5.71 (30)	229.9 ± 8.83 (31)	P=0.158
53	312.3 ± 6.13 (28)	295.8 ± 7.27 (30)	291.6 ± 10.18 (31)	P=0.135
60	366.6 ± 6.44 (28)	354.9 ± 7.27 (30)	347.1 ± 11.46 (31)	P=0.224
67	405.7 ± 7.38 (28)	399.7 ± 7.73 (30)	390.9 ± 12.44 (31)	P=0.431
74	448.8 ± 7.39 (28)	438.8 ± 8.57 (30)	428.1 ± 13.55 (31)	P=0.280
81	482.2 ± 8.05 (28)	469.0 ± 8.65 (30)	461.9 ± 14.53 (31)	P=0.309
88	505.6 ± 8.76 (28)	493.9 ± 9.09 (30)	485.3 ± 15.35 (31)	P=0.298
95	528.9 ± 9.06 (28)	516.2 ± 9.35 (30)	507.4 ± 15.86 (31)	P=0.293
<u>FEMALE</u>				
32	103.9 ± 2.27 (30)	101.5 ± 2.61 (33)	95.6 ± 2.98 (33)	P=0.082
39	145.2 ± 2.44 (30)	141.3 ± 2.63 (33)	137.5 ± 3.26 (33)	P=0.149
46	175.4 ± 2.87 (30)	171.7 ± 2.83 (33)	167.1 ± 3.12 (33)	P=0.099
53	201.5 ± 3.61 (30)	200.2 ± 3.65 (33)	194.1 ± 3.40 (33)	P=0.179
60	227.5 ± 3.83 (30)	223.6 ± 3.97 (33)	217.2 ± 3.66 (33)	P=0.115
67	245.5 ± 3.95 (30)	244.3 ± 4.39 (33)	236.2 ± 3.94 (33)	P=0.173
74	261.0 ± 4.62 (30)	261.9 ± 4.57 (33)	253.8 ± 4.40 (33)	P=0.374
81	273.0 ± 5.05 (30)	275.8 ± 5.12 (33)	269.7 ± 4.64 (33)	P=0.790
88	279.4 ± 5.53 (30)	280.2 ± 5.11 (33)	273.8 ± 4.58 (33)	P=0.482
95	291.0 ± 5.85 (30)	290.7 ± 5.67 (33)	281.8 ± 4.91 (33)	P=0.313

a Mean ± standard error (number of animals).

b Each dose group is compared to the control with Shirley's test when a trend is present (P<0.01 from Jonckheere's trend test), otherwise Dunn's test is applied (*= P<0.05).

TABLE 1A-2
 F1: SUMMARY OF ADOLESCENT MORTALITY DATA
 TWO GENERATION REPRODUCTION TOXICITY STUDY OF PROPYLTHIURACIL WHEN ADMINISTERED TO SPRAGUE-DAWLEY RATS IN THE DRINKING WATER

	DOSE GROUP ^a			
	0.0000 %	0.0001 %	0.0004 %	0.0015 %
	<u>MALE</u>			
FOUND DEAD	0/28	0/30	0/31	6/29
MORIBUND KILLS	0/28	0/30	0/31	17/29
REMOVED FROM STUDY	0/28	0/30	0/31	6/29
	<u>FEMALE</u>			
FOUND DEAD	0/30	0/33	0/33	8/26
MORIBUND KILLS	0/30	0/33	0/33	12/26
REMOVED FROM STUDY	0/30	0/33	0/33	6/26

a Number of animals found dead, moribund, or removed / total number of animals in group.

TABLE 1-1A
 F₁: SUMMARY OF MALE BODY WEIGHTS (GRAMS)
 TWO GENERATION REPRODUCTION TOXICITY STUDY OF PROPYLTHIOURACIL WHEN ADMINISTERED TO SPRAGUE-DAWLEY RATS IN THE DRINKING WATER

WEEK OF STUDY	DOSE GROUP ^a			TREND ^b
	0.0000 %	0.0001 %	0.0004 %	
1	552.8 ± 10.88 (20)	531.6 ± 12.86 (20)	531.4 ± 16.64 (20)	P=0.191
2	567.4 ± 10.66 (20)	547.0 ± 13.45 (20)	547.6 ± 16.60 (20)	P=0.249
3	587.0 ± 10.82 (20)	564.8 ± 14.29 (20)	564.5 ± 17.33 (20)	P=0.186
4	605.5 ± 11.00 (20)	581.6 ± 15.26 (20)	585.0 ± 17.43 (20)	P=0.212
5	619.1 ± 11.42 (20)	591.9 ± 15.25 (20)	597.6 ± 17.60 (20)	P=0.184
6	633.3 ± 11.70 (20)	607.3 ± 16.10 (20)	613.8 ± 18.18 (20)	P=0.255
7	647.7 ± 12.39 (20)	621.7 ± 16.07 (20)	629.6 ± 19.42 (20)	P=0.306
8	662.5 ± 13.36 (20)	635.2 ± 16.38 (20)	643.4 ± 19.99 (20)	P=0.309
9	672.0 ± 13.44 (20)	646.6 ± 16.90 (20)	652.4 ± 20.60 (20)	P=0.296
10	681.1 ± 13.49 (20)	659.6 ± 17.26 (20)	663.5 ± 21.41 (20)	P=0.303
11	689.9 ± 14.38 (19)	669.6 ± 17.78 (20)	671.1 ± 21.65 (20)	P=0.310
12	708.7 ± 26.52 (9)	620.0 ± 19.93 (10)	685.1 ± 33.63 (10)	P=0.401

a Mean ± standard error (number of animals).

b Each dose group is compared to the control with Shirley's test when a trend is present (P<0.01 from Jonckheere's trend test), otherwise Dunn's test is applied (*= P<0.05).

TABLE 1-1B
F₁: SUMMARY OF FEMALE BODY WEIGHTS (GRAMS)
TWO GENERATION REPRODUCTION TOXICITY STUDY OF PROPYLTHIOURACIL WHEN ADMINISTERED TO SPRAGUE-DAWLEY RATS IN THE DRINKING WATER

WEEK OF STUDY	DOSE GROUP ^a			TREND ^b
	0.0000 %	0.0001 %	0.0004 %	
1	295.9 ± 7.90 (20)	295.3 ± 7.71 (20)	283.0 ± 6.97 (20)	P=0.332
2	330.7 ± 7.21 (20)	322.2 ± 7.44 (20)	311.0 ± 6.84 (20)	P=0.068
3	359.9 ± 7.56 (20)	348.7 ± 7.53 (20)	335.5 ± 8.19 (20)	P=0.042
4	407.6 ± 10.54 (20)	394.0 ± 10.16 (20)	375.6 ± 12.75 (20)	P=0.139
7	353.3 ± 7.23 (20)	352.0 ± 8.00 (20)	336.4 ± 7.33 (20)	P=0.179
8	327.9 ± 8.82 (20)	329.3 ± 9.15 (20)	317.9 ± 7.15 (20)	P=0.592
9	341.8 ± 7.42 (20)	341.5 ± 8.27 (20)	324.8 ± 5.88 (20)	P=0.095
10	355.8 ± 7.71 (20)	345.5 ± 7.50 (20)	334.9 ± 6.56 (20)	P=0.065
11	363.2 ± 8.28 (20)	350.9 ± 7.91 (20)	340.3 ± 7.43 (20)	P=0.055
12	366.1 ± 13.76 (10)	364.2 ± 11.50 (10)	355.4 ± 8.07 (10)	P=0.776

a Mean ± standard error (number of animals).

b Each dose group is compared to the control with Shirley's test when a trend is present (P<0.01 from Jonckheere's trend test), otherwise Dunn's test is applied (*= P<0.05).

TABLE 1-2A
 F₁: SUMMARY OF DAILY FEED CONSUMPTION (G/KG BODY WEIGHT/DAY)
 TWO GENERATION REPRODUCTION TOXICITY STUDY OF PROPYLTHIOURACIL WHEN ADMINISTERED TO SPRAGUE-DAWLEY RATS IN THE DRINKING WATER

WEEK OF STUDY	DOSE GROUP ^a			TREND ^b
	0.0000 %	0.0001 %	0.0004 %	
<u>MALE</u>				
3	55.6 ± 0.95 (20)	59.2 ± 2.24 (20)	56.1 ± 0.95 (20)	P=0.724
4	51.7 ± 0.79 (20)	49.7 ± 1.01 (20)	50.8 ± 0.98 (20)	P=0.306
5	50.9 ± 0.80 (20)	51.6 ± 0.68 (20)	51.9 ± 0.93 (20)	P=0.578
6	49.6 ± 0.71 (20)	49.9 ± 0.64 (20)	49.7 ± 0.51 (20)	P=0.704
7	51.3 ± 0.80 (20)	51.4 ± 0.71 (19)	50.1 ± 0.83 (20)	P=0.206
8	49.3 ± 1.31 (19)	49.1 ± 0.71 (20)	48.6 ± 1.01 (20)	P=0.626
9	48.0 ± 0.92 (20)	48.1 ± 0.64 (20)	49.6 ± 2.52 (20)	P=0.139
10	46.2 ± 0.77 (19)	46.5 ± 0.85 (20)	46.3 ± 0.62 (20)	P=0.540
<u>FEMALE</u>				
3	76.1 ± 2.23 (20)	76.2 ± 2.27 (20)	75.5 ± 2.28 (20)	P=0.903
10	68.0 ± 1.76 (20)	64.8 ± 1.41 (20)	65.1 ± 1.35 (20)	P=0.150

a Mean ± standard error (number of animals).

b Each dose group is compared to the control with Shirley's test when a trend is present (P<0.01 from Jonckheere's trend test), otherwise Dunn's test is applied (*= P<0.05).

TABLE 1-2B
 F₁: SUMMARY OF DAILY FEED CONSUMPTION (G/ANIMAL/DAY)
 TWO GENERATION REPRODUCTION TOXICITY STUDY OF PROPYLTHIOURACIL WHEN ADMINISTERED TO SPRAGUE-DAWLEY RATS IN THE DRINKING WATER

WEEK OF STUDY	DOSE GROUP ^a			TREND ^b
	0.0000 %	0.0001 %	0.0004 %	
	<u>MALE</u>			
3	32.5 ± 0.56 (20)	33.2 ± 1.05 (20)	31.5 ± 0.76 (20)	P=0.220
4	31.2 ± 0.57 (20)	*28.8 ± 0.65 (20)	29.5 ± 0.75 (20)	P=0.067
5	31.5 ± 0.59 (20)	30.4 ± 0.70 (20)	30.8 ± 0.75 (20)	P=0.356
6	31.4 ± 0.62 (20)	30.2 ± 0.62 (20)	30.4 ± 0.89 (20)	P=0.166
7	33.2 ± 0.73 (20)	31.7 ± 0.73 (19)	31.4 ± 0.86 (20)	P=0.071
8	32.5 ± 0.98 (19)	31.1 ± 0.73 (20)	31.1 ± 0.95 (20)	P=0.330
9	32.2 ± 0.75 (20)	31.0 ± 0.76 (20)	31.9 ± 1.09 (20)	P=0.447
10	31.4 ± 0.74 (19)	30.6 ± 0.79 (20)	30.5 ± 0.78 (20)	P=0.404
	<u>FEMALE</u>			
3	27.3 ± 0.84 (20)	26.5 ± 0.77 (20)	25.4 ± 1.04 (20)	P=0.147
10	24.1 ± 0.72 (20)	22.3 ± 0.44 (20)	21.8 ± 0.70 (20)	P=0.028

a Mean ± standard error (number of animals).

b Each dose group is compared to the control with Shirley's test when a trend is present (P<0.01 from Jonckheere's trend test), otherwise Dunn's test is applied (*= P<0.05).

TABLE 1-3A
 F₁: SUMMARY OF DAILY WATER CONSUMPTION (G/KG BODY WEIGHT/DAY)
 TWO GENERATION REPRODUCTION TOXICITY STUDY OF PROPYLTHIURACIL WHEN ADMINISTERED TO SPRAGUE-DAWLEY RATS IN THE DRINKING WATER

WEEK OF STUDY	DOSE GROUP ^a			TREND ^b
	0.0000 %	0.0001 %	0.0004 %	
<u>MALE</u>				
3	64.7 ± 1.91 (20)	66.1 ± 2.96 (20)	66.2 ± 2.34 (20)	P=0.735
4	63.4 ± 2.05 (20)	65.5 ± 2.87 (20)	67.3 ± 3.81 (20)	P=0.431
5	60.0 ± 2.20 (20)	64.5 ± 3.28 (20)	62.6 ± 2.90 (20)	P=0.533
6	59.5 ± 2.16 (20)	60.6 ± 3.04 (20)	59.3 ± 2.06 (20)	P=1.000
7	57.8 ± 1.78 (20)	59.7 ± 3.29 (19)	57.6 ± 2.19 (20)	P=0.900
8	54.1 ± 1.96 (20)	58.4 ± 2.77 (19)	54.7 ± 2.38 (20)	P=0.878
9	50.8 ± 1.72 (20)	53.5 ± 2.18 (20)	53.1 ± 2.89 (20)	P=0.674
10	50.5 ± 1.49 (19)	53.7 ± 2.16 (20)	52.8 ± 2.23 (20)	P=0.461
<u>FEMALE</u>				
3	107.1 ± 10.20 (20)	122.5 ± 5.13 (20)	121.3 ± 6.61 (20)	P=0.342
10	88.2 ± 3.38 (20)	87.9 ± 4.11 (20)	89.1 ± 3.25 (20)	P=0.946

a Mean ± standard error (number of animals).

b Each dose group is compared to the control with Shirley's test when a trend is present (P<0.01 from Jonckheere's trend test), otherwise Dunn's test is applied (*= P<0.05).

TABLE 1-3B
 F₁: SUMMARY OF DAILY WATER CONSUMPTION (G/ANIMAL/DAY)
 TWO GENERATION REPRODUCTION TOXICITY STUDY OF PROPYLTHIOURACIL WHEN ADMINISTERED TO SPRAGUE-DAWLEY RATS IN THE DRINKING WATER

WEEK OF STUDY	DOSE GROUP ^a			TREND ^b
	0.0000 %	0.0001 %	0.0004 %	
	<u>MALE</u>			
3	37.9 ± 1.12 (20)	37.3 ± 1.76 (20)	36.9 ± 1.08 (20)	P=0.729
4	38.2 ± 1.02 (20)	37.9 ± 1.67 (20)	38.6 ± 1.51 (20)	P=0.802
5	36.9 ± 1.12 (20)	38.0 ± 1.91 (20)	36.9 ± 1.24 (20)	P=0.729
6	37.4 ± 1.06 (20)	36.5 ± 1.74 (20)	36.0 ± 0.93 (20)	P=0.424
7	37.2 ± 1.01 (20)	37.1 ± 1.93 (19)	35.7 ± 0.93 (20)	P=0.388
8	35.6 ± 1.07 (20)	37.3 ± 1.73 (19)	34.6 ± 0.84 (20)	P=0.536
9	34.0 ± 1.06 (20)	34.5 ± 1.57 (20)	33.9 ± 1.10 (20)	P=0.679
10	34.2 ± 0.97 (19)	35.4 ± 1.58 (20)	34.4 ± 0.96 (20)	P=0.911
	<u>FEMALE</u>			
3	38.2 ± 3.55 (20)	42.3 ± 1.50 (20)	40.7 ± 2.34 (20)	P=0.914
10	31.3 ± 1.33 (20)	30.1 ± 1.30 (20)	29.6 ± 0.99 (20)	P=0.272

a Mean ± standard error (number of animals).

b Each dose group is compared to the control with Shirley's test when a trend is present (P<0.01 from Jonckheere's trend test), otherwise Dunn's test is applied (*= P<0.05).

TABLE 1-4
F₁: SUMMARY OF ESTIMATED DOSAGE (MG/KG BODY WEIGHT/DAY)
TWO GENERATION REPRODUCTION TOXICITY STUDY OF PROPYLTHIOURACIL WHEN ADMINISTERED TO SPRAGUE-DAWLEY RATS IN THE DRINKING WATER

WEEK OF STUDY	DOSE GROUP ^a	
	0.0001 %	0.0004 %
	<u>MALE</u>	
3	0.1 ± 0.00 (20)	0.3 ± 0.01 (20)
4	0.1 ± 0.00 (20)	0.3 ± 0.02 (20)
5	0.1 ± 0.00 (20)	0.3 ± 0.01 (20)
6	0.1 ± 0.00 (20)	0.2 ± 0.01 (20)
7	0.1 ± 0.00 (19)	0.2 ± 0.01 (20)
8	0.1 ± 0.00 (19)	0.2 ± 0.01 (20)
9	0.1 ± 0.00 (20)	0.2 ± 0.01 (20)
10	0.1 ± 0.00 (20)	0.2 ± 0.01 (20)
	<u>FEMALE</u>	
3	0.1 ± 0.01 (20)	0.5 ± 0.03 (20)
10	0.1 ± 0.00 (20)	0.4 ± 0.01 (20)

a Mean ± standard error (number of animals).

TABLE 1-5
F₁: SUMMARY OF REPRODUCTIVE PERFORMANCE OF COHABITED PAIRS
TWO GENERATION REPRODUCTION TOXICITY STUDY OF PROPYLTHIOURACIL WHEN ADMINISTERED TO SPRAGUE-DAWLEY RATS IN THE DRINKING WATER

REPRODUCTIVE PARAMETER	DOSE GROUP ^a						TREND
	0.0000 %		0.0001 %		0.0004 %		
Pregnancy Index ^{b,e,g}	15/20	(75)	15/20	(75)	14/20	(70)	P=0.500
Live Pups Per Litter ^{c,h,i}							
Male	7.1 ± 0.51	(15)	6.1 ± 0.77	(15)	7.0 ± 0.66	(14)	P=0.803
Female	8.1 ± 0.51	(15)	6.4 ± 0.61	(15)	6.9 ± 0.75	(14)	P=0.154
Combined	15.2 ± 0.63	(15)	12.5 ± 1.06	(15)	13.9 ± 0.61	(14)	P=0.100
Proportion Pups Born Alive ^{c,h,i}	1.00 ± 0.003	(15)	0.98 ± 0.012	(15)	0.98 ± 0.012	(14)	P=0.236
Sex of Pups Born Alive (Males/Total) ^{c,j,i}	0.47 ± 0.028	(15)	0.48 ± 0.040	(15)	0.51 ± 0.048	(14)	P=0.412
Live Pup Weight (grams) ^{c,j,i}							
Male	6.83 ± 0.137	(14)	6.74 ± 0.191	(15)	6.93 ± 0.157	(13)	P=0.412
Female	6.34 ± 0.123	(14)	6.41 ± 0.189	(15)	6.55 ± 0.135	(13)	P=0.166
Combined	6.57 ± 0.139	(14)	6.52 ± 0.163	(15)	6.75 ± 0.142	(13)	P=0.128
Adjusted Live Pup Weight (grams) ^{c,k,l}							OVERALL
Male	6.94 ± 0.147	(14)	6.63 ± 0.142	(15)	6.94 ± 0.149	(13)	P=0.240
Female	6.43 ± 0.144	(14)	6.32 ± 0.140	(15)	6.55 ± 0.147	(13)	P=0.520
Combined	6.67 ± 0.135	(14)	6.42 ± 0.131	(15)	6.76 ± 0.137	(13)	P=0.191
Average Dam Weight (grams) ^{c,m,i}	356.9 ± 9.64	(15)	352.2 ± 11.10	(15)	342.3 ± 10.34	(14)	P=0.598
Average Sire Weight (grams) ^{n,i}	614.6 ± 10.52	(15)	579.4 ± 16.54	(15)	586.8 ± 23.39	(11)	P=0.235
Average Days to Litter ^{c,m,i}	25.1 ± 0.62	(15)	25.0 ± 0.83	(15)	25.3 ± 0.37	(14)	P=0.298

a Statistical significance for comparisons of dosed groups to controls: * = P<0.05.

b Only those pairs surviving cohabitation were included for statistical analysis of data.

c Only those females surviving to the end of the parturition period were included for statistical analysis of the data.

d Number of females with vaginal sperm / number of cohabiting pairs (percent with vaginal sperm).

e Number of fertile pairs / number of cohabiting pairs (percent pregnant).

f Number of fertile pairs / number of females with vaginal sperm (percent fertile).

g P-value from Cochran-Armitage trend test. Dosed groups compared to control group by Chi-squared test. If fertility index > 1, then it is set equal to 1 for statistical analysis.

h Mean ± standard error (number of pregnant pairs).

i Each dose group is compared to the control group by Shirley's test if P<0.01 from Jonckheere's trend test. Otherwise, Dunn's test is applied.

j Mean ± standard error (number of pregnant pairs producing live pups).

k Least squares estimate of mean pup weight adjusted for average litter size ± standard error (number of pregnant pairs producing live pups).

l Overall differences tested with an F-test, pairwise comparisons from Dunnett's test.

m Mean ± standard error (number of dams).

n Mean ± standard error (number of sires).

TABLE 1-6
F₁: SUMMARY OF DAM WEIGHTS (GRAMS) DURING LACTATION OF THE F₂ LITTER
TWO GENERATION REPRODUCTION TOXICITY STUDY OF PROPYLTHIOURACIL WHEN ADMINISTERED TO SPRAGUE-DAWLEY RATS IN THE DRINKING WATER

AGE (DAYS)	DOSE GROUP ^a			TREND ^b
	0.0000 %	0.0001 %	0.0004 %	
1	356.9 ± 9.64 (15)	352.2 ± 11.10 (15)	342.3 ± 10.34 (14)	P=0.598
4	366.4 ± 8.96 (14)	353.7 ± 10.85 (15)	347.5 ± 8.08 (13)	P=0.188
7	363.7 ± 8.82 (10)	358.3 ± 10.65 (15)	359.1 ± 8.13 (14)	P=0.826
14	366.6 ± 7.00 (15)	361.5 ± 9.04 (15)	360.8 ± 6.71 (14)	P=0.880
21	336.4 ± 6.19 (15)	341.2 ± 8.95 (15)	335.5 ± 7.47 (14)	P=0.813

a Mean ± standard error (number of dams).

b Each dose group is compared to the control with Shirley's test when a trend is present (P<0.01 from Jonckheere's trend test), otherwise Dunn's test is applied (* = P<0.05).

TABLE 1-7A
 F₁: SUMMARY OF DAM DAILY FEED CONSUMPTION (G/KG BODY WEIGHT/DAY) DURING LACTATION OF THE F₂ LITTER
 TWO GENERATION REPRODUCTION TOXICITY STUDY OF PROPYLTHIOURACIL WHEN ADMINISTERED TO SPRAGUE-DAWLEY RATS IN THE DRINKING WATER

POSTNATAL DAYS	DOSE GROUP ^a			TREND ^b
	0.0000 %	0.0001 %	0.0004 %	
	<u>FEMALE</u>			
1-4	90.3 ± 6.93 (14)	86.2 ± 5.19 (15)	111.5 ± 17.62 (13)	P=0.791
4-7	149.5 ± 5.77 (10)	135.3 ± 7.31 (15)	141.5 ± 7.90 (13)	P=0.397
11-14	194.3 ± 3.88 (14)	184.2 ± 9.60 (14)	193.5 ± 6.29 (14)	P=0.818
18-21	243.7 ± 7.49 (15)	219.6 ± 12.14 (15)	222.0 ± 8.07 (14)	P=0.172

a Mean ± standard error (number of animals).

b Each dose group is compared to the control with Shirley's test when a trend is present (P<0.01 from Jonckheere's trend test), otherwise Dunn's test is applied (*= P<0.05).

TABLE 1-7B
 F₁: SUMMARY OF DAM DAILY FEED CONSUMPTION (G/ANIMAL/DAY) DURING LACTATION OF THE F₂ LITTER
 TWO GENERATION REPRODUCTION TOXICITY STUDY OF PROPYLTHIOURACIL WHEN ADMINISTERED TO SPRAGUE-DAWLEY RATS IN THE DRINKING WATER

POSTNATAL DAYS	DOSE GROUP ^a			TREND ^b
	0.0000 %	0.0001 %	0.0004 %	
	<u>FEMALE</u>			
1-4	33.1 ± 2.13 (15)	29.9 ± 1.28 (15)	38.5 ± 5.92 (13)	P=0.608
4-7	54.3 ± 1.31 (11)	*47.9 ± 2.29 (15)	50.1 ± 2.36 (13)	P=0.141
7-11	63.7 ± 0.97 (11)	*56.5 ± 2.43 (15)	61.3 ± 1.98 (14)	P=0.440
11-14	71.0 ± 0.87 (14)	65.4 ± 3.10 (14)	69.6 ± 2.11 (14)	P=0.764
14-18	68.1 ± 2.03 (15)	65.7 ± 2.89 (15)	69.1 ± 1.97 (14)	P=0.755
18-21	81.7 ± 2.27 (15)	74.2 ± 3.71 (15)	*74.0 ± 2.28 (14)	P=0.027

a Mean ± standard error (number of animals).

b Each dose group is compared to the control with Shirley's test when a trend is present (P<0.01 from Jonckheere's trend test), otherwise Dunn's test is applied (*= P<0.05).

TABLE 1-8A
 F₁: SUMMARY OF DAM DAILY WATER CONSUMPTION (G/KG BODY WEIGHT/DAY) DURING LACTATION OF THE F₂ LITTER
 TWO GENERATION REPRODUCTION TOXICITY STUDY OF PROPYLTHIOURACIL WHEN ADMINISTERED TO SPRAGUE-DAWLEY RATS IN THE DRINKING WATER

POSTNATAL DAYS	DOSE GROUP ^a			TREND ^b
	0.0000 %	0.0001 %	0.0004 %	
	<u>FEMALE</u>			
1-4	135.8 ± 6.55 (13)	123.3 ± 6.12 (15)	132.9 ± 8.78 (13)	P=0.765
4-7	190.0 ± 8.11 (10)	163.8 ± 10.76 (14)	173.9 ± 8.70 (13)	P=0.189
11-14	245.8 ± 5.07 (15)	226.7 ± 14.21 (15)	253.9 ± 14.30 (14)	P=0.957
18-21	333.2 ± 11.72 (15)	289.7 ± 20.13 (15)	304.6 ± 16.33 (14)	P=0.216

a Mean ± standard error (number of animals).

b Each dose group is compared to the control with Shirley's test when a trend is present (P<0.01 from Jonckheere's trend test), otherwise Dunn's test is applied (*= P<0.05).

TABLE 1-8B
 F₁: SUMMARY OF DAM DAILY WATER CONSUMPTION (G/ANIMAL/DAY) DURING LACTATION OF THE F₂ LITTER
 TWO GENERATION REPRODUCTION TOXICITY STUDY OF PROPYLTHIOURACIL WHEN ADMINISTERED TO SPRAGUE-DAWLEY RATS IN THE DRINKING WATER

POSTNATAL DAYS	DOSE GROUP ^a			TREND ^b
	0.0000 %	0.0001 %	0.0004 %	
	<u>FEMALE</u>			
1-4	49.2 ± 1.67 (14)	*43.0 ± 1.70 (15)	45.9 ± 2.80 (13)	P=0.184
4-7	69.7 ± 2.32 (11)	*58.3 ± 3.40 (14)	*61.5 ± 2.48 (13)	P=0.027
7-11	79.5 ± 1.61 (11)	*68.6 ± 3.41 (15)	75.3 ± 3.19 (14)	P=0.279
11-14	89.8 ± 1.41 (15)	*81.0 ± 4.51 (15)	91.3 ± 4.97 (14)	P=0.377
14-18	85.0 ± 3.64 (15)	85.6 ± 4.90 (15)	90.0 ± 3.52 (14)	P=0.525
18-21	111.6 ± 3.39 (15)	97.6 ± 6.13 (15)	101.6 ± 5.17 (14)	P=0.135

a Mean ± standard error (number of animals).

b Each dose group is compared to the control with Shirley's test when a trend is present (P<0.01 from Jonckheere's trend test), otherwise Dunn's test is applied (*= P<0.05).

TABLE 1-9
F₁: SUMMARY OF ESTIMATED DOSAGE (MG/KG BODY WEIGHT/DAY) DURING LACTATION OF THE F₂ LITTER
TWO GENERATION REPRODUCTION TOXICITY STUDY OF PROPYLTHIOURACIL WHEN ADMINISTERED TO SPRAGUE-DAWLEY RATS IN THE DRINKING WATER

POSTNATAL DAYS	DOSE GROUP ^a	
	0.0001 %	0.0004 %
	<u>FEMALE</u>	
1-4	0.1 ± 0.01 (15)	0.5 ± 0.04 (13)
4-7	0.2 ± 0.01 (14)	0.7 ± 0.03 (13)
11-14	0.2 ± 0.01 (15)	1.0 ± 0.06 (14)
18-21	0.3 ± 0.02 (15)	1.2 ± 0.07 (14)

a Mean ± standard error (number of animals).

TABLE 1-10
F₁: SUMMARY OF ANOGENITAL DISTANCE AND PUP WEIGHT FOR THE F₂ LITTER
TWO GENERATION REPRODUCTION TOXICITY STUDY OF PROPYLTHIOURACIL WHEN ADMINISTERED TO SPRAGUE-DAWLEY RATS IN THE DRINKING WATER

Endpoint	DOSE GROUPS ^a			TREND ^b
	0.0000 %	0.0001 %	0.0004 %	
<u>MALE</u>				
Anogenital Distance (PND1)	2.72 ± 0.041 (15)	*2.51 ± 0.034 (15)	*2.52 ± 0.055 (14)	P=0.001
Average Pup Weight (PND1)	6.79 ± 0.132 (15)	6.81 ± 0.177 (15)	6.87 ± 0.154 (14)	P=0.575
Ratio AGD/Pup Weight (PND1)	0.40 ± 0.007 (15)	0.37 ± 0.011 (15)	*0.37 ± 0.009 (14)	P=0.021
<u>FEMALE</u>				
Anogenital Distance (PND1)	1.20 ± 0.020 (15)	1.20 ± 0.024 (15)	1.22 ± 0.016 (14)	P=0.738
Average Pup Weight (PND1)	6.31 ± 0.119 (15)	6.36 ± 0.173 (15)	6.50 ± 0.123 (14)	P=0.126
Ratio AGD/Pup Weight (PND1)	0.19 ± 0.005 (15)	0.19 ± 0.006 (15)	0.19 ± 0.004 (14)	P=0.355

a Mean ± standard error (number of dams).

b Each dose group is compared to the control with Shirley's test when a trend is present (P<0.01 from Jonckheere's trend test), otherwise Dunn's test is applied (* = P<0.05).

TABLE 1-11
 F₁: SUMMARY OF AVERAGE PUP WEIGHTS FOR THE F₂ LITTER
 TWO GENERATION REPRODUCTION TOXICITY STUDY OF PROPYLTHIOURACIL WHEN ADMINISTERED TO SPRAGUE-DAWLEY RATS IN THE DRINKING WATER

AGE (DAYS)	DOSE GROUP ^a			TREND ^b
	0.0000 %	0.0001 %	0.0004 %	
<u>MALE</u>				
1	6.83 ± 0.137 (14)	6.74 ± 0.191 (15)	6.93 ± 0.157 (13)	P=0.412
4	9.55 ± 0.298 (14)	9.94 ± 0.356 (15)	9.84 ± 0.345 (13)	P=0.350
7	14.15 ± 0.790 (10)	15.13 ± 0.718 (15)	14.26 ± 0.362 (14)	P=0.707
14	28.14 ± 0.866 (15)	29.23 ± 1.367 (15)	27.27 ± 0.727 (14)	P=0.813
21	44.31 ± 1.528 (15)	45.66 ± 2.359 (15)	41.79 ± 1.197 (14)	P=0.338
<u>FEMALE</u>				
1	6.34 ± 0.123 (14)	6.41 ± 0.189 (15)	6.55 ± 0.135 (13)	P=0.166
4	8.95 ± 0.286 (14)	9.31 ± 0.335 (15)	9.45 ± 0.341 (13)	P=0.090
7	13.26 ± 0.785 (10)	14.30 ± 0.669 (15)	13.70 ± 0.364 (14)	P=0.415
14	26.44 ± 0.995 (15)	27.87 ± 1.386 (15)	26.29 ± 0.558 (14)	P=0.643
21	42.02 ± 1.697 (15)	43.18 ± 2.169 (15)	39.89 ± 1.075 (14)	P=0.458

a Mean ± standard error (number of litters).

b Each dose group is compared to the control with Shirley's test when a trend is present (P<0.01 from Jonckheere's trend test), otherwise Dunn's test is applied (* = P<0.05).

TABLE 1-12
 F₁: SUMMARY OF PUP SURVIVAL FOR THE F₂ LITTER
 TWO GENERATION REPRODUCTION TOXICITY STUDY OF PROPYLTHIOURACIL WHEN ADMINISTERED TO SPRAGUE-DAWLEY RATS IN THE DRINKING WATER

AGE (DAYS)	DOSE GROUP ^a			TREND ^b
	0.0000 %	0.0001 %	0.0004 %	
<u>MALE</u>				
4	0.99 ± 0.012 (14)	0.99 ± 0.007 (15)	0.99 ± 0.007 (13)	P=0.979
7	0.98 ± 0.017 (10)	0.99 ± 0.007 (15)	0.99 ± 0.006 (14)	P=0.758
14	0.98 ± 0.012 (15)	0.98 ± 0.010 (15)	0.99 ± 0.010 (14)	P=1.000
21	0.97 ± 0.014 (15)	0.98 ± 0.010 (15)	0.97 ± 0.019 (14)	P=0.962
<u>FEMALE</u>				
4	1.00 ± 0.000 (14)	0.99 ± 0.008 (15)	0.97 ± 0.017 (13)	P=0.043
7	0.99 ± 0.011 (10)	0.98 ± 0.017 (15)	0.97 ± 0.016 (14)	P=0.337
14	0.98 ± 0.012 (15)	0.98 ± 0.017 (15)	0.97 ± 0.016 (14)	P=0.879
21	0.98 ± 0.012 (15)	0.98 ± 0.018 (15)	0.97 ± 0.016 (14)	P=0.848
<u>COMBINED</u>				
4	0.93 ± 0.066 (15)	0.99 ± 0.008 (15)	0.98 ± 0.010 (13)	P=0.300
7	0.99 ± 0.008 (11)	0.98 ± 0.015 (15)	0.98 ± 0.009 (14)	P=0.402
14	0.98 ± 0.008 (15)	0.98 ± 0.015 (15)	0.97 ± 0.012 (14)	P=0.934
21	0.97 ± 0.009 (15)	0.97 ± 0.015 (15)	0.97 ± 0.012 (14)	P=0.858

a Mean of (number of live pups/number of pups born alive) ± standard error (number of litters). Ratios greater than 1 or increases in survival over time indicate missexing of pups.

b Each dose group is compared to the control with Shirley's test when a trend is present (P<0.01 from Jonckheere's trend test), otherwise Dunn's test is applied (* = P<0.05).

TABLE 1-13
F₁: SUMMARY OF NIPPLE RETENTION FOR THE F₂ LITTER
TWO GENERATION REPRODUCTION TOXICITY STUDY OF PROPYLTHIURACIL WHEN ADMINISTERED TO SPRAGUE-DAWLEY RATS IN THE DRINKING WATER

AGE (DAYS)	DOSE GROUP ^a		
	0.0000 %	0.0001 %	0.0004 %
12	0.00 ± 0.00 (15)	0.00 ± 0.00 (15)	0.00 ± 0.00 (14)
13	0.00 ± 0.00 (15)	0.00 ± 0.00 (15)	0.00 ± 0.00 (14)

a Mean ± standard error (number of animals).

TABLE 1-14
F₁: SUMMARY OF PINNA DETACHMENT AND EYE OPENING FOR THE F₂ LITTER
TWO GENERATION REPRODUCTION TOXICITY STUDY OF PROPYLTHIOURACIL WHEN ADMINISTERED TO SPRAGUE-DAWLEY RATS IN THE DRINKING WATER

ENDPOINT ^b	DOSE GROUP ^a			TREND ^c
	0.0000 %	0.0001 %	0.0004 %	
DAY OF EYE OPENING	16.3 ± 0.18 (15)	16.1 ± 0.17 (15)	16.9 ± 0.25 (14)	P=0.045
DAY OF PINNA DETACHMENT	5.2 ± 0.17 (15)	5.5 ± 0.24 (15)	4.9 ± 0.27 (14)	P=0.431

a Mean ± standard error (number of animals).

b Mean = age (days)

c Each dose group is compared to the control with Shirley's test when a trend is present (P<0.01 from Jonckheere's trend test), otherwise Dunn's test is applied (* = P<0.05).

TABLE 1-15
 F₁: SUMMARY OF F₂ MALE PND 21 NECROPSY GROSS OBSERVATIONS
 TWO GENERATION REPRODUCTION TOXICITY STUDY OF PROPYLTHIOURACIL WHEN ADMINISTERED TO SPRAGUE-DAWLEY RATS IN THE DRINKING WATER

	GROUP:	1	2	3
	DOSE (%):	0.0000	0.0001	0.0004
	NUMBER OF ANIMALS ON STUDY:	45	42	42
	NUMBER OF ANIMALS COMPLETED:	(45)	(42)	(42)
RIGHT TESTIS;				
SUBMITTED.....	(45)	(42)	(42)	(42)
NO VISIBLE LESIONS.....	45	42	42	42
SEMINAL VESICLES W/ COAGULATING GLANDS;				
SUBMITTED.....	(45)	(42)	(42)	(42)
NO VISIBLE LESIONS.....	45	42	42	42
LIVER;				
SUBMITTED.....	(45)	(42)	(42)	(42)
NO VISIBLE LESIONS.....	45	42	42	42
ENLARGEMENT; ALL LOBES	1	0	0	0
LEFT TESTES W/EPIDIDYMIDES;				
SUBMITTED.....	(45)	(42)	(42)	(42)
NO VISIBLE LESIONS.....	45	42	42	42
THYROID/PARATHYROID;				
SUBMITTED.....	(45)	(42)	(42)	(42)
NO VISIBLE LESIONS.....	45	42	42	42
ADRENAL GLAND(S);				
SUBMITTED.....	(45)	(42)	(42)	(42)
NO VISIBLE LESIONS.....	45	42	42	42
BRAIN;				
SUBMITTED.....	(1)	(1)	(4)	(4)
NO VISIBLE LESIONS.....	45	42	42	42
KIDNEY(S);				
SUBMITTED.....	(45)	(42)	(42)	(42)
NO VISIBLE LESIONS.....	45	42	42	42

TABLE 1-15 (CONTINUED)
 F₁: SUMMARY OF F₂ MALE PND 21 NECROPSY GROSS OBSERVATIONS
 TWO GENERATION REPRODUCTION TOXICITY STUDY OF PROPYLTHIOURACIL WHEN ADMINISTERED TO SPRAGUE-DAWLEY RATS IN THE DRINKING WATER

	GROUP:	1	2	3
	DOSE (%):	0.0000	0.0001	0.0004
PITUITARY GLAND;				
SUBMITTED.....		(45)	(42)	(41)
NO VISIBLE LESIONS.....		45	42	42
SPLEEN;				
SUBMITTED.....		(1)	(2)	(4)
NO VISIBLE LESIONS.....		45	42	42
THYMUS;				
SUBMITTED.....		(18)	(6)	(8)
NO VISIBLE LESIONS.....		45	42	42
VENTRAL PROSTATE;				
SUBMITTED.....		(45)	(42)	(42)
NO VISIBLE LESIONS.....		45	42	42
DORSOLATERAL PROSTATE;				
SUBMITTED.....		(45)	(42)	(42)
NO VISIBLE LESIONS.....		45	42	42

TABLE 1-16
 F₁: SUMMARY OF F₂ MALE PND 21 ABSOLUTE ORGAN WEIGHTS AND ORGAN-TO-BODY WEIGHT RATIOS
 TWO GENERATION REPRODUCTION TOXICITY STUDY OF PROPYLTHIOURACIL WHEN ADMINISTERED TO SPRAGUE-DAWLEY RATS IN THE DRINKING WATER

ORGAN	DOSE GROUP			TREND ^a
	0.0000 %	0.0001 %	0.0004 %	
	<u>ORGAN WEIGHTS (g)^b</u>			
BODY	44.9 ± 0.95 (45)	44.7 ± 1.17 (42)	43.2 ± 0.78 (42)	P=0.264
BRAIN	1.477 ± 0.0092 (45)	1.466 ± 0.0096 (42)	1.479 ± 0.0096 (42)	P=0.884
SPLEEN	0.190 ± 0.0046 (45)	*0.216 ± 0.0048 (42)	0.201 ± 0.0048 (42)	P=0.105
THYMUS	0.191 ± 0.0041 (45)	0.187 ± 0.0042 (42)	0.180 ± 0.0042 (42)	P=0.064
	<u>ORGAN-TO-BODY WEIGHT RATIOS^c</u>			
BRAIN	33.479 ± 0.5867 (45)	33.632 ± 0.7650 (42)	34.479 ± 0.5796 (42)	P=0.289
SPLEEN	4.270 ± 0.1124 (45)	*4.799 ± 0.1460 (42)	4.406 ± 0.1209 (42)	P=0.601
THYMUS	4.295 ± 0.0782 (45)	4.211 ± 0.1069 (42)	4.020 ± 0.0940 (42)	P=0.073

a Each dose group is compared with the control group by Shirley's test if P<0.01 from Jonckheere's trend test, otherwise Dunn's test is applied (*= P<0.05).

b Mean organ weight ± standard error (number of animals).

c Mean ratio (mg/g body weight) ± standard error (number of animals).

Note: Absolute organ weights were covaried by body weight.

TABLE 1-17
 F₁: SUMMARY OF F₂ FEMALE PND 21 NECROPSY GROSS OBSERVATIONS
 TWO GENERATION REPRODUCTION TOXICITY STUDY OF PROPYLTHIOURACIL WHEN ADMINISTERED TO SPRAGUE-DAWLEY RATS IN THE DRINKING WATER

	GROUP:	1	2	3
	DOSE (%):	0.0000	0.0001	0.0004
	NUMBER OF ANIMALS ON STUDY:	45	43	40
	NUMBER OF ANIMALS COMPLETED:	(45)	(43)	(40)
LIVER;				
SUBMITTED.....		(45)	(43)	(40)
NO VISIBLE LESIONS.....		45	43	40
THYROID/PARATHYROID;				
SUBMITTED.....		(45)	(43)	(40)
NO VISIBLE LESIONS.....		45	43	40
ADRENAL GLAND(S);				
SUBMITTED.....		(45)	(43)	(40)
NO VISIBLE LESIONS.....		45	43	40
BRAIN;				
SUBMITTED.....		(1)	(1)	(4)
NO VISIBLE LESIONS.....		45	43	40
CERVIX;				
SUBMITTED.....		(45)	(43)	(40)
NO VISIBLE LESIONS.....		45	43	40
KIDNEY(S);				
SUBMITTED.....		(45)	(43)	(40)
NO VISIBLE LESIONS.....		45	43	40
OVARY(IES);				
SUBMITTED.....		(45)	(43)	(40)
NO VISIBLE LESIONS.....		45	43	40
PITUITARY GLAND;				
SUBMITTED.....		(45)	(43)	(40)
NO VISIBLE LESIONS.....		45	43	40

TABLE 1-17 (CONTINUED)
 F₁: SUMMARY OF F₂ FEMALE PND 21 NECROPSY GROSS OBSERVATIONS
 TWO GENERATION REPRODUCTION TOXICITY STUDY OF PROPYLTHIOURACIL WHEN ADMINISTERED TO SPRAGUE-DAWLEY RATS IN THE DRINKING WATER

	GROUP:	1	2	3
	DOSE (%):	0.0000	0.0001	0.0004
SPLEEN;				
SUBMITTED.....		(0)	(1)	(4)
NO VISIBLE LESIONS.....		45	43	40
THYMUS;				
SUBMITTED.....		(16)	(8)	(9)
NO VISIBLE LESIONS.....		45	43	40
UTERUS;				
SUBMITTED.....		(45)	(43)	(40)
NO VISIBLE LESIONS.....		45	43	40
VAGINA;				
SUBMITTED.....		(45)	(43)	(40)
NO VISIBLE LESIONS.....		45	43	40

TABLE 1-18
 F₁: SUMMARY OF F₂ FEMALE PND 21 ABSOLUTE ORGAN WEIGHTS AND ORGAN-TO-BODY WEIGHT RATIOS
 TWO GENERATION REPRODUCTION TOXICITY STUDY OF PROPYLTHIOURACIL WHEN ADMINISTERED TO SPRAGUE-DAWLEY RATS IN THE DRINKING WATER

ORGAN	0.0000 %	DOSE GROUP		TREND ^a
		0.0001 %	0.0004 %	
		<u>ORGAN WEIGHTS (g)^b</u>		
BODY	42.8 ± 1.06 (45)	42.6 ± 1.27 (43)	41.3 ± 0.72 (40)	P=0.348
BRAIN	1.403 ± 0.0089 (45)	1.418 ± 0.0091 (43)	1.432 ± 0.0095 (40)	P=0.027
SPLEEN	0.198 ± 0.0047 (45)	0.211 ± 0.0049 (43)	0.196 ± 0.0050 (40)	P=0.789
THYMUS	0.194 ± 0.0041 (45)	0.192 ± 0.0042 (43)	*0.177 ± 0.0044 (40)	P=0.007
		<u>ORGAN-TO-BODY WEIGHT RATIOS^c</u>		
BRAIN	33.554 ± 0.6693 (45)	34.285 ± 0.8345 (43)	34.804 ± 0.5544 (40)	P=0.121
SPLEEN	4.615 ± 0.1508 (45)	4.878 ± 0.1657 (43)	4.493 ± 0.1067 (40)	P=0.643
THYMUS	4.559 ± 0.0917 (45)	4.537 ± 0.0994 (43)	*4.145 ± 0.1106 (40)	P=0.012

a Each dose group is compared with the control group by Shirley's test if P<0.01 from Jonckheere's trend test, otherwise Dunn's test is applied (*= P<0.05).

b Mean organ weight ± standard error (number of animals).

c Mean ratio (mg/g body weight) ± standard error (number of animals).

Note: Absolute organ weights were covaried by body weight.

TABLE 1-19
F₁: SUMMARY OF STAGES AND LENGTH OF ESTROUS CYCLE
TWO GENERATION REPRODUCTION TOXICITY STUDY OF PROPYLTHIOURACIL WHEN ADMINISTERED TO SPRAGUE-DAWLEY RATS IN THE DRINKING WATER

DOSE %	ESTROUS STAGE (%)					CYCLE LENGTH ^{a,d} (DAYS)	NUMBER OF CYCLES ^{a,d}	NUMBER OF CYCLING FEMALES ^{b,c}	NUMBER OF FEMALES W/REGULAR CYCLE ^{c,e}
	P	E	M	D	N				
0.0000	17.5	33.9	20.7	27.5	0.4	4.2 ± 0.10 (18) ^f	2.4 ± 0.12 (18)	18/20 (90)	17/20 (85)
0.0001	16.1	35.0	18.2	30.7	0.0	4.5 ± 0.21 (18) ^g	2.4 ± 0.16 (18)	18/20 (90)	15/20 (75)
0.0004	18.6	33.9	19.6	27.9	0.0	4.3 ± 0.16 (19) ^h	2.5 ± 0.18 (19)	19/20 (95)	14/20 (70)

P=Proestrus, E=Estrus, M=Metestrus, D=Diestrus, N=Not Clear, Poor Quality, or Insufficient Number of Cells

a Mean ± standard error (number of animals)

b Number of cycling females / number of females sampled (percent cycling).

c Number of females with a regular cycle / number of females sampled (percent regular).

d Each dose group is compared to the control group by Shirley's test if P<0.01 from Jonckheere's trend test. Otherwise, Dunn's test is applied.

e Dose group compared to the control group using the Chi-squared test.

f Estrous cycle ten days or longer, not cycling, not available, or unclear in 2 of 20 animals.

g Estrous cycle ten days or longer, not cycling, not available, or unclear in 2 of 20 animals.

h Estrous cycle ten days or longer, not cycling, not available, or unclear in 1 of 20 animals.

NOTE: There is no evidence of any differences between the dosed and control groups in cycle length, in number of cycles, or in relative length of time spent in estrous stages.

TABLE 1-20
 F₁: SUMMARY OF THYROID HORMONES
 TWO GENERATION REPRODUCTION TOXICITY STUDY OF PROPYLTHIOURACIL WHEN ADMINISTERED TO SPRAGUE-DAWLEY RATS IN THE DRINKING WATER

ENDPOINT	DOSE GROUP ^a			TREND ^b
	0.0000 %	0.0001 %	0.0004 %	
<u>MALE</u>				
T3 (NG/DL)	94.34 ± 4.686 (19)	92.87 ± 4.819 (20)	87.07 ± 3.762 (20)	P=0.278
T4 (UG/DL)	3.53 ± 0.207 (19)	4.58 ± 0.234 (20)	*1.37 ± 0.158 (20)	P<0.001
TSH (NG/ML)	5.13 ± 0.642 (19)	5.47 ± 0.704 (20)	*13.93 ± 1.542 (20)	P<0.001
<u>FEMALE</u>				
T3 (NG/DL)	111.00 ± 5.190 (20)	109.43 ± 5.600 (20)	111.77 ± 5.084 (20)	P=0.989
T4 (UG/DL)	2.25 ± 0.187 (20)	2.61 ± 0.306 (20)	*0.92 ± 0.189 (20)	P<0.001
TSH (NG/ML)	3.73 ± 0.565 (20)	4.77 ± 0.503 (20)	*9.09 ± 1.340 (20)	P<0.001

a Mean ± standard error (number of animals).

b Each dose group is compared to the control with Dunnett's one-tailed low test for T3 and T4 or Dunnett's one-tailed high test for TSH.
 (* = P<0.05 from Jonckheere's trend test.)

TABLE 1-21
 F₁: SUMMARY OF MALE GROSS OBSERVATIONS
 TWO GENERATION REPRODUCTION TOXICITY STUDY OF PROPYLTHIOURACIL WHEN ADMINISTERED TO SPRAGUE-DAWLEY RATS IN THE DRINKING WATER

	GROUP: DOSE (%):	1 0.0000	2 0.0001	3 0.0004	
		NUMBER OF ANIMALS ON STUDY:	19	20	20
		NUMBER OF ANIMALS COMPLETED:	(19)	(20)	(20)
RIGHT TESTIS;					
SUBMITTED.....		(19)	(20)	(20)	
NO VISIBLE LESIONS.....		19	20	20	
VENTRAL PROSTATE;					
SUBMITTED.....		(19)	(20)	(20)	
NO VISIBLE LESIONS.....		19	20	20	
THYROID/PARATHYROID;					
SUBMITTED.....		(19)	(20)	(20)	
NO VISIBLE LESIONS.....		19	20	20	
ADRENAL GLAND(S);					
SUBMITTED.....		(19)	(20)	(20)	
NO VISIBLE LESIONS.....		19	20	20	
BRAIN;					
SUBMITTED.....		(19)	(20)	(20)	
NO VISIBLE LESIONS.....		19	20	20	
KIDNEY(S);					
SUBMITTED.....		(19)	(20)	(20)	
NO VISIBLE LESIONS.....		19	20	20	
LIVER;					
SUBMITTED.....		(19)	(20)	(20)	
NO VISIBLE LESIONS.....		19	20	20	
PITUITARY GLAND;					
SUBMITTED.....		(19)	(20)	(20)	
NO VISIBLE LESIONS.....		19	20	20	
SPLEEN;					
SUBMITTED.....		(19)	(20)	(20)	
NO VISIBLE LESIONS.....		19	20	20	

TABLE 1-21 (CONTINUED)
 F₁: SUMMARY OF MALE GROSS OBSERVATIONS
 TWO GENERATION REPRODUCTION TOXICITY STUDY OF PROPYLTHIOURACIL WHEN ADMINISTERED TO SPRAGUE-DAWLEY RATS IN THE DRINKING WATER

	GROUP:	1	2	3
	DOSE (%):	0.0000	0.0001	0.0004
URINARY BLADDER;				
SUBMITTED.....		(0)	(1)	(0)
NO VISIBLE LESIONS.....		0	0	0
DISTENSION; RED		0	1	0
DORSOLATERAL PROSTATE;				
SUBMITTED.....		(19)	(20)	(20)
NO VISIBLE LESIONS.....		19	20	20
LEFT TESTIS W/ EPIDIDYMIS;				
SUBMITTED.....		(19)	(20)	(20)
NO VISIBLE LESIONS.....		19	20	20
SEMINAL VESICLES W/COAGULATING GLANDS;				
SUBMITTED.....		(19)	(20)	(20)
NO VISIBLE LESIONS.....		19	20	20
RIGHT CAUDA EPIDIDYMIS;				
SUBMITTED.....		(19)	(20)	(20)
NO VISIBLE LESIONS.....		19	20	20
RIGHT EPIDIDYMIS;				
SUBMITTED.....		(19)	(20)	(20)
NO VISIBLE LESIONS.....		19	20	20

TABLE 1-22
 F₁: SUMMARY OF MALE ABSOLUTE ORGAN WEIGHTS (G) AND ORGAN-TO-BODY WEIGHT RATIOS
 TWO GENERATION REPRODUCTION TOXICITY STUDY OF PROPYLTHIOURACIL WHEN ADMINISTERED TO SPRAGUE-DAWLEY RATS IN THE DRINKING WATER

ORGAN	DOSE GROUP ^a			TREND ^b
	0.0000 %	0.0001 %	0.0004 %	
	<u>ORGAN WEIGHTS (g)^b</u>			
BODY	698.7 ± 15.78 (19)	673.6 ± 17.60 (20)	678.9 ± 21.57 (20)	P=0.307
ADRENALS	0.0556 ± 0.00269 (19)	0.0574 ± 0.00262 (20)	0.0581 ± 0.00268 (19)	P=0.505
BRAIN	2.242 ± 0.0233 (19)	2.208 ± 0.0227 (20)	2.190 ± 0.0232 (19)	P=0.118
KIDNEYS	4.2506 ± 0.07942 (19)	4.2579 ± 0.07725 (20)	4.1637 ± 0.07903 (19)	P=0.436
LIVER	23.0832 ± 0.46462 (19)	23.1898 ± 0.45191 (20)	22.2020 ± 0.46234 (19)	P=0.182
PITUITARY	0.0146 ± 0.00063 (19)	0.0146 ± 0.00061 (20)	0.0145 ± 0.00063 (19)	P=0.868
DORSOLAT PROSTATE	0.7059 ± 0.04331 (19)	0.6811 ± 0.04213 (20)	0.7130 ± 0.04310 (19)	P=0.903
VENTRAL PROSTATE	0.6347 ± 0.03540 (19)	0.5541 ± 0.03443 (20)	0.5874 ± 0.03523 (19)	P=0.357
RIGHT CAUDA EPIDIDYMIS	0.3464 ± 0.01178 (19)	0.3334 ± 0.01146 (20)	0.3774 ± 0.01172 (19)	P=0.073
RIGHT EPIDIDYMIS	0.7560 ± 0.01754 (19)	0.7413 ± 0.01706 (20)	0.7902 ± 0.01745 (19)	P=0.174
RIGHT TESTIS	1.8022 ± 0.05495 (19)	1.7661 ± 0.05345 (20)	1.8889 ± 0.05486 (19)	P=0.266
SEMINAL VESICLES	2.1614 ± 0.10701 (19)	2.0554 ± 0.10408 (20)	2.2805 ± 0.10648 (19)	P=0.430
SPLEEN	0.931 ± 0.0268 (19)	0.937 ± 0.0261 (20)	0.940 ± 0.0267 (19)	P=0.803
THYROID	0.0306 ± 0.00241 (19)	0.0379 ± 0.00234 (20)	*0.0551 ± 0.00240 (19)	P<0.001
	<u>ORGAN-TO-BODY WEIGHT RATIOS^c</u>			
ADRENALS	0.0808 ± 0.00363 (19)	0.0849 ± 0.00421 (20)	0.0848 ± 0.00346 (19)	P=0.653
BRAIN	3.243 ± 0.0740 (19)	3.310 ± 0.0844 (20)	3.266 ± 0.0956 (19)	P=0.825
KIDNEYS	6.1934 ± 0.13949 (19)	6.2876 ± 0.11873 (20)	6.1221 ± 0.11526 (19)	P=0.674
LIVER	33.6967 ± 0.84311 (19)	33.6435 ± 0.66433 (20)	32.3431 ± 0.50029 (19)	P=0.251
PITUITARY	0.0212 ± 0.00093 (19)	0.0216 ± 0.00084 (20)	0.0216 ± 0.00098 (19)	P=0.412
DORSOLATERAL PROSTATE	1.0222 ± 0.05691 (19)	1.0112 ± 0.06289 (20)	1.0654 ± 0.07331 (19)	P=0.792
VENTRAL PROSTATE	0.9213 ± 0.05007 (19)	0.8288 ± 0.05947 (20)	0.8743 ± 0.05290 (19)	P=0.462
RIGHT CAUDA EPIDIDYMIS	0.5001 ± 0.01232 (19)	0.4984 ± 0.02502 (20)	0.5639 ± 0.02312 (19)	P=0.043
RIGHT EPIDIDYMIS	1.0933 ± 0.02508 (19)	1.1123 ± 0.04720 (20)	1.1794 ± 0.03828 (19)	P=0.091
RIGHT TESTIS	2.6065 ± 0.05803 (19)	2.6431 ± 0.12859 (20)	2.8194 ± 0.10059 (19)	P=0.076
SEMINAL VESICLES	3.1351 ± 0.17383 (19)	3.0741 ± 0.19020 (20)	3.3748 ± 0.12505 (19)	P=0.182
SPLEEN	1.356 ± 0.0420 (19)	1.376 ± 0.0319 (20)	1.384 ± 0.0416 (19)	P=0.705
THYROID	0.0446 ± 0.00229 (19)	*0.0563 ± 0.00399 (20)	*0.0819 ± 0.00440 (19)	P<0.001

a Each dose group is compared with the control group by Shirley's test if P<0.01 from Jonckheere's trend test, otherwise Dunn's test is applied (*= P<0.05).

b Mean organ weight ± standard error (number of animals).

c Mean ratio (mg/g body weight) ± standard error (number of animals).

TABLE 1-23A
F₁: SUMMARY OF SPERM ANALYSIS DATA
TWO GENERATION REPRODUCTION TOXICITY STUDY OF PROPYLTHIOURACIL WHEN ADMINISTERED TO SPRAGUE-DAWLEY RATS IN THE DRINKING WATER

ENDPOINT	DOSE GROUP ^a			TREND ^b
	0.0000 %	0.0001 %	0.0004 %	
EPIDIDYMAL SPERM DENSITY ^c	399.62 ± 14.314 (19)	394.59 ± 24.371 (20)	396.44 ± 12.458 (19)	P=0.937
SPERM MORPHOLOGY (% ABNORMAL) ^d	0.5 ± 0.20 (19)	0.7 ± 0.13 (19)	0.4 ± 0.07 (20)	P=0.944
SPERMATIDS/MG TESTIS ^e	61.72 ± 3.023 (19)	52.56 ± 3.526 (20)	54.62 ± 2.802 (20)	P=0.198
TOTAL SPERMATIDS/TESTIS ^f	105.03 ± 4.630 (19)	90.63 ± 5.962 (20)	95.25 ± 4.000 (20)	P=0.168
TOTAL SPERM/CAUDA ^f	138.28 ± 4.963 (19)	135.51 ± 8.874 (20)	146.46 ± 4.630 (20)	P=0.333

a Endpoint mean ± standard error (number of animals).

b Each dose group is compared with the control group by Shirley's test if P<0.01 from Jonckheere's trend test, otherwise Dunn's test is applied (* = P<0.05).

c Sperm density expressed as 1000 sperm per mg right caudal tissue.

d Dose group means and standard errors are computed only from samples with at least 200 sperm.

e Spermatid heads per mg testis (x 1,000) = (total number of spermatid heads per right testis) / (right testis weight in mg)

f Total number of spermatid heads or sperm (x 1,000,000) = (average number of spermatid heads or sperm x 2.5 x 100) / 0.0001

TABLE 1-23B
F₁: SUMMARY OF HAMILTON-THORNE SPERM ANALYSIS DATA
TWO GENERATION REPRODUCTION TOXICITY STUDY OF PROPYLTHIOURACIL WHEN ADMINISTERED TO SPRAGUE-DAWLEY RATS IN THE DRINKING WATER

ENDPOINT	DOSE GROUP ^a			TREND ^b
	0.0000 %	0.0001 %	0.0004 %	
AMPLITUDE	15.6 ± 0.55 (18)	15.2 ± 0.59 (19)	14.4 ± 0.70 (18)	P=0.237
BEAT/CROSS FREQUENCY (hz/sec)	16.9 ± 0.47 (18)	17.1 ± 0.52 (19)	17.2 ± 0.48 (18)	P=0.716
LINEARITY	35.7 ± 0.81 (18)	37.6 ± 0.48 (19)	37.7 ± 0.53 (18)	P=0.068
MOTILE PERCENTAGE	72.3 ± 3.79 (18)	80.9 ± 4.65 (19)	74.6 ± 4.24 (18)	P=0.526
PROGRESSIVE VELOCITY	82.5 ± 2.71 (18)	89.8 ± 2.80 (19)	86.0 ± 3.52 (18)	P=0.250
PATH VELOCITY	136.8 ± 3.56 (18)	145.9 ± 4.64 (19)	137.5 ± 5.10 (18)	P=0.865
STRAIGHTNESS	60.3 ± 1.22 (18)	60.9 ± 0.56 (19)	61.3 ± 0.91 (18)	P=0.889
TRACK SPEED	240.7 ± 6.69 (18)	248.1 ± 8.34 (19)	234.6 ± 8.14 (18)	P=0.688

a Endpoint mean ± standard error (number of animals).

b Each dose group is compared with the control group by Shirley's test if P<0.01 from Jonckheere's trend test, otherwise Dunn's test is applied (* = P<0.05).

Note: Absolute organ weights were covaried by body weight.

TABLE 1-24
 F₁: SUMMARY OF FEMALE GROSS OBSERVATIONS
 TWO GENERATION REPRODUCTION TOXICITY STUDY OF PROPYLTHIOURACIL WHEN ADMINISTERED TO SPRAGUE-DAWLEY RATS IN THE DRINKING WATER

	GROUP:	1	2	3
	DOSE (%):	0.0000	0.0001	0.0004
	NUMBER OF ANIMALS ON STUDY:	20	20	20
	NUMBER OF ANIMALS COMPLETED:	(20)	(20)	(20)
THYROID/PARATHYROID;				
SUBMITTED.....	(20)	(20)	(20)	(20)
NO VISIBLE LESIONS.....	20	20	20	20
ADRENAL GLAND(S);				
SUBMITTED.....	(20)	(20)	(20)	(20)
NO VISIBLE LESIONS.....	20	20	20	20
BRAIN;				
SUBMITTED.....	(20)	(20)	(20)	(20)
NO VISIBLE LESIONS.....	20	20	20	20
KIDNEY(S);				
SUBMITTED.....	(20)	(20)	(20)	(20)
NO VISIBLE LESIONS.....	20	20	20	20
LIVER;				
SUBMITTED.....	(20)	(20)	(20)	(20)
NO VISIBLE LESIONS.....	20	20	20	20
OVARY (IES);				
SUBMITTED.....	(20)	(20)	(20)	(20)
NO VISIBLE LESIONS.....	20	20	20	20
PITUITARY GLAND;				
SUBMITTED.....	(20)	(20)	(20)	(20)
NO VISIBLE LESIONS.....	20	20	20	20
SPLEEN;				
SUBMITTED.....	(20)	(20)	(20)	(20)
NO VISIBLE LESIONS.....	20	20	20	20
VAGINA/CERVIX/UTERUS;				
SUBMITTED.....	(20)	(20)	(20)	(20)
NO VISIBLE LESIONS.....	20	20	20	20

TABLE 1-25
 F₁: SUMMARY OF FEMALE ABSOLUTE ORGAN WEIGHTS (G) AND ORGAN-TO-BODY WEIGHT RATIOS
 TWO GENERATION REPRODUCTION TOXICITY STUDY OF PROPYLTHIOURACIL WHEN ADMINISTERED TO SPRAGUE-DAWLEY RATS IN THE DRINKING WATER

ORGAN	DOSE GROUP			TREND ^a
	0.0000 %	0.0001 %	0.0004 %	
	<u>ORGAN WEIGHTS (g)^b</u>			
BODY	367.1 ± 8.97 (20)	354.4 ± 7.79 (20)	347.2 ± 8.03 (20)	P=0.158
ADRENALS	0.0772 ± 0.00284 (19)	0.0703 ± 0.00274 (20)	0.0703 ± 0.00276 (20)	P=0.091
BRAIN	1.962 ± 0.0233 (19)	1.998 ± 0.0224 (20)	2.013 ± 0.0226 (20)	P=0.125
KIDNEYS	2.2830 ± 0.04624 (19)	2.2549 ± 0.04458 (20)	2.2726 ± 0.04496 (20)	P=0.879
LIVER	12.7539 ± 0.28576 (19)	12.5759 ± 0.27553 (20)	12.3024 ± 0.27785 (20)	P=0.261
OVARIES	0.1234 ± 0.00540 (19)	0.1104 ± 0.00521 (20)	0.1183 ± 0.00525 (20)	P=0.533
PITUITARY	0.0165 ± 0.00081 (19)	0.0171 ± 0.00078 (20)	0.0156 ± 0.00078 (20)	P=0.441
SPLEEN	0.590 ± 0.0208 (19)	0.604 ± 0.0200 (20)	0.610 ± 0.0202 (20)	P=0.489
THYROID	0.0234 ± 0.00148 (19)	0.0268 ± 0.00143 (20)	*0.0403 ± 0.00144 (20)	P<0.001
UTERUS	1.012 ± 0.0493 (19)	0.997 ± 0.0475 (20)	1.053 ± 0.0479 (20)	P=0.553
	<u>ORGAN-TO-BODY WEIGHT RATIOS^c</u>			
ADRENALS	0.2144 ± 0.00896 (19)	0.2004 ± 0.00957 (20)	0.2028 ± 0.00640 (20)	P=0.243
BRAIN	5.463 ± 0.1544 (19)	5.675 ± 0.1063 (20)	5.831 ± 0.1331 (20)	P=0.053
KIDNEYS	6.4075 ± 0.11635 (19)	6.3893 ± 0.14526 (20)	6.4343 ± 0.12564 (20)	P=0.846
LIVER	35.9446 ± 0.72829 (19)	35.4875 ± 0.81424 (20)	34.3619 ± 0.77643 (20)	P=0.085
OVARIES	0.3428 ± 0.01956 (19)	0.3144 ± 0.01731 (20)	0.3460 ± 0.01426 (20)	P=0.781
PITUITARY	0.0465 ± 0.00218 (19)	0.0481 ± 0.00211 (20)	0.0436 ± 0.00228 (20)	P=0.373
SPLEEN	1.654 ± 0.0577 (19)	1.713 ± 0.0671 (20)	1.730 ± 0.0443 (20)	P=0.303
THYROID	0.0660 ± 0.00394 (19)	0.0757 ± 0.00383 (20)	*0.1150 ± 0.00437 (20)	P<0.001
UTERUS	2.816 ± 0.1479 (19)	2.841 ± 0.1466 (20)	3.056 ± 0.1679 (20)	P=0.156

a Each dose group is compared with the control group by Shirley's test if P<0.01 from Jonckheere's trend test, otherwise Dunn's test is applied (*= P<0.05).

b Mean organ weight ± standard error (number of animals).

c Mean ratio (mg/g body weight) ± standard error (number of animals).

Note: Absolute organ weights were covaried by body weight.

TABLE 1-26
 F₁: SUMMARY OF MORTALITY DATA
 TWO GENERATION REPRODUCTION TOXICITY STUDY OF PROPYLTHIOURACIL WHEN ADMINISTERED TO SPRAGUE-DAWLEY RATS IN THE DRINKING WATER

	DOSE GROUP ^a		
	0.0000 %	0.0001 %	0.0004 %
<u>MALE</u>			
FOUND DEAD	1/20	0/20	0/20
MORIBUND KILLS	0/20	0/20	0/20
REMOVED FROM STUDY	0/20	0/20	0/20
<u>FEMALE</u>			
FOUND DEAD	0/20	0/20	0/20
MORIBUND KILLS	0/20	0/20	0/20
REMOVED FROM STUDY	0/20	0/20	0/20

a Number of animals found dead, moribund, or removed / total number of animals in group.