ISSUES AND CHALLENGES IN MODELING CHILDREN'S LONGITUDINAL EXPOSURES: AN OZONE CASE STUDY

Halûk Özkaynak, Jianping Xue, Valerie Zartarian U.S.EPA/ORD National Exposure Research Laboratory



ABSTRACT

Odeling children's exposures is a complicated, data-intensive process. Modeling longitudinal exposures, which
are important for regulatory decision making, especially
for most air toxics, adds another level of complexity and data
requirements. Because it is difficult to model inter- and intrapersonal variability for exposure model inputs, there is potential for
inaccurate estimation of upper percentiles of longitudinal exposure
prediction model, we need to resolve how to do the following: (1)
obtain longitudinal data needs for time activity and pollutant measurements; (2) separate intra- and inter- person variability of model
puts; (3) his, injust from different data sources and if those data
as inputs for the model, which will preserve its variance-covariance
structure; and (4) use cross-sectional data to simulate longitudinal
data, which is difficult and expensive to collect.

In this reseenantion, we address these issues by anothyine both an

structure; and (4) use cross-ectional data to simulate longitudinal data, which is difficult and expensive to collect.

In this presentation, we address these issues by applying both an existing and a new technique for estimating personal coone exposures of school-age children living in two California communities. The first modeling analysis employed commonly used methods for estimating exposures using a microenvironmental exposure model that used independent distributions in simulations, which were fit to observed inputs for the model. In the new modeling methodology, various warrance components derived from the underlying data usually in the control of the ariance-covariance structures of inputs and output in a sin ut also can accurately predict high percentiles of longitud xposures that are important for regulatory evaluations.



NTRODUCTION

General Modeling Questions

- How can we best determine children's exporonmental agents by different life stages?
- Which locations, sources, media, routes and pathways contribute greatest to exposures of interest?
- What activities and behaviors influence the pollutant contact rates, exposures, uptake and dose?

- rates, exposures, uptake and dose?
 What are the measures of exposures of health relevance (e.g., duration, intensity, timing, frequency)?
 How do exposure and physical factors vary by age, gender, susceptibility, personal habits (e.g., cooking and cleaning exposures second hand smoke), housing type, consumer product usage?
- How do subject-specific longitudinal correlations of personal exposures with outdoor concentrations vary? How are microenvironmental concentrations of PM., ozone, air
- toxics correlated?
- Ambient central site vs. outdoor
- Indoor residential vs. outdoor Personal vs. outdoor

Challenges

- · Lack of longitudinal spatial and temporal concentration data · Lack of longitudinal activity data
- How to simulate height and weight, activity patterns and breath ing rates for children as they grow older?
- · How to use cross-sectional data to simulate longitudinal data?
- How to separate intra- and inter- person variability of model inputs?
- How to preserve variance-covariance structure of inputs when combining different data sources in exposure models (see Oza Case Study)?



SSUES

Pollutant Sources of Concern

- Nearby ETS and PM generating activities
- VOCs from proximity to mobile sources/refueling/attached
- Air toxics from consumer products and hobbies
- Indoor and outdoor aeroallergens · High outdoor ozone/photochemical smog

Concentration-Related

- Do children spend more time indoors (schools, daycare) or in other microenvironments than the adults in the population (see
- Do children engage in fewer pollution generating activities than the other age groups?
- Do personal exposures of children correlate well with outdoor

Figure 1. Relevant Indoor Microenvironments for Children



Figure 2. Relevant Outdoor/In-Vehicle Microenvironments for Children



Activity-Related

- Where do children spend their time with respect to relevant pollutant-containing microenvironments (see Figure 3 for an example of a time-activity diary)?
- Do children spend more time indoors (schools, daycare) or in other special microenvironments than the rest of the population? Do children engage in fewer pollution generating activities than other cohorts?
- Are time-activity profiles of children appreciably different than those for the general population?
- Do children engage in different types of personal activities which generate elevated indoor or personal exposures to PM, combus-tion gases and air toxics?
- How do children's exposures differ as a function of childhood life stages?

Fig 3. Example Human Activity Diary from EPA's Consolidated Human Activities Database (CHAD) (http://www.epa.gov/chadnet1/)

Age-Grouping for Children

- EPA guidance on age grouping is supported by an underlying scientific rationale (Figure 4)
- Use in exposure and risk assessments for consistency Foundation for future exposure factors and/or monitoring data
- generation/anajusa

 U.S. EPA (2005). Guidance on selecting age groups for monitoring and assessing childhood exposures to environmental contaminants. Risk Assessment Forum, Office of Research and Development, Washington, D.C. EPA/650P-03/003F. http://cfpub.epa.
 gov/ncca/cfm/recordisplay.cfm?deid=14658

Fig. 4. EPA Guidance on Age Groups for

METHODS

- Empirical models
 Developed using empirical data from personal monitoring studies
- Example: Regression equations that explain measured exposures
- Deterministic/mechanistic models
- Based on known or assumed physical relationships
- Example: Air quality models, pharmacokinetic models

 $E = \sum_{i} E_{i} = \frac{1}{T} \left(\sum_{i} \overline{C}_{i} t_{i} \right)$

- Stochastor/mechanistic models

 Input data are distributions, not single value

 Input distributions characterize
 variability/uncertainty in the data

 Use Monte Carlo or other random sampling techniques

 Produces output distributions (variability/uncertainty)

Modeling Elements

- Identify cohorts and microenvironments of concern
- Collect available time-activity data

 Locations spent time in and the activities performed while in a location

 Age, gender, region, etc.
- Collect available microenvironmental mea
- Concentrations outdoors, indoors, in-vehicle
- Estimate physical factors
- Source use and emission
- Penetration, Infiltration, re-suspension, volatilization, decay Estimate exposure factors
- Contact/transfer/uptake/PBPK rates or parameters
- Confluctualistic updated by Kanacs of parameters.
 Apply data and algorithms using a selected modeling structure to predict exposure distributions (see Figure 5 for PM₂₅ and Figure 6 for benzene exposures generated by the EPA's SHEDS model)
 Compare estimates to available personal exposure measurements

Inhalation Exposure Pathways



W

Figure 5. Distribution of Daily-average Total PM_{2,5} Exposure and Uncertainty for Selected Percentiles in Philadelphia, PA Predicted by the SHEDS-PM Model





Figure 6. Daily Inhalation Exposures to Benzene in Houston, Texas predicted by the SHEDS-Air-Toxics Model



Ozone Case Study

- Extend an earlier analysis conducted by Xue et al. 2005 by applying both an existing and a new technique for estimating personal ozone exposures of school-age children living in two California communities
- The first modeling analysis employed commonly used methods for estimating exposures using a microenvironmental exposure model that used independent distributions in simulations, fit to observed indoor, outdoor and time-activity data
- observed indoor, outdoor and time-activity data in the new modeling methodology, different variance components derived from the underlying data were put back into the model, so that proper variance-covariance relationships in model inputs maintainated $\mathcal{O}_{\rm ind} = \mathcal{O}_{\rm inst} = \mathcal$

- Intra-personal, inter-personal, seasonal, and area/location related variances predicted by the new model were close to those derived from the original data, while intra-personal variance was overly initiated by the basis method which did not consider proper decomposition of the variance (Table 1)
- New exposure simulation method successfully retains the varinew exposure simination incurous directs stury terms in the vari-ance-covariance structure of different inputs and outputs, and can accurately predict both the annual average and the high percentiles of longitudinal exposures that are important for scientific and regu-latory evaluations (Figures 7 and 8)

Table 1. Percent of Variance Explained by Intra-personal, Inter-personal, Season and Area Terms for Personal Ozone and Outdoor Time

variables	Variance decompostion (%)			
	intra	inter	season	are
observed personal ozone concentration	41.8%	13.3%	44.0%	0.89
predicted personal ozone with observed data	32.6%	10.6%	50.2%	6.79
predicted personal ozone*	91.3%	8.6%	0.1%	0.0%
predicted personal ozone with simulated inputs**	37.7%	18.7%	39.3%	4.39
actual outdoor time near home	59.8%	33.4%	6.8%	0.09

Simulations with decomposition of major variance terms

Figure 7. Measured and Modeled Annual Aver-





Figure 8. Distributions of 95th Percentile Values of 7-Day AveragePersonal Ozone for each child over a Year



DISCUSSION

- sure-related changes over the first 6 years of life Children are exposed to different sources and concentrations of pollutants in various indoor, outdoor and commuting environ-ments
- For modeling long-term exposures to air pollutants, age-specific longitudinal time-activity data and information on key determinants and components (e.g., intra- and inter-personal factors) of the variance of personal exposures are needed
- Probabilistic exposure simulation modeling with decomposition of variance terms is a promising approach that addresses these
- A case study based on data from Southern California, showed A case study based on data from Southern Cantonina, showed that by proper apportionment of the variance-covariance rela-tionships in model inputs, it is possible to accurately predict the central values and the high percentiles of longitudinal personal ozone exposures of children

Burke, J.M., Zufall, M.J and Özkaynak, H. A population exposure model for partic ulate matter: case study results for PM2.5 in Philadelphia, PA, Journal of Exposure Analysis and Environmental Epidemiology, 11: 470-489, 2001

EU 21, ZAUG 1, S.V.,Özkaynak, H.,Spengler, J.D. Parameter evaluation and of ozone exposure assessment using Harvard southern Califo udy data. Journal of the Air & Waste Management Associati

A lthough this work was reviewed by EPA and approved for publication, it may not necessarily reflect official Agency policy.