Modeling Deterioration of Running Emissions

Phil Enns EPA Office of Mobile Sources

Modeling Deterioration of Running Emissions

- ♦ Data Sources
- Models
- ♦ Results

Data Sources

- IM240
 Ohio
 Arizona
 Colorado
 FTP
 - Emission Factors Database
 - Manufacturers
 - API

Ohio IM 240

- 240,000 tests, model years 1975-96, from Dayton (no previous IM program)
- ◆ Fast Pass scores used to simulate 240
- Simulated 240 correlated with Running LA4

Conclusions

- No indication of accelerated deterioration
- Decision: May use Ohio truck data for model years not adequately represented by FTP data

Deterioration of Running LA4 Emissions

- ♦ FTP Data (AAMA, EPA, API)
 - used for
 - * carbureted vs. fuel injected
 - model year groups
 - -1981-82
 - -1983-85
 - -1986-89
 - -1990-93
 - * cars vs. trucks

Fitted Models

♦ Linear


- Emissions = A + B*Miles
- Constrained Linear
 - Emissions = ZML + B*Miles
- Constrained Nonlinear
 - Emissions = ZML + K*(Miles)^B

Proposed Solution


Piecewise function

- Constant emissions at low mileage
- Constrained linear function through low mileage point, connected to...
- Unconstrained linear function for high mileage












MILES (x1000)




MILES (x1000)



MILES (x1000)




MILES (x1000)




DRAFT Oct. 1, 1997


RUNNING LA4 HC: OPEN LOOP AND CARBURETED CARS




RUNNING LA4 HC: FUEL INJECTED CARS



RUNNING LA4 CO: OPEN LOOP AND CARBURETED CARS




RUNNING LA4 CO: FUEL INJECTED CARS



DRAFT October 1, 1997

RUNNING LA4 NOX: OPEN LOOP AND CARBURETED CARS



RUNNING LA4 NOX: FUEL INJECTED CARS

