


Abnormal Development of Retinal Vasculature and Retinal Neovascularization During Oxygen-Induced Ischemic Retinopathy in Endoglin Heterozygous Mice

Nader Sheibani, Ph.D.


*University of Wisconsin
School of Medicine and Public Health
Departments of Ophthalmology and
Visual Sciences, and Pharmacology*

Retinal Vascular Development


P0








P2






P7



P22



Normal Retinal Vascularization Occurs Along the Network of Astrocytes





Blue: PECAM


Red: NG2

Green: GFAP



Oxygen-Induced Ischemic Retinopathy



Abnormal Development of Retinal Vasculature in PECAM-1^{-/-} Mice




Attenuation of Retinal Neovascularization in PECAM-1^{-/-} Mice



Wholemout Staining of Retinal Vessels


WT P8

PECAM-1^{-/-} P8






Endoglin

Increased Expression of Endoglin in Retinal Vasculature during OIR




Wholemout Retinal Trypsin Digests



Retinal Trypsin Digests

WT P42


Eng^{+/-} P42



Wholemout Staining of Retinal Vasculature

WT P7

Eng^{+/-} P7




Endoglin

Arteriovenous Malformations in Eng^{+/-} Mice

WT P7

Eng^{+/-} P7




PECAM-1

Wholemout Staining of Retinal Vasculature

WT P7

Eng^{+/-} P7









Collagen IV

Wholemout Staining of Retinal Vasculature


WT P7

Eng^{+/-} P7



Smooth Muscle Actin

Attenuation of Retinal Neovascularization in Eng^{+/-} Mice



SUMMARY

- Endoglin plays an important role during retinal vascular development and its expression is essential for maturation of retinal blood vessels.
- Endoglin is expressed in retinal vessels and its expression is increased during active neovascularization.
- The ability of Eng^{+/-} mice to undergo retinal neovascularization is abrogated during OIR.
- Endoglin may provide an important target for modulation of angiogenesis.

Acknowledgements

Sheibani Lab

Yongji Wang, PhD

Shoujian Wang, MD

Elizabeth Scheef, MS

Terri DiMaio, BS

Carol Kiekhaefer, MD

Yixin Tang, MS

Robert Wirka, BS

Kirsten Austad, BS

Sources of Support

-NEI/NIH EY013700

-Retina Research Foundation

-Career Development Award from RPB

-ADA and JDRF

Collaborators

Dean Y. Li (University of Utah)

C.M. Sorenson, PhD (University of Wisconsin)

R.L. Engerman, PhD

J.L. Gottlieb, MD