

## 10.0 REFERENCES

- Adgate JL, Barr DB, Clayton CA, Eberly LE, Freeman NCG, Lioy PJ, Needham LL, Pellizzari ED, Quackenboss JJ, Roy A, Sexton K. (2001) Measurement of children's exposure to pesticides: analysis of urinary metabolite levels in a probability-based sample. *Environ Health Perspect* 109:583-590.
- Adgate JL, Kukowski A, Stroebel C, Shubat PJ, Morrell S, Quackenboss JJ, Whitmore RW, Sexton K. (2000) Pesticide storage and use patterns in Minnesota households with children. *J Expo Anal Environ Epidemiol* 10:159-167.
- Agency for Toxic Substances and Disease Registry (ATSDR). (2003) Toxicological Profile for Pyrethrins and Pyrethroids. US Department of Health and Human Services, Agency for Toxic Substances and Disease Registry. Atlanta, GA. <<http://www.atsdr.cdc.gov/toxprofiles/tp155.pdf>>.
- American Society for Testing and Materials (ASTM). (1993) Standard Practice for Collection of Floor Dust for Chemical Analysis (D 5438-94). Philadelphia, PA: ASTM.
- Aprea C, Colosio C, Mammone T, Minoia C, Maroni M. (2002) Biological monitoring of pesticide exposure: a review of analytical methods. *J Chromatogr B Analyt Technol Biomed Life Sci* 769(2):191-219.
- Baker SE, Barr DB, Driskell WJ, Beeson MD, Needham LL. (2000) Quantification of selected pesticide metabolites in human urine using isotope dilution high-performance liquid chromatography/tandem mass spectrometry. *J Expo Anal Environ Epidemiol* 10(6 Pt 2):789-98.
- Baker SE, Olsson AO, Barr DB. (2004) Isotope dilution high-performance liquid chromatography–tandem mass spectrometry method for quantifying urinary metabolites of synthetic pyrethroid insecticides. *Arch Environ Contam Toxicol* 46:281–288.
- Barr DB, Bravo R, Weerasekera G, Caltabiano LM, Whitehead RD Jr, Olsson AO, Caudill SP, Schober SE, Pirkle JL, Sampson EJ, Jackson RJ, Needham LL. (2004) Concentrations of dialkyl phosphate metabolites of organophosphorus pesticides in the U.S. population. *Environ Health Perspect* 112(2):186-200.
- Bartelt N, Hubbell J. (1987) Percutaneous absorption of topically applied <sup>14</sup>C-permethrin in volunteers: Final medical report. Burroughs Wellcome Co. Cited in: National Research Council. 1994. *Health Effects of Permethrin-Impregnated Army Battle-Dress Uniforms*. Washington, DC: National Academy Press.
- Bass JK, Ortega L, Rosales C, Petersen NJ, Philen RM. (2001) What's being used at home: a household pesticide survey. *Pan Am J Public Hlth* 9(3):138-144.
- Beeson MD, Driskell WJ, Barr DB. (1999) Isotope dilution high-performance liquid chromatography/tandem mass spectrometry method for quantifying urinary metabolites of atrazine, malathion, and 2,4-dichlorophenoxyacetic acid. *Anal Chem* 71(16):3526-30.
- Bennett DH, Furtaw Jr EJ. (2004) Fugacity-based indoor residential pesticide fate model. *Environ Sci Technol* 38(7):2142-52.
- Berry MR, Cohen Hubal EA, Fortmann RC, Melnyk LJ, Sheldon LS, Stout DM, Tulve NS, Whitaker DA. (2001) Draft Protocol for Measuring Children's Non-Occupational Exposure to Pesticides by all Relevant Pathways. U.S. Environmental Protection Agency, Office of Research and Development, Research Triangle Park, NC. EPA/600/R-03/026.
- Blancato JN, Power FW, Brown RN, Dary CC. (2004) Exposure Related Dose Estimating Model (ERDEM) for Assessing Human Exposure and Dose. U.S. Environmental Protection Agency, Office of Research and Development, Washington, DC. EPA/600/R-04/060.
- Bradman A, Whitaker D, Quiros L, Castorina R, Henn BC, Nishioka M, Morgan J, Barr DB, Harnly M, Brisbin JA, Sheldon LS, McKone TE, Eskenazi B. (2006) Pesticides and their metabolites in the homes and urine of farmworker children living in the Salinas Valley, CA. *J Expo Sci Environ Epidemiol* [May 31 Epub ahead of print].
- Brouwer DH, Kroese R, Van Hemmen JJ. (1999) Transfer of contaminants from surface to hands: experimental assessment of linearity of the exposure process, adherence to the skin, and area exposed during fixed

- pressure and repeated contact with surfaces contaminated with a powder. *Appl Occup Environ Hyg* 14(4):231-9.
- Butte W, Heinzel B. (2002) Pollutants in house dust as indicators of indoor contamination. *Rev Environ Contam Toxicol* 175:1-46.
- Byrne SL, Saunders DG, Sherdut BA. (1998) Potential chlorpyrifos exposure following standard crack and crevice treatment. *Environ Health Perspect* 106(11):725-31.
- Camann DE, Majumdar TK, Harding HJ, Ellenson WD, Lewis RG. (1996) Transfer efficiency of pesticides from carpets to saliva-moistened hands. In: Measurement of toxic and related air pollutants; Proceedings of an International Specialty Conference, Research Triangle Park, NC, May 7-9. VIP-64. Pittsburgh, PA: Air and Waste Management Association. Pp. 532-40.
- Carlton EJ, Moats HL, Feinberg M, Shepard P, Garfinkel R, Whyatt R, Evans D. (2004) Pesticide sales in low-income, minority neighborhoods. *J Community Hlth* 29(3):231-244.
- Clayton C, Pellizzari E, Whitmore RW, Quackenboss JJ. (2003) Distributions, associations, and partial aggregate exposure of pesticides and polynuclear aromatic hydrocarbons in the Minnesota Children's Pesticide Exposure Study (MNCPES). *J Expo Anal Environ Epidemiol* 13(2):100-111.
- Clothier JM. (2000) Dermal Transfer Efficiency of Pesticides from New, Vinyl Sheet Flooring to Dry and Wetted Palms. U.S. Environmental Protection Agency, Office of Research and Development, Washington, D.C. EPA/600/R-00/029.
- Coble J, Arbuckle T, Lee W, Alavanja M, Dosemeci M. (2005) The validation of a pesticide exposure algorithm using biological monitoring results. *J Occup Environ Hyg* 2(3):194-201.
- Cohen Hubal EA, Egeghy PP, Leovic KW, Akland GG. (2006) Measuring potential dermal transfer of a pesticide to children in a child care center. *Environ Health Perspect* 114(2):264-9.
- Cohen Hubal EA, Sheldon LS, Burke JM, McCurdy TR, Berry MR, Rigas ML, Zartarian VG, Freeman NCG. (2000a) Children's exposure assessment: a review of factors influencing children's exposure, and the data available to characterize and assess that exposure. *Environ Health Perspect* 108(6):475-486.
- Cohen Hubal EA, Sheldon LS, Zufall MJ, Burke JM, and Thomas K. (2000b) The challenge of assessing children's residential exposure to pesticides. *J Expo Anal Environ Epidemiol* 10:638-649.
- Colt JS, Zahm SH, Camann DE, Hartge P. (1998) Comparison of pesticides and other compounds in carpet dust samples collected from used vacuum cleaner bags and from a high-volume surface sampler. *Environ Health Perspect* 106(11):721-724.
- Curl CL, Fenske RA, Kissel JC, Shirai JH, Moate TF, Griffith W, Coronado G, Thompson B. (2002) Evaluation of take-home organophosphorus pesticide exposure among agricultural workers and their children. *Environ Health Perspect* 110(12):A787-92.
- Curwin B, Sanderson W, Reynolds S, Hein M, Alavanja M. (2002) Pesticide use and practices in an Iowa farm family pesticide exposure study. *J Agr Safe Hlth* 8(4):423-433.
- Davis JR, Brownson RC, Garcia R. (1992) Family pesticide use in the home, garden, orchard, and yard. *Arch Environ Contam Toxicol* 22:260-266.
- Droz PO, Wu MM, Cumberland WG, Berode M. (1989) Variability in biological monitoring of solvent exposure. I. Development of a population physiological model. *Br J Ind Med* 46(7):447-60.
- Eadsforth CV, Baldwin MK. (1983) Human dose-excretion studies with the pyrethroid insecticide, cypermethrin. *Xenobiotica* 13(2):67-72.
- Eadsforth CV, Bragt PC, van Sittert NJ. (1988) Human dose-excretion studies with pyrethroid insecticides cypermethrin and alpha-cypermethrin: relevance for biological monitoring. *Xenobiotica* 18(5):603-14.
- Edwards RD, Lioy PJ. (1999) The EL sampler: a press sampler for the quantitative estimation of dermal exposure to pesticides in house dust. *J Expo Anal Environ Epidemiol* 9:521-529.

- Feldmann RJ, Maibach HI. (1974) Percutaneous penetration of some pesticides and herbicides in man. *Toxicol Appl Pharmacol* 28(1):126-32.
- Fenske, RA. (1993) Dermal exposure assessment techniques. *Ann Occup Hyg* 37(6):687-706.
- Fenske RA, Black KG, Elkner KP, Lee CL, Methner MM, Soto R. (1990) Potential exposure and health risks of infants following indoor residential pesticide applications. *Am J Public Health* 80(6):689-693.
- Fenske RA, Kissel JC, Lu C, Kalman DA, Simcox NJ, Allen EH, Keifer MC. (2000) Biologically based pesticide dose estimates for children in an agricultural community. *Environ Health Perspect* 108:515-520.
- Fortune CR. (1997) Round-Robin Testing of Methods for Collecting Dislodgeable Residues from Carpets. Final Report. U.S. Environmental Protection Agency, Office of Research and Development, Research Triangle Park. EPA/600/R-97/119.
- Freeman N, Hore P, Black K, Jimenez M, Sheldon L, Tulve N, and Lioy P. (2005) Contributions of children's activities to pesticide hand loadings following residential pesticide applications. *J Expo Anal Environ Epidemiol* 15:81-88.
- Freeman NCG, Jimenez M, Reed KJ, Gurunathan S, Edwards RD, Roy A, Adgate JL, Pellizzari ED, Quackenboss JJ, Sexton K, Lioy PJ. (2001) Quantitative analysis of children's microactivity patterns: The Minnesota Children's Pesticide Exposure Study. *J Expo Anal Environ Epidemiol* 11(6):501-509.
- Freeman NCG, Shalat SL, Black K, Jimenez M, Kirby C, Donnelly KC, Calvin A, Ramirez J. (2004) Seasonal pesticide use in a rural community on the US/Mexico border. *J Expo Anal Environ Epidemiol* 14:473-478.
- Freeman NCG, Wainman T, Lioy PJ. (1996) Field testing of the LWW dust sampler and association of observed household factors with dust loadings. *Appl Occup Environ Hyg* 11:476-483.
- Garfitt SJ, Jones K, Mason HJ, Cocker, J. (2002) Exposure to the organophosphate diazinon: data from a human volunteer study with oral and dermal doses. *Toxicol Lett* 134(1-3):105-13.
- Griffin P, Mason H, Heywood K, Cocker J. (1999) Oral and dermal absorption of chlorpyrifos: a human volunteer study. *Occup Environ Med* 56(1):10-3.
- Gurunathan S, Robson M, Freeman N, Buckley B, Roy A, Meyer R, Bukowski J, Lioy PJ. (1998) Accumulation of chlorpyrifos on residential surfaces and toys accessible to children. *Environ Health Perspect* 106(1):9-16.
- Hill RH Jr, Head SL, Baker S, Gregg M, Shealy DB, Bailey SL, Williams CC, Sampson EJ, Needham LL. (1995) Pesticide residues in urine of adults living in the United States: reference range concentrations. *Environ Res* 71(2):99-108.
- Hore P. (2003) Pesticide Accumulation Patterns for Child Accessible Surfaces and Objects and Urinary Metabolite Excretion by Children for Two Weeks after a Professional Crack and Crevice Application. Ph.D. Dissertation, The State University of New Jersey, Rutgers, New Brunswick, New Jersey.
- Hore P, Robson M, Freeman N, Zhang J, Wartenberg D, Ozkaynak H, Tulve N, Sheldon L, Needham L, Barr D, Lioy PJ. (2005) Chlorpyrifos accumulation patterns for child-accessible surfaces and objects and urinary metabolite excretion by children for 2 weeks after crack-and-crevice application. *Environ Health Perspect* 113(2):211-9.
- Hornung RW, Reed LD. (1990) Estimation of average concentration in the presence of nondetectable values. *Appl Occup Environ Hyg* 5(1):46-51.
- Hsu JP, Camann DE, Schattenberg H, Wheeler B, Villolobos K, Kyle M, Quardeer S, Lewis RG. (1990) New dermal exposure sampling technique. In: Proceedings of 1990 EPA/AWMA International Symposium on Measurement of Toxic and Related Air Pollutants. VIP-17. Pittsburgh, PA: Air & Waste Management Association 489-497.
- Hu YA, Akland GG, Pellizzari ED, Berry MR, Melnyk LJ. (2004) Use of pharmacokinetic modeling to design studies for pathway-specific exposure model evaluation. *Environ Health Perspect* 112(17):1697-1703.
- Hulka BS, Margolin BH. (1992) Methodological issues in epidemiologic studies using biological markers. *Am J Epidemiol* 135(2):200-9.

- Karalliedde L, Feldman RJ, Henderson LO, Marrs T. (2001) Organophosphates and Health. London: Imperial College Press.
- Kissel JC, Richter KY, Fenske RA. (1996) Factors affecting soil adherence to skin in hand-press trials. *Bull Environ Contam Toxicol* 56(5):722-8.
- Krieger RI, Bernard CE, Dinoff TM, Fell L, Osimitz TG, Ross JH, Ongsinthusak, T. (2000) Biomonitoring and whole body cotton dosimetry to estimate potential human dermal exposure to semivolatile chemicals. *J Expo Anal Environ Epidemiol* 10(1):50-7.
- Krieger RI, Bernard CE, Dinoff TM, Ross JH, Williams RL. (2001) Biomonitoring of persons exposed to insecticides used in residences. *Ann Occup Hyg* 45(Suppl 1):S143-53.
- LaGoy PK. (1987) Estimated soil ingestion rates for use in risk assessment. *Risk Anal* 7(3):355-9.
- Landrigan PJ, Claudio L, Markowitz SB, Berkowitz GS, Brenner BL, Romero H, Wetmur JG, Matte TD, Gore AC, Godbold JH, Wolff MS. (1999) Pesticides and inner-city children: exposures, risks, and prevention. *Environ Health Perspect* 107 (Suppl 3):431-7.
- Lanphear BP, Emond M, Jacobs DE, Weitzman M, Tanner M, Winter NL, Yakir B, Eberly S. (1995) A side-by-side comparison of dust collection methods for sampling lead-contaminated house dust. *Environ Res* 68(2):114-23.
- Laskowski DA. (2002) Physical and chemical properties of pyrethroids. *Rev Environ Contam Toxicol* 174:49-170.
- Leidy RB, Wright CG, Dupree Jr HE. (1996) Distribution of chlorpyrifos in air and on surfaces in rooms after a crack and crevice application. *Environ Monit Assess* 42:253-263.
- Leng G, Leng A, Kuhn KH, Lewalter J, Pauluhn J. (1997) Human dose-excretion studies with the pyrethroid insecticide cyfluthrin: urinary metabolite profile following inhalation. *Xenobiotica* 27(12):1273-83.
- Lewis RG. (2001) Residential post-application pesticide exposure monitoring. In: Occupational and Incidental Residential Exposure Assessment. Worgan JP, Franklin CA (eds). Sussex: John Wiley & Sons, Ltd.
- Lewis RG, Fortune CR, Blanchard FT, Camann DE. (2001) Movement and deposition of two organophosphorus pesticides within a residence after interior and exterior applications. *J Air Waste Manag Assoc* 51(3):339-51.
- Lewis RG, Fortmann RC, Camann DE. (1994) Evaluation of methods for monitoring the potential exposure of small children to pesticides in the residential environment. *Arch Environ Contam Toxicol* 26(1):37-46.
- Lewis RG, Fortune CR, Willis RD, Camann DE, Antley JT. (1999) Distribution of pesticides and polycyclic aromatic hydrocarbons in house dust as a function of particle size. *Environ Health Perspect* 107(9):721-6.
- Lioy PJ, Edwards RD, Freeman N, Gurunathan S, Pellizzari E, Adgate JL, Quackenboss J, Sexton K. (2000) House dust levels of selected insecticides and a herbicide measured by the EL and LWW samplers and comparisons to hand rinses and urine metabolites. *J Expo Anal Environ Epidemiol* 10:327-340.
- Lioy PJ, Wainman T, Weisel C. (1993) A wipe sampler for the quantitative measurement of dust on smooth surfaces: laboratory performance studies. *J Expo Anal Environ Epidemiol* 3:315-330.
- Lowry LK. (1986) Biological exposure index as a complement to the TLV. *J Occup Med* 28(8):578-82.
- Lu C, Fenske RA. (1998) Air and surface chlorpyrifos residues following residential broadcast and aerosol pesticide applications. *Environ Sci Technol* 32(10):1386-1390.
- Lu C, Fenske RA. (1999) Dermal transfer of chlorpyrifos residues from residential surfaces: comparison of hand press, hand drag, wipe, and polyurethane foam roller measurements after broadcast and aerosol pesticide applications. *Environ Health Perspect* 107(6):463-7.
- Lu C, Kedan G, Fisker-Andersen J, Kissel JC, Fenske RA. (2004) Multipathway organophosphorus pesticide exposures of preschool children living in agricultural and nonagricultural communities. *Environ Res* 96(3):283-9.

- Maibach HI, Feldman RJ, Milby TH, Serat WF. (1971) Regional variation in percutaneous penetration in man. Pesticides. Arch Environ Health 23(3):208-11.
- Meuling WJ, Ravensberg LC, Roza L, van Hemmen JJ. (2005) Dermal absorption of chlorpyrifos in human volunteers. Int Arch Occup Environ Health 78(1):44-50.
- Morgan MK, Sheldon LS, Croghan CW, Jones PA, Robertson GL, Chuang JC, Wilson NK, Lyu CW. (2005) Exposures of preschool children to chlorpyrifos and its degradation product 3,5,6-trichloro-2-pyridinol in their everyday environments. J Expo Anal Environ Epidemiol 15:297–309.
- Morgan MK, Sheldon LS, Croghan CW, Chuang JC, Lordo RA, Wilson NK, Lyu C, Brinkman M, Morse N, Chou YL, Hamilton C, Finegold JK, Hand K, Gordon SM. (2004) A Pilot Study of Children's Total Exposure to Persistent Pesticides and Other Persistent Organic Pollutants (CTEPP). U.S. Environmental Protection Agency, Office of Research and Development, Washington, DC. EPA/600/R-04/193.
- Morgan MK, Stout II DM. (2001) Feasibility study of the potential for human exposure to pet-borne diazinon residues following lawn applications. Bull Environ Contam Toxicol 66:295-300.
- Ness SA. (1994) Surface and Dermal Monitoring for Toxic Exposures. New York: John Wiley and Sons.
- Nolan RJ, Rick DL, Freshour NL, Saunders JH. (1984) Chlorpyrifos: pharmacokinetics in human volunteers. Toxicol Appl Pharmacol 73(1):8-15.
- National Research Council (NRC). (1993) Pesticides in the Diets of Infants and Children. National Research Council, Commission on Life Sciences. Washington, DC: National Academy Press. 386 pp.
- Oros DR, Werner I. (2005) Pyrethroid Insecticides: An Analysis of Use Patterns, Distributions, Potential Toxicity and Fate in the Sacramento-San Joaquin Delta and Central Valley. White Paper for the Interagency Ecological Program. SFEI Contribution 415. San Francisco Estuary Institute, Oakland, CA, 113 pp.
- O'Rourke MK, Lizardi PS, Rogan SP, Freeman NC, Aguirre A, Saint CG. (2000) Pesticide exposure and creatinine variation among young children. J Expo Anal Environ Epidemiol 10(6 Pt2):672-81.
- Ozkaynak H. (2005) Modeling children's exposures to pesticides: Issues and challenges. EPA's Workshop on the Analysis of Children's Measurements Data. Research Triangle Park, NC; September 27, 2005.
- Reed K, Jimenez M, Freeman N, Lioy P. (1999) Quantification of children's hand and mouthing activities through a videotaping methodology. J Expo Anal Environ Epidemiol 9(5):513-20.
- Rigas ML, Okino MS, Quackenboss JJ. (2001) Use of a pharmacokinetic model to assess chlorpyrifos exposure and dose in children, based on urinary biomarker measurements. Toxicol Sci 61:374-381.
- Roberts JW, Budd WT, Ruby MG, Camann DE, Fortmann RC, Sheldon LS, Lewis RG. (1991) Development and field testing of a high volume sampler for pesticides and toxics in dusts. J Expo Anal Environ Epidemiol 1(2):143-155.
- Rodes CE, Newsome JR, Vanderpool RW, Antley JT, Lewis RG. (2001) Experimental methodologies and preliminary transfer factor data for estimation of dermal exposures to particles. J Expo Anal Environ Epidemiol 11(2):123-39.
- Rohrer CA, Hieber T, Melnyk LJ, Berry MR. (2003) Transfer efficiencies of pesticides to household flooring surfaces. J Expo Anal Environ Epidemiol 13:454 - 464.
- Rust MK, Owens JM, Reierson DA (eds.). (1995) Understanding and Controlling the German Cockroach. New York: Oxford University Press.
- Savage EP, Keefe TJ. (1981) Household pesticide usage in the US. Arch Environ Health 36:304- 309.
- Sexton K, Adgate JL, Eberly LE, Clayton CA, Whitmore RW, Pellizzari ED, Lioy PJ, Quackenboss JJ. (2003) Predicting children's short-term exposure to pesticides: results of a questionnaire screening approach. Environ Health Perspect 110:123–128.
- Shah PV, Fisher HL, Sumler MR, Monroe RJ, Chernoff N, Hall LL. (1987) Comparison of the penetration of 14 pesticides through the skin of young and adult rats. J Toxicol Environ Health 21(3):353-66.

- Stout II DM, Mason M. (2003) The distribution of chlorpyrifos following a crack and crevice type application in the US EPA indoor air quality research house. *Atmos Environ* 37:5539-5549.
- Tomlin C. (1994) The Pesticide Manual: Incorporating the Agrochemicals Handbook, 10th Edition. Crop Protection Publication, United Kingdom.
- Tulve NS, Driver J, Egeghy PP, Evans J, Fortmann RC, Kissel J, McMillan N, Melnyk LJ, Morgan MK, Starr J, Stout II DM, Strynar M. (2006) The U.S. EPA National Exposure Research Laboratory's (NERL's) Workshop on the Analysis of Children's Measurement Data: Workshop Summary Report. U.S. Environmental Protection Agency, Office of Research and Development, Washington, DC. EPA 600/R-06/026
- Tulve NS, Jones PA, Nishioka M, Fortmann RC, Croghan CW, Zhou JY, Fraser A, Cave C, Friedman W. (2006) Pesticide Measurements from the First National Environmental Health Survey of Child Care Centers Using a Multi-Residue GC/MS Analysis Method. *Environ Sci Technol* 40(20):6269-74.
- Tulve NS, Suggs JC, McCurdy T, Cohen Hubal EA, Moya J. (2002) Frequency of mouthing behavior in young children. *J Expo Anal Environ Epidemiol* 12:259-264.
- U.S. Consumer Product Safety Commission (US CPSC). (1999) CPSC Staff Study of Safety Hazards in Child Care Settings. U.S. Consumer Product Safety Commission, Washington, DC.
- US EPA. (1980) National Household Pesticide Usage Study, 1976-1977. Final Report. Office of Pesticide Programs, Washington, D.C.
- US EPA. (1992) National Home and Garden Pesticide Use Survey. No. RTI/5100/17-01f. 400pp. Office of Pesticides and Toxic Substances. Washington, D.C.
- US EPA. (2001) Draft Protocol for Measuring Children's Non-Occupational Exposure to Pesticides by all Relevant Pathways. Office of Research and Development. Research Triangle Park, NC. EPA/600/R-03/026.
- US EPA. (2002) Child-specific exposure factors handbook. National Center for Environmental Assessment, Washington, DC; EPA/600/P-00/002B. Available from: National Information Service, Springfield, VA; PB2003-101678 and <<http://www.epa.gov/ncea>>.
- US EPA. (2004) Pesticide Industry Sales and Usage: 2000 and 2001 Market Estimates. Office of Pesticide Programs, Washington, D.C.
- Wessels D, Barr DB, Mendola P. (2003) Use of biomarkers to indicate exposure of children to organophosphate pesticides: implications for a longitudinal study of children's environmental health. *Environ Health Perspect* 111(16):1939-46.
- Wester RC, Bucks DA, Maibach HI. (1994) Human in vivo percutaneous absorption of pyrethrin and piperonyl butoxide. *Food Chem Toxicol* 32(1):51-3.
- Whitmore R, Clayton A, Akland G. (2003) California Portable Classrooms Study. Phase II: Main Study. Final Report. Vol II. Research Triangle Park, NC: RTI International. Contract No. 00-317.
- Whitmore RW, Immerman FW, Camann DE, Bond AE, Lewis RG, Schaum JL. (1994) Non-occupational exposures to pesticides for residents of two U.S. cities. *Arch Environ Contam Toxicol* 26:47-59.
- Whyatt RM, Camann DE, Kinney PL, Reyes A, Ramirez J, Dietrich J, Diaz D, Holmes D, Perera FP. (2002) Residential pesticide use during pregnancy among a cohort of urban minority women. *Environ Health Perspect* 110(5):A256-7.
- Wilson NK, Chuang JC, Lyu C, Menton R, Morgan MK. (2003) Aggregate exposures of nine preschool children to persistent organic pollutants at day care and at home. *J Expo Anal Environ Epidemiol* 13(3):187-202.
- Woolen BH, Marsh JR, Laird WJ, Lesser JE. (1992) The metabolism of cypermethrin in man::differences in urinary metabolite profiles following oral and dermal administration. *Xenobiotica* 22(8):983-91.
- World Health Organization. (1989) Environmental Health Criteria 82: Cypermethrin. Geneva.
- Wright CG, Leidy RB. (1978) Chlorpyrifos residues in air after application to crevices in rooms. *Bull Environ Contam Toxicol* 20:340-344.

- Zartarian V, Bahadori T, McKone T. (2005) Adoption of an official ISEA glossary. *J Expo Anal Environ Epidemiol* 15(1):1-5.
- Zartarian V, Ferguson A, and Leckie J. (1998) Quantified dermal activity data from a four-child pilot field study. *J Expo Anal Environ Epidemiol* 8(1):109.
- Zartarian VG, Ferguson AC, Ong CG, Leckie JO. (1997) Quantifying videotaped activity patterns: video translation software and training methodologies. *J Expo Anal Environ Epidemiol* 7:535-542.
- Zartarian VG, Ozkaynak H, Burke JM, Zufall MJ, Rigas ML, Furtaw EJ Jr. (2000) A modeling framework for estimating children's residential exposure and dose to chlorpyrifos via dermal residue contact and nondietary ingestion. *Environ Health Perspect* 108(6):505-14.