




REVIEW OF THE NATIONAL AMBIENT AIR QUALITY STANDARDS FOR SULFUR OXIDES:

UPDATED ASSESSMENT OF SCIENTIFIC AND TECHNICAL INFORMATION

ADDENDUM TO THE 1982 OAQPS STAFF PAPER



Strategies and Air Standards Division
Office of Air Quality Planning and Standards
U.S. Environmental Protection Agency
Research Triangle Park, N.C. 27711

December 1986


Cover Illustration. Individual dose-response curves of percentage (normalized) changes in airway resistance (SRaw), adjusted for response to clean air exposure, as a function of SO₂ exposure for asthmatic subjects. (A) 6 subjects with response at < 0.5 ppm, (B) 9 subjects with response between 0.5 and 1.0 ppm, (C) 8 subjects with response between 1.0 and 2.0 ppm, and (D) 4 subjects with response at > 2.0 ppm SO₂. Data are not included for 0.0 ppm since they were used to adjust for exercise-induced bronchoconstriction. The interrupted horizontal line represents 100% increase in SRaw and the SO₂ concentration corresponding to its point of intercept with each subject's curve was defined as PC(SO₂) (Horstman et al. 1986). The substantial variability in sensitivity to peak SO₂ exposures among asthmatics is an important consideration in the review of the sulfur oxides standards.

REVIEW OF THE NATIONAL AMBIENT AIR QUALITY STANDARDS FOR SULFUR OXIDES:
UPDATED ASSESSMENT OF SCIENTIFIC AND TECHNICAL INFORMATION

ADDENDUM TO THE 1982 OAQPS STAFF PAPER

Strategies and Air Standards Division
Office of Air Quality Planning and Standards
U.S. Environmental Protection Agency
Research Triangle Park, N.C. 27711

December 1986



Acknowledgements

This staff paper is the product of the Office of Air Quality Planning and Standards (OAQPS). The principal authors include Jeff Cohen, John Bachmann, and Henry Thomas. The report incorporates comments from OAQPS, the Office of Research and Development, the Office of Policy, Planning, and Evaluation, and the Office of General Counsel within EPA and was formally reviewed by the Clean Air Scientific Advisory Committee.

Helpful comments and suggestions were also submitted by independent scientists, by officials from the California Air Resources Board, and by environmental and industry groups including the Natural Resources Defense Council, the Environmental Defense Fund, the American Mining Congress, and several member companies, the Utility Air Regulatory Group, the Air Pollution Institute, Standard Oil Company, Phelps Dodge Corporation, and Kennecott Company.

The authors wish to thank Teresa Clemons and Tricia Holland for word processing, and Dick Atherton for graphics assistance.




TABLE OF CONTENTS

	<u>Page</u>
List of Figures	iv
List of Tables	iv
Executive Summary	v
I. Introduction	1
A. Purpose	1
B. Background	1
C. Approach	3
II. Air Quality Considerations	5
A. Peak to Mean Ratios	5
B. Factors Affecting Assessment of Peak Air Quality Levels . .	7
III. Critical Elements in the Review of the Primary Standards . . .	9
A. Mechanisms	9
B. Concentration/Response Information	12
IV. Factors to be Considered in Selecting Primary Standards for Sulfur Oxides	33
A. Levels and Averaging Times of the Standards	33
B. Analysis of Relative Protection Afforded by Alternative Standards	50
C. Summary of Staff Conclusions and Recommendations	58
Appendix A. Analysis of Dose Response Relationships from Controlled SO ₂ Exposure Studies on Asthmatics	A-1
Appendix B. CASAC Closure Memorandum	B-1
References	

LIST OF TABLES

<u>Number</u>	<u>Title</u>	<u>Page</u>
1	Updated Staff Assessment of Key Controlled Human Studies	vii
2	Updated Staff Assessment of Short-Term Epidemiological Studies	x
3-1	Summary of Recent (1982-86) Epidemiological Studies Providing Most Useful Concentration Response Information for Short-Term SO ₂ Exposures	24
3-2	Summary of Recent (1982-86) Epidemiological Studies Providing Most Useful Concentration Response Information on Long-Term Exposures to SO ₂	30
4-1	Updated Staff Assessment of Key Controlled Human Studies	35
4-2	Updated Staff Assessment of Short-Term Epidemiological Studies	45
A-1	Summary of Data from SO ₂ Controlled Asthmatic Studies Used in Dose-Response Analysis	A-2

LIST OF FIGURES

3-1	Combined SO ₂ Dose Response Relationships	19
3-2	Distribution of Individual Airway Sensitivity to SO ₂	20
4-1	Gradiation of Physiological Responses to SO ₂	38
4-2	Expected Percentage of Asthmatics with One Exposure of Concern Per Year. 1) Current Emissions 2) Current NAAQS	55
4-3	Expected Percentage of Asthmatics with One Exposure of Concern Per Year. Alternative One Hour Standards	57

EXECUTIVE SUMMARY

This paper evaluates and interprets the updated scientific and technical information that the EPA staff believes is most relevant to the review of primary (health) national ambient air quality standards (NAAQS) for sulfur oxides* and represents an update of the 1982 sulfur oxides staff paper. This paper assesses what the staff believes should be considered in selecting appropriate averaging times and levels for the primary sulfur oxides standards, updating and supplementing previous staff conclusions and recommendations in these areas to incorporate more recent information. The assessment in this staff paper addendum is intended to help bridge the gap between the scientific review contained in the EPA criteria document addendum "Second Addendum Air Quality Criteria for Particulate Matter and Sulfur Oxides (1982): Assessment of Newly Available Health Effects Information" and the judgments required of the Administrator in setting ambient standards for sulfur oxides. The staff paper and this addendum are, therefore, an important element in the standards review process and provide an opportunity for public comment on proposed staff recommendations before they are presented to the Administrator. The focus of this paper is on sulfur dioxide (SO₂), alone and in combination with other pollutants.

SO₂ is a rapidly diffusing reactive gas that is quite soluble in water. It is emitted principally from combustion or processing of sulfur-containing fossil fuels and ores. SO₂ occurs in the atmosphere

*The current standards for sulfur dioxide (SO₂) are: primary, 0.03 ppm (80 µg/m³) annual arithmetic mean and 0.14 ppm (365 µg/m³) 24-hour average not to be exceeded more than once per year; and, secondary, 0.5 ppm (1300 µg/m³) 3-hour average not to be exceeded more than once per year.

with a variety of particles and other gases, and undergoes chemical and physical interactions with them forming sulfates and other transformation products.

Because much of the recently available health effects information on SO₂ is related to short-term exposures, the staff paid particular attention to updating information on short-term peak concentrations. The staff found that:

1) Maximum 5 minute to hourly SO₂ concentrations are found near major point sources. The newer information tends to support earlier conclusions that near such sources, the 5 to 10 minute peak SO₂ concentration is likely to be within a factor of 1.4 to 2.4 times the hourly average. Maximum peak to mean ratios can be higher.

2) Short duration peaks (less than 30 seconds to 2 minutes) in excess of 0.5 ppm appear likely to occur near numerous smaller sources of SO₂. None of the recently published assessments of the health effects of SO₂ has addressed exposures of such limited duration. Due to limitations of the monitoring instruments, it is not presently possible to assess the extent to which such peaks may be occurring in particular urban locations.

Updated Assessment of the Primary Standards

Conclusions and recommendations based on the updated staff assessment of the information in the criteria document addendum are summarized below.

1) The present staff assessment of the more recent studies reinforce the earlier conclusion reached in the 1982 staff assessment that the most striking acute response to SO₂ is reflex bronchoconstriction, or airway narrowing, in exercising asthmatics and others with hyperreactive airways.

2) a) The updated staff assessment of key controlled human studies of peak (minutes to an hour) SO₂ exposures is summarized in Table 1. Both

SO ₂ Concentration (5-60 minutes)	Observed Effects ¹	Comments/Implications
1-2 ppm	Substantial changes in 8 of 12 subjects (Δ SRaw 100-600%) exposed to 2 ppm. At 1 ppm, functional changes (Δ SRaw 170-200%), symptoms in free breathing asthmatics at moderate exercise ²	Effects range from moderate to incapacitating for some individuals. At 2 ppm, 80% of mild asthmatics could experience at least a doubling of SRaw. Some might not tolerate exposure at moderate exercise. Approx. 60% at 1 ppm could experience at least a doubling of SRaw. ³ Some asthmatic mouth breathers have significant bronchoconstriction at 2 ppm even at light activity.
0.6-0.75 ppm	Functional changes (Δ SRaw 120-260%), symptoms in free breathing asthmatics at light-moderate exercise ⁴	Effects indicative of clinical significance; on average, changes were mild to moderate although severe for some individuals; 25-50% of mild, free-breathing asthmatics at moderate exercise could experience at least a doubling of airway resistance. ³
0.5 ppm	Significant functional changes (Δ SRaw 50-100%), symptoms in free breathing asthmatics at moderate, but not at light exercise. ⁵ At heavy exercise, Δ SRaw 220-240%. ⁶	On average, mild responses at moderate or higher exercise, symptoms possibly of clinical significance; severe responses for some individuals. About 20-25% could experience at least a doubling in airway resistance.
0.4 ppm	Functional changes (Δ SRaw 70%), symptoms in free breathing asthmatics at moderate-heavy exercise ⁷	Lowest level of clinically significant response for some free breathers. Approx. 10% of mild, free breathing asthmatics could experience a doubling in airway resistance. ³
0.1-0.3 ppm	No effects in free breathing asthmatics at light exercise. Slight but not significant functional changes in free-breathing subjects at moderate-heavy exercise (0.25 ppm) ⁶ , but not at lower levels. ⁷	Significant effects unlikely at moderate exercise. Effects of SO ₂ indistinguishable at heavy exercise. Possibility of more significant responses in small percentage of sensitive asthmatics at 0.28 ppm. ³

¹Specific Airway Resistance (SRaw) is the lung function measure most often reported in SO₂ studies. Unless otherwise noted, (Δ SRaw %) reflects group mean increase over clean air control at rest. Light, moderate, heavy exercise refers to ventilation rates approximating < 35 L/min, 40-45 L/min, and ≥ 50 L/min, respectively. Effects reflect results from range of moderate temperature/humidity conditions (i.e., 7-26°C, 36-90% RH). Studies at 0.5-0.6 ppm indicate that exercise-induced bronchoconstriction associated with cold and/or dry air exacerbates response to SO₂ while warm, humid air mitigates asthmatic responses relative to moderate conditions.

²Schacter et al. (1984); Roger et al. (1985); Horstman et al. (1986).

³Horstman et al., (1986).

⁴Hackney et al. (1984); Schacter et al. (1984); Linn et al. (1983a,b, 1984a,b,c, 1985a).

⁵Kirkpatrick et al. (1982); Linn et al. (1984b); Roger et al. (1985); Schacter et al. (1984).

⁶Bethel et al. (1983a,b; 1985).

⁷Linn et al. (1983b, 1984a).

recently published studies and those assessed in the 1982 staff paper are included. The table focuses on those studies involving free breathing (chamber) or facemask exposures, which provide the closest approximation of natural breathing. After account is made for differences in ventilation rates and oral/nasal breathing patterns, consistent results are derived from the various studies including even those that used mouthpiece exposures. The major effects observed in these studies are increases in airway resistance and decreases in other functional measures indicative of significant bronchoconstriction in sensitive asthmatic or atopic subjects. At 0.4 ppm SO₂, changes in functional measures are accompanied by mild increases in perceptible symptoms such as wheezing, chest tightness, and coughing. At higher concentrations, effects are more pronounced and the fraction of asthmatic subjects who respond increase, with clearer indications of clinically or physiologically significant effects at 0.6-0.75 ppm and above.

b) Significant bronchconstriction has been observed in asthmatics after 5-10 minutes of exposure and usually diminishes within one hour once either exposure or exercise alone is discontinued. Responses are mitigated with repeated exposures within one hour but not with continuous exposure, nor with subsequent exposures 5-24 hours later. Recent work indicates that the combined effect of SO₂ and cold, dry air further exacerbates the asthmatic response while warm, humid conditions mitigate SO₂ effects.

c) Given practical considerations related to monitoring, modeling, data manipulation and storage, and implementation, the staff previously recommended consideration of a 1-hour averaging time to protect against the responses to short-term peak (5-10 minute) SO₂ exposures observed in the controlled human studies. Based on this updated staff assessment,

the range of potential 1-hour levels of interest is revised from 0.25 to 0.75 ppm to 0.2 to 0.5 ppm (525 to 1300 $\mu\text{g}/\text{m}^3$). The lower bound represents a 1-hour level for which the maximum 5 to 10 minute peak exposures are unlikely to exceed 0.4 ppm, which is the lowest level where potentially significant responses in free (oronasal) breathing asthmatics have been reported in the criteria document addendum. The upper bound of the range represents a 1-hour level for which 5 to 10 minute peak concentrations are unlikely to exceed 1 ppm, a concentration at which the risk of significant functional and symptomatic responses in exposed sensitive asthmatics and atopics appears high. In evaluating these laboratory data in the context of decision making on possible 1-hour standards, the following considerations are important: (a) the significance of the observed or anticipated responses to health, (b) the relative effect of SO_2 compared to normal day to day variations in asthmatics from exercise and other stimuli, (c) the low probability of exposures of exercising asthmatics to peak levels, and (d) five to ten minute peak exposures may be a factor of two greater than hourly averages.

d) Independent of frequency of exposure considerations, the upper bound of the range contains little or no margin of safety for exposed sensitive individuals. The limited geographical areas likely to be affected and low frequency of peak exposures to active asthmatics if the standard is met add to the margin of safety. The widespread use of medication among asthmatics that prevents or rapidly relieves bronchoconstrictive effects due to natural and commonly encountered stimuli (e.g., exercise, cold air) further adds to the margin of safety. The data do not suggest other groups that are more sensitive than asthmatics to single peak exposures, but qualitative data suggest repeated peaks might produce

effects of concern in other sensitive individuals. Potential interactions of SO₂ and O₃ have not been investigated in asthmatics. The qualitative data, potential pollution interactions, and other considerations listed above should be considered in determining the need for and evaluating the margin of safety provided by alternative 1-hour standards.

3) Based on a staff assessment of the recent short-term epidemiological data summarized in Table 2, the original staff range of 24-hour SO₂ levels of interest - 0.14 to 0.19 ppm (365 to 500 µg/m³) - still appears appropriate, although some consideration could be given to the findings of physiological changes of uncertain significance at levels as low as 0.1 ppm. Earlier staff conclusions and recommendations concerning retaining the present 24-hour standard remain appropriate.

Table 2. UPDATED STAFF ASSESSMENT OF SHORT-TERM EPIDEMIOLOGICAL STUDIES

Effects/ Study	Measured SO ₂ - µg/m ³ (ppm) - 24 hour mean			
	Daily Mortality in London ¹	Aggravation of Bronchitis ²	Small, Reversible Declines in Children's Lung Function ³	Combined Effects Levels
Effects Likely	500-1000 (0.19-0.38)	500-600 (0.19-0.23)	-	500 (0.19)
Effects Possible	-	<500 (0.19)	250-450 (0.10-0.18)	250 (0.10)
No Effects Observed	-	-	100-200 (0.04-0.08)	<200 (.08)

¹Deviations in daily mortality during London winters (1958-1972). Early winters dominated by high smoke and SO₂, principally from coal combustion emissions, and with frequent fogs (Martin and Bradley, 1960; Ware et al., 1981; Mazumdar et al., 1981, 1982; Schwartz and Marcus, 1986).

²Examination of symptoms reported by bronchitics in London. Studies conducted from the mid-1950's to the early 1970's (Lawther et al., 1970).

³Studies of children in Steubenville (1978-80) and in the Netherlands (1985-86) before, during, and after pollution episodes characterized by high particle and SO₂ levels (Dockery et al., 1982; Dassen et al., 1986).

4) The previous staff assessment concluded that although the possibility of effects from continuous lower level exposures to SO_2 cannot be ruled out, no quantitative rationale could be offered to support a specific range of interest for an annual standard. The more recent epidemiological data, indicating associations between respiratory illnesses and symptoms and persistent exposures to SO_2 in areas with long-term averages exceeding .04 ppm ($100 \mu\text{g}/\text{m}^3$), provide additional support for the original recommendation for retaining an annual standard at or near the current level of 0.03 ppm ($80 \mu\text{g}/\text{m}^3$). This recommendation was based in part on a finding that alternative short-term standards (1, 3, and 24-hour) would not prevent annual levels in excess of the current standard in a limited number of heavily populated urban areas. In addition, recent evidence suggests smaller sources in urban areas may produce short duration (< 1 minute) peaks of potential concern. The long-term standard often serves to limit the emissions of numerous smaller sources in such areas. Given the additional information and the possibility of both chronic and acute effects from a large increase in population exposure, the staff recommends maintaining the primary annual standard at its current level.

5) Analyses of alternative averaging times and population exposures suggest that:

- a) The current standards provide substantial protection against the effects identified as being associated with 24 hour and annual exposures.
- b) The current standards - as reflected by current emissions or emissions when the standards are just met with somewhat less restrictive implementation assumptions -- also provide some limit on peak SO_2 exposures of concern for asthmatics. In some cases,

however, up to 1 to 14% of the sensitive population in the vicinity of major sources could be exposed once a year to levels at or above 0.5 ppm for 5 minutes, while at elevated ventilation.

- c) The range of 1-hour standards analyzed (0.25 to 0.5 ppm) provides increased protection against such exposures, limiting the fraction of asthmatics exposed living near certain major point sources to less than 4%, although very short-term (<2 minutes) exposures greater than 0.5 ppm around smaller facilities would not be eliminated.

The relative protection afforded by current vs. alternative standards as indicated by current and ongoing exposure analyses is an important consideration in determining what, if any, standard revisions may be necessary.

REVIEW OF THE NATIONAL AMBIENT AIR QUALITY STANDARDS FOR SULFUR OXIDES:
UPDATED ASSESSMENT OF SCIENTIFIC AND TECHNICAL INFORMATION

ADDENDUM TO THE 1982 OAQPS STAFF PAPER

I. INTRODUCTION

A. Purpose

This paper evaluates and interprets the most relevant scientific and technical information reviewed in the draft EPA document, Second Addendum to Air Quality Criteria for Particulate Matter and Sulfur Oxides (1982): Assessment of Newly Available Health Effects Information (EPA, 1986a) and represents an update of the 1982 sulfur oxides staff paper (EPA, 1982a). This staff paper addendum is intended to help bridge the gap between the scientific review of recent health effects information contained in the criteria document (CD) addendum and the judgments required of the Administrator in setting primary national ambient air quality standards (NAAQS) for sulfur oxides. As such, particular emphasis in this paper is placed on conclusions, recommendations, and uncertainties regarding the averaging times and levels for the primary standards. While the paper should be of use to all parties interested in the standards review, it is written for those decision makers, scientists, and staff who have some familiarity with the technical discussions contained in the criteria document addendum.

B. Background

1. Legislative Requirements

Since 1970 the Clean Air Act as amended has provided authority and guidance for the listing of certain ambient air pollutants which may endanger

public health or welfare and the setting and revising of NAAQS for those pollutants. Primary standards must be based on health effects criteria and provide an adequate margin of safety to ensure protection of public health. As several recent judicial decisions have made clear, the economic and technological feasibility of attaining primary standards are not to be considered in setting them, although such factors may be considered to a degree in the development of State plans to implement the standards (D.C. Cir., 1980, 1981). Further guidance provided in the legislative history of the Act indicates that the standards should be set at "the maximum permissible ambient air level . . . which will protect the health of any (sensitive) group of the population." Also, margins of safety are to be provided such that the standards will afford "a reasonable degree of protection . . . against hazards which research has not yet identified." (Committee on Public Works, 1974). In the final analysis, the EPA Administrator must make a policy decision in setting the primary standards, based on his judgment regarding the implications of all the health effects evidence and the requirement that an adequate margin of safety be provided.

2. Original Sulfur Oxides Standards and Review to Date

The current primary standards for sulfur oxides (to protect public health) are 0.03 parts per million (ppm) or 80 micrograms per cubic meter ($\mu\text{g}/\text{m}^3$), annual arithmetic mean, and 0.14 ppm ($365 \mu\text{g}/\text{m}^3$), maximum 24-hour concentration not to be exceeded more than once per year. The current secondary standard for sulfur oxides (to protect public welfare) is 0.5 ppm ($1300 \mu\text{g}/\text{m}^3$), maximum 3-hour concentration, not to be exceeded more than once per year. For both primary and secondary standards, sulfur oxides are measured as sulfur dioxide (SO_2). Thus, SO_2 is the current indicator for the sulfur oxides standards.

The formal review of the original SO₂ criteria and standards was initiated in 1978. The Clean Air Scientific Advisory Committee (CASAC) closed on the criteria document (which also addressed particulate matter) in January 1982. The first addendum to the criteria document (CD), which summarized the recent controlled human studies on the health effects of SO₂, was issued the same year. A staff paper, which identified critical issues and summarized the staff's interpretation of key studies, received verbal closure at a CASAC meeting in August 1982 and formal written closure in August 1983 (See Appendix A for Executive Summary of staff paper). The decision to produce a second addendum to the combined PM/SO₂ criteria document and this sulfur oxides staff paper addendum was taken in context of the recommendations to review certain new studies on the effects of particulate matter and announced on April 1, 1986 [51 FR 11058].

A preliminary draft of this paper was reviewed by the CASAC in October 1986. This final product incorporates the suggestions and recommendations of the CASAC as well as other comments received on the initial draft. The CASAC closure memorandum (Lippmann, 1987) is reprinted in Appendix B.

C. Approach

The approach in this paper is to address the newly available health effects information in the second criteria document addendum (CD addendum or CDA; EPA, 1986a) in the context of those critical elements which the staff believes have implications for previous conclusions reached on the primary sulfur oxides standards. Particular attention is drawn to judgments related to the ranges of interest for the primary standards. Previous staff conclusions related to the secondary standard, and the form of the standards will not be addressed here.

Because sulfur oxides are often studied in combination with particulate matter, much of the more important literature has already been assessed in the companion staff paper and staff paper addendum on particulate matter (EPA, 1982c; 1986b). Where possible, pertinent references are made to those papers, with only summaries presented in this paper.

The principal focus of this paper is on the effects of SO_2 , alone and in combination with other pollutants. Other sulfur oxide vapors (e.g., SO_3) are not commonly found in the atmosphere. The effects of the principal atmospheric transformation products of SO_2 (i.e., sulfuric acid and sulfates) are discussed in the companion staff paper on particulate matter (EPA, 1982c) and will be further examined in a forthcoming document on acid aerosols.

Section II provides an update of air quality information on sulfur oxides to support discussions of the primary standards. Section III addresses those essential elements that require re-examination in light of the new information reviewed in the CD addendum; these elements include identification of possible mechanisms of toxicity and discussion of controlled human and community studies relating level(s) and duration(s) of exposure to indicators of health effects.

Drawing from the discussion in Sections II and III, Section IV identifies and assesses the factors the staff believes should be considered in selecting the averaging times and levels of primary standards. Updated staff findings and recommendations on the alternative policy options in these areas are also presented.

II. AIR QUALITY CONSIDERATIONS

The major chemical and physical properties of SO₂ in the atmosphere and characterization of ambient concentrations at U.S. sites are presented in the 1982 staff paper ("SP"; EPA 1982a) and discussed in more detail in Chapters 2 and 5 of the CD (EPA, 1982b). Because most of the recently available health effects information on SO₂ is related to short-term (5 to 10 minute) exposures, this section will update information on short-term peak-to-mean ratios and related issues. This information is relevant in estimating human exposures and examining relationships among different standard averaging times.

A. Peak-to-Mean Ratios

The shortest averaging period retained in many monitoring data banks and produced by atmospheric models is one hour. The 1982 staff paper summarized the available information on the variance of 5 to 10 minute peak concentration within particular hourly periods. That assessment concluded that, based on typical distributions, the 5 to 10 minute peak is likely to be within a factor of 1.4 to 2.4 times the hourly average (Larsen, 1968; Burton and Thrall, 1982).

Recent work (Thrall et al.; 1982, Rote and Lee, 1983; Armstrong, 1985, 1986) on peak-to-mean ratio appears consistent with the earlier assessment. Thrall et al. (1982) analyzed monitoring data taken from a dense (18 site) network around the Kincaid (Illinois) power plant. The network was established as a part of an Electric Power Research Institute (EPRI) model validation study. Kincaid is an isolated 1300 MWe, base load, coal-fired plant with a single 187 meter stack. A 23-week sample (March-August 1980) was examined. The maximum hourly value in this sample was approximately 0.34 ppm and the maximum 5-minute value was 0.56 ppm. Thrall

et al. found that the peak-to-mean ratios tend to fall as the hourly average increases. Thus, although the overall ratio of 5-minute peak to hourly mean was $2.3 \pm 1.3^*$ for all hours, the ratio for hours over 0.1 ppm was only 1.8. The overall 10-minute peak to hourly mean ratio was $2.0 \pm 0.96^*$, dropping to 1.6 for hours over 0.1 ppm.

Thrall and coworkers considered the situation of an isolated fuel combustion source. Rote and Lee (1983) provides a similar analysis for urban areas. In this case, the Regional Air Pollution Study (RAPS) data base was used. RAPS was a two year (1975-1976) study of air pollution in St. Louis which included 13 SO₂ monitoring sites. Unfortunately, the instruments were spanned to 1.0 ppm and for 10 sites, as many as 6% of the 1-minute values exceeded 1.0 ppm. Analyzing a large random sample of station hours (40,000), Rote and Lee found that the overall ratio of 5 minute peak to hourly mean concentration was $1.5 \pm 0.48^*$ while the 10-minute peak-to-mean ratio was $1.4 \pm 0.39^*$. These ratios for all hours combined were found to be unaffected by hours containing out-of-range 1-minute values. At higher mean concentrations, the ratios also tended to be lower. However, in this case Rote and Lee found evidence that, for hours > 0.5 ppm, the apparent decline in ratio with increasing mean concentration was in part due to the spanning of the instruments.

Recent air quality analyses of sites near two primary copper smelters in Arizona estimated six minute peak-to-one-hour mean ratios (Armstrong, 1985, 1986). Although the ratios found at the Magma - San Manuel smelter were in the range of those found at Kincaid and other sites, the ratios at a second smelter (Phelps-Dodge, Douglas) were higher, with a 6 minute peak to hourly mean ratio of 3.3.

*Standard deviation

B. Factors Affecting Assessment of Peak Air Quality Levels

The 1982 staff paper concluded that short-term peak levels in excess of 0.5 ppm were most likely in the vicinity of major SO₂ point sources. Recent theoretical work on low persistence meteorological events (Huber and Pooler, 1985) as well as analyses of ambient data (Kilkelly and Roberson, 1985) have raised questions regarding both the impact of smaller sources of SO₂ and the adequacy of monitoring data to assess such impacts. A staff assessment of these issues found that small sources with less than Good Engineering Practice (GEP) stack height may also produce SO₂ peaks in excess of 0.5 ppm (EPA, 1986c). Most of these peaks are due to building downwash, are of limited area and extent (usually within 0.5 km of the stack), and are of very short duration (usually 30 seconds to 2 minutes). Based on the analyses noted above, it appears that very short duration peaks in excess of 0.5 ppm may occur on the order of 1000 per year at a fixed location. No accurate determination of how many sources may be subject to downwash appears feasible. Preliminary, but very rough, calculations indicate that the numbers may be quite significant. In addition, small sources regardless of stack height, may also produce comparable short duration peaks due to looping plumes. These exceedances would likely be found within 3 km of the stack and occur on the order of 10 times per year (EPA, 1986c).

A review of Kilkelly and Roberson (1985) and related strip charts permits several insights regarding the monitoring of very short-term (2-3 minute) peak SO₂ concentrations. The data in question were recorded near facilities with short stacks and are reported to show evidence of building downwash (Docket No. A-83-49, Item IV.H.39). Staff examination showed that the instruments were spanned to 1 ppm and frequently hit this limit for short-time periods. This means that the true peaks can not be readily

estimated but were presumably in excess of 1 ppm. This "peak lopping" does not appear to affect significantly the hourly averages at the sites in question because the area under the curve at the peak is quite small. Clearly, for peaks of longer duration (\geq 5-10 minutes), peak lopping would lead to a significant underestimate of the hourly average. Peaks in excess of the spanned value for 5-10 minutes were seen at some of the facilities in the Kilkelly set and around some TVA facilities (Lott, 1985). In such cases, it is possible that hourly averages may be underestimated due to spanning. Peak lopping, if it occurs, would also bias any analysis of peak to mean ratios. EPA monitoring guidance calls for SO₂ instruments to be spanned to 0.5 ppm with a requirement that they be respanned if the limit is reached.

A related concern examined by the recent staff assessment (EPA, 1986c) is instrument response time. Many SO₂ instruments now in wide use require 4-5 minutes to reach 95% of scale. Thus, if the actual peak lasts only 30 seconds to 1 minute, most instruments would not respond fast enough to register the true peak.

In summary, the recent staff assessment of short-term peaks and smaller sources prompts the following conclusions:

- 1) Peaks well in excess of 0.5 ppm appear likely to occur around numerous small sources of SO₂. Although of very limited duration and areal extent, they can occur with relatively high frequency. None of the recently published assessments of the health effects of SO₂ has addressed exposures of such limited duration (< 30 seconds to 2 minutes); and

- 2) It appears that, due to spanning and instrument response time, most monitored data are not accurately measuring very short-term peaks. It is therefore not presently possible to assess the extent to which such peaks may be occurring.

III. CRITICAL ELEMENTS IN THE REVIEW OF THE PRIMARY STANDARDS

This section summarizes relevant aspects of recent information in the CD addendum on the mechanisms by which SO₂ may cause airway reactions and concentration/response relationships derived from controlled human and community studies of SO₂ effects. A comprehensive discussion of these and other critical elements including mechanisms of toxicity, effects of concern, and sensitive populations is contained in Section V of the 1982 staff paper (EPA, 1982a). The present summary provides a basis for later discussions of the implications of the more recent studies for the standards review.

A. Mechanisms

The previous staff assessment found that the most striking acute response to SO₂ for asthmatics and others with hyper-reactive airways is rapid bronchoconstriction (airway narrowing), usually evidenced in increased airway resistance, decreased expiratory flow rates, and the occurrence of symptoms such as wheezing, chest tightness, and shortness of breath. Several of the more recent studies discussed in the CD addendum contribute further to understanding mechanisms and factors that affect these responses (CDA, Section 4.4). The discussion below highlights insights from the CD addendum with respect to the impact of breathing mode, temperature/humidity conditions, and the time course of exposure and recovery.

1. Inhalation Patterns and Airway Cooling/Drying

The penetration of SO₂ to sensitive portions of the respiratory tract is largely determined by the efficiency of the oral or nasal mucosa in absorbing SO₂, which in turn depends on the mode of breathing (nasal, oral, or oronasal) and the rate of airflow. Newly published controlled SO₂ exposure studies on asthmatics confirm previous findings that at

comparable SO_2 concentrations, bronchoconstriction effects increase with both increased ventilation rates and as the relative contribution of oral ventilation to total ventilation increases, as seen by comparing oral-only (i.e., mouthpiece) breathing with oronasal breathing (Bethel et al., 1983b, 1985; Roger et al., 1985; Koenig et al., 1985).

The CD addendum notes that increased oral ventilation not only allows more direct penetration of SO_2 but may also result in airway drying and alterations in airway surface liquid that further affects SO_2 absorption and penetration (CDA, pp. 4-41 to 4-42). Evaporation of airway surface liquid and perhaps convective cooling of the airways caused by cold, dry air can act as direct bronchoconstrictive stimuli in asthmatics (Deal et al., 1979; Strauss et al., 1977; Anderson, 1985). Recent studies indicate that the combined effect of SO_2 and cold, dry air further exacerbates the asthmatic response (Bethel et al., 1984; Sheppard et al., 1984; Linn et al., 1984a,b, 1985a). It has been suggested that reduced water content and not cold per se could be responsible for much of this effect. This is consistent with other recent findings that the bronchoconstrictive effects of SO_2 are reduced under warm, humid conditions (Linn et al., 1985a). It appears that the interactive effects of breathing SO_2 and dry (or cold) air range from less than additive to synergistic depending on whether oral airway geometry is altered by use of mouthpieces, preventing any initial conditioning of inspired air in the mouth (e.g., warming, humidifying) (CDA, p. 4-42).

2. Time Course of Response, Recovery and Adaption

The time required for SO_2 exposure to elicit significant bronchoconstriction in exercising asthmatics is brief. Exposure durations as short as 3 minutes have produced significant responses in a mouthpiece study (Sheppard et al., 1984)

with the majority of studies using 5-10 minute exposure durations. Little enhancement of response is apparent with prolonged exposure beyond 5 minutes, although some suggestion of an increase is seen with continuous exercise between 10 and 30 minutes (Kehrl et al., 1986). On mechanistic grounds, it would appear possible for some response to occur with exposures of less than 5 minutes with high enough concentrations. The relationship between concentration, time and response for such periods has not, however, been systematically examined.

Following a single SO₂ exposure during exercise, airway resistance in asthmatics appears to almost fully recover within one hour, even if low-level SO₂ exposure is continued without exercise (Hackney et al., 1984). A reduced response is observed if SO₂ exposure is repeated within 15-60 minutes, but not with subsequent exposures 5-24 hours later (Sheppard et al., 1983; Roger et al., 1985; Kehrl et al., 1986; Linn et al., 1984c; Snashall and Baldwin, 1982). Similar attenuation of airway constriction, induced by exercise or hyperventilation of cold, dry air, is observed when the exercise exposure is repeated at short-time intervals, with a refractory period that persists for 2-4 hours (Stearns et al., 1981; Bar-Yishay et al., 1983). Significantly, while repeated short exercise periods over a 1-hour period result in reduced response, 30 minutes of continued exercise results in responses that equal or exceed those observed after a single 10 minute period (Kehrl et al., 1986).

The CD addendum discusses several possible mechanisms that might account for the mitigated responses to SO₂ over time (e.g., decreased responsiveness of airway smooth muscle or vagal reflex pathways due to mediator depletion or inhibition of SO₂-receptors) (CDA, p. 4-43).

Since continuous exercise apparently prevents a recovery period, Kehrl

et al. (1986) suggest that the mechanism for "adaptation" to rapidly repeated SO₂ exposures may be increased production and/or secretion of airway surface liquid during recovery following an SO₂ challenge. This would act to decrease relative SO₂ penetration in subsequent exposures.

B. Concentration/Response Information

The following review summarizes key results from those recent studies cited by the CD addendum as providing the most reliable quantitative information as well as some that provide reasonable evidence of concentration-response relationships without allowing derivation of specific levels. Responses to SO₂, alone or in combination with other pollutants, are examined in three time scales: 1) peak exposures (minutes-hours), 2) short-term exposures (hours-days) and 3) long-term exposures (months-years). A further assessment of these studies as applied to selecting alternative levels for air quality standards is presented in Section IV.

1. Peak Exposures

Information on the effects of relatively brief (minutes-hours) peak exposures to SO₂ is derived from studies of humans under controlled laboratory conditions. The importance and limitations of controlled human exposure studies are discussed in the CD and CD addendum as well as the 1982 staff paper (EPA, 1982a,b; 1986a). Recent controlled exposure studies confirm that "normal", healthy subjects, even at moderately heavy exercise, do not experience significant effects on pulmonary function due to peak SO₂ exposures in the range of 0.25 to 2 ppm (CDA, p. 4-10). A single recent chamber study of chronic obstructive pulmonary disease patients was conducted under conditions that the CD addendum states are unlikely to produce effects even in sensitive individuals. Thus, the preponderance of newly available exposure-response information on peak SO₂ exposures is for exercising asthmatic subjects.

The results of the recent studies of asthmatic subjects are summarized in Table 7 of the CD addendum, which organizes the data according to concentration. Most of the data reflect 5 to 10 minute exposures to young, relatively healthy, mild non-smoking volunteers with no symptoms and fairly stable pulmonary function at the time of exposure. The following discussion of anticipated responses associated with particular concentrations is drawn from that tabular summary.

a) 1.0 to 2.0 ppm

Recent studies by two separate research laboratories of the effects of 1 ppm SO₂ on freely breathing, mild asthmatics at moderate exercise are qualitatively consistent with each other as well as with previous studies that administered exposures through mouthpieces. All found statistically and potentially clinically significant* changes in respiratory mechanics, most pronounced within minutes after exercise had ceased, followed by gradual recovery (within 1 hour). When reported, associated symptoms (e.g., shortness of breath, chest discomfort) also increased significantly (Schacter et al., 1984; Roger et al., 1985). Group mean functional changes were increases in specific airway resistance (SRaw) (170 to 230%) and declines in FEV₁ (CDA, Table 7). A third laboratory found consistent reductions in FEV₁ (-23%) using mouthpiece exposures (Koenig et al., 1983b). Individual variability is illustrated by the Roger et al. results. One subject could not be tested at 1.0 ppm because he required medication following exposure to a lower concentration. Another was removed after the second exercise period due to pronounced wheezing and chest tightness and a 10-fold increase period in SRaw. Two other subjects had a greater than 500% increase. The responses in asthmatics observed by Kehrl et al.

*Unless otherwise modified (as in this case), the use of "significant" with respect to measured changes could be understood as "statistically" significant and not necessarily clinically or medically significant.

(1986) appear to be greatest after 30 minutes of continuous exercise although the increase in airway resistance was statistically no greater than the changes observed after 10 minute exposure (233% vs. 172% increase over baseline). Successive exercise periods separated by 15 minute intervals resulted in attenuated responses even to 1 ppm SO₂ (Roger et al., 1985; Kehrl et al., 1986).

Horstman et al. (1986) report that 12 (of 27) subjects in the Roger et al. (1985) study, whose SRaw values did not increase by 100% at 1 ppm or lower levels, were also exposed to 2 ppm using the same protocol. At this level, 7 of these less sensitive asthmatics had SRaw increases of 100 to over 600%.

b) 0.75 ppm

Recently published studies of moderately exercising asthmatics exposed to 0.75 ppm SO₂ for 10 minutes (Linn et al., 1983a; Hackney et al., 1984; Schacter et al., 1984) replicate earlier results, finding significant increases in airway resistance (group mean SRaw increase was 186 to 263%), substantial decreases in FEV_{1.0}, and significantly increased reports of lower airway symptoms. In contrast to functional measurements, the increase in symptom scores were not significantly greater when SO₂ was administered through mouthpieces compared to freebreathing in a chamber.

c) 0.6 ppm

Highly consistent and significant bronchoconstrictive responses have been observed in freely breathing mild asthmatics exposed to 0.6 ppm for 5 minutes while exercising at fairly high levels (minute ventilation, V_e, ~ 50 L/min) under a wide range of temperature and humidity conditions (Linn et al., 1983b; 1984a,b; 1985a). Increases in airway resistance and symptom scores were most pronounced (~ 207% over control) in either cold or dry air (-6°C, 20% RH) compared with more humid, warmer

conditions (e.g., 39% increase in SRaw in 38°C, 80% RH). Even under moderate conditions (~ 22°C, 85% RH), Linn et al. (1984a) found that typical respiratory symptoms were sufficient to impair subjective well-being and "normal performance capabilities". Three of the 23 subjects in this study required medications to relieve symptoms following exposures and four had SRaw increases in excess of 250%. In this and a subsequent study (Linn et al., 1984c), these investigators examined symptoms during the week after SO₂ exposures. In the latter study, they reported a tendency toward less favorable clinical states (i.e., increased symptoms) in the week following exposures on two successive days to 0.6 ppm and that three (of 14) subjects reported experiencing an asthma attack during the week after SO₂ exposure, whereas no subject reported such an attack after clean air exposure. In contrast, two subjects reported a need for extra bronchodilator medication after the SO₂ exposures while four reported such a need after clean air exposures. The authors concluded that these post-exposure effects "were small and inconsistent enough that they might be attributable to chance, or to preferential recall of symptoms after the clinically stressful SO₂ exposure experience." Comparable findings have not been noted in other studies.

d) 0.5 ppm

Recent studies of airway responses in free breathing mild asthmatics exposed at exercise to 0.5 ppm SO₂ for durations of 5, 10, and 20 minutes indicate that significant bronchoconstriction occurs at moderate to heavy exercise rates ($V_e \sim 40-60$ L/min) (Bethel et al., 1983a, b; Koenig et al., 1983; Roger et al., 1985) but not at lower exercise rates ($V_e \sim 27-40$ L/min) (Schacter et al., 1984; Bethel et al., 1983b).

Roger et al. (1985) examined both repeated exposures and subject variability. Responses to SO_2 were mitigated after repeated, free-breathing exposures separated by 15-minute intervals, although they remained significant. Elevations in airway resistance over baseline averaged 93% after the first exercise period and 52% after the third exercise period. Cumulative frequency distributions of the subjects' S_{Raw} values at rest and at exercise in clean air and after 10-minute exercise in 0.5 ppm SO_2 indicate that exercise and SO_2 each contributed about equally to the overall increase in S_{Raw} . As in other studies, there was a wide range in the magnitude of the induced bronchoconstriction in various subjects. For example, after exercise in 0.5 ppm SO_2 , 25% of the subjects had a S_{Raw} increase of ~ 170% over baseline compared to the mean of 93%, while 25% had negligible changes. In addition, while significant increases in symptoms were not reported for the group as a whole, three subjects had S_{Raw} increases of over 320% and one, who was removed before completion of the full protocol, had an eight-fold S_{Raw} increase after 10-minutes of 0.5 ppm and an 11-fold increase after the 2nd exposure, with audible wheezing and chest tightness.

e) 0.4 ppm

Mild asthmatics performing moderately heavy exercise ($V_e = 48$ L/min) while freely breathing 0.4 ppm SO_2 for 5 minutes had statistically significant increases in S_{Raw} (group mean 69% increase vs. 35% in clean air) and mild increases in several symptoms (e.g., cough, wheeze, chest tightness) after 5 minute exposure (Linn et al., 1983b). One subject (of 23) was reported to have experienced "clinically significant bronchoconstriction" after this exposure and required medication to relieve asthma symptoms. As part of another study discussed previously, a group of mild asthmatics exercising at a similar level (~ 50 L/min) at a much colder temperature (5°C),

responded with apparent increases in airway resistance and respiratory symptoms at 0.4 ppm SO₂ under both high and low humidity (Linn et al., 1984a).

f) 0.1 - 0.3 ppm

Most recent chamber exposures found no clearly significant increases in airway resistance among freely breathing mild asthmatics exercising at moderate to high levels (35-50 L/min) below 0.4 ppm (Linn et al., 1984a,b; 1983b; Roger et al., 1985; Schacter et al., 1984). At 0.25 ppm with heavy exercise (60 L/min), Bethel et al. (1985), found apparently significant responses although the application of a more appropriate statistical test did not confirm this (CDA, p. 4-27). Even here, a significant increase over exercise control was not observed with 0.25 ppm in the same study at an even higher ventilation rate (80-90 L/min), suggesting that the bronchoconstriction induced by exercise alone overshadowed any effects of SO₂ (Bethel et al., 1985). Although some minimal increases in symptom scores were reported even as low as 0.2 ppm, the clinical significance of these changes is questionable (Linn et al., 1983b; 1984a). The fact that some hyper-reactive individuals may be responsive to such low SO₂ levels cannot be dismissed, however, given that an SO₂ concentration of 0.25 ppm was sufficient to nearly double SRaw over baseline in the most sensitive subject (Bethel et al., 1985).

g) Combined Relationships/Subject Variability

A number of the more recent studies developed exposure/response relationships over various concentration and ventilation ranges while others examined the influence of various subject-related and environmental factors. Although individual studies fix various important factors to permit within study comparisons, it is more difficult to compare directly the results from different investigations. One approach suggested in an earlier staff assessment (Cohen, 1983) and used by Kleinman (1984) and Linn et al. (1983b),

normalizes studies according to effective oral dose rate. The results of such an analysis applied to both recent and earlier SO_2 studies are shown in Figure 3-1. As illustrated, reasonably consistent results are derived from the various controlled SO_2 asthmatic studies when adjustments are made for differences in ventilation rates and oral/nasal breathing patterns by expressing the results in terms of the oral dose rates of SO_2 . Earlier analyses also found a good consistency among then available studies using similar surrogates of "effective dose" (Kleinman, 1984; Linn et al., 1983b).

This relationship can be used to estimate responses for exposures of interest not yet tested. For example, it is of interest to determine whether large responses might occur in asthmatics at high concentrations, e.g., 2 ppm, while at lower ventilation rates typical of everyday activities. Assuming oronasal ventilation for "mouth" breathers (Ninamäa et al., 1981), oral V_e would be about 4 to 7 L/min at rest to light activity and the predicted mean increase in S_{Raw} for 2 ppm would be approximately 0 to 70%.

The consistency among group mean responses represented in Figure 3-1 masks the substantial variability among individual asthmatics, both within and among studies. Among the most useful studies for examining this variability is the work of Roger et al. (1985) and companion analysis by Horstman et al. (1986). The study covers a wide range of concentrations (0.25 to 2 ppm), includes a substantial number of subjects (28) who were not pre-selected for SO_2 sensitivity, and presents individual exposure-response data. The highlighting in Figure 3-1 shows that the group mean results from Roger et al. are representative of the range of values for all SO_2 /asthmatic studies. The range of subject responses from this work is illustrated in Figure 3-2, reproduced from the Horstman et al. (1986) report. The points represent a logarithmic linear interpolation of exposure-response relationships




Figure 3-1. Combined SO₂ dose/response relationships. Includes recent and earlier controlled studies of exercising or hyperventilating asthmatics with SRAW reported. Points represent group mean responses significantly different from baseline vs. normalized oral-dose rate (SO₂ concentration x oral V_e). Oral V_e estimates for freebreathing from Niinimaa et al. (1981). The results of Roger et al. (1985) are highlighted at specific concentrations (V_e total = 42 l/min) for orientation. Despite uncertainties in estimating the oral V_e the results suggest reasonably good consistency among three laboratories with no apparent difference between 5 and 10 minute exposures and no tendency for mouthpiece studies to overstate response when appropriate adjustments for ventilation are made.

- Cohen, J. (1983) Analysis of SO₂ Controlled Human Data. Memorandum to John Haines, Ambient Standards Branch, Office of Air Quality Planning and Standards, U.S. EPA, Research Triangle Park, N.C., September 6, 1983.
- Cohen, J. (1984) Assessing Asthmatic Responses to SO₂: Summary of Expert Opinions. Memorandum to John Bachmann, Ambient Standards Branch, Office of Air Quality Planning and Standards, U.S. EPA, Research Triangle Park, N.C., February 27, 1984.
- Committee on Public Works, U.S. Senate (1974) A Legislative History of the Clean Air Amendments. Volume 1. Serial no. 93-18. U.S. Government Printing Office, Washington, D.C. prepared by the Environmental Policy Division of the Congressional Research Services of the Library of Congress.
- CEC [Commission of the European Communities] (1983) Report on the EC epidemiology survey on the relationship between air pollution and respiratory health in primary school children. Brussels, Belgium: Environmental Research Programme.
- Dassen, W.; Brunekreef, B.; Hoek, G.; Hofschreuder, P.; Staatsen, B.; deGroot, H.; Schouten, E.; Biersteker, K. (1986) Decline in children's pulmonary function during an air pollution episode. *J. Air Pollut. Control Assoc.* (in press).
- D.C. Cir. (1980) *Lead Industries Association, Inc. v. EPA*, F. 2d, 14 ERC 1906 (D.C. Cir.) Cert. Denied 49 U.S.L.W. 3428 December 8, 1980.
- D.C. Cir. (1981) *American Petroleum Institute v. Costle*, Nos. 79-1104 et. al. (D.C. Cir.) September 3, 1981.
- Deal, E. C.; McFadden, E. R.; Ingram, R. H.; Jaeger, J. J. (1979b) Hyperpnea and heat flux: initial reaction sequence in exercise-induced asthma. *J. Appl. Physiol.* 46: 476-483.
- Dockery, D. W.; Ware, J. H.; Ferris, B. G., Jr.; Speizer, F. E.; Cook, N. R.; Herman, S. M. (1982) Change in pulmonary function in children associated with air pollution episodes. *J. Air Pollut. Control Assoc.* 32: 937-942.
- Dodge, R. (1983) The respiratory health and lung function of Anglo-American children in a smelter town. *Am. Rev. Respir. Dis.* 127: 158-161.
- Dodge, R.; Solomon, P.; Moyers, J.; Hayes, C. (1985) A longitudinal study of children exposed to sulfur oxides. *Am. J. Epidemiol.* 121: 720-736.
- EPA [U.S. Environmental Protection Agency] (1982a) Review of the National Ambient Air Quality Standards for Sulfur Oxides: Assessment of Scientific and Technical Information (OAQPS Staff Paper) Office of Air Quality Planning and Standards, Research Triangle Park, N.C. EPA-450/5-82-007. Available from NTIS, Springfield, VA; PB 84-102920.

for each subject to determine the SO_2 exposure producing a 100% increase in SRaw over exercise in clean air (termed $\text{PC} [\text{SO}_2]$). The resulting cumulative plot is useful in estimating the likelihood of a possibly clinically significant response (doubling of SRaw) in mild asthmatics exposed at moderate exercise (or ventilation) to particular SO_2 concentrations.

h) Recent Epidemiological Evidence

The literature contains little epidemiologic evidence regarding the relationship between peak SO_2 levels and asthma. Early epidemiological analyses of asthma incidence reported in the 1982 criteria document were based on daily averages of SO_2 and substantially confounded by the presence of other pollutants. A more recent study (Goldstein and Weinstein, 1986), however, has examined the relationship between incidence of emergency room visits for asthma in three New York City hospitals from 1969 to 1972 and hourly SO_2 peaks. Adjustments were made for seasonal, epidemic, day-of-week, and lag effects, as well as for the long-term downward trend in SO_2 levels in N.Y.C. over the study years. Temperature, other pollutants, and pollen counts were not included. No significant association was found between days with "high" hourly SO_2 measurements (> 0.1 , 0.3 , and rarely 0.5 ppm) and days with elevated asthma visits. There are several factors that would have made detecting any associations difficult, including 1) (as noted by the authors) centralized, rooftop monitors may represent population exposure too crudely to detect an effect; 2) hourly SO_2 levels may not detect rapid responses such as those observed in human studies in which exercising asthmatics respond quickly to brief 5-10 minute SO_2 peaks whose effects diminish within an hour; 3) less than 2% of all days had hourly SO_2 levels higher than 0.5 ppm, which substantially limits the statistical power related to examining high exposure situations; 4) actual hour of the day for emergency room visits was not readily available, so the analysis was

done on a daily level that overlooked the timing of the SO₂ peaks and asthma visits within the days. The authors note that visits not influenced by a pollution peak on a day would have been included among all visits examined in relation to that peak thus serving to attenuate any relation that may be present; and 5) confounding influences of normally encountered agents or stresses in the urban environment that asthmatics are susceptible to could not have been readily controlled as in controlled exposure studies. Additional uncertainties regarding the definition of elevated asthma days, the use of 14-day averages to detrend the time series structure of hospital admissions, and controls for day-of-the-week effect require further examination. The authors conclude that the results do not rule out a relationship between asthma and ambient SO₂ and that additional study is needed on the individual exposure-response level over time in order to determine whether the effects observed in the controlled laboratory studies can be detected in free-living populations.

2. Short-term Exposures

The principal basis for developing quantitative assessments of acute effects of ambient exposures of SO₂ on a daily basis remains community epidemiological studies. Such studies can provide strong evidence for the existence of pollution effects resulting from community exposures. The major limitations of the epidemiological studies are discussed in the CD, CD addendum, as well as the 1982 staff paper.

Recognizing these limitations, the discussion in the 1982 staff paper outlined those studies cited by the CD as providing the most reliable quantitative information as well as other studies that provide useful

information on the relative importance of SO₂ without allowing derivation of specific levels. These included a set of British studies of mortality and morbidity. The CD addendum identifies several more recent analyses of the London mortality data and one U.S. morbidity study as providing the most useful new information on the short-term SO₂ exposures. These studies are summarized in Table 3-1. The more full description and evaluation of these studies contained in Section III of the companion draft PM staff paper addendum (EPA, 1986b) will not be repeated here. The discussion will focus on the relative importance of SO₂ as compared with particulate matter in producing the observed effects.

With respect to the daily mortality studies, the CD addendum states that:

"the following conclusions appear warranted based on the earlier criteria review (U.S. EPA, 1982b) and present evaluation of newly available analyses of the London mortality experience: (1) Markedly increased mortality occurred, mainly among the elderly and chronically ill, in association with BS and SO₂ concentrations above 1000 µg/m³, especially during episodes when such pollutant elevations occurred for several consecutive days; (2) During such episodes coincident high humidity or fog was also likely important, possibly by providing conditions leading to formation of H₂SO₄ or other acidic aerosols; (3) Increased risk of mortality is associated with exposure to BS and SO₂ levels in the range of 500 to 1000 µg/m³, for SO₂ most clearly at concentrations in excess of ~ 700 µg/m³; (4) Convincing evidence indicates that relatively small but statistically significant increases in the risk of mortality exist at BS (but not SO₂) levels below 500 µg/m³, with no indications of any specific threshold level having been demonstrated at lower concentrations of BS (e.g., at < 150 µg/m³). However, precise quantitative specification of the lower PM levels associated with mortality is not possible, nor can one rule out potential contributions of other possible confounding variables at these low PM levels" (CDA, p. 3-9).

Because of the high colinearity between BS and SO₂ levels during the study period, it has been difficult to readily separate the effects of the two pollutants on mortality. The CD addendum states that attempts by Mazumdar et al. (1982) using nested quartile analysis were only partially successful given the substantial covariation that remained between the

TABLE 3-1. SUMMARY OF RECENT (1982-86) EPIDEMIOLOGICAL STUDIES PROVIDING MOST USEFUL CONCENTRATION/RESPONSE INFORMATION FOR SHORT-TERM SO₂ EXPOSURES

Observed Effects	Time	Observed PM ($\mu\text{g}/\text{m}^3$)	Concentration Range SO ₂ ($\mu\text{g}/\text{m}^3$)	Comments	Study
Increases in daily mortality in metropolitan London	1958-1972 winters	<500 BS* 24-hr averages	>500	Recently published studies reinforce 1982 CD, SP conclusions regarding likelihood of increased mortality at 500 to 1000 $\mu\text{g}/\text{m}^3$ for BS and SO ₂ , with no clearly defined threshold for BS in the range of 150 to 500 $\mu\text{g}/\text{m}^3$. Nature of relationships vary significantly with model. Mazumdar et al. infer no association < 700 $\mu\text{g}/\text{m}^3$ SO ₂ , but methodology for separating pollutants questioned in CDA.	Mazumdar et al. 1982, 1983; Ostro 1984
Short-term reductions in lung function in 330 school children, Steubenville, OH	Four separate study periods of 3 weeks following pollution "episodes" in 1978-1980	1) 420 TSP 2) 270 TSP 3) 220 TSP 4) 160 TSP (max 24 averages for "alert" or "sham" episode)	280 460 170 190	Recent unpublished analyses confirm major findings of the published studies with advanced statistical techniques accounting for auto-correlation and temperature effects. Schwartz and Marcus findings suggest significant association for BS at lowest levels (<100 $\mu\text{g}/\text{m}^3$ BS), but not for SO ₂ below about 500 $\mu\text{g}/\text{m}^3$.	Shumway et al. 1983, Schwartz and Marcus, 1986
Short-term reductions in lung function in 179 school children in the Netherlands (Ijmond)	Before, during, after pollution episode Nov. 1984-Feb. 1985	200-250 TSP and RSP (D50 < 3.5 μm) 24-hr average	200-250	First 3 episodes: small (2%-3%) but significant reversible declines in FVC up to 2-3 weeks after peak. Less consistent results for FEV ₁ . No significant effects after 4th "sham" episode. Baseline measurements for 1st, 4th taken on days with high pollution. Linear regression of pooled data for 330 children indicate significantly more negative slopes in functions vs. TSP and SO ₂ across ranges (10-270 $\mu\text{g}/\text{m}^3$, 0-280 $\mu\text{g}/\text{m}^3$, respectively). Higher response in some children.	Dockery et al. 1982
				Small (3-5%) reversible declines in several measures of airway function (FVC, FEV ₁ , MEF) during episode and 5 days later. No effect after 26 days or shortly after a day when TSP, RSP and SO ₂ levels all averaged 100-150 $\mu\text{g}/\text{m}^3$. Separate sub-groups of children tested on each day. Peak TSP levels possibly understated.	Dassen et al., 1986

*British Smoke (BS) is a pseudo-mass indicator related to small particle (size less than a nominal 4.5 μm) darkness (CD, pp. 1-88 to 1-90).

pollutants in the highest and lowest quartiles (CDA, p. 3-5). Based on regression analyses using the highest pollution days, the authors concluded, however, that the mortality/pollution association was almost entirely due to smoke with a possible SO₂ effect at higher concentrations (above 700 µg/m³).

Schwartz and Marcus (1986)* examined further the suggestion that the effects of smoke are separable from those of SO₂ in this data set. In regressions involving both pollutants, the colinearity between the two tended to deflate the apparent significance of both. However, the overall results for all years combined and for those individual years with lower correlations between BS and SO₂ ($r < 0.9$) show that the mortality effects of BS remain significant and relatively large even when SO₂ is included in the model, while the inclusion of BS in the model reduces the SO₂ coefficients to insignificant values. This analysis cannot, however, be used to exclude an independent effect of SO₂ at higher concentrations.

Besides the uncertainties that remain in separating the effects of SO₂ and PM, various issues are still unresolved regarding these London data including a possible threshold for PM-mortality associations, varying coefficients obtained with different subsets of data and models, the effects of unmeasured variables such as demographic change over time and indoor air pollution, and the appropriate statistical methods to account for long-term seasonal trends in mortality (Wyzga et al., 1985).

*This paper and a summary memorandum (Marcus and Schwartz, 1986), are reprinted in full as Appendix A to the Criteria Document Addendum. Although not published, the paper was presented to the CASAC and the public for review at the October 15-16, 1986 meeting. Copies were made available to the public at the time of the meeting. Subsequently, EPA received and considered comments on this study from industry and environmental groups and from members of the scientific community.

While the possibility of small increases in the risk of mortality at SO_2 levels less than the "likely effects level" ($500 \mu\text{g}/\text{m}^3$ or 0.19 ppm) cannot be dismissed conclusively, the available analyses of London mortality data provide little basis to determine whether 24-hour concentrations of SO_2 below this level may have accounted for any of the observed association between mortality and pollution. Because significant quantities of SO_2 are unlikely to penetrate to the tracheobronchial region at lower concentrations without increased ventilation, the mechanisms by which SO_2 could contribute to excess mortality in ill or otherwise sensitive populations are limited. Peak levels in London at the time of these studies were undoubtedly well in excess of the 24-hour values, but at lower daily concentrations were less likely to affect even individuals with hyperreactive airways. The capacity of fog particles to "carry" untransformed SO_2 is limited. At present, it appears more likely that the role of SO_2 , in the presence of smoke, involved transformation products such as acidic fine particles.

Other recent studies discussed in the CD addendum and in the PM staff paper addendum examined pollutant/mortality relationships in more contemporary atmospheres in New York City, Pittsburgh, and Athens, Greece. The Ozkaynak et al. (1985) reanalysis of 14 years of N.Y.C. data (1963-76) found significant associations between excess daily mortality and PM, SO_2 and temperature. Differences in the rate of change of SO_2 and PM indicators during the study period allowed estimation of their separate effects. In joint regression analysis across all years, PM indicators (coefficient of haze and visibility extinction coefficient) together accounted for significantly greater excess mortality than did SO_2 . As the CD addendum notes, however, these findings must be considered preliminary for risk assessment purposes.

The work of Mazumdar and Sussman (1983) in Pittsburgh and that of Hatzakis et al. (1986) in Athens, however, found conflicting results. The

first found a significant association between particulate matter and excess deaths in Pittsburgh, but no effect of SO₂, while the Athens study found an association with SO₂ but not with smoke measurements. The CD addendum points out that limitations in both studies with respect to measurement of particulate matter as well as methodological difficulties prevent drawing meaningful conclusions from these studies with respect to the effects of particulate matter and SO₂.

b) Morbidity

Previous conclusions regarding morbidity effects of short-term PM/SO₂ exposures were primarily based on studies of bronchitic subjects in London from the 1950's through the early 1970's. Findings related to more contemporary conditions are presented by Dockery et al. (1982) and Dassen et al. (1986) and summarized in Table 3-1. The CD addendum concludes that the repeated measurements of lung function by Dockery et al. showed statistically significant but physiologically small and apparently reversible declines of FVC and FEV_{0.75} levels associated with short-term increases in PM and SO₂ air pollution (CDA, pp. 3-16, 3-19). The small, reversible decrements appear to persist for up to 3-4 weeks after episodic exposures to these pollutants across a wide range of concentrations with no clear delineation of a threshold defined by the authors or by the CD addendum. A staff assessment of that study is contained in the draft PM staff paper addendum (EPA, 1986b). The following additional points are relevant in assessing the implications of Dockery et al. (1982) for SO₂ concentration/response relationships.

1) Of the 4 study periods in Steubenville, the most significant declines in FEV_{0.75} (4% on average) were observed following the episode with the highest SO₂ level (455 µg/m³, 24 hr. avg). This observation is, however, confounded

because pollution levels during baseline measurements for this period were among the lowest for any of the four study periods.

2) No significant effects on lung function were reported in the Fall 1980 study, when 24-hour SO_2 levels reached $190 \mu\text{g}/\text{m}^3$. Significant lung function declines were measured following a pollution episode in the Spring 1980 study when SO_2 was lower ($169 \mu\text{g}/\text{m}^3$ maximum), suggesting any pollution related effect was more attributable to particles.

3) When data for all 4 study periods were pooled and lung function was regressed on TSP and SO_2 levels - assuming the relationship was linear across all studies -- similar results were obtained for both pollutants.

A similar study of the effects on children of episodic exposures to particulate matter and SO_2 conducted in the Netherlands by Dassen et al. (1986) produced results similar to those of Dockery et al. Pulmonary function values measured during an air pollution episode in which both 24-hour average PM (as TSP or RSP*) and SO_2 levels reached 200 to $250 \mu\text{g}/\text{m}^3$ were significantly lower (3-5%) than baseline values measured 1-2 months earlier in a group of Dutch school children. Lung function parameters that showed significant declines included FVC and FEV, as well as measures of small airway function (e.g., maximum mid-expiratory flow, maximum flow at 50% of vital capacity). Declines from baseline were observed 16 days after the episode in a different subset of children, but not after 25 days in yet a third subgroup. Shortly before the last set of measurements, 24-hour averages of both PM (as TSP or RSP) and SO_2 reached between $100\text{-}150 \mu\text{g}/\text{m}^3$, suggesting that these levels were not associated with observable functional effects (CDA, p. 3-17).

*Respirable Suspended Particles, reportedly $D_{50} \leq 3.5 \mu\text{m}$ by cyclone sampler.

Although comparison of the Netherlands study with the Steubenville episodes must be done with considerable caution, the absolute magnitude of functional response in the Netherlands episode was comparable or greater than that for any of the four Steubenville episodes, while the peak SO_2 levels (200 to 250 $\mu\text{g}/\text{m}^3$) were lower than two of those episodes. Thus, the relative magnitude of the effect appears to be better related to the concentration of small particles (EPA, 1986b).

In summary, the more quantitative epidemiological evidence from London suggests that effects may occur at SO_2 levels at or above 0.19 ppm (500 $\mu\text{g}/\text{m}^3$), 24-hour average, in combination with elevated particle levels. Additional evidence suggests the possibility of short-term, reversible declines in lung function at SO_2 levels above approximately 250-450 $\mu\text{g}/\text{m}^3$ (0.10-.18 ppm). Whether any of these effects are due (in part) to SO_2 alone, formation of sulfuric acid or other irritant aerosols, particles alone, or peak SO_2 values well above the daily mean cannot be determined unequivocally.

3. Chronic Exposures

Table 3-2 summarizes the most useful of the recent studies that have examined the long-term effects of exposures to SO_2 , in the presence of particles, on respiratory mechanics, symptoms, and illness. Other, less reliable, studies are evaluated in Appendix B of the PM staff paper addendum (CEC, 1983; Muhling et al., 1985; Wojtyniak et al., 1986). Several cross-sectional studies report significant associations between long-term SO_2 exposures and effects in populations of adults and children (PAARC, 1982a,b; Chapman et al., 1985; Ware et al., 1986; Dodge, 1983; Dodge et al., 1985).

The CD addendum (p. 3-49 to 3-50) concludes that these new studies provide evidence for: 1) increased respiratory symptoms among young adults in association with annual average SO_2 levels of 115 $\mu\text{g}/\text{m}^3$ (Chapman et al.,

Table 3-2. SUMMARY OF RECENT (1982-86) EPIDEMIOLOGICAL STUDIES PROVIDING MOST USEFUL CONCENTRATION/RESPONSE INFORMATION ON LONG-TERM EXPOSURES TO SO₂

Observed Effects	Time	Population	Observed Concentration Range SO ₂ (µg/m ³)	PM (µg/m ³)	Comments	Study
a) Communities Dominated by Large Point Sources						
Increased prevalence of chronic phlegm, cough	1976	~ 5,600 adults in 4 Utah towns at varying distances from large copper smelter	115	70 TSP/14 SO ₄ (5 yr average - 1971-76)	3 cleaner communities had 5-yr. SO ₂ between 11-35 µg/m ³ , 5-8 µg/m ³ SO ₄ ; little gradient across towns in TSP (50-70 µg/m ³). Results more consistent in non-smokers, women; consistent with previous 1970 survey. No lung function measurements. Any effects of SO ₂ likely attributable to high short-term peaks.	Chapman et al., 1985
Increased prevalence of cough	1979-82	~ 700 3rd-6th graders in 2 smelter and 2 control communities in Arizona	103	52 TSP/9-10 SO ₄ (3 yr average)	High short-term peaks in 1 smelter town (repeated 3-hr. avgs. ~ 1.0 ppm), as well as 2nd smelter area with elevated cough (avg. 24 hr max ~ 440 µg/m ³ ; 3 yr avg. SO ₂ ~ 50 µg/m ³ , 28 µg/m ³ TSP). 2 control areas had 3-yr SO ₂ < 14 µg/m ³ , 4-7 µg/m ³ SO ₄ , 58-60 µg/m ³ TSP. No trend with pollution in shortness of breath, wheeze, sputum production, lung function (FEV ₁).	Dodge, 1983; Dodge et al., 1985
b) Multiple Urban Area Comparisons						
Increased prevalence of cough, expectoratory illness in men. Upper respiratory disease in children. Reduced lung function in adults and children.	1974-76	~ 20,000 adults and children in 20 areas of 7 cities and 1 industrial region in France	13-127	20-150 Smoke 45-240 "Dust" (3 yr average)	Significant effects found only with SO ₂ ; PM measurements of questionable comparability (CDA, p. 3-43). Inconsistent trends within cities. No control for parental smoking in children; uncertain control for season; apparently incomplete statistical analysis.	PAARC 1982a,b

1983); 2) increased prevalence of cough in children (but not lung function changes) being associated with intermittent exposures to mean peak 3-hr SO_2 levels of ~ 1.0 ppm or annual average levels of $\approx 103 \mu\text{g}/\text{m}^3$ (Dodge et al., 1985); and 3) symptoms of lower respiratory disease and decrements in lung function in adults possibly associated with annual average SO_2 levels ranging without evident threshold from about 25 to $130 \mu\text{g}/\text{m}^3$ (PAARC, 1982a,b). In addition, the PAARC study suggests that upper respiratory disease and lung function decrements in children may also be associated with annual average SO_2 levels across the above range.

Some questions must be raised regarding the PAARC analysis, however: (1) SO_2 and PM indices were only tested in separate regressions resulting in potentially confounded results, especially given the remarkably low colinearity in the 2 pollutants; (2) The positive associations between SO_2 and lung function were significant for only one of the two SO_2 measurement methods used and are apparently dominated by a large difference in Rouen (an industrial city) between the SO_2 levels as measured by the two methods; (3) The large within-city and between-city differences as separate sources of variability were not assessed, possibly greatly reducing the statistical significance of estimated effects in this very large study. These and other uncertainties related to aerometry, the lack of control for parental smoking (for children), in controls for seasonal effects, and the counter-intuitive results for NO_2 further limit the confidence to be placed in the present results.

Correlations, and conclusions, from the Ware et al. (1986) study are weakened by the relatively low illness rates in one area (Carondolet, St. Louis) during periods of relatively high SO_2 levels and by the fact that after SO_2 levels declined there (from $184 \mu\text{g}/\text{m}^3$ in 1976 to $88 \mu\text{g}/\text{m}^3$ in 1977) and TSP

dropped only slightly ($125 \mu\text{g}/\text{m}^3$ to $104 \mu\text{g}/\text{m}^3$), illness rates increased slightly. Otherwise, reduced ventilatory function has been found to be significantly related to elevated SO_2 levels in only the PAARC study and possibly in the recent van der Lende et al. (1986) report, although the latter findings are considered too preliminary for risk assessment purposes. Similarly, the Schenker et al. (1983) study suggests increased risk of wheeze (but not cough or phlegm) associated with elevated SO_2 concentrations but specific effect levels are difficult to identify (CDA, p. 3-40).

Many of these studies in which high long-term SO_2 concentrations have been measured and correlated with health effects were conducted in areas around major point sources of SO_2 emissions (e.g., copper smelters, coal-fired power plants). It is therefore likely that the populations studied were exposed to repeated high short-term peak concentrations of SO_2 , primary sulfuric acid, and other stack related particles. In light of the controlled human and animal exposure studies on SO_2 and sulfuric acid discussed previously in this paper and in the 1982 PM staff paper (EPA, 1982c), it appears likely that the effects associated with SO_2 in these studies were at least in part related to intermittent, acute respiratory insults. None of these studies, however, have attempted to separately analyze those individuals expected to be most responsive to short-term SO_2 or other exposures, i.e., asthmatics and atopics.

IV. FACTORS TO BE CONSIDERED IN SELECTING PRIMARY STANDARDS FOR SULFUR OXIDES

This section, drawing upon the previous summary of newly available scientific information, enumerates key factors that should be considered by the Administrator in decisions on the primary standards for sulfur oxides. The staff conclusions and recommendations on the most appropriate policy options update and supplement those made in the 1982 staff assessment. Where the original conclusions and recommendations and supporting rationale are unchanged by the newly available information, they are summarized without restating the supporting discussion. Particular emphasis is placed on aspects of the new information that amend or revise the original assessment. The key standard components discussed are the levels and averaging times for the primary standards. In addition, a summary assessment of the relative protection afforded by alternative standard combinations is presented.

A. Levels and Averaging Times of the Standards

1. General Considerations

The major scientific basis for selecting SO₂ standards that have an adequate margin of safety comes from controlled human exposures and community epidemiological studies, with mechanistic support from toxicological, deposition, and air chemistry investigations. The limitations of available controlled human studies for quantitative evaluation of ambient exposures of populations are summarized in the CD and in the CD addendum. Such studies provide accurate measurements of specific pollutant exposures, but are limited in exposure regimes, numbers and sensitivity of subjects, and severity of effects tested, and may involve artifacts not representative of ambient exposures. Community epidemiological studies, while representing real world conditions, can only provide associations between a complex

pollutant mix and a particular set of observable health endpoints. It follows that, although the scientific literature provides substantial information on the potential health risks associated with various levels and exposure patterns of SO₂, selection of appropriate levels, form, and averaging times remains largely a public health policy judgment.

The following sections present a brief staff assessment of how the concentration/response relationships suggested by the most significant controlled human and epidemiological studies in the CD addendum supplement the quantitative information previously assessed in the 1982 staff paper, and indicate how these studies may be applied in decision-making on standards for SO₂. The presentation also outlines a qualitative assessment of the key factors that affect the margin of safety associated with the ranges of standards derived from these studies. This assessment includes identification of those aspects of the qualitative literature that should be considered in establishing standards that provide an adequate margin of safety. Peak (< 1-hour), short-term (≤ 24-hour), and long-term (annual), exposures are discussed separately.

2. Peak (< 1-hour) Exposures

a) Derivation of Ranges of Interest from Controlled Human Exposure Studies.

Table 4-1 presents an updated staff assessment of the controlled human studies most useful in developing a range of interest for selecting a 1-hour SO₂ standard. Both recently published studies and those assessed in the 1982 staff paper are included. The table focuses on those studies involving free breathing (chamber) or facemask exposures, which provide the closest approximation of natural breathing. Studies in which subjects breathed through mouthpieces have also been considered. Although caution is necessary in extrapolating mouthpiece study results to ambient conditions,

Table 4-1. UPDATED STAFF ASSESSMENT OF KEY CONTROLLED HUMAN STUDIES

SO ₂ Concentration (5-60 minutes)	Observed Effects ¹	Comments/Implications
1-2 ppm	Substantial changes in 8 of 12 subjects (Δ SRaw 100-600%) exposed to 2 ppm. At 1 ppm, functional changes (Δ SRaw 170-200%), symptoms in free breathing asthmatics at moderate exercise ²	Effects range from moderate to incapacitating for some individuals. At 2 ppm, 80% of mild asthmatics could experience at least a doubling of SRaw. Some might not tolerate exposure at moderate exercise. Approx. 60% at 1 ppm could experience at least a doubling of SRaw. ³ Some asthmatic mouth breathers have significant bronchoconstriction at 2 ppm even at light activity.
0.6-0.75 ppm	Functional changes (Δ SRaw 120-260%), symptoms in free breathing asthmatics at light-moderate exercise ⁴	Effects indicative of clinical significance on average, changes were mild to moderate although severe for some individuals; 25-50% of mild, free-breathing asthmatics at moderate exercise could experience at least a doubling of airway resistance. ³
0.5 ppm	Significant functional changes (Δ SRaw 50-100%), symptoms in free breathing asthmatics at moderate, but not at light exercise. ⁵ At heavy exercise, Δ SRaw 220-240%. ⁶	On average, mild responses at moderate or higher exercise, symptoms possibly of clinical significance; severe responses for some individuals. About 20-25% could experience at least a doubling in airway resistance.
0.4 ppm	Functional changes (Δ SRaw 70%), symptoms in free breathing asthmatics at moderate-heavy exercise ⁷	Lowest level of clinically significant response for some free breathers. Approx. 25% of mild, free breathing asthmatics could experience a doubling in airway resistance. ³
0.1-0.3 ppm	No effects in free breathing asthmatics at light exercise. Slight but not significant functional changes in free-breathing subjects at moderate-heavy exercise (0.25 ppm) ⁶ , but not at lower levels. ⁷	Significant effects unlikely at moderate exercise. Effects of SO ₂ indistinguishable at heavy exercise. Possibility of more significant responses in small percentage of sensitive asthmatics at 0.28 ppm. ³

¹Unless otherwise noted, (Δ SRaw %) reflects group mean increase over clean air control at rest. Light, moderate, heavy exercise refers to ventilation rates approximating < 35 L/min, 40-45 L/min, and > 50 L/min, respectively. Effects reflect results from range of moderate temperature/humidity conditions (i.e., 7-26°C, 36-90% RH). Studies at 0.5-0.6 ppm indicate that exercise-induced bronchoconstriction associated with cold and/or dry air exacerbates response to SO₂ while warm, humid air mitigates asthmatic responses relative to moderate conditions.

²Schacter et al. (1984); Roger et al. (1985); Horstman et al. (1986).

³Horstman et al., (1986).

⁴Hackney et al. (1984); Schacter et al. (1984); Linn et al. (1983a,b, 1984a,b,c, 1985a).

⁵Kirkpatrick et al. (1982); Linn et al. (1984b); Roger et al. (1985); Schacter et al. (1984).

⁶Bethel et al. (1983a,b; 1985).

⁷Linn et al. (1983b, 1984a).

it does not appear that substantial differences exist in SO_2 -induced responses for the different breathing modes when account is made for the partitioning of oral and nasal airflow components in oronasal breathing (see Appendix A). Inferences made in the "implications" column are derived from observations made by the investigators or in the CD addendum. The percentage of asthmatics showing a potentially clinically significant increase in airway resistance (100%) is derived from Horstman et al. (1986) (See Figure 3-2).

Table 4-1 indicates that functional changes and symptoms are likely in a large percentage of freely breathing asthmatics exposed to 5 to 10 minute peaks of SO_2 between 1 and 2 ppm while involved in light to moderate exercise ($V_e \sim 30\text{-}50$ L/min), comparable to daily activities such as climbing stairs and light bicycling or jogging. At comparable exercise rates ($V_e \sim 40$ to 48 L/min), Linn et al. (1983a,b) found "clinically and physiologically significant responses" in free breathing young adult asthmatics exposed to 0.75 ppm and to 0.6 ppm SO_2 . Several studies report significant asthmatic responses at 0.5 ppm with oronasal (free or facemask) breathing at moderate-heavy exercise ($V_e \sim 40\text{-}60$ L/min) (Kirkpatrick et al., 1982; Bethel et al., 1983b; Roger et al., 1985) but no substantial symptomatic or functional effects at lower ventilation rates (27-40 L/min) (Linn et al., 1982; Bethel et al., 1983b; Roger et al., 1985; Schacter et al., 1984).

Asthmatics exposed to 0.4 ppm SO_2 at a moderate to heavy exercise rate ($V_e \sim 48$ L/min) showed on average a moderate increase in S_{Raw} and a mild increase in group mean symptom score, with substantial bronchoconstriction in some individuals and one subject requiring medication to relieve symptoms (Linn et al., 1983b). Studies of free breathing exposures at lower

concentrations (0.1 to 0.3 ppm) suggest marginal, if any, group responses only with 0.25 ppm at heavy exercise (50-60 L/min). Any effect of SO₂ is negligible compared to exercise at these levels (Linn et al., 1984b; Bethel et al., 1985). The CD addendum concludes from these observations that "some SO₂-sensitive asthmatics are at risk of experiencing clinically significant (i.e., symptomatic) bronchoconstriction requiring termination of activity and/or medical intervention when exposed to SO₂ concentrations of 0.4 to 0.5 ppm or greater when this exposure is accompanied by at least moderate activity" (CDA, pp. 5-10).

The 1982 staff paper outlined several considerations that are important in evaluating these results in the context of decision making on a standard to limit peak (5-10 minute) SO₂ exposures. The following discussion represents an update of those considerations.

1) Health Significance of the Observed or Anticipated Effects

Although little controversy exists that a full asthma attack represents an adverse health effect, the relative significance of some of the less severe responses observed in the above controlled human studies is open to question. Based on the 1982 CD discussion of these matters, the staff paper concluded that the results of these studies begin to be of some concern when bronchoconstriction is accompanied by noticeable symptoms. This is an imprecise criterion, however, as not all studies report symptoms and symptom reports are not always a reliable indicator of clinical status. The CD Addendum identifies at least four variables frequently measured that can be used to classify the medical significance of responses observed in the studies (see Figure 4-1). These variables are a) change in SRaw; b) duration of effect of SO₂; c) changes in spirometry, chiefly FEV_{1.0}; and d) types of symptoms and relative discomfort. As noted in the CD Addendum, "This table is

GRADE OF RESPONSE	NONE	MILD	MODERATE	SEVERE	INCAPACITATING
CHANGE IN SRAW	NO CHANGE	INCREASE LESS THAN 100%	INCREASE UP TO 200%	INCREASES MORE THAN 200%	INCREASES >> 200%
DURATION OF EFFECT	NA	SPONTANEOUS RECOVERY < 30 MIN	SPONTANEOUS RECOVERY < 1 HR	BRONCHODILATOR REQUIRED TO RESOLVE SYMPTOMS	EMERGENCY TREATMENT REQUIRED
CHANGE IN SPIROMETRY FEV _{1.0} , FVC	NO CHANGE	NO CHANGE	DECREASE BUT < 15%	DECREASE > 15%	DECREASE >> 15%
SYMPTOMS	NO RESPIRATORY SYMPTOMS	MILD SYMPTOMS NO WHEEZE OR CHEST TIGHTNESS	SOME WHEEZE* OR CHEST TIGHTNESS	OBVIOUS WHEEZE* MARKED CHEST TIGHTNESS BREATHING DISTRESS	SEVERE BREATHING DISTRESS*

* STATISTICALLY SIGNIFICANT CHANGE

Figure 4-1. Gradation of physiological responses to SO₂ (CDA, Figure 7).

not intended to provide a quantitative description of what does or does not constitute an adverse health effect but is primarily intended to demonstrate that there are an array of responses and to assist the reader in judging the relative severity of the different responses which are described" (CDA, p. 4-8).

The scientific literature does not provide sufficient information to specify an SO₂ concentration at which the observed effects can themselves be considered adverse or serve as indicators of potentially more serious consequences. In making such a judgment, the Administrator should consider, among other factors, the following:

a) In almost all cases, the bronchoconstriction and symptoms observed appear to have been transient and reversible. Sheppard et al. (1983), however, reported that for two subjects, exposure to 0.5 ppm SO₂ via mouth-piece with hyperventilation produced severe bronchoconstriction that lasted longer than 45 minutes. In other studies, asthmatic subjects have been removed from the chambers because of severe responses accompanying free breathing exposures at 0.5 ppm and higher (Roger et al., 1985). In other studies, some subjects needed bronchodilator medication after exposure to 0.4-0.6 ppm (Linn et al., 1983b, 1984a,b). Although direct evidence of long-term consequences from repeated peak exposures is not available, the possibility of such effects cannot be ruled out.

b) At concentrations less than 0.4 ppm with free breathing, group mean functional changes were moderate to small (Δ SRaw ~ 0 to 70% over baseline) and within the range of variability observed for day to day changes in many asthmatics. At 0.6-0.75 ppm, group mean effects were more substantial (Δ SRaw ~ 200% over baseline).

c) Most studies utilized mild, young adult or adolescent, non-smoking asthmatic volunteers. Furthermore, the subjects were exposed only when they were asymptomatic and without apparent respiratory tract infections or allergic responses. Even among the otherwise well defined groups of relatively mild asthmatics studied, there was great variability in the magnitude of bronchoconstriction induced by SO_2 . As illustrated by the data derived from Roger et al. (1985) in Figure 3-2, the SO_2 concentration necessary to increase S_{Raw} by 100% or more in freely breathing asthmatics at 42 L/min was 0.75 ppm for 50% of the subjects, and ranged between approximately 0.3 and 1.4 ppm in 80% of the subjects. Even more sensitive individuals may exist in the population of "mild" asthmatics. Individuals with more severe asthma may also be more sensitive to SO_2 -induced bronchoconstriction, but the evidence on this issue is inadequate. The consequences of any particular functional change in a more severe asthmatic is thought to be of greater concern than in a mild asthmatic. However, more severe asthmatics may be somewhat protected from SO_2 because of their greater reliance on medication and their reduced tolerance to sustained levels of moderate to high exercise.

d) Although the reported responses are not generally interpreted as overt asthma attacks, the combination of bronchoconstriction and symptoms might be perceived by some subjects as a "mild" attack; this could result in discomfort, the need for medication, and curtailment of desired physical activities. According to Linn et al. (1983b), the responses of their subjects at 0.6 ppm "might be judged to show adversity in that the subjects

sense of well being was clearly diminished, their degree of air-flow obstruction seemed to impair physical performance meaningfully, and drug treatment was clinically indicated in a few. On the other hand, possibly arguing against a judgment of adversity, are the observations that the effects were quickly reversible, were similar to effects produced by exercise even in clean air, and did not prevent the subjects from carrying out their duties (completing the exposure protocol)."

The staff obtained additional guidance on the physiological or health significance of asthmatic responses in the controlled exposure studies through discussions with a number of experts in the field (Cohen, 1984). Some experts felt that the relatively mild, transient, and reversible effects are not of physiological significance given the current widespread use of effective medication. In contrast, others felt that despite asthmatics' sensory accommodation and learning to manage attacks through medication or altered activity, even subtle functional changes are significant and potentially serious especially when accompanied by symptoms. Several pointed out that there may be persistent, undetected effects (e.g., residual obstruction) associated with even "mild" episodes which may increase airway reactivity and predispose the individual to further insults (e.g., infections, other bronchoconstrictive agents). Furthermore, these experts agreed that any asthmatic experience is alarming and in different degrees, disabling. They felt that the effects observed at 0.5 ppm SO₂ would, at a minimum, affect an individual's lifestyle by causing discomfort, an increase in their medication usage, or discontinuance or restriction of their activity.

2) Relative Effect of SO₂ Exposure Compared to Exercise, Other Stimuli; Exercise-induced bronchoconstriction, without pollutant exposure, is a relatively common occurrence for many asthmatics and SO₂ represents one of many potentially encountered stimuli that can cause asthmatic reactions (see SP, pp. 66-67). Consistent with previous findings discussed in the 1982 staff paper, recent studies find that SO₂ enhances the bronchoconstrictive effects due to exercise. Roger et al. (1985) report that the effects of moderate exercise ($V_e \sim 42$ L/min) in inducing bronchoconstriction is roughly equal to that of 0.5 ppm SO₂ while the effects of 0.25 ppm SO₂ on asthmatics are insignificant compared to those caused by moderate-heavy exercise. The exercise rate in this study is roughly equivalent to light jogging or climbing several flights of stairs (SP, Appendix A).

Cold (< 6°C) and/or dry air has been found to exacerbate the effects of SO₂ in exercising asthmatics, producing effects greater than those seen at normal temperatures. SO₂ at concentrations as low as 0.3 ppm may measurably potentiate the effect of cold air (Linn et al., 1984b) which may be possible in ambient winter conditions in the U.S. On the other hand, effects with warm, humid temperatures are less than those seen in conditions typical of most laboratory studies.

3) Exposure Considerations

Peak 1-hour SO₂ levels in excess of 0.5 ppm are rare with current U.S. air quality, and almost always occur only in the vicinity of major point sources. Shorter term (5 to 10 minute) peaks at these levels are somewhat more common, but no systematic data exist. Moreover, indoor SO₂ levels are almost always substantially lower than outdoor levels (EPA, 1982b; pp. 5-117). Thus, effects appear likely for situations involving

asthmatics undergoing light to moderate exercise outdoors relatively near (< 10 km) sources of SO₂ in conditions resulting in peak (> 0.5 ppm, 5 to 10 minutes) SO₂ levels. Staff estimates of the probability of such exposures near large sources under alternative standards are summarized in the next section (IV.B). Asthmatics may also be exposed to more frequent peaks of limited durations (< 30 seconds to 2 minutes) around numerous smaller industrial and commercial sources (Section IIB). It is not currently possible to determine whether exposures of such limited duration would produce effects approaching those seen at the 5 to 10 minute exposures used in most of the studies to date.

To the extent such sources produce repeated frequent short-term peaks, the findings of temporary adaptation response may be of some significance. Within a single day, repeated episodes of exercise with elevated SO₂ concentrations would be expected to produce mitigated responses. Since tolerance appears to be short-lived (<5 hrs.), however, it would not afford protection against SO₂ on subsequent days, nor necessarily on the same day.

Some data suggest that rapid rises in SO₂ levels, such as those involved in many of the controlled studies, are more likely to produce effects than are more gradual rises. As discussed in the 1982 staff paper, however, a rapid rise could result from a) movement from indoors to outdoors, b) onset of exercise resulting in a rapid rise in SO₂ at sensitive respiratory tract receptors, c) movement into an area of peak levels (by vehicle or otherwise), as well as, d) an actual rapid increase in ambient levels at a point.

4) Variance about the 1-hour average

The controlled studies discussed in Section III indicate that effects occur within 5-10 minutes but do not necessarily worsen with continued exposure to SO₂ over the course of an hour. Five and ten minute averages will vary about the 1-hour mean. Thus, for an area just attaining a 1-hour standard of 0.5 ppm, 5 or 10 minute peaks would be higher.

Analyses of recent data (Section II), indicate that the peak is likely to be within a factor of 2 (1.5 to 1.8 of the mean) or less than 1.0 ppm.

Based on the above evaluation of the more recent studies and related factors, the staff revises its original recommended range of interest for a possible 1-hour SO₂ standard to 0.2 to 0.5 ppm. The lower bound of 0.2 ppm represents a 1-hour level for which maximum 5 to 10 minute peak exposures are not likely to exceed 0.4 ppm, the lowest level at which the CD Addendum indicates a risk of clinically significant responses for asthmatics engaged in moderate (or higher) activity levels. Based on normal air quality variations, a 1-hour standard at the upper bound of the range of 0.5 ppm would permit 5-10 minute peaks as high as 1.0 ppm during the peak hours, and would permit multiple hours in which the 5-10 minute peak would exceed 0.5 ppm, even when the 1-hour average is within this range. The risk of substantial effects with such exposures is higher.

Independent of frequency of exposure considerations, 1-hour concentrations at the high end of the above range may not provide a substantial margin of safety for exercising asthmatics. The low frequency with which such peak values would occur in the presence of active sensitive subjects is, however, a mitigating factor that should be examined in determining the margin of safety provided by alternative standards.

b) Additional Factors to be Considered in Evaluating Margin of Safety and Risks--Peak Exposures

The data do not suggest other groups that are more sensitive than asthmatics and atopics to single peak exposures. To the extent that the suggested range is protective of asthmatics and atopics, the risk of functional effects in other sensitive individuals appears small. Other effects of concern (aggravation of bronchitis, increased respiratory illnesses) have not been evaluated adequately in controlled human studies, but epidemiological

evidence suggests that they may result from repeated peak exposures over longer time periods. Potential interactions of SO₂ and ozone have not been investigated in the more recent literature.

The potential pollutant interactions and other considerations listed above should be considered in determining the need for and evaluating the margin of safety provided by alternative 1-hour standards.

3. Short-Term (24-hour) Exposures

a) Derivation of Ranges of Interest from Epidemiological Studies

An updated staff assessment of the most useful epidemiological studies for deriving ranges of interest for 24-hour standards is summarized in Table 4-2 and discussed below.

Table 4-2. UPDATED STAFF ASSESSMENT OF SHORT-TERM EPIDEMIOLOGICAL STUDIES

Effects/ Study	Measured SO ₂ - µg/m ³ (ppm) - 24 hour mean			
	Daily Mortality in London ¹	Aggravation of Bronchitis ²	Small, Reversible Declines in Children's Lung Function ³	Combined Effects Levels
Effects Likely	500-1000 (0.19-0.38)	500-600 (0.19-0.23)	-	500 (0.19)
Effects Possible	-	<500 (0.19)	250-450 (0.10-0.18)	250 (0.10)
No Effects Observed	-	-	100-200 (0.04-0.08)	<200 (.08)

¹Deviations in daily mortality during London winters (1958-1972). Early winters dominated by high smoke and SO₂, principally from coal combustion emissions, and with frequent fogs (Martin and Bradley, 1960; Ware et al., 1981; Mazumdar et al., 1981, 1982; Schwartz and Marcus, 1986).

²Examination of symptoms reported by bronchitics in London. Studies conducted from the mid-1950's to the early 1970's (Lawther et al., 1970).

³Studies of children in Steubenville (1978-80) and in the Netherlands (1985-86) before, during, and after pollution episodes characterized by high particle and SO₂ levels (Dockery et al., 1982; Dassen et al., 1986).

The "effects likely" row in Table 4-2 is unchanged from the 1982 assessment. The CD addendum relies on the original London mortality and bronchitic studies as those most appropriate in concluding that notable increases in excess mortality and exacerbation of bronchitic symptoms may occur above $500 \mu\text{g}/\text{m}^3$ BS and SO_2 . With regard to increased mortality, greater certainty with respect to effects occurs when both pollutants exceed about $750 \mu\text{g}/\text{m}^3$. These estimates represent judgments of the most scientifically reliable "effects levels" for daily smoke and SO_2 at least in the context of historical London pollution episodes.

Because of the severity of the health endpoints in these studies and the need to provide an adequate margin of safety in standard setting, it is important to determine whether the data suggest the possibility of health risks below these "effects likely levels". As discussed in the CD addendum, the London mortality studies and reanalyses support the possibility of effects due to particles below $500 \mu\text{g}/\text{m}^3$ with no obvious threshold.

The situation with respect to SO_2 , however, is less clear. The 1982 CD notes that results from a selected group of subjects suggested that $500 \mu\text{g}/\text{m}^3$ SO_2 (and $250 \mu\text{g}/\text{m}^3$ BS) may not be absolute thresholds for the most sensitive bronchitis patients, although the lead author of the study strongly objects to this interpretation (Lawther, 1986). On the other hand, the 1982 staff assessment previously concluded that the available evidence on daily mortality did not suggest a significant risk of increased mortality for exposures to SO_2 alone at concentrations below the likely effects levels.

The recent London mortality reanalyses provide differing results regarding the effects of SO_2 . Shumway et al. (1983) did not attempt to separate the effects of the two pollutants and found that their association

with daily increases in mortality were nearly identical with no apparent threshold. Mazumdar et al. (1982) found no consistent trend in mortality with increasing SO_2 below $700 \mu\text{g}/\text{m}^3$ (0.27 ppm) and that the component of London mortality explained by pollution in the 1958-72 winters is almost entirely due to smoke across all levels considered. For days with BS and SO_2 below $500 \mu\text{g}/\text{m}^3$, the association between mortality and pollution persisted for smoke and not SO_2 . Schwartz and Marcus (1986) in joint regressions for these winters found BS was significantly associated with mortality, independent of SO_2 effects. While the effects of SO_2 and BS cannot clearly be separated due to the high degree of their covariance in this data set, it does not appear that the recently published analyses suggest a revision to the previous assessment, which concluded there was not a significant risk of increased daily mortality with SO_2 alone below the effects likely levels.

The studies of school children in Table 4-2 exposed to peak SO_2 and particle concentrations during pollution episodes suggest small, but significant, reversible declines in lung function. The studies suggest the possibility of effects below the low end of the original range of interest ($365 \mu\text{g}/\text{m}^3$ or 0.14 ppm) down to levels as low as approximately $250 \mu\text{g}/\text{m}^3$ (0.10 ppm) with more certainty at levels around $450 \mu\text{g}/\text{m}^3$ (0.18 ppm). Again, it is difficult to distinguish the effects of the two pollutants though a more consistent trend of reduced lung function with higher TSP, and not SO_2 , was reported by Dockery et al. (1982). Given that SO_2 alone has not been observed to cause altered clearance or lung function in animals or humans

in controlled laboratory conditions without very high short-term peaks (> 1-5 ppm) (EPA, 1982a,b), it may be that the observed declines in lung function during and after the pollution episodes were due to the elevated particle levels (including the transformation products of SO₂) either acting alone or in the presence of SO₂, rather than SO₂ alone. Alternatively, very high peak SO₂ concentrations on the order of minutes may have accounted for the lung function decrements, though this does not seem likely.

Therefore, caution should be applied in using the recent episode studies in the context of evaluating the range of interest for SO₂ alone. While effects may be associated with levels between 250-450 µg/m³ (0.10-0.18 ppm), it is questionable to what extent SO₂ was a factor in causing the observed responses.

In summary, the available data indicate that the upper bound for the range of interest for 24-hour SO₂ standards remains at 500 µg/m³ where effects appear to be likely. Although consideration should be given to a lower bound of 250 µg/m³ (0.10 ppm), it is not clear whether important effects are caused by SO₂ at levels below the current standard level (365 µg/m³, 0.14 ppm) which was previously judged - and still appears - to provide adequate protection.

b) Summary of Factors to be Considered in Evaluating Margin of Safety -- Short-Term Exposures

The 1982 staff paper identified a number of factors to be considered in developing a 24-hour standard with a margin of safety. The staff finds that this original discussion (SP, pp. 75-78) is still appropriate. In summary, the factors include:

- 1) Interaction with ozone, particles, and other pollutants as well as fog.
- 2) Relative exposure in the U.S. compared to the British studies.

- 3) Risk for other sensitive groups and effects not evaluated in the more quantitative data, and
- 4) Whether the 24-hour standard acts alone or in concert with a new 1-hour standard.

4. Long-Term (Annual) Exposures

Based on the 1982 assessment, the staff concluded that although the possibility of effects from continuous low-level exposures to SO₂ could not be ruled out, no quantitative rationale could be offered to support a specific range of interest for an annual standard given the inconclusive nature of the available epidemiological data. As discussed in Section III, several recent community studies suggest increased risk of respiratory symptoms (cough, phlegm production, wheeze) in populations (children and adults) exposed to high (>100 µg/m³), long-term levels of SO₂, with and without high particle concentrations. The majority of these studies were conducted in areas subjected to intermittent short-term peak SO₂ concentrations resulting from point source emissions (Chapman et al., 1985; Dodge, 1983; Dodge et al., 1985; Schenker et al., 1983). A major concern, therefore, is whether repeated SO₂ peaks permitted by 24-hour or 1-hour standard ranges in area-source dominated population centers might, after some long time period, result in increased risk of the effects noted, along with other effects suggested by animal data (EPA 1982a,b).

One recent study (PAARC, 1982a,b) demonstrating associations between SO₂ and respiratory health effects did not focus on point-source dominated exposures. Increased respiratory symptoms and disease in adults and children were associated with SO₂, but not particles, across a range from 25 to 130 µg/m³ with no apparent threshold (CDA, p. 3-55). In addition, unlike in any other study, associations between SO₂ and reduced lung function

were detected. As noted in Section III, a number of questions regarding the aerometry, statistical analyses, and interpretation of this work limit the reliance that can be placed on the conclusions of this study at present.

While no single study may provide strong evidence for substantial risks, there does appear to be some consistency across results indicating a possibility of respiratory impacts associated with either low-level, long-term exposures to SO_2 or, more certainly, with repeated exposures to peak SO_2 levels over long periods. In essence, the recent studies do add support to previous staff recommendations to retain an annual primary standard. This recommendation was in part based on the finding that elimination or substantial relaxation of the current annual standard would result in increased exposures to large numbers of people in several heavily populated urban centers (Frank and Thrall, 1982). Such exposures could lead to increased risks of health effects that are not readily measured in controlled studies but for which there is qualitative evidence, summarized in the 1982 staff paper. These possible effects include effects on clearance and other host defense systems, and to a lesser extent, potential mutagenicity or co-carcinogenicity of SO_2 (SP, pp. 78-79). In addition, the long-term standard serves to limit emissions from numerous smaller sources that have recently been found to produce brief short-term peaks (< 30 seconds to 2 minutes) that are of potential concern to asthmatics (see Section IIB above). Pending resolution of the issues raised by the new studies, the staff recommends maintaining an annual standard at about the current level of 0.03 ppm ($80 \mu\text{g}/\text{m}^3$).

B. Analysis of Relative Protection Afforded by Alternative Standards

An essential consideration in evaluating potential standards is the relative protection afforded by standards with different averaging times

and levels. A preliminary staff assessment of this issue is presented in the 1982 SP (pp. 79-83, Appendix D). This assessment, based on analysis of air quality data (Frank and Thrall, 1982; Johnson, 1982), air quality modeling (Burton et al., 1982), and source/population information (Anderson, 1982), found that no single averaging time (annual, 24-hr, 3-hr, 1-hr) would provide the same degree of protection and control afforded by the other averaging times in all situations. The current 24-hour standard would significantly limit 1-hour peaks in the range of interest from occurring in most population oriented sites, but would allow multiple exceedances of these values in many point source oriented sites. Similarly, the 24-hour standard limits high annual values in most, but not all sites of interest. The current 3-hour secondary standard limits 1-hour peaks even more than the 24-hour standard, but does not materially affect long-term urban values. In essence, based on that preliminary analysis of alternative averaging times, the staff concluded "that implementation of the current suite of SO₂ standards (annual, 24-hour, 3-hour) provides substantial protection against the direct effects of SO₂ identified in the scientific literature" (SP, pp. 82-83).

Since closure on the 1982 staff paper, the staff has continued to analyze relationships among averaging times and relative protection afforded. Based on the above updated assessment of effects associated with both 24-hour and annual exposures, the staff finds that the above conclusions concerning protection provided by the current standards remain demonstrably valid. The staff has found the most critical aspect of examining the relative (or alternative) standards to be in relation to peak exposures associated with effects in asthmatics. Over the past several years, the staff has developed tools to permit analysis of substantially greater detail than previously possible. These tools and the results of their application

to examine 1) current standards, 2) emissions typical of current conditions, and 3) alternative standards are presented in detail in separate reports (EPA, 1986c; 1986d). The following discussion summarizes the major findings from these reports.

Population exposure simulations require detailed analyses of both air quality and population patterns. The EPA (1986d) describes a population exposure study around four utility power plants each located in or near an urban area. The decision to focus that analysis on power plants was guided by earlier studies (Frank and Thrall, 1982; Burton et al., 1982) which showed they were the source category most likely to produce high, short-term levels of SO_2 in populated areas. Other large sources, such as smelters or Kraft pulp mills, however, can also produce such peaks. A complete risk assessment would combine exposure results with detailed exposure-response functions. To reduce the complexity of this analytic problem to a manageable size, the staff developed a benchmark called an Exposure of Concern (EOC). This benchmark permitted fixing a concentration, averaging period, and exercise rate above which effects of concern could be expected in some fraction of asthmatics. Based on the health studies and analyses described above, the benchmark EOC most often used was defined as an asthmatic exposed at or above 0.5 ppm SO_2 for 5 minutes while at an activity level associated with a ventilation rate at or above 35 L/minute. At these levels, on the order of 25% of asthmatics might experience a doubling of airway resistance (Figure 3-2). In some of the work, other concentration levels and averaging periods were also examined. The EOC defined above is not intended to define a threshold of response, but rather as a level where a significant fraction of individuals so exposed might experience potentially adverse effects.

The air quality modeling estimated the probability that the 5-minute peak equalled or exceeded 0.5 ppm. These probability estimates of exceeding a target level (0.5 ppm) provided the air quality basis for the exposure calculation. The EPA's NAAQS Exposure Model (NEM) (Biller et al., 1981) was modified to incorporate these probability estimates. NEM is designed to simulate daily population movement around an urban area accounting for travel patterns, activity levels, and microenvironment (e.g., indoor vs. outdoor). The population and travel data were specific to the urban areas being studied. The activities which are defined in NEM as "High" correspond to ventilation ≥ 35 L/minute. The use of air quality probability estimates meant that it was possible to express the exposure results as a probability weighted distribution and allowed estimation of the expected number of exposures.

The findings of the exposure analysis are subject to a number of uncertainties inherent in both air quality modeling and large population simulations. The results are conditioned by the analytic assumptions made. The exposure analysis identifies some 16 separate sources of uncertainty and error. Among the more significant are: 1) Lacking activity pattern and residential location data for asthmatics, it was assumed that the geographic distributions, and activity patterns and ventilation rates for asthmatics are the same as for the general population. Although this may not be an unreasonable assumption for most mild asthmatics, it undoubtedly overstates the time spent at elevated ventilation rates for more severe cases; 2) Power plants were assumed to operate at 100% capacity. Sensitivity analyses indicate that exposures are overestimated because of this assumption; and 3) Although care was taken to select a representative sample of plants/exposure regimes, only four power plants were modeled. Nonetheless, despite

the caveats noted above as well as others in the reports, the results do provide an indication of both current exposures and those which might occur under alternative standards around large utility power plants.

The exposure analysis results in EPA (1986d) include air quality levels, the expected number and percent of asthmatics living in the vicinity of each plant that experiences one or more EOC per year, and the highest probability of an EOC for any single asthmatic. Because of variations in population around plants and the tendency for the maximum probability of exposure to approach one under a variety of scenarios, the fraction (%) of asthmatics with one or more EOC/yr is the most useful metric for comparing results around different plants. This number is, however, somewhat sensitive to the size of the area modeled (EPA, 1986d).

The results of the analysis of the fraction of asthmatics with an EOC under 1) current emissions, and 2) maximum emissions assuming the current standards are just met, are displayed in Figure 4-2. With current emissions, approximately 0.2 to 13% of resident asthmatics are predicted to experience at least one 5-minute exposure to 0.5 ppm per year while at moderate or higher exercise. With the exception of Eddystone, this represents on the order of one to four thousand asthmatics (assuming 4% of the population is asthmatic) for each plant. With the exception of Eddystone, the maximum probability of an EOC for "most exposed" individual approaches unity at all plants. The results for just meeting the current standards are comparable to the "current" case but with 3 of 4 plants showing increases in predicted EOC fraction. In part, such increases are due to assumptions regarding implementation, which reflect current practice in some areas of the country, but are less restrictive than more strict compliance requirements in practice in other areas. The 3-hour standard tends to be controlling for large, more isolated plants, while the 24-hour standard controls in more urban locations.




Figure 4-2. Expected percentage of asthmatics with one exposure of concern (> 0.5 ppm SO_2 for 5 minutes, "high" activity) per year in vicinity of four selected power plants. For 1) current emissions and 2) just meeting current SO_2 NAAQS assuming a 30 day emission limit (EPA, 1986d). The degree of protection varies significantly with implementation approach.

The results of the exposure analysis for alternative 1-hour standards selected from the range of interest are illustrated in Figure 4-3. Standards in this range would reduce the EOC fraction to under 4% for all plants modeled, but still do not eliminate all such exposures. A standard of 0.4 ppm, for example, would protect over 98% of potentially exposed asthmatics from an EOC. The maximum probability of an individual EOC for the range illustrated is 0.1 to 0.9.

The results of the exposure analysis for utilities should be viewed in light of the assumptions and uncertainties noted above. In addition, although utility power plants account for the majority of SO₂ emissions in the U.S., recent work has shown that other smaller sources may also produce peak exposures (Section II). Around smaller sources (e.g., industrial or commercial boilers), limited duration peaks in excess of 0.5 ppm are due either to low persistence meteorological events or, if the facility has a short stack, may be due to the phenomenon of building downwash. In either event, the peaks are likely to be of very short duration (less than 30 seconds to two minutes). Because the meteorological events causing the peaks are not well characterized and are not normally addressed in standard EPA dispersion models, a complete analysis of the situation around smaller plants is not feasible. Very rough estimates indicate that the populations at risk of an exposure in such situations may be large. However, given the very short duration of most such peaks, their health significance for exercising asthmatics is uncertain. Furthermore, it is not clear that a 1-hour standard would further limit such exposures.

In summary, the staff analysis of relative protection afforded by alternative standards results in the following conclusions:



1-HOUR NAAGS

Figure 4-3. Expected percentage of asthmatics with one exposure of concern (> 0.5 ppm SO₂ for 5 minutes, "high activity") per year in vicinity of four selected power plants. For just meeting alternative one hour standards (EPA, 1986d).

- 1) The current standards provide substantial protection against the effects identified as being associated with 24-hour and annual exposures.
- 2) The current standards - as reflected by current emissions or emissions when the standards are just met with somewhat less restrictive implementation assumptions - also provide some limit on peak SO₂ exposures of concern for asthmatics. In some cases, however, up to 10 to 15% of the sensitive population could be exposed once a year to levels ≥ 0.5 ppm for 5 minutes, while at elevated ventilation.
- 3) The range of 1-hour standards analyzed (0.25 to 0.5 ppm) would provide increased protection against such exposures, limiting the fraction of asthmatics exposed to less than 4%.

C. Summary of Staff Conclusions and Recommendations

The major updated staff conclusions and recommendations made in Section IV, A-B are briefly summarized below:

- 1) The more recent data provide additional support for the earlier staff recommendations regarding consideration of a new 1-hour SO₂ standard. Based on an updated staff assessment of controlled human exposures to peak (minutes to hours) SO₂ concentrations, the staff has revised the range of potential 1-hour levels of interest to 0.2 to 0.5 ppm (525 to 1300 $\mu\text{g}/\text{m}^3$). The lower bound represents a 1-hour level for which the maximum 5- to 10-minute peak exposures are unlikely to exceed 0.4 ppm, which is the lowest level where potentially significant responses in free (oronasal) breathing asthmatics have been reported in the CD addendum. The upper bound of the range represents a 1-hour level for which 5- to 10-minute peak concentrations

are unlikely to exceed 1 ppm, a concentration at which the risk of significant functional and symptomatic responses in exposed sensitive asthmatics and atopics appears high. In evaluating these laboratory data in the context of decision making on possible 1-hour standards, the following considerations are important: (a) the significance of the observed or anticipated responses to health, (b) the relative effect of SO₂ compared to normal day to day variations in asthmatics from exercise and other stimuli, (c) the low probability of exposures of exercising asthmatics to peak levels, and (d) 5- to 10-minute peak exposures may be a factor of two greater than hourly averages.

Independent of frequency of exposure consideration, the upper bound of the range contains little or no margin of safety for exposed sensitive individuals. The limited geographical areas likely to be affected and low frequency of peak exposure to active asthmatics if the standard is met add to the margin of safety. The data do not suggest other groups that are more sensitive than asthmatics to single peak exposures, but qualitative data suggest repeated peaks might produce effects of concern in other sensitive individuals. Potential interactions of SO₂ and O₃ have not been investigated in asthmatics. The qualitative data, potential pollution interactions, and other considerations listed above should be considered in determining the need for and evaluating the margin of safety provided by alternative 1-hour standards.


- 2) Based on a staff assessment of the recent short-term epidemiological data, the original range of 24-hour SO₂ levels of interest - 0.14 to 0.19 ppm (365 to 500 µg/m³) - still appears appropriate, although some

consideration could be given to the findings of physiological changes of uncertain significance at levels as low as 0.1 ppm. Earlier staff conclusions and recommendations concerning a 24-hour standard (SP, pp. 85-86) remain appropriate.

- 3) The previous staff assessment concluded that although the possibility of effects from continuous lower level exposures to SO₂ cannot be ruled out, no quantitative rationale could be offered to support a specific range of interest for an annual standard. The more recent epidemiological data, indicating associations between respiratory illnesses and symptoms and persistent exposures to SO₂ in areas with long-term averages exceeding 100 µg/m³, provide additional support for the original recommendation for retaining an annual standard at or near the current level 0.03 ppm (80 µg/m³). This recommendation was based in part on a finding that alternative short-term standards would not prevent annual levels in excess of the current standard in a limited number of heavily populated urban areas. In addition, recent evidence suggests smaller sources in urban areas may produce short duration peaks of potential concern. The long-term standard often serves to limit the emissions of numerous smaller sources in such areas. Given the additional information and the possibility of both chronic and acute effects from a large increase in population exposure, the staff recommends maintaining the primary annual standard at its current level.
- 4) Analyses of alternative averaging times and population exposures suggest that:
 - a) The current standards provide substantial protection against the effects identified as being associated with 24-hour and annual exposures.

- b) The current standards - as reflected by current emissions or emissions when the standards are just met with somewhat less restrictive implementation assumptions - also provide some limit on peak SO₂ exposures of concern for asthmatics. In some cases, however, up to 10 to 15% of the sensitive population in the vicinity of major sources could be exposed once a year to levels at or above 0.5 ppm for 5 minutes, while at elevated ventilation.
- c) The range of 1-hour standards analyzed (0.25 to 0.5 ppm) provides increased protection against such exposures, limiting the fraction of asthmatics exposed to less than 4%.

The relative protection afforded by current vs. alternative standards as indicated by current and ongoing exposure analyses is an important consideration in determining what, if any, standard revisions may be necessary.



APPENDIX A. ANALYSIS OF DOSE-RESPONSE RELATIONSHIPS FROM CONTROLLED SO₂ EXPOSURE STUDIES ON ASTHMATICS

The following discussion describes the analyses used to generate Figure 3-1, which plots results from the various controlled SO₂ exposure studies on mild asthmatics.

1) The studies used are summarized in Table A-1. To standardize comparisons, only studies that reported changes in specific airway resistance (SRaw). Unfortunately, several studies reporting significant declines only for other lung function parameters could not be represented (e.g., Koenig et al., 1983b, 1985a; Schacter et al., 1984). Studies involving unusual temperature and/or humidity conditions (i.e., < 6°C, RH < 40% or > 90%) were also excluded to avoid the interactive effects of airway drying or cooling in contributing to bronchoconstriction. In addition, results at low SO₂ exposure levels (generally \leq 0.25 ppm) where changes in SRaw were not statistically different from changes due to exercise alone were eliminated from the analysis. This would not be expected to bias the analysis in the domain where SRaw increases significantly with increased SO₂ exposure. The regression line in Figure 3-1 should not be extrapolated to zero dose, since at SO₂ levels below 0.25 ppm (~ 20 µg/min, oral airway dose rate) exercise-induced constriction dominates.

2) The studies involved either 5- or 10-minute exposure periods with one exception. Although total dose is a less satisfactory predictor of response than dose rate when considering longer exposure times (e.g., 1-hour) (Linn et al., 1982), no consistent trend can be seen in comparing responses to 3- vs. 5-minute vs. 10-minute exposures, which supports findings of Linn et al., (1983b).

Table A-1. Summary of Data from SO₂ Controlled Asthmatic Studies Used in Dose-Response Analysis

CONC. (ug/m ³)	CONC. ppm	DURATION MINUTES	TEMP C/RH	N	EXPOSURE M ³ /min.	V _e M ³ /min.	ORAL V _e M ³ /min.	ORAL V _e MOUTH	ORAL SO ₂ ug/min.	ORAL SO ₂ MOUTH	DESERATE ug/min.	% STRAW CLEARAIR SO ₂ / EXERCISE CLEARAIR	INCREASE SEE OVER CLEARAIR	REFERENCE
1300	0.5	5	23/80	6	M	0.04	0.04	0.04	52	52		15	141	KIRKPATRICK (1982)
1300	0.5	5	23/80	7	M	0.041	0.041	0.041	53.3	53.3		5	231	BETHEL (1983b)
1970	0.75	10	23/80	23	M	0.04	0.04	0.04	78.8	78.8		52	373	LINN (1982b)
630	0.25	10	23/80	7	M	0.03	0.03	0.03	18.9	18.9		0	30	SHEPPARD (1981a)
2600	1	5	23/80	6	M	0.031	0.031	0.031	80.6	80.6		0	300	SHEPPARD (1981a)
1300	0.5	3	23/82	8	M	0.035	0.035	0.035	45.5	45.5		0	104	SHEPPARD (1983)
1300	0.5	10	23/80	7	M	0.035	0.035	0.035	45.5	45.5		0	115	SHEPPARD (1981a)
1300	0.5	10	22/85	24	M	0.027	0.027	0.027	35.1	35.1		35	96	LINN (1983a)
1300	0.5	5	23/80	9	M	0.061	0.061	0.061	79.3	79.3		25	306	BETHEL (1983b)
1970	0.75	10	23/90	14	M	0.04	0.04	0.04	78.8	78.8		54	321	LINN (1983a)
1300	0.5	5	23/80	6	F	0.04	0.019	0.026	24.7	33.8		15	69	KIRKPATRICK (1982)
1300	0.5	5	23/90	9	F	0.061	0.038	0.042	49.4	54.5		25	219	BETHEL (1983b)
2600	1	10	26/70	27	C	0.042	0.02	0.027	52	70.2		47	190	ROGER (1985)
2600	1	10	26/70	10	C	0.041	0.02	0.027	52	70.2		48	172	KEHRL (1986)
1570	0.6	5	22/85	24	C	0.05	0.027	0.033	42.39	51.81		27	132	LINN (1984a)
1300	0.5	10	26/70	28	C	0.042	0.02	0.027	26	35.1		47	93	ROGER (1985)
1570	0.6	5	21/80	24	C	0.05	0.027	0.033	42.39	51.81		28	150	LINN (1985)
1040	0.4	5	23/85	23	C	0.048	0.023	0.032	23.92	33.28		35	69	LINN (1984b)
800	0.3	5	7/80	24	C	0.05	0.027	0.033	21.6	26.4		59	87	LINN (1983b)
800	0.3	5	21/80	24	C	0.05	0.027	0.033	21.6	26.4		28	59	LINN (1984b)
1570	0.6	5	21/20	22	C	0.05	0.027	0.033	42.39	51.81		20	206	LINN (1984b)
1570	0.6	5	21/80	22	C	0.05	0.027	0.033	42.39	51.81		10	157	LINN (1985a)
630	0.25	5	23/36	19	C	0.06	0.032	0.042	20.16	26.46		77	134	LINN (1985a)
1570	0.6	5	5/85	24	C	0.05	0.027	0.033	42.39	51.81		38	182	BETHEL (1985)
1300	0.5	5	23/41	10	C	0.06	0.032	0.042	41.6	54.6		39	238	LINN (1984a)
2600	1	30	26/70	10	C	0.041	0.02	0.027	52	70.2		50	233	BETHEL (1983a)
1570	0.6	5	7/80	24	C	0.05	0.027	0.033	42.39	51.81		58	207	KEHRL (1986)
1570	0.6	5	23/85	23	C	0.048	0.023	0.032	36.11	50.24		64	120	LINN (1984b)
1970	0.75	10	23/90	23	C	0.04	0.019	0.026	37.43	51.22		54	166	LINN (1983b)
1970	0.75	10	22/85	23	C	0.04	0.019	0.026	37.43	51.22		55	241	LINN (1983a)
1970	0.75	10	22/85	17	C	0.045	0.025	0.03	44.325	59.1		?	263	LINN (1982b)
														HOCKNEY (1984)

¹M = mouthpiece, F = facemask, C = chamber (freebreathing)

²V_e = total ventilation rate. Oral V_e is oral component of ventilation for normal habitual mouthbreathers. Oral SO₂ dose rate calculated separately for these 2 populations (see text).

3) Given the almost complete absorption of SO_2 that occurs in the moist surfaces of the nasal airways, the oral component of ventilation is critical in determining the SO_2 dose that penetrates to the airways where bronchomotor responses are triggered (Kleinman, 1984). Data on the partitioning between oral and nasal breathing under different exercise levels (Niinimaa et al., 1981; see 1982 staff paper, Appendix A) were used to estimate the oral component of ventilation given the ventilation rates (V_e) reported by the investigators. For example, Kirkpatrick et al. (1982) exposed asthmatics via mouthpiece to 0.5 ppm while exercising at about 40 L/min. Because a mouthpiece forces inspired air through the oral cavity thereby bypassing the nasal airways, it can be assumed that the oral V_e was 40 L/min resulting in an estimated SO_2 dose delivered via the oral airways of $[1300 \mu\text{g}/\text{m}^3 (0.5 \text{ ppm}) \times 0.04 \text{ m}^3/\text{min} (40 \text{ L/min})]$, or 52 $\mu\text{g}/\text{min}$. The asthmatics in the Kehrl et al. (1986) study were exposed free-breathing 1.0 ppm SO_2 while exercising at a ventilation rate of approximately 41 L/min. At this exercise level, most normal healthy individuals breathing unencumbered augment the amount of air entering the nasal passages by inhaling some air via the mouth so that the oral V_e would be approximately 20 L/min (Niinimaa et al., 1981). The oral airway SO_2 dose is estimated as $[2600 \mu\text{g}/\text{m}^3 (1.0 \text{ ppm}) \times 0.02 \text{ m}^3/\text{min} (20 \text{ L/min})]$, or 52 $\mu\text{g}/\text{min}$, which is identical to that in the Kirkpatrick study. Interestingly, the increases in airway resistance over clean air/exercise control in these studies were almost identical (126% vs. 124%).

For all calculations on free breathing experiments, typical oral/nasal breathing patterns were used as determined by Niinimaa et al., (1981) (see 1982 staff paper, Appendix A). By assuming that all of the freebreathing subjects were normal augmentors, some underestimation of SO_2 dose likely

results, especially given indications of increased frequency of allergic rhinitis and nasal congestion in asthmatics resulting in obligatory mouth breathing. Variability in such conditions between different groups of subjects may explain observed differences in responses between studies, as evidenced by the failure to fully replicate the Kirkpatrick et al. (1982) results under similar conditions but with fewer subjects with nasal disorders (Bethel et al., 1983b). An alternative approach is taken by Kleinman (1984) who estimates population-weighted oral V_e at different activity levels. A separate analysis (not illustrated), which used the same group of data assuming subjects were habitual mouthbreathers, produced no apparent improvements ($r^2 = 0.76$).

For the facemask experiment included in Bethel et al. (1983b), actual measurements of oral airflow through the masks were provided and roughly matched Niinimaa et al.'s prediction for oronasal breathing. In the Kirkpatrick et al. (1982) facemask study, it was assumed that free, oronasal breathing was simulated.

5) Changes in SRaw in response to SO_2 exposure while at exercise over baseline measurements were used as opposed to changes in SRaw over increases due to exercise alone in clean air. Again, separate analysis (not shown) using the latter measure yielded nearly identical results.

6) A simple linear regression was fit to the data. As mentioned, the linear relationship should not be extended to lower SO_2 exposure levels down to zero.

UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D. C. 20460

February 19, 1987

OFFICE OF
THE ADMINISTRATORThe Honorable Lee M. Thomas
Administrator
U.S. Environmental Protection
Agency
Washington, DC 20460

Dear Mr. Thomas:

The Clean Air Scientific Advisory Committee (CASAC) has completed its review of the 1986 Addendum to the 1982 Staff Paper on Sulfur Oxides (Review of the National Ambient Air Quality Standards for Sulfur Oxides: Updated Assessment of Scientific and Technical Information) prepared by the Agency's Office of Air Quality Planning and Standards (OAQPS).

The Committee unanimously concludes that this document is consistent in all significant respects with the scientific evidence presented and interpreted in the combined Air Quality Criteria Document for Particulate Matter/Sulfur Oxides (1982) and its 1986 Addendum, on which CASAC issued its closure letter on December 15, 1986. The Committee believes that the 1986 Addendum to the 1982 Staff Paper on Sulfur Oxides provides you with the kind and amount of technical guidance that will be needed to make appropriate decisions with respect to the standards. The Committee's major findings and conclusions concerning the various scientific issues and studies discussed in the Staff Paper Addendum are contained in the attached report.

Thank you for the opportunity to present the Committee's views on this important public health and welfare issue.

Sincerely,

A handwritten signature in cursive script, appearing to read "Morton Lippmann".

Morton Lippmann, Ph.D.

Chairman

Clean Air Scientific Advisory
Committeecc: A. James Barnes
Gerald Emison
Lester Grant
Vaun Newill
John O'Connor
Craig Potter
Terry Yosie

SUMMARY OF MAJOR SCIENTIFIC ISSUES AND CASAC
CONCLUSIONS ON THE 1986 DRAFT ADDENDUM
TO THE 1982 SULFUR OXIDES STAFF PAPER

The Committee found the technical discussions contained in the Staff Paper Addendum to be scientifically thorough and acceptable, subject to minor editorial revisions. This document is consistent in all significant respects with the scientific evidence presented in the 1982 combined Air Quality Criteria Document for Particulate Matter/Sulfur Oxides and its 1986 Addendum, on which the Committee issued its closure letter on December 15, 1986.

Scientific Basis for Primary Standards

The Committee addressed the scientific basis for a 1-hour, 24-hour, and annual primary standards at some length in its August 26, 1983 closure letter on the 1982 Sulfur Oxides Staff Paper. That letter was based on the scientific literature which had been published up to 1982. The present review has examined the more recently published studies.

It is clear that no single study of SO₂ can fully address the range of public health issues that arise during the standard setting process. The Agency has completed a thorough analysis of the strengths and weaknesses of various studies and has derived its recommended ranges of interest by evaluating the weight of the evidence. The Committee endorses this approach.

The Committee wishes to comment on several major issues concerning the scientific data that are available. These issues include:

- Recent studies more clearly implicate particulate matter than SO₂ as a longer-term public health concern at low exposure levels.
- A majority of Committee members believe that the effects reported in the clinical studies of asthmatics represent effects of significant public health concern.
- The exposure uncertainties associated with a 1-hour standard are quite large. The relationship between the frequency of short-term peak exposures and various scenarios of asthmatic responses is not well understood. Both EPA and the electric power industry are conducting further analyses of a series of exposure assessment issues. Such analyses have the potential to increase the collective understanding of the relationship between SO₂ exposures and responses observed in subgroups of the general population.
- The number of asthmatics vulnerable to peak exposures near electric power plants, given the protection afforded by the current standards, represents a small number of people. Although the Clean Air Act requires that sensitive population groups receive protection, the size of such groups has not been defined. CASAC believes that this issue represents a legal/policy matter and has no specific scientific advice to provide on it.

CASAC's advice on primary standards for three averaging times is presented below:

1-Hour Standard - It is our conclusion that a large, consistent data base exists to document the bronchoconstrictive response in mild to moderate asthmatics subjected in clinical chambers to short-term, low levels of sulfur dioxide while exercising. There is, however, no scientific basis at present to support or dispute the hypothesis that individuals participating in the SO₂ clinical studies are surrogates for more sensitive asthmatics. Estimates of the size of the asthmatic population that experience exposures to short-term peaks of SO₂ (0.2 - 0.5 parts per million (ppm) SO₂ for 5-10 minutes) during light to moderate exercise, and that can be expected to exhibit a bronchoconstrictive response, varies from 5,000 to 50,000.

The majority of the Committee believes that the scientific evidence supporting the establishment of a new 1-hour standard is stronger than it was in 1983. As a result, and in view of the significance of the effects reported in these clinical studies, there is strong, but not unanimous support for the recommendation that the Administrator consider establishing a new 1-hour standard for SO₂ exposures. The Committee agrees that the range suggested by EPA staff (0.2 - 0.5 ppm) is appropriate, with several members of the Committee suggesting a standard from the middle of this range. The Committee concludes that there is not a scientifically demonstrated need for a wide margin of safety for a 1-hour standard.


24-Hour Standard - The more recent studies presented and analyzed in the 1986 Staff Paper Addendum, in particular, the episodic lung function studies in children (Dockery et al., and Dassen et al.) serve to strengthen our previous conclusion that the rationale for reaffirming the 24-hour standard is appropriate.

Annual Standard - The Committee reaffirms its conclusion, voiced in its 1983 closure letter, that there is no quantitative basis for retaining the current annual standard. However, a decision to abolish the annual standard must be considered in the light of the total protection that is to be offered by the suite of standards that will be established.

The above recommendations reflect the consensus position of CASAC. Not all CASAC reviewers agree with each position adopted because of the uncertainties associated with the existing scientific data. However, a strong majority supports each of the specific recommendations presented above, and the entire Committee agrees that this letter represents the consensus position.

Secondary Standards

The 3-hour secondary standard was not addressed at this review.



REFERENCES

- Anderson, G.E. (1982) Systems Applications, Inc., San Rafael, CA. Personal communications with Henry Thomas (July 1982).
- Anderson, S. D. (1985) Issues in exercise-induced asthma. *J. Allergy Clin. Immunol.* 76: 763-772.
- Armstrong, B. (1985) Peak/Mean Ratios for Phelps Dodge Douglas, Inter-Office Memorandum thru David O. Chelgren, Manager, Compliance Unit, Arizona Department of Health Services, December 4, 1985.
- Armstrong, B. (1986) Peak/Mean Ratios for Magma, Inter-Office Memorandum thru David O. Chelgren, Manager, Compliance Unit, Arizona Department of Health Services, January 20, 1986.
- Bar-Yishay, E.; Ben-Dov, I.; Godfrey, S. (1983) Refractory period after hyperventilation-induced asthma. *Am. Rev. Respir. Dis.* 127: 572-74.
- Bethel, R. A.; Epstein, J.; Sheppard, D.; Nadel, J. A.; Boushey, H. A. (1983a) Sulfur dioxide-induced bronchoconstriction in freely breathing, exercising, asthmatic subjects. *Am. Rev. Respir. Dis.* 129: 987-990.
- Bethel, R. A.; Erle, D. J.; Epstein, J.; Sheppard, D.; Nadel, J. A.; Boushey, H. A. (1983b) Effect of exercise rate and route of inhalation on sulfur-dioxide-induced bronchoconstriction in asthmatic subjects. *Am. Rev. Respir. Dis.* 128: 592-596.
- Bethel, R. A.; Sheppard, D.; Epstein, J.; Tam, E.; Nadel, J. A.; Boushey, H. A. (1984) Interaction of sulfur dioxide and cold dry air in causing bronchoconstriction in asthmatic subjects. *J. Appl. Physiol.: Respir. Environ. Exercise Physiol.* 57: 419-423.
- Bethel, R. A., Sheppard, D.; Geffroy, B.; Tam, E.; Nadel, J. A.; Boushey, H. A. (1985) Effect of 0.25 ppm sulfur dioxide on airway resistance in freely breathing, heavily exercising, asthmatic subjects. *Am. Rev. Respir. Dis.* 131: 659-661.
- Biller, W.F.; Fegans, T.B.; Johnson, T.R.; Duggan, G.M.; Paul, R.A.; McCurdy, T.; Thomas, H.C. (1981) A general model for estimating exposure associated with alternative NAAQS. Presented at 74th Annual meeting of the Air Pollution Control Assoc., Philadelphia, PA, June 21-26, 1981.
- Burton, C.S.; Nordin, J.; Stoeckenius, T. (1982) On a New Short-Term Standard for Sulfur Dioxide: The Protection from Calculated Peak 1-Hour Concentrations Provided by the Existing 3- and 24-Hour NAAQS in the Vicinity of a Hypothetical Power Plant. Systems Applications, Inc., San Rafael, CA.
- Burton, C.S.; Thrall, A.D. (1982) A Brief Review of Peak to One-Hour Average Ratios Observed in the Vicinity of Power Plants. Systems Applications, Inc., San Rafael, CA.
- Chapman, R. S.; Calafiore, D. C.; Hasselblad, V. (1985) Prevalence of persistent cough and phlegm in young adults in relation to long-term ambient sulfur oxide exposure. *Am. Rev. Respir. Dis.* 132: 261-267.

- Cohen, J. (1983) Analysis of SO₂ Controlled Human Data. Memorandum to John Haines, Ambient Standards Branch, Office of Air Quality Planning and Standards, U.S. EPA, Research Triangle Park, N.C., September 6, 1983.
- Cohen, J. (1984) Assessing Asthmatic Responses to SO₂: Summary of Expert Opinions. Memorandum to John Bachmann, Ambient Standards Branch, Office of Air Quality Planning and Standards, U.S. EPA, Research Triangle Park, N.C., February 27, 1984.
- Committee on Public Works, U.S. Senate (1974) A Legislative History of the Clean Air Amendments. Volume 1. Serial no. 93-18. U.S. Government Printing Office, Washington, D.C. prepared by the Environmental Policy Division of the Congressional Research Services of the Library of Congress.
- CEC [Commission of the European Communities] (1983) Report on the EC epidemiology survey on the relationship between air pollution and respiratory health in primary school children. Brussels, Belgium: Environmental Research Programme.
- Dassen, W.; Brunekreef, B.; Hoek, G.; Hofschreuder, P.; Staatsen, B.; deGroot, H.; Schouten, E.; Biersteker, K. (1986) Decline in children's pulmonary function during an air pollution episode. *J. Air Pollut. Control Assoc.* (in press).
- D.C. Cir. (1980) *Lead Industries Association, Inc. v. EPA*, F. 2d, 14 ERC 1906 (D.C. Cir.) Cert. Denied 49 U.S.L.W. 3428 December 8, 1980.
- D.C. Cir. (1981) *American Petroleum Institute v. Costle*, Nos. 79-1104 et. al. (D.C. Cir.) September 3, 1981.
- Deal, E. C.; McFadden, E. R.; Ingram, R. H.; Jaeger, J. J. (1979b) Hyperpnea and heat flux: initial reaction sequence in exercise-induced asthma. *J. Appl. Physiol.* 46: 476-483.
- Dockery, D. W.; Ware, J. H.; Ferris, B. G., Jr.; Speizer, F. E.; Cook, N. R.; Herman, S. M. (1982) Change in pulmonary function in children associated with air pollution episodes. *J. Air Pollut. Control Assoc.* 32: 937-942.
- Dodge, R. (1983) The respiratory health and lung function of Anglo-American children in a smelter town. *Am. Rev. Respir. Dis.* 127: 158-161.
- Dodge, R.; Solomon, P.; Moyers, J.; Hayes, C. (1985) A longitudinal study of children exposed to sulfur oxides. *Am. J. Epidemiol.* 121: 720-736.
- EPA [U.S. Environmental Protection Agency] (1982a) Review of the National Ambient Air Quality Standards for Sulfur Oxides: Assessment of Scientific and Technical Information (OAQPS Staff Paper) Office of Air Quality Planning and Standards, Research Triangle Park, N.C. EPA-450/5-82-007. Available from NTIS, Springfield, VA; PB 84-102920.


- EPA [U.S. Environmental Protection Agency] (1982b) Air Quality Criteria for Particulate Matter and Sulfur Oxides. Environmental Criteria and Assessment Office. Research Triangle Park, N.C. EPA-600/8-82-029. Available from NTIS, Springfield, VA; PB 54-156801/REB.
- EPA [U.S. Environmental Protection Agency] (1982c) Review of the National Ambient Air Quality Standards for Particulate Matter: Assessment of Scientific and Technical Information (OAQPS Staff Paper) Office of Air Quality Planning and Standards. Research Triangle Park, N.C. EPA-450/5-82-001. Available from NTIS, Springfield, VA; PB 82-177874.
- EPA [U.S. Environmental Protection Agency] (1986a) Second Addendum to Air Quality Criteria for Particulate Matter and Sulfur Oxides (1982): Assessment of Newly Available Health Effects Information. Environmental Criteria and Assessment Office, Office of Research and Development, Research Triangle Park, N.C. EPA-600/8-86-020F.
- EPA [U.S. Environmental Protection Agency] (1986b) Review of the National Ambient Air Quality Standards for Particulate Matter: Updated Assessment of Scientific and Technical Information (Addendum to the 1982 JAQPS Staff Paper). Office of Air Quality Planning and Standards. Research Triangle Park, N.C., December 1986.
- EPA [U.S. Environmental Protection Agency] (1986c) Staff Assessment of Recent Comments on the Short-term Sulfur Dioxide Exposure Analysis, Strategies and Air Standards Division, Office of Air Quality Planning and Standards, Research Triangle Park, N.C., July 1986.
- EPA [U.S. Environmental Protection Agency] (1986d) An Analysis of Short-term Sulfur Dioxide Population Exposures in the Vicinity of Utility Power Plants. Strategies and Air Standards Division, Office of Air Quality Planning and Standards, Research Triangle Park, N.C., September 12, 1986.
- Frank, N.H.; Thrall, A.D. (1982) Relationships Among SO₂ Averaging Times and Ambient Standards. Monitoring and Data Analysis Division Report. Office of Air Quality Planning and Standards, Research Triangle Park, N.C.
- Goldstein, I.F.; Weinstein, A.L. (1986) Air pollution and asthma: effects of exposures to short-term sulfur dioxide peaks. *Env. Res.* 40: 332-345.
- Hackney, J. D.; Linn, W. S.; Bailey, R. M.; Spier, C. E.; Valencia, L. M. (1984) Time course of exercise-induced bronchoconstriction in asthmatics exposed to sulfur dioxide. *Environ. Res.* 34: 321-327.
- Hatzakis, A.; Katsouyanni, K.; Kalandidi, A.; Day, N.; Trichopoulos, D. (1986) Short-term effects of air pollution on mortality in Athens. *Int. J. Epidemiol.* 15: 73-81.
- Huber, A.H.; Pooler, F. (1985) Comments on Peak Ground Level Concentrations Due to Building Downwash Relative to Peak Concentrations Under Atmospheric Dispersion Processes. Atmospheric Sciences Research Laboratory, U.S. EPA, Research Triangle Park, N.C.

- Horstman, D. H.; Roger, L. J.; Kehrl, H. R.; Hazucha, M. J. (1986) Airway sensitivity of asthmatics to sulfur dioxide. *Toxicol. Ind. Health.* 2:289-298.
- Kehrl, H. R.; Roger, L. J.; Hazucha, M. J.; Horstman, D.H. (1986) Differing response of asthmatics to SO₂ exposure with continuous and intermittent exercise. *Am. Rev. Respir. Dis.:* [in press].
- Kilkelly, G.T.; Roberson R.L. (1985) Analysis of Ambient SO₂ Data During Downwash Events. Contractor report prepared for Utilities Air Regulatory Group by Kilkelly Environmental Associates, Raleigh, N.C.
- Kirkpatrick, M. B.; Sheppard, D.; Nadel, J. A.; Boushey, H. A. (1982) Effect of the oronasal breathing route on sulfur dioxide-induced bronchoconstriction in exercising asthmatic subjects. *Am. Rev. Respir. Dis.* 125: 627-631.
- Kleinman, M. T. (1984) Sulfur dioxide and exercise: relationships between response and absorption in upper airways. *J. Air Pollut. Control Assoc.* 34: 32-37.
- Koenig, J. Q.; Pierson, W. E.; Horike, M.; Frank, R. (1983) A comparison of the pulmonary effects of 0.5 ppm versus 1.0 ppm sulfur dioxide plus sodium chloride droplets in asthmatic adolescents. *J. Toxicol. Environ. Health* 11: 129-139.
- Koenig, J. Q.; Morgan, M. S.; Horike, M.; Pierson, W. E. (1985) The effects of sulfur oxides on nasal and lung function in adolescents with extrinsic asthma. *J. Allergy Clin. Immunol.* 76: 813-818.
- Larsen, R.I. (1968) A New Mathematical Model of Air Pollution Concentration, Averaging Time and Frequency (Presented at 61st Annual Meeting of the Air Pollution Control Association, St. Paul, Minnesota).
- Lawther, P.J. (1986) Letter to John Bachmann, Office of Air Quality Planning and Standards, U.S. Environmental Protection Agency, Research Triangle Park, N.C., August 22, 1986.
- Lawther, P. J.; Waller, R. E.; Henderson, M. (1970) Air pollution and exacerbations of bronchitis. *Thorax* 25: 525-539.
- Linn, W. S.; Bailey, R. M.; Shamoo, D. A.; Venet, T. G.; Wightman, L. H.; Hackney, J. D. (1982) Respiratory responses of young adult asthmatics to sulfur dioxide exposure under simulated ambient conditions. *Environ. Res.* 29: 220-232.
- Linn, W. S.; Shamoo, D. A.; Spier, C. E.; Valencia, L. M.; Anzar, U. T.; Venet, T. G.; Hackney, J. D. (1983a) Respiratory effects of 0.75 ppm sulfur dioxide in exercising asthmatics: influence of upper-respiratory defenses. *Environ. Res.* 30: 340-348.
- Linn, W. S.; Venet, T. G.; Shamoo, D. A.; Valencia, L. M.; Anzar, U. T.; Spier, C. E.; Hackney, J. D. (1983b) Respiratory effects of sulfur dioxide in heavily exercising asthmatics: a dose-response study. *Am. Rev. Respir. Dis* 127: 278-283.

- Linn, W. S.; Shamoo, D. A.; Venet, T. G.; Bailey, R. M.; Wightman, L. H.; Hackney, J. D. (1984a) Comparative effects of sulfur dioxide exposures at 5°C and 22°C in exercising asthmatics. *Am. Rev. Respir. Dis.* 129: 234-239.
- Linn, W. S.; Shamoo, D. A.; Vinet, T. G.; Spier, C. E.; Valencia, L. M.; Anzar, U. T.; Hackney, J. D. (1984b) Combined effect of sulfur dioxide and cold in exercising asthmatics. *Arch Environ. Health* 39: 339-346.
- Linn, W. S.; Avol, E. L.; Shamoo, D. A.; Venet, T. G.; Anderson, K. R.; Whynot, J. D.; Hackney, J. D. (1984c) Asthmatics' responses to 6-hr sulfur dioxide exposures on two successive days. *Arch. Environ. Health* 39: 313-319.
- Linn, W. S.; Shamoo, D. A.; Anderson, K. R.; Whynot, J. D.; Avol, E. L.; Hackney, J. D. (1985a) Effects of heat and humidity on the responses of exercising asthmatics to sulfur dioxide exposure. *Am. Rev. Respir. Dis.* 131: 221-225.
- Lott, R.A. (1985) SO₂ concentrations near tall stacks. *Atmos. Environ.* 19: 1589-1599.
- Martin, A. E. (1964) Mortality and morbidity statistics and air pollution. *Proc. R. Soc. Med.* 57: 969-975.
- Martin, A. E.; Bradley, W. H. (1960) Mortality, fog and atmospheric pollution-- An investigation during the winter of 1958-59. *Mon. Bull. Minist. Health Public Health Lab. Serv.* 19: 56-72.
- Mazumdar, S.; Sussman, N. (1981) Relationships of air pollution to health: Results from the Pittsburgh Study. Presented at: 74th annual meeting of the Air Pollution Control Association; June, Philadelphia, PA. Pittsburgh, PA; Air Pollution Control Association.
- Mazumdar, S.; Sussman, N. (1983) Relationships of air pollution to health: results from the Pittsburgh study. *Arch. Environ. Health* 38: 17-24.
- Mazumdar, S.; Schimmel, H.; Higgins, I. T. T. (1982) Relation of daily mortality to air pollution: an analysis of 14 London winters, 1958/59-1971/72. *Arch. Environ. Health* 37: 213-220.
- Muhling, P.; Bory, J.; Haupt, H. (1985) Einfluss der Luftbelastung auf Atemwegserkrankungen. Untersuchungen bei Sauglingen und Kleinkindern [The influence of air pollution on respiratory diseases: studies of babies and infants]. *Staub Reinhalt. Luft* 45: 35-38.
- Niinimaa, V.; Cole, P.; Mintz, S.; Shephard, R. J. (1981) Oronasal distribution of respiratory airflow. *Respir. Physiol.* 43: 69-75.
- Ozkaynak, H.; Spengler, J. D. (1985) Analysis of health effects resulting from population exposures to acid precipitation precursors. *EHP Environ. Health Perspect.* 63: 45-55.
- Ozkaynak, H.; Schatz, A. D.; Thurston, G. D.; Isaacs, R. G.; Husar, R. B. (1985) Relationships between aerosol extinction coefficient derived from airport visual range observations and alternative measures of airborne particle mass. *J. Air Pollut. Control Assoc.* 35: 1176-1185.

- PAARC Cooperative Group. (1982a) Atmospheric pollution and chronic or recurrent respiratory diseases. I. Methods and material. *Bull. Eur. Physiopathol. Respir.* 18: 87-99.
- PAARC Cooperative Group. (1982b) Atmospheric pollution and chronic or recurrent respiratory disease. II. Results and discussion. *Bull. Eur. Physiopathol. Respir.* 18: 101-116.
- Roger, L. J.; Kehrl, H. R.; Hazucha, M.; Horstman, D. H. (1985) Bronchoconstriction in asthmatics exposed to sulfur dioxide during repeated exercise. *J. Appl. Physiol.* 59: 784-791.
- Rote, D.M.; Lee, J.L. (1983) Discussion and Interpretation of Results of the Analysis of the St. Louis SO₂ Data: Final Report. Energy and Environmental Systems Division, Argonne National Laboratory, Argonne, IL.
- Roth, H.D.; Wyzga, R.E.; Hayter, A.J. (1986) Methods and Problems in Estimating Health Risks from Particulates. In *Aerosols* (S.D. Lee, T. Schneider, L.D. Grant, P.J. Verkerk, eds.) Lewis Publishers, Chelsea, MI, pp. 837-857.
- Saric, M.; Fugas, M.; Hrustic, O. (1981) Effects of urban air pollution on school-age children. *Arch. Environ. Health* 36: 101-108.
- Schachter, E. N.; Witek, T. J., Jr.; Beck, G. J.; Hosein, H. R.; Colice, G.; Leaderer, B. P.; Cain, W. (1984) Airway effects of low concentrations of sulfur dioxide: dose-response characteristics. *Arch Environ. Health* 39: 34-42.
- Schenker, M. B.; Samet, J. M.; Speizer, F. E.; Gruhl, J.; Batterman, S. (1983) Health effects of air pollution due to coal combustion in the Chestnut Ridge region of Pennsylvania: results of cross-sectional analysis in adults. *Arch. Environ. Health* 38: 325-330.
- Sheppard, D.; Epstein, J.; Bethel, R. A.; Nadel, J. A.; Boushey, H. A. (1983) Tolerance to sulfur dioxide-induced bronchoconstriction in subjects with asthma. *Environ. Res.* 30: 412-419.
- Sheppard, D.; Eschenbacher, W. L.; Boushey, H. A.; Bethel, R. A. (1984) Magnitude of the interaction between the bronchomotor effects of sulfur dioxide and those of dry (cold) air. *Am. Rev. Respir. Dis.* 130: 52-55.
- Shumway, R. H.; Tai, R. Y.; Tai, L. P.; Pawitan, Y. (1983) Statistical analysis of daily London mortality and associated weather and pollution effects. Sacramento, CA: California Air Resources Board; contract no. A1-154-33.
- Snashall, P. D.; Baldwin, C. (1982) Mechanisms of sulphur dioxide induced bronchoconstriction in normal and asthmatic man. *Thorax* 37: 118-123.

- Stearns, D. R.; McFadden, E. R., Jr.; Breslin, F. J.; Ingram, R. H., Jr. (1981) Reanalysis of the refractory period in exertional asthma. *J. Appl. Physiol.: Respir. Environ. Exercise Physiol.* 50: 503-508.
- Stebbing, J. H., Jr.; Fogleman, D. G. (1979) Identifying a susceptible subgroup: effects of the Pittsburgh air pollution episode upon school children. *Am. J. Epidemiol.* 110: 27-40.
- Strauss, R. H.; McFadden, E. R., Jr.; Ingram, R. H., Jr.; Jaeger, J. J. (1977) Enhancement of exercise-induced asthma by cold air. *New Engl. J. Med.* 297: 743-747.
- Thrall, A.D.; Langstaff, J.E.; Liu, M.K.; Burton, C.S. (1982) On the Variability in Peak Sulfur Dioxide Concentrations Contained in Longer-Term Averages. An Empirical Study of an Isolated Power Plant. System Applications, Inc., San Rafael, CA, Pub. No. 82262, October, 1982.
- van der Lende, R.; Schouten, J. P.; Rijcken, B.; van der Meulen, A. (1986) Longitudinal epidemiologic studies on effects of air pollution in The Netherlands. In: Lee, S.D.; Schneider, T.; Grant, L.D.; Verkerk, P.J., eds. *Aerosols: research, risk assessment and control strategies, proceedings of the second U.S.-Dutch international symposium*; May 1985; Williamsburg, VA; Chelsea, MI, Lewis Publishers, Inc; pp. 731-742.
- Ware, J. H.; Thibodeau, L. A.; Speizer, F. E.; Colome, S.; Ferris, B. G., Jr. (1981) Assessment of the health effects of atmospheric sulfur oxides and particulate matter: Evidence from observational studies. *Environ. Health Perspect.* 41: 255-276.
- Ware, J. H.; Ferris, B. G., Jr.; Dockery, D. W.; Spengler, J. D.; Stram, D. O.; Speizer, F. E. (1986) Effects of ambient sulfur oxides and suspended particles on respiratory health of preadolescent children. *Am. Rev. Respir. Dis.* 133: 834-842.
- Wojtyniak, B.; Krzyzanowski, M.; Jedrychowski, W. (1984) Importance of urban air pollution in chronic respiratory problems = Die Bedeutung staedterischer Luftverunreinigung fuer chronische Atemwegsbeschwerden. *Z. Erkr. Atmungsorgane* 163: 274-284.



TECHNICAL REPORT DATA
(Please read Instructions on the reverse before completing)

1. REPORT NO. EPA-450/5-86-13		2.	3. RECIPIENT'S ACCESSION NO.	
4. TITLE AND SUBTITLE Review of the National Ambient Air Quality Standards for Sulfur Oxides: Updated Assessment of Scientific and Technical Information Addendum to the 1982 OAQPS Staff Paper			5. REPORT DATE December 1986	
			6. PERFORMING ORGANIZATION CODE	
7. AUTHOR(S)			8. PERFORMING ORGANIZATION REPORT NO.	
9. PERFORMING ORGANIZATION NAME AND ADDRESS Office of Air and Radiation Office of Air Quality Planning and Standards U. S. Environmental Protection Agency Research Triangle Park, NC 27711			10. PROGRAM ELEMENT NO.	
			11. CONTRACT/GRANT NO.	
12. SPONSORING AGENCY NAME AND ADDRESS			13. TYPE OF REPORT AND PERIOD COVERED Final	
			14. SPONSORING AGENCY CODE	
15. SUPPLEMENTARY NOTES				
16. ABSTRACT This paper evaluates and interprets the updated scientific and technical information that EPA staff believes is most relevant to the review of primary (health) national ambient air quality standards for sulfur oxides and represents an update of previous staff conclusions and recommendations in the 1982 sulfur oxides staff paper to incorporate more recent information. This assessment is intended to bridge the gap between the scientific review in the EPA criteria document second addendum for particulate matter and sulfur oxides and the judgments required of the Administrator in setting ambient air quality standards for sulfur oxides. The major recommendations of the staff paper addendum include the following: (1) that the health data support the need for sulfur dioxide (SO ₂) standards; (2) that new data from controlled human exposure studies on asthmatics and atopics warrant consideration of a new short-term (1-hour) standard; (3) that the current primary and secondary standards (annual, 24-hour, and 3-hour) provide substantial protection against effects associated with 24-hour and annual exposures, and some limit on peak exposures of concern for asthmatics; (4) that the relative protection afforded by current vs. alternative standards is an important consideration in determining, what, if any, standards revisions may be necessary.				
17. KEY WORDS AND DOCUMENT ANALYSIS				
a. DESCRIPTORS		b. IDENTIFIERS/OPEN ENDED TERMS		c. COSATI Field/Group
Sulfur Oxides Sulfur Dioxide Air Pollution Particulate Matter		Air Quality Standards		
18. DISTRIBUTION STATEMENT Release to Public		19. SECURITY CLASS (This Report) Unclassified		21. NO. OF PAGES 88
		20. SECURITY CLASS (This page)		22. PRICE

