

Table V-4
CERVIX UTERI CANCER (Invasive)

SEER INCIDENCE^a AND U.S. DEATH^b RATES, AGE-ADJUSTED AND AGE-SPECIFIC RATES, BY RACE

	Incidence ^a			US Death ^b		
	All Races, Females	White Females	Black Females	All Races, Females	White Females	Black Females
<u>AGE-ADJUSTED RATES, 2001-2005</u>						
AGE AT DIAGNOSIS/DEATH:						
All ages	8.4	8.2	10.8	2.5	2.3	4.7
Under 65	7.8	7.7	8.8	1.9	1.8	3.1
65 and over	13.0	11.4	24.6	6.6	5.6	15.4
All ages(IARC world std) ^c	6.6	6.5	8.2	1.8	1.7	3.3
<u>AGE-SPECIFIC RATES, 2001-2005</u>						
AGE AT DIAGNOSIS/DEATH:						
<1	-	-	-	-	-	-
1-4	-	-	-	-	-	-
5-9	-	-	-	-	-	-
10-14	-	-	-	-	-	-
15-19	-	-	-	-	-	-
20-24	1.5	1.5	1.8	0.1	0.1	-
25-29	5.8	6.2	5.7	0.6	0.5	1.0
30-34	11.3	12.2	9.7	1.4	1.4	1.9
35-39	13.3	14.2	11.9	2.4	2.3	3.4
40-44	15.3	15.3	16.0	3.4	3.2	5.3
45-49	14.6	13.8	18.3	4.2	3.9	6.8
50-54	13.9	13.5	17.6	4.6	4.2	7.4
55-59	14.1	13.4	18.3	5.2	4.8	8.7
60-64	14.3	12.6	24.2	5.5	4.9	10.5
65-69	15.1	13.7	23.9	5.8	5.0	12.8
70-74	13.1	11.9	22.2	6.0	5.1	13.2
75-79	12.5	10.5	25.2	6.3	5.4	15.2
80-84	11.4	9.8	27.4	7.7	6.7	18.7
85+	10.7	9.0	26.5	8.5	7.2	22.2

^a SEER 17 areas. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130), unless noted.

^b US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130), unless noted.

^c Rates are per 100,000 and are age-adjusted to the IARC world standard population.

- Statistic not shown. Rate based on less than 16 cases for the time interval.