

Web Table Only: Incidence Rates^a for Selected Common Cancer Sites^b in Adults by Age, Race/Ethnicity and Sex
in the Selected Areas^c in the United States, 1999-2003.

Cancer Site	Age Group	Hispanic ^d			Non-Hispanic White			Non-Hispanic Black		
		No. ^a	Rate ^a	95% CI	No. ^a	Rate ^a	95% CI	No. ^a	Rate ^a	95% CI
Male										
Colon and Rectum										
	20-39	891	3.1	(2.9 - 3.3)	4,398	4.1	(3.9 - 4.2) *	848	4.4	(4.1 - 4.7) *
	40-49	1,811	18.4	(17.5 - 19.2)	14,581	22.4	(22.1 - 22.8) *	2,813	29.9	(28.8 - 31.0) *
	50-64	5,693	81.3	(79.2 - 83.5)	65,958	97.7	(97.0 - 98.5) *	9,700	125.3	(122.8 - 127.8) *
	65-74	5,184	231.6	(225.3 - 238.1)	77,198	279.0	(277.1 - 281.0) *	7,929	301.7	(295.0 - 308.4) *
	75+	4,415	350.5	(340.1 - 361.1)	98,593	445.3	(442.5 - 448.1) *	7,167	443.2	(432.9 - 453.7) *
Lung and Bronchus										
	20-39	279	1.0	(0.9 - 1.1)	2,081	1.9	(1.8 - 2.0) *	454	2.4	(2.2 - 2.6) *
	40-49	994	10.2	(9.5 - 10.8)	15,551	23.8	(23.5 - 24.2) *	4,011	42.6	(41.3 - 44.0) *
	50-64	4,939	71.1	(69.1 - 73.1)	98,876	146.5	(145.6 - 147.4) *	16,838	218.0	(214.7 - 221.3) *
	65-74	5,634	253.0	(246.4 - 259.7)	131,755	476.2	(473.7 - 478.8) *	14,131	536.4	(527.5 - 545.4) *
	75+	4,979	389.0	(378.1 - 400.1)	127,268	563.7	(560.6 - 566.8) *	10,292	620.3	(608.3 - 632.5) *
Prostate										
	20-49	1,237	4.5	(4.3 - 4.8)	13,713	7.4	(7.2 - 7.5) *	4,477	16.8	(16.3 - 17.3) *
	50-64	15,049	218.9	(215.4 - 222.4)	204,240	302.6	(301.3 - 303.9) *	40,128	521.9	(516.8 - 527.1) *
	65-74	19,039	847.4	(835.4 - 859.6)	254,328	918.6	(915.0 - 922.1) *	35,545	1337.9	(1324.0 - 1352.0) *
	75+	10,846	833.8	(817.9 - 849.9)	182,263	806.6	(802.9 - 810.4) *	19,867	1204.9	(1188.0 - 1221.9) *
Female										
Breast										
	20-39	5,049	20.6	(20.0 - 21.1)	28,255	26.3	(26.0 - 26.6) *	6,513	30.5	(29.8 - 31.3) *
	40-49	11,493	118.7	(116.6 - 120.9)	100,504	154.1	(153.2 - 155.1) *	15,603	144.2	(141.9 - 146.5) *
	50-64	16,486	210.9	(207.6 - 214.1)	220,057	311.8	(310.5 - 313.1) *	24,829	258.1	(254.9 - 261.3) *
	65-74	8,240	286.8	(280.6 - 293.1)	146,400	447.6	(445.3 - 449.9) *	12,688	338.0	(332.2 - 344.0) *
	75+	6,054	297.8	(290.4 - 305.4)	166,665	446.6	(444.4 - 448.7) *	11,366	356.5	(350.0 - 363.1) *
Cervix										
	20-39	2,874	11.3	(10.8 - 11.7)	9,786	9.2	(9.0 - 9.3) *	2,049	9.4	(9.0 - 9.9) *
	40-49	2,578	26.4	(25.3 - 27.4)	8,857	13.7	(13.4 - 14.0) *	2,173	20.1	(19.2 - 20.9) *
	50-64	2,118	27.0	(25.8 - 28.1)	8,553	12.2	(11.9 - 12.4) *	2,188	22.7	(21.8 - 23.7) *
	65-74	823	28.4	(26.5 - 30.5)	3,794	11.6	(11.3 - 12.0) *	1,141	30.4	(28.6 - 32.2)
	75+	500	24.6	(22.5 - 26.8)	3,924	10.4	(10.1 - 10.8) *	1,047	32.8	(30.8 - 34.8) *

Web Table Only: Incidence Rates^a for Selected Common Cancer Sites^b in Adults by Age, Race/Ethnicity and Sex
in the Selected Areas^c in the United States, 1999-2003.

Cancer Site	Age Group	Hispanic ^d			Non-Hispanic White			Non-Hispanic Black		
		No. ^a	Rate ^a	95% CI	No. ^a	Rate ^a	95% CI	No. ^a	Rate ^a	95% CI
Colon and Rectum										
	20-39	812	3.3	(3.0 - 3.5)	3,941	3.7	(3.6 - 3.8) *	916	4.3	(4.0 - 4.6) *
	40-49	1,657	17.2	(16.4 - 18.0)	12,563	19.2	(18.9 - 19.6) *	2,968	27.5	(26.5 - 28.5) *
	50-64	4,459	57.4	(55.7 - 59.1)	47,006	66.4	(65.8 - 67.0) *	9,244	96.4	(94.5 - 98.4) *
	65-74	4,176	146.2	(141.8 - 150.7)	61,565	187.5	(186.1 - 189.0) *	7,928	212.2	(207.5 - 216.9) *
	75+	5,168	255.9	(248.9 - 263.0)	127,773	334.1	(332.2 - 335.9) *	10,850	338.1	(331.8 - 344.5) *
Lung and Bronchus										
	20-39	266	1.1	(0.9 - 1.2)	2,333	2.2	(2.1 - 2.3) *	457	2.2	(2.0 - 2.4) *
	40-49	888	9.2	(8.6 - 9.8)	13,911	21.3	(20.9 - 21.6) *	2,895	26.8	(25.8 - 27.8) *
	50-64	3,300	42.7	(41.2 - 44.2)	77,418	109.0	(108.2 - 109.8) *	9,898	103.5	(101.5 - 105.6) *
	65-74	3,576	125.2	(121.2 - 129.4)	100,408	306.3	(304.4 - 308.2) *	9,146	244.3	(239.4 - 249.4) *
	75+	3,472	171.0	(165.3 - 176.8)	111,591	302.2	(300.4 - 304.0) *	7,789	246.3	(240.8 - 251.8) *

Source: SEER and NPCR areas reported by the North American Association of Central Cancer Registries as meeting high quality standards for 1999-2003.

^a Age categories: No. = number; Rate is age-adjusted within the age group.

^b Common cancers selected based on available tests for diagnosis in early stages of disease progression.

^c The data from 38 cancer registries [Alabama, Alaska, California, Colorado, Connecticut, Delaware, District Of Columbia, Florida, Georgia, Hawaii, Idaho, Illinois, Indiana, Iowa, Kentucky, Louisiana, Maine, Maryland, Massachusetts, Michigan, Minnesota, Missouri, Montana, Nebraska, Nevada, New Hampshire, New Jersey, New York, Oklahoma, Oregon, Pennsylvania, Rhode Island, South Carolina, Texas, Utah, Washington, West Virginia, Wisconsin] were included covering 82% of U.S., 90% of the Hispanic, 81% of NHW and 79% of NHB populations.

^d NHIA derived Hispanic origin.

* Rate is statistically different from Hispanic rate.