

Table 11 (Web Supplement). Counts for the Stage Distributions<sup>a</sup> for Selected Common Cancer Sites<sup>b</sup> and Age Groups by Race/Ethnicity<sup>c</sup>, County Poverty<sup>d</sup>, and Sex in Selected Areas<sup>e</sup> of the United States, 2001-2003.

Cancer Site	Hispanic County Poverty			Non-Hispanic White County Poverty			Non-Hispanic Black County Poverty		
	<10%	10-19%	≥20%	<10%	10-19%	≥20%	<10%	10-19%	≥20%
Stage <sup>a</sup>	No.	No.	No.	No.	No.	No.	No.	No.	No.
<b>Male</b>									
Colon and Rectum (Aged 50 and Older)									
All	1,542	6,094	1,775	55,740	74,029	8,463	2,206	9,023	2,865
Localized	541	1,995	578	21,757	28,618	3,090	714	3,114	893
Regional	603	2,424	671	21,191	27,500	3,253	807	3,179	1,020
Distant	305	1,078	332	9,005	12,168	1,380	529	1,969	648
Unknown	93	597	194	3,787	5,743	740	156	761	304
Lung and Bronchus (Aged 20 and Older)									
All	1,635	6,396	1,937	79,862	118,704	14,114	3,711	16,448	4,884
Localized	233	806	249	13,520	20,143	2,384	503	2,371	607
Regional	369	1,330	411	20,285	29,143	3,487	957	4,165	1,185
Distant	884	3,411	958	38,339	55,472	6,494	1,901	8,324	2,528
Unknown	149	849	319	7,718	13,946	1,749	350	1,588	564
Prostate (Aged 50 and Older)									
All	4,686	17,565	5,172	157,178	193,599	19,343	9,334	33,472	10,893
Localized	3,725	13,450	3,677	127,702	152,211	15,121	7,440	25,710	8,244
Regional	479	1,736	490	14,089	17,787	1,634	829	2,992	761
Distant	202	866	286	5,129	6,795	807	466	2,103	766
Unknown	280	1,513	719	10,258	16,806	1,781	599	2,667	1,122
<b>Female</b>									
Breast (Aged 40 and Older)									
All	4,666	16,521	4,363	153,237	191,175	19,715	5,831	23,483	7,365
Localized	2,736	9,064	2,245	98,155	120,965	12,071	3,144	12,136	3,751
Regional	1,597	5,686	1,530	43,346	53,526	5,704	2,045	8,372	2,553
Distant	185	802	258	6,345	8,194	927	375	1,730	586
Unknown	148	969	330	5,391	8,490	1,013	267	1,245	475
Cervix (Aged 20 and Older)									
All	810	3,437	934	7,366	10,707	1,246	636	2,917	1,087
Localized	430	1,718	402	3,915	5,460	663	263	1,277	440
Regional	283	1,195	343	2,269	3,355	371	255	1,096	399
Distant	59	300	77	716	1,044	112	76	316	137
Unknown	38	224	112	466	848	100	42	228	111
Colon and Rectum (Aged 50 and Older)									
All	1,374	5,391	1,540	54,782	71,347	8,276	2,269	10,107	3,456
Localized	452	1,943	503	20,173	26,524	2,917	750	3,382	1,087
Regional	596	2,031	597	21,977	27,598	3,267	899	3,815	1,296
Distant	238	896	288	8,621	11,290	1,360	460	2,022	676
Unknown	88	521	152	4,011	5,935	732	160	888	397
Lung and Bronchus (Aged 20 and Older)									
All	1,280	4,447	1,184	69,095	96,273	10,538	2,663	10,838	3,595
Localized	221	743	179	14,033	18,853	1,950	438	1,765	529
Regional	299	968	256	16,865	22,844	2,477	678	2,715	869
Distant	634	2,176	548	31,248	42,448	4,744	1,310	5,274	1,764
Unknown	126	560	201	6,949	12,128	1,367	237	1,084	433

Source: SEER and NPCR areas reported by the North American Association of Central Cancer Registries as meeting high quality standards for 1999-2003.

<sup>a</sup> The 2000 SEER Summary Staging System.

<sup>b</sup> Common cancers for age groups selected based on available tests for diagnosis in early stages of disease progression. Cases excluded when based on death certificate information only.

<sup>c</sup> NHIA derived Hispanic origin.

<sup>d</sup> Percent (%) of county population below poverty in 1999.

<sup>e</sup> The data from 36 cancer registries [Alabama, Alaska, California, Colorado, Connecticut, Delaware, District Of Columbia, Florida, Georgia, Hawaii, Idaho, Illinois, Indiana, Iowa, Kentucky, Louisiana, Maine, Massachusetts, Michigan, Minnesota, Missouri, Montana, Nebraska, Nevada, New Hampshire, New Jersey, New York, Oklahoma, Oregon, Pennsylvania, Rhode Island, South Carolina, Texas, Utah, Washington, West Virginia, Wisconsin, but excludes Maryland and Hawaii data] were included covering 80% of U.S., 89% of the Hispanic, 79% NHW, and 75% NHB populations.