

Table I-18  
US COMPLETE PREVALENCE COUNTS, INVASIVE CANCERS ONLY, JANUARY 1, 2003<sup>a</sup>  
BY AGE AT PREVALENCE

Age at Prevalence	Age Specific								
	All Ages <sup>c</sup>	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70+
<u>Site/Sex</u>									
All Sites									
Males	4,692,382	16,647	37,774	65,287	145,909	311,209	617,614	1,063,685	2,434,257
Females	5,803,603	14,602	31,271	70,291	210,983	572,039	1,012,808	1,208,754	2,682,855
Oral Cavity & Pharynx									
Males	150,051	57	455	1,232	3,429	14,435	31,522	36,748	62,173
Females	85,582	97	431	1,580	3,259	8,175	14,170	18,342	39,528
Esophagus									
Males	18,104	0	0	46	184	1,124	3,858	5,475	7,418
Females	5,961	0	12	11	44	264	807	1,363	3,459
Stomach									
Males	34,299	5	12	94	520	2,064	4,793	8,224	18,587
Females	25,554	11	35	86	493	1,731	2,949	4,431	15,818
Colon & Rectum									
Males	514,789	11	46	768	4,757	19,996	59,515	114,125	315,571
Females	553,409	11	58	823	4,550	18,576	50,643	93,617	385,131
Liver & Intrahep									
Males	10,850	264	363	253	245	1,279	3,041	2,380	3,026
Females	6,495	337	304	242	342	684	993	1,368	2,224
Pancreas									
Males	13,302	0	47	34	361	1,237	2,846	3,532	5,245
Females	14,386	0	64	114	291	1,325	2,347	3,407	6,839
Larynx									
Males	78,071	0	12	33	262	2,949	11,097	21,300	42,418
Females	19,242	0	0	55	189	1,322	2,564	5,483	9,630
Lung & Bronchus									
Males	173,431	34	64	264	1,096	6,579	23,655	50,459	91,279
Females	181,558	0	22	388	1,445	7,503	25,227	49,299	97,674
Melanoma of the Skin									
Males	320,178	55	537	4,417	16,136	46,262	72,977	70,992	108,800
Females	342,255	68	886	8,787	30,219	65,632	79,188	60,803	96,672

<sup>a</sup> US 2003 cancer prevalence counts are based on 2003 cancer prevalence proportions from the SEER 9 registries (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta) and 1/1/2003 US population estimates based on the average of 2002 and 2003 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

<sup>b</sup> Cases diagnosed more than 28 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000).

<sup>c</sup> Due to rounding, the sum of the age specific estimates may not equal the all ages estimate.

Table I-18 - continued  
US COMPLETE PREVALENCE COUNTS, INVASIVE CANCERS ONLY, JANUARY 1, 2003<sup>a</sup>  
 BY AGE AT PREVALENCE

Age at Prevalence	Age Specific								
	All Ages <sup>c</sup>	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70+
<u>Site/Sex</u>									
Breast									
Males	12,241	0	12	0	138	501	2,018	3,220	6,352
Females	2,356,795	0	58	1,897	35,796	202,103	463,141	551,858	1,101,941
Cervix									
Females	253,781	0	12	2,238	19,634	45,855	54,849	48,649	82,544
Corpus & Uterus, NOS									
Females	570,806	0	68	451	4,374	22,425	69,334	115,436	358,717
Ovary <sup>d</sup>									
Females	171,840	57	1,036	2,924	7,728	21,508	35,022	38,235	65,331
Prostate									
Males	1,937,798	34	69	102	287	14,311	158,820	480,773	1,283,404
Urinary Bladder									
Males	372,313	28	168	630	2,626	13,116	40,053	82,767	232,925
Females	133,452	11	34	312	1,451	4,424	12,280	25,854	89,087
Kidney & Renal Pelvis									
Males	136,080	1,562	2,410	1,944	3,920	12,023	25,151	34,383	54,686
Females	94,068	1,692	2,445	2,393	3,237	8,407	14,562	20,318	41,015
Hodgkin Lymphoma									
Males	76,517	175	1,937	8,639	17,052	20,887	15,363	7,794	4,669
Females	70,870	78	1,774	8,645	17,642	19,549	13,169	5,097	4,917
Non-Hodgkin Lymphoma									
Males	189,637	628	3,276	5,999	11,545	24,272	37,160	41,448	65,309
Females	174,848	318	1,345	3,202	7,858	16,562	29,315	36,241	80,005
Myeloma									
Males	28,818	0	5	17	532	1,989	6,003	7,918	12,354
Females	23,112	0	0	33	229	1,199	4,167	5,911	11,573
Leukemia									
Males	112,324	5,971	10,583	8,752	7,749	9,840	14,402	19,623	35,403
Females	86,689	4,951	8,862	8,196	5,790	6,752	9,025	12,414	30,700
Acute Lymphocytic Leuk									
Males	28,078	5,125	9,066	6,824	4,440	1,343	635	402	243
Females	22,861	4,345	7,539	5,903	2,846	1,161	567	236	263

<sup>a</sup> US 2003 cancer prevalence counts are based on 2003 cancer prevalence proportions from the SEER 9 registries (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta) and 1/1/2003 US population estimates based on the average of 2002 and 2003 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

<sup>b</sup> Cases diagnosed more than 28 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000).

<sup>c</sup> Due to rounding, the sum of the age specific estimates may not equal the all ages estimate.

<sup>d</sup> Ovary excludes borderline cases or histologies 8442, 8451, 8462, 8472, and 8473.