

Table II-2  
ALL CANCER SITES (Invasive)

AGE ADJUSTED SEER INCIDENCE<sup>a</sup> RATES BY YEAR, RACE AND SEX

	<u>All Races</u>			<u>Whites</u>			<u>Blacks</u>		
	<u>Total</u>	<u>Males</u>	<u>Females</u>	<u>Total</u>	<u>Males</u>	<u>Females</u>	<u>Total</u>	<u>Males</u>	<u>Females</u>
<u>AGE-ADJUSTED RATES</u>									
<u>YEAR OF DIAGNOSIS:</u>									
1975	400.4	466.8	365.8	402.2	468.7	369.6	426.2	525.0	356.6
1976	407.3	481.1	367.3	409.7	481.8	373.2	427.3	552.9	340.4
1977	407.7	486.5	363.6	409.3	487.1	367.8	442.5	577.6	351.8
1978	407.3	488.4	362.1	409.8	489.9	367.9	436.3	570.4	343.9
1979	412.5	496.5	365.4	414.0	497.6	369.6	451.4	589.8	358.7
1980	417.9	505.7	367.9	419.4	505.2	373.2	467.0	614.6	366.1
1981	425.2	510.9	377.0	427.7	510.9	383.8	472.5	633.6	362.8
1982	424.4	511.0	376.1	427.7	512.4	383.4	465.1	621.9	359.4
1983	431.1	519.9	380.7	434.4	520.8	388.0	486.7	645.6	381.6
1984	439.8	525.4	392.3	443.4	527.4	399.3	487.5	643.7	385.1
1985	448.7	531.1	403.0	453.1	533.2	411.2	488.1	642.6	387.4
1986	451.2	538.1	402.0	455.7	542.7	408.8	493.1	642.9	396.0
1987	467.9	563.0	413.2	474.7	568.4	423.2	500.8	665.8	393.9
1988	463.6	558.2	409.0	470.0	564.8	417.6	503.5	656.9	404.2
1989	467.4	566.8	408.0	474.7	573.2	418.2	491.9	660.3	384.2
1990	481.4	592.2	415.0	488.3	598.6	424.3	520.2	698.2	407.3
1991	502.5	637.0	418.3	509.8	643.7	427.9	548.7	764.5	412.2
1992	510.0	657.3	414.4	516.0	662.0	423.1	567.4	806.9	412.4
1993	492.8	621.3	407.9	496.3	619.2	416.7	566.8	815.6	403.2
1994	482.7	588.8	412.8	486.5	586.6	422.2	555.4	779.2	407.6
1995	475.8	570.1	413.9	480.8	568.4	425.0	539.8	743.1	403.6
1996	477.7	570.7	417.0	483.2	571.5	427.0	537.2	722.5	414.4
1997	484.6	573.3	426.9	491.6	573.8	439.6	535.1	730.4	407.6
1998	486.2	570.2	432.2	494.5	573.4	445.2	529.7	713.4	411.2
1999	488.3	578.1	428.5	495.8	579.2	441.4	534.7	714.4	416.8
2000	482.8	575.5	419.4	493.2	579.1	435.4	520.6	703.8	397.9
2001	484.6	574.1	422.7	495.9	580.1	438.5	509.4	681.5	392.7
2002	479.9	565.1	421.3	489.5	570.3	434.3	522.5	682.1	415.0
2003	465.0	547.3	408.3	474.2	550.1	422.5	503.9	658.4	399.8
2004	457.8	534.2	405.0	466.8	537.9	417.9	493.1	637.2	396.6
1975-2004	460.5	552.8	402.9	465.5	554.7	411.9	505.5	676.4	392.3

<sup>a</sup> SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta). Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

- Statistic not shown. Rate based on less than 16 cases for the time interval.

Table II-3  
ALL CANCER SITES (Invasive)

AGE-ADJUSTED U.S. DEATH<sup>a</sup> RATES BY YEAR, RACE AND SEX

	<u>All Races</u>			<u>Whites</u>			<u>Blacks</u>		
	<u>Total</u>	<u>Males</u>	<u>Females</u>	<u>Total</u>	<u>Males</u>	<u>Females</u>	<u>Total</u>	<u>Males</u>	<u>Females</u>
<u>AGE-ADJUSTED RATES</u>									
<u>YEAR OF DEATH:</u>									
1975	199.1	258.4	160.0	196.3	254.0	159.0	235.5	320.5	174.2
1976	202.3	262.2	162.7	199.4	257.5	161.7	241.8	328.4	179.1
1977	203.0	264.2	162.9	199.7	258.8	161.6	246.4	337.9	181.6
1978	204.4	267.1	163.8	201.1	261.6	162.5	247.2	340.9	181.5
1979	204.5	268.7	163.1	201.2	263.2	161.7	249.0	344.7	182.7
1980	207.0	271.1	165.8	203.3	265.0	164.2	256.2	353.9	189.2
1981	206.4	269.8	165.9	202.6	263.3	164.6	256.8	358.6	187.5
1982	208.3	272.9	167.4	204.5	266.3	166.1	260.4	364.9	190.2
1983	209.2	274.1	168.3	205.2	267.2	166.7	263.8	369.3	193.2
1984	210.9	275.1	170.5	206.7	267.6	169.0	268.1	377.9	195.7
1985	211.3	275.7	171.0	207.3	268.2	169.6	268.1	377.8	196.3
1986	211.8	276.0	171.6	207.7	268.8	170.0	269.4	378.9	199.1
1987	211.9	276.3	171.5	207.8	269.0	169.9	271.1	383.2	199.9
1988	212.6	276.9	172.5	208.5	269.5	170.9	271.6	384.2	200.4
1989	214.3	278.5	174.3	210.0	270.2	172.9	275.9	395.3	200.9
1990	214.9	279.8	174.7	210.4	271.5	172.9	279.3	399.1	205.4
1991	215.1	279.1	175.3	210.6	271.2	173.4	279.0	395.9	207.4
1992	213.5	276.5	174.4	209.2	268.9	172.6	274.9	389.2	205.8
1993	213.4	275.9	174.6	209.2	267.9	172.8	277.1	394.8	206.3
1994	211.7	272.1	174.1	207.8	264.7	172.6	272.0	385.2	204.1
1995	209.9	268.5	173.4	206.1	261.5	171.7	269.5	379.0	204.5
1996	207.0	263.7	171.2	203.4	256.8	169.8	265.3	373.7	200.9
1997	203.6	258.1	169.0	199.9	251.6	167.3	262.1	363.2	202.5
1998	200.8	253.6	166.9	197.6	247.6	165.4	255.7	353.1	198.4
1999	200.7	252.8	167.2	197.8	247.1	165.9	253.8	349.2	197.5
2000	198.7	248.2	166.6	196.3	243.1	165.9	248.1	341.0	193.2
2001	195.9	244.1	164.3	193.6	239.4	163.5	244.1	332.4	191.9
2002	193.7	240.2	162.9	191.7	236.3	162.0	240.3	323.0	191.1
2003	190.0	234.1	160.5	188.3	230.6	159.7	234.4	311.3	188.1
2004	185.7	228.3	157.0	184.1	224.8	156.4	228.1	303.5	182.8
1975-2004	205.5	263.7	168.2	202.2	257.6	166.9	258.5	359.1	195.1

<sup>a</sup> NCHS public use data file for the total US. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

- Statistic not shown. Rate based on less than 16 cases for the time interval.

Table III-2  
 BRAIN AND OTHER NERVOUS SYSTEM CANCER (Invasive)

AGE ADJUSTED SEER INCIDENCE<sup>a</sup> RATES BY YEAR, RACE AND SEX

	All Races			Whites			Blacks		
	Total	Males	Females	Total	Males	Females	Total	Males	Females
<u>AGE-ADJUSTED RATES</u>									
YEAR OF DIAGNOSIS:									
1975	5.9	6.8	5.0	6.2	7.3	5.3	4.1	4.3	3.9
1976	5.8	7.1	4.7	6.2	7.5	5.0	3.3	5.4	-
1977	6.2	7.8	4.9	6.6	8.3	5.3	3.5	5.2	2.5
1978	5.8	6.8	4.9	6.1	7.2	5.2	3.9	4.3	3.5
1979	6.1	7.4	5.0	6.6	8.0	5.4	3.7	4.2	3.2
1980	6.3	7.6	5.2	6.8	8.2	5.6	3.1	3.3	2.9
1981	6.5	8.1	5.2	6.9	8.4	5.6	5.0	7.2	3.2
1982	6.4	7.9	5.3	6.9	8.6	5.6	3.7	4.4	3.1
1983	6.3	7.6	5.2	6.9	8.2	5.8	2.7	3.8	1.8
1984	6.1	7.2	5.2	6.5	7.6	5.5	4.5	5.3	3.9
1985	6.9	8.5	5.6	7.4	9.1	6.1	4.9	5.6	4.2
1986	6.8	8.1	5.8	7.3	8.6	6.3	4.1	4.7	3.5
1987	7.0	8.1	6.0	7.6	8.7	6.7	4.2	5.5	3.1
1988	6.8	7.9	5.9	7.4	8.6	6.4	4.4	4.5	4.3
1989	6.8	8.5	5.4	7.4	9.2	5.9	4.9	6.0	4.1
1990	7.0	8.1	6.2	7.6	8.8	6.6	4.3	5.2	3.6
1991	6.9	8.6	5.6	7.6	9.3	6.2	3.6	4.5	3.0
1992	7.0	8.7	5.6	7.6	9.5	6.2	3.6	4.0	3.3
1993	6.8	8.1	5.7	7.4	8.9	6.1	4.0	4.1	3.8
1994	6.6	7.9	5.4	7.1	8.5	5.9	4.2	4.5	4.0
1995	6.5	7.9	5.3	7.0	8.6	5.8	3.9	5.0	3.2
1996	6.6	8.2	5.3	7.3	9.0	5.8	4.0	5.1	3.3
1997	6.7	8.1	5.6	7.4	8.9	6.1	3.9	5.1	3.1
1998	6.6	7.7	5.7	7.3	8.4	6.3	4.0	4.3	3.8
1999	6.9	8.4	5.6	7.6	9.2	6.3	3.7	5.2	2.6
2000	6.7	8.2	5.6	7.4	9.0	6.1	4.3	5.4	3.4
2001	6.5	7.9	5.4	7.2	8.6	6.0	4.1	5.3	3.3
2002	6.7	7.9	5.6	7.4	8.8	6.2	4.2	5.2	3.5
2003	6.5	7.4	5.7	7.0	8.1	6.1	4.6	4.4	4.6
2004	6.5	7.6	5.5	7.2	8.3	6.1	3.9	4.6	3.4
1975-2004	6.6	7.9	5.5	7.1	8.5	6.0	4.0	4.9	3.4

<sup>a</sup> SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta). Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).  
 - Statistic not shown. Rate based on less than 16 cases for the time interval.

Table III-3  
BRAIN AND OTHER NERVOUS SYSTEM CANCER (Invasive)  
 AGE-ADJUSTED U.S. DEATH<sup>a</sup> RATES BY YEAR, RACE AND SEX

	<u>All Races</u>			<u>Whites</u>			<u>Blacks</u>		
	<u>Total</u>	<u>Males</u>	<u>Females</u>	<u>Total</u>	<u>Males</u>	<u>Females</u>	<u>Total</u>	<u>Males</u>	<u>Females</u>
<u>AGE-ADJUSTED RATES</u>									
<u>YEAR OF DEATH:</u>									
1975	4.1	5.0	3.4	4.3	5.2	3.6	2.4	3.2	1.8
1976	4.3	5.3	3.6	4.5	5.5	3.7	2.7	3.5	2.0
1977	4.4	5.4	3.6	4.6	5.7	3.8	2.5	3.0	2.2
1978	4.5	5.5	3.7	4.8	5.8	3.9	2.7	3.4	2.2
1979	4.3	5.2	3.5	4.5	5.5	3.6	2.6	3.3	2.1
1980	4.4	5.4	3.5	4.6	5.8	3.7	2.5	3.0	2.0
1981	4.4	5.3	3.6	4.6	5.6	3.8	2.6	3.3	2.0
1982	4.4	5.4	3.7	4.7	5.7	3.9	2.6	3.4	2.1
1983	4.4	5.3	3.6	4.7	5.6	3.9	2.5	2.9	2.2
1984	4.6	5.6	3.7	4.8	5.9	4.0	2.6	3.3	1.9
1985	4.6	5.6	3.7	4.8	5.9	4.0	2.5	3.2	2.0
1986	4.5	5.6	3.7	4.8	5.9	3.9	2.8	3.5	2.3
1987	4.7	5.7	3.9	5.0	6.0	4.2	2.9	3.6	2.4
1988	4.7	5.9	3.8	5.0	6.2	4.0	2.9	3.6	2.4
1989	4.7	5.8	3.9	5.0	6.2	4.1	2.7	3.5	2.2
1990	4.9	6.0	4.0	5.2	6.3	4.2	3.0	3.7	2.4
1991	4.9	6.0	4.1	5.3	6.4	4.4	2.8	3.4	2.4
1992	4.9	5.9	4.0	5.2	6.3	4.3	2.8	3.4	2.4
1993	4.8	5.9	3.9	5.1	6.2	4.2	2.8	3.6	2.3
1994	4.8	5.9	4.0	5.2	6.3	4.3	2.7	3.2	2.3
1995	4.7	5.7	3.9	5.0	6.0	4.1	2.8	3.4	2.3
1996	4.7	5.7	3.9	5.1	6.1	4.2	2.6	3.2	2.2
1997	4.7	5.7	3.9	5.0	6.0	4.1	2.9	3.3	2.6
1998	4.7	5.7	3.8	5.0	6.1	4.1	2.7	3.4	2.2
1999	4.6	5.7	3.8	5.0	6.0	4.1	2.7	3.3	2.3
2000	4.5	5.6	3.7	4.9	6.0	3.9	2.7	3.3	2.2
2001	4.4	5.5	3.6	4.8	5.9	3.9	2.6	3.1	2.2
2002	4.5	5.5	3.6	4.8	5.8	3.9	2.6	3.2	2.2
2003	4.4	5.4	3.6	4.7	5.7	3.9	2.7	3.4	2.1
2004	4.3	5.2	3.5	4.6	5.6	3.8	2.5	3.0	2.0
1975-2004	4.6	5.6	3.8	4.9	5.9	4.0	2.7	3.3	2.2

<sup>a</sup> NCHS public use data file for the total US. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

- Statistic not shown. Rate based on less than 16 cases for the time interval.

Table IV-3  
BREAST CANCER (Invasive)

AGE ADJUSTED SEER INCIDENCE<sup>a</sup> RATES BY YEAR, RACE AND SEX

	All Races			Whites			Blacks		
	Total	Males	Females	Total	Males	Females	Total	Males	Females
<u>AGE-ADJUSTED RATES</u>									
YEAR OF DIAGNOSIS:									
1975	58.1	0.9	105.1	59.5	0.9	107.4	52.2	-	93.7
1976	56.4	1.1	101.9	58.1	1.1	104.8	48.0	-	85.3
1977	56.0	1.0	100.8	57.5	0.9	103.4	49.7	-	87.0
1978	55.8	0.9	100.6	57.5	0.9	103.6	48.3	-	86.1
1979	56.8	1.0	102.1	58.3	0.9	104.8	49.8	-	86.8
1980	56.8	0.9	102.1	58.5	1.0	105.0	50.5	-	89.2
1981	59.3	1.0	106.3	61.2	0.9	109.9	54.4	-	94.2
1982	59.2	1.0	106.4	61.1	0.9	109.9	53.9	-	93.5
1983	61.9	1.0	111.1	63.7	1.1	114.5	58.8	-	103.4
1984	64.4	0.9	115.9	66.5	0.9	119.8	58.1	-	102.0
1985	69.0	1.0	124.1	71.0	1.0	127.8	63.9	-	111.4
1986	70.4	0.9	126.8	72.3	1.0	130.4	66.2	-	114.9
1987	74.8	1.0	134.4	77.8	1.0	140.3	63.0	-	109.3
1988	73.0	1.2	131.2	75.6	1.2	136.3	68.9	2.8	118.6
1989	70.7	1.1	127.1	73.3	1.1	132.3	60.4	-	104.7
1990	73.1	1.1	131.7	75.3	1.1	136.2	68.3	-	118.2
1991	74.0	1.1	133.7	76.6	1.2	138.8	67.3	-	117.3
1992	72.9	1.0	131.8	74.7	1.0	135.6	71.3	-	123.5
1993	71.2	1.1	129.1	73.1	1.0	133.1	68.8	-	119.4
1994	72.0	1.1	130.8	74.2	1.1	135.6	70.3	-	122.0
1995	72.7	1.0	132.4	75.0	0.9	137.4	71.7	-	123.9
1996	73.2	1.3	133.4	75.2	1.3	137.9	71.1	-	123.3
1997	75.3	1.1	137.5	77.5	1.1	142.4	71.7	-	123.9
1998	77.1	1.1	141.0	79.5	1.1	146.4	70.3	-	122.5
1999	76.9	1.0	140.8	79.3	1.0	146.4	72.2	2.2	124.7
2000	74.0	1.2	135.8	77.0	1.2	142.6	67.2	2.9	115.7
2001	74.8	1.2	137.4	77.6	1.2	143.8	65.4	1.5	113.3
2002	72.7	1.2	133.9	75.0	1.1	139.6	69.2	2.1	119.8
2003	67.6	1.3	124.6	69.5	1.3	129.3	69.4	-	120.6
2004	67.3	1.2	124.3	68.7	1.2	128.2	68.6	1.6	119.3
1975-2004	68.6	1.1	124.5	70.6	1.1	128.5	64.3	1.6	111.9

<sup>a</sup> SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta). Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

- Statistic not shown. Rate based on less than 16 cases for the time interval.

Table IV-4  
BREAST CANCER (Invasive)

AGE-ADJUSTED U.S. DEATH<sup>a</sup> RATES BY YEAR, RACE AND SEX

	<u>All Races</u>			<u>Whites</u>			<u>Blacks</u>		
	<u>Total</u>	<u>Males</u>	<u>Females</u>	<u>Total</u>	<u>Males</u>	<u>Females</u>	<u>Total</u>	<u>Males</u>	<u>Females</u>
<u>AGE-ADJUSTED RATES</u>									
<u>YEAR OF DEATH:</u>									
1975	17.8	0.4	31.4	18.0	0.4	31.8	16.8	0.4	29.5
1976	18.1	0.4	31.8	18.2	0.4	32.2	17.6	0.6	30.5
1977	18.5	0.4	32.5	18.6	0.4	32.7	18.9	0.5	32.8
1978	18.0	0.4	31.7	18.1	0.3	31.9	18.6	0.6	32.1
1979	17.7	0.3	31.2	17.9	0.3	31.5	17.9	0.5	30.8
1980	18.0	0.3	31.7	18.1	0.3	31.9	18.4	0.4	31.7
1981	18.2	0.3	31.9	18.3	0.3	32.1	19.0	0.5	32.6
1982	18.4	0.3	32.2	18.4	0.3	32.3	19.7	0.6	33.7
1983	18.3	0.3	32.1	18.4	0.3	32.2	19.6	0.5	33.5
1984	18.8	0.3	32.9	18.8	0.3	32.9	21.0	0.5	35.9
1985	18.8	0.3	33.0	18.9	0.3	33.1	20.3	0.4	34.9
1986	18.8	0.3	32.9	18.8	0.3	32.9	20.7	0.4	35.4
1987	18.7	0.3	32.7	18.6	0.3	32.6	21.5	0.5	36.7
1988	19.0	0.3	33.2	18.9	0.3	33.1	22.4	0.9	37.8
1989	19.0	0.3	33.2	18.9	0.3	33.2	21.6	0.5	36.6
1990	18.9	0.3	33.1	18.8	0.3	33.0	22.5	0.5	38.0
1991	18.7	0.3	32.7	18.5	0.3	32.4	22.6	0.4	38.3
1992	18.1	0.3	31.6	17.9	0.3	31.4	21.9	0.4	37.1
1993	18.0	0.4	31.4	17.7	0.3	31.1	22.6	0.7	38.0
1994	17.7	0.4	30.9	17.4	0.3	30.6	22.4	0.6	37.7
1995	17.4	0.4	30.6	17.1	0.3	30.1	22.7	0.8	38.2
1996	16.8	0.3	29.5	16.5	0.3	29.0	22.1	0.7	37.1
1997	16.1	0.3	28.2	15.7	0.3	27.6	22.3	0.6	37.4
1998	15.7	0.3	27.5	15.3	0.3	27.0	21.1	0.6	35.5
1999	15.2	0.3	26.6	14.8	0.3	26.0	20.9	0.7	35.2
2000	15.2	0.4	26.6	14.9	0.4	26.2	20.5	0.6	34.4
2001	14.7	0.3	26.0	14.3	0.3	25.4	20.5	0.6	34.5
2002	14.5	0.3	25.6	14.1	0.3	25.0	20.3	0.6	34.1
2003	14.2	0.3	25.2	13.8	0.3	24.6	20.2	0.5	34.1
2004	13.7	0.3	24.4	13.3	0.3	23.8	19.1	0.5	32.3
1975-2004	17.2	0.3	30.2	17.1	0.3	30.1	20.6	0.6	35.1

<sup>a</sup> NCHS public use data file for the total US. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

- Statistic not shown. Rate based on less than 16 cases for the time interval.

Table IV-5  
FEMALE BREAST CANCER (Invasive)

AGE-ADJUSTED SEER INCIDENCE<sup>a</sup> RATES BY YEAR, RACE AND AGE

	<u>All Races, Females</u>			<u>White Females</u>			<u>Black Females</u>		
	<u>All</u>	<u>&lt;50</u>	<u>50+</u>	<u>All</u>	<u>&lt;50</u>	<u>50+</u>	<u>All</u>	<u>&lt;50</u>	<u>50+</u>
<u>SEER INCIDENCE RATES</u>									
<u>YEAR OF DIAGNOSIS:</u>									
1975	105.1	40.6	273.9	107.4	40.7	281.8	93.7	42.6	227.5
1976	101.9	40.0	264.0	104.8	40.9	272.2	85.3	37.8	209.8
1977	100.8	39.1	262.4	103.4	39.4	271.0	87.0	37.6	216.4
1978	100.6	38.8	262.3	103.6	39.2	272.0	86.1	40.2	206.1
1979	102.1	38.0	269.9	104.8	38.6	278.0	86.8	37.7	215.3
1980	102.1	37.7	270.7	105.0	38.3	279.8	89.2	37.4	224.9
1981	106.3	39.2	282.0	109.9	40.1	292.5	94.2	38.9	239.3
1982	106.4	40.5	279.0	109.9	41.3	289.8	93.5	37.9	239.0
1983	111.1	40.2	296.8	114.5	40.9	307.2	103.4	41.2	266.0
1984	115.9	42.3	308.5	119.8	43.1	320.8	102.0	44.5	252.4
1985	124.1	44.3	333.1	127.8	44.5	345.9	111.4	45.3	284.4
1986	126.8	44.4	342.4	130.4	45.2	353.5	114.9	45.3	297.0
1987	134.4	45.4	367.5	140.3	46.5	385.8	109.3	43.1	282.7
1988	131.2	44.0	359.7	136.3	45.0	375.3	118.6	42.9	316.6
1989	127.1	43.4	346.4	132.3	44.4	362.3	104.7	42.2	268.5
1990	131.7	45.6	357.2	136.2	45.4	373.9	118.2	48.8	299.7
1991	133.7	47.1	360.3	138.8	47.2	378.7	117.3	50.9	291.0
1992	131.8	43.4	363.5	135.6	43.6	376.6	123.5	46.9	324.1
1993	129.1	42.7	355.1	133.1	43.2	368.4	119.4	45.2	313.6
1994	130.8	42.3	362.5	135.6	42.6	378.8	122.0	47.2	317.8
1995	132.4	43.1	366.3	137.4	43.7	382.9	123.9	45.4	329.4
1996	133.4	43.4	369.1	137.9	43.8	384.3	123.3	43.9	331.1
1997	137.5	43.3	384.0	142.4	44.0	400.0	123.9	43.9	333.6
1998	141.0	45.3	391.4	146.4	45.6	410.2	122.5	46.1	322.7
1999	140.8	43.5	395.6	146.4	44.8	412.7	124.7	43.3	338.0
2000	135.8	43.2	378.3	142.6	43.9	401.2	115.7	44.1	303.2
2001	137.4	43.4	383.4	143.8	44.4	404.2	113.3	39.2	307.4
2002	133.9	42.2	374.1	139.6	42.9	392.8	119.8	39.8	329.4
2003	124.6	42.5	339.6	129.3	43.2	354.5	120.6	42.0	326.4
2004	124.3	43.5	335.7	128.2	43.4	350.2	119.3	44.9	314.3
1975-2004	124.5	42.8	338.4	128.5	43.3	351.5	111.9	43.3	291.5

<sup>a</sup> SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta). Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

- Statistic not shown. Rate based on less than 16 cases for the time interval.

Table IV-6  
 FEMALE BREAST CANCER (Invasive)

AGE-ADJUSTED U.S. DEATH<sup>a</sup> RATES BY YEAR, RACE AND AGE

	All Races, Females			White Females			Black Females		
	All	<50	50+	All	<50	50+	All	<50	50+
<u>U.S. DEATH RATES</u>									
YEAR OF DEATH:									
1975	31.4	9.1	89.9	31.8	9.0	91.4	29.5	10.7	78.8
1976	31.8	8.7	92.2	32.2	8.6	93.8	30.5	10.2	83.5
1977	32.5	8.9	94.3	32.7	8.7	95.3	32.8	10.5	91.1
1978	31.7	8.7	92.1	31.9	8.5	93.3	32.1	11.4	86.5
1979	31.2	8.6	90.5	31.5	8.3	92.1	30.8	11.4	81.8
1980	31.7	8.6	92.2	31.9	8.4	93.6	31.7	10.8	86.3
1981	31.9	8.5	93.3	32.1	8.2	94.8	32.6	11.5	87.7
1982	32.2	8.4	94.4	32.3	8.2	95.5	33.7	11.2	92.8
1983	32.1	8.2	94.7	32.2	7.9	95.8	33.5	11.2	92.1
1984	32.9	8.6	96.6	32.9	8.2	97.5	35.9	12.5	97.2
1985	33.0	8.6	96.9	33.1	8.3	98.1	34.9	12.0	94.6
1986	32.9	8.6	96.3	32.9	8.3	97.5	35.4	12.9	94.6
1987	32.7	8.4	96.1	32.6	8.0	96.9	36.7	12.9	99.0
1988	33.2	8.4	98.2	33.1	8.0	98.8	37.8	12.7	103.4
1989	33.2	8.4	98.2	33.2	8.1	99.2	36.6	12.4	100.0
1990	33.1	8.3	98.1	33.0	8.0	98.5	38.0	12.0	106.1
1991	32.7	8.2	96.9	32.4	7.7	97.2	38.3	12.3	106.3
1992	31.6	7.8	94.2	31.4	7.4	94.5	37.1	12.0	102.7
1993	31.4	7.4	94.2	31.1	7.0	94.2	38.0	11.7	107.1
1994	30.9	7.4	92.6	30.6	6.9	92.5	37.7	11.7	105.9
1995	30.6	7.3	91.5	30.1	6.8	91.1	38.2	11.7	107.5
1996	29.5	6.9	88.5	29.0	6.4	88.3	37.1	11.7	103.6
1997	28.2	6.7	84.5	27.6	6.2	83.6	37.4	11.0	106.6
1998	27.5	6.3	83.1	27.0	5.8	82.5	35.5	10.6	100.7
1999	26.6	5.7	81.2	26.0	5.2	80.4	35.2	10.3	100.5
2000	26.6	5.9	81.0	26.2	5.4	80.6	34.4	9.9	98.3
2001	26.0	5.9	78.7	25.4	5.4	77.9	34.5	9.9	99.1
2002	25.6	5.6	77.8	25.0	5.1	77.1	34.1	10.0	97.2
2003	25.2	5.6	76.5	24.6	5.1	75.6	34.1	9.7	97.9
2004	24.4	5.2	74.6	23.8	4.6	74.0	32.3	9.3	92.4
1975-2004	30.2	7.4	90.0	30.1	7.0	90.4	35.1	11.1	97.7

<sup>a</sup> NCHS public use data file for the total US. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

- Statistic not shown. Rate based on less than 16 cases for the time interval.



Table IV-7  
 FEMALE BREAST CANCER (*In Situ*)

AGE ADJUSTED SEER INCIDENCE<sup>a</sup> RATES BY YEAR, RACE AND AGE

	All Races, Females			White Females			Black Females		
	All	<50	50+	All	<50	50+	All	<50	50+
<u>SEER INCIDENCE RATES</u>									
YEAR OF DIAGNOSIS:									
1975	5.8	3.8	11.2	5.9	3.8	11.3	3.9	-	8.3
1976	5.3	2.7	12.0	5.4	2.9	11.8	3.3	-	9.4
1977	4.3	2.6	8.7	4.3	2.6	8.9	3.0	-	-
1978	4.4	2.4	9.7	4.6	2.5	9.9	3.3	-	-
1979	4.6	2.8	9.2	4.7	2.9	9.4	4.8	-	9.6
1980	4.9	3.0	9.7	5.0	3.1	10.2	3.7	-	-
1981	5.0	2.6	11.0	4.9	2.5	11.3	4.1	-	8.7
1982	5.3	2.7	12.0	5.3	2.7	12.2	5.6	2.7	12.9
1983	6.1	3.0	14.1	6.2	3.0	14.5	5.4	-	13.2
1984	8.6	4.8	18.7	9.2	5.2	19.6	6.3	2.4	16.8
1985	11.5	5.9	26.2	12.1	6.3	27.4	8.2	3.5	20.6
1986	13.6	6.8	31.6	14.3	7.2	32.8	9.2	5.1	20.0
1987	17.4	7.8	42.4	18.2	8.3	44.3	12.3	5.5	30.1
1988	17.6	8.2	42.2	18.1	8.5	43.2	16.1	7.4	38.9
1989	16.5	7.0	41.4	17.1	7.3	42.7	14.5	5.5	38.1
1990	19.1	8.4	47.2	19.9	8.8	49.2	15.1	5.5	40.0
1991	19.8	8.8	48.6	20.4	8.9	50.8	15.3	7.3	36.3
1992	21.4	9.0	54.0	22.2	9.2	56.2	19.1	6.8	51.2
1993	20.7	8.4	53.1	21.4	8.8	54.2	16.5	6.1	43.9
1994	22.3	8.8	57.8	22.9	9.1	59.2	20.1	7.5	53.3
1995	24.7	9.6	64.1	25.0	9.9	64.6	22.6	6.4	65.2
1996	25.4	9.8	66.1	25.9	9.8	68.1	22.4	7.7	61.0
1997	28.4	11.1	73.6	29.0	11.2	75.4	28.0	11.4	71.4
1998	32.8	12.0	87.3	33.8	12.3	89.9	30.3	10.8	81.5
1999	32.8	12.0	87.2	33.2	12.2	88.0	29.7	10.4	80.2
2000	32.8	11.8	87.9	33.9	12.5	89.9	29.3	8.0	85.1
2001	33.5	11.5	91.2	34.1	12.0	92.0	31.5	8.9	90.7
2002	33.8	11.8	91.6	34.3	12.2	92.2	28.4	8.6	80.5
2003	32.0	11.8	84.9	32.9	12.3	87.1	29.3	9.2	81.9
2004	32.5	12.5	85.1	33.2	12.7	86.9	30.2	9.6	84.0
1975-2004	19.6	8.3	49.2	19.7	8.4	49.3	18.0	6.7	47.4

<sup>a</sup> SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta). Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

- Statistic not shown. Rate based on less than 16 cases for the time interval.

Table V-2  
CERVIX UTERI CANCER (Invasive)

AGE-ADJUSTED SEER INCIDENCE<sup>a</sup> RATES BY YEAR, RACE AND AGE

	<u>All Races, Females</u>			<u>White Females</u>			<u>Black Females</u>		
	<u>All</u>	<u>&lt;50</u>	<u>50+</u>	<u>All</u>	<u>&lt;50</u>	<u>50+</u>	<u>All</u>	<u>&lt;50</u>	<u>50+</u>
<u>SEER INCIDENCE RATES</u>									
<u>YEAR OF DIAGNOSIS:</u>									
1975	14.8	9.7	28.2	13.2	8.9	24.8	33.1	17.6	73.7
1976	14.2	9.5	26.7	12.9	8.8	23.7	29.3	15.8	64.4
1977	13.0	8.4	25.0	11.7	7.5	22.5	28.3	15.9	60.8
1978	12.5	8.3	23.6	11.4	7.8	20.9	24.1	12.4	55.0
1979	12.7	8.7	23.4	11.0	7.6	19.9	28.8	16.0	62.4
1980	12.2	8.1	23.0	11.0	7.3	20.6	23.5	13.6	49.4
1981	10.8	7.4	19.7	9.6	6.6	17.3	24.0	13.6	51.2
1982	10.6	7.1	19.8	9.4	6.7	16.5	21.8	10.2	52.4
1983	10.5	7.3	18.8	9.6	6.8	17.0	18.8	11.9	37.0
1984	11.0	8.0	19.0	10.0	7.4	16.8	21.0	12.4	43.3
1985	10.2	7.1	18.5	9.1	6.7	15.4	19.0	9.5	43.8
1986	10.8	7.9	18.6	9.7	7.3	16.0	18.4	10.7	38.5
1987	10.0	7.1	17.6	9.0	6.9	14.6	18.7	8.7	45.0
1988	10.6	7.9	17.7	9.5	7.4	15.0	19.0	10.4	41.3
1989	10.7	7.9	18.0	10.0	7.9	15.8	15.7	8.0	35.9
1990	10.6	8.0	17.5	9.9	7.7	15.7	16.7	9.6	35.2
1991	10.0	7.0	17.9	9.2	6.8	15.6	16.3	9.2	35.0
1992	9.9	7.5	16.4	9.4	7.2	15.2	14.1	9.9	25.0
1993	9.6	7.1	16.4	9.2	7.0	14.8	13.7	8.9	26.5
1994	9.4	6.9	16.2	8.7	6.8	13.7	14.0	6.9	32.6
1995	8.8	6.7	14.3	7.9	6.5	11.7	13.5	7.7	28.5
1996	9.6	7.3	15.5	8.7	7.1	13.0	14.2	8.0	30.6
1997	9.2	7.1	14.7	8.4	6.9	12.3	13.6	8.1	28.1
1998	9.1	7.1	14.4	8.4	6.8	12.5	13.0	8.9	23.9
1999	8.2	6.5	12.8	7.8	6.5	11.0	13.4	8.0	27.3
2000	7.7	5.8	12.6	7.4	5.9	11.4	10.2	6.4	20.1
2001	7.9	5.8	13.3	7.4	5.7	11.6	11.0	5.0	26.9
2002	7.3	5.6	11.8	7.0	5.7	10.4	10.2	6.3	20.3
2003	7.2	5.7	11.2	6.7	5.5	9.8	10.5	6.1	22.1
2004	7.0	5.4	11.1	6.7	5.6	9.8	10.3	6.3	20.7
1975-2004	10.0	7.2	17.2	9.2	6.9	15.1	16.5	9.2	35.5

<sup>a</sup> SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta). Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

- Statistic not shown. Rate based on less than 16 cases for the time interval.

Table V-3  
CERVIX UTERI CANCER (Invasive)

AGE-ADJUSTED U.S. DEATH<sup>a</sup> RATES BY YEAR, RACE AND AGE

	<u>All Races, Females</u>			<u>White Females</u>			<u>Black Females</u>		
	<u>All</u>	<u>&lt;50</u>	<u>50+</u>	<u>All</u>	<u>&lt;50</u>	<u>50+</u>	<u>All</u>	<u>&lt;50</u>	<u>50+</u>
<u>U.S. DEATH RATES</u>									
<u>YEAR OF DEATH:</u>									
1975	5.6	2.4	13.9	4.7	1.9	12.1	13.3	5.7	33.3
1976	5.4	2.2	13.9	4.7	1.9	12.1	12.9	4.9	33.7
1977	5.0	2.0	12.7	4.2	1.6	11.0	12.2	5.0	31.1
1978	4.8	2.1	12.0	4.2	1.8	10.4	11.6	4.8	29.6
1979	4.6	1.9	11.6	3.9	1.6	10.2	10.6	4.5	26.7
1980	4.5	2.0	11.0	3.8	1.6	9.5	10.8	4.6	26.9
1981	4.3	1.8	10.9	3.7	1.6	9.4	9.9	3.7	26.0
1982	4.1	1.8	10.2	3.5	1.5	8.7	9.8	3.6	25.9
1983	4.0	1.8	9.9	3.4	1.5	8.4	9.8	3.9	25.2
1984	3.9	1.8	9.5	3.4	1.5	8.1	9.3	3.9	23.3
1985	3.8	1.7	9.4	3.3	1.5	7.9	9.0	3.4	23.7
1986	3.8	1.8	9.2	3.2	1.5	7.8	9.2	4.0	22.9
1987	3.6	1.7	8.6	3.1	1.5	7.4	8.2	3.6	20.3
1988	3.6	1.7	8.6	3.1	1.5	7.2	8.2	3.0	21.7
1989	3.6	1.7	8.5	3.0	1.5	7.1	8.6	3.3	22.6
1990	3.7	1.8	8.6	3.2	1.6	7.4	7.8	3.4	19.2
1991	3.5	1.7	8.2	3.0	1.5	7.0	7.6	3.2	19.4
1992	3.5	1.7	8.2	3.0	1.5	6.9	8.1	3.4	20.4
1993	3.4	1.7	8.0	2.9	1.5	6.8	7.7	3.1	19.6
1994	3.4	1.7	7.9	3.0	1.5	6.9	6.7	2.8	16.8
1995	3.2	1.6	7.5	2.8	1.5	6.4	7.0	3.0	17.5
1996	3.2	1.6	7.5	2.9	1.5	6.5	6.2	2.5	15.8
1997	3.1	1.6	7.2	2.8	1.4	6.2	6.5	2.8	16.0
1998	3.0	1.6	6.7	2.6	1.5	5.8	5.9	2.5	14.7
1999	2.8	1.4	6.7	2.6	1.3	5.8	5.4	1.9	14.6
2000	2.8	1.4	6.5	2.5	1.2	5.7	5.4	2.2	14.0
2001	2.7	1.4	6.1	2.4	1.3	5.4	4.8	1.9	12.4
2002	2.6	1.3	5.9	2.3	1.2	5.1	5.0	2.1	12.6
2003	2.5	1.2	5.7	2.2	1.2	5.1	4.7	1.9	11.8
2004	2.4	1.3	5.5	2.2	1.2	4.8	4.5	1.9	11.3
1975-2004	3.6	1.7	8.6	3.1	1.5	7.4	7.7	3.1	19.9

<sup>a</sup> NCHS public use data file for the total US. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

- Statistic not shown. Rate based on less than 16 cases for the time interval.

Table VI-3  
COLON AND RECTUM CANCER (Invasive)

AGE ADJUSTED SEER INCIDENCE<sup>a</sup> RATES BY YEAR, RACE AND SEX

	All Races			Whites			Blacks		
	Total	Males	Females	Total	Males	Females	Total	Males	Females
<u>AGE-ADJUSTED RATES</u>									
YEAR OF DIAGNOSIS:									
1975	59.5	68.4	53.6	60.2	69.8	54.0	56.8	59.2	54.7
1976	61.4	71.7	54.1	62.1	72.7	55.0	55.1	58.5	51.3
1977	62.4	73.5	55.0	63.2	74.3	55.8	60.6	75.8	51.3
1978	62.0	72.4	55.2	62.8	73.9	55.6	62.2	65.7	60.6
1979	62.4	74.1	54.5	62.9	75.2	54.9	58.6	61.6	56.3
1980	63.7	74.5	56.5	64.0	74.9	56.8	70.6	79.4	64.0
1981	64.2	75.8	56.5	64.9	76.7	57.1	62.8	72.8	55.9
1982	62.8	75.2	54.5	63.5	76.3	55.1	61.3	69.0	55.9
1983	63.7	75.6	55.6	64.3	76.6	56.1	66.7	75.5	61.1
1984	64.8	77.7	56.1	65.8	79.3	56.7	62.2	70.1	58.0
1985	66.3	79.1	57.4	67.2	80.1	58.4	63.8	74.0	56.3
1986	64.2	77.8	54.7	64.6	78.9	54.7	64.4	72.7	58.6
1987	62.7	77.5	52.7	62.9	78.2	52.6	66.2	77.4	59.0
1988	61.4	75.2	51.8	61.5	75.7	51.7	63.1	72.4	56.9
1989	61.7	74.9	52.4	62.0	75.4	52.7	64.2	79.0	54.5
1990	60.7	74.3	51.3	60.6	74.7	51.0	66.6	74.7	61.5
1991	59.5	72.9	50.0	59.3	73.1	49.7	65.6	76.3	58.2
1992	58.0	70.4	49.2	57.8	70.7	48.7	65.4	78.2	57.3
1993	56.8	68.4	48.3	56.3	68.1	47.8	64.3	77.9	55.1
1994	55.6	66.3	47.7	55.4	66.4	47.1	64.2	74.9	57.9
1995	54.0	63.7	46.9	53.6	63.3	46.6	62.0	71.2	56.1
1996	54.7	65.4	46.7	54.7	65.8	46.4	58.3	66.5	52.6
1997	56.3	68.0	47.9	56.3	67.9	48.0	64.0	76.7	55.7
1998	56.6	66.9	49.2	56.6	67.1	48.9	62.6	73.5	55.9
1999	55.4	65.6	47.8	55.2	65.6	47.1	63.0	71.9	57.8
2000	53.9	63.5	46.5	53.7	63.0	46.3	63.8	73.5	57.4
2001	53.2	62.5	45.9	52.6	62.0	45.1	63.5	74.0	56.7
2002	52.5	60.6	46.1	51.7	59.4	45.5	61.6	72.4	54.8
2003	49.9	58.5	43.0	49.4	57.6	42.7	60.4	72.9	52.5
2004	48.2	56.7	41.7	47.5	56.2	40.7	59.0	70.0	51.7
1975-2004	58.3	69.3	50.4	58.5	69.7	50.5	62.7	72.6	56.3

<sup>a</sup> SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta). Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).  
- Statistic not shown. Rate based on less than 16 cases for the time interval.

Table VI-4  
COLON CANCER (Invasive)

AGE ADJUSTED SEER INCIDENCE<sup>a</sup> RATES BY YEAR, RACE AND SEX

	All Races			Whites			Blacks		
	Total	Males	Females	Total	Males	Females	Total	Males	Females
<u>AGE-ADJUSTED RATES</u>									
YEAR OF DIAGNOSIS:									
1975	41.3	45.4	38.9	41.7	46.1	39.1	42.8	44.0	41.6
1976	43.1	47.8	39.9	43.4	48.2	40.4	43.1	43.6	42.0
1977	44.0	49.6	40.4	44.7	50.1	41.2	45.4	58.4	37.6
1978	43.8	49.1	40.6	44.1	49.9	40.6	47.8	48.8	47.9
1979	43.8	50.2	39.6	44.1	51.0	39.7	44.7	45.6	44.0
1980	45.6	50.7	42.3	45.6	50.9	42.3	55.0	58.0	52.7
1981	45.5	51.3	41.7	45.9	52.0	42.1	46.9	50.6	44.4
1982	45.8	53.5	40.7	46.1	54.1	40.9	48.3	52.5	45.2
1983	46.1	53.0	41.6	46.4	53.5	41.9	50.2	58.5	45.2
1984	46.5	54.6	41.3	47.0	55.6	41.6	48.6	54.0	45.8
1985	47.9	55.0	43.0	48.4	55.6	43.7	49.8	57.2	44.2
1986	46.8	54.4	41.6	47.1	55.2	41.5	49.2	55.0	45.5
1987	45.2	54.4	39.1	45.1	54.5	39.1	51.3	59.7	45.7
1988	44.3	52.6	38.7	44.3	52.8	38.4	48.9	54.0	45.6
1989	44.3	52.4	38.9	44.5	52.5	39.2	48.8	60.3	41.4
1990	44.2	51.9	38.9	44.0	51.9	38.8	51.9	57.7	48.5
1991	43.6	51.8	37.9	43.3	51.9	37.4	51.7	57.7	47.7
1992	42.3	49.7	37.2	42.1	49.8	36.7	50.7	60.1	45.2
1993	41.6	48.9	36.3	41.1	48.6	35.7	51.1	60.4	45.0
1994	40.8	46.9	36.3	40.6	47.1	35.8	49.8	56.6	46.2
1995	39.6	45.0	35.9	39.3	44.6	35.5	48.7	54.8	45.4
1996	39.3	45.3	34.9	39.3	45.7	34.4	44.4	48.3	41.9
1997	41.0	48.3	35.9	41.1	48.2	36.1	48.8	58.4	42.3
1998	41.0	46.7	37.0	40.8	47.0	36.5	47.4	54.1	43.9
1999	39.7	45.3	35.6	39.5	45.2	35.1	48.3	54.9	44.5
2000	39.2	44.3	35.3	38.9	43.9	35.0	50.6	56.9	46.6
2001	38.9	43.7	35.3	38.5	43.4	34.7	48.4	55.0	44.2
2002	38.2	42.7	34.7	37.5	41.6	34.2	48.3	55.8	43.5
2003	36.3	40.7	32.7	36.0	40.0	32.6	46.1	54.7	40.8
2004	35.1	40.0	31.5	34.6	39.7	30.7	44.9	51.6	40.5
1975-2004	42.1	48.2	37.7	42.1	48.4	37.7	48.4	54.8	44.3

<sup>a</sup> SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta). Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).  
- Statistic not shown. Rate based on less than 16 cases for the time interval.

Table VI-5  
RECTUM CANCER (Invasive)

AGE ADJUSTED SEER INCIDENCE<sup>a</sup> RATES BY YEAR, RACE AND SEX

	All Races			Whites			Blacks		
	Total	Males	Females	Total	Males	Females	Total	Males	Females
<u>AGE-ADJUSTED RATES</u>									
YEAR OF DIAGNOSIS:									
1975	18.2	23.0	14.8	18.5	23.6	14.9	14.0	15.2	13.0
1976	18.3	24.0	14.2	18.7	24.5	14.6	12.0	14.9	9.4
1977	18.4	23.9	14.6	18.5	24.3	14.6	15.3	17.3	13.8
1978	18.2	23.2	14.7	18.7	24.0	15.0	14.4	16.9	12.8
1979	18.6	23.9	15.0	18.8	24.2	15.2	13.9	16.1	12.3
1980	18.2	23.9	14.2	18.4	24.0	14.5	15.6	21.4	11.3
1981	18.8	24.5	14.8	19.0	24.8	15.0	15.9	22.2	11.6
1982	17.0	21.7	13.8	17.4	22.2	14.2	13.0	16.5	10.7
1983	17.6	22.6	14.0	17.9	23.1	14.2	16.5	17.0	15.9
1984	18.3	23.0	14.8	18.8	23.7	15.1	13.7	16.1	12.1
1985	18.4	24.1	14.4	18.8	24.6	14.7	14.0	16.8	12.1
1986	17.4	23.4	13.1	17.5	23.7	13.2	15.2	17.7	13.2
1987	17.5	23.0	13.5	17.7	23.7	13.5	14.9	17.6	13.3
1988	17.1	22.6	13.1	17.3	22.9	13.3	14.2	18.4	11.3
1989	17.3	22.5	13.5	17.5	22.9	13.5	15.4	18.8	13.1
1990	16.5	22.4	12.4	16.6	22.7	12.3	14.6	17.0	13.0
1991	15.9	21.0	12.1	16.0	21.2	12.3	13.9	18.5	10.5
1992	15.7	20.8	12.0	15.8	20.9	12.0	14.6	18.1	12.1
1993	15.2	19.5	12.0	15.3	19.4	12.1	13.2	17.5	10.1
1994	14.8	19.3	11.4	14.7	19.3	11.3	14.4	18.3	11.7
1995	14.4	18.7	11.1	14.4	18.7	11.1	13.2	16.3	10.7
1996	15.4	20.1	11.8	15.4	20.1	12.0	13.8	18.2	10.7
1997	15.3	19.6	12.0	15.3	19.6	11.9	15.2	18.3	13.4
1998	15.7	20.2	12.2	15.7	20.1	12.4	15.2	19.4	12.0
1999	15.6	20.3	12.1	15.7	20.4	12.0	14.7	17.0	13.2
2000	14.7	19.2	11.2	14.8	19.2	11.3	13.2	16.6	10.8
2001	14.2	18.8	10.6	14.1	18.6	10.4	15.0	19.0	12.4
2002	14.3	18.0	11.4	14.2	17.9	11.3	13.3	16.6	11.3
2003	13.6	17.7	10.3	13.4	17.5	10.1	14.3	18.2	11.7
2004	13.1	16.7	10.3	12.9	16.5	9.9	14.1	18.4	11.2
1975-2004	16.2	21.0	12.6	16.4	21.2	12.8	14.3	17.7	11.9

<sup>a</sup> SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta). Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).  
- Statistic not shown. Rate based on less than 16 cases for the time interval.

Table VI-6  
COLON AND RECTUM CANCER (Invasive)

AGE-ADJUSTED U.S. DEATH<sup>a</sup> RATES BY YEAR, RACE AND SEX

	All Races			Whites			Blacks		
	Total	Males	Females	Total	Males	Females	Total	Males	Females
<u>AGE-ADJUSTED RATES</u>									
YEAR OF DEATH:									
1975	28.1	32.8	25.0	28.3	33.2	25.1	26.8	30.1	24.6
1976	28.6	33.3	25.4	28.7	33.8	25.5	28.2	31.2	26.3
1977	28.2	33.2	24.9	28.3	33.5	25.0	27.6	31.4	24.9
1978	28.5	33.7	25.1	28.7	34.1	25.2	27.9	31.2	25.7
1979	28.1	33.6	24.6	28.2	33.9	24.6	28.2	31.6	26.0
1980	28.1	33.7	24.4	28.1	34.0	24.3	28.8	33.2	25.9
1981	27.5	33.0	24.0	27.5	33.2	23.9	28.9	32.8	26.3
1982	27.2	33.0	23.6	27.2	33.1	23.4	29.1	33.7	26.1
1983	27.1	33.1	23.2	27.1	33.3	23.1	28.7	32.9	25.8
1984	27.3	33.0	23.6	27.3	33.1	23.5	29.6	34.6	26.2
1985	26.9	32.7	23.1	26.8	32.7	22.8	30.5	34.1	28.0
1986	26.2	32.1	22.3	26.0	32.0	22.1	29.8	35.4	26.3
1987	25.9	31.9	22.0	25.7	31.8	21.6	29.8	35.1	26.5
1988	25.3	31.1	21.5	25.1	31.0	21.2	29.2	34.3	25.9
1989	25.0	30.8	21.2	24.7	30.5	20.8	30.1	36.4	26.1
1990	24.6	30.8	20.6	24.3	30.4	20.2	30.8	37.6	26.6
1991	24.0	29.6	20.3	23.6	29.4	19.8	30.0	35.8	26.5
1992	23.6	29.3	19.9	23.2	29.0	19.4	29.5	36.2	25.4
1993	23.3	28.6	19.8	22.9	28.2	19.3	29.8	35.8	26.0
1994	22.9	28.3	19.3	22.5	27.9	18.8	29.0	35.7	24.9
1995	22.6	27.7	19.1	22.1	27.2	18.6	29.5	36.3	25.5
1996	21.9	26.7	18.5	21.4	26.3	18.0	28.6	34.5	24.9
1997	21.5	26.2	18.1	20.9	25.7	17.6	28.8	35.1	24.8
1998	21.2	25.6	18.0	20.7	25.1	17.5	28.3	33.9	24.8
1999	20.9	25.5	17.8	20.4	25.0	17.2	28.5	34.5	24.7
2000	20.7	25.0	17.5	20.1	24.4	17.0	28.1	34.9	23.9
2001	20.1	24.2	17.1	19.5	23.7	16.5	27.7	33.2	24.2
2002	19.6	23.8	16.5	19.1	23.2	16.0	26.9	33.4	22.8
2003	19.0	23.0	16.0	18.5	22.4	15.5	26.4	32.1	22.7
2004	17.9	21.6	15.2	17.4	21.1	14.7	24.8	30.2	21.2
1975-2004	23.9	28.9	20.5	23.7	28.7	20.2	28.6	34.0	25.2

<sup>a</sup> NCHS public use data file for the total US. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

- Statistic not shown. Rate based on less than 16 cases for the time interval.

Table VII-2  
CORPUS AND UTERUS, NOS CANCER (Invasive)

AGE-ADJUSTED SEER INCIDENCE<sup>a</sup> RATES BY YEAR, RACE AND AGE

	All Races, Females			White Females			Black Females		
	All	<50	50+	All	<50	50+	All	<50	50+
<u>SEER INCIDENCE RATES</u>									
YEAR OF DIAGNOSIS:									
1975	35.5	6.5	111.4	37.3	6.8	117.2	19.9	3.6	62.5
1976	34.4	6.2	108.1	36.2	6.3	114.7	17.6	3.3	55.1
1977	31.7	5.7	99.5	33.1	5.9	104.4	20.1	3.5	63.4
1978	30.0	6.1	92.8	31.7	6.3	98.0	18.6	3.7	57.5
1979	27.9	5.3	87.0	29.2	5.3	91.8	16.8	3.9	50.5
1980	27.3	5.0	85.8	28.5	5.0	90.2	16.2	-	52.1
1981	27.3	5.6	84.0	28.3	5.5	88.3	17.4	2.9	55.5
1982	26.8	5.0	83.8	28.2	5.2	88.5	17.1	2.5	55.2
1983	26.5	4.9	83.3	27.8	4.9	87.7	18.8	3.4	59.2
1984	25.7	4.6	80.9	27.1	4.8	85.5	17.4	-	58.4
1985	25.2	4.6	79.2	26.5	4.8	83.5	17.5	2.2	57.5
1986	24.4	4.8	75.6	25.6	5.0	79.6	16.8	2.5	54.1
1987	24.7	5.0	76.4	26.1	5.3	80.8	15.8	-	51.8
1988	23.6	4.5	73.4	24.6	4.5	77.3	16.9	3.0	53.3
1989	24.4	4.5	76.6	25.5	4.8	79.7	19.4	1.8	65.4
1990	25.1	4.6	78.6	26.5	4.8	83.4	16.9	2.3	55.3
1991	24.6	4.4	77.5	26.0	4.5	82.3	17.4	3.1	54.7
1992	24.8	4.8	77.1	26.2	4.8	82.0	16.9	2.8	54.1
1993	24.3	5.2	74.3	25.7	5.1	79.5	17.1	3.6	52.6
1994	25.1	5.0	77.8	26.3	5.0	81.9	17.7	2.5	57.6
1995	25.3	4.7	79.3	26.5	4.6	83.6	18.7	2.7	60.6
1996	24.9	4.5	78.3	26.0	4.4	82.5	19.3	3.2	61.5
1997	25.8	5.0	80.4	27.3	5.0	85.6	16.5	2.5	53.1
1998	25.7	4.8	80.3	27.0	4.7	85.3	19.4	3.0	62.3
1999	25.3	5.2	78.0	26.7	5.1	83.5	17.5	2.8	55.9
2000	24.7	5.2	75.9	26.2	5.3	80.7	17.9	3.0	57.1
2001	25.2	5.1	77.8	26.7	5.0	83.5	18.9	3.9	58.1
2002	24.7	5.0	76.3	25.1	4.5	78.9	23.1	4.7	71.2
2003	24.0	5.2	73.3	25.2	5.2	77.5	19.6	3.6	61.6
2004	24.4	5.3	74.4	25.4	5.1	78.7	19.1	3.3	60.5
1975-2004	26.2	5.0	81.8	27.6	5.1	86.7	18.1	3.0	57.7

<sup>a</sup> SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta). Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

- Statistic not shown. Rate based on less than 16 cases for the time interval.



Table VII-3  
CORPUS AND UTERUS, NOS CANCER (Invasive)

AGE-ADJUSTED U.S. DEATH<sup>a</sup> RATES BY YEAR, RACE AND AGE

	All Races, Females			White Females			Black Females		
	All	<50	50+	All	<50	50+	All	<50	50+
<u>U.S. DEATH RATES</u>									
YEAR OF DEATH:									
1975	5.3	0.5	17.7	5.1	0.5	17.1	7.8	0.9	25.8
1976	5.3	0.5	17.9	5.1	0.5	17.2	8.1	0.9	27.0
1977	5.2	0.5	17.5	5.0	0.5	16.7	8.4	0.8	28.0
1978	5.1	0.5	17.1	4.8	0.5	16.2	8.2	0.8	27.3
1979	5.0	0.4	17.0	4.7	0.4	16.2	7.9	0.7	26.7
1980	5.1	0.4	17.4	4.8	0.4	16.5	8.4	0.9	28.1
1981	4.9	0.4	16.6	4.7	0.4	15.9	7.3	0.6	25.1
1982	4.9	0.4	16.5	4.6	0.4	15.7	7.6	0.4	26.4
1983	4.8	0.4	16.4	4.6	0.3	15.8	7.3	0.6	24.8
1984	4.7	0.4	16.0	4.4	0.3	15.2	7.7	0.7	26.1
1985	4.6	0.4	15.7	4.4	0.3	15.0	7.4	0.5	25.3
1986	4.4	0.3	15.2	4.2	0.3	14.4	7.3	0.6	24.9
1987	4.4	0.3	15.1	4.2	0.3	14.3	7.3	0.5	25.1
1988	4.3	0.3	14.9	4.1	0.3	14.0	7.4	0.5	25.6
1989	4.3	0.3	14.5	4.0	0.3	13.6	7.3	0.5	25.0
1990	4.3	0.3	14.8	4.1	0.3	13.9	7.4	0.4	25.6
1991	4.2	0.3	14.2	3.9	0.3	13.3	7.3	0.5	25.1
1992	4.2	0.3	14.4	4.0	0.3	13.6	7.0	0.4	24.1
1993	4.2	0.3	14.3	3.9	0.3	13.3	7.3	0.4	25.2
1994	4.2	0.3	14.3	3.9	0.3	13.4	7.1	0.3	24.7
1995	4.1	0.3	14.1	4.0	0.3	13.5	6.5	0.3	22.5
1996	4.1	0.3	14.1	3.9	0.3	13.2	7.2	0.4	25.0
1997	4.0	0.3	13.7	3.8	0.3	12.9	6.7	0.3	23.6
1998	4.1	0.3	14.0	3.9	0.3	13.2	6.8	0.4	23.6
1999	4.1	0.3	13.9	3.9	0.3	13.1	6.8	0.4	23.7
2000	4.1	0.4	13.9	3.8	0.3	13.0	7.2	0.4	25.1
2001	4.2	0.4	14.2	3.9	0.3	13.4	7.0	0.5	24.0
2002	4.2	0.3	14.1	3.9	0.3	13.1	7.3	0.4	25.4
2003	4.1	0.3	14.0	3.9	0.3	13.2	7.0	0.5	24.2
2004	4.1	0.4	13.9	3.9	0.4	13.0	7.0	0.5	24.0
1975-2004	4.4	0.4	15.1	4.2	0.3	14.3	7.3	0.5	25.1

<sup>a</sup> NCHS public use data file for the total US. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

- Statistic not shown. Rate based on less than 16 cases for the time interval.

Table VIII-2  
ESOPHAGUS CANCER (Invasive)

AGE ADJUSTED SEER INCIDENCE<sup>a</sup> RATES BY YEAR, RACE AND SEX

	<u>All Races</u>			<u>Whites</u>			<u>Blacks</u>		
	<u>Total</u>	<u>Males</u>	<u>Females</u>	<u>Total</u>	<u>Males</u>	<u>Females</u>	<u>Total</u>	<u>Males</u>	<u>Females</u>
<u>AGE-ADJUSTED RATES</u>									
<u>YEAR OF DIAGNOSIS:</u>									
1975	4.1	6.8	2.1	3.5	5.6	2.0	10.9	19.9	3.7
1976	4.3	6.9	2.3	3.7	5.9	2.1	10.7	16.9	5.6
1977	4.1	6.6	2.2	3.3	5.3	1.9	12.0	20.9	5.4
1978	4.1	7.0	1.9	3.4	5.7	1.7	13.0	24.4	4.3
1979	4.4	7.4	2.2	3.7	6.2	2.0	12.9	23.0	5.2
1980	4.3	6.9	2.3	3.5	5.7	1.9	12.3	18.8	7.4
1981	4.1	6.5	2.4	3.3	5.2	1.9	13.5	21.3	7.5
1982	4.3	6.8	2.3	3.5	5.5	2.0	13.6	22.7	6.4
1983	4.3	7.4	2.0	3.6	6.1	1.8	12.2	22.2	4.3
1984	4.2	6.7	2.3	3.5	5.6	1.9	12.0	19.9	6.0
1985	4.5	7.4	2.2	3.7	6.3	1.9	12.7	22.1	5.8
1986	4.6	7.5	2.5	3.9	6.2	2.2	14.1	24.0	6.3
1987	4.5	7.5	2.3	3.9	6.4	2.0	11.4	20.1	4.8
1988	4.4	7.3	2.2	3.9	6.3	2.0	11.2	19.5	5.0
1989	4.4	7.3	2.2	3.8	6.2	2.0	10.5	17.3	5.4
1990	4.8	8.3	2.2	4.2	7.2	1.9	12.5	21.4	5.8
1991	4.6	7.4	2.4	4.2	6.7	2.2	10.2	17.8	4.9
1992	4.6	8.0	2.0	4.2	7.4	1.8	10.6	18.0	5.3
1993	4.6	7.7	2.2	4.2	6.9	2.1	10.0	17.5	4.6
1994	4.5	7.7	2.1	4.2	7.2	1.9	9.2	14.7	5.1
1995	4.4	7.1	2.2	4.1	6.6	2.1	8.4	13.7	4.4
1996	4.8	8.1	2.2	4.4	7.5	2.0	9.7	16.4	4.8
1997	4.7	7.8	2.2	4.6	7.7	2.0	7.1	10.8	4.5
1998	4.7	8.2	2.0	4.5	7.9	1.9	7.8	12.5	4.2
1999	4.9	8.4	2.1	4.9	8.4	2.0	7.2	11.5	4.1
2000	4.8	8.0	2.2	4.7	8.0	2.0	7.0	10.6	4.4
2001	4.8	8.4	1.9	4.8	8.4	1.9	6.3	11.3	2.6
2002	4.6	7.9	2.0	4.6	8.0	1.8	6.3	10.0	3.5
2003	4.6	8.0	1.9	4.7	8.2	1.9	6.3	10.0	3.7
2004	5.2	8.7	2.3	5.3	9.0	2.3	6.0	8.8	3.9
1975-2004	4.5	7.6	2.2	4.1	6.9	2.0	9.7	16.3	4.8

<sup>a</sup> SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta). Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).  
 - Statistic not shown. Rate based on less than 16 cases for the time interval.

Table VIII-3  
ESOPHAGUS CANCER (Invasive)

AGE-ADJUSTED U.S. DEATH<sup>a</sup> RATES BY YEAR, RACE AND SEX

	<u>All Races</u>			<u>Whites</u>			<u>Blacks</u>		
	<u>Total</u>	<u>Males</u>	<u>Females</u>	<u>Total</u>	<u>Males</u>	<u>Females</u>	<u>Total</u>	<u>Males</u>	<u>Females</u>
<u>AGE-ADJUSTED RATES</u>									
<u>YEAR OF DEATH:</u>									
1975	3.7	6.4	1.7	3.1	5.4	1.5	9.7	17.0	4.1
1976	3.7	6.4	1.8	3.1	5.3	1.5	10.2	17.7	4.3
1977	3.7	6.3	1.8	3.1	5.2	1.5	10.1	17.6	4.4
1978	3.8	6.4	1.8	3.2	5.4	1.6	10.0	17.6	4.3
1979	3.8	6.4	1.8	3.1	5.3	1.5	10.3	18.6	4.2
1980	3.8	6.6	1.8	3.2	5.5	1.5	10.6	18.3	4.9
1981	3.8	6.6	1.8	3.2	5.4	1.5	10.4	18.7	4.3
1982	3.8	6.5	1.8	3.2	5.4	1.6	10.5	18.6	4.5
1983	3.9	6.6	1.8	3.2	5.6	1.5	10.2	17.9	4.5
1984	3.9	6.7	1.9	3.3	5.6	1.6	10.6	18.8	4.6
1985	3.9	6.7	1.7	3.3	5.7	1.5	9.9	17.7	4.3
1986	3.9	6.8	1.8	3.3	5.8	1.6	10.2	18.0	4.5
1987	4.0	6.9	1.8	3.4	6.0	1.5	9.7	17.1	4.4
1988	4.0	7.0	1.8	3.5	6.0	1.6	10.0	18.4	3.9
1989	4.1	7.1	1.8	3.6	6.2	1.6	9.7	17.3	4.3
1990	4.1	7.2	1.8	3.6	6.3	1.5	9.3	16.3	4.4
1991	4.1	7.2	1.8	3.6	6.4	1.6	9.2	16.5	4.1
1992	4.2	7.3	1.8	3.7	6.6	1.6	8.6	15.1	4.1
1993	4.2	7.3	1.8	3.7	6.6	1.6	9.0	16.0	4.1
1994	4.3	7.6	1.8	3.9	6.9	1.6	8.6	15.3	3.9
1995	4.3	7.6	1.8	3.9	7.0	1.6	8.0	14.4	3.5
1996	4.3	7.5	1.9	4.0	7.1	1.7	7.8	13.6	3.7
1997	4.3	7.5	1.8	4.0	7.1	1.6	7.3	12.8	3.4
1998	4.4	7.7	1.8	4.1	7.3	1.7	7.1	12.2	3.6
1999	4.3	7.7	1.8	4.2	7.4	1.6	6.5	11.6	3.1
2000	4.4	7.7	1.8	4.3	7.6	1.7	6.4	10.9	3.3
2001	4.4	7.8	1.8	4.3	7.7	1.7	6.4	11.3	3.0
2002	4.4	7.8	1.7	4.3	7.8	1.6	5.9	10.2	2.9
2003	4.4	7.7	1.7	4.3	7.7	1.7	5.7	9.7	2.9
2004	4.4	7.7	1.7	4.3	7.8	1.6	5.4	9.1	2.9
1975-2004	4.1	7.2	1.8	3.7	6.5	1.6	8.5	14.9	3.8

<sup>a</sup> NCHS public use data file for the total US. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

- Statistic not shown. Rate based on less than 16 cases for the time interval.

Table IX-2  
HODGKIN LYMPHOMA

AGE ADJUSTED SEER INCIDENCE<sup>a</sup> RATES BY YEAR, RACE AND SEX

	All Races			Whites			Blacks		
	Total	Males	Females	Total	Males	Females	Total	Males	Females
<u>AGE-ADJUSTED RATES</u>									
YEAR OF DIAGNOSIS:									
1975	3.1	3.9	2.4	3.2	3.9	2.6	2.6	4.4	-
1976	2.8	3.4	2.2	2.9	3.4	2.4	2.6	3.9	-
1977	3.0	3.6	2.5	3.3	3.9	2.7	1.7	1.9	-
1978	2.8	3.4	2.3	3.1	3.7	2.6	1.4	1.9	-
1979	2.9	3.9	2.1	3.1	4.0	2.3	2.9	4.7	-
1980	2.8	3.3	2.3	3.0	3.5	2.5	1.8	2.0	1.5
1981	2.9	3.6	2.3	3.2	3.9	2.6	1.4	1.9	-
1982	2.9	3.5	2.4	3.1	3.7	2.7	2.6	3.2	2.0
1983	3.0	3.7	2.4	3.3	4.0	2.7	2.1	3.8	-
1984	3.1	3.7	2.5	3.4	4.0	2.8	2.1	3.3	-
1985	3.0	3.5	2.5	3.3	3.8	2.8	1.7	2.3	-
1986	2.7	3.3	2.2	2.9	3.5	2.4	2.3	2.8	1.9
1987	3.0	3.7	2.5	3.3	4.0	2.8	2.0	2.8	1.4
1988	3.1	3.6	2.7	3.4	4.0	2.9	1.9	1.9	1.9
1989	3.1	3.7	2.5	3.3	4.0	2.8	2.2	3.2	1.4
1990	3.1	3.5	2.6	3.3	3.8	2.8	3.0	3.2	2.7
1991	3.0	3.5	2.6	3.3	3.8	2.9	2.4	2.6	2.1
1992	2.9	3.1	2.7	3.0	3.2	2.9	2.6	2.8	2.5
1993	2.8	3.3	2.5	3.1	3.5	2.7	2.9	3.7	2.4
1994	2.8	3.1	2.6	3.1	3.5	2.8	2.0	2.3	1.8
1995	2.8	3.2	2.4	3.0	3.5	2.6	2.3	2.8	1.8
1996	2.8	3.3	2.4	3.1	3.6	2.7	2.5	3.1	1.9
1997	2.8	2.9	2.7	3.1	3.2	2.9	2.5	2.4	2.7
1998	2.8	3.1	2.5	3.1	3.4	2.8	2.1	2.2	2.0
1999	2.8	3.1	2.7	3.1	3.2	3.0	2.5	3.0	2.0
2000	2.8	3.3	2.3	3.0	3.5	2.6	2.4	3.0	1.9
2001	2.5	2.9	2.2	2.7	3.1	2.4	2.4	3.1	1.9
2002	2.9	3.2	2.7	3.1	3.4	2.9	2.9	3.2	2.6
2003	2.7	3.1	2.3	2.9	3.4	2.6	2.4	2.7	2.2
2004	2.9	2.9	2.8	3.1	3.1	3.1	2.9	2.9	2.8
1975-2004	2.9	3.4	2.5	3.1	3.6	2.7	2.3	2.9	1.9

<sup>a</sup> SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta). Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).  
- Statistic not shown. Rate based on less than 16 cases for the time interval.

Table IX-3  
HODGKIN LYMPHOMAAGE-ADJUSTED U.S. DEATH<sup>a</sup> RATES BY YEAR, RACE AND SEX

	All Races			Whites			Blacks		
	Total	Males	Females	Total	Males	Females	Total	Males	Females
<u>AGE-ADJUSTED RATES</u>									
YEAR OF DEATH:									
1975	1.3	1.7	1.0	1.4	1.7	1.0	1.1	1.5	0.7
1976	1.2	1.5	0.9	1.2	1.5	0.9	1.0	1.4	0.6
1977	1.1	1.5	0.9	1.2	1.5	0.9	0.9	1.2	0.7
1978	1.1	1.3	0.8	1.1	1.4	0.9	0.8	1.2	0.4
1979	1.0	1.3	0.7	1.0	1.3	0.7	0.8	1.1	0.5
1980	1.0	1.2	0.8	1.0	1.3	0.8	0.7	1.1	0.5
1981	0.9	1.1	0.7	1.0	1.2	0.8	0.7	0.9	0.6
1982	0.9	1.1	0.7	0.9	1.1	0.7	0.7	0.8	0.6
1983	0.8	1.1	0.6	0.9	1.1	0.7	0.6	0.9	0.4
1984	0.9	1.2	0.6	0.9	1.2	0.6	0.8	1.2	0.5
1985	0.8	1.0	0.6	0.8	1.0	0.6	0.7	1.0	0.5
1986	0.8	1.0	0.6	0.8	1.0	0.6	0.7	0.8	0.5
1987	0.7	1.0	0.6	0.8	1.0	0.6	0.7	0.9	0.5
1988	0.7	0.9	0.5	0.7	0.9	0.5	0.6	0.8	0.5
1989	0.7	0.9	0.5	0.7	0.9	0.6	0.7	0.9	0.5
1990	0.7	0.9	0.5	0.7	0.9	0.5	0.6	0.8	0.4
1991	0.7	0.8	0.5	0.7	0.8	0.5	0.7	0.9	0.5
1992	0.6	0.8	0.5	0.7	0.9	0.5	0.6	0.8	0.5
1993	0.6	0.8	0.5	0.6	0.8	0.5	0.6	0.8	0.4
1994	0.6	0.7	0.5	0.6	0.7	0.5	0.6	0.7	0.4
1995	0.5	0.7	0.4	0.6	0.7	0.5	0.5	0.7	0.4
1996	0.5	0.6	0.4	0.5	0.7	0.5	0.5	0.6	0.4
1997	0.5	0.6	0.5	0.5	0.6	0.5	0.5	0.7	0.4
1998	0.5	0.6	0.4	0.5	0.6	0.4	0.5	0.7	0.3
1999	0.5	0.6	0.4	0.5	0.7	0.4	0.4	0.6	0.3
2000	0.5	0.5	0.4	0.5	0.6	0.4	0.4	0.4	0.3
2001	0.5	0.6	0.4	0.5	0.6	0.4	0.4	0.5	0.4
2002	0.5	0.6	0.4	0.5	0.6	0.4	0.4	0.6	0.3
2003	0.5	0.5	0.4	0.5	0.6	0.4	0.4	0.5	0.3
2004	0.4	0.5	0.3	0.4	0.6	0.3	0.4	0.5	0.3
1975-2004	0.7	0.9	0.5	0.7	0.9	0.6	0.6	0.8	0.4

<sup>a</sup> NCHS public use data file for the total US. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

- Statistic not shown. Rate based on less than 16 cases for the time interval.

Table X-4  
KAPOSI SARCOMA

AGE ADJUSTED SEER INCIDENCE<sup>a</sup> RATES BY YEAR, RACE AND SEX

	<u>All Races</u>			<u>Whites</u>			<u>Blacks</u>		
	<u>Total</u>	<u>Males</u>	<u>Females</u>	<u>Total</u>	<u>Males</u>	<u>Females</u>	<u>Total</u>	<u>Males</u>	<u>Females</u>
<u>AGE-ADJUSTED RATES</u>									
<u>YEAR OF DIAGNOSIS:</u>									
1975	0.2	0.6	-	0.2	0.6	-	-	-	-
1976	0.2	0.5	-	0.2	0.5	-	-	-	-
1977	0.3	0.5	-	0.3	0.6	-	-	-	-
1978	0.2	0.3	0.2	0.2	0.3	-	-	-	-
1979	0.2	0.3	-	0.2	0.3	-	-	-	-
1980	0.3	0.6	-	0.3	0.6	-	-	-	-
1981	0.3	0.5	-	0.3	0.6	-	-	-	-
1982	0.7	1.2	0.2	0.7	1.3	0.2	-	-	-
1983	1.1	2.2	-	1.2	2.4	-	-	-	-
1984	1.8	3.6	-	2.0	3.9	-	1.3	2.9	-
1985	2.5	4.9	0.2	2.8	5.5	0.2	1.1	2.2	-
1986	3.3	6.5	-	3.7	7.3	-	1.6	3.6	-
1987	3.8	7.6	-	4.3	8.5	-	2.1	4.3	-
1988	4.4	8.9	0.1	4.8	9.6	0.1	3.3	7.2	-
1989	4.8	9.5	0.1	5.1	10.2	-	4.6	9.7	-
1990	4.6	9.2	0.2	5.0	10.0	0.2	3.6	7.5	-
1991	4.7	9.4	0.2	5.0	10.0	0.2	4.6	9.5	-
1992	4.4	8.7	0.2	4.6	9.0	0.2	4.9	10.4	-
1993	3.9	7.6	0.2	4.0	8.0	0.1	4.3	8.6	-
1994	3.8	7.5	0.2	3.9	7.6	0.1	4.1	8.4	-
1995	3.3	6.4	0.2	3.3	6.4	0.2	4.9	10.2	-
1996	2.2	4.4	0.2	2.0	4.0	-	4.6	9.1	-
1997	1.1	2.1	0.1	0.9	1.8	-	2.6	5.0	-
1998	0.8	1.6	-	0.7	1.3	-	1.8	3.8	-
1999	0.8	1.5	0.1	0.7	1.2	-	1.9	3.6	-
2000	0.8	1.5	-	0.7	1.3	-	2.0	4.0	-
2001	0.8	1.5	0.2	0.7	1.2	0.2	1.9	3.7	-
2002	0.7	1.3	0.1	0.5	1.0	-	1.5	3.0	-
2003	0.6	1.2	-	0.6	1.0	-	1.4	3.0	-
2004	0.7	1.2	-	0.5	1.0	-	1.5	3.1	-
1975-2004	2.0	3.9	0.1	2.1	4.0	0.1	2.3	4.7	0.2

<sup>a</sup> SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta). Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

- Statistic not shown. Rate based on less than 16 cases for the time interval.

Table XI-2  
 KIDNEY AND RENAL PELVIS CANCER (Invasive)

AGE ADJUSTED SEER INCIDENCE<sup>a</sup> RATES BY YEAR, RACE AND SEX

	All Races			Whites			Blacks		
	Total	Males	Females	Total	Males	Females	Total	Males	Females
<u>AGE-ADJUSTED RATES</u>									
YEAR OF DIAGNOSIS:									
1975	7.1	10.3	4.5	7.3	10.8	4.6	6.2	8.2	4.3
1976	8.0	11.2	5.5	8.1	11.5	5.6	7.7	11.6	4.7
1977	8.0	11.4	5.5	8.1	11.7	5.5	8.5	11.1	6.4
1978	7.8	11.8	4.8	8.0	12.1	4.9	8.0	11.0	5.7
1979	7.6	11.1	5.0	7.8	11.6	5.1	7.3	8.2	6.3
1980	8.1	11.7	5.4	8.4	12.2	5.6	6.0	8.0	4.4
1981	8.5	12.8	5.3	8.6	13.0	5.3	10.0	15.8	5.8
1982	8.3	11.8	5.7	8.5	12.0	5.8	7.6	11.2	4.9
1983	8.9	13.2	5.7	9.2	13.8	5.8	8.8	13.5	5.5
1984	9.2	13.1	6.2	9.5	13.6	6.3	9.0	12.0	6.9
1985	8.9	13.1	5.8	9.2	13.6	5.9	8.6	11.6	6.2
1986	9.7	13.8	6.6	9.8	14.1	6.7	10.1	15.0	6.6
1987	9.9	14.1	6.7	10.1	14.4	6.9	11.1	15.6	7.5
1988	9.9	14.0	7.0	10.1	14.3	7.0	11.6	15.4	8.8
1989	10.3	14.5	7.1	10.6	15.0	7.2	10.8	15.1	7.8
1990	10.4	14.7	7.1	10.7	15.0	7.4	10.5	15.9	6.6
1991	10.6	15.1	7.2	10.8	15.2	7.3	12.2	18.3	7.7
1992	10.8	15.3	7.4	11.1	15.6	7.7	11.0	16.4	7.2
1993	10.7	14.7	7.6	10.8	14.7	7.7	12.8	17.3	9.4
1994	11.3	15.6	7.9	11.5	16.0	8.0	12.7	18.0	8.7
1995	11.1	15.5	7.6	11.1	15.4	7.7	14.2	21.5	9.2
1996	11.3	15.8	7.9	11.5	16.2	7.8	13.7	17.2	11.1
1997	10.9	15.1	7.6	11.0	15.0	7.7	13.5	19.4	9.2
1998	11.8	16.4	8.2	12.1	16.7	8.4	12.6	17.2	9.1
1999	11.4	15.9	7.8	11.6	16.2	7.9	13.7	18.7	10.0
2000	12.4	17.3	8.5	12.7	17.9	8.5	14.7	19.1	11.8
2001	12.5	17.4	8.5	12.7	17.6	8.7	15.5	22.9	10.2
2002	12.7	17.6	8.7	13.0	17.8	8.9	15.1	22.0	10.2
2003	13.2	17.8	9.5	13.5	18.0	9.8	14.2	21.2	9.1
2004	13.1	17.8	9.2	13.4	18.2	9.4	15.1	21.6	10.3
1975-2004	10.4	14.6	7.1	10.5	14.9	7.2	11.7	16.5	8.1

<sup>a</sup> SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta). Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).  
 - Statistic not shown. Rate based on less than 16 cases for the time interval.

Table XI-3  
 KIDNEY AND RENAL PELVIS CANCER (Invasive)

AGE-ADJUSTED U.S. DEATH<sup>a</sup> RATES BY YEAR, RACE AND SEX

	All Races			Whites			Blacks		
	Total	Males	Females	Total	Males	Females	Total	Males	Females
<u>AGE-ADJUSTED RATES</u>									
YEAR OF DEATH:									
1975	3.6	5.2	2.4	3.7	5.3	2.5	2.8	4.1	1.8
1976	3.6	5.2	2.4	3.7	5.4	2.5	2.9	4.2	1.9
1977	3.7	5.4	2.4	3.8	5.5	2.5	2.9	4.0	2.0
1978	3.7	5.4	2.4	3.8	5.5	2.5	3.2	4.8	1.9
1979	3.6	5.3	2.4	3.7	5.5	2.4	2.9	4.2	2.0
1980	3.7	5.5	2.4	3.8	5.6	2.4	3.0	4.2	2.0
1981	3.7	5.4	2.5	3.8	5.5	2.6	3.2	4.6	2.1
1982	3.9	5.7	2.5	3.9	5.8	2.6	3.4	4.8	2.4
1983	3.8	5.6	2.6	3.9	5.7	2.7	3.3	4.9	2.1
1984	3.9	5.8	2.6	4.0	5.9	2.6	3.5	4.9	2.4
1985	4.0	5.8	2.7	4.0	5.8	2.7	3.8	6.0	2.4
1986	4.0	5.8	2.8	4.1	5.9	2.8	3.7	5.3	2.5
1987	4.1	6.0	2.8	4.2	6.1	2.8	3.7	5.5	2.4
1988	4.0	5.9	2.7	4.1	5.9	2.8	3.8	5.8	2.4
1989	4.2	6.1	2.8	4.2	6.2	2.8	4.0	6.1	2.6
1990	4.2	6.2	2.8	4.2	6.2	2.8	4.2	6.4	2.8
1991	4.3	6.2	2.9	4.3	6.2	3.0	4.3	6.4	3.0
1992	4.3	6.2	2.9	4.4	6.3	2.9	4.0	5.9	2.6
1993	4.2	6.1	2.8	4.2	6.2	2.8	4.1	6.1	2.7
1994	4.3	6.2	2.9	4.3	6.3	2.9	4.2	6.4	2.8
1995	4.3	6.2	3.0	4.4	6.3	3.0	4.4	6.3	3.1
1996	4.3	6.2	2.8	4.3	6.3	2.9	4.1	5.9	2.9
1997	4.3	6.2	2.9	4.3	6.2	2.9	4.3	6.5	2.8
1998	4.3	6.2	2.8	4.3	6.3	2.9	4.0	6.3	2.5
1999	4.1	5.9	2.7	4.1	6.0	2.7	4.2	6.1	2.8
2000	4.2	6.1	2.8	4.3	6.2	2.8	4.1	6.3	2.7
2001	4.3	6.2	2.8	4.3	6.3	2.8	4.3	6.4	2.8
2002	4.2	6.1	2.8	4.3	6.3	2.8	4.1	5.8	2.8
2003	4.2	6.1	2.7	4.3	6.2	2.8	4.1	6.1	2.7
2004	4.1	5.9	2.7	4.2	6.1	2.8	4.0	5.8	2.7
1975-2004	4.1	5.9	2.7	4.1	6.0	2.8	3.8	5.6	2.6

<sup>a</sup> NCHS public use data file for the total US. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

- Statistic not shown. Rate based on less than 16 cases for the time interval.



Table XII-2  
LARYNX CANCER (Invasive)

AGE ADJUSTED SEER INCIDENCE<sup>a</sup> RATES BY YEAR, RACE AND SEX

	<u>All Races</u>			<u>Whites</u>			<u>Blacks</u>		
	<u>Total</u>	<u>Males</u>	<u>Females</u>	<u>Total</u>	<u>Males</u>	<u>Females</u>	<u>Total</u>	<u>Males</u>	<u>Females</u>
<u>AGE-ADJUSTED RATES</u>									
<u>YEAR OF DIAGNOSIS:</u>									
1975	5.0	9.5	1.5	5.0	9.4	1.5	7.1	13.4	-
1976	5.2	10.0	1.4	5.2	9.9	1.4	7.3	14.6	-
1977	5.0	9.7	1.4	5.0	9.7	1.3	7.5	13.9	2.4
1978	5.3	9.8	1.8	5.3	9.8	1.9	6.9	13.2	2.2
1979	5.4	10.1	1.7	5.4	10.2	1.6	7.5	13.2	3.1
1980	5.2	10.0	1.5	5.2	9.9	1.5	7.3	14.5	-
1981	5.4	9.8	1.9	5.3	9.6	2.0	8.1	16.3	-
1982	5.3	9.8	1.8	5.3	9.8	1.8	7.9	14.5	2.9
1983	5.3	9.9	1.7	5.4	10.2	1.7	6.8	12.2	2.6
1984	5.1	9.6	1.6	5.0	9.6	1.5	8.5	14.9	3.4
1985	5.4	9.9	1.9	5.4	10.0	1.9	8.4	15.4	3.3
1986	5.0	9.4	1.6	4.9	9.2	1.7	7.7	15.4	2.1
1987	5.2	9.4	1.9	5.1	9.3	1.9	8.3	15.2	3.1
1988	5.2	9.6	1.9	5.2	9.6	1.9	8.1	14.8	3.2
1989	5.0	8.9	2.0	5.0	8.9	1.9	7.9	14.3	3.3
1990	5.1	9.2	1.9	5.0	9.0	1.9	9.5	17.8	3.6
1991	4.7	8.5	1.8	4.8	8.5	1.9	6.6	12.3	2.3
1992	5.0	9.1	1.8	4.9	8.9	1.7	8.6	15.7	3.4
1993	4.4	8.1	1.5	4.4	8.0	1.5	6.9	14.1	1.7
1994	4.7	8.3	1.8	4.5	8.1	1.7	8.9	16.0	3.7
1995	4.4	7.9	1.6	4.4	7.8	1.6	6.8	13.1	2.2
1996	4.3	7.6	1.6	4.3	7.6	1.6	6.5	11.5	2.8
1997	4.2	7.5	1.6	4.1	7.3	1.6	7.2	13.3	2.7
1998	3.8	6.8	1.6	3.7	6.5	1.5	6.5	11.8	2.7
1999	4.2	7.1	1.8	4.1	7.0	1.8	7.1	12.2	3.6
2000	4.0	7.3	1.4	4.1	7.2	1.5	6.0	11.4	2.2
2001	3.6	6.5	1.3	3.6	6.3	1.3	6.1	12.0	1.8
2002	3.5	6.3	1.3	3.5	6.1	1.3	6.3	11.2	2.7
2003	3.3	5.9	1.2	3.3	5.8	1.3	5.0	9.5	1.8
2004	3.4	6.0	1.3	3.4	5.9	1.3	5.8	10.9	2.3
1975-2004	4.6	8.4	1.6	4.6	8.4	1.6	7.2	13.4	2.6

<sup>a</sup> SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta). Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).  
 - Statistic not shown. Rate based on less than 16 cases for the time interval.

Table XII-3  
 LARYNX CANCER (Invasive)

AGE-ADJUSTED U.S. DEATH<sup>a</sup> RATES BY YEAR, RACE AND SEX

	All Races			Whites			Blacks		
	Total	Males	Females	Total	Males	Females	Total	Males	Females
<u>AGE-ADJUSTED RATES</u>									
YEAR OF DEATH:									
1975	1.7	3.3	0.5	1.6	3.2	0.4	2.6	5.0	0.8
1976	1.7	3.4	0.5	1.6	3.3	0.5	2.6	5.1	0.7
1977	1.7	3.4	0.5	1.6	3.3	0.4	2.7	5.5	0.7
1978	1.7	3.4	0.5	1.6	3.2	0.5	2.9	5.9	0.7
1979	1.7	3.3	0.5	1.6	3.2	0.4	2.6	5.2	0.6
1980	1.6	3.2	0.5	1.5	3.0	0.4	2.9	5.7	0.9
1981	1.7	3.3	0.5	1.5	3.1	0.5	3.0	5.6	1.1
1982	1.7	3.2	0.5	1.6	3.0	0.5	2.8	5.5	0.8
1983	1.6	3.2	0.5	1.5	3.0	0.5	2.7	5.3	0.9
1984	1.6	3.2	0.5	1.5	2.9	0.5	3.0	6.2	0.7
1985	1.6	3.0	0.5	1.4	2.8	0.5	3.0	5.8	0.9
1986	1.6	3.1	0.5	1.5	2.8	0.5	3.1	6.1	1.0
1987	1.6	3.1	0.5	1.5	2.9	0.5	2.9	5.9	0.9
1988	1.6	3.0	0.5	1.4	2.7	0.5	3.2	6.7	0.8
1989	1.6	3.0	0.6	1.4	2.7	0.6	3.2	6.6	0.8
1990	1.6	3.0	0.5	1.4	2.7	0.5	3.0	5.8	1.1
1991	1.7	3.1	0.6	1.5	2.8	0.6	3.3	6.4	1.2
1992	1.6	3.0	0.6	1.5	2.7	0.6	3.3	6.4	1.1
1993	1.6	3.0	0.6	1.5	2.7	0.5	3.2	6.4	1.1
1994	1.6	2.9	0.6	1.4	2.6	0.5	3.2	6.4	1.0
1995	1.5	2.8	0.5	1.4	2.6	0.5	3.0	6.0	1.0
1996	1.5	2.8	0.6	1.4	2.5	0.5	3.2	6.3	1.0
1997	1.4	2.7	0.5	1.3	2.4	0.5	3.0	6.1	0.9
1998	1.4	2.6	0.6	1.3	2.3	0.5	2.8	5.6	0.9
1999	1.4	2.5	0.5	1.3	2.3	0.5	2.6	5.1	0.9
2000	1.4	2.5	0.5	1.3	2.3	0.5	2.8	5.5	1.0
2001	1.3	2.5	0.5	1.2	2.3	0.5	2.6	5.0	0.9
2002	1.3	2.3	0.5	1.2	2.2	0.5	2.5	4.8	0.8
2003	1.3	2.4	0.5	1.2	2.2	0.4	2.5	5.1	0.7
2004	1.2	2.2	0.4	1.2	2.1	0.4	2.2	4.6	0.7
1975-2004	1.5	2.9	0.5	1.4	2.7	0.5	2.8	5.7	0.9

<sup>a</sup> NCHS public use data file for the total US. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

- Statistic not shown. Rate based on less than 16 cases for the time interval.

Table XIII-2

LEUKEMIAAGE ADJUSTED SEER INCIDENCE<sup>a</sup> RATES BY YEAR, RACE AND SEX

	<u>All Races</u>			<u>Whites</u>			<u>Blacks</u>		
	<u>Total</u>	<u>Males</u>	<u>Females</u>	<u>Total</u>	<u>Males</u>	<u>Females</u>	<u>Total</u>	<u>Males</u>	<u>Females</u>
<u>AGE-ADJUSTED RATES</u>									
<u>YEAR OF DIAGNOSIS:</u>									
1975	12.8	16.9	9.9	13.1	17.5	10.1	11.5	16.2	8.5
1976	13.5	18.5	10.2	13.8	19.1	10.3	11.3	14.2	9.4
1977	12.7	16.3	10.2	13.2	17.0	10.5	10.2	13.6	7.7
1978	12.9	17.2	9.8	13.5	18.3	10.2	9.2	11.4	7.5
1979	12.6	16.8	9.7	13.0	17.5	10.0	10.6	13.6	8.6
1980	12.9	17.4	9.9	13.3	17.9	10.1	12.2	16.0	9.5
1981	12.4	16.5	9.6	12.7	17.1	9.8	12.2	14.5	10.3
1982	13.2	18.3	9.7	13.5	18.8	10.0	10.7	15.2	7.9
1983	13.1	17.8	9.9	13.5	18.4	10.1	12.5	16.0	10.1
1984	13.0	17.1	10.1	13.4	17.9	10.3	10.7	11.9	9.6
1985	13.3	17.7	10.4	13.7	18.2	10.6	12.2	16.8	9.3
1986	12.8	17.5	9.5	13.4	18.3	9.9	10.2	13.1	8.1
1987	13.3	17.7	10.3	13.8	18.4	10.6	12.6	17.6	9.3
1988	12.9	17.5	9.7	13.4	18.2	9.9	11.6	14.0	9.4
1989	13.4	17.6	10.4	13.9	18.2	10.9	11.3	15.9	8.0
1990	12.8	17.2	9.7	13.4	18.1	10.1	11.3	14.8	8.8
1991	13.0	16.8	10.2	13.6	17.7	10.5	10.9	12.0	10.2
1992	13.0	17.4	9.7	13.5	18.5	9.9	10.7	13.9	8.4
1993	13.0	16.9	10.0	13.5	17.8	10.3	11.6	15.1	9.1
1994	12.9	16.8	10.0	13.5	17.7	10.5	9.8	12.6	7.8
1995	13.4	17.6	10.3	14.2	18.6	11.0	9.7	13.4	7.2
1996	12.9	16.7	10.2	13.5	17.4	10.6	10.7	13.9	8.7
1997	12.9	17.3	9.8	13.7	18.3	10.4	10.3	13.8	8.1
1998	12.9	16.8	10.1	13.7	17.7	10.7	9.7	14.1	7.0
1999	12.5	16.7	9.4	13.1	17.5	9.8	10.0	13.6	7.8
2000	13.1	16.9	10.2	13.7	17.8	10.6	11.3	13.3	9.8
2001	13.4	17.5	10.4	14.3	18.4	11.1	11.1	13.5	9.4
2002	12.5	16.5	9.5	13.5	17.8	10.2	8.0	10.2	6.4
2003	12.4	16.2	9.4	13.0	17.1	9.8	10.0	12.8	7.9
2004	12.0	15.4	9.5	12.5	15.9	9.8	10.4	13.8	8.1
1975-2004	12.9	17.0	9.9	13.4	17.8	10.3	10.7	13.9	8.5

<sup>a</sup> SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta). Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

- Statistic not shown. Rate based on less than 16 cases for the time interval.

Table XIII-3  
ACUTE LYMPHOCYTIC LEUKEMIA

AGE ADJUSTED SEER INCIDENCE<sup>a</sup> RATES BY YEAR, RACE AND SEX

	<u>All Races</u>			<u>Whites</u>			<u>Blacks</u>		
	<u>Total</u>	<u>Males</u>	<u>Females</u>	<u>Total</u>	<u>Males</u>	<u>Females</u>	<u>Total</u>	<u>Males</u>	<u>Females</u>
<u>AGE-ADJUSTED RATES</u>									
<u>YEAR OF DIAGNOSIS:</u>									
1975	0.9	1.1	0.8	1.0	1.2	0.8	-	-	-
1976	1.1	1.3	0.9	1.1	1.3	0.9	0.8	-	-
1977	1.1	1.3	0.9	1.2	1.4	1.0	-	-	-
1978	1.0	1.2	0.8	1.1	1.3	0.9	-	-	-
1979	1.0	1.2	0.9	1.1	1.2	1.0	-	-	-
1980	1.1	1.4	0.9	1.2	1.5	0.9	0.7	-	-
1981	1.2	1.3	1.2	1.3	1.4	1.2	1.0	-	-
1982	1.2	1.5	1.0	1.3	1.6	1.0	-	-	-
1983	1.3	1.5	1.1	1.4	1.7	1.2	0.8	-	-
1984	1.3	1.5	1.2	1.3	1.5	1.2	0.9	-	-
1985	1.3	1.5	1.1	1.4	1.6	1.1	-	-	-
1986	1.3	1.7	1.0	1.4	1.8	1.0	0.8	-	-
1987	1.3	1.5	1.2	1.3	1.5	1.2	0.9	-	1.1
1988	1.4	1.8	1.1	1.4	1.8	1.1	1.0	-	-
1989	1.6	1.8	1.4	1.7	1.9	1.5	1.0	-	-
1990	1.4	1.7	1.2	1.5	1.8	1.3	0.6	-	-
1991	1.4	1.7	1.1	1.5	1.9	1.1	0.9	-	-
1992	1.4	1.6	1.1	1.4	1.7	1.1	0.9	-	-
1993	1.3	1.4	1.1	1.4	1.6	1.1	0.8	-	-
1994	1.2	1.5	0.9	1.3	1.6	1.0	0.7	-	-
1995	1.5	1.8	1.2	1.6	1.9	1.3	1.0	1.3	-
1996	1.4	1.6	1.2	1.4	1.7	1.2	0.8	-	-
1997	1.3	1.5	1.2	1.5	1.6	1.3	0.8	0.9	-
1998	1.5	1.6	1.3	1.6	1.7	1.4	0.9	1.2	-
1999	1.3	1.5	1.1	1.4	1.6	1.2	0.7	-	-
2000	1.4	1.5	1.3	1.5	1.6	1.5	0.9	-	-
2001	1.4	1.6	1.3	1.6	1.8	1.3	0.7	-	-
2002	1.5	1.8	1.2	1.6	2.0	1.3	0.6	-	-
2003	1.4	1.6	1.2	1.4	1.7	1.3	0.8	1.0	-
2004	1.5	1.6	1.3	1.6	1.8	1.3	1.0	-	1.3
1975-2004	1.3	1.5	1.1	1.4	1.6	1.2	0.8	0.8	0.7

<sup>a</sup> SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta). Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).  
 - Statistic not shown. Rate based on less than 16 cases for the time interval.

Table XIII-4  
 CHRONIC LYMPHOCYTIC LEUKEMIA

AGE ADJUSTED SEER INCIDENCE<sup>a</sup> RATES BY YEAR, RACE AND SEX

	All Races			Whites			Blacks		
	Total	Males	Females	Total	Males	Females	Total	Males	Females
<u>AGE-ADJUSTED RATES</u>									
YEAR OF DIAGNOSIS:									
1975	4.3	5.9	3.3	4.5	6.1	3.4	4.8	6.9	3.2
1976	4.6	6.8	3.1	4.8	7.2	3.2	4.3	6.6	3.0
1977	4.2	5.7	3.1	4.4	6.0	3.3	3.8	5.8	2.5
1978	4.5	5.9	3.4	4.7	6.4	3.6	3.3	3.6	3.1
1979	4.0	5.7	2.9	4.3	6.1	3.1	3.1	4.2	-
1980	4.5	6.3	3.3	4.7	6.5	3.5	4.5	7.1	2.9
1981	4.3	6.3	2.9	4.5	6.6	3.0	3.8	5.4	2.4
1982	4.3	6.2	3.0	4.5	6.4	3.2	3.8	6.7	-
1983	4.5	6.4	3.1	4.7	6.8	3.2	4.2	6.1	3.0
1984	4.4	6.3	3.1	4.7	6.8	3.3	3.5	4.5	2.7
1985	4.4	6.1	3.3	4.7	6.5	3.5	4.1	6.1	2.9
1986	4.5	6.3	3.2	4.8	6.8	3.5	2.8	4.2	-
1987	4.6	6.7	3.1	4.9	7.2	3.3	4.0	6.1	2.5
1988	4.5	6.4	3.1	4.7	6.8	3.2	4.4	5.3	3.5
1989	4.6	6.4	3.4	4.9	6.8	3.6	3.9	5.3	2.9
1990	4.5	6.2	3.3	4.8	6.7	3.5	4.5	6.1	3.4
1991	4.5	6.1	3.4	4.9	6.7	3.6	3.4	3.6	3.2
1992	4.5	6.6	3.1	4.9	7.3	3.4	3.2	5.0	2.1
1993	4.4	6.0	3.1	4.6	6.4	3.3	4.4	6.6	2.9
1994	4.4	5.9	3.3	4.8	6.4	3.6	3.2	4.1	2.3
1995	4.4	6.3	3.0	4.8	6.8	3.3	3.3	5.2	2.1
1996	4.5	6.1	3.4	4.9	6.5	3.7	3.5	5.4	2.2
1997	4.3	5.8	3.1	4.6	6.3	3.4	3.3	4.9	2.4
1998	4.0	5.6	2.8	4.3	6.1	3.1	2.7	3.4	2.2
1999	3.9	5.7	2.5	4.2	6.1	2.8	3.4	5.5	2.2
2000	4.0	5.7	2.8	4.5	6.3	3.0	3.0	3.8	2.4
2001	4.5	6.3	3.1	4.9	6.9	3.4	3.3	4.1	2.7
2002	4.2	5.9	2.9	4.7	6.6	3.2	2.3	3.1	1.7
2003	4.4	6.2	3.1	4.8	6.7	3.3	3.3	4.6	2.3
2004	4.0	5.5	2.9	4.3	5.8	3.2	3.0	4.8	1.8
1975-2004	4.3	6.1	3.1	4.7	6.5	3.3	3.5	5.0	2.5

<sup>a</sup> SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta). Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).  
 - Statistic not shown. Rate based on less than 16 cases for the time interval.

Table XIII-5  
ACUTE MYELOID LEUKEMIA

AGE ADJUSTED SEER INCIDENCE<sup>a</sup> RATES BY YEAR, RACE AND SEX

	All Races			Whites			Blacks		
	Total	Males	Females	Total	Males	Females	Total	Males	Females
<u>AGE-ADJUSTED RATES</u>									
YEAR OF DIAGNOSIS:									
1975	3.4	4.2	3.0	3.5	4.3	3.0	2.5	3.4	2.0
1976	3.5	4.4	2.9	3.5	4.5	2.9	2.6	-	2.6
1977	3.5	4.2	3.1	3.6	4.4	3.1	1.9	-	-
1978	3.4	4.4	2.7	3.5	4.6	2.8	2.7	3.5	2.1
1979	3.3	4.1	2.8	3.4	4.2	2.9	3.4	3.5	3.4
1980	3.1	4.1	2.4	3.2	4.1	2.5	2.9	3.3	2.5
1981	3.2	4.0	2.6	3.2	4.1	2.7	3.1	3.2	2.9
1982	3.4	4.8	2.6	3.5	4.9	2.6	2.5	3.2	2.1
1983	3.2	4.2	2.5	3.2	4.3	2.4	2.9	3.0	2.9
1984	3.3	4.1	2.7	3.4	4.2	2.8	2.4	2.1	2.4
1985	3.3	4.1	2.9	3.3	4.0	2.8	3.7	4.9	3.0
1986	3.0	3.9	2.4	2.9	3.9	2.4	3.2	4.0	2.7
1987	3.0	4.0	2.5	3.0	4.0	2.5	3.0	4.7	2.0
1988	3.0	3.8	2.5	3.1	3.9	2.5	2.8	3.1	2.5
1989	3.2	4.0	2.6	3.3	4.0	2.8	2.2	3.8	-
1990	3.0	3.9	2.4	3.1	4.0	2.5	2.5	3.0	2.1
1991	3.3	3.9	2.9	3.4	3.9	3.0	3.3	4.4	2.6
1992	3.2	3.9	2.8	3.3	4.0	2.7	3.0	3.0	3.0
1993	3.5	4.3	2.9	3.5	4.5	2.9	2.5	2.6	2.5
1994	3.5	4.1	3.0	3.5	4.2	3.0	2.9	3.2	2.7
1995	3.7	4.2	3.3	3.8	4.3	3.4	2.5	2.6	2.6
1996	3.5	4.3	2.9	3.5	4.3	2.9	3.2	4.1	2.7
1997	3.5	4.6	2.8	3.6	4.8	2.9	3.0	3.8	2.6
1998	3.9	4.9	3.2	4.0	5.1	3.3	3.1	4.6	2.1
1999	3.7	4.5	3.2	3.8	4.6	3.3	3.0	3.4	2.8
2000	4.0	4.9	3.4	4.0	4.9	3.4	4.4	4.9	4.0
2001	4.0	4.9	3.4	4.2	5.0	3.6	3.5	4.1	3.1
2002	3.6	4.5	3.0	3.7	4.7	3.1	2.9	3.2	2.7
2003	3.5	4.4	2.9	3.6	4.6	3.0	3.1	3.4	2.7
2004	3.3	4.0	2.8	3.4	4.0	2.9	3.2	4.0	2.8
1975-2004	3.4	4.3	2.8	3.5	4.3	2.9	3.0	3.6	2.6

<sup>a</sup> SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta). Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).  
- Statistic not shown. Rate based on less than 16 cases for the time interval.

Table XIII-6  
CHRONIC MYELOID LEUKEMIA

AGE ADJUSTED SEER INCIDENCE<sup>a</sup> RATES BY YEAR, RACE AND SEX

	<u>All Races</u>			<u>Whites</u>			<u>Blacks</u>		
	<u>Total</u>	<u>Males</u>	<u>Females</u>	<u>Total</u>	<u>Males</u>	<u>Females</u>	<u>Total</u>	<u>Males</u>	<u>Females</u>
<u>AGE-ADJUSTED RATES</u>									
<u>YEAR OF DIAGNOSIS:</u>									
1975	2.0	2.9	1.3	2.0	3.0	1.3	1.9	-	-
1976	1.9	2.5	1.6	1.9	2.4	1.5	1.9	2.6	-
1977	1.7	2.1	1.4	1.7	2.1	1.4	1.8	-	-
1978	1.7	2.3	1.3	1.7	2.3	1.4	1.4	-	-
1979	1.9	2.5	1.4	1.9	2.5	1.5	1.9	2.8	-
1980	1.7	2.1	1.5	1.7	2.1	1.5	1.5	-	-
1981	1.6	2.0	1.4	1.6	1.9	1.4	2.4	3.0	1.8
1982	1.7	2.2	1.4	1.7	2.2	1.3	1.7	-	1.8
1983	1.7	2.2	1.3	1.6	2.1	1.3	2.5	3.4	1.8
1984	1.7	2.4	1.2	1.6	2.4	1.1	2.6	2.6	2.6
1985	2.0	2.8	1.4	2.0	2.8	1.5	2.1	2.9	-
1986	1.7	2.2	1.3	1.7	2.3	1.3	1.5	-	-
1987	1.9	2.3	1.6	1.9	2.4	1.6	2.1	2.9	-
1988	1.9	2.5	1.5	1.9	2.6	1.5	2.1	2.9	-
1989	1.7	2.2	1.3	1.7	2.2	1.3	2.2	3.1	-
1990	1.8	2.5	1.2	1.7	2.5	1.2	2.6	3.5	2.0
1991	1.9	2.4	1.5	1.9	2.6	1.4	1.6	-	1.9
1992	1.8	2.2	1.5	1.8	2.2	1.5	1.5	1.7	-
1993	1.9	2.5	1.5	2.0	2.5	1.6	2.3	3.6	-
1994	1.9	2.5	1.4	1.9	2.5	1.5	1.7	2.5	-
1995	1.9	2.5	1.4	2.0	2.7	1.4	1.5	2.0	-
1996	1.8	2.4	1.3	1.8	2.5	1.3	2.0	2.4	1.8
1997	1.9	2.7	1.3	1.9	2.7	1.3	1.7	2.5	-
1998	1.6	1.9	1.4	1.7	1.9	1.5	1.5	2.3	-
1999	1.8	2.5	1.3	1.8	2.4	1.3	1.7	2.2	1.3
2000	1.8	2.3	1.3	1.8	2.3	1.4	1.5	1.7	1.3
2001	1.7	2.2	1.3	1.8	2.2	1.4	1.8	2.6	1.2
2002	1.5	1.9	1.2	1.6	2.0	1.3	0.9	-	-
2003	1.5	2.0	1.1	1.5	2.1	1.1	1.2	1.8	-
2004	1.6	2.2	1.1	1.6	2.3	1.1	1.3	1.9	-
1975-2004	1.8	2.3	1.3	1.8	2.3	1.4	1.8	2.3	1.4

<sup>a</sup> SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta). Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).  
 - Statistic not shown. Rate based on less than 16 cases for the time interval.

Table XIII-7

LEUKEMIAAGE-ADJUSTED U.S. DEATH<sup>a</sup> RATES BY YEAR, RACE AND SEX

	<u>All Races</u>			<u>Whites</u>			<u>Blacks</u>		
	<u>Total</u>	<u>Males</u>	<u>Females</u>	<u>Total</u>	<u>Males</u>	<u>Females</u>	<u>Total</u>	<u>Males</u>	<u>Females</u>
<u>AGE-ADJUSTED RATES</u>									
<u>YEAR OF DEATH:</u>									
1975	8.1	11.0	6.2	8.3	11.3	6.3	6.6	8.6	5.1
1976	8.1	10.9	6.2	8.3	11.2	6.3	6.8	8.5	5.5
1977	8.1	10.8	6.3	8.3	11.1	6.3	6.8	8.8	5.5
1978	8.0	10.8	6.1	8.2	11.1	6.3	6.7	9.2	4.9
1979	8.3	11.1	6.4	8.4	11.4	6.5	7.0	8.9	5.7
1980	8.4	11.2	6.5	8.5	11.5	6.6	7.0	9.1	5.7
1981	8.1	10.9	6.2	8.2	11.2	6.3	7.1	8.9	5.9
1982	8.2	11.0	6.3	8.3	11.3	6.4	7.1	9.4	5.7
1983	8.1	10.9	6.3	8.2	11.2	6.3	7.2	9.2	5.9
1984	8.1	10.7	6.4	8.2	10.9	6.5	7.1	9.2	5.7
1985	8.1	10.8	6.3	8.2	11.0	6.4	7.1	9.6	5.4
1986	8.0	10.7	6.2	8.1	11.0	6.2	7.3	9.3	5.9
1987	7.8	10.3	6.1	7.9	10.5	6.2	7.2	9.4	5.9
1988	7.8	10.6	6.0	8.0	10.8	6.1	7.2	9.8	5.6
1989	8.0	10.6	6.2	8.1	10.8	6.3	7.4	10.3	5.5
1990	8.0	10.7	6.2	8.1	10.9	6.3	7.4	10.0	5.8
1991	8.0	10.5	6.3	8.1	10.6	6.4	7.4	10.0	5.8
1992	8.0	10.7	6.1	8.1	10.9	6.2	7.2	9.4	5.7
1993	7.9	10.7	6.1	8.0	10.8	6.2	7.4	9.8	5.8
1994	7.9	10.6	6.0	8.0	10.8	6.1	7.1	9.6	5.5
1995	7.9	10.8	6.0	8.0	11.0	6.0	7.2	9.8	5.5
1996	7.8	10.5	6.1	8.0	10.7	6.1	7.5	10.1	5.9
1997	7.7	10.4	5.9	7.9	10.7	6.0	6.8	9.2	5.3
1998	7.6	10.1	5.8	7.7	10.3	5.9	7.0	9.4	5.6
1999	7.7	10.2	6.0	7.9	10.5	6.1	6.7	8.8	5.4
2000	7.7	10.3	5.9	7.9	10.5	6.0	6.7	9.1	5.3
2001	7.6	10.1	5.9	7.8	10.4	6.0	6.7	8.8	5.4
2002	7.5	10.1	5.7	7.7	10.4	5.8	6.7	8.8	5.4
2003	7.4	9.9	5.6	7.6	10.3	5.7	6.5	8.4	5.3
2004	7.2	9.7	5.5	7.4	9.9	5.6	6.7	8.8	5.3
1975-2004	7.9	10.6	6.1	8.0	10.8	6.2	7.0	9.2	5.6

<sup>a</sup> NCHS public use data file for the total US. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

- Statistic not shown. Rate based on less than 16 cases for the time interval.



Table XIV-2  
LIVER AND INTRAHEPATIC BILE DUCT CANCER (Invasive)

AGE ADJUSTED SEER INCIDENCE<sup>a</sup> RATES BY YEAR, RACE AND SEX

	<u>All Races</u>			<u>Whites</u>			<u>Blacks</u>		
	<u>Total</u>	<u>Males</u>	<u>Females</u>	<u>Total</u>	<u>Males</u>	<u>Females</u>	<u>Total</u>	<u>Males</u>	<u>Females</u>
<u>AGE-ADJUSTED RATES</u>									
<u>YEAR OF DIAGNOSIS:</u>									
1975	2.6	3.9	1.6	2.2	3.2	1.4	4.7	6.5	3.2
1976	2.7	3.8	1.8	2.3	3.2	1.6	4.0	6.2	-
1977	2.7	4.2	1.6	2.2	3.6	1.3	3.5	5.0	-
1978	2.7	4.2	1.6	2.2	3.3	1.4	4.2	6.8	2.1
1979	2.7	4.1	1.7	2.2	3.3	1.5	4.1	7.1	-
1980	2.6	3.9	1.7	2.2	3.3	1.5	3.6	5.0	2.5
1981	2.9	4.6	1.7	2.5	3.9	1.5	4.6	7.1	2.5
1982	2.9	4.4	1.9	2.4	3.5	1.6	3.9	5.4	2.6
1983	2.9	4.7	1.6	2.4	3.7	1.4	4.1	6.3	2.3
1984	2.9	4.3	1.9	2.3	3.4	1.6	3.3	5.3	1.9
1985	3.2	5.1	1.7	2.6	4.2	1.5	4.0	7.6	-
1986	3.2	4.8	2.0	2.6	3.9	1.7	5.0	7.6	3.1
1987	3.4	4.8	2.2	2.7	3.6	1.9	5.3	7.7	3.6
1988	3.4	5.4	1.8	2.7	4.2	1.6	4.8	8.9	-
1989	3.6	5.6	2.2	2.9	4.3	1.9	5.3	9.1	2.6
1990	3.9	6.0	2.3	3.1	4.7	1.9	6.5	10.6	3.5
1991	4.3	6.7	2.6	3.5	5.5	2.1	6.0	9.2	3.5
1992	4.0	6.1	2.4	3.2	4.8	1.9	5.6	8.4	3.4
1993	4.5	7.1	2.5	3.7	5.9	2.1	6.2	10.1	3.1
1994	4.5	7.0	2.5	3.6	5.6	2.1	5.1	9.6	1.9
1995	4.5	6.8	2.8	3.6	5.4	2.2	6.0	8.6	3.9
1996	5.4	8.3	3.1	4.4	6.7	2.6	6.7	10.8	3.6
1997	5.4	8.1	3.2	4.4	6.5	2.6	7.5	11.5	4.4
1998	5.5	8.3	3.2	4.5	6.9	2.6	6.5	9.6	4.1
1999	5.8	8.8	3.3	4.7	7.2	2.7	7.7	12.1	4.4
2000	5.6	8.5	3.2	4.7	7.0	2.7	6.9	11.6	3.4
2001	5.6	8.5	3.1	4.3	6.5	2.5	7.9	13.7	3.5
2002	5.7	8.7	3.2	4.6	6.9	2.6	7.2	11.9	3.6
2003	6.0	9.2	3.3	4.9	7.5	2.6	7.6	12.3	3.9
2004	6.1	9.5	3.2	4.9	7.6	2.5	9.0	14.8	4.4
1975-2004	4.2	6.5	2.4	3.4	5.2	2.0	5.9	9.5	3.2

<sup>a</sup> SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta). Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).  
 - Statistic not shown. Rate based on less than 16 cases for the time interval.

Table XIV-3  
LIVER AND INTRAHEPATIC BILE DUCT CANCER (Invasive)  
 AGE-ADJUSTED U.S. DEATH<sup>a</sup> RATES BY YEAR, RACE AND SEX

	<u>All Races</u>			<u>Whites</u>			<u>Blacks</u>		
	<u>Total</u>	<u>Males</u>	<u>Females</u>	<u>Total</u>	<u>Males</u>	<u>Females</u>	<u>Total</u>	<u>Males</u>	<u>Females</u>
<u>AGE-ADJUSTED RATES</u>									
<u>YEAR OF DEATH:</u>									
1975	2.8	3.8	2.0	2.6	3.5	2.0	4.3	6.3	2.7
1976	2.7	3.8	2.0	2.5	3.5	1.9	4.5	6.8	2.7
1977	2.8	3.8	2.0	2.6	3.4	1.9	4.5	7.2	2.5
1978	2.7	3.9	1.9	2.5	3.5	1.8	4.6	7.0	2.8
1979	2.8	3.9	2.0	2.6	3.5	2.0	4.4	6.6	2.7
1980	2.8	3.9	2.0	2.6	3.5	1.9	4.5	6.8	2.8
1981	2.9	4.1	2.0	2.7	3.8	1.9	4.4	6.7	2.7
1982	2.9	4.2	2.1	2.7	3.8	1.9	4.7	7.2	2.9
1983	2.9	4.2	2.0	2.6	3.7	1.9	4.8	7.3	3.0
1984	3.0	4.3	2.1	2.7	3.8	2.0	4.8	7.9	2.7
1985	3.1	4.4	2.2	2.8	3.9	2.1	4.6	7.2	2.7
1986	3.1	4.5	2.2	2.9	4.1	2.0	4.7	7.3	2.9
1987	3.2	4.6	2.2	2.9	4.1	2.0	4.8	6.9	3.2
1988	3.3	4.8	2.2	3.0	4.4	2.1	4.8	7.2	3.0
1989	3.4	5.0	2.3	3.1	4.5	2.2	5.1	8.0	3.1
1990	3.6	5.3	2.4	3.3	4.7	2.2	5.3	7.8	3.6
1991	3.7	5.3	2.5	3.4	4.8	2.3	5.3	8.0	3.5
1992	3.9	5.7	2.6	3.6	5.2	2.4	5.4	8.5	3.3
1993	4.0	5.8	2.7	3.7	5.3	2.6	5.5	8.1	3.6
1994	4.1	6.0	2.7	3.8	5.4	2.6	5.7	8.9	3.5
1995	4.4	6.3	2.9	4.0	5.7	2.7	6.2	9.3	4.1
1996	4.4	6.5	2.9	4.1	5.9	2.7	5.9	9.1	3.7
1997	4.5	6.5	3.0	4.1	5.9	2.8	6.0	8.9	3.9
1998	4.6	6.7	2.9	4.2	6.1	2.7	6.4	10.1	3.7
1999	4.5	6.5	2.9	4.1	6.0	2.7	5.9	8.8	3.8
2000	4.6	6.8	2.9	4.2	6.1	2.7	6.1	9.5	3.6
2001	4.7	6.9	3.0	4.3	6.4	2.7	6.3	9.1	4.1
2002	4.9	7.2	3.0	4.5	6.5	2.8	6.3	10.0	3.6
2003	5.0	7.4	3.1	4.6	6.7	2.8	6.7	10.3	4.0
2004	5.1	7.4	3.2	4.7	6.7	3.0	6.9	10.7	4.1
1975-2004	3.8	5.5	2.5	3.5	5.0	2.4	5.5	8.4	3.4

<sup>a</sup> NCHS public use data file for the total US. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

- Statistic not shown. Rate based on less than 16 cases for the time interval.

Table XV-3  
LUNG AND BRONCHUS CANCER (Invasive)

AGE ADJUSTED SEER INCIDENCE<sup>a</sup> RATES BY YEAR, RACE AND SEX

	All Races			Whites			Blacks		
	Total	Males	Females	Total	Males	Females	Total	Males	Females
<u>AGE-ADJUSTED RATES</u>									
YEAR OF DIAGNOSIS:									
1975	52.3	89.6	24.5	52.0	89.2	24.9	64.6	114.9	24.7
1976	55.4	93.8	27.3	54.6	92.8	27.3	72.3	131.0	28.6
1977	56.7	95.5	28.3	55.9	94.7	28.1	73.6	124.8	33.9
1978	57.8	96.7	29.7	57.3	95.9	29.9	74.5	130.5	31.6
1979	58.6	95.8	31.6	58.0	95.2	31.6	74.5	127.6	34.1
1980	60.7	100.0	32.2	59.4	97.7	32.4	86.6	151.5	38.3
1981	62.0	99.4	35.1	61.6	98.5	35.5	82.2	143.0	37.2
1982	63.3	100.7	36.7	63.1	99.7	37.7	80.4	141.1	35.7
1983	63.4	99.5	37.9	63.0	97.6	39.1	86.6	151.3	39.5
1984	65.5	102.1	39.5	64.3	99.9	39.6	93.1	159.2	44.8
1985	64.6	98.7	40.2	64.0	96.9	40.9	89.6	149.7	46.0
1986	65.8	99.1	42.3	65.2	97.4	42.9	92.9	156.2	48.6
1987	67.9	101.5	44.2	68.1	100.7	45.5	84.8	143.7	43.2
1988	68.0	98.7	46.3	68.2	97.9	47.7	88.1	144.9	48.1
1989	67.5	97.9	46.1	67.5	97.0	47.2	89.0	142.8	51.4
1990	68.0	96.9	47.7	68.2	96.1	48.9	86.7	137.0	52.0
1991	69.2	97.3	49.6	69.0	95.9	50.6	93.2	145.0	56.9
1992	69.4	97.2	49.8	68.9	94.8	51.0	93.9	149.0	55.8
1993	67.7	93.9	49.2	67.8	92.5	50.7	86.3	135.4	52.4
1994	67.1	90.9	50.5	67.3	90.1	51.6	86.9	131.1	56.2
1995	66.8	89.7	50.4	67.0	87.9	52.4	85.7	138.6	50.0
1996	66.4	88.0	51.1	66.7	86.7	52.9	84.0	127.9	55.1
1997	66.5	86.2	52.5	67.2	84.8	54.9	82.1	129.4	50.4
1998	67.5	88.0	52.9	68.0	86.5	55.0	87.0	130.5	57.7
1999	65.7	84.5	52.4	66.2	83.2	54.3	84.2	121.7	58.7
2000	63.9	81.8	51.1	64.6	80.6	53.2	79.1	112.8	56.5
2001	63.8	80.8	51.4	64.8	80.6	53.4	79.6	113.4	56.7
2002	63.5	79.8	51.9	64.2	79.2	53.7	80.5	111.1	60.1
2003	63.8	79.9	52.2	64.7	78.5	54.9	79.3	116.4	54.3
2004	60.0	73.6	50.2	60.6	72.7	52.0	73.0	100.0	54.9
1975-2004	64.3	91.8	44.5	64.3	90.7	45.4	83.2	132.1	48.8

<sup>a</sup> SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta). Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).  
- Statistic not shown. Rate based on less than 16 cases for the time interval.

Table XV-4  
 SMALL CELL LUNG AND BRONCHUS CANCER (Invasive)

AGE ADJUSTED SEER INCIDENCE<sup>a</sup> RATES BY YEAR, RACE AND SEX

	All Races			Whites			Blacks		
	Total	Males	Females	Total	Males	Females	Total	Males	Females
<u>AGE-ADJUSTED RATES</u>									
YEAR OF DIAGNOSIS:									
1975	6.6	10.4	3.8	6.9	10.7	4.0	5.9	8.9	3.5
1976	7.1	11.4	3.8	7.1	11.5	3.9	8.2	12.8	4.8
1977	7.7	12.4	4.2	7.8	12.8	4.2	8.1	11.2	5.8
1978	8.6	13.4	5.0	8.9	13.7	5.3	7.8	13.9	3.1
1979	8.7	13.1	5.3	9.0	13.6	5.6	7.6	11.8	4.5
1980	9.1	14.2	5.2	9.3	14.4	5.5	10.0	17.5	4.4
1981	9.7	14.5	6.1	10.0	14.8	6.5	9.2	15.9	4.2
1982	10.4	14.9	7.1	10.7	15.3	7.6	9.6	15.9	4.9
1983	10.4	14.9	7.1	10.7	15.0	7.6	9.1	15.1	4.9
1984	10.7	15.2	7.4	11.0	15.6	7.8	10.4	15.0	6.9
1985	10.4	15.0	7.1	10.8	15.3	7.5	11.1	17.8	6.3
1986	11.2	15.5	8.0	11.5	16.0	8.3	12.3	17.7	8.4
1987	10.6	14.1	8.2	11.2	14.6	8.8	9.5	14.5	6.0
1988	11.4	15.3	8.5	11.9	15.6	9.2	10.5	17.9	5.3
1989	11.2	14.8	8.7	11.7	15.2	9.2	12.1	17.7	8.4
1990	10.7	13.8	8.4	11.3	14.5	9.1	9.4	13.7	6.2
1991	11.3	14.0	9.4	11.7	14.3	9.8	12.4	16.0	10.0
1992	10.5	13.4	8.4	11.0	13.8	9.0	10.2	13.4	8.0
1993	10.0	12.3	8.3	10.5	12.6	8.9	9.4	13.2	6.7
1994	9.8	11.7	8.4	10.2	12.0	9.1	9.9	13.6	7.4
1995	9.8	12.2	8.1	10.4	12.6	8.8	9.4	13.7	6.6
1996	9.7	12.1	8.0	10.3	12.4	8.6	8.8	12.0	6.8
1997	9.5	10.9	8.6	10.1	11.1	9.5	7.9	10.8	6.1
1998	9.3	10.8	8.1	9.9	11.3	8.8	9.4	12.1	7.4
1999	8.8	10.0	7.9	9.3	10.2	8.7	8.1	10.9	6.0
2000	8.7	10.4	7.6	9.3	10.8	8.1	9.4	11.5	7.8
2001	8.3	9.8	7.3	8.9	10.2	7.9	8.4	10.8	6.8
2002	8.2	9.3	7.5	8.9	9.8	8.3	7.2	9.7	5.7
2003	8.1	9.0	7.5	8.6	9.2	8.3	7.8	9.4	6.7
2004	7.9	8.7	7.3	8.3	9.0	7.8	8.4	9.7	7.6
1975-2004	9.5	12.4	7.3	9.9	12.8	7.8	9.2	13.2	6.4

<sup>a</sup> SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta). Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).  
 - Statistic not shown. Rate based on less than 16 cases for the time interval.

Table XV-5  
 NON-SMALL CELL LUNG AND BRONCHUS CANCER (Invasive)

AGE ADJUSTED SEER INCIDENCE<sup>a</sup> RATES BY YEAR, RACE AND SEX

	All Races			Whites			Blacks		
	Total	Males	Females	Total	Males	Females	Total	Males	Females
<u>AGE-ADJUSTED RATES</u>									
YEAR OF DIAGNOSIS:									
1975	45.6	79.2	20.8	45.1	78.5	20.9	58.7	106.0	21.2
1976	48.4	82.5	23.5	47.5	81.3	23.4	64.1	118.2	23.9
1977	49.0	83.1	24.1	48.0	82.0	23.8	65.4	113.6	28.1
1978	49.2	83.3	24.6	48.4	82.2	24.6	66.7	116.6	28.5
1979	49.9	82.7	26.2	49.0	81.6	26.0	66.9	115.8	29.6
1980	51.6	85.8	27.0	50.1	83.3	26.8	76.7	133.9	33.9
1981	52.3	84.9	29.0	51.5	83.6	29.1	73.0	127.1	33.0
1982	52.9	85.8	29.6	52.3	84.4	30.2	70.7	125.1	30.8
1983	53.1	84.6	30.8	52.3	82.6	31.5	77.5	136.3	34.6
1984	54.8	86.9	32.0	53.3	84.3	31.9	82.6	144.2	38.0
1985	54.2	83.7	33.1	53.2	81.7	33.4	78.5	131.9	39.8
1986	54.6	83.5	34.3	53.6	81.4	34.7	80.6	138.5	40.3
1987	57.3	87.4	36.0	56.9	86.1	36.7	75.2	129.2	37.2
1988	56.6	83.4	37.8	56.3	82.3	38.5	77.6	127.1	42.8
1989	56.3	83.1	37.4	55.9	81.8	38.0	77.0	125.1	43.0
1990	57.4	83.1	39.3	56.9	81.7	39.8	77.3	123.3	45.8
1991	57.9	83.3	40.2	57.3	81.6	40.8	80.8	129.0	46.9
1992	58.9	83.8	41.4	57.9	81.0	42.1	83.7	135.5	47.8
1993	57.7	81.6	40.8	57.3	79.8	41.7	76.9	122.1	45.7
1994	57.4	79.2	42.1	57.0	78.1	42.6	76.9	117.5	48.7
1995	56.9	77.5	42.3	56.7	75.3	43.7	76.2	124.9	43.4
1996	56.7	75.9	43.1	56.5	74.2	44.3	75.2	115.9	48.2
1997	57.0	75.3	43.9	57.0	73.6	45.4	74.3	118.6	44.3
1998	58.2	77.1	44.7	58.1	75.3	46.1	77.7	118.4	50.2
1999	56.9	74.4	44.4	56.9	72.9	45.6	76.0	110.8	52.6
2000	55.2	71.5	43.5	55.3	69.9	45.1	69.7	101.3	48.7
2001	55.4	71.1	44.1	55.9	70.3	45.4	71.1	102.6	49.9
2002	55.2	70.4	44.4	55.3	69.4	45.4	73.2	101.4	54.4
2003	55.7	70.8	44.7	56.0	69.3	46.6	71.5	107.0	47.6
2004	52.1	64.8	42.9	52.3	63.7	44.1	64.5	90.3	47.3
1975-2004	54.8	79.4	37.2	54.4	78.0	37.6	74.0	118.9	42.4

<sup>a</sup> SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta). Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).  
 - Statistic not shown. Rate based on less than 16 cases for the time interval.

Table XV-6  
LUNG AND BRONCHUS CANCER (Invasive)

AGE-ADJUSTED U.S. DEATH<sup>a</sup> RATES BY YEAR, RACE AND SEX

	All Races			Whites			Blacks		
	Total	Males	Females	Total	Males	Females	Total	Males	Females
<u>AGE-ADJUSTED RATES</u>									
YEAR OF DEATH:									
1975	42.6	76.4	17.6	42.1	75.5	17.7	49.3	91.0	17.3
1976	44.2	78.5	19.0	43.7	77.6	19.1	51.1	94.0	18.5
1977	45.5	80.4	20.0	44.8	79.1	20.1	54.6	100.8	19.9
1978	46.9	82.1	21.4	46.3	80.8	21.6	55.3	101.6	20.7
1979	47.7	82.9	22.3	47.1	81.6	22.4	56.4	102.9	22.2
1980	49.4	84.7	24.1	48.8	83.3	24.2	59.2	106.7	24.6
1981	50.0	85.2	24.9	49.3	83.5	25.1	60.6	109.7	25.1
1982	51.4	86.8	26.5	50.8	85.2	26.7	62.3	112.5	26.3
1983	52.4	87.0	28.1	51.7	85.4	28.3	63.7	112.8	28.7
1984	53.4	88.2	28.9	52.6	86.2	29.3	65.1	118.0	28.0
1985	54.3	88.5	30.4	53.7	86.7	30.8	65.8	117.5	29.6
1986	55.0	89.0	31.4	54.4	87.0	31.8	67.0	119.2	31.0
1987	56.2	90.0	32.7	55.6	88.1	33.0	68.6	121.8	32.5
1988	57.0	89.9	34.1	56.4	87.9	34.6	69.2	122.2	33.5
1989	57.9	89.8	35.8	57.2	87.6	36.3	71.0	124.7	34.9
1990	58.9	90.6	36.8	58.2	88.4	37.3	72.1	125.2	36.5
1991	59.0	89.9	37.6	58.4	87.8	38.1	71.6	123.6	37.2
1992	58.9	88.0	38.7	58.3	86.0	39.2	71.4	121.1	38.6
1993	59.1	87.6	39.3	58.7	85.5	40.0	71.1	122.0	37.7
1994	58.5	85.7	39.6	58.2	83.8	40.3	70.2	118.6	38.6
1995	58.4	84.4	40.3	58.1	82.6	41.0	69.3	116.1	38.8
1996	57.9	82.8	40.4	57.7	81.2	41.2	68.3	113.2	39.1
1997	57.5	81.3	40.8	57.3	79.8	41.5	67.9	110.2	40.5
1998	57.1	79.9	41.0	56.9	78.5	41.7	66.9	107.4	40.7
1999	55.4	77.0	40.1	55.3	75.7	40.9	65.0	103.6	40.0
2000	55.8	76.5	41.1	56.0	75.4	42.1	63.8	100.9	39.7
2001	55.3	75.2	41.0	55.5	74.3	42.0	62.7	99.4	38.8
2002	55.0	73.5	41.6	55.3	72.8	42.6	62.2	95.7	40.3
2003	54.2	71.9	41.2	54.5	71.2	42.2	61.1	93.1	40.3
2004	53.3	70.3	40.9	53.6	69.6	41.9	60.0	90.6	40.0
1975-2004	54.5	82.9	34.2	54.1	81.5	34.6	64.4	109.2	33.7

<sup>a</sup> NCHS public use data file for the total US. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

- Statistic not shown. Rate based on less than 16 cases for the time interval.

Table XVI-2  
MELANOMA OF THE SKIN (Invasive)

AGE ADJUSTED SEER INCIDENCE<sup>a</sup> RATES BY YEAR, RACE AND SEX

	<u>All Races</u>			<u>Whites</u>			<u>Blacks</u>		
	<u>Total</u>	<u>Males</u>	<u>Females</u>	<u>Total</u>	<u>Males</u>	<u>Females</u>	<u>Total</u>	<u>Males</u>	<u>Females</u>
<u>AGE-ADJUSTED RATES</u>									
<u>YEAR OF DIAGNOSIS:</u>									
1975	7.9	8.5	7.4	8.7	9.4	8.2	-	-	-
1976	8.1	8.7	7.7	9.0	9.6	8.5	1.7	-	-
1977	8.9	9.8	8.2	9.8	10.8	9.1	-	-	-
1978	8.9	9.5	8.7	10.1	10.7	9.8	-	-	-
1979	9.5	10.6	8.9	10.8	11.9	10.1	-	-	-
1980	10.5	11.8	9.6	12.0	13.4	11.1	-	-	-
1981	11.1	12.1	10.4	12.6	13.6	12.0	1.4	-	-
1982	11.2	12.5	10.3	12.8	14.1	11.8	-	-	-
1983	11.1	12.7	10.0	12.6	14.4	11.4	-	-	-
1984	11.4	12.9	10.4	13.0	14.8	11.9	-	-	-
1985	12.8	15.1	11.2	14.3	17.0	12.6	-	-	-
1986	13.3	15.7	11.6	15.1	17.8	13.2	-	-	-
1987	13.7	15.9	12.2	15.6	18.1	13.9	1.3	-	-
1988	12.9	15.0	11.4	14.9	17.4	13.2	1.1	-	-
1989	13.7	16.4	11.9	15.8	18.9	13.7	1.2	-	-
1990	13.8	16.6	11.8	16.1	19.2	13.9	1.2	-	-
1991	14.6	17.6	12.5	17.0	20.4	14.7	1.2	-	-
1992	14.8	18.3	12.2	17.1	21.1	14.3	-	-	-
1993	14.6	18.1	12.1	17.0	20.9	14.2	-	-	-
1994	15.6	19.8	12.6	18.3	23.0	14.8	1.1	-	-
1995	16.4	20.2	13.7	19.2	23.4	16.3	1.7	3.0	-
1996	17.2	21.6	14.1	20.3	25.2	16.8	1.1	-	-
1997	17.7	21.6	14.8	20.8	25.1	17.8	1.1	-	-
1998	17.8	22.1	14.8	21.2	26.0	17.9	1.2	-	-
1999	18.1	22.7	14.9	21.6	26.7	18.1	1.1	-	-
2000	18.8	23.7	15.3	22.6	28.0	18.8	0.8	-	-
2001	19.5	24.2	16.2	23.6	28.8	20.0	0.8	-	-
2002	18.6	23.1	15.5	22.4	27.3	19.1	1.0	-	-
2003	18.9	23.2	16.0	22.7	27.3	19.5	0.8	-	-
2004	19.6	24.1	16.5	23.8	28.9	20.3	0.8	-	-
1975-2004	14.6	17.6	12.5	16.9	20.3	14.6	1.0	1.2	0.9

<sup>a</sup> SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta). Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

- Statistic not shown. Rate based on less than 16 cases for the time interval.

Table XVI-3  
MELANOMA OF THE SKIN (Invasive)

AGE-ADJUSTED U.S. DEATH<sup>a</sup> RATES BY YEAR, RACE AND SEX

	<u>All Races</u>			<u>Whites</u>			<u>Blacks</u>		
	<u>Total</u>	<u>Males</u>	<u>Females</u>	<u>Total</u>	<u>Males</u>	<u>Females</u>	<u>Total</u>	<u>Males</u>	<u>Females</u>
<u>AGE-ADJUSTED RATES</u>									
<u>YEAR OF DEATH:</u>									
1975	2.1	2.6	1.6	2.3	2.9	1.7	0.5	0.6	0.4
1976	2.2	2.9	1.8	2.5	3.1	1.9	0.5	0.7	0.4
1977	2.3	2.9	1.8	2.5	3.2	1.9	0.5	0.7	0.4
1978	2.3	3.0	1.7	2.5	3.3	1.9	0.5	0.6	0.5
1979	2.4	3.2	1.8	2.7	3.5	2.0	0.5	0.6	0.5
1980	2.3	3.1	1.7	2.6	3.4	1.9	0.5	0.5	0.6
1981	2.4	3.1	1.9	2.7	3.4	2.1	0.6	0.6	0.6
1982	2.5	3.3	1.8	2.7	3.6	2.0	0.5	0.6	0.5
1983	2.5	3.3	1.8	2.7	3.6	2.0	0.5	0.6	0.4
1984	2.5	3.4	1.8	2.8	3.8	2.0	0.5	0.5	0.5
1985	2.6	3.5	1.9	2.8	3.8	2.1	0.5	0.6	0.5
1986	2.6	3.5	1.9	2.9	3.9	2.1	0.5	0.7	0.3
1987	2.6	3.7	1.8	2.9	4.1	2.0	0.6	0.6	0.5
1988	2.6	3.7	1.9	3.0	4.1	2.1	0.5	0.5	0.4
1989	2.7	3.7	1.9	3.0	4.2	2.1	0.6	0.6	0.5
1990	2.8	3.8	2.0	3.1	4.2	2.2	0.5	0.6	0.4
1991	2.7	3.9	1.8	3.0	4.3	2.0	0.6	0.6	0.5
1992	2.7	3.8	1.8	3.0	4.3	2.1	0.5	0.6	0.4
1993	2.7	3.9	1.9	3.0	4.3	2.1	0.5	0.4	0.6
1994	2.7	3.8	1.8	3.0	4.3	2.0	0.4	0.4	0.4
1995	2.7	3.9	1.8	3.0	4.3	2.0	0.5	0.5	0.4
1996	2.8	4.0	1.9	3.1	4.5	2.1	0.4	0.5	0.4
1997	2.7	3.9	1.8	3.1	4.4	2.1	0.6	0.5	0.6
1998	2.8	4.1	1.8	3.1	4.6	2.0	0.5	0.5	0.5
1999	2.6	3.8	1.7	3.0	4.3	2.0	0.4	0.5	0.3
2000	2.7	3.8	1.8	3.0	4.3	2.0	0.5	0.6	0.5
2001	2.7	3.9	1.7	3.0	4.4	2.0	0.4	0.3	0.4
2002	2.6	3.8	1.7	3.0	4.2	2.0	0.4	0.5	0.3
2003	2.7	3.9	1.7	3.0	4.4	2.0	0.4	0.4	0.4
2004	2.7	3.9	1.7	3.0	4.5	1.9	0.4	0.5	0.4
1975-2004	2.6	3.6	1.8	2.9	4.0	2.0	0.5	0.5	0.5

<sup>a</sup> NCHS public use data file for the total US. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

- Statistic not shown. Rate based on less than 16 cases for the time interval.



Table XVII-2

MESOTHELIOMAAGE ADJUSTED SEER INCIDENCE<sup>a</sup> RATES BY YEAR, RACE AND SEX

	<u>All Races</u>			<u>Whites</u>			<u>Blacks</u>		
	<u>Total</u>	<u>Males</u>	<u>Females</u>	<u>Total</u>	<u>Males</u>	<u>Females</u>	<u>Total</u>	<u>Males</u>	<u>Females</u>
<u>AGE-ADJUSTED RATES</u>									
<u>YEAR OF DIAGNOSIS:</u>									
1975	0.6	1.0	0.3	0.6	1.0	0.3	-	-	-
1976	0.7	1.1	0.3	0.7	1.2	0.4	-	-	-
1977	0.6	1.0	0.3	0.6	1.1	0.3	-	-	-
1978	0.7	1.4	0.2	0.8	1.5	0.2	-	-	-
1979	0.8	1.5	0.3	0.8	1.6	0.3	-	-	-
1980	1.0	1.8	0.4	1.1	2.0	0.4	-	-	-
1981	0.8	1.4	0.3	0.8	1.5	0.3	-	-	-
1982	0.8	1.4	0.4	0.9	1.5	0.4	-	-	-
1983	1.0	1.9	0.5	1.1	2.0	0.5	-	-	-
1984	1.1	2.0	0.4	1.1	2.2	0.4	-	-	-
1985	0.9	1.7	0.4	1.0	1.7	0.4	-	-	-
1986	0.9	1.7	0.4	1.0	1.8	0.5	-	-	-
1987	1.0	1.7	0.4	1.0	1.8	0.5	-	-	-
1988	1.0	1.8	0.4	1.1	2.0	0.4	-	-	-
1989	1.1	2.2	0.4	1.2	2.4	0.4	-	-	-
1990	1.1	2.2	0.5	1.2	2.3	0.5	-	-	-
1991	1.1	2.1	0.4	1.2	2.2	0.5	-	-	-
1992	1.2	2.5	0.4	1.3	2.7	0.4	-	-	-
1993	1.1	2.0	0.4	1.1	2.2	0.4	-	-	-
1994	1.2	2.2	0.4	1.2	2.3	0.5	1.1	-	-
1995	1.2	2.2	0.5	1.2	2.3	0.5	-	-	-
1996	1.1	2.1	0.5	1.3	2.4	0.5	-	-	-
1997	1.0	1.9	0.4	1.1	2.1	0.4	0.9	-	-
1998	1.1	2.1	0.4	1.2	2.2	0.5	-	-	-
1999	1.1	2.2	0.4	1.2	2.3	0.4	-	-	-
2000	1.1	2.1	0.3	1.2	2.4	0.4	-	-	-
2001	1.1	2.0	0.4	1.2	2.2	0.4	-	-	-
2002	1.0	1.9	0.4	1.2	2.2	0.4	-	-	-
2003	0.9	1.7	0.4	1.0	1.8	0.4	-	-	-
2004	1.1	2.0	0.4	1.2	2.2	0.5	-	-	-
1975-2004	1.0	1.9	0.4	1.1	2.0	0.4	0.6	1.1	0.3

<sup>a</sup> SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta). Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

- Statistic not shown. Rate based on less than 16 cases for the time interval.

Table XVIII-2

MYELOMAAGE ADJUSTED SEER INCIDENCE<sup>a</sup> RATES BY YEAR, RACE AND SEX

	<u>All Races</u>			<u>Whites</u>			<u>Blacks</u>		
	<u>Total</u>	<u>Males</u>	<u>Females</u>	<u>Total</u>	<u>Males</u>	<u>Females</u>	<u>Total</u>	<u>Males</u>	<u>Females</u>
<u>AGE-ADJUSTED RATES</u>									
<u>YEAR OF DIAGNOSIS:</u>									
1975	4.9	6.4	3.9	4.6	6.2	3.6	9.7	10.9	8.6
1976	5.0	6.1	4.4	4.8	5.8	4.1	9.2	11.2	7.9
1977	5.1	6.4	4.2	4.8	6.2	3.9	9.1	11.0	7.7
1978	4.7	5.9	3.9	4.4	5.6	3.7	9.6	11.0	8.7
1979	4.9	6.0	4.1	4.5	5.4	3.8	11.0	15.3	8.2
1980	4.8	5.8	4.1	4.4	5.5	3.7	9.9	10.8	9.0
1981	5.0	6.0	4.3	4.6	5.5	4.0	11.0	15.2	8.2
1982	5.3	6.3	4.7	4.9	5.9	4.3	11.6	13.9	9.7
1983	5.3	6.9	4.2	4.7	6.3	3.7	13.7	16.2	11.9
1984	5.4	7.0	4.3	5.0	6.4	4.1	10.8	15.3	7.7
1985	5.2	6.3	4.3	4.9	5.9	4.2	9.3	12.6	7.0
1986	5.3	6.3	4.7	4.9	6.0	4.1	11.1	11.6	10.5
1987	5.9	7.4	5.0	5.7	7.2	4.7	11.1	13.5	9.7
1988	5.3	6.6	4.5	5.0	6.4	4.2	10.8	12.2	9.7
1989	5.4	6.7	4.5	5.1	6.4	4.2	10.7	13.5	9.0
1990	5.6	7.2	4.6	5.2	6.9	4.1	11.9	14.6	10.2
1991	6.0	7.7	4.8	5.6	7.2	4.4	12.8	15.9	10.6
1992	5.9	7.3	5.1	5.6	7.0	4.7	11.5	13.0	10.4
1993	5.6	7.3	4.3	5.1	6.8	3.9	12.7	15.6	10.6
1994	5.7	6.8	5.0	5.2	6.3	4.4	12.8	15.2	11.8
1995	5.7	7.2	4.7	5.3	6.8	4.3	12.0	15.0	10.1
1996	5.8	7.1	4.7	5.4	6.8	4.3	12.6	13.4	11.9
1997	6.1	7.3	5.2	5.7	7.0	4.7	12.6	14.2	11.9
1998	5.9	7.2	4.8	5.5	7.0	4.4	11.5	13.0	10.5
1999	5.6	7.0	4.5	5.1	6.6	4.1	11.9	14.9	9.9
2000	5.9	7.4	4.9	5.7	7.4	4.4	10.6	10.5	10.5
2001	5.8	7.1	4.8	5.4	6.8	4.4	11.6	13.9	10.2
2002	5.8	7.6	4.4	5.5	7.3	4.1	10.7	14.0	8.6
2003	5.5	7.0	4.5	5.3	6.8	4.2	10.3	13.4	8.2
2004	5.3	6.7	4.3	4.8	6.2	3.9	11.3	14.1	9.3
1975-2004	5.5	6.9	4.5	5.1	6.5	4.2	11.3	13.6	9.7

<sup>a</sup> SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta). Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

- Statistic not shown. Rate based on less than 16 cases for the time interval.

Table XVIII-3

MYELOMAAGE-ADJUSTED U.S. DEATH<sup>a</sup> RATES BY YEAR, RACE AND SEX

	<u>All Races</u>			<u>Whites</u>			<u>Blacks</u>		
	<u>Total</u>	<u>Males</u>	<u>Females</u>	<u>Total</u>	<u>Males</u>	<u>Females</u>	<u>Total</u>	<u>Males</u>	<u>Females</u>
<u>AGE-ADJUSTED RATES</u>									
<u>YEAR OF DEATH:</u>									
1975	2.9	3.7	2.4	2.7	3.5	2.2	5.3	6.4	4.4
1976	3.1	3.9	2.6	2.9	3.6	2.4	5.6	6.9	4.6
1977	3.1	3.9	2.6	2.9	3.6	2.4	5.6	7.2	4.4
1978	3.2	3.9	2.7	2.9	3.7	2.5	5.7	7.0	4.9
1979	3.3	4.1	2.7	3.0	3.9	2.5	5.9	7.4	4.9
1980	3.3	4.1	2.7	3.0	3.8	2.5	6.1	7.4	5.2
1981	3.3	4.1	2.8	3.0	3.7	2.5	6.6	8.2	5.5
1982	3.4	4.2	2.9	3.1	3.8	2.6	6.7	8.6	5.4
1983	3.4	4.3	2.9	3.1	4.0	2.6	6.8	8.7	5.6
1984	3.5	4.3	3.0	3.2	3.9	2.7	6.6	8.4	5.5
1985	3.6	4.5	3.0	3.3	4.1	2.8	6.9	8.6	5.8
1986	3.6	4.5	3.0	3.3	4.1	2.8	7.2	9.1	6.0
1987	3.6	4.6	3.0	3.3	4.2	2.7	7.1	9.2	5.7
1988	3.6	4.6	3.0	3.4	4.3	2.7	6.9	8.1	6.1
1989	3.7	4.6	3.1	3.4	4.3	2.8	7.1	8.9	5.9
1990	3.8	4.8	3.1	3.5	4.4	2.9	7.3	9.6	5.9
1991	3.9	4.8	3.3	3.5	4.5	2.9	7.8	9.6	6.6
1992	3.8	4.7	3.2	3.5	4.4	2.9	7.5	9.3	6.3
1993	4.0	4.9	3.4	3.6	4.6	3.0	7.8	9.3	6.9
1994	4.0	5.0	3.3	3.7	4.7	3.0	7.7	9.7	6.5
1995	4.0	4.9	3.4	3.7	4.6	3.0	7.8	9.4	6.9
1996	3.9	4.9	3.2	3.6	4.6	2.9	7.7	9.6	6.5
1997	3.9	4.9	3.2	3.6	4.6	2.9	7.5	9.2	6.5
1998	3.8	4.8	3.2	3.5	4.4	2.9	7.7	9.5	6.8
1999	3.8	4.7	3.2	3.5	4.4	2.9	7.5	8.8	6.6
2000	3.8	4.7	3.3	3.5	4.4	3.0	7.5	8.8	6.7
2001	3.8	4.7	3.2	3.5	4.4	2.9	7.2	8.7	6.3
2002	3.8	4.8	3.1	3.6	4.5	2.9	6.9	8.3	6.0
2003	3.7	4.6	3.1	3.4	4.3	2.8	7.1	8.4	6.3
2004	3.6	4.4	3.0	3.3	4.1	2.7	6.8	8.2	6.0
1975-2004	3.6	4.6	3.0	3.4	4.2	2.8	7.0	8.6	6.0

<sup>a</sup> NCHS public use data file for the total US. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

- Statistic not shown. Rate based on less than 16 cases for the time interval.

Table XIX-2  
NON-HODGKIN LYMPHOMA

AGE ADJUSTED SEER INCIDENCE<sup>a</sup> RATES BY YEAR, RACE AND SEX

	All Races			Whites			Blacks		
	Total	Males	Females	Total	Males	Females	Total	Males	Females
<u>AGE-ADJUSTED RATES</u>									
YEAR OF DIAGNOSIS:									
1975	11.1	12.9	9.6	11.6	13.5	10.1	6.4	7.3	5.3
1976	11.2	12.8	10.0	11.6	13.2	10.5	6.0	7.4	4.9
1977	11.2	12.9	9.8	11.7	13.5	10.2	6.6	7.2	6.1
1978	11.9	14.3	10.1	12.5	14.8	10.7	7.2	9.5	5.2
1979	12.5	14.2	11.2	13.0	14.6	11.7	8.4	11.8	5.9
1980	12.6	15.0	10.8	13.0	15.3	11.2	8.8	10.9	7.2
1981	13.6	15.9	11.8	14.0	16.3	12.3	7.4	10.5	5.0
1982	13.4	15.6	11.6	14.0	16.4	12.0	10.3	10.2	10.0
1983	14.0	16.6	11.9	14.6	17.5	12.4	10.3	10.9	9.6
1984	15.2	18.3	12.8	15.9	19.1	13.5	9.4	12.5	6.9
1985	15.5	18.4	13.2	16.3	19.2	13.9	9.9	12.0	8.3
1986	15.9	19.4	13.0	16.8	20.5	13.7	10.6	13.1	8.5
1987	16.7	20.8	13.4	17.7	22.1	14.1	10.4	10.4	10.1
1988	17.2	21.3	14.1	18.2	22.4	14.9	11.6	15.8	8.5
1989	17.3	21.6	13.8	18.3	23.0	14.5	11.6	14.2	9.8
1990	18.5	23.0	14.8	19.4	24.0	15.5	13.8	17.5	10.9
1991	18.8	23.7	14.6	19.7	24.8	15.3	14.0	18.8	10.3
1992	18.6	23.0	15.0	19.6	23.9	15.9	13.6	18.7	9.6
1993	18.8	23.5	15.0	19.6	24.3	15.7	14.0	19.3	9.6
1994	19.9	24.9	15.8	20.6	25.6	16.5	14.3	21.4	8.6
1995	19.9	25.3	15.3	20.6	26.0	15.9	16.5	23.4	11.3
1996	19.3	24.2	15.4	20.2	25.2	16.1	15.1	18.9	11.9
1997	19.9	24.4	16.4	20.8	25.1	17.4	16.1	23.7	10.9
1998	19.4	22.8	16.6	20.4	23.9	17.6	13.9	16.3	12.0
1999	19.8	24.4	16.1	20.7	25.3	17.1	13.8	17.8	10.8
2000	19.6	24.1	15.9	20.7	25.3	16.9	13.9	18.2	10.6
2001	19.7	24.1	16.1	20.5	25.2	16.7	15.4	17.7	13.3
2002	19.8	23.7	16.7	20.9	25.0	17.6	14.1	16.9	12.0
2003	20.2	24.1	17.1	21.2	25.3	18.1	15.5	18.8	13.1
2004	20.4	24.7	17.1	21.3	25.5	17.9	17.4	22.8	13.4
1975-2004	17.2	21.0	14.2	17.9	21.8	14.8	12.6	16.1	9.8

<sup>a</sup> SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta). Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).  
- Statistic not shown. Rate based on less than 16 cases for the time interval.

Table XIX-3  
NON-HODGKIN LYMPHOMA

AGE-ADJUSTED U.S. DEATH<sup>a</sup> RATES BY YEAR, RACE AND SEX

	All Races			Whites			Blacks		
	Total	Males	Females	Total	Males	Females	Total	Males	Females
<u>AGE-ADJUSTED RATES</u>									
YEAR OF DEATH:									
1975	5.6	6.9	4.6	5.8	7.2	4.8	3.7	4.8	2.9
1976	5.7	6.9	4.8	5.9	7.1	5.0	3.5	4.5	2.8
1977	5.8	6.9	4.9	6.0	7.1	5.1	3.9	4.9	3.1
1978	5.9	7.3	4.9	6.2	7.5	5.1	3.9	5.4	2.8
1979	5.9	7.4	4.9	6.2	7.7	5.1	3.6	4.5	3.0
1980	6.2	7.5	5.3	6.5	7.7	5.5	4.0	5.1	3.1
1981	6.2	7.4	5.3	6.4	7.7	5.5	4.0	5.0	3.3
1982	6.6	8.0	5.5	6.8	8.3	5.7	4.1	5.2	3.3
1983	6.7	8.1	5.6	6.9	8.4	5.8	4.3	5.8	3.2
1984	6.8	8.2	5.8	7.1	8.5	6.0	4.2	5.3	3.4
1985	7.1	8.8	5.8	7.3	9.2	6.0	4.6	5.6	3.9
1986	7.3	9.0	6.0	7.6	9.3	6.3	4.9	6.6	3.7
1987	7.3	8.8	6.1	7.6	9.1	6.4	4.7	6.3	3.6
1988	7.5	9.3	6.2	7.8	9.6	6.5	5.0	6.8	3.7
1989	7.8	9.7	6.4	8.1	10.0	6.7	5.3	7.1	4.1
1990	7.9	10.0	6.3	8.2	10.3	6.6	5.4	7.1	4.2
1991	8.2	10.1	6.7	8.5	10.5	7.0	5.7	7.2	4.6
1992	8.2	10.3	6.7	8.5	10.7	7.0	5.6	7.2	4.5
1993	8.2	10.1	6.9	8.6	10.5	7.2	5.7	7.5	4.4
1994	8.6	10.8	7.1	9.0	11.2	7.4	5.5	6.8	4.6
1995	8.7	10.8	7.2	9.1	11.2	7.5	5.8	7.6	4.5
1996	8.8	10.9	7.1	9.1	11.3	7.5	6.2	8.2	4.7
1997	8.9	11.0	7.3	9.2	11.4	7.6	6.3	8.4	4.9
1998	8.7	10.9	7.1	9.0	11.3	7.4	6.0	7.5	4.9
1999	8.3	10.4	6.9	8.7	10.7	7.2	5.7	7.4	4.4
2000	8.2	10.2	6.7	8.5	10.6	7.0	5.2	6.6	4.2
2001	7.9	9.9	6.4	8.2	10.3	6.7	5.6	7.3	4.5
2002	7.6	9.6	6.2	7.9	10.0	6.4	5.4	6.9	4.4
2003	7.3	9.3	5.9	7.6	9.7	6.1	4.9	5.8	4.3
2004	7.0	8.8	5.7	7.3	9.2	5.9	5.0	6.2	4.1
1975-2004	7.5	9.3	6.2	7.8	9.7	6.5	5.1	6.5	4.0

<sup>a</sup> NCHS public use data file for the total US. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

- Statistic not shown. Rate based on less than 16 cases for the time interval.

Table XX-2  
CANCER OF THE ORAL CAVITY AND PHARYNX (Invasive)

AGE ADJUSTED SEER INCIDENCE<sup>a</sup> RATES BY YEAR, RACE AND SEX

	<u>All Races</u>			<u>Whites</u>			<u>Blacks</u>		
	<u>Total</u>	<u>Males</u>	<u>Females</u>	<u>Total</u>	<u>Males</u>	<u>Females</u>	<u>Total</u>	<u>Males</u>	<u>Females</u>
<u>AGE-ADJUSTED RATES</u>									
<u>YEAR OF DIAGNOSIS:</u>									
1975	13.2	21.2	7.1	13.3	21.7	6.9	13.4	20.2	8.2
1976	13.3	21.0	7.4	13.2	21.0	7.4	15.2	23.8	8.4
1977	12.7	20.1	7.0	12.6	19.9	7.0	14.4	23.9	6.6
1978	13.4	20.9	7.7	13.2	20.8	7.6	15.8	26.0	8.1
1979	14.0	22.0	8.0	13.7	21.6	7.7	18.4	28.2	10.6
1980	13.3	20.3	7.9	13.0	20.0	7.6	16.8	25.4	9.6
1981	13.6	20.8	7.9	13.4	20.5	7.9	16.7	28.2	7.5
1982	13.3	20.3	7.9	13.1	19.9	7.9	17.1	27.4	8.9
1983	13.3	21.2	7.2	13.3	21.2	7.3	16.1	26.1	8.3
1984	13.5	20.8	7.9	13.0	20.1	7.6	18.9	30.0	10.1
1985	13.3	19.8	8.2	13.3	19.6	8.3	16.0	25.7	8.5
1986	12.5	19.1	7.2	12.3	18.9	7.3	16.2	28.1	7.0
1987	13.2	20.5	7.5	13.1	20.2	7.7	16.8	30.0	7.0
1988	12.1	18.1	7.3	11.8	17.7	7.1	16.4	25.6	9.1
1989	12.1	18.3	7.3	11.8	17.7	7.3	15.7	27.2	7.0
1990	12.6	19.4	7.3	12.4	18.8	7.4	16.0	27.8	6.9
1991	12.2	18.3	7.2	12.1	18.1	7.3	14.4	23.7	7.3
1992	12.0	18.3	6.9	11.8	18.0	6.9	14.8	25.1	6.9
1993	12.4	18.7	7.3	12.2	18.4	7.2	15.5	24.9	8.1
1994	11.7	17.7	6.8	11.4	16.8	6.9	15.7	26.8	7.1
1995	11.5	16.9	7.2	11.5	16.7	7.3	13.5	22.5	6.7
1996	11.7	17.7	7.0	11.4	17.1	6.9	14.9	24.3	7.5
1997	11.5	17.3	6.8	11.5	17.1	6.9	11.4	18.2	6.2
1998	11.2	16.6	6.7	11.2	16.4	6.9	13.0	22.4	6.2
1999	10.5	15.5	6.4	10.4	15.4	6.3	11.6	19.5	5.8
2000	10.7	16.0	6.3	10.7	15.6	6.4	11.6	20.6	4.8
2001	10.7	15.3	6.9	10.8	15.4	6.9	11.9	18.8	6.9
2002	11.0	16.1	6.6	11.2	16.3	6.7	11.0	16.8	6.5
2003	10.3	15.3	6.1	10.3	15.3	5.9	11.4	16.3	7.6
2004	10.4	15.5	6.0	10.6	15.7	6.0	9.9	15.7	5.6
1975-2004	12.1	18.3	7.1	12.0	18.2	7.1	14.2	23.1	7.2

<sup>a</sup> SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta). Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

- Statistic not shown. Rate based on less than 16 cases for the time interval.

Table XX-3  
CANCER OF THE ORAL CAVITY AND PHARYNX (Invasive)  
 AGE-ADJUSTED U.S. DEATH<sup>a</sup> RATES BY YEAR, RACE AND SEX

	<u>All Races</u>			<u>Whites</u>			<u>Blacks</u>		
	<u>Total</u>	<u>Males</u>	<u>Females</u>	<u>Total</u>	<u>Males</u>	<u>Females</u>	<u>Total</u>	<u>Males</u>	<u>Females</u>
<u>AGE-ADJUSTED RATES</u>									
<u>YEAR OF DEATH:</u>									
1975	4.3	6.9	2.3	4.1	6.6	2.3	5.8	9.9	2.5
1976	4.2	6.9	2.2	4.0	6.6	2.2	6.1	10.6	2.6
1977	4.3	7.0	2.3	4.1	6.7	2.2	6.2	10.9	2.6
1978	4.2	6.8	2.3	4.0	6.4	2.2	6.1	10.7	2.7
1979	4.2	6.7	2.4	4.0	6.3	2.3	6.6	11.5	3.0
1980	4.2	6.7	2.3	3.9	6.1	2.2	7.0	12.5	2.8
1981	4.1	6.5	2.2	3.8	6.0	2.2	6.5	11.4	2.8
1982	4.0	6.5	2.2	3.7	6.0	2.1	6.6	11.5	2.9
1983	3.9	6.3	2.2	3.6	5.7	2.1	6.6	11.8	2.8
1984	3.9	6.2	2.2	3.6	5.6	2.1	6.4	11.7	2.4
1985	3.8	5.9	2.1	3.5	5.4	2.0	6.2	11.0	2.6
1986	3.7	5.8	2.2	3.4	5.3	2.1	6.5	11.4	2.9
1987	3.5	5.5	2.1	3.3	5.0	2.0	5.7	10.0	2.5
1988	3.6	5.6	2.0	3.3	5.1	2.0	6.0	10.7	2.5
1989	3.5	5.3	2.1	3.2	4.8	2.0	5.9	10.4	2.6
1990	3.6	5.6	2.0	3.3	5.1	2.0	6.1	11.0	2.5
1991	3.5	5.3	2.0	3.2	4.9	2.0	5.8	10.1	2.7
1992	3.3	5.1	1.9	3.1	4.7	1.9	5.6	10.0	2.3
1993	3.3	5.2	1.9	3.1	4.7	1.8	5.7	10.1	2.4
1994	3.2	4.8	1.8	2.9	4.4	1.8	5.3	9.3	2.4
1995	3.2	4.9	1.8	2.9	4.4	1.8	5.4	9.3	2.5
1996	3.0	4.6	1.8	2.8	4.2	1.7	4.9	8.7	2.1
1997	3.0	4.5	1.8	2.8	4.2	1.7	4.7	8.1	2.2
1998	3.0	4.5	1.7	2.8	4.2	1.7	4.5	7.9	2.1
1999	2.7	4.1	1.6	2.5	3.8	1.5	4.4	7.5	2.1
2000	2.7	4.1	1.6	2.5	3.7	1.6	4.1	7.3	1.8
2001	2.7	4.1	1.6	2.6	3.9	1.5	3.9	6.7	1.8
2002	2.7	4.1	1.5	2.6	3.9	1.5	3.7	6.3	1.7
2003	2.6	4.1	1.5	2.5	3.8	1.4	3.9	6.9	1.6
2004	2.6	4.0	1.5	2.5	3.7	1.5	3.8	6.9	1.6
1975-2004	3.4	5.3	1.9	3.2	4.9	1.9	5.3	9.4	2.3

<sup>a</sup> NCHS public use data file for the total US. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

- Statistic not shown. Rate based on less than 16 cases for the time interval.

Table XXI-2  
OVARY CANCER (Invasive)

AGE-ADJUSTED SEER INCIDENCE<sup>a</sup> RATES BY YEAR, RACE AND AGE

	<u>All Races, Females</u>			<u>White Females</u>			<u>Black Females</u>		
	All	<65	65+	All	<65	65+	All	<65	65+
<u>SEER INCIDENCE RATES</u>									
<u>YEAR OF DIAGNOSIS:</u>									
1975	16.3	11.6	48.9	16.8	12.0	49.8	11.7	7.4	41.6
1976	15.9	11.2	47.9	16.6	11.9	49.5	11.4	7.4	39.3
1977	15.6	11.2	46.2	16.2	11.6	48.1	11.3	8.7	29.7
1978	15.4	10.8	47.2	16.2	11.4	49.3	10.0	6.1	36.8
1979	15.4	10.7	47.8	15.8	10.9	49.9	11.9	9.1	31.8
1980	15.4	10.6	48.6	16.2	11.2	50.9	11.7	7.4	41.6
1981	15.4	10.5	49.5	16.0	10.9	51.5	11.7	7.6	40.1
1982	15.6	10.1	53.4	16.1	10.5	55.1	12.3	7.4	46.2
1983	16.0	10.4	54.2	16.4	10.7	55.8	13.1	9.2	39.9
1984	16.2	11.0	52.8	17.1	11.7	54.7	11.0	5.9	46.3
1985	16.5	10.8	56.3	17.4	11.4	59.1	11.3	7.3	39.3
1986	15.0	9.9	50.4	15.7	10.5	51.8	10.6	6.2	41.5
1987	16.2	10.2	57.6	17.0	10.9	59.8	11.8	6.3	50.3
1988	15.3	9.6	54.7	16.0	10.1	57.0	11.6	6.6	45.6
1989	15.5	9.5	56.8	16.5	10.3	59.5	10.9	5.4	49.5
1990	15.4	9.4	56.5	16.3	10.1	59.2	10.1	5.6	41.7
1991	15.7	9.7	57.7	16.5	10.3	59.7	11.0	5.5	49.5
1992	14.9	9.3	53.4	15.6	9.7	56.6	11.2	7.8	35.2
1993	15.2	9.1	57.4	16.0	9.6	60.4	11.5	6.4	46.3
1994	14.5	8.9	53.1	15.0	9.2	55.3	12.4	6.5	53.5
1995	14.6	8.7	54.8	15.3	9.3	57.0	10.3	4.9	47.7
1996	14.1	8.4	53.6	15.3	9.2	57.1	9.0	4.5	40.1
1997	14.7	9.2	52.7	15.4	9.6	55.1	10.6	5.6	44.9
1998	14.3	8.8	52.3	15.0	9.2	55.3	10.7	6.0	43.4
1999	14.6	8.7	55.8	15.4	8.9	60.1	10.7	6.3	41.4
2000	14.2	9.0	50.3	15.1	9.6	53.4	10.6	5.4	46.5
2001	14.4	9.1	51.2	15.4	9.7	55.4	9.4	6.2	31.2
2002	13.5	7.8	53.3	14.2	8.2	55.8	10.2	5.5	42.7
2003	13.3	8.2	48.4	13.8	8.6	49.3	11.8	5.9	52.4
2004	12.6	7.7	47.0	13.3	7.8	50.8	9.5	5.9	34.0
1975-2004	15.0	9.5	52.7	15.7	10.0	55.1	11.0	6.4	42.8

Ovary excludes borderline cases or histologies 8442, 8451, 8462, 8472, and 8473.

<sup>a</sup> SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta). Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

- Statistic not shown. Rate based on less than 16 cases for the time interval.



Table XXI-3  
OVARY CANCER (Invasive)

AGE-ADJUSTED U.S. DEATH<sup>a</sup> RATES BY YEAR, RACE AND AGE

	<u>All Races, Females</u>			<u>White Females</u>			<u>Black Females</u>		
	<u>All</u>	<u>&lt;65</u>	<u>65+</u>	<u>All</u>	<u>&lt;65</u>	<u>65+</u>	<u>All</u>	<u>&lt;65</u>	<u>65+</u>
<u>U.S. DEATH RATES</u>									
<u>YEAR OF DEATH:</u>									
1975	9.8	5.8	37.5	10.2	6.0	38.8	7.2	4.6	24.6
1976	10.0	5.7	39.7	10.3	5.9	40.6	8.0	4.6	31.5
1977	9.6	5.5	38.0	9.9	5.6	39.1	7.4	4.3	29.2
1978	9.6	5.4	38.0	9.8	5.6	38.9	7.4	4.2	29.3
1979	9.3	5.1	38.6	9.6	5.2	39.7	7.5	4.4	28.9
1980	9.3	5.0	39.1	9.5	5.1	40.0	7.9	4.2	33.2
1981	9.2	4.9	38.9	9.5	5.1	39.8	7.4	3.7	32.4
1982	9.2	4.8	39.8	9.5	5.0	40.7	7.4	3.6	33.5
1983	9.2	4.7	40.3	9.4	4.8	41.4	7.7	4.1	32.3
1984	9.1	4.6	40.1	9.4	4.8	40.9	7.6	3.8	33.8
1985	9.1	4.6	40.2	9.3	4.7	41.3	7.3	3.7	31.9
1986	9.2	4.6	41.6	9.5	4.7	42.6	7.4	3.7	33.4
1987	9.2	4.5	41.9	9.5	4.6	42.8	7.6	3.5	35.5
1988	9.3	4.4	43.2	9.6	4.6	44.1	7.9	3.7	36.9
1989	9.2	4.3	43.2	9.5	4.5	44.3	7.4	3.3	36.2
1990	9.3	4.3	44.4	9.6	4.5	45.2	7.9	3.3	39.2
1991	9.5	4.3	45.4	9.7	4.5	46.1	8.4	3.4	42.5
1992	9.5	4.2	45.5	9.7	4.4	46.5	8.2	3.6	40.0
1993	9.1	4.0	44.1	9.3	4.2	44.9	7.7	3.0	40.4
1994	9.4	4.1	45.6	9.7	4.3	46.7	7.8	3.2	39.7
1995	9.1	4.0	44.8	9.5	4.2	46.1	7.4	2.9	38.0
1996	8.9	3.8	43.7	9.1	4.0	44.8	7.3	2.9	37.9
1997	8.9	3.8	44.7	9.2	3.9	45.6	7.9	3.1	41.2
1998	8.7	3.7	43.6	9.0	3.8	45.0	7.4	3.2	36.1
1999	8.8	3.7	43.6	9.1	3.9	44.8	7.4	3.1	37.3
2000	8.9	3.7	45.0	9.2	3.8	46.6	7.1	2.9	36.5
2001	9.0	3.7	45.4	9.3	3.9	46.9	7.5	3.2	37.4
2002	9.0	3.8	45.2	9.4	4.0	46.7	7.4	3.1	37.3
2003	8.8	3.7	44.6	9.2	3.8	46.1	7.5	3.1	37.4
2004	8.8	3.5	44.8	9.1	3.6	46.4	7.6	3.1	38.9
1975-2004	9.2	4.3	42.9	9.5	4.5	44.0	7.6	3.5	36.1

<sup>a</sup> NCHS public use data file for the total US. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

- Statistic not shown. Rate based on less than 16 cases for the time interval.

Table XXII-2  
PANCREAS CANCER (Invasive)

AGE ADJUSTED SEER INCIDENCE<sup>a</sup> RATES BY YEAR, RACE AND SEX

	<u>All Races</u>			<u>Whites</u>			<u>Blacks</u>		
	<u>Total</u>	<u>Males</u>	<u>Females</u>	<u>Total</u>	<u>Males</u>	<u>Females</u>	<u>Total</u>	<u>Males</u>	<u>Females</u>
<u>AGE-ADJUSTED RATES</u>									
<u>YEAR OF DIAGNOSIS:</u>									
1975	11.8	15.6	9.1	11.7	15.6	8.9	15.9	18.1	13.9
1976	12.0	14.8	9.9	11.8	14.5	9.8	17.6	22.7	14.0
1977	11.9	14.8	9.8	11.6	14.6	9.5	17.1	19.4	15.2
1978	11.1	14.1	8.9	10.8	13.9	8.6	15.1	20.0	11.5
1979	11.5	14.2	9.7	11.2	13.8	9.4	16.1	19.1	13.7
1980	11.5	14.2	9.6	11.1	13.7	9.2	17.9	21.8	15.5
1981	11.7	13.5	10.2	11.4	13.3	9.9	17.2	20.9	14.1
1982	11.7	13.8	10.2	11.4	13.5	9.9	16.9	21.0	14.0
1983	12.2	14.7	10.4	11.7	14.1	10.1	18.1	23.1	14.9
1984	12.2	13.9	10.9	12.0	13.9	10.6	17.2	18.4	16.3
1985	12.1	14.3	10.5	11.7	13.6	10.3	18.7	25.0	14.6
1986	11.7	13.8	10.2	11.4	13.5	9.8	17.3	20.2	15.6
1987	11.6	13.8	10.1	11.1	13.4	9.5	18.4	19.0	17.9
1988	11.7	13.7	10.2	11.1	13.4	9.6	18.5	19.4	17.4
1989	11.3	12.9	10.1	11.0	12.7	9.6	15.0	16.0	14.0
1990	11.3	12.8	10.1	11.0	12.5	9.8	15.2	18.8	12.6
1991	11.4	12.9	10.2	11.0	12.6	9.8	17.0	18.2	16.2
1992	11.6	13.2	10.3	11.3	12.9	10.0	17.3	19.8	15.8
1993	11.0	12.5	9.7	10.5	12.0	9.3	16.8	18.7	15.1
1994	11.4	13.3	9.9	10.9	12.3	9.7	17.4	21.5	14.5
1995	11.1	12.7	10.0	10.7	12.3	9.6	18.1	20.3	16.3
1996	11.3	12.9	9.9	10.9	12.5	9.6	17.1	19.8	14.9
1997	11.5	13.0	10.2	11.0	12.5	9.7	17.9	19.4	16.6
1998	11.5	13.1	10.2	11.2	13.0	9.8	14.7	16.2	13.9
1999	11.1	12.6	9.8	10.7	12.2	9.3	16.7	19.5	14.7
2000	11.4	13.3	9.8	11.1	12.9	9.6	15.7	18.3	13.2
2001	11.3	13.1	9.9	11.0	13.0	9.5	14.1	15.2	13.2
2002	11.6	13.0	10.4	11.5	13.2	10.0	15.8	14.9	15.8
2003	11.4	12.6	10.4	11.2	12.4	10.1	15.8	16.6	15.2
2004	11.4	13.3	9.8	11.3	13.0	9.8	15.1	18.3	12.5
1975-2004	11.5	13.4	10.0	11.2	13.1	9.7	16.6	19.0	14.8

<sup>a</sup> SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta). Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).  
 - Statistic not shown. Rate based on less than 16 cases for the time interval.

Table XXII-3  
PANCREAS CANCER (Invasive)

AGE-ADJUSTED U.S. DEATH<sup>a</sup> RATES BY YEAR, RACE AND SEX

	<u>All Races</u>			<u>Whites</u>			<u>Blacks</u>		
	<u>Total</u>	<u>Males</u>	<u>Females</u>	<u>Total</u>	<u>Males</u>	<u>Females</u>	<u>Total</u>	<u>Males</u>	<u>Females</u>
<u>AGE-ADJUSTED RATES</u>									
<u>YEAR OF DEATH:</u>									
1975	10.7	13.8	8.4	10.5	13.6	8.3	12.8	16.3	10.1
1976	10.7	13.6	8.6	10.5	13.4	8.4	12.9	16.0	10.5
1977	10.9	13.9	8.8	10.7	13.6	8.6	13.2	17.1	10.3
1978	10.8	13.6	8.8	10.6	13.5	8.6	12.9	15.9	10.7
1979	10.7	13.4	8.8	10.5	13.3	8.6	13.4	15.9	11.6
1980	10.6	13.1	8.8	10.4	12.8	8.6	13.8	16.5	11.8
1981	10.7	13.2	8.9	10.5	12.9	8.8	13.4	17.0	10.8
1982	10.6	13.1	8.8	10.3	12.8	8.6	14.1	17.1	11.9
1983	10.7	12.8	9.1	10.4	12.4	8.9	14.8	18.1	12.5
1984	10.8	13.0	9.2	10.5	12.6	8.9	15.0	17.5	13.1
1985	10.6	12.9	9.0	10.3	12.5	8.8	14.4	17.3	12.4
1986	10.6	12.5	9.2	10.3	12.2	8.9	14.6	16.7	13.0
1987	10.6	12.5	9.2	10.3	12.2	8.9	14.6	17.4	12.6
1988	10.5	12.5	9.0	10.1	12.2	8.6	14.5	17.1	12.7
1989	10.6	12.6	9.2	10.3	12.2	8.8	15.0	17.7	13.2
1990	10.7	12.6	9.3	10.3	12.2	8.9	15.2	18.1	13.2
1991	10.7	12.6	9.3	10.4	12.3	9.0	14.5	16.9	12.8
1992	10.7	12.6	9.3	10.3	12.2	8.9	15.4	18.2	13.4
1993	10.7	12.4	9.4	10.3	12.1	9.0	15.0	17.6	13.3
1994	10.6	12.4	9.3	10.3	12.0	9.0	15.2	17.7	13.4
1995	10.4	12.1	9.2	10.2	11.9	8.9	13.9	15.8	12.5
1996	10.5	12.1	9.2	10.2	11.8	8.9	14.4	16.4	13.0
1997	10.5	12.2	9.1	10.2	12.0	8.8	14.6	16.4	13.1
1998	10.5	12.3	9.2	10.2	12.0	8.8	14.5	16.5	13.1
1999	10.6	12.3	9.3	10.4	12.1	9.0	14.2	16.6	12.4
2000	10.5	12.1	9.3	10.3	12.0	9.0	14.0	15.8	12.7
2001	10.5	12.1	9.3	10.4	12.0	9.0	13.7	15.1	12.5
2002	10.5	12.2	9.2	10.3	12.0	8.9	13.7	15.4	12.3
2003	10.5	12.0	9.2	10.3	11.9	9.0	13.8	15.5	12.4
2004	10.6	12.3	9.2	10.4	12.2	9.0	13.6	15.6	12.1
1975-2004	10.6	12.6	9.1	10.4	12.4	8.9	14.3	16.7	12.5

<sup>a</sup> NCHS public use data file for the total US. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

- Statistic not shown. Rate based on less than 16 cases for the time interval.

Table XXIII-2  
PROSTATE CANCER (Invasive)

AGE ADJUSTED SEER INCIDENCE<sup>a</sup> RATES BY YEAR, RACE AND AGE

	<u>All Races, Males</u>			<u>White Males</u>			<u>Black Males</u>		
	<u>All</u>	<u>&lt;65</u>	<u>65+</u>	<u>All</u>	<u>&lt;65</u>	<u>65+</u>	<u>All</u>	<u>&lt;65</u>	<u>65+</u>
<u>SEER INCIDENCE RATES</u>									
<u>YEAR OF DIAGNOSIS:</u>									
1975	94.0	13.6	649.5	92.2	13.0	639.6	141.0	26.8	930.5
1976	97.9	14.6	674.1	97.3	13.9	673.9	140.8	29.4	910.9
1977	100.5	15.0	691.0	98.6	14.4	680.7	159.2	29.2	1,057.2
1978	99.4	15.2	681.2	97.7	14.8	671.1	148.1	26.8	986.4
1979	103.4	14.9	715.3	102.3	14.6	708.4	162.2	26.2	1,102.1
1980	106.0	15.5	731.7	104.9	14.6	729.1	161.0	33.6	1,041.4
1981	108.9	16.8	745.5	107.9	15.9	743.6	161.7	34.4	1,041.9
1982	108.2	16.3	743.3	107.3	15.7	740.1	167.6	31.3	1,109.8
1983	111.6	17.1	764.6	110.8	16.6	762.2	170.8	34.1	1,116.4
1984	111.7	17.2	764.4	110.1	16.2	759.0	178.8	37.1	1,158.1
1985	115.5	17.8	790.5	114.6	17.5	785.6	170.2	31.4	1,129.4
1986	119.0	18.5	813.8	119.0	18.2	815.2	168.0	33.8	1,095.3
1987	133.6	21.5	908.7	134.4	21.1	917.8	189.2	36.1	1,247.6
1988	137.6	22.2	935.3	138.7	22.3	943.1	190.1	34.6	1,264.9
1989	145.3	24.1	983.0	146.0	23.9	989.7	192.0	37.3	1,261.3
1990	170.8	28.7	1,153.0	172.6	28.8	1,166.7	221.4	44.3	1,446.0
1991	214.6	38.8	1,429.5	216.1	39.2	1,439.3	288.5	57.2	1,887.7
1992	237.1	49.3	1,534.6	237.8	49.4	1,540.2	327.9	77.0	2,062.0
1993	209.1	50.4	1,306.3	204.1	49.0	1,276.2	343.1	93.6	2,067.8
1994	180.0	48.5	1,088.9	174.0	46.8	1,053.5	311.5	94.2	1,813.6
1995	168.6	49.9	989.5	163.7	48.0	963.6	279.8	97.7	1,538.1
1996	168.6	53.2	966.5	164.1	51.7	941.1	281.1	100.1	1,532.6
1997	172.7	55.0	986.6	169.0	53.8	965.2	279.7	96.1	1,548.9
1998	169.8	54.7	965.8	166.0	52.8	948.9	282.1	101.6	1,529.9
1999	182.5	60.6	1,024.8	177.7	58.8	1,000.0	286.7	110.4	1,505.6
2000	181.3	61.4	1,009.9	177.5	59.5	993.5	287.9	112.7	1,498.8
2001	182.8	63.8	1,005.4	180.8	61.9	1,002.4	265.8	113.0	1,322.1
2002	179.6	64.5	975.5	176.5	62.7	963.1	282.2	115.1	1,437.4
2003	166.6	60.6	898.9	162.3	58.1	882.5	252.2	113.4	1,211.8
2004	159.5	59.5	850.8	155.3	57.4	832.2	238.2	100.8	1,187.6
1975-2004	150.7	37.3	934.9	147.9	35.7	923.3	232.4	69.8	1,356.4

<sup>a</sup> SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta). Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

- Statistic not shown. Rate based on less than 16 cases for the time interval.

Table XXIII-3  
PROSTATE CANCER (Invasive)

AGE ADJUSTED U.S. DEATH<sup>a</sup> RATES BY YEAR, RACE AND AGE

	<u>All Races, Males</u>			<u>White Males</u>			<u>Black Males</u>		
	<u>All</u>	<u>&lt;65</u>	<u>65+</u>	<u>All</u>	<u>&lt;65</u>	<u>65+</u>	<u>All</u>	<u>&lt;65</u>	<u>65+</u>
<u>U.S. DEATH RATES</u>									
<u>YEAR OF DEATH:</u>									
1975	31.0	2.5	227.5	29.1	2.2	214.9	55.5	6.6	394.0
1976	31.8	2.7	232.9	29.9	2.3	220.8	55.6	6.9	392.2
1977	31.8	2.7	233.3	29.9	2.3	220.6	56.4	6.9	398.7
1978	32.7	2.6	240.3	30.6	2.2	226.8	58.5	6.8	416.1
1979	32.8	2.6	242.1	30.8	2.2	228.2	59.2	6.6	422.9
1980	33.1	2.7	243.1	30.8	2.3	227.4	62.0	6.6	445.0
1981	33.2	2.7	244.0	30.8	2.3	227.5	63.6	7.1	454.4
1982	33.4	2.7	245.5	31.1	2.3	229.8	62.4	6.6	447.7
1983	33.9	2.8	249.3	31.6	2.4	233.2	64.6	7.0	462.5
1984	34.1	2.7	250.8	31.6	2.4	233.5	65.9	6.7	475.1
1985	33.9	2.7	249.4	31.3	2.3	231.3	67.3	7.2	482.7
1986	34.9	2.8	257.0	32.5	2.5	239.7	67.4	6.6	487.4
1987	35.1	2.7	259.1	32.6	2.4	241.3	68.9	6.7	499.0
1988	35.9	2.8	264.4	33.4	2.5	247.1	69.6	7.0	502.3
1989	37.1	3.0	273.0	34.5	2.6	255.2	71.7	7.1	518.5
1990	38.6	3.0	284.2	35.7	2.6	264.0	78.0	7.4	565.6
1991	39.3	2.9	291.1	36.5	2.5	271.4	78.2	7.1	570.2
1992	39.2	2.9	290.6	36.3	2.4	270.3	79.8	7.7	578.7
1993	39.3	2.7	292.5	36.3	2.3	270.8	81.9	7.0	599.1
1994	38.5	2.7	286.5	35.6	2.3	266.0	80.0	6.8	585.9
1995	37.3	2.6	277.2	34.4	2.2	256.9	78.2	6.7	572.5
1996	36.0	2.5	267.6	33.0	2.1	246.4	78.8	6.5	579.2
1997	34.2	2.3	254.3	31.3	1.9	234.6	74.3	6.3	544.2
1998	32.6	2.3	242.5	29.9	1.9	223.4	72.8	6.3	532.4
1999	31.6	2.1	235.5	28.9	1.7	216.9	70.1	5.8	514.8
2000	30.3	2.0	226.1	27.7	1.7	207.7	68.7	5.4	505.9
2001	29.2	1.9	217.4	26.7	1.6	199.7	66.5	5.4	489.0
2002	28.2	1.9	209.7	25.9	1.6	193.9	63.0	5.4	461.3
2003	26.6	1.9	197.0	24.5	1.6	182.7	58.0	5.4	421.4
2004	25.4	1.8	188.7	23.4	1.6	174.6	56.1	5.0	409.5
1975-2004	33.6	2.5	248.4	31.1	2.2	231.0	68.4	6.5	496.7

<sup>a</sup> NCHS public use data file for the total US. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

- Statistic not shown. Rate based on less than 16 cases for the time interval.

Table XXIV-2  
STOMACH CANCER (Invasive)

AGE ADJUSTED SEER INCIDENCE<sup>a</sup> RATES BY YEAR, RACE AND SEX

	<u>All Races</u>			<u>Whites</u>			<u>Blacks</u>		
	<u>Total</u>	<u>Males</u>	<u>Females</u>	<u>Total</u>	<u>Males</u>	<u>Females</u>	<u>Total</u>	<u>Males</u>	<u>Females</u>
<u>AGE-ADJUSTED RATES</u>									
<u>YEAR OF DIAGNOSIS:</u>									
1975	11.7	17.1	7.8	10.7	16.0	7.0	17.0	24.8	11.5
1976	12.2	17.5	8.3	11.2	16.0	7.7	15.8	23.5	10.1
1977	11.5	16.6	7.9	10.3	15.0	7.0	17.3	24.7	11.6
1978	11.5	16.5	7.9	10.3	15.0	7.0	17.6	22.9	13.5
1979	12.0	17.4	8.2	10.8	15.6	7.5	18.5	28.2	11.7
1980	11.3	17.1	7.2	10.3	15.7	6.6	15.1	26.3	7.3
1981	11.1	16.6	7.4	10.1	15.1	6.7	17.6	27.8	10.4
1982	10.9	16.3	7.1	9.7	14.6	6.3	19.0	31.2	10.3
1983	10.9	15.8	7.5	9.5	13.7	6.6	18.0	27.8	11.2
1984	10.6	15.3	7.3	9.4	13.9	6.4	16.5	21.8	12.3
1985	10.2	15.1	6.9	8.9	13.3	6.0	16.4	23.5	11.3
1986	10.2	15.0	6.8	9.1	13.5	5.9	16.5	23.2	11.9
1987	10.2	15.1	6.8	9.0	13.2	6.0	17.1	26.7	10.7
1988	10.2	15.5	6.5	8.8	13.6	5.5	16.2	24.8	10.1
1989	10.0	15.2	6.3	8.7	13.5	5.3	15.8	23.2	10.7
1990	9.3	13.7	6.2	8.0	11.9	5.2	14.4	21.8	9.4
1991	9.7	14.2	6.6	8.3	12.3	5.5	16.5	25.6	10.6
1992	9.2	13.3	6.1	7.9	11.6	5.2	14.0	19.9	9.8
1993	9.0	13.6	5.8	7.7	11.6	4.9	14.4	23.7	8.4
1994	9.0	13.6	5.7	7.7	11.8	4.8	15.0	24.7	8.8
1995	8.3	12.6	5.2	7.3	11.3	4.4	13.3	18.8	9.5
1996	8.5	12.6	5.5	7.3	11.0	4.7	14.5	22.6	8.9
1997	8.6	12.7	5.6	7.2	10.8	4.5	15.6	22.1	10.9
1998	8.6	12.2	5.9	7.2	10.5	4.7	14.9	20.1	11.4
1999	8.6	12.1	6.0	7.4	10.5	5.0	13.0	18.2	9.8
2000	8.1	11.7	5.4	6.8	9.9	4.5	12.8	19.8	8.1
2001	7.7	11.2	5.1	6.4	9.6	4.0	12.9	18.3	9.0
2002	7.9	11.2	5.4	6.6	9.8	4.3	12.7	15.4	10.6
2003	7.6	10.7	5.3	6.5	9.3	4.3	12.8	17.5	9.7
2004	7.6	10.9	5.1	6.7	9.7	4.5	10.5	15.8	7.1
1975-2004	9.5	13.9	6.3	8.3	12.3	5.4	15.0	22.2	10.1

<sup>a</sup> SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta). Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).  
 - Statistic not shown. Rate based on less than 16 cases for the time interval.

Table XXIV-3  
STOMACH CANCER (Invasive)

AGE-ADJUSTED U.S. DEATH<sup>a</sup> RATES BY YEAR, RACE AND SEX

	<u>All Races</u>			<u>Whites</u>			<u>Blacks</u>		
	<u>Total</u>	<u>Males</u>	<u>Females</u>	<u>Total</u>	<u>Males</u>	<u>Females</u>	<u>Total</u>	<u>Males</u>	<u>Females</u>
<u>AGE-ADJUSTED RATES</u>									
<u>YEAR OF DEATH:</u>									
1975	8.5	12.3	5.9	7.9	11.4	5.6	13.9	21.0	8.6
1976	8.3	12.0	5.7	7.7	11.2	5.4	13.5	20.2	8.6
1977	7.9	11.5	5.4	7.3	10.7	5.0	13.4	19.5	9.0
1978	7.7	11.0	5.5	7.2	10.2	5.1	12.6	18.7	8.3
1979	7.6	11.0	5.2	7.0	10.2	4.8	12.9	18.7	8.7
1980	7.4	10.6	5.1	6.8	9.8	4.8	12.2	18.3	8.0
1981	7.3	10.6	5.0	6.7	9.8	4.6	12.5	18.1	8.6
1982	7.0	10.3	4.9	6.5	9.4	4.5	12.2	19.2	7.4
1983	6.8	9.9	4.8	6.2	9.0	4.4	12.3	18.4	8.1
1984	6.8	9.9	4.7	6.2	9.1	4.3	11.9	17.9	7.9
1985	6.5	9.5	4.4	6.0	8.8	4.1	11.5	17.0	7.8
1986	6.4	9.4	4.2	5.8	8.6	3.8	11.4	17.4	7.4
1987	6.2	9.1	4.2	5.6	8.3	3.8	11.3	17.0	7.5
1988	6.1	8.8	4.2	5.5	7.9	3.8	11.0	17.0	7.1
1989	6.2	9.2	4.2	5.6	8.3	3.8	11.5	17.7	7.3
1990	6.1	8.9	4.2	5.4	8.0	3.7	11.5	17.2	7.7
1991	6.0	8.9	4.0	5.4	8.0	3.6	11.5	17.6	7.5
1992	5.6	8.1	3.9	5.0	7.3	3.4	10.7	15.6	7.5
1993	5.6	8.2	3.8	5.0	7.3	3.4	10.7	16.3	7.1
1994	5.4	7.9	3.7	4.8	7.0	3.3	10.3	15.4	7.0
1995	5.3	7.7	3.7	4.7	6.9	3.2	10.4	14.9	7.4
1996	5.1	7.4	3.5	4.5	6.6	3.1	10.0	15.1	6.8
1997	4.9	7.2	3.4	4.3	6.3	2.9	9.7	14.6	6.5
1998	4.8	6.8	3.4	4.3	6.1	3.0	9.3	13.7	6.4
1999	4.6	6.6	3.3	4.0	5.8	2.8	9.2	13.7	6.5
2000	4.5	6.4	3.2	4.0	5.6	2.8	8.9	12.8	6.5
2001	4.4	6.1	3.1	3.8	5.4	2.7	8.3	12.0	5.8
2002	4.2	5.9	3.0	3.7	5.3	2.5	8.4	12.0	6.1
2003	4.1	5.7	3.0	3.6	5.0	2.6	7.8	11.4	5.4
2004	4.0	5.5	2.8	3.5	4.9	2.5	7.5	11.2	5.2
1975-2004	5.8	8.4	4.0	5.3	7.6	3.6	10.7	15.9	7.2

<sup>a</sup> NCHS public use data file for the total US. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

- Statistic not shown. Rate based on less than 16 cases for the time interval.

Table XXV-2  
TESTIS CANCER (Invasive)

AGE-ADJUSTED SEER INCIDENCE<sup>a</sup> RATES BY YEAR, RACE AND AGE

	<u>All Races, Males</u>			<u>White Males</u>			<u>Black Males</u>		
	<u>All</u>	<u>&lt;50</u>	<u>50+</u>	<u>All</u>	<u>&lt;50</u>	<u>50+</u>	<u>All</u>	<u>&lt;50</u>	<u>50+</u>
<u>SEER INCIDENCE RATES</u>									
<u>YEAR OF DIAGNOSIS:</u>									
1975	3.7	4.2	2.4	4.2	4.7	2.6	-	-	-
1976	3.4	4.2	1.5	3.8	4.5	1.7	-	-	-
1977	4.3	5.1	2.2	4.9	5.8	2.4	-	-	-
1978	3.6	4.2	1.8	4.0	4.7	2.1	-	-	-
1979	3.9	4.5	2.2	4.3	5.1	2.3	-	-	-
1980	4.3	5.3	1.9	4.9	6.0	2.1	-	-	-
1981	4.2	5.1	1.8	4.8	5.8	2.1	-	-	-
1982	4.4	5.2	2.3	5.0	6.0	2.4	-	-	-
1983	4.6	5.5	2.2	5.2	6.3	2.4	-	-	-
1984	4.4	5.4	1.9	5.0	6.1	2.1	-	-	-
1985	4.5	5.5	1.8	5.1	6.3	2.1	-	-	-
1986	4.8	5.8	2.2	5.6	6.8	2.4	-	-	-
1987	5.0	6.3	1.9	5.7	7.1	1.9	-	-	-
1988	4.6	5.8	1.5	5.5	6.9	1.7	-	-	-
1989	5.5	6.7	2.2	6.3	7.8	2.5	-	-	-
1990	5.1	6.3	2.1	6.0	7.4	2.3	-	-	-
1991	5.1	6.2	2.1	5.7	7.0	2.3	-	-	-
1992	5.1	6.4	1.8	6.1	7.6	2.0	-	-	-
1993	5.1	6.4	1.6	6.0	7.5	1.9	-	-	-
1994	5.5	6.7	2.1	6.3	7.8	2.3	1.2	1.5	-
1995	4.5	5.6	1.8	5.2	6.4	2.1	1.5	1.9	-
1996	5.2	6.6	1.7	6.2	7.8	1.9	-	-	-
1997	5.4	6.5	2.4	6.3	7.7	2.7	1.3	-	-
1998	5.6	7.1	1.6	6.6	8.4	1.8	1.5	2.0	-
1999	5.4	6.8	1.8	6.3	7.9	2.0	1.6	2.0	-
2000	5.7	7.2	2.0	6.7	8.4	2.3	1.5	1.9	-
2001	5.5	6.8	2.0	6.5	8.1	2.2	1.7	1.8	-
2002	5.7	7.1	1.9	6.9	8.6	2.3	1.3	1.6	-
2003	5.2	6.5	1.8	6.3	8.0	2.0	1.1	-	-
2004	5.7	7.1	2.2	7.0	8.7	2.5	1.0	-	-
1975-2004	4.9	6.1	2.0	5.7	7.0	2.2	1.1	1.3	0.6

<sup>a</sup> SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta). Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).  
 - Statistic not shown. Rate based on less than 16 cases for the time interval.



Table XXV-3  
TESTIS CANCER (Invasive)

AGE-ADJUSTED U.S. DEATH<sup>a</sup> RATES BY YEAR, RACE AND AGE

	<u>All Races, Males</u>			<u>White Males</u>			<u>Black Males</u>		
	All	<50	50+	All	<50	50+	All	<50	50+
<u>U.S. DEATH RATES</u>									
YEAR OF DEATH:									
1975	0.7	0.7	0.7	0.8	0.8	0.8	0.2	-	-
1976	0.7	0.7	0.7	0.8	0.8	0.7	0.4	0.2	-
1977	0.6	0.7	0.6	0.7	0.7	0.6	0.2	-	-
1978	0.6	0.6	0.5	0.6	0.6	0.6	0.2	0.2	-
1979	0.5	0.5	0.6	0.5	0.5	0.6	0.4	0.3	-
1980	0.5	0.4	0.5	0.5	0.5	0.5	0.2	-	-
1981	0.4	0.4	0.4	0.4	0.4	0.4	0.2	-	-
1982	0.4	0.4	0.4	0.4	0.4	0.4	0.2	-	-
1983	0.4	0.4	0.4	0.4	0.4	0.4	0.2	0.2	-
1984	0.3	0.3	0.4	0.4	0.4	0.4	0.2	-	-
1985	0.4	0.3	0.4	0.4	0.4	0.4	0.2	-	-
1986	0.3	0.3	0.3	0.3	0.3	0.4	0.2	-	-
1987	0.3	0.4	0.3	0.4	0.4	0.3	0.2	0.2	-
1988	0.3	0.3	0.4	0.3	0.3	0.4	0.2	-	-
1989	0.3	0.3	0.3	0.3	0.3	0.3	0.2	0.2	-
1990	0.3	0.3	0.2	0.3	0.3	0.2	-	-	-
1991	0.3	0.3	0.3	0.3	0.3	0.3	0.1	-	-
1992	0.3	0.3	0.3	0.3	0.3	0.3	0.1	0.1	-
1993	0.3	0.3	0.3	0.3	0.3	0.3	0.2	0.1	-
1994	0.3	0.3	0.2	0.3	0.3	0.2	0.1	0.1	-
1995	0.2	0.2	0.2	0.3	0.3	0.2	-	-	-
1996	0.3	0.3	0.3	0.3	0.3	0.3	0.1	-	-
1997	0.2	0.2	0.2	0.3	0.3	0.3	0.2	0.2	-
1998	0.3	0.3	0.3	0.3	0.3	0.3	0.2	0.1	-
1999	0.3	0.3	0.3	0.3	0.3	0.3	0.1	0.1	-
2000	0.2	0.3	0.2	0.3	0.3	0.2	0.2	0.2	-
2001	0.2	0.2	0.3	0.3	0.3	0.2	0.2	-	-
2002	0.3	0.3	0.3	0.3	0.3	0.3	0.2	0.2	-
2003	0.2	0.2	0.3	0.3	0.3	0.3	0.2	0.1	-
2004	0.2	0.2	0.3	0.3	0.3	0.3	0.1	0.1	-
1975-2004	0.3	0.3	0.3	0.4	0.4	0.4	0.2	0.1	0.3

<sup>a</sup> NCHS public use data file for the total US. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

- Statistic not shown. Rate based on less than 16 cases for the time interval.

Table XXVI-2  
THYROID CANCER (Invasive)

AGE ADJUSTED SEER INCIDENCE<sup>a</sup> RATES BY YEAR, RACE AND SEX

	<u>All Races</u>			<u>Whites</u>			<u>Blacks</u>		
	<u>Total</u>	<u>Males</u>	<u>Females</u>	<u>Total</u>	<u>Males</u>	<u>Females</u>	<u>Total</u>	<u>Males</u>	<u>Females</u>
<u>AGE-ADJUSTED RATES</u>									
<u>YEAR OF DIAGNOSIS:</u>									
1975	4.9	3.1	6.4	4.7	2.8	6.4	3.0	-	4.2
1976	4.8	2.9	6.6	4.7	2.9	6.4	3.4	-	5.0
1977	5.4	3.5	7.3	5.3	3.4	7.2	3.7	-	5.4
1978	5.1	3.1	6.9	4.9	3.0	6.8	2.9	-	4.0
1979	4.5	2.7	6.2	4.4	2.6	6.2	3.2	-	4.3
1980	4.3	2.4	6.1	4.3	2.3	6.2	2.4	-	3.4
1981	4.4	2.5	6.2	4.4	2.5	6.2	2.5	-	3.6
1982	4.6	3.0	6.1	4.6	2.9	6.1	3.3	-	4.3
1983	4.7	2.8	6.5	4.7	2.6	6.7	3.0	2.1	3.7
1984	4.8	2.6	6.9	4.7	2.6	6.8	3.0	-	4.1
1985	5.1	3.1	7.1	5.1	3.1	7.0	2.8	-	3.6
1986	5.3	3.1	7.5	5.2	3.1	7.3	3.3	-	5.1
1987	5.0	2.8	7.1	5.0	2.9	7.1	2.8	-	4.3
1988	4.9	2.9	6.9	4.9	3.0	6.8	2.5	-	3.2
1989	5.4	3.0	7.7	5.5	3.2	7.7	2.5	-	4.0
1990	5.5	2.9	7.9	5.5	3.0	8.1	3.8	2.3	4.9
1991	5.5	3.2	7.6	5.6	3.3	7.9	3.2	-	4.2
1992	5.9	3.5	8.1	5.9	3.5	8.2	4.2	3.1	5.1
1993	5.6	3.6	7.6	5.7	3.7	7.8	3.4	2.1	4.6
1994	6.1	3.4	8.7	6.2	3.4	9.0	3.8	1.8	5.3
1995	6.2	3.4	8.9	6.4	3.6	9.2	3.0	-	4.4
1996	6.5	3.5	9.4	6.7	3.6	9.8	3.9	-	5.8
1997	6.7	3.7	9.7	7.0	3.7	10.3	3.9	2.4	5.2
1998	6.9	3.7	10.0	7.2	4.0	10.3	3.9	1.6	5.8
1999	7.3	3.9	10.6	7.5	4.1	10.9	4.6	1.6	6.9
2000	7.6	4.0	11.0	8.0	4.3	11.7	4.6	2.3	6.4
2001	8.2	4.3	12.0	8.6	4.6	12.6	4.8	2.7	6.5
2002	9.1	4.8	13.2	9.6	5.2	14.1	5.3	2.7	7.4
2003	9.4	4.7	14.1	10.1	5.0	15.2	5.2	2.7	7.4
2004	9.8	5.1	14.4	10.4	5.4	15.6	6.0	3.3	8.3
1975-2004	6.2	3.4	8.8	6.3	3.5	8.9	3.8	1.9	5.2

<sup>a</sup> SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta). Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

- Statistic not shown. Rate based on less than 16 cases for the time interval.

Table XXVI-3  
THYROID CANCER (Invasive)

AGE-ADJUSTED U.S. DEATH<sup>a</sup> RATES BY YEAR, RACE AND SEX

	<u>All Races</u>			<u>Whites</u>			<u>Blacks</u>		
	<u>Total</u>	<u>Males</u>	<u>Females</u>	<u>Total</u>	<u>Males</u>	<u>Females</u>	<u>Total</u>	<u>Males</u>	<u>Females</u>
<u>AGE-ADJUSTED RATES</u>									
<u>YEAR OF DEATH:</u>									
1975	0.5	0.4	0.6	0.5	0.4	0.6	0.5	0.3	0.6
1976	0.6	0.5	0.6	0.6	0.5	0.6	0.5	0.4	0.6
1977	0.6	0.5	0.6	0.6	0.5	0.6	0.7	0.5	0.7
1978	0.5	0.4	0.6	0.6	0.5	0.6	0.5	0.3	0.6
1979	0.5	0.4	0.6	0.5	0.5	0.6	0.5	0.3	0.6
1980	0.5	0.4	0.5	0.5	0.4	0.5	0.4	0.3	0.5
1981	0.5	0.4	0.5	0.5	0.4	0.5	0.5	0.3	0.6
1982	0.5	0.4	0.5	0.5	0.4	0.5	0.5	0.4	0.5
1983	0.4	0.3	0.5	0.4	0.3	0.5	0.4	0.3	0.5
1984	0.5	0.4	0.5	0.5	0.4	0.5	0.5	-	0.6
1985	0.4	0.4	0.5	0.4	0.4	0.5	0.5	0.3	0.6
1986	0.5	0.4	0.5	0.5	0.4	0.5	0.5	0.3	0.6
1987	0.5	0.4	0.5	0.4	0.4	0.5	0.5	0.3	0.6
1988	0.4	0.4	0.5	0.4	0.4	0.4	0.4	0.3	0.5
1989	0.4	0.4	0.4	0.4	0.4	0.4	0.5	0.4	0.6
1990	0.4	0.4	0.5	0.4	0.4	0.5	0.4	0.3	0.5
1991	0.4	0.4	0.5	0.4	0.4	0.4	0.5	0.4	0.6
1992	0.5	0.4	0.5	0.5	0.4	0.5	0.4	0.3	0.5
1993	0.5	0.4	0.5	0.5	0.4	0.5	0.5	0.3	0.6
1994	0.4	0.4	0.4	0.4	0.4	0.4	0.4	0.3	0.4
1995	0.4	0.4	0.4	0.4	0.4	0.4	0.4	0.3	0.5
1996	0.5	0.4	0.5	0.5	0.4	0.5	0.4	0.3	0.5
1997	0.5	0.4	0.5	0.5	0.5	0.4	0.5	0.4	0.6
1998	0.4	0.4	0.5	0.4	0.4	0.5	0.4	0.3	0.5
1999	0.5	0.4	0.5	0.4	0.4	0.4	0.4	0.4	0.5
2000	0.5	0.5	0.5	0.5	0.5	0.5	0.5	0.3	0.6
2001	0.5	0.5	0.5	0.5	0.5	0.5	0.5	0.4	0.6
2002	0.5	0.4	0.5	0.5	0.5	0.5	0.5	0.4	0.5
2003	0.4	0.4	0.5	0.5	0.4	0.5	0.4	0.3	0.4
2004	0.5	0.5	0.5	0.5	0.5	0.5	0.4	0.4	0.5
1975-2004	0.5	0.4	0.5	0.5	0.4	0.5	0.5	0.3	0.5

<sup>a</sup> NCHS public use data file for the total US. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

- Statistic not shown. Rate based on less than 16 cases for the time interval.

Table XXVII-2  
 URINARY BLADDER CANCER (Invasive and *In Situ*)

AGE ADJUSTED SEER INCIDENCE<sup>a</sup> RATES BY YEAR, RACE AND SEX

	All Races			Whites			Blacks		
	Total	Males	Females	Total	Males	Females	Total	Males	Females
<u>AGE-ADJUSTED RATES</u>									
YEAR OF DIAGNOSIS:									
1975	19.3	34.3	8.9	20.2	36.4	9.2	10.6	16.3	6.3
1976	19.7	34.4	9.4	20.7	36.7	9.7	12.5	18.2	8.6
1977	19.0	33.4	8.9	20.0	35.6	9.2	13.5	22.7	7.1
1978	20.0	35.4	9.4	21.1	37.9	9.7	11.7	19.4	6.4
1979	20.0	35.3	9.4	21.3	38.2	9.8	10.3	15.6	6.5
1980	20.4	37.1	8.9	21.5	39.8	9.1	13.3	18.2	10.0
1981	20.7	36.3	9.7	22.0	38.8	10.3	11.5	20.6	5.5
1982	20.1	35.8	9.5	21.4	38.3	10.2	12.2	21.8	5.6
1983	20.1	36.0	9.0	21.3	38.7	9.4	12.2	18.7	7.6
1984	20.8	37.7	9.2	22.3	40.9	9.6	11.4	18.6	6.2
1985	20.7	36.4	9.9	22.1	39.3	10.3	13.0	20.3	8.0
1986	21.0	38.0	9.4	22.4	40.8	9.9	12.9	22.2	6.6
1987	21.6	39.5	9.2	23.2	42.8	9.6	13.3	21.8	7.5
1988	20.9	38.1	9.0	22.7	41.9	9.7	11.3	18.0	6.4
1989	21.0	37.4	9.6	22.7	40.9	10.2	11.3	17.3	7.2
1990	21.1	37.5	9.6	22.6	40.7	10.0	12.7	18.6	8.7
1991	20.9	37.6	9.3	22.7	41.2	9.9	12.9	19.7	8.5
1992	21.2	37.5	9.8	22.9	40.7	10.5	12.5	20.8	6.9
1993	21.2	37.6	9.7	22.8	40.5	10.4	14.1	24.4	7.8
1994	20.8	36.8	9.4	22.6	40.2	10.2	12.3	20.2	6.9
1995	20.6	36.2	9.6	22.4	39.5	10.3	11.8	18.1	7.4
1996	20.8	36.5	9.5	22.6	39.9	10.3	12.2	19.6	7.3
1997	21.0	37.0	9.5	22.9	40.5	10.2	12.4	19.8	7.9
1998	21.5	38.4	9.4	23.5	42.0	10.2	11.8	20.2	6.4
1999	21.7	38.1	10.0	23.6	41.5	10.6	14.5	22.4	9.4
2000	21.7	38.7	9.5	23.8	42.3	10.2	13.1	20.2	8.5
2001	21.5	37.9	9.6	23.8	41.8	10.5	11.3	18.1	7.0
2002	21.0	36.6	9.6	23.1	40.1	10.5	12.2	17.4	8.6
2003	21.1	36.9	9.5	23.2	40.6	10.4	12.3	20.3	6.9
2004	20.6	36.3	9.1	22.6	39.8	9.9	12.6	20.2	8.0
1975-2004	20.8	36.9	9.4	22.4	40.1	10.0	12.4	19.8	7.5

<sup>a</sup> SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta). Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).  
 - Statistic not shown. Rate based on less than 16 cases for the time interval.

Table XXVII-3  
URINARY BLADDER CANCER (Invasive)

AGE-ADJUSTED U.S. DEATH<sup>a</sup> RATES BY YEAR, RACE AND SEX

	All Races			Whites			Blacks		
	Total	Males	Females	Total	Males	Females	Total	Males	Females
<u>AGE-ADJUSTED RATES</u>									
YEAR OF DEATH:									
1975	5.5	9.8	2.8	5.6	10.1	2.7	5.0	6.9	3.6
1976	5.6	9.8	2.9	5.6	10.0	2.8	5.3	7.8	3.5
1977	5.5	9.7	2.9	5.6	9.9	2.8	5.2	7.6	3.6
1978	5.4	9.5	2.9	5.5	9.8	2.8	4.9	7.2	3.4
1979	5.2	9.3	2.7	5.3	9.6	2.7	5.0	7.6	3.3
1980	5.2	9.2	2.7	5.2	9.5	2.7	4.7	6.2	3.7
1981	5.1	9.0	2.7	5.1	9.3	2.6	4.9	7.2	3.4
1982	5.0	8.9	2.6	5.0	9.2	2.5	4.9	6.8	3.6
1983	4.9	8.8	2.5	4.9	9.0	2.5	4.7	7.2	3.2
1984	4.7	8.5	2.5	4.8	8.7	2.4	4.7	6.9	3.4
1985	4.7	8.4	2.4	4.7	8.7	2.4	4.5	6.9	3.0
1986	4.5	8.1	2.4	4.6	8.4	2.3	4.2	5.8	3.3
1987	4.4	7.9	2.3	4.5	8.1	2.3	4.4	6.5	3.1
1988	4.4	7.8	2.4	4.5	8.0	2.3	4.3	5.9	3.3
1989	4.5	8.0	2.4	4.5	8.2	2.3	4.6	7.0	3.2
1990	4.5	8.0	2.4	4.6	8.2	2.4	4.4	6.4	3.2
1991	4.4	7.9	2.3	4.5	8.1	2.3	4.4	6.6	3.1
1992	4.5	7.9	2.4	4.5	8.1	2.4	4.3	6.7	3.1
1993	4.5	8.1	2.3	4.5	8.3	2.3	4.2	6.4	2.9
1994	4.5	7.9	2.4	4.5	8.1	2.4	4.3	6.4	3.2
1995	4.4	7.8	2.3	4.4	8.1	2.2	4.0	5.8	3.1
1996	4.4	7.8	2.4	4.5	8.1	2.3	4.2	5.9	3.1
1997	4.4	7.6	2.4	4.5	7.9	2.4	4.0	5.9	3.0
1998	4.4	7.7	2.3	4.5	8.0	2.3	3.9	5.6	3.0
1999	4.4	7.7	2.3	4.5	7.9	2.2	4.0	5.8	3.0
2000	4.3	7.6	2.3	4.5	7.9	2.3	3.7	5.7	2.7
2001	4.3	7.6	2.3	4.5	7.9	2.2	3.7	5.1	2.9
2002	4.4	7.5	2.4	4.5	7.8	2.3	3.9	5.5	3.0
2003	4.3	7.4	2.2	4.4	7.7	2.2	3.7	5.4	2.7
2004	4.4	7.6	2.2	4.5	8.0	2.2	3.6	5.0	2.8
1975-2004	4.6	8.2	2.4	4.7	8.5	2.4	4.3	6.3	3.2

<sup>a</sup> NCHS public use data file for the total US. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

- Statistic not shown. Rate based on less than 16 cases for the time interval.