

Table B2.1. Winter flounder commercial landings (metric tons) for the Gulf of Maine stock (U.S. statistical reporting areas 512 to 515). Landings from 1964-1981 is taken directly from SARC 21, 1982-1993 is re-estimated from the wodets, data and 1994-2001 is estimated using prorated dealer and VTR data.

| Year | metric tons |
|------|-------------|
| 1964 | 1,081 |
| 1965 | 665 |
| 1966 | 785 |
| 1967 | 803 |
| 1968 | 864 |
| 1969 | 975 |
| 1970 | 1,092 |
| 1971 | 1,113 |
| 1972 | 1,085 |
| 1973 | 1,080 |
| 1974 | 885 |
| 1975 | 1,181 |
| 1976 | 1,465 |
| 1977 | 2,161 |
| 1978 | 2,194 |
| 1979 | 2,021 |
| 1980 | 2,437 |
| 1981 | 2,406 |
| 1982 | 2,793 |
| 1983 | 2,096 |
| 1984 | 1,699 |
| 1985 | 1,582 |
| 1986 | 1,188 |
| 1987 | 1,140 |
| 1988 | 1,250 |
| 1989 | 1,253 |
| 1990 | 1,116 |
| 1991 | 1,008 |
| 1992 | 825 |
| 1993 | 611 |
| 1994 | 552 |
| 1995 | 796 |
| 1996 | 600 |
| 1997 | 618 |
| 1998 | 637 |
| 1999 | 253 |
| 2000 | 382 |
| 2001 | 571 |

Table B2.2. Percent commercial landings by gear for Gulf of Maine winter flounder.

| Year | otter trawl | shrimp trawl | gillnet | other |
|------|-------------|--------------|---------|-------|
| 1964 | 96% | | 1% | 3% |
| 1965 | 95% | - | 2% | 3% |
| 1966 | 98% | - | 1% | 2% |
| 1967 | 99% | - | - | 1% |
| 1968 | 98% | - | - | 2% |
| 1969 | 99% | - | - | 1% |
| 1970 | 99% | - | 1% | - |
| 1971 | 95% | - | 4% | 1% |
| 1972 | 95% | - | 4% | 1% |
| 1973 | 97% | - | 2% | - |
| 1974 | 95% | - | 5% | - |
| 1975 | 92% | 4% | 1% | 3% |
| 1976 | 87% | 2% | 6% | 5% |
| 1977 | 93% | 1% | 3% | 3% |
| 1978 | 89% | - | 3% | 9% |
| 1979 | 94% | - | 1% | 5% |
| 1980 | 95% | - | 1% | 4% |
| 1981 | 92% | 3% | 1% | 3% |
| 1982 | 89% | 5% | 2% | 4% |
| 1983 | 87% | 7% | 3% | 4% |
| 1984 | 85% | 8% | 2% | 6% |
| 1985 | 91% | 4% | 1% | 4% |
| 1986 | 77% | 6% | 14% | 4% |
| 1987 | 74% | 8% | 12% | 5% |
| 1988 | 81% | 5% | 13% | 1% |
| 1989 | 80% | 5% | 11% | 4% |
| 1990 | 77% | 2% | 19% | 2% |
| 1991 | 86% | 2% | 9% | 2% |
| 1992 | 77% | 2% | 19% | 2% |
| 1993 | 75% | - | 23% | 2% |
| 1994 | 78% | - | 21% | 1% |
| 1995 | 66% | - | 32% | 3% |
| 1996 | 72% | - | 27% | 1% |
| 1997 | 72% | - | 27% | 1% |
| 1998 | 73% | - | 27% | 1% |
| 1999 | 65% | - | 33% | 1% |
| 2000 | 73% | - | 26% | 1% |
| 2001 | 77% | - | 22% | 1% |

Table B2.3. Percent commercial landings by state for Gulf of Maine winter flounder.

| Year | ME | NH | MA | RI |
|------|-----|----|-----|----|
| 1964 | 3% | - | 97% | - |
| 1965 | 7% | - | 93% | - |
| 1966 | 6% | - | 94% | - |
| 1967 | 6% | - | 94% | - |
| 1968 | 3% | - | 97% | - |
| 1969 | 4% | - | 96% | - |
| 1970 | 13% | - | 87% | - |
| 1971 | 6% | - | 93% | 1% |
| 1972 | 12% | - | 88% | - |
| 1973 | 9% | - | 91% | - |
| 1974 | 13% | - | 87% | - |
| 1975 | 20% | - | 80% | - |
| 1976 | 12% | - | 88% | - |
| 1977 | 9% | - | 91% | - |
| 1978 | 14% | - | 86% | - |
| 1979 | 21% | - | 79% | - |
| 1980 | 23% | - | 77% | - |
| 1981 | 27% | 2% | 71% | - |
| 1982 | 32% | 4% | 64% | - |
| 1983 | 31% | 4% | 65% | - |
| 1984 | 23% | 6% | 71% | - |
| 1985 | 21% | 5% | 74% | 1% |
| 1986 | 22% | 4% | 73% | - |
| 1987 | 19% | 8% | 72% | 1% |
| 1988 | 22% | 9% | 69% | - |
| 1989 | 18% | 9% | 72% | - |
| 1990 | 14% | 7% | 78% | - |
| 1991 | 16% | 7% | 76% | - |
| 1992 | 14% | 7% | 79% | - |
| 1993 | 8% | 6% | 86% | - |
| 1994 | 5% | 7% | 88% | - |
| 1995 | 3% | 4% | 93% | - |
| 1996 | 1% | 5% | 94% | - |
| 1997 | 3% | 2% | 95% | - |
| 1998 | 1% | 2% | 97% | - |
| 1999 | - | 3% | 97% | - |
| 2000 | - | 4% | 95% | 1% |
| 2001 | 1% | 3% | 96% | - |

Table B2.4. Percent commercial landings by statistical area for Gulf of Maine winter flounder.

| Year | 511 | 512 | 513 | 514 | 515 |
|------|-----|-----|-----|-----|-----|
| 1964 | - | 2% | 1% | 96% | - |
| 1965 | - | 1% | 6% | 92% | 1% |
| 1966 | - | 2% | 7% | 90% | - |
| 1967 | - | 1% | 6% | 94% | - |
| 1968 | - | 2% | 1% | 97% | - |
| 1969 | - | 1% | 4% | 95% | - |
| 1970 | - | 1% | 12% | 87% | - |
| 1971 | - | 1% | 6% | 93% | - |
| 1972 | - | 1% | 12% | 87% | - |
| 1973 | - | 1% | 8% | 91% | - |
| 1974 | - | 2% | 11% | 87% | - |
| 1975 | 1% | 2% | 18% | 79% | - |
| 1976 | - | 1% | 13% | 86% | - |
| 1977 | - | 2% | 9% | 89% | - |
| 1978 | - | 3% | 13% | 83% | - |
| 1979 | 2% | 4% | 18% | 77% | - |
| 1980 | 1% | 3% | 20% | 76% | 1% |
| 1981 | - | 3% | 27% | 69% | 1% |
| 1982 | 3% | 5% | 27% | 62% | 2% |
| 1983 | 2% | 4% | 29% | 64% | 1% |
| 1984 | 1% | 3% | 27% | 68% | 1% |
| 1985 | 4% | 2% | 21% | 70% | 2% |
| 1986 | 4% | 5% | 26% | 64% | 2% |
| 1987 | 2% | 3% | 25% | 69% | 1% |
| 1988 | 4% | 6% | 22% | 67% | 1% |
| 1989 | 1% | 5% | 24% | 69% | 2% |
| 1990 | 4% | 3% | 21% | 71% | 1% |
| 1991 | 2% | 1% | 23% | 68% | 5% |
| 1992 | 1% | 3% | 21% | 73% | 3% |
| 1993 | 1% | - | 17% | 81% | 2% |
| 1994 | - | 2% | 14% | 81% | 2% |
| 1995 | 2% | 9% | 8% | 80% | 1% |
| 1996 | - | - | 9% | 90% | 1% |
| 1997 | - | - | 9% | 90% | 1% |
| 1998 | - | - | 4% | 96% | - |
| 1999 | - | - | 3% | 94% | 2% |
| 2000 | 1% | - | 5% | 94% | - |
| 2001 | - | - | 4% | 95% | - |

Table B2.5. Percent commercial landings by quarter for Gulf of Maine winter flounder.

| year | 1 | 2 | 3 | 4 |
|------|-----|-----|-----|-----|
| 1964 | 21% | 31% | 22% | 27% |
| 1965 | 22% | 27% | 11% | 40% |
| 1966 | 21% | 23% | 8% | 48% |
| 1967 | 15% | 35% | 8% | 42% |
| 1968 | 12% | 39% | 17% | 32% |
| 1969 | 23% | 37% | 15% | 26% |
| 1970 | 19% | 40% | 11% | 30% |
| 1971 | 25% | 33% | 19% | 22% |
| 1972 | 23% | 34% | 18% | 25% |
| 1973 | 24% | 27% | 16% | 33% |
| 1974 | 22% | 30% | 7% | 41% |
| 1975 | 18% | 25% | 17% | 40% |
| 1976 | 22% | 18% | 18% | 42% |
| 1977 | 24% | 19% | 13% | 44% |
| 1978 | 21% | 32% | 12% | 35% |
| 1979 | 13% | 28% | 17% | 42% |
| 1980 | 17% | 30% | 16% | 37% |
| 1981 | 23% | 28% | 14% | 34% |
| 1982 | 24% | 28% | 9% | 38% |
| 1983 | 28% | 31% | 12% | 30% |
| 1984 | 29% | 27% | 8% | 36% |
| 1985 | 26% | 31% | 10% | 33% |
| 1986 | 33% | 32% | 7% | 29% |
| 1987 | 29% | 34% | 7% | 30% |
| 1988 | 30% | 29% | 7% | 34% |
| 1989 | 27% | 39% | 8% | 27% |
| 1990 | 27% | 38% | 10% | 26% |
| 1991 | 26% | 32% | 9% | 32% |
| 1992 | 26% | 36% | 7% | 32% |
| 1993 | 18% | 37% | 11% | 34% |
| 1994 | 13% | 38% | 11% | 38% |
| 1995 | 22% | 38% | 15% | 25% |
| 1996 | 20% | 38% | 10% | 32% |
| 1997 | 18% | 34% | 16% | 31% |
| 1998 | 16% | 44% | 13% | 28% |
| 1999 | 13% | 44% | 17% | 25% |
| 2000 | 15% | 39% | 17% | 29% |
| 2001 | 9% | 41% | 17% | 32% |

Table B2.6. Percent commercial landings by market category for Gulf of Maine winter flounder.

| year | unclassified | small | medium | large |
|------|--------------|-------|--------|-------|
| 1964 | 77% | - | - | 23% |
| 1965 | 66% | - | - | 34% |
| 1966 | 68% | - | - | 32% |
| 1967 | 78% | - | - | 22% |
| 1968 | 70% | - | - | 30% |
| 1969 | 71% | - | - | 29% |
| 1970 | 75% | - | - | 25% |
| 1971 | 71% | - | - | 29% |
| 1972 | 64% | - | - | 36% |
| 1973 | - | 40% | - | 60% |
| 1974 | - | 38% | - | 62% |
| 1975 | - | 31% | - | 69% |
| 1976 | - | 42% | - | 58% |
| 1977 | - | 53% | - | 47% |
| 1978 | - | 50% | - | 50% |
| 1979 | - | 51% | - | 49% |
| 1980 | - | 49% | - | 50% |
| 1981 | 3% | 47% | - | 50% |
| 1982 | 12% | 41% | 2% | 44% |
| 1983 | 15% | 48% | 3% | 35% |
| 1984 | 15% | 46% | 7% | 33% |
| 1985 | 11% | 41% | 17% | 31% |
| 1986 | 17% | 39% | 16% | 29% |
| 1987 | 22% | 36% | 20% | 23% |
| 1988 | 19% | 42% | 17% | 22% |
| 1989 | 20% | 35% | 20% | 25% |
| 1990 | 22% | 34% | 15% | 29% |
| 1991 | 15% | 34% | 22% | 29% |
| 1992 | 16% | 33% | 23% | 29% |
| 1993 | 14% | 32% | 29% | 25% |
| 1994 | 14% | 33% | 28% | 26% |
| 1995 | 12% | 46% | 18% | 25% |
| 1996 | 10% | 56% | 17% | 18% |
| 1997 | 10% | 46% | 25% | 20% |
| 1998 | 29% | 44% | 18% | 9% |
| 1999 | 42% | 32% | 18% | 7% |
| 2000 | 36% | 41% | 14% | 9% |
| 2001 | 36% | 30% | 28% | 6% |

Table B2.7. Estimated number (000's) and weight (mt) of winter flounder caught, landed, and discarded in the recreational fishery, Gulf of Maine stock.

| Year | Numbers (000's) | | | | Metric Tons |
|------|-----------------|--------|----------|--------------|-------------|
| | Catch | Landed | Released | 15 % Release | Landed |
| | A+B1+B2 | A+B1 | B2 | Mortality | A+B2 |
| 1981 | 6,200 | 5,433 | 767 | 115 | 2,554 |
| 1982 | 8,207 | 7,274 | 933 | 140 | 1,876 |
| 1983 | 2,169 | 1,988 | 181 | 27 | 868 |
| 1984 | 2,477 | 2,285 | 191 | 29 | 1,300 |
| 1985 | 3,694 | 3,220 | 474 | 71 | 1,896 |
| 1986 | 946 | 691 | 255 | 38 | 523 |
| 1987 | 3,070 | 2,391 | 679 | 102 | 1,809 |
| 1988 | 953 | 841 | 111 | 17 | 345 |
| 1989 | 1,971 | 1,678 | 294 | 44 | 620 |
| 1990 | 786 | 652 | 134 | 20 | 370 |
| 1991 | 213 | 154 | 59 | 9 | 91 |
| 1992 | 186 | 137 | 48 | 7 | 90 |
| 1993 | 396 | 249 | 147 | 22 | 140 |
| 1994 | 232 | 145 | 87 | 13 | 83 |
| 1995 | 150 | 82 | 68 | 10 | 39 |
| 1996 | 184 | 98 | 86 | 13 | 56 |
| 1997 | 192 | 64 | 129 | 19 | 43 |
| 1998 | 109 | 65 | 44 | 7 | 30 |
| 1999 | 115 | 67 | 48 | 7 | 34 |
| 2000 | 177 | 75 | 102 | 15 | 42 |
| 2001 | 172 | 72 | 100 | 15 | 43 |

Table B2.8. Gulf of Maine winter flounder recreational landings (mt) by state.

| Year | ME | NH | MA | total |
|------|-----|----|-------|-------|
| 1981 | 45 | 55 | 2,455 | 2,554 |
| 1982 | 2 | 20 | 1,855 | 1,876 |
| 1983 | 11 | 36 | 821 | 868 |
| 1984 | 5 | 68 | 1,227 | 1,300 |
| 1985 | 4 | 28 | 1,864 | 1,896 |
| 1986 | 112 | 21 | 390 | 523 |
| 1987 | 1 | 12 | 1,796 | 1,809 |
| 1988 | 0 | 15 | 329 | 345 |
| 1989 | 197 | 20 | 402 | 620 |
| 1990 | 265 | 5 | 100 | 370 |
| 1991 | 23 | 0 | 68 | 91 |
| 1992 | 16 | 13 | 61 | 90 |
| 1993 | 37 | 9 | 94 | 140 |
| 1994 | 2 | 12 | 68 | 83 |
| 1995 | 0 | 4 | 35 | 39 |
| 1996 | 0 | 5 | 51 | 56 |
| 1997 | 17 | 6 | 20 | 43 |
| 1998 | 1 | 12 | 18 | 30 |
| 1999 | 0 | 6 | 27 | 34 |
| 2000 | 0 | 4 | 37 | 42 |
| 2001 | 1 | 7 | 36 | 43 |

Table B2.9. Percent Gulf of Maine winter flounder recreational landings (mt) by state.

| Year | ME | NH | MA |
|------|-----|-----|-----|
| 1981 | 2% | 2% | 96% |
| 1982 | 0% | 1% | 99% |
| 1983 | 1% | 4% | 95% |
| 1984 | 0% | 5% | 94% |
| 1985 | 0% | 1% | 98% |
| 1986 | 21% | 4% | 75% |
| 1987 | 0% | 1% | 99% |
| 1988 | 0% | 4% | 95% |
| 1989 | 32% | 3% | 65% |
| 1990 | 72% | 1% | 27% |
| 1991 | 25% | 0% | 75% |
| 1992 | 18% | 14% | 67% |
| 1993 | 27% | 6% | 67% |
| 1994 | 3% | 15% | 82% |
| 1995 | 0% | 11% | 89% |
| 1996 | 0% | 9% | 91% |
| 1997 | 40% | 13% | 46% |
| 1998 | 2% | 38% | 60% |
| 1999 | 0% | 19% | 81% |
| 2000 | 0% | 10% | 90% |
| 2001 | 1% | 15% | 83% |

Table B2.10. Gulf of Maine winter flounder recreational landing (mt) by halfyear.

| Year | halfyear 1 | halfyear 2 | total |
|------|------------|------------|-------|
| 1981 | 1,407 | 1,148 | 2,554 |
| 1982 | 517 | 1,359 | 1,876 |
| 1983 | 455 | 413 | 868 |
| 1984 | 599 | 701 | 1,300 |
| 1985 | 1,742 | 154 | 1,896 |
| 1986 | 485 | 39 | 523 |
| 1987 | 415 | 1,393 | 1,809 |
| 1988 | 211 | 134 | 345 |
| 1989 | 127 | 493 | 620 |
| 1990 | 52 | 318 | 370 |
| 1991 | 39 | 52 | 91 |
| 1992 | 24 | 66 | 90 |
| 1993 | 50 | 91 | 140 |
| 1994 | 38 | 45 | 83 |
| 1995 | 27 | 13 | 39 |
| 1996 | 39 | 17 | 56 |
| 1997 | 32 | 11 | 43 |
| 1998 | 15 | 15 | 30 |
| 1999 | 23 | 11 | 34 |
| 2000 | 14 | 28 | 42 |
| 2001 | 26 | 17 | 43 |

Table B2.11. Percent Gulf of Maine winter flounder recreational landing by halfyear.

| year | halfyear 1 | halfyear 2 |
|------|------------|------------|
| 1981 | 55% | 45% |
| 1982 | 28% | 72% |
| 1983 | 52% | 48% |
| 1984 | 46% | 54% |
| 1985 | 92% | 8% |
| 1986 | 93% | 7% |
| 1987 | 23% | 77% |
| 1988 | 61% | 39% |
| 1989 | 20% | 80% |
| 1990 | 14% | 86% |
| 1991 | 43% | 57% |
| 1992 | 27% | 73% |
| 1993 | 36% | 64% |
| 1994 | 46% | 54% |
| 1995 | 68% | 32% |
| 1996 | 69% | 31% |
| 1997 | 74% | 26% |
| 1998 | 50% | 50% |
| 1999 | 67% | 33% |
| 2000 | 33% | 67% |
| 2001 | 60% | 40% |

Table B2.12. Number of lengths, samples, and metric tons per sample for Gulf of Maine winter flounder. Number of samples and calculations of metric tons per samples does not include observer data or gillnet landings from 1990-2001. * = redistributed according to market category and halfyear proportions. Bold are lengths from observer trawl data.

| year | Qtr | Number of lengths. | | | | total | Number of samples | | | | | total | mt/samples | | | | | | | |
|------|-----|--------------------|-----|------|-----|-------|-------------------|----|-----|----|-------|-------|------------|-----|-----|----|-------|----|--|-----|
| | | lg | sm | med | un | | lg | sm | Med | un | total | | lg | sm | med | un | total | | | |
| 1982 | 1 | | | | 296 | | | | | 3 | | | | | | | | | | |
| | 2 | 102 | 101 | | 159 | | 1 | 1 | | 1 | | 838 | 453 | | | | | 46 | | |
| | 3 | 84 | 81 | | 106 | | 1 | 1 | | 1 | | | | | | | | | | |
| | 4 | | | | | 929 | | | | | 9 | 396 | 691 | | | | 231 | | | 310 |
| 1983 | 1 | 80 | | 99 | | | 1 | | | 1 | | | | | | | | | | |
| | 2 | 300 | 100 | | 407 | | 3 | 1 | | 4 | | 120 | 510 | | | | | 53 | | |
| | 3 | 108 | 388 | | | | 1 | 3 | | | | | | | | | | | | |
| | 4 | 107 | 956 | | 106 | 2651 | 1 | 8 | | 1 | 24 | 125 | 44 | 64 | 95 | | | | | 87 |
| 1984 | 1 | 201 | 209 | | | | 2 | 2 | | | | | | | | | | | | |
| | 2 | 237 | 294 | | 221 | | 3 | 2 | | 2 | | 74 | 95 | | | | | | | |
| | 3 | | 123 | | | | | 1 | | | | | | | | | | | | |
| | 4 | 126 | 690 | 100 | | 2201 | 1 | 5 | | 1 | 19 | 189 | 67 | 114 | 124 | | | | | 89 |
| 1985 | 1 | 273 | 565 | | | | 3 | 3 | | | | | | | | | | | | |
| | 2 | 392 | 170 | | | | 3 | 2 | | | | 54 | | | | | | | | |
| | 3 | 105 | | | | | 1 | | | | | | | | | | | | | |
| | 4 | 116 | | | 80 | 1701 | 1 | | | 1 | 14 | 87 | | 182 | 176 | | | | | 113 |
| 1986 | 1 | | | | 266 | | | | | 3 | | | | | | | | | | |
| | 2 | 237 | 109 | 109 | | | 3 | 1 | 1 | | | | 242 | 126 | 48 | | | | | |
| | 3 | | 111 | 86 | | | | 1 | 1 | | | | | | | | | | | |
| | 4 | | 389 | 107 | 89 | 1503 | 1 | 5 | 1 | 1 | 17 | 113 | 37 | 31 | 56 | | | | | 70 |
| 1987 | 1 | | | | 113 | | | | | 1 | | | | | | | | | | |
| | 2 | | | | | | | | | | | | | | | | | | | |
| | 3 | | 95 | | | | | 1 | | | | | | | | | | | | |
| | 4 | 47 | 156 | 272 | | 683 | 1 | 2 | | 3 | 8 | 257 | 137 | 75 | 249 | | | | | 143 |
| 1988 | 1 | | 258 | 311 | | | | 3 | 3 | | | | | | | | | | | |
| | 2 | 102 | | 395* | | | 1 | | 4* | | | | 108 | 23 | | | | | | |
| | 3 | | | | | | | | | | | | | | | | | | | |
| | 4 | | 169 | 107* | | 1342 | 1 | 2 | 1* | | 14 | 340 | 164 | 96 | | | | | | 89 |
| 1989 | 1 | | | | 100 | | | | | 1 | | | | | | | | | | |
| | 2 | 113 | | 91 | 134 | | 1 | | 1 | | | | | 168 | | | | | | |
| | 3 | | 95 | 120 | 32 | | | 1 | 1 | | | | | | | | | | | |
| | 4 | | | 100 | | 785 | 1 | | | 1 | 6 | 313 | 435 | 42 | 254 | | | | | 209 |
| 1990 | 1 | 328 | 301 | | | | 3 | 4 | | | | | | | | | | | | |
| | 2 | | | | 102 | | | | | 1 | | 64 | 48 | | | | | | | |
| | 3 | | | | | | | | | | | | | | | | | | | |
| | 4 | 117 | 197 | 97 | | 1142 | 1 | 2 | 1 | | 12 | 83 | 90 | 138 | 118 | | | | | 75 |

Table B2.12 Continued.

| Year | qtr | Number of lengths. | | | | total | Number of samples | | | | | total | mt/samples | | | | |
|------|-----|--------------------|-----|------------|------------|-------|-------------------|----|-----|----|-------|-------|------------|-----|-----|-----|-------|
| | | lg | sm | med | un | | lg | sm | med | un | total | | lg | sm | med | un | total |
| 1991 | 1 | 100 | 51 | 105 | 101 | 1375 | 1 | 1 | 1 | 1 | 15 | 65 | 92 | 72 | 95 | 115 | |
| | 2 | 88 | 203 | 100 | 42 | | 2 | 1 | 2 | 1 | | | | | | | |
| | 3 | | 95 | | | | 3 | | 1 | | | | | | | | |
| | 4 | 236 | 254 | | | | 4 | 3 | 3 | | | | | | | | |
| 1992 | 1 | 110 | | | 107 | 930 | 1 | 1 | | | 10 | 67 | 47 | 119 | 84 | | |
| | 2 | 136 | 100 | 93 | | | 2 | 2 | 1 | 1 | | | | | | | |
| | 3 | | | | | | 3 | | | | | | | | | | |
| | 4 | 57 | 74 | 253 | | | 4 | 1 | 1 | 3 | | | | | | | |
| 1993 | 1 | 100 | | | | 822 | 1 | 1 | | | 8 | 59 | 83 | 16 | | | |
| | 2 | | | 288 | | | 2 | | | 3 | | | | | | | |
| | 3 | | 55 | | 91 | | 3 | | 1 | | | | | | | | |
| | 4 | 80 | | 157 | 51 | | 4 | 1 | | 2 | | | | | | | |
| 1994 | 1 | | | | | 594 | 1 | | | | 7 | 62 | 112 | 143 | 15 | 60 | |
| | 2 | | 71 | 92 | 102 | | 2 | | 1 | 1 | | | | | | | 1 |
| | 3 | | | | | | 3 | | | | | | | | | | |
| | 4 | 94 | | 235 | | | 4 | 1 | | 3 | | | | | | | |
| 1995 | 1 | 101 | | 175 | 63 | 1661 | 1 | 1 | | 2 | 10 | 55 | 134 | 42 | | | |
| | 2 | | | 299 | | | 2 | | | 3 | | | | | | | |
| | 3 | | | 414 | | | 3 | | | 4 | | | | | | | |
| | 4 | | | | 609 | | 4 | | | | | | | | | | |
| 1996 | 1 | | 77 | | | 1637 | 1 | | 1 | | 15 | 29 | 80 | 16 | 18 | | |
| | 2 | | 231 | | | | 2 | | 2 | | | | | | | | |
| | 3 | | 355 | 252 | | | 3 | | 2 | 3 | | | | | | | |
| | 4 | 84 | 440 | 86 | 112 | | 4 | 1 | 5 | 1 | | | | | | | |
| 1997 | 1 | | 204 | | | 1709 | 1 | | 2 | | 23 | 19 | 25 | 11 | 14 | | |
| | 2 | | 127 | 75* | | | 2 | | 2 | 1* | | | | | | | |
| | 3 | | 220 | 218 | | | 3 | | 2 | 3 | | | | | | | |
| | 4 | 307 | 502 | 56* | | | 4 | 4 | 8 | 1* | | | | | | | |
| 1998 | 1 | | 148 | 79 | | 1504 | 1 | | 2 | 1 | 19 | 25 | 65 | 14 | 30 | | |
| | 2 | | 151 | 201* | | | 2 | | 3 | 2* | | | | | | | |
| | 3 | | 583 | | | | 3 | | 7 | | | | | | | | |
| | 4 | 69 | 163 | 110* | | | 4 | 1 | 2 | 1* | | | | | | | |
| 1999 | ↑ | | | | | 763 | ↑ | | | | 5 | 34 | | 26 | 10 | | |
| | 1 | | | 104 | | | 1 | | | 1 | | | | | | | |
| | 2 | | | 171 | | | 2 | | | 2 | | | | | | | |
| | 3 | | 28 | | | | 3 | | 1 | | | | | | | | |
| 4 | | 52 | | 408 | | 4 | | 1 | | | | | | | | | |

Table B2 . 12. Continued.

| | | Number of lengths. | | | | | Number of samples | | | | | mt/samples | | | | | |
|------|-----|--------------------|------|-----|-------------|-------|-------------------|----|-----|----|-------|------------|----|-----|----|-------|----|
| year | qtr | lg | sm | med | un | total | lg | sm | med | un | total | lg | sm | med | un | total | |
| 2000 | 1 | | 866 | 143 | 480 | 5827 | 1 | 12 | | 2 | 64 | | | | | 4 | |
| | 2 | | 3441 | 51 | 554 | | 2 | 45 | | 1 | | | | 1 | | | |
| | 3 | | 102 | | 50 | | 3 | 2 | | | | | | | | | |
| | 4 | | 114 | | 26 | | 4 | 2 | | | | | | 12 | 13 | | |
| 2001 | 1 | | | 187 | 172 | 3644 | 1 | | | 2 | 14 | | | | | 32 | |
| | 2 | 99 | 157 | 189 | 630 | | 2 | 1 | 2 | 3 | | | | 37 | 10 | | |
| | 3 | | 100 | 52 | 399 | | 3 | | 1 | 1 | | | | | | | |
| | 4 | | 154 | 198 | 1307 | | 4 | | 2 | 2 | | | | 26 | 21 | | 24 |

Table B2.13. Number of kept observer lengths, trips, and gillnet metric tons landed per 100 lengths sampled for Gulf of Maine winter flounder.

| Year | half | gillnet | | | mt/100 lengths |
|------|------|---------|-------|---------------|----------------|
| | | lengths | trips | landings (mt) | |
| 1990 | 1 | 539 | 90 | 184 | |
| | 2 | 78 | 1 | 29 | |
| | | 617 | 91 | 214 | 35 |
| 1991 | 1 | 126 | 6 | 81 | |
| | 2 | 30 | 8 | 13 | |
| | | 156 | 14 | 94 | 60 |
| 1992 | 1 | 1950 | 39 | 134 | |
| | 2 | 172 | 25 | 26 | |
| | | 2122 | 64 | 160 | 8 |
| 1993 | 1 | 2004 | 63 | 96 | |
| | 2 | 375 | 20 | 42 | |
| | | 2379 | 83 | 138 | 6 |
| 1994 | 1 | 330 | 22 | 101 | |
| | 2 | 206 | 10 | 15 | |
| | | 536 | 32 | 115 | 21 |
| 1995 | 1 | 1116 | 20 | 217 | |
| | 2 | 306 | 23 | 35 | |
| | | 1422 | 43 | 253 | 18 |
| 1996 | 1 | 1275 | 26 | 146 | |
| | 2 | 118 | 17 | 19 | |
| | | 1393 | 43 | 164 | 12 |
| 1997 | 1 | 793 | 18 | 139 | |
| | 2 | 42 | 4 | 27 | |
| | | 835 | 22 | 166 | 20 |
| 1998 | 1 | 1162 | 19 | 141 | |
| | 2 | 431 | 8 | 32 | |
| | | 1593 | 27 | 173 | 11 |
| 1999 | 1 | 747 | 5 | 78 | |
| | 2 | 526 | 12 | 7 | |
| | | 1273 | 17 | 85 | 7 |
| 2000 | 1 | 911 | 8 | 85 | |
| | 2 | 261 | 4 | 15 | |
| | | 1172 | 12 | 100 | 9 |
| 2001 | 1 | 862 | 15 | 94 | |
| | 2 | 42 | 2 | 32 | |
| | | 904 | 17 | 126 | 14 |

Table B2 . 14. Gulf of Maine winter flounder numbers of fish aged.

| Year | NEFSC | | | MA DMF | |
|------|---------------------|--------|------|--------|------|
| | Commercial landings | Spring | Fall | Spring | Fall |
| 1982 | 483 | 68 | 94 | 133 | |
| 1983 | 1182 | 150 | 104 | 159 | |
| 1984 | 908 | 63 | 150 | 139 | |
| 1985 | 318 | 135 | 160 | 97 | |
| 1986 | 344 | 84 | 62 | 57 | |
| 1987 | 130 | 118 | 67 | 125 | |
| 1988 | 249 | 127 | 68 | 104 | 7 |
| 1989 | 148 | 60 | 88 | 320 | |
| 1990 | 241 | 122 | 111 | 224 | |
| 1991 | 262 | 174 | 179 | 333 | |
| 1992 | 270 | 144 | 148 | 362 | |
| 1993 | 183 | 91 | 107 | 172 | |
| 1994 | 139 | 122 | 134 | 253 | 149 |
| 1995 | 248 | 170 | 55 | 213 | 221 |
| 1996 | 246 | 97 | 181 | 324 | |
| 1997 | 295 | 103 | 189 | 286 | |
| 1998 | 341 | 122 | 75 | 135 | |
| 1999 | 149 | 171 | 194 | 146 | |
| 2000 | 883 | 176 | 216 | 160 | |
| 2001 | 246 | 154 | 118 | 166 | |

Table B2.15. Gulf of Maine winter flounder discard ratios and number of trips/tows in the observer and VTR data for the large mesh, small mesh and gillnet fishery.

| Year | Half-year | Large Mesh Otter Trawl | | | | | Small Mesh Otter Trawl | | | | | Gillnet | | | | |
|------|-----------|------------------------|-------|----------|-----------|--------------|------------------------|-------|----------|-----------|-----------|---------|-------|--------------|-----------|-----------|
| | | # trips | #tows | SS ratio | VTR trips | VTR ratio | # trips | #tows | SS ratio | VTR trips | VTR ratio | # trips | #tows | SS ratio | VTR trips | VTR ratio |
| 1989 | Jan-Jun | 15 | 44 | 0.130 | | | 2 | 3 | 0.200 | | | | | | | |
| | Jul-Dec | 7 | 16 | 0.071 | | | 10 | 25 | 0.290 | | | 26 | 62 | 0.084 | | |
| 1990 | Jan-Jun | 5 | 6 | 0.167 | | | | | | | | 50 | 164 | 0.166 | | |
| | Jul-Dec | 6 | 14 | 0.287 | | | 2 | 3 | 0.333 | | | 33 | 63 | 0.223 | | |
| 1991 | Jan-Jun | 8 | 25 | 0.072 | | | 4 | 14 | 0.029 | | | 73 | 164 | 0.164 | | |
| | Jul-Dec | 23 | 103 | 0.055 | | | 8 | 18 | 1.152 | | | 321 | 618 | 0.142 | | |
| 1992 | Jan-Jun | 21 | 48 | 0.098 | | | 1 | 1 | 0.000 | | | 257 | 617 | 0.130 | | |
| | Jul-Dec | 6 | 22 | 0.039 | | | 3 | 11 | 0.068 | | | 224 | 397 | 0.114 | | |
| 1993 | Jan-Jun | 1 | 1 | 0.600 | | | | | | | | 196 | 576 | 0.150 | | |
| | Jul-Dec | 4 | 12 | 0.080 | | | 3 | 10 | 0.153 | | | 97 | 198 | 0.107 | | |
| 1994 | Jan-Jun | 1 | 1 | 0.000 | 445 | 0.053 | | | | | | 43 | 101 | 0.174 | 249 | 0.229 |
| | Jul-Dec | | | | 1422 | 0.062 | | | | | | 15 | 35 | 0.103 | 648 | 0.091 |
| 1995 | Jan-Jun | 4 | 15 | 1.101 | 2417 | 0.048 | | | | | | 18 | 54 | 0.285 | 907 | 0.150 |
| | Jul-Dec | 3 | 52 | 0.011 | 1149 | 0.037 | 22 | 57 | | | | 19 | 52 | 0.201 | 548 | 0.388 |
| 1996 | Jan-Jun | 2 | 5 | 0.068 | 2196 | 0.044 | 1 | 1 | | | | 17 | 62 | 0.128 | 589 | 0.159 |
| | Jul-Dec | 2 | 19 | 0.013 | 1227 | 0.035 | 26 | 93 | 3.344 | | | 18 | 39 | 0.066 | 364 | 0.553 |
| 1997 | Jan-Jun | 3 | 13 | 0.231 | 1700 | 0.034 | 1 | 4 | 0.218 | | | 18 | 56 | 0.245 | 470 | 0.112 |
| | Jul-Dec | | | | 887 | 0.023 | | | | | | 10 | 22 | 0.272 | 291 | 0.087 |
| 1998 | Jan-Jun | 5 | 16 | 0.233 | 1809 | 0.046 | | | | | | 27 | 87 | 0.109 | 543 | 0.144 |
| | Jul-Dec | | | | 939 | 0.030 | | | | | | 35 | 66 | 0.049 | 329 | 0.117 |
| 1999 | Jan-Jun | | | | 942 | 0.038 | | | | | | 14 | 41 | 0.141 | 285 | 0.136 |
| | Jul-Dec | 15 | 35 | 0.015 | 1148 | 0.038 | 13 | 35 | | | | 23 | 60 | 0.100 | 359 | 0.090 |
| 2000 | Jan-Jun | 35 | 78 | 0.041 | 1240 | 0.060 | 7 | 10 | 0.123 | | | 27 | 74 | 0.137 | 378 | 0.094 |
| | Jul-Dec | 6 | 8 | 0.000 | 1418 | 0.032 | 6 | 13 | 0.170 | | | 18 | 39 | 0.098 | 472 | 0.088 |
| 2001 | Jan-Jun | 27 | 61 | 0.100 | 1289 | 0.029 | | | | | | 13 | 27 | 0.061 | 340 | 0.095 |
| | Jul-Dec | 51 | 129 | 0.037 | 1272 | 0.045 | 2 | 3 | 0.000 | | | 9 | 21 | 0.101 | 523 | 0.107 |

Table B2.16. Gulf of Maine winter flounder discard lengths from observer data. MADMF observer length data in the small mesh otter trawl was also added to the table (6 tows, 2 trips, and 213 lengths in 1994; 55 tows, 20 trips, and 891 lengths in 1999; 20 tows, 8 trips, and 637 lengths in 2000).

| YEAR | large-mesh trawl | | | small mesh otter trawl | | | shrimp fishery | | | gillnet | | | | | | |
|------|------------------|-----|-----|------------------------|----|------|----------------|------|--|---------|-----|------|-----|-----|-----|-----|
| | H1 | H2 | | H1 | H2 | | H1 | H2 | | H1 | H2 | | | | | |
| 1989 | tows | 13 | | 13 | | | 7 | 7 | | 12 | 2 | 14 | | 1 | 1 | |
| | trips | 9 | | 9 | | | 4 | 4 | | 6 | 1 | 7 | | 1 | 1 | |
| | lengths | 116 | | 116 | | | 239 | 239 | | 347 | 79 | 426 | | 2 | 2 | |
| 1990 | tows | | | 0 | | | 0 | 3 | | 3 | | | 20 | 1 | 21 | |
| | trips | | | 0 | | | 0 | 3 | | 3 | | | 10 | 1 | 11 | |
| | lengths | | | 0 | | | 0 | 126 | | 126 | | | 313 | 18 | 331 | |
| 1991 | tows | 1 | | 1 | | | 0 | 32 | | 32 | | | 3 | 2 | 5 | |
| | trips | 1 | | 1 | | | 0 | 15 | | 15 | | | 3 | 1 | 4 | |
| | lengths | 9 | | 9 | | | 0 | 1144 | | 1144 | | | 20 | 2 | 22 | |
| 1992 | tows | | 1 | 1 | | | 0 | 72 | | 72 | | | 39 | 9 | 48 | |
| | trips | | 1 | 1 | | | 0 | 24 | | 24 | | | 30 | 7 | 37 | |
| | lengths | | 18 | 18 | | | 0 | 1026 | | 1026 | | | 352 | 32 | 384 | |
| 1993 | tows | | 2 | 2 | | | 3 | 3 | | 132 | 2 | 134 | | 35 | 20 | 55 |
| | trips | | 2 | 2 | | | 2 | 2 | | 53 | 1 | 54 | | 20 | 14 | 34 |
| | lengths | | 12 | 12 | | | 43 | 43 | | 1685 | 2 | 1687 | | 400 | 38 | 438 |
| 1994 | tows | | | 0 | | | 6 | 6 | | 106 | 3 | 109 | | 18 | 4 | 22 |
| | trips | | | 0 | | | 2 | 2 | | 49 | 3 | 52 | | 10 | 3 | 13 |
| | lengths | | | 0 | | | 213 | 213 | | 1002 | 5 | 1007 | | 136 | 6 | 142 |
| 1995 | tows | 2 | 9 | 11 | | | 21 | 21 | | 85 | 13 | 98 | | 23 | 12 | 35 |
| | trips | 1 | 2 | 3 | | | 12 | 12 | | 45 | 7 | 52 | | 14 | 8 | 22 |
| | lengths | 28 | 18 | 46 | | | 264 | 264 | | 1118 | 34 | 1152 | | 377 | 38 | 415 |
| 1996 | tows | | 2 | 2 | 1 | 59 | 60 | | | 36 | 6 | 42 | | 16 | 2 | 18 |
| | trips | | 1 | 1 | 1 | 21 | 22 | | | 17 | 3 | 20 | | 7 | 2 | 9 |
| | lengths | | 5 | 5 | 1 | 250 | 251 | | | 197 | 105 | 302 | | 89 | 2 | 91 |
| 1997 | tows | 1 | | 1 | | | 0 | 13 | | 13 | | | | 9 | | 9 |
| | trips | 1 | | 1 | | | 0 | 7 | | 7 | | | | 3 | | 3 |
| | lengths | 2 | | 2 | | | 0 | 155 | | 155 | | | | 67 | | 67 |
| 1998 | tows | | | 0 | | | 0 | | | 0 | | | | 17 | 2 | 19 |
| | trips | | | 0 | | | 0 | | | 0 | | | | 9 | 2 | 11 |
| | lengths | | | 0 | | | 0 | | | 0 | | | | 70 | 5 | 75 |
| 1999 | tows | | | 0 | | 71 | 71 | | | 0 | | | | 10 | 15 | 25 |
| | trips | | | 0 | | 30 | 30 | | | 0 | | | | 5 | 7 | 12 |
| | lengths | | | 0 | | 1195 | 1195 | | | 0 | | | | 163 | 53 | 216 |
| 2000 | tows | 5 | | 5 | 3 | 21 | 24 | | | 0 | | | | 11 | 1 | 12 |
| | trips | 3 | | 3 | 3 | 9 | 12 | | | 0 | | | | 6 | 1 | 7 |
| | lengths | 90 | | 90 | 9 | 640 | 649 | | | 0 | | | | 219 | 1 | 220 |
| 2001 | tows | 1 | 9 | 10 | | | 0 | | | 0 | | | | 5 | | 5 |
| | trips | 1 | 4 | 5 | | | 0 | | | 0 | | | | 3 | | 3 |
| | lengths | 8 | 184 | 192 | | | 0 | | | 0 | | | | 42 | | 42 |

Table B2 . 17. Discard ratios and estimated discards (mt) for large mesh trawl VTR data and gillnet observer data. A 50% mortality rate was applied to the total discard estimate. Discard estimates using the survey method for otter trawl is also shown for comparison. Gillnet ratio from 1986-1988 is the average from 1989-1993.

| year | large mesh trawl vtr ratio | vtr trawl discards (mt) | survey trawl discards (mt) | observer Gillnet ratio | gillnet discards (mt) |
|------|-------------------------------|----------------------------|-------------------------------|---------------------------|-----------------------------|
| 1982 | - | - | 343 | - | - |
| 1983 | - | - | 112 | - | - |
| 1984 | - | - | 67 | - | - |
| 1985 | - | - | 93 | - | - |
| 1986 | - | - | 63 | 0.136 | 11 |
| 1987 | - | - | 81 | 0.136 | 9 |
| 1988 | - | - | 106 | 0.136 | 11 |
| 1989 | - | - | 86 | 0.084 | 6 |
| 1990 | - | - | 81 | 0.173 | 18 |
| 1991 | - | - | 84 | 0.152 | 7 |
| 1992 | - | - | 56 | 0.129 | 10 |
| 1993 | - | - | 11 | 0.144 | 10 |
| 1994 | 0.061 | 13 | 65 | 0.165 | 9 |
| 1995 | 0.043 | 11 | 100 | 0.257 | 32 |
| 1996 | 0.040 | 8 | 72 | 0.119 | 10 |
| 1997 | 0.028 | 6 | 62 | 0.247 | 20 |
| 1998 | 0.038 | 9 | 53 | 0.100 | 8 |
| 1999 | 0.038 | 3 | 13 | 0.127 | 5 |
| 2000 | 0.041 | 6 | 19 | 0.133 | 7 |
| 2001 | 0.036 | 8 | 39 | 0.065 | 4 |

Table B2.18. Gulf of Maine winter flounder estimated discard ratios in the shrimp fishery (total discard kg / total days fished estimated from NEFSC and MA Observer data by shrimp season). Ratio for 1982-1988 is the average ratio from 1989-1992. Total shrimp fishery days fished estimated by Wigley et al 1999 and estimated discards are also shown. A 50% mortality is used for estimating dead discards. Dotted line indicates the introduction of the Nordmore grate.

| Year | trips | tows | ratio | Shrimp df | discard wt (mt) | dead discards (mt) |
|------|-------|------|--------|-----------|-----------------|--------------------|
| 1982 | | | 22.225 | 970.1 | 22 | 11 |
| 1983 | | | 22.225 | 1156.9 | 26 | 13 |
| 1984 | | | 22.225 | 1754.0 | 39 | 19 |
| 1985 | | | 22.225 | 2081.4 | 46 | 23 |
| 1986 | | | 22.225 | 2395.1 | 53 | 27 |
| 1987 | | | 22.225 | 3708.2 | 82 | 41 |
| 1988 | | | 22.225 | 2815.2 | 63 | 31 |
| 1989 | 12 | 24 | 13.361 | 2839.5 | 38 | 19 |
| 1990 | 25 | 53 | 24.070 | 3204.6 | 77 | 39 |
| 1991 | 38 | 94 | 27.720 | 2587.7 | 72 | 36 |
| 1992 | 72 | 225 | 23.749 | 2313.3 | 55 | 27 |
| 1993 | 63 | 178 | 10.730 | 1902.2 | 20 | 10 |
| 1994 | 63 | 183 | 7.320 | 1982.3 | 15 | 7 |
| 1995 | 58 | 136 | 7.382 | 3375.7 | 25 | 12 |
| 1996 | 40 | 92 | 6.290 | 3242.9 | 20 | 10 |
| 1997 | 21 | 55 | 12.511 | 3661.2 | 46 | 23 |
| 1998 | 3 | 6 | 10.559 | 2204.0 | 23 | 12 |
| 1999 | 4 | 5 | 5.645 | 1217.4 | 7 | 3 |
| 2000 | 4 | 10 | 10.927 | 792.9 | 9 | 4 |
| 2001 | 3 | 6 | 9.749 | 672.8 | 7 | 3 |

Table B2.19. Gulf of Maine winter flounder commercial numbers (000's) at age.

| Year | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10 | 11 | 12 | 13 |
|------|---|-----|-------|-------|-----|-----|-----|----|----|----|----|----|----|
| 1982 | | 550 | 2,025 | 1,288 | 733 | 482 | 181 | 22 | | | | | |
| 1983 | 5 | 366 | 1,026 | 1,311 | 632 | 282 | 109 | 68 | 21 | 13 | 7 | 2 | 1 |
| 1984 | | 599 | 1,512 | 982 | 384 | 235 | 152 | 76 | 44 | 7 | | | 1 |
| 1985 | | 25 | 573 | 1,164 | 759 | 263 | 82 | 64 | 26 | 5 | 5 | | |
| 1986 | | 310 | 629 | 512 | 303 | 199 | 58 | 28 | 12 | 4 | 1 | | |
| 1987 | | 283 | 821 | 422 | 356 | 141 | 25 | 35 | 2 | 0 | | | |
| 1988 | | 327 | 745 | 725 | 217 | 94 | 49 | 46 | 5 | 1 | | | |
| 1989 | | 37 | 840 | 733 | 602 | 102 | 8 | 7 | | | | | |
| 1990 | | 102 | 478 | 690 | 446 | 145 | 43 | 11 | 5 | 2 | | | |
| 1991 | | 175 | 735 | 519 | 191 | 104 | 45 | 28 | 1 | | | | |
| 1992 | | 188 | 609 | 511 | 174 | 57 | 20 | 7 | 2 | | | | |
| 1993 | 2 | 105 | 605 | 545 | 77 | 46 | 4 | | | | | | |
| 1994 | | 4 | 386 | 557 | 130 | 31 | 7 | | | | | | |
| 1995 | | 8 | 267 | 680 | 456 | 162 | 21 | 14 | 2 | | | | |
| 1996 | | 107 | 693 | 347 | 61 | 11 | 1 | 2 | 1 | | | | |
| 1997 | | 93 | 512 | 455 | 105 | 27 | 4 | 2 | | | | | |
| 1998 | | 25 | 217 | 458 | 321 | 105 | 34 | 4 | 1 | | | | |
| 1999 | | | 49 | 158 | 143 | 59 | 19 | 5 | 4 | | | | |
| 2000 | | 1 | 57 | 212 | 173 | 50 | 14 | 7 | | 1 | | | |
| 2001 | | 2 | 27 | 287 | 390 | 175 | 63 | 26 | 6 | 3 | | | |

Table B2.20. Gulf of Maine winter flounder commercial weight (kg) at age.

| Year | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10 | 11 | 12 | 13 |
|------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|
| 1982 | | 0.351 | 0.454 | 0.502 | 0.617 | 0.817 | 0.901 | 1.087 | 1.330 | | | | |
| 1983 | 0.293 | 0.281 | 0.403 | 0.528 | 0.667 | 0.814 | 0.970 | 1.062 | 1.238 | 1.415 | 1.467 | 1.224 | 1.422 |
| 1984 | | 0.294 | 0.301 | 0.392 | 0.550 | 0.763 | 0.971 | 1.124 | 1.124 | 1.275 | | | 1.578 |
| 1985 | | 0.307 | 0.366 | 0.449 | 0.572 | 0.802 | 1.020 | 1.121 | 1.183 | 1.071 | 1.462 | | |
| 1986 | | 0.412 | 0.470 | 0.534 | 0.699 | 0.842 | 0.940 | 1.231 | 1.387 | 0.479 | 2.996 | | |
| 1987 | | 0.380 | 0.437 | 0.586 | 0.650 | 0.843 | 1.107 | 1.272 | 1.684 | | | | |
| 1988 | | 0.510 | 0.524 | 0.530 | 0.669 | 0.620 | 0.976 | 1.082 | 1.132 | 2.338 | 1.619 | | |
| 1989 | | 0.286 | 0.434 | 0.542 | 0.592 | 1.034 | 1.155 | 1.264 | | | | | |
| 1990 | | 0.435 | 0.482 | 0.541 | 0.646 | 0.780 | 1.039 | 1.261 | 1.214 | 1.310 | | | |
| 1991 | | 0.393 | 0.487 | 0.626 | 0.624 | 0.725 | 0.741 | 0.896 | 1.810 | | | | |
| 1992 | | 0.364 | 0.447 | 0.569 | 0.653 | 0.787 | 1.075 | 1.461 | 1.745 | | | | |
| 1993 | 0.125 | 0.336 | 0.396 | 0.457 | 0.701 | 0.607 | 1.331 | | | | | | |
| 1994 | | 0.274 | 0.402 | 0.489 | 0.669 | 0.829 | 1.324 | 1.558 | | | | | |
| 1995 | | 0.305 | 0.369 | 0.437 | 0.552 | 0.653 | 1.030 | 1.181 | 1.447 | 2.572 | | | |
| 1996 | | 0.387 | 0.451 | 0.546 | 0.634 | 0.915 | 1.452 | 1.694 | 2.177 | 2.663 | | | |
| 1997 | | 0.412 | 0.451 | 0.540 | 0.701 | 0.847 | 0.998 | 1.479 | | | | | |
| 1998 | | 0.371 | 0.426 | 0.482 | 0.598 | 0.750 | 0.991 | 1.709 | 2.149 | 2.459 | | | |
| 1999 | | | 0.431 | 0.503 | 0.564 | 0.735 | 0.962 | 1.102 | 1.236 | 2.941 | | | |
| 2000 | | 0.449 | 0.400 | 0.480 | 0.560 | 0.711 | 0.930 | 1.178 | 1.467 | 1.555 | | | |
| 2001 | | 0.175 | 0.373 | 0.468 | 0.546 | 0.693 | 0.869 | 0.953 | 1.215 | 1.562 | | | |

Table B2.21. Gulf of Maine winter flounder recreational numbers (000's) at age.

| Year | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10 | 11 | 12 | 13 |
|------|----|-------|-------|-------|-------|-----|-----|----|----|----|----|----|----|
| 1982 | 40 | 1,546 | 2,526 | 2,180 | 669 | 135 | 95 | 22 | 38 | 6 | 5 | 7 | 3 |
| 1983 | 89 | 381 | 654 | 488 | 224 | 80 | 49 | 12 | 4 | | 6 | | |
| 1984 | 12 | 166 | 423 | 847 | 468 | 112 | 159 | 50 | 37 | | 10 | | |
| 1985 | | 112 | 762 | 875 | 1,163 | 136 | 136 | 37 | | | | | |
| 1986 | | 18 | 102 | 301 | 56 | 154 | 44 | 18 | | | | | |
| 1987 | | 28 | 805 | 739 | 436 | 170 | 113 | 37 | 52 | 9 | | | |
| 1988 | 2 | 10 | 103 | 320 | 142 | 153 | 75 | 30 | 3 | | | 3 | |
| 1989 | | 124 | 469 | 729 | 172 | 110 | 43 | 21 | 7 | 2 | | | |
| 1990 | | 111 | 228 | 236 | 37 | 25 | 5 | 5 | 3 | 2 | 1 | | |
| 1991 | | 9 | 31 | 47 | 34 | 12 | 9 | 7 | 3 | 1 | | | |
| 1992 | | 10 | 29 | 50 | 26 | 9 | 5 | 1 | 3 | 3 | | | |
| 1993 | | 21 | 54 | 79 | 66 | 20 | 5 | | 3 | | | | |
| 1994 | | 4 | 32 | 55 | 30 | 13 | 7 | 5 | | | | | |
| 1995 | | 2 | 22 | 27 | 19 | 8 | 3 | 2 | | | | | |
| 1996 | | | 17 | 40 | 17 | 11 | 7 | 5 | | 1 | | | |
| 1997 | | | 8 | 20 | 18 | 5 | 5 | 5 | 3 | 1 | | | |
| 1998 | | 2 | 19 | 32 | 8 | 4 | | | | | | | |
| 1999 | | | 8 | 23 | 17 | 11 | 4 | 5 | 1 | | | | |
| 2000 | | | 10 | 23 | 26 | 11 | 4 | | 1 | 1 | | | |
| 2001 | | | 8 | 22 | 16 | 14 | 12 | | | | | | |

Table B2.22. Gulf of Maine winter flounder recreational mean weights (kg) at age.

| Year | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10 | 11 | 12 | 13 |
|------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|
| 1982 | 0.109 | 0.197 | 0.339 | 0.479 | 0.571 | 0.746 | 1.025 | 1.522 | 1.929 | 2.801 | 3.431 | 3.963 | 5.187 |
| 1983 | 0.131 | 0.258 | 0.331 | 0.444 | 0.578 | 0.730 | 0.893 | 0.959 | 1.395 | | 1.365 | | |
| 1984 | 0.098 | 0.256 | 0.349 | 0.419 | 0.539 | 0.594 | 0.745 | 1.073 | 0.932 | | 1.784 | | |
| 1985 | | 0.196 | 0.293 | 0.456 | 0.592 | 0.823 | 0.872 | 1.047 | | | | | |
| 1986 | | 0.201 | 0.312 | 0.497 | 0.563 | 0.776 | 1.090 | 1.187 | | | | | |
| 1987 | | 0.138 | 0.417 | 0.510 | 0.724 | 0.871 | 1.062 | 1.195 | 1.252 | 1.784 | | | |
| 1988 | 0.098 | 0.254 | 0.372 | 0.464 | 0.620 | 0.838 | 1.053 | 1.359 | 1.600 | 0.000 | | 0.976 | |
| 1989 | | 0.277 | 0.432 | 0.630 | 0.762 | 0.981 | 1.179 | 1.298 | 1.781 | 1.547 | 0.000 | | |
| 1990 | | 0.268 | 0.425 | 0.644 | 0.642 | 0.770 | 0.678 | 1.317 | 1.078 | 1.257 | 1.199 | | |
| 1991 | | 0.360 | 0.375 | 0.460 | 0.569 | 0.708 | 0.916 | 0.993 | 1.307 | 0.616 | | | |
| 1992 | | 0.224 | 0.358 | 0.466 | 0.636 | 0.886 | 1.013 | 1.199 | 1.576 | 1.365 | | | |
| 1993 | | 0.282 | 0.381 | 0.482 | 0.626 | 0.848 | 0.997 | | 1.453 | | | | |
| 1994 | | 0.275 | 0.386 | 0.477 | 0.558 | 0.701 | 0.908 | 1.009 | | | | | |
| 1995 | | 0.284 | 0.393 | 0.446 | 0.552 | 0.621 | 0.644 | 0.872 | | | | | |
| 1996 | | 0.317 | 0.398 | 0.434 | 0.516 | 0.616 | 0.766 | 0.958 | 0.000 | 1.744 | | | |
| 1997 | | 0.271 | 0.428 | 0.426 | 0.471 | 0.545 | 0.619 | 0.690 | 0.765 | 0.869 | | | |
| 1998 | | 0.293 | 0.325 | 0.419 | 0.572 | 0.753 | | | | | | | |
| 1999 | | | 0.383 | 0.446 | 0.520 | 0.595 | 0.666 | 0.922 | 0.669 | | | | |
| 2000 | | | 0.449 | 0.496 | 0.529 | 0.567 | 0.668 | 0.616 | 0.983 | 1.047 | | | |
| 2001 | | | 0.347 | 0.405 | 0.521 | 0.640 | 0.689 | | | | | | |

Table B2.23. Gulf of Maine winter flounder recreational discards (000's) at age.

| Year | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10 | 11 | 12 | 13 |
|------|----|-----|----|---|---|---|---|---|---|----|----|----|----|
| 1982 | 25 | 105 | 9 | | | | | | | | | | |
| 1983 | 17 | 7 | 3 | | | | | | | | | | |
| 1984 | 5 | 14 | 10 | | | | | | | | | | |
| 1985 | 12 | 30 | 28 | 1 | | | | | | | | | |
| 1986 | 20 | 13 | 4 | 1 | | | | | | | | | |
| 1987 | 29 | 39 | 32 | 2 | | | | | | | | | |
| 1988 | 3 | 6 | 7 | 1 | | | | | | | | | |
| 1989 | 13 | 23 | 7 | 1 | | | | | | | | | |
| 1990 | 3 | 14 | 4 | | | | | | | | | | |
| 1991 | 2 | 4 | 3 | 1 | | | | | | | | | |
| 1992 | 3 | 2 | 1 | | | | | | | | | | |
| 1993 | 5 | 12 | 4 | 1 | | | | | | | | | |
| 1994 | 2 | 7 | 3 | 1 | | | | | | | | | |
| 1995 | 2 | 4 | 3 | 1 | | | | | | | | | |
| 1996 | 3 | 5 | 3 | 1 | | | | | | | | | |
| 1997 | 2 | 9 | 6 | 2 | | | | | | | | | |
| 1998 | 2 | 3 | 2 | | | | | | | | | | |
| 1999 | 2 | 3 | 2 | 1 | | | | | | | | | |
| 2000 | 4 | 6 | 4 | 2 | | | | | | | | | |
| 2001 | 3 | 4 | 5 | 3 | 1 | | | | | | | | |

Table B2.24. Gulf of Maine winter flounder recreational discards (kg) at age.

| Year | 1 | 2 | 3 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 11 | 12 | 13 |
|------|-------|-------|-------|-------|-------|---|---|---|---|---|----|----|----|
| 1982 | 0.041 | 0.084 | 0.116 | | | | | | | | | | |
| 1983 | 0.071 | 0.087 | 0.128 | | | | | | | | | | |
| 1984 | 0.072 | 0.072 | 0.117 | | | | | | | | | | |
| 1985 | 0.041 | 0.083 | 0.171 | 0.210 | | | | | | | | | |
| 1986 | 0.078 | 0.161 | 0.209 | 0.258 | 0.295 | | | | | | | | |
| 1987 | 0.043 | 0.088 | 0.216 | 0.307 | | | | | | | | | |
| 1988 | 0.059 | 0.120 | 0.177 | 0.279 | | | | | | | | | |
| 1989 | 0.055 | 0.158 | 0.228 | 0.285 | 0.325 | | | | | | | | |
| 1990 | 0.043 | 0.123 | 0.199 | 0.259 | 0.325 | | | | | | | | |
| 1991 | 0.055 | 0.108 | 0.210 | 0.288 | 0.325 | | | | | | | | |
| 1992 | 0.048 | 0.132 | 0.236 | 0.277 | 0.307 | | | | | | | | |
| 1993 | 0.048 | 0.108 | 0.184 | 0.286 | 0.293 | | | | | | | | |
| 1994 | 0.059 | 0.111 | 0.201 | 0.251 | 0.299 | | | | | | | | |
| 1995 | 0.055 | 0.127 | 0.207 | 0.239 | 0.325 | | | | | | | | |
| 1996 | 0.046 | 0.117 | 0.217 | 0.268 | 0.271 | | | | | | | | |
| 1997 | 0.042 | 0.092 | 0.170 | 0.247 | 0.287 | | | | | | | | |
| 1998 | 0.037 | 0.114 | 0.190 | 0.269 | 0.325 | | | | | | | | |
| 1999 | 0.051 | 0.103 | 0.207 | 0.245 | 0.314 | | | | | | | | |
| 2000 | 0.074 | 0.158 | 0.211 | 0.272 | 0.297 | | | | | | | | |
| 2001 | 0.042 | 0.098 | 0.208 | 0.261 | 0.285 | | | | | | | | |

Table B2.25. Gulf of Maine winter flounder commercial large mesh trawl discards (000's) at age using vtr ratios.

| Year | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10 | 11 | 12 | 13 |
|------|----|-----|-----|-----|----|---|---|---|---|----|----|----|----|
| 1982 | 40 | 642 | 697 | 18 | | | | | | | | | |
| 1983 | 18 | 124 | 249 | 36 | | | | | | | | | |
| 1984 | 3 | 87 | 97 | 59 | 3 | | | | | | | | |
| 1985 | 4 | 59 | 196 | 77 | 3 | | | | | | | | |
| 1986 | 1 | 77 | 143 | 23 | 9 | | | | | | | | |
| 1987 | 1 | 20 | 236 | 49 | 1 | | | | | | | | |
| 1988 | 3 | 61 | 233 | 107 | 3 | 1 | | | | | | | |
| 1989 | 2 | 118 | 105 | 71 | 19 | 6 | | | | | | | |
| 1990 | 1 | 86 | 162 | 49 | 17 | | | | | | | | |
| 1991 | 5 | 70 | 147 | 89 | 5 | | | | | | | | |
| 1992 | 2 | 56 | 105 | 45 | 8 | | | | | | | | |
| 1993 | 1 | 14 | 20 | 9 | 2 | | | | | | | | |
| 1994 | 1 | 10 | 22 | 13 | 4 | | | | | | | | |
| 1995 | 1 | 5 | 21 | 14 | 1 | | | | | | | | |
| 1996 | 2 | 7 | 12 | 8 | 1 | | | | | | | | |
| 1997 | | 5 | 9 | 6 | 2 | | | | | | | | |
| 1998 | | 7 | 14 | 9 | 3 | | | | | | | | |
| 1999 | | 2 | 5 | 3 | 1 | | | | | | | | |
| 2000 | 0 | 3 | 7 | 5 | 3 | 1 | | | | | | | |
| 2001 | | 2 | 8 | 10 | 4 | 2 | | | | | | | |

Table B2.26. Gulf of Maine winter flounder commercial large mesh trawl discards weight (kg) at age using vtr ratios.

| Year | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10 | 11 | 12 | 13 |
|------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|----|----|
| 1982 | 0.095 | 0.212 | 0.282 | 0.368 | 0.560 | 0.640 | 0.943 | 1.259 | 1.625 | 2.284 | | | |
| 1983 | 0.122 | 0.247 | 0.264 | 0.370 | 0.514 | 0.458 | 0.648 | 1.252 | | | 1.422 | | |
| 1984 | 0.091 | 0.223 | 0.278 | 0.322 | 0.350 | 0.595 | 0.699 | 0.954 | 1.014 | | | | |
| 1985 | 0.114 | 0.221 | 0.273 | 0.318 | 0.414 | 0.595 | 0.761 | 1.093 | 1.713 | | | | |
| 1986 | 0.038 | 0.182 | 0.275 | 0.317 | 0.301 | 0.508 | 0.815 | 1.014 | 1.422 | | | | |
| 1987 | 0.045 | 0.125 | 0.260 | 0.324 | 0.424 | 0.699 | 1.038 | 1.362 | 1.612 | | | | |
| 1988 | 0.068 | 0.210 | 0.249 | 0.314 | 0.388 | 0.410 | 0.768 | 1.029 | 1.432 | 1.619 | | | |
| 1989 | 0.056 | 0.229 | 0.280 | 0.289 | 0.351 | 0.336 | 0.594 | 1.249 | 0.000 | | | | |
| 1990 | 0.040 | 0.216 | 0.254 | 0.300 | 0.353 | 0.468 | 0.949 | 1.178 | 0.949 | 1.248 | | | |
| 1991 | 0.101 | 0.220 | 0.264 | 0.305 | 0.379 | 0.411 | 0.589 | 0.876 | 1.349 | 1.746 | | | |
| 1992 | 0.067 | 0.202 | 0.264 | 0.315 | 0.332 | 0.419 | 0.824 | 1.258 | 1.617 | | | | |
| 1993 | 0.069 | 0.202 | 0.243 | 0.306 | 0.348 | 0.494 | 0.751 | 1.377 | 1.533 | | | | |
| 1994 | 0.060 | 0.160 | 0.255 | 0.320 | 0.345 | 0.518 | 0.956 | | | | | | |
| 1995 | 0.045 | 0.152 | 0.249 | 0.319 | 0.390 | 0.499 | 0.249 | 1.351 | 1.515 | | | | |
| 1996 | 0.077 | 0.214 | 0.286 | 0.333 | 0.359 | 0.507 | 0.642 | 1.176 | | | | | |
| 1997 | 0.046 | 0.174 | 0.277 | 0.312 | 0.346 | 0.514 | 0.538 | 0.751 | | | | | |
| 1998 | 0.030 | 0.146 | 0.261 | 0.328 | 0.363 | 0.542 | 0.890 | 1.106 | | | | | |
| 1999 | 0.061 | 0.157 | 0.280 | 0.339 | 0.395 | 0.481 | 1.033 | 1.195 | 1.457 | | | | |
| 2000 | 0.094 | 0.205 | 0.270 | 0.309 | 0.367 | 0.382 | 0.468 | | 0.878 | 1.105 | | | |
| 2001 | 0.038 | 0.159 | 0.292 | 0.329 | 0.354 | 0.368 | 0.527 | 0.592 | 0.813 | 1.333 | | | |

Table B2.27. Gulf of Maine winter flounder gillnet discards (000's) at age.

| Year | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10 | 11 | 12 | 13 |
|------|---|---|----|----|----|---|---|---|---|----|----|----|----|
| 1986 | | 3 | 26 | 9 | 3 | | | | | | | | |
| 1987 | | | 27 | 6 | | | | | | | | | |
| 1988 | | | 27 | 13 | | | | | | | | | |
| 1989 | | | 14 | 7 | | | | | | | | | |
| 1990 | | 1 | 39 | 28 | 2 | | | | | | | | |
| 1991 | | 2 | 17 | 7 | 1 | | | | | | | | |
| 1992 | | 3 | 28 | 6 | | | | | | | | | |
| 1993 | | 1 | 25 | 10 | 1 | | | | | | | | |
| 1994 | | 1 | 22 | 11 | 2 | | | | | | | | |
| 1995 | | 6 | 37 | 23 | 12 | 5 | 3 | 1 | | | | | |
| 1996 | | 2 | 21 | 10 | 2 | | | | | | | | |
| 1997 | | 1 | 26 | 30 | 13 | | | | | | | | |
| 1998 | | 3 | 14 | 8 | 2 | | 1 | | | | | | |
| 1999 | | | 2 | 2 | 1 | 2 | 1 | 1 | | | | | |
| 2000 | | 1 | 8 | 7 | 4 | 1 | | | | | | | |
| 2001 | | | 4 | 5 | 2 | 1 | | | | | | | |

Table B2.28. Gulf of Maine winter flounder gillnet discard weight (kg) at age.

| Year | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10 | 11 | 12 | 13 |
|------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|----|----|----|
| 1986 | | 0.182 | 0.276 | 0.294 | 0.274 | 0.593 | | | | | | | |
| 1987 | | 0.154 | 0.265 | 0.306 | 0.503 | 0.693 | | | | | | | |
| 1988 | | 0.106 | 0.261 | 0.292 | 0.476 | 0.543 | | | | | | | |
| 1989 | | 0.122 | 0.259 | 0.295 | 0.363 | 0.346 | 0.693 | | | | | | |
| 1990 | | 0.143 | 0.249 | 0.278 | 0.338 | | | | | | | | |
| 1991 | | 0.200 | 0.269 | 0.298 | 0.341 | | | | | | | | |
| 1992 | | 0.196 | 0.283 | 0.311 | 0.360 | 0.409 | | | | | | | |
| 1993 | | 0.174 | 0.264 | 0.287 | 0.307 | 0.631 | | | | | | | |
| 1994 | | 0.172 | 0.246 | 0.295 | 0.313 | 0.538 | | | | | | | |
| 1995 | 0.112 | 0.246 | 0.285 | 0.358 | 0.546 | 0.636 | 0.600 | 0.824 | | | | | |
| 1996 | | 0.207 | 0.268 | 0.286 | 0.309 | 0.793 | 0.812 | | | | | | |
| 1997 | | 0.222 | 0.265 | 0.299 | 0.333 | | | | | | | | |
| 1998 | | 0.172 | 0.232 | 0.305 | 0.475 | 0.568 | 0.761 | 0.693 | | | | | |
| 1999 | | 0.184 | 0.277 | 0.372 | 0.540 | 0.684 | 0.793 | 0.786 | 1.132 | 1.484 | | | |
| 2000 | | 0.185 | 0.260 | 0.296 | 0.363 | 0.403 | 0.607 | 0.837 | 0.789 | | | | |
| 2001 | | | 0.267 | 0.315 | 0.323 | 0.401 | 0.812 | | 0.812 | 0.812 | | | |

Table B2.29. Gulf of Maine winter flounder commercial shrimp fishery discards (000's) at age.

| Year | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10 | 11 | 12 | 13 |
|------|----|-----|-----|----|---|---|---|---|---|----|----|----|----|
| 1982 | 13 | 65 | 16 | 1 | | | | | | | | | |
| 1983 | 17 | 62 | 37 | 4 | | | | | | | | | |
| 1984 | 15 | 83 | 55 | 19 | 1 | | | | | | | | |
| 1985 | 39 | 94 | 57 | 7 | | | | | | | | | |
| 1986 | 62 | 137 | 32 | 8 | 2 | | | | | | | | |
| 1987 | 48 | 182 | 110 | 7 | | | | | | | | | |
| 1988 | 44 | 103 | 101 | 13 | | | | | | | | | |
| 1989 | 42 | 136 | 45 | 4 | | | | | | | | | |
| 1990 | 35 | 53 | 86 | 33 | 7 | | | | | | | | |
| 1991 | 36 | 145 | 62 | 12 | 1 | | | | | | | | |
| 1992 | 46 | 177 | 30 | 3 | | | | | | | | | |
| 1993 | 38 | 67 | 17 | 4 | 1 | | | | | | | | |
| 1994 | 30 | 73 | 11 | 1 | | | | | | | | | |
| 1995 | 41 | 70 | 19 | 4 | | | | | | | | | |
| 1996 | 52 | 52 | 13 | 5 | 1 | | | | | | | | |
| 1997 | 34 | 171 | 44 | 7 | | | | | | | | | |
| 1998 | 41 | 61 | 16 | 3 | 1 | | | | | | | | |
| 1999 | 16 | 18 | 4 | 1 | | | | | | | | | |
| 2000 | 19 | 22 | 11 | 2 | 1 | | | | | | | | |
| 2001 | 17 | 16 | 5 | 2 | | | | | | | | | |

Table B2.30. Gulf of Maine winter flounder shrimp fishery weight (kg) at age.

| Year | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10 | 11 | 12 | 13 |
|------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|----|----|----|
| 1982 | 0.025 | 0.093 | 0.212 | 0.341 | 0.429 | | | | | | | | |
| 1983 | 0.023 | 0.074 | 0.183 | 0.322 | 0.505 | 0.400 | | 0.522 | | | | | |
| 1984 | 0.016 | 0.067 | 0.151 | 0.273 | 0.357 | 0.502 | 0.453 | | | | | | |
| 1985 | 0.034 | 0.094 | 0.188 | 0.293 | 0.470 | 0.000 | | | | | | | |
| 1986 | 0.035 | 0.107 | 0.234 | 0.308 | 0.316 | 0.469 | | | | | | | |
| 1987 | 0.028 | 0.081 | 0.197 | 0.343 | 0.470 | 0.519 | | | | | | | |
| 1988 | 0.028 | 0.078 | 0.170 | 0.291 | 0.400 | 0.353 | | | | | | | |
| 1989 | 0.029 | 0.079 | 0.191 | 0.277 | 0.393 | | | | | | | | |
| 1990 | 0.039 | 0.093 | 0.201 | 0.316 | 0.397 | 0.442 | | | | | | | |
| 1991 | 0.040 | 0.106 | 0.208 | 0.297 | 0.336 | 0.460 | | | | | | | |
| 1992 | 0.028 | 0.097 | 0.217 | 0.296 | 0.361 | 0.076 | | | | | | | |
| 1993 | 0.025 | 0.064 | 0.187 | 0.295 | 0.427 | 0.621 | 0.953 | | | | | | |
| 1994 | 0.026 | 0.066 | 0.145 | 0.286 | 0.413 | 0.603 | 0.767 | | | | | | |
| 1995 | 0.042 | 0.091 | 0.186 | 0.224 | 0.579 | 0.426 | 0.221 | 0.795 | | | | | |
| 1996 | 0.029 | 0.084 | 0.214 | 0.299 | 0.277 | 0.377 | | | | | | | |
| 1997 | 0.043 | 0.076 | 0.155 | 0.245 | 0.329 | 0.117 | 0.170 | | | | | | |
| 1998 | 0.037 | 0.088 | 0.162 | 0.299 | 0.440 | 0.568 | 0.687 | 0.974 | | | | | |
| 1999 | 0.033 | 0.078 | 0.196 | 0.219 | 0.400 | 0.569 | 0.866 | 0.810 | 0.933 | | | | |
| 2000 | 0.031 | 0.065 | 0.122 | 0.258 | 0.355 | 0.424 | 0.633 | 0.937 | 0.943 | | | | |
| 2001 | 0.032 | 0.068 | 0.163 | 0.240 | 0.300 | 0.431 | 0.683 | 0.931 | 0.751 | 0.920 | | | |

Table B2.31. Gulf of Maine winter flounder composition of the catch by number.

| year | Landings | | Discards | | | | Total |
|------|--------------|------------|--------------|---------|---------|--------|--------|
| | recreational | commercial | recreational | gillnet | lg mesh | shrimp | |
| 1982 | 7,274 | 5,282 | 140 | 0 | 1,397 | 96 | 14,188 |
| 1983 | 1,988 | 3,842 | 27 | 0 | 428 | 120 | 6,406 |
| 1984 | 2,285 | 3,992 | 29 | 0 | 249 | 174 | 6,729 |
| 1985 | 3,220 | 2,965 | 71 | 0 | 340 | 197 | 6,793 |
| 1986 | 691 | 2,055 | 38 | 41 | 253 | 240 | 3,318 |
| 1987 | 2,391 | 2,086 | 102 | 34 | 308 | 346 | 5,266 |
| 1988 | 841 | 2,210 | 17 | 40 | 406 | 262 | 3,775 |
| 1989 | 1,678 | 2,329 | 44 | 21 | 321 | 227 | 4,620 |
| 1990 | 652 | 1,922 | 20 | 70 | 315 | 214 | 3,193 |
| 1991 | 154 | 1,799 | 9 | 26 | 315 | 257 | 2,559 |
| 1992 | 137 | 1,567 | 7 | 36 | 216 | 256 | 2,220 |
| 1993 | 249 | 1,384 | 22 | 36 | 45 | 127 | 1,863 |
| 1994 | 145 | 1,116 | 13 | 36 | 49 | 116 | 1,475 |
| 1995 | 82 | 1,609 | 10 | 85 | 42 | 134 | 1,963 |
| 1996 | 98 | 1,224 | 13 | 35 | 31 | 123 | 1,524 |
| 1997 | 64 | 1,198 | 19 | 70 | 23 | 257 | 1,630 |
| 1998 | 65 | 1,166 | 7 | 29 | 33 | 123 | 1,423 |
| 1999 | 67 | 437 | 7 | 9 | 11 | 39 | 571 |
| 2000 | 75 | 516 | 15 | 22 | 20 | 54 | 701 |
| 2001 | 72 | 980 | 15 | 13 | 26 | 41 | 1,146 |

Table B2.32. Gulf of Maine winter flounder composition of the catch by weight (mt).

| year | Landings | | Discards | | | | Total |
|------|--------------|------------|--------------|---------|---------|--------|-------|
| | recreational | commercial | recreational | gillnet | lg mesh | shrimp | |
| 1982 | 1,876 | 2,793 | 11 | | 343 | 11 | 5,034 |
| 1983 | 868 | 2,096 | 2 | | 112 | 13 | 3,091 |
| 1984 | 1,300 | 1,699 | 2 | | 67 | 19 | 3,089 |
| 1985 | 1,896 | 1,582 | 8 | | 93 | 23 | 3,602 |
| 1986 | 523 | 1,188 | 5 | 11 | 63 | 27 | 1,817 |
| 1987 | 1,809 | 1,140 | 12 | 9 | 81 | 41 | 3,091 |
| 1988 | 345 | 1,250 | 2 | 11 | 106 | 31 | 1,745 |
| 1989 | 620 | 1,253 | 6 | 6 | 86 | 19 | 1,989 |
| 1990 | 370 | 1,116 | 3 | 18 | 81 | 39 | 1,626 |
| 1991 | 91 | 1,008 | 1 | 7 | 84 | 36 | 1,227 |
| 1992 | 90 | 825 | 1 | 10 | 56 | 27 | 1,009 |
| 1993 | 140 | 611 | 3 | 10 | 11 | 10 | 785 |
| 1994 | 83 | 552 | 2 | 9 | 13 | 7 | 666 |
| 1995 | 39 | 796 | 1 | 32 | 11 | 12 | 892 |
| 1996 | 56 | 600 | 2 | 10 | 8 | 10 | 686 |
| 1997 | 43 | 618 | 2 | 20 | 6 | 23 | 712 |
| 1998 | 30 | 637 | 1 | 8 | 9 | 12 | 697 |
| 1999 | 34 | 253 | 1 | 5 | 3 | 3 | 300 |
| 2000 | 42 | 382 | 2 | 7 | 6 | 4 | 443 |
| 2001 | 43 | 571 | 2 | 4 | 8 | 3 | 632 |

Table B2.33. Gulf of Maine winter flounder total catch at age (000's).

| Year | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8+ |
|------|-----|-------|-------|-------|-------|-----|-----|-----|
| 1982 | 118 | 2,909 | 5,274 | 3,487 | 1,402 | 617 | 276 | 104 |
| 1983 | 146 | 941 | 1,970 | 1,839 | 857 | 362 | 158 | 133 |
| 1984 | 36 | 949 | 2,097 | 1,907 | 856 | 348 | 312 | 225 |
| 1985 | 54 | 320 | 1,617 | 2,124 | 1,925 | 398 | 218 | 136 |
| 1986 | 83 | 557 | 936 | 852 | 373 | 353 | 102 | 62 |
| 1987 | 78 | 553 | 2,031 | 1,224 | 794 | 311 | 138 | 136 |
| 1988 | 52 | 507 | 1,215 | 1,179 | 361 | 248 | 123 | 89 |
| 1989 | 56 | 439 | 1,480 | 1,545 | 793 | 218 | 51 | 38 |
| 1990 | 39 | 366 | 997 | 1,037 | 509 | 170 | 48 | 29 |
| 1991 | 43 | 405 | 995 | 674 | 232 | 116 | 55 | 40 |
| 1992 | 52 | 436 | 802 | 615 | 208 | 67 | 24 | 16 |
| 1993 | 46 | 220 | 725 | 647 | 147 | 66 | 9 | 3 |
| 1994 | 33 | 98 | 477 | 638 | 166 | 44 | 14 | 5 |
| 1995 | 43 | 95 | 367 | 749 | 488 | 174 | 27 | 18 |
| 1996 | 57 | 174 | 758 | 413 | 83 | 23 | 8 | 9 |
| 1997 | 37 | 279 | 605 | 519 | 139 | 32 | 9 | 11 |
| 1998 | 44 | 100 | 283 | 511 | 335 | 109 | 36 | 5 |
| 1999 | 18 | 23 | 70 | 188 | 162 | 71 | 24 | 16 |
| 2000 | 23 | 33 | 97 | 251 | 206 | 62 | 18 | 11 |
| 2001 | 20 | 24 | 58 | 329 | 412 | 192 | 76 | 35 |

Table B2.34. Gulf of Maine winter flounder mean weight at age (kg).

| Year | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8+ |
|------|-------|-------|-------|-------|-------|-------|-------|-------|
| 1982 | 0.081 | 0.223 | 0.375 | 0.487 | 0.595 | 0.802 | 0.943 | 2.037 |
| 1983 | 0.115 | 0.252 | 0.357 | 0.502 | 0.644 | 0.795 | 0.946 | 1.164 |
| 1984 | 0.059 | 0.257 | 0.305 | 0.400 | 0.543 | 0.708 | 0.855 | 1.115 |
| 1985 | 0.041 | 0.169 | 0.311 | 0.447 | 0.584 | 0.809 | 0.927 | 1.122 |
| 1986 | 0.045 | 0.291 | 0.408 | 0.510 | 0.664 | 0.813 | 1.005 | 1.221 |
| 1987 | 0.034 | 0.240 | 0.390 | 0.527 | 0.690 | 0.858 | 1.070 | 1.284 |
| 1988 | 0.034 | 0.376 | 0.421 | 0.487 | 0.648 | 0.753 | 1.022 | 1.204 |
| 1989 | 0.036 | 0.197 | 0.412 | 0.570 | 0.623 | 0.989 | 1.175 | 1.397 |
| 1990 | 0.040 | 0.271 | 0.398 | 0.538 | 0.631 | 0.778 | 1.003 | 1.247 |
| 1991 | 0.048 | 0.256 | 0.429 | 0.563 | 0.609 | 0.722 | 0.771 | 0.965 |
| 1992 | 0.031 | 0.229 | 0.405 | 0.539 | 0.638 | 0.799 | 1.064 | 1.468 |
| 1993 | 0.031 | 0.226 | 0.380 | 0.454 | 0.658 | 0.680 | 1.148 | 1.453 |
| 1994 | 0.029 | 0.096 | 0.379 | 0.481 | 0.637 | 0.790 | 1.128 | 1.052 |
| 1995 | 0.043 | 0.127 | 0.345 | 0.431 | 0.552 | 0.651 | 0.929 | 1.186 |
| 1996 | 0.029 | 0.279 | 0.437 | 0.520 | 0.593 | 0.768 | 0.851 | 1.381 |
| 1997 | 0.043 | 0.191 | 0.415 | 0.514 | 0.630 | 0.802 | 0.798 | 0.859 |
| 1998 | 0.036 | 0.170 | 0.384 | 0.471 | 0.594 | 0.749 | 0.984 | 1.814 |
| 1999 | 0.035 | 0.088 | 0.391 | 0.490 | 0.559 | 0.713 | 0.907 | 1.062 |
| 2000 | 0.039 | 0.108 | 0.345 | 0.470 | 0.549 | 0.676 | 0.869 | 1.187 |
| 2001 | 0.033 | 0.090 | 0.317 | 0.454 | 0.542 | 0.685 | 0.840 | 1.055 |

Table B2 . 35. Gulf of Maine winter flounder catch at age construction summary.

| Catch at age component | years | halfyear | length data | age data |
|--|-------|---------------|--|---|
| Trawl and other commercial landings | 82-01 | mix | commercial and observer (unclassified) | commercial |
| gillnet commercial Landings | 90-01 | whole year | observer (kept) | commercial |
| recreational Landings | 82-01 | halfyear | MRFSS | combine NEFSC and MA DMF ages by halfyear |
| recreational Discards | 82-01 | halfyear | spr & fall MA DMF | combine NEFSC and MA DMF ages by halfyear |
| Large mesh trawl discards (survey) | 82-93 | whole year | survey method (spr & fall MA DMF) | combine NEFSC spr & fall survey |
| Large mesh trawl discards (vtr/survey) | 94-01 | whole year | survey method (spr & fall MA DMF) | combine NEFSC spr & fall survey |
| gillnet discards | 86-01 | whole year | observer (discards) | combine spr NEFSC and MA DMF ages |
| shrimp discards | 82-01 | shrimp season | observer (discards) | combine spr NEFSC and MA DMF ages |

Table B2 . 36. NEFSC and MADMF stratified mean survey indices of abundance for Gulf of Maine winter flounder. NEFSC indices use offshore strata (26,27,38-40) and inshore strata (58-61,65,66). NEFSC indices are calculated with trawl door conversion factors where appropriate. MADMF uses strata 25-36.

| year | NEFSC spring | | NEFSC fall | | MADMF spring | | MADMF fall | |
|------|--------------|--------|------------|--------|--------------|--------|------------|--------|
| | number | weight | number | weight | number | weight | number | weight |
| 1978 | | | | | 86.805 | 18.373 | 43.360 | 9.887 |
| 1979 | 9.063 | 3.218 | 6.003 | 2.602 | 64.952 | 14.407 | 119.506 | 28.978 |
| 1980 | 11.284 | 4.447 | 13.141 | 6.553 | 66.231 | 17.494 | 74.684 | 15.940 |
| 1981 | 13.051 | 3.946 | 4.179 | 3.029 | 100.569 | 28.370 | 47.342 | 13.228 |
| 1982 | 7.670 | 3.022 | 4.201 | 1.924 | 60.719 | 14.687 | 106.053 | 23.635 |
| 1983 | 12.367 | 5.653 | 10.304 | 3.519 | 108.508 | 27.233 | 88.143 | 15.772 |
| 1984 | 5.155 | 1.979 | 7.732 | 3.106 | 66.271 | 15.977 | 35.956 | 10.817 |
| 1985 | 3.469 | 1.418 | 7.638 | 2.324 | 48.651 | 13.594 | 44.564 | 7.381 |
| 1986 | 2.343 | 0.998 | 2.502 | 0.938 | 62.356 | 14.724 | 41.914 | 6.603 |
| 1987 | 5.609 | 1.503 | 1.605 | 0.488 | 83.171 | 17.648 | 50.426 | 7.227 |
| 1988 | 6.897 | 1.649 | 3.000 | 1.031 | 52.733 | 10.617 | 33.063 | 7.173 |
| 1989 | 3.717 | 1.316 | 6.402 | 2.013 | 63.595 | 13.317 | 33.983 | 7.462 |
| 1990 | 5.415 | 2.252 | 3.527 | 1.177 | 74.131 | 12.966 | 67.874 | 13.452 |
| 1991 | 4.517 | 1.436 | 7.035 | 1.467 | 49.265 | 11.587 | 88.777 | 15.473 |
| 1992 | 3.933 | 1.160 | 10.447 | 3.096 | 74.146 | 13.938 | 77.350 | 13.471 |
| 1993 | 1.556 | 0.353 | 7.559 | 1.859 | 80.133 | 12.390 | 92.476 | 14.996 |
| 1994 | 3.481 | 0.891 | 4.870 | 1.319 | 71.710 | 10.036 | 67.351 | 13.560 |
| 1995 | 12.185 | 3.149 | 4.765 | 1.446 | 87.848 | 14.560 | 84.768 | 17.250 |
| 1996 | 2.736 | 0.732 | 10.099 | 3.116 | 77.249 | 12.823 | 74.295 | 13.031 |
| 1997 | 2.806 | 0.664 | 10.008 | 2.950 | 95.918 | 14.796 | 74.347 | 14.316 |
| 1998 | 2.001 | 0.528 | 3.218 | 0.987 | 91.466 | 15.756 | 93.889 | 14.934 |
| 1999 | 6.510 | 1.982 | 10.921 | 3.269 | 77.941 | 14.198 | 117.648 | 22.672 |
| 2000 | 10.383 | 2.885 | 12.705 | 5.065 | 169.291 | 35.453 | 101.633 | 25.693 |
| 2001 | 5.242 | 1.666 | 8.786 | 3.131 | 90.153 | 23.891 | 80.978 | 18.367 |
| 2002 | 12.066 | 3.693 | 10.691 | 4.003 | 87.376 | 21.404 | | |

Table B2 . 37. NEFSC spring stratified mean number per tow at age for Gulf of Maine winter flounder (offshore strata 26,27,38-40 and inshore 58-61,65,66).

| Year | 0 | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10 | 11 | 12 | 13 | 14 | total |
|------|------|------|------|------|------|------|------|------|------|------|------|----|----|----|----|-------|
| 1980 | 0.10 | 3.28 | 4.73 | 1.79 | 0.96 | 0.31 | 0.06 | 0.06 | 0.05 | | | | | | | 11.28 |
| 1981 | 1.05 | 5.36 | 2.05 | 3.14 | 0.92 | 0.39 | 0.09 | 0.04 | | | | | | | | 13.05 |
| 1982 | 0.16 | 1.92 | 3.40 | 0.85 | 1.00 | 0.11 | 0.06 | 0.10 | | 0.03 | | | | | | 7.67 |
| 1983 | 0.42 | 0.88 | 3.65 | 3.06 | 1.88 | 1.00 | 1.21 | 0.23 | 0.02 | | 0.02 | | | | | 12.37 |
| 1984 | 0.23 | 1.13 | 1.37 | 1.17 | 0.61 | 0.08 | 0.35 | 0.03 | 0.16 | | 0.02 | | | | | 5.15 |
| 1985 | 0.01 | 0.53 | 1.41 | 0.65 | 0.57 | 0.10 | 0.14 | 0.04 | | 0.01 | | | | | | 3.47 |
| 1986 | 0.03 | 0.75 | 0.42 | 0.58 | 0.14 | 0.31 | 0.10 | 0.02 | | | | | | | | 2.34 |
| 1987 | 0.19 | 1.58 | 2.65 | 0.61 | 0.23 | 0.14 | 0.12 | 0.05 | 0.03 | | | | | | | 5.61 |
| 1988 | 0.65 | 1.36 | 3.04 | 1.42 | 0.26 | 0.11 | 0.03 | 0.03 | | | | | | | | 6.90 |
| 1989 | 0.06 | 0.49 | 1.39 | 1.13 | 0.31 | 0.13 | 0.10 | 0.11 | | | | | | | | 3.72 |
| 1990 | 0.04 | 0.61 | 1.63 | 1.54 | 0.78 | 0.34 | 0.04 | 0.17 | 0.14 | 0.14 | | | | | | 5.42 |
| 1991 | 0.09 | 1.26 | 1.52 | 1.01 | 0.47 | 0.10 | 0.04 | 0.01 | 0.01 | 0.01 | | | | | | 4.52 |
| 1992 | 0.31 | 1.16 | 1.01 | 0.96 | 0.34 | 0.10 | 0.03 | 0.01 | 0.01 | | | | | | | 3.93 |
| 1993 | 0.01 | 0.53 | 0.59 | 0.28 | 0.11 | 0.02 | 0.01 | | | | | | | | | 1.56 |
| 1994 | 0.02 | 1.00 | 1.28 | 0.78 | 0.29 | 0.08 | 0.01 | 0.01 | | | | | | | | 3.48 |
| 1995 | 0.59 | 2.89 | 5.45 | 2.20 | 0.68 | 0.20 | 0.14 | 0.02 | | | | | | | | 12.19 |
| 1996 | 0.05 | 0.59 | 1.05 | 0.74 | 0.23 | 0.06 | 0.01 | | | | | | | | | 2.74 |
| 1997 | 0.04 | 0.69 | 0.81 | 0.71 | 0.41 | 0.09 | 0.04 | 0.01 | | | | | | | | 2.81 |
| 1998 | 0.10 | 0.59 | 0.60 | 0.48 | 0.21 | 0.01 | | | 0.01 | | | | | | | 2.00 |
| 1999 | 0.31 | 1.17 | 2.28 | 1.68 | 0.71 | 0.36 | | | | | | | | | | 6.51 |
| 2000 | 0.16 | 1.50 | 3.76 | 2.41 | 1.56 | 0.75 | 0.17 | | 0.04 | 0.02 | | | | | | 10.38 |
| 2001 | 0.07 | 0.52 | 1.41 | 1.49 | 0.83 | 0.60 | 0.22 | 0.09 | 0.02 | | | | | | | 5.24 |
| 2002 | 0.20 | 1.59 | 2.98 | 3.57 | 2.29 | 0.92 | 0.34 | 0.11 | 0.07 | | | | | | | 12.07 |

Table B2 . 38. NEFSC fall stratified mean number per tow at age for Gulf of Maine winter flounder (offshore strata 26,27,38-40 and inshore 58-61,65,66).

| year | 0 | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10 | 11 | 12 | 13 | 14 | total |
|------|------|------|------|------|------|------|------|------|------|------|----|------|------|----|------|-------|
| 1980 | | 0.57 | 4.36 | 5.34 | 1.85 | 0.74 | 0.18 | | | 0.05 | | 0.05 | | | | 13.14 |
| 1981 | | 0.07 | 0.71 | 1.76 | 0.78 | 0.12 | 0.37 | 0.08 | 0.12 | 0.08 | | | 0.41 | | 0.04 | 4.18 |
| 1982 | | 0.30 | 1.21 | 1.68 | 0.40 | 0.32 | 0.08 | 0.21 | | | | | | | | 4.20 |
| 1983 | | 2.14 | 3.60 | 3.12 | 1.01 | 0.27 | 0.11 | 0.07 | | | | | | | | 10.30 |
| 1984 | | 0.45 | 2.34 | 1.67 | 2.17 | 0.59 | 0.22 | 0.17 | 0.11 | | | | | | | 7.73 |
| 1985 | | 1.30 | 2.74 | 1.92 | 1.15 | 0.33 | 0.10 | 0.10 | | | | | | | | 7.64 |
| 1986 | | 0.02 | 0.73 | 1.15 | 0.49 | 0.05 | 0.02 | 0.01 | 0.02 | | | | | | | 2.50 |
| 1987 | | 0.08 | 0.46 | 0.84 | 0.19 | 0.03 | | | | 0.01 | | | | | | 1.61 |
| 1988 | | 0.49 | 0.96 | 0.60 | 0.71 | 0.15 | 0.06 | 0.03 | | | | | | | | 3.00 |
| 1989 | | 0.46 | 3.60 | 1.42 | 0.77 | 0.08 | 0.07 | | | 0.01 | | | | | | 6.40 |
| 1990 | | 0.10 | 1.86 | 1.09 | 0.41 | 0.04 | 0.02 | 0.02 | | | | | | | | 3.53 |
| 1991 | 0.03 | 2.60 | 2.83 | 1.09 | 0.39 | 0.03 | 0.05 | 0.03 | | | | | | | | 7.04 |
| 1992 | | 1.92 | 3.70 | 2.40 | 1.63 | 0.75 | 0.01 | 0.03 | | | | | | | | 10.45 |
| 1993 | | 1.66 | 3.16 | 1.82 | 0.69 | 0.23 | 0.01 | | | | | | | | | 7.56 |
| 1994 | | 0.43 | 2.32 | 1.29 | 0.65 | 0.12 | 0.03 | 0.03 | | | | | | | | 4.87 |
| 1995 | | 0.47 | 1.83 | 1.51 | 0.63 | 0.19 | 0.14 | | | | | | | | | 4.77 |
| 1996 | 0.01 | 1.77 | 2.37 | 2.57 | 2.63 | 0.60 | 0.13 | 0.01 | | | | | | | | 10.10 |
| 1997 | | 0.41 | 4.32 | 3.19 | 1.47 | 0.57 | 0.03 | | | | | | | | | 10.01 |
| 1998 | | 0.19 | 0.92 | 1.13 | 0.78 | 0.14 | 0.06 | | | | | | | | | 3.22 |
| 1999 | | 0.81 | 2.77 | 3.65 | 2.85 | 0.68 | 0.15 | 0.01 | | | | | | | | 10.92 |
| 2000 | | 0.62 | 2.03 | 4.00 | 3.54 | 1.41 | 0.96 | 0.15 | | | | | | | | 12.70 |
| 2001 | | 0.36 | 1.66 | 2.59 | 2.80 | 0.96 | 0.36 | 0.04 | 0.01 | | | | | | | 8.79 |

Table B2 . 39. MADMF spring stratified mean number per tow at age for Gulf of Maine winter flounder (strata 25-36).

| year | 0 | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10 | 11 | 12 | 13 | 14 | total |
|------|------|-------|-------|-------|-------|-------|------|------|------|------|------|------|----|------|------|--------|
| 1982 | | 7.51 | 30.59 | 8.96 | 8.80 | 2.57 | 0.90 | 1.33 | 0.02 | 0.04 | | | | | | 60.72 |
| 1983 | 0.07 | 14.01 | 32.31 | 30.65 | 18.11 | 8.82 | 2.36 | 1.02 | 0.84 | 0.28 | | | | | 0.02 | 108.51 |
| 1984 | | 5.80 | 26.27 | 16.96 | 11.65 | 3.94 | 0.38 | 0.83 | 0.08 | 0.31 | | 0.04 | | | | 66.27 |
| 1985 | | 9.47 | 7.29 | 15.34 | 11.28 | 3.57 | 1.39 | 0.25 | 0.03 | 0.03 | | | | | | 48.65 |
| 1986 | | 9.35 | 19.78 | 20.97 | 10.29 | 1.22 | 0.46 | 0.06 | 0.04 | 0.19 | | | | | | 62.36 |
| 1987 | | 16.93 | 18.71 | 32.69 | 11.54 | 0.72 | 1.74 | 0.33 | 0.02 | 0.49 | | | | | | 83.17 |
| 1988 | 0.08 | 7.47 | 15.76 | 18.87 | 9.37 | 0.61 | 0.38 | 0.00 | 0.04 | 0.10 | | | | 0.05 | | 52.73 |
| 1989 | | 9.15 | 23.03 | 17.39 | 9.10 | 3.72 | 0.71 | 0.13 | 0.23 | 0.15 | | | | | | 63.59 |
| 1990 | | 14.31 | 18.33 | 27.47 | 10.04 | 2.04 | 1.35 | 0.39 | 0.08 | 0.08 | 0.02 | 0.04 | | | | 74.13 |
| 1991 | | 4.82 | 19.21 | 13.00 | 7.84 | 3.17 | 0.50 | 0.24 | 0.17 | 0.11 | 0.15 | 0.04 | | | | 49.27 |
| 1992 | | 19.96 | 32.12 | 12.31 | 6.70 | 1.97 | 0.69 | 0.16 | 0.07 | 0.08 | 0.07 | | | | | 74.15 |
| 1993 | | 17.86 | 37.10 | 15.09 | 6.46 | 2.03 | 1.09 | 0.34 | 0.02 | 0.11 | 0.04 | | | | | 80.13 |
| 1994 | | 14.33 | 36.11 | 15.44 | 4.66 | 0.79 | 0.12 | 0.17 | 0.08 | | 0.02 | | | | | 71.71 |
| 1995 | 0.06 | 20.76 | 36.25 | 22.59 | 6.02 | 1.33 | 0.54 | 0.15 | 0.11 | 0.02 | 0.02 | | | | | 87.85 |
| 1996 | | 14.96 | 34.59 | 17.79 | 7.04 | 1.88 | 0.73 | 0.19 | 0.08 | | | | | | | 77.25 |
| 1997 | | 15.04 | 39.94 | 22.78 | 10.72 | 5.34 | 1.08 | 0.58 | 0.26 | 0.09 | 0.06 | 0.03 | | | | 95.92 |
| 1998 | | 10.23 | 32.61 | 29.11 | 13.26 | 4.12 | 1.15 | 0.81 | 0.17 | | | | | | | 91.47 |
| 1999 | | 14.31 | 25.96 | 21.79 | 9.02 | 4.66 | 1.14 | 0.57 | 0.44 | 0.05 | | | | | | 77.94 |
| 2000 | | 28.67 | 69.85 | 33.39 | 18.16 | 11.00 | 5.83 | 1.79 | 0.37 | 0.22 | | | | | | 169.29 |
| 2001 | | 14.37 | 11.22 | 29.56 | 19.47 | 7.23 | 4.79 | 2.34 | 0.68 | 0.33 | 0.16 | | | | | 90.15 |
| 2002 | | 9.59 | 23.85 | 19.60 | 19.52 | 7.59 | 4.97 | 1.64 | 0.25 | 0.27 | 0.09 | | | | | 87.38 |

Table B2 .40. MADMF fall stratified mean number per tow at age for Gulf of Maine winter flounder (strata 25-36).

| year | 0 | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10 | 11 | 12 | 13 | 14 | total |
|------|------|-------|-------|-------|-------|------|------|------|------|---|----|----|----|----|----|--------|
| 1980 | 0.13 | 27.26 | 31.13 | 14.18 | 1.54 | 0.38 | 0.01 | 0.04 | | | | | | | | 74.68 |
| 1981 | 0.13 | 13.05 | 21.14 | 11.46 | 1.31 | 0.02 | 0.19 | 0.04 | | | | | | | | 47.34 |
| 1982 | 0.44 | 42.30 | 39.70 | 19.00 | 3.62 | 0.63 | 0.30 | 0.04 | 0.02 | | | | | | | 106.05 |
| 1983 | 0.00 | 49.19 | 23.26 | 11.70 | 2.80 | 1.11 | 0.07 | 0.01 | | | | | | | | 88.14 |
| 1984 | 0.06 | 8.29 | 11.63 | 6.41 | 6.89 | 1.80 | 0.59 | 0.25 | 0.02 | | | | | | | 35.96 |
| 1985 | 0.28 | 22.32 | 12.36 | 6.14 | 2.66 | 0.54 | 0.21 | 0.05 | | | | | | | | 44.56 |
| 1986 | 0.23 | 16.68 | 14.78 | 8.44 | 1.46 | 0.24 | 0.00 | 0.04 | 0.04 | | | | | | | 41.91 |
| 1987 | 0.50 | 17.29 | 19.40 | 11.68 | 1.34 | 0.10 | 0.11 | 0.02 | | | | | | | | 50.43 |
| 1988 | 0.16 | 11.96 | 12.69 | 3.87 | 3.09 | 0.80 | 0.34 | 0.11 | 0.04 | | | | | | | 33.06 |
| 1989 | | 12.17 | 14.59 | 5.29 | 1.41 | 0.31 | 0.19 | 0.03 | | | | | | | | 33.98 |
| 1990 | | 8.35 | 45.03 | 11.72 | 2.54 | 0.18 | 0.03 | 0.03 | | | | | | | | 67.87 |
| 1991 | 2.41 | 40.54 | 23.35 | 16.65 | 4.92 | 0.58 | 0.22 | 0.12 | | | | | | | | 88.78 |
| 1992 | 0.65 | 38.61 | 18.43 | 10.65 | 5.87 | 2.58 | 0.11 | 0.44 | | | | | | | | 77.35 |
| 1993 | 0.32 | 34.29 | 38.90 | 13.55 | 3.82 | 1.37 | 0.17 | 0.06 | | | | | | | | 92.48 |
| 1994 | 0.12 | 17.93 | 28.24 | 14.66 | 5.00 | 1.08 | 0.14 | 0.14 | 0.05 | | | | | | | 67.35 |
| 1995 | 0.29 | 29.32 | 30.17 | 17.27 | 6.04 | 0.91 | 0.49 | 0.22 | 0.05 | | | | | | | 84.77 |
| 1996 | 1.01 | 33.45 | 16.23 | 13.19 | 8.53 | 1.51 | 0.37 | | | | | | | | | 74.30 |
| 1997 | 0.47 | 20.04 | 29.06 | 17.89 | 5.25 | 1.54 | 0.10 | | | | | | | | | 74.35 |
| 1998 | 0.34 | 38.17 | 28.88 | 16.86 | 7.30 | 1.71 | 0.63 | | | | | | | | | 93.89 |
| 1999 | 1.17 | 30.34 | 42.82 | 23.00 | 15.01 | 4.10 | 1.15 | 0.06 | | | | | | | | 117.65 |
| 2000 | 0.30 | 25.54 | 30.64 | 23.79 | 13.65 | 4.34 | 2.43 | 0.94 | | | | | | | | 101.63 |
| 2001 | 0.20 | 27.85 | 17.67 | 14.22 | 14.96 | 4.13 | 1.71 | 0.22 | 0.01 | | | | | | | 80.98 |

Table B2 .41. Seabrook spring mean number per tow at age for Gulf of Maine winter flounder.

| year | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10 | 11 | total |
|------|------|------|------|------|------|------|------|------|------|------|----|-------|
| 1985 | 1.16 | 0.49 | 0.40 | 0.21 | 0.08 | 0.04 | 0.02 | | | | | 2.39 |
| 1986 | 1.65 | 1.06 | 0.52 | 0.23 | 0.06 | 0.01 | | | | | | 3.53 |
| 1987 | 1.60 | 1.47 | 1.08 | 0.15 | 0.01 | 0.08 | 0.03 | | 0.01 | | | 4.43 |
| 1988 | 0.88 | 1.18 | 1.52 | 0.31 | 0.02 | 0.02 | | | | | | 3.92 |
| 1989 | 3.73 | 1.30 | 1.35 | 0.37 | 0.06 | 0.03 | 0.01 | | | | | 6.85 |
| 1990 | 1.63 | 1.06 | 0.93 | 0.40 | 0.08 | 0.02 | | | | 0.01 | | 4.14 |
| 1991 | 2.66 | 1.19 | 1.19 | 0.37 | 0.12 | 0.02 | | | | | | 5.55 |
| 1992 | 0.58 | 1.00 | 0.34 | 0.16 | 0.02 | | | | | | | 2.11 |
| 1993 | | | | | | | | | | | | |
| 1994 | 0.81 | 1.16 | 0.32 | 0.05 | | | | | | | | 2.33 |
| 1995 | 0.97 | 0.97 | 0.38 | 0.09 | 0.02 | 0.01 | | | | | | 2.44 |
| 1996 | 1.38 | 1.35 | 0.63 | 0.11 | 0.03 | 0.01 | | | | | | 3.51 |
| 1997 | 0.94 | 1.29 | 0.59 | 0.21 | 0.08 | 0.02 | 0.01 | 0.01 | | | | 3.15 |
| 1998 | 1.39 | 2.62 | 1.67 | 0.56 | 0.17 | 0.04 | 0.01 | 0.01 | 0.02 | | | 6.50 |
| 1999 | 3.13 | 3.94 | 2.49 | 0.39 | 0.12 | 0.02 | 0.01 | 0.03 | | | | 10.14 |
| 2000 | 3.32 | 6.72 | 1.53 | 0.38 | 0.23 | 0.10 | 0.03 | 0.01 | 0.01 | | | 12.31 |
| 2001 | 2.74 | 0.97 | 1.76 | 0.32 | 0.06 | 0.03 | 0.02 | | | | | 5.91 |

Table B2 .42. Age and length at 50% maturity for Gulf of Maine winter flounder in the spring NEFSC, MADMF, and combined surveys with the sexes combined.

| time period | NEFSC | | | MADMF | | | Both | | |
|-------------|---------|------|-----|---------|------|-----|---------|------|-----|
| | total N | L50 | A50 | total N | L50 | A50 | total N | L50 | A50 |
| 81-85 | 456 | 23.7 | 2.5 | 479 | 29.1 | 3.5 | 935 | 26.6 | 2.9 |
| 86-90 | 510 | 21.3 | 2.3 | 763 | 28.5 | 3.4 | 1,273 | 25.4 | 3.0 |
| 91-95 | 700 | 24.2 | 2.8 | 1,312 | 28.4 | 3.2 | 2,012 | 26.8 | 3.0 |
| 96-01 | 823 | 22.8 | 2.6 | 1,212 | 27.7 | 3.3 | 2,035 | 25.3 | 3.0 |
| 81-01 | 2,489 | 23.1 | 2.6 | 3,766 | 28.3 | 3.3 | 6,255 | 26.0 | 3.0 |

Table B2 .43. Age at 50% maturity by sex and sexes combined for Gulf of Maine winter flounder in the Spring NEFSC, MADMF, and combined surveys.

| time period | sex | NEFSC | | MADMF | | Both | |
|-------------|----------|---------|-----|---------|-----|---------|-----|
| | | total N | A50 | total N | A50 | total N | A50 |
| 81-01 | male | 948 | 2.5 | 1,406 | 3.3 | 2,354 | 2.9 |
| | female | 1,601 | 2.6 | 2,533 | 3.4 | 4,134 | 3.1 |
| | Combined | 2,489 | 2.6 | 3,766 | 3.3 | 6,255 | 3.0 |

Table B2 .44. Comparison of length and age at 50% maturity for Gulf of Maine winter flounder in the spring NEFSC and MADMF surveys with the sexes combined. NEFSC data was limited to inshore Gulf of Maine Massachusetts strata (58-66) which overlap with the MADMF survey (25-36).

| time period | NEFSC | | | MADMF | | |
|-------------|---------|------|-----|---------|------|-----|
| | total N | L50 | A50 | total N | L50 | A50 |
| 81-85 | 209 | 24.0 | 2.4 | 479 | 29.1 | 3.5 |
| 86-90 | 248 | 21.0 | 2.1 | 763 | 28.5 | 3.4 |
| 91-95 | 493 | 25.0 | 2.8 | 1,312 | 28.4 | 3.2 |
| 96-01 | 577 | 23.0 | 2.5 | 1,212 | 27.7 | 3.3 |
| 81-01 | 1,527 | 23.5 | 2.5 | 3,766 | 28.3 | 3.3 |

Table B2.45. Virtual Population Analysis for Gulf of Maine winter flounder, 1982-2001.

Fisheries Assessment Toolbox gom wf total catch Run Number 1 12/3/2002 12:55:40 PM
 FACT Version 1.5.0
 gom wf total catch 1982 - 2002
 Input Parameters and Options Selected

 Natural mortality is a matrix below
 Oldest age (not in the plus group) is 7
 For all years prior to the terminal year (20), backcalculated
 stock sizes for the following ages used to estimate
 total mortality (Z) for age 7 : 5 6 7
 This method for estimating F on the oldest age is generally used when a
 flat-topped partial recruitment curve is thought to be characteristic of the stock.
 F for age 8 + is then calculated from the following
 ratios of F[age 8 +] to F[age 7]

| | |
|------|---|
| 1982 | 1 |
| 1983 | 1 |
| 1984 | 1 |
| 1985 | 1 |
| 1986 | 1 |
| 1987 | 1 |
| 1988 | 1 |
| 1989 | 1 |
| 1990 | 1 |
| 1991 | 1 |
| 1992 | 1 |
| 1993 | 1 |
| 1994 | 1 |
| 1995 | 1 |
| 1996 | 1 |
| 1997 | 1 |
| 1998 | 1 |
| 1999 | 1 |
| 2000 | 1 |
| 2001 | 1 |

Stock size of the 8 + group is then calculated using
 the following method: CATCH EQUATION

Partial recruitment estimate for 2002

| | |
|---|------|
| 1 | 0.02 |
| 2 | 0.04 |
| 3 | 0.15 |
| 4 | 0.57 |
| 5 | 1 |
| 6 | 1 |
| 7 | 1 |

The Indices that will be used in this run are:

| | |
|----|---------|
| 1 | NEC_S11 |
| 2 | NEC_S22 |
| 3 | NEC_S33 |
| 4 | NEC_S44 |
| 5 | NEC_S55 |
| 6 | NEC_S66 |
| 7 | NEC_S77 |
| 8 | NEC_S88 |
| 9 | NEC_F23 |
| 10 | NEC_F34 |
| 11 | NEC_F45 |
| 12 | NEC_F56 |
| 13 | NEC_F67 |
| 14 | MA_S11 |
| 15 | MA_S22 |
| 16 | MA_S33 |
| 17 | MA_S44 |
| 18 | MA_S55 |
| 19 | MA_S66 |
| 20 | MA_S77 |
| 21 | MA_S88 |
| 22 | MA_F01 |
| 23 | MA_F12 |
| 24 | MA_F23 |
| 25 | MA_F34 |
| 26 | MA_F45 |
| 27 | SEA_S11 |
| 28 | SEA_S22 |
| 29 | SEA_S33 |
| 30 | SEA_S44 |
| 31 | SEA_S55 |
| 32 | SEA_S66 |
| 33 | SEA_S77 |

Table B2.45. Continued.

STOCK NUMBERS (Jan 1) in thousands

| | 1982 | 1983 | 1984 | 1985 | 1986 | 1987 | 1988 |
|----|-------|-------|-------|-------|-------|-------|-------|
| 1 | 11761 | 8778 | 6269 | 9277 | 7686 | 6125 | 4482 |
| 2 | 14415 | 9522 | 7055 | 5100 | 7547 | 6218 | 4944 |
| 3 | 11100 | 9170 | 6945 | 4917 | 3886 | 5675 | 4590 |
| 4 | 6207 | 4316 | 5725 | 3788 | 2563 | 2334 | 2808 |
| 5 | 3058 | 1927 | 1869 | 2962 | 1180 | 1327 | 804 |
| 6 | 1177 | 1235 | 802 | 756 | 683 | 628 | 368 |
| 7 | 571 | 405 | 683 | 342 | 259 | 240 | 233 |
| 8 | 212 | 337 | 486 | 209 | 156 | 232 | 166 |
| 1+ | 48500 | 35690 | 29834 | 27351 | 23959 | 22779 | 18395 |
| | 1989 | 1990 | 1991 | 1992 | 1993 | 1994 | 1995 |
| 1 | 4043 | 4242 | 4542 | 3322 | 3240 | 4519 | 7503 |
| 2 | 3622 | 3259 | 3438 | 3680 | 2673 | 2611 | 3670 |
| 3 | 3589 | 2569 | 2337 | 2448 | 2618 | 1989 | 2049 |
| 4 | 2659 | 1599 | 1201 | 1013 | 1279 | 1488 | 1197 |
| 5 | 1232 | 779 | 371 | 373 | 273 | 461 | 641 |
| 6 | 331 | 291 | 177 | 94 | 117 | 91 | 228 |
| 7 | 77 | 74 | 85 | 40 | 16 | 36 | 34 |
| 8 | 56 | 44 | 60 | 26 | 05 | 13 | 22 |
| 1+ | 15610 | 12857 | 12211 | 10996 | 10221 | 11208 | 15343 |
| | 1996 | 1997 | 1998 | 1999 | 2000 | 2001 | 2002 |
| 1 | 7588 | 7249 | 8967 | 10080 | 7474 | 7391 | 6274 |
| 2 | 6104 | 6161 | 5902 | 7301 | 8237 | 6099 | 6033 |
| 3 | 2919 | 4840 | 4792 | 4742 | 5957 | 6714 | 4971 |
| 4 | 1345 | 1704 | 3415 | 3667 | 3819 | 4789 | 5444 |
| 5 | 302 | 728 | 925 | 2334 | 2832 | 2899 | 3624 |
| 6 | 83 | 172 | 470 | 454 | 1764 | 2132 | 2001 |
| 7 | 29 | 47 | 112 | 286 | 308 | 1388 | 1572 |
| 8 | 32 | 57 | 15 | 190 | 188 | 638 | 1558 |
| 1+ | 18402 | 20958 | 24598 | 29055 | 30578 | 32050 | 31477 |

Table B2.45. Continued.

| FISHING MORTALITY | | | | | | | |
|--------------------|------|------|------|------|------|------|------|
| | 1982 | 1983 | 1984 | 1985 | 1986 | 1987 | 1988 |
| 1 | 0.01 | 0.02 | 0.01 | 0.01 | 0.01 | 0.01 | 0.01 |
| 2 | 0.25 | 0.12 | 0.16 | 0.07 | 0.09 | 0.10 | 0.12 |
| 3 | 0.74 | 0.27 | 0.41 | 0.45 | 0.31 | 0.50 | 0.35 |
| 4 | 0.97 | 0.64 | 0.46 | 0.97 | 0.46 | 0.87 | 0.62 |
| 5 | 0.71 | 0.68 | 0.71 | 1.27 | 0.43 | 1.08 | 0.69 |
| 6 | 0.87 | 0.39 | 0.65 | 0.87 | 0.85 | 0.79 | 1.36 |
| 7 | 0.76 | 0.56 | 0.70 | 1.22 | 0.57 | 1.01 | 0.88 |
| 8 | 0.76 | 0.56 | 0.70 | 1.22 | 0.57 | 1.01 | 0.88 |
| | 1989 | 1990 | 1991 | 1992 | 1993 | 1994 | 1995 |
| 1 | 0.02 | 0.01 | 0.01 | 0.02 | 0.02 | 0.01 | 0.01 |
| 2 | 0.14 | 0.13 | 0.14 | 0.14 | 0.10 | 0.04 | 0.03 |
| 3 | 0.61 | 0.56 | 0.64 | 0.45 | 0.37 | 0.31 | 0.22 |
| 4 | 1.03 | 1.26 | 0.97 | 1.11 | 0.82 | 0.64 | 1.18 |
| 5 | 1.24 | 1.28 | 1.17 | 0.96 | 0.90 | 0.51 | 1.84 |
| 6 | 1.30 | 1.03 | 1.29 | 1.56 | 0.97 | 0.77 | 1.87 |
| 7 | 1.31 | 1.26 | 1.26 | 1.09 | 0.95 | 0.55 | 2.03 |
| 8 | 1.31 | 1.26 | 1.26 | 1.09 | 0.95 | 0.55 | 2.03 |
| | 1996 | 1997 | 1998 | 1999 | 2000 | 2001 | |
| 1 | 0.01 | 0.01 | 0.01 | 0.00 | 0.00 | 0.00 | |
| 2 | 0.03 | 0.05 | 0.02 | 0.00 | 0.00 | 0.00 | |
| 3 | 0.34 | 0.15 | 0.07 | 0.02 | 0.02 | 0.01 | |
| 4 | 0.41 | 0.41 | 0.18 | 0.06 | 0.08 | 0.08 | |
| 5 | 0.36 | 0.24 | 0.51 | 0.08 | 0.08 | 0.17 | |
| 6 | 0.37 | 0.23 | 0.30 | 0.19 | 0.04 | 0.10 | |
| 7 | 0.37 | 0.24 | 0.44 | 0.10 | 0.07 | 0.06 | |
| 8 | 0.37 | 0.24 | 0.44 | 0.10 | 0.07 | 0.06 | |
| 5,6 | | | | | | | |
| Average F for 5,6 | | | | | | | |
| | 1982 | 1983 | 1984 | 1985 | 1986 | 1987 | 1988 |
| 5,6 | 0.79 | 0.53 | 0.68 | 1.07 | 0.64 | 0.94 | 1.02 |
| | 1989 | 1990 | 1991 | 1992 | 1993 | 1994 | 1995 |
| 5,6 | 1.27 | 1.16 | 1.23 | 1.26 | 0.94 | 0.64 | 1.85 |
| | 1996 | 1997 | 1998 | 1999 | 2000 | 2001 | |
| 5,6 | 0.36 | 0.23 | 0.40 | 0.13 | 0.06 | 0.14 | |
| Biomass Weighted F | | | | | | | |
| | 1982 | 1983 | 1984 | 1985 | 1986 | 1987 | 1988 |
| | 0.60 | 0.33 | 0.42 | 0.70 | 0.30 | 0.55 | 0.40 |
| | 1989 | 1990 | 1991 | 1992 | 1993 | 1994 | 1995 |
| | 0.74 | 0.64 | 0.54 | 0.49 | 0.41 | 0.39 | 0.51 |
| | 1996 | 1997 | 1998 | 1999 | 2000 | 2001 | |
| | 0.20 | 0.17 | 0.14 | 0.05 | 0.05 | 0.07 | |

Table B2.45. Continued.

| BACKCALCULATED PARTIAL RECRUITMENT | | | | | | | |
|--|-------|------|------|------|------|------|------|
| | 1982 | 1983 | 1984 | 1985 | 1986 | 1987 | 1988 |
| 1 | 0.01 | 0.03 | 0.01 | 0.01 | 0.01 | 0.01 | 0.01 |
| 2 | 0.26 | 0.17 | 0.23 | 0.06 | 0.10 | 0.10 | 0.09 |
| 3 | 0.77 | 0.40 | 0.58 | 0.36 | 0.37 | 0.47 | 0.25 |
| 4 | 1.00 | 0.94 | 0.65 | 0.76 | 0.54 | 0.80 | 0.46 |
| 5 | 0.73 | 1.00 | 1.00 | 1.00 | 0.51 | 1.00 | 0.50 |
| 6 | 0.89 | 0.58 | 0.93 | 0.69 | 1.00 | 0.73 | 1.00 |
| 7 | 0.79 | 0.83 | 1.00 | 0.96 | 0.68 | 0.93 | 0.64 |
| 8 | 0.79 | 0.83 | 1.00 | 0.96 | 0.68 | 0.93 | 0.64 |
| | 1989 | 1990 | 1991 | 1992 | 1993 | 1994 | 1995 |
| 1 | 0.01 | 0.01 | 0.01 | 0.01 | 0.02 | 0.01 | 0.00 |
| 2 | 0.11 | 0.10 | 0.11 | 0.09 | 0.10 | 0.06 | 0.01 |
| 3 | 0.46 | 0.44 | 0.49 | 0.29 | 0.38 | 0.40 | 0.11 |
| 4 | 0.78 | 0.98 | 0.75 | 0.71 | 0.84 | 0.83 | 0.58 |
| 5 | 0.95 | 1.00 | 0.91 | 0.62 | 0.93 | 0.66 | 0.91 |
| 6 | 0.99 | 0.81 | 1.00 | 1.00 | 1.00 | 1.00 | 0.92 |
| 7 | 1.00 | 0.98 | 0.98 | 0.70 | 0.98 | 0.72 | 1.00 |
| 8 | 1.00 | 0.98 | 0.98 | 0.70 | 0.98 | 0.72 | 1.00 |
| | 1996 | 1997 | 1998 | 1999 | 2000 | 2001 | |
| 1 | 0.02 | 0.01 | 0.01 | 0.01 | 0.04 | 0.02 | |
| 2 | 0.08 | 0.13 | 0.04 | 0.02 | 0.05 | 0.03 | |
| 3 | 0.82 | 0.36 | 0.13 | 0.09 | 0.22 | 0.06 | |
| 4 | 1.00 | 1.00 | 0.35 | 0.31 | 0.90 | 0.46 | |
| 5 | 0.87 | 0.58 | 1.00 | 0.42 | 1.00 | 1.00 | |
| 6 | 0.88 | 0.56 | 0.58 | 1.00 | 0.47 | 0.61 | |
| 7 | 0.88 | 0.58 | 0.86 | 0.51 | 0.80 | 0.37 | |
| 8 | 0.88 | 0.58 | 0.86 | 0.51 | 0.80 | 0.37 | |
| MEAN BIOMASS (using catch mean weights at age) | | | | | | | |
| | 1982 | 1983 | 1984 | 1985 | 1986 | 1987 | 1988 |
| 1 | 859 | 907 | 334 | 344 | 312 | 188 | 137 |
| 2 | 2586 | 2058 | 1522 | 755 | 1911 | 1287 | 1591 |
| 3 | 2693 | 2611 | 1588 | 1124 | 1242 | 1589 | 1489 |
| 4 | 1782 | 1468 | 1677 | 1000 | 958 | 757 | 932 |
| 5 | 1196 | 826 | 668 | 907 | 581 | 516 | 346 |
| 6 | 581 | 741 | 382 | 375 | 344 | 342 | 140 |
| 7 | 345 | 268 | 385 | 169 | 181 | 149 | 146 |
| 8 | 277 | 275 | 357 | 125 | 132 | 173 | 122 |
| 1+ | 10319 | 9153 | 6914 | 4798 | 5662 | 5000 | 4903 |
| | 1989 | 1990 | 1991 | 1992 | 1993 | 1994 | 1995 |
| 1 | 131 | 153 | 197 | 93 | 90 | 118 | 292 |
| 2 | 604 | 751 | 746 | 714 | 523 | 223 | 417 |
| 3 | 1014 | 716 | 680 | 729 | 760 | 591 | 577 |
| 4 | 873 | 452 | 399 | 304 | 364 | 484 | 280 |
| 5 | 407 | 256 | 123 | 141 | 109 | 211 | 151 |
| 6 | 170 | 130 | 67 | 35 | 47 | 46 | 63 |
| 7 | 47 | 39 | 34 | 24 | 11 | 29 | 13 |
| 8 | 41 | 29 | 31 | 22 | 05 | 10 | 11 |
| 1+ | 3285 | 2527 | 2275 | 2062 | 1909 | 1710 | 1802 |
| | 1996 | 1997 | 1998 | 1999 | 2000 | 2001 | |
| 1 | 199 | 282 | 292 | 320 | 264 | 221 | |
| 2 | 1520 | 1041 | 901 | 581 | 805 | 496 | |
| 3 | 986 | 1696 | 1615 | 1667 | 1847 | 1920 | |
| 4 | 523 | 656 | 1338 | 1584 | 1569 | 1898 | |
| 5 | 137 | 371 | 393 | 1138 | 1354 | 1313 | |
| 6 | 49 | 112 | 278 | 268 | 1060 | 1259 | |
| 7 | 19 | 31 | 82 | 225 | 235 | 1026 | |
| 8 | 34 | 40 | 21 | 175 | 196 | 592 | |
| 1+ | 3466 | 4228 | 4918 | 5957 | 7328 | 8724 | 00 |

Table B2.45. Continued.

SSB AT THE START OF THE SPAWNING SEASON -MALES AND FEMALES (MT) (using SSB mean weights)

| | 1982 | 1983 | 1984 | 1985 | 1986 | 1987 | 1988 |
|----|------|------|------|------|------|------|------|
| 1 | 00 | 00 | 00 | 00 | 00 | 00 | 00 |
| 2 | 00 | 00 | 00 | 00 | 00 | 00 | 00 |
| 3 | 454 | 368 | 265 | 189 | 144 | 257 | 204 |
| 4 | 1685 | 1307 | 1578 | 898 | 744 | 714 | 857 |
| 5 | 1255 | 867 | 778 | 991 | 549 | 571 | 376 |
| 6 | 665 | 733 | 437 | 383 | 362 | 370 | 180 |
| 7 | 390 | 292 | 449 | 194 | 193 | 165 | 167 |
| 8 | 339 | 325 | 433 | 164 | 157 | 220 | 153 |
| 1+ | 4790 | 3890 | 3941 | 2820 | 2149 | 2298 | 1936 |
| | 1989 | 1990 | 1991 | 1992 | 1993 | 1994 | 1995 |
| 1 | 00 | 00 | 00 | 00 | 00 | 00 | 00 |
| 2 | 00 | 00 | 00 | 00 | 00 | 00 | 00 |
| 3 | 185 | 95 | 103 | 107 | 107 | 82 | 54 |
| 4 | 824 | 450 | 365 | 302 | 366 | 444 | 295 |
| 5 | 474 | 323 | 151 | 167 | 124 | 208 | 198 |
| 6 | 183 | 149 | 82 | 42 | 58 | 51 | 87 |
| 7 | 50 | 51 | 46 | 25 | 12 | 26 | 17 |
| 8 | 54 | 38 | 40 | 28 | 06 | 11 | 15 |
| 1+ | 1769 | 1106 | 787 | 672 | 672 | 823 | 666 |
| | 1996 | 1997 | 1998 | 1999 | 2000 | 2001 | |
| 1 | 00 | 00 | 00 | 00 | 00 | 00 | |
| 2 | 00 | 00 | 00 | 00 | 00 | 00 | |
| 3 | 96 | 241 | 194 | 185 | 157 | 189 | |
| 4 | 421 | 596 | 1180 | 1283 | 1315 | 1521 | |
| 5 | 133 | 373 | 428 | 1116 | 1369 | 1335 | |
| 6 | 47 | 107 | 285 | 268 | 1022 | 1211 | |
| 7 | 19 | 33 | 85 | 219 | 227 | 980 | |
| 8 | 39 | 44 | 24 | 188 | 208 | 630 | |
| 1+ | 754 | 1395 | 2197 | 3260 | 4298 | 5866 | |

Table B2.45b. VPA retrospective analysis for Gulf of Maine winter flounder.

Fishing Mortality

Terminal year

| | 1982 | 1983 | 1984 | 1985 | 1986 | 1987 | 1988 | 1989 | 1990 | 1991 | 1992 | 1993 | 1994 | 1995 | 1996 | 1997 | 1998 | 1999 | 2000 | 2001 |
|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|
| 1995 | 0.79 | 0.53 | 0.68 | 1.07 | 0.64 | 0.94 | 1.02 | 1.26 | 1.14 | 1.17 | 1.05 | 0.59 | 0.29 | 0.72 | | | | | | |
| 1996 | 0.79 | 0.53 | 0.68 | 1.07 | 0.64 | 0.94 | 1.02 | 1.27 | 1.15 | 1.22 | 1.21 | 0.85 | 0.52 | 1.05 | 0.07 | | | | | |
| 1997 | 0.79 | 0.53 | 0.68 | 1.07 | 0.64 | 0.94 | 1.02 | 1.27 | 1.15 | 1.22 | 1.22 | 0.87 | 0.55 | 1.19 | 0.14 | 0.09 | | | | |
| 1998 | 0.79 | 0.53 | 0.68 | 1.07 | 0.64 | 0.94 | 1.02 | 1.27 | 1.16 | 1.22 | 1.23 | 0.88 | 0.56 | 1.27 | 0.16 | 0.09 | 0.23 | | | |
| 1999 | 0.79 | 0.53 | 0.68 | 1.07 | 0.64 | 0.94 | 1.02 | 1.27 | 1.16 | 1.23 | 1.25 | 0.91 | 0.6 | 1.54 | 0.23 | 0.13 | 0.21 | 0.09 | | |
| 2000 | 0.79 | 0.53 | 0.68 | 1.07 | 0.64 | 0.94 | 1.02 | 1.27 | 1.16 | 1.23 | 1.25 | 0.93 | 0.63 | 1.73 | 0.30 | 0.19 | 0.27 | 0.08 | 0.06 | |
| 2001 | 0.79 | 0.53 | 0.68 | 1.07 | 0.64 | 0.94 | 1.02 | 1.27 | 1.16 | 1.23 | 1.26 | 0.94 | 0.64 | 1.85 | 0.36 | 0.23 | 0.40 | 0.13 | 0.06 | 0.14 |

Spawning Stock Biomass

Terminal year

| | 1982 | 1983 | 1984 | 1985 | 1986 | 1987 | 1988 | 1989 | 1990 | 1991 | 1992 | 1993 | 1994 | 1995 | 1996 | 1997 | 1998 | 1999 | 2000 | 2001 |
|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|
| 1995 | 4790 | 3890 | 3941 | 2821 | 2150 | 2299 | 1939 | 1776 | 1121 | 831 | 804 | 910 | 1283 | 1759 | | | | | | |
| 1996 | 4790 | 3890 | 3941 | 2820 | 2149 | 2298 | 1937 | 1770 | 1108 | 795 | 695 | 735 | 1080 | 1373 | 2108 | | | | | |
| 1997 | 4790 | 3890 | 3941 | 2820 | 2149 | 2298 | 1937 | 1770 | 1108 | 794 | 690 | 722 | 957 | 1046 | 1510 | 2530 | | | | |
| 1998 | 4790 | 3890 | 3941 | 2820 | 2149 | 2298 | 1936 | 1770 | 1108 | 793 | 688 | 715 | 934 | 1008 | 1417 | 2274 | 2956 | | | |
| 1999 | 4790 | 3890 | 3941 | 2820 | 2149 | 2298 | 1936 | 1769 | 1106 | 789 | 678 | 688 | 868 | 799 | 1137 | 2082 | 2799 | 4038 | | |
| 2000 | 4790 | 3890 | 3941 | 2820 | 2149 | 2298 | 1936 | 1769 | 1106 | 788 | 674 | 678 | 839 | 719 | 873 | 1753 | 2616 | 3601 | 4808 | |
| 2001 | 4790 | 3890 | 3941 | 2820 | 2149 | 2298 | 1936 | 1769 | 1106 | 787 | 672 | 672 | 823 | 666 | 754 | 1395 | 2197 | 3260 | 4298 | 5866 |

Population Numbers Age1:

Terminal year

| | 1982 | 1983 | 1984 | 1985 | 1986 | 1987 | 1988 | 1989 | 1990 | 1991 | 1992 | 1993 | 1994 | 1995 | 1996 | 1997 | 1998 | 1999 | 2000 | 2001 | 2002 |
|------|-------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|-------|-------|------|------|
| 1995 | 11762 | 8779 | 6271 | 9285 | 7698 | 6150 | 4556 | 4377 | 4717 | 5296 | 6200 | 6700 | 6302 | 8273 | 6222 | | | | | | |
| 1996 | 11761 | 8778 | 6269 | 9278 | 7688 | 6129 | 4496 | 4096 | 4336 | 5330 | 5327 | 6547 | 6324 | 7084 | 6987 | 6895 | | | | | |
| 1997 | 11761 | 8778 | 6269 | 9278 | 7688 | 6127 | 4499 | 4067 | 4390 | 4811 | 4419 | 4909 | 6072 | 7098 | 7490 | 7043 | 7090 | | | | |
| 1998 | 11761 | 8778 | 6269 | 9278 | 7688 | 6127 | 4497 | 4061 | 4380 | 4723 | 4402 | 4662 | 5446 | 6768 | 7060 | 7347 | 8617 | 11412 | | | |
| 1999 | 11761 | 8778 | 6269 | 9277 | 7687 | 6126 | 4487 | 4052 | 4283 | 4657 | 3598 | 4474 | 5794 | 7011 | 7774 | 7883 | 9687 | 13335 | 16197 | | |
| 2000 | 11761 | 8778 | 6269 | 9277 | 7686 | 6125 | 4484 | 4045 | 4262 | 4567 | 3482 | 3425 | 5692 | 7749 | 7257 | 7352 | 9106 | 10817 | 8113 | 6990 | |
| 2001 | 11761 | 8778 | 6269 | 9277 | 7686 | 6125 | 4482 | 4043 | 4242 | 4542 | 3322 | 3240 | 4519 | 7503 | 7588 | 7249 | 8967 | 10080 | 7474 | 7391 | 6274 |

Table B2.46. VPA Bootstrap results: precision of estimates.

The number of bootstraps: 500

Bootstrap Output Variable: N hat

| | NLLS ESTIMATE | BOOTSTRAP MEAN | BOOTSTRAP StdError | C.V. FOR NLLS SOLN |
|-----|---------------|----------------|--------------------|--------------------|
| N 1 | 6274 | 6578 | 2984 | 0.48 |
| N 2 | 6033 | 6313 | 1951 | 0.32 |
| N 3 | 4971 | 5148 | 1277 | 0.26 |
| N 4 | 5444 | 5544 | 1043 | 0.19 |
| N 5 | 3624 | 3711 | 674 | 0.19 |
| N 6 | 2001 | 2043 | 394 | 0.20 |
| N 7 | 1572 | 1576 | 273 | 0.17 |
| N 8 | 1068 | 1077 | 170 | 0.16 |

| | BIAS ESTIMATE | BIAS STD ERROR | PERCENT BIAS | NLLS EST CORRECTED FOR BIAS | C.V. FOR CORRECTED ESTIMATE | LOWER 80%CI | UPPER 80%CI |
|-----|---------------|----------------|--------------|-----------------------------|-----------------------------|-------------|-------------|
| N 1 | 304 | 133 | 4.85 | 5969 | 0.499901 | 3546 | 11460 |
| N 2 | 280 | 87 | 4.65 | 5752 | 0.339112 | 3677 | 8478 |
| N 3 | 177 | 57 | 3.56 | 4794 | 0.266440 | 3559 | 6826 |
| N 4 | 100 | 47 | 1.83 | 5344 | 0.195187 | 4245 | 6919 |
| N 5 | 88 | 30 | 2.42 | 3536 | 0.190486 | 2818 | 4483 |
| N 6 | 42 | 18 | 2.11 | 1959 | 0.201157 | 1523 | 2548 |
| N 7 | 04 | 12 | 0.27 | 1568 | 0.173815 | 1286 | 1984 |
| N 8 | 10 | 08 | 0.90 | 1058 | 0.160299 | 874 | 1312 |

Bootstrap Output Variable: F t

| | NLLS ESTIMATE | BOOTSTRAP MEAN | BOOTSTRAP StdError | C.V. FOR NLLS SOLN |
|-------|---------------|----------------|--------------------|--------------------|
| Age 1 | 0.0030 | 0.0032 | 0.0011 | 0.37 |
| Age 2 | 0.0044 | 0.0045 | 0.0011 | 0.25 |
| Age 3 | 0.0096 | 0.0097 | 0.0018 | 0.19 |
| Age 4 | 0.0790 | 0.0795 | 0.0139 | 0.18 |
| Age 5 | 0.1708 | 0.1730 | 0.0311 | 0.18 |
| Age 6 | 0.1048 | 0.1074 | 0.0180 | 0.17 |
| Age 7 | 0.0624 | 0.0633 | 0.0098 | 0.16 |
| Age 8 | 0.0624 | 0.0633 | 0.0098 | 0.16 |

| | BIAS ESTIMATE | BIAS STD ERROR | PERCENT BIAS | NLLS EST CORRECTED FOR BIAS | C.V. FOR CORRECTED ESTIMATE | LOWER 80%CI | UPPER 80%CI |
|-------|---------------|----------------|--------------|-----------------------------|-----------------------------|-------------|-------------|
| Age 1 | 0.0001690 | 0.0000491 | 5.641 | 0.0028262 | 0.39 | 0.0021 | 0.0048 |
| Age 2 | 0.0001079 | 0.0000490 | 2.477 | 0.0042508 | 0.26 | 0.0032 | 0.0061 |
| Age 3 | 0.0001559 | 0.0000812 | 1.625 | 0.0094377 | 0.19 | 0.0075 | 0.0122 |
| Age 4 | 0.0005479 | 0.0006197 | 0.694 | 0.0784051 | 0.18 | 0.0640 | 0.0998 |
| Age 5 | 0.0021128 | 0.0013886 | 1.237 | 0.1687286 | 0.18 | 0.1359 | 0.2178 |
| Age 6 | 0.0025929 | 0.0008055 | 2.474 | 0.1022216 | 0.18 | 0.0836 | 0.1266 |
| Age 7 | 0.0009146 | 0.0004373 | 1.465 | 0.0614994 | 0.16 | 0.0506 | 0.0752 |
| Age 8 | 0.0009146 | 0.0004373 | 1.465 | 0.0614994 | 0.16 | 0.0506 | 0.0752 |

Bootstrap Output Variable: Mean Biomass

| | NLLS ESTIMATE | BOOTSTRAP MEAN | BOOTSTRAP StdError | C.V. FOR NLLS SOLN |
|--|---------------|----------------|--------------------|--------------------|
| | 8723.9106 | 8873.3264 | 775.7433 | 0.09 |

| | BIAS ESTIMATE | BIAS STD ERROR | PERCENT BIAS | NLLS EST CORRECTED FOR BIAS | C.V. FOR CORRECTED ESTIMATE | LOWER 80%CI | UPPER 80%CI |
|--|---------------|----------------|--------------|-----------------------------|-----------------------------|-------------|-------------|
| | 149.4158 | 34.6923 | 1.71 | 8574.4947 | 0.09 | 7730.5482 | 9603.4137 |

Bootstrap Output Variable: SSB spawn t

| | NLLS ESTIMATE | BOOTSTRAP MEAN | BOOTSTRAP StdError | C.V. FOR NLLS SOLN |
|--|---------------|----------------|--------------------|--------------------|
| | 5865.7415 | 5945.3298 | 554.7207 | 0.09 |

| | BIAS ESTIMATE | BIAS STD ERROR | PERCENT BIAS | NLLS EST CORRECTED FOR BIAS | C.V. FOR CORRECTED ESTIMATE | LOWER 80%CI | UPPER 80%CI |
|--|---------------|----------------|--------------|-----------------------------|-----------------------------|-------------|-------------|
| | 79.59 | 24.81 | 1.36 | 5786.15 | 0.10 | 5203.0726 | 6580.6435 |

Table B2.47. Yield Per Recruit analysis for Gulf of Maine winter flounder.

```

The NEFC Yield and Stock Size per Recruit Program - PDBYPRC
PC Ver.2.0 [Method of Thompson and Bell (1934)] 1-Jan-1999
-----
Run Date: 3-10-2002; Time: 12:05:35.00
gulf of Maine Winter Flounder - 1999-01 PR, Mean Weights at Age from
-----
Proportion of F before spawning: 0.2500
Proportion of M before spawning: 0.2500
Natural Mortality is Constant at: 0.200
Initial age is: 1; Last age is: 15
Last age is a TRUE Age;
Original age-specific PRs, Mats, and Mean Wts from file:
==> C:\Program Files\FACT\wv\ypr\gomwfy3.dat
-----
Age-specific Input data for Yield per Recruit Analysis
-----

```

| Age | Fish Mort Pattern | Nat Mort Pattern | Proportion Mature | Average Weights Catch Stock | |
|-----|----------------------|---------------------|----------------------|--------------------------------|-------|
| 1 | 0.0300 | 1.0000 | 0.0000 | 0.036 | 0.021 |
| 2 | 0.0400 | 1.0000 | 0.0000 | 0.095 | 0.059 |
| 3 | 0.1300 | 1.0000 | 0.1600 | 0.351 | 0.206 |
| 4 | 0.5700 | 1.0000 | 0.8600 | 0.471 | 0.420 |
| 5 | 1.0000 | 1.0000 | 1.0000 | 0.550 | 0.512 |
| 6 | 1.0000 | 1.0000 | 1.0000 | 0.691 | 0.626 |
| 7 | 1.0000 | 1.0000 | 1.0000 | 0.872 | 0.788 |
| 8 | 1.0000 | 1.0000 | 1.0000 | 0.993 | 0.993 |
| 9 | 1.0000 | 1.0000 | 1.0000 | 1.091 | 1.091 |
| 10 | 1.0000 | 1.0000 | 1.0000 | 1.171 | 1.171 |
| 11 | 1.0000 | 1.0000 | 1.0000 | 1.234 | 1.234 |
| 12 | 1.0000 | 1.0000 | 1.0000 | 1.284 | 1.284 |
| 13 | 1.0000 | 1.0000 | 1.0000 | 1.323 | 1.323 |
| 14 | 1.0000 | 1.0000 | 1.0000 | 1.353 | 1.353 |
| 15 | 1.0000 | 1.0000 | 1.0000 | 1.377 | 1.377 |

```

-----
Summary of Yield per Recruit Analysis:
-----
Slope of the Yield/Recruit Curve at F=0.00: --> 2.0105
F level at slope=1/10 of the above slope (F0.1): -----> 0.258
Yield/Recruit corresponding to F0.1: -----> 0.1970
F level to produce Maximum Yield/Recruit (Fmax): -----> 0.687
Yield/Recruit corresponding to Fmax: -----> 0.2201
F level at 40 % of Max Spawning Potential (F40): -----> 0.261
SSB/Recruit corresponding to F40: -----> 0.8333
-----
1
Listing of Yield per Recruit Results for:
-----

```

| | FMORT | TOTCTHN | TOTCTHW | TOTSTKN | TOTSTKW | SPNSTKN | SPNSTKW | % MSP |
|------|-------|---------|---------|---------|---------|---------|---------|--------|
| | 0.00 | 0.00000 | 0.00000 | 5.2420 | 2.4078 | 2.6476 | 2.0834 | 100.00 |
| | 0.10 | 0.17406 | 0.12996 | 4.5658 | 1.6980 | 1.9691 | 1.3773 | 66.11 |
| | 0.20 | 0.26851 | 0.18214 | 4.1562 | 1.3009 | 1.5634 | 0.9877 | 47.41 |
| F0.1 | 0.26 | 0.30487 | 0.19700 | 3.9894 | 1.1500 | 1.4000 | 0.8411 | 40.37 |
| F40% | 0.26 | 0.30682 | 0.19770 | 3.9802 | 1.1419 | 1.3911 | 0.8333 | 40.00 |
| | 0.30 | 0.32662 | 0.20421 | 3.8874 | 1.0616 | 1.3007 | 0.7557 | 36.27 |
| | 0.40 | 0.36623 | 0.21387 | 3.6983 | 0.9070 | 1.1185 | 0.6074 | 29.16 |
| | 0.50 | 0.39537 | 0.21807 | 3.5575 | 0.8010 | 0.9848 | 0.5067 | 24.32 |
| | 0.60 | 0.41803 | 0.21972 | 3.4476 | 0.7243 | 0.8823 | 0.4345 | 20.85 |
| Fmax | 0.69 | 0.43413 | 0.22009 | 3.3697 | 0.6733 | 0.8108 | 0.3869 | 18.57 |
| | 0.70 | 0.43638 | 0.22010 | 3.3588 | 0.6664 | 0.8009 | 0.3805 | 18.26 |
| | 0.80 | 0.45170 | 0.21982 | 3.2847 | 0.6211 | 0.7343 | 0.3387 | 16.26 |
| | 0.90 | 0.46481 | 0.21920 | 3.2215 | 0.5846 | 0.6787 | 0.3053 | 14.66 |
| | 1.00 | 0.47624 | 0.21839 | 3.1666 | 0.5544 | 0.6314 | 0.2781 | 13.35 |
| | 1.10 | 0.48637 | 0.21747 | 3.1180 | 0.5288 | 0.5905 | 0.2553 | 12.25 |
| | 1.20 | 0.49545 | 0.21650 | 3.0745 | 0.5069 | 0.5547 | 0.2359 | 11.32 |
| | 1.30 | 0.50368 | 0.21549 | 3.0352 | 0.4878 | 0.5230 | 0.2193 | 10.52 |
| | 1.40 | 0.51121 | 0.21446 | 2.9992 | 0.4708 | 0.4947 | 0.2047 | 9.83 |
| | 1.50 | 0.51816 | 0.21343 | 2.9660 | 0.4557 | 0.4693 | 0.1919 | 9.21 |
| | 1.60 | 0.52462 | 0.21238 | 2.9352 | 0.4421 | 0.4463 | 0.1805 | 8.67 |
| | 1.70 | 0.53064 | 0.21134 | 2.9065 | 0.4297 | 0.4253 | 0.1703 | 8.18 |
| | 1.80 | 0.53630 | 0.21029 | 2.8795 | 0.4183 | 0.4060 | 0.1611 | 7.73 |
| | 1.90 | 0.54163 | 0.20924 | 2.8541 | 0.4078 | 0.3883 | 0.1527 | 7.33 |
| | 2.00 | 0.54668 | 0.20819 | 2.8300 | 0.3981 | 0.3719 | 0.1451 | 6.96 |

Table B2.48. Stock-recruitment model comparison for Gulf of Maine winter flounder.

| | Prior | Prior | Prior | Prior | Prior | Prior | Prior | Prior | Prior | Prior |
|----------------------------------|-------------------------------------|--|--------|--|--------|--|-------------|-------------|--|--|
| | 1 | 0 | 1 | 0 | 1 | 0 | 0 | 0 | 0 | 0 |
| | BH | ABH | PBH | PABH | PRBH | PRABH | RK | ARK | PRK | PARK |
| Posterior Probability | 0.36 | 0.00 | 0.32 | 0.00 | 0.31 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 |
| Odds Ratio for Most Likely Model | 1.00 | | 1.12 | | 1.16 | | | | | |
| Normalized Likelihood | 0.363 | 0.000 | 0.323 | 0.000 | 0.313 | 0.000 | 0.000 | 0.000 | 0.000 | 0.000 |
| Model AIC Ratio | 1.160 | 0 | 1.033 | 0 | 1.000 | 0 | 0 | 0 | 0 | 0 |
| | BH | ABH | PBH | PABH | PRBH | PRABH | RK | ARK | PRK | PARK |
| Number_of_data_points | 20 | 20 | 20 | 20 | 20 | 20 | 20 | 20 | 20 | 20 |
| Number_of_parameters | 3 | 4 | 3 | 4 | 3 | 4 | 3 | 4 | 3 | 4 |
| Fit_negloglikelihood | 41.146 | 33.566 | 41.263 | 33.724 | 41.295 | 33.732 | 43.534 | 34.926 | 52.285 | 37.530 |
| Penalty_steepness | 0 | 0 | -0.810 | -1.087 | 0 | 0 | 0 | 0 | 0 | 0 |
| Penalty_slope | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 3.160 | -0.774 |
| Penalty_unfished_R | 0 | 0 | 0 | 0 | 2.085 | 1.809 | 0 | 0 | 0 | 0 |
| Negative_loglikelihood | 41.146 | 33.566 | 40.452 | 32.637 | 43.380 | 35.541 | 43.534 | 34.926 | 55.445 | 36.756 |
| Bias-corrected_AIC | 89.792 | 77.799 | 90.025 | 78.115 | 90.090 | 78.130 | 94.568 | 80.519 | 112.070 | 85.726 |
| Diagnostic Comments | Most likely parametric model | Power spectrum dominant frequency exceeds 1/2 time series length | | Power spectrum dominant frequency exceeds 1/2 time series length | | Power spectrum dominant frequency exceeds 1/2 time series length | Fmsy>> Fmax | Fmsy>> Fmax | no stock recruit data at SSB where density dependence is predicted | Power spectrum dominant frequency exceeds 1/2 time series length |

Table B2.48. Continued.

| Parameter Point Estimate | BH | ABH | PBH | PABH | PRBH | PRABH | RK | ARK | PRK | PARK |
|---------------------------------|-----------|------------|------------|-------------|-------------|--------------|-----------|------------|------------|-------------|
| MSY | 1.543 | 1.587 | 1.596 | 1.623 | 1.640 | 1.771 | 1.753 | 1.836 | 2.153 | 0.568 |
| FMSY | 0.430 | 0.415 | 0.405 | 0.380 | 0.410 | 0.395 | 0.745 | 0.705 | 0.375 | 0.240 |
| SMSY | 4.104 | 4.359 | 4.484 | 4.830 | 4.554 | 5.087 | 2.871 | 3.154 | 6.485 | 2.594 |
| Alpha | 7.706 | 8.051 | 8.167 | 8.579 | 8.365 | 9.161 | 2.043 | 1.982 | 1.296 | 0.828 |
| expected_alpha | 8.084 | 8.422 | 8.574 | 8.998 | 8.783 | 9.612 | 2.171 | 2.097 | 1.500 | 1.431 |
| Beta | 0.387 | 0.473 | 0.516 | 0.698 | 0.516 | 0.636 | -0.359 | -0.323 | -0.134 | -0.281 |
| Steepness | 0.923 | 0.911 | 0.905 | 0.881 | 0.907 | 0.896 | | | | |
| R_at_input_SMAX | 7.302 | 7.542 | 7.606 | 7.800 | 7.791 | 8.398 | 4.388 | 5.310 | 10.032 | 2.233 |
| expected_R_at_input_SMAX | 7.661 | 7.889 | 7.985 | 8.182 | 8.180 | 8.811 | 4.663 | 5.618 | 11.611 | 3.862 |
| unfished_S | 18.138 | 18.883 | 19.118 | 19.925 | 19.594 | 21.389 | 8.144 | 8.863 | 16.247 | 6.058 |
| unfished_R | 7.544 | 7.855 | 7.952 | 8.288 | 8.150 | 8.897 | 3.387 | 3.686 | 6.758 | 2.520 |
| Sigma | 0.310 | 0.300 | 0.312 | 0.309 | 0.312 | 0.310 | 0.349 | 0.336 | 0.541 | 1.047 |
| Phi | | 0.720 | | 0.734 | | 0.736 | | 0.749 | | 0.973 |
| Sigmaw | | 0.208 | | 0.210 | | 0.210 | | 0.222 | | 0.240 |
| last_residual_R | | -1.177 | | -1.392 | | -1.991 | | -0.141 | | 3.699 |
| last_logresidual_R | | -0.172 | | -0.200 | | -0.276 | | -0.022 | | 0.890 |
| expected_lognormal_error_term | 1.049 | 1.046 | 1.050 | 1.049 | 1.050 | 1.049 | 1.063 | 1.058 | 1.157 | 1.729 |
| prior_mean_steepness | | | 0.80 | 0.80 | | | | | | |
| prior_se_steepness | | | 0.09 | 0.09 | | | | | | |
| prior_mean_slope | | | | | | | | | 0.79 | 0.79 |
| prior_se_slope | | | | | | | | | 0.18 | 0.18 |
| prior_mean_unfished_R | | | | | 10.09 | 10.09 | | | | |
| prior_se_unfished_R | | | | | 2.06 | 2.06 | | | | |

Table B2.49. Input parameters and stochastic projection results for Gulf of Maine winter flounder using recruitment predicted from the Beverton-Holt stock-recruitment model and an estimated $F_{msy} = 0.43$.

| Age | Stock Size on 1 Jan 2002 (000s) | Fishing Mortality Pattern | Proportion Landed | Proportion mature | Mean Weights Spawning Stock | Mean Weights Landings | Mean Weights Discards |
|-----|--|---------------------------------|----------------------|----------------------|--------------------------------------|-----------------------------|-----------------------------|
| 1 | 6274 | 0.030 | 0.000 | 0.000 | 0.021 | 0.000 | 0.036 |
| 2 | 6033 | 0.040 | 0.040 | 0.000 | 0.059 | 0.000 | 0.089 |
| 3 | 4971 | 0.130 | 0.710 | 0.160 | 0.203 | 0.399 | 0.229 |
| 4 | 5444 | 0.570 | 0.940 | 0.860 | 0.419 | 0.480 | 0.306 |
| 5 | 3624 | 1.000 | 0.980 | 1.000 | 0.512 | 0.553 | 0.389 |
| 6 | 2001 | 1.000 | 0.980 | 1.000 | 0.626 | 0.696 | 0.468 |
| 7 | 1572 | 1.000 | 0.990 | 1.000 | 0.788 | 0.875 | 0.694 |
| 8+ | 1558 | 1.000 | 0.990 | 1.000 | 1.100 | 1.105 | 0.867 |

F2002 is assumed equal to F2001; F during 2003-2013 = $F_{msy} = 0.43$.

| Forecast Medians (50% probability level) | | | | | | | | | | | |
|--|------|------|-----|----------------|------|------|-----|----------------|------|------|-----|
| 2002 | | | | 2003 | | | | 2013 | | | |
| 000s Metric tons | | | | | | | | | | | |
| F | Land | Disc | SSB | F | Land | Disc | SSB | F | Land | Disc | SSB |
| 0.14 | 0.9 | <0.1 | 7.6 | $F_{msy}=0.43$ | 2.9 | 0.1 | 7.8 | $F_{msy}=0.43$ | 1.5 | 0.1 | 4.3 |

F2002 is assumed $0.85 \cdot F_{2001}$ (15% decrease in F from 2001 to 2002); F during 2003-2013 = $F_{msy} = 0.43$.

| Forecast Medians (50% probability level) | | | | | | | | | | | |
|--|------|------|-----|----------------|------|------|-----|----------------|------|------|-----|
| 2002 | | | | 2003 | | | | 2013 | | | |
| 000s Metric tons | | | | | | | | | | | |
| F | Land | Disc | SSB | F | Land | Disc | SSB | F | Land | Disc | SSB |
| 0.12 | 0.8 | <0.1 | 7.7 | $F_{msy}=0.43$ | 2.9 | 0.1 | 7.9 | $F_{msy}=0.43$ | 1.6 | 0.1 | 4.3 |