

Click here for  
[DISCLAIMER](#)

Document starts on next page

Amy L. Leaberry

United States  
Environmental Protection  
Agency

Office of Water  
Regulations and Standards  
Criteria and Standards Division  
Washington DC 20460

EPA 440/5-86-005  
September 1986




---

Water

---

# Ambient Water Quality Criteria for

## Chlorpyrifos - 1986



AMBIENT AQUATIC LIFE WATER QUALITY CRITERIA FOR  
CHLORPYRIFOS

U.S. ENVIRONMENTAL PROTECTION AGENCY  
OFFICE OF RESEARCH AND DEVELOPMENT  
ENVIRONMENTAL RESEARCH LABORATORIES  
DULUTH, MINNESOTA  
NARRAGANSETT, RHODE ISLAND

## NOTICES

This document has been reviewed by the Criteria and Standards Division, Office of Water Regulations and Standards, U.S. Environmental Protection Agency, and approved for publication.

Mention of trade names or commercial products does not constitute endorsement or recommendation for use.

This document is available to the public through the National Technical Information Service (NTIS), 5285 Port Royal Road, Springfield, VA 22161.

## FOREWORD

Section 304(a)(1) of the Clean Water Act of 1977 (P.L. 95-217) requires the Administrator of the Environmental Protection Agency to publish water quality criteria that accurately reflect the latest scientific knowledge on the kind and extent of all identifiable effects on health and welfare that might be expected from the presence of pollutants in any body of water, including ground water. This document is a revision of proposed criteria based upon consideration of comments received from other Federal agencies, State agencies, special interest groups, and individual scientists. Criteria contained in this document replace any previously published EPA aquatic life criteria for the same pollutant(s).

The term "water quality criteria" is used in two sections of the Clean Water Act, section 304(a)(1) and section 303(c)(2). The term has a different program impact in each section. In section 304, the term represents a non-regulatory, scientific assessment of ecological effects. Criteria presented in this document are such scientific assessments. If water quality criteria associated with specific stream uses are adopted by a State as water quality standards under section 303, they become enforceable maximum acceptable pollutant concentrations in ambient waters within that State. Water quality criteria adopted in State water quality standards could have the same numerical values as criteria developed under section 304. However, in many situations States might want to adjust water quality criteria developed under section 304 to reflect local environmental conditions and human exposure patterns before incorporation into water quality standards. It is not until their adoption as part of State water quality standards that criteria become regulatory.

Guidelines to assist States in the modification of criteria presented in this document, in the development of water quality standards, and in other water-related programs of this Agency, have been developed by EPA.

William A. Whittington  
Director  
Office of Water Regulations and Standards

## ACKNOWLEDGMENTS

Loren J. Larson  
(freshwater author)  
University of Wisconsin-Superior  
Superior, Wisconsin

Jeffrey L. Hyland  
Robert E. Hillman  
(saltwater authors)  
Battelle New England Laboratory  
Duxbury, Massachusetts

Charles E. Stephan  
(document coordinator)  
Environmental Research Laboratory  
Duluth, Minnesota

David J. Hansen  
(saltwater coordinator)  
Environmental Research Laboratory  
Narragansett, Rhode Island

Clerical Support: Shelley A. Heintz  
Nancy J. Jordan  
Terry L. Highland  
Diane L. Spehar

## CONTENTS

	<u>Page</u>
Foreword . . . . .	iii
Acknowledgments . . . . .	iv
Tables . . . . .	vi
Introduction . . . . .	1
Acute Toxicity to Aquatic Animals . . . . .	4
Chronic Toxicity to Aquatic Animals . . . . .	6
Toxicity to Aquatic Plants . . . . .	7
Bioaccumulation . . . . .	8
Other Data . . . . .	9
Unused Data . . . . .	13
Summary . . . . .	15
National Criteria . . . . .	17
References . . . . .	41

TABLES

	<u>Page</u>
1. Acute Toxicity of Chlorpyrifos to Aquatic Animals . . . . .	18
2. Chronic Toxicity of Chlorpyrifos To Aquatic Animals . . . . .	24
3. Ranked Genus Mean Acute Values with Species Mean Acute-Chronic Ratios . . . . .	27
4. Toxicity of Chlorpyrifos to Aquatic Plants . . . . .	30
5. Bioaccumulation of Chlorpyrifos by Aquatic Organisms . . . . .	31
6. Other Data on Effects of Chlorpyrifos on Aquatic Organisms . . . . .	33



## Introduction\*

Chlorpyrifos\*\* is one of several organophosphorus compounds developed in the 1960s to replace persistent organochlorine pesticides. It is the active ingredient in various products designed to control a variety of pests including fire ants, turf and ornamental plant insects, mosquitos, cockroaches, termites, lice, and hornflies. In the agricultural industry in the United States, chlorpyrifos is used primarily to control pests on cotton, peanuts, and sorghum. In the past it was directly applied to aquatic environments in mosquito, midge, and blackfly abatement projects, but the current label states that it is not to be applied directly to bodies of water. Gray (1965) and Marshall and Roberts (1978) have reviewed its composition and physical and chemical properties.

Chlorpyrifos is available for pesticide applications as emulsifiable concentrates, wettable powders, dusts, granules, and controlled-release polymers. The resulting concentrations of chlorpyrifos in water and its persistence varies from one form to another. When applied to water, emulsifiable concentrates and wettable powders generally produce a large increase in chlorpyrifos concentrations immediately after application. The concentration in water rapidly declines as chlorpyrifos is sorbed onto sediments and suspended organics. Granules and controlled-release forms do not produce as rapid an increase in the concentration in water, but the resulting concentration has a longer duration.

---

\* An understanding of the "Guidelines for Deriving Numerical National Water Quality Criteria for the Protection of Aquatic Organisms and Their Uses" (Stephan et al. 1985), hereafter referred to as the Guidelines, and the response to public comment (U.S. EPA 1985a) is necessary in order to understand the following text, tables, and calculations.

\*\*Dursban® and Lorsban® are trade names owned by the Dow Chemical Company, Midland, MI for chlorpyrifos.

The percentage of active ingredient in available formulations varies considerably, both between formulations and within a single formulation over time as manufacturers' specifications change. Such variations presumably result in large changes in the amount, and possibly the identity, of the unspecified ingredients. These ingredients are considered inert, although technical-grade chlorpyrifos has generally been found to be more toxic than an equal quantity of active ingredient in a formulation (Darwazeh and Mulla 1974; Jarvinen and Tanner 1982; Siefert et al. 1984). For this reason, the effects of the inert ingredients can not be discounted. Furthermore, under normal application conditions, the commercial formulations are often combined with petroleum products, such as No. 2 diesel oil and kerosene, to increase the rate of dispersal. Such solvents have been shown to have significant toxic effects in addition to those associated with chlorpyrifos (Jamnback and Frempong-Boadu 1966; Wallace et al. 1973).

The toxicity of chlorpyrifos is probably the result of metabolic conversion to its oxygen analogue, chlorpyrifos-oxon, and its subsequent inhibition of various enzymes (e.g., cholinesterases, carboxylases, acetylcholinesterases, and mitochondrial oxidative phosphorylases). Interference with acetylcholinesterase (AChE) is generally accepted to be the major mode of action of organophosphorus pesticides. Inhibition of AChE results in accumulation of the neurotransmitter, acetylcholine, in synapses and disruption of normal neural transmission. Although even substantial reductions in brain AChE activity have not always been fatal to fish, the effect of this condition on such functions as feeding and reproduction in nature is not known.

Chlorpyrifos enters both freshwater and saltwater ecosystems primarily as drift from spraying, and on particles eroded from treated areas. Because

of its affinity for organic soils, little leaching occurs. When unbound chlorpyrifos enters an aquatic system, it appears to be rapidly sorbed to suspended organics and sediment, although some is removed by volatilization and degradation. Its penetration into sediment appears to be shallow, with most occurring in the upper several millimeters. Menzie (1969) reported that chlorpyrifos remained stable for long periods of time under the acidic conditions (pH = 5 to 6) found in some salt marshes.

Use of slow-release polymers in water probably results in differential exposure, both in concentration and duration, between benthic and pelagic organisms. Organisms inhabiting the water-sediment interface probably receive larger and more sustained concentrations than free-swimming organisms. Evans (1977) found concentrations of chlorpyrifos that were still toxic to mosquito larvae one year after application of a slow-release polymer formulation to a natural pond.

Because chlorpyrifos is rapidly metabolized by fish, with 3,5,6-trichloro-2-pyridinol being the major product (Marshall and Roberts 1978), concentrations in wild fishes (Clark et al. 1984; Mulla et al. 1973) and cultured or experimental fishes (Macek et al. 1972; Siefert et al. 1984; Winterlin et al. 1968) are generally low. Braun and Frank (1980), Hughes (1977), Hughes et al. (1980), Hurlbert et al. (1970), Macalady and Wolfe (1985), Nelson and Evans (1973), Rawn et al. (1978), Siefert et al. (1984), and Winterlin et al. (1968) have reported concentrations of chlorpyrifos in natural sediment and water samples.

Unless otherwise noted, all concentrations reported herein are expressed as chlorpyrifos, not as the material tested. Whenever adequately justified, a national criterion may be replaced by a site-specific criterion (U.S. EPA 1983a), which may include not only

site-specific criterion concentrations (U.S. EPA 1983b), but also site-specific durations of averaging periods and site-specific frequencies of allowed excursions (U.S. EPA 1985b). The latest comprehensive literature search for information for this document was conducted in July, 1986; some more recent information might have been included.

#### Acute Toxicity to Aquatic Animals

Data, which are usable according to the Guidelines, on the acute toxicity of chlorpyrifos to freshwater animals are available for seven fish species and eleven invertebrate species (Table 1). Invertebrates are among the most sensitive and most resistant species, and the nine most sensitive species are arthropods. Although data are available for eighteen species, no data are available for a planktonic crustacean, and the snail, Aplexa hypnorum, is the only one of the eighteen that is not an arthropod or a fish. Within arthropods and fishes separately, and within all species combined, there appears to be an inverse relationship between size and sensitivity to chlorpyrifos.

Species Mean Acute Values (Table 1) were calculated as geometric means of the available acute values, and then Genus Mean Acute Values (Table 3) were calculated as geometric means of the available freshwater Species Mean Acute Values. The most sensitive genus, Gammarus, is more than 4,300 times more sensitive than the most resistant genera, Aplexa, Carassius, and Ictalurus, but the four most sensitive genera are within a factor of 4, and all are invertebrates. Acute values are available for more than one species in each of two genera, and the range of Species Mean Acute Values within each genus is less than a factor of 3. The freshwater Final Acute Value for chlorpyrifos was calculated to be 0.1669  $\mu\text{g/L}$

using the procedure described in the Guidelines and the Genus Mean Acute Values in Table 3. Thus the freshwater Final Acute Value is higher than the Species Mean Acute Value for one of three amphipods in the genus Gammarus.

Tests of the acute toxicity of chlorpyrifos to saltwater animals have been conducted with five species of invertebrates and ten species of fish (Table 1). The range of acute values extends from 0.01 µg/L for adult Korean shrimp, Palaemon macrodactylus, (Earnest 1970) to 1,991 µg/L for larvae of the eastern oyster, Crassostrea virginica, (Borthwick and Walsh 1981). Four species of saltwater arthropods have been tested and they were all more sensitive than the most sensitive fish species. The range of acute toxicity values for fish is narrower than for invertebrates, with LC50s extending from 0.4 µg/L for 14-day-old larvae of the tidewater silverside, Menidia peninsulae, (Borthwick et al. 1985) to 520 µg/L for juveniles of the gulf toadfish, Opsanus beta, (Hansen et al. 1986).

Borthwick et al. (1985) conducted a series of 96-hr acute tests under both static and flow-through conditions with four different ages of larvae of three estuarine fishes (Table 1). LC50s ranged from 0.4 to 5.5 µg/L for all tests. In static tests, 14-day-old larvae were more sensitive than newly hatched or 28-day larvae of all species. In flow-through tests, relative sensitivities of the ages were similar to those in static tests for tidewater silverside, decreased with age for Atlantic silverside, and differed little for California grunion (Table 1).

Of the twelve genera for which saltwater Genus Mean Acute Values are available, the most sensitive genus, Mysidopsis, is about 57,000 times more sensitive than the most resistant genus, Crassostrea (Table 3). Acute values are available for more than one species in each of two genera, and the range of Species Mean Acute Values within each genus is

less than a factor of 5.7. The saltwater Final Acute Value was calculated to be 0.02284 µg/L, which is lower than the lowest Genus Mean Acute Value.

#### Chronic Toxicity to Aquatic Animals

Usable data on the chronic toxicity of chlorpyrifos are available for only one freshwater species, the fathead minnow. Chronic values for technical-grade and encapsulated material were 2.26 µg/L and 3.25 µg/L, respectively, in early life-stage tests (Jarvinen and Tanner 1982). Growth over the 32-day test was the most sensitive parameter with technical-grade chlorpyrifos, whereas with the encapsulated formulation, growth and survival were equally sensitive. In a life-cycle test with the same species (Jarvinen et al. 1983), unacceptable effects occurred at 0.41 µg/L in the first generation and at 0.12 µg/L in the second generation, showing rather poor agreement between the early life-stage tests and the life-cycle test. Based on these results, the acute-chronic ratio for chlorpyrifos is greater than 1,417 with the fathead minnow. Jarvinen et al. (1983) also estimated the chronic effect of chlorpyrifos on the viable biomass recruitment of a natural population of the fathead minnow.

Data on the chronic toxicity of chlorpyrifos to saltwater animals are available for the mysid, Mysidopsis bahia, and six fishes. In the 28-day life-cycle test with the mysid, survival and reproduction were reduced at 42 µg/L, and growth was significantly reduced at a nominal concentration of 0.004 µg/L (McKenney et al. 1981). This nominal concentration, which was less than the limit of detection of the analytical method used, is likely representative of the actual concentration because the test concentrations that could be measured averaged  $10 \pm 2.5\%$  of the nominal

concentrations. Of the six saltwater fishes exposed to chlorpyrifos in early life-stage toxicity tests, the California grunion was the most sensitive. Decreased weight was the most sensitive endpoint for this species (Goodman et al. 1985a), the sheepshead minnow (Cripe et al. 1986), and the gulf toadfish (Hansen et al. 1986). Decreased survival was the most sensitive endpoint with the three species of Menidia, although growth was also affected with two of these species (Goodman et al. 1985b).

The Species Mean Acute-Chronic Ratios for the seven saltwater species range from 1.374 to 228.5, whereas that for the freshwater fathead minnow is greater than 1,417. However, the ratios for the five sensitive species only range from 1.374 to 12.50. Thus it seems reasonable to calculate the Final Acute-Chronic Ratio for chlorpyrifos as the geometric mean of these five. Division of the freshwater and saltwater Final Acute Values by the Final Acute-Chronic Ratio of 4.064 results in Final Chronic Values of 0.04107  $\mu\text{g/L}$  and 0.005620  $\mu\text{g/L}$ , respectively. The freshwater value is about a factor of three lower than the 0.12  $\mu\text{g/L}$  that affected the fathead minnow, which is an acutely insensitive species. The saltwater value is a factor of two higher than the chronic value for the most acutely sensitive saltwater species, Mysidopsis bahia.

#### Toxicity to Aquatic Plants

Several field studies have examined the effects of chlorpyrifos on phytoplankton under more or less natural conditions (Brown et al. 1976; Butcher et al. 1975,1977; Hughes et al. 1980; Hurlbert 1969; Hurlbert et al. 1972; Papst and Boyer 1980). All used an emulsifiable concentrate of chlorpyrifos, which makes them inappropriate for inclusion in Table 4, but the general trends identified are germane to a discussion of the

effects of chlorpyrifos under natural conditions. With the exception of Brown et al. (1976), all observed increased phytoplankton numbers after application of chlorpyrifos. This change is generally accepted not to be a direct effect, but rather a result of changes in the herbivore-algal relationship caused by large reductions in herbivorous zooplankton populations. Papst and Boyer (1980) attempted to substantiate this hypothesis experimentally by monitoring concentrations of pheopigments, the major chlorophyll degradation product of herbivory, after chlorpyrifos application. Although they found reductions in pheopigments, the effect was delayed. They did observe rapid increases in the numbers of microzooplankton (e.g., rotifers) immediately after chlorpyrifos application, presumably due to reduced competition with macrozooplankton. Other studies have also observed an increase in microzooplankton after chlorpyrifos treatment (Hughes 1977; Hurlbert et al. 1970, 1972; Siefert et al. 1984). Although increased phytoplankton numbers might be explained by release from herbivory, another possible factor is increased phosphate concentration from decomposition of chlorpyrifos and from decomposition of intoxicated organisms (Butcher et al. 1977).

The concentrations of chlorpyrifos reducing growth or survival of six saltwater species of phytoplankton range from 138 to 10,000  $\mu\text{g/L}$  (Tables 4 and 6). These concentrations are well above those that are acutely lethal to saltwater animals. Therefore, criteria derived using data on the toxicity of chlorpyrifos to saltwater animals will probably also protect saltwater plants.

#### Bioaccumulation

Although chlorpyrifos is hydrophobic, which would suggest its accumulation in tissues, this is offset by its rapid metabolism (Kenaga



and Goring 1980; Marshall and Roberts 1978). With the fathead minnow, Jarvinen et al. (1983) found a mean bioconcentration factor (BCF) of 1,673 after 60 days (Table 5). In a review, Kenaga and Goring (1980) cite results of an unpublished study reporting a BCF in an unnamed fish of 450. In an experimental outdoor stream, Eaton et al. (1985) reported average BCFs for a fathead minnow and a bluegill of 590 and 100, respectively, in exposures ranging from 18 to 33 days.

Data on the uptake of chlorpyrifos are available for five species of saltwater fish (Table 5). In two early life-stage tests with the gulf toadfish, Opsanus beta, the BCF increased from 100 to 5,100 as the concentration of chlorpyrifos in the test solution increased from 1.4 to 150 µg/L (Hansen et al. 1986). Cripe et al. (Manuscript) found that the BCF with the sheepshead minnow depended on the availability of food as well as the concentration of chlorpyrifos in water (Tables 5 and 6).

No U.S. FDA action level or other maximum acceptable concentration in tissue is available for chlorpyrifos, and, therefore, no Final Residue Value can be calculated.

#### Other Data

Additional data on the lethal and sublethal effects of chlorpyrifos on aquatic organisms are given in Table 6. Because both chlorpyrifos and the mosquitofish are used in mosquito control programs, several studies have been conducted on the effects of chlorpyrifos on the survival and effectiveness of mosquitofish as a predator of mosquito larvae. Hansen et al. (1972) reported a 24-hr LC50 of 4,000 µg/L for this fish. A 36-hr LC50 of 215 to 230 µg/L was reported by Ferguson et al. (1966), whereas a 72-hr LC50 of 0.19 to 0.22 µg/L was reported by Ahmed (1977). After a

24-hr exposure to 5.0 µg/L, Johnson (1977a,1978a) observed a decreased thermal tolerance in mosquitofish. Hansen et al. (1972) found that mosquitofish chose clean water when given a choice between clean water and 100 µg chlorpyrifos/L in laboratory experiments.

For rainbow trout, the 96-hr LC50s at 1.6, 7.2, and 12.7°C were 51, 15, and 7.1 µg/L, respectively (Macek et al. 1969). Increased toxicity of chlorpyrifos with increased temperature was thought to be the result of either increased metabolism producing lower concentrations of dissolved oxygen and higher metabolic wastes, or increased enzyme activity converting chlorpyrifos to its more toxic form, chlorpyrifos-oxon. In a 24-hr exposure to 100 µg/L, Atlantic salmon had a 4°C lower temperature preference (Peterson 1976).

Eaton et al. (1985) conducted studies on chlorpyrifos in outdoor experimental streams containing native invertebrates, stocked fathead minnows (Pimephales promelas) and bluegills (Lepomis macrochirus), and wild white suckers (Catostomus commersoni). Two dosing strategies were used in separate streams. One was a continuous exposure; the other was a pulsed exposure. Concentrations were adjusted so that equal total amounts of chlorpyrifos were applied to each stream. Continuous exposure for 100 days to an average chlorpyrifos concentration of 0.35 µg/L (range 0.12 to 0.83 µg/L) produced no significant effect on the fishes. The invertebrate community was greatly effected, with a decrease in species diversity and a strong shift from an amphipod- to an isopod-dominated benthic community. Chironomids were greatly reduced. BCFs for the fathead minnow and bluegill were 590 and 100, respectively. Continuous and pulsed exposures gave similar results for most biological measurements.

Because of the previous use of chlorpyrifos as a mosquito larvicide, many toxicity studies have used various species of mosquito larvae as test organisms. Unfortunately, many studies have followed guidelines set forth by the World Health Organization on testing of pesticides. These guidelines prescribe a 24-hr test duration, making results unusable for derivation of water quality criteria.

As would be expected, chlorpyrifos is highly toxic to mosquitos. Rettich (1977) reported 24-hr LC50s of 0.5 to 3.5  $\mu\text{g/L}$  for 4th instars of 6 species of the genus Aedes. For A. aegypti, a species not tested by Rettich, Saleh et al. (1981) cited 24-hr LC50s of 0.0011 and 0.0014  $\mu\text{g/L}$  for 2nd and 4th instars, respectively. Reports of 24-hr LC50s for 4th instars of various Culex species range from 0.41 to 2.0  $\mu\text{g/L}$  (Ahmed 1977; Helson et al. 1979; Rettich 1977). For C. pipiens, Saleh et al. (1981) found a 24-hr LC50 of 0.0052  $\mu\text{g/L}$ .

Chlorpyrifos was also used to control noxious midge populations (Ali and Mulla 1978a,1980; Mulla and Khasawinah 1969; Mulla et al. 1971; Thompson et al. 1970). The 24-hr LC50s for various midges range from 0.5 to 40  $\mu\text{g/L}$  (Ali and Mulla 1978a,1980; Mulla and Khasawinah 1969) although a value of 1,470  $\mu\text{g/L}$  was reported for Cricotopus decorus (Ali and Mulla 1980).

Ahmed (1977) determined 24-hr LC50s with 6 species of aquatic coleoptera and observed a range of 4.6 to 52  $\mu\text{g/L}$ . Levy and Miller (1978) observed the delayed effects of 24-hr exposures to 1.0 and 4.0  $\mu\text{g/L}$  on a planarian, Dugesia dorotocephala, over 108 hr. They reported no significant effects at either concentration.

Winner et al. (1978) used a single concentration of an emulsifiable concentrate in a study of the effects on a mermithid nematode parasite of mosquito larvae. They examined toxicity to infectious, parasitic, post-parasitic, and embryo stages of the nematode. Rawn et al. (1978) investigated the effect of various sediments on the toxicity of chlorpyrifos to larvae of a mosquito in artificial ponds. They found lower toxicity and lower concentrations in water in sod-lined ponds compared to sand-lined ponds at equal application rates. Macek et al. (1972) conducted a field study that included analyses of fish brain AChE activity, fish stomach contents, residues in fish and water, numbers of larval insects, and numbers of emerging insects. Siefert et al. (1984) conducted an extensive survey of changes within a natural pond after chlorpyrifos was applied using methods employed by pest control authorities. Their study included analysis of water quality, fish and invertebrate populations, and associated laboratory studies.

Schaefer and Dupras (1970) examined the effect of polluted waters on the stability of chlorpyrifos in the field. Zepp and Schlotzhauer (1983) studied the effect of algae on photolysis of chlorpyrifos. As part of a laboratory study, El-Refai et al. (1976) tested the effectiveness of a simulated water treatment facility in lowering toxicity of Nile River water spiked with chlorpyrifos. They found a 33% decrease in toxicity with alum treatment, and no significant change with sand filtration.

Jamback and Frempong-Boadu (1966), Mohsen and Mulla (1981), and Muirhead-Thomson (1978,1979) observed delayed effects after short exposures.

Among saltwater species, juvenile brown shrimp, Penaeus aztecus, were the most sensitive with a 48-hr EC50 of 0.32 µg/L (U.S. Bureau of Commercial Fisheries 1965; Lowe et al. 1970). Other 48-hr EC50s are 1.5

$\mu\text{g/L}$  for juvenile grass shrimp, Palaemonetes pugio, 2.4  $\mu\text{g/L}$  for the pink shrimp, Penaeus duorarum, and 5.2  $\mu\text{g/L}$  for the blue crab, Callinectes sapidus. For the eastern oyster the 96-hr EC50s based on shell deposition range from 270 to 340  $\mu\text{g/L}$ .

The 48-hr LC50s for fish ranged from 3.2  $\mu\text{g/L}$  for the longnose killifish, Fundulus similis, to over 1,000  $\mu\text{g/L}$  for the sheepshead minnow, Cyprinodon variegatus. The most sensitive effect on a fish species was a reduction in growth of the California grunion, Leuresthes tenuis, exposed to 0.62  $\mu\text{g/L}$  for 26 days (Goodman et al. 1985a). Acetylcholinesterase activity in the brain of adult mummichogs, Fundulus heteroclitus, was inhibited by 2.1  $\mu\text{g/L}$  (Thirugnanam and Forgash 1977).

The effect of chlorpyrifos on benthic communities was investigated by Tagatz et al. (1982). In communities exposed in the laboratory for 8 weeks during colonization, total faunal species richness and the abundance of arthropods and molluscs were reduced by concentrations from 0.1 to 8.5  $\mu\text{g/L}$ . For communities previously colonized in the field, the number of arthropods was reduced by 5.9  $\mu\text{g/L}$ , but not by 1.0  $\mu\text{g/L}$ .

#### Unused Data

Some data on the effects of chlorpyrifos on aquatic organisms were not used because the studies were conducted with species that are not resident in North America (e.g., Moorthy et al. 1982). Results of tests reported by Ali (1981), Ferguson et al. (1966), Naqvi (1973), and Nelson and Evans (1973) were not used because the test organisms probably had been previously exposed to pesticides or other pollutants. Chiou et al. (1977), Dean and Ballantyne (1985), Kenaga (1980), Marshall and Roberts (1978),

Ramke (1969), Yoshioka et al. (1986), and Zarogian et al. (1985) only contain data that have been published elsewhere.

Data were not used if the test was on a commercial formulation (e.g., Atallah and Ishak 1971; Birmingham and Colman 1977; Chang and Lange 1967; Hurlbert et al. 1970; Ledieu 1978; Muirhead-Thomson 1970; Mulla et al. 1973; Rettich 1979; Roberts and Miller 1971; Scirocchi and D'Erme 1980; Siefert et al. 1984; Smith et al. 1966) or if the source of the chlorpyrifos was not adequately described (e.g., Ali and Mulla 1976,1977; Boike and Rathburn 1969; Gillies et al. 1974; Johnson 1977b,1978b; Kenaga et al. 1965; Micks and Rougeau 1977; Muirhead-Thomson and Merryweather 1969; Ruber and Kocor 1976; Thayer and Ruber 1976; Wilder and Schaefer 1969; Zboray and Gutierrez 1979). Data were not used if the organisms were exposed to chlorpyrifos by injection or gavage or in food (e.g., Herin et al. 1978; Wilton et al. 1973), if chlorpyrifos was a component of a mixture (Meyer 1981), or if the organisms were fed during exposure in short term tests (Karnak and Collins 1974).

The concentration of solvent was too high in the tests of Al-Khatib (1985) and Davey et al. (1976). Barton (1970) conducted a static chronic test with mosquito larvae. Because polyethylene sorbs chlorpyrifos (Brown et al. 1976; Hughes 1977; Hughes et al. 1980), toxicity tests conducted in polyethylene test chambers were not used if the concentration of chlorpyrifos was not measured (e.g., Brown and Chow 1975; Darwazen and Mulla 1974; Dixon and Brust 1971; Hughes 1977; Miller et al. 1973; Roberts et al. 1973a,b). Results of some laboratory tests were not used because the tests were conducted in distilled or deionized water without addition of appropriate salts (e.g., Jones et al. 1976; Nelson and Evans 1973; Rongsriyam et al. 1968; Steelman et al. 1969). High control mortalities

occurred in tests reported by Khudairi and Ruber (1974). Test procedures were inadequately described by Mellon and Georghiou (1985) and Ruber and Baskar (1969).

BCFs obtained from microcosm or model ecosystem studies were not used if the concentration of chlorpyrifos in water decreased with time or if the exposure was too short (e.g., Metcalf 1974). Data from field studies and measurements of chlorpyrifos in wild organisms were not used if the concentrations of chlorpyrifos in water were not measured (e.g., Ali and Mulla 1976, 1977, 1978a, b; Axtell et al. 1979; Best 1969; Campbell and Denno 1976; Carter and Graves 1972; Chang and Lange 1967; Chatterji et al. 1979; Cooney and Pickard 1974; Evans et al. 1975; Fitzpatrick and Sutherland 1978; Frank and Sjogren 1978; Hazeleur 1971; Holbrook and Agun 1984; Hoy et al. 1972; Jamnback 1969; Lembright 1968; Linn 1968; Marganian and Wall 1972; McNeill et al. 1968; Moore and Breeland 1967; Mulla and Khasawinah 1969; Mulla et al. 1971; Nelson et al. 1976a, b; Polls et al. 1975; Roberts et al. 1984; Steelman et al. 1969; Stewart 1977; Tawfik and Gooding 1970; Taylor and Schoof 1971; Thompson et al. 1970; Wallace et al. 1973; Washino et al. 1968, 1972a, b; Wilkinson et al. 1971; Winterlin et al. 1968; Yap and Ho 1977) or if the concentration in water was not uniform enough (e.g., Macek et al. 1972).

### Summary

The acute values for eighteen freshwater species in fifteen genera range from 0.11  $\mu\text{g/L}$  for an amphipod to greater than 806  $\mu\text{g/L}$  for two fishes and a snail. The bluegill is the most acutely sensitive fish species with an acute value of 10  $\mu\text{g/L}$ , but seven invertebrate genera are more sensitive. Smaller organisms seem to be more acutely sensitive than larger ones.

Chronic toxicity data are available for one freshwater species, the fathead minnow. Unacceptable effects occurred in second generation larvae at 0.12 µg/L, which was the lowest concentration tested. The resulting acute-chronic ratio was greater than 1,417.

Little information is available on the toxicity of chlorpyrifos to freshwater plants, although algal blooms frequently follow field applications of chlorpyrifos. The only available bioconcentration test on chlorpyrifos with a freshwater species was with the fathead minnow and resulted in a bioconcentration factor of 1,673.

The acute toxicity of chlorpyrifos has been determined with 15 species of saltwater animals in 12 genera, and the acute values ranged from 0.01 µg/L for the Korean shrimp, Palaemon macrodactylus, to 1,911 µg/L for larvae of the eastern oyster, Crassostrea virginica. Arthropods are particularly sensitive to chlorpyrifos. Among the 10 species of fish tested, the 96-hr LC50s range from 0.58 µg/L for striped bass to 520 µg/L for gulf toadfish; larvae are more sensitive than other life stages. Growth of the mysid, Mysidopsis bahia, was reduced at 0.004 µg/L in a life-cycle test. In early life-stage tests, the California grunion, Leuresthes tenuis, was the most sensitive of the six fishes, with growth being reduced at 0.30 µg/L. Of the seven acute-chronic ratios that have been determined with saltwater species, the five lowest range from 2.388 to 12.50, whereas the highest is 228.5.

Concentrations of chlorpyrifos affecting six species of saltwater phytoplankton range from 138 to 10,000 µg/L. BCFs ranged from 100 to 5,100 when the gulf toadfish was exposed to concentrations increasing from 1.4 to 150 µg/L. Steady-state BCFs averaged from 100 to 757 for five fishes exposed in early life-stage tests.



## National Criteria

The procedures described in the "Guidelines for Deriving Numerical National Water Quality Criteria for the Protection of Aquatic Organisms and Their Uses" indicate that, except possibly where a locally important species is very sensitive, freshwater aquatic organisms and their uses should not be affected unacceptably if the four-day average concentration of chlorpyrifos does not exceed 0.041  $\mu\text{g/L}$  more than once every three years on the average and if the one-hour average concentration does not exceed 0.083  $\mu\text{g/L}$  more than once every three years on the average.

The procedures described in the "Guidelines for Deriving Numerical National Water Quality Criteria for the Protection of Aquatic Organisms and Their Uses" indicate that, except possibly where a locally important species is very sensitive, saltwater aquatic organisms and their uses should not be affected unacceptably if the four-day average concentration of chlorpyrifos does not exceed 0.0056  $\mu\text{g/L}$  more than once every three years on the average and if the one-hour average concentration does not exceed 0.011  $\mu\text{g/L}$  more than once every three years on the average.

Three years is the Agency's best scientific judgment of the average amount of time aquatic ecosystems should be provided between excursions (U.S. EPA 1985b). The resiliences of ecosystems and their abilities to recover differ greatly, however, and site-specific allowed excursion frequencies may be established if adequate justification is provided.

Use of criteria for developing water quality-based permit limits and for designing waste treatment facilities requires selection of an appropriate wasteload allocation model. Dynamic models are preferred for the application of these criteria (U.S. EPA 1985b). Limited data or other considerations might make their use impractical, in which case one must rely on a steady-state model (U.S. EPA 1986).

Table 1. Acute Toxicity of Chlorpyrifos to Aquatic Animals

<u>Species</u>	<u>Method</u> <sup>#</sup>	<u>Chemical</u> <sup>**</sup>	<u>LC50 or EC50 (µg/L)</u> <sup>***</sup>	<u>Species Mean Acute Value (µg/L)</u>	<u>Reference</u>
<u>FRESHWATER SPECIES</u>					
<u>Snail (adult), Aplexa hypnorum</u>	F, M	Technical (98.7%)	>806	>806	Phipps and Holcombe 1985a,b
<u>Amphipod, Gammarus fasciatus</u>	S, U	Technical	0.32	0.32	Sanders 1972
<u>Amphipod (2 mo. old), Gammarus lacustris</u>	S, U	Technical (97%)	0.11	0.11	Sanders 1969; Johnson and Finley 1980
<u>Amphipod, Gammarus pseudolimnaeus</u>	F, M	Encapsu- lated****	0.18	0.18	Siefert et al. 1984
<u>Crayfish (1.8 g), Orconectes immunis</u>	F, M	Technical (98.7%)	6	6	Phipps and Holcombe 1985a,b
<u>Stonefly (naïve), Pteronarcella badla</u>	S, U	Technical (97%)	0.38	0.38	Sanders and Cope 1968
<u>Stonefly (naïve), Pteronarcys californica</u>	S, U	Technical (97%)	10	10	Sanders and Cope 1968; Johnson and Finley 1980
<u>Stonefly (naïve), Claassenia sabulosa</u>	S, U	Technical (97%)	0.57	0.57	Sanders and Cope 1968; Johnson and Finley 1980
<u>Trichopteran, Leptoceridae sp.</u>	S, M	Encapsu- lated****	0.77	0.77	Siefert et al. 1984
<u>Pygmy backswimmer, Neoplea striola</u>	S, M	Encapsu- lated****	1.22	-	Siefert et al. 1984
<u>Pygmy backswimmer, Neoplea striola</u>	S, M	Encapsu- lated****	1.56	1.38	Siefert et al. 1984
<u>Crawling water beetle (adult), Peltodytes sp.</u>	S, U		0.8	0.8	Federle and Collins 1976
<u>Cutthroat trout (1.4 g), Salmo clarkii</u>	S, U	Technical (97%)	18	18	Johnson and Finley 1980

Table 1. (continued)

<u>Species</u>	<u>Method<sup>a</sup></u>	<u>Chemical<sup>b</sup></u>	<u>LC50 or EC50 (<math>\mu\text{g/L}</math>)<sup>c</sup></u>	<u>Species Mean Acute Value (<math>\mu\text{g/L}</math>)</u>	<u>Reference</u>
<u>Rainbow trout (0.6-1.5 g), Salmo gairdneri</u>	S, U	Technical	7.1	-	Macek et al. 1969; Johnson and Finley 1980
<u>Rainbow trout (juvenile), Salmo gairdneri</u>	F, M	Technical (99.9%)	8.0	-	Holcombe et al. 1982
<u>Rainbow trout (3.0 g), Salmo gairdneri</u>	F, M	Technical (98.7%)	9	8.485	Phipps and Holcombe 1985a,b
<u>Lake trout (2.3 g), Salvelinus namaycush</u>	S, U	Technical (97%)	98	98	Johnson and Finley 1980
<u>Goldfish (10.7 g), Carassius auratus</u>	F, M	Technical (98.7%)	>806	>806	Phipps and Holcombe 1985a,b
<u>Fathead minnow, Pimephales promelas</u>	S, M	Technical (98.7%)	170	-	Jarvinen and Tanner 1982
<u>Fathead minnow (juvenile), Pimephales promelas</u>	F, M	Technical (99.9%)	203	-	Holcombe et al. 1982
<u>Fathead minnow (0.5 g), Pimephales promelas</u>	F, M	Technical (98.7%)	542	331.7	Phipps and Holcombe 1985a,b
<u>Channel catfish (0.8 g), Ictalurus punctatus</u>	S, U	Technical (97%)	280	-	Johnson and Finley 1980
<u>Channel catfish (7.9 g), Ictalurus punctatus</u>	F, M	Technical (98.7%)	806	806	Phipps and Holcombe 1985a,b
<u>Bluegill (0.6 g), Lepomis macrochirus</u>	S, U	Technical (97%)	2.4	-	Johnson and Finley 1980
<u>Bluegill (0.8 g), Lepomis macrochirus</u>	F, M	Technical (98.7%)	10	10	Phipps and Holcombe 1985a,b

Table 1. (Continued)

<u>Species</u>	<u>Method*</u>	<u>Chemical**</u>	<u>Salinity (g/kg)</u>	<u>LC50 or EC50 (µg/L)***</u>	<u>Species Mean Acute Value (µg/L)</u>	<u>Reference</u>
<u>SALTWATER SPECIES</u>						
<u>Eastern oyster (larva), Crassostrea virginica</u>	S, U	Technical (92%)	20	1,991	1,991	Borthwick and Walsh 1981
<u>Mysid (juvenile), Mysidopsis bahia</u>	S, U	Technical (92%)	20	0.056	-	Borthwick and Walsh 1981
<u>Mysid (juvenile), Mysidopsis bahia</u>	F, M	Technical (92%)	26.7	0.035	0.035	Schimmel et al. 1983
<u>Amphipod, Ampelisca abdita</u>	R, U	Technical (92%)	32	0.16	0.16	Scott and Redmond 1986a
<u>Amphipod, Rhepoxynius abronius</u>	R, U	Technical (92%)	32	0.07	-	Scott and Redmond 1986b
<u>Amphipod, Rhepoxynius abronius</u>	R, U	Technical (92%)	32	0.14	0.0990	Scott and Redmond 1986b
<u>Korean shrimp (adult), Palaemon macrodactylus</u>	S, U	(99%)	15	0.25	-	Earnest 1970
<u>Korean shrimp (adult), Palaemon macrodactylus</u>	F, U	(99%)	24	0.01	0.05	Earnest 1970
<u>Gulf toadfish (juvenile), Opsanus beta</u>	R, M	Technical (92%)	29-30	520	520	Hansen et al. 1986
<u>Sheepshead minnow (juvenile), Cyprinodon variegatus</u>	S, U	Technical (92%)	20	270	-	Borthwick and Walsh 1981
<u>Sheepshead minnow (juvenile), Cyprinodon variegatus</u>	F, M	Technical (92%)	10.3	136	136	Schimmel et al. 1983
<u>Mummichog (adult), Fundulus heteroclitus</u>	S, U	Technical (99.5%)	20-25	4.65	4.65	Thirugnanam and Forqash 1977

Table 1. (Continued)

<u>Species</u>	<u>Method<sup>#</sup></u>	<u>Chemical<sup>**</sup></u>	<u>Salinity (g/kg)</u>	<u>LC50 or EC50 (<math>\mu</math>g/L)<sup>***</sup></u>	<u>Species Mean Acute Value (<math>\mu</math>g/L)</u>	<u>Reference</u>
<u>Longnose killifish (juvenile), Fundulus similis</u>	F, M	Technical (92%)	25.9	4.1	4.1	Schimmel et al. 1983
<u>California grunion (day 0 larva), Leuresthes tenuis</u>	S, U	Technical (92%)	25	5.5	-	Borthwick et al. 1985
<u>California grunion (day 7 larva), Leuresthes tenuis</u>	S, U	Technical (92%)	25	2.7	-	Borthwick et al. 1985
<u>California grunion (day 14 larva), Leuresthes tenuis</u>	S, U	Technical (92%)	25	1.8	-	Borthwick et al. 1985
<u>California grunion (day 28 larva), Leuresthes tenuis</u>	S, U	Technical (92%)	25	2.6	-	Borthwick et al. 1985
<u>California grunion (day 0 larva), Leuresthes tenuis</u>	F, M	Technical (92%)	25	1.0	-	Borthwick et al. 1985
<u>California grunion (day 7 larva), Leuresthes tenuis</u>	F, M	Technical (92%)	25	1.0	-	Borthwick et al. 1985
<u>California grunion (day 14 larva), Leuresthes tenuis</u>	F, M	Technical (92%)	25	1.0	-	Borthwick et al. 1985
<u>California grunion (day 28 larva), Leuresthes tenuis</u>	F, M	Technical (92%)	25	1.3	1.068	Borthwick et al. 1985
<u>Inland silverside (juvenile), Menidia beryllina</u>	F, M	Technical	5.0	4.2	4.2	Clark et al. 1985
<u>Atlantic silverside (day 0 larva), Menidia menidia</u>	S, U	Technical (92%)	20	4.5	-	Borthwick et al. 1985

Table 1. (Continued)

<u>Species</u>	<u>Method*</u>	<u>Chemical**</u>	<u>Salinity (g/kg)</u>	<u>LC50 or EC50 (µg/L)***</u>	<u>Species Mean Acute Value (µg/L)</u>	<u>Reference</u>
<u>Atlantic silverside (day 7 larva), Menidia menidia</u>	S, U	Technical (92%)	20	2.8	-	Borthwick et al. 1985
<u>Atlantic silverside (day 14 larva), Menidia menidia</u>	S, U	Technical (92%)	20	2.4	-	Borthwick et al. 1985
<u>Atlantic silverside (day 28 larva), Menidia menidia</u>	S, U	Technical (92%)	20	4.1	-	Borthwick et al. 1985
<u>Atlantic silverside (juvenile) Menidia menidia</u>	F, M	Technical (92%)	24.3	1.7	-	Schimmel et al. 1983
<u>Atlantic silverside (day 0 larva), Menidia menidia</u>	F, M	Technical (92%)	20	0.5	-	Borthwick et al. 1985
<u>Atlantic silverside (day 7 larva), Menidia menidia</u>	F, M	Technical (92%)	20	1.0	-	Borthwick et al. 1985
<u>Atlantic silverside (day 14 larva), Menidia menidia</u>	F, M	Technical (92%)	20	1.1	-	Borthwick et al. 1985
<u>Atlantic silverside (day 28 larva), Menidia menidia</u>	F, M	Technical (92%)	20	3.0	1.229	Borthwick et al. 1985
<u>Tidewater silverside (day 0 larva), Menidia peninsulae</u>	S, U	Technical (92%)	20	4.2	-	Borthwick et al. 1985
<u>Tidewater silverside (day 7 larva), Menidia peninsulae</u>	S, U	Technical (92%)	20	2.0	-	Borthwick et al. 1985
<u>Tidewater silverside (day 14 larva), Menidia peninsulae</u>	S, U	Technical (92%)	20	1.8	-	Borthwick et al. 1985

Table 1. (Continued)

<u>Species</u>	<u>Method*</u>	<u>Chemical**</u>	<u>Salinity (g/kg)</u>	<u>LC50 or EC50 (µg/L)***</u>	<u>Species Mean Acute Value (µg/L)</u>	<u>Reference</u>
<u>Tidewater silverside (day 28 larva), Menidia peninsulae</u>	S, U	Technical (92%)	20	3.9	-	Borthwick et al. 1985
<u>Tidewater silverside (day 0 larva), Menidia peninsulae</u>	F, M	Technical (92%)	20	1.0	-	Borthwick et al. 1985
<u>Tidewater silverside (day 7 larva), Menidia peninsulae</u>	F, M	Technical (92%)	20	0.5	-	Borthwick et al. 1985
<u>Tidewater silverside (day 14 larva), Menidia peninsulae</u>	F, M	Technical (92%)	20	0.4	-	Borthwick et al. 1985
<u>Tidewater silverside (day 28 larva), Menidia peninsulae</u>	F, M	Technical (92%)	20	0.9	-	Borthwick et al. 1985
<u>Tidewater silverside (juvenile), Menidia peninsulae</u>	F, M	Technical	19.3	1.3	0.7479	Clark et al. 1985
<u>Striped bass (juvenile), Morone saxatilis</u>	F, U	(99%)	30	0.58	0.58	Earnest 1970; Korn and Earnest 1974
<u>Striped mullet (juvenile), Mull cephalus</u>	F, M	Technical (92%)	24.7	5.4	5.4	Schimmel et al. 1983

\* S = static; R = renewal; F = flow-through; U = unmeasured; M = measured.

\*\* Percent purity is given in parentheses when available.

\*\*\* If the concentrations were not measured and the published results were not reported to be adjusted for purity, the published results were multiplied by the purity if it was reported to be less than 97%.

\*\*\*\* The test material was dissolved from an encapsulated form of chlorpyrifos.

Table 2. Chronic Toxicity of Chlorpyrifos to Aquatic Animals

<u>Species</u>	<u>Test*</u>	<u>Chemical**</u>	<u>Salinity (g/kg)</u>	<u>Limits (µg/L)***</u>	<u>Chronic Value (µg/L)</u>	<u>Reference</u>
<u>FRESHWATER SPECIES</u>						
<u>Fathead minnow, Pimephales promelas</u>	ELS	Technical (98.7%)	-	1.6-3.2	2.263	Jarvinen and Tanner 1982
<u>Fathead minnow, Pimephales promelas</u>	ELS	Encapsulated <sup>†</sup>	-	2.2-4.8	3.250	Jarvinen and Tanner 1982
<u>Fathead minnow, Pimephales promelas</u>	LC	Encapsulated <sup>†</sup>	-	<0.12 <sup>††</sup>	<0.12	Jarvinen et al. 1983
<u>SALTWATER SPECIES</u>						
<u>Mysid, Mysidopsis bahia</u>	LC	Technical (92%)	19-28	0.002-0.004	0.0028	McKenney et al. 1981
<u>Gulf toadfish, Opsanus beta</u>	ELS	Technical (92%)	25-34	1.4-3.7	2.276	Hansen et al. 1986
<u>Sheepshead minnow, Cyprinodon variegatus</u>	ELS	Technical (92%)	9-28	1.7-3.0	2.258	Cripe et al. 1986
<u>Sheepshead minnow, Cyprinodon variegatus</u>	ELS	Technical (92%)	9-28	1.7-3.0	2.258	Cripe et al. 1986
<u>Sheepshead minnow, Cyprinodon variegatus</u>	ELS	Technical (92%)	9-28	1.7-3.0	2.258	Cripe et al. 1986
<u>California grunion, Leuresthes tenuis</u>	ELS	Technical (92%)	24.5-34.0	0.14-0.30	0.2049	Goodman et al. 1985a
<u>Inland silverside, Menidia beryllina</u>	ELS	Technical (92%)	4-6	0.75-1.8	1.162	Goodman et al. 1985b
<u>Atlantic silverside, Menidia menidia</u>	ELS	Technical (92%)	18-27	0.28-0.48	0.3666	Goodman et al. 1985b



Table 2. (Continued)

<u>Species</u>	<u>Test*</u>	<u>Chemical**</u>	<u>Salinity (g/kg)</u>	<u>Limits (µg/L)***</u>	<u>Chronic Value (µg/L)</u>	<u>Reference</u>
Tidewater silverside, <u>Menidia peninsulae</u>	ELS	Technical (92%)	18-25	0.38-0.78	0.5444	Goodman et al. 1985b

\* LC = life-cycle or partial life-cycle; ELS = early life-stage.

\*\* Percent purity is given in parentheses when available.

\*\*\* Results are based on measured concentrations of chlorpyrifos.

† The test material was dissolved from an encapsulated form of chlorpyrifos.

†† Unacceptable effects occurred at all tested concentrations.

Table 2. (Continued)

<u>Species</u>	<u>Acute-Chronic Ratio</u>		<u>Ratio</u>
	<u>Acute Value</u> ( $\mu\text{g/L}$ )	<u>Chronic Value</u> ( $\mu\text{g/L}$ )	
<u>Fathead minnow,</u> <u>Pimephales promelas</u>	170	<0.12*	>1,417
<u>Mysid,</u> <u>Mysidopsis bahia</u>	0.035	0.0028	12.50
<u>Gulf toadfish,</u> <u>Opsanus beta</u>	520	2.276	228.5
<u>Sheepshead minnow,</u> <u>Cyprinodon variegatus</u>	136	2.258	60.23
<u>Sheepshead minnow,</u> <u>Cyprinodon variegatus</u>	136	2.258	60.23
<u>Sheepshead minnow,</u> <u>Cyprinodon variegatus</u>	136	2.258	60.23
<u>California grunion,</u> <u>Leuresthes tenuis</u>	1.068	0.2049	5.212
<u>Inland silverside,</u> <u>Menidia beryllina</u>	4.2	1.162	3.614
<u>Atlantic silverside,</u> <u>Menidia menidia</u>	1.229	0.3666	3.352
<u>Tidewater silverside,</u> <u>Menidia peninsulae</u>	0.7479	0.5444	1.374

\* Results of the early life-stage tests are not used because results of a life-cycle test are available.

Table 3. Ranked Genus Mean Acute Values with Species Mean Acute-Chronic Ratios

<u>Rank*</u>	<u>Genus Mean Acute Value (µg/L)</u>	<u>Species</u>	<u>Species Mean Acute Value (µg/L)**</u>	<u>Species Mean Acute-Chronic Ratio***</u>
<u>FRESHWATER SPECIES</u>				
15	>806	Snail, <u>Aplexa hypnorum</u>	>806	-
14	>806	Goldfish, <u>Carassius auratus</u>	>806	-
13	806	Channel catfish, <u>Ictalurus punctatus</u>	806	-
12	331.7	Fathead minnow, <u>Pimephales promelas</u>	331.7	>1,417
11	98	Lake trout, <u>Salvelinus namaycush</u>	98	-
10	12.36	Cutthroat trout, <u>Salmo clarki</u>	18	-
		Rainbow trout, <u>Salmo gairdneri</u>	8,485	-
9	10	Stonefly, <u>Pteronarcys californica</u>	10	-
8	10	Bluegill, <u>Lepomis macrochirus</u>	10	-
7	6	Crayfish, <u>Orconectes immunis</u>	6	-
6	1.38	Pygmy backswimmer, <u>Neoplea striola</u>	1.38	-
5	0.8	Crawling water beetle, <u>Peltodytes sp.</u>	0.8	-
4	0.77	Trichoptera <u>Leptoceridae sp.</u>	0.77	-

Table 3. (continued)

<u>Rank<sup>a</sup></u>	<u>Genus Mean Acute Value (µg/L)</u>	<u>Species</u>	<u>Species Mean Acute Value (µg/L)<sup>aa</sup></u>	<u>Species Mean Acute-Chronic Ratio<sup>aaa</sup></u>
3	0.57	Stonefly, <u>Clasenia sabulosa</u>	0.57	-
2	0.38	Stonefly, <u>Pteronarcella badia</u>	0.38	-
1	0.1850	Amphipod, <u>Gammarus fasciatus</u>	0.32	-
		Amphipod, <u>Gammarus lacustris</u>	0.11	-
		Amphipod, <u>Gammarus pseudolimnaeus</u>	0.18	-
<u>SALTWATER SPECIES</u>				
12	1,991	Eastern oyster, <u>Crassostrea virginica</u>	1,991	-
11	520	Gulf toadfish, <u>Opsanus beta</u>	520	228.5
10	136	Sheepshead minnow, <u>Cyprinodon variegatus</u>	136	60.23
9	5.4	Striped mullet, <u>Mugil cephalus</u>	5.4	-
8	4.366	Mummichog, <u>Fundulus heteroclitus</u>	4.65	-
		Longnose killifish, <u>Fundulus similis</u>	4.1	-
7	1.569	Inland silverside, <u>Menidia beryllina</u>	4.2	3.614
		Atlantic silverside, <u>Menidia menidia</u>	1.229	3.352
		Tidewater silverside, <u>Menidia peninsulae</u>	0.7479	1.374

Table 3. (continued)

<u>Rank*</u>	<u>Genus Mean Acute Value (µg/L)</u>	<u>Species</u>	<u>Species Mean Acute Value (µg/L)**</u>	<u>Species Mean Acute-Chronic Ratio***</u>
6	1.068	California grunion, <u>Leuresthes tenuis</u>	1.068	5.212
5	0.58	Striped bass, <u>Morone saxatilis</u>	0.58	-
4	0.16	Amphipod, <u>Ampelisca abdita</u>	0.16	-
3	0.0990	Amphipod, <u>Rhepoxynius abronius</u>	0.0990	-
2	0.05	Korean shrimp, <u>Palaemon macrodactylus</u>	0.05	-
1	0.035	Mysid, <u>Mysidopsis bahia</u>	0.035	12.50

\* Ranked from most resistant to most sensitive based on Genus Mean Acute Value. Inclusion of "greater than" values does not necessarily imply a true ranking, but does allow use of all genera for which data are available so that the Final Acute Value is not unnecessarily lowered.

\*\* From Table 1.

\*\*\* From Table 2.

Fresh water

Final Acute Value = 0.1669 µg/L

Criterion Maximum Concentration = (0.1669 µg/L) / 2 = 0.08345 µg/L

Final Acute-Chronic Ratio = 4.064 (see text)

Final Chronic Value = (0.1669 µg/L) / 4.064 = 0.04107 µg/L

Salt water

Final Acute Value = 0.02284 µg/L

Criterion Maximum Concentration = (0.02284 µg/L) / 2 = 0.01142 µg/L

Final Acute-Chronic Ratio = 4.064 (see text)

Final Chronic Value = (0.02284 µg/L) / 4.064 = 0.005620 µg/L

Table 4. Toxicity of Chlorpyrifos to Aquatic Plants

<u>Species</u>	<u>Chemical*</u>	<u>Duration (days)</u>	<u>Salinity (g/kg)</u>	<u>Effect</u>	<u>Concentration (µg/L)**</u>	<u>Reference</u>
<u>SALTWATER SPECIES</u>						
<u>Golden-brown alga, <i>Isochrysis galbana</i></u>	Technical (92%)	4	30	EC50 (population growth)	138	Borthwick and Walsh 1981
<u>Diatom, <i>Skeletonema costatum</i></u>	Technical (92%)	4	30	EC50 (population growth)	297.8***	Borthwick and Walsh 1981
<u>Diatom, <i>Thalassiosira pseudonana</i></u>	Technical (92%)	4	30	EC50 (population growth)	148	Borthwick and Walsh 1981

\* Percent purity is given in parentheses when available.

\*\* If the concentrations were not measured and the published results were not reported to be adjusted for purity, the published results were multiplied by the purity if it was reported to be less than 97%.

\*\*\* Geometric mean of five values.

Table 5. Bioconcentration of Chlorpyrifos by Aquatic Organisms

<u>Species</u>	<u>Chemical*</u>	<u>Concentration in Water (µg/L)**</u>	<u>Duration (days)</u>	<u>Tissue</u>	<u>BCF or BAF***</u>	<u>Reference</u>
<u>FRESHWATER SPECIES</u>						
<u>Fathead minnow, Pimephales promelas</u>	Encapsulated†	0.12-2.68	60	Whole body	1,673	Jarvinen et al. 1983
<u>Fathead minnow, Pimephales promelas</u>	-	0.14	21	Whole body	570	Eaton et al. 1985
<u>Fathead minnow, Pimephales promelas</u>	-	0.15	18	Whole body	260	Eaton et al. 1985
<u>Fathead minnow, Pimephales promelas</u>	-	0.32	21	whole body	780	Eaton et al. 1985
<u>Fathead minnow, Pimephales promelas</u>	-	0.46	33	Whole body	760	Eaton et al. 1985
<u>Bluegill, Lepomis macrochirus</u>	-	0.41	33	whole body	100	Eaton et al. 1985
<u>SALTWATER SPECIES</u>						
<u>Gulf toadfish, Opsanus beta</u>	Technical (92%)	1.4	49	Whole body	100	Hansen et al. 1986
<u>Sheepshead minnow, Cyprinodon variegatus</u>	Technical (92%)	0.41 0.78	28	Whole body (n=6)	118††	Cripe et al. 1986
<u>California grunion, Leuresthes tenuis</u>	Technical (92%)	0.14-0.63	35	Whole body (n=3)	757†† (from ELS test)	Goodman et al. 1985a
<u>California grunion, Leuresthes tenuis</u>	Technical (92%)	0.28-1.3	26	Whole body (n=3)	153†† (from fry test)	Goodman et al. 1985a
<u>Inland silverside, Menidia beryllina</u>	Technical (92%)	0.18-1.8	28	Whole body (n=4)	186††	Goodman et al. 1985b

Table 5. (continued)

<u>Species</u>	<u>Chemical*</u>	<u>Concentration in Water (µg/L)**</u>	<u>Duration (days)</u>	<u>Tissue</u>	<u>BCF or BAF***</u>	<u>Reference</u>
<u>Tidewater silverside, Menidia peninsulae</u>	Technical (92%)	0.093-0.38	28	Whole body (n=3)	456 <sup>††</sup>	Goodman et al. 1985b

\* Percent purity is given in parentheses when available.

\*\* Measured concentration of chlorpyrifos.

\*\*\* Bioconcentration factors (BCFs) and bioaccumulation factors (BAFs) are based on measured concentrations of chlorpyrifos in water and tissue.

† The test material was dissolved from an encapsulated form of chlorpyrifos.

†† Geometric mean of values from the listed concentrations in water.



Table 6. Other Data on Effects of Chlorpyrifos on Aquatic Organisms

<u>Species</u>	<u>Chemical<sup>a</sup></u>	<u>Duration</u>	<u>Effect</u>	<u>Concentration (<math>\mu\text{g/L}</math>)<sup>b,c</sup></u>	<u>Reference</u>
<u>FRESHWATER SPECIES</u>					
Diatoms, Unidentified	-	10 days	Reduced growth	400	Roberts and Miller 1970
Planarian, <u>Dugesia dorotocephala</u>	-	24 hr	None	4.0	Levy and Miller 1978
Cladoceran, <u>Daphnia</u> sp.	Encapsulated <sup>***</sup>	4 hr	LC50	0.88	Siefert et al. 1984
Amphipod, <u>Hyalella azteca</u>	Encapsulated <sup>***</sup>	24 hr	LC50	1.28	Siefert et al. 1984
Mayfly, <u>Ephemereilla</u> sp.	Encapsulated <sup>***</sup>	72 hr	LC50	0.33	Siefert et al. 1984
Pygmy backswimmer, <u>Neopisra striola</u>	Encapsulated <sup>***</sup>	144 hr	LC50	0.97	Siefert et al. 1984
Giant water bug (adult), <u>Belostomatidae</u> sp.	Technical	24 hr	LC50	15	Ahmed 1977
Predaceous diving beetle (adult), <u>Hygrotus</u> sp.	Technical	24 hr	LC50	40	Ahmed 1977
Predaceous diving beetle (adult), <u>Laccophilus decipiens</u>	Technical	24 hr	LC50	4.6	Ahmed 1977
Predaceous diving beetle (adult), <u>Thermonectus basillaris</u>	Technical	24 hr	LC50	6	Ahmed 1977
Water scavenger beetle (adult), <u>Berosus styliferus</u>	Technical	24 hr	LC50	9	Ahmed 1977
Water scavenger beetle (larva), <u>Hydrophilus triangularis</u>	Technical	24 hr	LC50	20	Ahmed 1977

Table 6. (continued)

<u>Species</u>	<u>Chemical*</u>	<u>Duration</u>	<u>Effect</u>	<u>Concentration (µg/L)**</u>	<u>Reference</u>
Water scavenger beetle (adult), <u>Hydrophilus triangularis</u>	Technical	24 hr	LC50	30	Ahmed 1977
Water scavenger beetle (larva), <u>Tropisternus lateralis</u>	Technical	24 hr	LC50	52	Ahmed 1977
Water scavenger beetle (adult), <u>Tropisternus lateralis</u>	Technical	24 hr	LC50	8	Ahmed 1977
Mosquito (3rd and 4th instar), <u>Aedes aegypti</u>	Technical	18 hr	LT50	10	Verma and Rahman 1984
Mosquito (2nd instar), <u>Aedes aegypti</u>	Technical (96%)	24 hr	LC50	0.0011	Saleh et al. 1981
Mosquito (4th instar), <u>Aedes aegypti</u>	Technical (96%)	24 hr	LC50	0.0014	Saleh et al. 1981
Mosquito (4th instar), <u>Aedes cantans</u>	Technical	24 hr	LC50	1.1	Rettich 1977
Mosquito (4th instar), <u>Aedes communis</u>	Technical	24 hr	LC50	3.5	Rettich 1977
Mosquito (4th instar), <u>Aedes excrucians</u>	Technical	24 hr	LC50	3.3	Rettich 1977
Mosquito (4th instar), <u>Aedes punctor</u>	Technical	24 hr	LC50	2.7	Rettich 1977
Mosquito (4th instar), <u>Aedes sticticus</u>	Technical	24 hr	LC50	0.5	Rettich 1977
Mosquito (4th instar), <u>Aedes vexans</u>	Technical	24 hr	LC50	1.0	Rettich 1977
Mosquito (larva), <u>Anopheles freeborni</u>	Technical	24 hr	LC50	3	Ahmed 1977

Table 6. (continued)

<u>Species</u>	<u>Chemical*</u>	<u>Duration</u>	<u>Effect</u>	<u>Concentration (<math>\mu\text{g/L}</math>)**</u>	<u>Reference</u>
<u>Mosquito (4th instar), Anopheles freeborni</u>	Technical	24 hr	LC50	1.3 0.9 2.5 1.2 1.2 1.9 1.3 1.2 2.8 3.3 7.0	Wameldorf et al. 1970
<u>Mosquito (3rd and 4th instar), Anopheles stephensi</u>	Technical	6.5 hr	LT50	25	Verma and Rahman 1984
<u>Phantom midge, Chaoborus sp.</u>	Encapsulated***	18 hr 42 hr 114 hr	LC50 LC50 LC50	2.36 1.29 0.85	Siefert et al. 1984
<u>Mosquito (4th instar), Culex pipiens</u>	Technical (99%)	24 hr	EC50	0.46	Helson et al. 1979
<u>Mosquito (2nd instar; DDT susceptible), Culex pipiens</u>	Technical (96%)	24 hr	LC50	0.0022	Saleh et al. 1981
<u>Mosquito (4th instar; DDT susceptible), Culex pipiens</u>	Technical (96%)	24 hr	LC50	0.0052	Saleh et al. 1981
<u>Mosquito (4th instar), Culex pipiens</u>	Technical	24 hr	LC50	1.2	Rettich 1977
<u>Mosquito (4th instar), Culex pipiens</u>	Technical	24 hr	LC50	1.6	Rettich 1977
<u>Mosquito (3rd and 4th instar), Culex quinquefasciatus</u>	Technical	5 hr	LT50	10	Verma and Rahman 1984

Table 6. (continued)

<u>Species</u>	<u>Chemical<sup>a</sup></u>	<u>Duration</u>	<u>Effect</u>	<u>Concentration (<math>\mu\text{g/L}</math>)<sup>b,c</sup></u>	<u>Reference</u>
Mosquito (4th Instar), <u>Culex restuans</u>	Technical (99%)	24 hr	LC50	0.41	Helson et al. 1979
Mosquito (larva), <u>Culex tarsalis</u>	Technical	-	LC50	2	Ahmed 1977
Mosquito (4th Instar), <u>Culiseta annulata</u>	Technical	24 hr	LC50	3.5	Rettich 1977
Midge (4th Instar), <u>Chironomus</u> sp.	Technical	24 hr	LC50	0.42	Mulla and Khasawinah 1969
Midge (4th Instar), <u>Chironomus decorus</u>	Technical	24 hr	LC50	7.0	All and Mulla 1978a
Midge (4th Instar), <u>Chironomus utahensis</u>	Technical	24 hr	LC50	1.2	All and Mulla 1978a
Midge (4th Instar), <u>Cricotopus decorus</u>	Technical	24 hr	LC50	1,470	All and Mulla 1980
Midge (4th Instar), <u>Dicretendipes californicus</u>	Technical	24 hr	LC50	40.0	All and Mulla 1980
Midge (4th Instar), <u>Goeldichironomus holoprassinus</u>	Technical	24 hr	LC50	0.97	Mulla and Khasawinah 1969
Midge (4th Instar), <u>Procladius</u> spp.	Technical	24 hr	LC50	0.5	All and Mulla 1978a
Midge (4th Instar), <u>Tanypus grodhausi</u>	Technical	24 hr	LC50	29.0	All and Mulla 1980
Midge (4th Instar), <u>Tanypus grodhausi</u>	Technical	24 hr	LC50	0.5	Mulla and Khasawinah 1969
Midge (4th Instar), <u>Tanygus grodhausi</u>	Technical	24 hr	LC50	5.7	Mulla and Khasawinah 1969
Rainbow trout, <u>Salmo gairdneri</u>	Technical	96 hr	LC50 (7.2°C) (1.6°C)	15 51	Macek et al. 1969

Table 6. (continued)

<u>Species</u>	<u>Chemical<sup>a</sup></u>	<u>Duration</u>	<u>Effect</u>	<u>Concentration (<math>\mu\text{g/L}</math>)<sup>b,c</sup></u>	<u>Reference</u>
<u>Atlantic salmon (juvenile), Salmo salar</u>	Technical (>95%)	24 hr	Temperature selection	100	Peterson 1976
<u>Golden shiner, Notemigonus crysoleucas</u>	Technical (99%)	36 hr	LC50	45	Ferguson et al. 1966
<u>Golden shiner, Notemigonus crysoleucas</u>	Technical (99%)	36 hr	LC50	35	Ferguson et al. 1966
<u>Fathead minnow, Pimephales promelas</u>	Technical****	96 hr	LC50	150	Jarvinen and Tanner 1982
<u>Fathead minnow, Pimephales promelas</u>	Encapsulated***	96 hr	LC50	130	Jarvinen and Tanner 1982
<u>Fathead minnow, Pimephales promelas</u>	Encapsu- lated***, ****	96 hr	LC50	280	Jarvinen and Tanner 1982
<u>Fathead minnow (larva), Pimephales promelas</u>	-	7 day	Reduced growth	5.2	Norberg and Mount 1985
<u>Mosquitofish, Gambusia affinis</u>	-	-	Avoidance	100	Hansen et al. 1972
<u>Mosquitofish (DDT susceptible), Gambusia affinis</u>	Technical	48 hr	LC50	1,018	Culley and Ferguson 1969
<u>Mosquitofish (DDT resistant), Gambusia affinis</u>	Technical	48 hr	LC50	1,291	Culley and Ferguson 1969
<u>Mosquitofish, Gambusia affinis</u>	-	24 hr	Decreased thermal tolerance	5	Johnson 1978a
<u>Mosquitofish (adult), Gambusia affinis</u>	Technical	72 hr	LC50	0.22 0.20 0.20 0.19 0.20 0.20	Ahmed 1977
<u>Mosquitofish, Gambusia affinis</u>	Technical (99%)	36 hr	LC50	230 215	Ferguson et al. 1966

Table 6. (continued)

<u>Species</u>	<u>Chemical*</u>		<u>Duration</u>	<u>Effect</u>	<u>Concentration (<math>\mu\text{g/L}</math>)**</u>	<u>Reference</u>
<u>Mosquitofish, Gambusia affinis</u>	-		24 hr	LC50	4,000	Hansen et al. 1972
<u>Guppy, Poecilia reticulata</u>	Technical		24 hr	LC50	220	Rongsrityam et al. 1968
<u>Green sunfish, Lepomis cyanellus</u>	Technical (99%)		36 hr	LC50	37.5 22.5	Ferguson et al. 1966
<u>Species</u>	<u>Chemical*</u>	<u>Salinity (g/kg)</u>	<u>Duration</u>	<u>Effect</u>	<u>Concentration (<math>\mu\text{g/L}</math>)**</u>	<u>Reference</u>
<u>SALTWATER SPECIES</u>						
<u>Green alga, Chlorococcum sp.</u>	-	27	48 hr	Reduced growth	10,000	Maly and Ruber 1983
<u>Diatom, Skeletonema costatum</u>	Technical (92%)	30	48 hr	100% mortality	5,000	Walsh 1981, 1983
<u>Diatom, Amphiprora sp.</u>	-	27	48 hr	Reduced growth	10,000	Maly and Ruber 1983
<u>Dinoflagellate, Gonyaulax sp.</u>	-	27	48 hr	Reduced growth	10,000	Maly and Ruber 1983
<u>Benthic macrofauna</u>	Technical (92%)	26.5 (14.5-34.0)	8 wk	Significant reduction in total faunal species richness and in abundance of arthropods and molluscs for laboratory-colonized benthos	0.1	Tagatz et al. 1982
<u>Benthic macrofauna</u>	Technical (92%)	27.5 (18.0-32.5)	1 wk	Significant reduction in abundance of arthropods for field-colonized benthos	5.9	Tagatz et al. 1982

Table 6. (continued)

<u>Species</u>	<u>Chemical<sup>a</sup></u>	<u>Salinity (g/kg)</u>	<u>Duration</u>	<u>Effect</u>	<u>Concentration (<math>\mu</math>g/L)<sup>b,c</sup></u>	<u>Reference</u>
Eastern oyster (juvenile), <u>Crassostrea virginica</u>	(99%)	24	96 hr	EC50 (shell deposition)	270	U.S. Bureau of Commercial Fisheries 1965; Lowe et al. 1970
Eastern oyster (juvenile), <u>Crassostrea virginica</u>	(99%)	28	96 hr	EC50 (shell deposition)	34	U.S. Bureau of Commercial Fisheries 1967
Amphipod, <u>Ampelisca abdita</u>	(92%)	32	96 hr	EC50 (with sediment)	0.34	Scott and Redmond 1986a
Brown shrimp (juvenile), <u>Penaeus aztecus</u>	(99%)	26	48 hr	EC50 (mortality and loss of equilibrium)	0.20	U.S. Bureau of Commercial Fisheries 1965; Lowe et al. 1970
Pink shrimp (juvenile), <u>Penaeus duorarum</u>	(99%)	26	48 hr	EC50	2.4	U.S. Bureau of Commercial Fisheries 1967
Grass shrimp (juvenile), <u>Palaemonetes pugio</u>	(99%)	26	48 hr	EC50 (mortality and loss of equilibrium)	1.5	U.S. Bureau of Commercial Fisheries 1967)
Grass shrimp (adult), <u>Palaemonetes pugio</u>	(99%)	20	1 hr	No avoidance of pesticide	0.01-1.0	Hansen et al. 1973
Blue crab (juvenile), <u>Callinectes sapidus</u>	(99%)	20	48 hr	EC50	5.2	U.S. Bureau of Commercial Fisheries 1967
Atlantic salmon (juvenile), <u>Salmo salar</u>	(>95%)	-	24 hr	Altered preferred temperature	100-250	Peterson 1976
Sheepshead minnow (juvenile), <u>Cyprinodon variegatus</u>	Technical (92%)	9-30	28 days	BCF = 42-660 (low food); 69-1,000 (med. food); 120-1,830 (high food)	0.41-52	Cripe et al. 1986
Sheepshead minnow (adult), <u>Cyprinodon variegatus</u>	(99%)	20	1 hr	Avoidance of pesticide	100-250	Hansen 1969, 1970

Table 6. (continued)

<u>Species</u>	<u>Chemical*</u>	<u>Salinity (g/kg)</u>	<u>Duration</u>	<u>Effect</u>	<u>Concentration (µg/L)**</u>	<u>Reference</u>
Sheepshead minnow (juvenile), <u>Cyprinodon variegatus</u>	(99%)	24	48 hr	LC50	>1,000	U.S. Bureau of Commercial Fisheries 1967
Gulf toadfish, <u>Opsanus beta</u>	(92%)	24-34	49 days	BCF = 100- 5,100	1.4-150	Hansen et al. 1986
Mummichog (adult), <u>Fundulus heteroclitus</u>	Technical (99.5%)	20-25	24 hr	100% inhibition of acetylcholin- esterase activity in brain	>2.1	Thirugnanam and Forgash 1977
Longnose killifish (juvenile), <u>Fundulus similis</u>	(99%)	24	48 hr	LC50	3.2	U.S. Bureau of Commercial Fisheries 1967; Lowe et al. 1970
California grunion, <u>Leuresthes tenuis</u>	Technical (92%)	24.5- 31.5	26 days	Significantly reduced growth of fry	0.62	Goodman et al. 1985a
Spot (juvenile), <u>Leiostomus xanthurus</u>	(99%)	26	48 hr	LC50	7	U.S. Bureau of Commercial Fisheries 1965

\* Percent purity is given in parentheses when available.

\*\* If the concentrations were not measured and the published results were not reported to be adjusted for purity, the published results were multiplied by the purity if it was reported to be less than 97%.

\*\*\* The test material was dissolved from an encapsulated form of chlorpyrifos.

\*\*\*\* Aged 11 weeks.



#### REFERENCES

- Ahmed, W. 1977. The effectiveness of predators of rice field mosquitoes in relation to pesticide use in rice culture. Ph.D. thesis. University of California-Davis, Davis, CA. Available from University Microfilms, Ann Arbor, MI. Order No. 77-6323.
- Ali, A. 1981. Laboratory evaluation of organophosphate and new synthetic pyrethroid insecticides against pestiferous chironomid midges of central Florida. *Mosq. News* 41:157-161.
- Ali, A. and M.S. Mulla. 1976. Insecticidal control of chironomid midges in the Santa Ana River water system, Orange County, California. *J. Econ. Entomol.* 69:509-513.
- Ali, A. and M.S. Mulla. 1977. The IGR diflubenzuron and organophosphorus insecticides against nuisance midges in man-made residential-recreational lakes. *J. Econ. Entomol.* 70:571-577.
- Ali, A. and M.S. Mulla. 1978a. Declining field efficacy of chlorpyrifos against chironomid midges and laboratory evaluation of substitute larvicides. *J. Econ. Entomol.* 71:778-782.
- Ali, A. and M.S. Mulla. 1978b. Effects of chironomid larvicides and diflubenzuron on nontarget invertebrates in residential-recreational lakes. *Environ. Entomol.* 7:21-27.
- Ali, A. and M.S. Mulla. 1980. Activity of organophosphate and synthetic pyrethroid insecticides against pestiferous midges in some southern California flood control channels. *Mosq. News* 40:593-597.

Al-Khatib, Z.I. 1985. Isolation of an organophosphate susceptible strain of Culex quinquefasciatus from a resistant field population by discrimination against esterase-2-phenotypes. J. Am. Mosq. Control Assoc. 1:105-107.

Atallah, Y.H. and M.M. Ishak. 1971. Toxicity of some commonly used insecticides to the snail Biomphalaria alexandrina, intermediate host of Schistoma monsoni in Egypt. Z. Angew. Entomol. 69:102-106.

Axtell, R.C., J.C. Dukes and T.D. Edwards. 1979. Field tests of diflubenzuron, methoprene, Flit MLO and chlorpyrifos for the control of Aedes taeniorhynchus larvae in diked dredged spoil areas. Mosq. News 39: 520-527.

Barton, L.C. 1970. The effect of sublethal concentrations of Dursban on immature Culex pipiens quinquefasciatus. AD712316. National Technical Information Service, Springfield, VA.

Best, D.W. 1969. Dursban effective for mosquito control in creek bottoms and duck ponds. Proc. Pap. Annu. Conf. Calif. Mosq. Control Assoc. 37:133-134.

Birmingham, B.C. and B. Colman. 1977. The effect of two organophosphate insecticides on the growth of freshwater algae. Can. J. Bot. 55:1453-1456.

Boike, A.H. and C.B. Rathburn. 1969. Laboratory tests of the susceptibility of mosquito larvae to insecticides in Florida, 1968. Mosq. News 29:392-395.

Borthwick, P.W. and G.E. Walsh. 1981. Initial toxicological assessment of Ambush, Bolero, Bux, Dursban, Fentrifanil, Larvin, and Pydrin: Static acute

- toxicity tests with selected estuarine algae, invertebrates and fish.  
EPA 600/4-81-076. National Technical Information Service, Springfield, VA.
- Borthwick, P.W., J.M. Patrick, Jr., and D.P. Middaugh. 1985. Comparative acute sensitivities of early life stages of atherinid fishes to chlorpyrifos and thiobencarb. Arch. Environ. Contam. Toxicol. 14:465-473.
- Braun, H.E. and R. Frank. 1980. Organochlorine and organophosphorus insecticides: Their use in eleven agricultural watersheds and their loss to stream waters in southern Ontario, Canada, 1975-1977. Sci. Total. Environ. 15:169-192.
- Brown, J.R. and L.Y. Chow. 1975. The effect of Dursban on micro-flora in non-saline waters. In: Environmental quality and safety supplement. Vol. III. Pesticides. Coulston, P. Fredrick and F. Korte (Eds.). International Union of Pure and Applied Chemistry, Helsinki, Finland pp. 774-779.
- Brown, J.R., L.Y. Chow and C.B. Deng. 1976. The effect of Dursban upon fresh water phytoplankton. Bull. Environ. Contam. Toxicol. 15:437-444.
- Butcher, J., M. Boyer and C.D. Fowle. 1975. Impact of Dursban and Abate on microbial numbers and some chemical properties of standing ponds. Water Pollut. Res. Can. 10:33-41.
- Butcher, J.E., M.G. Boyer and C.D. Fowle. 1977. Some changes in pond chemistry and photosynthetic activity following treatment with increasing concentrations of chlorpyrifos. Bull. Environ. Contam. Toxicol. 17:752-758.
- Campbell, B.C. and R.F. Denno. 1976. The effect of temephos and chlorpyrifos on the aquatic insect community of a New Jersey salt marsh. Environ. Entomol. 5:477-483.

- Carter, F.L. and J.B. Graves. 1972. Measuring effects of insecticides on aquatic animals. *La. Agric.* 16:14-15.
- Chang, V.C. and W.H. Lange. 1967. Laboratory and field evaluations of selecting pesticides for control of the red crayfish in California rice fields. *J. Econ. Entomol.* 60:473-477.
- Chatterji, S.M., J.P. Kulshroshta and S. Rajamani. 1979. Some promising insecticides for the control of the rice gall midge, Orseolia oryzae. *J. Entomol. Res.* 3:168-171.
- Chiou, C.T., V.H. Freed, D.W. Schmedding and R.L. Kohnert. 1977. Partition coefficient and bioaccumulation of selected organic chemicals. *Environ. Sci. Technol.* 11:475-478.
- Clark, J.R., D. DeVault, R.J. Bowden and J.A. Weishaar. 1984. Contaminant analysis of fillets from Great Lakes coho salmon. 1980. *J. Great Lakes Res.* 10:38-47.
- Clark, J.R., J.M. Patrick, Jr., D.P. Middaugh and J.C. Moore. 1985. Relative sensitivity of six estuarine fishes to carbophenothion, chlorpyrifos, and fenvalerate. *Ecotoxicol. Environ. Safety* 10:382-390.
- Cooney, J.C. and E. Pickard. 1974. Field tests with Abate and Dursban insecticides for control of floodwater mosquitoes in the Tennessee Valley region. *Mosq. News* 34:12-22.
- Cripe, G.M., D.J. Hansen, S.F. Macauley and J. Forester. 1986. Effects of diet quantity on sheepshead minnows (Cyprinodon variegatus)

during early life-stage exposures to chlorpyrifos. In: Aquatic toxicology and environmental fate: Ninth volume. ASTM STP 921. Poston, T.M. and R. Purdy (Eds.). American Society for Testing and Materials, Philadelphia, PA.

Culley, D.D. and D.E. Ferguson. 1969. Patterns of insecticide resistance in the mosquitofish, Gambusia affinis. J. Fish. Res. Board Can. 26:2395-2401.

Darwazeh, H.A. and M.S. Mulla. 1974. Toxicity of herbicides and mosquito larvicides to the mosquitofish Gambusia affinis. Mosq. News 34:214-219.

Davey, R.B., M.V. Meisch and F.L. Carter. 1976. Toxicity of five rice field pesticides to the mosquitofish, Gambusia affinis, and green sunfish, Lepomis cyanellus, under laboratory and field conditions in Arkansas. Environ. Entomol. 5:1053-1056.

Dean, J.H. and M.A. Ballantyne. 1985. Human health and aquatic life literature search and data base evaluation for chlorpyrifos. Draft Report. Battelle Columbus Laboratories, Columbus, OH.

Dixon, R.D. and R.A. Brust. 1971. Field testing of insecticides used in mosquito control, and a description of the bioassay technique used in temporary pools. J. Econ. Entomol. 64:11-14.

Earnest, R. 1970. Effects of pesticides on aquatic animals in the estuarine and marine environment. In: Progress in sport fisheries research 1970. Resource Publication 106. U.S. Bureau of Sport Fisheries and Wildlife, Washington, DC. pp. 10-13.

Eaton, J., J. Arthur, R. Hermanutz, R. Kiefer, L. Mueller, R. Anderson, R. Erickson, B. Nordling, J. Rogers and H. Pritchard. 1985. Biological effects of continuous and intermittent dosing of outdoor experimental streams

with chlorpyrifos. In: Aquatic toxicology and hazard assessment. ASTM STP 891. Bahner, R.C. and D.J. Hansen (Eds.). American Society for Testing and Materials, Philadelphia, PA. pp. 85-118.

El-Refai, A., F.A. Fahmy, M.F. Abdel-Lateef and A-K Imam. 1976. Toxicity of three insecticides to two species of fish. Int. Pest Control 18:4-8.

Evans, E.S. 1977. Field evaluation of the extended mosquito larvicidal activity of a controlled-release chlorpyrifos polymer in a woodland pool habitat, March 1974-October 1976. Entomological Special Study No. 44-0364-77. U.S. Army Environmental Hygiene Agency, Aberdeen Proving Ground, MD.

Evans, E.S., J.H. Nelson, N.E. Pennington and W.W. Young. 1975. Larvicidal effectiveness of a controlled-release formulation of chlorpyrifos in a woodland pool habitat. Mosq. News 35:343-350.

Federle, P.F. and W.J. Collins. 1976. Insecticide toxicity to three insects from Ohio ponds. Ohio J. Sci. 76:19-24.

Ferguson, D.E., D.T. Gardner and A.L. Lindley. 1966. Toxicity of Dursban to three species of fish. Mosq. News 26:80-82.

Fitzpatrick, G. and D.J. Sutherland. 1978. Effects of the organophosphorus insecticides temephos (Abate) and chlorpyrifos (Dursban) on populations of the salt-marsh snail, Melampus bidentatus. Mar. Biol. (Berl.) 46:23-28.

Frank, A.M. and R.D. Sjogren. 1978. Effects of temephos and chlorpyrifos on crustacea. Mosq. News 38:138-139

Gillies, P.A., D.J. Womeldorf, E.P. Zboray, and K.E. White. 1974. Insecticide susceptibility of mosquitoes in California: Status of organophosphorus

resistance in larval Aedes nigromaculis and Culex tarsalis through 1973. Proc. Pap. Annu. Conf. Calif. Mosq. Control Assoc. 42:107-112.

Goodman, L.R., D.J. Hansen, G.M. Cripe, D.P. Middaugh and J.C. Moore. 1985a. A new early life-stage toxicity test using the California grunion (Leuresthes tenuis) and results with chlorpyrifos. Ecotoxicol. Environ. Safety 10:12-21.

Goodman, L.R., D.J. Hansen, D.P. Middaugh, G.M. Cripe and J.C. Moore. 1985b. Method for early life-stage toxicity tests using three atherinid fishes and results with chlorpyrifos. In: Aquatic toxicology and hazard assessment: 7th symposium. Cardwell, R.D., R. Purdy and R.C. Bahner (Eds.). ASTM STP 854. American Society for Testing and Materials, Philadelphia, PA. pp. 145-154.

Gray, H.E. 1965. Dursban, a new organophosphorus insecticide. Down Earth 21:26-27.

Hansen, D.J. 1969. Avoidance of pesticide by untrained sheepshead minnows. Trans. Am. Fish. Soc. 98:426-429.

Hansen, D.J. 1970. Behavior of estuarine organisms. Progress report. Contribution No. 98:23-28. Bureau of Commercial Fisheries Center for Estuarine and Menhaden Research, Pesticide Field Station, Gulf Breeze, FL.

Hansen, D.J., E. Matthews, S.L. Nali and D.P. Dumas. 1972. Avoidance of pesticides by untrained mosquitofish, Gambusia affinis. Bull. Environ. Contam. Toxicol. 8:46-51.

- Hansen, D.J., S.C. Schimmel and J.M. Keltner, Jr. 1973. Avoidance of pesticides by grass shrimp (Palaemonetes pugio). Bull. Environ. Contam. Toxicol. 9:129-133.
- Hansen, D.J., L.R. Goodman, G.M. Cripe and S.F. Macauley. 1986. Early life-stage toxicity test methods for gulf toadfish (Opsanus beta) and results using chlorpyrifos. Ecotoxicol. Environ. Safety 11:15-22.
- Hazeleur, W.C. 1971. Use of Dursban for mosquito control in log ponds in the Shasta Mosquito Abatement District. Proc. Pap. Annu. Conf. Calif. Mosq. Control Assoc. 39:47.
- Helson, B.V., G.A. Surgeoner and W.E. Ralley. 1979. Susceptibility of Culex spp. and Aedes spp. larvae (Diptera: Culicidae) to temephos and chlorpyrifos in southern Ontario. Proc. Entomol. Soc. Ont. 110: 79-83.
- Herin, R.A., J.E. Suggs, E.M. Lores, L.T. Heiderscheit, J.D. Farmer and D. Prather. 1978. Correlation of salt gland function with levels of chlorpyrifos in the feed of mallard ducklings. Pestic. Biochem. Physiol. 9:157-163.
- Holbrook, F.R. and S.K. Agun. 1984. Field trials of pesticides to control larval Culicoides variipennis (Ceratopogonidae). Mosq. News 44:233-235.
- Holcombe, G.W., G.L. Phipps and D.K. Tanner. 1982. The acute toxicity of kelthane, Dursban, disulfoton, pydrin, and permethrin to fathead minnows Pimephales promelas and rainbow trout Salmo gairdneri. Environ. Pollut. (Ser. A.) 29:167-178.



Hoy, J.B., E.E. Kauffman, and A.G. O'Berg. 1972. A large-scale field test of Gambusia affinis and chlorpyrifos for mosquito control. Mosq. News 32:161-171.

Hughes, D.N. 1977. The effects of three organophosphorus insecticides on zooplankton and other invertebrates in natural and artificial ponds. M.S. dissertation. York University, Toronto, Canada.

Hughes, D.N., M.G. Boyer, M.H. Papst, C.D. Fowle, G.A. Rees and P. Baulu. 1980. Persistence of three organophosphorus insecticides in artificial ponds and some biological implications. Arch. Environ. Contam. Toxicol. 9:269-279.

Hurlbert, S.H. 1969. The impact of Dursban on pond ecosystems. Proc. Pap. Annu. Conf. Calif. Mosq. Control Assoc. 37:8.

Hurlbert, S.H., M.S. Mulla, J.O. Keith, W.E. Westlake and M.E. Dusch. 1970. Biological effects and persistence of Dursban in freshwater ponds. J. Econ. Entomol. 63:43-52.

Hurlbert, S.H., M.S. Mulla and H.R. Wilson. 1972. Effects of an organophosphorus insecticide on the phytoplankton, zooplankton, and insect populations of freshwater ponds. Ecol. Monogr. 42:269-299.

Jamback, H. 1969. Field tests with larvicides other than DDT for control of blackfly (Diptera: Simuliidae) in New York. Bull. W.H.O. 40:635-638.

Jamback, H. and J. Frempong-Boadu. 1966. Testing blackfly larvicides in the laboratory and in streams. Bull. W.H.O. 34:405-421.

Jarvinen, A.W. and D.K. Tanner. 1982. Toxicity of selected controlled release and corresponding unformulated technical grade pesticides to the fathead minnow Pimephales promelas. Environ. Pollut. (Series A) 27:179-195.

Jarvinen, A.W., B.R. Nordling and M.E. Henry. 1983. Chronic toxicity of Dursban (chlorpyrifos) to the fathead minnow (Pimephales promelas) and the resultant acetylcholinesterase inhibition. Ecotoxicol. Environ. Safety 7:423-434.

Johnson, C.R. 1977a. The effects of field applied rates of five organophosphorus insecticides on thermal tolerance, orientation, and survival in Gambusia affinis affinis. Proc. Pap. Annu. Conf. Calif. Mosq. Vector Control Assoc. 45:56-58.

Johnson, C.R. 1977b. The effect of exposure to the organophosphorus insecticide chlorpyrifos on the feeding rate in the mosquitofish, Gambusia affinis. Proc. Pap. Annu. Conf. Calif. Mosq. Vector Control Assoc. 45:69-70.

Johnson, C.R. 1978a. The effects of sublethal concentrations of five organophosphorus insecticides on temperature tolerance, reflexes, and orientation in Gambusia affinis affinis. Zool. J. Linn. Soc. 64:63-70.

Johnson, C.R. 1978b. The effect of five organophosphorus insecticides on survival and temperature tolerance in the copepod, Macrocyclus albidus. Zool. J. Linn. Soc. 64:59-62.

Johnson, W.W. and M.T. Finley. 1980. Handbook of acute toxicity of chemicals to fish and aquatic invertebrates. Resource Publication 137. U.S. Fish and Wildlife Service, Washington, DC. p. 21.

- Jones, G.E., D.F. Carroll and W. Wills. 1976. Susceptibility of Pennsylvania mosquito larvae to Abate, Dursban and Baytex. Proc. 63rd. Annu. Meet. N.J. Mosq. Exterm. Assoc. pp. 161-164.
- Karnak, R.E. and W.J. Collins. 1974. The susceptibility of selected insecticides and acetylcholinesterase activity in a laboratory colony of midge larvae, Chironomus tentans (Diptera: Chironomidae). Bull. Environ. Contam. Toxicol. 12:62-69.
- Kenaga, E.E. 1980. Correlation of bioconcentration factors of chemicals in aquatic and terrestrial organisms with their physical and chemical properties. Environ. Sci. Technol. 14:553-556.
- Kenaga, E.E. and C.A. Goring. 1980. Relationship between water solubility, soil sorption, octanol-water partitioning, and concentration of chemicals in biota. In: Aquatic toxicology. Eaton, J.G., P.R. Parrish and A.C. Hendricks (Eds.). ASTM STP 707. American Society for Testing and Materials, Philadelphia, PA. pp. 78-115.
- Kenaga, E.E., W.K. Whitney, J.L. Hardy and A.E. Doty. 1965. Laboratory tests with Dursban insecticide. J. Econ. Entomol. 58:1043-1050.
- Khudairi, S.Y. and E. Ruber. 1974. Survival and reproduction of ostracods as affected by pesticides and temperature. J. Econ. Entomol. 67:22-24.
- Korn, S. and R. Earnest. 1974. Acute toxicity of 20 insecticides to a striped bass (Morone saxatilis). Calif. Fish Game 60:128-131.
- Ledieu, M.S. 1978. Candidate insecticides for the control of larvae of Mamestra brassicae (Lepidoptera) (Noctuidae). Ann. Appl. Biol. 88:251-255.

- Lembright, H.W. 1968. Dosage studies with low volume applications of Dursban insecticide. *Down Earth* 24:16-19.
- Levy, R. and T.W. Miller, Jr. 1978. Tolerance of the planarian Dugesia dorotocephala to high concentrations of pesticides and growth hormones. *Entomophaga* 23:31-34.
- Linn, J.D. 1968. Effects of low volume aerial spraying of Dursban and fenthion on fish. *Down Earth* 24:28-30.
- Lowe, J.T., P.D. Wilson and R.B. Davison. 1970. Laboratory bioassays. In: Progress report for fiscal year 1969. Circular 335. U.S. Fish and Wildlife Service, Pesticide Field Station, Gulf Breeze, FL.
- Macalady, D.L. and N.L. Wolfe. 1985. Effects of sediment sorption and abiotic hydrolyses. 1. Organophosphorothioate esters. *J. Agric. Food Chem.* 33:167-173.
- Macek, K.J., C. Hutchinson and O.B. Cope. 1969. The effects of temperature on the susceptibility of bluegills and rainbow trout to selected pesticides. *Bull. Environ. Contam. Toxicol.* 4:174-183.
- Macek, K.J., D.F. Walsh, J.W. Hogan and D.D. Holz. 1972. Toxicity of the insecticide Dursban to fish and aquatic invertebrates in ponds. *Trans. Am. Fish. Soc.* 101:420-427.
- Maly, M. and E. Ruber. 1983. Effects of pesticides on pure and mixed species cultures of salt marsh pool algae. *Bull. Environ. Contam. Toxicol.* 30:464-472.

Marganian, V.M. and W.J. Wall, Jr. 1972. Dursban and diazinon residues in biota following treatment of intertidal plots on Cape Cod - 1967-69. Pestic. Monit. J. 6:160-165.

Marshall, W.K. and J.R. Roberts. 1978. Ecotoxicology of chlorpyrifos. NRCC 16079. National Research Council of Canada, Ottawa, Ontario, Canada.

McKenney, C., Jr., E. Matthews and D. Lawrence. 1981. Mysid life-cycle tests. Progress Report, FY81. Experimental Environments Branch, U.S. EPA, Gulf Breeze, FL.

McNeill, J.C., W.O. Miller and C.M. Wleczyk. 1968. Evaluation of Dursban as a larvicide in septic ditches. Mosq. News. 28:160-161.

Mellon, R.B. and G.P. Georghiou. 1985. Rotational use of insecticides in mosquito control programs. Proc. Pap. Annu. Conf. Calif. Mosq. Vector Control Assoc. 52:65-67.

Menzie, C.M. 1969. Metabolism of pesticides. Special Scientific Report - Wildlife No. 127. Bureau of Sport Fisheries and Wildlife, Washington, DC.

Metcalf, R.L. 1974. A laboratory model ecosystem to evaluate compounds producing biological magnification. In: Essays in toxicology-V. Hayes, W.J. (Ed.). Academic Press, New York, NY. pp. 17-38.

Meyer, F.P. 1981. Influences of contaminants on toxicity of lampricides. Quarterly report of progress, April-June, 1981. U.S. Fish and Wildlife Service, National Fisheries Research Laboratory, LaCrosse, WI and S.E. Fish Control Laboratory, Warm Springs, GA.

Micks, D.W. and D. Rougeau. 1977. Organophosphorus tolerance in Culex quinquefasciatus in Texas. Mosq. News 37:233-239.

Miller, T.A., L.L. Nelson, W.W. Young, L.W. Roberts, D.R. Roberts and R.N. Wilkinson. 1973. Polymer formulations of mosquito larvicides. I. Effectiveness of polyethylene and polyvinyl chloride formulations of chlorpyrifos applied to artificial field pools. Mosq. News. 33:148-155.

Mohsen, Z.H. and M.S. Mulla. 1981. Toxicity of blackfly larvicidal formulations to some aquatic insects in the laboratory. Bull. Environ. Contam. Toxicol. 26:696-703.

Moore, J.B. and S.G. Breeland. 1967. Field evaluation of two mosquito larvicides, Abate and Dursban, against Anopheles quadrimaculatus and associated Culex species. Mosq. News 27:105-111.

Moorthy, M.V., S. Chandrasekhar and V.R. Chandran. 1982. A note on acute toxicity of chlorpyrifos to the freshwater fish Thilapia mossambica. Pesticides 16:32.

Muirhead-Thomson, R.C. 1970. The potentiating effect of pyrethrins and pyrethroids on the action of organophosphorus larvicides in simulium control. Trans. R. Soc. Trop. Med. Hyg. 64:895-906.

Muirhead-Thomson, R.C. 1978. Relative susceptibility of stream macroinvertebrates to temephos and chlorpyrifos, determined in laboratory continuous-flow systems. Arch. Environ. Contam. Toxicol. 7:129-137.

Muirhead-Thomson, R.C. 1979. Experimental studies on macroinvertebrate predator-prey impact of pesticides. The reactions of Rhyacophila and

Hydropsyche (Trichoptera) larvae to simulium larvicides. Can. J. Zool. 57:2264-2270.

Muirhead-Thomson, R.C. and J. Merryweather. 1969. Effects of larvicides on simulium eggs. Nature 22:858-859.

Mulla, M.S. and A.M. Khasawinah. 1969. Laboratory and field evaluation of larvicides against chironomid midges. J. Econ. Entomol. 62:37-41.

Mulla, M.S., R.L. Norland, D.M. Fanara, H.A. Darwazeh and D.W. McKean. 1971. Control of chironomid midges in recreational lakes. J. Econ. Entomol. 64:300-307.

Mulla, M.S., R.L. Norland, W.E. Westlake, B. Dell and J. St. Amant. 1973. Aquatic midge larvicides, their efficacy and residues in water, soil, and fish in a warm water lake. Environ. Entomol. 2:58-65.

Naqvi, S.M. 1973. Toxicity of twenty-three insecticides to a tubificid worm Branchiura sowerbyi from the Mississippi delta. J. Econ. Entomol. 66:70-74.

Nelson, J.H. and E.S. Evans, Jr. 1973. Field evaluation of larvicidal effectiveness, effects on nontarget species and environmental residues of a slow-release polymer formulation of chlorpyrifos, March-October 1973. AD/A-002054. National Technical Information Service, Springfield, VA.

Nelson, J.H., D.L. Stoneburner, E.S. Evans, N.E. Pennington and M.V. Meisch. 1976a. Diatom diversity as a function of insecticidal treatment with a controlled release formulation of chlorpyrifos. Bull. Environ. Contam. Toxicol. 15: 630-634.

Nelson, J.H., N.E. Pennington and M.V. Meisch. 1976b. Use of a controlled release material for control of rice field mosquitoes. Arkansas Farm Res. 25:9.

Norberg, T.J. and D.I. Mount. 1985. A new fathead minnow (Pimephales promelas) subchronic toxicity test. Environ. Toxicol. Chem. 4:711-718.

Papst, M.H. and M.G. Boyer. 1980. Effects of two organophosphorus insecticides on the chlorophyll a and pheopigment concentrations of standing ponds. Hydrobiologia 69:245-250.

Peterson, R.H. 1976. Temperature selection of juvenile Atlantic salmon (Salmo salar) as influenced by various toxic substances. J. Fish. Res. Board Can. 33:1722-1730.

Phipps, G.L. and G.W. Holcombe. 1985a. A method for aquatic multiple species toxicant testing: Acute toxicity of 10 chemicals to 5 vertebrates and 2 invertebrates. Environ. Pollut. (Series A) 38:141-157.

Phipps, G.L. and G.W. Holcombe. 1985b. U.S. EPA, Duluth, MN. (Memorandum to C.E. Stephan, U.S. EPA, Duluth, MN. October 22.)

Polls, I., B. Greenberg and C. Lue-Hing. 1975. Control of nuisance midges in a channel receiving treated municipal sewage. Mosq. News 35:533-537.

Ramke, D. 1969. Development of organophosphorus resistant Aedes nigromaculis in the Tulane Mosquito Abatement District. Proc. Pap. Annu. Conf. Calif. Mosq. Control Assoc. 37:63.



Rawn, G.P., G.R. Webster and G.M. Findley. 1978. Effect of pool bottom substrate on residues and bioactivity of chlorpyrifos against larvae of Culex tarsalis (Diptera: Culicidae). Can. Entomol. 110:1269-1276.

Rettich, F. 1977. The susceptibility of mosquito larvae to eighteen insecticides in Czechoslovakia. Mosq. News 37:252-257.

Rettich, F. 1979. Laboratory and field investigations in Czechoslovakia with fenitrothion, pirimiphos-methyl, temephos and other organophosphorus larvicides applied as sprays for control of Culex pipiens molestus and Aedes cantans. Mosq. News 39: 320-328.

Roberts, D. and T.A. Miller. 1970. The effects of diatoms on the larvicidal activity of Dursban, November 1969-March 1970. AD-724647. National Technical Information Service, Springfield, VA.

Roberts, D.R. and T.A. Miller. 1971. Effects of polymer formulations of Dursban and Abate on nontarget organism populations, April-October 1970. AD-729342. National Technical Information Service, Springfield, VA.

Roberts, L.W., D.R. Roberts, T.A. Miller, L.L. Nelson and W.W. Young. 1973a. Polymer formulations of mosquito larvicides. II. Effects of a polyethylene formulation of chlorpyrifos on Culex populations naturally infesting artificial field pools. Mosq. News 33:155-161.

Roberts, D.R., L.W. Roberts, T.A. Miller, L.L. Nelson and W.W. Young. 1973b. Polymer formulations of mosquito larvicides. III. Effects of a polyethylene formulation of chlorpyrifos on non-target populations naturally infesting artificial field pools. Mosq. News 33:165-172.

- Roberts, R.H., W.B. Kottkamp and M.V. Meisch. 1984. Larvicide evaluations against the rice field mosquito Psorophora columbiae. Mosq. News 44:84-86.
- Rongsriyam, Y., S. Prownebon and S. Hirakoso. 1968. Effects of insecticides on the feeding activity of a guppy, a mosquito-eating fish in Thailand. Bull. W.H.O. 39:977-980.
- Ruber, E. and J. Baskar. 1969. Sensitivities of selected microcrustacea to eight mosquito toxicants. Proc. 55th Annu. Meet. N.J. Mosq. Exterm. Assoc. pp. 99-103.
- Ruber, E. and R. Kocor. 1976. The measurement of upstream migration in a laboratory stream as an index of potential side-effects of temephos and chlorpyrifos on Gammarus fasciatus. Mosq. News 36:424-429.
- Saleh, M.S., I.A. Gaaboub and M.I. Kassem. 1981. Larvicidal effectiveness of three controlled release formulations of Dursban and dimilin on Culex pipiens and Aedes aegypti. J. Agric. Sci. 97:87-96.
- Sanders, H.O. 1969. Toxicity of pesticides to the crustacean Gammarus lacustris. Technical Paper No. 25. U.S. Fish and Wildlife Service, Columbia, MO.
- Sanders, H.O. 1972. Toxicity of some insecticides to four species of malacostracan crustaceans. Technical Paper No. 66. U.S. Fish and Wildlife Service, Washington, DC.
- Sanders, H.O. and O.B. Cope. 1968. The relative toxicities of several pesticides to naiads of three species of stoneflies. Limnol. Oceanogr. 13:112-117.

- Schaefer, C.H. and E.F. Dupras, Jr. 1970. Factors affecting the stability of Dursban in polluted waters. *J. Econ. Entomol.* 63:701-705.
- Schimmel, S.C., R.L. Garnas, J.M. Patrick, Jr. and J.C. Moore. 1983. Acute toxicity, bioconcentration, and persistence of AC 222,705, benthocarb, chlorpyrifos, fenvalerate, methyl parathion, and permethrin in the estuarine environment. *J. Agric. Food Chem.* 31:104-113.
- Scott, J. and M.S. Redmond. 1986a. U.S. EPA, Narragansett, RI. (Memorandum to D.J. Hansen, U.S. EPA, Narragansett, RI.)
- Scott, J. and M.S. Redmond. 1986b. U.S. EPA, Narragansett, RI. (Memorandum to D.J. Hansen, U.S. EPA, Narragansett, RI.)
- Scirocchi, A. and A. D'Erme. 1980. Toxicity of seven insecticides on some species of freshwater fishes. *Riv. Parassitol.* 41:113-121.
- Siefert, R.E., C.F. Kleiner, B.R. Nordling, L.H. Mueller, D.K. Tanner, A.W. Jarvinen, J.A. Zischke, N. Larson and R.L. Anderson. 1984. Effects of Dursban (chlorpyrifos) on nontarget aquatic organisms in a natural pond undergoing mosquito control treatment. Progress Report. U.S. EPA, Duluth, MN.
- Smith, G.N., B.S. Watson and F.S. Fischer. 1966. The metabolism of [<sup>14</sup>C]O,O-diethyl-O-(3,5,6-trichloro-2-pyridyl)phosphorothiole (Dursban) in fish. *J. Econ. Entomol.* 59:1464-1475.
- Steelman, C.D., B.R. Craven and E.J. Vallavaso. 1969. Control of southern house mosquito larvae in Louisiana papermill log ponds. *J. Econ. Entomol.* 62:1152-1154.

Stephan, C.E., D.I. Mount, D.J. Hansen, J.H. Gentile, G.A. Chapman and W.A. Brungs. 1985. Guidelines for deriving numerical national water quality criteria for the protection of aquatic organisms and their uses. PB85-227049. National Technical Information Service, Springfield, VA.

Stewart, J.P. 1977. Synergism of chlorpyrifos by DEF in the control of organophosphorus resistant Culex pipiens quinquefasciatus larvae, with notes on synergism of parathion and fenthion. Proc. Pap. Annu. Conf. Calif. Mosq. Vector Control Assoc. 45:132-133.

Tagatz, M.E., N.R. Gregory and G.R. Plaia. 1982. Effects of chlorpyrifos on field- and laboratory-developed estuarine benthic communities. J. Toxicol. Environ. Health 10:411-421.

Tawfik, M.S. and R.H. Gooding. 1970. Dursban and Abate clay granules for larval mosquito control in Alberta. Mosq. News 30:461-464.

Taylor, R.T. and H.F. Schoof. 1971. Experimental field treatment with larvicides for control of Anopheles, Aedes, and Culex mosquitoes. J. Econ. Entomol. 64:1173-1176.

Thayer, A. and E. Ruber. 1976. Previous feeding history as a factor in the effect of temephos and chlorpyrifos on migration of Gammarus fasciatus. Mosq. News 36:429-432.

Thirugnanam, M. and A.J. Forgash. 1977. Environmental impact of mosquito pesticides: Toxicity and anticholinesterase activity of chlorpyrifos to fish in a salt marsh habitat. Arch. Environ. Contam. Toxicol. 5:415-425.

Thompson, A.H., C.L. Barnes and D.A. Mathews. 1970. Injection of Dursban spray emulsion at half mile intervals controls mosquitoes and chironomid larvae in large drainage channels. Proc. Pap. Annu. Conf. Calif. Mosq. Control Assoc. 38:76-77.

U.S. Bureau of Commercial Fisheries. 1965. Unpublished laboratory data.  
U.S. EPA, Environmental Research Laboratory, Gulf Breeze, FL.

U.S. Bureau of Commercial Fisheries. 1967. Unpublished laboratory data.  
U.S. EPA, Environmental Research Laboratory, Gulf Breeze, FL.

U.S. EPA. 1983a. Water quality standards regulation. Fed. Regist. 48:  
51400-51413. November 8.

U.S. EPA. 1983b. Water quality standards handbook. Office of Water Regulations and Standards, Washington, DC.

U.S. EPA. 1985a. Appendix B - Response to public comments on "Guidelines for deriving numerical national water quality criteria for the protection of aquatic organisms and their uses." Fed. Regist. 50:30793-30796. July 29.

U.S. EPA. 1985b. Technical support document for water-quality based toxics control. Office of Water, Washington, DC. September.

U.S. EPA. 1986. Chapter 1 - Stream design flow for steady-state modeling. In: Book VI - Design Conditions. In: Technical guidance manual for performing wasteload allocations. Office of Water, Washington, DC.

Verma, K.V. and S.J. Rahman. 1984. Determination of minimum lethal time of commonly used mosquito larvicides. J. Commun. Dis. 16:162-164.

- Wallace, R.R., A.S. West, A.E. Downe and H.B. Hynes. 1973. The effects of experimental blackfly (Diptera: Simuliidae) larviciding with Abate, Dursban and methoxychlor on stream invertebrates. Can. Entomol. 105: 817-831.
- Walsh, G.E. 1981. Effects of pesticides and industrial wastes on unicellular algae and seagrass. Part IIA. In: "Research and Development: Experimental Environments Branch, Progress Report for Fiscal Year 1981." Unpublished laboratory data. U.S. EPA, Environmental Research Laboratory, Gulf Breeze, FL.
- Walsh, G.E. 1983. Cell death and inhibition of population growth of marine unicellular algae by pesticides. Aquatic Toxicol. 3:209-214.
- Washino, R.K., K.G. Whitesell and D.J. Womeldorf. 1968. The effect of low volume application of Dursban on nontarget organisms. Down Earth 24:21-22.
- Washino, R.K., K.G. Whitesell, E.J. Sherman, M.C. Kramer and R.J. McKenna. 1972a. Rice field mosquito control studies with low volume Dursban sprays in Colusa County, California. III. Effects upon the target organisms. Mosq. News 32:375-382.
- Washino, R.K., W. Ahmed, J.D. Linn and K.G. Whitesell. 1972b. Rice field mosquito control studies with low volume Dursban sprays in Colusa County, California IV. Effects upon aquatic nontarget organisms. Mosq. News 32:531-537.
- Wilder, W.H. and C.S. Schaefer. 1969. Organophosphorus resistance levels in adults and larvae of the pasture mosquito, Aedes nigromaculis in the San Joaquin Valley of California. Proc. Pap. Annu. Conf. Calif. Mosq. Control Assoc. 37:64-67.

Wilkinson, R.N., W.W. Barnes, A.R. Gillogly and C.D. Minnemeyer. 1971. Field evaluation of slow-release mosquito larvicides. J. Econ. Entomol. 64:1-3.

Wilton, D.P., L.E. Fetzer and R.W. Fay. 1973. Insecticide baits for anopheline larvae. Mosq. News 33:198-203.

Winner, R.A., C.D. Steelman and P.E. Schilling. 1978. Effects of selected insecticides on Romanomermis culicivorax, a mermithid nematode parasite of mosquito larvae. Mosq. News 38:546-553.

Winterlin, W.L., K. Moilanen and W.E. Burgoyne. 1968. Residues of Dursban insecticide following mosquito control applications. Down Earth 24:34-37.

Womeldorf, D.J., R.K. Washino, K.E. White and P.A. Gieke. 1970. Insecticide susceptibility of mosquitoes in California: Response of Anopheles freeborni larvae to organophosphorus compounds. Mosq. News 30:375-382.

Yap, H.H. and S.C. Ho. 1977. Evaluation of Dursban and Dowco 214 as mosquito larvicides in rice-fields. Southeast Asian J. Trop. Med. Public Health 8:63-70.

Yoshioka, Y., T. Mizuno, Y. Ose and T. Sato. 1986. The estimation for toxicity of chemicals on fish by physio-chemical properties. Chemosphere 15:195-203.

Zarogian, G., J.F. Heltshe and M. Johnson. 1985. Estimation of toxicity to marine species with structure-activity models developed to estimate toxicity to freshwater fish. Aquat. Toxicol. 6:251-270.

Zboray, E.P. and M.C. Gutierrez. 1979. Insecticide susceptibility of mosquitoes in California: Status of organophosphorus resistance in larval Culex Larsalis through 1978, with notes on mitigating the problem. Proc. Pap. Annu. Conf. Calif. Mosq. Vector Control Assoc. 47:26-28.

Zepp, R.G. and P.F. Schlotzhauer. 1983. Influence of algae on photolysis rates of chemicals in water. Environ. Sci. Technol. 17:462-468.