



Madden-Julian Oscillation: Recent Evolution, Current Status and Predictions

**Update prepared by
Climate Prediction Center / NCEP
June 16, 2008**



Outline

- **Overview**
- **Recent Evolution and Current Conditions**
- **MJO Index Information**
- **MJO Index Forecasts**
- **MJO Composites**



Overview


- **The MJO has weakened with the enhanced phase centered across the eastern Maritime Continent.**
- **Based on the latest observations, the MJO is expected to have minimal impacts over the next 1-2 weeks. MJO dynamical forecast tools have not performed well with the current MJO activity.**
- **During Week 1, enhanced convection is expected for parts of India, extreme southeast China and the Bay of Bengal. Drier than average conditions are anticipated for Central America and the eastern Pacific Ocean.**

Additional potential impacts across the global tropics are available at:
<http://www.cpc.ncep.noaa.gov/products/precip/CWlink/ghazards/ghaz.shtml>



850-hPa Vector Wind Anomalies (m s^{-1})

Note that shading denotes the magnitude of anomalous wind vectors




The Indian monsoon flow has weakened slightly during the last five days.

Easterly anomalies decreased across the Maritime Continent and the western Pacific.

Westerly anomalies weakened across the eastern Pacific during the last five days.



850-hPa Zonal Wind Anomalies (m s^{-1})



Westerly anomalies (orange/red shading) represent anomalous west-to-east flow

Easterly anomalies (blue shading) represent anomalous east-to-west flow

Moderate-to-strong MJO activity is evident from late November to mid-late February as shown by westerly anomalies shifting eastward from the Indian Ocean across Indonesia and a weakening of the easterlies at the Date Line during early mid-January and mid-February.


MJO activity was weaker during much of March and April.

During mid-May, easterlies weakened near the Date Line associated with the latest MJO activity.

Most recently, winds are closer to average across much of the Pacific Ocean.



OLR Anomalies: Last 30 days



Drier-than-normal conditions, positive OLR anomalies (red shading)
Wetter-than-normal conditions, negative OLR anomalies (blue shading)


Suppressed convection developed across the Indian Ocean and western Indonesia in late May.

At the same time and in early June, wet conditions became widespread across the Atlantic Ocean and equatorial Africa.

During early-mid June, enhanced convection developed across the Indian Ocean, Arabian Sea, India and Indonesia as the MJO propagated eastward.



Outgoing Longwave Radiation (OLR) Anomalies (7.5°S-7.5°N)



Drier-than-normal conditions, positive OLR anomalies (blue shading)

Wetter-than-normal conditions, negative OLR anomalies (yellow/red shading)

(Courtesy of the Bureau of Meteorology - Australia)

Moderate-to-strong MJO activity was evident from mid-November to mid-February with coherent eastward propagation of enhanced (suppressed) convection indicated by the dashed (dotted) lines.

Weak MJO activity was evident during mid-late March.

Strong suppressed convection organized across the Indian Ocean and shifted eastward during mid-late May.


A weaker area of normal-to-enhanced convection shifted through the Indian Ocean and Maritime Continent. Enhanced convection is more evident north of the equator.



IR Temperatures (K) / 200-hPa Velocity Potential Anomalies

Positive anomalies (brown contours) indicate unfavorable conditions for precipitation

Negative anomalies (green contours) indicate favorable conditions for precipitation



The large-scale wave structure in anomalous velocity potential has become less coherent during the past week. Upper-level convergence is present across much of the Pacific with upper-level divergence observed over parts of the Indian Ocean and Southeast Asia.




200-hPa Velocity Potential Anomalies (5°S-5°N)

Positive anomalies (brown shading) indicate unfavorable conditions for precipitation

Negative anomalies (green shading) indicate favorable conditions for precipitation

Time



Moderate-to-strong MJO activity developed in mid-November and continued into mid-February.

Weak MJO activity was evident during parts of March.

The MJO was largely incoherent during the month of April.


The MJO strengthened during May as velocity potential anomalies increased with eastward propagation evident.

Longitude



200-hPa Vector Wind Anomalies (m s^{-1})


Note that shading denotes the magnitude of anomalous wind vectors



Westerly anomalies over the central Pacific Ocean have increased during the last five-ten days.



200-hPa Zonal Wind Anomalies (m s^{-1})



Westerly anomalies (orange/red shading) represent anomalous west-to-east flow

Easterly anomalies (blue shading) represent anomalous east-to-west flow


MJO activity is evident in the upper-levels by eastward propagation of easterly (westerly) anomalies by dashed (dotted) lines globally from late November to mid-February.

During March, MJO activity was evident as easterly anomalies propagated eastward from the western hemisphere to the Maritime continent.

Easterly anomalies increased during early May over Indonesia and have shifted eastward. Now westerly anomalies are centered over the central Pacific.



Weekly Heat Content Evolution in the Equatorial Pacific



Kelvin wave activity (downwelling phases indicated by dashed lines) was observed from May 2007 to February 2008 and affected sub-surface temperature departures at varying degrees across the Pacific Ocean. The strongest wave occurred during May and June 2007.

During September and October, negative heat content anomalies increased markedly across the eastern Pacific Ocean and continued until February 2008.

Beginning in March, increasingly positive anomalies have developed across parts of the western and central Pacific.

Since April, a region of positive anomalies has strengthened in the eastern Pacific.



MJO Index -- Information

- The MJO index illustrated on the next several slides is the CPC version of the Wheeler and Hendon index (2004, hereafter WH2004).


Wheeler M. and H. Hendon, 2004: An All-Season Real-Time Multivariate MJO Index: Development of an Index for Monitoring and Prediction, *Monthly Weather Review*, 132, 1917-1932.

- The methodology is nearly identical to that described in WH2004 but small deviations from the BMRC figure are possible at times due to differences in input data and methodology. These typically occur during weak MJO periods.
- The index is based on a combined Empirical Orthogonal Function (EOF) analysis using fields of near-equatorially-averaged 850-hPa and 200-hPa zonal wind and outgoing longwave radiation (OLR).



MJO Index -- Recent Evolution



- The axes (RMM1 and RMM2) represent daily values of the principal components from the two leading modes
- The triangular areas indicate the location of the enhanced phase of the MJO
- Counter-clockwise motion is indicative of eastward propagation
- Distance from the origin is proportional to MJO strength
- Line colors distinguish different months



The MJO index has decreased in amplitude since mid-May, but has continued to propagate into the Maritime Continent region.



MJO Index – Historical Daily Time Series



Time series of daily MJO index amplitude from 1995 to present

Plots put current MJO activity in historical context



Ensemble GFS MJO Forecasts


Yellow Lines – 20 Individual Members
Green Line – Ensemble Mean

RMM1 and RMM2 values for the most recent 40 days and forecasts from the ensemble Global Forecast System (GEFS) for the next 15 days

light gray shading: 90% of forecasts
dark gray shading: 50% of forecasts

The GEFS predicts the MJO signal to further weaken during the next week with some eastward propagation.

The GEFS has consistently underestimated the eastward propagation and strength of the recent MJO activity.






Ensemble Mean GFS MJO Forecast


Figures below show MJO associated OLR anomalies only (reconstructed from RMM1 and RMM2) and do not include contributions from other modes (*i.e.*, ENSO, monsoons)

Spatial map of OLR anomalies for the next 15 days



The forecast from the GEFS for MJO-associated convection indicates weak wet conditions for western Indonesia slowly shifting northeastward.

Time-longitude section of (7.5°S-7.5°N) OLR anomalies for the last 180 days and for the next 15 days






Statistical MJO Forecast

Figure below shows MJO associated OLR anomalies only (reconstructed from RMM1 and RMM2) and do not include contributions from other modes (*i.e.*, ENSO, monsoons)

Spatial map of OLR anomalies and 850-hPa wind vectors for the next 20 days

(Courtesy of the Bureau of Meteorology Research Centre - Australia)

Weak MJO activity is forecast. Dry conditions are forecast to develop in the Indian Ocean during the period.





MJO Composites – Global Tropics

Precipitation Anomalies (May-Sep)

850-hPa Wind Anomalies (May-Sep)

