

Chapter 4 – List of Preparers

4.1 List of Preparers

The following people were the primary authors of this EIS.

<p><i>Vince Archer</i> US Forest Service TEAMS Enterprise Unit B.S. Physical Science, M.S. Soils Soil Scientist</p>	<p><i>Barbara A. F. Ott</i> US Forest Service TEAMS Enterprise Unit M.S. Management, B.A. Business Administration Social Scientist</p>
<p><i>Sandy Caveny</i> US Forest Service TEAMS Enterprise Unit B.S Forest Management Recreation Specialist</p>	<p><i>Janice Schultz</i> US Forest Service TEAMS Enterprise Unit Writer/Editor – Project Record</p>
<p><i>Allison Kuehl</i> US Forest Service TEAMS Enterprise Unit M.S. Range Management, B.S. Wildlife Wildlife Biologist</p>	<p><i>Carol Thornton</i> US Forest Service TEAMS Enterprise Unit B.S. Geology, M.S. Hydrology/Hydro- geology Hydrologist</p>
<p><i>Julie Laufman</i> US Forest Service TEAMS Enterprise Unit PhD. Rangeland Ecosystem Science Botanist</p>	<p><i>Chris Tootell</i> US Forest Service TEAMS Enterprise Unit B.S. Natural Resources Conservation MS Forest Science Interdisciplinary Team Leader – NEPA</p>
<p><i>Bill Overland</i> US Forest Service TEAMS Enterprise Unit B.S. Forestry and Natural Resources GIS</p>	<p><i>Tiffany Vandosall</i> US Forest Service TEAMS Enterprise Unit B.S. Biology and Chemistry M.S. Aquatic Ecology Fisheries Biologist</p>
<p><i>Scott Reitz</i> US Forest Service TEAMS Enterprise Unit A.A Natural Resources and Conservation B.S. Wildlife Sciences Certified Wildlife Biologist</p>	<p><i>Rochelle Desser</i> US Forest Service TEAMS Enterprise Unit A.A. Geo-technology Regional Oversight</p>

4.2 Consultation with Regulatory Agencies

The Forest Service has initiated consultation with the US Fish and Wildlife Service and NOAA Fisheries regarding potential adverse effects on Endangered Species. The consultation is tiered to programmatic consultation at the Regional Scale.

A Biological Assessment will be prepared for the Preferred Alternative. A Record of Decision will not be signed prior to receiving a Letter of Concurrence from the regulatory agencies supporting determinations of Not Likely to Adversely Affect and/or they issue a Biological Opinion describing terms and conditions associated with a determination of Likely to Adversely Affect.

4.3 Consultation with Tribes

Letters were sent to Tribal leaders of the Nez Perce, Confederate Tribes of the Umatilla Indian Reservation (CTUIR), and Confederate Tribes of the Warm Springs Reservation in April of 2006. None of tribes responded to the letter. The Forest had meetings with various tribal resource staff. During these meetings the tribes were supportive of the Forest's efforts to treat invasive plants and being able to use all the tools/methods described in the Proposed Action. The experience of the CTUIR using aerial treatments for yellowstar thistle is described as successful when intergrated with other land owners. The Nez Perce felt that biological treatments should be an intergrated approach used on the landscape. All tribes have a concern about coordinating herbicide treatments with traditional gathering activities and areas. A process will be developed for notifying each tribe when herbicides are being used as required by the Project Design Features in Chapter 2.

4.4 Consultation with Counties

The Forest Service has worked closely with the County Weed Boards. County staff have presented information to the Forest Service (Available in the Project Record) and participated in field visits. The Counties often implement projects for the Forest Service and other land managers in the area and fully support this project.

4.5 Consultation with Others

Many people within and outside the Forest Service helped the team develop and analyze the project. Managers and specialists from the National Forest reviewed analysis documentation and suggested changes.

Public scoping has occurred on this project since 2006. The public has been apprised of project progress through the newspaper, direct mailings, Notices of Intent published in the Federal Register in 2006, the Forest Schedule of Proposed Actions, informal meetings and discussions, and other media.

Many organizations and individuals have expressed interest in the project; everyone who commented during scoping was offered a hard copy or CD containing the DEIS and Appendices.

The full DEIS and Appendices is also available electronically by website:

<http://www.fs.fed.us/w-w/projects/> or on request (see cover page for more information to request a CD or hard copy).

Hard copies are available for review at Forest Service offices throughout the area. The DEIS has been sent to the Environmental Protection Agency (who commented during scoping) and other federal and state agencies. The following is a list of individuals, tribal governments, government officials, county programs, and individual organizations to whom this DEIS was sent:

Individuals	
Andrew and Mercedes Hansen	Kent Weaver
Dorene McCarthy	Lawson Denny
Fred and Anna May Warnock	Lyle and Margaret Defrees
Gene Walker	Michael Riggs
Greg K. Johnson	Michael D. Rutherford
Jacob and Wendy Bingham	Neal J. and Anna Dow
Jeanne M. Rachau	Robert E. and Edna Loftus
Jeff Dawson	Rod George
Jennifer Schemm	Spencer Scott Lewis
Joann Anderson	Teddy and Virginia Daggett
Jon Croghan	Thomas Thomayer
Kathryn M. Krieger	Willard A. Godfrey
Ken Witty	William and Donna Tsiatsos
Kenneth Grant	
Tribes	
Confederated Tribes of the Umatilla Indian Reservation	
Honorable Antone Minthorn, Chairman Board of Trustees	
Eric Quaempts, Natural Resources Director	
Nez Perce Tribe	
Honorable Sam Penney, Chairman Nez Perce Tribal Executive Committee	
Brooklyn Baptiste, Chairman Natural Resources Subcommittee	
Government Officials	
U.S. Representative Bill Sahli, Todd Snodderly Field Representative	
U.S. Representative Greg Walden, Colby Marshall Field Representative	
U.S. Senator Gordon Smith, Larry Bartee Field Representative	
U.S. Senator Larry Craig, Jeff Sayre Regional Director	
U.S. Senator Mike Crapo, Mitch Silvers Regional Director	
U.S. Senator Ron Wyden, Kathleen Cathey Field Representative	
Counties	
Adams County Commissioners	
Adams County Weed Control, Ray McKinney	
Baker County Commissioners	

Baker County Weed Coordinator, Arnie Grammon
Idaho County Commissioners
Idaho County Weed Management, Carl Crabtree
Nez Perce County Commissioners
Nez Perce County Weed Control, Hugh Jacobs
Union County Commissioners
Union County Weed Control, Gary Dade
Wallowa County Commissioners
Wallowa County Weed Coordinator, Allen Schnetzky
Individual Organizations
Bell Family Partners,
Blue Mountain Biodiversity Project, Karen Coulter
Blue Mountain Forum, Jennie Tucker
Bruno Ranch, Maria Bruno
Eastern Oregon All Terrain Vehicle Association, Tim Mahan
Eastern Oregon Mining Association, Ed Hardt
Flying O Cattle Company, Janet DeLong
Grande Ronde Model Watershed Program, Jeff Oveson
Hells Canyon Preservation Council, Greg Dyson
Hurricane Creek Grange #608
Noxious Weed Management Coordinator, Don Sharrat
Oregon Hunters Association, Charlie Brinton
Oregon Sportsmen Halfway Chapter, Marion Crow
Oregon Wild, Doug Heiken
Pacific Northwest 4-Wheel Drive Association, Art Waugh
Rock'n "J" Properties
Rocky Mountain Elk Foundation, Steve Martin
Sams Orchard, Dan Sams
Sierra Club, Asante Riverwind
The Wilderness Society, Brad Brooks
Tri-County Cooperative Weed Management, Gary Winans
Upper Burnt River Weed Control District, Ross Shumway
VFW Auxiliary, Agnes Roberts
Wallowa Resources, Nils Christoffersen
Wallowa Valley Trail Riders Association, Randall & Velma Parmelee

4.6 Glossary

Active ingredient (a.i.) - In any pesticide product, the component (a chemical or biological substance) that kills or otherwise controls the target pests - Pesticides are regulated primarily on the basis of active ingredients. The remaining ingredients are called “inerts.”

Acute effect - An adverse effect on any living organism in which severe symptoms develop rapidly and often subside after the exposure stops.

Acute exposure - A single exposure or multiple brief exposures occurring within a short time (e.g., 24 hours or less in humans). The classification of multiple brief exposures as “acute” is dependant on the life span of the organism. (See also, chronic exposure and cumulative exposure.)

Acute toxicity - Any harmful effect produced in an organism through an acute exposure to one or more chemicals.

Adaptation - Changes in an organism's physiological structure or function or habits that allow it to survive in new surroundings.

Adapted - How well organisms are physiologically or structurally suited for survival, growth, and resistance to pests and diseases in a particular environment.

Additive effect - A situation in which the combined effects of exposure to two chemicals simultaneously is equal to the sum of the effect of exposure to each chemical given alone. The effect most commonly observed when an organism is exposed to two chemicals together is an additive effect.

Adaptive management - A continuing process of action-based planning, monitoring, researching, evaluating, and adjusting with the objective of improving implementation and achieving the goals of the standards and guidelines

Adjuvant(s) - Chemicals that are added to pesticide products to enhance the toxicity of the active ingredient or to make the active ingredient easier to handle or mix.

Administratively Withdrawn Areas (AWA) - Areas removed from the suitable timber base through agency direction and land management plans.

Adsorption - The tendency of one chemical to adhere to another material such as soil.

Aerobic - Life or processes that require, or are not destroyed by, the presence of oxygen. (See also, anaerobic.)

Affected Environment - Existing biological, physical, social, and economic conditions of an area subject to change, both directly and indirectly, as the result of a proposed human action.

Agent - Any substance, force, radiation, organism, or influence that affects the body. The effects may be beneficial or injurious.

Agency for Toxic Substances and Disease Registry (ATSDR) - Federal agency within the Public Health Service charged with carrying out the health-related analyses under

Comprehensive Environmental Response, Compensation, and Liability Act (CERCLA) and Superfund Amendments and Reauthorization Act (SARA).

Alien species - “With respect to a particular ecosystem, any species, including its seeds, eggs, spores, or other biological material capable of propagating that species, that is not native to that ecosystem” (Executive Order 13122, 2/3/99). (See also, invasive, noxious, and weed species.)

Allelopathy - The suppression of growth of one plant species due to the release of toxic substances by another plant.

Alluvial - Relating to clay, silt, sand, gravel, or similar detrital material deposited by flowing water. Alluvial deposits may occur after a heavy rain storm.

Ambient - Usual or surrounding conditions.

Amphibian - Any of a class of cold-blooded vertebrates (including frogs, toads, or salamanders) intermediate in many characteristics between fishes and reptiles and having gilled aquatic larvae and air-breathing adults.

Anadromous - Fish that spend their adult life in the sea but swim upriver to fresh water spawning grounds to reproduce.

Anaerobic - Life or process that occurs in, or is not destroyed by, the absence of oxygen. (See also, aerobic.)

Anions - Negatively charged ions in solution e.g., hydroxyl or OH⁻ ion. (See also, cations.)

Annual - A plant that endures for not more than a year. A plant which completes its entire life cycle from germinating seedling to seed production and death within a year.

Annuity - Payment or receipt of a series of equal amounts at stated intervals for a specified number of time periods. An “annuity due” is a series of equal value outputs or inputs occurring for N equal time periods with “payments” made at the beginning of each period.

Anoxia - Literally, "without oxygen." A deficiency of oxygen reaching the tissues of the body especially of such severity as to result in permanent damage.

Aquatic Influence Zone – The inner half of a Riparian Reserve.

Aqueous - Describes a water-based solution or suspension.

Aquifer - An underground geological formation, or group of formations, containing usable amounts of groundwater that can supply wells and springs.

Arid - A terrestrial region lacking moisture, or a climate in which the rainfall is not sufficient to support the growth of most vegetation.

Background level - In pollution, the level of pollutants commonly present in ambient media (air, water, soil.)

Bacteria - Microscopic living organisms that metabolize organic matter in soil, water, or other environmental media. Some bacteria can also cause human, animal and plant health problems.

Basal application - In pesticides, the spreading of a chemical on stems or trunks of plants just above the soil line.

Base - Substances that (usually) liberate hydroxyl (OH⁻) anions when dissolved in water and weaken a strong acid.

Benchmark - A dose associated with a defined effect level or designated as a no effect level.

Benthic region - The bottom layer of a body of water.

Benthos - The plants and animals that inhabit the bottom layer of a water body.

Best Management Practices (BMP) - A practice or combination of practices determined by a state or an agency to be the most effective and practical means (technological, economic, and institutional) of controlling point and non-point source pollutants at levels compatible with environmental quality.

Bioaccumulation - The increase in concentration of a substance in living organisms as they take in contaminated air, water, or food because the substance is very slowly metabolized or excreted (often concentrating in the body fat.)

Bioassay - (1) To measure the effect of a substance, factor, or condition using living organisms. (2) A test to determine the toxicity of an agent to an organism.

Bioconcentration - The accumulation of a chemical in tissues of a fish or other organism to levels greater than in the surrounding water or environment.

Bioconcentration Factor (BCF) - The concentration of a compound in an aquatic organism divided by the concentration in the ambient water of the organism.

Biodegradability - Susceptibility of a substance to decomposition by microorganisms; specifically, the rate at which compounds may be chemically broken down by bacteria and/or natural environmental factors.

Biodiversity or biological diversity - The diversity of living things (species) and of life patterns and processes (ecosystem structures and functions). Includes genetic diversity, ecosystem diversity, landscape and regional diversity, and biosphere diversity.

Biological control - The use of natural enemies, including invertebrate parasites and predators (usually insects, mites, and nematodes,) and plant pathogens to reduce populations of nonnative, invasive plants.

Biological magnification - The process whereby certain substances such as pesticides or heavy metals increase in concentration as they move up the food chain.

Biologically sensitive - A term used to identify a group of individuals who, because of their developmental stage or some other biological condition, are more susceptible than the general population to a chemical or biological agent in the environment.

Biomass - The amount of living matter.

Biota or Biome - All living organisms of a region or system.

Body Burden - The amount of a chemical stored in the body at a given time, especially a potential toxin in the body as the result of exposure.

Broadcast application - Herbicide treatment method generally used along roads; boom truck spray is directed at target species. Broadcast methods are used for larger infestations where spot treatments would not be effective.

Bryophytes - Plants of the phylum Bryophyta, including mosses, liverworts, and hornworts; characterized by the lack of true roots, stems, and leaves.

Buffer Zone - A strip of untreated land that separates a waterway or other environmentally sensitive area from an area being treated with pesticides.

Candidate species - Those plant and animal species that, in the opinion of the Fish and Wildlife Service (FWS) or National Oceanic and Atmospheric Administration (NOAA) Fisheries, may qualify for listing as “endangered” or “threatened.” The FWS recognizes two categories of candidates. Category 1 candidates are taxa for which the FWS has on file sufficient information to support proposals for listing. Category 2 candidates are taxa for which information available to the FWS indicates that proposing to list is possibly appropriate, but for which sufficient data are not currently available to support proposed rules.

Capillary fringe - The zone above the water table within which the soil or rock is saturated by water under less than atmospheric pressure.

Carcinogen - A chemical capable of inducing cancer.

Carrier - A non-pesticidal substance added to a commercial pesticide formulation to make it easier to handle or apply.

Chemical Abstracts Service (CAS) Registry Number - An assigned number used to identify a chemical. Chemical Abstracts Service is an organization that indexes information published in Chemical Abstracts by the American Chemical Society and that provides index guides to help locate information about particular substances in the abstracts. Sequentially assigned CAS numbers identify specific chemicals. The numbers have no chemical significance. The CAS number is a concise, unique means of chemical identification.

Cations - Positively charged ions in a solution. (See also, anion.)

Characteristic Landscape - The naturally established landscape within a scene or scenes being viewed.

Chemical Control - The use of naturally derived or synthetic chemicals called herbicides to eliminate or control the growth of invasive plants.

Chronic exposure - Exposures that extend over the average lifetime or for a significant fraction of the lifetime of the species (for a rat, chronic exposure is typically about two years). Chronic exposure studies are used to evaluate the carcinogenic potential of chemicals and other long-term health effects. (See also, acute and cumulative exposure.)

Chronic Reference Dose (RfD) - An estimate of a lifetime daily exposure level (in mg/kg/day) for the human population, including sensitive subpopulations, that is likely to be without an

appreciable risk of deleterious effects. Chronic RfDs are specifically developed to be protective for long-term exposure to a compound (seven years to lifetime.)

Chronic toxicity - The ability of a substance or mixture of substances to cause harmful effects over an extended period, usually upon repeated or continuous exposure sometimes lasting for the entire life of the exposed organism

Code of Federal Regulations (CFR) - Document that codifies all rules of the executive departments and agencies of the federal government. It is divided into fifty volumes, known as titles. Title 40 of the CFR (referenced as 40 CFR) lists all environmental regulations, including regulations for EPA pesticide programs (40 CFR Parts 150-189).

Competitive seeding - Treatment method; most effective after weed populations have been reduced by other control actions.

Congressionally Designated Areas - Areas that require Congressional enactment for their establishment, such as National Parks, Wild and Scenic Rivers, National Recreation Areas, National Monuments, and Wilderness. Also referred to as Congressional Reserves. Includes similar areas established by Executive Order, such as National Monuments.

Conifer - An order of the Gymnospermae, comprising a wide range of trees and a few shrubs, mostly evergreens that bear cones and have needle-shaped or scale-like leaves. Conifer timber is commercially identified as softwood.

Connected actions - Exposure to other chemical and biological agents, in addition to exposure to a specific pesticide formulation in a field application to control pest organisms.

Contaminants - For chemicals, impurities present in a commercial grade chemical. For biological agents, other agents that may be present in a commercial product.

Control - Means, as appropriate, eradicating, suppressing, reducing, or managing invasive plant species populations, preventing spread of invasive plants from areas where they are present, and taking steps such as restoration of native plant species and habitats to reduce the effects of invasive plants and to prevent further invasions (Executive Order 13122, 2/3/99).

Cultural control - The establishment or maintenance of competitive vegetation, use of fertilizing, mulching, prescribed burning, or grazing animals to control or eliminate invasive plants. Prescribed burning and grazing animals are cultural control treatments that would not be utilized under this project; nor are they covered under this EIS.

Cumulative Effect (CE) - The impact on the environment that results from the incremental impact of the action when added to other past, present, and reasonably foreseeable future actions—regardless of what agency (federal or non-federal) or person undertakes such other actions. Cumulative impacts can result from individually minor, but collectively significant, actions taking place over a period of time (40 CFR 1508.7).

Cumulative exposure - Exposure resulting from one or more activities that are repeated over a period of time. (See also, acute and chronic exposure.)

Detritus - Loose fragments, particles, or grains formed by the disintegration of organic matter or rocks.

Discount - In economics, discounting is the process of carrying an end value backward in time at compound interest.

Distance Zones - Landscape areas denoted by specified distances from the observer. Used as a frame of reference in which to discuss landscape attributes or the scenic effect of human activities in a landscape.

Disturbance - An effect of a planned human management activity, or unplanned native or exotic agent or event that changes the state of a landscape element, landscape pattern, or regional composition.

Dosage/Dose - (1) The actual quantity of a chemical administered to an organism or to which it is exposed. (2) The amount of a substance that reaches a specific tissue (e.g. the liver). (3) The amount of a substance available for interaction with metabolic processes after crossing the outer boundary of an organism.

Dose Rate - In exposure assessment, dose per time unit (e.g. mg/day); also called dosage.

Dose Response - Changes in toxicological responses of an individual (such as alterations in severity of symptoms) or populations (such as alterations in incidence) that are related to changes in the dose of any given substance.

Drift - The portion of a sprayed chemical that is moved by wind off of a target site.

Emergent Vegetation - Plants growing out of or standing in water, in contrast to “submerged aquatic vegetation (SAV),” which grows entirely underneath the waters’ surface.

Endangered Species - Any species listed in the Federal Register as being in danger of extinction throughout all, or a significant portion, of its range.

Endangered Species Act (ESA) - A law passed in 1973 to conserve species of wildlife and plants, determined by the Director of the U.S. Fish and Wildlife Service or the NOAA Fisheries to be endangered or threatened with extinction in all or a significant portion of its range. Among other measures, ESA requires all federal agencies to conserve these species and consult with the Fish and Wildlife Service or NOAA Fisheries on federal actions that may affect these species or their designated critical habitat.

Endemic - A species or other taxonomic group that is restricted to a particular geographic region due to factors such as isolation or response to soil or climatic conditions. (Compare to “Indigenous” and “Native.”)

Environmental justice - Executive Order 12898 of February 11, 1994 requires federal agencies, to the greatest extent practicable and permitted by law, to make achieving environmental justice part of its mission by identifying and addressing, as appropriate, disproportionately high and adverse human health or environmental effects of its programs, policies, and activities on minority populations and low-income populations in the United States and its territories and possessions, the District of Columbia, the Commonwealth of Puerto Rico, and the commonwealth of the Mariana Islands.

Exposure assessment - The process of estimating the amount of contact with a chemical or biological agent that an individual or a population of organisms will receive from a pesticide application conducted under specific, stated circumstances.

Exotic – Non-native species; introduced from elsewhere, but not completely naturalized. (See also alien and introduced species.)

Extirpate - To destroy completely; wipe out.

Extrapolation - The use of a model to make estimates of values of a variable in an unobserved interval from values within an already observed interval.

Fauna - The animals of a specified region or time.

Federally listed species - Formally listed as a threatened or endangered species under the Endangered Species Act. Designations are made by the Fish and Wildlife Service or the National Marine Fisheries Service.

Federal Insecticide and Rodenticide Act (FIFRA) Pesticide Ingredient - An ingredient of a pesticide that must be registered with EPA under the Federal Insecticide, Fungicide, and Rodenticide Act. Products making pesticide claims must submit required information to EPA to register under FIFRA and may be subject to labeling and use requirements.

Fertilization - Treatment method involving adding of nutrients, which could improve the success of desirable species; may be limited, depending on species/soil characteristics.

Flora - Plant life, especially all the plants found in a particular country, region, or time regarded as a group. Also, a systematic set of descriptions of all the plants of a particular place or time.

Foaming - Hot foam is a mechanical method that is effective on seedlings and annuals and can be applied under certain weather conditions, including wind and light rain.

Food chain - A hierarchical sequence of organisms, each of which feeds on the next, lower member of the sequence

Forage - Food for animals. In this document, term applies to both availability of plant material for wildlife and domestic livestock.

Formulation - A commercial preparation of a chemical including any inerts and/or contaminants.

Fungi - Molds, mildews, yeasts, mushrooms, and puffballs, a group of organisms that lack chlorophyll and therefore are not photosynthetic. They are usually non-mobile, filamentous, and multi-cellular.

Game fish - Species like trout, salmon, or bass, caught for sport. Many of them show more sensitivity to environmental change than non-game fish.

Grazing animals - Treatment method which requires matching the invasive plant species with the appropriate grazer for best success.

Groundwater Loading Effects of Agricultural Management Systems (GLEAMS) – A model which displays herbicide concentrations in streams under a variety of conditions.

Groundwater - The supply of fresh water found beneath the Earth's surface, usually in aquifers, which often supply wells and springs.

Habitat - The place where a population (e.g., human, animal, plant, microorganism) lives and its surroundings, both living and non-living.

Halftime or half-life - The time required for the concentration of the chemical to decrease by one-half.

Hand/Selective application - Herbicide treatment of individual plants through wicking, wiping, injecting stems, etc., with low likelihood of drift or delivery of herbicides away from treatment sites. This method ensures no herbicide directly contacts soil.

Hand-pulling/Grubbing - Treatment method which is labor-intensive but effective on single plants or on small, low-density infestations.

Hazard Quotient (HQ) - The Hazard Quotient is the amount of herbicide or additives to which an organism may be exposed over a specified period, divided by the estimated daily exposure level at which no adverse health effects are likely to occur. An HQ less than or equal to one indicates an extremely low level of risk; therefore, an HQ less than or equal to one is presumed to indicate a level of exposure below the threshold of concern for adverse health effects.

Hazard identification - The process of identifying the array of potential effects that an agent may induce in an exposed of humans or other organisms.

Herbaceous - A plant that does not develop persistent woody tissue above the ground (annual, biennial, or perennial.) Herbaceous vegetation includes grasses and grass-like vegetation, and broadleaved forbs.

Herbicide - A chemical preparation designed to kill plants, especially weeds, or to otherwise inhibit their growth.

Humus - Organic portion of the soil remaining after prolonged microbial decomposition.

Tribal and Treaty Rights - Native American treaty and other rights or interests recognized by treaties, statutes, laws, executive orders, or other government action, or federal court decisions.

Indian Tribe - Any American Indian or Alaska Native tribe, band, nation, pueblo, community, rancheria, colony, or group meeting the provisions of the Code of Federal Regulations Title 25, Section 83.7 (25 FR 83.7), or those recognized in statutes or treaties with the United States.

Indigenous - An indigenous species is any which were or are native or inherent to an area. (See also, native.)

Inerts - Anything other than the active ingredient in a pesticide product; not having pesticide properties.

Infested area - A contiguous area of land occupied by, in this case, invasive plant species. An infested area of land is defined by drawing a line around the actual perimeter of the infestation as defined by the canopy cover of the plants, excluding areas not infested. Generally, the smallest area of infestation mapped will be 1/10th (0.10) of an acre or 0.04 hectares.

Integrated Weed Management (IWM) - An interdisciplinary weed management approach for selecting methods for preventing, containing, and controlling noxious weeds in coordination with other resource management activities to achieve optimum management goals and objectives

Interdisciplinary Team (IDT) - A group of individuals with varying areas of specialty assembled to solve a problem or perform a task. The team is assembled out of recognition that no one scientific discipline is sufficiently broad enough to adequately analyze the problem and propose action.

Introduced species - An alien or exotic species that has been intentionally or unintentionally released into an area as a result of human activity. (See also exotic, invasive, and noxious.)

Introduction - “The intentional or unintentional escape, release, dissemination, or placement of a species into an ecosystem as a result of human activity” (Executive Order 13122, 2/3/99).

Invasive plant species - An alien plant species whose introduction does or is likely to cause economic or environmental harm or harm to human health (Executive Order 13122, 2/3/99) (See also exotic and introduced species)

Irreversible effect - Effect characterized by the inability of the body to partially or fully repair injury caused by a toxic agent.

Irritant - Non-corrosive material that causes a reversible inflammatory effect on living tissue by chemical action at the site of contact as a function of concentration or duration of exposure.

LC50 (Lethal Concentration50) - A calculated concentration of a chemical in air or water to which exposure for a specific length of time is expected to cause death in 50 percent of a defined experimental animal population.

LD50 (Lethal Dose50) - The dose of a chemical calculated to cause death in 50 percent of a defined experimental animal population over a specified observation period. The observation period is typically 14 days.

Label - All printed material attached to, or part of, the pesticide container.

Land allocation - Commitment of a given area of land or a resource to one or more specific uses (e.g. wilderness). In the Northwest Forest Plan, one of the seven allocations of Congressionally Withdrawn Areas, Late-Successional Reserves, Adaptive Management Areas, Managed Late-Successional Areas, Administratively Withdrawn Areas, Riparian Reserves, or Matrix.

Landscape - An area composed of interacting ecosystems that are repeated because of geology, land form, soils, climate, biota, and human influences throughout the area. Landscapes are generally of a size, shape, and pattern which is determined by interacting ecosystems.

Landscape Character - Particular attributes, qualities, and traits of a landscape that give it an image and make it identifiable or unique.

Landscape Setting - The context and environment in which a landscape is set; a landscape backdrop. It is the combination of land use, landform, and vegetation patterns that distinguish an area in appearance and character from other areas.

Leachate - Water that collects chemicals as it trickles through soil or other porous media containing the chemicals.

Leaching - The process by which chemicals on or in soil or other porous media are dissolved and carried away by water, or are moved into a lower layer of soil.

Level of Concern (LOC) - The concentration in media or some other estimate of exposure above which there may be effects.

Lichens - Complex thallophytic plants comprised of an alga and a fungus growing in symbiotic association on a solid surface (such as a rock.)

Littoral zone - (1) That portion of a body of fresh water extending from the shoreline lakeward to the limit of occupancy of rooted plants. (2) The strip of land along the shoreline between the high and low water levels.

Lowest-Observed-Adverse-Effect Level (LOAEL) - The lowest dose of a chemical in a study, or group of studies, that produces statistically or biologically significant increases in frequency or severity of adverse effects between the exposed and control populations.

Manual Control - The use of any non-mechanized approach to control or eliminate invasive plants (i.e. hand-pulling, grubbing)

Material Safety Data Sheet (MSDS) - A compilation of information required under the OSHA Communication Standard on the identity of hazardous chemicals, health and physical hazards, exposure limits, and precautions.

Mechanical Control - The use of any mechanized approach to control or eliminate invasive plants (i.e. mowing, weed whipping, hot foam.)

Microorganisms - A generic term for all organisms consisting only of a single cell, such as bacteria, viruses, protozoa and some fungi.

Minimum tool - Use of a weed treatment alternative that would accomplish management objectives and have the least impact on resources

Modification - A visual quality objective meaning human activities may dominate the characteristic landscape but must, at the same time, utilize naturally established form, line, color, and texture. It should appear as a natural occurrence when viewed in foreground or middleground.

Mollusks - Invertebrate animals (such as slugs, snails, clams, or squids) that have a soft, unsegmented body, usually enclosed in a calcareous shell; representatives found on National Forest System land include snails, slugs, and clams.

Monitoring - A process of collecting information to evaluate if objectives and anticipated or assumed results of a management plan are being realized or if implementation is proceeding as planned.

Morbidity - Rate of disease, injury or illness.

Mowing - Invasive plant treatment method which is limited to level/gently-sloping smooth-surface terrain. Treatment timing is critical, and must be conducted for several consecutive years.

National Environmental Policy Act (NEPA) - An Act passed in 1969 to declare a national policy that encourages productive and enjoyable harmony between humankind and the environment, promotes efforts that prevent or eliminate damage to the environment and biosphere, stimulates the health and welfare of humanity, enriches the understanding of the

ecological systems and natural resources important to the nation, and establishes a Council on Environmental Quality.

National Forest Management Act (NFMA) - A law passed in 1976 as an amendment to the Forest and Rangeland Renewable Resources Planning Act, requiring preparation of Forest Plans and the preparation of regulations to guide that development.

National Marine Fisheries Service (NMFS) - The federal agency that is the listing authority for marine mammals and anadromous fish under the ESA.

National Pollutant Discharge Elimination System (NPDES) - As authorized by the Clean Water Act, the National Pollutant Discharge Elimination System (NPDES) permit program controls water pollution by regulating point sources that discharge pollutants into waters of the United States. Point sources are discrete conveyances such as pipes or man-made ditches. Individual homes that are connected to a municipal system, use a septic system, or do not have a surface discharge do not need an NPDES permit; however, industrial, municipal, and other facilities must obtain permits if their discharges go directly to surface waters.

National Visitor Use Monitoring (NVUM) - A permanent, ongoing sampling system which measures national forest visitor demographics, experiences, preferences, and impressions. A stratified random sample is done for 25% of the National Forest system each year according to a national research protocol. NVUM responds to the need to better understand the use and importance of, and satisfaction with, national forest system recreation opportunities.

National Wilderness Preservation System (NWPS) - The Wilderness Act of 1964 established the national Wilderness Preservation System to ensure that certain federally owned areas in the United States would be preserved and protected in their natural condition. The Act defines a wilderness area, in part, as an area which generally appears to have been affected primarily by the forces of nature, with the imprint of man's work substantially unnoticeable. Areas included in the system are administered for the use and enjoyment of the American people in such manner as to leave them unimpaired for future use and enjoyment as wilderness.

Native species - With respect to a particular ecosystem, a species that, other than as a result of an introduction, historically occurred or currently occurs in that ecosystem (Executive Order 13122, 2/3/99).

Naturalized - Applied to a species that originally was imported from another country but that now behaves like a native in that it maintains itself without further human intervention and has invaded native populations.

Non-local native - This term has two meanings: (1) a population of a native plant species which does not occur naturally in the local ecosystem and/or (2) plant material of a native species that does not originate from genetically local sources.

Non-target species - Any plant or animal that is not the intended organism to be controlled by a pesticide treatment.

No-Observed-Adverse-Effect level (NOAEL) - Exposure level at which there are no statistically or biologically significant differences in the frequency or severity of any adverse effect in the exposed or control populations

No-Observed-Effect-Level (NOEL) - Exposure level at which there are no statistically or biological significant differences in the frequency or severity of any effect in the exposed or control populations.

Not Likely to Adversely Affect (NLAA) - Determinations are applied to those species that had very little habitat on National Forests in Region Six, were not in habitats susceptible to invasive plants, or were known to tolerate herbicide treatments without effects.

Noxious weed - “Any living stage (including but not limited to, seeds and reproductive parts) of any parasitic or other plant of a kind, or subdivision of a kind, which is of foreign origin, is new to or not widely prevalent in the United States, and can directly or indirectly injure crops, other useful plants, livestock, or poultry or other interests of agriculture, including irrigation, or navigation or the fish and wildlife resources of the United States or the public health” (Public Law 93-629, January 3, 1975, Federal Noxious Weed Act of 1974).

Outstandingly Remarkable Value (ORV) - A characteristic of rivers or sections of rivers in the national Wild and Scenic River System. In order for a river to be included in the system, it must possess at least one “outstandingly remarkable” value, such as scenic, recreational, geologic, fish, wildlife, historic, cultural, or other similar features. ORV’s are values or opportunities in a river corridor which are directly related to the river and which are rare, unique, or exemplary from a **regional or national perspective**.

Partial Retention - A visual quality objective which in general means human activities may be evident but must remain subordinate to the characteristic landscape.

Pathogen - A living organism, typically a bacteria or virus, that causes adverse effects in another organism.

Percolation - Downward flow or filtering of water through pores or spaces in rock or soil.

Perennial - A plant species having a life span of more than two years.

Periphyton - Microscopic plants and animals that are firmly attached to solid surfaces under water such as rocks, logs, pilings and other structures.

Persistence - Refers to the length of time a compound, once introduced into the environment, stays there.

Personal Protective Equipment (PPE) - Clothing and equipment worn by pesticide mixers, loaders and applicators and re-entry workers, hazmat emergency responders, workers cleaning up Superfund sites, et. al., which is worn to reduce their exposure to potentially hazardous chemicals and other pollutants.

Pest - An insect, rodent, nematode, fungus, weed or other form of terrestrial or aquatic plant or animal life that is classified as undesirable because it is injurious to health or the environment.

Pesticide - Any substance used for controlling, preventing, destroying, repelling, or mitigating any pest. Includes fungicides, herbicides, fumigants, insecticides, nematicides, rodenticides, desiccants, defoliants, plant growth regulators, etc.

Pesticide tolerance - The amount of pesticide residue allowed by law to remain in or on a harvested crop.

pH - The negative log of the hydrogen ion concentration. A high pH (greater than seven) is alkaline or basic and a low pH (less than seven) is acidic.

Population - A group of individuals of the same species in an area.

Population at Risk - A population subgroup that is more likely to be exposed to a chemical, or is more sensitive to the chemical, than is the general population.

Porosity - Degree to which soil, gravel, sediment, or rock is permeated with pores or cavities through which water or air can move.

Potable Water - Water that is considered safe for drinking and cooking.

Project Design Features (PDF) - A set of implementation Design Features/features applied to projects to ensure that the project is done according to environmental standards and adverse effects are within the scope of those predicted in this Environmental Impact Statement.

Proposed species - Any plant or animal species that is proposed by the Fish and Wildlife Service or NOAA Fisheries in a Federal Register notice to be listed as threatened or endangered.

Potential Vegetation Type (PVT) - The term Potential Vegetation Type is used to represent the combination of species that could occupy the site in the absence of disturbance.

Protozoa - Single-celled, microorganisms without cell walls containing visibly evident nuclei and organelles. Most protozoa are free-living although many are parasitic.

Recreational Rivers - A classification within the national Wild and Scenic River System. Recreational rivers are those rivers, or sections of rivers, that are readily accessible by road or railroad, that may have some development along their shorelines, and that may have undergone some impoundment or diversion in the past.

Reference Dose (RfD) - The RfD is a numerical estimate of a daily exposure to the human population, including sensitive subgroups such as children, that is not likely to cause harmful effects during a lifetime. RfDs are generally used for health effects that are thought to have a threshold or minimum dose for producing effects.

Registered Pesticides - Pesticide products which have been approved for the uses listed on the label.

Registration - Formal licensing with EPA of a new pesticide before it can be sold or distributed. Under the Federal Insecticide, Fungicide, and Rodenticide Act, EPA is responsible for registration (pre-market licensing) of pesticides on the basis of data demonstrating no unreasonable adverse effects on human health or the environment when applied according to approved label directions.

Restoration - Ecological restoration is the process of assisting the recovery and management of ecological integrity. Ecological integrity includes a critical range of variability in biodiversity, ecological processes and structures, regional and historical context, and sustainable cultural practices.

Retention - A visual quality objective which in general means human activities are not evident to the casual forest visitor.

Revegetation - The re-establishment of plants on a site - The term does not imply native or nonnative; does not imply that the site can ever support any other types of plants or species and is not at all concerned with how the site 'functions' as an ecosystem.

Riparian Area - A geographic area containing an aquatic ecosystem and adjacent upland areas that directly affect it.

Riparian Reserves - Areas along live and intermittent streams, wetlands, ponds, lakes, and unstable and potentially unstable areas where riparian-dependent resources receive primary emphasis. Riparian Reserves are important to the terrestrial ecosystem as well, serving as dispersal habitat for certain terrestrial species.

Risk Assessment - An analytic process that is firmly based on scientific considerations, but also requires judgments to be made when the available information is incomplete. These judgments inevitably draw on both scientific and policy considerations.

Risk - The chance of an adverse or undesirable effect, often measured as a percentage.

Risk assessment - The qualitative and quantitative evaluation performed in an effort to estimate the risk posed to human health and/or the environment by the presence or potential presence and/or use of specific chemical or biological agents.

Saturated zone - A subsurface area in which all pores and cracks are filled with water under pressure equal to or greater than that of the atmosphere.

Scenery Management - The art and science of arranging, planning, and designing landscape attributes relative to the appearance of places and expanses in outdoor settings.

Scenic - Of or relating to landscape scenery; pertaining to natural or natural-appearing scenery; constituting or affording pleasant views of natural landscape attributes or positive cultural elements.

Scenic Rivers - A classification within the national Wild and Scenic River System. Scenic rivers are those rivers, or sections of rivers, that are free of impoundments, with shorelines or watersheds still largely primitive and shorelines largely undeveloped, but accessible in places by roads.

Seen Area - The total landscape area observed based upon landform screening. Seen-areas may be divided into zones of immediate foreground, foreground, middleground, and background. Some landscapes are seldom seen by the public.

Sensitive species - Species identified by the Regional Forester for which population variability is a concern, as evidenced by significant current or predicted downward trend in population numbers or density; or significant current or predicted downward trends in habitat capability that would reduce a species existing distribution.

Species of Local Interest (SOLI) - Threatened, endangered and proposed species; Regional Forester's Sensitive species, management indicator species, and other rare or endemic species of concern.

Species - "A group of organisms, all of which have a high degree of physical and genetic similarity, generally interbreed only among themselves, and show persistent differences from members of allied groups of organisms." (Executive Order 13122, 2/3/99).

Spot application - Herbicide treatment involving use of a backpack sprayer or other means. Application is aimed at specific target species, with methods of prevention (such as barriers,) to control damage to non-target species.

Standards and guidelines - The rules and limits governing actions, as well as the principles specifying the environmental conditions or levels to be achieved and maintained

Sub-chronic exposure - An exposure duration that can last for different periods of time (5 to 90 days), with 90 days being the most common test duration for mammals. The sub-chronic study is usually performed in two species (rat and dog) by the route of intended use or exposure.

Sub-chronic toxicity - The ability of one or more substances to cause effects over periods from about 90 days but substantially less than the lifetime of the exposed organism. Sub-chronic toxicity only applies to relatively long-lived organisms such as mammals.

Submerged Aquatic Vegetation (SAV) - Vegetation that lives at or below the water surface; an important habitat for young fish and other aquatic organisms. In contrast to "emergent vegetation," which is growing out of or standing in water.

Substrate - With reference to enzymes, the chemical that the enzyme acts upon

Surface water - All water naturally open to the atmosphere (rivers, lakes, reservoirs, streams, impoundments, seas, estuaries, etc.) and all springs, wells, or other collectors which are directly influenced by surface water.

Surfactant - A surface active agent; usually an organic compound whose molecules contain a hydrophilic group at one end and a lipophilic group at the other. Promotes solubility of a chemical, or lathering, or reduces surface tension of a solution.

Synergistic effect - Situation in which the combined effects of exposure to two chemicals simultaneously is much greater than the sum of the effect of exposure to each chemical given alone.

Take - "The term 'take' means to harass, harm, pursue, hunt, shoot, wound, kill, trap, capture, or collect, or to attempt to engage in any such conduct." (Title 16, Chapter 35, Section 1532, Endangered Species Act of 1973)

Threatened species - Plant or animal species likely to become endangered throughout all, or a significant portion of, its range within the foreseeable future. A plant or animal identified and defined in accordance with the 1973 Endangered Species Act and published in the Federal Register.

Threshold - The maximum dose or concentration level of a chemical or biological agent that will not cause an effect in the organism.

Tolerances - Permissible residue levels for pesticides in raw agricultural produce and processed foods. Whenever a pesticide is registered for use on a food or a feed crop, a tolerance (or exemption from the tolerance requirement) must be established. EPA establishes the tolerance

levels, which are enforced by the Food and Drug Administration and the Department of Agriculture.

Toxicity - The inherent ability of an agent to affect living organisms adversely. Toxicity is the degree to which a substance or mixture of substances can harm humans or animals.

Toxicology - The study of the nature, effects, and detection of poisons in living organisms. Also, substances that are otherwise harmless but prove toxic under particular conditions. The basic assumption of toxicology is that there is a relationship among the dose (amount), the concentration at the affected site, and the resulting effects.

Treatment Area - An infested area where weeds have been treated or retreated by an acceptable method for the specific objective of controlling their spread or reducing their density.

U.S. Fish and Wildlife Service (US FWS) - The federal agency that is the listing authority for species other than marine mammals and anadromous fish under the ESA.

U.S. Forest Service (USDA FS or USFS) - The federal agency responsible for management of the Nation's National Forest system lands.

Variety Class - A particular level of visual variety or diversity of landscape character.

Viability - Ability of a wildlife or plant population to maintain sufficient size to persist over time in spite of normal fluctuations in numbers, usually expressed as a probability of maintaining a specific population for a specified period.

Viable Population - A wildlife or plant population that contains an adequate number of reproductive individuals appropriately distributed on the planning area to ensure the long-term existence of the species.

Viewshed - Total visible area from a single observer position, or the total visible area from multiple observer position. Viewsheds are accumulated seen-areas from highways, trails, campgrounds, towns, cities, or other viewer locations. Examples are corridor, feature, or basin viewsheds.

Visual Quality Objective - A desired level of excellence based on physical and sociological characteristics of an area. Refers to degree of acceptable alteration of the characteristic landscape.

Well-distributed - Distribution sufficient to permit normal biological function and species interactions, considering life history characteristics of the species and the habitats for which it is specifically adapted.

Wetland - An area that is regularly saturated by surface or ground water and subsequently is characterized by a prevalence of vegetation that is adapted for life in saturated soil conditions. Examples include swamps, bogs, fens, marshes, and estuaries.

Wild and Scenic River System - The Wild and Scenic Rivers Act of 1968 established a system of selected rivers in the United States, which possess outstandingly remarkable values, to be preserved in free-flowing condition. Within the national system of rivers, three classifications define the general character of designated rivers: Wild, Scenic, and Recreational.

Classifications reflect levels of development and natural conditions along a stretch of river. Classifications are used to help develop management goals for the river.

Wilderness - Areas designated by Congressional action under the 1964 Wilderness Act. Wilderness is defined as undeveloped federal land retaining its primeval character and influence without permanent improvements or human habitation. Wilderness areas are protected and managed to preserve their natural conditions, which generally appear to have been affected primarily by the forces of nature with the imprint of human activity substantially unnoticeable; have outstanding opportunities for solitude or for a primitive and confined type of recreation; include at least 5,000 acres, or are of sufficient size to make practical their preservation, enjoyment, and use in an unimpaired condition; and may contain features of scientific, educational, scenic, or historical value as well as ecological and geologic interest.

Wild Rivers - A classification within the national Wild and Scenic River System. Wild rivers are those rivers, or sections of rivers that are free of impoundments and generally inaccessible except by trail, with watersheds or shorelines essentially primitive and waters unpolluted.

4.7 References

- Alt, D., and D.W. Hyndman. 1978. Roadside Geology of Oregon. Mountain Press Publishing Co. Missoula, MT. 280p.
- Altman, B. 2000a. Conservation strategy for landbirds of the east-slope of the Cascade Mountains in Oregon and Washington. Version 1.0. June. Oregon-Washington Partners in Flight. 128pp.
- Altman, B. 2000b. Conservation strategy for landbirds in the northern Rocky Mountains of eastern Oregon and Washington. Version 1.0. May. Oregon-Washington Partners in Flight. 121pp.
- Altman, R. 2000. Conservation strategy for landbirds of the east-slope of the Cascade Mountains in Washington and Oregon. ver. 1.0. Prepared for Oregon-Washington Partners in Flight, 81+p.
- Andelman, S.J. and A. Stock. 1994. Management, research, and monitoring priorities for conservation of neotropical migratory landbirds that breed in Oregon state. Wash. Nat. Heritage Prog., Wash. Dept. Nat. Resources, Olympia.
- Anthony, R.G., and F.B. Isaacs. 1989. Characteristics of bald eagle nest sites in Oregon. *Journal Wildlife Management* 53: p.148-159
- Anthony, R.G., R.L. Knight, G.T. Allen, B. R. McClelland, and J.I. Hodges. 1982. Habitat use by nesting and roosting bald eagles in the Pacific Northwest. *Trans. North American Wildlife Natural Resources Conference*, 47:332-342p
- Archer, A. J. 2001. *Taeniatherum caput-medusae* [Online]. Available: <http://www.fs.fed.us/database/feis>. [Accessed Aug 11, 2003].
- Arthur, M.A., and Wang, Y. 1999. Soil nutrients and microbial biomass following weed control treatments in a Christmas tree plantation. *Soil Science Society of America Journal* 63(3): p.629-37.
- Asher, Jerry and C. Spurrier (1998). "The Spread of Invasive Weeds in Western Wildlands: A State of Biological Emergency." Report to the Governor's Idaho Weed Summit, Boise, ID. May 19, 1998.
- Atkinson, D., Abernathy, W., and Crisp, C. 1980. The effect of several herbicides on moss establishment in orchards. In *Proceedings of the 1980 British Crop Protection Conference*. British Crop Protection Society, London. pp. 297-302. British Crop Protection Society, London. pp. 297-302.
- ATSDR (Agency for Toxic Substances and Disease Registry). 2004. Guidance manual for the assessment of joint toxic action of chemical mixtures. Available from U.S. Department of Health and Human Services, Public Health Service, ATSDR, Division of Toxicology.
- Aubry, Keith B. and Jeffrey C. Lewis. 2003. Extirpation and reintroduction of fishers in Oregon: implications for their conservation in the Pacific states. USDA Forest Service, Pacific Northwest Research Station, 3625 93rd Avenue SW, Olympia, WA 98512, USA

- Aubry, K.B. and C. M. Raley. 2002. The Pileated woodpecker as a keystone habitat modifier in the Pacific Northwest. Pp. 257-274 in Proceedings of the symposium on the ecology and management of dead wood in western forests. 1999 November 2-4; Reno, NV. Gen. Tech. Rep. PSW-GTR-181. Albany, CA: U.S. Department of Agriculture, Forest Service, Pacific Southwest Research Station.
- Bais, H.P., R.Vepachedu, S. Gilroy, R.M. Callaway, J.M. Vivanco. 2003. Allelopathy and exotic plant invasion: from molecules and genes to species interactions. *Science*. 301: 1377-1380
- Baker, J.G. 1974. Patterns of plant invasions in North America. Pages 44-57 in H.A. Mooney and J.A. Drake, eds. *Ecology of biological invasions of North America and Hawaii*. Springer Verlag, New York.
- Bakke, D. 2003. Analysis of issues surrounding the use of spray adjuvants with herbicides. unpublished report by the Forest Service Pacific Southwest Regional Pesticide Use Specialist.
- Bakke, D. 2003. Human and ecological risk assessments of nonylphenol polyethoxylate-based (NPE) surfactants in Forest Service herbicide applications. Unpublished report by the Forest Service Pacific Southwest Regional Pesticide Use Specialist.
- Balcerkiewicz, S., and Rusińska, A. 1987. Expansion of bryophytes on areas treated with herbicides. *Symposium Biologica Hungarica* 35:285-293.
- Barnett, J.K. and J.A. Crawford. 1994. Pre-laying nutrition of sage grouse hens in Oregon. *Journal of Range Management* 47: 114-118.
- Barnhart R. A. 1986. Species Profiles: Life Histories and Environmental Requirements of Coastal Fishes and Invertebrates (Pacific Southwest)--Steelhead. U.S. Fish and Wildlife Service Biological Report. 82(11.60): 21.
- Barrows, C. W. 1996. Tamarisk control and common sense. In: Proceedings of the California Exotic Pest Plant Council; October 4-6; San Diego, CA.
- Bashkin, M., T.J. Stohlgren, Y. Otsuki, M. Lee, P. Evangelista, and J. Belnap. 2003. Soil characteristics and plant exotic species invasion in the Grand Staircase-Escalante national Monument, Utah, USA. *Applied soil Ecology*. 22(1) 67-77
- Batt, P.E. 1996. State of Idaho bull trout conservation plan. Office of the Governor. Boise, ID. 20pp.
- Bautista, S. 2005. [Personal communication]. Portland, OR U.S. Department of Agriculture, Forest Service, Pacific Northwest Region, Wildlife Biologist.
- Bautista, S. 2006. [Personal communication]. Portland, OR. U.S. Department of Agriculture, Forest Service, Pacific Northwest Region, Wildlife Biologist R-6 Invasive Plant Team. Also unpublished report: Analysis Methods and Issue Identification Regarding Effects to Wildlife for the Invasive Plant EIS

- Bautista, Shawna 2006. [Personal communication]. Portland, OR: U.S. Department of Agriculture, Forest Service, Pacific Northwest Region, Wildlife Biologist R-6 Invasive Plant Team.
- Baxter, C.V., C.A. Frissell, and F.R. Hauer. 1999. Geomorphology, Logging Roads, and the Distribution of Bull Trout Spawning in a Forested River Basin: Implications for Management and Conservation. *Transactions of the American Fisheries Society* 128:854-867.
- Baxter, J.S., and J.D. McPhail. 1999. The influence of redd site selection, groundwater upwelling, and over-winter incubation temperature on survival of bull trout (*Salvelinus confluentus*) from egg to alevin. *Canadian Journal of Zoology/Revue Canadien de Zoologie* 77:1233-1239.
- Bedunah, D.; Carpenter, J. 1989. Plant community response following spotted knapweed (*Centaurea maculosa*) control on three elk winter ranges in western Montana. 1998 Knapweed Symposium; Bozeman, MT.: Plant & Soil Department and Extension Service, Montana State University.
- Beedy, E.C. and W.J. Hamilton, III. 1999. Tricolored Blackbird (*Agelaius tricolor*). In A. Poole and F. Gill(eds.), *The Birds of North America*, No. 423. Philadelphia, PA: The Birds of North America, Inc
- Behnke R. J. 1992. Native Trout of Western North America. *American Fisheries Society Monograph*(6): 275.
- Belsky, A.J., and J.L. Gelbard. 2000. Livesotck grazing and weed invasions in the arid west. Oregon Natural Desert Association, Bend, OR.
- Benefield, C.B., DiTomaso, J.M., Kyser, G.B., Orloff, S.B., Churches, K.R. and others. 1999. Success of mowing to control yellow starthistle depends on timing and plant's branching form. *California Agriculture*, University of California, Division of Agriculture and Natural Resources 53:17-21.
- Berg, N. 2004. Assessment of Herbicide Best Management Practices: Status of Our Knowledge of BMP Effectiveness. Pacific Southwest Research Station, USDA Forest Service, Albany, CA.
- Berg, N. 2005. Assessment of Herbicide Best Management Practices: Status of Our Knowledge of BMP Effectiveness. Pacific Southwest Research Station, USDA Forest Service, Albany, CA.
- Berrill, M., S., Bertram, and B., Pauli. 1997. Effects of Pesticides on Amphibian Embryos and Larvae. *Herpetological Conservation* 1: p.233-45.
- Berrill, M., S., Bertram, L., McGillivray, M., Kolohon, and B., Pauli . 1994. Effects of low concentrations of forest-use pesticides on frog embryos and tadpoles. *Environmental Toxicology and Chemistry* 13(4): p.657-64.
- Beschta, R.L., R.E. Bilby, G.W. Brown, L.B. Holtby, and T.D. Hofstra. 1987. Stream temperature and aquatic habitat: fisheries and forestry interactions. Pages 191-232 in E.D. Salo and T.W. Cundy (eds). *Streamside Management Forestry and Fisheries*

- Interactions. Institute of Forest Resources, University of Washington, Seattle, Washington, Contribution No. 57.
- Blossey, B., C. Luke, J. Skinner and J. Taylor. 2001. Impact and management of purple loosestrife (*Lythrum salicaria*) in North America. *Biodiversity and Conservation* 10:1787-1807.
- Blus, L.J., C.S. Staley, C.J. Henny, G.W. Pendleton, T.H. Craig, E.H. Craig, and K.K. Halford. 1989. Effects of organophosphorus insecticides on sage grouse in southeastern Idaho. *Journal of Wildlife Management* 53: 1139-1146.
- Boag, T.D. 1987. Food habits of bull char, *Salvelinus confluentus*, and rainbow trout, *Salmo gairdneri*, coexisting in a foothills stream in northern Alberta. *Canadian Field-Naturalist* 101: 56-62.
- Bollag, J.M., S.Y. Liu. 1990. Biological Transformation processes of Pesticides. In *Pesticides in the Soil Environment*. H.H. Cheng. (ed.) Soil Science Society of America book series, no. 2. Madison, Wis., USA: Soil Science Society of America. p. 169-211
- Bond, C.E. 1992. Notes on the nomenclature and distribution of the bull trout and the effects of human activity on the species. Pages 1-4 in P.J. Howell, and D.V. Buchanan (eds). *Proceedings of the Gearhart Mountain Bull Trout Workshop, Oregon Chapter of the American Fisheries Society, Corvallis, Oregon*.
- Bond, T.E.T. 1976. *Polytrichum* spp. in an apple orchard on herbicide treated soil. *Proceedings of the Bristol Nature Society*. 35:69-72
- Boutin, C., Elmegaard, N., Kjaer, C. 2004. Toxicity testing of fifteen non-crop plant species with six herbicides in a greenhouse experiment: implications for risk assessment. *Ecotoxicology* 13:349-369.
- Braun, C.E. 1987. Current issues in sage grouse management. *Proceedings of the Western Association of Fish and Wildlife Agencies* 67: 134-144.
- Braun, C.E. 1998. Sage grouse declines in western North America: what are the problems? *Proceedings of the Western Association of Fish and Wildlife Agencies* 78: 139-156.
- Bridges, C.M. & Semlitsch, R.D. 2000. Variation in pesticide tolerance of tadpoles among and within species of Ranidae and patterns of amphibian decline. *Conservation Biology* 14: 1490-1499.
- Bridges, C.M. 1999a. The effects of a chemical stressor on amphibian larvae: individual, population and species level responses. Dissertation, University of Missouri, Columbia, Missouri, USA.
- Bridges, C.M. (1999b) Predator-prey interactions between two amphibian species: effects of insecticide exposure. *Aquatic Ecology* 33: 205-211.
- Brotherson, J. D.; Field, D. 1987. Tamarix: impacts of a successful weed. *Rangelands*. 9: 110-112.

- Brown, L.G. 1992. On the zoogeography and life history of Washington native char Dolly Varden (*Salvelinus malma*) and bull trout (*Salvelinus confluentus*). Washington Department of Wildlife, Fisheries Management Division Report, Olympia, Washington.
- Brown, M.L., Duncan, C.A., and Halstvedt, M.B. 2001. Cost and efficacy of spotted knapweed management and with integrated methods. *Techline* (May): p. 2-5.
- Buchanan, D.V., M.L Hanson, and R.M. Hooton. 1997. Status of Oregon's bull trout. Oregon Department of Fish and Wildlife, Portland, Oregon. 168 pp.
- Buchmann, S., and G. Nabhan, G.P. 1996. *The Forgotten Pollinators*. Washington. D.C. Island Press.
- Bulkin, Stephen 2006. [Personal communication]. Medford, OR: U.S. Department of Agriculture, Forest Service, RR-SISK Supervisor's Office, R6 Regional Office Invasive Plant EIS Implementation.
- Bull, E.L., and R.S. Holthausen. 1992. Habitat use and management of pileated woodpeckers in northeastern Oregon. *Journal of Wildlife Management* 57(2):335-345.
- Bull, E.L., and R.S. Holthausen. 1992. Habitat use and management of pileated woodpeckers in northeastern Oregon. *Journal of Wildlife Management* 57(2):335-345.
- Bulthuis, M. 2006. Range Specialist. Umatilla and Walllowa Whitman National Forests. Baker City, OR. Personal communication April – November, 2006.
- Burgner R. L., J. T. Light, L. Margolis, T. Okazaki, A. Tautz and S. Ito. 1992. Distribution and Origins of Steelhead Trout (*Oncorhynchus mykiss*) in Offshore Waters of the North Pacific Ocean. *International North Pacific Fish Commission Bulletin*.(51): 92.
- Burke, I.C., W.K. Lauenroth, and D.P. Coffin. 2005. Soil organic matter recovery in semiarid grasslands: implications for the conservation reserve program. *Ecological Applications*. 5(3): 793-801
- Burke, I.C., W.K. Lauenroth, M.A. Vinton, P.B. Hook, R.H. Kelly, H.E. Epstein, M.R. Aguiar, M.D. Robles, M.O. Aguilera, K.L. Murphy, and R.A. Gill. 1998. Plant-Soil interactions in temperate grasslands. *Biogeochemistry*. 42: 121-143
- Burkey, T.V. 1989. Extinction in nature reserves: the effect of fragmentation and the importance of migration between reserve fragments. *Oikos* 55:75-81.
- Burkey, T.V. 1995. Extinction rates in archipelagoes: Implications for populations in fragmented habitats. *Conservation Biology* 9: 527-541.
- Busby P. J., O. W. Johnson, T. C. Wainwright, F. W. Waknitz and R. S. Waples. 1993. Status Review for Oregon S Illinois River Winter Steelhead. U.S. Department of Commerce, NOAA Technical Memorandum.
- Busby P. J., T. C. Wainwright, G. J. Bryant, L. Leirheimer, R. S. Waples, F. W. Waknitz and I. V. Lagomarsino. 1996. Status Review of West Coast Steelhead from Washington, Idaho, Oregon, and California. NOAA Technical Memorandum. U.S. Department of Commerce: 281.

- Buskohl, C. 2008. [Personal communication]. Discussion and follow-up correspondence for risk of leaching on fractured bedrock for Wallowa Whitman NF. Zoned Forest Soil Scientist for Wallowa-Whitman NF and Umatilla NF. [Oct 24, 2008]
- Busse, M., G. Fiddler, and A. Ratcliff 2004. Ectomycorrhizal formation in herbicide treated soils of differing clay and organic matter content. *Water, Air and Soil Pollution* 152:23-24.
- Busse, M.D., A.W. Ratcliff, C.J. Shestak, and R.F. Powers. 2001. Glyphosate toxicity and the effects of long-term vegetation control on soil microbial communities. *Soil Biology and Biochemistry*. 33: 1777-1789
- Byrne, M. W. 2002. Habitat use by female Greater Sage Grouse in relation to fire at Hart Mountain National Antelope Refuge, Oregon. M.S. thesis. Oregon State University, Corvallis, Oregon. 50 pages. Calabrese, E.J. and L.A. Baldwin. 1993. *Performing Ecological Risk Assessments*. Lewis Publishers. Ann Arbor, Michigan. 257pp
- Callaway, R., G.C. Thelen, A. Rodriguez, and W.E. Holben. 2004. Soil biota and exotic plant invasion. *Nature*. 427: 431-433
- Callaway, R.M., T.H. DeLuca, and W.M. Belliveau. 1999. Biological-control herbivores may increase competitive ability of the noxious weed *Centaurea maculosa*. *Ecology* 80:1196-1201.
- Carroll, R.N. 2001. Carnivores as Focal Species for Conservation Planning in the Rocky Mountain Region. *Ecological Applications*, Vol. 11, No. 4. pp. 961-980
- Cavender, T.M. 1978. Taxonomy and distribution of the bull trout, *Salvelinus confluentus*, from the American northwest. *California Fish and Game* 64:139-174.
- CDFG (California Department of Fish and Game). 1995. Letter to M. Schiewe for the ESA Administrative Record for West Coast Steelhead, Dated 30 March 1995, 10 P. Plus Attachments. Available from Environmental and Technical Services Division, National Marine Fisheries Service, 525 Ne Oregon Street, Suite 500, Portland, Oregon 97232.
- Chamberlain, T. W., R. D. Harr, and F. H. Everest. 1991. Timber harvesting, silviculture and watershed processes. Pages 181-205 in W. R. Meehan (ed). *Influences of forest and rangeland management on salmonid fishes and their habitats*. American Fisheries Society Special Publication 19.
- Chapman D. W. 1986. Salmon and Steelhead Abundance in the Columbia River in the Nineteenth Century. *Transactions of the American Fisheries Society*. 115: 662-670.
- Chew, F. S. 1981. Coexistence and local extinction in two *Pieris* butterflies. *American Naturalist*. 118: 655-672.
- Chornesky, E.A.; A.M. Bartuska; G.H. Aplet; K.O. Britton; J. Cummings-Carlson; F.W. Davis; J. Eskow; D.R. Gordon; K.W. Gottschalk; and R.A. Haack. 2005. Science priorities for reducing the threat of invasive species to sustainable forestry. *BioScience* 55(4): 335-348.

- Choudhury, H.; J. Cogliano; R. Hertzberg; D. Mukerjee; G. Rice; L. Teuschler; E. Doyle; and R. Schoeny. 2000. Supplementary guidance for conducting Health Risk Assessment of chemical mixtures. Risk Assessment Forum, U.S. Environmental Protection Agency, Washington, D.C. 143 pp. + appendices.
- Cichosz T., D. Saul, A. Davidson, W. Warren, D. Rollins, J. Willey, T. Tate, T. Papanicolaou and S. Juul. 2001. Clearwater Subbasin Summary. Draft Submitted to the Northwest Power Planning Council. 477.
- Cipollini, M.L., B.G. Drake, and D. Whigham. 1993. Effects of elevated CO₂ on growth and carbon/nutrient balance in the deciduous wood shrub *Lindera benzoin* (L.) Blume (Lauraceae). *Oecologia* 96: 339-346.
- Climate Change Science Program. 2008. Preliminary review of adaptation options for climate-sensitive ecosystems and resources. A Report by the U.S. Climate Change Science Program and the Subcommittee on Global Change Research [Julius, S.H.; J.M. West (eds.), J.S. Baron, B. Griffith, L.A. Joyce, P. Kareiva, B.D. Keller, M.A. Palmer, C.H. Peterson, and J.M. Scot (Authors)]. U.S. Environmental Protection Agency, Washington, D.C. 873 pp.
- Cole, E.C., W. C. McComb, M. Newton, , C.L. Chambers and J.P. Leeming. 1997. Response of amphibians to clearcutting, burning, and glyphosate application in the Oregon Coast Range. *Journal of Wildlife Management*. 61:p.656-64.
- Cole, E.C., W. C. McComb, M. Newton, J.P. Leeming, and C.L. Chambers. 1998. Response of small mammals to clearcutting, burning, and glyphosate application in the Oregon Coast Range. *Journal of Wildlife Management*. 62(4): p. 1207-16.
- Colony Collapse Disorder Steering Committee. 2007. Colony collapse disorder action plan. USDA Agricultural Research Service. Downloaded on June 16, 2008 from: http://www.ars.usda.gov/is/br/ccd/ccd_actionplan.pdf.
- Colorado Weed Management Associations (CWMA) 2007. Diffuse knapweed. http://www.cwma.org/nx_plants/diff.htm
- Connelly J.W., H.W. Browsers, R.J. Gates. 1988. Seasonal movements of sage grouse in southeastern Idaho. *J. Wildl. Manage.* 52:116–122.
- Connelly, J. W., JR. 1982. An ecological study of sage grouse in southeastern Idaho. Dissertation, Washington State University, Pullman, USA.
- Connelly, J.W. and O.D. Markham. 1983. Movements and radio nuclide concentrations of sage grouse in southeastern Idaho. *Journal of Wildlife Management* 47:169-177.
- Connelly, J.W., M. A. Schroeder, A. R. Sands, C. E. Braun. 2000. Guidelines to manage sage grouse populations and their habitats. *Wildlife Society Bulletin* 28(4): 967-985.
- Connelly, J.W.; M.A. Schroeder; A.R. Sands; and C.E. Braun. 1991. Sage grouse use of nest sites in southeastern Idaho. *Journal of Wildlife Management* 55: 521-524.
- Copeland, J.P. 1996. Biology of the wolverine in central Idaho. M.S. Thesis, University of Idaho, 138 pp.

- Copeland, J.P. 1996. Biology of the Wolverine in Central Idaho. Masters Thesis. 152 pp.
- Corkran, C. C., and C. R. Thoms. 2006. Amphibians of Oregon, Washington, and British Columbia. Lone Pine Publishing, Auburn, WA.
- Cowman, D.F., and L.E. Mazanti. 2000. Ecotoxicology of “new generation” pesticides to amphibians. Pp. 233-268 In D.W. Sparling, G. Linder, and C.A. Bishop, eds. Ecotoxicology of Amphibians and Reptiles. Society of Environmental Toxicology and Chemistry. Pensacola, Florida. 904pp.
- Cox, C. 2000. Tricylorpyr Fact Sheet. Journal of Pesticide Reform 20:12-19.
http://72.14.253.104/search?q=cache:JGuFu_ZIEO4J:www.pesticide.org/triclopyr.pdf+herbicide+impacts+to+lichens,+bryophytes,+and+fungi&hl=en&gl=us&ct=clnk&cd=13. Accessed 11/2006.
- Cox-Foster, D.L.; S. Conlan; E.C. Holmes; G. Palacios; J.D. Evans, et al. 2007. A Metagenomic survey of microbes in honey bee colony collapse disorder. Science 318: 283-287.
- Craig, S.D., and R.C. Wissmar. 1993. Habitat conditions influencing a remnant bull trout spawning population, Gold Creek, Washington (draft report). Fisheries Research Institute, University of Washington. Seattle, Washington.
- Cramer S. P. 1995. The Status of Steelhead Populations in California in Regards to the Endangered Species Act. Document Prepared for Association of California Water Agencies. Environmental and Technical Services Division, National Marine Fisheries Service.
- Crawford, J.A. 1982. History of Sage Grouse in Oregon. Oregon Wildlife. 37(3):3-6.
- Crawford, J.A., and R.S. Lutz. 1985. Sage grouse population trends in Oregon, 1941-1983. Murrelet 66 (3): 69-74.
- Crawford, J.A., R.A. Olson, N.E. West, J.C. Mosley, M.A. Schroeder; T.D. Whitson; R.F. Miller; M.A. Gregg; and C.S. Boyd. 2004. Ecology and management of sage-grouse and sage-grouse habitat. Journal of Range Management 57: 2-19.
- Csuti, B., A. J. Kimerling, T. A. O’Neil, M. M. Shaughnessy, E. P. Gaines, and J. C. Hak 2001. Atlas of Oregon wildlife: distribution, habitat, and natural history. Oregon State University Press, Corvallis, OR. 492 pp.
- Curtis, P.S., B.G. Drake, and D.F. Whigham. 1989. Nitrogen and carbon dynamics in C3 and C4 estuarine marsh plants grown under elevated CO2 in situ. Oecologia 78: 297-301.
- D’Antonio, C.M., N.E. Jackson, C.C. Horvitz, and Rob Hedberg. 2004. Invasive plants in wildland ecosystems: merging the study of invasion processes and management needs. Front Ecol Environ 2(10): 513-521
- Dabbert, C.B., R.B. Mitchell, and D. Oberheu. 1997. Northern bobwhite egg hatchability and chick immunocompetence following field application of clopyralid. Bulletin of Environmental Contamination and Toxicology. 58: 801-806
- D’Antonio, C. M.; Vitousek, P. 1992. Biological invasions by exotic grasses, the grass/fire cycle, and global change. Annual Review of Ecological Systems. 23: 63-87.

- Dauble D. D., R. L. Johnson and A. P. Garcia. 1999. Fall Chinook Salmon Spawning in the Tailraces of Lower Snake River Hydroelectric Projects. *Transaction of the American Fisheries Society*. 128(4): 672-679.
- Dawson, L. 2007. Personal Communication. Discussion on current locations of weed infestations. Wallowa Mountains District Botany and Invasive Plants Staff/Manager.
- Dent and Robben. 2000. Oregon Department of Forestry: aerial pesticide applicaton monitoring final report. Oregon Department of Forestry, Salem OR.
http://oregon.gov/ODF/PRIVATE_FORESTS/docs/fp/ChemAppFinal.pdf
- DiTomaso, J. 2001. Yellow starthistle management possible with planning and integrated approach. *Techline* May: 6-8.
- DiTomaso, J.M, G.B.Kyser and M.J. Pitcarin. 2006. Yellow Starthistle Management Guide. California Invasive Plant Council. 74 pp.
- DiTomaso, J.M. 2000. Invasive weeds in rangelands: Species, impacts and management. *Weed Science* 48:255-265.
- Donald, D.B., and D.J. Alger. 1993. Geographic distribution, species displacement, and niche overlap for lake trout and bull trout in mountain lakes. *Canadian Journal of Zoology/Revue Canadien de Zoologie* 71:238-247.
- Dremann, C.C. and M. Shaw. 2002. Releasing the Native Seedbank: An Innovative Approach to Restoring a Coastal California Ecosystem *Ecological Restoration* 20: 103-107.
- Drewek, J. 1970. Population characteristics and behavior of introduced bighorn sheep in Owyhee County, Idaho. M.S. thesis. Univ. Idaho, Moscow. 46 p.
- Dudley, D. R. 2000. Wicked weed of the west. *California Wild*. 53:32-35.
- Dudley, T. L. 2000a. Noxious wildland weeds of California: *Arundo Donax*. In: Bossard, C. C.; Randall, J. M.; Hoshousky, M. C., eds. *Invasive plants of California wildlands*. Berkeley, CA: Univ. of California Press: 53-58.
- Dukes, J.S. and H.A. Mooney. 1999. Does global change increase the success of biological invaders? *TREE* 14(4): 135-139.
- Duncan, C. A., and J. K. Clark. eds. 2005. *Invasive Plants of Range and Wildlands and their Environmental, Economic, and Societal Impacts*. Lawrence, KS: Weed Science Society of America. In press.
- Dunham, J.B., and B.E. Rieman. 1999. Metapopulation structure of bull trout: Influences of physical, biotic, and geometrical landscape characteristics. *Ecological Applications* 9:642-655.
- Ehrlich, P.R., Dobkin, D. S., D.E. Wheye. 1988. *The Birders Handbook, A field guide to the natural history of North American birds*. Simon and Schuster Inc. New York, NY. 785p
- Elliot, W.J., D.E. Hall and S.R. Graves. 1999. Predicting sedimentation from forest roads. *Journal of Forestry*. 97(8): 23-29

- Enyeart, G. 1956. Responses of sage grouse to grass reseeding in the Pines area, Garfield County, Utah. Thesis, Utah State Agricultural College, Logan, USA.
- Erickson, V. 2006. [Personal communication] Pendleton, OR. U. S. Department of Agriculture, Forest Service, Umatilla National Forest, Genetecist
- Erickson, V., J. Wood, and S. Riley. 2003. Guidelines for Revegetation of Invasive Weed Sites and Other Disturbed Areas on National Forests and Grasslands in the Pacific Northwest. http://www.fs.fed.us/gpnf/04projects/pinshotprojects/documents/1-gp_nsa_deis.pdf (accessed 12-2006)
- Erickson, Vickie J., et al. 2003. Guidelines for Revegetation of Invasive Weed Sites and Other Disturbed Areas on National Forests and Grasslands in the Pacific Northwest.
- Everest F. H. 1973. Ecology and Management of Summer Steelhead in the Rogue River. Oregon State Game Commission, Fishery Research Report.(7): 48.
- Executive Order 13112 1999a. <http://www.fs.fed.us/r6/weeds/execorder.htm>. - accessed 11/06.
- Fagerstone, K.A., H.P. Tietjen, and G.K. LaVoie. 1977. Effects of range treatment with 2,4-D on prairie dog diet. *Journal of Range Management* 30(1): 57-60.
- Finch D. M, Stangel P. W 1993. Status and Management of Neotropical Migratory Birds. U.S. Department of Agriculture, Forest Service General Technical Report RM-229.
- Fischer, R.A.; K.P. Reese; and J.W. Connelly. 1996. Influence of vegetal moisture content and nest fate on timing of female sage grouse migration. *Condor* 98: 868-872.
- Flatter, B. 1998. Life history and population status of migratory bull trout (*Salvelinus confluentus*) in Arrowrock Reservoir, Idaho. Prepared for U.S. Bureau of Reclamation by Idaho Department of Fish and Game, Nampa, Idaho.
- Fletcher, J.S., J.E. Nellessen, and T.G. Pflieger. 1994. Literature review and evaluation of the EPA food-chain (Kenaga) nomogram, and instrument for estimating pesticide residue on plants. *Environ. Toxicol. And Chemistry* 13(9):p.1383-91.
- Fraley, J.J., and B.B. Shepard. 1989. Life history, ecology and population status of migratory bull trout (*Salvelinus confluentus*) in the Flathead Lake and river system, Montana. *Northwest Science* 63:133-143.
- Frissell, C.A. 1997. A spatial approach to species viability: Conservation of fishes in the Columbia River Basin. Biological Station Open File Report Number 101-97. Flathead Lake Biological Station, University of Montana, Polson, MT.
- Fuller, T.C., and Barbe, G.D. 1985. The Bradley Method of Eliminating Exotic Plants from Natural Reserves.:2p.
- Fulton L. A. 1968. Spawning Areas and Abundance of Chinook Salmon, *Oncorhynchus tshawytscha*, in the Columbia River Basin--Past and Present. U.S. Fish and Wildlife Service Special Science Report.
- Furniss, M.J., T.D. Roelofs, and C.S. Yee. 1991. Road construction and maintenance. *American Fisheries Society Special Publication* 19:297-323.

- Geist, V. 1971. Mountain sheep: A Study in Behavior and Evolution. University of Chicago Press, Chicago
- Gelbard, J.L. and J.Belnap. 2003. Roads as conduits for exotic plant invasions in a semiarid landscape. *Conservation Biology*. 17(2): 420-432
- Germaine, S. S.; Rosenstock, S. S.; Schweinsburg, R. E; Richardson, W. S. August 1998. Relationships among breeding birds, habitat, and residential development in greater Tucson, Arizona. *Ecological Applications*. 8(3): 680-691.
- Gilbert C. H. 1912. Age at Maturity of Pacific Coast Salmon of the Genus *Oncorhynchus*. *Bulletin of the U.S. Fisheries Commission*.(32): 57-70.
- Gilpin, M., University of California. 1997. Bull trout connectivity on the Clark Fork River, letter to Shelly Spalding, Montana Department of Fish, Wildlife and Parks, Helena, Montana. 5 pp
- Goetz, F. 1989. Biology of the bull trout, *Salvelinus confluentus*, literature review. U.S. Department of Agriculture, U.S. Forest Service, Willamette National Forest, Eugene, Oregon.
- Goodwin, K., R. Sheley, and J. Clark. 2002. Integrated Noxious Weed management after Wildfires. Department of Land Resources and Environmental Sciences, Montana State University Extension Service. 26p.
- Goodwin, K.M., and Sheley, R.L. 2001. What To Do When Fires Fuel Weeds: A step-by-stepguide for managing invasive plants after a wildfire. *Rangelands* 23(6): p.15-21
- Graham, R. T. and T. B. Jain. 1998. Silviculture's role in managing boreal forests. *Conservation Ecology* [online] 2(2): 8. Available from the Internet. URL: <http://www.consecol.org/vol2/iss2/art8/>
- Grinnell, J., J. S. Dixon, and J. M. Linsdale. 1937. Fur-bearing mammals of California. 2 Vols. Univ. California Press, Berkeley. 777pp.
- Hall, R.J. 1980. Effects of environmental contaminants on reptiles: a review. USDI Fish and Wildlife Service, Special Scientific Report, Wildlife No. 228. Washington, D.C. 12pp.
- Hall, R.J. and D.R. Clark, Jr. 1982. Responses of the iguanid lizard *Anolis carolinensis* to four organophosphorus pesticides. *Environmental Pollution (Series A)* 28: 45-52.
- Hall, R.J. and P.F.P. Henry. 1992. Review: Assessing effects of pesticides on amphibians and reptiles: status and needs. *Herpetological J.* 2: 65-71.
- Hanf J.M., P.A. Schmidt, E.B. Groshend. 1994. Sage grouse in the high desert of Central Oregon: results of a study, 1988–1993. U.S.D.I.-Bur. Land Manage., Series P-SG-01, Prineville, Ore.
- Hayes, M.P. & Jennings, M.R. 1986. Decline of ranid frog species in western North America: are bullfrogs (*Rana catesbeiana*) responsible? *Journal of Herpetology* 20: 490–509.
- Hayes, Marc, J.D. Engler, R.D. Haycock, D.H. Knopp, W.P. Leonard, K.R. McAllister, and L.L. Todd. 1997. Status of the Oregon spotted frog (*Rana pretiosa*) across its geographical

- range. In *The Spotted Frog of Oregon*. Oregon Chapter of The Wildlife Society, Corvallis, OR. August 15.
- Healey M. C. 1983. Coastwide Distribution and Ocean Migration Patterns of Stream- and Ocean Type Chinook Salmon, *Oncorhynchus tshawytscha*. *Canadian Field Naturalist*.(97): 427-433.
- Healey M. C. 1986. Optimum Size and Age at Maturity in Pacific Salmon and Effects of Size Selective Fisheries. *Canadian Special Publication on Fisheries and Aquatic Sciences*.(89): 39-52.
- Healey M. C. 1991. Life History of Chinook Salmon (*Oncorhynchus tshawytscha*). In: C. Groot and L. Margolis (Ed). *Pacific Salmon Life Histories*. Vancouver, British Columbia, University of British Columbia Press.
- Hellamann, J.J.; J.E. Byers; B.G. Bierwagen; and J.S. Dukes. 2008. Five potential consequences of climate change for invasive species. *Conservation Biology* 22(3): 534-543.
- Henjum, M.G., J.R. Karr, D.L. Bottom, D.A. Perry, J.C. Bednarz, S.G. Wright, S.A. Beckwitt, and E. Beckwitt. 1994. Interim protection for late-successional forests, fisheries, and watersheds. National forests east of the Cascade Crest, Oregon, and Washington. A report to the Congress and President of the United States Eastside Forests Scientific Society Panel. American Fisheries Society, American Ornithologists Union Incorporated, The Ecological Society of America, Society for Conservation Biology, The Wildlife Society. *The Wildlife Society Technical Review* 94-2.
- Higby, L. W. 1969. A summary of the Longs Creek sagebrush control project. *Proceedings, Biennial Western States Sage Grouse Workshop* 6:164–168.
- Hoelscher, B., and T.C. Bjornn. 1989. Habitat, density and potential production of trout and char in Pend Oreille Lake tributaries. Project F-71-R-10, Subproject III, Job No. 8. Idaho Department of Fish and Game, Boise, Idaho.
- Hooper, D.U. and P.M. Vitousek. 1998. Effects of plant composition and diversity on nutrient cycling. *Ecological Monographs*. 68(1): 121-149
- Hornocker, M. and J. Hash. 1981. Ecology of the wolverine in Northwestern Montana. *Canadian Journal of Zoology* 59:1286-1301.
- Horton, J.S. 1977. The development and perpetuation of the permanent tamarisk type in the phreatophyte zone of the southwest. *Gen. Tech. Rep. 43*. U.S. Department of Agriculture, Forest Service, Rocky Mountain Forestry and Range Experiment Station. p. 124-127.
- Howell, P.J., and D.V. Buchanan. 1992. *Proceedings of the Gearhart Mountain Bull Trout Workshop*. Oregon Chapter of the American Fisheries Society, Corvallis, Oregon. <http://www.nwfsc.noaa.gov/publications/techmemos/index.cfm>
- Huenneke, L.F., S.P. Hamburg, R. Koide, H.A. Mooney, P.M. Vitousek. 1990. Effects of soil resources on plant invasion and community structure in Californian serpentine grassland. *Ecology* 71:478 491.

- Huibregtse, D. 2007. Rangeland Management Specialist Bureau of Land Management ,Coeur d'Alene District, Cottonwood Field Office, Cottonwood, Idaho. Personal communication
- IDFG (Idaho Department of Fish and Game). 1994. Documents Submitted to the Esa Administrative Record for West Coast Steelhead by E. Leitzinger, 18 October 1994. Available from Environmental and Technical Services Division, National Marine Fisheries Service. Idaho Department of Fish and Game in litt., 1995; M
- Irving J.S. and T. Bjornn. 1991. A Forecast of Abundance of Snake River Fall Chinook Salmon. Prepared for U.S. Fish & Wildlife Service. Available from Idaho Cooperative Fishery Research Unit.
- Jakle, M. D.; Gatz, T. A. 1985. Herpetofaunal use of four habitats of the middle Gila River drainage, Arizona. Paper Presented at the North American Riparian Conference; April 16-18, 1985; Tucson, AZ.
- Jakober, M. 1995. Autumn and winter movement and habitat use of resident bull trout and west slope cutthroat trout in Montana. M.S. Thesis, Montana State University, Bozeman, Montana.
- Janzen, D. 1984. Dispersal of Small Seeds by Big Herbivores: Foliage is the Fruit
- Johnson, C.G., and Swanson, D.K. 2005. Bunchgrass plant communities of the Blue and Ochoco Mountains: a guide for managers. USDA Forest Service, Pacific Northwest Research Station, PNW-GTR-641, 119 pp.
- Johnson, D. H., and T. A. O'Neil, Managing Directors. 2001. Wildlife-habitat relationships in Oregon and Washington. Oregon State University Press, Corvallis, OR. 736 pp.
- Johnson, D.R., and Hansen, R.M. 1969. Effects of range treatment with 2,4-D on rodent populations. *Journal of Wildlife Management* 33: 1125-132.
- Johnson, H.B., H.W. Polley, and H.S. Mayeux. 1993. Increasing CO₂ and plant-plant interactions: effects on natural vegetation. *Vegetatio* 104/105: 157-170.
- Johnson, K.H.; Olson, R.A.; Whitson, T.D.; Swearingen, R.J.; Jurz, G. L. 1994. Ecological implications of Russian knapweed infestation: small mammal and habitat associations. In: *Proceedings of the Western Society for Weed Science*. [Place of publication unknown.] 47: 98-101.
- Johnson, M. L. 1995. Reptiles in the state of Washington (1954). *Northwest Fauna* 3:5-80. The Society of Northwestern Vertebrate Biology, Olympia, WA.
- Joyce, L.A.; G.M. Blate; J.S. Littell; S.G. McNulty; C.I. Millar; S.C. Moser; R.P. Neilson; K. O'Halloran; and D.L. Peterson. 2008. National Forests. Pp. 3-1 to 3-127 In Preliminary review of adaptation options for climate-sensitive ecosystems and resources. A Report by the U.S. Climate Change Science Program and the Subcommittee on Global Change Research [Julius, S.H.; J.M. West (eds.), J.S. Baron, B. Griffith, L.A. Joyce, P. Kareiva, B.D. Keller, M.A. Palmer, C.H. Peterson, and J.M. Scot (Authors)]. U.S. Environmental Protection Agency, Washington, D.C. 873 pp.

- Kaminski, T., and J. Hansen. 1984. Wolves of central Idaho. Mont. Coop. Wildl. Res. Unit. U.S. Fish and Wildl. Sev., Missoula, MT 196p.
- Kearns, C., D. Inouye, and N. Waser. 1998. Endangered Mutualisms: The Conservation of Plant-Pollinator Interactions. Annual Review of Ecology and Systematics, Vol. 29: 83-112 [http://links.jstor.org/sici?sici=0066-4162\(1998\)29%3C83%3AEMTCOP%3E2.0.CO%3B2-6](http://links.jstor.org/sici?sici=0066-4162(1998)29%3C83%3AEMTCOP%3E2.0.CO%3B2-6) (accessed 12-2006)
- Kenaga, E.E. 1973. Factors to be considered in the evaluation of the toxicity of pesticides to birds in their environment. In: Environmental Quality and Safety: Global Aspects of Chemistry, Toxicology and Technology as Applied to the Environment. Vol. II. New York, NY: George Thieme Publishers, Stuttgart Academic Press, Inc.
- Kesner W. D. and R. A. Barnhart. 1972. Characteristics of the Fall-Run Steelhead Trout (*Salmo Gairdneri* *Gairdneri*) of the Klamath River System with Emphasis on the Half-Pounder. California Fish and Game. 58(3): 204-220.
- Kiviat, E. 1996. Short communications: American goldfinch nests in purple loosestrife. Wilson Bulletin. 108(1): 182-186.
- Lacey J.R., R. Wallender, and K. Olson-Rutz. 1992. Recovery, germinability, and viability of leafy spurge (*Euphorbia esula*) seeds ingested by sheep and goats. Weed Technol. 6:559-602.
- Lacey, C.A, Lacey, J.R, Fay, P.K., Story, J.M., and Zamora, D.L. 1997. Controlling knapweed on Montana rangeland. Montana State University Extension Service Bulletin, Circular 311p.
- Lacey, J. R., Olson, B. E. 1991. Environmental and economic impacts of noxious range weeds. In: James, L. F.; Evans, J. O.; Ralphs, M. H.; Child, R. D., eds. Noxious range weeds. Boulder, CO: Westview Press
- Lacey, J.R., C.B. Marlow and J.R. Lane. 1989. Influence of spotted knapweed (*Centaurea maculosa*) on surface water runoff and sediment yield. Weed Technology 3: 627-631.
- Lacey, J.R., Marlow, C.B., and Lane, J.R. 1989. Influence of spotted knapweed (*Centaurea maculosa*) on surface runoff and sediment yield. Weed Technology 3:627-31.
- Laufman, J. 2007. Wallowa-Whitman National Forest Invasive Plants Treatment Botany Report.
- Leathe, S.A., and P. Graham. 1982. Flathead Lake fish food habits study. Environmental Protection Agency, through Steering Committee for the Flathead River Basin Environmental Impact Study. Contract R008224-01-4 to Montana Department of Fish, Wildlife and Parks
- Lee D. S., C. R. Gilbert, C. H. Hocutt, R. E. Jenkins, D. E. McAllister, J. R. Stauffer Jr. 1980. Atlas of North American freshwater fishes. North Carolina State Museum of Natural History, Raleigh. 854 p.
- Lefcort, H., R. A. Meguire, L. H. Wilson, W. F. Ettinger. 1998. Heavy metals alter the survival, growth, metamorphosis, and antipredatory behavior of Columbia spotted frog (*Rana*

- luteiventris) tadpoles. *Archives of Environmental Contamination and Toxicology* 15:447–456.
- LeJeune, K.D., Seastedt, T.R. 2001. *Centaurea* species: the forb that won the west. *Conserv Biol.* 15:1568-1574
- Levine, J.M. and C.M. D'Antonio. 1999. Elton revisited: a review of evidence linking diversity and invisibility. *Oikos.* 87: 15-26
- Levine, J.M. Vila, C. D'Antonio, J. Duke, K. Grigulis and S. Lavorel. 2003. Review Paper
- Light, J., L. Herger, and M. Robinson. 1996. Upper Klamath basin bull trout conservation strategy, a conceptual framework for recovery. Part one. The Klamath Basin Bull Trout Working Group
- Lor, S.K. 1999. Habitat use and population status of marsh birds in western New York. Ithaca, NY: Department of Natural Resources, Cornell University. 135 p. Thesis.
- Mabb, L.P. 1989. Uptake and effects of herbicides on the lawn moss *Rhytidiadelphus squarrosus*. Ph.D. thesis, University of London, London. Found in Newmaster et al 1999.
- Mack R.N. and J.N. Thompson. 1982. Evolution in steppe with few large, hooved mammals. *American Midland Naturalist* 119: 757-773.
- Mack, R. N. 1981. Invasion of *Bromus tectorum* L. into western North America: an ecological chronicle. *Agro-Ecosystems.* 7: 145-165.
- Mallet J. 1974. Inventory of Salmon and Steelhead Resources, Habitats, Use and Demands. Performance Report. Project F-58-R-1. Idaho Department of Fish and Game. Boise. Idaho.
- Mann, R.M. and J.R. Bidwell. 2001. The acute toxicity of agricultural surfactants to the tadpoles of four Australian and two exotic frogs. *Environmental Pollution* 114: 195-205.
- Mann, R.M. and J.R. Bidwell. 2000. Application of the FETAX protocol to assess the developmental toxicity of nonylphenol ethoxylate to *Xenopus laevis* and two Australian frogs. *Aquatic Toxicology* 51: 19-29.
- Marco, A., C. Quilchano, and A. R. Blaustein. 1999. Sensitivity to nitrate and nitrite in pond breeding amphibians from the Pacific Northwest, USA. *Environmental Toxicology and Chemistry* 18:2836–2839.
- Marrs, R.J., C.T. Williams, A.J. Frost, and R.A. Plant. 1989. Assessment of the Effects of Herbicide Spray Drift on a Range of Plant Species of Conservation Interest. *Environmental Pollution* 59:71-86
- Marshall, D. B., M. G. Hunter, and A. L. Contreras, Eds. 2003. *Birds of Oregon: a general reference.* Oregon State University Press, Corvallis, OR. 768 pp.
- Marshall, J.S., and L.W. Vandruff. 2002. Impact of Selective Herbicide Right-of-Way Vegetation on Birds. *Environmental Management* 30(6): p. 801-6.

- Maser C. 1998. Mammals of the Pacific Northwest: from the coast to the high Cascades. Oregon State Univ. Press, Corvallis. 406 p.
- Maser, Chris; Mate, Bruce R.; Franklin, Jerry F.; Dyrness, C. T. 1981. Natural history of Oregon coast mammals. Gen. Tech. Rep. PNW-133. Portland, OR.: USDA Forest Service, Pacific Northwest Range and Experiment Station. 496 p.
- Matthews G. M. and R. S. Waples. 1991. Status Review for Snake River Spring and Summer Chinook Salmon. NOAA Technical Memorandum. U.S. Department of Commerce, U.S. Department of Commerce. Available at:
<http://www.nwfsc.noaa.gov/publications/techmemos/tm200/tm200.htm>
- Mazzu L.C. 2005. Common Control Measures for Pacific Northwest Invasive Plants. 58 pp.
<http://www.fs.fed.us/r6/invasiveplant-eis/Region-6-Inv-Plant-Toolbox/#link03>, accessed 11/2006. Mechanisms underlying the impacts of exotic plant invasions. Ecology Letters. 7:975–989.
- McAllister, K.R., and W. P. Leonard. 1997. Washington state status report for the Oregon Spotted Frog. Washington
- McAllister, K.R., W.P. Leonard, D.W. Hays, and R.C. Friesz. 1999. Washington State status report for the northern leopard frog. Washington Dept. of Fish and Wildl., Olympia. 36p
- McAllister, K.R., W.P. Leonard, D.W. Hays, and R.C. Friesz. 1999. Washington State status report for the northern leopard frog. Washington Dept. of Fish and Wildl., Olympia. 36p
- McEwan D. and T. A. Jackson. 1996. Steelhead Restoration and Management Plan for California. California Department of Fish and Game.
- McIntosh, B.A., J.R. Sedell, J.E. Smith, R.C. Wissmar, S.E. Clarke, G.H. Reeves, and L.A. Brown. 1994. Management history of eastside ecosystems: Changes in fish habitat over 50 years, 1935 to 1992. U.S. Forest Service, Pacific Northwest Research Station, General Technical Report. PNW-GTR 321.
- McMurray, S.T., R.L. Lochmiller, J.F. Boggs, D.M. Leslie Jr., Engle and others. 1993b. Opportunistic Foraging of Eastern Woodrats (*Neotoma floridana*) in Manipulated Habitats. American Midland Naturalist (130): p. 325-37..
- McMurray, S.T., R.L. Lochmiller, J.F. Boggs, D.M. Leslie Jr., Engle and others. 1993a. Woodrat Population Dynamics Following Modification of Resource Availability. American Midland Naturalist (129): p. 48-56.
- McPhail J. D. and C. C. Lindsey. 1970. Freshwater Fishes of Northwestern Canada and Alaska. Bulletin of the Fisheries Research Board of Canada.(173): 381.
- McPhail, J.D., and J.S. Baxter. 1996. A review of bull trout (*Salvelinus confluentus*) life-history and habitat use in relation to compensation and improvement opportunities. Department of Zoology, University of British Columbia. Fisheries Management Report No. 104. Vancouver, British Columbia, Canada.
- Mechanisms underlying the impacts of exotic plant invasions. Ecology Letters. 7:975–989.

- Meehan, W.R. 1991. Influences of Forest and Rangeland Management on Salmonid Fishes and Their Habitats. American Fisheries Society Special Publication 19.
- Meffe, G.K., and C.R. Carroll. 1994. Principles of conservation biology. Sinauer Associate, Inc. Sunderland, Massachusetts.
- Michael, Jerry Lee. 2002. Impact of herbicides on the forest ecosystem, aquatic ecosystems, and wildlife. The US Experience Special Issue 6: 593-608.
- Mills, G. S.; Dunning Jr., J. B.; Bates, J. M. 1989. Effects of urbanization on breeding bird community structure in Southwestern desert habitats. The Condor. (91): 416-28.
- Mitchell, J. 2006. Invasive weed specialist, USDA Forest Service, Umatilla National Forest, Walla Walla Ranger District, WA. Personal communication.
- Montana Bull Trout Scientific Group. 1995a. Upper Clark Fork River drainage bull trout status report (including Rock Creek). Prepared for Montana Bull Trout Restoration Team. Helena, Montana.
- Montana Bull Trout Scientific Group. 1995b. Bitterroot River drainage bull trout status report. Prepared for Montana Bull Trout Restoration Team. Helena, Montana.
- Montana Bull Trout Scientific Group. 1995c. Blackfoot River drainage bull trout status report. Prepared for Montana Bull Trout Restoration Team. Helena, Montana.
- Montana Bull Trout Scientific Group. 1995d. Flathead River drainage bull trout status report (including Flathead Lake, the North and Middle forks of the Flathead River and the Stillwater and Whitefish River). Prepared for Montana Bull Trout Restoration Team. Helena, Montana.
- Montana Bull Trout Scientific Group. 1995e. South Fork Flathead River drainage bull trout status report (upstream of Hungry Horse Dam). Prepared for Montana Bull Trout Restoration Team. Helena, Montana.
- Montana Bull Trout Scientific Group. 1996a. Swan River drainage bull trout status report (including Swan Lake). Prepared for Montana Bull Trout Restoration Team. Helena, Montana.
- Montana Bull Trout Scientific Group. 1996b. Lower Clark Fork River drainage bull trout status report (Cabinet Gorge Dam to Thompson Falls). Prepared for Montana Bull Trout Restoration Team. Helena, Montana.
- Montana Bull Trout Scientific Group. 1996c. Middle Clark Fork River drainage bull trout status report (from Thompson Falls to Milltown, including the lower Flathead River to Kerr Dam). Prepared for Montana Bull Trout Restoration Team. Helena, Montana.
- Montana Bull Trout Scientific Group. 1996d. Lower Kootenai River drainage bull trout status report (below Kootenai Falls). Prepared for Montana Bull Trout Restoration Team. Helena, Montana.
- Montana Bull Trout Scientific Group. 1996e. Middle Kootenai River drainage bull trout status report (between Kootenai Falls and Libby Dam). Prepared for Montana Bull Trout Restoration Team. Helena, Montana.

- Montana Bull Trout Scientific Group. 1996f. Upper Kootenai River drainage bull trout status report (including Lake Koocanusa, upstream of Libby Dam). Prepared for Montana Bull Trout Restoration Team. Helena, Montana.
- Morse L.J. Kartesz and L. Kutner 2006. Native Vascular Plants. <http://biology.usgs.gov/s+t/noframe/j085.htm>. accessed 12/2006.
- Mote, P.W. 2004. How and why is Northwest climate changing? Pp. 11-22 In D.L Peterson, J.L. Innes, and K. Obrian (eds.). Climate change, carbon, and forestry in Northwestern North America: Proceedings of a Workshop. November 14-15, 2001. Orcas Island, Washington. USDA Forest Service, Pacific Northwest Research Station, Gen. Tech. Rept. PNW-GTR-614.
- Mote, P.W. and 18 co-authors. 1999. Impacts of climate variability and change: Pacific Northwest. A report of the Pacific Northwest Regional Assessment Group for the U.S. Global Change Research Program. JISAO/SMA Climate Impacts Group, University of Washington, Seattle.
- Moyle, P.B. 2002. Inland Fishes of California, Revised and Expanded. University of California
- Myers J. M., R. G. Kope, G. J. Bryant, D. Teel, L. J. Lierheimer, T. C. Wainwright, W. S. Grand, F. W. Waknitz, K. Neely, S. T. Lindley and R. S. Waples. 1998. Status Review of Chinook Salmon from Washington, Idaho, Oregon, and California. U.S. Department of Commerce, NOAA Technical Memorandum. 443. Available at:
- National Assessment Synthesis Team. 2000. Climate change impacts on the United States: The potential consequences of climate variability and change. US Global Change Research Program. Washington, D.C. 154pp.
- NatureServe Explorer: An online encyclopedia of life [web application]. 2006. Version 1.6. Arlington, VA, USA: NatureServe. Available: <http://www.natureserve.org/explorer/>.
- NatureServe Explorer: An online encyclopedia of life [web application]. 2008. Version 1.6. Arlington, VA, USA: NatureServe. Available: <http://www.natureserve.org/explorer/>. Accessed 9/15/08.
- Nehlsen, W., J. Williams, and J. Lichatowich. 1991. Pacific salmon at the crossroads: stocks at risk from California, Oregon, Idaho, and Washington. Fisheries 16(02):4-21.
- Newmaster, S.G., F. W. Bell, and D. H. Vitt. 1999. The effects of glyphosate and triclopyr on common bryophytes and lichens in northwestern Ontario Canadian Journal of Forest Research 29:1101-1111.
- Newton, J.A., and S. Pribyl. 1994. Bull trout population summary: Lower Deschutes River subbasin. Oregon Department of Fish and Wildlife, The Dalles, Oregon. Oregon administrative rules, proposed amendments to OAR 340-41-685 and OAR 340-41-026. January 11, 1996
- Nolte, K. R.; Fulbright, T. E. 1997. Plant, small mammal, and avian diversity following control of honey mesquite. Journal of Range Management. 50(2): 205-212.

- Norton, J.B., T.A. Monaco, J.M. Norton, D.A. Johnson, and T.A. Jones. 2004. Soil morphology and organic Matter dynamics under cheatgrass and sagebrush-steppe plant communities. *Journal of Arid Environments* 57: 445-466
- Nussbaum, R. A.; Brodie Jr., E. D.; Storm R. M. 1983. Amphibians and reptiles of the Pacific Northwest. Moscow, ID: The University Press of Idaho.
- ODFW (Oregon Department of Fish and Wildlife). 1991. Grand Ronde River Subbasin Salmon and Steelhead Plan. Prepared for Northwest Power Planning Council.
- Office of the President (1994). Executive Order 12898, Federal Actions to Address Environmental Justice in Minority Populations and Low-Income Populations, Federal Register. 59.
- Olson, B.E. 1999. Impacts of noxious weeds on ecologic and economic systems. In: Sheley, R.L.; Petroff, J.K., eds. *Biology and management of noxious rangeland weeds*. Corvallis, Oregon: Oregon State Univ. Press. p. 4-18.
- ONRC (Oregon Natural Resources Council). 1992. Siskiyou Regional Education Project, Federation of Fly Fishers, Kalmiopsis Audubon Society, Siskiyou Audubon Society, Klamath/Siskiyou Coalition, Headwaters, the Wilderness Society, North Coast Environmental Center, Oregon Chapter of the Sierra Club, and the National Wildlife Federation, Petition for a Rule to List the Illinois River Winter Steelhead as Threatened or Endangered under the Endangered Species Act and to Designate Critical Habitat. Unpublished Manuscript, 16 P. (Document Submitted to USDOC NOAA NMFS Northwest Region, Seattle, Washington, May 1992).
- Oregon Natural Heritage Program. 2004. <http://oregonstate.edu/ornhic/ornhp.html>. Accessed 11/2006.
- Ortega, Y.K. 2005. Winter Range Restoration on the Lolo National Forest: Vegetation Response to Herbicide Treatments. Draft Unpublished Report. Rocky Mountain Research Station, Forestry Sciences Laboratory, Missoula, MT. 11p.
- Ortega, Y.K. and D.E. Pearson. 2005. Weak vs. strong invaders of natural plant communities: assessing invisibility and impact. *Ecological Applications*. 15(2): 651-661
- Ortega, Y.K., D.E. Pearson, K.McKelvey, and D.L. Six. 2005. Evaluating effects of herbicide for restoration of arthropod and songbird communities. Unpublished report. FS-PIAP RM-7 Final Report. Rocky Mountain Research Station, Forestry Sciences Laboratory, Missoula, MT. 16p.
- Page, L.M. and B.R. Burr. 1991. *A Field Guide to Freshwater Fishes: North America North of Mexico*. Houghton Mufflin.
- Pagel, J. 2006. Peregrine falcon nest site data, 1983-2006 collected for PNW Interagency Peregrine Falcon Program, unpublished data. Ashland OR
- Paschke, M.W., T. McLendon, and E.F. Redente. 2000. Nitrogen availability and old-field succession in a shortgrass steppe. *Ecosystems*. 3: 144-158

- Pauli, B.D., and S. Money. 2000. Ecotoxicology of pesticides in reptiles. pp. 269-324 In D.W. Sparling, G. Linder, and C.A. Bishop, eds. *Ecotoxicology of Amphibians and Reptiles*. Society of Environmental Toxicology and Chemistry. Pensacola, Florida. 904pp.
- Pearson, D.E., K.S. McKelvey, and L.F. Roggiro. 2000. Non-target effects of an introduced biological control agent on deer mouse ecology. *Oecologia* 122:121-128.
- Pellant, M. 1990. The cheatgrass-wildfire cycle – are there any solutions? In. McArthur, E. Durant, E.M. Romney, S.D. Smith, P.T. Tueller. *Proceedings of the Symposium on cheatgrass invasion, shrub die-off, and other aspects of shrub biology and management: 1989 april 5-7. Las Vegas, NV. USDA Forest Service Gen. Tech. Rep. INT-276. Ogden, UT.*
- Perkins, Peggy J., Boermans, Jerman H., Stephenson, Gerald R. 2000. Toxicity of Glyphosate and Triclopyr using the frog embryo. *Teratogenesis Assay-Xenopus. Environmental Toxicol and Chemistry* 19(4): p.940-5.
- Peterson, J.G. 1970. The food habits and summer distribution of juvenile sage grouse in central Montana. *Journal of Wildlife Management* 34: 147-155.
- Pettis, J., Vanengelsdorp, D., Cox-Foster, D. 2007. Colony collapse disorder working group pathogen sub-group progress report. *Amer.Bee J.* 147:595-597.
- Pfleeger, T.G., A. Fong, R. Hayes, H. Ratsch, and C. Wickliff. 1996. Field evaluation of the EPA (Kenaga) nomogram, a method for estimating wildlife exposure to pesticide residue on plants. *Environmental Toxicology and Chemistry* 15(4): p. 535-43.
- Pickart, A.J. and J.O. Sawyer. 1998. *Ecology and restoration of Northern California coastal dunes. California Native Plant Society. Sacramento, CA. 152 pgs*
- PIF (Partner In Flight). 2000. *Conservation Strategy for Landbirds in the Columbia Plateau of Eastern Oregon and Washington. Version 1.0. March 2000. 144 pp.*
- Pihakaski, S., and Pihakaski, K. 1980. Effects of glyphosate on ultrastructure and photosynthesis of *Pellia epiphylla*. *Annals of Botany* 46: 181–192.
- Pokorny, M.L., R.L. Sheley, C.A. Zabinski, R.E. Engel, T.J. Svejcar and J.J. Borkowski. 2005. Plant functional group diversity as a mechanism for invasion resistance. *Restoration Ecology*. 13: 448-458
- Pokorny, M.L., R.L. Sheley, T.J. Svejcar, and R.E. Engel. 2004. Plant species diversity in a grassland plant community: evidence for forbs as a critical management consideration. *Western North American Naturalist*. 64(2): 219-230
- Pope, J. 2006. Pesticide applicator and Valley Helicopter Service, Clarkston, ID. Personal communication via telephone.
- Pratt, K.L. 1985. Pend Oreille trout and char life history study. Idaho Department of Fish and Game in cooperation with the Pend Oreille Idaho Club.
- Pratt, K.L. 1992. A review of bull trout life history. Pages 5-9 in P.J. Howell, and D.V. Buchanan, eds. *Proceedings of the Gearhart Mountain Bull Trout Workshop. Oregon Chapter of the American Fisheries Society, Corvallis, Oregon.*

- Pratt, K.L., and J.E. Huston. 1993. Status of bull trout (*Salvelinus confluentus*) in Lake Pend Oreille and the lower Clark Fork River. Draft report. Prepared for the Washington Water Power Company, Spokane, Washington.
- Pyke, C.R.; R. Thomas; R.D. Porter; J.J. Hellmann; J.S. Duke; D.M. Lodge; and G. Chavarria. 2008. Current practices and future opportunities for policy on climate change and invasive species. *Conservation Biology* 22(3): 585-592.
- Quigley, T.M. and S.J. Arbelbide, tech. editors. 1997. An assessment of ecosystem components in the interior Columbia basin and portions of the Klamath and Great basins: volume III. General Technical Report PNW-GTR-4-5. U.S. Department of Agriculture, Forest Service, and U.S. Department of Interior, Bureau of Land Management.
- Quigley, T.M., and S.J. Arbelbide. Tech Eds., 1997. An Assessment of Ecosystem Components in the Interior Columbia Basin and Portions of the Klamath and Great Basins. Gen. Tech. Rep. PNW-GTR-405. Portland, OR. USDA, Forest Service, Pacific Northwest Research Station. Vol. 4.
- Raloff, J. 1998. Botanical 'velcro' entraps hummingbirds- burrs cause bird fatalities. [Online]. Available: http://www.findarticles.com/p/articles/mi_m1200/is_n16_v154/ai_21250276. Accessed July 2004.
- Randall, J.M. 1996. Weed control for the preservation of biological diversity. *Weed Technology*. 10: 370-383.
- Rashin and Graber. 1993. Effectiveness of best management practices for aerial application of forest pesticides. TFW-WQ1-93-001-127 pp.
- Ratliff, D.E., and P.J. Howell. 1992. The status of bull trout populations in Oregon. Pages 10-17 in: P.J. Howell and D.V. Buchanan (eds). *Proceedings of the Gearhart Mountain bull trout workshop*. Oregon Chapter of the American Fisheries Society, Corvallis.
- Rawinski T. J. 1982. The ecology and management of purple loosestrife (*Lythrum salicaria* L.) in central New York. Ithaca, N.Y.: Cornell University, ix: 88.
- Rawinski, T. J.; Malecki, R. A. 1984. Ecological relationships among purple loosestrife, cattail and wildlife at the Montezuma National Wildlife Refuge. *New York Fish and Game Journal*. 31(1): 81-87.
- Relyea, R.A. 2005. The impact of insecticides and herbicides on the biodiversity and productivity of aquatic communities. *Ecological Applications* 15(2): p. 618-27.
- Reynolds, R. T., R. T. Graham, M. H. Reiser; and others. 1992. Management recommendations for the northern goshawk in the southwestern United States. General Technical Report RM-217. Ft. Collins, CO: U. S. Department of Agriculture, Forest Service, Rocky Mountain Forest and Range Experiment Station. 184 p.
- Rice, P. M.; Toney, C.; Bedunah, D. J.; Carlson, C. E. 1997. Plant community diversity and growth form responses to herbicide applications for control of *Centaurea maculosa*. *Journal of Applied Ecology*. 34: 1397-1412.

- Rice, Peter. 2000. Restoration of native plant communities infested by invasive weeds--Sawmill Creek Research Natural Area. In: Smith, Helen Y., ed. *The Bitterroot Ecosystem Management Research Project: what we have learned: Symposium proceedings; 1999 May 18-20; Missoula, MT. Proceedings RMRS-P-17.* Ogden, UT: U.S. Department of Agriculture, Forest Service, Rocky Mountain Research Station: 29-30.
http://www.fs.fed.us/rm/ecopartner/vegrecent_papers.shtml (accessed 11/2007).
- Rich, C.F., Jr. 1996. Influence of abiotic and biotic factors on occurrence of resident bull trout in fragmented habitats, western Montana. M.S. Thesis, Montana State University, Bozeman, Montana.
- Rieman, B.E., and J.B. Dunham. 2000. Metapopulations and salmonids: a synthesis of life history patterns and empirical observations. *Ecology of Freshwater Fish* 9:1-2.
- Rieman, B.E., and J.D. McIntyre. 1993. Demographic and habitat requirements for conservation of bull trout. General Technical Report INT-302. U.S. Department of Agriculture, Forest Service, Intermountain Research Station, Ogden, Utah.
- Rieman, B.E., and J.D. McIntyre. 1995. Occurrence of bull trout in naturally fragmented habitat patches of varied size. *Transactions of the American Fisheries Society* 124(3):285-296.
- Rieman, B.E., and J.D. McIntyre. 1996. Spatial and temporal variability in bull trout redd counts. *North American Journal of Fisheries Management* 16:132-141.
- Rieman, B.E., D.C. Lee, and R.F. Thurow. 1997. Distribution, status, and likely future trends of bull trout within the Columbia River and Klamath River basins. *North American Journal of Fisheries Management*. 17:1111-1125.
- Rieman, B.E., D.C. Lee, and R.F. Thurow. 1997. Distribution, status, and likely future trends of bull trout within the Columbia River and Klamath River basins. *North American Journal of Fisheries Management*. 17:1111-1125.
- Robbins, C. S., D. K. Dawson, and B. A. Dowell. 1989. Habitat requirements of breeding birds of the Middle Atlantic States. *Wildlife Monographs* 103:1-34.
- Roelofs T. D. 1983. Current Status of California Summer Steelhead (*Salmo Gairdneri*) Stocks and Habitat, and Recommendations for Their Management. Submitted to USDA Forest Service, Region 5. Environmental and Technical Services Division, National Marine Fisheries Service.
- Rogers, Jr., H.H.; D.H. Gjerstad ; G.B. Runion; S.A. Prior; A.J. Price; E. Van Santen; H.A. Torbert III. 2005. Effects of elevated atmospheric CO₂ on invasive weed species in managed terrestrial ecosystems of the southeastern U.S. Southeastern Regional Center, National Institute for Global Environmental Change, Annual Report for Fiscal Year 2004. University of Alabama, Tuscaloosa, AL. 4 pp.
- Rogers, Jr., H.H.; G.B. Runion; S.A. Prior; A.J. Price; H.A. Torbert III; and D.H. Gjerstad. 2008a. Effects of elevated atmospheric CO₂ on invasive plants: comparison of purple and yellow nutsedge (*Cyperus rotundus* L. and *C. esculentus* L.). *Journal of Environmental Quality* 37: 395-400.

- Ronoprawiro, S. 1975. Control of mosses in tea. In Asian Pacific Weed Science Society Conference. Asian Pacific Weed Science, Beijing. pp. 365–369.
- Rosenshield, M.L., Jofre, M.B., and W.H. Karasov. 1999. Effects of polychlorinated biphenyl 126 on green frog (*Rana clamitans*) and leopard frog (*Rana pipiens*) hatching success, development, and metamorphosis. *Environmental Toxicology and Chemistry* 18:2478-2486.
- Rowland, M.M., M.J. Wisdom, B.J. Johnson, and J.G. Kie. 2000. Elk distribution and modeling in relation to roads. *J. Wildl. Manage.* 64(3): 672-684.
- Rudolph, H., and Samland, J. 1985. Occurrence and metabolism of Sphagnum acid in cell walls of bryophytes. *Phytochemistry*, 24: 745–749.
- Ruediger, B., J. Claar, S. Gniadek, B. Holt, L. Lewis, S. Mighton, B. Naney, G. Patton, T. Rinaldi, J. Trick, A. Vandehey, F. Wahl, N. Warren, D. Wenger, and A. Williamson. 2000. Canada lynx conservation assessment and strategy. USDA Forest Service, USDI Fish and Wildlife Service, USDI Bureau of Land Management, and USDI National Park Service. Forest Service Publication #R1-00-53, Missoula, MT. 142 pp. (<http://www.fs.fed.us/r1/wildlife/carnivore>)
- Ruggiero, L. F., K. B. Aubry, S. W. Buskirk, G. M. Koehler, C. J. Krebs, K. S. McKelvey, and J. R. Squires. 2000. The scientific basis for lynx conservation: qualified insights. Pages 443-454 in Ruggiero, L. F., K. B. Aubry, S. W. Buskirk, G. M. Koehler, C. J. Krebs, K. S. McKelvey, and J. R. Squires, editors. 2000. *Ecology and conservation of lynx in the United States*. University Press of Colorado, Boulder, CO. 480 pp.
- Ruggiero, Leonard F., K. B. Aubry, S. W. Buskirk, G. M. Koehler, C. J. Krebs, K. S. McKelvey, and J. R. Squires. 1999. *Ecology and Conservation of Lynx in the United States*. Univ. Press of Colorado. Boulder, CO and USDA, Forest Service, Rocky Mountain Research Station. General Technical Report, RMRS-GTR-30WWW, October. 480p
- Saab, V., and T. Rich. 1997. Large-scale conservation assessment for neotropical migratory landbirds in the Interior Columbia River Basin. USDA Forest Service General Technical Report PNW-GTR-399, Portland, Oregon, USA.
- Sallabanks, R., B. G. Marcot, R. A. Riggs, E. B. Arnett, and C. A. Mehl. 2001. Wildlife of eastside (interior) forests and woodlands. In: D. H. Johnson and T. A. O'Neil, managing directors. *Wildlife-habitat relationships in Oregon and Washington*. Corvallis, OR: Oregon State University Press; 213–238.
- Sasich, J. 2006. Landtype Associations of Blue Mountains Province (Malheur, Wallowa-Whitman, and Umatilla NF. USDA Forest Service, Portland, OR. 23p.
- Saunders, D.A., R.J. Hobbs, and C.R. Margules. 1991. Biological consequences of ecosystem fragmentation: A review. *Conservation Biology* 5:18-32.
- Schempf, P. F., and M. White. 1977. Status of six furbearer populations in the mountains of northern California. U.S. Dep. Agric., For. Serv., San Francisco, Calif. 51pp.
- Schill, D.J. 1992. River and stream investigations. Job Performance Report, Project F-73-R-13. Idaho Department of Fish and Game, Boise, Idaho.

- Schmidt, K.A., and C.J. Whelan. 1999. Effects of exotic *Lonicera* and *Rhamnus* on songbird nest predation. Volume 13.6. p. 1502-6.
- Scholz N. L., N. K. Truelove, B. L. French, B. A. Berejikian, T. P. Quinn, E. Casillas and T. K. Collier. 2000. Diazinon Disrupts Antipredator and Homing Behaviors in Chinook Salmon (*Oncorhynchus tshawytscha*). *Canadian Journal of Fisheries and Aquatic Sciences*. 57(9): 1911-1918.
- Schommer, Tim 2007. [Personal communication]. Baker City, OR: U.S. Department of Agriculture, Forest Service, Wallowa-Whitman National Forest, Forest Wildlife Biologist.
- Sedell, J.R., and F.H. Everest. 1991. Historic changes in pool habitat for Columbia River Basin salmon under study for TES listing. Draft U.S. Department of Agriculture Report, Pacific Northwest Research Station, Corvallis, Oregon.
- SERA (Syracuse Environmental Research Associated, Inc.). 2003. All risk assessments associated with herbicides can be found in the Regional FEIS USDA 2005a Pacific Northwest Regions Invasive Plant Program, Preventing and Managing Invasive Plants Final Environmental Impact Statement and appendices <http://www.fs.fed.us/r6/invasiveplant-eis/Region-6-Inv-Plant-Toolbox/#link12>. Accessed 11/2006.
- SERA (Syracuse Environmental Research Associates Inc.). 2003a. Glyphosate- Human Health and Ecological Risk Assessment Final Report. SERA TR 02-43-09-04a. March 1, 2003. http://www.fs.fed.us/foresthealth/pesticide/risk_assessments/04a03_glyphosate.pdf.
- SERA (Syracuse Environmental Research Associates Inc.). 2003a. Glyphosate- Human Health and Ecological Risk Assessment Final Report. SERA TR 02-43-09-04a. March 1, 2003. http://www.fs.fed.us/foresthealth/pesticide/risk_assessments/04a03_glyphosate.pdf
- SERA (Syracuse Environmental Research Associates, Inc.). 1999. Imazapyr - Human Health and Ecological Risk Assessment Preliminary Draft - Program Description. SERA TR 98- 21-14-01b. May 6, 1999.
- SERA (Syracuse Environmental Research Associates, Inc.). 2001. Impazapic [Plateau and Plateau DG] - Human Health and Ecological Risk Assessment Final Report. SERA TR 00-21-28-01e. January 28, 2001.
- SERA (Syracuse Environmental Research Associates, Inc.). 2001. Sethoxydim [Poast] - Human Health and Ecological Risk Assessment Final Report. SERA TR 01-43-01-01c. October 31, 2001. http://www.fs.fed.us/foresthealth/pesticide/risk_assessments/100202_sethoxydim_ra.PDF
- SERA (Syracuse Environmental Research Associates, Inc.). 2003. Chlorsulfuron - Human Health and Ecological Risk Assessment Preliminary Draft - Introduction and Program Description. SERA TR 02-43-18-01a. April 10, 2003.

- SERA (Syracuse Environmental Research Associates, Inc.). 2003. Metsulfuron methyl - Human Health and Ecological Risk Assessment Preliminary Draft - Introduction and Program Description. SERA TR 02-43-17-01a. April 14, 2003.
- SERA (Syracuse Environmental Research Associates, Inc.). 2003. Sulfometuron methyl - Human Health and Ecological Risk Assessment Preliminary Draft - Introduction and Program Description. SERA TR 02-43-17-02a. October 16, 2003.
- SERA (Syracuse Environmental Research Associates, Inc.). 2003b. Picloram - Revised Human Health and Ecological Risk Assessment Final Report. SERA TR 03-43-26-01b. June 30, 2003.
http://www.fs.fed.us/foresthealth/pesticide/risk_assessments/03431601b_picloram.pdf.
- SERA (Syracuse Environmental Research Associates, Inc.). 2003d. Triclopyr - Revised Human Health and Ecological Risk Assessments Final Report. SERA TR 02-43-13-03b. March 15, 2003.
http://www.fs.fed.us/foresthealth/pesticide/risk_assessments/0303_triclopyr.pdf.
- SERA (Syracuse Environmental Research Associates, Inc.). 2004a. Chlorsulfuron - Human Health and Ecological Risk Assessment – Final Report. SERA TR 04-43-18-01c. November 21, 2004.
http://www.fs.fed.us/foresthealth/pesticide/risk_assessments/112104_chlorsulf.pdf.
- SERA (Syracuse Environmental Research Associates, Inc.). 2004b. Clopyralid - Human Health and Ecological Risk Assessment - Final Report. SERA TR 04-43-17-03c. December 5, 2004.
http://www.fs.fed.us/foresthealth/pesticide/risk_assessments/120504_clopyralid.pdf.
- SERA (Syracuse Environmental Research Associates, Inc.). 2004c. Impazapic - Human Health and Ecological Risk Assessment - Final Report. SERA TR 04-43-17-04b. December 23, 2004.
http://www.fs.fed.us/foresthealth/pesticide/risk_assessments/122304_Imazapic.pdf.
- SERA (Syracuse Environmental Research Associates, Inc.). 2004d. Imazapyr - Human Health and Ecological Risk Assessment – Final Report. SERA TR 04-43-17-05b. December 18, 2004.
http://www.fs.fed.us/foresthealth/pesticide/risk_assessments/121804_Imazapyr.pdf.
- SERA (Syracuse Environmental Research Associates, Inc.). 2004e. Metsulfuron methyl - Human Health and Ecological Risk Assessment Final Report. SERA TR 02-43-17-01b. December 9, 2004.
http://www.fs.fed.us/foresthealth/pesticide/risk_assessments/120804_Metsulfuron.pdf.

- SERA (Syracuse Environmental Research Associates, Inc.). 1997a. Effects of Surfactants on the Toxicity of Glyphosate, with Specific Reference to RODEO. SERA TR 97-206-1b February 6, 1997.
- SERA (Syracuse Environmental Research Associates, Inc.). 1997b. Use and Assessment of Marker Dyes Used With Herbicides. SERA TR 96-21-07-03b December 21, 1997.
- SERA (Syracuse Environmental Research Associates, Inc.). 1999. Clopyralid (Transline)- Final Report. SERA TR 99-21-11/12-01c. February 27, 1999.
- SERA (Syracuse Environmental Research Associates, Inc.). 1999b. Imazapyr - Human Health and Ecological Risk Assessment Preliminary Draft - Program Description. SERA TR 98- 21-14-01b. May 6, 1999.
- SERA (Syracuse Environmental Research Associates, Inc.). 2001. Sethoxydim [Poast] - Human Health and Ecological Risk Assessment Final Report. SERA TR 01-43-01-01c. October 31, 2001.
http://www.fs.fed.us/foresthealth/pesticide/risk_assessments/100202_sethoxydim_ra.PDF
- SERA (Syracuse Environmental Research Associates, Inc.). 2001b. Preparation of Environmental Documentation of Risk Assessments. SERA MD 2001-01a. July 18, 2001.
- SERA (Syracuse Environmental Research Associates, Inc.). 2001c. Sethoxydim [Poast] - Human Health and Ecological Risk Assessment Final Report. SERA TR 01-43-01-01c. October 31, 2001.
http://www.fs.fed.us/foresthealth/pesticide/risk_assessments/100202_sethoxydim_ra.pdf
- SERA (Syracuse Environmental Research Associates, Inc.). 2003b. Picloram - Revised Human Health and Ecological Risk Assessment Final Report. SERA TR 03-43-26-01b. June 30, 2003.
http://www.fs.fed.us/foresthealth/pesticide/risk_assessments/03431601b_picloram.pdf
- SERA (Syracuse Environmental Research Associates, Inc.). 2003c. Sulfometuron methyl - Human Health and Ecological Risk Assessment – Final Report. SERA TR 03-43-17-02c. December 14, 2004.
http://www.fs.fed.us/foresthealth/pesticide/risk_assessments/121404_Sulfometuron.pdf
- SERA (Syracuse Environmental Research Associates, Inc.). 2003d. Triclopyr - Revised Human Health and Ecological Risk Assessments Final Report. SERA TR 02-43-13-03b. March 15, 2003.
http://www.fs.fed.us/foresthealth/pesticide/risk_assessments/0303_triclopyr.pdf
- SERA (Syracuse Environmental Research Associates, Inc.). 2004a. Chlorsulfuron - Human Health and Ecological Risk Assessment – Final Report. SERA TR 04-43-18-01c. November 21, 2004.
http://www.fs.fed.us/foresthealth/pesticide/risk_assessments/112104_chlorsulf.pdf

- SERA (Syracuse Environmental Research Associates, Inc.). 2004b. Clopyralid - Human Health and Ecological Risk Assessment - Final Report. SERA TR 04-43-17-03c. December 5, 2004.
http://www.fs.fed.us/foresthealth/pesticide/risk_assessments/120504_clopyralid.pdf
- SERA (Syracuse Environmental Research Associates, Inc.). 2004c. Impazapic - Human Health and Ecological Risk Assessment - Final Report. SERA TR 04-43-17-04b. December 23, 2004.
http://www.fs.fed.us/foresthealth/pesticide/risk_assessments/122304_Imazapic.pdf
- SERA (Syracuse Environmental Research Associates, Inc.). 2004d. Imazapyr - Human Health and Ecological Risk Assessment – Final Report. SERA TR 04-43-17-05b. December 18, 2004.
http://www.fs.fed.us/foresthealth/pesticide/risk_assessments/121804_Imazapyr.pdf
- SERA (Syracuse Environmental Research Associates, Inc.). 2004e. Metsulfuron methyl - Human Health and Ecological Risk Assessment Final Report. SERA TR 02-43-17-01b. December 9, 2004.
http://www.fs.fed.us/foresthealth/pesticide/risk_assessments/120804_Metsulfuron.pdf
- Sexauer, H.M., and P.W. James. 1997. Microhabitat use by juvenile trout in four streams located in the eastern Cascades, Washington. Pages 361-370 in W.C. Mackay, M.K. Brewin and M. Monita (eds). Friends of the Bull Trout Conference Proceedings. Bull Trout Task Force (Alberta), c/o Trout Unlimited Calgary, Alberta, Canada.
- Sheley, R.L., and Petroff, J.K., eds. 1999. Biology and management of noxious rangeland weeds. Corvallis OR: Oregon State University Press. 438p.
- Sheley, R.L., T.J. Svejcar, B.D. Maxwell, and J.S. Jacobs. 1996. Successional Rangeland Weed Management. Rangelands. 18:155-159.
- Shen, Miaoqing; Xiaolong Yang; D. Cox-Foster; Liwang Cui. 2005. The Role of varroa mites in infection of Kashmir bee virus (KBV) and deformed wing virus (DMV) in honey bees. Virology 342: 141-149.
- Shepherd, M.D., S.L. Buchmann, M. Vaughan, and S.H. Black. 2003. Pollinator Conservation Handbook: A Guide to Understanding, Protecting, and Providing Habitat for Native Pollinator Insects. The Xerces Society in Association with The Bee Works. Portland: The Xerces Society. 145 pp. www.pollinator.org/Resources/Pollination%20Handbook.pdf (accessed 11/2007).
- Sherry, T. W., & R. T. Holmes. 1992. Population fluctuations in a long-distance Neotropical migrant: demographic evidence for the importance of breeding season events in the American Redstart. Pp. 431-442 in Ecology and Conservation of Neotropical Migrant Landbirds (J.M. Hagan and D. W. Johnston, eds.), Smithsonian Press, Wash. D.C.
- Simpson, J.C., and R.L. Wallace. 1982. Fishes of Idaho. University Press of Idaho. Moscow, Idaho.
- Smith, G.R. 2001. Effects of acute exposure to a commercial formulation of glyphosate on the tadpoles of two species of Anurans. Bull. Environ. Contam. Toxicol. 67: 483-488.
107Source <http://www.ag.ndsu.edu/pubs/plantsci/weeds/a657w.htm#factors>

- Smith, S.D.; T.E. Huxman; S.F. zitzer; T.N. Charlet; D.C. Housman; J.S. Coleman; L.K. Fenstermaker; J.R. Seemann; and R.S. Nowak. 2000. Elevated CO₂ increases productivity and invasive species success in an arid ecosystem. *Nature* 408:79-82.
- Soil Conservation Service (SCS). 1982. Southeast Washington cooperative river basin study, Snake River watershed. Draft. Spokane, Washington.
- Spahr, R., L. Armstrong, D. Atwood, and M. Rath. 1991. Threatened, Endangered, and Sensitive Species of the Intermountain Region. USDA Forest Service, Intermountain Region, Ogden, Utah.
- Spruell, P., B.E. Rieman, K.L. Knudsen, F.M. Utter, and F.W. Allendorf. 1999. Genetic population structure within streams: microsatellite analysis of bull trout populations. *Ecology of Freshwater Fish* 8:114-121.
- Squires, J. R., and R. T. Reynolds. 1997. Northern Goshawk. No. 298. In *The birds of North America*. The Academy of Natural Sciences, Philadelphia, Pennsylvania, and The American Ornithologists' Union, Washington, DC, USA.
- St. John, Alan. 2002. *Reptiles of the Northwest: California to Alaska – Rockies to the Coast*. Lone Pine Publishing. Renton, WA. 272p
- Stebbins, R. C. 1985. *The Peterson Field Guide Series: A Field Guide to Western Reptiles and Amphibians*. Houghton Mifflin Company, Boston, MA. 336 pp.
- Stein, B.A., and S.R. Flack, eds. 1996. *America's Least Wanted: Alien Species Invasion of U.S. Ecosystems*. The Nature Conservancy, Arlington, Virginia. 31 p.
- Stjernquist, I. 1981. Photosynthesis, growth and competitive ability of some coniferous forest mosses and the influence of herbicides and heavy metals (Cu, Zn). Unpublished thesis, University of Lund, Lund, Sweden. From Newmaster et al 1999.
- Sullivan, T., and D. Sullivan. 2003. Vegetation management and ecosystem disturbance: impact of glyphosate herbicide on plant and animal diversity in terrestrial systems *Environmental review* 11:37-58.
- Sullivan, T.P., C. Nowotny, and R.A. Lautenschlager. 1998. Silvicultural use of herbicide in sub-boreal spruce forest: implications for small mammal population dynamics. *Journal of Wildlife Management* 62(4): p. 1196-206.
- Sussmann, P. 2006. Deschutes NF, Ochoco NF and Crooked River National Grasslands Invasive Plants Environmental Impact Statement. USDA Forest Service, Region 6, Deschutes NF. Bend, OR. 35p.
- Sutherland, S. 2004. What makes a weed a weed: life history traits of native and exotic plants in the USA. *Oecologia* 141: 24-39
- Swanberg, T. R. 1997. Movements of and habitat use by fluvial bull trout in the Blackfoot River, Montana. *Transactions of the American Fisheries Society* 126:735-746.
- Swenson, J.E.; C.A. Simmons, and C.D. Eustace. 1987. Decrease of sage grouse *Centrocercus urophasianus* after ploughing of sagebrush steppe. *Biological Conservation* 41: 125-132.

- Talsma, A. 2006. Western Idaho Project Manager. The Nature Conservancy. Boise ID. Personal communication..
- Talsma, A., M. Lucia, and J. Karl. 2006. Starthistle control using helicopter application of transline and tordon. Unpublished internal document. The Nature Conservancy. Boise, ID. 2p.
- Tausch, R.J. 2008. Invasive plants and climate change. (May 20, 2008). U.S. Department of Agriculture, Forest Service, Climate Change Resource Center. <http://www.fs.fed.us/ccrc/topics/invasive-plants.shtml>
- Taylor, K., and C. Potvin. 1997. Understanding the long-term effect of CO₂ enrichment on a pasture: the importance of disturbance. *Canadian Journal of Botany* 75(10): 1621-1627.
- Terrestrial Ecosystem Unit Inventory. Unpublished. Region 6, Wallowa Whitman NF, Baker City, OR.
- Thayer, T.P., and C.E. Brown. 1966a. Geologic map of Aldrich Mountain quadrangle. US Geological Survey. Grant County, OR. Map CQ-438.
- Thayer, T.P., and C.E. Brown. 1966b. Geologic map of Mount Vernon quadrangle. US Geological Survey. Grant County, OR. Map DQ-548.
- The Nature Conservancy (TNC). 2006. Starthistle Control Using Helicopter Application of Transline and Tordon.--Aerial Monitoring and Application of Herbicide to Control Yellow Starthistle in Rugged Range and Large Landscapes. Unpublished report.
- The Nature Conservancy 2005. The Global Invasive Species Initiative. <http://tncweeds.ucdavis.edu/> accessed 11/2006.
- The Research Group (2000). Economic Analysis of Containment Programs, Damages, and Production Losses From Noxious Weeds in Oregon. Corvallis, Oregon, Oregon Department of Agriculture Plant Division, Noxious Weed Control Program.
- Thomas, G. 1992. Status of bull trout in Montana. Report prepared for Montana Department of Fish, Wildlife and Parks, Helena, Montana.
- Thomas, J. W., H. Black, Jr., R. J. Scherzinger, and R. J. Petersen. 1979. Wildlife Habitats in Managed Forests: The Blue Mountains of Oregon and Washington. [USDA Agriculture Handbook 553]. Washington, D.C. (Pages 60-77).
- Thomas, J. W.; Maser, C.; Rodiek, J. E. 1979. Wildlife habitats in managed rangelands - the Great Basin of southeastern Oregon: riparian zones. Portland, OR.: USDA Forest Service. Gen. Tech. Rep. PNW-80. 18 p.
- Thompson, D. Q., Stuckey, R. L.; Thompson, E. B. 1987. Spread, impact, and control of purple loosestrife (*Lythrum salicaria*) in North American wetlands. Washington, D.C.: USDI Fish and Wildlife Service; Research Report No. 2. 55 p. <http://www.npwrc.usgs.gov/resource/plants/loosstrf/index.htm>
- Timchalk C; Finco DR; Quast JF. 1997. Evaluation of renal function in rhesus monkeys and comparison to beagle dogs following oral administration of the organic acid triclopyr (3,5,6-trichloro-2-pyridinyloxyacetic acid). *Fundam Appl Toxicol.* 36(1): 47-53.

- Timchalk C; Nolan RJ. 1997. Pharmacokinetics of triclopyr (3,5,6-trichloro-2-pyridinyloxyacetic acid) in the beagle dog and rhesus monkey: perspective on the reduced capacity of dogs to excrete this organic acid relative to the rat, monkey, and human. *Toxicology and Applied Pharmacology*. 144 (2): 268-278.
- Trammell, M. A.; Butler, J. L. 1995. Effects of exotic plants on native ungulate use of habitat. *Journal of Wildlife Management*. 59(4): 808-816.
- Tu, M., C. Hurd, and J.M Randall. 2001. *Weed Control Methods Handbook: Tools and Techniques for Use in Natural Areas*. <http://tncweeds.ucdavis.edu/handbook.html>. April 2001 revision.
- Tyser, R.W. and C.H. Key. 1988. Spotted knapweed in natural area fescue grasslands: an ecological assessment. *Northwest Science* 62: 151-158
- U. S. Fish and Wildlife Service. 2007. 50CFR Part 17. Endangered and Threatened Wildlife and Plants; Removing the Bald Eagle in the Lower 48 States from the List of Endangered and Threatened Wildlife; Final Rule. *Federal Register* Vol. 72, No. 130. July 9, 2007.
- U.S. Fish and Wildlife Service. 1986. Recovery Plan for the Pacific Bald Eagle. U.S. Department of the Interior, Fish and Wildlife Service. Portland, OR. 160 pp.
- U.S. Fish and Wildlife Service. 1995. eagle.
- U.S. Fish and Wildlife Service. 1999. Proposed rule: delist bald eagle (*Haliaeetus leucocephalus*) in the lower 48 states. *Federal Register* 64:36453-36464.
- U.S. Fish and Wildlife Service. 2004. 12 month finding for a petition to list the west coast distinct population segment of the fisher (*Martes pennanti*); Proposed Rule. 24 pp.
- U.S. Fish and Wildlife Service. 2005. Lynx Conservation Agreement between the U.S. Forest Service and U.S. Fish and Wildlife Service. 9 pp.
- U.S. Fish and Wildlife Service. 2005. Lynx Recovery Outline. Contiguous United States Distinct Population Segment of the Canada Lynx. 21 pp.
- U.S. Fish and Wildlife Service. 2007. National Bald Eagle Management Guidelines. U.S. Fish and Wildlife Service. 23 p.
- U.S. Forest Service. 1990. Land and Resource Management Plan, Umatilla National Forest. USDA, Forest Service, Pacific Northwest Region (6), Wallowa-Whitman National Forest.
- University of Georgia Invasive Weeds Task Force. 2005. Japanese Knotweed. Invasive and Exotic Species. Joint project of The Bugwood Anetwork, USDA Forest Service, USDA APHIS PPQ, and The University of Georgia. Available [ONLINE] @ <http://www.invasive.org/browse/subject.cfm?sub=3414> [Feb 2008].
- US Census Bureau (2000). 2006. P4. Hispanic or Latino, Not Hispanic or Latino by Race.
- US Census Bureau (2000). Total Population. 2006.

- US Department of Commerce, NOAA Fisheries. 1993. Endangered and Threatened Species; Illinois River Winter Steelhead in Oregon. Federal Register [Docket 930517-3117, 20 May 1993].
- US Department of Commerce, NOAA Fisheries. 1997. Status Review Update for West Coast Steelhead from Washington, Idaho, Oregon, and California. Memorandum from the Biological Review Team to the NMFS Northwest Regional Office.
- US Department of Commerce, NOAA Fisheries. 2005. Endangered Species Act Section 7 Formal Consultation and Magnuson-Stevens Fishery Conservation and Management Act Essential Fish Habitat Consultation for the Pacific Northwest Region Invasive Plant Program, Oregon and Washington. NOAA Fisheries Northwest Region, Seattle, WA.
- US Environmental Protection Agency. 2004. Overview of the Ecological Risk Assessment Process in the Office of Pesticide Programs, U.S. Environmental Protection Agency. Office of Prevention, Pesticides and Toxic Substances, Office of Pesticide Programs. Washington, D.C. January 23, 2004. 92 pp.
- USDA and USDI. 1995. Decision Notice/Decision Record Finding of No Significant Impact, Environmental Assessment for the Interim Strategies for Managing Anadromous Fish-producing Watersheds in Eastern Oregon, and Washington, Idaho, and portions of California (PACFISH). http://www.fs.fed.us/r6/uma/blue_mtn_planrevision/documents/2006_05_08_SDA_public.pdf accessed 11/2006
- USDA and USDI. 1996. Status of the Interior Columbia Basin, Summary of Scientific Findings.
- USDA and USDI. 1997. Interior Columbia River Basin Ecosystem Management Project, Upper Columbia River Basin Draft Environmental Impact Statement. Vol I., Vol II, Vol III.
- USDA Forest Service. 2003. A Framework for the Development of the National Invasive Species Management Strategy for the USDA Forest Service, 2003. US Department of Agriculture, Forest Service, Washington, D.C.
- USDA Forest Service. 2008. "Invasive Plant Biological Assessment, Umatilla and Wallowa-Whitman National Forests" September 8, 2008.
- USDA Forest Service. 1988. Vegetation Management EIS and Record of Decision.
- USDA Forest Service. 1989. Mediated Agreement focused on competing vegetation in forest plantations
- USDA Forest Service. 1990. Land and Resource Management Plan, Wallowa-Whitman National Forest. USDA, Forest Service, Pacific Northwest Region (6), Wallowa-Whitman National Forest.
- USDA Forest Service. 1990a. Forest Service Manual (FSM) 2600, Wildlife, Fish and Sensitive Plant Habitat Management. Washington Office Amendment 2600-90-1, effective 6/1/90.

- USDA Forest Service. 1992. The Wallowa-Whitman National Forest Environmental Assessment (EA) for the Management of Noxious Weeds and Forest Plan Amendment 4. Wallowa-Whitman National Forest. 1992.
- USDA Forest Service. 1994a. The 1994 Wallowa-Whitman Management of Noxious Weeds Environmental Assessment. Wallowa-Whitman National Forest. 1994.
- USDA Forest Service. 1994b. The scientific basis for conserving forest carnivores; American marten, fisher, lynx, and wolverine, in the western United States. GTR, RM-254. U.S. Dept. Of Agriculture, Forest Service, Rocky Mountain Forest and Range Experiment Station. Fort Collins, Colorado. September. 184p
- USDA Forest Service. 1994c. Pesticide-Use Management and Coordination Handbook. Forest Service Handbook (FSH) 2109.14-94-1. USDA Forest Service, Washington D.C. December 6, 1994.
- USDA Forest Service. 1995a. Forest Service Manual (FSM) 2600, Wildlife, Fish and Sensitive Plant Habitat Management. R-6 Supplement 2600-95-3, effective 6/29/95.
- USDA Forest Service. 1995b. Forest Service Manual 2080, Noxious Weed Management. Washington, D.C. <http://www.fs.fed.us/im/directives/fsm/2000/2080.txt>.
- USDA Forest Service. 1998. Stemming the Invasive Tide: Forest Service Strategy for Noxious and Nonnative Invasive Plant Management. National Office, Washington, D.C.
- USDA Forest Service. 1999b. Noxious Weed Strategy. Pacific Northwest Region, Portland, Oregon.
- USDA Forest Service. 1999c. North Fork John Day Ecosystem Analysis. Umatilla National Forest. Umatilla National Forest, Pendleton, OR.
- USDA Forest Service. 2000. Landbird Strategic Plan. Washington Office. Washington, D.C. (FS-648)
- USDA Forest Service. 2001a. Phillips-Gordon Ecosystem Analysis, Umatilla National Forest. Pendleton, OR. 126p.
- USDA Forest Service. 2001b. Guide to Noxious Weed Prevention Practices. U.S. Department of Agriculture, Forest Service, Washington, D.C. www.fs.fed.us/rangelands/ftp/invasives/ - accessed 11/2006
- USDA Forest Service. 2003a. (HFQLG FSEIS) Herger-Feinstein Quincy Library Group Forest Recovery Act. Final Supplemental Environmental Impact Statement and Record of Decision. Lassen, Plumas, and Tahoe National Forests, Pacific Southwest Region.
- USDA Forest Service. 2003b. Human and Ecological Risk Assessment of NPE Surfactants in Forest Service Herbicide Applications, 2003. David Bakke. Pacific Southwest Region. Vellego, California.
- USDA Forest Service. 2003c. Land and Resource Management Plan, Wallowa-Whitman National Forest; Hells Canyon Comprehensive Management Plan (CMP) (Forest Plan Amendment #29). US Forest Service Wallowa-Whitman National Forest. 2003.

- USDA Forest Service. 2004a. Blue Mountains Forest Plan Revision: Draft Current Management Situation Report. Malheur, Umatilla, and Wallowa-Whitman National Forests.
- USDA Forest Service. 2004b. Implementing the National Fire Plan Using Livestock Grazing to Manage Vegetation (Official Letter).
- USDA Forest Service. 2004c. National Strategy and Implementation Plan for Invasive Species Management FS-805.
http://www.fs.fed.us/invasivespecies/documents/Final_National_Strategy_100804.pdf accessed 11/06.
- USDA Forest Service. 2004d. Regional Forester Sensitive Species List (Update). Forest Service Manual 2670. Portland, OR. USDA Forest Service. Pacific NW Region
- USDA Forest Service. 2005d. Biological Assessment for the USDA Forest Service, Pacific Northwest Region, Invasive Plant Program. USDA Forest Service, Pacific Northwest Region, Portland, OR. June 17, 2005. 452pp + appendices.
- USDA Forest Service. 2005a. Pacific Northwest Region Invasive Plant Program. Preventing and Managing Invasive Weeds Final Environmental Impact Statement. R6-NR-FHP-PR-02-05.
- USDA Forest Service. 2005b. Pacific Northwest Region Invasive Plant Program Record of Decision. USDA Forest Service, Pacific Northwest Region, Portland, OR. October 2005, R6-NR-FHP-PR-02-05. 39 pp. + appendices.
- USDA Forest Service. 2005c. Fisheries Biological Assessment for USDA Forest Service, Pacific Northwest Region, Invasive Plant Program, Environmental Impact Statement. USDA Forest Service, Pacific Northwest Region, Portland, OR. 221pp.
- USDA Forest Service. 2006a. Umatilla National Forest Invasive Plant Treatment Project, Botany Report and Biological Evaluation. Umatilla National Forest.
- USDA Forest Service. 2006b. Specially Designated Areas in Forest Plan Revision. Blue Mountain Forest Revision, Draft.
http://www.fs.fed.us/r6/uma/blue_mtn_planrevision/documents/2006_05_08_SDA_public.pdf accessed 11/2006
- USDA Forest Service. 2006c. Weed management, Custer National Forest Draft EIS, Appendix N. Drift Model Results. http://www.fs.fed.us/r1/custer/projects/Planning/weed_eis.
- USDA Forest Service. 2006e. Blue Mountain Land Exchange-Oregon, Draft Environmental Impact Statement. <http://www.fs.fed.us/r6/w-w/projects/bmle/deis/>
- USDA Forest Service. 2007a. Wallowa-Whitman National Forest Invasive Plants Treatment Project, Botany Report and Biological Evaluation.
- USDA Forest Service. 2007b. Wallowa-Whitman National Forest website: Forest Facts. Wallowa-Whitman National Forest; <http://www.fs.fed.us/r6/w-w/> 2007.

- USDA Forest Service. 1995. Inland Native Fish Strategy, Environmental Assessment, Decision Notice and Finding of No Significant Impact. United States Department of Agriculture, Intermountain, Northern and Pacific Northwest Regions.
- USDA Forest Service. 2005d. Biological Assessment for the USDA Forest Service Pacific Northwest Region Invasive Plant Program. Pacific Northwest Region, US Forest Service. 453 pp. plus appendices.
- USDA Forest Service. 2005d. Pacific Northwest Region Invasive Plant Program Biological Assessment. USDA Forest Service, Pacific Northwest Region, Portland, OR. February 2005. June 2005. 141 pp.
- USDA Forest Service. 2005e. Pacific Northwest Region Invasive Plant Program Appendix P. Summary of Herbicide Effects to Wildlife. USDA Forest Service, Pacific Northwest Region, Portland, OR. February 2005. 141 pp.
- USDA Forest Service. 2005f. Pacific Northwest Region Invasive Plant Program Appendix J. The Effects of Non-herbicidal Methods of Invasive Plant Treatments on Wildlife, Fish and Plants. USDA Forest Service, Pacific Northwest Region, Portland, OR. 27 pp.
- USDA Forest Service. 2006. Lynx Conservation Agreement between USDA Forest Service and USDI Fish and Wildlife Service. 9 pp.
- USDA Forest Service. 2007. Water Erosion Prediction Project (WEPP: Disturbed WEPP). [Online]. Available: <http://forest.moscowfsl.wsu.edu/fswepp/> [Feb 2007].
- USDA Forest Service. July 21, 2004. Regional Forester Sensitive Species List (Update). Forest Service Manual 2670. Portland, OR.: USDA Forest Service, Pacific Northwest Region.
- USDA Natural Resource Conservation Service. 1999. American Elk (*Cervus Elaphus*). Fish and Wildlife Management Leaflet. Wildlife Management Institute. 8 pp.
- USDA Forest Service. 1994. The scientific basis for conserving forest carnivores; American marten, fisher, lynx, and wolverine, in the western United States. GTR, RM-254. U.S. Dept. Of Agriculture, Forest Service, Rocky Mountain Forest and Range Experiment Station. Fort Collins, Colorado. 184pp.
- USDA Forest Service 2008. Source and Effectiveness of Project Design Features/Criteria for Herbicide Use in Invasive Plant Treatment Projects Forest Service Region Six. Unpublished report. 17pp.
- USDI BLM, Fish and Wildlife Service, USDA Forest Service, Oregon Dept. of Fish and Wildlife, Oregon Dept. of State Lands. 2000. Greater sage-grouse and sagebrush-steppe ecosystems: management guidelines. Unpublished document. 27 pp.
- USDI Bureau of Land Management, USDI Fish and Wildlife Service, USDA Forest Service, Oregon Department of Fish and Wildlife, and Oregon Department of State Lands. 2000. Greater sage-grouse and sagebrush-steppe ecosystems; management guidelines. USDI Bureau of Land Management, Oregon and Washington. Portland, OR. 27 pp.

- USDI Fish and Wildlife Service. 2004c. Endangered and Threatened Wildlife and Plants; 12-month Finding for a Petition To List the West Coast Distinct Population Segment of the Fisher (*Martes pennanti*); Proposed Rule . Federal Register 69 (68): 18770-18792
- USDI Fish and Wildlife Service (J. Dillon). 1981. Bald Eagle Management Guidelines Oregon - Washington . U.S. Fish and Wildlife Service. Portland, OR .
- USDI Fish and Wildlife Service and USDC NOAA Fisheries. 1996. Policy regarding the distinct vertebrate population. Federal Register 61:4721-4725
- USDI Fish and Wildlife Service. 1986. Recovery plan for the Pacific Bald Eagle. U.S. Department of the Interior, Fish and Wildlife Service. Portland, OR. 160 pp.
- USDI Fish and Wildlife Service. 1987. Northern Rocky Mountain Wolf Recovery Plan. U.S. Fish and Wildlife Service, Denver, Colorado. 119 pp.
- USDI Fish and Wildlife Service. 1996. Endangered and Threatened Wildlife and Plants: Review of Plant and Animal Taxa that are Candidates for Listing as Endangered or Threatened Species. Federal Register 61(40):7596-7613.
- USDI Fish and Wildlife Service. 1998a. Proposed rule: threatened status for fish *Salvelinus confluentus* (bull trout) for the Coastal-Puget Sound, Jarbridge River, and St. Mary-Belly River populations. Federal Register 63:31693-31710.
- USDI Fish and Wildlife Service. 1998b. Final rule: Threatened status for fish *Salvelinus confluentus* (bull trout) for the Columbia River and Klamath River populations. Federal Register 63:31647-31674.
- USDI Fish and Wildlife Service. 2000. Determination of Threatened status for the contiguous United States distinct population segment of the Canada Lynx and related rule; final rule. Federal Register 65 (58) 16052-16086.
- USDI Fish and Wildlife Service. 2002a. 25 Chapters. In: Bull trout (*Salvelinus confluentus*) draft recovery plan. U.S. Fish and Wildlife Service, Portland, Oregon.
- USDI Fish and Wildlife Service. 2002b. Proposed rule: Critical habitat for fish *Salvelinus confluentus* (bull trout) for Klamath River and Columbia River DPSs. Federal Register 67:71235-71438.
- USDI Fish and Wildlife Service. 2003a. Final rule to reclassify and remove the gray wolf from the list of endangered and threatened wildlife in portions of the conterminous United States. Federal Register 68 (62):15804-15882, 50CFR Part 17, April 1, 2003.
- USDI Fish and Wildlife Service. 2003b. Rocky Mountain Wolf Recovery 2002 Annual Report. <http://www.fws.gov/mountain-prairie/species/mammals/wolf/annualrpt03>.
- USDI Fish and Wildlife Service. 2004a. Draft recovery plan for the Coastal-Puget Sound distinct population segment of bull trout (*Salvelinus confluentus*), Volume I and II. Portland, OR
- USDI Fish and Wildlife Service. 2004b. Final rule: critical habitat for fish *Salvelinus confluentus* (bull trout) for Klamath River and Columbia River DPSs. Federal Register 69:59996-60075.

<http://a257.g.akamaitech.net/7/257/2422/14mar20010800/edocket.access.gpo.gov/2003/pdf/03-16664.pdf>

- USDI Fish and Wildlife Service. 1995. Endangered and Threatened Species; Bald eagle reclassification; final rule. Federal Register 60(133): 36000-36010 .
- USDI Fish and Wildlife Service. 1996. Endangered and Threatened Wildlife and Plants: Review of Plant and Animal Taxa that are Candidates for Listing as Endangered or Threatened Species. Federal Register 61(40):7596-7613.
- USDI Fish and Wildlife Service. 2003c. Endangered and threatened wildlife and plants; notice of remanded determination of status for the contiguous United States distinct population segment of the Canada lynx; clarification of findings; final rule. U.S. Fish and Wildlife Service. Federal Register 50 Code of Federal Regulations, Part 17, Volume 68(128):68400766840101.
- USDI Fish and Wildlife Service. 2003d. Endangered and Threatened Wildlife and plants 90day finding on a petition to list the Western Sage Grouse . Federal Register 68(26): 6500-6504 .
- USDI Fish and Wildlife Service. 2007. Endangered and Threatened Wildlife and Plants; Removing the bald eagle in the lower 48 States from the list of Endangered and Threatened Wildlife, Final Rule; Draft post-delisting and monitoring plan for the bald eagle (*Haliaeetus leucocephalus*) and proposed information collection; Notice. Federal Register 72 (130): 37346-37347.
- Valdez, R. and P.R. Krausman. 1999. Description, distribution and abundance of mountain sheep in North America. Pages 3 - 22 in Mountain sheep of North American, eds. R. Valdez and P.R. Krausman. The University of Arizona Press, Tucson, AZ.
- Vallentine, J. F. 1990. Grazing management. Academic, San Diego, California, USA.
- Van Dyke, W.A., A. Sands, J. Yoakum, A. Polenz, and J. Blaisdell. 1983. Wildlife habitats in managed rangelands- Bighorn Sheep. General Technical Report #159. 37 p.
- Verts, B. J., and L. N. Carraway. 1998. Land Mammals of Oregon. University of California Press, Berkeley, CA. 668 pp.
- Vinton, M.V. and I.C. Burke. 1995. Interactions between individual plant species and soil nutrient status in shortgrass steppe. Ecology. 76(4): 1116-1133
- Vitousek P. M., D'Antonio C.M., Loope L.L., Rejmanek M., Westerbrooks R. 1997. Introduced species: a significant component of human-caused global change. New Zealand Journal of Ecology. 21 (1): 1-16.
- Washington Department of Fish and Wildlife. 1997. Grandy Creek trout hatchery biological assessment. FishPro Inc., and Beak Consultants.
- Washington Department of Fish and Wildlife. 2003. Game Management Plan: July 2003-June 2009.
- Washington Department of Fish and Wildlife. 2003. Game Management Plan: July 2003-June 2009. http://wdfw.wa.gov/wlm/game/management/final_gmp_01jan03.pdf

- Washington Department of Fish and Wildlife. 2005. Guidelines for Timing of In-Water Work Periods
- Washington Natural Heritage Program, Washington Dept. of Fish & Wildlife, and USDI Bureau of Land Management. 2005. Washington Herp Atlas. [Online]. Available: <http://www.dnr.wa.gov/nhp/refdesk/herp>. [Access date unknown.]
- Watson, A.K., and Renney, A.J. 1974. The biology of Canadian weeds, *Centaurea diffusa* and *C. maculosa*. Canadian Journal of Plant Science 54: p.678-701
- Watson, G., and T. W. Hillman. 1997. Factors affecting the distribution and abundance of bull trout: An investigation at hierarchical scales. North American Journal of Fisheries Management 17:237-252.
- Weber, K. T., C. L. Marcum, M. G. Burcham, and L. J. Lyon. 2000. Landscape influences on elk vulnerability to hunting. Intermountain Journal of Science 6:86-94.
- Weihe, P. E.; Neely, R. K. 1997. The effects of shading on competition between purple loosestrife and broad-leaved cattail. Aquatic Botany, 59:127-138.
- Weiher, E. I.C. Wisheu, P.A. Keddy, and D.R.J. Moore. 1996. Establishment, persistence, and management implications of experimental wetland plant communities. Wetlands, 16(2):208-18.
- Weiher, E., Wisheu, I.C., Keddy, P.A., and Moore, D.R.J. 1996. Establishment, Persistence, and Management Implications of Experimental Wetland Plant Communities. Wetlands 16(2): p.208-18.
- Whisenant, S.G. 1990. Changing fire frequencies on Idaho's Snake river plains: ecological and management implications. In: McArthur, E.D.; Romney, E. M.; Smith, S. D.; Tueller, P. T. eds. Proceedings of a Symposium on Cheatgrass Invasion, Shrub Die-off, and Other Aspects of Shrub Biology and Management. U.S. Forest Service Gen. Tech. Rep. INT-276. Ogden, UT.: Intermountain Forest and Range Experiment Station: 4-10.
- Whiteley, A. R., P. Spruell, and F.W. Allendorf. 2003. Population genetics of Boise Basin bull trout (*Salvelinus confluentus*). Final report to Rocky Mountain Research Station, Contract:RMRS # 00-JV-1122014-561.
- Wiemeyer, S.N., Bunck, C.M., and Stafford, C.J. 1993. Environmental Contaminants in Bald Eagle Eggs - 1980-84 and further interpretations of relationships to productivity and shell thickness. Archives of Environmental Contamination and Toxicoology 24: p.214-27.
- Wilde, S. B., S. deKozlowski, and T. Murphy. 2004. Invasive Weeds, Blue-Green Algae, Coots, and Bald Eagles: a deadly trophic cascade. Aquatic Nuisance Species Digest, Vol. 5, No. 2.
- Winfree, R.; N.M. Williams; J. Dushoff; and C. Kremen. 2007. Native bees provide insurance against ongoing honey bee losses. Ecology Letters 10: 1105-1113.
- Wisdom, M.J., R.S. Holthausen, B.C. Wales, C.D. Hargis, V.A. Saab, D.C. Lee, W.J. Hann, T.D. Rich, M.M. Rowland, W.J. Murphy and M.R. Earnes. 2000. Source habitats for

- terrestrial vertebrates of focus in the Interior Columbia Basin: Broad-scale trends and management implications, Volume 2 – Group level results. Gen. Tech. Rep. Threatened and Endangered Species, Sensitive Species and Management Indicator Species and the level of analysis required. PNW-GTR-485. pp. 181-190; 199-208; 219-223; 237-241; 242-248; 258-265; 293-297.
- Wissmar, R. C., J. E. Smith, B. A. McIntosh, H. W. Li, G. H. Reeves, and J. R. Sedell. 1994. A history of resource use and disturbance in riverine basins of eastern Oregon and Washington (early 1800s-1990s). *Northwest Science* 68:1-35.
- Withler, R.E. 1988. Genetic consequences of fertilizing chinook salmon (*Oncorhynchus tshawytscha*) eggs with pooled milt. *Aquaculture* 68(1):15-25.
- Witmer, G. W., S. K. Martin, and R. D. Sayler. 1998. Forest carnivore conservation and management in the interior Columbia Basin: Issues and environmental correlates. General Technical Report GTR-PNW-420. Portland, OR: USDA Forest Service, Pacific Northwest Research Station, Portland, OR. 51 pp.
- Wood, Tamara. 2001. Herbicide Use in the Management of Roadside Vegetation, Western Oregon, 1999–2000: Effects on the Water Quality of Nearby Streams, USGS, Water- Resources Investigations Report 01–4065.
(<http://www.ecy.wa.gov/programs/wq/plants/weeds/aqua011.html>).
- Wright, H.A., L. F. Neuenschwander, C. M. Britton. 1979. The role and use of fire in sagebrush-grass and pinyon-juniper plant communities. A state of the art review. U.S. Dep. Agric. For. Serv. Res. Pap. INT-58. 48pp.
- Wydoski, R.S. and R.R. Whitney. 1979. *Inland Fishes of Washington State*. University of Washington Press.
- Wydoski, R.S. and R.R. Whitney. 2003. *Inland Fishes of Washington State, Revised and Expanded*. University of Seattle Press, Seattle, WA
- Yates, G. 2006 and 2007. [Forest Botanist, Wallowa-Whitman Forest, Supervisor's Office, Baker, Oregon. Zavaleta, E. 2000. Valuing ecosystems services lost to Tamarix invasion in the United States. In: H.A. Mooney, R.J. Hobbs, eds. *Invasive species in a changing world*. Washington, D.C.: Island Press: 261-300.
- Ziller, J. S. 1992. Distribution and relative abundance of bull trout in the Sprague River subbasin, Oregon. Pages 18-29 in P.J. Howell, and D.V. Buchanan, eds. *Proceedings of the Gearhart Mountain Bull Trout Workshop*. Oregon Chapter of the American Fisheries Society, Corvallis, Oregon
- Ziska, L.H. 2003. Evaluation of the growth response of six invasive species to past, present and future atmospheric carbon dioxide. *Journal of Experimental Botany* 54(381): 395-404.
- Ziska, L.H., S. Faulkner, and J. Lydon. 2004. Changes in biomass and root:shoot ration of field-grown Canada thistle (*Cirsium arvense*), a noxious, invasive weed, with elevated CO₂: implications for control with glyphosate. *Weed Science* 52:584-588.

4.8 Index

A

Alternatives

- Alternative A, 21, 24, 80, 81, 85, 104, 130, 133, 135, 137, 142, 143, 144, 271, 272, 281, 323, 351, 368, 369, 378, 382, 397, 401, 402, 407, 409, 410, 411, 414, 416, 417
- Alternative B, 16, 21, 25, 78, 79, 80, 81, 85, 104, 108, 133, 134, 135, 137, 138, 139, 140, 143, 145, 146, 262, 270, 272, 273, 274, 275, 282, 293, 324, 349, 351, 370, 378, 383, 398, 399, 401, 411, 412, 413, 414, 415, 416, 417
- Alternative C, 77, 85, 104, 134, 138, 139, 145, 262, 274, 275, 293, 349, 351, 377, 382, 383, 399, 401, 415, 417, 421
- Alternative D, 22, 78, 85, 104, 108, 134, 139, 140, 262, 275, 293, 349, 351, 377, 382, 384, 399, 415, 416, 417, 421
- No Action, 21, 24, 80, 81, 85, 104, 130, 131, 132, 133, 136, 138, 141, 262, 270, 271, 272, 281, 318, 323, 324, 368, 369, 378, 382, 397, 398, 399, 401, 402, 407, 408, 411, 412, 421
- preferred alternative, 424
- proposed action, i, xiii, 8, 13, 15, 21, 25, 36, 70, 79, 81, 85, 89, 104, 131, 132, 133, 135, 143, 249, 272, 282, 294, 298, 324, 326, 327, 352, 353, 355, 356, 357, 370, 378, 382, 383, 384, 398, 399, 401, 421, 424
- amphibians, 59, 80, 156, 451, 454, 463
- aquatic species, 11, 17, 80, 83, 90, 129, 257, 298, 318, 321, 325, 341, 353, 355

C

- Clean Water Act, xii, 9, 19, 254, 255, 345, 437
- consultation, 64, 326, 424, 474
- critical habitat, 19, 299, 300, 302, 303, 304, 305, 308, 341, 343, 344, 346, 352, 353, 354, 362, 432, 478
- cultural resources, 13, 67, 375

E

- economics, xi, 432
- Endangered Species Act, xii, 9, 19, 36, 117, 124, 300, 308, 327, 332, 336, 338, 339, 340, 343, 355, 412, 432, 433, 437, 441, 442, 474
- Environmental Justice, 13, 410, 413, 415, 416, 432, 462
- erosion, 39, 69, 115, 249, 250, 253, 256, 257, 264, 265, 266, 267, 271, 273, 274, 275, 282, 286, 289, 323, 324, 325, 336, 341, 353
- Essential Fish Habitat, xii, 355, 356, 357, 474

H

habitat

- critical habitat, 19, 299, 300, 302, 303, 304, 305, 308, 341, 343, 344, 346, 352, 353, 354, 362, 432, 478

- Essential Fish Habitat, xii, 355, 356, 357, 474
- wildlife, 472, 479
- heritage resources, 13, 67, 375

I

- issues, 14, 15, 18, 19, 21, 22, 79, 100, 248, 251, 276, 324, 396, 416, 420, 422, 424, 445, 481

M

- Management Indicator Species, xii, 299, 440

N

- National Forest Management Act, xii, 9, 19, 437
- National Historic Preservation Act, 13, 417

P

- Proposed action, 81, 124, 248, 261, 297, 319, 324, 325, 338, 339, 340, 343, 349, 351, 352, 354, 355, 357, 401, 424
- public involvement, 14
- Purpose and need, 95, 135

R

- range allotments, 392, 393
- riparian, 11, 14, 16, 17, 18, 36, 38, 51, 52, 58, 59, 60, 77, 80, 82, 83, 86, 87, 100, 116, 134, 139, 254, 257, 262, 276, 282, 294, 300, 302, 304, 305, 317, 324, 325, 326, 327, 332, 339, 340, 347, 350, 352, 353, 354, 357, 358, 362, 363, 376, 377, 382, 383, 399, 400, 415, 420, 421, 440, 472
- Riparian Habitat Conservation Areas (RHCAs), xii, 52, 57, 72, 81, 138, 139, 248, 254, 256, 257, 259, 261, 262, 274, 275, 279, 281, 282, 283, 284, 286, 287, 290, 291, 293, 295, 297, 323, 330, 346, 347, 349, 352, 407
- road, 140, 266, 298, 362, 363

S

- sediment*, 17, 254, 255, 256, 257, 262, 275, 278, 281, 286, 289, 293, 311, 317, 319, 324, 325, 340, 341, 343, 344, 345, 347, 349, 350, 351, 352, 354, 356, 439
- sera*, xiii, 53, 55, 57, 58, 59, 60, 69, 92, 127, 133, 137, 248, 249, 267, 273, 277, 283, 284, 318, 319, 321, 322, 323, 326, 329, 330, 332, 338, 341, 350, 354, 398, 467, 468, 469
- soil*
 - compaction, 253, 325
 - erosion, 39, 69, 115, 249, 250, 253, 256, 257, 264, 265, 266, 267, 271, 273, 274, 275, 282, 286, 289, 323, 324, 325, 336, 341, 353
 - nutrients*, 17, 38, 109, 252, 253, 255, 324, 344, 346, 433, 444

SOLI, 60, 62, 63, 117, 440
stream flow, 254, 276, 280, 288, 310, 311, 337, 341

W

water quality, 1, 11, 14, 15, 17, 38, 58, 248, 254, 255,
256, 276, 278, 279, 281, 287, 300, 302, 303, 304,
305, 317, 337, 342, 344, 345, 352, 357, 362, 376,
415, 420, 421
watershed, 258, 259, 275, 282, 286, 292, 293, 294,
295, 333, 348, 353
wetlands, 53, 55, 57, 59, 71, 77, 262, 279, 280, 290,
291, 326, 344, 346, 352, 353, 356, 440, 442, 480

Wild and Scenic Rivers, 80, 81, 89, 115, 357, 360,
361, 366, 375, 378, 382, 404, 405, 408, 412, 431,
442
wildlife, 11, 16
amphibians, 66, 83, 428, 447
aquatic species, 11, 17, 80, 83, 90, 129, 257, 298,
318, 321, 325, 341, 353, 355
federally listed, 55, 60, 61, 62, 117, 118, 124, 249,
299, 319, 327, 328, 329, 343, 351, 352, 355, 433
Management Indicator Species, xii, 299, 440
sensitive, 12, 117, 320, 440