References and Included Studies (appearing in text and evidence tables)

- Macera CA, Jones DA, Yore MM, et al. Prevalence of physical activity, including lifestyle activities among adults--United States, 2000-2001. MMWR. 2003;52(32):764-8.
- Centers for Disease Control and Prevention (CDC). Prevalence of Physical activity, including lifestyle activities among adults - United States, 2000-2001. Morbidity and Mortality Weekly Report. 2003;52(32):764-9.
- Grunbaum JA, Kann L, Kinchen SA, et al. Youth Risk Behavior Surveillance - United States, 2001. Morbidity and Mortality Weekly Review. 2002;51(SS-4):1-66.
- Ries LAG, Eisner MP, Kosary CL, et al. SEER Cancer Statistics Review. National Cancer Institute, Bethesda, MD. Available at: http://seer.cancer.gov/ csr/1975 2000. Accessed December 22, 2003.
- McTiernan A, Ulrich C, Kumai C, et al. Anthropometric and hormone effects of an eight-week exercise-diet intervention in breast cancer patients: Results of a pilot study. Cancer Epidemiology, Biomarkers & Prevention. 1998;7(6):477-81.
- Friedenreich CM. Physical activity and cancer prevention: from observational to intervention research. Cancer Epidemiol Biomarkers Prev. 2001;10(4):287-301.
- Courneya KS, Friedenreich CM. Physical exercise and quality of life following cancer diagnosis: a literature review. Annals of Behavioral Medicine. 1999;21(2):171-9.
- Brown J. Nutrition and Physical Activity During and After Cancer Treatment: An American Cancer Society Guide for Informed Choices. CA Cancer J Clin. 2003;53:268-91.
- Casperson CJ, Powell KE, Christenson GM. Physical activity, exercise, and physical fitness: Definitions and distinctions for health-related research. Public Health Reports. 1985;100(2):126-31.
- National Coalition for Cancer Survivorship. Glossary. Available at: http://www.canceradvocacy.org/ resources/glossary.aspx#C. Accessed December 22, 2003.
- 11. ES [computer program]. Version 1.01. St. Paul: Assessment Systems Corporation; 1996.

- US Department of Health and Human Services. Physical activity and health: A report of the Surgeon General: National Center for Chronic Disease Prevention and Health Promotion; 1996.
- Courneya KS, Freidenreich CM. Framework PEACE: An organizational model for examining physical exercise across the cancer experience. Annals of Behavioral Medicine. 2001;23(4):263-72.
- Friedenreich CM. Physical activity and cancer prevention: from observational to intervention research. Cancer Epidemiology, Biomarkers & Prevention. 2001;10(4):287-301.
- World Health Organization, ed. Weight control and physical activity. Lyon: IARCPress; 2002. Vainio H, Bianchini F, eds. IARC Handbooks of Cancer Prevention; No. 6.
- Eden KB, Orleans CT, Mulrow CD, et al. Does counseling by clinicians improve physical activity? A summary of the evidence for the U.S. Preventive Services Task Force. Ann Intern Med. 2002;137(3):208-15.
- Kahn EB, Ramsey LT, Brownson RC, et al. The effectiveness of interventions to increase physical activity. A systematic review. Am J Prev Med. 2002;22(4 Suppl):73-107.
- US Preventive Services Task Force. Behavioral counseling in primary care to promote physical activity: recommendation and rationale. Ann Intern Med. 2002;137(3):205-7.
- Task Force on Community Preventive Services. Recommendations to increase physical activity in communities. Am J Prev Med. 2002;22(4 Suppl):67-72.
- Stone EJ, McKenzie TL, Welk GJ, et al. Effects of physical activity interventions in youth. Review and synthesis. Am J Prev Med. 1998;15(4):298-315.
- 21. King AC, Rejeski WJ, Buchner DM. Physical activity interventions targeting older adults. A critical review and recommendations. Am J Prev Med. 1998;15(4):316-33.
- 22. Taylor WC, Baranowski T, Young DR. Physical activity interventions in low-income, ethnic minority, and populations with disability. Am J Prev Med. 1998;15(4):334-43.
- Dishman RK, Oldenburg B, O'Neal H, et al. Worksite physical activity interventions. Am J Prev Med. 1998;15(4):344-61.

- Marcus BH, Owen N, Forsyth LH, et al. Physical activity interventions using mass media, print media, and information technology. Am J Prev Med. 1998;15(4):362-78.
- Sallis JF, Bauman A, Pratt M. Environmental and policy interventions to promote physical activity. Am J Prev Med. 1998;15(4):379-97.
- Dunn AL, Andersen RE, Jakicic JM. Lifestyle physical activity interventions. History, short- and long-term effects, and recommendations. Am J Prev Med. 1998;15(4):398-412.
- Simons-Morton DG, Calfas KJ, Oldenburg B, et al. Effects of interventions in health care settings on physical activity or cardiorespiratory fitness. Am J Prev Med. 1998;15(4):413-30.
- Ashenden R, Silagy C, Weller D. A systematic review of the effectiveness of promoting lifestyle change in general practice. Fam Pract. 1997;14(2):160-76.
- Baranowski T, Anderson C, Carmack C. Mediating variable framework in physical activity interventions. How are we doing? How might we do better? Am J Prev Med. 1998;15(4):266-97.
- Dishman RK, Buckworth J. Increasing physical activity: a quantitative synthesis. Med Sci Sports Exerc. 1996;28(6):706-19.
- Eaton CB, Menard LM. A systematic review of physical activity promotion in primary care office settings. Br J Sports Med. 1998;32(1):11-6.
- Hillsdon M, Thorogood M. A systematic review of physical activity promotion strategies. Br J Sports Med. 1996;30(2):84-9.
- Ketola E, Sipila R, Makela M. Effectiveness of individual lifestyle interventions in reducing cardiovascular disease and risk factors. Ann Med. 2000;32(4):239-51.
- Lawlor DA, Hanratty B. The effect of physical activity advice given in routine primary care consultations: a systematic review. J Public Health Med. 2001;23(3):219-26.
- Lewis BA, Marcus BH, Pate RR, et al. Psychosocial mediators of physical activity behavior among adults and children. Am J Prev Med. 2002;23(2 Suppl):26-35.
- Marshall SJ, Biddle SJ. The transtheoretical model of behavior change: a meta-analysis of applications to physical activity and exercise. Ann Behav Med. 2001;23(4):229-46.

- Petrella RJ, Lattanzio C. Does counseling help patients get active? Systematic review of the literature. Canadian Family Physician. 2002;48:72-80.
- Shepard R. Fitness and exercise programs: A review of methodology and health impact. American Journal of Health Promotion. 1996;10(6):436-52.
- Conn VS, Minor MA, Burks KJ, et al. Integrative review of physical activity intervention research with aging adults. Journal of the American Geriatrics Society. 2003;51:1159-68.
- Friedenreich CM, Courneya KS. Exercise as rehabilitation for cancer patients. Clinical Journal of Sport Medicine. 1996;6(4):237-44.
- Rock CL, Demark-Wahnefried W. Can lifestyle modification increase survival in women diagnosed with breast cancer? Journal of Nutrition. 2002;132(11 Suppl):3504S-7S.
- 42. Stone P. The measurement, causes and effective management of cancer-related fatigue. International Journal of Palliative Nursing. 2002;8(3):120-8.
- Chlebowski RT, Aiello E, McTiernan A. Weight loss in breast cancer patient management. Journal of Clinical Oncology. 2002;20(4):1128-43.
- 44. Briss PA, Zaza S, Pappaioanou M, et al. Developing an evidence-based Guide to Community Prevention Services--methods. The Task Force on Community Preventive Services. American Journal of Preventive Medicine. 2000;18(1 Suppl):35-43.
- 45. Shephard RJ. Worksite fitness and exercise programs: A review of methodology and health impact. American Journal of Health Promotion. 1996;10(6):436-52.
- 46. Zaza S, Wright-De Aguero L, Briss PA, et al. Task Force on Community Preventive Services. Data collection instrument and procledure for systematic reviews in the Guide to Community Preventive Services. Am J Prev Med. 2000;18(1S):44-74.
- Chen SC, Chen MF. Timing of shoulder exercise after modified radical mastectomy: a prospective study. Changgeng Yi Xue Za Zhi. 1999;22(1):37-43.
- Follmann D, Elliott P, Cutler J. Variance imputation for overviews of clinical trials with continuous response. J Clinical Epidemiology. 1992;45(7)769-773.
- Luepker RV, Murray DM, Jacobs DRJ, et al. Community education for cardiovascular disease prevention: risk factor changes in the Minnesota Heart Health Program. Am J Public Health. 1994;84(9):1383-93.

- Eaton CB, Lapane KL, Garber CE, et al. Effects of a community-based intervention on physical activity: the Pawtucket Heart Health Program. Am J Public Health. 1999;89(11):1741-4.
- 51. Bauer RL, Heller RF, Challah S. United Kingdom Heart Disease Prevention Project: 12-year follow-up of risk factors. Am J Epidemiol. 1985;121(4):563-9.
- 52. Nader PR, Stone EJ, Lytle LA, et al. Three-year maintenance of improved diet and physical activity: the CATCH cohort. Child and Adolescent Trial for Cardiovascular Health. Arch Pediatr Adolesc Med. 1999;153(7):695-704.
- Luepker RV, Perry CL, McKinlay SM, et al. Outcomes of a field trial to improve children's dietary patterns and physical activity. The Child and Adolescent Trial for Cardiovascular Health. CATCH collaborative group. JAMA. 1996;275(10):768-76.
- Goran MI, Poehlman ET. Endurance training does not enhance total energy expenditure in healthy elderly persons. Am J Physiol. 1992;263(5 Pt 1):E950-7.
- Cohen J. Statistical Power Analysis in the Behavioral Sciences. 2nd ed. Hillsdale, NY: Lawrence Erlbaum; 1992.
- 56. Howard JK, Bindler RM, Synoground G, et al. A cardiovascular risk reduction program for the classroom. J Sch Nurs. 1996;12(4):4-11.
- Bull FC, Jamrozik K, Blanksby BA. Tailored advice on exercise--does it make a difference? Am J Prev Med. 1999;16(3):230-9.
- Steptoe A, Doherty S, Rink E, et al. Behavioural counselling in general practice for the promotion of healthy behaviour among adults at increased risk of coronary heart disease: randomised trial. BMJ. 1999;319(7215):943-7; discussion 7-8.
- Halbert JA, Silagy CA, Finucane PM, et al. Physical activity and cardiovascular risk factors: effect of advice from an exercise specialist in Australian general practice. Med J Aust. 2000;173(2):84-7.
- Kerse NM, Flicker L, Jolley D, et al. Improving the health behaviours of elderly people: randomised controlled trial of a general practice education programme. Bmj. 1999;319(7211):683-7.
- Green BB, McAfee T, Hindmarsh M, et al. Effectiveness of telephone support in increasing physical activity levels in primary care patients. Am J Prev Med. 2002;22(3):177-83.

- Stevens W, Hillsdon M, Thorogood M, et al. Costeffectiveness of a primary care based physical activity intervention in 45-74 year old men and women: a randomised controlled trial. Br J Sports Med. 1998;32(3):236-41.
- Gillett P, White A, Caserta M. Effect of exercise and/or fitness education on fitness in older, sedentary, obese women. J Aging Phys Activity. 1996;4:42-55.
- Pereira MA, Kriska AM, Day RD, et al. A randomized walking trial in postmenopausal women: effects on physical activity and health 10 years later. Arch Intern Med. 1998;158(15):1695-701.
- Burke V, Milligan RA, Thompson C, et al. A controlled trial of health promotion programs in 11year-olds using physical activity "enrichment" for higher risk children. J Pediatr. 1998;132(5):840-8.
- 66. Dale D, Corbin CB. Physical activity participation of high school graduates following exposure to conceptual or traditional physical education. Res Q Exerc Sport. 2000;71(1):61-8.
- 67. O'Loughlin J, Renaud L, Paradis G, et al. Screening school personnel for cardiovascular disease risk factors: short-term impact on behavior and perceived role as promoters of heart health. Prev Med. 1996;25(6):660-7.
- Gemson DH, Sloan RP. Efficacy of computerized health risk appraisal as part of a periodic health examination at the worksite. Am J Health Promot. 1995;9(6):462-6.
- Lombard DN, Lombard TN, Winett RA. Walking to meet health guidelines: the effect of prompting frequency and prompt structure. Health Psychol. 1995;14(2):164-70.
- Mutrie N, Carney C, Blamey A, et al. "Walk in to Work Out": a randomised controlled trial of a self help intervention to promote active commuting. J Epidemiol Community Health. 2002;56(6):407-12.
- Linenger JM, Chesson CVn, Nice DS. Physical fitness gains following simple environmental change. Am J Prev Med. 1991;7(5):298-310.
- Perkio-Makela M. Influence of exercise-focused group activities on the physical activity, functional capacity, and work ability of female farmers--a threeyear follow-Up. Int J Occup Saf Ergon. 1999;5(3):381-94.
- Marcus BH, Bock BC, Pinto BM, et al. Efficacy of an individualized, motivationally-tailored physical activity intervention. Ann Behav Med. 1998;20(3):174-80.

- Belisle M, Roskies E, Levesque JM. Improving adherence to physical activity. Health Psychol. 1987;6(2):159-72.
- Steptoe A, Rink E, Kerry S. Psychosocial predictors of changes in physical activity in overweight sedentary adults following counseling in primary care. Prev Med. 2000;31(2 Pt 1):183-94.
- Blalock SJ, Currey SS, DeVellis RF, et al. Effects of educational materials concerning osteoporosis on women's knowledge, beliefs, and behavior. Am J Health Promot. 2000;14(3):161-9.
- Owen N, Lee C, Naccarella L, et al. Exercise by mail: A media-based behavior change program for aerobic exercise. J Sport Psychol. 1987;9:346-57.
- Miller YD, Trost SG, Brown WJ. Mediators of physical activity behavior change among women with young children. Am J Prev Med. 2002;23(2 Suppl):98-103.
- 79. Gillett PA, Caserta MS. Changes in aerobic power, body composition, and exercise adherence in obese post-menopausal women six months after exercise training. Menopause. 1996;3(3):126-32.
- Chalmers I, Adams M, Dickersin K, et al. A cohort study of summary reports of controlled trials. JAMA. 1990;263(10):1401-5.
- Djuric Z, DiLaura NM, Jenkins I, et al. Combining weight-loss counseling with the weight watchers plan for obese breast cancer survivors. Obesity Research. 2002;10(7):657-65.
- 82. Courneya KS, Friedenreich CM, Sela RA, et al. Correlates of adherence and contamination in a randomized controlled trial of exercise in cancer survivors: an application of the theory of planned behavior and the five factor model of personality. Annals of Behavioral Medicine. 2002;24(4):257-68.
- Courneya KS, Friedenreich CM, Sela RA, et al. The Group Psychotherapy and Home-Based Physical Exercise (Group-Hope) Trial in Cancer Survivors: Physical Fitness and Quality of Life Outcomes. Psycho-Oncology. 2003;12:357-74.
- Segar ML, Katch VL, Roth RS, et al. The effect of aerobic exercise on self-esteem and depressive and anxiety symptoms among breast cancer survivors. Oncology Nursing Forum. 1998;25(1):107-13.
- Cunningham BA, Morris G, Cheney CL, et al. Effects of resistive exercise on skeletal muscle in marrow transplant recipients receiving total parenteral nutrition. Journal of Parenteral & Enteral Nutrition. 1986;10(6):558-63.

- Burnham TR, Wilcox A. Effects of exercise on physiological and psychological variables in cancer survivors. Medicine & Science in Sports & Exercise. 2002;34(12):1863-7.
- Segal R, Evans W, Johnson D, et al. Structured exercise improves physical functioning in women with stages I and II breast cancer: results of a randomized controlled trial. Journal of Clinical Oncology. 2001;19(3):657-65.
- MacVicar MG, Winningham ML, Nickel JL. Effects of aerobic interval training on cancer patients' functional capacity. Nursing Research. 1989;38(6):348-51.
- Winningham ML, MacVicar MG. The effect of aerobic exercise on patient reports of nausea. Oncol Nurs Forum. 1988;15(4):447-50.
- Courneya KS, Mackey JR, Bell GJ, et al. Randomized controlled trial of exercise training in postmenopausal breast cancer survivors: cardiopulmonary and quality of life outcomes.[comment]. Journal of Clinical Oncology. 2003;21(9):1660-8.
- Segal RJ, Reid RD, Courneya KS, et al. Resistance exercise in men receiving androgen deprivation therapy for prostate cancer. Journal of Clinical Oncology. 2003;21(9):1653-9.
- 92. Berglund G. Starting Again--A comparison study of a group rehabilitation program for cancer patients. Acta Oncologica. 1993;32(1):15-21.
- Berglund G, Bolund C, Gustafsson UL, et al. Oneyear follow-up of the 'Starting Again' group rehabilitation programme for cancer patients. European Journal of Cancer. 1994;30A(12):1744-51.
- 94. Mock V, Dow KH, Meares CJ, et al. Effects of exercise on fatigue, physical functioning, and emotional distress during radiation therapy for breast cancer. Oncology Nursing Forum. 1997;24(6):991-1000.
- 95. Mock V, Pickett M, Ropka ME, et al. Fatigue and quality of life outcomes of exercise during cancer treatment. Cancer Practice. 2001;9(3):119-27.
- Pickett M, Mock V, Ropka ME, et al. Adherence to moderate-intensity exercise during breast cancer therapy. Cancer Practice. 2002;10(6):284-92.
- 97. Hillsdon M, Thorogood M, White I, et al. Advising people to take more exercise is ineffective: a randomized controlled trial of physical activity promotion in primary care. Int J Epidemiol. Aug 2002;31(4):808-15.

- Mock V, Ropka ME, Rhodes VA, et al. Establishing mechanisms to conduct multi-institutional researchfatigue in patients with cancer: an exercise intervention. Oncology Nursing Forum. 1998;25(8):1391-7.
- Mock V, Burke MB, Sheehan P, et al. A nursing rehabilitation program for women with breast cancer receiving adjuvant chemotherapy. Oncology Nursing Forum. 1994;21(5):899-907.
- 100. Fairey AS, Courneya KS, Field CJ, et al. Effects of Exercise Training on Fasting Insulin, Insulin Resistance, Insulin-like Growth Factors, and Binding Proteins in Postmenopausal Breast Cancer Survivors: A Randomized Controlled Trial. Cancer Epidemiology, Biomarkers & Prevention. 2003;12:721-7.
- 101. Dimeo F, Fetscher S, Lange W, et al. Effects of aerobic exercise on the physical performance and incidence of treatment-related complications after high-dose chemotherapy. Blood. 1997;90(9):3390-4.
- 102. Dimeo FC, Tilmann MH, Bertz H, et al. Aerobic exercise in the rehabilitation of cancer patients after high dose chemotherapy and autologous peripheral stem cell transplantation. Cancer. 1997;79(9):1717-22.
- 103. Nieman DC, Cook VD, Henson DA, et al. Moderate exercise training and natural killer cell cytotoxic activity in breast cancer patients. International Journal of Sports Medicine. 1995;16(5):334-7.
- 104. Hayes SC, D. R, Davies PS, et al. Immunological changes after cancer treatment and participation in an exercise program. Medicine & Science in Sports & Exercise. 2003;35(1):2-9.
- 105. Na YM, Kim MY, Kim YK, et al. Exercise therapy effect on natural killer cell cytotoxic activity in stomach cancer patients after curative surgery. Archives of Physical Medicine & Rehabilitation. 2000;81(6):777-9.
- 106. Dimeo FC, Stieglitz RD, Novelli-Fischer U, et al. Effects of physical activity on the fatigue and psychologic status of cancer patients during chemotherapy. Cancer. 1999;85(10):2273-7.
- 107. McKenzie DC, Kalda AL. Effect of upper extremity exercise on secondary lymphedema in breast cancer patients: A pilot study. Journal of Clinical Oncology. 2003;21(3):463-6.
- 108. Winningham ML, MacVicar MG, Bondoc M, et al. Effect of aerobic exercise on body weight and composition in patients with breast cancer on adjuvant chemotherapy. Oncology Nursing Forum. 1989;16(5):683-9.

- 109. Cohen LA, Boylan E, Epstein M, et al. Voluntary exercise and experimental mammary cancer. In: Jacobs MM, ed. Exercise, Fat, and Cancer. New York: Plenum Press; 1992.
- 110. Cohen LA, Choi K, Wang C-X. Influence of dietary fat, caloric restriction, and voluntary exercise on N-Nitrosomethylurea-induced mammary tumorigenesis in rats. Cancer Research. 1988;48:4276-83.
- Hoffman-Goetz L. Exercise, natural immunity, and tumor metastasis. Med Sci Sports Exerc. 1994;26:157-63.
- 112. MacVicar MG, Winningham ML. Promoting the functional capacity of cancer patients. The Cancer Bulletin. 1986;38(5):235-9.
- 113. Bull FC, Jamrozik K. Advice on exercise from a family physician can help sedentary patients to become active. Am J Prev Med. 1998;15(2):85-94.
- 114. Bock BC, Marcus BH, Pinto BM, et al. Maintenance of physical activity following an individualized motivationally tailored intervention. Ann Behav Med. 2001;23(2):79-87.
- 115. Caserta MS, Gillett PA. Older women's feelings about exercise and their adherence to an aerobic regimen over time. Gerontologist. 1998;38(5):602-9.
- 116. Godin G, Desharnais R, Jobin J, et al. The impact of physical fitness and health-age appraisal upon exercise intentions and behavior. J Behav Med. 1987;10(3):241-50.
- 117. Graham-Clarke P, Oldenburg B. The effectiveness of a general practice-based physical activity intervention on patient physical activity status. Behav Change. 1994;11:132-44.
- 118. Edmundson E, Parcel GS, Feldman HA, et al. The effects of the Child and Adolescent Trial for Cardiovascular Health upon psychosocial determinants of diet and physical activity behavior. Prev Med. 1996;25(4):442-54.
- 119. Perry CL, Sellers DE, Johnson C, et al. The Child and Adolescent Trial for Cardiovascular Health (CATCH): intervention, implementation, and feasibility for elementary schools in the United States. Health Educ Behav. 1997;24(6):716-35.
- Simons-Morton BG, McKenzie TJ, Stone E, et al. Physical activity in a multiethnic population of third graders in four states. Am J Public Health. 1997;87(1):45-50.
- 121. Stone EJ, Osganian SK, McKinlay SM, et al. Operational design and quality control in the CATCH multicenter Trial. Prev Med. 1996;25(4):384-99.

- 122. Nader PR, Sellers DE, Johnson CC, et al. The effect of adult participation in a school-based family intervention to improve Children's diet and physical activity: the Child and Adolescent Trial for Cardiovascular Health. Prev Med. 1996;25(4):455-64.
- 123. McKenzie TL, Stone EJ, Feldman HA, et al. Effects of the CATCH physical education intervention: teacher type and lesson location. Am J Prev Med. 2001;21(2):101-9.
- 124. McKenzie TL, Nader PR, Strikmiller PK, et al. School physical education: effect of the Child and Adolescent Trial for Cardiovascular Health. Prev Med. 1996;25(4):423-31.
- 125. McKenzie TL, Strikmiller PK, Stone EJ, et al. CATCH: physical activity process evaluation in a multicenter trial. Health Educ Q. 1994;Suppl 2:S73-89.
- 126. Hearn M. Involving Families in Cardiovascular Health Promotion: The CATCH Feasibility Study. Journal of Health Education. 1992;23(1):22-31.
- 127. McKenzie TL, Feldman H, Woods SE, et al. Children's activity levels and lesson context during third-grade physical education. Res Q Exerc Sport. 1995;66(3):184-93.
- 128. Nader PR, Sallis JF, Rupp J, et al. San Diego family health project: reaching families through the schools. J Sch Health. 1986;56(6):227-31.
- 129. Nader PR, Sallis JF, Patterson TL, et al. A family approach to cardiovascular risk reduction: results from the San Diego Family Health Project. Health Educ Q. 1989;16(2):229-44.
- 130. Hilton S, Doherty S, Kendrick SK, et al. Promotion of healthy behaviour among adults at increased risk of coronary heart disease in general practice: methodology and baseline data from the Change of Heart Study. Health Education Journal. 1999;58:3-16.
- 131. Steptoe A, Kerry S, Rink E, et al. The impact of behavioral counseling on stage of change in fat intake, physical activity, and cigarette smoking in adults at increased risk of coronary heart disease. Am J Public Health. 2001;91(2):265-9.
- 132. Halbert JA, Silagy CA, Finucane P, et al. Recruitment of older adults for a randomized, controlled trial of exercise advice in a general practice setting. J Am Geriatr Soc. 1999;47(4):477-81.
- Kreuter MW, Chheda SG, Bull FC. How does physician advice influence patient behavior? Evidence for a priming effect. Arch Fam Med. 2000;9(5):426-33.

- 134. Bull FC, Kreuter MW, Scharff DP. Effects of tailored, personalized and general health messages on physical activity. Patient Educ Couns. 1999;36(2):181-92.
- 135. Harland J, White M, Drinkwater C, et al. The Newcastle exercise project: a randomised controlled trial of methods to promote physical activity in primary care. Bmj. 1999;319(7213):828-32.
- 136. Eckstrom E, Hickam DH, Lessler DS, et al. Changing physician practice of physical activity counseling. J Gen Intern Med. 1999;14(6):376-8.
- Rose G. A proposed trial of heart disease prevention in industry. Trans Soc Occup Med. 1970;20(3):109-11.
- 138. Rose G, Heller RF, Pedoe HT, et al. Heart disease prevention project: a randomised controlled trial in industry. Br Med J. 1980;280(6216):747-51.
- 139. Gomel M, Oldenburg B, Simpson JM, et al. Work-site cardiovascular risk reduction: a randomized trial of health risk assessment, education, counseling, and incentives. Am J Public Health. 1993;83(9):1231-8.
- 140. Gomel MK, Oldenburg B, Simpson JM, et al. Composite cardiovascular risk outcomes of a worksite intervention trial. Am J Public Health. 1997;87(4):673-6.
- 141. Carlaw RW, Mittelmark MB, Bracht N, et al. Organization for a community cardiovascular health program: experiences from the Minnesota Heart Health Program. Health Education Quarterly. 1984;11(3):243-52.
- 142. Jacobs DR, Jr., Luepker RV, Mittelmark MB, et al. Community-wide prevention strategies: evaluation design of the Minnesota Heart Health Program. J Chronic Dis. 1986;39(10):775-88.
- Luepker RV, Jacobs DR, Gillum RF, et al. Population risk of cardiovascular disease: the Minnesota Heart Survey. J Chronic Dis. 1985;38(8):671-82.
- 144. Mittelmark MB, Luepker RV, Jacobs DR, et al. Community-wide prevention of cardiovascular disease: education strategies of the Minnesota Heart Health Program. Prev Med. 1986;15(1):1-17.
- 145. Chen AH, Sallis JF, Castro CM, et al. A home-based behavioral intervention to promote walking in sedentary ethnic minority women: project WALK. Womens Health. 1998;4(1):19-39.
- 146. Dale D, Corbin C, Cuddihy T. Can conceptual physical education promote physically active lifestyles? Ped Exerc Sci. 1998;10:97-109.

- 147. Edye BV, Mandryk JA, Frommer MS, et al. Evaluation of a worksite programme for the modification of cardiovascular risk factors. Med J Aust. 1989;150(10):574, 6-8, 81.
- 148. Elder JP, Williams SJ, Drew JA, et al. Longitudinal effects of preventive services on health behaviors among an elderly cohort. Am J Prev Med. 1995;11(6):354-9.
- 149. Elder JP, McGraw SA, Stone EJ, et al. CATCH: process evaluation of environmental factors and programs. Health Educ Q. 1994;Suppl 2:S107-27.
- 150. Keyserling TC, Samuel-Hodge CD, Ammerman AS, et al. A randomized trial of an intervention to improve self-care behaviors of African-American women with type 2 diabetes: impact on physical activity. Diabetes Care. 2002;25(9):1576-83.
- 151. Knutsen SF, Knutsen R. The Tromso Heart Study: family approach to intervention on CHD. Feasibility of risk factor reduction in high-risk persons--project description. Scand J Soc Med. 1989;17(1):109-19.
- 152. Knutsen SF, Knutsen R. Wives of coronary high-risk men--are they also at higher risk? The Tromso Heart Study. J Intern Med. 1990;228(4):333-7.
- 153. Knutsen SF, Knutsen R. The Tromso Survey: the Family Intervention study--the effect of intervention on some coronary risk factors and dietary habits, a 6year follow-up. Prev Med.1991;20(2):197-212.
- 154. Thelle D, Forde O, Try K, et al. The Tromso Heart Study: Methods and Main Results of the Crosssectional study. Acta Med Scand. 1976;200:107-18.
- 155. Kreuter MW, Strecher VJ. Do tailored behavior change messages enhance the effectiveness of health risk appraisal? Results from a randomized trial. Health Educ Res. 1996;11(1):97-105.
- 156. Lovibond SH, Birrell PC, Langeluddecke P. Changing coronary heart disease risk-factor status: the effects of three behavioral programs. J Behav Med. 1986;9(5):415-37.
- 157. MacKeen P, Rosenberg J, Slater J, et al. A 13-year follow-up of a coronary heart disease risk factor screening and exercise program for 40-to-59-year-old men: exercise habit maintenance and physiologic status. J Cardiopul Rehabil. 1985;5:510-23.
- 158. Remington RD, Taylor HL, Buskirk ER. A method for assessing volunteer bias and its application to a cardiovascular disease prevention progamme involving physical activity. J Epidemiol Community Health. 1978;32(4):250-5.

- 159. Taylor HL, Buskirk ER, Remington RD. Exercise in controlled trials of the prevention of coronary heart disease. Fed Proc. 1973;32(5):1623-7.
- 160. Ostwald SK. Changing employees' dietary and exercise practices: an experimental study in a small company. J Occup Med. 1989;31(2):90-7.
- 161. Owen N, Lee C, Sedgwick A. Exercise maintenance: developing self-management guidelines for community fitness courses. Aust J Sci Med Sport. 1987;1:8-12.
- 162. Kriska AM, Bayles C, Cauley JA, et al. A randomized exercise trial in older women: increased activity over two years and the factors associated with compliance. Med Sci Sports Exerc. 1986;18(5):557-62.
- 163. Sherman JB, Clark L, McEwen MM. Evaluation of a worksite wellness program: impact on exercise, weight, smoking, and stress. Public Health Nurs. 1989;6(3):114-9.
- 164. Smith BJ, Bauman AE, Bull FC, et al. Promoting physical activity in general practice: a controlled trial of written advice and information materials. Br J Sports Med. 2000;34(4):262-7.
- 165. Carleton RA, Lasater TM, Assaf A, et al. The Pawtucket Heart Health Program: I. An experiment in population-based disease prevention. R I Med J. 1987;70(12):533-8.
- 166. Carleton RA, Lasater TM, Assaf AR, et al. The Pawtucket Heart Health Program: community changes in cardiovascular risk factors and projected disease risk. Am J Public Health. 1995;85(6):777-85.
- 167. Marcus BH, Banspach SW, Lefebvre RC, et al. Using the stages of change model to increase the adoption of physical activity among community participants. Am J Health Promot. 1992;6(6):424-9.
- Levin S, Gans KM, Carleton RA. The Evolution of a Physical Activity Campaign. Fam Community Health. 1998;21(1):65-77.
- 169. McGraw SA, McKinlay SM, McClements L. Methods in Program Evaluation: The Process Evaluation System of the Pawtucket Heart Health Program. Evaluation Review. 1989;13:459-83.
- 170. Godin G, Jobin J, Bouillon J. Assessment of leisure time exercise behavior by self-report: A concurrent validity study. Can J Public Health. 1986;77:359-61.
- 171. Godin G, Shepard RJ. A simple method to assess exercise behavior in the community. Can J Appl Sport Sci. 1985;10:141-6.

- 172. American College of Sports Medicine. Guidelines for Exercise Testing and Prescription. 6th ed. Philadelphia, PA: Lea & Febiger; 2000.
- 173. Bhambhani Y, Singh M. The effects of three training intensities on VO2 max and VE/VO2 ratio. Respiration. 1985;47:120-8.
- 174. McGavin CR, Gupta SP, McHardy GJR. Twelveminute walking test for assessing disability in chronic bronchitis. BMJ. 1976;46(1):822-3.
- 175. Jette M, Sidney K, Landry F, et al. Blood pressure response to a progressive step test in normotensive males and females. Can J Appl Physiol. 1994;19:421-31.
- 176. Invergo JJ, Ball TE, Looney M. Relationship of pushups and absolute muscular endurance to bench press strengt. J Appl Sport Sci Res. 1991;5:121-5.
- 177. Baumgartner TA, Jackson AS. Measurement for Evaluation in Physical Education and Exercise Science. Dubuque, IA: W.C. Brown; 1995.
- 178. Sutherland HJ, Walker P, Till JE. The development of a method for determining oncology patient's emotional distress using linear analogue scales. Cancer Nursing. 1988;11:303-8.
- 179. Yellen BS, Cella DF, Webster J, et al. Measuring fatigue and other anemia-related symptoms with the Functional Assessment of Cancer Therapy (FACT) measurement system. J Pain Symptom Manage. 1997;13:63-74.
- Shacham S. A shortened version of the Profile of Mood Status States. J Pers Assess. 1983;47:305-6.
- 181. CIPS. International Skalen fur Psychiatrie. 2nd ed. Beltz: Weinheim; 1981.
- 182. McNair PM, Lorr M, Droppleman LF. Profile of Mood States Manual. San Diego, CA: Educational and Industrial Testing Service; 1971.
- 183. Piper BF, Dibble E, Gilliss N, et al. Revised Piper Fatigue Scale: psychometric evaluation in women with breast cancer. Oncol Nurs Forum. 1998;25:677-84.
- 184. Mock VL. Body image in women treated for breast cancer: Doctoral Dissertation, Catholic University of America, 1988. Dissertation Abstracts International. 48(12) B edition. (University Microfilms No.88-03894); 1988.
- 185. Mock VL. Body image in women treated for breast cancer. Nursing Research. 1993;42:153-7.

- Roid GH, Fitts WH. Tennessee Self-Concept Scale: Revised Manual. Los Angeles: Western Psychological Services; 1988.
- 187. Padilla GV, Presant MM, Grant G, et al. Quality of Life Index for patients with cancer. Rew Nurs Health. 1983;6:117-26.
- 188. Cella DF, Tulsky DS, Gray G, et al. The functional assessment of cancer therapy scale: Development and validation of the general measure. J Clin Oncol. 1993;11:570-9.
- Brady MJ, Cella DF, Mo F, et al. Reliability and validity of the Functional Assessment of Cancer Therapy-Breast Quality of Life instrument. J Clin Oncol. 1997;15:974-86.
- 190. Ware JE, Snow KK, Kosinski MK, et al. SF-36 Health Survey Manual and Interpretation Guide. Boston, MA: Nimrod Press; 1993.
- 191. Ware JE, Donald-Sherbourne C. The MOS 36-item short form health survey (SF-36): Conceptual framework and item selection. Med Care. 1992;30:473-83.
- 192. Mor V, Laliberte L, Morris JN, et al. The Karnofsky Performance Status Scale-An examination of its reliability and validity in a research setting. Cancer. 1984;53:2002-7.
- 193. Cella DF. Manual of the Functional Assessment of Chronic Illness Therapy (FACIT) Scales. Chicago, IL: Center on Outcomes Research and Education, Northwestern University; 1997.
- 194. Rosenberg M. Society and Adolescent Self-Image. Princeton, NJ: Princeton University Press; 1995.
- 195. Curbow B, Somerfield M. Use of the Rosenberg Esteem Scale with cancer patients. J Psychosoc Oncol. 1991;9:113-31.
- 196. Fordyce MW. A review of research on the happiness measures: A sixty second index of happiness and mental health. Soc Indic Res. 1988;20:355-81.
- 197. Diener E, Emmons RA, Larsen RJ, et al. The satisfaction with life scale. J Personality Assess. 1985;49:71-5.
- 198. Herth K. Abbreviated instrument to measure hope: Development and psychometric evaluation. J of Advanc Nurs. 1992;17:1251-9.
- 199. Barrett EAM. Development of an instrument to measure power as knowing participation in change. 1987, Unpublished manuscript.

- 200. Durin JVGA, Womersley J. Body fat assessed from total body density and its estimation from skinfold thickness: Measurements on 481 men and women aged from 16 to 72 years. Br J Nutr. 1974;32:77-97.
- 201. Grant A. Nutritional Assessment Guidelines. 2nd ed. Seattle, WA; 1979.
- 202. Barale K, Cheney C, Lenssen P, et al. Anthropometric measurement in a protective environment. J Am Diet Assoc. 1981;78:359-61.
- 203. Frisancho AR. New norms of upper limb fat and muscle areas for assessment of nutritional status. Am J Clin Nutr. 1981;34(11):2540-5.
- 204. Farncombe M, Daniels G, Cross L. Lymphedema: the seemingly forgotten complication. J Pain Symptom Manage. 1994;9:269-76.
- 205. Derogatis LR. SCL-90-R manual I. Administration, scoring and procedures manual for the revised version. Baltimore, MD: Clinical Psychometrics Research Unit, Johns Hopkins University School of Medicine; 1977.
- 206. Zigmond AS, Snaith RP. The hospital anxiety and depression scale. Acta Psychiat Scand. 1983;67:361-70.
- 207. Radloff LS. The use of the center or epidemiologic studies depression scale in adolescents and young adults. J Youth Adolesc. 1977;20:149-66.
- 208. Spielberger CD, Gorusch RL, Lushene RF. Manual for the State-Trait Anxiety Inventory. Palo Alto, CA: Consulting Psychologists Press; 1970.
- Greer S, Watson M. Mental adjustment to cancer: its measurement and prognostic importance. Cancer Surv. 1987;6:439-53.
- Derogatis LR. The psychosocial adjustment to illness scale (PAIS). Journal of psychosomatic research. 1986;30(1):77-91.
- Derogatis LR, Spencer PM. The Brief Symptom Inventory (BSI): Administration, scoring, and procedures manual. Baltimore, Clinical Psychometric Research, 1993.
- 212. Beck AT. Depression: Causes and treatment. Philadelphia: University of Pennsylvania Press; 1972.
- 213. Wall LM. Changes in hope and power in lung cancer patients who exercise. Nursing Science Quarterly. 2000;13(3):234-42.
- 214. Sharkey AM, Carey AB, Heise CT, et al. Cardiac rehabilitation after cancer therapy in children and young adults. American Journal of Cardiology. 1993;71(16):1488-90.

- 215. Schwartz AL. Fatigue mediates the effects of exercise on quality of life. Quality of Life Research. 1999;8(6):529-38.
- 216. Schwartz AL. Exercise and weight gain in breast cancer patients receiving chemotherapy. Cancer Practice. 2000;8(5):231-7.
- 217. Schwartz AL. Daily Fatigue Patterns and Effect of Exercise in Women with Breast Cancer. Cancer Practice. 2000;8(1):16-24.
- 218. Schwartz AL, Mori M, Gao R, et al. Exercise reduces daily fatigue in women with breast cancer receiving chemotherapy. Medicine & Science in Sports & Exercise. 2001;33(5):718-23.
- 219. Decker WA, Turner-McGlade J, Fehir KM. Psychosocial aspects and the physiological effects of a cardiopulmonary exercise program in patients undergoing bone marrow transplantation (BMT) for acute leukemia (AL). Transplantation Proceedings. 1989;21(1 Pt 3):3068-9.
- 220. Dimeo F, Bertz H, Finke J, et al. An aerobic exercise program for patients with haematological malignancies after bone marrow transplantation. Bone Marrow Transplantation. 1996;18(6):1157-60.
- 221. Young-McCaughan S, Mays MZ, Arzola SM, et al. Research and commentary: Change in exercise tolerance, activity and sleep patterns, and quality of life in patients with cancer participating in a structured exercise program. Oncology Nursing Forum. Online. 2003;30(3):441-54.
- 222. Durak EP, Lilly PC. The application of an exercise and wellness program for cancer patients: A preliminary outcomes report. Journal of Strength and Conditioning Research. 1998;12(1):3-6.
- 223. Kolden GG, Strauman TJ, Ward A, et al. A pilot study of group exercise training (GET) for women with primary breast cancer: feasibility and health benefits. Psycho-Oncology. 2002;11(5):447-56.
- 224. Dimeo F, Rumberger BG, Keul J. Aerobic exercise as therapy for cancer fatigue. Medicine & Science in Sports & Exercise. 1998;30(4):475-8.
- 225. Porock D, Kristjanson LJ, Tinnelly K, et al. An exercise intervention for advanced cancer patients experiencing fatigue: a pilot study. Journal of Palliative Care. 2000;16(3):30-6.
- 226. Peters C, Lotzerich H, Niemeier B, et al. Influence of a moderate exercise training on natural killer cytotoxicity and personality traits in cancer patients. Anticancer Research. 1994;14:1033-6.

- 227. Ahmed RL, Schmitz K, Yee D. Effects of resistance training on lymphedema symptoms in recent breast cancer survivors. Paper presented at: Proceedings of the American Association for Cancer Research, 2003.
- 228. Harris SR, Niesen-Vertommen SL. Challenging the myth of exercise-induced lymphedema following breast cancer: A series of case reports. J Surg Oncol. 2000;74:95-9.
- 229. Petrek JA, Senie RT, Peters M, et al. Lymphedema in a cohort of breast carcinoma survivors 20 years after diagnosis. Cancer. 2001;92:1368-77.
- 230. Cox CE, Bass SS, McCann CR, et al. Lymphatic mapping and sentinel lymph node biopsy in patients with breast cancer. Ann Rev Med. 2000;51:525-42.

- 231. Jemal A, Murray T, Samuels A, et al. Cancer statistics 2003. CA Cancer J Clin. 2002;53:5-26.
- 232. Keyserling TC, Ammerman AS, Samual-Hodge CD, Ingram AF, Skelly AH, Elasy TA, Johnston LF, Cole AS, Henrequez-Roldan, CF. A diabetes management program for African American women with type 2 diabetes. The Diabetes Educator. 2000;26(5):796-805.
- 233. Mayer JA, Jermanovich A, Wright BL, et al. Changes in health behaviors of older adults: the San Diego Medicare Preventive Health Project. Prev Med. Mar 1994;23(2):127-33.