SOLVD ALTERATION IN STUDY DRUG DOSAGE FORM

RAND ID:	FORM: S D C VISIT:
	SEQUENCE NUMBER:
INSTRUCTIONS:	This form is to be used whenever a dosage change is needed. This form is to be used between SOLVD visits. The visit number entered should be the last SOLVD visit attended by the participant. The sequence number is needed to indicate the number of times this form has been used between any two visits. Sequence number should start with 01 the first time the form is used for the participant for a specific visit number. Print clearly when entering a response in the appropriate boxes. For multiple choice questions, circle the one appropriate letter corresponding to the response chosen. Specific instructions for various questions are enclosed in boxes directly below the question. See the SOLVD General Instructions for Completing Forms for details.
. IDENTIFYING I	
. Today's Date	Month Day Year
2.2. First Maco	
2.3. Hiddle Nam	
B. INITIALS OF COMPLETING T	PERSON HIS FORM

C. FORMER STUDY MEDICATION (Medication the participant is taking up until the use of this form.)	D. CURRENT (NEW) MEDICATION
4. Pills dispensed/returned:	5. Pills dispensed today:
Instructions: Enter the following information for each type dispensed either at the last SOLVD visit or 1 use of this form: 8 pills dispensed, dose (Q=QD=on B=BID=twice daily), pills returned and 8 days since pills were dispensed.	dispensed today, enter the \$ pills dispensed and
# Pills (Circle: # Pills Pill previously Q=QD or returned type dispensed B=BID) today	# days # Pills (Circle: since dispensed Q=QD or last visit today B=BID)
2.5 eq b) 0 c)	a) a) B
5.0 eg f) g)	b) c) d) Q B
10.0 mg (i) (i) (k)	1) e) f) Q 10.0. eq B
SOLVD ALTERATION IN STUDY DRUG DOSAGE FOR	H (screen 3 of 4) (SDC page 2 of 3)
6. Type of change in dosageIncrease I Decrease D	F. REASON FOR DECREASING DOSE B.1. Side effects?
If Decrease (D), go to section F. REASON FOR DECREASING DOSE, Question 8.1.	No N If No, go to Question 9. on page 3. If Yes, indicate the
	following side effects: Yes No
E. REASON FOR INCREASING DOSE 7.1. Increase toward prescribed maintenance dose following dose reduction	8.2. Symptomatic hypotension Y N 8.3. Taste abnormalities Y N
7.2. Increase toward prescribed saintenance dose by protocol Y	8.4. Skin rash Y N
7.3. Other Y	B.S. Azotemia Y N
If No (Other), EXIT THE FORM.	8.6. Other Y N
If Yes (Other), specify:	If No (Other), go to Question 9. on page 3.
	If Yes (Other), specify:
EXIT THE FORM.	

SOLVO ALTERATION IN STUDY DRUG DOSAGE FORM (screen 4 of 4) (SDC page 3 of 3)

9. Pyocardial Infarction	Yes	Y		Yes	No
	No	N	11. Cardiac transplant	Y	N
			12. Moncardiac surgery	Y	N
10.1. Cardiac surgery other than transplant		Y	13. Worsening CHF with need for treatment with "open label" medication identical or		
	No	N	similar to the study drug	¥	N
If No, go to Question 11. 10.2. If hes (cardiac surgery), spec	ıfy:		14. Requested by the referring physician	Y	N
			15. Requested by participant	Y	N
			16. Other	¥	*
			If No (Other), EXIT THE FORM.		
			If Yes (Other), specify:		