395	*metrizamide, The Sanofi-Aventis Group.
396	[†] disulfiram, Odyssey Pharmaceuticals, Inc.
397	
398	Medication Guide
399	Antidepressant Medicines, Depression and Other Serious Mental Illnesses, and Suicidal
400	Thoughts or Actions
401	PARNATE [®] (PAR-nate) (tranylcypromine sulfate) Tablets
402	
403	Read the Medication Guide that comes with you or your family member's antidepressant
404	medicine. This Medication Guide is only about the risk of suicidal thoughts and actions with
405	antidepressant medicines. Talk to your, or your family member's, healthcare provider
406	about:
407	• All risks and benefits of treatment with antidepressant medicines
408	• All treatment choices for depression or other serious mental illness
409	
410	What is the most important information I should know about antidepressant medicines,
411	depression and other serious mental illnesses, and suicidal thoughts or actions?
412	· · · · · · · · · · · · · · · · · · ·
413	1. Antidepressant medicines may increase suicidal thoughts or actions in some children,
414	teenagers, and young adults within the first few months of treatment.
415	2. Depression and other serious mental illnesses are the most important causes of suicidal
416	thoughts and actions. Some people may have a particularly high risk of having suicidal
417	thoughts or actions. These include people who have (or have a family history of) bipolar
418	illness (also called manic-depressive illness) or suicidal thoughts or actions.
419	3. How can I watch for and try to prevent suicidal thoughts and actions in myself or a
420	family member?
421	• Pay close attention to any changes, especially sudden changes, in mood, behaviors,
422	thoughts, or feelings. This is very important when an antidepressant medicine is started or
423	when the dose is changed.
424	• Call the healthcare provider right away to report new or sudden changes in mood,
425	behavior, thoughts, or feelings.
426	• Keep all follow-up visits with the healthcare provider as scheduled. Call the healthcare
427	provider between visits as needed, especially if you have concerns about symptoms.
428	
429	Call a healthcare provider right away if you or your family member has any of the
430	following symptoms, especially if they are new, worse, or worry you:
431	• Thoughts about suicide or dying
432	Attempts to commit suicide
433	New or worse depression
434	• New or worse anxiety

435	٠	Feeling very agitated or restless	
436	٠	Panic attacks	
437	•	Trouble sleeping (insomnia)	
438	•	New or worse irritability	
439	•	Acting aggressive, being angry, or violent	
440	•	Acting on dangerous impulses	
441	•	An extreme increase in activity and talking (mania)	
442	•	Other unusual changes in behavior or mood	
443			
444	W	Vhat else do I need to know about antidepressant medicines?	
445	•	Never stop an antidepressant medicine without first talking to a healthcare	provider.
446		Stopping an antidepressant medicine suddenly can cause other symptoms.	
447	•	Antidepressants are medicines used to treat depression and other illnesses.	lt is
448		important to discuss all the risks of treating depression and also the risks of not t	reating it.
449		Patients and their families or other caregivers should discuss all treatment choice	es with the
450		healthcare provider, not just the use of antidepressants.	
451	٠	Antidepressant medicines have other side effects. Talk to the healthcare provi	der about the
452		side effects of the medicine prescribed for you or your family member.	
453	٠	Antidepressant medicines can interact with other medicines. Know all of the	medicines
454		that you or your family member takes. Keep a list of all medicines to show the h	ealthcare
455		provider. Do not start new medicines without first checking with your healthcare	provider.
456	٠	Not all antidepressant medicines prescribed for children are FDA approved	for use in
457		children. Talk to your child's healthcare provider for more information.	
458			
459	Th	his Medication Guide has been approved by the U.S. Food and Drug Administration	on for all
460	an	ntidepressants.	
461			
462	M	Iay 2007 Pl	RT:2MG
463			
	Q	gsk GlaxoSmithKline	
464		Glaxosmithkline	
165	CI		
465		BlaxoSmithKline	
466 467		Research Triangle Park, NC 27709 2007, GlaxoSmithKline. All rights reserved.	
468	01	2007, Glaxoomanikine. 7 in rights reserved.	
	٦	4 av. 2007	DDT.71DI
469	IVI	1ay 2007	PRT:71PI
470			