



APR 15 1993

Principal Display Panel-carton

**Infants' Motrin
CONCENTRATED**

Drops

For ages 6 months to 23 months

IBUPROFEN ORAL SUSPENSION

50 mg per 1.25 mL (dropperful)

Fever Reducer/Pain Reliever

Lasts up to 8 hours

Use Only With Enclosed Dropper

See New Label

BERRY-FLAVORED DROPS

Alcohol Free

1/2 FL OZ (15 mL)

Back Panel

Important: Read all product information before using. Keep this box for important information. This product is intended for use in children ages 6 months to 23 months of age.

Drug Facts

Active ingredient (in each 1.25 mL)

Purposes

Ibuprofen 50 mg.....Fever reducer/pain reliever

Uses temporarily:

- reduces fever
- relieves minor aches and pains due to the common cold, flu, sore throat, headaches and toothaches

Drug Facts (continued)

Warnings

Allergy alert: ibuprofen may cause a severe allergic reaction which may include:

- hives
- facial swelling
- asthma (wheezing)
- shock

Sore throat warning: severe or persistent sore throat or sore throat accompanied by high fever, headache, nausea, and vomiting may be serious. Consult doctor promptly. Do not use more than 2 days or administer to children under 3 years of age unless directed by doctor.

Do not use

- if the child has ever had an allergic reaction to any pain reliever/fever reducer

Ask a doctor before use if the child has

- not been drinking fluids
- lost a lot of fluid due to continued vomiting or diarrhea
- stomach pain
- problems or serious side effects from taking fever reducers or pain relievers

Ask a doctor or pharmacist before use if the child is

- under a doctor's care for any serious condition
- taking any other drug
- taking any other product that contains ibuprofen, or any other pain reliever/fever reducer

When using this product give with food or milk if stomach upset occurs

Stop use and ask a doctor if

- an allergic reaction occurs. Seek medical help right away.
- fever or pain gets worse or lasts more than 3 days
- the child does not get any relief within first day (24 hours) of treatment
- stomach pain or upset gets worse or lasts
- redness or swelling is present in the painful area
- any new symptoms appear

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- do not give more than directed
- shake well before using
- find the right dose on chart below. If possible, use weight to dose; otherwise use age.
- use only with enclosed dropper. Fill to dose level. Do not use any other dosing device.
- dispense liquid slowly into the child's mouth, toward inner cheek
- replace original bottle cap to maintain child resistance
- if needed, repeat dose every 6-8 hours
- do not use more than 4 times a day

Dosing Chart

| Weight (lb) | Age (mos) | Dose (mL) |
|-------------|-----------|----------------------------|
| Under 6 mos | | call a doctor |
| 12 - 17 lbs | 6-11 mos | 1 dropperful (1.25 mL) |
| 18 - 23 lbs | 12-23 mos | 1 ½ dropperfuls (1.875 mL) |

Attention: Specifically designed for use with enclosed dropper. Use only enclosed dropper to dose this product. Do not use any other dosing device.

Other information store at 20-25°C (68-77°F)

Inactive ingredients artificial flavors, citric acid, corn starch, FD&C Red #40, glycerin, polysorbate 80, purified water, sodium benzoate, sorbitol, sucrose, xanthan gum

Questions or Comments?

Call toll-free 1-800-962-5357

Or ask your Pharmacist, Doctor or Health Care Professional
See bottom of box for lot number and expiration date

McNeil Consumer Healthcare
DIVISION OF McNeil-PPC, Inc.
FORT WASHINGTON, PA 19034 USA
U.S. Patent No. 5,374,659